

X-ray crystal structure of *trans*-bis(pyridin-3-yl)-ethylene: comparing the supramolecular structural features among the symmetrical bis(*n*-pyridyl)-ethylenes (*n* = 2, 3, or 4) constitutional isomers

Jay Quentin,^a Eric W. Reinheimer^b and Leonard R. MacGillivray^{a*}

Received 13 October 2020

Accepted 16 November 2020

Edited by G. Díaz de Delgado, Universidad de Los Andes, Venezuela

Keywords: crystal structure; bis(pyridin-3-yl)ethylene; olefin.**CCDC reference:** 1985201**Supporting information:** this article has supporting information at journals.iucr.org/e^aDepartment of Chemistry, University of Iowa, 305 Chemistry Building, Iowa City, IA 52242-1290, USA, and ^bRigaku Oxford Diffraction, 9009 New Trails Dr., The Woodlands, TX 77381, USA. *Correspondence e-mail: len-macgillivray@uiowa.edu

The molecular structure of *trans*-bis(pyridin-3-yl)ethylene (**3,3'-bpe**), C₁₂H₁₀N₂, as determined by single-crystal X-ray diffraction is reported. The molecule self-assembles into two dimensional arrays by a combination of C—H···N hydrogen bonds and edge-to-face C—H··· π interactions that stack in a herringbone arrangement perpendicular to the crystallographic *c*-axis. The supramolecular forces that direct the packing of **3,3'-bpe** as well as its packing assembly within the crystal are also compared to those observed within the structures of the other symmetrical isomers *trans*-1,2-bis(*n*-pyridyl)ethylene (***n,n'*-bpe**, where *n* = *n'* = 2 or 4).

1. Chemical context

Bis(pyridyl)ethylenes have arisen as somewhat of a natural extension of cinnamic acid as a series of molecules capable of undergoing [2+2] photodimerization in the solid state to generate cyclobutanes. Foundational work by Schmidt and coworkers on *trans*-cinnamic acids led to the formation of the 'Topochemical Postulate', which dictated that olefins within 4.2 Å of one another are capable of undergoing the photodimerization process. Unlike cinnamic acid, which crystallizes in such a way that the olefins are rendered photoactive (olefins within 4.2 Å of one another), the native crystalline forms of bis(pyridyl)-ethylenes are photostable (olefins separated by distances > 4.2 Å in the crystal). To achieve photoreactivity of these olefins, it often becomes necessary to use a 'molecular template' that can interact with the olefin-containing bi-pyridine *via* supramolecular interactions such as hydrogen bonding, halogen bonding, argento- and aurophilic interactions, and dative N→B interactions. Analyses of the crystal structures of symmetric bis(pyridyl)ethylenes derivatives such as the *trans*-bis(*n*-pyridyl)ethylenes series of isomers (*n* = 2, 3 or 4) is necessary to understand the forces that govern their crystallization, why they are photostable, and why use templates to achieve photoreactivity (Campillo-Alvarado *et al.*, 2019; Chanthapally *et al.*, 2014; MacGillivray *et al.*, 2008; Pahari *et al.*, 2019; Sezer *et al.*, 2017; Volodin *et al.*, 2018).





Table 1

Structural features of the n,n' -bpe series of constitutional isomers.

The twist angle is defined as the angle between the plane defined by the four alkene atoms and the plane defined by either pyridine ring.

Compound	2,2'-bpe	3,3'-bpe	4,4'-bpe
Twist angle φ ($^\circ$)	7.43	5.17	9.14
Solid-state packing assembly	corrugated chains	approximately planar sheets	planar sheets
Assembly forces	edge-to-face C—H $\cdots$ π	edge-to-face C—H $\cdots$ π , C—H $\cdots$ N	C—H $\cdots$ N, face-to-face π — π
Nearest-neighbor alkene separation ($\text{\AA}$)	6.09	5.50	5.72

2. Structural commentary

The alkene **3,3'-bpe** crystallizes in the centrosymmetric monoclinic space group $P2_1/n$ (Fig. 1). The asymmetric unit consists of one-half molecule of **3,3'-bpe** with the C=C bond sitting on a crystallographic center of inversion. The pyridyl rings adopt an *anti*-conformation with respect to each other (Fig. 1).

3. Supramolecular features

Adjacent **3,3'-bpe** molecules interact primarily *via* edge-to-face C—H $\cdots$ π [$d(\text{C6} \cdots \text{pyr})$ 3.58 $\text{\AA}$; $\angle(\text{C6—H6} \cdots \text{pyr})$ 131.8 $^\circ$] forces between pyridyl rings (Fig. 2). Those rings also participate in C—H $\cdots$ N [$d(\text{C4} \cdots \text{N1})$ 3.59 $\text{\AA}$; $\angle(\text{C4—H4} \cdots \text{N1})$ 139.5 $^\circ$] hydrogen bonds (Fig. 2). The forces generate nearly




Figure 1

Single crystal structure for *trans*-bis(pyridin-3-yl)ethylene (**3,3'-bpe**) with anisotropic displacement ellipsoids at 50% probability.




Figure 2

C—H $\cdots$ N and edge-to-face C—H $\cdots$ π intermolecular interactions (both yellow dotted lines) highlighting nearest-neighbor alkene separations (red dashed arrow) (view along *a*).

planar sheets (Fig. 3), which aggregate into a herringbone arrangement of adjacent sheets (Fig. 4). Nearest-neighbor alkene C=C bonds of **3,3'-bpe** between adjacent sheets reveals a parallel, but offset orientation of the neighboring alkenes relative to one another at a distance of 5.50 $\text{\AA}$. The distance exceeds the inter-alkene separation of Schmidt for photodimerization and suggests that **3,3'-bpe** is photostable (Schmidt, 1971).

4. Database survey

For the n,n' -bpe (where: $n = n' = 2, 3$, or 4) series of symmetric alkenes, all three adopt nearly planar conformations (Table 1), with the pyridyl rings of **3,3'-bpe** and **2,2'-bpe** adopting *anti*-conformations with respect to each other. The packings of the symmetric alkenes are defined by combinations of C—H $\cdots$ π and/or C—H $\cdots$ N hydrogen bonds (Table 1) to form either one-dimensional chain (**2,2'-bpe**, Fig. 5) or two-dimensional sheet (**3,3'-bpe** and **4,4'-bpe**) structures (Fig. 6). Similar to **3,3'-bpe**, the alkene C=C bonds of **2,2'-bpe** (6.09 $\text{\AA}$; Vansant *et al.*,



Figure 3

Edge-on view of sheets encompassing neighboring molecules of **3,3'-bpe** supported by C—H $\cdots$ N and C—H $\cdots$ π intermolecular interactions.




Figure 4

Herringbone arrangement of neighboring sheets of **3,3'-bpe** molecules.




Figure 6
Planar, two-dimensional sheets of **4,4'-bpe**.




Figure 5
Corrugated, one-dimensional chains of **2,2'-bpe**.

1980) and **4,4'-bpe** (5.72 Å; Tinnemans *et al.*, 2018) (Table 1) are beyond the separation distance of Schmidt (1971).

5. Synthesis and crystallization

The alkene **3,3'-bpe** was prepared as described (Quentin *et al.*, 2020; Gordillo *et al.*, 2007, 2013) via a one-pot, aqueous Pd-catalyzed Hiyama–Heck cross-coupling between 3-bromopyridine and triethoxyvinylsilane (2:1 molar ratio) (Fig. 7). Flash chromatography (SiO₂, 10% MeOH/CH₂Cl₂) furnished **3,3'-bpe** as yellow crystals: 222.3 mg (23%). A portion of **3,3'-bpe** was dissolved in CHCl₃ and allowed to slowly evaporate at room temperature. Single crystals in the form of colorless plates suitable for single crystal X-ray diffraction formed within seven days.




Figure 7
Synthesis of **3,3'-bpe** via Pd-catalyzed Hiyama–Heck cross-coupling.

Table 2
Experimental details.

Crystal data	
Chemical formula	C ₁₂ H ₁₀ N ₂
<i>M</i> _r	182.22
Crystal system, space group	Monoclinic, <i>P</i> 2 ₁ / <i>n</i>
Temperature (K)	296
<i>a</i> , <i>b</i> , <i>c</i> (Å)	7.4591 (7), 5.5045 (6), 11.7803 (12)
β (°)	99.638 (5)
<i>V</i> (Å ³)	476.86 (8)
<i>Z</i>	2
Radiation type	Mo <i>K</i> α
μ (mm ^{−1})	0.08
Crystal size (mm)	0.18 × 0.12 × 0.06
Data collection	
Diffractometer	Bruker Nonius KappaCCD
Absorption correction	Multi-scan (<i>SADABS</i> ; Krause <i>et al.</i> , 2015)
<i>T</i> _{min} , <i>T</i> _{max}	0.989, 0.995
No. of measured, independent and observed [<i>I</i> > 2σ(<i>I</i>)] reflections	2410, 836, 587
<i>R</i> _{int}	0.034
Refinement	
<i>R</i> [<i>F</i> ² > 2σ(<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.050, 0.137, 1.07
No. of reflections	836
No. of parameters	84
H-atom treatment	All H-atom parameters refined
$\Delta\rho_{\text{max}}$, $\Delta\rho_{\text{min}}$ (e Å ^{−3})	0.13, −0.16

Computer programs: *COLLECT* (Nonius, 1988), *HKL DENZO* and *SCALEPACK* (Otwinowski & Minor, 1997), *SHELXT* (Sheldrick, 2015a), *SHELXL* (Sheldrick, 2015b) and *OLEX2* (Dolomanov *et al.*, 2009).

6. Refinement

Crystal data, data collection and structure refinement details for **3,3'-bpe** are summarized in Table 2. All non-hydrogen atoms were refined anisotropically. Hydrogen atoms were located in the difference-Fourier map and freely refined with 0.93 < C–H < 0.99 Å. Refinement of the hydrogen atoms led to a data-to-parameter ratio of ~10. The single-crystal data were collected at room temperature to best reflect conditions under which photochemical reactions are typically conducted. Room-temperature data can also lead to fewer reflections and/or scaling anomalies.

Funding information

Funding for this research was provided by: National Science Foundation (grant No. DMR-1708673 to L. R. MacGillivray).

References

- Campillo-Alvarado, G., Li, C., Swenson, D. C. & MacGillivray, L. R. (2019). *Cryst. Growth Des.* **19**, 2511–2518.
- Chanthapally, A., Oh, W. T. & Vittal, J. J. (2014). *Chem. Commun.* **50**, 451–453.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Gordillo, A., de Jesús, E. & López-Mardomingo, C. (2007). *Chem. Commun.* 4056–4058.
- Gordillo, A., Ortuño, M. A., López-Mardomingo, C., Lledós, A., Ujaque, G. & de Jesús, E. (2013). *J. Am. Chem. Soc.* **135**, 13749–13763.
- Krause, L., Herbst-Irmer, R., Sheldrick, G. M. & Stalke, D. (2015). *J. Appl. Cryst.* **48**, 3–10.

- MacGillivray, L. R., Papaefstathiou, G. S., Friščić, T., Hamilton, T. D., Bučar, D.-K., Chu, Q., Varshney, D. B. & Georgiev, I. G. (2008). *Acc. Chem. Res.* **41**, 280–291.
- Nonius (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Pahari, G., Bhattacharya, B., Reddy, C. M. & Ghoshal, D. (2019). *Chem. Commun.* **55**, 12515–12518.
- Quentin, J. & MacGillivray, L. R. (2020). *ChemPhysChem*, **21**, 154–163.
- Schmidt, G. M. J. (1971). *Pure Appl. Chem.* **27**, 647–678.
- Sezer, G. G., Yeşilel, O. Z. & Büyükgüngör, O. (2017). *J. Mol. Struct.* **1137**, 562–568.
- Sheldrick, G. M. (2015a). *Acta Cryst.* **A71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst.* **C71**, 3–8.
- Tinnemans, P. & Brugman, S. (2018). Private communication (deposition number CCDC 1843770. CCDC, Cambridge, England. <https://doi.org/10.5517/ccdc.csd.cc1zwlxg>).
- Vansant, J., Smets, G., Declercq, J. P., Germain, G. & Van Meerssche, M. (1980). *J. Org. Chem.* **45**, 1557–1565.
- Volodin, A. D., Korlyukov, A. A., Zorina-Tikhonova, E. N., Chistyakov, A. S., Sidorov, A. A., Eremenko, I. L. & Vologzhanina, A. V. (2018). *Chem. Commun.* **54**, 13861–13864.

supporting information

Acta Cryst. (2020). E76, 1859-1862 [https://doi.org/10.1107/S2056989020015303]

X-ray crystal structure of *trans*-bis(pyridin-3-yl)ethylene: comparing the supramolecular structural features among the symmetrical bis(*n*-pyridyl)ethylenes (*n* = 2, 3, or 4) constitutional isomers

Jay Quentin, Eric W. Reinheimer and Leonard R. MacGillivray

Computing details

Data collection: *HKL SCALEPACK* (Otwinowski & Minor, 1997); cell refinement: *COLLECT* (Nonius, 1998); data reduction: *HKL DENZO* and *SCALEPACK* (Otwinowski & Minor, 1997); program(s) used to solve structure: *ShelXT* (Sheldrick, 2015a); program(s) used to refine structure: *SHELXL* (Sheldrick, 2015b); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *OLEX2* (Dolomanov *et al.*, 2009).

trans-1,2-Bis(pyridin-3-yl)ethene

Crystal data

C₁₂H₁₀N₂
 $M_r = 182.22$
 Monoclinic, *P*2₁/*n*
 $a = 7.4591$ (7) Å
 $b = 5.5045$ (6) Å
 $c = 11.7803$ (12) Å
 $\beta = 99.638$ (5)°
 $V = 476.86$ (8) Å³
 $Z = 2$

$F(000) = 192$
 $D_x = 1.269$ Mg m⁻³
 Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 1169 reflections
 $\theta = 1.0$ – 26.7°
 $\mu = 0.08$ mm⁻¹
 $T = 296$ K
 Plate, colourless
 $0.18 \times 0.12 \times 0.06$ mm

Data collection

Bruker Nonius KappaCCD
 diffractometer
 Radiation source: fine-focus sealed tube
 CCD phi and ω scans
 Absorption correction: multi-scan
 (SADABS; Krause *et al.*, 2015)
 $T_{\min} = 0.989$, $T_{\max} = 0.995$
 2410 measured reflections

836 independent reflections
 587 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 3.0^\circ$
 $h = -8 \rightarrow 8$
 $k = -6 \rightarrow 6$
 $l = -13 \rightarrow 13$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.137$
 $S = 1.07$
 836 reflections
 84 parameters
 0 restraints

Primary atom site location: dual
 Hydrogen site location: difference Fourier map
 All H-atom parameters refined
 $w = 1/[\sigma^2(F_o^2) + (0.0703P)^2 + 0.056P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.13$ e Å⁻³
 $\Delta\rho_{\min} = -0.16$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.5400 (2)	0.7577 (3)	0.63093 (15)	0.0609 (6)
C2	0.2479 (2)	0.5639 (3)	0.57302 (15)	0.0459 (5)
C3	0.3752 (3)	0.7464 (4)	0.56601 (17)	0.0537 (6)
C6	0.2998 (3)	0.3821 (4)	0.65272 (17)	0.0529 (6)
C4	0.5835 (3)	0.5788 (4)	0.70678 (19)	0.0564 (6)
C1	0.0695 (3)	0.5737 (4)	0.49890 (16)	0.0509 (6)
C5	0.4688 (3)	0.3894 (4)	0.71993 (19)	0.0556 (6)
H4	0.705 (3)	0.590 (3)	0.7528 (19)	0.062 (6)*
H3	0.345 (3)	0.875 (4)	0.507 (2)	0.068 (6)*
H5	0.504 (3)	0.265 (4)	0.7803 (18)	0.063 (6)*
H6	0.215 (3)	0.250 (4)	0.6607 (17)	0.066 (6)*
H1	0.051 (3)	0.706 (4)	0.4498 (19)	0.071 (7)*

Atomic displacement parameters ($\text{\AA}^2$)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0566 (11)	0.0569 (11)	0.0674 (11)	−0.0077 (8)	0.0048 (9)	0.0012 (9)
C2	0.0493 (11)	0.0476 (11)	0.0416 (10)	−0.0010 (9)	0.0103 (8)	−0.0024 (8)
C3	0.0562 (13)	0.0522 (13)	0.0519 (12)	−0.0045 (9)	0.0068 (10)	0.0027 (10)
C6	0.0491 (12)	0.0522 (13)	0.0585 (13)	−0.0029 (9)	0.0120 (10)	0.0048 (10)
C4	0.0465 (12)	0.0671 (14)	0.0551 (12)	0.0010 (10)	0.0069 (10)	−0.0019 (11)
C1	0.0553 (12)	0.0526 (12)	0.0448 (11)	−0.0034 (8)	0.0085 (9)	0.0020 (10)
C5	0.0517 (12)	0.0591 (13)	0.0570 (12)	0.0067 (9)	0.0117 (10)	0.0095 (10)

Geometric parameters ($\text{\AA}$, $^\circ$)

N1—C3	1.336 (3)	C6—H6	0.98 (2)
N1—C4	1.333 (3)	C4—C5	1.374 (3)
C2—C3	1.395 (3)	C4—H4	0.98 (2)
C2—C6	1.382 (3)	C1—C1 ⁱ	1.320 (4)
C2—C1	1.465 (3)	C1—H1	0.93 (2)
C3—H3	0.99 (2)	C5—H5	0.99 (2)
C6—C5	1.372 (3)		
C4—N1—C3	116.54 (18)	N1—C4—C5	123.4 (2)
C3—C2—C1	119.85 (19)	N1—C4—H4	115.2 (11)
C6—C2—C3	116.44 (19)	C5—C4—H4	121.4 (11)
C6—C2—C1	123.71 (18)	C2—C1—H1	115.3 (13)
N1—C3—C2	124.8 (2)	C1 ⁱ —C1—C2	127.1 (3)

N1—C3—H3	116.6 (12)	C1 ⁱ —C1—H1	117.4 (13)
C2—C3—H3	118.6 (12)	C6—C5—C4	119.1 (2)
C2—C6—H6	119.4 (12)	C6—C5—H5	120.0 (11)
C5—C6—C2	119.80 (19)	C4—C5—H5	120.8 (11)
C5—C6—H6	120.8 (12)		
N1—C4—C5—C6	−0.5 (3)	C6—C2—C3—N1	−0.7 (3)
C2—C6—C5—C4	0.2 (3)	C6—C2—C1—C1 ⁱ	4.7 (4)
C3—N1—C4—C5	0.1 (3)	C4—N1—C3—C2	0.5 (3)
C3—C2—C6—C5	0.3 (3)	C1—C2—C3—N1	178.62 (17)
C3—C2—C1—C1 ⁱ	−174.6 (2)	C1—C2—C6—C5	−178.96 (17)

Symmetry code: (i) $-x, -y+1, -z+1$.