



The Role of XPB/Xsl2 dsDNA Translocase Processivity in Transcription Start-site Scanning

Eric J. Tomko¹, Olivia Luyties², Jenna K. Rimel², Chi-Lin Tsai⁴, Jill O. Fuss⁵, James Fishburn³, Steven Hahn³, Susan E. Tsutakawa⁵, Dylan J. Taatjes² and Eric A. Galburt^{1*}

1 - Biochemistry and Molecular Biophysics, Washington University School of Medicine, St. Louis, MO 63110, USA

2 - Dept. of Biochemistry, University of Colorado, Boulder, CO 80303, USA

3 - Division of Basic Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA 98109, USA

4 - Department of Molecular and Cellular Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA

5 - Molecular Biophysics and Integrated Bioimaging, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

Correspondence to Eric A. Galburt: egalburt@wustl.edu (E.A. Galburt)

<https://doi.org/10.1016/j.jmb.2021.166813>

Edited by Anna Pyle

Abstract

The general transcription factor TFIID contains three ATP-dependent catalytic activities. TFIID functions in nucleotide excision repair primarily as a DNA helicase and in Pol II transcription initiation as a dsDNA translocase and protein kinase. During initiation, the XPB/Xsl2 subunit of TFIID couples ATP hydrolysis to dsDNA translocation facilitating promoter opening and the kinase module phosphorylates Pol II to facilitate the transition to elongation. These functions are conserved between metazoans and yeast; however, yeast TFIID also drives transcription start-site scanning in which Pol II scans downstream DNA to locate productive start-sites. The ten-subunit holo-TFIID from *S. cerevisiae* has a processive dsDNA translocase activity required for scanning and a structural role in scanning has been ascribed to the three-subunit TFIID kinase module. Here, we assess the dsDNA translocase activity of ten-subunit holo- and core-TFIID complexes (i.e. seven subunits, lacking the kinase module) from both *S. cerevisiae* and *H. sapiens*. We find that neither holo nor core human TFIID exhibit processive translocation, consistent with the lack of start-site scanning in humans. Furthermore, in contrast to holo-TFIID, the *S. cerevisiae* core-TFIID also lacks processive translocation and its dsDNA-stimulated ATPase activity was reduced ~5-fold to a level comparable to the human complexes, potentially explaining the reported upstream shift in start-site observed *in vitro* in the absence of the *S. cerevisiae* kinase module. These results suggest that neither human nor *S. cerevisiae* core-TFIID can translocate efficiently, and that the *S. cerevisiae* kinase module functions as a processivity factor to allow for robust transcription start-site scanning.

© 2021 Elsevier Ltd. All rights reserved.

Introduction

The general transcription factor TFIID plays a pivotal role in both RNA polymerase II (Pol II) transcription initiation and DNA repair.^{1–3} The core TFIID complex from *H. sapiens* and other meta-

zoans has seven subunits: XPB, XPD, p62, p52, p44, p34, and p8. The core is functional in nucleotide excision repair (NER) and is dependent on the XPB and XPD nucleic acid motor proteins to unwind DNA around a site of damage for excision and repair.⁴ For transcription initiation, core-TFIID

associates with a kinase module comprised of three proteins: cyclin-dependent kinase 7 (CDK7), Cyclin H and MAT1, thus forming holo-TFIH. In metazoans, the TFIH kinase module is also called the CDK activating kinase (CAK), because CDK7 activates other cellular CDKs.^{3,5} The orthologous 3-subunit TFIH kinase module in yeast (*S. cerevisiae*) is called TFIK. Similar to metazoan TFIH, TFIK associates with core-TFIH to form a 10-subunit holo-TFIH complex. During transcription initiation and promoter escape, CDK7 phosphorylates the C-terminal domain of the Pol II Rpb1 subunit.^{6,7} Unlike in DNA repair, XPD only plays a structural role in the formation of the pre-initiation complex (PIC),⁸ while XPB activity is required for promoter unwinding.⁹ While these molecular activities appear to be generally conserved across Eukaryotes, in yeast exclusively (*S. cerevisiae*, and a host of recently identified related yeasts), TFIH has been shown to drive transcription start-site (TSS) scanning at TATA-dependent promoters, generating a distinct TSS profile genome-wide compared to other Eukaryotes.^{10–13}

The TSS-scanning activity of the yeast Pol II PIC results in downstream TSS being favored compared to the upstream site of initial DNA unwinding. More specifically, the *S. cerevisiae* PIC utilizes start-sites 40–150 bp downstream of the TATA box whereas metazoans initiate transcription 25–30 bp downstream of the TATA sequence.^{10,14–16} Yeast and human Pol II initiation complexes share a high degree of structural conservation with no apparent distinctions that explain differences in TSS-scanning propensity.^{17,18} In studies of the *S. cerevisiae* homologue of XPB, Ssl2, we previously showed that in the context of TFIH, scanning depends on an ATP-dependent double-stranded DNA (dsDNA) translocase activity which leads to a partial DNA opening.^{19,20} This translocation has also been observed directly in single-molecule optical tweezers assays.²¹ Furthermore, mutations in the Pol II trigger loop (causing faster or slower transcription kinetics) shift the TSS upstream or downstream, respectively.^{13,22} Taken together, these observations suggest that an interplay between the kinetics of Pol II initial transcription and the kinetics of TFIH dsDNA translocation determines TSS usage.^{13,20} This model, which we refer to as the kinetic competition model, has several possible versions involving different aspects of the kinetics of each motor. However, in general, it predicts that if TFIH is quicker to engage, exhibits faster translocation, and is more processive, downstream scanning will occur. Conversely, if TFIH translocation is slower to engage, exhibits slower translocation, and is less processive, initiation at upstream sites will be favored.

To test this hypothesis, we determined the parameters of dsDNA translocation of both *H. sapiens* and *S. cerevisiae* TFIH complexes in the presence and absence of their respective kinase

modules. Consistent with the model, we find that human TFIH is a poor dsDNA translocase compared to *S. cerevisiae* TFIH and that the yeast kinase module specifically stimulates processive translocation.

Results

HsTFIIH XPB ATPase is stimulated by double-stranded (ds) DNA

As described above, *S. cerevisiae* TFIH has been shown to have an ATP-dependent dsDNA translocase activity which is required for DNA unwinding and scanning.¹⁹ In contrast, scanning does not appear to occur in metazoans at TATA-dependent promoters despite structural conservation between *S. cerevisiae* and *H. sapiens* Pol II PICs.^{17,18} One possible explanation for this lack of scanning could be a difference in the dsDNA translocase kinetics of TFIH. To see if the *H. sapiens* TFIH (*HsTFIIH*) dsDNA translocase kinetics differs from *S. cerevisiae* TFIH (*ScTFIIH*), we quantitated the steady-state ATPase kinetics as a function of DNA length as done previously with the *S. cerevisiae* enzyme.¹⁹

TFIIH ATPase activity was measured using a continuous fluorescence assay in plate reader format in which ATP turnover is coupled to NADH reduction through the action of pyruvate kinase and lactate dehydrogenase (Figure 1A and Methods). For every equivalent of ADP generated, an equivalent of NADH is oxidized to NAD, resulting in a decrease in fluorescence.^{23,24} The continuous assay is advantageous, yielding more data points in a lower reaction volume and adapting the assay to a plate format allows for multiple experiments to be performed in parallel. Core- or holo-*HsTFIIH* was preincubated in the absence of DNA or the presence of either plasmid DNA (5500 bp) or linear dsDNA in *Hs*-ATPase buffer at 30 °C for 10 min. ATP and phosphoenol pyruvate (PEP) were added to initiate the reaction. No ATPase activity was detectable in the absence of DNA, but in the presence of DNA, the fluorescence signal decreased linearly over time (Figure 1B). This fluorescence signal was converted to moles of ADP using a standard curve (collected during each experiment, Figure 1C) and fit to a line to measure steady-state ADP production (Figure 1D). The plasmid-concentration dependence of the core-*HsTFIIH* steady-state rate increased hyperbolically, displaying Michaelis-Menten kinetics with $K_m = 12 \pm 3 \mu\text{M-bp}$ and $V_{\max} = 0.8 \pm 0.1 \text{ ATP/sec/TFIIH}$ (Figure 1E). Holo-*HsTFIIH* produced similar linear ATPase time courses at saturating plasmid concentration (Figure 1D); however, the $V_{\max}$ was approximately 2-fold lower ($V_{\max} = 0.5 \pm 0.02 \text{ ATP/sec/TFIIH}$). Fit parameters are listed in Supplementary Table 1.

To distinguish between the ATPase activities of XPD and XPB, we used the XPB-specific inhibitor




Figure 1. Human TFIIH core ATPase is stimulated by dsDNA and is XBP-dependent. (A) Diagram of the continuous, fluorescent, enzyme-coupled ATPase assay used to measure TFIIH steady-state ATPase activity. TFIIH ATP hydrolysis was monitored by following the oxidation of NADH to NAD through the regeneration of ADP to ATP through the glycolytic enzymes pyruvate kinase and lactate dehydrogenase. NADH fluorescence at 460 nm is quenched upon NADH oxidation to NAD. (B) Observed fluorescence decrease relative to the initial value ($F_{obs} - F_{int}$) time-course of core-HsTFIIH ATPase activity in the presence of 100 μ M-bp plasmid DNA after subtracting a control-time course of core-HsTFIIH in the absence of DNA. (C) ADP standard curve for converting observed NADH fluorescence to moles of ADP. (D) Representative ATPase time courses for core- and holo-HsTFIIH. The core-HsTFIIH, time course is the same time course in (B) converted to ADP/HsTFIIH using the standard curve in (C) and the moles of HsTFIIH determined by SDS-PAGE as described in Materials and Methods. The slope of a linear fit is the observed rate of ATP hydrolysis. (E) The rate of core-HsTFIIH ATP hydrolysis as a function of plasmid DNA concentration. The trends are fit with the Michaelis-Menten equation (solid line), see Materials and Methods. (F) Triptolide inhibition curves for core-HsTFIIH ATPase in the presence of plasmid (circles) and linear dsDNA (80 bp, triangles).

triptolide²⁵ (Supplemental Figure 1A). Core-HsTFIIH was incubated with 100 μ M-bp plasmid DNA in the absence or presence of increasing amounts of triptolide in Hs-ATPase buffer at 30 °C for 20 minutes prior to reaction. As judged by fitting the data to an inhibitory binding curve, $81 \pm 4\%$ of the observed ATPase activity was inhibited by a sat-

urating concentration of triptolide (100 μ M), indicating the plasmid DNA-stimulated ATPase activity is XBP-dependent (Figure 1F). An 80 bp linear dsDNA was also tested since the presence of a free DNA end, although blunt, could stimulate loading of XPD, resulting in ATP hydrolysis.²⁶ As with the plasmid template, triptolide inhibited the majority

(80 ± 10%) of the observed ATPase activity on the 80 bp dsDNA (Figure 1F) although the inhibition curve is shifted to higher triptolide concentrations (saturation at 10 mM). As TFIIH was incubated with triptolide in the presence of DNA, this suggested that DNA may competitively inhibit triptolide binding to core-*HsTFIIH* and that core-*HsTFIIH* has a higher affinity for the 80 bp linear DNA than the plasmid DNA.

Structures of XPB in the contexts of core-TFIIH and the PIC provide insight into how triptolide disrupts its ATPase activity and how DNA binding may compete with triptolide binding. Mass spectroscopic analysis of triptolide inactivated-*HsTFIIH* showed cysteine 342 of XPB to be covalently modified.²⁷ Inspection of the apo core-*HsTFIIH* structure²⁸ reveals cysteine 342 to be exposed to solvent and positioned at the back of a pocket surrounded by residues that surround and form the ATP binding site in nucleic-acid-stimulated ATPases.²⁹ However, in the DNA-bound XPB sub-structure from the human PIC, cysteine 342 becomes less solvent exposed due the presence of the bound DNA¹⁷ (Supplemental Figure 1B). This can be seen directly by aligning the apo and DNA bound XPB structures and is consistent with our data which suggests that bound DNA limits triptolide access to cysteine 342 (Supplemental Figure 1C). Speculatively, in turn, a triptolide adduct at cysteine 342 would be predicted to disrupt both ATP- and DNA-binding by XPB.

Both core- and holo-*HsTFIIH* are poor dsDNA translocases

The dsDNA-length dependence of the steady-state ATPase kinetic parameters, K_m and V_{max} , can provide insight into the translocation mechanism of nucleic acid motors.^{19,30,31} In particular, the probability of the motor to take a step along the DNA rather than dissociate, quantitated by the processivity (P), can be determined from the DNA length dependence of V_{max} .³¹ For processive motors, V_{max} will increase as the DNA length increases past the average length translocated per binding event, and plateau at significantly longer lengths. In contrast, a DNA-length independent V_{max} provides evidence of either no processivity or a processivity significantly less than the lowest DNA length tested. We measured the steady-state ATPase rate for core-*HsTFIIH*, titrating the concentration of a series of linear dsDNA substrates with different lengths (20, 30, 60, 80 and 100 bp). These data were compared with plasmid DNA, representing an infinitely long template. The steady-state rate increased hyperbolically with DNA concentration for each DNA length tested (Figure 2A). Each titration was fit to the Michaelis-Menten equation to determine V_{max} and K_m . In contrast to a length-dependent V_{max} previously observed using holo-*ScTFIIH*,¹⁹ V_{max} was DNA length-independent with

core-*HsTFIIH* (Figure 2B, blue). The K_m was similar for lengths 20, 30, 60, 80, and 100 bp, but plasmid DNA K_m was 2-fold higher, suggesting a lower affinity toward plasmid DNA (Figure 2C).

We wondered if the lack of length-dependence could be explained by the lack of the kinase module, which has been suggested to play a role in yeast TSS-scanning. To test this idea, we determined the V_{max} for holo-*HsTFIIH* at saturating DNA concentration (1000 μ M-bp) for each DNA length. However, V_{max} for holo-*HsTFIIH* still DNA length independent (Figure 2B, orange). The lack of DNA-length dependence for both core- and holo-*HsTFIIH*, suggests that, in contrast to holo-*ScTFIIH*, human TFIIH complexes are not processive dsDNA translocases. Importantly, all subunits of holo-*HsTFIIH* are present in the preparation (Figure 2D) and the complex was active in both transcription and kinase assays (Figure 2E and F), as expected. In contrast, despite having the expected subunits and displaying ATPase activity, Core-*HsTFIIH* was not active in transcription assays (Figure 2E). Furthermore, when commercially available purified CAK (Millipore, cat: 14–476 M, CDK7-6xHis/CCNH/GST-MAT1) was added to reactions with core-*HsTFIIH* the “core + CAK” activity did not match that of purified intact holo-TFIIH (Figure 2E, F), suggesting that it was not incorporated into a *bona fide* holo-complex.

Removal of the *S. cerevisiae* Kinase Module Reduces Both ATPase Rate and dsDNA Translocation Processivity

As described above, the kinase module had little effect on *HsTFIIH* dsDNA translocase activity. However, the removal of the *S. cerevisiae* kinase module has been shown result in upstream shifts in TSS *in vitro*.³² In fact, this shift leads to the usage of a start-site position similar to that found in metazoans. While the shift is not absolute, the result suggests a significant reduction in the prevalence of TSS scanning. Furthermore, adding back an enzymatically dead kinase module restores usage of downstream start-sites, suggesting that the module plays a structural role, as opposed to a catalytic one, in facilitating TSS scanning.³² We hypothesized that if scanning activity is due to processive TFIIH translocation, removal of the kinase module should also reduce (or eliminate) *ScTFIIH* dsDNA translocase activity.

To test this, we affinity purified core-*ScTFIIH* (Figure 3A) and measured the steady-state dsDNA-stimulated ATPase activity of core- and holo-*ScTFIIH* at saturating DNA concentrations on DNA templates of varying lengths as described above. As the Tfb3/MAT1 subunit of the kinase module anchors the kinase module to core-TFIIH,³³ core-*ScTFIIH* was purified from a strain containing Tfb3 with an auxin degron tag (see Meth-




Figure 2. Core- and holo-*HsTFIIH* are poor dsDNA translocases. (A) DNA concentration dependence of core-*HsTFIIH* steady-state ATPase rate for different linear dsDNA lengths (lengths: 20, 30, 60, 80 and 100 bp) and plasmid DNA. Data was fit with the Michaelis-Menten equation (solid lines) to determine K_m and V_{max} for each DNA length. (B) Core-*HsTFIIH* and holo ATPase V_{max} as a function of DNA length. A V_{max} independent of DNA length suggests very low to no processivity for dsDNA translocase. (C) Core-*HsTFIIH* ATPase K_m for DNA as a function of DNA length. (D) SDS-PAGE sypro ruby stain of purified core and holo-*HsTFIIH* complexes and commercially purchased CAK. (E) Plot shows quantitation of *in vitro* transcription of *HsTFIIH* complexes with or without added CAK, with bars representing the standard error of the mean calculated from at least 7 experiments (No *TFIIH*: N = 8, holo-*HsTFIIH*: N = 9, core-*HsTFIIH*: N = 7, CAK: N = 9, core-*HsTFIIH* + CAK: N = 9). For all comparisons with holo-*HsTFIIH*, p-values were 2.2×10^{-4} or smaller. Shown below the plot are runoff RNA transcripts (7% polyacrylamide-urea gel) from a representative experiment with the contrast enhanced for visualization. (F) *In vitro* kinase assay of *HsTFIIH* complexes with or without added CAK. Both a hyperphosphorylated (IIO) and hypo-phosphorylated (IIA) CTD is observed.

ods). Both core and holo complexes also contained an ATPase-dead Rad3 subunit (homologous to XPD) to ensure that any ATPase activity was from

Ssl2 (homologous to XPB). Furthermore, both holo- and core-*ScTFIIH* were active in transcription where PIC formed with core-*ScTFIIH* was able to




Figure 3. Purified core-ScTFIIH displays impaired TSS-scanning. (A) SDS-PAGE silver stain of purified holo- and core-ScTFIIH. **(B)** *In vitro* run-off transcription with holo- and core-ScTFIIH on the HIS4 promoter template. Asterisks denote RNA products initiated from the metazoan (blue) and yeast primary endogenous (green) transcription start-sites. Secondary TSSs from the yeast reaction are obscured by a TFIIH-independent band also observed in the no TFIIH control lane.

initiate transcription upstream at or near the metazoan start site with less initiation at downstream sites as previously observed (Figure 3B).³² As expected from our previous work, the ATPase time courses for holo-ScTFIIH increase linearly (Supplemental Figure S3) and $V_{\max}$ increases with DNA

length, plateauing at lengths greater than 100 bp (Figure 4B, orange).¹⁹ A non-linear least-squares analysis of the DNA-length dependence of $V_{\max}$ yielded a processivity of 0.91 ± 0.01 equivalent to on average 11 bps translocated per binding event.

In contrast, core-ScTFIIH time courses increase linearly at a slower rate (Supplemental Figure S3) and the measured $V_{\max}$ values taken under saturating DNA concentrations do not show DNA-length dependence (Figure 4A and 4B, blue). This result is consistent with core-ScTFIIH lacking observable dsDNA translocase activity similar to the behavior of the human complexes. We also measured the core-ScTFIIH DNA concentration dependence of the ATPase activity to confirm that we were measuring $V_{\max}$ in the presence of saturating DNA and to eliminate alternative models of translocation that predict a length-dependence in K_m instead of $V_{\max}$.³¹ Each DNA length tested displayed a hyperbolic dependence on DNA concentration which was fit to the Michaelis-Menten equation to determine K_m (Figure 4A). The core-ScTFIIH K_m was similar for each length (Figure 4C) and higher than holo-ScTFIIH ($K_m = 1.5 \pm 0.3 \mu\text{M-bp}$) determined previously (Figure 4C dashed line),¹⁹ suggesting the core-ScTFIIH has a lower affinity for DNA than holo-ScTFIIH. Interestingly, the core-ScTFIIH $V_{\max}$ is 5-fold lower than the holo-ScTFIIH $V_{\max}$ at long DNA lengths revealing that the kinase module not only promotes dsDNA translocation processivity but also appears to enhance the absolute dsDNA stimulated ATPase rate. Taken together, these results further support a mechanism in which TFIIH dsDNA translocation is the underlying activity that produces TSS-scanning in *S. cerevisiae*.

Discussion

The general transcription factor TFIIH plays a pivotal role in Pol II transcription initiation, facilitating promoter DNA opening by translocating downstream DNA into Pol II. As TBP binds upstream DNA, this translocation generates torsional strain which then promotes DNA melting around the TSS. This process is conserved among Eukaryotes and allows ssDNA to enter the Pol II active site. However, in some yeast species, TFIIH also facilitates TSS-scanning in which, after initial DNA unwinding, the PIC scans downstream DNA and utilizes TSS located 40–120 bp downstream of TATA. This scanning mechanism requires Ssl2/XPB, the ATP-dependent dsDNA translocase of TFIIH. The model for scanning proposes a kinetic competition between Ssl2/XPB dsDNA translocation and Pol II initial transcription. This model is supported by the requirement of TFIIH dsDNA translocation and the observation that loss-of-function mutations in *S. cerevisiae* Pol II (i.e. slower polymerase) shift the TSS downstream, whereas gain-of-function mutations

(i.e. faster polymerase) shift the TSS upstream.¹³ Interestingly, metazoan Pol II PICs do not appear to scan on TATA containing promoters and instead utilize TSS that are 25–30 bp downstream of the TATA sequence.^{10,14,16} In the context of the kinetic

competition model, a lack of scanning would be predicted in the case of a shift in the relative kinetics between the Pol II and Ssl2/XPB motors such that Pol II initial transcription is faster and/or more processive while Ssl2/XPB dsDNA translocation is slower and/or less processive. Therefore, we undertook the work described here to test the hypothesis that metazoan TFIIF is a slower and/or less processive dsDNA translocase compared with *S. cerevisiae* TFIIF.

We previously showed that *S. cerevisiae* holo-TFIIF processively translocates along dsDNA using a discontinuous ATPase assay.¹⁹ We repeated those experiments using the continuous fluorescent ATPase assay used to measure core- and holo-*Hs*TFIIF in this study and obtained results consistent with our previous study. Analysis of the DNA-length dependence of the ATPase $V_{\max}$ yielded a translocation processivity of 0.91 (11 bp per binding event on average), indicating holo-*Sc*TFIIF translocates about one turn of dsDNA, on average, before dissociation from DNA (Figure 4B). In contrast, neither core- nor holo-*Hs*TFIIF showed any length-dependence of $V_{\max}$ for the DNA lengths tested. This result indicates that their processivity, if any, cannot be detected by this assay (Figure 2B). However, we can place an upper limit on the processivity of 2 bp by comparing the data to $V_{\max}$ values calculated from Eq. (1) for different DNA lengths and varying processivities. This analysis shows that processivities of 1 and 2 bp are within the error of the observed $V_{\max}$ values (Supplemental Figure 2). Crucially, the affinity purified holo-*Hs*TFIIF was active in transcription and kinase assays *in vitro*, suggesting the lack of processivity is a *bona fide* feature of the complex. Furthermore, the observed ATPase activity in *Hs*TFIIF was inhibited by triptolide, establishing that XPB, and not XPD, was the motor stimulated in our assays. These observations reveal that holo-*Hs*TFIIF is a poor DNA translocase compared to holo-*Sc*TFIIF, consistent with the predictions of our model (Figure 5A, B).




Figure 4. *Sc*TFIIF kinase module promotes dsDNA translocation processivity and enhances ATPase activity. (A) DNA concentration dependence of core-*Sc*TFIIF steady-state ATPase. Data was fit with the Michaelis-Menten equation (solid lines) to determine $V_{\max}$ and K_m . (B) Holo- and core-*Sc*TFIIF ATPase $V_{\max}$ as a function of DNA length. Holo-*Sc*TFIIF $V_{\max}$ as a function of DNA length was fit with Eq. (1) to determine the translocation processivity. Core-*Sc*TFIIF core $V_{\max}$ is independent of DNA length, indicating the translocase has very low to no processivity. (C) Core-*Sc*TFIIF K_m as a function of DNA length. Dashed line is the average K_m (1.5 ± 0.3 μM-bp) previously observed for holo-*Sc*TFIIF¹⁹ under the same solution conditions.




Figure 5. TFIID dsDNA translocase processivity limits TSS-scanning. (A, B) A model for TSS-scanning comparing the *S. cerevisiae* and *H. sapiens* systems. In each case, ATP hydrolysis by TFIID (i.e., either Ssl2 or XPB) leads to initial DNA unwinding. In this particular model, TFIID engagement precludes Pol II activity and vice-versa. When TFIID dissociates from the DNA, Pol II attempts to initiate by forming a 10-mer RNA. The probability of a translocation run of a certain length is governed by the processivity of TFIID and the probability of successfully initiating is governed by the processivity of Pol II. See Material and Methods for details on the model and simulations. **(A)** In *S. cerevisiae*, Ssl2 processivity ($P = 0.9$) leads to downstream DNA being drawn into the complex and the formation of a dsDNA loop. **(B)** In *H. sapiens* the low (or nonexistent) XPB processivity ($P = 0.5$) supports DNA unwinding, but not start-site scanning. **(C)** Cumulative distribution functions of TSSs from model simulations with different processivities for TFIID ($P = 0.5, 0.9, 0.99$) and a constant Pol II processivity (0.98). **(D)** Cumulative distribution functions of TSSs from model simulations with different processivities for Pol II ($P = 1, 0.83, 0.67$) and a constant TFIID processivity (0.9).

There are many possible versions of the general kinetic competition model, especially in terms of how the two motors may be coupled. In other words, to what extent does the engagement of one motor prohibit the engagement of the other? Alternatively, can the engagement of one motor interrupt a cycle of the other? As our data do not comment on these aspects of the model, we considered a simplified model where either TFIID (i.e., Ssl2 or XPB) or Pol II can be engaged with the DNA productively at any given time. In addition, the binding of one motor must wait until the other motor dissociates. We performed kinetic simulations using rates of translocation and dissociation for each motor to define their processivities. After a cycle of TFIID translocation, Pol II may attempt to initiate transcription. If successful, a TSS is scored; if not successful,

another cycle of TFIID translocation moves the system further downstream (Figure 5A, B). We first defined a Pol II processivity of 0.98 representing a probability of initiation less than unity and varied the processivity of TFIID and plotted TSS distributions that result. As one expects intuitively, the distribution expands downstream with increasing translocase processivity (Figure 5C). Second, we used a fixed TFIID processivity of 0.9 (similar to that measured for *S. cerevisiae* Ssl2/TFIID) and varied the processivity of Pol II. Again, as one expects, the TSS distribution expands downstream as Pol II processivity is reduced as more cycles of TFIID translocation per initiation event are produced (Figure 5D). We would like to stress that the processivity defines the probability of taking a forward step instead of dissociating and leads to

an exponential distribution of run lengths. Thus, longer runs than the average occur with probabilities considerably higher than that expected from a normal distribution. For example, with a processivity of 0.9, 10% of binding events would be expected to lead to runs longer than 22 bp and 0.1 % would be expected to translocate at least 66 bp (Figure 5A, Supplemental Figure S4).

Intriguingly, PICs formed with core-ScTFIIH have been reported to have reduced ability to scan downstream to endogenous TSS *in vitro* and instead initiate upstream near the TSS observed in metazoans.³² This result suggested that removal of the kinase module may reduce the processivity and/or translocase activity of ScTFIIH. Our assays bore this out: core-ScTFIIH complexes displayed a DNA-length independent V_{max} and a 5-fold lower rate than holo-ScTFIIH at long DNA lengths (Figure 4B). In addition, PICs formed with core-ScTFIIH in our assay displayed upstream shifted TSS consistent with previous *in vitro* observations (Figure 3B).³² These observations support the kinetic competition model of TSS scanning.

The TFIIH kinase module itself is not well-resolved in the current PIC and TFIIH structural models.^{18,28,34,35} However, a number of chemical cross-links have been observed between Tfb3/MAT1, the main component of the kinase module that connects it to core-ScTFIIH, and Ssl2/XBP.³³ These cross-links are located away from the Ssl2 DNA binding cleft and similar cross-links were observed in HsTFIIH. Thus, while a Tfb3/MAT1 interaction with the Ssl2/XBP N-terminal region could potentially stimulate *S. cerevisiae* Ssl2 ATPase and translocase processivity through an allosteric mechanism, the presence of these contacts in both species argues that these contacts do not play direct roles in TSS scanning.

Based on Endonuclease III digestions in the presence and absence of the kinase module, it has been suggested that the kinase module affects the PIC footprint on the promoter DNA.³² Whether this is via direct DNA binding or indirect changes in the footprints of core-TFIIH or the polymerase is not known. However, if this conformational change prevents the *S. cerevisiae* PIC from initiating at the upstream metazoan site and/or stimulates the processivity of Ssl2, it may partially underlie the observed shift in TSS.

The effect of the *S. cerevisiae* kinase module on dsDNA translocation described here adds to observations in other contexts that the nucleic acid motors within core-TFIIH can be activated through factor-specific interactions. More specifically, core-HsTFIIH DNA translocation activity was shown to be enhanced by the NER factor XPA although XPA-dependent activation was not unambiguously assigned to either XPB or XPD.³⁶ Relatedly, structures of core-HsTFIIH in the presence and absence of a partial kinase module (MAT1) revealed that core subunits interact with

XPB and may block its DNA binding sites, suggesting that conformational rearrangements stimulated by protein-protein interactions, are needed to activate XPD.^{28,35}

While the removal of the kinase domain from ScTFIIH results in loss of a processivity signal in our assay (Figure 4B), the core-ScTFIIH still retains some ability to scan to the endogenous sites as judged from the *in vitro* transcription reaction itself (Figure 3B). This suggests that the activities of TFIIH may be modulated by its incorporation into the PIC. One consistent interpretation of these data is that the processivities of both core- and holo-ScTFIIH are stimulated by protein-protein contacts with PIC general transcription factors. Future studies will be needed to determine how interactions between XPB/Ssl2, the kinase module, and other general transcription factors modulate the molecular activities we have observed here.

Material and Methods

Protein Purification. Core-HsTFIIH complex (XPB-preScission-GFP, XPD, p62, p52, p44, p34, and p8) was cloned into MacroBac vector 438a.³⁷ The protein was expressed in Sf9 cells supplemented with 1 mM L-cysteine and 0.1 mM ferric ammonium citrate. Based on the XPB fusion with GFP, TFIIH was purified anaerobically using GFP-nanobody binder³⁸ covalently linked to agarose-beads (NHS agarose, Pierce/Thermo); eluted with preScission protease; and purified using Superpose 6 (10/300) in 25 mM HEPES, pH 7.8, 150 mM NaCl, 50 mM KCl, 3% glycerol, and 3 mM β -mercaptoethanol. The elution with TFIIH core complex (confirmed by SDS-PAGE gel and 420 nm absorption peak) was stored at -80°C until use. The holo-HsTFIIH purification was completed as described.³⁹

Core-ScTFIIH was purified from strain SHY1290 (genotype: *mat a ade2::hisG his3 Δ 200 leu2 Δ 0 lys2 Δ 0 met15 Δ 0 trp1 Δ 63 ura3 Δ 0 rad3 Δ ::KanMx tfb6 Δ ::HPH pGPD1-OSTIR::HIS3 TFB3-3xV5-IAA7::URA3/pJF82 (ars cen LEU2 rad3 (E236Q) – (HA)1-Tap tag). This strain contains a Tap-tagged ATPase dead Rad3/XPB (E236Q); a deletion of the TFB6 gene to facilitate complex purification; and Tfb3 with an auxin degron tag to facilitate elimination of the kinase module. Cells (6 liters) were grown at 30°C in YPD + adenine media (3% final Glucose) to Abs600 = 3–4. 3-indoleacetic acid (IAA) in DMSO was added to 0.5 mM final for 30 min to deplete Tfb3 protein (the protein was undetectable by Western blot after this treatment, Supplemental Figure S5). Core-ScTFIIH and holo-ScTFIIH with an ATPase dead Rad3/XPB were purified by Tap-tag as previously described.¹⁹ Core-ScTFIIH was additionally purified by chromatography on a 0.28 ml Source 15Q column equilibrated in 20 mM HEPES, pH 7.6, 10% (v/v)*

glycerol, 200 mM KOAc, 1 mM EDTA, and 1 mM DTT. Protein was eluted over 30 column volumes with a KOAc gradient from 200 mM to 1.2 M. Fractions with core-ScTFIIH were then concentrated on a 0.1 mL calmodulin affinity column, aliquoted, and stored at -80°C previously described.¹⁹ The concentration of the TFIIH complexes were determined by comparing the purified protein with a titrated BSA standard on SDS-PAGE gel (HsTFIIH) or by quantitative Western (ScTFIIH).

The *S. cerevisiae* Pol II RNA polymerase and the other general transcription factors (TBP, TFIIB, TFIIF, and TFIIE) were purified as previously described.⁴⁰ The human factors (Pol II, TBP, TFIIB, TFIIF, TFIIE, and TFIIH) were purified as described.³⁹

DNA preparation. DNA oligos of random sequence (Supplemental Table 2) and corresponding complementary oligos were purchased from IDT (Integrated DNA Technologies, Inc., Coralville, Iowa, USA) and resuspended in 10 mM Tris-HCl, pH 8.0. For a given DNA length oligos were annealed in a 1:1 molar ratio in Annealing buffer (10 mM Tris-HCl, pH 8.0, 50 mM NaCl). The DNA solution was heated at 95°C for 5 min then slowly cooled (~ 4 – 5 hrs) to room temperature ($\sim 23^{\circ}\text{C}$) in an insulated heating block. Annealed DNAs were verified by native-PAGE following standard protocols. The annealed DNA concentrations were determined from their absorbance at 260 nm using the calculated molar extinction coefficient for the duplex DNA based on the DNA sequence (nearest neighbor). A midi prep scale of plasmid DNA (pGO10GG, 5500 bp, sequence available upon request) was purified following manufacture protocol (Qiagen). The promoter template for in vitro transcription with the human Pol II system was prepared from the native human HSPA1B gene. The HSPA1B gene was initially amplified from HeLa genomic DNA by PCR (forward primer: CTCCTT CCCATT AAGACG GAAAAA ACATCC GGGAGA GCCGGT CCG; reverse primer: ACCTTG CCGTGT TGGAAC ACCCCC ACGCAG GAGTAG GTGGTG CCCAGGTC) and cloned into a TOPO vector. The -500 to $+216$ region corresponding to the HSPA1B promoter was amplified from the plasmid by PCR with Phusion polymerase (Thermo-Fisher #F530S) and purified via E.Z.N.A Gel Extraction Kit (Omega BioTek #D2500). The DNA was ethanol precipitated, resuspended to 100 nM in milliQ water, and stored at -80°C in 10 μL single-use aliquots. The promoter template for in vitro transcription with the *S. cerevisiae* Pol II system was prepared from the native His4 gene. A 236 bp DNA region, spanning -74 to $+107$, was PCR amplified from plasmid (pSH515) containing the native His4 gene promoter and purified by PCR cleanup kit (Qiagen). DNA was eluted from the

spin column with 10 mM Tris-HCl, pH 8.0 and stored at -20°C .

Fluorescent coupled ATPase assay. Steady-state ATP hydrolysis was measured by coupling the production of ADP to the reduction of NADH via the glycolytic enzymes pyruvate kinase and lactate dehydrogenase as depicted in Figure 1A.²³ ATPase reactions with HsTFIIH were carried out in hTRxn buffer (10 mM Tris-HCl pH 7.9, 10 mM HEPES pH 7.9, 50 mM KCl, 4 mM MgCl_2 , 10% (v/v) glycerol, 0.1 mg/ml BSA, and 1 mM DTT) at 30°C . ATPase reactions with ScTFIIH were carried out in yTRxn buffer (10 mM HEPES pH 7.7, 100 mM KCl, 10 mM MgOAc_2 , 3.5% (v/v) glycerol, 0.05 mg/ml BSA, and 1 mM DTT) at 25°C . All reagents were diluted and or prepared fresh in the appropriate TRxn buffer the day of the experiment and kept on ice. All reactions were prepared in a total volume of 10 μL the reported concentrations are the final concentrations after mixing. In a 8.75 μL volume, 0.133 mg/ml NADH (β -nicotinamide adenine dinucleotide reduced, Sigma, cat: N4505), 5.6 U/ml pyruvate kinase/ 7.8 U/ml lactate dehydrogenase (Sigma, cat: P0294), TFIIH (1.2 nM yeast variants or 6 nM human variants), and DNA were mixed and incubated on ice for 30 min. The reactions were then transferred to a black, 384 low volume, round bottom well plate (Corning, cat: 4514); covered with optical film to prevent evaporation (Applied Biosystems, cat: 4360954); and placed in the Synergy2 fluorescent plate reader (Bio Tek, Winooski, VT) and incubated for 10 min at the reaction temperature. After an initial scan of the plate (excitation: 340 nm, emission: 460 nm), the film was temporarily removed and the reaction initiated by adding 0.31 mg/ml PEP (phosphoenolpyruvic acid, Sigma, cat: 860077) and 0.5 mM ATP in a 1.25 μL volume using a multichannel pipet. The plate was then rapidly (~ 5 – 10 sec) resealed and placed back in the plate reader. Every two minutes the plate was scanned over a period of 75 mins. Triptolide inhibition of the ATPase activity was assayed by incubating TFIIH:DNA with varying amounts of triptolide (Sigma, cat: T3652) at 30°C for 30 min before initiating the ATPase reaction with PEP and ATP. Triptolide was dissolved in DMSO and stored at -20°C . The fluorescent signal was calibrated for each plate of reactions by setting up a series of reactions with known amounts of ADP without TFIIH and DNA. All reactions were prepared in triplicate.

In vitro transcription assays. Human pre-initiation complexes (PICs) were assembled on an activated HSPA1B promoter template (5 nM in 10 μL) by incubating 400 nM HSF1 with promoter template for 30 minutes at 30°C in transcription buffer (20 mM HEPES pH 7.6, 1 mM DTT, 8 mM MgCl_2) and enough DB (100) buffer (10% glycerol, 10 mM Tris pH 7.9, 180 mM KCl, 1 mM DTT) to ensure the final reaction volume is 20 μL . Pre-initiation complex (PIC) mix (TFIIB, TBP, TFIIF,

and Pol II) was then added (with or without the TFIIH variants as noted) and incubated for 15 minutes at 30 °C to permit PIC assembly. PIC factor concentration was optimized via titration, with each factor at approximately 20–50 nM as previously described.³⁹ *S. cerevisiae* PICs were assembled on the His4 promoter template (10 nM) in yTRxn buffer in a final volume of 20 µL at 25 °C (223 nM TBP, 31 nM TFIIIB, 13 nM TFIIIF, 10 nM Pol II, 2 nM TFIIIE, and 3 nM TFIIH). PIC components along with either holo- or core-ScTFIIH, at equivalent concentrations, are added to the promoter DNA and incubated for 30 min to permit PIC assembly.

Transcription was initiated by the addition of 2 µL of NTP mix (25 mM A/G/UTP, 5 mM CTP, 0.1 mCi α -³²P-CTP (3000 Ci/mmol), brought to 40 µL with DB (100) or 20 µL yTRxn buffer and allowed to proceed for 30 minutes at either 30 °C for the human or 25 °C for the yeast system. Reactions were stopped with the addition of 150 µL of STOP buffer (20 mM EDTA, 200 mM NaCl, 1% SDS) and 350 µL of cold 100% ethanol. RNA was precipitated at –20 °C overnight, resuspended in formamide loading buffer (95% deionized formamide, 0.025% w/v Bromophenol blue, 0.025% w/v Xylene cyanol FF, 5 mM EDTA), boiled, and run on a 40 cm 7% acrylamide sequencing gel (National Diagnostics SequaGel UreaGel 29:1 Denaturing Gel System). RNA was quantitated through pixel-density analysis of the ³²P signal in the sequencing gel using ImageJ. Bands corresponding to 216 nt were regarded as products of successful human Pol II transcription and selected for quantitation with background correction. Data from ImageJ was analyzed and graphed with GraphPad Prism8 software. For the yeast system bands corresponding to 138 nt are products initiated from the metazoan start-site while bands corresponding to 107, 96, and 94 nt are products initiated from the native downstream start-sites.

In vitro kinase assay. The kinase assays evaluating CDK7 activity with the HsTFIIH variants were performed with equivalent concentrations (2.5 nM) of holo-TFIIH, core-TFIIH, and the CAK with the Pol II CTD (*M. musculus*⁴¹ as the substrate and 100 µM ATP and ³²P-γ ATP (3000 Ci/mmol, 5 mCi/ml EasyTide, 250 µCi) for 30 minutes at 30 °C. These conditions were chosen to remain consistent with the *in vitro* transcription assays. The CAK complex was purchased from Millipore (CDK7-6xHis/CCNH/GST-MAT1; CAK complex, 250 µg cat: 14–476 M) and the Pol II GST-CTD was expressed in BL21 cells and purified as described.⁴¹ Kinase reactions were quenched with 4X Laemmli buffer, boiled for 5 minutes, and run on a 4–20% gradient protein gel (BioRad 4–20% Mini-PROTEAN® TGX™ Gel, 15 well, 15 µl cat#456-1096). Gels were dried, exposed to a phosphor imaging plate. Plates were scanned using a Typhoon imager and bands were quantitated

using ImageJ. The measured kinase activity was normalized to TFIIH holo and the standard deviation was reported.

Kinetic analysis and modeling. The initial ATPase steady-state rate was determined from the slope of the linear portion of the ATPase time courses. The ATPase steady-state kinetic parameters V_{max} and K_m for a given DNA substrate was determined by fitting the initial ATPase steady-state rate as a function of DNA concentration to the Michaelis-Menten equation. The steady-state ATPase rate for a translocating motor along a lattice will display Michaelis-Menten kinetics where the V_{max} will change as a function of lattice length.³¹ The expression for the DNA length dependent change in V_{max} is given in Eq. (1) and was previously used to determine the translocation processivity for holo-ScTFIIH.¹⁹

$$V_{max} = A \left(\frac{\left(\frac{L-d}{m} \right) - \frac{P \left(1 - P \left(\frac{L-d}{m} \right) \right)}{(1-P)}}{\left(1 + \left(\frac{L-d}{m} \right) \right)} \right) \quad (1)$$

In Eq. (1), P is the processivity, L , the DNA length, d , the translocase contact size, and m , the translocase step size per cycle. The parameter A is a product combination of microscopic parameters for the translocation model and the contact size and step size were constrained to 16 bp and 1 bp, respectively as done previously¹⁹ while the processivity was allowed to float in a non-linear least-squares analysis of the DNA length dependence of the V_{max} using Conlin.⁴²

Sequence-independent distributions of transcription start-sites were simulated using the scanning model depicted in Figure 5A. In this model, TFIIH engagement precludes Pol II activity and vice-versa. When TFIIH dissociates from the DNA, Pol II attempts to initiate by forming a 10-mer RNA. The probability of a TFIIH translocation run of a certain length is governed by the processivity of TFIIH and the probability of successfully initiating is governed by the processivity of Pol II. If Pol II successfully transcribes a 10-mer before dissociating, a transcription start-site is scored and the system resets; otherwise, the short RNA is released and TFIIH is able to rebind and perform another cycle of translocation. Gillespie simulations were performed in Matlab until 10,000 transcription start-sites were scored for different TFIIH processivities ($P = 0.5, 0.91, 0.99$; defined by $k_{translocate} = 10 \text{ s}^{-1}$, and $k_{dissociate} = 10 \text{ s}^{-1}, 1 \text{ s}^{-1}, 0.1 \text{ s}^{-1}$ respectively) and a constant Pol II processivity ($P = 0.98$; defined by $k_{NTP} = 50 \text{ s}^{-1}$ and $k_{dissociate} = 1 \text{ s}^{-1}$). In addition, simulations were performed with different Pol II processivities ($P = 1, 0.83, 0.67$; defined by $k_{NTP} = 50 \text{ s}^{-1}$ and $k_{dissociate} = 0 \text{ s}^{-1}, 10.2 \text{ s}^{-1}, 24.6 \text{ s}^{-1}$) and a constant TFIIH processivity ($P = 0.91$; defined by $k_{translocate} = 10 \text{ s}^{-1}$ and $k_{dissociate} = 1 \text{ s}^{-1}$).

$k_{\text{translocate}}$ was estimated using the observed ATPase V_{max} assuming TFIIH takes a single step with each ATP hydrolysis. k_{NTP} was estimated based on *in vitro* Pol II elongation rates. A dissociation rate constant for the polymerase was selected to give a processivity that was less than unity since Pol II stability on the DNA template is directly related to DNA:RNA hybrid length which is shorter during initiation.

Author Contributions

CLT, JOF, and SET cloned, expressed, and purified human core TFIIH. OL completed *in vitro* transcription assays with purified factors. JKR purified human TFIIH and performed kinase assays. JF purified yeast ScTFIIH derivatives and performed *in vitro* transcription assays. SH generated yeast strains for TFIIH core purification. EJT performed ATPase assays, analyzed the data, and did the modeling. EAG, DJT, SET, and SH supervised research and provided funding support. EJT and EAG designed the research and wrote the manuscript with input from all the other authors.

CRedit authorship contribution statement

Eric J. Tomko: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Writing - original draft, Visualization. **Olivia Luyties:** Investigation, Visualization. **Jenna K. Rimel:** Resources, Investigation, Visualization. **Chi-Lin Tsai:** Resources. **Jill O. Fuss:** Resources. **James Fishburn:** Resources. **Steven Hahn:** Resources, Writing - review & editing, Funding acquisition. **Susan E. Tsutakawa:** Resources, Writing - review & editing, Funding acquisition. **Dylan J. Taatjes:** Resources, Writing - review & editing, Funding acquisition. **Eric A. Galburt:** Conceptualization, Methodology, Formal analysis, Writing - original draft, Visualization, Funding acquisition, Project administration.

Acknowledgements

This work was supported by NCI P01 CA092584 (to SET and JOF); F31 CA254478 (to OL); F31 CA250432 (to JKR); NSF MCB-1818147 (to DJT); NIGMS R01GM110387 (to SET); NIGMS R01GM110064 (to DJT); NIGMS 2R01GM053451 (to SH); and NIGMS R01GM120559 (to EAG).

Declaration of Competing Interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: DJT is a member of the SAB at Dewpoint Therapeutics.

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jmb.2021.166813>.

Received 9 November 2020;

Accepted 4 January 2021;

Available online 13 January 2021

Keywords:

transcription initiation;

pre-initiation complex;

RNA polymerase II;

kinetics;

TFIIH double-stranded DNA translocation

References

- Schier, A.C., Taatjes, D.J., (2020). Structure and mechanism of the RNA polymerase II transcription machinery. *Gene Dev.*, **34**, 465–488. <https://doi.org/10.1101/gad.335679.119>.
- Tsutakawa, S.E., Tsai, C.-L., Yan, C., Bralić, A., Chazin, W. J., Hamdan, S.M., Schärer, O.D., Ivanov, I., Tainer, J.A., (2020). Envisioning how the prototypic molecular machine TFIIH functions in transcription initiation and DNA repair 102972 *Dna Repair.*, **96** <https://doi.org/10.1016/j.dnarep.2020.102972>.
- Rimel, J.K., Taatjes, D.J., (2018). The essential and multifunctional TFIIH complex. *Protein Sci.*, **27**, 1018–1037. <https://doi.org/10.1002/pro.3424>.
- Spivak, G., (2015). Nucleotide excision repair in humans. *Dna Repair.*, **36**, 13–18. <https://doi.org/10.1016/j.dnarep.2015.09.003>.
- Rimel, J.K., Poss, Z.C., Erickson, B., Maas, Z.L., Ebmeier, C.C., Johnson, J.L., Decker, T.-M., Yaron, T.M., Bradley, M.J., Hamman, K.B., Hu, S., Malojcic, G., Marineau, J.J., White, P.W., Brault, M., Tao, L., DeRoy, P., Clavette, C., Nayak, S., Damon, L.J., Kalthauer, I.H., Bunch, H., Cantley, L.C., Geyer, M., Iwasa, J., Dowell, R.D., Bentley, D.L., Old, W.M., Taatjes, D.J., (2020). Selective inhibition of CDK7 reveals high-confidence targets and new models for TFIIH function in transcription. *Gene Dev.*, <https://doi.org/10.1101/gad.341545.120>.
- Glover-Cutter, K., Larochele, S., Erickson, B., Zhang, C., Shokat, K., Fisher, R.P., Bentley, D.L., (2009). TFIIH-associated Cdk7 kinase functions in phosphorylation of C-terminal domain Ser7 residues, promoter-proximal pausing, and termination by RNA polymerase II ∇ †. *Mol. Cell. Biol.*, **29**, 5455–5464. <https://doi.org/10.1128/mcb.00637-09>.
- Larochele, S., Amat, R., Glover-Cutter, K., Sansó, M., Zhang, C., Allen, J.J., Shokat, K.M., Bentley, D.L., Fisher, R.P., (2012). Cyclin-dependent kinase control of the initiation-to-elongation switch of RNA polymerase II. *Nat. Struct. Mol. Biol.*, **19**, 1108–1115. <https://doi.org/10.1038/nsmb.2399>.
- Kuper, J., Braun, C., Elias, A., Michels, G., Sauer, F., Schmitt, D.R., Poterszman, A., Egly, J.-M., Kisker, C., (2014). In TFIIH, XPD helicase is exclusively devoted to

- DNA repair e1001954 *Plos Biol.*, **12** <https://doi.org/10.1371/journal.pbio.1001954>.
9. Kolesnikova, O., Radu, L., Poterszman, A., (2019). TFIIH: A multi-subunit complex at the cross-roads of transcription and DNA repair. *Adv. Protein Chem. Str.*, **115**, 21–67. <https://doi.org/10.1016/bs.apcsb.2019.01.003>.
 10. Lu, Z., Lin, Z., (2020). The origin and evolution of a distinct mechanism of transcription initiation in yeasts pp. 04.025502 *Biorxiv.*, **2020** (04) <https://doi.org/10.1101/2020.04.04.025502>.
 11. Giardina, C., Lis, J.T., (1993). DNA melting on yeast RNA polymerase II promoters. *Science (New York, NY)*, **261**, 759–762 <http://www.sciencemag.org/cgi/reprint/261/5122/759>.
 12. Fishburn, J., Galburt, E., Hahn, S., (2016). Transcription start site scanning and the requirement for ATP during transcription initiation by RNA polymerase II. *J. Biol. Chem.*, **291**, 13040–13047. <https://doi.org/10.1074/jbc.m116.724583>.
 13. Qiu, C., Jin, H., Vvedenskaya, I., Llenas, J.A., Zhao, T., Malik, I., Visbisky, A.M., Schwartz, S.L., Cui, P., Čabart, P., Han, K.H., Lai, W.K.M., Metz, R.P., Johnson, C.D., Sze, S.-H., Pugh, B.F., Nickels, B.E., Kaplan, C.D., (2020). Universal promoter scanning by Pol II during transcription initiation in *Saccharomyces cerevisiae*. *Genome Biol.*, **21**, 132. <https://doi.org/10.1186/s13059-020-02040-0>.
 14. Hampsey, M., (1998). Molecular genetics of the RNA polymerase II general transcriptional machinery. *Microbiol. Mol. Biol. Rev. Mmbr.*, **62**, 465–503.
 15. Carninci, P., Sandelin, A., Lenhard, B., Katayama, S., Shimokawa, K., Ponjavic, J., Semple, C.A.M., Taylor, M.S., Engström, P.G., Frith, M.C., Forrest, A.R.R., Alkema, W. B., Tan, S.L., Plessy, C., Kodzius, R., Ravasi, T., Kasukawa, T., Fukuda, S., Kanamori-Katayama, M., Kitazume, Y., Kawaji, H., Kai, C., Nakamura, M., Konno, H., Nakano, K., Mottagui-Tabar, S., Arner, P., Chesi, A., Gustincich, S., Persichetti, F., Suzuki, H., Grimmond, S.M., Wells, C.A., Orlando, V., Wahlestedt, C., Liu, E.T., Harbers, M., Kawai, J., Bajic, V.B., Hume, D.A., Hayashizaki, Y., (2006). Genome-wide analysis of mammalian promoter architecture and evolution. *Nat. Genet.*, **38**, 626–635. <https://doi.org/10.1038/ng1789>.
 16. Li, C., Lenhard, B., Luscombe, N.M., (2018). Integrated analysis sheds light on evolutionary trajectories of young transcription start sites in the human genome. *Genome Res.*, **28**, 676–688. <https://doi.org/10.1101/gr.231449.117>.
 17. He, Y., Yan, C., Fang, J., Inouye, C., Tjian, R., Ivanov, I., Nogales, E., (2016). Near-atomic resolution visualization of human transcription promoter opening. *Nature*, **533**, 359–365. <https://doi.org/10.1038/nature17970>.
 18. Schilbach, S., Hantsche, M., Tegunov, D., Dienemann, C., Wigge, C., Urlaub, H., Cramer, P., (2017). Structures of transcription pre-initiation complex with TFIIH and mediator. *Nature*, **551**, 204–209. <https://doi.org/10.1038/nature24282>.
 19. Fishburn, J., Tomko, E., Galburt, E., Hahn, S., (2015). Double-stranded DNA translocase activity of transcription factor TFIIH and the mechanism of RNA polymerase II open complex formation. *Proc. Natl. Acad. Sci.*, **112** <https://doi.org/10.1073/pnas.1417709112>.
 20. Tomko, E.J., Fishburn, J., Hahn, S., Galburt, E.A., (2017). TFIIH generates a six-base-pair open complex during RNAP II transcription initiation and start-site scanning. *Nat. Struct. Mol. Biol.*, **24**, 1139–1145. <https://doi.org/10.1038/nsmb.3500>.
 21. Fazal, F.M., Meng, C.A., Murakami, K., Kornberg, R.D., Block, S.M., (2015). Real-time observation of the initiation of RNA polymerase II transcription. *Nature*, **525**, 274–277. <https://doi.org/10.1038/nature14882>.
 22. Malik, I., Qiu, C., Snively, T., Kaplan, C.D., (2017). Widespread and unexpected consequences of altered Pol II catalytic activity in vivo. *Nucleic Acids Res.*, **45**, gkx037. <https://doi.org/10.1093/nar/gkx037>.
 23. Kiianitsa, K., Solinger, J.A., Heyer, W.-D., (2003). NADH-coupled microplate photometric assay for kinetic studies of ATP-hydrolyzing enzymes with low and high specific activities. *Anal. Biochem.*, **321**, 266–271. [https://doi.org/10.1016/s0003-2697\(03\)00461-5](https://doi.org/10.1016/s0003-2697(03)00461-5).
 24. Khaki, A.R., Field, C., Malik, S., Niedziela-Majka, A., Leavitt, S.A., Wang, R., Hung, M., Sakowicz, R., Brendza, K.M., Fischer, C.J., (2010). The macroscopic rate of nucleic acid translocation by hepatitis C virus helicase NS3h is dependent on both sugar and base moieties. *J. Mol. Biol.*, **400**, 354–378. <https://doi.org/10.1016/j.jmb.2010.04.065>.
 25. Titov, D.V., Gilman, B., He, Q.-L., Bhat, S., Low, W.-K., Dang, Y., Smeaton, M., Demain, A.L., Miller, P.S., Kugel, J. F., Goodrich, J.A., Liu, J.O., (2011). XPB, a subunit of TFIIH, is a target of the natural product triptolide. *Nat. Chem. Biol.*, **7**, 182–188. <https://doi.org/10.1038/nchembio.522>.
 26. Rudolf, J., Rouillon, C., Schwarz-Linek, U., White, M.F., (2010). The helicase XPD unwinds bubble structures and is not stalled by DNA lesions removed by the nucleotide excision repair pathway. *Nucleic Acids Res.*, **38**, 931–941. <https://doi.org/10.1093/nar/gkp1058>.
 27. He, Q., Titov, D.V., Li, J., Tan, M., Ye, Z., Zhao, Y., Romo, D., Liu, J.O., (2015). Covalent modification of a cysteine residue in the XPB subunit of the general transcription factor TFIIH through single epoxide cleavage of the transcription inhibitor triptolide. *Angewandte Chemie Int. Ed.*, **54**, 1859–1863. <https://doi.org/10.1002/anie.201408817>.
 28. Yan, C., Dodd, T., He, Y., Tainer, J.A., Tsutakawa, S.E., Ivanov, I., (2019). Transcription preinitiation complex structure and dynamics provide insight into genetic diseases. *Nat. Struct. Mol. Biol.*, **26**, 397–406. <https://doi.org/10.1038/s41594-019-0220-3>.
 29. Hall, M.C., Matson, S.W., (1999). Helicase motifs: the engine that powers DNA unwinding. *Mol. Microbiol.*, **34**, 867–877. <https://doi.org/10.1046/j.1365-2958.1999.01659.x>.
 30. Fischer, C.J., Saha, A., Cairns, B.R., (2007). Kinetic model for the ATP-dependent translocation of *Saccharomyces cerevisiae* RSC along double-stranded DNA. *Biochemistry-US*, **46**, 12416–12426. <https://doi.org/10.1021/bi700930n>.
 31. Young, M.C., Kuhl, S.B., von Hippel, P.H., (1994). Kinetic theory of ATP-driven translocases on one-dimensional polymer lattices. *J. Mol. Biol.*, **235**, 1436–1446. <https://doi.org/10.1006/jmbi.1994.1099>.
 32. Murakami, K., Mattei, P.-J., Davis, R.E., Jin, H., Kaplan, C. D., Kornberg, R.D., (2015). Uncoupling promoter opening from start-site scanning. *Mol. Cell.*, **59**, 1–7. <https://doi.org/10.1016/j.molcel.2015.05.021>.
 33. Luo, J., Cimermancic, P., Viswanath, S., Ebmeier, C.C., Kim, B., Dehecq, M., Raman, V., Greenberg, C.H., Pellarin, R., Sali, A., Taatjes, D.J., Hahn, S., Ranish, J., (2015).

- Architecture of the human and yeast general transcription and DNA repair factor TFIIH. *Mol. Cell.*, **59**, 794–806. <https://doi.org/10.1016/j.molcel.2015.07.016>.
34. Nogales, E., Louder, R.K., He, Y., (2017). Structural insights into the eukaryotic transcription initiation machinery. *Annu. Rev. Biophys.*, **46**, 59–83. <https://doi.org/10.1146/annurev-biophys-070816-033751>.
35. Greber, B.J., Toso, D.B., Fang, J., Nogales, E., (2019). The complete structure of the human TFIIH core complex e44771 *Elife*, **8** <https://doi.org/10.7554/elife.44771>.
36. Kovic, G., Chernev, A., Tegunov, D., Dienemann, C., Urlaub, H., Cramer, P., (2019). Structural basis of TFIIH activation for nucleotide excision repair. *Nat. Commun.*, **10**, 2885. <https://doi.org/10.1038/s41467-019-10745-5>.
37. Gradia, S.D., Ishida, J.P., Tsai, M.-S., Jeans, C., Tainer, J. A., Fuss, J.O., (2017). Chapter one MacroBac: New technologies for robust and efficient large-scale production of recombinant multiprotein complexes. *MethodsEnzymol.*, **592**, 1–26. <https://doi.org/10.1016/bs.mie.2017.03.008>.
38. Meek, K., Lees-Miller, S.P., Modesti, M., (2012). N-terminal constraint activates the catalytic subunit of the DNA-dependent protein kinase in the absence of DNA or Ku. *Nucleic Acids Res.*, **40**, 2964–2973. <https://doi.org/10.1093/nar/gkr1211>.
39. Fant, C.B., Levandowski, C.B., Gupta, K., Maas, Z.L., Moir, J., Rubin, J.D., Sawyer, A., Esbin, M.N., Rimel, J.K., Luyties, O., Marr, M.T., Berger, I., Dowell, R.D., Taatjes, D. J., (2020). TFIID enables RNA polymerase II promoter-proximal pausing. *Mol. Cell.*, **78**, 785–793.e8. <https://doi.org/10.1016/j.molcel.2020.03.008>.
40. Fishburn, J., Hahn, S., (2012). Architecture of the yeast RNA polymerase II open complex and regulation of activity by TFIIF. *Mol. Cell. Biol.*, **32**, 12–25. <https://doi.org/10.1128/mcb.06242-11>.
41. Ebmeier, C.C., Erickson, B., Allen, B.L., Allen, M.A., Kim, H., Fong, N., Jacobsen, J.R., Liang, K., Shilatfard, A., Dowell, R.D., Old, W.M., Bentley, D.L., Taatjes, D.J., (2017). Human TFIIH kinase CDK7 regulates transcription-associated chromatin modifications. *Cell. Rep.*, **20**, 1173–1186. <https://doi.org/10.1016/j.celrep.2017.07.021>.
42. Williams, D.J., Hall, K.B., (2000). Monte Carlo applications to thermal and chemical denaturation experiments of nucleic acids and proteins. *MethodsEnzymol.*, **321**, 330–352. [https://doi.org/10.1016/s0076-6879\(00\)21201-9](https://doi.org/10.1016/s0076-6879(00)21201-9).