

Review Paper

Cite this article: Pihl E *et al.* (2021). Ten new insights in climate science 2020 – a horizon scan. *Global Sustainability* 4, e5, 1–18. <https://doi.org/10.1017/sus.2021.2>

Received: 28 October 2020
Revised: 16 December 2020
Accepted: 13 January 2021




Key words:

climate anxiety; climate feedbacks; climate governance; climate impacts; climate litigation; climate mitigation; climate models; climate policy; environmental economics; future earth; risk governance; thermokarst; urban transformations; water stress

Author for correspondence:

Erik Pihl, E-mail: erik.pihl@futureearth.org

Ten new insights in climate science 2020 – a horizon scan

Erik Pihl¹ , Eva Alfredsson², Magnus Bengtsson³, Kathryn J. Bowen⁴, Vanesa Cástan Broto⁵, Kuei Tien Chou⁶, Helen Cleugh⁷, Kristie Ebi⁸, Clea M. Edwards⁹ , Eleanor Fisher¹⁰, Pierre Friedlingstein¹¹, Alex Godoy-Faúndez¹² , Mukesh Gupta¹³, Alexandra R. Harrington^{14,15,16}, Katie Hayes¹⁷, Bronwyn M. Hayward^{18,19}, Sophie R. Hebden¹, Thomas Hickmann²⁰, Gustaf Hugelius²¹, Tatiana Ilyina²², Robert B. Jackson^{23,24}, Trevor F. Keenan^{25,26}, Ria A. Lambino²⁷, Sebastian Leuzinger²⁸, Mikael Malmaeus²⁹, Robert I. McDonald³⁰, Celia McMichael³¹, Clark A. Miller³², Matteo Muratori³³, Nidhi Nagabhatla^{34,35}, Harini Nagendra³⁶, Cristian Passarello³⁷, Josep Penuelas³⁸, Julia Pongratz^{22,39}, Johan Rockström⁴⁰, Patricia Romero-Lankao^{41,42}, Joyashree Roy^{43,44}, Adam A. Scaife^{45,46}, Peter Schlosser⁴⁷, Edward Schuur^{48,49}, Michelle Scobie⁵⁰, Steven C. Sherwood⁵¹, Giles B. Sioen^{52,53}, Jakob Skovgaard⁵⁴, Edgardo A. Sobenes Obregon⁵⁵, Sebastian Sonntag⁵⁶, Joachim H. Spangenberg⁵⁷, Otto Spijkers⁵⁸, Leena Srivastava⁵⁹, Detlef B. Stammer⁶⁰, Pedro H. C. Torres⁶¹, Merritt R. Turetsky⁶², Anna M. Ukkola⁶³, Detlef P. van Vuuren^{20,64}, Christina Voigt⁶⁵, Chadia Wannous^{66,67} and Mark D. Zelinka⁶⁸

¹Future Earth Global Secretariat, Stockholm, Sweden; ²Division of Sustainable Development and Environmental Science and Engineering, KTH Royal Institute of Technology, Stockholm, Sweden; ³Hot or Cool Institute, Berlin, Germany; ⁴Climate Change Institute, Australian National University, Canberra, Australia; ⁵Urban Institute, University of Sheffield, Sheffield, UK; ⁶Risk Society and Policy Research Center, National Taiwan University, Taipei, Taiwan; ⁷Commonwealth Scientific and Industrial Research Organisation (CSIRO), Canberra, Australia; ⁸Department of Global Health, Center for Health and the Global Environment, University of Washington, Seattle, USA; ⁹Arizona State University, Global Futures Laboratory, Tempe, USA; ¹⁰The Nordic Africa Institute, Uppsala, Sweden; ¹¹College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter, UK; ¹²Sustainability Research Centre, Universidad del Desarrollo, Santiago de Chile, Chile; ¹³Department of Environmental Sciences, Asian University for Women, Chittagong, Bangladesh; ¹⁴Center for International Sustainable Development Law, Montreal, Canada; ¹⁵Fulbright Canada Foundation, Ottawa, Canada; ¹⁶Albany Law School, Albany, USA; ¹⁷Health Canada, Ottawa, Canada; ¹⁸Department of Political Science, University of Canterbury, Christchurch, New Zealand; ¹⁹The Centre for the Understanding of Sustainable Prosperity, University of Surrey, Guildford, UK; ²⁰Copernicus Institute of Sustainable Development, Utrecht University, Utrecht, The Netherlands; ²¹Department of Physical Geography, Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden; ²²Max Planck Institute for Meteorology, Hamburg, Germany; ²³Earth System Science Department, Woods Institute for the Environment, Stanford University, Stanford, USA; ²⁴Precourt Institute for Energy, Stanford University, Stanford, USA; ²⁵Department of Environmental Science, Policy and Management, UC Berkeley, Berkeley, USA; ²⁶Earth and Environmental Science Area, Lawrence Berkeley National Lab., Berkeley, USA; ²⁷Future Earth Asia Regional Center, Research Institute for Humanity and Nature, Kyoto, Japan; ²⁸Auckland University of Technology, School of Science, Auckland, New Zealand; ²⁹IVL Swedish Environmental Research Institute, Stockholm, Sweden; ³⁰Center for Sustainability Science, The Nature Conservancy, Arlington, USA; ³¹School of Geography, The University of Melbourne, Melbourne, Australia; ³²College of Global Futures, Arizona State University, Tempe, USA; ³³National Renewable Energy Laboratory, Golden, USA; ³⁴United Nations University, Institute for Water, Environment and Health (UNU INWEH), Hamilton, Canada; ³⁵McMaster University, Hamilton, Canada; ³⁶Center for Climate Change and Sustainability, Azim Premji University, Bangalore, India; ³⁷Future Earth Global Secretariat, Paris, France; ³⁸Global Ecology Unit CREAF-CSIC-UAB, CSIC and CREAF, Madrid, Spain; ³⁹Department of Geography, Ludwig-Maximilians-Universität Munich, Munich, Germany; ⁴⁰University of Potsdam, Institute for Climate Impact Research, Potsdam, Germany; ⁴¹Center for Integrated Mobility Sciences, National Renewable Energy Laboratory, Golden, USA; ⁴²Mansueto Institute for Urban Innovation, University of Chicago, Chicago, USA; ⁴³Department of Energy, Environment and Climate Change, Asian Institute of Technology, Klong Nueng, Thailand; ⁴⁴Department of Economics, Jadavpur University, Kolkata, India; ⁴⁵Met Office Hadley Centre, Exeter, UK; ⁴⁶Department of Mathematics, University of Exeter, Exeter, UK; ⁴⁷Julie Ann Wrigley Global Futures Laboratory, Global Institute of Sustainability and Innovation, Arizona State University, Tempe, USA; ⁴⁸Center for Ecosystem Science and Society, Northern Arizona University, Flagstaff, USA; ⁴⁹Department of Biological Sciences, Northern Arizona University, Flagstaff, USA; ⁵⁰Institute of International Relations, The University of the West Indies, Saint Augustine, Trinidad & Tobago; ⁵¹Climate Change Research Centre, UNSW Sydney, Sydney, Australia; ⁵²Future

© The Author(s), 2021. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

Earth Global Secretariat, Tokyo, Japan; ⁵³National Institute for Environmental Studies, Tsukuba, Japan; ⁵⁴Department of Political Science, Lund University, Lund, Sweden; ⁵⁵Independent Consultant in Public International Law, The Hague, The Netherlands; ⁵⁶Climate Service Center Germany (GERICS), Helmholtz-Zentrum Geesthacht, Hamburg, Germany; ⁵⁷SERI Germany, Sustainable Europe Research Institute, Cologne, Germany; ⁵⁸China Institute of Boundary and Ocean Studies (CIBOS), Research Institute of Environmental Law (RIEL) and International Water Law Academy (IWLA), Wuhan University, Wuhan, China; ⁵⁹International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria; ⁶⁰Centrum für Erdsystemforschung und Nachhaltigkeit (CEN), Universität Hamburg, Hamburg, Germany; ⁶¹Institute of Energy and Environment, University of São Paulo, São Paulo, Brazil; ⁶²Ecology and Evolutionary Biology Department, Institute of Arctic and Alpine Research, University of Colorado Boulder, Boulder, USA; ⁶³ARC Centre of Excellence for Climate Extremes and Research, School of Earth Sciences, Australian National University, Canberra, Australia; ⁶⁴Department of Climate, Air and Energy, PBL Netherlands Environmental Assessment Agency, Den Haag, The Netherlands; ⁶⁵Department of Public and International Law, University of Oslo, Oslo, Norway; ⁶⁶Towards A Safer World Network (TASW), Stockholm, Sweden; ⁶⁷Future-Earth Health Knowledge Action Network, Stockholm, Sweden and ⁶⁸Lawrence Livermore National Laboratory, Livermore, USA

Non-technical summary. We summarize some of the past year's most important findings within climate change-related research. New research has improved our understanding of Earth's sensitivity to carbon dioxide, finds that permafrost thaw could release more carbon emissions than expected and that the uptake of carbon in tropical ecosystems is weakening. Adverse impacts on human society include increasing water shortages and impacts on mental health. Options for solutions emerge from rethinking economic models, rights-based litigation, strengthened governance systems and a new social contract. The disruption caused by COVID-19 could be seized as an opportunity for positive change, directing economic stimulus towards sustainable investments.

Technical summary. A synthesis is made of ten fields within climate science where there have been significant advances since mid-2019, through an expert elicitation process with broad disciplinary scope. Findings include: (1) a better understanding of equilibrium climate sensitivity; (2) abrupt thaw as an accelerator of carbon release from permafrost; (3) changes to global and regional land carbon sinks; (4) impacts of climate change on water crises, including equity perspectives; (5) adverse effects on mental health from climate change; (6) immediate effects on climate of the COVID-19 pandemic and requirements for recovery packages to deliver on the Paris Agreement; (7) suggested long-term changes to governance and a social contract to address climate change, learning from the current pandemic; (8) updated positive cost–benefit ratio and new perspectives on the potential for green growth in the short- and long-term perspective; (9) urban electrification as a strategy to move towards low-carbon energy systems and (10) rights-based litigation as an increasingly important method to address climate change, with recent clarifications on the legal standing and representation of future generations.

Social media summary. Stronger permafrost thaw, COVID-19 effects and growing mental health impacts among highlights of latest climate science.

1. Introduction

From mental health distress to severe water crisis, societies around the world are experiencing the impacts of human-induced climate

change. In 2020, alongside these impacts, we experienced the COVID-19 pandemic. COVID-19 has served to expose our societal vulnerabilities while also providing unique opportunities to act for a fair and climate-friendly world. With research supporting the need to develop carbon-neutral societies by 2050 to safely achieve the goals of the Paris Agreement, the need for transformative change is urgent (Head, 2020; Otto *et al.*, 2020). Enhanced understanding of the challenges facing Earth's systems – and of the social and economic consequences – contributes to identifying appropriate action.

In this paper, we carry out a horizon scan of important insights emerging from advances in integrated research related to climate change over the past year (focusing on findings published in 2019–2020). The objective of this horizon scan is two-fold. First, through expert elicitation we attempt to identify the 10 most important new scientific insights over the past year. Second, this horizon scan constitutes an effort to provide an integrated synthesis of key research outputs and how these add up into broader science-based insights that should guide climate policy. This scientific horizon scan forms the basis of a wider research synthesis report on the 10 New Insights in Climate Science (10NICS) produced annually and officially handed over to the United Nations Framework Convention on Climate Change Secretariat in connection with the Conferences of Parties. Taken together, we are not claiming this to be a top-10 climate science ranking, but rather an effort of scanning the wide interdisciplinary arena of climate research and identifying key insights – and which provide evidence that is of critical importance for evidence-based policymaking.

The 10 insights begin by considering climate modelling advances and the improvements in our understanding of climate sensitivity, and regional climate predictability. In doing so, we are better placed to understand future risks and to plan for change. We then draw attention to evidence on thawing permafrost in the Arctic, which stores one-third of the world's soil carbon in a location that is responding quickly to climate change. We turn to carbon uptake by land sinks, which respond to anthropogenic change with consequences for their potential to mitigate carbon emissions. We consider how climate change will exacerbate the water crises already felt in many places, underlining how impacts depend on and contribute to social inequality. We also bring to the fore growing evidence that changing climatic conditions are adversely affecting mental health, an issue garnering attention in 2020 as it is exacerbated by the COVID-19 pandemic. We reflect on the most urgent task for a post-COVID-19 era, namely making 2020 a turning point for reduction in global greenhouse gas (GHG) emissions. How our economies and societies contribute to emissions and the changes that can occur has been brought into stark relief by the lockdowns initiated to control spread of the pandemic. Following that, again in response to the pandemic, we consider the potential for COVID-19 to catalyse a new social compact for a just and climate-friendly world through strengthening inclusive forms of governance. We also underline how greening the economy through sustainable investments is cost effective and gives substantial co-benefits. This is vital given evidence showing that a primary focus on economic growth, which puts climate mitigation as a secondary goal, jeopardizes our last chance of achieving the Paris Agreement. From this we turn to energy, outlining evidence that shows how urban electrification provides a strategy to move towards low-carbon sustainable energy systems. Finally, our last insight draws attention to rights-based litigation, which clarifies the international legal

standing and representation of the rights and interests of future generations in a healthy environment.

2. Methodology

The horizon scan has been overseen by an expert panel 'Editorial Board' with 10 researchers appointed by Future Earth, The Earth League and World Climate Research Programme (WCRP). The 10 'insights' were identified through an expert elicitation process beginning with an open call for inputs, through an open-ended questionnaire for suggesting new topics. A link to the form was sent directly to 221 international experts covering a broad array of disciplines. It was also distributed to members of The Earth League, WCRP (Secretariat, Joint Scientific Committee, Core Project Chairs and Grand Challenge leaders), to international project offices and development teams of Future Earth Global Research Projects and Future Earth Knowledge-Action Networks, to Future Earth National Committees and Networks and posted on the Future Earth Open Network and Future Earth website. The questions posed to questionnaire respondents were: 'What [do] you think are the 1–3 most important new discoveries or advancements in your overarching field of research since 1st July 2019 and the key articles and reports highlighting them[?]'.

The questionnaire resulted in 73 individual responses suggesting 128 topics. Additional 18 topics were suggested by 11 researchers via email, of which eight were unique respondents who had not answered the questionnaire. The suggested topics were summarized in 20 candidate 'insights'. The Editorial Board identified the 10 insights that best satisfied the requirements for novelty, relevance and sufficient scientific evidence. Each insight was written by two or more experts selected from the questionnaire and the Future Earth, The Earth League and WCRP networks, based on qualifications and quality of topic suggestions. All authors were approved by the Editorial Board.

Further details on methodology can be found in Supplementary materials.

3. New insights

3.1. Climate sensitivity and predictability are now better understood

At the centre of international climate change negotiations are the rising concentration of carbon dioxide (CO₂) in the atmosphere. CO₂ is the most significant anthropogenic GHG being emitted into the atmosphere, reducing emissions of terrestrial radiation to space and causing global temperatures to rise. Although this understanding pre-dates the 20th century, the quantitative relationship between CO₂ levels and global warming has remained uncertain for decades, hampering efforts to understand future risks and plan for change.

The 'likely range' (at least a 66% chance of being within this range) of equilibrium climate sensitivity – the long-term global rise in air temperature expected as a result of doubling atmospheric CO₂ concentrations – was estimated to be 1.5–4.5°C by Intergovernmental Panel on Climate Change (IPCC, 2013) in its Fifth Assessment Report (AR5); these figures remained unchanged since the Charney report of 1979.

Larger climate sensitivity is suggested by global-scale climate change experiments carried out using the latest Earth System Models and coordinated under the Coupled Model




Fig. 1. Climate sensitivity ranges from recent sources. First entry shows 66% (thick bar) and 90% (thin bar) probability ranges from Sherwood et al. (2020), with hatched extensions bounding the span of these ranges under plausible alternative assumptions. The second bar shows the 66%-or-greater (thick bar) and 90%-or-greater (thin bar) probability ranges from IPCCs AR5 report in 2013. The third and fourth bars show the full span of values predicted by the previous and current generation of global climate models respectively. All values are 'effective' climate sensitivities except that IPCCs is formally given as an equilibrium value; the particular definition has a 5%-or-less impact on probability ranges (Sherwood et al., 2020).

Intercomparison Project Phase 6 (CMIP6), which exhibit sensitivity values ranging from 1.8 to 5.6°C. The values of 10 models exceeded the upper end of the aforementioned likely range. The higher climate sensitivity seen in many models is due primarily to stronger amplifying cloud feedbacks from low clouds at middle and high latitudes (Flynn & Mauritsen, 2020; Zelinka et al., 2020). This may be related to improvements in how models decide whether cold clouds are made of liquid or ice water, a difficult problem (Bodas-Salcedo et al., 2019; Gettelman et al., 2019). But even though *a priori* these changes seem to be improvements, many high-sensitivity models overestimate recent warming trends (Nijse et al., 2020; Tokarska et al., 2020), suggesting that the higher sensitivity models should be treated with caution.

Indeed, the likely range of climate sensitivity has now been narrowed to 2.3–4.5°C by a new, comprehensive WCRP analysis of the broader evidence (Sherwood et al., 2020). This took a three-pronged approach of examining climate feedback processes, the historical record and the palaeoclimate record, which all provided evidence against the high model climate sensitivities (Sherwood et al., 2020). In particular, they find that sensitivities above 4.5°C are hard to reconcile with palaeoclimate evidence, also noted by Zhu et al. (2020). On the other hand, Sherwood et al. (2020) find that the likely range does not extend below 2.3°C, which discounts the lower end of the IPCC AR5 range. This conclusion was supported by all lines of evidence and indicates that moderate emissions reduction scenarios are less likely to meet the Paris temperature targets than previously anticipated (Figure 1).

On regional scales, climate models are also becoming better at simulating temperature and hydrological extremes (Di Luca et al., 2020; Kim et al., 2020), including the intensity of heavy rainfall events (Kim et al., 2020) and hot and cold extremes (Di Luca et al., 2020). Models are now able to simulate rainfall droughts well, particularly at the seasonal scale and the projections of drought duration and frequency are becoming more consistent

over many regions (Ukkola et al., 2020). Although regional changes in mean rainfall remain uncertain, this provides new opportunities for water resource management.

In the near term, climate models are better able to predict the observed evolution of regional climate than previously thought possible, particularly around the Atlantic Basin. Decadal predictions of the atmospheric circulation and regional temperature and rainfall all now show encouraging levels of skill and this offers great promise for the utility of regional climate predictions. However, climate models also show a spuriously low ratio of predictable signal strength to internal noise variability. This means that newfound decadal prediction skill can only be realized by averaging large ensembles of hundreds of simulations and it could affect the attribution and prediction of quantitative changes in extratropical climate using current models (Scaife & Smith, 2018; Smith et al., 2020).

3.2. Greenhouse gas emissions from permafrost will be larger due to abrupt thaw processes

Thawing permafrost in the Arctic is expected to release significant quantities of GHGs over the coming decades, enough to merit consideration in climate negotiations. Recent research shows it will be larger than earlier projections due to abrupt permafrost thaw processes (Turetsky et al., 2020).

Permafrost is a perpetually frozen layer beneath the seasonally thawed surface layer of the ground. The northern permafrost region covers 18 million km² and stores 1460–1600 petagrams of carbon (PgC) – one-third of the world's soil carbon (Meredith et al., 2019). The Arctic is responding quickly to climate change, with air temperatures warming more than twice as fast as the global average. Unusually warm summers – such as the record-breaking 2020 heatwave in Siberia and Svalbard – are happening more often (Ciavarella et al., 2020). This is causing Arctic permafrost to thaw in some northern regions almost a century earlier than some climate models projected (Farquharson et al., 2019).

Abrupt permafrost thaw happens when melting ground ice causes the ground surface above to collapse. This liberates previously frozen soil carbon, creating a so-called 'thermokarst' landscape of slumps and gullies in upland areas and collapse-scar wetlands and lakes in less well-drained areas. Satellite observations of these landscape-scale changes have shown an acceleration in abrupt thaw processes over the past two decades; they are expected to substantially increase this century as climate warms (Lewkowicz & Way, 2019).

Although climate models do include gradual permafrost thaw, they do not include the more complex thermokarst-inducing processes. When thermokarst is included, by the year 2100 up to three times more carbon becomes exposed assuming a moderate emission scenario at Representative Concentration Pathway (RCP) 4.5 and up to 12 times more carbon is exposed under a high emission scenario of RCP8.5 (Nitzbon et al., 2020).

Abrupt permafrost thaw increases thaw rates and also causes ecosystem shifts to conditions more conducive to producing strong GHG emissions, notably methane. The IPCC Special Report 1.5 estimated 27 PgC of cumulative carbon emissions from permafrost thaw and wetlands by 2100 for low emission scenarios (where PgC is carbon loss in CO₂ equivalents). The more recent studies indicate that under moderate and high emission scenarios (RCP4.5–RCP8.5), emissions from abrupt thaw processes would approximately double the projected cumulative

carbon emissions compared to estimates of gradual thaw alone (Gasser et al., 2018; Turetsky et al., 2020). Increased losses through abrupt thaw may also apply to emission scenarios consistent with 1.5- or 2-degree warming targets but these more aggressive climate change mitigation pathways could halve abrupt thaw carbon losses compared to high emission pathways (Figure 2).

Peatlands have year-round waterlogged conditions that slow plant decomposition, allowing peat to accumulate – one of the largest natural carbon stores on land. Nearly half of northern peatlands are underlain by permafrost. Abrupt thaw could shift the entire northern hemisphere peatland carbon sink into a net source of global warming, dominated by methane, lasting several centuries (Hugelius et al., 2020).

Most of the methane emissions from thawing permafrost are fuelled by recently stored carbon, rather than carbon sequestered thousands of years ago (Dean, 2020; Dean et al., 2020). A study of atmospheric methane over the past million years of Earth's history, using ice cores from Antarctica, found no evidence for substantial releases of methane due to the destabilization of old permafrost carbon stores (Dyonisius et al., 2020). This is because when methane is produced at depth in thawing soils or sediments, microorganisms living in the soil or water columns above oxidize most of the methane before it reaches the surface, instead releasing it as CO₂ (Dean, 2020).

An ecological feedback associated with permafrost thaw that is not yet included in global climate models is a priming effect on soil respiration, caused by an increase in root activity. This amplifies soil carbon loss, with an additional 40 PgC loss projected from Arctic permafrost by 2100 for RCP8.5 (Keuper et al., 2020).

In summary, when adding new knowledge on abrupt thaw to what's currently modelled for gradual thaw, the expected carbon emissions from permafrost could as much as double by year 2100. The carbon emissions from permafrost regions could be even higher when including effects on root activity which increase soil decomposition. Accounting for these effects will impose tighter restrictions on the remaining anthropogenic carbon emission budgets.

3.3. Carbon uptake by land sinks – potentials and limits

Land ecosystems remove about 30% of the CO₂ emitted through fossil fuels and land-use change (LUC) emissions, an ecosystem service referred to as the '(natural) land sink' (see Figure 3; Friedlingstein et al., 2019). This serves to slow the growth rate of atmospheric CO₂, and consequently reduces the rate of climate change. The amount of CO₂ absorbed by the land has also increased rapidly over the past few decades, likewise to anthropogenic CO₂ emissions it has more than doubled since 1960 (Friedlingstein et al., 2019) with extensive greening reported (Piao et al., 2020) as well as large associated changes in the effect vegetation has on local and global climate (Forzieri et al., 2020). The increased land sink has occurred despite an increased prevalence of large-scale natural disruptions to ecosystems (McDowell et al., 2020) and evidence that some of the largest carbon sinks of the planet have already saturated (Hubau et al., 2020). Its increase is stronger than changes in emissions from LUCs (Friedlingstein et al., 2019) but largely undermined by the impact of LUC on tropical ecosystems (Tagesson et al., 2020). The natural land sink is not constant, however, and responds directly to environmental changes, such as heatwaves and droughts (Bastos et al., 2020), and anthropogenic interventions such as deforestation



Fig. 2. Thawing coastal permafrost in Arctic Canada with person for scale. Credit: G. Hugelius.

and LUC (Brando et al., 2020). The dynamic nature of terrestrial carbon uptake makes understanding the regional hotspots of source or sink potential – and the processes that dictate the likelihood of continued increased uptake in those regions – essential for adequate policy design.

CO₂ fertilization is widely reported to be the primary cause of the increased land sink (Tharammal et al., 2019; Walker et al., 2020). Rising atmospheric CO₂ increases leaf-scale photosynthesis and resource-use efficiencies, which can lead to increased plant growth, vegetation biomass and soil organic matter. However, due to the complexity and heterogeneity of ecosystems, the resulting impact of CO₂ on carbon uptake is context dependent. Particularly, nutrient availability constrains the ability of global ecosystems to translate increased photosynthesis into increased biomass and thus carbon storage (Terrer et al., 2019). The CO₂ fertilization and other effects beneficial for carbon uptake are further offset by the detrimental impact of warming on soil carbon (Vaughn & Torn, 2019) and permafrost (Wang et al., 2020a), and regional increases in forest mortality due to changes in the frequency of extreme events (McDowell et al., 2020). A recent report suggests that CO₂ fertilization effects on vegetation photosynthesis are globally declining as a result of these and other offsetting factors such as water and nutrient limitations (Wang et al., 2020c).

The processes that offset CO₂ fertilization are highly regionally specific, and emerging evidence suggests that many tropical

regions are at or near sink saturation (Hubau et al., 2020), while boreal and temperate zones continue to increase their sink capacity (Tagesson et al., 2020). LUC impacts explain much of the regional differences with deforestation in tropical regions (Brando et al., 2020) and increased wood harvesting in Europe (Ceccherini et al., 2020). Moreover, unprecedented carbon losses also occurred due to fires in Australia, California, the Amazon and the Arctic, with fire impacts predicted to worsen as a result of anthropogenic climate change (Bowman et al., 2020; Witze 2020). Although results for the world's drylands are currently inconclusive, recent reports suggest that previous long-term aridity-change projections overestimated dryland aridification (Yang et al., 2019).

Several knowledge gaps exist regarding the future potential of the land sink to offset carbon emissions. Although the effect of CO₂ on global ecosystem productivity is now widely acknowledged, the estimated magnitude of the effect spans an order of magnitude across studies (Walker et al., 2020), which greatly hinders the ability of models to project future expected changes. On large scales the natural land sink can be measured only concurrently with CO₂ sinks and sources due to land-use activities. Better quantification of the net LUC flux is thus key for a better understanding of the natural land sink. Land management is an important unknown, and practices that co-deliver food security, climate change mitigation and combat land-degradation and




Fig. 3. Long-term trajectories of the residual land sink, along with the atmospheric CO₂ growth rate and emissions from fossil fuel burning and land use. The inset attributes the long-term changes in the sink to the percentage contribution of CO₂ fertilization, LUC and other (e.g. N-deposition, ozone and phenology) factors (data from Tharammal *et al.*, 2019).

desertification are needed (Smith *et al.*, 2019). Although effective practices could potentially achieve up to 30% of mitigation targets needed to limit warming to 1.5°C (Roe *et al.*, 2019), proposed approaches based on widespread afforestation need to recognize the potential negative impacts of tree planting, such as habitat loss and interference with naturally treeless ecosystems (Bond *et al.*, 2019). Funding mechanisms for such natural solution approaches are also needed, with renewed calls for carbon tax strategies to support land-based mitigation (Barbier *et al.*, 2020).

In summary, although we currently see a substantial and slightly increasing land carbon sink, there is evidence of a weakening sink capacity as the effects of drought and warming start to outweigh CO₂ fertilization effects (Figure 3).

3.4. Climate change will severely exacerbate the water crisis

Climate change is already causing extreme events in many watersheds, impacting communities (Madakumbura *et al.*, 2019). Changes in extreme precipitation are likely to be stronger than changes in mean precipitation, with extreme events increasing in intensity and frequency (Myhre *et al.*, 2019). Extreme precipitation will increase over all climate regions, but with greater intensity in humid and semi-humid regions compared to semi-arid areas, with a corresponding change in flood risk – overall flood intensity is also projected to increase for most areas (Tabari *et al.*, 2019). Changes in precipitation impact spatio-temporal distribution and water availability, with seasonally variable rainfall regimes anticipated to become even more variable, whereas regimes with low seasonal variation will receive more rainfall in the monsoon (Konapala *et al.*, 2020). There is likely to be an increase in the aridity of 72% of land area which, even when accounting for vegetation response to the increased CO₂ levels, is expected to have deleterious effects on ecosystems and their ability to sustain life. This particularly affects the Middle East, North Africa, south Europe and Australia (Tabari *et al.*, 2019). Urbanization is further altering regional climate patterns – for instance increasing the magnitude and recurrence of extreme

precipitation events in large urban areas in China (Li *et al.*, 2019). Climate hazards will drive water scarcity due to physical shortage, or scarcity in access due to the failure of institutions to ensure a regular supply or because of a lack of adequate infrastructure (Empinotti *et al.*, 2019).

Extreme events are very important drivers of water crises, however, current practice in general circulation models may understate the potential for significant changes in the hydrological cycle including the risk of extreme events (Hamstead & Coseo, 2020; Lomba-Fernández *et al.*, 2019; Nicklin *et al.*, 2019), for instance by focusing on the ensemble mean and variance (Teegne & Melesse, 2020). Changes in extreme precipitation require greater attention in climate modelling and prediction research. There is greatest global uncertainty in tropical and subtropical regions because of a combination of the difficulty in modelling convective rainstorms and the sparsity of weather observation networks for model validation and refinement (Tabari *et al.*, 2019).

Climate change coupled with socioeconomic drivers can also impact water quality – for instance shifts in monsoon timings can lead to dilution or concentration of nitrogen, phosphorus and other pollutants (Whitehead *et al.*, 2019). Conversely, water quality and pollution levels can impact the ability of sensitive ecosystems such as coral reefs to recover from extreme climate events (MacNeil *et al.*, 2019).

The Cape Town water crisis has been a clear example of a water insecurity event that is indicative of how extreme climatic events are exacerbated by climate change. In 2018 Cape Town went through a severe water crisis as a result of a multi-year drought. Shepherd (2019) reviewed how the city responded to the threat of ‘Day Zero’ for the urban supply, the moment when the reservoirs might run dry. The water crisis in Cape Town has complex political and social ramifications, both reinforcing existing inequalities and increasing competition between water users, but also opening up new potentials for solidarity and collective action. Water conservation efforts, particularly the city’s creative campaign to reduce demand among residents and businesses, reduced the severity of water scarcity (Simpson *et al.*, 2020; Van Zyl & Jooste, 2020).

The impacts of water crises and climate risks are highly unequal, driven by social inequality (Craig *et al.*, 2019; Roshan & Kumar, 2020; WWAP, 2020). A review of water, sanitation and hygiene (WASH) and gender linkages shows that this highly unequal impact of inadequate water supply is the rule (Pouramin *et al.*, 2020). Inadequate WASH resources disproportionately affect women and girls, leading to negative health outcomes in 71% of the studies reviewed.

Finally, there is increasing policy recognition that water-related extreme events are also contributing to the migration and displacement of millions of people. A new United Nations (UN) report documents these cases and suggests that rather than trying to prevent climate-driven migration, the international policy community should begin considering migration as a potential adaptation strategy, one that can help in achievement of the Sustainable Development Goals (SDGs) (Nagabhatla *et al.*, 2020). Migration, urbanization and climate change are disruptors that can catalyse shifts in values towards water use and management (IPBES, 2019). Integrated climate change mitigation and adaptation strategies could be a win-win policy: it could concurrently combat both the causes and impacts of climate change, thus help tackle water crises and disaster risk in tandem (WWAP, 2020).

3.5. Climate change can profoundly affect our mental health

Climate change is contributing to increased injuries, illnesses and deaths, with health risks projected to increase as temperatures, precipitation and other climatic variables continue to change (Haines et al., 2014). There is growing evidence that changing climatic conditions are adversely affecting mental health including states of mental wellness, emotional resilience and psychosocial well-being (see e.g. Basu et al., 2018; Hanigan, Schirmer and Niyonsenga, 2018). These affects can become severe when people experience the consequences of cascading and compounding risks, such as heatwaves coincident with wildfires. Climate hazards can result in new or worsened stress and clinical disorders such as trauma, anxiety, post-traumatic stress disorder and depression (Hayes, Berry & Ebi, 2019; Middleton et al., 2020; Wu, Snell & Samji, 2020). Some studies describe increased risk of suicide related to exposure to warming temperatures (Burke et al., 2018b).

In 2016, it was estimated that mental and addictive disorders affected more than 1 billion people globally (Rehm & Shield, 2019) but accurate statistics are lacking. Growing public awareness of the current impacts and future risks of changing climate and weather patterns, wildfires, sea level rise and ocean acidification are increasing the prevalence of emotional responses, especially among youth concerned about the future (Clayton, 2020). Terms used to describe this phenomenon include eco-anxiety, biospheric concern and solastalgia (Cianconi, Betrò & Janiri, 2020). It is expected that rising sea levels and coastal erosion and other climate impacts will contribute to relocation, displacement and migration away from high-risk human settlements (McMichael et al., 2020; Palinkas & Wong, 2020). The associated disruption of community networks, livelihoods and place attachment can lead to heightened psychosocial risks (Hayes et al., 2020).

Tackling climate-related mental health issues requires proactive planning with (inter)national agreements, preparedness building activities but also displacement, migration and mental health support for those on the move or 'left behind' (Matias, 2020; Schwerdtle, Bowen & McMichael, 2018). Health-system resilience also needs to be strengthened to include mental health support for survivors of climate-related disasters, including the mental health impacts that can last years from living in temporary shelters over prolonged periods of time or enduring the lengthy reconstruction of settlements (Schwartz et al., 2017; Yokoyama et al., 2014). Figure 4 shows a comprehensive list with factors influencing mental health, and how mental health, well-being and emotional resilience can be improved (Hayes, Berry & Ebi, 2019). In order to better understand the risks to mental health arising from climate change, it is important to support transdisciplinary research and practice collaborations (Hayes, Berry & Ebi, 2019).

A large body of research identifies strategies for addressing mental health and improving emotional resilience (Hayes & Poland, 2018; Hayes et al., 2018). Such strategies will need to be harnessed and adapted to address current and future mental health risks and impacts of climate change. Concrete actions include communicating with individuals and populations about climate change and mental health; advocacy for GHG reductions and adaptation measures that enable populations to cope with, prepare for and respond to climatic risks. In this regard, governmental acknowledgment of mental ill-health as a worrying and increasing burden of disease is growing (McIver et al., 2016; Rehm & Shield, 2019).

Policies and measures to protect and strengthen blue and green spaces (i.e. visible waters and greenery, respectively) are important

as the ecosystem services they provide are associated with positive mental health and well-being outcomes (Bratman et al., 2019). For example, the presence of green space during childhood has been associated with better mental health later in life (Engemann et al., 2019). Likewise, short, frequent walks or time spent in blue spaces have proven mental health benefits (Vert et al., 2020). Ecosystem service assessments and policies, land-use decisions and climate change resilience plans need to include psychosocial well-being considerations. Such considerations are also fundamental components of climate-resilient development and have multiple benefits – for human health and the health of our natural environment.

In sum, climate change can profoundly affect mental health. Cascading and compounding risks are projected to increase, contributing to anxiety and distress. There are opportunities to address the mental health consequences of climate change, including by implementing and communicating effective mitigation and adaptation strategies (such as blue and green spaces), protecting ecosystems and biodiversity with resultant co-benefits for human health, as well as developing mental health support strategies.

This insight is further elaborated on in the Supplementary material.

3.6. Many governments are missing the opportunity to use COVID-19 recovery spending for decarbonization

In the first half of 2020, the COVID-19 pandemic led to widespread confinement and human mobility restrictions, resulting in economic contraction and reduced emissions of GHGs and air pollutants. For this period, global CO₂ emissions were estimated to decline by 8.8% compared to 2019 (Liu et al., 2020). This included a CO₂ emission decline of 17% at days of peak lockdown (Le Quéré et al., 2020) and a nitrogen oxides (NO_x) decline of 30% (Forster et al., 2020) in April. The transport sector was responsible for roughly half of the decline, while industry and the power sector yielded another 43%. Declines in single countries were even greater than the global total, averaging one-quarter at respective peak confinement. Significant air quality improvements were also observed, especially in urban areas, attributable to a reduction in car use, factory production and construction activities (Wang 2020b).

Despite large reductions during lockdowns, global carbon emissions have bounced back and are expected to decline by 'only' 7% in 2020 as a whole (Friedlingstein et al., 2020). Emissions from cars and other vehicles have returned and are close to 2019 levels as economies are opening up, while emissions from air travel are still down by almost half. To make 2020 a turning point in global emissions, the 7% emission reductions expected for 2020 will need to be repeated year on year to reach net zero by mid-century (Hepburn et al., 2020). Climate change response strategies with accelerated systemic changes in energy sources, technology, personal choices and additional policies (Barbier, 2020) are essential to stay on a low-carbon path.

Some major economies like the United States, Japan and Germany are implementing recovery packages amounting to nearly 15% of their GDP (Sovacool et al., 2020a). The size of these packages means that they can lock the world into more or less green trajectories. The global investment requirement for a Paris-compatible pathway has been estimated to be 1.4 trillion USD per year in the period 2020–2024, a modest sum compared

Factors that Influence the Psychosocial Health Impacts of Climate Change




Fig. 4. Factors that influence the psychosocial health impacts of climate change. A framework showing the mental health consequences of climate change and how these consequences are mediated by the social and ecological determinants of health, response interventions and factors that influence psychosocial adaptation when they are in place or when absent act as barriers to psychosocial adaptation. Adapted with permission from Hayes, Berry, and Ebi (2019).

to the global stimulus funds (Andrijevic et al., 2020) amounting to more than 12 trillion USD. Investments in areas like clean physical infrastructure, building efficiency retrofits, education and training, natural capital and clean R&D can achieve both economic revitalization and climate goals simultaneously (Engström et al., 2020; Gawel & Lehmann, 2020; Hepburn et al., 2020; Malliet et al., 2020). However, governments are not taking the opportunity to decarbonize about 3.7 trillion USD of stimulus funds being allocated to environmentally relevant sectors suitable for such green investments (Vivid Economics, 2020). Instead, G20 governments are committing 233 billion USD to fossil fuel-based ('brown') activities, compared to only 146 billion USD for green activities, as of November 2020 (SEI et al.,

2020). This will lock in brown activities for years or even decades (Barbier, 2020; Hepburn et al., 2020) and also reinforce the power structures favouring fossil fuel companies, including their ability to hinder climate policy (Kuzemko et al., 2020; Mildenberger, 2020).

Although the GHG emission reductions caused by mobility restrictions were mostly temporary, governmental economic recovery efforts invested in low-carbon solutions could reduce global warming by 0.3°C by 2050 and put the world on track to meet the Paris Agreement goals (Forster et al., 2020). Unfortunately, based on the stimulus plans announced at time of as of this writing, most governments are still on crisis mode and so far appear to be missing this unique and critically

important opportunity for green investments (Climate Action Tracker, 2020). The period following the containment of the pandemic, when additional recovery packages will be designed and released, will be crucial for the global climate.

3.7. The COVID-19 pandemic demonstrates the need for a new social compact for a just and climate-friendly world

The world has a unique opportunity to reshape the future in new directions. COVID-19, coupled with divided and divisive national responses, abruptly exposed the weaknesses of international cooperation in an era of climate crisis (Oldekop et al., 2020). COVID-19 has laid bare governance deficiencies in many countries and led to a disruptive new normal – the tackling of which requires transformative strategies and collaborations. COVID-19 and climate change are transboundary risks that affect all regions indiscriminately. No government, community or company can unilaterally address the systemic risks posed by COVID-19 or climate change to human well-being and economic and environmental security. Finding new ways to act together is crucial, and this requires the strengthening of capacity and inclusive forms of governance.

Systemic risks will continue to grow (Renn et al., 2019). Throughout 2020, climate disasters and actions to address COVID-19 have together imposed difficult economic and social hardships around the world, especially on marginalized communities, in many cases increasing inequality (Howarth et al., 2020). Yet, at the same time, responses to COVID-19, coupled with activism by social movements surrounding climate change, opens up new possibilities for transformation and underlines the need to develop a new global social compact for a more just and sustainable future (Dixon-Decleve et al., 2020). People everywhere are increasingly aware of their vulnerabilities to emergent transnational risks and the threats they pose to global systems and supply chains for energy, food, water, transport and material goods (Laborde et al., 2020). Short-term political expediency is being challenged as communities demand effective, long-term and just solutions to global risks, particularly for the most vulnerable (Leach et al., 2018).

In 2020, the climate and pandemic crises catalysed the emergence of an informal, yet increasingly powerful global commitment to change:

- (1) Youth, labour and indigenous climate movements redoubled their commitments to creating and sharing knowledge and pressuring governments and the private sector to act decisively, even in the face of significant hurdles (Hayward, 2020; Whyte, 2020). In July, the United Nations created the Youth Advisory Group on Climate Change.
- (2) Public health researchers, the private sector and health officials urgently collaborated to develop effective responses to COVID-19 (Rourke et al., 2020), and the climate science community continued to advocate for action to address climate change and other systemic risks.
- (3) Transnational networks of businesses, cities, regions and countries collaborated to fight COVID-19 and to set targets and develop strategies for achieving carbon-neutral or even carbon-negative economies by mid-century (Bai et al., 2020). In October, the International Energy Agency (IEA) acknowledged that net-zero carbon by 2050 was the new standard for a clean energy transition and laid out a roadmap for the world to get there (IEA, 2020a).

Year 2020 witnessed a public willing to tackle systemic risks by transforming the intertwined social, economic and technological systems (Sovacool et al., 2020b) that have created overlapping crises of sustainability, health, equality and democracy (Miller, 2019). Generating shifts in human values and new ways of thinking and acting are often harder to achieve than technical solutions. Yet, as public responses to COVID-19 have demonstrated, when motivated, people can and do change. New narratives and forms of imagination are emerging to guide transformative change and to facilitate a transition towards new, more sustainable and equitable models for the economy, more socially and environmentally responsible technological innovation, and more just systems of governance (Eschrich & Miller, 2019; Iwaniec et al., 2020).

Translating this emergent global compact into stronger forms of international collaboration for the planet is the key to effective long-term responses to COVID-19 and climate change. Systemic risks will require innovative, adaptive, reflexive, transparent, participatory and accountable approaches to governance (Brown & Scobie, 2020; Chou et al., 2020). Rapid, networked, transformative responses that foster greater trust and more just relationships between diverse actors will be indispensable to creating a thriving and equitable global future for all (Scobie et al., 2020).

3.8. Economic stimulus focused primarily on growth would jeopardize the Paris Agreement

An increasing number of studies provide solid evidence that there are substantial economic benefits of climate action in the short as well as long term. Climate mitigation has substantial co-benefits, here and now, in terms of local economic, environmental and health benefits (Karlsson et al., 2020; Rauner et al., 2020). Recent research insights show that economically ‘optimal’ abatement could very well be in line with the UN climate targets of limiting global warming to well below 2°C and to actively pursue a 1.5°C limit (Burke et al., 2018a; Glanemann et al., 2020; Hänsel et al., 2020). As the remaining carbon budget is limited it is essential to use it on investments that lead to high net CO₂ savings, that is, have a high return on investment in terms of CO₂-emission reductions (Alfredsson & Malmaeus, 2019).

An important driver for the changing cost landscape is the significant drop in costs being realized for renewable energy, battery storage and electric mobility. The global average levelized cost of electricity has fallen by 82% for solar photovoltaics, and 29 and 40% for offshore and onshore wind power respectively, between 2010 and 2019 (IRENA, 2020). Of all the newly commissioned utility-scale renewable power generation projects, 56% (by capacity) had a levelized cost lower than the cheapest new source of fossil fuel-fired power (IRENA, 2020). Batteries for electric vehicles in the United States have dropped in average price from more than 1100 USD/kWh in 2010 to 156 USD/kWh in 2019 (IEA, 2020b).

There is a risk, however, that gains from growing clean energy sources are offset by rapid growth of economic activity that increase the overall demand for energy slowing down system-wide decarbonization. Dyrstad et al. (2019) have shown that this has happened in OECD countries since 1980. A large body of literature finds that there has generally been – at a global level – a strong coupling between GDP growth, resource use and GHG emissions (Haberl et al., 2020; Parrique et al., 2019; Vadén et al., 2020). In high-income countries there is, in terms of GHG emissions and if measured from a production perspective,

evidence of a small absolute decoupling. Several countries have shown that it is possible to combine (low) economic growth with decreasing CO₂ emissions, also for consumption-based emissions, when there are targeted policies (Le Quéré et al., 2019). Still, current policies are insufficient to reduce emissions globally at the rate needed to achieve the goals of the Paris Agreement (Roelfsema et al., 2020).

The decarbonization rate and mitigation costs not only depend on technology development, but also on the rate and type of economic development. In order to stay below 2°C, modelling scenarios with high growth often require CO₂ removal (CDR) at quantities that threaten several sustainability goals. Van Vuuren et al. (2018) show that in a scenario with moderate growth, the needs for CDR can be greatly reduced when combining a technology transition with substantial behavioural changes.

Weighing in the critical time factor, recent scientific evidence shows that if the economic recovery after COVID-19 has a primary focus on economic growth, with sustainability and climate mitigation as a secondary goal, it could jeopardize our last chance of achieving the Paris Agreement and safeguarding people's health, well-being and a prosperous economic development.

This insight is further elaborated on in the Supplementary material.

3.9. Electrification increasingly pivotal for just sustainability transitions and urban areas are at the forefront

Urban electrification has accelerated in recent years (World Economic Forum, 2020). However, although the decarbonization impacts of electrification are well documented in industrial and transport sectors (Alarfaj et al., 2020; Arabzadeh et al. (2020); Lah et al., 2020; Madeddu et al., 2020; Mai et al., 2020), comprehensive analyses of the role that urban electrification can play are lacking (Fuso Nerini et al., 2019). The sustainable energy transition relies on a concurrent global urban transition (IRENA, 2019; IPCC et al., 2018). Urban electrification offers opportunities to examine the challenges and harness the opportunities of urbanization and decarbonization in tandem (Romero-Lankao et al., 2019; Allam et al., 2020); it opens up new areas of discussion for bridging urban and energy planning, which require interdisciplinary dialogues. Cities will need to develop new solutions, including fundamental structural and systemic changes, to cope with expected urbanization trends (Salvucci & Tattini, 2019) and other emerging technological innovations, like e-commerce and e-ride hailing.

The current wave of electrification is mainly driven by urban buildings and on-road transportation, especially battery electric vehicles, heat pumps and cookstoves (Romero-Lankao et al., 2019). Electric utilities and investors see these changes as new sources of growth, as can be seen from the global trends in investment in electricity networks. Rates of decline in carbon intensity are forecast to be faster in cities and with municipally owned utilities, due to their renewable targets, unique regulatory structures and prominent role in regional, state and national economies (REN21, 2019). Electrification via micro-grids can support the development of a small- and medium-sized enterprise-based industry (Ganguly et al., 2020), that shares economic benefits throughout communities (Westman, Moores & Burch, 2021), and is linked with improvements in per capita income (Akin et al., 2018). However, there is hardly any effort to examine those possibilities in urban contexts.

The expectation is that urban electrification can help leapfrog societies towards low-carbon sustainable energy systems and facilitate broad-based, just changes in the urban environment, thereby aligning adaptation and mitigation with the SDGs (IPCC, 2018). Reductions in local air pollution and improvements to health and quality of life are some tangible co-benefits of urban electrification (REN21, 2019). Cities, including government, community and private actors, are at the forefront of innovation and adoption of technologies and thereby can be hubs of accelerated and equitable energy transitions (Bai et al., 2018; de Chalendar, 2019; Kern, 2019; Romero-Lankao, 2018; Ryan, 2015). Cities are also places of informal settlements, environmental inequalities and energy poverty; and adaptation may increase energy demand (Gielen et al., 2019).

Urban electrification opens up opportunities to provide access to clean and affordable energy from renewable sources (e.g. Stewart et al., 2018) to over a billion people in the world who lack access to electricity, many of whom live in rapidly urbanizing areas or urbanized areas where access to electricity is highly uneven (de Collaço et al., 2019).

Urban electrification can help democratize electricity provision (Burke & Stephens, 2018). Decentralized energy systems, for example, can facilitate a transition away from exclusively centralized high-carbon electricity systems (Adil & Ko, 2016), returning control to citizens over energy systems. Notable risks stem from significant inequalities of access to decision-making on investments and technologies; unmitigated, these factors could deepen the divide between those who benefit and can afford low-carbon systems and those who do not, or who bear the negative impacts (Korkovelos et al., 2020). This electrification divide is a question that has not yet received sufficient attention in the academic literature.

Many actions can help realize the potential of urban electrification. Communities, local officials and utilities are introducing decentralized power systems such as distributed energy generation, micro-grids and smart grids (Adil & Ko, 2016; Pullins, 2019). City officials are promoting the use of renewables in their government-owned facilities and also integrating them into their building codes (Schmid et al., 2020). Infrastructure is also being deployed to support end-use electrification, like electric vehicle charging solutions (IEA, 2020b).

It is, increasingly, grassroots movements that drive actions at the city level, involving diverse stakeholders. Youth climate activists, community actors and transnational networks are engaging in a variety of actions from working on urban planning, green transport and grid integration, to challenging existing power relationships around current energy regimes as well as the actors and political authorities who maintain them (Szulecki, 2018). Urban electrification benefits, including reduction of GHG emissions, will be realized only if the demands of the built environment, institutional constraints and the carbon intensity of energy sources are addressed (Castan Broto, 2019; Romero-Lankao et al., 2019).

This insight is further elaborated on in the Supplementary material.

3.10. Rights-based litigation as an essential tool in climate action

Litigation is an essential tool to urge action to prevent dangerous climate change and support the goals of the Paris Agreement (Gerrard, 2019; Setzer & Byrnes, 2019; International Bar

Association, 2020; Mitkidis & Valkanou, 2020; Wegener, 2020). Most climate cases are public interest litigation against a government (e.g. *Urgenda Foundation v. State of the Netherlands*, 2020) although claims are also brought against private actors such as oil companies (e.g. *Milieudefensie et al., v. Royal Dutch Shell plc.*, filed in 2019), and can be initiated before domestic courts and international courts, tribunals or human rights treaty bodies or non-compliance mechanisms (Spijkers, 2020). Developing climate policy is typically the domain of the legislative branch of the State, but given the urgency to act and the absence of adequate climate action or enforcement, the courts come in as 'lawmakers' (Spijkers & Oosterhuis, 2020; Voigt, 2019). This challenges conventional interpretations of the balance of power and features a critical interplay between scientific evidence and adjudication.

Climate cases have been based primarily on alleged human rights violations, around which litigation in developing countries, particularly in Latin America, is growing in scale and extent (Peel & Lin, 2019). Such rights-based litigation appears to be a suitable and effective channel to clarify the content and scope of existing human rights, such as the right to life and the right to a private life, in light of climate change impacts (Rodríguez-Garavito, 2020). The human rights prism has also led to a more focused debate on the obligation of conduct that states have in order to avoid dangerous climate change. This in particular details the definition of due diligence and the requirement of states to reflect on their highest possible ambition in their national climate plans, policies and laws. Moreover, climate litigation plays an important role in defining the content of a human right to a clean and healthy environment, and how this relates to the duty to inform, the precautionary principle and other substantive and procedural principles of international environmental law (Peel & Osofsky, 2018).

Responsibility for extraterritorial emissions or harm is another critical issue addressed by climate litigation. One contentious issue in this context is whether states are responsible and should account for 'imported emissions' (which are produced elsewhere and cause emissions during those processes but are consumed 'at home') or 'exported emissions' (the result of exported oil and gas products that are refined and burned abroad). Extraterritoriality also applies to human rights violations due to climate impacts, where the cause of such impacts may have been in states other than those whose people are the victims of such harm.

Climate litigation clarifies the issue of international legal standing and representation of the rights and interests of future generations in a healthy environment. Standing is closely linked to establishing victimhood, which may involve future harm or harm to future generations. In some instances, children have initiated cases or similar proceedings as representatives of future generations. In September 2019, 16 children – representing 12 nationalities – filed complaints against five countries before the United Nations Committee on the Rights of the Child, and a group of Portuguese youth lodged an application in September 2020 at the European Court of Human Rights against 33 states to provoke legally binding climate action.

Furthermore, there has been an increase in climate-related cases yielding legal rights of nature. For example, *Asociación Civil por la Justicia Ambiental v. Province of Entre Ríos*, et al., filed 7 July 2020 in Argentina; and *Demanda Generaciones Futuras v. Minambiente*, (Republica de Colombia, 2018), which found the Amazon to have standing and be subject to protection. The number of different actors who can represent climate-related

cases has widened, such as an NGO (McGrath, 2019), ombuds-person, trustee, institution, governmental agency or a select group of individuals. Also, courts, compliance procedures and human rights treaty bodies are starting to be asked to recognize the standing and rights of those who leave their country because it no longer sustains their life – known as climate migrants, like the case of *Ioane Teitiota v. New Zealand* (24 October 2019).

During recent decades, states have considered International Courts and Tribunals (ICT) to be an appropriate forum for the settlement of their international environmental legal disputes. ICTs are increasingly recognized as a potentially powerful venue for adjudication on climate and the court's jurisdiction to advise. This is due also to the demonstrated influence and cross-fertilization among judges, courts and tribunals at domestic, regional and international levels (Saiger, 2019; Wegener, 2020). Recent decisions of ICT have highlighted the challenges of resolving environmental disputes, such as the assessment of the evidence and the complexity of reparation for the loss of environmental goods and services, including gas regulation and carbon sequestration.

In summary, important developments are seen in climate litigation concerning the expansion of who and what has legal standing in courts, who may represent interests such as that of future generations, how to address harm across national boundaries, the role of courts in mandating climate action and cross-fertilization between courts and tribunals across levels and scales.

This insight is further elaborated on in the Supplementary material.

4. Conclusions

Year 2020 will enter history as the year in which the COVID-19 pandemic ravaged our world and reshaped our lives. Global responses to the pandemic provide a unique opportunity for the crucial large-scale sustainable investments needed to reach the Paris Agreement goals; these are investments that in turn are needed for healthy, sustainable lifestyles and a prosperous economic development. Although our horizon scan identifies a continued amplification in key environmental impacts (e.g. emissions and permafrost thaw), it also points to opportunities that arise from new views on climate change economics and governance, partially in response to the pandemic.

The COVID-19 pandemic not only reinforces the links between human health and climate change, it also provides a strong manifestation of how global crises can emerge in the hyper-connected world of the Anthropocene. Furthermore, it provides a unique opportunity for positive change by stimulating new social contracts and narratives but it's critical that economic stimulus measures reduce the risk of climate change rather than to drive short-term economic growth. Ultimately, the most fundamental 2020 insight may be that the world's nations and citizens can act together in the face of global threats and, although we cannot yet be said to be on the right climate trajectory, we can draw upon science and evidence to shape a safe, equitable and resilient future.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/sus.2021.2>

Acknowledgements. The authors acknowledge support from Future Earth, The Earth League, World Climate Research Programme (WCRP), Arizona State University, Earth System Governance Project (ESG Project), Integrated Land Ecosystem-Atmosphere Processes Study (iLEAPS), Global Carbon

Project (GCP) and Future Earth's Knowledge-Action Networks for (a) Health, (b) Systems of Sustainable Consumption and Production and (c) Urban. We further acknowledge support from Clara Burgard at Helmholtz-Zentrum Geesthacht and The Earth League, Roy Yi Ling Ngerng, National Taiwan University and Maria Martin at University of Potsdam, Institute for Climate Impact Research.

Author contributions. HAC, EF, TH, SL, HN, JohR, PR-L, DBS, LS and PS constituted the Editorial Board, conceiving and designing the study and providing editorial oversight. EP coordinated and provided editorial oversight. CME, SRH, RAL, CP, EP and GBS coordinated writing. EA, MB, KJB, VCB, KTC, KE, CME, PF, AG-F, MG, ARH, KH, BMH, SRH, GH, TI, RBJ, TFK, RAL, MiM, MaM, RIM, CM, CAM, NN, EASO, CP, JoP, JuP, PR-L, JoyR, AAS, ES, MS, SCS, GBS, JaS, JHS, OS, PHCT, MRT, AMU, DPvV, CV, CW and MDZ performed investigations and writing.

Financial support. VCB was supported by European Union's Horizon 2020 research and innovation programme, Grant Agreement No. 804051- LOACT – ERC-2018-Stg. PF was supported by the European Union's Horizon 2020 research and innovation programme under 4C (Grant Agreement No. 821003). AGF was supported by ANID/FONDAP/15130015. SRH was supported by the European Space Agency (ESA). GH was supported by The European Union Horizon 2020 research and innovation project Nunataryuk (773421). TH acknowledges support from the European Union's Horizon 2020 research and innovation programme under grant agreement No. 821471 (ENGAGE). TI acknowledges support by European Union's Horizon 2020 research and innovation programme under grant agreement no. 821003 (4C) and European Union's Horizon 2020 research and innovation programme under grant agreement no. 641816 (CRESCENDO). TFK acknowledges support by the Reducing Uncertainties in Biogeochemical Interactions through Synthesis and Computation Scientific Focus Area (RUBISCO SFA), which is sponsored by the Regional and Global Model Analysis (RGMA) Program in the Climate and Environmental Sciences Division (CESD) of the Office of Biological and Environmental Research (BER) in the U.S. Department of Energy Office of Science. NN acknowledges support from Global Affairs Canada. PR-L acknowledges support by the National Renewable Energy Laboratory (NREL), operated by Alliance for Sustainable Energy, LLC, for the U.S. Department of Energy (DOE) under Contract No. DE-AC36-08GO28308. JoP was supported by the European Research Council Synergy grant (ERC-SyG-2013-610028 IMBALANCE-P), the Spanish Government grant PID2019-110521GB-I00 and the Catalan government grant SGR2017-1005. AAS was supported by the Met Office Hadley Centre Climate Programme funded by BEIS and Defra. ES acknowledges support from NSF PLR Arctic System Science Research Networking Activities (RNA) Permafrost Carbon Network: Synthesizing Flux Observations for Benchmarking Model Projections of Permafrost Carbon Exchange, Grant no. 1931333. (2019–2023). SCS's work was supported by the Australian Research Council grant FL150100035. GBS would like to acknowledge the International Research Fellow programme of Japan Society for the Promotion of Science. PHCT's work was supported by a share of a grant from The São Paulo Research Foundation – FAPESP (grant number 2018/06685-9). EF acknowledges support from the Nordic Africa Institute. AMU acknowledges support from the Australian Research Council Centre of Excellence for Climate Extremes (CE170100023). The effort of MDZ was performed under the auspices of the U.S. Department of Energy (DOE) by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344 and was supported by the Regional and Global Model Analysis Program of the Office of Science at the DOE.

Conflict of interest. Johan Rockström is the current Editor-in-Chief of *Global Sustainability* journal. All other authors declare no conflict of interest.

References

Adil, A. M., & Ko, Y. (2016). Socio-technical evolution of decentralized energy systems: A critical review and implications for urban planning and policy. *Renewable and Sustainable Energy Reviews*, 57, 1025–1037. <https://doi.org/10.1016/j.rser.2015.12.079>

- Aklin, M., Harish, S. P., & Urpelainen, J. (2018). A global analysis of progress in household electrification. *Energy Policy*, 122, 421–428. <https://doi.org/10.1016/j.enpol.2018.07.018>
- Alarfaj, A. F., Griffin, M., & Samaras, C. (2020). Decarbonizing US passenger vehicle transport under electrification and automation uncertainty has a travel budget. *Environmental Research Letters*, 15, 940–942. <https://doi.org/10.1088/1748-9326/ab7c89>
- Alfredsson, E., & Malmaeus, M. J. (2019). Real capital investments and sustainability – The case of Sweden. *Ecological Economics*, 161, 216–224. <https://doi.org/10.1016/j.ecolecon.2019.04.008>
- Allam, Z., Jones, D., & Thondoo, M. (2020). Decarbonization and Urban Sustainability. In: *Cities and Climate Change: Climate Policy, Economic Resilience and Urban Sustainability* (Allam, Z., Jones, D., & Thondoo, M. (eds.) pp. 33–54). Springer International Publishing. https://doi.org/10.1007/978-3-030-40727-8_2
- Andrijevic, M., Schleussner, C.-F., Gidden, M. J., McCollum, D. L., & Rogelj, J. (2020). COVID-19 recovery funds dwarf clean energy investment needs. *Science (New York, N.Y.)*, 370(6514), 298–300. <https://doi.org/10.1126/science.abc9697>
- Arabzadeh, V., Mikkola, J., Jasiūnas, J., & Lund, P. D. (2020). Deep decarbonization of urban energy systems through renewable energy and sector-coupling flexibility strategies. *Journal of Environmental Management*, 260, 110090. <https://doi.org/10.1016/j.jenvman.2020.110090>
- Bai, X., Dawson, R. J., Ürges-Vorsatz, D., Delgado, G. C., Salisu Barau, A., Dhakal, S., Dodman, D., Leonardsen, L., Masson-Delmotte, V., Roberts, D. C., Schultz, S. (2018). Six research priorities for cities and climate change. *Nature*, 555, 23–25. <https://doi.org/10.1038/d41586-018-02409-z>
- Bai, X., Nagendra, H., Shi, P., & Liu, H. (2020). Cities: Build networks and share plans to emerge stronger from COVID-19. *Nature*, 584(7822), 517–520. <https://doi.org/10.1038/d41586-020-02459-2>
- Barbier, E. B. (2020). Greening the post-pandemic recovery in the G20. *Environmental and Resource Economics*, 76, 685–703. <https://doi.org/10.1007/s10640-020-00437-w>
- Barbier, E. B., Lozano, R., Rodríguez, C. M., & Tröeng, S. (2020). Adopt a carbon tax to protect tropical forests. *Nature*, 578, 213–216. <https://doi.org/10.1038/d41586-020-00324-w>
- Bastos, A., Ciais, P., Friedlingstein, P., Sitch, S., Pongratz, J., Fan, L., Wigneron, J. P., Weber, U., Reichstein, M., Fu, Z., Anthoni, P., Arneth, A., Haverd, V., Jain, A. K., Joetzier, E., Knauer, J., Lienert, S., Loughran, T., McGuire, P. C., ... Zaehle, S. (2020). Direct and seasonal legacy effects of the 2018 heat wave and drought on European ecosystem productivity. *Science Advances*, 6(24), eaba2724. <https://doi.org/10.1126/sciadv.aba2724>
- Basu, R., Gavin, L., Pearson, D., Ebisu, K., & Malig, B. (2018). Examining the association between apparent temperature and mental health-related emergency room visits in California. *American Journal of Epidemiology*, 187(4), 726–735. <https://doi.org/10.1093/aje/kwx295>
- Bodas-Salcedo, A., Mulcahy, J. P., Andrews, T., Williams, R. D., Ringer, M. A., Field, P. R., & Elsaesser, G. S. (2019). Strong dependence of atmospheric feedbacks on mixed-phase microphysics and aerosol-cloud interactions in HadGEM3. *Journal of Advances in Modeling Earth Systems*, 11(6), 1735–1758. <https://doi.org/10.1029/2019MS001688>
- Bond, W. J., Stevens, N., Midgley, G. F., & Lehmann, C. E. R. (2019). The trouble with trees: Afforestation plans for Africa. *Trends in Ecology & Evolution*, 34(11), 963–965. <https://doi.org/10.1016/j.tree.2019.08.003>
- Bowman, D. M. J. S., Kolden, C. A., Abatzoglou, J. T., Johnston, F. H., van der Werf, G. R., & Flannigan, M. (2020). Vegetation fires in the Anthropocene. *Nature Reviews Earth & Environment*, 1, 500–515. <https://doi.org/10.1038/s43017-020-0085-3>
- Brando, P. M., Soares-Filho, B., Rodrigues, L., Assunção, A., Morton, D., Tuchsneider, D., Fernandes, E. C. M., Macedo, M. N., Oliveira, U., & Coe, M. T. (2020). The gathering firestorm in southern Amazonia. *Science Advances*, 6(2), 1632. <https://advances.sciencemag.org/content/6/2/eaay1632>
- Bratman, G. N., Anderson, C. B., Berman, M. G., Cochran, B., de Vries, S., Flanders, J., Folke, C., Frumkin, H., Gross, J. J., Hartig, T., Kahn Jr., P. H., Kuo, M., Lawler, J. J., Levin, P. S., Lindahl, T., Meyer-Lindenberg, A., Mitchell, R., Ouyang, Z., Roe, J., ... Daily, G. C. (2019). Nature and mental health: An ecosystem service perspective. *Science Advances*, 5(7), eaax0903. <https://doi.org/10.1126/sciadv.aax0903>

- Brown, C., & Scobie, M. (2020). Accountability in the governance of global change. In M. M. Betsill, T. M. Benney & A. Gerlak (eds), *Agency in Earth System Governance* (pp. 155–167). Cambridge, UK: Cambridge University Press. <https://doi.org/10.1017/9781108688277>
- Burke, M., Davis, W. M., & Diffenbaugh, N. S. (2018a). Large potential reduction in economic damages under UN mitigation targets. *Nature*, 557, 549–553. <https://doi.org/10.1038/s41586-018-0071-9>
- Burke, M., González, F., Baylis, P., Heft-Neal, S., Baysan, C., Basu, S., & Hsiang, S. (2018b). Higher temperatures increase suicide rates in the United States and Mexico. *Nature Climate Change*, 8(8), 723–729. <https://doi.org/10.1038/s41558-018-0222-x>
- Burke, M. J., & Stephens, J. C. (2018). Political power and renewable energy futures: A critical review. *Energy Research and Social Science*, 35, 78–93. <https://doi.org/10.1016/j.erss.2017.10.018e>
- Ceccherini, G., Duveiller, G., Grassi, G., Lemoine, G., Avitabile, V., Pilli, R., & Cescatti, A. (2020). Abrupt increase in harvested forest area over Europe after 2015. *Nature*, 583, 72–77. <https://doi.org/10.1038/s41586-020-2438-y>
- Castan Broto, V. (2019). *Urban Energy Landscapes*. Cambridge, UK: Cambridge University Press. <https://doi.org/10.1017/9781108297868>
- Cianconi, P., Betrò, S., & Janiri, L. (2020). The impact of climate change on mental health: A systematic descriptive review. *Frontiers in Psychiatry*, 11 (March), 1–15. <https://doi.org/10.3389/fpsyt.2020.00074>
- Ciavarella, A., Cotterill, D., Stott, P., Kew, S., Philip, S., van Oldenborgh, G. J., Skålevåg, A., Lorenz, P., Robin, Y., Otto, F., Hauser, M., Seneviratne, S. I., Lehner, F., Shirshov, P. P., & Zolina, O. (2020). Prolonged Siberian heat of 2020. <https://www.worldweatherattribution.org/wp-content/uploads/WWA-Prolonged-heat-Siberia-2020.pdf>
- Clayton, S. (2020). Climate anxiety: Psychological responses to climate change. *Journal of Anxiety Disorders*, 74, 102263. <https://doi.org/10.1016/j.janxdis.2020.102263>
- Climate Action Tracker. (2020) (accessed 30 November 2020); https://climateactiontracker.org/documents/829/CAT_2020-12-01_Briefing_GlobalUpdate_Paris5Years_Dec2020.pdf
- Collaço, F. M. de A., Dias, L. P., Simoes, S. G., Pukšec, T., Seixas, J., & Bermann, C. (2019). What if São Paulo (Brazil) would like to become a renewable and endogenous energy-based megacity? *Renewable Energy*, 138, 416–433. <https://doi.org/10.1016/j.renene.2019.01.073>
- Craig, C. A., Feng, S., & Gilbert, S. (2019). Water crisis, drought, and climate change in the southeast United States. *Land Use Policy*, 88, 104110. <https://doi.org/10.1016/j.landusepol.2019.104110>
- Dean, J. F. (2020). Old methane and modern climate change. *Science* (New York, N.Y.), 367(6480), 846–848. <https://doi.org/10.1126/science.aba8518>
- Dean, J. F., Meisel, O. H., Rosco, M. M., Marchesini, L. B., Garnett, M. H., Lenderink, H., van Logtestijn, R., Borges, A. V., Bouillon, S., Lambert, T., Röckmann, T., Maximov, T., Petrov, R., Karsanaev, S., Aerts, R., van Huissteden, J., Vonk, J. E., & Dolman, A. J. (2020). East Siberian Arctic inland waters emit mostly contemporary carbon. *Nature Communications*, 11, 1627. <https://doi.org/10.1038/s41467-020-15511-6>
- de Chalendar, J. A., Glynn, P. W., & Benson, S. M. (2019). City-scale decarbonization experiments with integrated energy systems. *Energy & Environmental Science*, 12, 1695–1707. <https://doi.org/10.1039/C8EE03706J>
- de Collaço, F. M. A., Dias, L. P., Simoes, S. G., Pukšec, T., Seixas, J., & Bermann, C. (2019). What if São Paulo (Brazil) would like to become a renewable and endogenous energy-based megacity? *Renewable Energy*, 138, 416–433. <https://doi.org/10.1016/j.renene.2019.01.073>
- Di Luca, A., Pitman, A. J., & de Elia, R. (2020). Decomposing temperature extreme errors in CMIP5 and CMIP6 models. *Geophysical Research Letters*, 47(14), e2020GL088031. <https://doi.org/10.1029/2020GL088031>
- Dixon-Decleve, S., Schellnhuber, H. J., & Raworth, K. (2020). Could COVID-19 give rise to a greener global future? World Economic Forum. <https://www.weforum.org/agenda/2020/03/a-green-reboot-after-the-pandemic>
- Dyonisius, M. N., Petrenko, V. V., Smith, A. M., Hua, Q., Yang, B., Schmitt, J., Beck, J., Seth, B., Bock, M., Hmiel, B., Vimont, I., Menking, J. A., Shackleton, S. A., Baggenstos, D., Bauska, T. K., Rhodes, R. H., Sperlich, P., Beaudette, R., Harth, C., ... Weiss, R. F. (2020). Old carbon reservoirs were not important in the deglacial methane budget. *Science* (New York, N.Y.), 367(6480), 907–910. <https://doi.org/10.1126/science.aaa0504>
- Dyrstad, J. M., Skonhoft, A., Christensen, M. Q., & Ødegaard, E. T. (2019). Does economic growth eat up environmental improvements? Electricity production and fossil fuel emission in OECD countries 1980–2014. *Energy Policy*, 125, 103–109. <https://doi.org/10.1016/j.enpol.2018.10.051>
- Empinotti, V. L., Budds, J., & Aversa, M. (2019). Governance and water security: The role of the water institutional framework in the 2013–15 water crisis in São Paulo, Brazil. *Geoforum; Journal of Physical, Human, and Regional Geosciences*, 98, 46–54. <https://doi.org/10.1016/j.geoforum.2018.09.022>
- Engemann, K., Pedersen, C. B., Arge, L., Tsirogiannis, C., Mortensen, P. B., & Svenning, J.-C. (2019). Residential green space in childhood is associated with lower risk of psychiatric disorders from adolescence into adulthood. *Proceedings of the National Academy of Sciences of the United States of America*, 116(11), 5188–5193. <https://doi.org/10.1073/pnas.1807504116>
- Engström, G., Gars, J., Jaakkola, N., Lindahl, T., Spiro, D., & van Benthem, A. A. (2020). What policies address both the coronavirus crisis and the climate crisis? *Environmental and Resource Economics*, 76, 789–810. <https://doi.org/10.1007/s10640-020-00451-y>
- Eschrich, J., & Miller, C. A. (2019). *The Weight of Light: A Collection of Solar Futures*. Tempe, AZ: Center for Science and the Imagination.
- Farquharson, L. M., Romanovsky, V. E., Cable, W. L., Walker, D. A., Kokelj, S. V., & Nicolsky, D. (2019). Climate change drives widespread and rapid thermokarst development in very cold permafrost in the Canadian High Arctic. *Geophysical Research Letters*, 46, 6681–6689. <https://doi.org/10.1029/2019GL082187>
- Flynn, C. M., & Mauritsen, T. (2020). On the climate sensitivity and historical warming evolution in recent coupled model ensembles. *Atmospheric Chemistry and Physics*, 20, 7829–7842. <https://doi.org/10.5194/acp-20-7829-2020>
- Forster, P. M., Forster, H. I., Evans, M. J., Gidden, M. J., Jones, C. D., Keller, C. A., Lamboll, R. D., Le Quéré, C., Rogelj, J., Rosen, D., Schleussner, C.-F., Richardson, T. B., Smith, C. J., & Turnock, S. T. (2020). Current and future global climate impacts resulting from COVID-19. *Nature Climate Change*, 10, 913–919. <https://doi.org/10.1038/s41558-020-0883-0>
- Forzieri, G., Miralles, D. G., Ciais, P., Alkama, R., Ryu, Y., Duveiller, G., Zhang, K., Robertson, E., Kautz, M., Martens, B., Jiang, C., Arneeth, A., Georgievski, G., Li, W., Ceccherini, G., Anthoni, P., Lawrence, P., Wiltshire, A., Pongratz, J., ... & Cescatti, A. (2020). Increased control of vegetation on global terrestrial energy fluxes. *Nature Climate Change*, 10, 356–362. <https://doi.org/10.1038/s41558-020-0717-0>
- Friedlingstein, P., Jones, M. W., O'Sullivan, M., Andrew, R. M., Hauck, J., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Bakker, D. C. E., Canadell, J. G., Ciais, P., Jackson, R. B., Anthoni, P., Barbero, L., Bastos, A., Bastrikov, V., Becker, M., ... Zaehle, S. (2019). Global carbon budget 2019. *Earth System Science Data*, 11(4), 1783–1838. <https://doi.org/10.5194/essd-11-1783-2019>
- Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S., Aragão, L. E. O. C., Arneeth, A., Arora, V., ... Zaehle, S. (2020). Global carbon budget 2020. *Earth System Science Data*, 12, 3269–3340. <https://essd.copernicus.org/articles/12/3269/2020/essd-12-3269-2020.html>
- Fuso Nerini, F., Slob, A., Ericsson Engström, R., & Trutnevyte, E. (2019). A research and innovation agenda for zero-emission European cities. *Sustainability*, 11(6), 1692. <https://doi.org/10.3390/su11061692>
- Ganguly, R., Jain, R., Sharma, K. R., & Shekhar, S. (2020). Mini grids and enterprise development: A study of aspirational change and business outcomes among rural enterprise owners in India. *Energy for Sustainable Development*, 56, 119–127. <https://doi.org/10.1016/j.esd.2020.04.004>
- Gasser, T., Kechiar, M., Ciais, P., Burke, E. J., Kleinen, T., Zhu, D., Huang, Y., Ekici, A., & Obersteiner, M. (2018). Path-dependent reductions in CO₂ emission budgets caused by permafrost carbon release. *Nature Geoscience*, 11(11), 830–835. <https://doi.org/10.1038/s41561-018-0227-0>
- Gawel, E., & Lehmann, P. (2020). Killing two birds with one stone? Green dead ends and ways out of the COVID-19 crisis. *Environmental & Resource Economics*, 76, 447–517. <https://doi.org/10.1007/s10640-020-00443-y>
- Gerrard, M. (2019). Overview of climate change litigation. *Proceedings of the ASIL Annual Meeting*, 113, 194–197. <https://doi.org/10.1017/amp.2019.183>
- Gettelman, A., Hannay, C., Bacmeister, J. T., Neale, R. B., Pendergrass, A. G., Danabasoglu, G., Lamarque, J.-F., Fasullo, J. T., Bailey, D. A., Lawrence, D.

- M., & Mills, M. J. (2019). High climate sensitivity in the community earth system model version 2 (CESM2). *Geophysical Research Letters*, 46(14), 8329–8337. <https://doi.org/10.1029/2019GL083978>
- Gielen, D., Boshell, F., Saygin, D., Bazilian, M. D., Wagner, N., & Gorini, R. (2019). The role of renewable energy in the global energy transformation. *Energy Strategy Reviews*, 24, 38–50. <https://doi.org/10.1016/j.esr.2019.01.006>
- Glanemann, N., Willner, S. N., & Levermann, A. (2020). Paris Climate agreement passes the cost–benefit test. *Nature Communications*, 11(110). <https://doi.org/10.1038/s41467-019-13961-1>
- Graham, R. L., Francis, J., & Bogacz, R. J. (2017). Challenges and Opportunities of Grid Modernization and Electric Transportation. US Department of Energy. Retrieved from https://www.energy.gov/sites/prod/files/2017/06/f34/Challenges_and_Opportunities_of_Grid_Modernization_and_Electric_Transportation.pdf
- Haberl, H., Wiedenhofer, D., Virág, D., Kalt, G., Plank, B., Brockway, P., Fishman, T., Hausknost, D., Krausmann, F., Leon-Gruchalski, B., Mayer, A., Pichler, M., Schaffartzik, A., Sousa, T., Streeck, J., & Creutzig, F. (2020). A systematic review of the evidence on decoupling of GDP, resource use and GHG emissions, part II: Synthesizing the insights. *Environmental Research Letters* 15, 065003. <https://iopscience.iop.org/article/10.1088/1748-9326/ab842a>
- Hamstead, Z., & Coseo, P. (2020). Building policies, plans, and cities to manage extreme weather events: Perspectives from urban planning and landscape architecture. In K. H. Smith & P. K. Ram (eds), *Transforming global health* (pp. 261–283). Springer, Cham, Switzerland. https://doi.org/10.1007/978-3-030-32112-3_17
- Hanigan, I. C., Schirmer, J., & Niyonsenga, T. (2018). Drought and distress in southeastern Australia. *EcoHealth*, 15(3), 642–655. <https://doi.org/10.1007/s10393-018-1339-0>
- Hayward, B. (2020). *Children, Citizenship and Environment: #SchoolStrike edition*. (1st ed.). London, UK: Routledge. <https://doi.org/10.4324/9781003000396>
- Haines, A., Ebi, K. L., Smith, K. R., & Woodward, A. (2014). Health risks of climate change: Act now or pay later. *The Lancet*, 384(9948), 1073–1075. [https://doi.org/10.1016/S0140-6736\(14\)61659-7](https://doi.org/10.1016/S0140-6736(14)61659-7)
- Hänsel, M. C., Drupp, M. A., Johansson, D. J. A., Nesje, F., Azar, C., Freeman, M. C., Groom, B., & Sterner, T. (2020). Climate economics support for the UN climate targets. *Nature Climate Change*, 10, 781–789. <https://doi.org/10.1038/s41558-020-0833-x>
- Hayes, K., Berry, P., & Ebi, K. L. (2019). Factors influencing the mental health consequences of climate change in Canada. *International Journal of Environmental Research and Public Health*, 16(9), 1583. <https://doi.org/10.3390/ijerph16091583>
- Hayes, K., Blashki, G., Wiseman, J., Burke, S., & Reifels, L. (2018). Climate change and mental health: Risks, impacts and priority actions. *International Journal of Mental Health Systems*, 12(1), 1–12. <https://doi.org/10.1186/s13033-018-0210-6>
- Hayes, K., & Poland, B. (2018). Addressing mental health in a changing climate: Incorporating mental health indicators into climate change and health vulnerability and adaptation assessments. *International Journal of Environmental Research and Public Health*, 15(9), 1806. <https://doi.org/10.3390/ijerph15091806>
- Hayes, K., Poland, B., Cole, D. C., & Agic, B. (2020). Psychosocial adaptation to climate change in High River, Alberta: Implications for policy and practice. *Canadian Journal of Public Health = Revue canadienne de sante publique*, 111, 880–889. Switzerland. <https://doi.org/10.17269/s41997-020-00380-9>
- Head, L. (2020). Transformative change requires resisting a new normal. *Nature Climate Change*, 10(3), 173–174. <https://doi.org/10.1038/s41558-020-0712-5>
- Hepburn, C., O'Callaghan, B., Stern, N., Stiglitz, J., & Zenghelis, D. (2020). Will COVID-19 fiscal recovery packages accelerate or retard progress on climate change? *Oxford Review of Economic Policy*, 36(1), 359–381. <https://doi.org/10.1093/oxrep/gra015>
- Howarth, C., Bryant, P., Corner, A., Fankhauser, S., Gouldson, A., Whitmarsh, L., & Willis, R. (2020). Building a social mandate for climate action: Lessons from COVID-19. *Environmental and Resource Economics*, 76, 1107–1115. <https://doi.org/10.1007/s10640-020-00446-9>
- Hubau, W., Lewis, S. L., Phillips, O. L., Affum-Baffoe, K., Breeckman, H., Cuní-Sánchez, A., Daniels, A. K., Ewango, C. E. N., Fauset, S., Mukinzi, J. M., Sheil, D., Sonké, B., Sullivan, M. J. P., Sunderland, T. C. H., Taedoum, H., Thomas, S. C., White, L. J. T., Abernethy, K. A., Adu-Bredu, S., ... Zemagho, L. (2020). Asynchronous carbon sink saturation in African and Amazonian tropical forests. *Nature*, 579, 80–87. <https://doi.org/10.1038/s41586-020-2035-0>
- Hugelius, G., Loisel, J., Chadburn, S., Jackson, R. B., Jones, M., MacDonald, G., Marushchak, M., Olefeldt, D., Packalen, M., Siewert, M. B., Treat, C., Turetsky, M., Voigt, C., & Yu, Z. (2020). Large stocks of peatland carbon and nitrogen are vulnerable to permafrost thaw. *Proceedings of the National Academy of Sciences of the United States of America*, 117(34), 20438–20446. <https://doi.org/10.1073/pnas.1916387117>
- IEA. (2020a). World energy outlook 2020. International Energy Agency. <https://www.iea.org/reports/world-energy-outlook-2020>
- IEA. (2020b). Global EV outlook 2020. International Energy Agency. <https://www.iea.org/reports/global-ev-outlook-2020>
- International Bar Association. (2020). Model statute for proceedings challenging government failure to act on climate change: An International Bar Association Climate Change Justice and Human Rights Task Force Report. <https://www.ibanet.org/Climate-Change-Model-Statute.aspx>
- IPBES. (2019). *Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*. E. S. Brondizio, J. Settele, S. Díaz, and H. T. Ngo (eds). IPBES secretariat, Bonn, Germany.
- IPCC. (2013). Summary for policymakers. In T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex & P. M. Midgley (eds), *Climate change 2013: The physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change*, 1535 pp, Cambridge, UK and New York, NY, USA: Cambridge University Press.
- IPCC. (2018). Summary for Policymakers. In: Masson-Delmotte, V., Zhai P., Pörtner, H.-O., Roberts, D., Skea, J., Shukla, P. R., Pirani, A., Moufouma-Okia, W., Péan, C., Pidcock, R., Connors, S., Matthews J. B. R., Chen, Y., Zhou, X., Gomis, M. I., Lonnoy, E., Maycock, T., Tignor, M. and Waterfield, T. (eds.) *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty*. World Meteorological Organization, Geneva, Switzerland, 32 pp. Retrieved from <https://www.ipcc.ch/sr15/>.
- IRENA. (2019). *Global energy transformation: A roadmap to 2050* (2019 edition), International Renewable Energy Agency.
- IRENA. (2020). *Renewable power generation costs in 2019*, International Renewable Energy Agency.
- Iwaniec, D. M., Cook, E. M., Davidson, M. J., Berbé-Blázquez, M., Georgescu, M., Krayenhoff, E. S., Middel, A., Sampson, D. A., & Grimm, N. B. (2020). The co-production of sustainable future scenarios. *Landscape and Urban Planning*, 197, 103744. <https://doi.org/10.1016/j.landurbplan.2020.103744>
- Karlsson, M., Alfredsson, E., & Westling, N. (2020). Climate policy co-benefits: A review. *Climate Policy*, 20(3), 292–316. <https://doi.org/10.1080/14693062.2020.1724070>
- Kern, K. (2019). Cities as leaders in EU multilevel climate governance: Embedded upscaling of local experiments in Europe. *Environmental Politics*, 28(1), 125–145. <https://doi.org/10.1080/09644016.2019.1521979>
- Keuper, F., Wild, B., Kumm, M., Beer, C., Blume-Werry, G., Fontaine, S., Gavazov, K., Gentsch, N., Guggenberger, G., Hugelius, G., Jalava, M., Koven, C., Krab, E. J., Kuhry, P., Monteux, S., Richter, A., Shahzad, T., Weedon, J. T., Dorrepaal, E.... (2020). Carbon loss from northern circumpolar permafrost soils amplified by rhizosphere priming. *Nature Geoscience*, 13, 560–565. <https://doi.org/10.1038/s41561-020-0607-0>
- Kim, Y.-H., Min, S.-K., Zhang, X., Sillmann, J., & Sandstad, M. (2020). Evaluation of the CMIP6 multi-model ensemble for climate extreme indices. *Weather and Climate Extremes*, 29, 100269. <https://doi.org/10.1016/j.wace.2020.100269>
- Konapala, G., Mishra, A. K., Wada, Y., & Mann, M. E. (2020). Climate change will affect global water availability through compounding changes in

- seasonal precipitation and evaporation. *Nature Communications*, 11(3044). <https://doi.org/10.1038/s41467-020-16757-w>
- Korkovelos, A., Zerriffi, H., Howells, M., Bazilian, M., Rogner, H.-H., & Fuso Nerini, F. A. (2020). Retrospective analysis of energy access with a focus on the role of mini-grids. *Sustainability*, 12, 1793. <https://doi.org/10.3390/su12051793>
- Kuzemko, C., Bradshaw, M., Bridge, G., Goldthau, A., Jewell, J., Overland, I., Scholten, D., Van de Graaf, T., & Westphal, K. (2020). COVID-19 and the politics of sustainable energy transitions. *Energy Research & Social Science*, 68, 101685. <https://doi.org/10.1016/j.erss.2020.101685>
- Laborde, D., Martin, W., Swinnen, J., & Vos, R. (2020). COVID-19 risks to global food security. *Science (New York, N.Y.)*, 369(6503), 500–502. <https://doi.org/10.1126/science.abc4765>
- Lah, O., Fulton, L., & Arioli, M. (2020). Decarbonization scenarios for transport and the role of urban mobility. *Sustainable Urban Mobility Pathways*, 65–80. <https://doi.org/10.1016/B978-0-12-814897-6.00003-X>
- Le Quéré, C., Jackson, R. B., Jones, M. W., Smith, A. J. P., Abernethy, S., Andrew, R. M., De-Gol, A. J., Willis, D. R., Shan, Y., Canadell, J. G., Friedlingstein, P., Creutzig, F., & Peters, G. P. (2020). Temporary reduction in daily global CO₂ emissions during the COVID-19 forced confinement. *Nature Climate Change*, 10(7), 647–653. <https://doi.org/10.1038/s41558-020-0797-x>
- Le Quéré, C., Korsbakken, J. I., Wilson, C., Tosun, J., Andrew, R., Andres, R. J., Canadell, J. G., Jordan, A., Peters, G. P., & van Vuuren, D. P. (2019). Drivers of declining CO₂ emissions in 18 developed economies. *Nature Climate Change*, 9(3), 213–217. <https://doi.org/10.1038/s41558-019-0419-7>
- Leach, M., Reyers, B., Bai, X., Brondizio, E. S., Cook, C., Díaz, S., Espindola, G., Scobie, M., Stafford-Smith, M., & Subramanian, S. M. (2018). Equity and sustainability in the Anthropocene: A social-ecological systems perspective on their intertwined futures. *Global Sustainability*, 1, e13. <https://doi.org/10.1017/sus.2018.12>
- Lewkowicz, A. G., & Way, R. G. (2019). Extremes of summer climate trigger thousands of thermokarst landslides in a High Arctic environment. *Nature Communications*, 10, 1329. <https://doi.org/10.1038/s41467-019-09314-7>
- Li, W., Zhao, M., Scaioni, M., Hosseini, S. R., Wang, X., Yao, D., Zhang, K., Gao, J., & Li, X. (2019). Extreme rainfall trends of 21 typical urban areas in China during 1998–2015 based on remotely sensed data sets. *Environmental Monitoring Assessment*, 191(709). <https://doi.org/10.1007/s10661-019-7900-7>
- Liu, Z., Ciais, P., Deng, Z., Lei, R., Davis, S. J., Feng, S., Zheng, B., Cui, D., Dou, X., Zhu, B., Guo, R., Ke, P., Sun, T., Lu, C., He, P., Wang, Y., Yue, X., Wang, Y., Lei, Y., ... Schellnhuber, H. J. (2020). Near-real-time monitoring of global CO₂ emissions reveals the effects of the COVID-19 pandemic. *Nature Communications*, 11(1), 5172. <https://doi.org/10.1038/s41467-020-18922-7>
- Lomba-Fernández, C., Hernantes, J., & Labaka, L. (2019). Guide for climate-resilient cities: An urban critical infrastructures approach. *Sustainability*, 11(17), 4727. <https://doi.org/10.3390/su11174727>
- MacNeil, M. A., Mellin, C., Matthews, S., Wolff, N. H., McClanahan, T. R., Devlin, M., Drovandi, C., Mengersen, K., & Graham, N. A. J. (2019). Water quality mediates resilience on the Great Barrier Reef. *Nature Ecology & Evolution*, 3(4), 620–627. <https://doi.org/10.1038/s41559-019-0832-3>
- Madakumbura, G. D., Kim, H., Utsumi, N., Shiogama, H., Fischer, E. M., Seland, Ø., Scinocca, J. F., Mitchell, D. M., Hirabayashi, Y., & Oki, T. (2019). Event-to-event intensification of the hydrologic cycle from 1.5°C to a 2°C warmer world. *Scientific Reports*, 9, 3483. <https://doi.org/10.1038/s41598-019-39936-2>
- Madeddu, S., Ueckerdt, F., Pehl, M., Peterseim, J., Lord, M., Kumar, K. A., Krüger, C., & Luderer, G. (2020). The CO₂ reduction potential for the European industry via direct electrification of heat supply (power-to-heat). *Environmental Research Letters*, 15(12). <https://doi.org/10.1088/1748-9326/abb0d2>
- Mai, T. T., Jadun, P., Logan, J., McMillan, C., Muratori, M., Steinberg, D., Vimmerstedt, L., Jones, R., Haley, B., & Nelson, B. (2020). Electrification futures study: Scenarios of electric technology adoption and power consumption for the United States (National Renewable Energy Laboratory, Golden, CO). <https://www.nrel.gov/docs/fy18osti/71500.pdf>
- Malliet, P., Reynès, F., Landa, G., Hamdi-Cherif, M., & Saussay, A. (2020). Assessing short-term and long-term economic and environmental effects of the COVID-19 crisis in France. *Environmental and Resource Economics*, 76, 867–883. <https://doi.org/10.1007/s10640-020-00488-z>
- Matias, D. M. S. (2020). Climate humanitarian visa: International migration opportunities as post-disaster humanitarian intervention. *Climatic Change*, 160, 143–156. <https://doi.org/10.1007/s10584-020-02691-9>
- McDowell, N. G., Allen, C. D., Anderson-Teixeira, K. A., Aukema, B. H., Bond-Lamberty, B., Chini, L., Clark, J. S., Dietze, M., Grossiord, C., Hanbury-Brown, A., Hurtt, G. C., Jackson, R. B., Johnson, D. J., Kueppers, L., Lichstein, J. W., Ogle, K., Poulter, B., Pugh, T. A. M., Seidl, R., ... Xu, C. (2020). Pervasive shifts in forest dynamics in a changing world. *Science (New York, N.Y.)*, 368(6494). <https://doi.org/10.1126/science.aaz9463>
- McGrath, C. (2019). Urgenda appeal is groundbreaking for ambitious climate litigation globally. *Environmental and Planning Law Journal*, 36(1), 90–94.
- McIver, L., Kim, R., Woodward, A., Hales, S., Spickett, J., Katscherian, D., Hashizume, M., Honda, Y., Kim, H., Iddings, S., Naicker, J., Bambrick, H., McMichael, A., & Ebi, J., & L. K. (2016). Health impacts of climate change in pacific island countries: A regional assessment of vulnerabilities and adaptation priorities. *Environmental Health Perspectives*, 124(11), 1707–1714. <https://doi.org/10.1289/ehp.1509756>
- McMichael, C., Dasgupta, S., Ayeb-Karlsson, S., & Kelman, I. (2020). A review of estimating population exposure to sea-level rise and the relevance for migration. *Environmental Research Letters*, 15(12). <https://doi.org/10.1088/1748-9326/abb398>
- Meredith, M., Sommerkorn, M., Cassotta, S., Derksen, C., Ekaykin, A., Hollowed, A., Kofinas, G., Mackintosh, A., Melbourne-Thomas, J., Muelbert, M.M.C., Ottersen, G., Pritchard, H., & Schuur E.A.G. (2019). Polar regions. In H.-O. Pörtner, D. C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama & N. M. Weyer (eds), *IPCC Special report on the ocean and cryosphere in a changing climate*. (pp. 203–320). World Meteorological Organization, Geneva, Switzerland. Retrieved from: <https://www.ipcc.ch/srocc/chapter/chapter-3-2/>
- Middleton, J., Cunsolo, A., Jones-Bitton, A., Shiwak, I., Wood, M., Pollock, N. J., Flowers, C., & Harper, S. (2020). ‘We’re people of the snow’: weather, climate change, and inuit mental wellness. *Social Science & Medicine*, 262, 113137. <https://doi.org/10.1016/j.socscimed.2020.113137>
- Mildenberger, M. (2020). *Carbon captured: how business and labor control climate politics*. Cambridge, Mass., USA and London, UK: The MIT Press.
- Miller, C. A. (2019). Sustainability, democracy, and the techno-human future. In J. Hoff, Q. Gausset & S. Lex (eds), *The role of non-state actors in the green transition: Building a sustainable future* (pp. 247–266). New York, USA: Routledge.
- Mitkidis, K. P., & Valkanov, T. (2020). Climate change litigation: Trends. *Policy Implications and the Way Forward. Transnational Environmental Law*, 9(1), 11–16. <https://doi.org/10.1017/S2047102519000414>
- Myhre, G., Alterskjær, K., Stjern, C. W., Hodnebrog, Ø., Marelle, L., Samset, B. H., Sillmann, J., Schaller, N., Fischer, E., Schulz, M., & Stohl, A. (2019). Frequency of extreme precipitation increases extensively with event rareness under global warming. *Scientific Reports*, 9(16062). <https://doi.org/10.1038/s41598-019-52277-4>
- Nagabhatla, N., Pouramin, P., Brahmabhatt, R., Fioret, C., Glickman, T., Newbold, K. B., & Smakhtin, V. (2020). Water and migration: A global overview. UNU-INWEH Report Series, Issue 10. United Nations University Institute for Water, Environment and Health.
- Nicklin, H., Leicher, A. M., Dieperink, C., & Van Leeuwen, K. (2019). Understanding the costs of inaction—an assessment of pluvial flood damages in two European cities. *Water*, 11(4), 801. <https://doi.org/10.3390/w11040801>
- Nijse, F. J. M. M., Cox, P. M., & Williamson, M. S. (2020). An emergent constraint on Transient Climate Response from simulated historical warming in CMIP6 models. *Earth System Dynamics*. <https://doi.org/10.5194/esd-2019-86>
- Nitzbon, J., Westermann, S., Langer, M., Martin, L. C. P., Strauss, J., Laboor, S., & Boike, J. (2020). Fast response of cold ice-rich permafrost in northeast Siberia to a warming climate. *Nature Communications*, 11, 2201. <https://doi.org/10.1038/s41467-020-15725-8>

- Oldekop, J. A., Horner, R., Hulme, D., Adhikari, R., Agarwal, B., Alford, M., Bakewell, O., Banks, N., Barrientos, S., Bastia, T., Bebbington, A. J., Das, U., Dimova, R., Duncombe, R., Enns, C., Fielding, D., Foster, C., Foster, T., Frederiksen, T., ... Zhang, Y.-F. (2020). COVID-19 and the case for global development. *World Development*, 134, 105044. <https://doi.org/10.1016/j.worlddev.2020.105044>
- Otto, M. I., Donges, J. F., Cremades, R., Bhowmik, A., Hewitt, R. J., Lucht, W., Rockström, J., Allerberger, F., McCaffrey, M., Doe, S. P., Lenferna, A., Morán, N., van Vuuren, D. P., & Schellnhuber, H. J. (2020). Social tipping dynamics for stabilizing Earth's climate by 2050. *Proceedings of the National Academy of Sciences of the United States of America*, 117(5), 2354–2365. <https://doi.org/10.1073/pnas.1900577117>
- Palinkas, L. A., & Wong, M. (2020). Global climate change and mental health. *Current Opinion in Psychology*, 32, 12–16. <https://doi.org/10.1016/j.copsyc.2019.06.023>
- Parrique T., Barth J., Briens F., Kerschner, C., Kraus-Polk, A., Kuokkanen, A., & Spangenberg J. H. (2019). *Decoupling debunked: Evidence and arguments against green growth as a sole strategy for sustainability*, Bruxelles, Belgium: European Environment Bureau.
- Peel, J., & Lin, J. (2019). Transnational climate litigation: The contribution of the global south. *American Journal of International Law*, 113(4), 679–726. <https://doi.org/10.1017/ajil.2019.48>
- Peel, J., & Osofsky, H. M. (2018). A rights turn in climate change litigation? *Transnational Environmental Law*, 7(1), 37–67. <https://doi.org/10.1017/S2047102517000292>
- Piao, S., Wang, X., Park, T., Chen, C., Lian, X., He, Y., Bjerke, J. W., Chen, A., Ciais, P., Tømmervik, H., Nemani, R. R., & Myneni, R. B. (2020). Characteristics, drivers and feedbacks of global greening. *Nature Reviews Earth & Environment*, 1, 14–27. <https://doi.org/10.1038/s43017-019-0001-x>
- Pouramin, P., Nagabhatla, N., & Miletto, M. (2020). A systematic review of water and gender interlinkages: Assessing the intersection with health. *Frontiers in Water*, 2(6), <https://doi.org/10.3389/frwa.2020.00006>
- Pullins, S. (2019, June 1). Why microgrids are becoming an important part of the energy infrastructure. *Electricity Journal*, 32, 17–21. <https://doi.org/10.1016/j.tej.2019.05.003>
- Rauner, S., Bauer, N., Dirnaichner, A., Van Dingenen, R., Mutel, C., & Luderer, G. (2020). Coal-exit health and environmental damage reductions outweigh economic impacts. *Nature Climate Change*, 10, 308–312. <https://doi.org/10.1038/s41558-020-0728-x>
- Rehm, J., & Shield, K. D. (2019). Global burden of disease and the impact of mental and addictive disorders. *Current Psychiatry Reports*, 21(10). <https://doi.org/10.1007/s11920-019-0997-0>
- REN21. (2019). Renewables 2019 global status report. Retrieved from <https://ren21.net/gsr-2019/>
- Republica de Colombia, Corte Supreme de Justicia, Radicacion no. 11001-22-03-000-2018-00319-01 (5 April 2018).
- Renn, O., Lucas, K., Haas, A., & Jaeger, C. (2019). Things are different today: The challenge of global systemic risks. *Journal of Risk Research*, 22(4), 401–415. <https://doi.org/10.1080/13669877.2017.1409252>
- Roe, S., Streck, C., Obersteiner, M., Frank, S., Griscom, B., Drouet, L., Fricko, O., Gusti, M., Harris, N., Hasegawa, T., Hausfather, Z., Havlik, P., House, J., Nabuurs, G.-J., Popp, A., Sánchez, M. J. S., Sanderman, J., Smith, P., Stehfest, E.... (2019). Contribution of the land sector to a 1.5°C world. *Nature Climate Change*, 9, 817–828. <https://doi.org/10.1038/s41558-019-0591-9>
- Rodríguez-Garavito, C. (2020). Human rights: The global south's route to climate litigation. *American Journal of International Law Unbound*, 114, 40–44. <https://doi.org/10.1017/aju.2020.4>
- Roelfsema, M., van Soest, H. L., Harmsen, M., van Vuuren, D. P., Bertram, C., den Elzen, M., Höhne, N., Jacobuta, G., Krey, V., Kriegler, E., Luderer, G., Riahi, K., Ueckerdt, F., Després, J., Drouet, L., Emmerling, J., Frank, S., Fricko, O., Gidden, M., ... Vishwanathan, S. S. (2020). Taking stock of national climate policies to evaluate implementation of the Paris Agreement. *Nature Communications*, 11, 2096. <https://doi.org/10.1038/s41467-020-15414-6>
- Romero-Lankao, P., Bulkeley, H., Pelling, M., Burch, S., Gordon, D. J., Gupta, J., Johnson, C., Kurian, P., Lecavalier, E., Simon, D., Tozer, L., Ziervogel, G., & Munshi, D. (2018). Urban transformative potential in a changing climate. In *Nature Climate Change*, 8(9), 754–756. <https://www.nature.com/articles/s41558-018-0264-0#citeas>
- Romero-Lankao, P., Wilson, A., Sperling, J., Miller, C., Zimny-Schmitt, D., Bettencourt, L., Wood, E., Young, S., Muratori, M., Arent, D., O'Malley, M., Sovacool, B. K., Brown, M. A., Southworth, F., Bazilian, M., Gearhart, C., Beukes, A., & Zund, D. (2019). Urban electrification: Knowledge pathway toward an integrated research and development agenda. *SSRN Electronic Journal*, 10. <https://doi.org/10.2139/ssrn.3440283>
- Roshan, A., & Kumar, M. (2020). Water end-use estimation can support the urban water crisis management: A critical review. *Journal of Environmental Management*, 268, 110663. <https://doi.org/10.1016/j.jenvman.2020.110663>
- Rourke, M., Eccleston-Turner, M., Phelan, A., & Gostin, L. (2020). Policy opportunities to enhance sharing for pandemic research. *Science (New York, N.Y.)*, 368(6492), 716–718. <https://doi.org/10.1126/science.abb9342>
- Ryan, D. (2015). From commitment to action: A literature review on climate policy implementation at city level. *Climatic Change*, 131, 519–529. <https://doi.org/10.1007/s10584-015-1402-6>
- Saiger, A.-J. (2019). Domestic courts and the Paris agreement's climate goals: The need for a comparative approach 9(1). *Transnational Environmental Law*, 9(1), 37–54. <https://doi.org/10.1017/S2047102519000256>
- Salvucci, R., & Tattini, J. (2019). Global outlook for the transport sector in energy scenarios. In B. Holst Jørgensen, K. Krogh Andersen & O. Anker Nielsen (eds), *DTU International energy report 2019: Transforming urban mobility* (pp. 21–27), Kongens Lyngby, Denmark: Technical University of Denmark. Retrieved from <https://www.dtu.dk/english/research/find-research/publications-and-research-data/dtu-international-energy-report>
- Scaife, A. A., & Smith, D. (2018). A signal-to-noise paradox in climate science. *npj Climate and Atmospheric Science*, 1, 28. <https://doi.org/10.1038/s41612-018-0038-4>
- Schmid, B., Meister, T., Klagge, B., & Seidl, I. (2020). Energy cooperatives and municipalities in local energy governance arrangements in Switzerland and Germany. *The Journal of Environment & Development*, 29(1), 123–146. <https://doi.org/10.1177/1070496519886013>
- Schwartz, R. M., Liu, B., Lieberman-Cribbin, W., & Taioli, E. (2017). Displacement and mental health after natural disasters. *The Lancet Planetary Health*, 1(8), e314. [https://doi.org/10.1016/S2542-5196\(17\)30138-9](https://doi.org/10.1016/S2542-5196(17)30138-9)
- Schwerdtle, P., Bowen, K., & McMichael, C. (2018). The health impacts of climate-related migration. *BMC Medicine*, 16(1), 1–7. <https://doi.org/10.1186/s12916-017-0981-7>
- Scobie, M., Benney, T. M., Brown, C., & Widerberg, O. E. (2020). Conceptualizing agency and agents in earth system governance. In M. M. Betsill, T. M. Benney & A. Gerlak (eds), *Agency in earth system governance* (pp. 25–37). Cambridge, UK: Cambridge University Press. <https://doi.org/10.1017/9781108688277.002>
- SEI, IISD, ODI, E3G, & UNEP. (2020). The production gap report: 2020 special report. <http://productiongap.org/2020report>
- Setzer, J., & Byrnes, R. (2019). Global trends in climate change litigation: 2019 snapshot. Grantham Research Institute on Climate Change and the Environment and Centre for Climate Change Economics and Policy, London School of Economics and Political Science. Retrieved from http://www.lse.ac.uk/granthaminstitute/wp-content/uploads/2019/07/GRI_Global-trends-in-climate-change-litigation-2019-snapshot-2.pdf
- Shepherd, N. (2019). Making sense of 'Day Zero': Slow catastrophes, Anthropocene futures, and the story of Cape Town's water crisis. *Water*, 11(9), 1744. <https://doi.org/10.3390/w11091744>
- Sherwood, S. C., Webb, M. J., Annan, J. D., Armour, K. C., Forster, P. M., Hargreaves, J. C., Hegerl, G., Klein, S. A., Marvel, K. D., Rohling, E. J., Watanabe, M., Andrews, T., Braconnot, P., Bretherton, C. S., Foster, G. L., Hausfather, Z., von der Heydt, A. S., Knutti, R., Mauritsen, T., ... Zelinka, M. D. (2020). An assessment of earth's climate sensitivity using multiple lines of evidence. *Reviews of Geophysics*, 58(4). <https://doi.org/10.1029/2019RG000678>
- Simpson, N. S., Shearing, C. D., & Dupont, B. (2020). Gated adaptation during the Cape Town drought: Mentalities, transitions and pathways to partial

- nodes of water security. *Society & Natural Resources*, 33(8), 1041–1049. <https://doi.org/10.1080/08941920.2020.1712756>
- Smith, P., Calvin, K., Nkem, J., Campbell, D., Cherubini, F., Grassi, G., Korotkov, V., Hoang, A. L., Lwasa, S., McElwee, P., Nkonya, E., Saigusa, N., Soussana, J.-F., Taboada, M. A., Manning, F. C., Nampanzira, D., Arias-Navarro, C., Vizzarri, M., House, J., ... Arneth, A. (2019). Which practices co-deliver food security, climate change mitigation and adaptation, and combat land degradation and desertification? *Global Change Biology*, 26(3), 1432–1575. <https://doi.org/10.1111/gcb.14878>
- Smith, D. M., Scaife, A. A., Eade, R., Athanasiadis, P., Bellucci, A., Bethke, I., Bilbao, R., Borchert, L. F., Caron, L.-P., Counillon, F., Danabasoglu, G., Delworth, T., Doblas-Reyes, F. J., Dunstone, N. J., Estella-Perez, V., Flavoni, S., Hermanson, L., Keenlyside, N., Kharin, V., ... Zhang, L. (2020). North Atlantic climate far more predictable than models imply. *Nature*, 583(7818), 796–800. <https://doi.org/10.1038/s41586-020-2525-0>
- Sovacool, B. K., Furszyfer Del Rio, D., & Griffiths, S. (2020a). Contextualizing the COVID-19 pandemic for a carbon-constrained world: Insights for sustainability transitions, energy justice, and research methodology. *Energy Research & Social Science*, 68, 101701. <https://doi.org/10.1016/j.erss.2020.101701>
- Sovacool, B. K., Hess, D. J., Amir, S., Geels, F. W., Hirsh, R., Medina, L. R., Miller, C., Palavicino, C. A., Phadke, R., Ryghaug, M., Schot, J., Silvast, A., Stephens, J., Stirling, A., Turnheim, B., van der Vleuten, E., van Lente, H., & Yearley, S. (2020b). Sociotechnical agendas: Reviewing future directions for energy and climate research. *Energy Research & Social Science*, 70, 101617. <https://doi.org/10.1016/j.erss.2020.101617>
- Spijkers, O. (2020). The Case Between Urgenda and the State of the Netherlands. *Hungarian Yearbook of International Law and European Law*, 8(1), <https://doi.org/10.5553/hyiel/266627012020008001012>.
- Spijkers, O., & Oosterhuis, S. (2020). The Dutch response to climate change: Evaluating the Netherlands' climate act and associated issues of importance. In T. Muinzer (ed), *National Climate Change Acts: The Emergence, Form and Nature of National Framework Climate Legislation*. The Hague, Netherlands: Hart Publishing.
- Stewart, I. D., Kennedy, C. A., Facchini, A., & Mele, R. (2018). The electric city as a solution to sustainable urban development. *Journal of Urban Technology*, 25(1), 3–20. <https://doi.org/10.1080/10630732.2017.1386940>
- Szulecki, K. (2018). Conceptualizing energy democracy. *Environmental Politics*, 27(1), 21–41. <https://doi.org/10.1080/09644016.2017.1387294>
- Tabari, H., Hosseinzadehtalaei, P., AghaKouchak, A., & Willems, P. (2019). Latitudinal heterogeneity and hotspots of uncertainty in projected extreme precipitation. *Environmental Research Letters*, 14(12), 124032. <https://doi.org/10.1088/1748-9326/ab55fd>
- Tagesson, T., Schurgers, G., Horion, S., Ciais, P., Tian, F., Brandt, M., Ahlström, A., Wigneron, J.-P., Ardö, J., Olin, S., Fan, L., Wu, Z., & Fensholt, R. (2020). Recent divergence in the contributions of tropical and boreal forests to the terrestrial carbon sink. *Nature Ecology & Evolution*, 4, 202–209. <https://doi.org/10.1038/s41559-019-1090-0>
- Teegne, G., & Melesse, A. M. (2020). Multimodel ensemble projection of hydro-climatic extremes for climate change impact assessment on water resources. *Water Resources Management*, 34, 3019–3035. <https://doi.org/10.1007/s11269-020-02601-9>
- Terrer, C., Jackson, R. B., Prentice, I. C., Keenan, T. F., Kaiser, C., Vicca, S., Fisher, J. B., Reich, P. B., Stocker, B. D., Hungate, B. A., Peñuelas, J., McCallum, I., Soudzilovskaia, N. A., Cernusak, L. A., Talheim, A. F., Van Sundert, K., Piao, S., Newton, P. C. D., Hovenden, M. J., ... Franklin, O. (2019). Nitrogen and phosphorus constrain the CO₂ fertilization of global plant biomass. *Nature Climate Change*, 9, 684–689. <https://doi.org/10.1038/s41558-019-0545-2>
- Tharammal, T., Bala, G., Devaraju, N., & Nemani, R. (2019). A review of the major drivers of the terrestrial carbon uptake: Model-based assessments, consensus, and uncertainties. *Environmental Research Letters*, 14(9). <https://doi.org/10.1088/1748-9326/ab3012>
- Tokarska, K. B., Stolpe, M. B., Sippel, S., Fischer, E. M., Smith, C. J., Lehner, F., & Knutti, R. (2020). Past warming trend constrains future warming in CMIP6 models. *Science Advances*, 6(12). <https://doi.org/10.1126/sciadv.aaz9549>
- Turetsky, M. R., Abbott, B. W., Jones, M. C., Anthony, K. W., Olefeldt, D., Schuur, E. A. G., Grosse, G., Kuhry, P., Hugelius, G., Koven, C., Lawrence, D. M., Gibson, C., Sannel, A. B. K., & McGuire, A. D. (2020). Carbon release through abrupt permafrost thaw. *Nature Geoscience*, 13, 138–143. <https://doi.org/10.1038/s41561-019-0526-0>
- Ukkola, A. M., De Kauwe, M. G., Roderick, M. L., Abramowitz, G., & Pitman, A. J. (2020). Robust future changes in meteorological drought in CMIP6 projections despite uncertainty in precipitation. *Geophysical Research Letters*, 47(11). <https://doi.org/10.1029/2020GL087820>
- Vadén, T., Lähde, V., Majave, A., Järvensivu, P., Toivanen, T., Hakala, E., & Eronen, J. T. (2020). Decoupling for ecological sustainability: A categorisation and review of research literature. *Environmental Science & Policy*, 112, 236–244. <https://doi.org/10.1016/j.envsci.2020.06.016>
- Van Vuuren, D. P., Stehfest, E., Gernaat, D. E. H. J., van den Berg, M., Bijl, D. L., Sytze de Boer, H., Daioglou, V., Doelman, J. C., Edelenbosch, O. Y., Harmsen, M., Hof, A. F., & van Sluisveld, M. A. E. (2018). Alternative pathways to the 1.5°C target reduce the need for negative emission technologies. *Nature Climate Change*, 8, 391–397. <https://doi.org/10.1038/s41558-018-0119-8>
- Van Zyl, A., & Jooste, J. L. (2020). Retaining and recycling water to address water scarcity in the City of Cape Town. *Development Southern Africa*. <https://doi.org/10.1080/0376835X.2020.1801387>
- Vaughn, L. J. S., & Torn, M. S. (2019). ¹⁴C Evidence that millennial and fast-cycling soil carbon are equally sensitive to warming. *Nature Climate Change*, 9, 467–471. <https://doi.org/10.1038/s41558-019-0468-y>
- Vert, C., Gascon, M., Ranzani, O., Márquez, S., Triguero-Mas, M., Carrasco-Turigas, G., Arjona, L., Koch, S., Llopis, M., Donaire-Gonzalez, D., Elliott, L. R., & Nieuwenhuijsen, M. (2020). Physical and mental health effects of repeated short walks in a blue space environment: A randomised crossover study. *Environmental Research*, 188, 109812. <https://doi.org/10.1016/j.envres.2020.109812>
- Vivid Economics. (2020). Greenness of stimulus index – an assessment of COVID-19 stimulus by G20 countries in relation to climate action and biodiversity goals. Accessed from https://www.vivideconomics.com/wp-content/uploads/2020/11/201028-GSI-report_October-release.pdf.
- Voigt, C. (ed) (2019). *International judicial practice on the environment – questions of legitimacy*. (Studies on International Courts and Tribunals). Cambridge: Cambridge University Press. <https://doi.org/10.1017/9781108684385>
- Walker, A. P., De Kauwe, M. G., Bastos, A., Belmecheri, S., Georgiou, K., Keeling, R., McMahon, S. M., Medlyn, B. E., Moore, D. J. P., Norby, R. J., Zaehle, S., Anderson-Teixeira, K. J., Battipaglia, G., Brien, R. J. W., Cabugao, K. G., Cailleret, M., Campbell, E., Canadell, J., Caias, P., ... Zuidema, P. A. (2020). Integrating the evidence for a terrestrial carbon sink caused by increasing atmospheric CO₂. *New Phytologist*, 229, 2413–2445. <https://doi.org/10.1111/NPH.16866>
- Wang, T., Yang, D., Yang, Y., Piao, S., Li, X., Cheng, G., & Fu, B. (2020a). Permafrost thawing puts the frozen carbon at risk over the Tibetan Plateau. *Science Advances*, 6(19). <https://doi.org/10.1126/sciadv.aaz3513>
- Wang, Y., Yuan, Y., Wang, Q., Liu, C., Zhi, Q., & Cao, J. (2020b). Changes in air quality related to the control of coronavirus in China: Implications for traffic and industrial emissions. *Science of the Total Environment*, 731, 139133. <https://doi.org/10.1016/j.scitotenv.2020.139133>
- Wang, S., Zhang, Y., Ju, W., Chen, J. M., Ciais, P., Cescatti, A., Sardans, J., Janssens, I. A., Wu, M., Berry, J. A., Campbell, E., Fernández-Martínez, M., Alkama, R., Sitch, S., Friedlingstein, P., Smith, W. K., Yuan, W., He, W., Lombardozzi, D., ... Peñuelas, J. (2020c). Recent global decline of CO₂ fertilization effects on vegetation photosynthesis. *Science (New York, N.Y.)*, 370, 1295–1300. <https://doi.org/10.1126/science.abb7772>
- Wegener, L. (2020). Can the Paris agreement help climate change litigation and vice versa? *Transnational Environmental Law*, 9(1), 17–36. <https://doi.org/10.1017/S2047102519000396>
- Westman, L., Moores, E., & Burch, S. L. (2021). Bridging the governance divide: The role of SMEs in urban sustainability interventions. *Cities (London, England)*, 108, 102944. <https://doi.org/10.1016/j.cities.2020.102944>
- Whitehead, P. G., Jin, L., Bussi, G., Voepel, H. E., Darby, S. E., Vasilopoulos, G., Manley, R., Rodda, H., Hutton, C., Hackney, C., & Van Pham Dang Tri, & Hung, N. N. (2019). Water quality modelling of the Mekong River basin:

- Climate change and socioeconomics drive flow and nutrient flux changes to the Mekong Delta. *Science of the Total Environment*, 673, 218–229. <https://doi.org/10.1016/j.scitotenv.2019.03.315>
- Whyte, K. (2020). Too late for indigenous climate justice: Ecological and relational tipping points. *WIREs Climate Change*, 11(1), e603. <https://doi.org/10.1002/wcc.603>
- Witze, A. (2020). The Arctic is burning like never before – and that's bad news for climate change. *Nature*, 585, 336–337. <https://www.nature.com/articles/d41586-020-02568-y>
- World Economic Forum. (2020). The future of the last-mile ecosystem: Transition roadmaps for public- and private-sector players.
- Wu, J., Snell, G., & Samji, H. (2020). Climate anxiety in young people: A call to action. *The Lancet Planetary Health*, 4(10), e435–e436. [https://doi.org/10.1016/S2542-5196\(20\)30223-0](https://doi.org/10.1016/S2542-5196(20)30223-0)
- WWAP (2020). *The united nations world water development report 2020: Water and climate change*. Paris: UNESCO.
- Yang, Y., Roderick, M. L., Zhang, S., McVicar, T. R., & Donohue, R. J. (2019). Hydrologic implications of vegetation response to elevated CO₂ in climate projections. *Nature Climate Change*, 9(1), 44–48. <https://www.nature.com/articles/s41558-018-0361-0>
- Yokoyama, Y., Otsuka, K., Kawakami, N., Kobayashi, S., Ogawa, A., Tannno, K., Onoda, T., Yaegashi, Y., & Sakata, K. (2014). Mental health and related factors after the great east Japan earthquake and tsunami. *PLoS ONE*, 9(7), 10. <https://doi.org/10.1371/journal.pone.0102497>
- Zelinka, M. D., Myers, T. A., McCoy, D. T., Po-Chedley, S., Caldwell, P. M., Ceppi, P., Klein, S. A., & Taylor, K. E. (2020). Causes of higher climate sensitivity in CMIP6 models. *Geophysical Research Letters*, 47(1), e2019GL085782. <https://doi.org/10.1029/2019GL085782>
- Zhu, J., Poulsen, C. J., & Otto-Bliesner, B. L. (2020). High climate sensitivity in CMIP6 model not supported by paleoclimate. *Nature Climate Change*, 10, 378–379. <https://doi.org/10.1038/s41558-020-0764-6>