

First measurement of T -odd moments in $D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ decays

K. Prasanth,²¹ J. Libby,²¹ I. Adachi,^{15,11} H. Aihara,⁷⁶ S. Al Said,^{69,34} D. M. Asner,⁵⁹ V. Aulchenko,^{4,58} T. Aushev,⁴⁹ R. Ayad,⁶⁹ V. Babu,⁷⁰ I. Badhrees,^{69,33} S. Bahinipati,¹⁹ A. M. Bakich,⁶⁸ V. Bansal,⁵⁹ E. Barberio,⁴⁶ M. Berger,⁶⁶ V. Bhardwaj,¹⁸ B. Bhuyan,²⁰ J. Biswal,²⁹ A. Bobrov,^{4,58} A. Bondar,^{4,58} G. Bonvicini,⁸² A. Bozek,⁵⁵ M. Bračko,^{44,29} T. E. Browder,¹⁴ D. Červenkov,⁵ V. Chekelian,⁴⁵ A. Chen,⁵² B. G. Cheon,¹³ K. Chilikin,^{40,48} R. Chistov,^{40,48} K. Cho,³⁵ S. -K. Choi,¹² Y. Choi,⁶⁷ D. Cinabro,⁸² N. Dash,¹⁹ S. Di Carlo,⁸² Z. Doležal,⁵ Z. Drásal,⁵ D. Dutta,⁷⁰ S. Eidelman,^{4,58} D. Epifanov,^{4,58} H. Farhat,⁸² J. E. Fast,⁵⁹ T. Ferber,⁸ B. G. Fulsom,⁵⁹ V. Gaur,⁸¹ N. Gabyshev,^{4,58} A. Garmash,^{4,58} R. Gillard,⁸² P. Goldenzweig,³¹ D. Greenwald,⁷² J. Haba,^{15,11} T. Hara,^{15,11} K. Hayasaka,⁵⁷ M. T. Hedges,¹⁴ W. -S. Hou,⁵⁴ K. Inami,⁵⁰ A. Ishikawa,⁷⁴ R. Itoh,^{15,11} Y. Iwasaki,¹⁵ W. W. Jacobs,²² I. Jaegle,⁹ H. B. Jeon,³⁸ Y. Jin,⁷⁶ D. Joffe,³² K. K. Joo,⁶ T. Julius,⁴⁶ A. B. Kaliyar,²¹ K. H. Kang,³⁸ G. Karyan,⁸ T. Kawasaki,⁵⁷ C. Kiesling,⁴⁵ D. Y. Kim,⁶⁵ J. B. Kim,³⁶ K. T. Kim,³⁶ M. J. Kim,³⁸ S. H. Kim,¹³ Y. J. Kim,³⁵ K. Kinoshita,⁷ P. Kodyš,⁵ S. Korpar,^{44,29} D. Kotchetkov,¹⁴ P. Krizan,^{41,29} P. Krokovny,^{4,58} T. Kuhr,⁴² R. Kulasiri,³² R. Kumar,⁶¹ T. Kumita,⁷⁸ A. Kuzmin,^{4,58} Y. -J. Kwon,⁸⁴ J. S. Lange,¹⁰ I. S. Lee,¹³ C. H. Li,⁴⁶ L. Li,⁶³ L. Li Gioi,⁴⁵ D. Liventsev,^{81,15} M. Lubej,²⁹ T. Luo,⁶⁰ M. Masuda,⁷⁵ T. Matsuda,⁴⁷ D. Matvienko,^{4,58} K. Miyabayashi,⁵¹ H. Miyata,⁵⁷ R. Mizuk,^{40,48,49} G. B. Mohanty,⁷⁰ S. Mohanty,^{70,80} H. K. Moon,³⁶ T. Mori,⁵⁰ R. Mussa,²⁷ K. R. Nakamura,¹⁵ M. Nakao,^{15,11} T. Nanut,²⁹ K. J. Nath,²⁰ Z. Natkaniec,⁵⁵ M. Nayak,^{82,15} M. Niiyama,³⁷ N. K. Nisar,⁶⁰ S. Nishida,^{15,11} S. Ogawa,⁷³ S. Okuno,³⁰ H. Ono,^{56,57} P. Pakhlov,^{40,48} G. Pakhlova,^{40,49} B. Pal,⁷ C. -S. Park,⁸⁴ H. Park,³⁸ S. Paul,⁷² L. Pesántez,³ R. Pestotnik,²⁹ L. E. Pilonen,⁸¹ C. Pulvermacher,¹⁵ M. Ritter,⁴² A. Rostomyan,⁸ Y. Sakai,^{15,11} M. Salehi,^{43,42} S. Sandilya,⁷ L. Santelj,¹⁵ T. Sanuki,⁷⁴ Y. Sato,⁵⁰ O. Schneider,³⁹ G. Schnell,^{1,17} C. Schwanda,²⁴ A. J. Schwartz,⁷ Y. Seino,⁵⁷ K. Senyo,⁸³ M. E. Sevir,⁴⁶ V. Shebalin,^{4,58} C. P. Shen,² T. -A. Shibata,⁷⁷ J. -G. Shiu,⁵⁴ B. Shwartz,^{4,58} F. Simon,^{45,71} R. Sinha,²⁶ A. Sokolov,²⁵ E. Solovieva,^{40,49} M. Starič,²⁹ J. F. Strube,⁵⁹ K. Sumisawa,^{15,11} T. Sumiyoshi,⁷⁸ M. Takizawa,^{64,16,62} U. Tamponi,^{27,79} K. Tanida,²⁸ F. Tenchini,⁴⁶ K. Trabelsi,^{15,11} M. Uchida,⁷⁷ S. Uehara,^{15,11} T. Uglov,^{40,49} Y. Unno,¹³ S. Uno,^{15,11} P. Urquijo,⁴⁶ C. Van Hulse,¹ G. Varner,¹⁴ A. Vinokurova,^{4,58} V. Vorobyev,^{4,58} A. Vossen,²² E. Waheed,⁴⁶ C. H. Wang,⁵³ M. -Z. Wang,⁵⁴ P. Wang,²³ M. Watanabe,⁵⁷ Y. Watanabe,³⁰ E. Widmann,⁶⁶ K. M. Williams,⁸¹ E. Won,³⁶ H. Yamamoto,⁷⁴ Y. Yamashita,⁵⁶ H. Ye,⁸ J. Yelton,⁹ Y. Yook,⁸⁴ C. Z. Yuan,²³ Y. Yusa,⁵⁷ Z. P. Zhang,⁶³ V. Zhilich,^{4,58} V. Zhukova,⁴⁸ V. Zhulanov,^{4,58} and A. Zupanc^{41,29}

(Belle Collaboration)

¹University of the Basque Country UPV/EHU, 48080 Bilbao²Beihang University, Beijing 100191³University of Bonn, 53115 Bonn⁴Budker Institute of Nuclear Physics SB RAS, Novosibirsk 630090⁵Faculty of Mathematics and Physics, Charles University, 121 16 Prague⁶Chonnam National University, Kwangju 660-701⁷University of Cincinnati, Cincinnati, Ohio 45221⁸Deutsches Elektronen-Synchrotron, 22607 Hamburg⁹University of Florida, Gainesville, Florida 32611¹⁰Justus-Liebig-Universität Gießen, 35392 Gießen¹¹SOKENDAI (The Graduate University for Advanced Studies), Hayama 240-0193¹²Gyeongsang National University, Chinju 660-701¹³Hanyang University, Seoul 133-791¹⁴University of Hawaii, Honolulu, Hawaii 96822¹⁵High Energy Accelerator Research Organization (KEK), Tsukuba 305-0801¹⁶J-PARC Branch, KEK Theory Center, High Energy Accelerator Research Organization (KEK), Tsukuba 305-0801¹⁷IKERBASQUE, Basque Foundation for Science, 48013 Bilbao¹⁸Indian Institute of Science Education and Research Mohali, SAS Nagar 140306¹⁹Indian Institute of Technology Bhubaneswar, Satya Nagar 751007²⁰Indian Institute of Technology Guwahati, Assam 781039²¹Indian Institute of Technology Madras, Chennai 600036²²Indiana University, Bloomington, Indiana 47408²³Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049²⁴Institute of High Energy Physics, Vienna 1050²⁵Institute for High Energy Physics, Protvino 142281²⁶Institute of Mathematical Sciences, Chennai 600113²⁷INFN-Sezione di Torino, 10125 Torino²⁸Advanced Science Research Center, Japan Atomic Energy Agency, Naka 319-1195

- ²⁹J. Stefan Institute, 1000 Ljubljana
³⁰Kanagawa University, Yokohama 221-8686
³¹Institut für Experimentelle Kernphysik, Karlsruher Institut für Technologie, 76131 Karlsruhe
³²Kennesaw State University, Kennesaw, Georgia 30144
³³King Abdulaziz City for Science and Technology, Riyadh 11442
³⁴Department of Physics, Faculty of Science, King Abdulaziz University, Jeddah 21589
³⁵Korea Institute of Science and Technology Information, Daejeon 305-806
³⁶Korea University, Seoul 136-713
³⁷Kyoto University, Kyoto 606-8502
³⁸Kyungpook National University, Daegu 702-701
³⁹École Polytechnique Fédérale de Lausanne (EPFL), Lausanne 1015
⁴⁰P.N. Lebedev Physical Institute of the Russian Academy of Sciences, Moscow 119991
⁴¹Faculty of Mathematics and Physics, University of Ljubljana, 1000 Ljubljana
⁴²Ludwig Maximilians University, 80539 Munich
⁴³University of Malaya, 50603 Kuala Lumpur
⁴⁴University of Maribor, 2000 Maribor
⁴⁵Max-Planck-Institut für Physik, 80805 München
⁴⁶School of Physics, University of Melbourne, Victoria 3010
⁴⁷University of Miyazaki, Miyazaki 889-2192
⁴⁸Moscow Physical Engineering Institute, Moscow 115409
⁴⁹Moscow Institute of Physics and Technology, Moscow Region 141700
⁵⁰Graduate School of Science, Nagoya University, Nagoya 464-8602
⁵¹Nara Women's University, Nara 630-8506
⁵²National Central University, Chung-li 32054
⁵³National United University, Miao Li 36003
⁵⁴Department of Physics, National Taiwan University, Taipei 10617
⁵⁵H. Niewodniczanski Institute of Nuclear Physics, Krakow 31-342
⁵⁶Nippon Dental University, Niigata 951-8580
⁵⁷Niigata University, Niigata 950-2181
⁵⁸Novosibirsk State University, Novosibirsk 630090
⁵⁹Pacific Northwest National Laboratory, Richland, Washington 99352
⁶⁰University of Pittsburgh, Pittsburgh, Pennsylvania 15260
⁶¹Punjab Agricultural University, Ludhiana 141004
⁶²Theoretical Research Division, Nishina Center, RIKEN, Saitama 351-0198
⁶³University of Science and Technology of China, Hefei 230026
⁶⁴Showa Pharmaceutical University, Tokyo 194-8543
⁶⁵Soongsil University, Seoul 156-743
⁶⁶Stefan Meyer Institute for Subatomic Physics, Vienna 1090
⁶⁷Sungkyunkwan University, Suwon 440-746
⁶⁸School of Physics, University of Sydney, New South Wales 2006
⁶⁹Department of Physics, Faculty of Science, University of Tabuk, Tabuk 71451
⁷⁰Tata Institute of Fundamental Research, Mumbai 400005
⁷¹Excellence Cluster Universe, Technische Universität München, 85748 Garching
⁷²Department of Physics, Technische Universität München, 85748 Garching
⁷³Toho University, Funabashi 274-8510
⁷⁴Department of Physics, Tohoku University, Sendai 980-8578
⁷⁵Earthquake Research Institute, University of Tokyo, Tokyo 113-0032
⁷⁶Department of Physics, University of Tokyo, Tokyo 113-0033
⁷⁷Tokyo Institute of Technology, Tokyo 152-8550
⁷⁸Tokyo Metropolitan University, Tokyo 192-0397
⁷⁹University of Torino, 10124 Torino
⁸⁰Utkal University, Bhubaneswar 751004
⁸¹Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061
⁸²Wayne State University, Detroit, Michigan 48202
⁸³Yamagata University, Yamagata 990-8560
⁸⁴Yonsei University, Seoul 120-749

(Received 17 March 2017; published 22 May 2017)

We report the first measurement of the T -odd moments in the decay $D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ from a data sample corresponding to an integrated luminosity of 966 fb^{-1} collected by the Belle experiment at the KEKB asymmetric-energy e^+e^- collider. From these moments we determine the CP -violation-sensitive

asymmetry $a_{CP}^{T\text{-odd}} = [-0.28 \pm 1.38(\text{stat.})_{-0.76}^{+0.23}(\text{syst.})] \times 10^{-3}$, which is consistent with no CP violation. In addition, we perform $a_{CP}^{T\text{-odd}}$ measurements in different regions of the $D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ phase space; these are also consistent with no CP violation.

DOI: [10.1103/PhysRevD.95.091101](https://doi.org/10.1103/PhysRevD.95.091101)

Standard Model (SM) CP violation, which is due to the Kobayashi-Maskawa mechanism [1], is very small [$\mathcal{O}(10^{-3})$] in interactions involving decays of charm hadrons. Hence, any enhancement with respect to the SM prediction can indicate new physics effects due to particles or interactions not included in the SM [2]. The decay $D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ has a self-conjugate final state that can be used for a precise test of CP symmetry. Due to its large branching fraction of 5.2% [3], one can isolate a sample of $\mathcal{O}(10^6)$ decays that allows a test at a precision of $\mathcal{O}(10^{-3})$. This decay has been studied once before [4] but with a sample of only 140 events. Here, we report the first measurement of the time-reversal (T) asymmetry in $D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ decays, which is sensitive to CP violation via the CPT theorem [5]. This is the first T asymmetry measurement for a D meson decay with two neutral particles in the final state, one of which is a π^0 meson.

For this measurement, we use the method described in Refs. [6–9]. Such T -violation-sensitive measurements are complementary to direct probes of CP violation because of the differing dependence on the strong-phase difference between the contributing amplitudes [10]. This method was used earlier by the FOCUS [11], BABAR [12,13], and LHCb [14] collaborations for similar studies in D^0 , D^+ , and D_s^+ decays. The measurement is performed by constructing the scalar triple product,

$$C_T = \mathbf{p}_1 \cdot (\mathbf{p}_2 \times \mathbf{p}_3), \quad (1)$$

where \mathbf{p}_1 , \mathbf{p}_2 , and \mathbf{p}_3 are the momenta of any three of the D^0 daughter particles. Similarly, \bar{C}_T is defined as the CP -conjugate observable with \bar{D}^0 daughter particles. There must be at least four particles in the final state for \mathbf{p}_1 to not be coplanar with \mathbf{p}_2 and \mathbf{p}_3 and allow nonzero C_T . We define two asymmetry parameters as

$$A_T = \frac{\Gamma(C_T > 0) - \Gamma(C_T < 0)}{\Gamma(C_T > 0) + \Gamma(C_T < 0)}, \quad (2)$$

$$\bar{A}_T = \frac{\Gamma(-\bar{C}_T > 0) - \Gamma(-\bar{C}_T < 0)}{\Gamma(-\bar{C}_T > 0) + \Gamma(-\bar{C}_T < 0)}, \quad (3)$$

for D^0 and \bar{D}^0 , respectively, with Γ being a partial decay rate. These asymmetries can be nonzero due to the final state interaction (FSI) effects [15]. These effects are eliminated by taking the difference between A_T and \bar{A}_T as

$$a_{CP}^{T\text{-odd}} = \frac{1}{2}(A_T - \bar{A}_T), \quad (4)$$

for which a nonzero value would be a clear signature of T violation [5].

In this paper, we also present measurements of $a_{CP}^{T\text{-odd}}$ in nine regions of the final state phase space. The regions are selected to isolate CP eigenstates such as $K_S^0 \omega$, vector-vector (VV) states such as $K^{*\pm} \rho^\mp$, Cabibbo-favored (CF) states such as $K^{*-} \pi^+ \pi^0$ and doubly Cabibbo-suppressed (DCS) states such as $K^{*+} \pi^- \pi^0$.

We reconstruct the final state in $e^+ e^- \rightarrow c \bar{c} \rightarrow D^{*+} X$ events [16], recorded by the Belle experiment, in which $D^{*+} \rightarrow D^0 \pi_{\text{slow}}^+$, $D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ and X is a collection of particles produced along with the D^{*+} meson. The π_{slow}^+ meson is so called because its momentum is low compared to the final state particles originating from the D^0 decay. We use the charge of π_{slow} to identify whether the accompanying candidate is a D^0 or a \bar{D}^0 meson.

The Belle detector [17] is located at the interaction region of the KEKB asymmetric-energy $e^+ e^-$ collider [18]. The analysis is performed with the full data sample corresponding to an integrated luminosity of 966 fb^{-1} collected at or near center-of-mass energies corresponding to the $\Upsilon(nS)$ ($n = 1, 2, 3, 4, 5$) resonances, where 74% of the sample is taken at the $\Upsilon(4S)$ peak. The subdetectors relevant to this measurement are: a tracking system comprising a silicon vertex detector (SVD) and a 50-layer central drift chamber (CDC), a particle identification system comprising a barrel like arrangement of time-of-flight (TOF) scintillation counters and an array of aerogel threshold Cherenkov counters (ACC), and a CsI(Tl) crystal-based electromagnetic calorimeter (ECL). These subdetectors are located inside a 1.5 T superconducting magnet.

Samples of Monte Carlo (MC) simulated data are used to optimize the selection criteria and to understand various types of background. The EvtGen [19] and Geant3 [20] software packages are used to generate the events and simulate the detector response, respectively. We also include initial and final state radiation effects [21] in the simulation study.

We require candidate π^\pm daughters of the D^0 and π_{slow}^\pm to have a distance of closest approach along and perpendicular to the e^+ beam direction of less than 3.0 and 0.5 cm; this removes tracks not originating from the interaction region. Furthermore, these track candidates need to be positively identified as pions based on the combined information from the CDC, TOF, and ACC. The pion identification

requirement has an efficiency of 88% [22] with the probability of misidentification of a kaon as a pion candidate of 8%. We select $K_S^0 \rightarrow \pi^+\pi^-$ candidates from pairs of oppositely charged tracks, both treated as pions. The two tracks are required to have a π - π invariant mass within $\pm 3\sigma$ of the K_S^0 mass [3], where σ is the mass resolution. The decay vertex of the K_S^0 candidates is required to be displaced from the e^+e^- interaction point by a transverse distance of greater than 0.22 cm for momenta greater than 1.5 GeV/c, and greater than 0.08 cm for momenta between 0.5 and 1.5 GeV/c [23]. We select π^0 meson candidates from pairs of photons reconstructed in the ECL. The photons have different minimum energy criteria of 50, 100, or 150 MeV, depending on whether they are reconstructed in the barrel, forward end cap, or backward end cap regions of the ECL, respectively. These criteria suppress the beam-related backgrounds, which are typically asymmetric in polar angle. A π^0 candidate is selected when the invariant mass of the photon pair lies between 115 and 145 MeV/ c^2 , which covers an asymmetric interval corresponding to 3σ about the nominal mass of the π^0 meson [3]. We require that π^0 candidates have momentum greater than 350 MeV/c to reduce combinatorial background from random combinations of particles not originating from $D^0 \rightarrow K_S^0\pi^+\pi^-\pi^0$ decays. We kinematically constrain the π^0 meson to its known mass [3] to improve the momentum resolution. We identify a $D^0 \rightarrow K_S^0\pi^+\pi^-\pi^0$ candidate if its reconstructed invariant mass (M_{D^0}) is between 1.80 and 1.95 GeV/ c^2 .

We select π_{slow}^+ candidates from the remaining pion candidates in the event that produce at least one hit in the SVD; this requirement reduces the multiplicity of candidates within an event. We form D^{*+} from the selected D^0 and π_{slow}^+ candidates. To eliminate D^* mesons from B decays, which have different kinematic and topological properties, we require the D^{*+} momentum in the center-of-mass frame to be greater than 2.5 GeV/c. A small contamination of 0.015% and 0.096% from B and B_s events, respectively, is found from MC simulation studies. We define the variable $\Delta M = M_{D^{*+}} - M_{D^0}$, where $M_{D^{*+}}$ is the mass of the D^{*+} candidate; this peaks at 145 MeV/ c^2 [3] for correctly reconstructed D^{*+} mesons. We require ΔM to be less than 150 MeV/ c^2 to suppress the combinatorial background. We perform kinematically constrained vertex fits for both the D^0 vertex (using the π^+ and π^- tracks, π^0 vertex, and K_S^0 momentum) and the D^{*+} vertex (using the D^0 momentum and π_{slow}^+ track). We remove very poorly reconstructed candidates whose vertex fit quality parameter exceeds 1000. We also apply a kinematically constrained mass fit for the D^0 meson candidates to improve the resolution of the momenta of D^0 daughters.

Selection criteria are chosen to maximize the significance $S/\sqrt{S+B}$, where $S(B)$ is the number of MC signal (background) events in the signal region, defined as

144–147 MeV/ c^2 for ΔM and 1.82–1.90 GeV/ c^2 for M_{D^0} . Two types of backgrounds are significant: (1) “combinatorial” and (2) “random π_{slow}^+ .” The latter consists of a correctly reconstructed $D^0 \rightarrow K_S^0\pi^+\pi^-\pi^0$ decay paired with a π_{slow}^+ candidate that is not from a common D^{*+} parent. The background contributions in the selected data sample are 55% and 1% for combinatorial and random π_{slow}^+ components, respectively. The signal purity is 79% in the signal region. The selection efficiency estimated from MC simulation is 4%, and the selected data sample contains 1,691,029 events.

The selection results in an average multiplicity of 1.5 D^* candidates per event. In events with two or more candidates, we retain for further analysis the one with the smallest χ^2 value of the D^* vertex. MC studies indicate that this requirement selects the correct candidate in 74% of the events with multiple candidates.

We define C_T in the D^0 rest frame as $\mathbf{p}_{K_S^0} \cdot (\mathbf{p}_{\pi^+} \times \mathbf{p}_{\pi^-})$ for D^0 events and \bar{C}_T for \bar{D}^0 as $\mathbf{p}_{K_S^0} \cdot (\mathbf{p}_{\pi^-} \times \mathbf{p}_{\pi^+})$; the values of $|C_T|$ and $|\bar{C}_T|$ with other combinations of final state particles are found to yield identical results. To determine $a_{CP}^{T\text{-odd}}$, we first divide the data sample into four categories using the C_T value and π_{slow} charge: (i) D^0 with $C_T > 0$, (ii) D^0 with $C_T < 0$, (iii) \bar{D}^0 with $-\bar{C}_T > 0$, and (iv) \bar{D}^0 with $-\bar{C}_T < 0$. We then perform a simultaneous maximum likelihood fit to the two-dimensional distributions of ΔM and M_{D^0} to determine $a_{CP}^{T\text{-odd}}$ and yields. The two yields [(i) and (iii)] and two asymmetry parameters (A_T and $a_{CP}^{T\text{-odd}}$) of the signal component are floated in the fit.

We model the signal component of the M_{D^0} distribution with a probability density function (PDF) that is the sum of a Crystal Ball (CB) function [24], a Landau distribution, and two Gaussian functions, with a common value for the Gaussian means and Landau central value. The combinatorial background component is parametrized with a first-order polynomial. The random π_{slow}^+ component is modeled by the signal PDF.

The ΔM signal component is described by a PDF formed from the sum of a CB function, two Gaussians, and an asymmetric Gaussian function. The combinatorial component is parametrized by a PDF that is the sum of an empirical threshold function and a Gaussian function. The threshold function has the form

$$f(\Delta M) = a(\Delta M - m_\pi)^\alpha \exp[-\beta(\Delta M - m_\pi)], \quad (5)$$

where a is the normalization parameter, α and β are shape parameters, and m_π is the mass of the charged pion [3]. We observe a small peaking structure in the signal region of the ΔM combinatorial background distribution that is due to partially reconstructed D^0 candidates associated with a genuine π_{slow}^+ , such as a correctly reconstructed $D^{*+} \rightarrow D^0\pi_{\text{slow}}^+$, $D^0 \rightarrow K_S^0\pi^+\pi^-$ event combined with a




FIG. 1. The signal-enhanced logarithmic distributions of (a) ΔM and (b) M_{D^0} for D^0 with $C_T > 0$, (c) ΔM and (d) M_{D^0} for D^0 with $C_T < 0$, (e) ΔM and (f) M_{D^0} for \bar{D}^0 with $-\bar{C}_T > 0$ and (g) ΔM and (h) M_{D^0} for \bar{D}^0 with $-\bar{C}_T < 0$; the ΔM distributions have a selection criteria on M_{D^0} in the signal region and vice versa. The black points with error bars are the data points and the solid blue curve is the projection of the total signal and background components. The dotted magenta and dashed red curves indicate combinatorial and random π_{slow}^+ backgrounds, respectively. The normalized residuals (pulls) and the χ^2/DoF , where DoF is the number of degrees of freedom, are shown above each plot.

low momentum π^0 from the rest of the event. We fix the Gaussian parameters and the fraction of Gaussian contribution of the ΔM combinatorial background PDF to those obtained from the MC sample. The random π_{slow}^+ component is modeled with the same threshold function as the combinatorial background.

We calculate signal yields via a two-dimensional unbinned maximum likelihood fit to the values ΔM and M_{D^0} . To perform this fit, we include a small correlation term in the PDFs between the width of ΔM and the value of M_{D^0} . We parametrize the width of the dominant signal-component Gaussian of ΔM as

$$\sigma(\Delta M) = \sigma(\Delta M)|_{m_{D^0}} + a_\sigma (M_{D^0} - m_{D^0})^2, \quad (6)$$

where a_σ is a constant and m_{D^0} is the known mass of the D^0 meson [3].

The background component yields for all four samples are floated independently, but the shape parameters are common for the four categories. In total, there are 21 free and nine fixed parameters in the fit. The parameters fixed from MC are one of the widths of the asymmetric Gaussian, the width and exponent of the CB PDFs in the ΔM signal component, the normalization parameter a in the threshold

PDF, three Gaussian parameters for the peaking structure in the combinatorial background, the relative contribution of the CB and Gaussian functions to the M_{D^0} PDF of the random π_{slow}^+ component, and the fraction of PDF that contains the correlation in the two-dimensional signal PDF of ΔM and M_{D^0} . The signal-enhanced ΔM and M_{D^0} distributions of the data for the four categories are shown in Fig. 1, along with the fit projections. The total signal yield obtained from the fit is $744,509 \pm 1,622$ and the asymmetries are $A_T = (11.60 \pm 0.19)\%$ and $a_{CP}^{T\text{-odd}} = (-0.28 \pm 1.38) \times 10^{-3}$, where the uncertainties are statistical. The nonuniform pull for the ΔM fits is due to the remaining correlation between ΔM and M_{D^0} . However, from MC studies we find that this correlation does not cause any bias in the signal yields, in A_T , nor in $a_{CP}^{T\text{-odd}}$. The large value for A_T is due to the FSI effects [15]. The value of $a_{CP}^{T\text{-odd}}$ is consistent with no CP violation.

We divide the $D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ phase space into nine exclusive regions according to the intermediate resonance contributions. These are (1) $K_S^0 \omega$ (CP eigenstate), (2) $K_S^0 \eta$ (CP eigenstate), (3) $K^{*-} \rho^+$ (VV CF state), (4) $K^{*+} \rho^-$ (VV DCS state), (5) $K^{*-} \pi^+ \pi^0$ (CF state), (6) $K^{*+} \pi^- \pi^0$ (DCS state), (7) $K^{*0} \pi^+ \pi^-$, (8) $K_S^0 \rho^+ \pi^-$ and (9) everything else.

TABLE I. A_T and $a_{CP}^{T\text{-odd}}$ values from different regions of $D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ phase space. $M_{ij[k]}$ indicates the invariant mass of mesons i and j [and k].

Bin	Resonance	Invariant mass requirement (MeV/ c^2)	$A_T (\times 10^{-2})$	$a_{CP}^{T\text{-odd}} (\times 10^{-3})$
1	$K_S^0 \omega$	$762 < M_{\pi^+ \pi^- \pi^0} < 802$	$3.6 \pm 0.5 \pm 0.5$	$-1.7 \pm 3.2 \pm 0.7$
2	$K_S^0 \eta$	$M_{\pi^+ \pi^- \pi^0} < 590$	$0.2 \pm 1.3 \pm 0.4$	$4.6 \pm 9.5 \pm 0.2$
3	$K^{*-} \rho^+$	$790 < M_{K_S^0 \pi^-} < 994$	$6.9 \pm 0.3^{+0.6}_{-0.5}$	$0.0 \pm 2.0^{+1.6}_{-1.4}$
		$610 < M_{\pi^+ \pi^0} < 960$		
4	$K^{*+} \rho^-$	$790 < M_{K_S^0 \pi^+} < 994$	$22.0 \pm 0.6 \pm 0.6$	$1.2 \pm 4.4^{+0.3}_{-0.4}$
		$610 < M_{\pi^- \pi^0} < 960$		
5	$K^{*-} \pi^+ \pi^0$	$790 < M_{K_S^0 \pi^-} < 994$	$25.5 \pm 0.7 \pm 0.5$	$-7.1 \pm 5.2^{+1.2}_{-1.3}$
6	$K^{*+} \pi^- \pi^0$	$790 < M_{K_S^0 \pi^+} < 994$	$24.5 \pm 1.0^{+0.7}_{-0.6}$	$-3.9 \pm 7.3^{+2.4}_{-1.2}$
7	$K^{*0} \pi^+ \pi^-$	$790 < M_{K_S^0 \pi^0} < 994$	$19.7 \pm 0.8^{+0.4}_{-0.5}$	$0.0 \pm 5.6^{+1.1}_{-0.9}$
8	$K_S^0 \rho^+ \pi^-$	$610 < M_{\pi^+ \pi^0} < 960$	$13.2 \pm 0.9 \pm 0.4$	$7.6 \pm 6.1^{+0.2}_{-0.0}$
9	Remainder	\dots	$20.5 \pm 1.0^{+0.5}_{-0.6}$	$1.8 \pm 7.4^{+2.1}_{-5.3}$

Due to the relatively small size of these samples in comparison with the combined one, we reduce the number of free shape parameters to six while fitting the distributions of ΔM and M_{D^0} in each bin. The remaining parameters are fixed to the values obtained from the fit to the combined data sample. The free parameters are the mean and the width of the ΔM signal component and the four CB parameters for the M_{D^0} signal component. The A_T and $a_{CP}^{T\text{-odd}}$ values in each bin are listed in Table I. The results for $a_{CP}^{T\text{-odd}}$ are all consistent with no CP violation. The values of A_T vary significantly due to the different resonance contributions. A value $A_T \approx 0$ indicates the presence of a single partial wave, as in bin 2 where the S -wave dominates. Values of $A_T > 0$ indicate a significant interference between even and odd partial waves as in bins 3 to 9 [25].

The sources of systematic uncertainties are the signal and background models, efficiency dependence on C_T , C_T resolution, and potential fit bias. The dominant contribution comes from modeling the signal and background PDFs. The fixed parameters in the fit not related to the peaking combinatorial background are varied by ± 1 standard deviation from their nominal value obtained from a simulation sample corresponding to the same integrated luminosity as the data; we assign the change in $a_{CP}^{T\text{-odd}}$ as a systematic uncertainty. Without having a suitable control sample to study the peaking component of the combinatorial background, we change the value of the fraction of Gaussian PDF to twice the value found in the MC sample and then to zero. The resulting changes $+0.02 \times 10^{-3}$ and -0.42×10^{-3} , respectively, for $a_{CP}^{T\text{-odd}}$ are assigned as a systematic uncertainty. These uncertainties are combined, accounting for correlations among the parameters, to give a total uncertainty of $^{+0.09}_{-0.73} \times 10^{-3}$.

To study the dependence of the efficiency on C_T , we calculate the efficiency in ten bins of C_T between $-0.05(\text{GeV}/c)^3$ and $0.05(\text{GeV}/c)^3$. We find a relative spread of 10% in efficiency across the bins that varies

quadratically as $c_2 C_T^2 + c_1 C_T + c_0$, where $c_1 = 0$ within its statistical limit. This dependence is due to a reduced reconstruction efficiency for low-momentum D^0 daughters, which tend to have C_T values close to zero. We correct the measured $a_{CP}^{T\text{-odd}}$ value for the efficiency dependence and see negligible change because of the symmetry implied by $c_1 = 0$. We introduce an artificial asymmetry by changing the value of c_1 by one standard deviation and perform the efficiency correction again. The change in $a_{CP}^{T\text{-odd}}$ of 0.05×10^{-3} is assigned as the systematic uncertainty due to the C_T efficiency dependence. The parameter c_2 is found to be different for D^0 and \bar{D}^0 but still compatible within uncertainties. We take the difference of 0.20×10^{-3} in $a_{CP}^{T\text{-odd}}$ when applying different efficiency corrections for D^0 and \bar{D}^0 as a systematic uncertainty. The C_T resolution follows a Cauchy distribution with zero mean and a half width at half maximum of $1.325 (\text{MeV}/c)^3$. We add a corresponding smearing to the C_T distribution to determine a systematic change in $a_{CP}^{T\text{-odd}}$ due to any asymmetric cross feed between the positive and negative C_T intervals. The variation in $a_{CP}^{T\text{-odd}}$ due to the migration is 0.02×10^{-3} , which is taken as a systematic uncertainty from this source. We obtain the fit bias systematic uncertainty, which is a multiplicative one, from a linearity test by giving different input values for $a_{CP}^{T\text{-odd}}$ in sets of simulated pseudo-experiments. We find a possible fit-bias uncertainty of 0.28×10^{-5} . We add all the individual systematic uncertainties in quadrature to obtain a total $a_{CP}^{T\text{-odd}}$ systematic uncertainty of $^{+0.76}_{-0.23} \times 10^{-3}$.

In addition to the systematic studies, we perform other cross-checks. There is an asymmetry between the number of D^0 and \bar{D}^0 events reconstructed in the data sample due to the forward-backward asymmetry (A_{FB}) generated by interference between the virtual photon and Z^0 boson [26]. This production asymmetry, coupled with the asymmetry of the Belle detector, may induce a different reconstruction

efficiency as a function of C_T for D^0 and \bar{D}^0 . This asymmetry is modeled in the MC samples and is found to introduce no bias to the measured value of $a_{CP}^{T\text{-odd}}$. We also measure $a_{CP}^{T\text{-odd}}$ in bins of $\cos\theta^*$, where θ^* is the polar angle of the D^{*+} with respect to the e^+ beam direction defined in the center-of-mass system, and find that the results are consistent with the integrated value. To check for any further systematic effect due to detector reconstruction asymmetry for particles of different charges, we compare the momentum and azimuthal angle distributions for D^0 and \bar{D}^0 daughters in data and MC samples and find no significant difference. Furthermore, we study the dependence of the C_T distribution on the D^{*+} momentum selection criterion by varying the latter value by ± 100 MeV/ c . No significant change in the shape of the C_T distribution is observed. In addition, we estimate the possible contamination from the decay $D^0 \rightarrow \pi^+\pi^-\pi^+\pi^-$, which is an irreducible background, and find that the contribution is negligible.

In summary, we report the first measurement of the T -odd moment asymmetry $a_{CP}^{T\text{-odd}} = (-0.28 \pm 1.38^{+0.23}_{-0.76}) \times 10^{-3}$ for $D^0 \rightarrow K_S^0\pi^+\pi^-\pi^0$, consistent with no CP violation. The results in various bins of $K_S^0\pi^+\pi^-\pi^0$ phase space also show no evidence for CP violation. This result constitutes one of the most precise tests of CP violation in the D meson system [3]. The measurement uncertainties are statistically dominated and thus can be improved further with the data from the upcoming Belle II experiment [27].

We thank the KEKB group for the excellent operation of the accelerator; the KEK cryogenics group for the efficient operation of the solenoid; and the KEK computer group, the National Institute of Informatics, and the PNNL/EMSL computing group for valuable computing and SINET5 network support. We acknowledge support from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan, the Japan Society for the Promotion of Science (JSPS), and the Tau-Lepton Physics

Research Center of Nagoya University; the Australian Research Council; Austrian Science Fund under Grant No. P26794-N20; the National Natural Science Foundation of China under Contracts No. 10575109, No. 10775142, No. 10875115, No. 11175187, No. 11475187, No. 11521505 and No. 11575017; the Chinese Academy of Science Center for Excellence in Particle Physics; the Ministry of Education, Youth and Sports of the Czech Republic under Contract No. LTT17020; the Carl Zeiss Foundation, the Deutsche Forschungsgemeinschaft, the Excellence Cluster Universe, and the VolkswagenStiftung; the Department of Science and Technology of India; the Istituto Nazionale di Fisica Nucleare of Italy; the WCU program of the Ministry of Education, National Research Foundation (NRF) of Korea Grants No. 2011-0029457, No. 2012-0008143, No. 2014R1A2A2A01005286, No. 2014R1A2A2A0102734, No. 2015R1A2A2A01003280, No. 2015H1A2A1033649, No. 2016R1D1A1B01010135, No. 2016K1A3A7A09005603, No. 2016K1A3A7A09005604, No. 2016R1D1A1B02012900, No. 2016K1A3A7A09005606, No. NRF-2013K1A3A7A06056592; the Brain Korea 21-Plus program, Radiation Science Research Institute, Foreign Large-size Research Facility Application Supporting project and the Global Science Experimental Data Hub Center of the Korea Institute of Science and Technology Information; the Polish Ministry of Science and Higher Education and the National Science Center; the Ministry of Education and Science of the Russian Federation and the Russian Foundation for Basic Research; the Slovenian Research Agency; Ikerbasque, Basque Foundation for Science and the Euskal Herriko Unibertsitatea (UPV/EHU) under program UFI 11/55 (Spain); the Swiss National Science Foundation; the Ministry of Education and the Ministry of Science and Technology of Taiwan; and the U.S. Department of Energy and the National Science Foundation.

-
- [1] M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* **49**, 652 (1973).
 - [2] Y. Grossman, A. L. Kagan, and Y. Nir, *Phys. Rev. D* **75**, 036008 (2007).
 - [3] C. Patrignani *et al.* (Particle Data Group Collaboration), *Chin. Phys. C* **40**, 100001 (2016).
 - [4] D. Coffman *et al.* (MARK III Collaboration), *Phys. Rev. D* **45**, 2196 (1992).
 - [5] G. Lüders, Det. Kong. Dansk. Vidensk. Selsk., Mat.-fys. Medd. **28**, 005 (1954).
 - [6] E. Golowich and G. Valencia, *Phys. Rev. D* **40**, 112 (1989).
 - [7] W. Bensalem and D. London, *Phys. Rev. D* **64**, 116003 (2001).
 - [8] W. Bensalem, A. Datta, and D. London, *Phys. Rev. D* **66**, 094004 (2002).
 - [9] W. Bensalem, A. Datta, and D. London, *Phys. Lett. B* **538**, 309 (2002).
 - [10] G. Valencia, *Phys. Rev. D* **39**, 3339 (1989).
 - [11] J. M. Link *et al.* (FOCUS Collaboration), *Phys. Lett. B* **622**, 239 (2005).
 - [12] P. del Amo Sanchez *et al.* (BABAR Collaboration), *Phys. Rev. D* **81**, 111103 (2010).
 - [13] J. P. Lees *et al.* (BABAR Collaboration), *Phys. Rev. D* **84**, 031103 (2011).
 - [14] R. Aaij *et al.* (LHCb Collaboration), *J. High Energy Phys.* **40** (2014) 005.

K. PRASANTH *et al.*

PHYSICAL REVIEW D **95**, 091101(R) (2017)

- [15] I. I. Bigi, [arXiv:hep-ph/0107102](#).
- [16] Here, and elsewhere in this paper, charge conjugation is implied.
- [17] A. Abashian *et al.* (Belle Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **479**, 117 (2002); also see the detector section in J. Brodzicka *et al.*, *Prog. Theor. Exp. Phys.* (**2012**) 04D001.
- [18] S. Kurokawa and E. Kikutani, *Nucl. Instrum. Methods Phys. Res., Sect. A* **499**, 1 (2003), and other papers included in this volume; T. Abe *et al.* *Prog. Theor. Exp. Phys.* (**2013**) 03A001 and following articles up to 03A011.
- [19] D. J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
- [20] R. Brun *et al.*, Report No. CERN-DD-EE-84-1.
- [21] E. Barbiero and Z. Wąs, *Comput. Phys. Commun.* **79**, 291 (1994).
- [22] E. Nakano, *Nucl. Instrum. Methods Phys. Res., Sect. A* **494**, 402 (2002).
- [23] K.-F. Chen *et al.* (Belle Collaboration), *Phys. Rev. D* **72**, 012004 (2005).
- [24] T. Skwarnicki, Ph.D. thesis, Cracow, INP, 1986.
- [25] G. Durieux and Y. Grossman, *Phys. Rev. D* **92**, 076013 (2015); R. Sinha (private communication).
- [26] F. A. Berends, K. J. F. Gaemers, and R. Gastmans, *Nucl. Phys.* **B63**, 381 (1973); R. W. Brown, K. O. Mikaelian, V. K. Cung, and E. A. Paschos, *Phys. Lett. B* **43**, 403 (1973); R. J. Cashmore, C. M. Hawkes, B. W. Lynn, and R. G. Stuart, *Z. Phys. C* **30**, 125 (1986).
- [27] T. Abe *et al.* (Belle II Collaboration), [arXiv:1011.0352](#).