

Mobility of a spatially modulated electron liquid on the helium surface

K. Moskovtsev* and M. I. Dykman

Department of Physics and Astronomy, Michigan State University, East Lansing, Michigan 48824, USA



(Received 18 March 2020; revised manuscript received 16 May 2020; accepted 9 June 2020; published 24 June 2020)

We present the results of large-scale numerical simulations of the mobility of a two-dimensional electron liquid on the helium surface in the presence of a one-dimensional periodic potential. Even where the potential is much weaker than the electron-electron interaction, it can strongly change the mobility. The effect depends on the interrelation between the potential period and the mean interelectron distance. It is most pronounced where the period is close to the period of the Wigner crystal that would form if the liquid were cooled to a lower temperature. The results suggest, in particular, that the correlation length in the electron liquid can be found by measuring the mobility in a weak periodic potential. The simulations are based on the microscopic model of the electron scattering by the excitations in helium.

DOI: [10.1103/PhysRevB.101.245435](https://doi.org/10.1103/PhysRevB.101.245435)

I. INTRODUCTION

Electrons floating above the surface of liquid helium form a peculiar two-dimensional (2D) condensed-matter system. Its major distinctive features are the absence of a disorder potential and the strong electron correlations [1,2]. The absence of disorder makes it the best-known condensed-matter conductor, with the electron momentum relaxation time $\sim 10^{-7}$ s for $T \lesssim 0.3$ K [3,4]. The strong correlations lead to Wigner crystallization for low temperatures [5,6], with the Wigner crystal having unusual and not yet entirely understood properties [7,8]. On the higher-temperature side of the transition, the electrons form a liquid, which displays anomalous classical and quantum magnetotransport [9–11] and numerous nontrivial nonequilibrium phenomena, see Refs. [12–14] and references therein. We emphasize that, even though the electron liquid is nondegenerate, its dynamics is determined entirely by the electron-electron interaction. The interaction is not a perturbation.

Since the electrons float in free space with no leads attached, a major way of studying them is by measuring the response to a low-frequency electric field or to microwaves. The response to a spatially uniform field is not directly affected by the electron-electron interaction, which preserves the total momentum [15]. However, this interaction modifies, sometimes dramatically, the short-wavelength electron scattering by helium excitations, in particular by the surface capillary waves (ripplons) and phonons. In turn, this changes the electron transport compared to the single-electron transport, as observed in the experiments mentioned above, cf. also Refs. [16,17]. While the general picture of the many-electron transport on helium is commonly accepted, we are not aware of a direct observation of correlations in the electron liquid.

In this paper, we argue that the electron correlations in the liquid can be directly revealed by studying electron transport in the presence of a one-dimensional periodic potential with period close to the mean interelectron distance. We provide the results of numerical simulations of the electron mobility that substantiate this claim. Since the mean interelectron distance on helium is ~ 1 μm , a potential with the corresponding period can be created by a conventionally grown grating of electrodes submerged beneath the helium surface, as sketched in Fig. 1.

A strong effect of the periodic potential on the electron transport is easy to see already in a single-electron picture. Here, in the classical regime, the mobility $\mu_{\perp}$ transverse to the potential troughs will be thermally activated, $\mu_{\perp} \propto \exp(-\Delta U/k_B T)$, where ΔU is the difference between the maximum and the minimum of the potential. Strong correlations in the electron liquid modify this picture in two ways. First, if the potential is much weaker than the electron-electron interaction and has a period very different from the mean distance between the electrons a_s , the effect of the potential is partly averaged out, as there are electrons both near the minima and near the maxima of the potential. On the other hand, even where the potential is weak, but its period is close to a_s , the electron density can become strongly modulated. In this case, the mobility should strongly depend on the correlation length in the electron liquid, since moving the system with a modulated density as a whole over the periodic potential barriers is impossible, in the limit of a large system. Indeed, our simulations show a strong dependence of $\mu_{\perp}$ on the correlation length.

Placing a system of interacting particles into a one-dimensional (1D) periodic potential is known to affect the transition from the liquid to the ordered phase, cf. Refs. [18,19]. The change of the critical temperature and of the very character of the transition is particularly strong in a system with the Coulomb coupling [20]. This is because such a system displays a true crystalline order if the period of the potential is equal to the lattice constant of the system [21].

*Present address: SiTime Corporation, 5451 Patrick Henry Drive Santa Clara, CA 95054, USA




FIG. 1. Left panel: Sketch of the electron liquid floating above the helium surface with a submerged periodic grating of nanowires that create a periodic one-dimensional potential for the electrons. The potential is shown in the right panel. It is essentially sinusoidal, if the electrodes are submerged by a depth that noticeably exceeds the interelectrode spacing.

Understanding the transport of a crystal in a periodic potential is interesting and challenging, cf. recent papers [22–24] and references therein. However, for electrons on helium the analysis is complicated by a specific mechanism of scattering by helium excitations [25,26]. Therefore here we do not study the dynamics of the electron crystal and do not consider the rich area of commensurate-incommensurate transitions in a periodic potential, cf. Refs. [27–29]. Our goal in this paper is to reveal the features of the electron liquid.

Our analysis is based on numerical simulations of a classical 2D electron system on helium. Simulations of 2D electron systems have attracted significant attention over the years, cf. Refs. [20,30–41]. The classical and quantum Monte Carlo as well as molecular dynamics (MD) methods have been employed. However, the MD simulations were done by adding friction forces and uncorrelated noises that drive individual electrons. Such phenomenological description does not describe the dynamics of electrons on helium. Rather, the major mechanism of electron scattering for low temperatures is scattering by surface capillary waves, ripplons [42,43]. Such scattering is quasielastic, since ripplons are very slow.

We simulate the mobility directly by taking into account the scattering by ripplons and studying the current induced by a weak electric field applied to the electron system. Such a field unavoidably heats up the electron system. Therefore it is necessary to incorporate a mechanism of energy exchange between the electrons and helium. An important mechanism of such exchange is scattering by phonons [21,44]. Even though the corresponding scattering rate is small compared to the rate of scattering by ripplons, it is sufficient to avoid electron overheating in the range of the fields we study.

The microscopic picture of the electron relaxation for an electron liquid takes into account the forces that the electrons exert on each other and that affect their scattering by ripplons and phonons [9]; it also relies on the assumption of the fast exchange of energy and momentum between the electrons [9,10,16]. This exchange is faster than the electron scattering by the helium excitations. The results of the analysis based on this picture are in a good agreement with the experiment, but in fact, the assumption has not been tested directly. As we show, numerical simulations suggest a way to carry out this test. However, the central results of the paper refer to the mobility in a periodic potential and its dependence on the amplitude of the potential and the temperature.

Below in Sec. II, we describe the model of the system, briefly outline the simulations and indicate the parameter range where they apply. In Sec. III, we present results on the mobility of a uniform electron liquid. In Sec. IV, we describe the mobility in a sinusoidal periodic potential with the period incommensurate with the period of the hexagonal Wigner crystal for the studied electron density; however, we emphasize that our system is not a crystal, the lattice constant of the crystal is used just as the spatial scale. Section V is the central part of the paper. It describes the correlations in the electron system and the dependence of the electron mobility on temperature and the amplitude of the potential where the potential is commensurate with the would-be Wigner crystal. In particular, there is discussed the activated dependence of the mobility on the potential amplitude and its relation to the correlation length in the electron liquid. Section VI briefly presents the results on the dependence of the electron mobility on the commensurability parameter. Section VII contains concluding remarks.

II. MANY-ELECTRON SYSTEM ON HELIUM

A. The Hamiltonian

The Hamiltonian of the electron system coupled to the helium excitations has the form

$$H = H_{ee} + H_U + H_{He} + H_i. \quad (1)$$

The term H_{ee} is a sum of the electron kinetic energy and the energy of the electron-electron interaction, whereas H_U is the electron energy in the external potential,

$$H_{ee} = \sum_n \frac{\mathbf{p}_n^2}{2m_e} + \frac{1}{2} \sum_{n \neq m} \frac{e^2}{|\mathbf{r}_n - \mathbf{r}_m|},$$

$$H_U = \sum_n U(\mathbf{r}_n), \quad U(\mathbf{r}) = -A \cos Qx. \quad (2)$$

Here, $\mathbf{r}_n = (x_n, y_n)$ and $\mathbf{p}_n = (p_{xn}, p_{yn})$ are the 2D coordinate and momentum of an n th electron, and $U(\mathbf{r})$ is the external periodic potential. The electrodes creating the potential, see Fig. 1, partly screen the electron-electron interaction. The screening depends on the specific geometry, and for reasonably deeply submerged thin electrodes is comparatively weak. The partial screening does not destroy the long-range nature of the electron-electron interaction. Therefore it should not qualitatively change the mobility of the electron liquid. In what follows it is disregarded. We note that, in the experiment, the effect of the screening can be independently tested by varying the height of the helium layer and simultaneously changing the electrode potential. If the distance to the electrodes significantly exceeds the interelectrode spacing, the potential $U(\mathbf{r}_n)$ is sinusoidal, as indicated in Eq. (2), with $2\pi/Q$ being the interelectrode spacing and A being the amplitude of the potential.

The term H_{He} in Eq. (1) is the Hamiltonian of the excitations in the liquid helium that are coupled to the electrons. These are vibrational modes, i.e., ripplons and phonons,

$$H_{He} = \sum_{\mathbf{q}, \alpha} \hbar \omega_{\mathbf{q}\alpha} \hat{a}_{\mathbf{q}\alpha}^\dagger \hat{a}_{\mathbf{q}\alpha}. \quad (3)$$

Here, $\mathbf{q}$ is the 2D wave vector of a mode, $\omega_{\mathbf{q}\alpha}$ is the mode frequency, and $a_{\mathbf{q}\alpha}^\dagger$ and $\hat{a}_{\mathbf{q}\alpha}$ are creation and annihilation operators. For phonons, the quantum number α is the wave number $q_z > 0$ of motion transverse to the surface. Ripplons are surface waves, in this case α is just the label of the vibrational branch.

The Hamiltonian of the electron coupling to the vibrational modes is

$$H_i = \sum_n \sum_{\mathbf{q}, \alpha} V_{\mathbf{q}\alpha} e^{i\mathbf{q}\mathbf{r}_n} (\hat{a}_{\mathbf{q}\alpha} + \hat{a}_{-\mathbf{q}\alpha}^\dagger). \quad (4)$$

The coupling parameters $V_{\mathbf{q}\alpha}$ are well-known. For completeness, they are given in Appendix A. As mentioned previously, electron scattering by ripplons is the major mechanism of the electron momentum relaxation, whereas the scattering by phonons is a major mechanism of the energy relaxation; the other energy relaxation mechanism is two-ripplon scattering. Since we allow for energy relaxation primarily to avoid electron heating when the electrons are additionally driven by an electric field, we consider only one energy relaxation mechanism, the phonon scattering.

To study the conductivity of the electron liquid, we add to the Hamiltonian the term

$$H_d = eE_d \sum_n x_n, \quad (5)$$

where E_d is the driving field. This field is weak, we chose it to be much smaller than the fluctuational field that drives an electron due to electron density fluctuations and is $\sim n_s^{3/4} (k_B T)^{1/2}$ [9,35], where n_s is the electron density. As we show, for the values of E_d we use, the electron heating is weak.

The major parameters of the electron liquid are (i) the characteristic short-wavelength plasma frequency ω_p , which is the value of the long-wavelength plasma frequency $\omega_l(q) = \omega_p q^{1/2} n_s^{-1/4}$ extended to the wave number $q = n_s^{1/2}$, cf. Ref. [6], and (ii) the plasma parameter Γ , which is the ratio of the typical interaction energy per electron E_C to the kinetic energy (the analog of r_s in degenerate systems),

$$\begin{aligned} \omega_p &= (2\pi e^2 n_s^{3/2} / m_e)^{1/2}, \quad \Gamma = E_C / k_B T, \\ E_C &= e^2 (\pi n_s)^{1/2}. \end{aligned} \quad (6)$$

Parameter ω_p gives the typical rate of the momentum and energy exchange between the electrons, whereas Γ shows how strong the electron correlations are. In the absence of the periodic potential, Monte Carlo simulations suggest that the classical electron liquid crystallizes into a Wigner crystal for $\Gamma \approx 140$, cf. Ref. [41] and references therein.

We note that sometimes, particularly in comparing the simulated temperature of the Wigner crystallization with the experiment, the conditions that the electron system is classical and nondegenerate are tacitly presumed to be equivalent. However, a nondegenerate electron system is not necessarily classical, see Fig. 2. In the absence of a transverse magnetic field, the characteristic temperature that separates the classical and quantum electron dynamics can be chosen as $\hbar\omega_p/k_B$, where ω_p is given by Eq. (6). This temperature is an analog of the Debye temperature of the Wigner crystal, but it also characterizes the electron dynamics in the liquid phase. It is shown by the dashed line in Fig. 2. The majority of experiments on Wigner crystallization on helium were done




FIG. 2. The characteristic boundaries of the classical motion (blue dashed line) and Wigner crystallization (red solid line) on the temperature/electron density plane. The plasma frequency $\omega_p = (2\pi e^2 n_s^{3/2} / m)^{1/2}$ is the characteristic frequency of electron vibrations about their quasiequilibrium positions in a liquid or solid with the electron density n_s . The temperature T_Γ is the temperature given by the relation $T = e^2 (\pi n_s)^{1/2} / k_B \Gamma$, where Γ is the plasma parameter and n_s is the electron density. Recent Monte-Carlo simulations of the classical system [41] set the value of Γ for crystallization in the classical limit at $\Gamma = 140$.

for the electron densities $n_s > 10^8 \text{ cm}^{-2}$, where the simulated transition temperature is essentially in the quantum regime. A comparison with classical simulations requires going to lower densities or higher temperatures. Our simulations refer to the temperatures $T > \hbar\omega_p/k_B$.

B. Simulations

We simulate the electron dynamics in a periodic potential and a superimposed uniform electric field by integrating the many-electron equations of motion. In contrast to the standard molecular dynamics simulations, to find the electron mobility and to describe the features of the many-electron dynamics on helium, we explicitly incorporate scattering by the helium excitations into the equations of motion. This is done by considering scattering as a random event in which the electron momentum and energy change. The probability of scattering of a given electron within a short time interval is determined by the instantaneous value of the electron momentum and by the temperature of helium.

Both ripplons and phonons are short-range scatterers, therefore the probability for an electron to scatter is independent of the state of other electrons, i.e., the electrons are scattered independently of each other. The scattering events are rare, whereas the dynamics between the scattering events is fully controlled by the electron-electron interaction. This approach was used [21] to describe self-diffusion and Wigner crystallization in the absence of a driving electric field.

It should be emphasized that the self-diffusion in an interacting system does not give the long-wavelength diffusion coefficient. Therefore it does not give the electron mobility. This is particularly clear from noting that self-diffusion arises in an isolated electron system at a finite temperature, whereas the mobility is limited by the momentum transfer from the electrons to external scatterers, in our case, to ripplons and phonons in helium.

Direct simulation of scattering is essential to separate the momentum and energy relaxation in the system. For electrons on helium, the quasielastic ripplon scattering rate is at least an order of magnitude higher than that of inelastic scattering processes. Separating scattering mechanisms allows us to capture such effects as electron heating in the driven system. However, qualitatively, several effects of a periodic potential described below can also be reproduced in the standard molecular dynamics approach based on solving Langevin equations of motion with a friction force proportional to the electron velocity. The phenomenological friction coefficient has to be assumed small, so that the electron motion is strongly underdamped.

To obtain reliable results we used a comparatively large system of 1600 electrons placed into a rectangular area, with periodic boundary conditions. The ratio of the sides of the rectangular area along the y and x axes was $L_y/L_x = 2/\sqrt{3}$, which allows a Wigner crystal with hexagonal symmetry to fit into the area [30] (however, we studied the parameter range where Wigner crystallization did not occur). Importantly, we allowed the system to form a stationary distribution for a long time of $> 10^6$ integration steps per electron and collected the data for more than 10^7 steps. More details of the simulations are given in Appendix A.

III. MOBILITY OF A UNIFORM SYSTEM

The conductivity of a strongly correlated electron system on helium is not described by the standard Boltzmann kinetic equation. In this equation, the momentum of an individual electron is randomized by successive short collisions with the vibrational excitations in helium and, for elastic scattering, the conductivity is given by the Drude expression $\sigma_{xx} = e^2 n_s \langle \tau \rangle / m_e$, where $\langle \tau \rangle$ is the momentum relaxation time averaged over the Boltzmann distribution of the single-electron energy.

We are not aware of a closed-form equation for the single-particle distribution function in the case of strong electron-electron interaction. In the electron liquid, after a collision with a vibrational excitation, and prior to the next collision, the momentum and energy of an individual electron are randomized by the electron-electron interaction. The long-wavelength many-electron conductivity can be calculated starting with the Kubo formula and using a transport equation for the many-electron Green function. In the case of elastic scattering, this equation was derived earlier [9,45], with the account taken of a strong effect of the electron-electron interaction on the scattering by vibrational excitations in the presence of a magnetic field.

The analysis [45] immediately extends to the case of inelastic scattering. The corresponding theory is developed in Ref. [46].

Even though the many-electron theory of magnetoconductivity was in an excellent agreement with the experiment with no adjustable parameters [11], the underlying picture of the many-electron relaxation could not be tested directly. In contrast, realistic simulations of the electron dynamics provide a means for such testing, as they allow one not only to calculate the (generally nonlinear) conductivity of the many-electron system, but also to find the electron energy distribution.




FIG. 3. The drift velocity v_d (a) and the electron temperature T_{eff} (b) vs the driving field E_d for a uniform strongly correlated electron liquid. For $E_d = 0$, where the electron system is in thermal equilibrium with the excitations in helium, $\Gamma = 90$. The filled circles are the simulation data. In (a) the black solid line is the velocity $v_d = (e\tau/m_e)E_d$ calculated with the many-electron relaxation time τ given by the analytical expressions (18) and (19) of Ref. [46], which are evaluated for the same coupling to ripples and phonons and the same electron density as in the simulations. The inset in (b) shows the distribution of the electron kinetic energies in the driven system measured in the frame moving with the drift velocity $\mathbf{v}_d$ for $E_d/n_s e = 4.24 \times 10^{-5}$; the solid line shows the Boltzmann distribution.

The results of the simulations of the drift velocity of the spatially uniform electron liquid are presented in Fig. 3(a) and are compared with the theory. The driving field E_d is scaled by the field $n_s e$, which is the field created by an electron at the distance on the order of the mean interelectron distance; the studied fields E_d are much smaller than $n_s e$. The velocity is scaled by the factor $\omega_p n_s^{-1/2}$, which is the velocity required to go over the characteristic interelectron distance over the time $1/\omega_p$, and again, $v_d \ll \omega_p n_s^{-1/2}$.

For the parameters used in the simulations, the mobility $\mu = v_d/E_d$ for small E_d is $3.5 \times 10^7 \text{ cm}^2 \text{ V}^{-1} \text{ s}^{-1}$. We note that the rate of quasielastic ripplon scattering is proportional to temperature. Therefore the value of μ depends not only on Γ , but also independently on T , with $\mu \propto 1/T$ for a fixed Γ . The above value and the plot in Fig. 3 refer to $T = 0.354 \text{ K}$, or equivalently, to $n_s \approx 1.15 \times 10^8 \text{ cm}^{-2}$ for $\Gamma = 90$. Strictly speaking, for such n_s one should also take into account the coupling to ripples due to the field that presses the electrons against the helium surface [1]; this would reduce the numerical value of the mobility, but will not affect the qualitative results discussed below.

The simulations demonstrated that the distribution over the electron energy is the Boltzmann distribution with the effective temperature T_{eff} that increases with the driving field. This increase is shown in Fig. 3(b). Such form of the distribution confirms the underlying assumption of the analytical theory. For the fields E_d used in the simulations, the electron heating is weak and has a negligible effect on the mobility.

IV. ELECTRON LIQUID IN A PERIODIC POTENTIAL

The effect of the periodic potential on the electron system strongly depends on the interrelation between the period of the potential $2\pi/Q$ and the mean interelectron distance. More




FIG. 4. The mobility $\mu_{\perp} \equiv \mu_{xx}$ transverse to the troughs of the sinusoidal periodic potential (2) with amplitude A scaled by $\mu_{\text{eff}} = e/m_e\omega_p$, see Eq. (11). The results refer to $\Gamma = 90$ and the commensurability parameter $p_c = 0.3$. The squares and the circles show the data for the electron liquid and the ideal electron gas, respectively.

specifically, one can think of the electrons forming a Wigner crystal with a triangular lattice and define the commensurability parameter

$$p_c = (\sqrt{3}/2n_s)^{1/2}Q/2\pi \equiv a_s\sqrt{3}Q/4\pi, \quad (7)$$

where $a_s = (2/\sqrt{3}n_s)^{1/2}$ is the interelectron distance in the Wigner crystal.

For the considered one-dimensional potential, p_c gives the ratio of the distance between the rows of the Wigner crystal $a_s\sqrt{3}/2$ and the period of the potential. The rows here have been chosen in such a way as to minimize the interelectron distance in a row, see Appendix B. Depending on p_c , the periodic potential can lead to Wigner crystallization for smaller values of the plasma parameter Γ (6), i.e., for higher temperatures than in a uniform system, or can impede crystallization [21].

The effect of a weak potential on crystallization is small if p_c is such that a crystal has to be strongly distorted to fit into the periodic potential [28]. For a classical Wigner crystal, the effect is also small if p_c is small, as the electron-electron interaction effectively screens the potential. Respectively, one expects that not only the mobility along the potential troughs, but also the mobility transverse to the troughs $\mu_{\perp}$ will be weakly affected for small p_c . This is indeed seen in Fig. 4.

As mentioned in the Introduction, the mobility of the electron liquid in a periodic potential should strongly differ from that of an ideal electron gas. This is also seen in Fig. 4. The scale of the barrier height $2A$ on which the single-electron transverse mobility $\mu_{\perp}$ exponentially falls off should be given by the temperature. In Fig. 4, on the abscissa, the scaled amplitude of the periodic potential A/E_C can be written as

$\Gamma^{-1}(A/k_B T)$, and as expected, for the studied $\Gamma = 90$ the exponential fall-off of $\mu_{\perp}$ with A starts for $A/E_C \gtrsim 0.01$.

V. “MAXIMALLY COMMENSURATE” POTENTIAL

The effect of the periodic potential is most pronounced in the case where $p_c = 1$. In this case, if the electrons formed a crystal, it would be fully commensurate with the potential, cf. Appendix B. We call such a potential maximally commensurate, with one electron row per one potential trough. Even where the amplitude of the potential A is much smaller than the electron interaction energy E_C and the temperature is well above the temperature of the Wigner crystallization in a uniform system, the electron liquid preferentially occupies the potential troughs. Respectively, the electron density can be comparatively strongly periodically modulated.

The strong effect of the maximally commensurate potential comes from the fact that the potential lifts the translational and orientational symmetries of the electron liquid and, rather than competing with the electron correlations, it constructively interferes with them. Therefore such a potential can significantly modulate the electron density without changing the small-amplitude fluctuations about quasiequilibrium electron positions in the liquid.

The latter can be seen from the following argument. A single electron localized at the minimum of a potential trough vibrates normal to the trough with frequency $(AQ^2/m_e)^{1/2}$. On the other hand, an electron in an unconfined electron liquid vibrates about its quasiequilibrium position with frequency $\sim\omega_p$ (6). For $p_c = 1$, the ratio of the squares of these frequencies is $(A/E_C)4\pi^{3/2}/\sqrt{3}$. In the range of the potential strengths and electron densities that we studied this ratio was very small (it is equal to ≈ 0.02 for $A/E_C = 0.0016$).

By the same token, a maximally commensurate potential may be expected to weakly affect the relative positions of the electrons. Those are characterized by the two-particle correlation function. For a system in a periodic potential, it can be defined as

$$g_U^{(2)}(\mathbf{r}', \mathbf{r}'') = \frac{1}{\rho_s(\mathbf{r}')\rho_s(\mathbf{r}'')} \sum_{n,m}' \delta(\mathbf{r}' - \mathbf{r}_n)\delta(\mathbf{r}'' - \mathbf{r}_m), \quad (8)$$

where $\rho_s(\mathbf{r})$ is the electron density. For $\rho_s(\mathbf{r}) = \text{const}$, Eq. (8) goes over into the standard expression for the pair correlation function of a spatially uniform system.

For the considered potential $U(\mathbf{r})$, the density $\rho_s(\mathbf{r})$ is periodic in x with period $2\pi/Q$ and is independent of y ; its average value is n_s ,

$$\rho_s(\mathbf{r}) = n_s \left[1 + \sum_m \alpha_m \cos(mQx) \right]. \quad (9)$$

The coefficients α_m quickly fall off with the increasing m for a weak potential, where they can be found by a perturbation theory in $A/k_B T$. They can be found also if the electrons are strongly confined within the troughs, see Appendix B.

An example of $\rho_s(\mathbf{r})$ is shown in Fig. 5. The result refers to a small ratio of the potential amplitude to temperature, $A/k_B T = \Gamma A/E_C \approx 0.14$. The modulation of $\rho_s(\mathbf{r})$ is much stronger than it would be in the single-electron picture ($\propto \exp[-(A/k_B T) \cos Qx]$). This is the result of the commensu-




FIG. 5. (a) The pair correlation function $\tilde{g}^{(2)}(\mathbf{r})$, Eq. (10), for the maximally commensurate periodic potential, $p_c = 1$. The scaled amplitude of the potential is $A/E_c = 0.00157$, and the plasma parameter is $\Gamma = 90$. (b) The pair correlation function in the absence of a potential for the same $\Gamma = 90$. (c) The averaged over time electron density along the direction x of the oscillations of the potential for the same parameters as in (a). (d) One-dimensional slices of the pair correlation function $\tilde{g}^{(2)}(\mathbf{r})$ in (a).

stability: the potential constructively interferes with the strong electron correlations and just lifts the translational and rotational degeneracy of the system.

The function $g_U^{(2)}(\mathbf{r}', \mathbf{r}'')$ defined by Eq. (8) depends on $y' - y''$ and does not change if both x' and x'' are incremented by the period of the potential $2\pi/Q$. Therefore, if written as a function of $x' - x''$ and $(x' + x'')/2$, it is a periodic function of $(x' + x'')/2$. In addition, $g_U^{(2)}(\mathbf{r}', \mathbf{r}'') \rightarrow 1$ for $|x' - x''|, |y' - y''| \rightarrow \infty$.

One can also consider the zeroth Fourier component of $g_U^{(2)}(\mathbf{r}', \mathbf{r}'')$ with respect to $(x' + x'')/2$ and introduce the function

$$\begin{aligned} \tilde{g}^{(2)}(\mathbf{r}) &= \frac{n_s}{N} \int d\mathbf{R} g_U^{(2)}\left(\mathbf{R} + \frac{1}{2}\mathbf{r}, \mathbf{R} - \frac{1}{2}\mathbf{r}\right) \\ &= \frac{n_s}{N} \sum'_{n,m} \delta[\mathbf{r} - (\mathbf{r}_n - \mathbf{r}_m)] / [\rho_s(\mathbf{r}_n)\rho_s(\mathbf{r}_m)], \end{aligned} \quad (10)$$

where N is the total number of electrons. Function $\tilde{g}^{(2)}(\mathbf{r})$ is an analog of the pair correlation function of a uniform system.

The correlation function $\tilde{g}^{(2)}(\mathbf{r})$ is shown in Fig. 5. Remarkably, even though the electron density is strongly periodically modulated by the weak periodic potential used in the calculation, $\tilde{g}^{(2)}$ is very similar to the pair correlation function of a spatially uniform system for the same Γ , which is also shown in Fig. 5. Both with and without the potential, the correlations in the electron liquid decay over several interelectron distances. However, because of the potential, the decay of $\tilde{g}^{(2)}(\mathbf{r})$ in the modulation direction (the x axis) is slightly slower. Moreover, as seen from Fig. 5(a), $\tilde{g}^{(2)}(\mathbf{r})$ shows sixfold symmetry, with the y axis and the directions




FIG. 6. Delaunay triangulation for a snapshot of the electron liquid in the periodic potential of amplitude $A/E_c = 0.00157$; $p_c = 1$. The electrons are represented by vertices. The plasma parameter is $\Gamma = 50$ and 90 for the left and right panels, respectively.

obtained by rotating this axis by 60° being the directions of the fastest decay.

Decay of electron correlations can be also seen from the Delaunay triangulation shown in Fig. 6. The short-range order is pronounced for the both values of the plasma parameter Γ in this figure. However, for larger Γ , the number of unbound vortices with 5 or 7 nearest neighbors is smaller. As Γ increases, such defects play an increasingly important role in the electron dynamics.

It is also seen from Fig. 6 that, for $\Gamma = 90$, already a single snapshot shows that the density is anisotropic and periodically modulated along the x axis. For the shown snapshot, the Fourier spectrum of the density as a function of x displays a pronounced peak at the wave number Q of the potential. The peak has height $\approx 0.25/n_s$ and width at half-maximum $\approx 0.05/a_s$. Visible in the spectrum is also a peak at $2Q$, but with a smaller amplitude and a larger width. The overall spectrum is noisy for a single snapshot, in contrast to the results of Fig. 5(c) obtained by averaging over many snapshots.

A. Electron mobility transverse to the potential troughs

Placing the electron system into a sufficiently strong commensurate periodic potential should strongly affect the electron mobility. In particular, the mobility transverse to the potential troughs $\mu_\perp$ should decrease. This decrease should sensitively depend on the amplitude of the potential and the temperature. In turn, as we show, this dependence may be used to reveal and characterize the correlations in the system.

It is instructive to compare the mobility $\mu_\perp$ with the self-diffusion coefficient. For a spatially uniform classical 2D electron liquid, the self-diffusion coefficient is of the order of $D_0 = k_B T / m_e \omega_p$ [21], an estimate close to the De Gennes estimate for a normal three-dimensional liquid [50]. In our simulations, we used a fixed electron density and varied the temperature. Then one can scale the self-diffusion coefficient and the mobility by the temperature-independent parameters D_{eff} and μ_{eff} , respectively,

$$D_{\text{eff}} = \omega_p / n_s = 2\sqrt{\pi} \Gamma D_0, \quad \mu_{\text{eff}} = e / m_e \omega_p. \quad (11)$$

For the electron liquid in the absence of a periodic potential and far from the crystallization transition, we expect $D \sim$




FIG. 7. The mobility of the electron liquid $\mu_{\perp}$ transverse to the troughs of the periodic potential (left y axis) and the coefficients of self-diffusion along ($D_{\parallel}$) and transverse ($D_{\perp}$) to the troughs (right y axis) as functions of the reciprocal temperature $T^{-1} \propto \Gamma$ (the electron density is fixed). The results refer to the maximally commensurate potential, $p_c = 1$. The amplitudes of the potential in the left and right panels are $A/E_C = 0.000628$ and 0.00157 , respectively.

$D_0 \ll D_{\text{eff}}$ and $\mu_{\perp} \sim e\tau/m_e \gg \mu_{\text{eff}}$, where τ is the electron relaxation time due to the scattering by ripples, $\tau \gg \omega_p^{-1}$.

When the electron liquid is in a 1D periodic potential, both the mobility and the self-diffusion become anisotropic. Similar to the case of the mobility, one can introduce the coefficients of self-diffusion along and transverse to the troughs, $D_{\parallel}$ and $D_{\perp}$, respectively. They are defined by the long-time mean-square displacements of an electron in the corresponding directions, $\langle [y_n(t) - y_n(0)]^2 \rangle = 2D_{\parallel}t$ and $\langle [x_n(t) - x_n(0)]^2 \rangle = 2D_{\perp}t$. Simulations of these coefficients are described in Ref. [21].

The mobility along the troughs is limited by the scattering by ripples and should be weakly affected by the potential. We have indeed seen this in the simulations and do not discuss this mobility. In contrast, self-diffusion is controlled by the electron correlations, and therefore the coefficient of self-diffusion along the troughs $D_{\parallel}$ is strongly affected by the commensurate potential. Moreover, when the system freezes into a Wigner crystal, the self-diffusion along the troughs vanishes [21], whereas the mobility along the troughs does not.

In Fig. 7, we show the dependence of the mobility transverse to the troughs $\mu_{\perp}$ and the self-diffusion coefficients $D_{\parallel}$ and $D_{\perp}$ on temperature for two values of the potential amplitude A . In Fig. 8, we show the dependence of these parameters on A for a fixed temperature. Self-diffusion monotonically decreases with the decreasing temperature (increasing Γ) and with the increasing potential amplitude. Moreover, the self-diffusion coefficients $D_{\parallel}$ and $D_{\perp}$ are close to each other for the considered weak potential in Fig. 7, $A \ll E_C$ (they become very different for a stronger potential [21]).

In contrast, for a very weak potential, Fig. 7(a), the mobility depends on T nonmonotonically, first increasing with the decreasing T for higher temperature. This increase is related to the decrease of the rate of electron scattering by ripples, which is $\propto T$. However, for still smaller T , the mobility decreases with the decreasing T (increasing Γ).

Figure 8 shows that $\mu_{\perp}$ monotonically decreases with the increasing potential amplitude A . This decrease is close to exponential. Moreover, for the value of the plasma parameter




FIG. 8. (a) The mobility of the electron liquid $\mu_{\perp}$ transverse to the potential troughs (left y-axis) and the coefficients of self-diffusion along ($D_{\parallel}$) and transverse ($D_{\perp}$) to the troughs (right y-axis) as functions of the potential amplitude. The results refer to the maximally commensurate potential, $p_c = 1$, and to the plasma parameter $\Gamma = 90$. Note the difference in the scales compared to Fig. 7(b). (b) The logarithm of the transverse mobility for the same $\Gamma = 90$.

$\Gamma = 90$ shown in the figure, the mobility falls off with the increasing A sharper than the self-diffusion coefficients.

Both the transverse mobility and the self-diffusion should vanish where electrons crystallize in a commensurate potential. The mobility is advantageous for finding the Wigner crystallization temperature as it is much easier to access in the experiment than the self-diffusion. However, measuring a very small mobility is complicated not only in the experiment, but also in the simulations, as seen from the error bars in Fig. 8.

We note the difference in the scales in Figs. 7(b) and 8(a). The scale for $\mu_{\perp}$ is larger whereas the scale for $D_{\parallel}$ and $D_{\perp}$ is smaller in 8(a). On the scale of Fig. 7, the difference between the parameter values where the mobility and the self-diffusion become close to zero is within the error bars.

To gain a better feeling for the scaling factors in Figs. 7 and 8, it is instructive to look at the Einstein ratio $\mu_{\perp}k_B T/eD$, which in the single-electron approximation should be equal to one. For our scaling factors, we have

$$\mu_{\text{eff}}k_B T(eD_{\text{eff}})^{-1} = (2\sqrt{\pi}\Gamma)^{-1}, \quad (12)$$

i.e., for $\Gamma = 10^2$, we have $\mu_{\text{eff}}k_B T/eD_{\text{eff}} \approx 1/300$. Therefore, for $\mu_{\perp}$ and D measured in the units of μ_{eff} and D_{eff} , if the Einstein relation held, the Einstein ratio would be ~ 300 . Instead, as seen from Figs. 7 and 8, it is $\sim 10^4$. This shows that, even though the mobility is reduced by the periodic potential, it is still orders of magnitude higher than what follows from the Einstein relation. We emphasize again that $D_{\perp}$ and $D_{\parallel}$ are the coefficients of self-diffusion, not of the long-wavelength diffusion, therefore the Einstein relation should not hold; it is just that the difference with this relation is as large as two orders of magnitude.

B. Many-electron nature of overcoming the barriers

The transverse drift in the commensurate potential involves overcoming the periodically repeated potential barriers. One can picture the transport of the electron liquid in a small electric field along the x axis as motion in a slightly tilted




FIG. 9. The mobility $\mu_{\perp}$ of the strongly correlated electron system as a function of the potential amplitude A for three values of Γ . The dashed lines show the least squares fit.

washboard potential. If the electron system were incompressible and formed a Wigner crystal, it would not move in the commensurate potential, as such motion would involve overcoming a barrier by all electrons at a time, i.e., the barrier height would be proportional to the system size. The very occurrence of the mobility is thus due to the correlation length of the electron system being finite. Still one would expect that the mobility will exponentially depend on the barrier height. The corresponding dependence has been indeed seen in the simulations and is shown in Fig. 9 for several values of Γ .

To understand the result one can think of a cartoon of the electron liquid as made up of “clusters” with a typical size given by the correlation length ξ . The electrons are reasonably well ordered within clusters. Such clusters can be seen in Fig. 6, particularly distinctly for $\Gamma = 90$. The number of electrons in a cluster is $\sim \pi(\xi/a_s)^2$. The mobility results from the electron motion in which the clusters rearrange and move while the correlation length is preserved. For the considered case of maximal commensurability, the ultimate result is that the areas with a typical size given by the correlation length go over the potential barriers of height $2A$. Such transitions are thermally activated. Thus the mobility should contain the Arrhenius factor,

$$\mu_{\perp} \propto \exp(-\gamma A/E_C), \quad \gamma \sim 2\pi \Gamma \xi^2/a_s^2. \quad (13)$$

We note that $\Gamma A/E_C \equiv A/k_B T$, so that the exponent in $\mu_{\perp}$ is $\gamma A/E_C \sim (2\pi \xi^2/a_s^2)A/k_B T$. It is important that, although the mobility is activated, $\gamma \propto \Gamma \propto 1/T$, the overall dependence of γ on temperature is more complicated than just $1/T$, because the correlation length depends on temperature. This leads to a nonexponential dependence of $\mu_{\perp}$ on $1/T$. Such a dependence has been seen in our simulations and can be inferred from Fig. 7.

The picture of a locally nonuniform transport implies internal friction in the electron liquid, with the momentum of the system as a whole being ultimately transferred to the periodic potential. If this is indeed the case, the mobility $\mu_{\perp}$ in the regime where it is activated, Eq. (13), should become largely independent of the ripplon scattering, which controls




FIG. 10. A comparison of the mobility $\mu_{\perp}$ of the electron liquid as a function of the potential amplitude A with and without electron scattering by ripples. The results refer to $\Gamma = 90$. The inset shows the results in the region $A/E_C > 3.8 \times 10^{-4}$, where the mobility is described by the activation law (13). Notice the different scales of $\mu_{\perp}$ in the main figure and the inset.

the mobility in the absence of the potential. This is indeed seen from Fig. 10.

It is important that, for the potential amplitudes we are studying, the interaction energy is much larger than the amplitude of the potential. Therefore the correlation length should be approximately the same as in the absence of the potential, the picture corroborated by Fig. 5. We determined the correlation length approximately from the radial distribution function $g(r)$ of the electron liquid in the absence of the potential,

$$g(r) = (2\pi r n_s N)^{-1} \sum_{n,m} \delta(r - |\mathbf{r}_n - \mathbf{r}_m|). \quad (14)$$

This expression immediately follows from Eq. (10) if $\rho_s(\mathbf{r}) = n_s$ and the system is isotropic.

The radial distribution function for an isolated electron system was studied earlier, cf. [30,33]. For the large system studied here it is shown in Fig. 11(a). It has a familiar form for a liquid form of decaying oscillations, with $g(r)$ approaching 1 for large r/a_s . If the decay of $g(r) - 1$ is exponential, the correlation length can be associated with the decay length. However, for not large r/a_s , the decay is not exponential and not a modulated exponential. On the other hand, for very large r/a_s the complicated form and the finite resolution of the calculated $g(r)$ do not allow us to characterize the decay quantitatively.

The procedure we adopted for estimating the correlation length ξ was to fit the heights of the resolved maxima of oscillating $g(r)$ by exponentials, starting with the second maximum, as shown by the dashed lines in Fig. 11(a). We then associate ξ with the corresponding decay length. We calculated ξ for Γ varying from $\Gamma = 30$ to $\Gamma = 110$ with step $\Delta\Gamma = 10$. In this range the values of ξ are increasing with the increasing Γ from $\approx 1.0 a_s$ to $\approx 1.7 a_s$.

Even after the first maximum of $g(r)$ is skipped, the actual decay of the heights of the maxima of $g(r)$ is still not strictly exponential. This leads to an uncertainty in ξ . To estimate this uncertainty we compared the values of ξ obtained by omitting,




FIG. 11. (a) Solid lines: the radial distribution function of the electron liquid in the absence of a periodic potential, Eq. (14), for several values of Γ . Dashed lines: the exponential fit to the envelope of the oscillations of $g(r)$. (b) Data points: the parameter γ in the activation law (13). It is extracted for several values of Γ from the falloff of $\mu_{\perp}$ with the increasing potential amplitude using Eq. (16). This parameter is plotted against the parameter γ_c obtained from the decay length ξ of the radial distribution function seen in (a). The straight solid line that goes through the origin shows the least squares fit to the relation $\gamma = \gamma_c$. From this fit we estimate $\kappa = 0.92 \pm 0.04$ in Eq. (15). The error bars are explained in the text.

along with the first, the second or the last visible maximum of $g(r)$. The results are shown by the horizontal error bars in Fig. 11(b). The vertical error bars show the error of the exponential fit of the mobility as a function of the amplitude of the periodic potential, see Fig. 9.

The found values of ξ are close to the mean interelectron distance a_s . Meanwhile, in Eq. (13) the correlation length was assumed to be large compared to a_s . One can, empirically, extrapolate the estimate of γ in Eq. (13) to the range $\xi \sim a_s$ by replacing $\xi \rightarrow \xi + a_s$, i.e., by setting $\gamma = \gamma_c$, with

$$\gamma_c = 2\pi\kappa\Gamma(\xi + a_s)^2/a_s^2. \quad (15)$$

Here, κ is a numerical coefficient, $\kappa \sim 1$.

Figure 11(b) shows a comparison between the factor γ_c , which is obtained from the pair correlation function of a spatially uniform electron liquid, with the factor

$$\gamma = -\frac{d \ln \mu_{\perp}}{d(A/E_C)}, \quad (16)$$

extracted from the entirely different simulations of the transverse mobility $\mu_{\perp}$ in a periodic potential. It is seen that, in a broad parameter range, γ as determined from the mobility is proportional to γ_c determined from the decay of correlations.

The proportionality of γ and γ_c provides a reasonably strong argument in favor of the proposed qualitative picture of the thermally activated mobility in a commensurate periodic potential. Moreover, and arguably more importantly, the results suggest that the correlation length in the electron liquid can be determined by measuring the transverse mobility in a weak maximally commensurate periodic potential.

VI. TRANSVERSE MOBILITY FOR DIFFERENT VALUES OF THE COMMENSURABILITY PARAMETER

The previous section described the mobility $\mu_{\perp}$ where the electron system is a liquid, but the electron crystal with the same density is maximally commensurate with the external




FIG. 12. The mobility $\mu_{\perp}$ of the strongly correlated electron system as a function of the commensurability ratio p_c for $\Gamma = 50$, 90, and 130. The scaled potential amplitude $u_s A/E_C = 0.00157$ for all three cases.

potential, the commensurability parameter (7) was $p_c = 1$. The electron density in this case was strongly modulated by even a weak potential that we used. One can expect that the electron density will be also modulated for $p_c > 1$, with the modulation becoming the strongest where the potential is commensurate, including $p_c = \sqrt{3}$ and $p_c = 2$, see Appendixes B and C. This should reduce the mobility $\mu_{\perp}$.

For electrons on helium, of interest are the values of p_c , which are not large. Indeed, the potential created by the electrodes in Fig. 1, exponentially falls off with the decreasing period $2\pi/Q$, for a given height of the helium layer. Therefore the results below are limited to $p_c \lesssim 2$.

In Fig. 12, we show how $\mu_{\perp}$ varies with the varying p_c . In the simulations, p_c is changed by changing the period of the potential, while keeping the mean interelectron distance a_s fixed. Our simulation cell is a rectangle that contains 1600 electrons and has the dimensions $L_y = 40a_s$ and $L_x = 40(\sqrt{3}/2)a_s$. To satisfy the periodic boundary conditions, the period of the potential should be a simple fraction of L_x , namely L_x/n with an integer n . Keeping in mind that, if the electrons crystallized, there would be 40 electron rows in the x direction, we chose the commensurability parameter to take on the values $n/40$. We used an integer n from 1 to 80.

It is seen from Fig. 12 that a small periodic potential weakly affects the mobility for $p_c \ll 1$. This is corroborated by the dependence of $\mu_{\perp}$ on the potential amplitude A shown in Fig. 4 for $p_c = 3/10$. It is also seen in Fig. 12 that for all $p_c \lesssim 0.5$ the mobility is close to that in the absence of a periodic potential in the studied range of A/E_C . The difference between the values of $\mu_{\perp}$ for small p_c and different Γ comes from the increase of the rate of scattering by ripplons with the increasing temperature, i.e., with the decreasing Γ for the given electron density.

As p_c approaches 1 from below, the mobility sharply drops. The decrease of $\mu_{\perp}$ becomes steeper with the increasing Γ . For p_c close to one, the strong electron correlations do not “compete” with the periodic potential, electrons can reason-

ably well fit into the repeated potential minima (perfectly fit, for $p_c = 1$). Essentially, the potential competes only with the disorder in the electron liquid. This is why it makes a strong effect even for $A/E_c \sim 10^{-2}$, provided $\Gamma = E_c/k_B T \gg 1$.

For $\Gamma = 130$ and $p_c = 1$, the electrons form a Wigner crystal and the mobility becomes zero. For this Γ , the electrons crystallize also for other commensurate values $p_c = \sqrt{3}$ and $p_c = 2$.

In the range $1 \lesssim p_c \lesssim 2$, the mobility in the liquid phase is smaller than for $p_c \ll 1$ even for an incommensurate potential. As seen from Fig. 12, the dependence on Γ is inverted compared to the case of small p_c : the larger Γ , the smaller the mobility. Such behavior suggests that the mobility is limited by the momentum transfer to the potential rather than to the ripples.

As electron clusters move and rearrange in the incommensurate potential, some electrons go “uphill” whereas the others go “downhill.” The energy gains and losses are not exactly locally balanced, which should lead to an exponential dependence of $\mu_\perp$ on A . Such a dependence has been indeed seen in the simulations, and for $p_c = 1.4$ it is presented in Appendix C. The rate of the decrease of $\log \mu_\perp$ with the increasing A is reduced for an incommensurate potential compared to the case $p_c = 1$. Also, as Fig. 12 shows, we did not find a pronounced dip in the mobility near $p_c = \sqrt{3}$ for $\Gamma \lesssim 90$.

It should be noted that the electron vibrations about their quasiequilibrium positions in the electron liquid do not average out the potential. Indeed, it follows from the estimate in Sec. V that the frequency of electron vibrations is $\sim \omega_p$ for $A/E_c \ll 1$ and $p_c \sim 1$. Then the mean-square displacement of an electron about its quasiequilibrium position $(k_B T / m_e \omega_p^2) \sim a_s^2 / \Gamma$ is small compared to the squared period of the potential for $p_c \sim 1$ and $\Gamma \gg 1$, cf. Appendix B.

VII. CONCLUSIONS

The presented numerical simulations of the electron liquid on helium have shown several effects of the strong electron correlations on the electron mobility. The results suggest a way to directly characterize these correlations in an experiment.

For a spatially uniform electron system, the simulations have confirmed the mechanism of transport inferred in the earlier work [9,10,16]. The underlying picture in that work is that the electron scattering by the helium excitations is weak, whereas the electron-electron interaction is strong. Yet in the classical regime $k_B T \gg \hbar \omega_p$ and in the absence of a magnetic field, the major effect of the electron-electron interaction on the mobility is the interelectron momentum exchange, which is faster than the momentum exchange with the helium excitations. This picture allowed calculating the mobility explicitly. The excellent agreement of our simulations with the analytical results provides a quantitative basis for the above picture. Moreover, the simulations have explicitly shown that, when the electron system is slightly heated by the external field, the distribution of the electron kinetic energy is of the Maxwell-Boltzmann form in the co-moving frame, with a temperature higher than the temperature of the helium.

The central results of the paper refer to the mobility of the electron liquid placed into a sinusoidal one-dimensional potential. We studied the parameter range where the potential amplitude A was three orders of magnitude smaller than the electron interaction energy E_c . Yet the effect of the potential can be strong. It depends on the interrelation between the mean interelectron spacing a_s and the period of the potential $2\pi/Q$. More specifically, it depends on whether the potential is commensurate with the electron crystal with the same spacing, even though the results refer to the region where the electrons form a liquid, not a crystal.

We found that, for $a_s Q / 2\pi \ll 1$, the effect of an incommensurate potential on the mobility is effectively washed out by strong electron correlations. Even where $A/k_B T \gtrsim 1$ and the single-electron mobility is strongly reduced, the many-electron mobility shows a very small change. This can be understood by noticing that, for $E_c \gg A$, the electron system averages out the potential. At the same time, the potential is smooth on the electron thermal wavelength, and therefore the potential does not add to the electron scattering.

A qualitatively different behavior is displayed in a commensurate potential. Even where the two-particle correlation function is weakly modified compared to the case of no potential, the electron density becomes periodically modulated with an amplitude significantly larger than in the single-electron picture. The mobility transverse to the potential troughs $\mu_\perp$ displays an exponential dependence on the potential amplitude A . We associate the very mechanism of the mobility with the absence of long-range order. In the picture suggested by the results, the mobility results from the correlated many-electron activated transitions within areas with the typical size given by the correlation length in the liquid.

We have found a simple relation between the correlation length in the electron system in the absence of the periodic potential and the activated fall-off of the mobility with the increasing A for $A \ll E_c$. This relation provides a means for measuring the electron correlation length in the experiment. We are not aware of other means to measure the correlation length of the nondegenerate electron liquid on the surface of helium.

The parameters used in the simulations are within the typical range of the parameters used in the experiments on electrons on helium, and the proposed weak one-dimensional potential with a period $\sim 1 \mu\text{m}$ can be implemented with conventional technology. The system is advantageous for studying the effect of commensurability given that the electron density can be easily varied. Our results show that these effects are strong even for a classical electron liquid. As a future direction, it would be interesting to study these effects where the electron dynamics becomes quantum, even though the system remains nondegenerate and therefore there is no need to have high electron densities, which significantly simplifies an experimental implementation.

ACKNOWLEDGMENTS

The research of K.M. was supported in part by the National Science Foundation, Grant DMR 1708331. M.I.D. was supported in part by the Grant DE-SC0020136 funded by the US Department of Energy, Office of Science.

APPENDIX A: OUTLINE OF THE SIMULATIONS

1. Matrix elements of coupling to the helium excitations

A key distinctive part of the simulations is the incorporation of the actual elastic and inelastic scattering into the electron equations of motion. The scattering is due to the coupling (4) to ripples and phonons in liquid helium. The coupling parameters $V_{q\alpha}$ are well-known [1,21,44,51]. We give them here for completeness.

We are interested in the case of a weak field that presses electrons to the surface. Then the coupling to ripples comes primarily from the ripplon-induced change of the image potential that attracts the electrons to the helium surface. The image potential is also changed by phonons, as they modulate the helium density and thus the dielectric constant. The corresponding changes of the potential energy of an electron at a distance z from the surface are, respectively,

$$\begin{aligned} V_q^{(\text{rp})}(z) &= \Lambda \frac{(\hbar q)^{1/2}}{(2\rho\omega_q S)^{1/2}} z^{-2} [1 - qzK_1(qz)], \\ V_{q,q_z}^{(\text{ph})}(z) &= -i\Lambda q(\hbar\omega_{q,q_z}/V v_{\text{He}}^2 \rho_{\text{He}})^{1/2} \\ &\quad \times \int_{-\infty}^0 dz' \frac{\sin(q_z z')}{z - z'} K_1(q(z - z')). \end{aligned} \quad (\text{A1})$$

Here, S is the area of the system, $\Lambda = e^2(\epsilon - 1)/4(\epsilon + 1)$ (ϵ is the dielectric constant of helium, $\epsilon \approx 1.057$), v_{He} is the speed of sound in helium, ρ_{He} is the helium density, and $K_1(x)$ is the Bessel function. The parameters $V_{q\alpha}$ of the coupling Hamiltonian (4) are obtained by calculating the diagonal matrix elements of the functions (A1) on the wave function $\psi_0(z)$ of the ground state of electron motion normal to the helium surface.

2. The calculation

The simulation procedure used in this work is described in detail in [21]. Here we briefly summarize the most important aspects. The simulations were conducted using HOOMD-BLUE [52,53] with a custom interaction potential, integrator, and external forces.

As indicated in the main text, we consider $N = 1600$ electrons placed into a rectangular cell with the aspect ratio of $L_x/L_y = \sqrt{3}/2$ and periodic boundary conditions. If an electron crosses the cell boundary, it is introduced back into the cell from the opposite side. Electrons interact via a long-range Coulomb force which is handled through the Ewald summation [30]. We consider an external potential (2) imposed on electrons, which is periodic along the x axis. The period is chosen so that the wave vector Q satisfies the condition $QL_x = 2\pi n$ with a positive integer n . The time step is chosen to be $\Delta t \approx (2\pi/\omega_p)/50$. We have checked that the results do not change upon reducing the step further.

We integrate equations of motion with the velocity Verlet algorithm modified for direct simulation of scattering events by excitations in helium. The inelastic scattering provides a thermal bath that maintains a set temperature in the electron system. However, since the inelastic scattering rate is low, the effective temperature of the system may deviate slightly from the bath temperature. We define the effective temperature in a




FIG. 13. Schematic of a triangular lattice placed in a 1D periodic potential $U = -A \cos Qx$. The dashed lines show the maxima of the potential. The left panel refers to what we call the maximally commensurate potential, $p_c = 1$, whereas the right panel refers to $p_c = \sqrt{3}$. This sketch illustrates the definition (7) of the commensurability parameter p_c . In simulations, we study the liquid phase with no crystalline order.

nonequilibrium state as follows:

$$T_{\text{eff}}(t) \equiv (m_e/2Nk_B) \sum_n (\mathbf{v}_n(t) - \mathbf{v}_d)^2, \quad (\text{A2})$$

where

$$\mathbf{v}_d \equiv \frac{1}{NN_t} \sum_{n=1}^N \sum_{s=0}^{N_t} \mathbf{v}_n(s\Delta t) \quad (\text{A3})$$

is the drift velocity of the system averaged over time. N_t is the number of time steps and s is the time step index.

In a typical simulation, we prepare the system in a triangular lattice with one side of the triangle oriented along the y axis. Initial velocities are randomly oriented with the absolute velocity values distributed according to the Boltzmann distribution for a set temperature. External drive electric field E_d is applied along the x axis. We allow the driven system to stabilize in a stationary state for 10^6 time steps before collecting data. This stabilization time is sufficient for the relatively high temperatures considered here.

We found that the results did not change when the system size changed from 1024 to 1600 electrons. Some results were also checked for different aspect ratios of the simulation box. The absence of changes indicates that the system size effects are within the simulation uncertainty.

The natural parameter of the state of the electron system in the classical regime is $\Gamma = e^2(\pi n_s)^{1/2}/k_B T$. However, the electron mobility depends on T separately. In our finite system, we cannot vary the density n_s continuously while maintaining a given ratio of the interelectron distance and the periodic potential. Therefore our results for different Γ are obtained by varying the temperature.

APPENDIX B: ELECTRON DENSITY IN THE MAXIMALLY COMMENSURATE POTENTIAL

Two simple types of the commensurate potentials are shown in Fig. 13. For the potential shown in the left panel, the distance between the electrons in the same potential trough is equal to the mean interelectron distance a_s and $p_c = 1$. For the potential shown in the right panel, this distance is $a_s\sqrt{3}$ and $p_c = \sqrt{3}$. Respectively, the distances between the




FIG. 14. Plots (a) and (b) show the pair correlation function of the electron liquid $\tilde{g}^{(2)}(\mathbf{r})$, Eq. (10), for $p_c = 1.4$ and $p_c = 1.725 \approx \sqrt{3}$, respectively. Plots (c) and (d) show the averaged over time electron density along the x direction for the same parameters as in (a) and (b), respectively. The scaled amplitude of the potential is $A/E_C = 0.00157$, and the plasma parameter is $\Gamma = 90$ in all panels.

troughs differ by the factor $\sqrt{3}$. Each of these potentials leads to a series of potentials with $p_c = n$ or $p_c = n\sqrt{3}$, into which the electron crystal can fit with no distortion. As mentioned in the text, we call the potential with $p_c = 1$ the maximally commensurate potential.

If the electrons are placed into a sufficiently strong maximally commensurate periodic potential ($p_c = 1$) but do not crystallize, one can approximate the density as uniform along the potential troughs and a sum of Gaussian peaks in the transverse direction,

$$\rho_s(\mathbf{r}) \approx \frac{n_s}{\varkappa Q} \sqrt{2\pi} \sum_n \exp[-(x - 2\pi n/Q)^2 / 2\varkappa^2]. \quad (\text{B1})$$

The distribution $\rho(\mathbf{r})$ corresponds to a set of “single-electron” wires: one wire per trough with the mean interelectron distance along the trough equal to the mean interelectron spacing a_s . The width of the peaks $\varkappa$ can be roughly estimated by assuming Boltzmann distribution about the minima of $U(\mathbf{r})$ and further assuming that the potential near a given minimum, in addition to $U(\mathbf{r})$, has a contribution from the electron “wires” localized in other minima, i.e., disregarding the short-range ordering along the wire. This gives $\varkappa = [k_B T / (A Q^2 + \frac{1}{3} \pi e^2 n_s Q)]^{1/2}$. For the considered case of maximum commensurability this can be also written as $\varkappa = (k_B T / Q^2)^{1/2} [A + E_C (\sqrt{3}/72\pi)^{1/2}]^{-1/2}$. According to this estimate, the major factor in the width of the peaks is the electron-electron interaction, for the considered range of the potential strength $A \ll E_C$.

From Eq. (B1), for a strongly confined liquid, the Fourier components α_m of the electron density in Eq. (9) are

$$\alpha_m = 2 \exp(-m^2 \varkappa^2 Q^2 / 2). \quad (\text{B2})$$




FIG. 15. The mobility transverse to the potential troughs as a function of the potential amplitude A for $p_c = 1.4$ on the linear (a) and logarithmic (b) scales. The plasma parameter $\Gamma = 90$.

They fall off exponentially fast with the increasing m for $m \gg 1/\varkappa Q$.

APPENDIX C: ELECTRON CORRELATIONS AND THE MOBILITY FOR $p_c > 1$

We defined p_c using the Wigner crystal with reciprocal lattice vectors $\mathbf{G}_1 = 2\pi a_s^{-1}(-1/\sqrt{3}, 1)$ and $\mathbf{G}_2 = 2\pi a_s^{-1}(2/\sqrt{3}, 0)$, so that the maximal commensurability $p_c = 1$ corresponded to the reciprocal lattice vector of the one-dimensional periodic potential $Q = G_{2x}$. Other commensurate values of Q for a one-dimensional potential are $\mathbf{Q} = n_1(2\mathbf{G}_1 + \mathbf{G}_2) + n_2\mathbf{G}_2$ with integer n_1, n_2 . Respectively, the smallest values of p_c that correspond to a commensurate lattice are $p_c = 1, \sqrt{3}, 2, \dots$

As shown in the main text, in the range $1 \lesssim p_c \lesssim 2$ the transverse mobility $\mu_{\perp}$ is significantly smaller than in the absence of the periodic potential. This is related to the partial adjustment of the electron liquid to the potential. To demonstrate this adjustment, in Fig. 14, we show the density modulation and the pair correlation function for $p_c = 1.4$, which lies between the two smallest values corresponding to the commensurability, and for $p_c = 1.725$, which is close to the commensurate case $p_c = \sqrt{3}$.

It is seen from Figs. 5 and 14 that the pair correlation function weakly depends on p_c for the considered weak periodic potential. In contrast, the density modulation is significantly stronger in the commensurate case than away from commensurability. Moreover, it is significantly stronger for $p_c = 1$ than for $p_c = \sqrt{3}$, i.e., for the electrons being spaced more closely within the potential trough and, respectively, for a larger distance between the troughs, cf. Fig. 13.

Figure 15 shows that the electron mobility displays an exponential dependence on the potential amplitude even away from the commensurability, where $p_c = 1.4$. However, the values of the mobility are much larger than in the maximally commensurate case $p_c = 1$ shown in Fig. 10. This is a consequence of the partial averaging of the incommensurate potential in the electron liquid.

- [1] E. Andrei, ed., *Two-Dimensional Electron Systems on Helium and Other Cryogenic Surfaces* (Kluwer Academic, Dordrecht, 1997).
- [2] Y. Monarkha and K. Kono, *Two-Dimensional Coulomb Liquids and Solids* (Springer, Berlin, 2004).
- [3] K. Shirahama, S. Ito, H. Suto, and K. Kono, Surface study of liquid ^3He using surface-state electrons, *J. Low Temp. Phys.* **101**, 439 (1995).
- [4] E. Collin, W. Bailey, P. Fozooni, P. G. Frayne, P. Glasson, K. Harrabi, M. J. Lea, and G. Papageorgiou, Microwave Saturation of the Rydberg States of Electrons on Helium, *Phys. Rev. Lett.* **89**, 245301 (2002).
- [5] C. C. Grimes and G. Adams, Evidence For A Liquid-to-Crystal Phase-Transition In A Classical, 2-Dimensional Sheet of Electrons, *Phys. Rev. Lett.* **42**, 795 (1979).
- [6] D. S. Fisher, B. I. Halperin, and P. M. Platzman, Phonon-Ripplon Coupling and the 2-Dimensional Electron Solid On A Liquid-Helium Surface, *Phys. Rev. Lett.* **42**, 798 (1979).
- [7] D. G. Rees, N. R. Beysengulov, J.-J. Lin, and K. Kono, Stick-Slip Motion of the Wigner Solid on Liquid Helium, *Phys. Rev. Lett.* **116**, 206801 (2016).
- [8] D. G. Rees, Sh.-Sh. Yeh, B.-C. Lee, K. Kono, and J.-J. Lin, Bistable transport properties of a quasi-one-dimensional wigner solid on liquid helium under continuous driving, *Phys. Rev. B* **96**, 205438 (2017).
- [9] M. I. Dykman and L. S. Khazan, Effect of the interaction between nondegenerate electrons localized in a thin surface layer on the cyclotron resonance and on the magnetoconductance, *JETP* **50**, 747 (1979) [*Zh. Eksp. Teor. Fiz.* **77**, 1488 (1979)].
- [10] M. I. Dykman, M. J. Lea, P. Fozooni, and J. Frost, Magnetoresistance In 2D Electrons On Liquid-Helium - Many-Electron Versus Single-Electron Kinetics, *Phys. Rev. Lett.* **70**, 3975 (1993).
- [11] M. J. Lea and M. I. Dykman, Magnetotransport in the two-dimensional electron fluid and solid on liquid helium, *Phys. B* **249**, 628 (1998).
- [12] D. Konstantinov, M. I. Dykman, M. J. Lea, Y. Monarkha, and K. Kono, Resonant Correlation-Induced Optical Bistability in an Electron System on Liquid Helium, *Phys. Rev. Lett.* **103**, 096801 (2009).
- [13] D. Konstantinov, Y. Monarkha, and K. Kono, Effect of Coulomb Interaction on Microwave-Induced Magnetoconductivity Oscillations of Surface Electrons on Liquid Helium, *Phys. Rev. Lett.* **111**, 266802 (2013).
- [14] A. D. Chepelianskii, M. Watanabe, K. Nasyedkin, K. Kono, and D. Konstantinov, An incompressible state of a photo-excited electron gas, *Nat. Commun.* **6**, 7210 (2015).
- [15] W. Kohn, Cyclotron resonance and de Haas-van Alphen oscillations of an interacting electron gas, *Phys. Rev.* **123**, 1242 (1961).
- [16] V. A. Buntar', Yu Z. Kovdrya, V. N. Grigor'ev, Yu P. Monarkha, and S. S. Sokolov, Studies of momentum relaxation rate and observation of complete control condition in two-dimensional electron system helium surface, *Fiz. Nizk. Temp.* **13**, 789 (1987) [*Sov. J. Low Temp. Phys.* **13**, 451 (1987)].
- [17] L. Menna, S. Yucel, and E. Y. Andrei, Giant Magnetic Suppression Of Tunneling Out Of A 2d Electron-System, *Phys. Rev. Lett.* **70**, 2154 (1993) T. Barabash, M. I. Dykman, and P. M. Platzman, Tunneling Transverse to a Magnetic Field and Its Occurrence in Correlated 2D Electron Systems, *ibid.* **84**, 2227 (2000).
- [18] Q.-H. Wei, C. Bechinger, D. Rudhardt, and P. Leiderer, Experimental Study of Laser-Induced Melting in Two-Dimensional Colloids, *Phys. Rev. Lett.* **81**, 2606 (1998).
- [19] L. Radzihovsky, E. Frey, and D. R. Nelson, Novel phases and reentrant melting of two-dimensional colloidal crystals, *Phys. Rev. E* **63**, 031503 (2001).
- [20] R. K. Kalia, P. Vashishta, S. D. Mahanti, and J. J. Quinn, Electrons on corrugated surfaces, *J. Phys. C: Solid State Phys.* **16**, L491 (1983).
- [21] K. Moskovtsev and M. I. Dykman, Self-diffusion in a spatially modulated system of electrons on helium, *J. Low Temp. Phys.* **195**, 266 (2019).
- [22] A. Bylinskii, D. Gangloff, I. Counts, and V. Vuletić, Observation of Aubry-type transition in finite atom chains via friction, *Nat. Mater.* **15**, 717 (2016).
- [23] T. Brazda, A. Silva, N. Manini, A. Vanossi, R. Guerra, E. Tosatti, and C. Bechinger, Experimental Observation of the Aubry Transition in Two-Dimensional Colloidal Monolayers, *Phys. Rev. X* **8**, 011050 (2018).
- [24] M. Y. Zakharov, D. Demidov, and D. L. Shepelyansky, Transport properties of a Wigner crystal in one- and two-dimensional asymmetric periodic potentials: Wigner crystal diode, *Phys. Rev. B* **99**, 155416 (2019).
- [25] M. I. Dykman and Y. G. Rubo, Bragg-Cherenkov Scattering and Nonlinear Conductivity of a Two-Dimensional Wigner Crystal, *Phys. Rev. Lett.* **78**, 4813 (1997).
- [26] W. F. Vinen, Non-linear electrical conductivity and sliding in a two-dimensional electron crystal on liquid helium, *J. Phys. Condens. Matter* **11**, 9709 (1999).
- [27] V. L. Pokrovsky and A. L. Talapov, The theory of two-dimensional incommensurate crystals, *Zh. Eksp. Teor. Fiz.* **78**, 269 (1980) [*JETP* **51**, 134 (1980)].
- [28] P. Bak, Commensurate phases, incommensurate phases and the Devil's staircase, *Rep. Prog. Phys.* **45**, 587 (1982).
- [29] S. Aubry, The twist map, the extended Frenkel-Kontorova model and the Devil's staircase, *Phys. Nonlinear Phenom.* **7**, 240 (1983).
- [30] R. C. Gann, S. Chakravarty, and G. V. Chester, Monte Carlo simulation of the classical two-dimensional one-component plasma, *Phys. Rev. B* **20**, 326 (1979).
- [31] J. P. Hansen, D. Levesque, and J. J. Weis, Self-Diffusion in the Two-Dimensional, Classical Electron Gas, *Phys. Rev. Lett.* **43**, 979 (1979).
- [32] R. H. Morf, Temperature Dependence of the Shear Modulus and Melting of the Two-Dimensional Electron Solid, *Phys. Rev. Lett.* **43**, 931 (1979).
- [33] R. K. Kalia, P. Vashishta, and S. W. de Leeuw, Melting of a two-dimensional electron lattice, *Phys. Rev. B* **23**, 4794 (1981).
- [34] K. J. Strandburg, Two-dimensional melting, *Rev. Mod. Phys.* **60**, 161 (1988).
- [35] C. Fang-Yen, M. I. Dykman, and M. J. Lea, Internal Forces in Nondegenerate Two-Dimensional Electron Systems, *Phys. Rev. B* **55**, 16272 (1997).
- [36] S. Muto and H. Aoki, Crystallization of a classical two-dimensional electron system: Positional and orientational orders, *Phys. Rev. B* **59**, 14911 (1999).

- [37] G. Piacente and F. M. Peeters, Pinning and depinning of a classic quasi-one-dimensional Wigner crystal in the presence of a constriction, *Phys. Rev. B* **72**, 205208 (2005).
- [38] B. K. Clark, M. Casula, and D. M. Ceperley, Hexatic and Mesoscopic Phases in a 2D Quantum Coulomb System, *Phys. Rev. Lett.* **103**, 055701 (2009).
- [39] P. F. Damasceno, C. J. DaSilva, J. P. Rino, and L. Cândido, Temperature and pinning effects on driving a 2D electron system on a helium film: A numerical study, *J. Low Temp. Phys.* **160**, 58 (2010).
- [40] D. G. Rees, I. Kuroda, C. A. Marrache-Kikuchi, M. Hofer, P. Leiderer, and K. Kono, Transport measurements of strongly correlated electrons on helium in a classical point-contact device, *J. Low Temp. Phys.* **166**, 107 (2012).
- [41] M. Mazars, The melting of the classical two-dimensional Wigner crystal, *EPL* **110**, 26003 (2015).
- [42] V. B. Shikin and Yu. P. Monarkha, On the interaction of surface electrons in liquid helium with oscillations of the vapor-liquid interface, *J. Low Temp. Phys.* **16**, 193 (1974).
- [43] M. W. Cole, Electronic surface states of liquid-helium, *Rev. Mod. Phys.* **46**, 451 (1974).
- [44] M. I. Dykman, P. M. Platzman, and P. Seddighrad, Qubits with electrons on liquid helium, *Phys. Rev. B* **67**, 155402 (2003).
- [45] M. I. Dykman, C. FangYen, and M. J. Lea, Many-electron transport in strongly correlated nondegenerate two-dimensional electron systems, *Phys. Rev. B* **55**, 16249 (1997).
- [46] The Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevB.101.245435> describes a theory of the transport of a strongly correlated nondegenerate two-dimensional electron liquid in the case where electrons experience elastic and inelastic scattering by vibrational modes of the medium. It also includes additional references [47–49].
- [47] M. I. Dykman, Cyclotron resonance of a two-dimensional Wigner crystal, *J. Phys. C: Solid State Phys.* **15**, 7397 (1982).
- [48] L.D. Landau and E. M. Lifshitz, *Statistical Physics. Part 1*, 3rd ed. (Pergamon Press, New York, 1980).
- [49] Yu. M. Vilk and Y. Monarkha, Correlation effects on relaxation processes in two-dimensional electron system at helium interface, *Fiz Nizk Temp.* **15**, 235 (1989) [*Sov. J. Low Temp. Phys.* **15**, 131 (1989)].
- [50] P. G. De Gennes, Liquid dynamics and inelastic scattering of neutrons, *Physica* **25**, 825 (1959).
- [51] D. I. Schuster, A. Fragner, M. I. Dykman, S. A. Lyon, and R. J. Schoelkopf, Proposal for Manipulating and Detecting Spin and Orbital States of Trapped Electrons on Helium Using Cavity Quantum Electrodynamics, *Phys. Rev. Lett.* **105**, 040503 (2010).
- [52] J. A. Anderson, C. D. Lorenz, and A. Travesset, General purpose molecular dynamics simulations fully implemented on graphics processing units, *J. Comput. Phys.* **227**, 5342 (2008).
- [53] J. Glaser, T. D. Nguyen, J. A. Anderson, P. Lui, F. Spiga, J. A. Millan, D. C. Morse, and S. C. Glotzer, Strong scaling of general-purpose molecular dynamics simulations on GPUs, *Comput. Phys. Commun.* **192**, 97 (2015).