

The Biology of Thermoacidophilic Archaea from the Order *Sulfolobales*

April M. Lewis¹, Alejandra Recalde², Christopher Bräsen³, James A. Counts¹, Phillip Nussbaum², Jan Bost², Larissa Schocke³, Lu Shen³, Daniel J. Willard¹, Tessa E. F. Quax⁴, Eveline Peeters⁵, Bettina Siebers³, Sonja-Verena Albers², and Robert M. Kelly^{1*}

¹*Department of Chemical and Biomolecular Engineering,
North Carolina State University, Raleigh, NC USA*

²*Institute for Biology,
Molecular Biology of Archaea,
University of Freiburg, Freiburg, Germany*

³*Department of Molecular Enzyme Technology and Biochemistry, Environmental Microbiology
and Biotechnology, and Centre for Water and Environmental Research,
University of Duisburg-Essen, Essen, Germany*

⁴*Archaeal Virus-Host Interactions, Faculty of Biology,
University of Freiburg, Freiburg, Germany*

⁵*Research Group of Microbiology,
Department of Bioengineering,
Vrije Universiteit, Brussels, Belgium*

Submitted to: *FEMS Microbiological Reviews* (July, 2020)

Revision to FEMSRE-20-07-0050 Submitted (November 2020)

Running Title: *Sulfolobales*

***Address Correspondence to:** Robert M. Kelly
Department of Chemical and Biomolecular Engineering
North Carolina State University
Raleigh, NC 27695
Phone: 919-515-6396
Email: rmkelly@ncsu

Abstract: Thermoacidophilic archaea belonging to the order *Sulfolobales* thrive in extreme biotopes, such as sulfuric hot springs and ore deposits. These microorganisms have been model systems for understanding life in extreme environments, as well as for probing the evolution of both molecular genetic processes and central metabolic pathways. Thermoacidophiles, such as the *Sulfolobales*, use typical microbial responses to persist in hot acid (e.g., motility, stress response, biofilm formation), albeit with some unusual twists. They also exhibit unique physiological features, including iron and sulfur chemolithoautotrophy, that differentiate them from much of the microbial world. Although first discovered more than 50 years ago, it was not until recently that genome sequence data and facile genetic tools have been developed for species in the *Sulfolobales*. These advances have not only opened up ways to further probe novel features of these microbes, but have also paved the way for potential biotechnological applications. Discussed here are the nuances of the thermoacidophilic lifestyle of the *Sulfolobales*, including their evolutionary placement, cell biology, survival strategies, genetic tools, metabolic processes, and physiological attributes together with how these characteristics make thermoacidophiles ideal platforms for specialized industrial processes.

1. Introduction

Thermoacidophiles are microorganisms that have developed mechanisms to successfully persist in unusually hot, acidic environments, with optimal conditions of $\text{pH} \leq 4$ and temperature $\geq 55^\circ\text{C}$. In fact, thermoacidophiles have been isolated from some of the most inhospitable environments on earth, such as acidic hot springs and volcanic solfataras. In 1972, Thomas Brock isolated the thermoacidophile *Sulfolobus acidocaldarius* from a sulfur hot spring in Yellowstone National Park (**Left image Figure 1**) and designated the genus *Sulfolobus* (Brock *et al.*, 1972). The natural habitat of this microbe, a member of the *Crenarchaeota*, was Locomotive Spring, an extremely hot acidic environment with a pH of 2.4 and temperature of 83°C . Likewise, in 1980 Wolfram Zillig described *Sulfolobus solfataricus* (renamed *Saccharolobus solfataricus*) isolated from a volcanic hot spring in Italy (Zillig *et al.*, 1980) and *Desulfurolobus ambivalens* (renamed *Acidianus ambivalens*) from a solfatara in Iceland in 1986 (Zillig *et al.*, 1986). Zillig also discovered the first Japanese isolate belonging to this group in 1990 - *Sulfolobus shibatae* (renamed *Saccharolobus shibatae*) (Grogan *et al.*, 1990). Beyond these discoveries, Zillig also isolated the first thermoacidophile virus (Martin *et al.*, 1984) (**Section 3**) and was the first to describe the eukaryotic-like archaeal RNA polymerase from *S. acidocaldarius* (Zillig *et al.*, 1979) (**Section 4**). In 1986, Karl Stetter established the genus *Acidianus* with the isolation of *Acidianus infernus* from a solfatara crater in Italy, which consequently led to the re-naming of *Sulfolobus brierleyi* as *Acidianus brierleyi* (Segerer *et al.*, 1986). Stetter also established the genus *Metallosphaera* with the isolation of *Metallosphaera sedula* in 1989 from a solfataric field in Italy (Huber *et al.*, 1989).

Thermoacidophiles not only thrive in thermal acidic biotopes but also encounter other biologically deleterious conditions, such as oxidative stress caused by high levels of metals in mining environments. For instance, *Metallosphaera prunae* was isolated from an uranium mine in Germany (Fuchs *et al.*, 1995) and uses an interesting stress response mechanism to withstand high levels of soluble uranium (**Section 7**). **Figure 1 (right)** shows the features of the isolation site of *M. prunae*. In addition to *Sa. shibatae*, several other thermoacidophiles have been isolated from hot springs in Japan, such as *Sulfurisphaera ohwakuensis* in 1988 (Kurosawa *et al.*, 1998), *Sulfolobus hakonensis* (renamed *Metallosphaera hakonensis*) in 1996 (Takayanagi *et al.*, 1996), and *Sulfolobus tokodaii* (renamed *Sulfurisphaera tokodaii*) in 2002 (Suzuki *et al.*, 2002), to name a few. It has become clear that thermoacidophiles are globally distributed in hot, acidic features; for example, recent isolates have come from the Copahue volcanic region in Argentina - *Acidianus copahuensis* in 2014 (Urbieta *et al.*, 2014), and Indonesian hot springs - *Sulfurisphaera javensis* in 2018 (Tsuboi *et al.*, 2018). Recently, *Saccharolobus caldissimus* was isolated from an acidic Japanese hot spring, establishing the *Saccharolobus* genus which, as mentioned above, led to the re-naming of both *Sulfolobus solfataricus* and *Sulfolobus shibatae* to *Saccharolobus solfataricus* and *Saccharolobus shibatae*, respectively (Sakai & Kurosawa, 2017). **Figure 1** illustrates the timeline of these thermoacidophile isolations. Many thermoacidophiles have leveraged the chemistry of metal and sulfur deposits for bioenergetic benefit through chemolithotrophy (**Section 8**). As such, chemolithotrophic metabolism in hot acid can be exploited for biomining applications (**Section 9**).

The study of thermoacidophiles was originally restricted to observational microbiology focused on phenotypic characteristics, such as cell morphology and growth physiology.

However, following the sequencing of the *Sa. solfataricus* genome in 2001 (She *et al.*, 2001), several other *Sulfolobales* genomes were reported, including *S. acidocaldarius* in 2005 (Chen *et al.*, 2005) (**Figure 1**). Genome sequences opened up prospects for transcriptomics (Auernik *et al.*, 2008, Ortmann *et al.*, 2008, Koerdt *et al.*, 2011, Kozubal *et al.*, 2011, Maezato *et al.*, 2012, Ulas *et al.*, 2012, Kouril *et al.*, 2013, Wolf *et al.*, 2016), proteomics (Ellen *et al.*, 2009, Koerdt *et al.*, 2011), metabolomics and systems biology (Ulas *et al.*, 2012, Kouril *et al.*, 2013, Wolf *et al.*, 2016), and metagenomics (Inskeep *et al.*, 2013, Campbell *et al.*, 2017) with these archaea, offering further insights into life in hot acid. The development of genetic systems for thermoacidophiles was challenging, given their unique characteristics and practical considerations related to their growth conditions. In 2003, soon after the availability of its genomic sequence, Paul Blum generated the first *Sulfolobales* mutant in *Sa. solfataricus* based on lactose autotrophy (Worthington *et al.*, 2003). Later in 2009, a genetic system was developed for *Sulfolobus islandicus* (renamed *Saccharolobus islandicus*), based on uracil auxotrophy and the ability to generate uracil through *pyrEF* as a selectable marker (She *et al.*, 2009). Similarly, in 2012, Wagner *et al.* developed a genetic system based on an uracil auxotrophic parent strain and 5-FOA toxicity that is widely used today (Wagner *et al.*, 2012) (**Section 9**). Genetic engineering capabilities for thermoacidophiles have expanded over the past decade that have not only supported fundamental microbiological studies but also fueled prospects for biotechnological processes.

While there are moderately thermoacidophilic bacteria (Norris *et al.*, 1996, Goto *et al.*, 2002, Johnson *et al.*, 2003), most thermoacidophiles are archaea. However, life in thermal, acidic environments is not limited to the order *Sulfolobales*. There are thermoacidophilic Euryarchaeota

belonging to the order *Thermoplasmatales*. Species in the genus *Picrophilus*, such as *Picrophilus oshimae* and *Picrophilus torridus* from solfataras in Japan (Schleper *et al.*, 1995), have an optimum growth temperature of 60°C and a pH optimum near 0. *Thermoplasma acidophilum*, isolated from a coal refuse pile, grows optimally at 59°C and pH of 1-2 (Darland *et al.*, 1970). Here, the focus will be on thermoacidophiles from the order *Sulfolobales* and an examination of what is currently known about their diversity, growth physiology, cell biology, and biotechnological prospects.

2. The Diversity of Thermoacidophilic Life

Thermoacidophilic biotopes are ubiquitously distributed across both terrestrial and marine environments, closely associated with volcanic outflows or calderas resulting from tectonic activity. In terrestrial realms, these environments are often isolated features, presenting as steam-saturated/super-heated discharges (fumaroles) in the form of geysers, solfatara, and pools, and on occasion mixing with soils to form mineral-heavy mud pots. In marine environments, these vents are distinguished by their rapid mixing with dramatically cooler, saline waters (Kelley *et al.*, 2002), resulting in sharp gradients of temperature, pH, oxygen, and solutes, and abrupt dislocated niches (Reysenbach *et al.*, 2000). In both environments, water chemistry is shaped by transformation of sulfur species from highly reduced metal sulfides and hydrogen sulfide to highly oxidized sulfate, with concomitant production of protons (i.e. acid) (Nordstrom *et al.*, 2009). Despite the incredibly exogenic nature of reduced inorganic species, their abiotic transformation at elevated temperatures and low pH is minimal (Chen & Morris, 1972), pointing

to the importance of sulfur biooxidizers in constructing and occupying this extremophilic niche (Odling-Smee *et al.*, 1996).

While sulfur oxidation is a potential bioenergetic source in these environments, strategies to handle thermal stress, acid stress, high levels of aqueous heavy metals, and minimal organic carbon availability must be employed (Section 7). These biotopes are dominated by archaeal chemolithoautotrophs (Inskeep *et al.*, 2013, Ward *et al.*, 2017) that have been intrinsically tailored by evolution to inhabit and thrive in these highly selective niches (Valentine, 2007, Colman *et al.*, 2018). In contrast, many bacterial and eukaryotic organisms in thermal, acidic biotopes are limited to acid- or temperature-tolerant microorganisms, as opposed to obligate/sustained thermoacidophily ($\text{pH} < 3.5$; $T > 65^\circ\text{C}$).

2.1. Diversity of Eukaryotic and Bacterial Thermoacidophiles

Previous efforts have identified the limitations of organisms at the cusp of thermoacidophily. Specifically, in Eukaryotes it appears that the boundary stems from a temperature limitation. As far back as the 1970's, exhaustive sampling and culturing has demonstrated the inability to cultivate eukaryotes (specifically, fungi and algae) from geothermal features in excess of 60°C , despite growth at slightly lower temperatures (Tansey & Brock, 1972). Further work showed that alga are limited to about 60°C (Boyd *et al.*, 2012), and protists to below 70°C (Brown & Wolfe, 2006). For unicellular organisms that inhabit more thermophilic locales, hydrogen sulfide levels can be inhibitory. For many more complex organisms that depend upon gaseous compounds for cellular processes, growth is limited by the solubility of

many gases (oxygen, carbon dioxide, *etc.*), which diminishes with rising temperatures (Rothschild & Mancinelli, 2001).

As is the case with the Eukarya, there are few lineages of Bacteria that are thermoacidophilic. Some bacteria grow at extreme temperatures, in excess of 70°C (*e.g.* the genera *Thermotoga*, *Caldicellulosiruptor*, *Aquifex*), albeit at neutralophilic conditions (Counts *et al.*, 2017). Conversely, there are also a number of acidophilic bacteria, primarily from the genera *Leptospirillum* and *Acidithiobacillus*, that are also autotrophic and are found in acidic features with low organic carbon concentrations. But these bacteria grow optimally at temperatures far below anything considered thermophilic (*i.e.*, 28-45°C); however, *Leptospirillum thermoferrooxidans* grows at temperatures up to 50°C (Kondrat'eva *et al.*, 2012, Dopson, 2016). As temperatures increase, bacteria from the thermotolerant and acidotolerant genera *Sulfbacillus*, *Alicyclobacillus*, and *Hydrogenobaculum* genera are most common (see **Figure 2**).

2.2. Archaeal Thermoacidophilic Diversity

In contrast to the other domains of life, the Archaea predominate in extremely thermoacidophilic features. While culture-independent techniques are rapidly expanding the number of recognized species in acidic hydrothermal environments (*e.g.*, candidate phyla *Geoarchaeota* (Kozubal *et al.*, 2013) and *Marsarchaeota* (Jay *et al.*, 2018)), most of the isolated thermoacidophiles to date originate from the archaeal phyla *Crenarchaeota* and *Euryarchaeota*. In both phyla, almost all isolated and named species are native to terrestrial hot acid environments. Currently, the only thermoacidophile with a validly published name from a marine environment is the deep-sea Euryarchaeon *Aciduliprofundum boonei*, an anaerobic heterotroph

growing optimally at 70°C and pH 4.2-4.8, that utilizes sulfur and iron as electron acceptors (Reysenbach *et al.*, 2006).

Other thermoacidophilic *Euryarchaeota* originate from the order *Thermoplasmatales*, which includes the thermotolerant *Acidiplasma*, as well as the moderately thermophilic *Picrophilus* and *Thermogymnomonas*. The genus *Acidiplasma* contains a few moderately thermophilic acidophiles (pH optimum 1-2, T_{opt} 45-55 °C), including the cell wall-lacking *Acidiplasma aeolicum* (Golyshina *et al.*, 2009). While these organisms grow chemoorganotrophically, they also oxidize iron; for example, *Acidiplasma cupricumulans* (f. *Ferroplasma cupricumulans*) originates from a copper mine heap (Hawkes *et al.*, 2006), and the recently sequenced *Acidiplasma* sp. strain MBA-1 originates from a pyrite-arsenopyrite gold bearing concentrate bioleaching reactor (Bulaev *et al.*, 2017). In contrast, the thermophilic genus *Picrophilus* contains two members: *Picrophilus oshimae* and *Picrophilus torridus*, which are aerobic, heterotrophic organisms from solfatara in Hokkaido Japan, exhibiting remarkable acid tolerance (pH optima less than 1.0), with optimal growth near 60°C (Schleper, 1996). While most organisms maintain a near circum-neutral intracellular pH, *P. oshimae* actually maintains an intracellular pH of approximately 4.6, making it a reservoir for acid-stable cytoplasmic proteins (van de Vossenberg *et al.*, 1998). Additionally, sequencing revealed that *Picrophilus* spp. have some of the smallest genomes (around 1.5 Mb) isolated from free-living organisms (Futterer *et al.*, 2004). Finally, in addition to the well-studied obligately aerobic heterotrophs from the genus *Thermoplasma*, *Thermoplasma acidophilum* and *Thermoplasma volcanium* (Segerer A, 1988), there is a cell wall-less species, *Thermogymnomonas acidicola*, that grows near 60°C, but at slightly higher pH (around 3.0 optimally) (Itoh *et al.*, 2007).

While the temperature optima of the Euryarchaeal thermoacidophiles is limited to around 60°C, the thermoacidophiles from the crenarchaeal phylum all grow at temperatures ranging from 65 – 88 °C. These organisms are composed of three major clades, spanning three orders: *Acidilobales*, *Sulfolobales*, and *Thermoproteales*. While the *Sulfolobales* are a well-studied archaeal lineage (over 30 named species, across 7 genera, and more than 20 distinct genomes), the other two lineages *Acidilobales* (containing *Caldisphaera* and *Acidilobus*) and *Thermoproteales* (only the *Caldivirga* are thermoacidophiles) contain just a few named, characterized strains. From the order *Acidilobales*, there are just two genera *Acidilobus* and *Caldisphaera*, belonging to families derived from the same names. Both groups consist of anaerobic heterotrophs, growing optimally at pH ranging from 2.5-4.0 (mostly moderate acidophiles) and temperatures around 70-75 °C for the *Caldisphaera* and slightly elevated temperatures of 50-80 °C for the *Acidilobus spp.* (Prokofeva *et al.*, 2000, Itoh *et al.*, 2003, Boyd *et al.*, 2007, Prokofeva *et al.*, 2009). Meanwhile, the genus *Caldivirga* is represented by a single member, *Caldivirga maquilingensis*, isolated from the Philippines, which is capable of anaerobic (and microaerophilic) growth on heterotrophic substrates at moderate pH (optimum 3.7-4.2) and extremely thermophilic conditions (85 °C) (Itoh *et al.*, 1999).

2.3. The *Sulfolobales*

As mentioned previously, one of the first archaeal lineages discovered was the *Sulfolobales*, named for their presence and perceived usage of sulfur by Thomas Brock from his excursions to Yellowstone in the 1960's (Brock *et al.*, 1972). Over the course of the following decades, a number of intriguing microorganisms emerged from terrestrial hot springs throughout

the world, representing the seven named genera today from the order: *Acidianus*, *Metallosphaera*, *Saccharolobus* (f. *Sulfolobus*), *Stygiolobus*, *Sulfodiicoccus*, *Sulfolobus*, *Sulfuracidifex* (f. *Sulfolobus*), and *Sulfurisphaera* (f. *Sulfolobus*) (Counts, 2020). These include organisms with a broad array of physiological traits, ranging from extreme to moderate acidophily (0.7-4.5), thermophily (65-88°C), obligate and facultative aerobes, obligate anaerobes, metal oxidizers, sulfur reducers/oxidizers, chemoheterotrophs and chemolithoautotrophs.

Species in the thermoacidophilic genus *Acidianus* grow anaerobically, reducing sulfur in its various forms, or aerobically, oxidizing sulfur (Segerer *et al.*, 1986). The genus contains the most acidophilic *Sulfolobales* member to date: *Acidianus sulfidivorans* ($\text{pH}_{\text{opt}} \sim 0.7$), and the most thermophilic member: *Acidianus infernus* ($T_{\text{opt}} \approx 88^{\circ}\text{C}$) (Segerer *et al.*, 1986, Plumb *et al.*, 2007). The order also contains several members with metal biooxidation capabilities (Huber *et al.*, 1989, Huber & Stetter, 1991) *Acidianus ambivalens* (f. *Desulfurolobus ambivalens*) has long-served as a model for the study of sulfur biotransformation within the *Sulfolobales* (Laska *et al.*, 2003, Müller *et al.*, 2004, Brito *et al.*, 2009, Protze *et al.*, 2011).

The genus *Metallosphaera* was named for the perceived ability of its members to biooxidize iron (and by proxy release other metals from ores, *e.g.* copper) (Huber *et al.*, 1989). The type species, *Metallosphaera sedula*, along with the recently isolated *Metallosphaera yellowstonensis*, serve as model systems for metal biooxidation by extremely thermoacidophilic archaea (Auernik *et al.*, 2008, Kozubal *et al.*, 2011). Further, *M. sedula* has also been examined for autotrophy catalyzed by the 3-hydroxypropionate/4-hydroxybutyrate cycle (Berg *et al.*,

2010), which has shown promise for metabolic engineering of biosynthetic pathways (Hawkins *et al.*, 2013, Keller *et al.*, 2013, Lian *et al.*, 2016, Straub *et al.*, 2018).

In contrast to the other genera of the *Sulfolobales*, the genus *Saccharolobus* appears to depend less on lithotrophic pathways and more, as its name suggests, on sugar catabolism (Sakai & Kurosawa, 2018). Members of the genus *Saccharolobus* are mostly aerobic and are among the most thermophilic ($T_{opt} \geq 80$ °C) and least acidophilic organisms in the order ($pH_{opt} \geq 3.0$) (Zillig *et al.*, 1980, Grogan, 1989, Sakai & Kurosawa, 2018). Their original taxonomical placement in *Sulfolobus* was changed following growing genomics information pointing to evolutionary divergence (Sakai & Kurosawa, 2018). In fact, a number of unnamed and informally named species, *e.g.* “*Sulfolobus islandicus*”, appear to be much more closely related to other members of the genus *Saccharolobus*, which is fitting given that they use pentoses, hexoses, and di-, tri-, and polysaccharides (Grogan, 1989).

The main representative of the genus *Sulfolobus*: *S. acidocaldarius*, has a much narrower range of carbohydrate utilization. *S. acidocaldarius* grows best at 75 °C and pH 3.0, using only amino acids, sucrose, dextrin, and starch (Grogan, 1989). This archaeon was originally named for its perceived capability to oxidize sulfur in the sulfur-rich pools of Yellowstone National Park (Brock *et al.*, 1972). Despite these early reports, sulfur biooxidation capacity in strains that are currently available from culture collections is limited. However, recent studies showed that sulfur oxidation can be restored in *S. acidocaldarius* DSM 639 by inserting genes encoding sulfur oxygenase reductase (SOR) and thiosulfate:quinone oxidoreductase (TQO) (Zeldes *et al.*, 2019), perhaps reflecting an evolutionary connection to this process. *S. acidocaldarius* has emerged as a tractable genetic platform to understand the physiological features of the

Sulfolobales, such as pili structure controlling motility (Albers & Jarrell, 2015), UV-stress response (Wagner *et al.*, 2012), biofilm formation (van Wolferen *et al.*, 2020), and cellular division (Pulschen *et al.*, 2020). In addition to *S. acidocaldarius*, *Saccharolobus solfataricus* and “*Sulfolobus islandicus*” are currently the only *Sulfolobales* with tractable genetic systems (Straub *et al.*, 2018).

The remaining genera are represented by only a few named species, but vary dramatically in some of their observed traits. For example, the genus *Stygiolobus* contains a single member, *Stygiolobus azoricus* (T_{opt} 80 °C and pH 2.5-3.0), and is the only obligate anaerobe from the order to date, capable of sulfur reduction in the presence of hydrogen (Segerer *et al.*, 1991). The genus *Sulfurisphaera* contains three species: *Sulfurisphaera javaensis*, *Sulfurisphaera tokodaii*, and *Sulfurisphaera ohwakuensis* (the genus type species), all of which are extremely thermophilic (optima 80-85 °C), but vary with respect to acidophily (optima 2.0-4.0) (Tsuboi *et al.*, 2018). *Sulfurisphaera* species are facultative anaerobes and oxidize sulfur and iron to varying extents, and grow on complex organic substrates (Kurosawa *et al.*, 1998, Tsuboi *et al.*, 2018). In contrast, the two current members of the genus *Sulfuracidifex*: *Sulfuracidifex* (f. *Sulfolobus*) *metallicus* and *Sulfuracidifex tepidarius*, are less thermophilic acidophiles (temperature optimum: 65 °C; pH optima: 2.0-3.5) and obligately aerobic chemolithoautotrophs, capable of mixotrophic growth in the presence of reduced sulfur compounds (Huber & Stetter, 1991, Itoh *et al.*, 2020). *Sulfura. metallicus* has served as a model system for metal biooxidation studies (Bathe & Norris, 2007). The genus *Sulfodiicoccus* is another single-member genus (type species *Sulfodiicoccus acidiphilus*), growing optimally at 65-70 °C and pH 3.0-3.5. This archaeon is different from other Sulfolobales in that it is not only unable to oxidize elemental sulfur, but is

possibly inhibited by it (Sakai & Kurosawa, 2017). Furthermore, *S. acidiphilus* also lacks key components for carbon dioxide fixation by the 3-hydroxypropionate/4-hydroxybutyrate cycle and apparently does not grow autotrophically (Sakai & Kurosawa, 2017, Sakai & Kurosawa, 2019). See **Table 1** for a listing of thermoacidophilic microorganisms,

3. Viruses and CRISPR Systems of Thermoacidophiles

Thermoacidophiles share their natural habitat with viruses (Munson-Mcgee *et al.*, 2018). A recent survey of the viral communities in thermal hot springs in Yellowstone National Park showed that > 60% of cells were infected by viruses and that the majority even contained two or more virus types at the same time (Munson-Mcgee *et al.*, 2018). Consequently, viruses represent an important evolutionary pressure in these archaeal dominated environments. The ongoing arms race between viruses and their hosts has led to the development of anti-viral defense strategies and mechanisms from viruses to circumvent them (Borges *et al.*, 2017, Hwang & Maxwell, 2019, Hampton *et al.*, 2020). However, the fact that in many cases cells carry multiple virus types suggests that viruses can also have beneficial relationships with their microbial hosts. Viruses shape microbial populations, are a major driver of microbial evolution, and impact host ecology. An excellent example is the virus-host mutualism by which chronically virus infected-*Sulfolobus* cells kill the virus-resistant cells in the population (DeWerff *et al.*, 2020).

3.1. Viruses of thermoacidophiles

In comparison with known bacterial and eukaryotic viruses, only a modest number of archaeal viruses have been isolated to date (Prangishvili *et al.*, 2017). However,

thermoacidophiles, especially members of the *Sulfolobales*, have proven to be a very rich source of archaeal viruses (Prangishvili *et al.*, 2001, Prangishvili *et al.*, 2017, Munson-Mcgee *et al.*, 2018). These viruses are characterized by a large genetic and morphological diversity, including many unique shapes that are not found in viruses infecting bacteria and eukaryotes (Pina *et al.*, 2011, Prangishvili *et al.*, 2017). The evolutionary origin of archaeal viruses is not clear, but the high diversity might have originated during the early stages of evolution of cellular life, maintained in Archaea, and lost in bacterial and eukaryotic lineages (Prangishvili *et al.*, 2006, Prangishvili, 2015, Prangishvili *et al.*, 2017). All isolated viruses from *Sulfolobales* have DNA genomes, and the majority of their gene products have unknown functions (Prangishvili *et al.*, 2017). Metagenomic analysis indicated the presence of viruses with RNA genomes in high-temperature acidic hot springs (Bolduc *et al.*, 2012). However, viral particles were not isolated, and the exact host remains unknown (Bolduc *et al.*, 2012, Stedman *et al.*, 2013).

Members of at least 8 different viral families infect thermoacidophilic archaea: bottle-shaped *Ampullaviridae* (Haring *et al.*, 2005), tailed *Bicaudaviridae* (Häring *et al.*, 2005), spindle-shaped *Fuselloviridae* (Schleper *et al.*, 1992), droplet-shaped *Guttaviridae* (Arnold *et al.*, 2000), filamentous *Lipotrixviridae* (Bettstetter *et al.*, 2003), polyhedral *Portogloboviridae* (Liu *et al.*, 2017), rod-shaped *Rudiviridae* (Prangishvili *et al.*, 1999), and the icosahedral *Turrriviridae* (Rice *et al.*, 2004) (**Figure 3**). The diversity of morphotypes encountered among viruses infecting thermoacidophilic *Crenarchaea* is in stark contrast to that found for euryarchaeal or bacterial viruses, which are dominated by head-tail morphologies (Pietilä *et al.*, 2014, Prangishvili *et al.*, 2017). Interestingly, recently available cryo-EM structures have shown that several viruses infecting members of the *Sulfolobales* package their dsDNA genome in A-form (DiMaio *et al.*,

2015, Wang *et al.*, 2019). A-form DNA was at first thought to be an artifact and have no biological significance, but the widespread usage of A-form DNA by archaeal viruses suggests that this packaging helps to protect the viral genomes against adverse conditions in thermal hot springs (Wang *et al.*, 2019).

Among the thermoacidophiles, *Sa. islandicus* and *Sa. solfataricus* are model systems for the study of virus-host interactions in *Crenarchaea* (Pina *et al.*, 2011, Prangishvili *et al.*, 2013, Dellas *et al.*, 2014). The valuable, but still limited, knowledge on infection strategies of crenarchaeal viruses mainly derives from viruses infecting these organisms. For initial attachment and entry into the cell, the various surface appendages with which *Sulfolobales* are covered (see **Section 6**) are important for multiple viruses. *Sulfolobus* turreted icosahedral virus (STIV1) binds with its turrets to thin filaments of unknown identity on the surface of *Sa. solfataricus* (Hartman *et al.*, 2019). *Sa. islandicus* rod-shaped virus (SIRV) particles use the three tail fibers that are present at the distal parts of the virion for attachment to adhesive type IV pili on the its surface (Quemin *et al.*, 2013, Deng *et al.*, 2014, Rowland *et al.*, 2020). Like SIRV, the *Sulfolobus* spindle-shaped virus SSV requires pili for infection, but the particles do not directly attach to the pili, and the role of pili in viral entry is unresolved (Rowland *et al.*, 2020). Primary attachment to filamentous surface structures is a common strategy of bacterial viruses, which can increase the chances of successful infection (Poranen *et al.*, 2002, Quemin & Quax, 2015). The mechanisms by which viruses move along archaeal filaments to the cell surface are unknown and are likely different from those of bacterial viruses, since the archaeal surface filaments have different structural organization (see **Section 6**) (Quemin & Quax, 2015, Chaudhury *et al.*, 2018).

Once virions have attached, their genomes can enter the cell. Some viruses, such as SSV, can integrate their genome into that of the host (Muskhelishvili *et al.*, 1993, Serre *et al.*, 2002, Clore & Stedman, 2007). Circularization and replication of the integrated SSV genome is induced by UV light (Schleper *et al.*, 1992, Fröls *et al.*, 2008). SSV has been employed to develop a genetic manipulation system for *Sa. solfataricus* (see **Section 9**). Several other viruses do not integrate, instead replicating directly after entry. The replication mechanism of only a few thermoacidophile model viruses has been studied. For example, replication of the dsDNA genome of *Acidianus* Filamentous Virus 1 (AFV1) relies on recombination events for initiation and termination and has a terminally bound protein (Pina *et al.*, 2014). Replication of the linear dsDNA genome of SIRV requires a virus-encoded dimeric Rep protein that initiates replication by making single stranded nicks and the virus-encoded holliday junction resolvase (Hjr) to resolve viral genome concatemers (Blum *et al.*, 2001, Peng *et al.*, 2001, Oke *et al.*, 2010, Oke *et al.*, 2011). SIRV Hjr interacts with proliferating cell nuclear antigen (PCNA), a key replication protein in archaea (Gardner *et al.*, 2014). Interestingly, SIRV forms a distinct replication focus in the cell to which viral and host replication proteins are specifically recruited (Martínez-Alvarez *et al.*, 2017).

After genome replication, virions are formed in the cytoplasm. Virion maturation can occur (i) before, (ii) during, or (iii) after release. (i) Several lytic viruses infecting *Sulfolobales*, such as STIV and SIRV, were shown to mature in the cytoplasm and employ an unusual lysis mechanism that relies on the formation of seven-fold symmetric pyramidal egress structures, of which a dozen form during viral infection on the host cell surface (Bize *et al.*, 2009, Brumfield *et al.*, 2009, Prangishvili & Quax, 2011, Quax *et al.*, 2011, Quax & Daum, 2017) (**Figure 3**). These

~150 nm structures consist of one viral protein and protrude through the protective S-layer (see **Section 6**) (Fu *et al.*, 2010, Quax *et al.*, 2010, Snyder *et al.*, 2011, Daum *et al.*, 2014). They open outwards at the end of the infection cycle to allow for the release of virions (Bize *et al.*, 2009, Brumfield *et al.*, 2009, Fu *et al.*, 2010, Daum *et al.*, 2014). (ii) SSV matures upon egress, as the virions are released via budding and are covered in a lipid layer during this process (Quemin *et al.*, 2016). In fact, this is the first case of budding observed for a prokaryotic virus. Budding viruses allow for a continuous release of virions and the cells remain alive throughout the infection cycle (Schleper *et al.*, 1992, Quemin *et al.*, 2016). The ESCRT-III system could play a role in the budding of archaeal viruses (**Figure 3**) (Liu *et al.*, 2017). (iii) *Acidianus* two-tailed virus (ATV) and *Sulfolobus* monocaudavirus (SMV1) are exceptional viruses for which virion maturation (the lengthening of the tails) happens outside and independent of the host cell, after viral release (Häring *et al.*, 2005, Prangishvili *et al.*, 2006, Scheele *et al.*, 2011, Uldahl *et al.*, 2016). These tails consist of helically arranged globular subunits that develop from the two pointed ends of the virion when it is outside the host cell (Prangishvili *et al.*, 2006). High temperatures are required for this morphological transformation. In summary, viruses of thermoacidophiles are unique because of their diverse morphologies and the model virus-host systems of the *Sulfolobales* have provided important insights into the infection strategies of crenarchaeal viruses in general.

3.2. CRISPR-Cas mediated viral defense in thermoacidophiles

The omnipresence of viruses in archaeal habitats has resulted in the development of several anti-viral defense strategies, of which CRISPR-Cas is without doubt the best known.

CRISPR (Clustered Regularly InterSpaced Palindromic Repeats) systems in *Sulfolobales* have been studied since the ‘early days’ of CRISPR research (Vestergaard *et al.*, 2008, Han & Krauss, 2009, Held & Whitaker, 2009, Lillestøl *et al.*, 2009, Garrett *et al.*, 2011, Zhang & White, 2013), just after these systems were suggested to play a role in defense against viruses in bacteria and archaea (Bolotin *et al.*, 2005, Mojica *et al.*, 2005, Pourcel *et al.*, 2005, Makarova *et al.*, 2006). CRISPR arrays consist of a series of ~30 bp genomic repeats, which are interspaced by unique sequences that can match foreign genetic elements (van der Oost J, 2014). CRISPR associated proteins (Cas) are usually encoded in the proximity of the CRISPR array. Upon a viral infection, new spacers, exactly matching the genome of the infecting virus, are integrated between two repeats. The arrays are then processed by Cas proteins, and the spacer is used as a guide to specifically target and interfere with the matching sequences in the viral genome (Barrangou *et al.*, 2007, Barrangou & Horvath, 2017, Jackson *et al.*, 2017). Thus, CRISPR-Cas provides specific and inheritable immunity (Barrangou *et al.*, 2007, Brouns *et al.*, 2008). CRISPR arrays can also be used as a map to track previous encounters with viruses and to indicate viral host range (Bautista *et al.*, 2017, Munson-Mcgee *et al.*, 2018, Pauly *et al.*, 2019).

CRISPR-Cas systems are present in ~40% of bacteria, most archaea (85%), and almost all extreme thermophiles (97%) (Makarova *et al.*, 2019). Based on the Cas proteins, the CRISPR systems have been divided into several different groups, and this division keeps evolving as new systems are being discovered (Makarova *et al.*, 2019). *Crenarchaea*, such as the *Sulfolobales*, usually harbor multiple CRISPR systems in their genome, and generally have longer CRISPR arrays than bacteria (Zhang & White, 2013). Crenarchaeal genomes are substantially enriched for type III systems of class 1 (Zhu *et al.*, 2018, Makarova *et al.*, 2019), which rely on transcription-

dependent (specific RNA binding and cleavage) and subsequent (non-specific) DNA degradation (Deng *et al.*, 2013, Goldberg *et al.*, 2014, Zhu *et al.*, 2018). Type III systems typically possess a Cas10 protein with a Palm polymerase domain that can cyclize ATP to generate cyclic oligoadenylylate (cOA) to act as a second messenger (Kazlauskiene *et al.*, 2017, Niewoehner *et al.*, 2017, Rouillon *et al.*, 2018). Formation of cOA leads to signal amplification that activates other defense mechanisms, including host and viral DNA degradation that results in immunity or cell dormancy (Rouillon *et al.*, 2018, Rostøl & Marraffini, 2019). In *Saccharolobus*, cyclic tetra-adenylate (cA₄) can be degraded by host-encoded ring-nucleases to reset the signal (Athukoralage *et al.*, 2018). Interestingly, some archaeal viruses encode a potent ring nuclease that acts as an anti-CRISPR (Arc) (Athukoralage *et al.*, 2020, Athukoralage *et al.*, 2020). More Arc's have been identified in archaeal viruses, such as those that bind and inhibit type III-B or I-D CRISPR systems (He *et al.*, 2018, Bhoobalan-Chitty *et al.*, 2019). Different viral families probably use different strategies to evade CRISPR-Cas immunity, as natural populations of *Sulfolobus* have developed CRISPR-Cas immunity with a different structure and diversity in response to SIRV and SSV viruses infections (Pauly *et al.*, 2019).

Despite the viral strategies to evade CRISPR-Cas mediated defense, this seems an effective immune system in extreme thermophiles ($T_{opt} \geq 70^\circ\text{C}$), illustrated by the presence of CRISPR-Cas in nearly all of their genomes. It remains to be seen why CRISPR-Cas systems are so ubiquitous, specifically in extremely thermophilic archaea. One possible explanation relates to the lower mutation rate of viruses in extreme environments that, combined with the lower population sizes of extreme thermophiles compared to mesophiles, gives extremely thermophilic

viruses limited possibility to escape immunity (Weinberger *et al.*, 2012, Prangishvili *et al.*, 2017).

Besides CRISPR-Cas, *Crenarchaea* are specifically enriched in Toxin-Antitoxin systems (TA), which play a role in abortive infection in bacteria (Koonin *et al.*, 2017). Viral infection of different *Sulfolobales* induces expression of TA systems (Ortmann *et al.*, 2008, Quax *et al.*, 2013, León-Sobrino *et al.*, 2016). Furthermore, several thermoacidophiles encode an Argonaute protein, which has been implicated in defense against foreign genetic elements (Makarova *et al.*, 2009, Swarts *et al.*, 2014, Swarts *et al.*, 2014, Willkomm *et al.*, 2017). However, it needs to be verified experimentally if archaeal TA or Argonaute systems provide immunity to viral infection. Certainly, thermoacidophiles encode novel viral defense mechanisms that are awaiting discovery (Doron *et al.*, 2018).

4. Genetic Mechanisms

Effective packaging and organization of genomic DNA into the confined space of the nucleus or nucleoid, while at the same time enabling a dynamic and reliable genome replication and gene expression, is essential for every living organism. The underlying molecular mechanisms (replication, transcription and translation) are central to life. The study of genetic mechanisms in archaea lags behind those focused on bacteria and eukaryotes. However, it is clear that archaeal information processing machineries are related to their eukaryotic counterparts, that gene regulation processes are bacteria-like, and that chromosome organization has both eukaryote-like and bacteria-like features (Peeters *et al.*, 2015, Blombach *et al.*, 2019, Lemmens *et al.*, 2019, Greci & Bell, 2020).

Thermoacidophilic *Crenarchaea* belonging to the *Sulfolobales* order have served as an archaeal model system to study chromosome organization, DNA replication, transcription, and translation processes. Although many insights can be extended further to the entire archaeal domain, there are also unique, lineage-specific aspects. For example, while most archaea harbor eukaryote-like histones involved in chromosome structuring, these are completely absent in thermoacidophilic archaea, namely in all *Crenarchaeota* and in *Thermoplasma acidophilum* (Peeters *et al.*, 2015, Hocher *et al.*, 2019).

4.1. Chromosome packaging and structuring

Thermoacidophilic archaea typically have a single circular, relatively small chromosome with a size between 1.5 and 3 Mbp (Chen & Morris, 1972, She *et al.*, 2001). This chromosome is packaged into a condensed and organized chromatin structure by the action of different types of chromatin proteins (**Figure 4**). There is a large evolutionary divergence in chromatin proteins present in archaea, including the *Sulfolobales*; while histone orthologs are absent in this lineage, an interplay exists between a variety of nucleoid-associated proteins (NAPs) with different levels of conservation (Peeters *et al.*, 2015). These proteins are small (between 7 and 10 kDa), basic and highly abundant in the cell, constituting up to 5% of soluble cellular protein (Mai *et al.*, 1998). They harbor DNA-binding motifs that are also found in specific transcription regulators (**see below, Section 4.3.2**), such as the winged helix-turn-helix (wHTH) motif, and bind DNA with low or no sequence specificity. In case of low sequence specificity, these NAPs typically prefer GC-rich sequences (Kalichuk *et al.*, 2016, Hocher *et al.*, 2019). The *Sulfolobales* harbor two paralogs of the archaea-universal NAP Alba, which was initially assumed to be an important

chromatin structuring protein (Bell *et al.*, 2002), with the extent of heterodimerization between the two paralogs determining the architectural effects (Laurens *et al.*, 2012). This hypothesis has recently been revisited as Alba was shown to be a general nucleic-acid binding protein interacting with RNA as well (Guo *et al.*, 2014). Besides Alba, the NAP Cren7 is highly conserved in not only the *Sulfolobales* but in all *Crenarchaeota*; Cren7 is a versatile architectural protein, bending and also bridging DNA, thereby forming highly condensed chromatin filaments (Guo *et al.*, 2008, Zhang *et al.*, 2019, Zhang *et al.*, 2020). Sul7d, analogous to Cren7, is a monomeric protein, which is capable of bending DNA; it has been found in many *Sulfolobales* genera: *Sulfolobus* and *Saccharolobus*, *Acidianus*, *Metallosphaera*, *Stygiolobus* and *Sulfurisphaera* (Kalichuk *et al.*, 2016). In addition, there are species-specific NAPs, such as the Sso10a paralogs in *S. solfataricus*, that are dimeric proteins capable of bending DNA and either bridging it or forming filaments (Driessens *et al.*, 2016). Finally, the euryarchaeal *T. acidophilum*, which also lacks histones, harbors a NAP that is homologous to the bacterial HU family (Hocher *et al.*, 2019).

The heterogeneity in the NAP protein machinery responsible for packaging DNA in the *Sulfolobales* (**Figure 4 B**) -when considering a single species- is hypothesized to accommodate differential local chromatin structuring when expression levels or posttranslational modifications (PTMs) of the individual NAPs are altered. This in turn might affect transcriptional expression in a polygenic manner (Peeters *et al.*, 2015). Indeed, PTMs have been observed for the NAPs Alba, Cren7, Sso7d and Sul7d (Vorontsov *et al.*, 2016). Early studies postulated that acetylation of the Lys16 residue of Alba constitutes a global gene regulation mechanism similar to eukaryotic histone modification (Bell *et al.*, 2002). This has later been refuted and determined to be an N-

terminal acetylation event instead that does not affect the nucleic-acid binding capacity of Alba and is possibly involved in protein turnover regulation (Ma *et al.*, 2016, Cao *et al.*, 2018).

Besides acetylation, methylation occurs widely on these NAPs (Niu *et al.*, 2013, Vorontsov *et al.*, 2016) and might be linked to thermostabilization of the chromatin, as well as to epigenetic mechanisms of gene regulation. The existence of such epigenetic mechanisms was recently demonstrated for a strain of *Sa. solfataricus* that was evolved in an adaptive laboratory evolution experiment and displayed a super-acid resistant phenotype. This evolved strain harbored no genomic changes with respect to the original strain (Payne *et al.*, 2018, Johnson *et al.*, 2019).

Instead, the acid resistance appeared to be mediated by a different methylation status of the NAPs Cren7 and Sso7d. Other than NAP methylation, methylation of the genomic DNA itself might also be responsible for epigenetic mechanisms of gene regulation. Recently, the DNA methylome of *S. acidocaldarius* was mapped and shown to consist of base methylations that are more than just a part of restriction modification defense systems (Couturier & Lindås, 2018). More specifically, N⁶-methyl-adenine methylations were found and hypothesized to be involved in the regulation of the cell cycle or other biological functions (Couturier & Lindås, 2018).

Besides NAP-mediated local structuring, the genome is also organized in domains at a higher level. This higher-order chromosome organization has recently been elucidated for *S. acidocaldarius* and *Sa. islandicus* using the Hi-C methodology, a combination of chromosome conformation capture (3C) and high-throughput sequencing (Takemata *et al.*, 2019). Similar to what has been observed in metazoan eukaryotes and very different from what is observed in bacteria, the *Sulfolobales* genome is organized in two distinct sub-Mbp compartments, each characterized by a different average level of transcription (Takemata *et al.*, 2019) (**Figure 4 A**).

While the A compartment, which harbors genes mainly involved in core metabolic processes such as protein biogenesis, is transcriptionally active, the B compartment appears to be in a more silent transcriptional state and harbors genes that function in diverse metabolic pathways and physiological processes. For example, gene expression of the B compartment is typically induced in response to environmental stress conditions, such as the archaellum motility apparatus and a fatty acid metabolism gene cluster. In addition, the B compartment is enriched in mobile genetic elements, such as CRISPR-Cas clusters. Although it is still unclear which proteins are responsible for the active structuring of the chromosome into the two compartments, a major role has been described for the novel chromatin structuring protein coalescin (ClsN) (Takemata *et al.*, 2019). While the *Sulfolobales* do not harbor a homolog of the typical condensin complex belonging to the Structural Maintenance of Chromosome (SMC) family, conserved in bacteria, eukaryotes and most other archaea (Kamada & Barillà, 2018), they possess the SMC-like ClsN instead, which is significantly smaller than the SMC subunits of condensin and possibly has a zinc hook domain instead of a hinge (Takemata *et al.*, 2019). There is an inverse correlation between the presence of ClsN and the transcriptional machinery, with ClsN being mainly associated with transcriptionally less active genes in the B compartment. It is hypothesized that the protein assists in the higher-level compartmentalization by mediating intra- and interdomain interactions (Takemata *et al.*, 2019).

4.2. DNA replication

One of the most important transactions undergone by genomic DNA is its replication as part of the cell division process. It is striking that, similar to eukaryotes, several archaeal lineages

are characterized by the chromosome harboring multiple replication origins, genomic sites at which replication is initiated. In contrast, bacteria only have a single origin. *Sulfolobales* are characterized by three replication origins (OriC1, 2 and 3), each of which accommodate a single replication initiation event during the cell cycle (Lundgren *et al.*, 2004, Robinson *et al.*, 2004, Robinson & Bell, 2007, Duggin *et al.*, 2008). Mutagenesis analysis indicated that, while none of the individual OriCs is essential, at least one is required (Samson *et al.*, 2013). The observation that all three OriCs are located in the transcriptionally active chromosome compartment A (**Figure 4 A**) might suggest that DNA replication processes are involved in higher-order chromatin structuring. However, this appears not to be the case as OriC mutant strains do not display any differences in their chromatin structure (Takemata *et al.*, 2019).

The archaeal replication machinery resembles the eukaryotic machinery, as exemplified by the well-described machinery of *Sa. solfataricus* (**Figure 4 C**) (Dionne *et al.*, 2003, Greci & Bell, 2020). Archaeal initiator proteins, responsible for OriC recognition and assembly of the replisome, are related to the eukaryotic Orc1 and Cdc6 replication initiation proteins, which are characterized by an N-terminal AAA+ fold and a C-terminal wHTH domain (Cunningham Dueber *et al.*, 2007). *Sulfolobales* encode three Orc1/Cdc6-like paralogs, with Orc1-1 specifically recognizing the origin recognition boxes (ORBs) in OriC1 (Samson *et al.*, 2016). Upon binding ATP, two inversely bound Orc1-1 proteins recruit two MCM homohexamers, which are the 3'-to-5' helicases (Samson *et al.*, 2016, Meagher *et al.*, 2019). Also, analogous to the eukaryotic system, the MCM helicase associates with additional replication proteins, forming the so-called CMG (Cdc45-MCM-GINS) replisome core (**Figure 4 C**). In *S. acidocaldarius* and *Sa. islandicus*, MCM recruits a Cdc45 ortholog and two GINS-like proteins Gins23 and Gins15,

each protein having a homodimeric composition in the complex (Xu *et al.*, 2016). In contrast, in *T. acidophilum* a homotetrameric GINS protein is part of the CMG complex (Ogino *et al.*, 2017). Although it is unclear how the melting of the DNA helix is accomplished in *Sulfolobales* after assembly of the replisome at the replication origin, the involvement of a replication-dedicated DNA-dependent RNA polymerase responsible for primer synthesis has been established. This DNA primase initiates leading strand synthesis, or the synthesis of Okazaki fragments for lagging strand synthesis, and is recruited by interacting with the GINS complex (Marinsek *et al.*, 2006), with the primase in *Sa. islandicus* being a heterotrimer PriSLX (Liu *et al.*, 2015). Finally, the enzyme responsible for DNA synthesis, DNA polymerase, has also been shown to be eukaryote-like. *Crenarchaeota* possess three different B-family DNA polymerases, with PolB1 being essential and PolB2 and PolB3 shown not to be required for cell viability and hypothesized to be involved in DNA damage repair (Greci & Bell, 2020). Prior to elongation a sliding clamp, constituted by a heterotrimeric PCNA protein (Dionne *et al.*, 2003), is loaded onto the DNA by replication factor C (RFC) and forms a ring-shaped structure. This clamp functions as a molecular platform to recruit the DNA polymerase and other replication-associated enzymes. In contrast to bacteria, which harbor site-specific mechanisms, replication termination appears to be mediated by passive-fork collision taking place halfway between the active replication origin(s) in a site-unspecific manner (Lundgren *et al.*, 2004, Duggin *et al.*, 2011, Samson *et al.*, 2013).

Sulfolobales have an organized cell cycle (**Section 6.4**), with well-defined gap phases and in which the process of DNA replication is temporally separated from the process of chromosome segregation (Bernander & Poplawski, 1997). Following the S phase in which the chromosome is replicated, there is a significant post-replicative period (G2 phase) in which sister

chromatids remain bound together to form hemicatenane structures (Robinson & Bell, 2007).

Next, chromosome segregation is accomplished by a bacterial-like ParAB-like system; in *Sa. solfataricus*, this system consists of SegA, an ortholog of the bacterial, Walker-type ParA ATPase protein, and an archaea-specific DNA-binding protein named SegB (Kallioma-Sanford *et al.*, 2012).

4.3. Transcription and its regulation

4.3.1. Basal transcription machinery in *Sulfolobales*

The small genomes of archaea share their genetic organization with bacteria, with an operonic transcription unit structure that is dense and characterized by short intergenic regions. Similar to other information processing steps, mechanisms of basal transcription have been extensively studied in archaeal species belonging to *Sulfolobales*, ranging from focused biochemical studies with *in vitro* reconstituted transcription systems (Qureshi *et al.*, 1997, Bell *et al.*, 1999, Blombach *et al.*, 2019) to high-resolution mapping of the transcriptome (Wurtzel *et al.*, 2010, Cohen *et al.*, 2016, Dar *et al.*, 2016). Not long after the isolation of the first thermoacidophilic crenarchaeal isolate *S. acidocaldarius*, Wolfram Zillig performed a biochemical analysis of its RNA polymerase, which is the key enzyme of transcription, thereby concluding that its subunit pattern resembles that of eukaryotic RNA polymerase (Zillig *et al.*, 1979). Much later, structural analysis of the *Sa. solfataricus* and *Sa. shibatae* RNA polymerases confirmed that they are complexes consisting of 13-subunit proteins that display an evolutionary relationship with the eukaryotic RNA polymerase II (Hirata *et al.*, 2008, Korkhin *et al.*, 2009). Transcription initiation requires a set of additional general transcription factors that are also

homologous to eukaryotic factors: TATA-binding protein (TBP), transcription factor B (TFB), and transcription factor E (TFE) (**Figure 4 D**). A typical promoter region in *Sa. solfataricus* is characterized by a core TATA-box region of which the center is located about 26 base pairs (bps) upstream of the transcription initiation site, which is directly preceded by a purine-rich factor B recognition element (BRE) (Wurtzel *et al.*, 2010). Besides these canonical archaeal promoter elements, an additional 6-bp, AT-rich, conserved promoter element was identified in *Sa. islandicus* just upstream of the transcription start site (TSS), named initiator (Inr) (Ao *et al.*, 2013).

Initiation of transcription proceeds by the stepwise assembly of the different components in the Pre-Initiation Complex (PIC), which was first studied with *Sa. shibatae* (Qureshi *et al.*, 1997, Bell *et al.*, 1999). First, the highly symmetrical TBP binds the TATA box region followed by the association of TFB to the TBP-DNA complex. By specific recognition of the BRE promoter element, TFB determines the correct orientation of the PIC (Bell *et al.*, 1999). Next, the RNA polymerase as well as TFE are recruited, with TFE being a heterodimeric protein consisting of TFE α and TFE β subunits in *Sulfolobales* (Blombach *et al.*, 2015). Although TFE is not absolutely required for *in vitro* transcription reactions to proceed, it stabilizes the PIC and facilitates DNA melting in the Inr promoter region, thereby assisting the formation of an open complex during the transitioning from the initiation to the elongation phase (Bell *et al.*, 2001, Blombach *et al.*, 2015). During elongation, TBP and TFB dissociate from the RNA polymerase, which is assisted by different transcription elongation factors. Most of these factors, such as the transcript cleavage factor TFS and the processivity factors Spt4/5 and Elf1, are also evolutionarily related to eukaryotic elongation factors (Fouqueau *et al.*, 2017, Blombach *et al.*,

2019). Finally, although transcription termination remains understudied in archaea, a transcriptome-wide Term-seq approach enabled the mapping of all RNA 3' termini in *S. acidocaldarius*, revealing a widespread occurrence of multiple terminators. This leads to alternative 3' isoforms, with U-rich terminator motifs retrieved for 53% of all transcription units (Dar *et al.*, 2016).

To some extent, components of the basal transcription machinery are capable of mediating a global regulation of the transcription initiation process. Certain archaea harbor multiple paralogs of TBP and TFB and it is hypothesized that these are employed for global gene regulation in a similar way as alternative sigma factors in bacteria (Facciotti *et al.*, 2007). *Sulfolobales* typically harbor a single TBP and three TFB paralogs, with TFB3 being a truncated form. The latter functions as a transcriptional activator in a *trans* dependent manner on the canonical TFB1 (Paytubi & White, 2009). TFB3 activates the expression of Ups pili and the Ced DNA import system in response to UV irradiation (Paytubi & White, 2009, Feng *et al.*, 2018, Schult *et al.*, 2018). On the other hand, TFE might be involved in global regulation in response to oxidative and heat shock stress, as the cellular protein levels were depleted under these stress conditions (Iqbal & Qureshi, 2010, Blombach *et al.*, 2015).

4.3.2. Regulatory transcription factors in *Sulfolobales*

The observation of extensive transcriptome-wide differential gene expression in response to stress conditions or shifts in nutritional conditions (**Sections 7 and 8**) indicates that transcription initiation is highly susceptible to regulation. It can be assumed that regulatory transcription factors (TFs) play an important role in this regard. In contrast to the eukaryotic

basal transcription machinery, archaeal regulatory TFs resemble bacterial regulators pointing to a shared ancestry (Aravind, 1999). One-component regulators are characterized by two domains: a N-terminal DNA-binding domain, with a wHTH or an HTH motif, and a C-terminal ligand-binding domain. The structural resemblance between TFs and wHTH-containing NAPs sometimes complicates their distinction, and dual-function DNA-binding proteins can be found within the entire spectrum between a specifically acting regulatory TF and a globally acting chromatin protein (Karr *et al.*, 2017, Dorman *et al.*, 2020). This is illustrated by the archaea-specific Lrs14 family of DNA-binding proteins, which is widespread in *Sulfolobales* and shown to bind DNA non-specifically and to regulate biofilm formation and motility in *S. acidocaldarius* (Orell *et al.*, 2013) (**Section 6.2.2**). Given the complete absence of typical bacterial two-component systems in *Crenarchaeota* including the *Sulfolobales* (Galperin *et al.*, 2018), these organisms are solely reliant on one-component regulators. Usually, TFs bind in the vicinity of the promoter elements of transcription units and interact with the different components of the PIC, thereby either repressing or activating transcription initiation (Peeters *et al.*, 2013) (**Table 2**). In some cases, a single TF can have a dual function, depending on the target gene or in a concentration-dependent manner, as has been shown for the *Sa. solfataricus* TF Ss-LrpB (Peeters *et al.*, 2013).

The functional understanding of the TFs in thermoacidophilic archaea is still limited and based on a relatively small number of case studies for individual TFs in model species, such as *Sa. solfataricus*, *Sa. islandicus* and *S. acidocaldarius* (**Table 2**). These TFs are involved in the regulation of various metabolic and physiological processes, such as motility, hetero- or autotrophic growth, metal resistance and detoxification mechanisms, typically in response to

interactions with small molecules, *e.g.* metabolites (**Sections 6.2.2, 7, and 8**). Unfortunately, a system-level approach for mapping TF-mediated gene regulatory networks in relation to common environmental stresses, similar to how it has been performed for the euryarchaeal model organism *Halobacterium salinarum* (Bonneau *et al.*, 2007), has not yet been undertaken for a thermoacidophilic archaeal species. An understanding of these networks would be valuable to gain insights into the physiology and stress adaptation of thermoacidophilic archaea and could be exploited for the engineering of metabolism for biotechnological purposes. Most of the characterized TFs in *Sulfolobales* belong to the dominant TF families, Lrp/AsnC and MarR, which together encompass more than 50% of all TFs in *Crenarchaeota* (Perez-Rueda *et al.*, 2018, Lemmens *et al.*, 2019). TFs belonging to the Lrp/AsnC family are responsive to amino acids or related small molecules and display either a global or specific regulatory function of central metabolic pathways (Vassart *et al.*, 2013, Liu *et al.*, 2014). BldR and BldR2 of *Sa. solfataricus* are prototypical MarR-family TFs involved in the detoxification of aromatic compounds (Di Fiore *et al.*, 2009, Fiorentino *et al.*, 2011). Finally, members of archaea-specific TF families are also found in *Sulfolobales*, such as the TrmB family that is typically involved in the regulation of sugar metabolism (Wagner *et al.*, 2014).

Intriguingly, archaeal genomes are predicted to harbor a lower fraction of TF-encoding genes as compared to bacterial genomes (Pérez-Rueda & Janga, 2010), and *Crenarchaeota* typically have even lower numbers of TFs than *Euryarchaeota* (Coulson *et al.*, 2007). These observations raise the question as to how thermoacidophilic *Crenarchaeota* are capable of efficiently regulating their transcriptome with a limited repertoire of TFs. While this might be partially explained by an extensive specialization to living in niche habitats, alternative

mechanisms are hypothesized to exist; for example, cross-interactions exist between paralogous TFs that lead to a combinatorial use of a limited set of regulators, as has been shown for members of the Lrp/AsnC family of TFs in *Sa. solfataricus* and *S. acidocaldarius* (Nguyen-Duc *et al.*, 2013, Liu *et al.*, 2016). Furthermore, additional layers of regulation might exist, for example at the post-transcriptional level (see below, Section 4.4), or by means of PTMs of TFs (see Section 5.4 and Figure 5.2). In this context, it is notable that phosphoproteomic studies have indicated the widespread occurrence of phosphorylation of TFs in *S. acidocaldarius* and *Sa. solfataricus* (Esser *et al.*, 2012, Reimann *et al.*, 2012). In *S. acidocaldarius*, phosphorylation has been shown to directly affect DNA binding of the Lrs14-type biofilm regulator AbfR1 (Li *et al.*, 2017), or ligand interaction in the case of the acyl-CoA-responsive TetR-family regulator FadR (Maklad *et al.*, 2020) (Section 5.4).

4.4. Translation and its regulation

4.4.1. Posttranscriptional regulatory mechanisms in *Sulfolobales*

An alternative explanation for the compensation of the rather limited repertoire of TFs in *Sulfolobales* is the existence of gene regulatory mechanisms at alternative levels of information processing, such as the posttranscriptional level (Lemmens *et al.*, 2019). RNA-based regulation is supported by a widespread occurrence of small non-coding RNAs (ncRNAs) in *Sa. solfataricus* and, to a lesser extent, in *S. acidocaldarius* (Tang *et al.*, 2005, Zago *et al.*, 2005, Wurtzel *et al.*, 2010, Cohen *et al.*, 2016). More than 300 ncRNAs were identified in *Sa. solfataricus*, 60% of which are *cis*-acting antisense transcripts (Wurtzel *et al.*, 2010). Possibly, these small RNAs assist in the stabilization of mRNA by RNA duplex formation,

which is relevant given the thermophilic lifestyle of the organism (Gomes-Filho & Randau, 2019). Nevertheless, given that most of these ncRNAs are conserved in closely related *Sa. islandicus* genomes (Reno *et al.*, 2009), they likely have functional roles and that these antisense ncRNAs regulate translation in a similar manner as RNA silencing mechanisms in eukaryotes. Besides the observation that they are overrepresented in coding regions of genes involved in ion transport and metabolism (Wurtzel *et al.*, 2010), the function of ncRNA-mediated regulation is still unclear as very few ncRNAs have been characterized thus far. A good example is RrrR in *S. acidocaldarius*, an antisense ncRNA that targets two mRNAs, including one that encodes a hypothetical membrane protein that was shown to influence biofilm formation (Orell *et al.*, 2018) (**Section 6.2.2**). In *Sa. solfataricus*, a small ncRNA has been shown to interact with the 3'-untranslated region (UTR) of its target mRNA in a phosphate-responsive gene regulatory process (Märtens *et al.*, 2013). This is a logical regulatory site given that most transcripts in thermoacidophilic *Crenarchaeota* are leaderless and lack a 5'-UTR (Brenneis *et al.*, 2007, Wurtzel *et al.*, 2010), which is the preferred site of small RNA-mediated posttranscriptional regulation in bacteria.

There is a huge variation in the small RNAs found in transcriptomes of *Sulfolobales*; these can have lengths as small as 20-25 nucleotides (nts) or up to 500 nts (Wurtzel *et al.*, 2010, Xu *et al.*, 2012). Besides the antisense ncRNAs that have a classic regulatory role, *Sa. solfataricus* harbors transposon-associated ncRNAs involved in transposition, CRISPR-associated small RNAs (**Section 3.2**), C/D box small nucleolar (sno) RNA that guide methylation sites in rRNAs and tRNAs (Zago *et al.*, 2005), and small RNAs associated with TTSs (TTSaRNAs) (Zaramela *et al.*, 2014). Different RNA-binding proteins act as chaperones of

ncRNAs, such as members of the L7As/L30 protein family and the Sm superfamily. These proteins are not only involved in the biogenesis and functioning of small RNAs, but also in other aspects of processing of rRNA, tRNA and mRNA species (Gomes-Filho & Randau, 2019).

4.4.2. Basal translation machinery in *Sulfolobales*

Like replication and transcription, the machinery of translation in archaea shows striking similarities to those in eukaryotes. Studies of translation initiation have been mainly performed in *Sa. solfataricus* as a model for thermoacidophilic archaea (La Teana *et al.*, 2013). Two different mechanisms have been discerned: 1) translation is initiated based on a canonical Shine-Dalgarno (SD)/anti-SD interaction analogous as in bacteria at internal cistrons of polycistronic mRNAs, or 2) direct pairing of the start codon with the anticodon of the initiator methionine-tRNA occurs at leaderless transcripts devoid of a SD sequence (Benelli *et al.*, 2003). The latter mechanism is the most prevalent one, given the observation that most monocistronic mRNAs and the proximal cistrons of polycistronic mRNAs in *Sulfolobales* are leaderless, lacking SD-harboring 5'-UTRs (Brenneis *et al.*, 2007, Wurtzel *et al.*, 2010). The translation initiation machinery has a complexity reminiscent of their eukaryotic counterpart and consists of multiple translation initiation factors (IFs), some of which have eukaryotic but no bacterial homologs (La Teana *et al.*, 2013). A crucial IF, especially for the translation initiation of leaderless transcripts, is aIF2, which is a heterotrimeric protein that forms a ternary complex with GTP and the methionine-loaded initiator tRNA, and binds the small ribosomal subunit. Other initiation factors are aIF1, aIF1a, aIF5b and aIF6. Despite the eukaryotic nature of the machinery, the operational

steps in the translation initiation process in *Sa. solfataricus* resemble those in bacteria (La Teana *et al.*, 2013).

5. Post-translational modification by reversible protein phosphorylation

All living cells, including thermoacidophiles, respond and adjust their cellular processes to a multitude of external and internal signals/cues. Information processing from gene to protein is also time and energy intensive. Therefore, post-translational-modifications (PTMs) are an efficient way to rapidly adjust protein function and coordinate the cellular response to changing needs. One of the best studied PTMs is the reversible protein phosphorylation mediated by protein kinases (PKs) and protein phosphatases (PPs) (Kennelly, 2003, Kennelly, 2014, Esser *et al.*, 2016, Papon & Stock, 2019). Phosphoproteins were first found in the Euryarchaeon *Halobacterium salinarum* (Spudich & Stoeckenius, 1980) but, despite this early report, little is still known about reversible phosphorylation in Archaea.

Two distinct phosphorylation systems exist. The two-component system (TCS) involves a histidine sensor kinase (HisK) and response regulator (RR); the covalent modification, i.e. addition of a phosphate group, takes place on histidine (His) and aspartate (Asp) residues, respectively (Galperin *et al.*, 2018). It was originally thought that TCS were specific for Bacteria, but it has since been shown that TCS occurs in all three domains of life (Loomis *et al.*, 1997, Kim & Forst, 2001, Schaller *et al.*, 2011). The first TCS reported in Archaea was CheA and CheY in the Euryarchaeon *Halobacterium salinarium* (Rudolph *et al.* 1995; Falke *et al.* 1997). In thermoacidophiles, TCS have only been identified in the *Thermoplasmata* within the *Euryarchaeota*, although their function is still unknown (Galperin *et al.*, 2018). In other archaeal

phyla, i.e. the *Crenarchaeota*, the *Korarchaeota* and the *Nanoarchaeota*, TCSs are largely absent and the organisms solely rely on canonical Hanks-type protein kinases for signal transduction (Hanks, 2003, Esser *et al.*, 2016, Galperin *et al.*, 2018). In these cases, autophosphorylation of the PK takes place on the amino acids Ser and Thr (eSTPKs) or tyrosine (Tyr). Often multiple PKs are interconnected, leading to a signal transduction cascade with a continuous hierarchical network structure. Hanks-type protein kinases can be broadly split into two groups, which have been well established in Archaea (Leonard *et al.*, 1998, Kennelly, 2014, Esser *et al.*, 2016, Hoffmann *et al.*, 2017): the conventional Hanks-type protein kinases (ePKs), and the non-canonical, atypical Hanks-type protein kinases (aPKs). The largest group, the ePKs, all share a conserved catalytic domain consisting of twelve subdomains (Stancik *et al.*, 2018). The other, smaller group, the aPKs, are distant members of the ePKs superfamily and share only some of the conserved subdomains (Esser *et al.*, 2016). Although Tyr phosphorylation has been demonstrated, so far only eSTPKs (phosphorylation on Ser/Thr residues) have been identified and characterized in Archaea, with the responsible tyrosine kinases yet unknown (Smith *et al.*, 1997, Kennelly, 2014).

The counterparts to PKs are PPs, which can remove the covalently linked phosphate residue, thereby making the process reversible. PPs can be classified into different subgroups, depending on their substrate specificity. In archaea, several phosphatases have been reported that differ in substrate specificity and cofactor requirement (Shi, 2009). Ser/Thr phosphatases (PPPs) and Mg^{2+} - or Mn^{2+} -dependent protein phosphatases (PPM) act on pSer and/or pThr, whereas protein tyrosine phosphatases (PTPs) are active on pTyr (Shi, 2009). Notably, dual-activity of the

PPM on pTyr and pSer/Thr was demonstrated *in vitro* in the thermoacidophile *Thermoplasma volcanium* (Dahche *et al.*, 2009).

Among the thermoacidophilic archaea, the protein phosphorylation pathways of the *Sulfolobales* are the best characterized. When considering protein phosphorylation in the *Sulfolobales*, it is important to know which PKs and PPs have been characterized so far, which proteins are targeted, and, most importantly, what is the physiological and cellular impact of reversible protein phosphorylation. First, homology searches (BlastXP) were performed based on the PKs and PPs identified in *S. acidocaldarius* and *Sa. solfataricus* (Leng *et al.*, 1995, Shi *et al.*, 1998, Lower *et al.*, 2000, Lower & Kennelly, 2002, Lower & Kennelly, 2003, Lower *et al.*, 2004, Kennelly, 2014, Ray *et al.*, 2015), *Sa. islandicus* (Huang *et al.*, 2017, Huang *et al.*, 2019) and *Sulf. tokodaii* (Wang *et al.*, 2010) (**Figure 5**). In surveying the *Sulfolobales* genomes, *Sulf. tokodaii* has 15 PKs, followed by *S. acidocaldarius* with 13 PKs, and *Sa. islandicus* and *Sa. solfataricus* both with 11 PKs. All strains harbor one typical ePK with an additional tetratricopeptide repeat (TPR-motif) that is known to mediate protein-protein interactions. In *Sa. islandicus*, the ePK was shown to act as a master PK, phosphorylating other PKs *in vivo*. Also, the homolog in *S. acidocaldarius* (ArnC) phosphorylates a variety of different target enzymes (Hoffmann *et al.*, 2017, Huang *et al.*, 2017, Knüppel *et al.*, 2018, Maklad *et al.*, 2020). Several PKs have an additional trans-membrane domain, implicating membrane-bound localization. In all *Sulfolobales* genomes, two Rio-like PKs (aPKs) are present. Rio kinases are an ancient conserved family that can be found in all three domains of life and are known to play a role in ribosome biogenesis (Esser & Siebers, 2013, LaRonde, 2014). All *Sulfolobales* are known to harbor a set of Rio B and Rio 2 PKs, except *S. acidocaldarius*, where Rio 1 and Rio 2

can be found (Esser & Siebers, 2013). Rio 1 and Rio 2 are quite similar, except that Rio 2 possesses an additional winged helix-turn-helix domain (wHTH), a structural motif enabling binding of DNA. Recently, further evidence for the role that Rio kinases play in the synthesis of the ribosomal small subunit in archaea was uncovered (Knüppel *et al.*, 2018). Furthermore, one piD261-like aPK, of which SSOPK5 is the only one studied in the *Sulfolobales*, one ABC1-like and one AQ578 aPK, both putative, can be found in *Sa. solfataricus*, *Sulf. tokodaii*, *Sa. islandicus* and *S. acidocaldarius* (**Figure 5**). Notably, in all of these genomes, only two PPs were found: one annotated as Tyr-phosphatase and one as Ser/Thr-phosphatase.

5.1. Protein phosphorylation in *S. solfataricus*

The first evidence of protein phosphorylation in the *Sulfolobales* was reported in *Sa. solfataricus* (Kennelly *et al.*, 1993). PP activity was detected in soluble extracts of *Sa. solfataricus* using ³²P-casein as the substrate and allowed the isolation of the PP from the soluble fraction. Since no activity could be detected with p-Tyr labeled substrates, it was concluded that the PP was a Ser/Thr phosphatase. The amino acid sequence of the SSO-PP was similar to the eukaryotic PP1/2A/2B superfamily (Leng *et al.*, 1995).

Sa. solfataricus was also the first *Sulfolobales* species in which PKs were characterized and now include four ePKs and one aPK (SSO-PK1, SSo-PK2, SSO-PK3, SSO-PK4 and SSO-PK5). Their activity on “artificial” substrates, such as histone, myelin basic protein, p53, casein, reduced carboxyamidomethylated, and maleylated lysozyme, in addition to their cofactor dependence, and inhibition by typical ePK inhibitors, have been analyzed (Lower *et al.*, 2000, Lower & Kennelly, 2002, Lower & Kennelly, 2003, Lower *et al.*, 2004, Haile & Kennelly, 2011,

Kennelly, 2014, Ray *et al.*, 2015, Esser *et al.*, 2016). SSO-PK4 (encoded by SSO3182), like its eukaryotic homologs, was shown to phosphorylate the *Sa. solfataricus* eukaryotic translation initiation factor (eIF2 α) homolog, aIF2 α , *in vitro*, but not on the conserved phosphorylation sites known for eIF2 α (Ray *et al.*, 2015). Inhibition was observed in the presence of 3',5'-cAMP *in vitro*, whereas a concentration dependent activation occurred in the presence of oxidized CoA, an indicator of oxidative stress in archaea. Additionally, the aPK of the pID261/Bud32 kinase family (SSO-PK5) was characterized (Haile & Kennelly, 2011). SSO-PK5 (encoded by SSO0433) phosphorylated artificial substrates, like p53 and casein, on Ser residues. Autophosphorylation of SSO-PK5 was shown to take place on both Ser and Thr residues and activation was observed in the presence of ADP-ribose (Haile & Kennelly, 2011). Phosphohexomutase (SSO0207), first identified in tryptic digests, was shown to be phosphorylated. Site-directed mutagenesis (S309D) was used to mimic the presence of a phosphoryl group, which drastically decreased the V_{max} -value of the enzyme. Therefore, it was suggested that protein phosphorylation is used *in vivo* to regulate the phosphohexomutase activity, but the corresponding PKs are yet unknown (Figure 6) (Ray *et al.*, 2005). This appears to be the first report on regulation of central metabolic enzymes by reversible protein phosphorylation in not only the *Sulfolobales*, but Archaea in general.

In 2012, Esser *et al.* performed the first phosphoproteome study in *Sa. solfataricus* where they compared the phosphoproteome of cells grown on D-glucose vs cells grown on tryptone (Esser *et al.*, 2012, Dopson, 2016). Using a precursor acquisition independent from ion count (PAcIFIC) approach, 540 phosphoproteins in 21 out of 26 arCOGs were found, highlighting the importance of regulation by reversible protein phosphorylation in *Sa. solfataricus*. Interestingly,

the phosphorylation profile was dependent on the respective carbon source and a high amount of Tyr phosphorylation (Ser/Thr/Tyr ratio of 26/21/54%) was detected. This was rather unexpected since, so far, no PKs with Tyr-phosphorylation activity had been reported. Due to the significant changes in the phosphoproteome in response to carbon source, it was concluded that reversible protein phosphorylation plays a major role in the regulation of central carbon metabolism (CCM) in *Sa. solfataricus* (Dopson, 2016).

5.2. Protein phosphorylation in *Sulf. tokodaii*

Detailed characterization of the ePK STK_15650, which comprises all signatures of a canonical Hanks type kinase in *Sulf. tokodaii*, showed that the important catalytic residues are all located in, or close to, the major functional domains of eSTKs (Wang *et al.*, 2010). It was the first time in Archaea that the interaction between an ePK and a target protein was characterized (Wang *et al.*, 2010). The interaction partner that is phosphorylated is the forkhead-associated (FHA) domain-containing protein (STK_00829). FHA domains are known to act as phosphorylation-dependent protein–protein interaction modules that can bind to pThr residues in their targets (England *et al.*, 2009). Additionally, specific interactions of both proteins were demonstrated *in vivo*. Important residues for the protein-protein interaction were identified. It was proposed that STK_00829 might be a transcriptional regulator and, therefore, phosphorylation might play a role in transcriptional regulation in Archaea (**Figure 6**) (Wang *et al.*, 2010, Duan & He, 2011). Subsequently, the homolog of STK_00829 ArnA was shown to be a repressor of archaellum expression and part of the archaellum regulatory network in *S. acidocaldarius* (See **Table 2**) (Reimann *et al.*, 2012).

5.3. Protein phosphorylation in *Sa. islandicus*

Autophosphorylation and cross-phosphorylation activities of the eleven PKs (three ePKs and eight aPKs) from *S. islandicus* REY15A revealed insights into the hierarchy of regulatory networks (Huang *et al.*, 2017). The seven PKs (SiRe_0101KD, SiRe_0171, SiRe_0181, SiRe_1570, SiRe_1810, SiRe_2030, and SiRe_2056KD) exhibited autophosphorylation activities, with the highest activity shown for the ePK SiRe_2056KD. Dephosphorylation assays revealed that autophosphorylation mainly proceeds on Ser/Thr residues (Huang *et al.*, 2017). To gain more insight into the cross-phosphorylation and cross-talk among the PKs, inactive ePKs were generated. Among the ePKs, SiRe_2030 and the truncated SiRe_2056KD were most active on phosphorylation of the other PKs. Next, to address the importance of ePKs, the effect of PK overexpression was evaluated. Only for SiRe_1531 and SiRe_2056 was there an obvious phenotype detectable, i.e. growth retardation (Huang *et al.*, 2017). Therefore, SiRe_2056 and SiRe_2030 are the master PKs, and SiRe_0101 is an accessory kinase at the apex of the phosphorylation hierarchy in *Sa. islandicus* that transduced the signal towards the other substrate kinases. However, more physiological information is needed to elucidate the complex signaling pathways in *Sa. islandicus* (Huang *et al.*, 2017).

With regards to targets in *Sa. islandicus* REY15A, phosphorylation of a conserved Holliday junction resolvase (Hjc), an enzyme employed in homologous recombination repair (HRR), by the PKs SiRe_0171 (Rio 1-like aPK), SiRe_2030 and SiRe_2056 (ePKs) was investigated (Huang *et al.*, 2019). These PKs phosphorylated different residues *in vitro* and the analysis of the respective phosphorylation-mimic mutants revealed that the phosphorylation of

S34 (phosphorylated by SiRe_0171) and S9 (phosphorylated by SiRe_2030) have a strong impact on Hjc activity. To elucidate the *in vivo* significance of Hjc protein phosphorylation, strains expressing the different phosphorylation-mimic mutants were tested for their sensitivity towards DNA damaging agents. The strain expressing S34E (mimicking phosphorylated Hjc) was less sensitive towards high doses of DNA-damaging agents (i.e. UV or cisplatin) indicating a higher DNA repair capability. In addition, deletion of the respective Rio 1 homolog SiRe_0171 (and thus preventing phosphorylation of the S34 residue) resulted in a strain with higher sensitivity towards DNA damaging agents, thus indicating that phosphorylation of S34 in Hjc enhances the DNA repair capability (**Figure 6**).

5.4. Protein phosphorylation in *S. acidocaldarius*

Protein phosphorylation in *S. acidocaldarius* was initially identified when several proteins were found to be phosphorylated in the presence of [γ -³²]ATP *in vivo* (Skórko, 1984). Three PKs, ArnC, ArnD and ArnS, are involved in the regulation of the best studied complex signal transduction system in archaea, the archaeallum regulatory network (Arn) (**Figure 6**). The genes encoding for the archaeal motility structure, the archaeallum, are arranged in an operon consisting of seven genes with two promoters. The main promoter upstream of the gene *arlB* (f. *flaB*) was induced upon starvation, whereas the second promoter upstream of *arlX* (f. *flaX*) revealed weak constitutive activity (Lassak *et al.*, 2012). The transcriptional regulators, ArnA and ArnB, repress *arlB* (f. *flaB*) expression. *arnA* harbors a zincfinger (ZnF) and a Forkhead-associated (FHA) domain and is a homolog to the previously identified regulator STK_00829 in *Sulf. tokodaii*, while ArnB possesses a vonWillebrand type A domain (vWA). Deletion of either

one of the two genes showed a hypermotile phenotype *in vivo* and protein levels of the archaellin ArlB (f. FlaB) were strongly enhanced compared to the wild type strain (Reimann *et al.*, 2012). The ePKs ArnC (Saci_1193), ArnD (Saci_1694), and ArnS (Saci_1181), and the Ser/Thr PP (Saci_0884) (Saci_PP2A), regulate the archaellum at the post-translational level by reversible phosphorylation. ArnC is able to phosphorylate both regulators ArnA and ArnB and deletion of the ePK resulted in reduced motility *in vivo*. In contrast, ArnD is only able to phosphorylate ArnB and its deletion resulted in a hypermotile phenotype. These divergent effects on motility suggested that the two PKs have different roles in the regulatory network. Both regulators were dephosphorylated by the addition of the Saci_PP2A (Reimann *et al.*, 2012). Since ArnA and ArnB interact with each other in a phosphorylation-dependent manner, protein phosphorylation seems to be the key for their regulatory function in the Arn (Reimann *et al.*, 2012). The deletion of the starvation induced ePK, ArnS, which is also located close to the archaellum operon, resulted in reduced motility, indicating an essential role of this ePK as well (Haurat *et al.*, 2017). Finally, the deletion of the Ser/Thr PP PPP2A also revealed a hypermotile phenotype, suggesting a negative regulation, although the respective relay mechanism and target protein(s) are still unknown. Another player of the Arn is the transcriptional regulator of the Lrs14 family, the archaeal biofilm regulator 1 (AbfR1) (Orell *et al.*, 2013, Liu *et al.*, 2017). AbfR1 binds to its own promoter, as well as the *arlB*, *arlX* (f. *flaB*, *flaX*, respectively) promoter, and has non-specific DNA binding activity suggesting a general chromatin structuring function. In the non-phosphorylated state, it binds to DNA and increases motility by expression of the archaellum operon and negatively regulates biofilm formation by decreasing the production of extracellular polymeric substance (EPS). Phosphorylation of two residues, S87 and Y84, in the wHTH domain

impair stable protein-DNA contacts *in vivo* (Li *et al.*, 2017). Thus, AbfR1 phosphorylation promotes the transition from a motile, planktonic growth to a sessile lifestyle in *S. acidocaldarius* (**Figure 6**). The PK(s) involved in phosphorylation of AbfR1 are still unknown.

The two Rio kinases, Rio 1 and Rio 2, of *S. acidocaldarius* exhibited ATP hydrolysis activity in a concentration-dependent manner (Knüppel *et al.*, 2018) and are non-essential in *S. acidocaldarius*, as demonstrated by creation of single deletion mutants (Hoffmann *et al.*, 2017). Both PPs in *S. acidocaldarius* were characterized in detail and the physiological role was addressed by comparison of the phosphoproteomes of the parent strain MW001 and the two PPs deletion mutants Δ Saci_ptp, Δ Saci_pp2a (Reimann *et al.*, 2013). Saci-PTP is a dual-specific phosphatase (active with pSer/pThr and pTyr), whereas Saci-PP2A exhibited specific pSer/pThr activity and could be inhibited by okadaic acid (Reimann *et al.*, 2013). The study revealed major differences in the phosphorylation and gene expression patterns of the two deletion strains, suggesting important roles for both phosphatases in signal transduction pathways (Reimann *et al.*, 2013). Interestingly, the ratio of pSer/pThr/pTyr, varied slightly between the different strains and revealed a high amount of pTyr, also seen in *Sa. solfataricus* (Reimann *et al.*, 2013, Dopson, 2016).

A very recent study highlighted the physiological effect of protein phosphorylation on central metabolism (**Figure 6**) (Maklad *et al.*, 2020). The transcription factor, FadR_{Sa}, represses a 30-kb gene cluster encoding enzymes involved in the lipid/fatty acid degradation in *S. acidocaldarius* (Wang *et al.*, 2019). Acyl-CoAs act as inducers leading to DNA dissociation of FadR_{Sa} and thus transcription of the gene cluster. FadR_{Sa} repressed the gene cluster by only four

binding sites; further studies revealed that FadR_{Sa} might have an additional function in organization of local chromatin architecture, as indicated by the interplay of FadR_{Sa} with the chromosome structuring factor coalescing (Takemata *et al.*, 2019). In previous phosphoproteome studies, FadR_{Sa} was found to be phosphorylated on three different residues (Y¹³³, T¹³⁴ and T¹³⁵) (Reimann *et al.*, 2013). Notably, all three residues are located within the binding pocket of acyl-CoA. *In vitro* phosphorylation studies with different PKs confirmed phosphorylation of FadR by ArnC and Saci_1041, whereas ArnD and Saci_0965 were not active. The constructed triple phosphomimetic mutant FadR_{Sa} (Y133D-T134E-T135E) was less sensitive for acyl-CoA. This implies that phosphorylation of FadR_{Sa} may act as an additional control mechanism that keeps the gene cluster in a condensed state and allows transcriptional repression of lipid/fatty acid metabolism in the presence of acyl-CoAs in *S. acidocaldarius* (Maklad *et al.*, 2020).

6. Cell Cycle and Modes of Growth

6.1. The planktonic lifestyle

6.1.1. The archaellum and other surface structures

Archaea form a multitude of different surface structures that play important roles in diverse processes, such as motility, adhesion, biofilm formation, DNA transfer, and probably many more (Chaudhury *et al.*, 2018). Many of these structures are type IV pili or type IV pilus-like, e.g. the archaellum, the motility structure of archaea (Albers & Pohlschröder, 2009, Jarrell & Albers, 2012, Makarova *et al.*, 2016). Archaeal type IV pili are similar to those of bacteria in that both pilin precursor proteins exhibit a class III signal peptide at their N-terminus, which is processed by a type IV prepilin signal peptidase (PibD in *Sulfolobales* and *Haloarchaea*) (Albers

et al., 2003, Tripepi *et al.*, 2010). Only after removal of the signal peptide can the pilins be assembled into the pilus filament by the assembly system which is formed by an integral membrane protein and an ATPase (**Figure 7**). The basic assembly mechanism of the archaellum resembles that of the type IV pilus. However, the associated accessory proteins (ArlFGH) enable the archaellum filament to rotate and therefore propel the cells forward (Jarrell & Albers, 2012, Albers & Jarrell, 2015).

Among the *Sulfolobales*, *S. acidocaldarius* is the best studied with respect to surface structures as it exhibits a variety of them (**Figure 7 EM, Model**): the Ups pili (UV-induced pili), the Aap pili (archaeal adhesive pili), the threads, and the archaellum (Fröls *et al.*, 2008, Henche *et al.*, 2012, Henche *et al.*, 2012, Tsai *et al.*, 2020). The threads, the only non-type IV pilus structure on the *S. acidocaldarius* surface, are formed by unknown proteins, but are used as binding sites for viruses (see **Section 3**) (Hartman *et al.*, 2019). The Ups pili are assembled after DNA double strand breakages and lead to species-specific cell-aggregation. During aggregation DNA is exchanged and subsequently used for DNA repair by homologous recombination (Fröls *et al.*, van Wolferen *et al.*, 2020). The species specificity is ensured by binding of the pilin subunits to the N-glycan trees on the S-layer protein that differ among species like *Sa. solfataricus*, *Sulf. tokodaii* and *S. acidocaldarius* (see **Section 6.3.1**) (van Wolferen *et al.*, 2020). The Aap pili are important for adhesion of *S. acidocaldarius* to surfaces and biofilm formation (see **Section 6.2**).

6.2. Biofilms

Biofilms are the most common form of microbial life. They consist of cells that are attached to a surface (which may or may not serve as a source of nutrients for the microorganism), embedded in extracellular polymeric substances (EPS) that form a matrix produced by the microbial population. This matrix consists of different types of polymers: lipids, polysaccharides, extracellular nucleic acids and proteins (Fröls, 2013, van Wolferen *et al.*, 2018). Biofilms can also be formed at air-liquid and liquid-solid interfaces.

Biofilms are dynamic communities where cells can leave the biofilm structure and swim free in a planktonic lifestyle or attach to another surface to colonize it (Koechler *et al.*, 2015). The first biofilm described for archaea was that of the extremely thermophilic euryarchaeon *Thermococcus litoralis* (Rinker & Kelly, 1996). Similarly to bacterial biofilms, biofilms protect archaea against diverse kinds of stress, such as changes in pH, desiccation, UV radiation, high salt and metal concentrations (Laplagia & Hartzell, 1997, Fröls, 2013). Biofilms also provide an advantage to cells because they form a microenvironment where they share nutrients, water channels, etc. (Petrova & Sauer, 2012). In *Sulfolobales* species, stress factors like changes in pH and temperature induce biofilm formation (Koerdt *et al.*, 2010), suggesting a role in protection against unfavorable conditions. A method to grow *Sulfolobales* biofilms in microtiter plates has been standardized (Koerdt *et al.*, 2010, Koerdt *et al.*, 2011). The method consists of growing the species in Brock media pH 3 in microtiter plates covered with a gas permeable membrane at ~75°C for 3 to 6 days without agitation. Using this method, biofilm from three *Sulfolobales* species were studied (*Sa. solfataricus*, *S. acidocaldarius*, and *Sulf. tokodaii*). Also, some *Sulfolobales* have been studied in the context of acid mine drainage (AMD) biofilms, for

example *Sulfura. metallicus* (Zhang *et al.*, 2015, Zhang *et al.*, 2019) and *Acidianus* spp. (Zhang *et al.*, 2015) on elemental sulfur and pyrite.

6.2.1. Biofilm formation process

There are three stages in biofilm formation: attachment of cells to the surface, formation of microcolonies and biofilm maturation, and finally, dispersion. In each of these stages, different structures are involved, and diverse morphologies had been described for *Sulfolobales* species (Summarized in **Table 3**). The first stage for biofilm formation is the attachment of cells to a surface. Different cell surface structures play a role in this process, such as type IV pili, in the later stages, the archaellum, and depending on the archaeal species, hami and fimbria are involved (van Wolferen *et al.*, 2018). *Sa. solfataricus* mutants that lack ArlB (formerly FlaB), the structural component of the archaellum, or either one of the Ups pili components, are defective in adhesion to surfaces like glass or pyrite (Zolghadr *et al.*, 2010). Likewise, the deletion of an Ups pili lead to less biofilm formation after 3 days of growth (Koerdt *et al.*, 2010). The Aap pili are absent in *Sa. solfataricus* but present in *S. acidocaldarius* and, along with the Ups pili, are important for surface adhesion (Henché *et al.*, 2012). However, the archaellum does not play a role in adhesion in *S. acidocaldarius*, but seems to be involved in biofilm maturation (Henché *et al.*, 2012).

Once attached to the surface, cells start to divide and produce EPS. In *S. acidocaldarius*, microcolony formation was seen after 36 h of inoculation. In *Sa. solfataricus*, *S. acidocaldarius*, and *Sulfura. tokodaii* during the first 3 days of growth, the secreted exopolysaccharides residues were mainly glucose and mannose, while from day five onward galactose and N-acetyl

glucosamine were predominant (Koerdt *et al.*, 2010). The complete path of exopolysaccharides biosynthesis is still unknown. However, an α -mannosidase encoded by SSO3006 in *Sa. solfataricus* affects EPS production (Koerdt *et al.*, 2012). The mutant strain *Sa. solfataricus* PBL2025, which lacks a 50 kB region including SSO3006, produced more EPS than the wild type strain and reverted back to the wild type phenotype when complemented with SSO3006 (Koerdt *et al.*, 2012). In *Sulfura. metallicus* DSM 6482(T), biofilm can form on elemental sulfur with microcolonies and cells clusters of up to 100 μm in diameter. This biofilm was rich in proteins and nucleic acids, in contrast to capsular EPS from planktonic cells, where the EPS mainly contained carbohydrates and proteins (Zhang *et al.*, 2015).

Mature biofilms formed by *S. acidocaldarius*, *Sa. solfataricus* and *Sulf. tokodaii* are morphologically different from each other. *Sa. solfataricus* forms biofilms with a carpet-like structure, with 20–30 μm thick covering the whole surface but with a low density of cells. *Sulf. tokodaii* forms biofilms of 25–35 μm thick and also exhibits a carpet-like structure, but with high cell density and, occasionally, cell aggregates. Finally, *S. acidocaldarius* forms 25–35 μm thick biofilms that contained a high density of cells and large aggregates, forming towering structures above the surface of attached cells (Koerdt *et al.*, 2010). Deletion mutant studies in *S. acidocaldarius* showed that the Ups pili and the Aap pili have profound impacts on the morphology of the biofilms (Henche *et al.*, 2012). While deletion of the Aap pilus led to a dense biofilm which was thinner than the wild type, the deletion of the Ups pili led to large aggregates of cells within a “fluffy” biofilm characterized by a single dense layer of cells at the surface. Clearly, both pili play an essential role in the optimal layering and distancing of the cells in the wild type biofilm (Henche *et al.*, 2012). Cell-cell connections were also seen extensively for *S.*

acidocaldarius and *Sulf. tokodaii*, and to a lesser degree in *Sa. solfataricus* (Koerdt *et al.*, 2010). Besides exopolysaccharides, proteins and extracellular (eDNA) can also be found in *S. acidocaldarius* biofilms. Composition analyses revealed several enzyme activities in EPS extracts, but most of them were cytoplasmic proteins (Jachlewski *et al.*, 2015), probably derived from cell lyses as these species secrete only small amounts of proteins (Ellen *et al.*, 2010).

Biofilm maturation was followed for several days in *S. acidocaldarius* (**Figure 8**, biofilm). From days 3-6, an increase in cell density was observed, and in days 7-8 dispersion of the cells was seen (Hence *et al.*, 2012, Koerdt *et al.*, 2012). In a $\Delta arlJ$ mutant, however, cell dispersion from the biofilm was decreased, showing that the archaellum is important for cells to leave the biofilm (Hence *et al.*, 2012). It is unknown what triggers dispersion of cells, but this process is important for colonizing other sites along the surface.

6.2.2. Regulation of Biofilm Processes

The biofilm and planktonic lifestyles differ, as inferred from transcriptomic and proteomic studies comparing both populations (Koerdt *et al.*, 2011), with the change from one to another depending on different regulators. In bacteria, it is known that secondary messenger and quorum sensing mechanisms are important for the regulation of biofilm formation, however these have not been described for *Sulfolobales*. Nonetheless, high throughput proteomics and transcriptomics allowed for a first glance at the factors that might be involved in biofilm formation in three *Sulfolobales* species and a species-specific response was found (Koerdt *et al.*, 2011). Among the common differentially regulated proteins were the archaea-specific Lrs14-like regulators (Leucine-responsive Regulator of *Sulfolobus*) (Koerdt *et al.*, 2011). Later, deletion

mutants confirmed that some of these Lrs14-like proteins are involved in biofilm regulation in *S. acidocaldarius* (see **Section 4.3.2, Table 2**). Knock out mutants for *Saci_1223* were impaired in biofilm formation, suggesting that this regulator promotes biofilm formation (Orell *et al.*, 2013). Furthermore, the deletion mutant of *Saci_0446* produced more EPS than the wild type, and also showed a non-motile phenotype, where expression levels of the archaellum were down-regulated and expression levels of *Aap* were increased. Therefore, it is thought to act as repressor of biofilm formation and named *AbfR1* for Archaeal Biofilm Regulator 1 (see Signal Transduction **Section 5, Figure 5**) (Orell *et al.*, 2013). *AbfR1* functions by stimulating motility through the induction of the archaellum by binding to the *arlB* promoter region, and repressing EPS production. Finally, detailed studies have shown that *AbfR1* is also regulated by phosphorylation through unknown mechanisms, and it cannot bind DNA when phosphorylated (Li *et al.*, 2017).

Some non-coding RNAs (ncRNA) also regulate biofilm formation. Sequencing of ncRNAs expressed in planktonic and biofilm cells was performed, and 29 ncRNA were differentially regulated in the latter (Orell *et al.*, 2018). One in particular, ncRNA239, named *RrrR* (RNase resistant RNA), was abundant in planktonic cells but further up-regulated in the biofilm. Moreover, deletion of this ncRNA led to impairment of biofilm formation. *RrrR* is a double stranded ncRNA, located in the intergenic region between *Saci_1004* and *Saci_1005*. The sense transcript of this RNA interacts with RNA-binding Lsm proteins, and its antisense RNA binds two mRNAs. The antisense transcript of this *RrrR* seems to stabilize the sense transcript (Orell *et al.*, 2018).

Polyphosphates (PolyP) are polymers of orthophosphate with roles in many cellular functions, including bacterial biofilm formation and related phenomena (Rashid *et al.*, Shi *et al.*, 2004, Grillo-Puertas *et al.*, 2012, Drozd *et al.*, 2014, Albi & Serrano, 2016). In *E. coli*, PolyP is involved in biofilm formation by triggering type II autoinducers (AI-II) synthesis in the stationary phase of growth through PolyP degradation (Grillo-Puertas *et al.*, 2012). In archaea, experiments in *Sa. solfataricus* and *S. acidocaldarius* demonstrated that this polymer is also involved in biofilm formation, adhesion, and motility (unpublished results), although the mechanism is still unclear.

6.3. Cell envelope of *Sulfolobales*

6.3.1. S-layer

The architecture of archaeal cell envelopes can be very diverse (Albers & Meyer, 2011, Klingl *et al.*, 2019). However, studied *Sulfolobales* species mainly have a cytoplasmic membrane surrounded by a proteinaceous coat, called the S-layer. In contrast to other archaea, which also have an S-layer as the main cell wall component, most *Sulfolobales* have two proteins that form the S-layer: SlaB, which is the membrane anchor, and SlaA forming the outer layer on top of the cell (Grogan, 1996, Veith *et al.*, 2009). As S-layers are ordered in 2D lattices, they are excellent targets for structural studies. In 1982, Amos and colleagues used electron microscopy to study the structure of the *S. acidocaldarius* S-layer (Taylor *et al.*, 1982) and related studies then showed that the architecture of the S-layers is species-specific (Prüschenk *et al.*, 1987). The *S. acidocaldarius* S-layer was found to be arranged in a conserved lattice with p3 symmetry, with 4.5 nm triangular and 8 nm hexagonal pores (Taylor *et al.*, 1982). By differential

solubilization, the SlaB subunits could be detached from the SlaA lattice, allowing cryoEM to pinpoint the placement of both subunits in the S-layer lattice (**Figure 9**) (Gambelli *et al.*, 2019).

As the only cell wall component, the S-layer provides stability to the cell and it has long been assumed that the S-layer is an essential component for *Sulfolobales* cells. However, a transposon library screen in *Sa. islandicus* indicated that both S-layer proteins can be deleted (Zhang *et al.*, 2018). When SlaB was deleted in *Sa. islandicus*, the cells were still able to assemble partial SlaA containing S-layer lattices. However, cells lacking SlaA were not only deformed and sometimes very large, but they also had an aberrant number of chromosomes, indicating that the coordinated assembly of the S-layer is important for cell division (Zhang *et al.*, 2019). Furthermore, the S-layer in *Sulfolobales* plays a role in anchoring other surface structures, such as the bindosome, involved in sugar binding and uptake (Zolghadr *et al.* 2011), and the archaeallum, where it is absolutely essential for torque generation (Tsai *et al.* 2020). The S-layer also acts as a sieve and as a surface for recognition that is conveyed by extensive N-glycosylation (discussed below).

6.3.2. Glycosylation

Many of the extracellular proteins of Archaea are glycosylated (this can be either O-glycosylation or N-glycosylation). Whereas we know very little about how proteins are O-glycosylated, the N-glycosylation pathway has been deciphered in halophiles, methanogens and *S. acidocaldarius* (Jarrell *et al.*, 2014). Many archaeal extracellular proteins have a so-called ST-linker, which is a stretch of many serine (Ser) and threonine (Thr) residues in a row (Albers *et al.*, 2004). In the *Halobacteria* S-layer, these residues were found to be O-glycosylated

(Lechner & Sumper, 1987, Sumper *et al.*, 1990) and, thus, it is thought that other extracellular proteins, like SlaB or the sugar binding proteins that also have these ST-linkers, are also O-glycosylated.

A number of proteins have been reported to be glycosylated in the *Sulfolobales*: the S-layer proteins (Peyfoon *et al.*, 2010), cytochromes (Zähringer *et al.*, 2000), sugar- and peptide binding proteins (Albers *et al.*, 1999, Elferink *et al.*, 2001, Gogliettino *et al.*, 2010), pilins (Wang *et al.*), the archaellum (Meyer *et al.*, 2015), and many hypothetical proteins (Palmieri *et al.*, 2013). Whereas proteins in other archaea have different glycans or the composition of the glycan changes due to environmental conditions, *Sulfolobales* have only one kind of glycan decorating all glycosylated proteins in one species. All determined N-glycans of *Sulfolobales* are in their basic structure quite similar to the eukaryotic N-glycan, as the glycans always starts with two N-acetylglucosamines and are branched (**Figure 10**) (Zähringer *et al.*, 2000, Palmieri *et al.*, 2013, van Wolferen *et al.*, 2020). Interestingly, they all contain a sulfated saccharide called 6-sulfoquinovose, a sugar otherwise only found in chloroplast and membranes of photosynthetic bacteria (Meyer *et al.*, 2011). However, except for this basic setup, the N-glycans of the different *Sulfolobus* species are diverse, as different numbers of mannose and glucose sugars are added to create a species-specific sugar labeling of extracellular proteins (see **Figure 10**). This is used for species-specific recognition during DNA exchange (Fröls *et al.*, 2008). During the pilin-dependent DNA exchange observed after double strand DNA breakages, *Sulfolobales* exchange homologous DNA, by a sugar specific binding of the pilin subunit to the S-layer of the aggregating cells (van Wolferen *et al.*, 2020). Although the differences are minor between the N-glycans of *Sa. solfataricus*, *Sulf. tokodaii* and *S. acidocaldarius* (see **Figure 10**), this is enough to

ensure that aggregation and subsequent DNA exchange only happens with the same species (van Wolferen *et al.*, 2020).

In contrast to methanogens and halophiles (Jarrell *et al.*, 2014), N-glycosylation seems to be essential in *Sa. islandicus* and *S. acidocaldarius*, as the enzyme attaching the N-glycan to the modified protein, the oligosaccharyl transferase AglB, cannot be deleted (Meyer & Albers, 2014, Zhang *et al.*, 2018). N-glycosylation is initiated by the assembly of the hexasaccharide N-glycan on short dolichol pyrophosphate carriers in *S. acidocaldarius* by AglH (Guan *et al.*, 2016, Meyer *et al.*, 2017). Deletion of the glycosyltransferase *agl16* led to a removal of a terminal hexose from the N-glycan, whereas deletion of *agl3*, the sulfoquinovose synthase, reduced the N-glycan to a trisaccharide (Meyer *et al.*, 2011, Meyer *et al.*, 2013). Both mutants had difficulties in adjusting to growth in media with elevated salt concentration. The N-glycosylation pathway of *S. acidocaldarius* is depicted in **Figure 10**.

6.4. *Sulfolobus* cell division

The most investigated archaeal cell division system is the ESCRT-III-based system in *Sulfolobales*. The proteins involved in cell division (Cdv) are organized in two groups. One group comprises the proteins CdvA, -B and -C organized in a gene cluster, while the other group consists of three CdvB paralogs located at different positions in the genome (Lindas *et al.*, 2008, Samson *et al.*, 2008, Samson *et al.*, 2011). CdvA is the only protein of these two groups that does not share homologies to the eukaryotic ESCRT-system or any known protein family. CdvA can bind membranes and polymerizes into helical filaments on the outside of liposomes composed of tetraether lipids isolated from *S. acidocaldarius*. Interestingly, the presence of lipids was

necessary for CdvA polymerization *in vitro* (Dobro *et al.*, 2013). Transcriptional analysis of the *cdv* operon showed that *cdvA* is up-regulated around 30 minutes before the *cdvB* genes are transcribed. Immunofluorescence with antibodies raised against CdvA displayed a ring-like localization pattern at mid-cell, but also outside the mid-cell region, before the genome was segregated (Lindas *et al.*, 2008, Samson *et al.*, 2011). These results suggest that CdvA is the earliest cell division protein localizing at the future division plane. However, it is unknown how CdvA is positioned. Remarkably, CdvA from *Metallosphaera sedula*, that was heterologously expressed and purified from *E. coli*, formed extended double-helical filaments which strongly interacted with the DNA of *E. coli* (Moriscot *et al.*, 2011). This suggests that CdvA participates in the chromosome segregation processes or functions as an inverse nucleoid-occlusion type localization mechanism for the archaeal divisome (Caspi & Dekker, 2018). However, the exact function of its DNA binding remains unknown. CdvA can recruit CdvB to preformed liposomes by direct interaction, resulting in extensively deformed vesicles. The β-strand forming C-terminal ESCRT-III-binding (E3B) peptide of CdvA interacts with the C-terminal incomplete winged-helix (wH) domain of CdvB. As such, the E3B peptide complements the “broken” part of the CdvB wH-domain, repairing the domain to a winged-helix related architecture. Importantly, the wH-domain of CdvB, necessary for interaction with CdvA and the recruitment of CdvB to the membrane, is not present in the other three ESCRT-III homologs found in *Sulfolobales* species. Also, no interaction of the other CdvB paralogs with CdvA has been reported (Samson *et al.*, 2011).

Interestingly, the CdvB paralogs can interact with themselves and the two other paralogs (Samson *et al.*, 2008). A recent study implied that CdvB does not constrict *in vivo* but functions

as a scaffold for CdvB1 and CdvB2 in *S. acidocaldarius*. The latter two proteins localize at the cell division plane, forming a ring-like structure after CdvB has already been recruited to mid-cell. Subsequent proteasomal degradation of CdvB allowed the formed CdvB1/B2 ring to constrict, completing cell division (Risa *et al.*, 2019). Furthermore, CdvB1/B2 were shown to be involved in other membrane-remodeling events in *Sulfolobales*, as they were found in secreted membrane vesicles (Ellen *et al.*, 2009). Deletion of *cdvB1* affected growth at 75°C only modestly and resulted in occasional failures in cell division. Remarkably, the $\Delta cdvB1$ strain was severely impaired at growth at 65°C, suggesting a more important role under stress conditions (Pulschen *et al.*, 2020). Additionally, deletion of *cdvB2* showed a loss of cell division symmetry and was severely affected in cell growth (Pulschen *et al.*, 2020). Analysis of cell division in the *SulfoScope* (an inverted fluorescent microscope with a heated chamber) showed that the CdvB and CdvB1 rings were correctly assembled in the $\Delta cdvB2$ strain, while after proteasome-mediated degradation of CdvB, CdvB1 rings were found at variable positions. This asymmetric division plane resulted in both ghost cells without DNA and in cells with a double amount of DNA. This suggested that CdvB2 fixes the position of the ring after the loss of the CdvB scaffold. It was proposed that CdvB2 and CdvB1 are recruited to the cell center by the CdvB ring and that, after proteasome mediated degradation of CdvB, they hold the division ring in position while it constricts (Pulschen *et al.*, 2020).

When *cdvB3* was deleted, growth was delayed, colonies on plates were very small, and cells had a division defect. In the *cdvB3* deletion strain, CdvB was not localized at mid-cell in a ring-like structure. Instead, a diffuse CdvB signal was detected in the cytoplasm. Furthermore, CdvA was mislocalized and formed distinct foci at the cell membrane of enlarged cells (Yang &

Driesssen, 2014). On the other hand, in *Sa. islandicus*, deletion of *cdvB1* and *cdvB2* was not successful. The reduction of intracellular CdvB1 protein levels led to a chain-like morphology of the cells, indicating that CdvB1 is important for the final abscission during cell division. In contrast to *S. acidocaldarius*, *cdvB3* deletion in *Sa. islandicus* lead to neither a cell division defect nor growth retardation. However, after infection with spindle-shaped virus 2 (STSV2), the $\Delta cdvB3$ *Sa. islandicus* strain no longer developed viral buds on the cell surface (Liu *et al.*, 2017). These findings indicate that, although all *Sulfolobus* species have four ESCRT-III homologs, their specific function is different among the *Sulfolobales*. Yet, CdvB seems to be the key cell division protein of the four ESCRT-III homologs, providing a platform for later cell division proteins. Recently, ESCRT-I and -II homologs were identified in *Lokiarchaeota*. Though the function of these other ESCRT homologs is unknown, they may be involved in vesicle formation rather than in cell division as ESCRT-III homologs are also present in *Lokiarchaeota* (Caspi & Dekker, 2018).

The third gene of the *cdv* operon encodes an ATPase (CdvC) that is homologous to the eukaryotic AAA-type (ATPase associated with various activities) ATPase Vps4 (Hobel *et al.*, 2008). In HeLa cell lines, Vps4 was observed to localize at the cell center during cytokinesis simultaneously with ESCRT-III. It was shown that Vps4 is important for the dynamic assembly and disassembly of ESCRT-III filaments, leading to the formation and constriction of intercellular bridges. As such, the dynamic turnover rate of ESCRT-III was dependent on the Vps4 ATPase activity (Mierzwa *et al.*, 2017, Caillat *et al.*, 2019). With regard to archaea, it is assumed that CdvC also remodels CdvB filaments and is thereby responsible for membrane invagination.

Coinciding with the other two Cdv proteins, CdvC localizes in a band-like structure to mid-cell during cytokinesis (Lindas *et al.*, 2008). Biochemical characterization of the CdvC protein from *Sa. solfataricus* showed (Samson *et al.*, 2008, Caspi & Dekker, 2018) that it assembles into single hexameric rings, comparable to Vps4 from *Saccharomyces cerevisiae* (Monroe *et al.*, 2014). Furthermore, it was shown that CdvC directly interacts with CdvB (Obita *et al.*, 2007). Important for this interaction is the N-terminal conserved MIT (microtubule interacting and trafficking) domain of CdvC that recognizes a C-terminal peptide motif of CdvB, called MIM2 (MIT domain interacting motif). The MIM2 domain is characterized by a proline-rich end part that is not present in the MIM2 domains of CdvB1 and CdvB2, resulting in a very weak binding affinity of the latter proteins to CdvC. Moreover, CdvB3 completely lacks the MIM2 domain and, hence, does not interact with CdvC (Samson *et al.*, 2008, Caspi & Dekker, 2018). This indicates that CdvB is the only ESCRT-III-like protein in *S. acidocaldarius*, recruiting CdvC to mid-cell. In contrast, a yeast-two-hybrid assay with ESCRT-III homologs and CdvC from *Sa. islandicus* showed interaction between CdvC and CdvB1/B2 (Liu *et al.*, 2017). Possibly, the lack of the proline-rich strand in CdvB1 and B2 is compensated by an additional interaction of the CdvB1/B2 protein with the MIT domain of CdvC (Caspi & Dekker, 2018). Moreover, the different interaction pattern of CdvC with the respective CdvB paralogs in *S. acidocaldarius* and *Sa. islandicus* might explain their different influence on cytokinesis. However, the mechanism of how CdvC is involved in cell division is still unknown. See **Figure 11** for the model of cell division in *S. acidocaldarius*.

7. Extreme Thermoacidophily and Stress Response

7.1. Thermoacidophily

Thermoacidophiles have specialized mechanisms to deal with stressors associated with their unusual temperature and pH optima and the fact that some thermal acidic environments are characterized by metal deposits. Although thermoacidophilic archaea largely do not have unique thermal or pH stress response mechanisms compared to thermophiles or acidophiles, some mechanisms play dual protective roles. Here, the major contributors to thermophily and acidophily will be discussed, with particular emphasis on where these stress responses overlap.

Because G-C bonds in DNA are more heat stable, it was initially hypothesized that thermophiles would have a higher G+C content than their mesophilic counterparts. Surprisingly, evaluation of DNA sequences from both thermophiles and mesophiles demonstrated that this was not the case (Tekaia *et al.*, 2002, Wang *et al.*, 2006, Zeldovich *et al.*, 2007). However, thermophiles have reduced mutation rates and have DNA repair mechanisms to combat biological damage at high temperatures. Though higher temperatures are correlated with an increase in mutations, base substitutions occur less frequently (Drake, 2009), or at an equivalent rate (Grogan *et al.*, 2001) in thermophiles compared to mesophiles. Furthermore, DNA repair can also occur by homologous recombination and DNA uptake through type IV pili (reviewed in **Section 6**).

Even though DNA stability does not depend on G+C content as it does not correlate with higher growth temperatures, thermophiles do have codon/amino acid preferences to support enzyme thermal stability. For instance, thermophiles favor codons AGG and AGA over CGN for arginine (Farias & Bonato, 2003, Singer & Hickey, 2003) and ATA for isoleucine, instead of

ATC or ATT (Singer & Hickey, 2003). In addition, thermophiles have a nucleotide bias for the more heat-stable purines A and G; these bases are more frequently neighbors in thermophile compared to mesophile genomes (Zeldovich *et al.*, 2007).

Not only do thermophiles have nucleotide preferences, but they also favor the use of certain amino acid residues. In general, thermophiles prefer amino acids that are charged or hydrophobic, while avoiding polar uncharged residues (Tekaia *et al.*, 2002). Intuitively, thermophilic proteins also avoid heat labile amino acids, such as histidine, glutamine, and threonine (Tekaia *et al.*, 2002, Singer & Hickey, 2003). Additionally, one study looking at the proteome amino acid composition of a variety of genomes discovered that the ratio (Glu + Lys)/(Gln + His) was lowest in mesophiles, higher in moderate thermophiles, and highest in extreme thermophiles ($T_{opt} \geq 70^{\circ}\text{C}$). This ratio was higher in thermostable chaperonins and DNA ligases in mesophiles compared to the rest of the proteome (Farias & Bonato, 2003). Furthermore, another study demonstrated that the occurrence of certain amino acids (IVYWREL) correlated with microbial optimal growth temperature and, thus, was a predictor of thermophily (Zeldovich *et al.*, 2007). Not only is the codon usage different in thermophiles, but their proteins averaged 283 aa compared to the 340 aa in mesophiles, possibly increasing the thermostability of the protein (Tekaia *et al.*, 2002).

Another thermal adaptation is the use of a reverse DNA gyrase, a DNA topoisomerase that introduces positive supercoiling to DNA strands, thereby increasing its heat stability. When the reverse DNA gyrase was deleted in *Thermococcus kodakarensis*, this extreme thermophile was more sensitive to higher temperatures, although the mutation was not shown to be lethal (Atomi *et al.*, 2004). However, though multiple methods were used to mutate the gyrase genes in

Sa. islandicus, no viable mutants were generated (Zhang *et al.*, 2013), indicating that this mutation can be lethal. In addition, a comparative genomic study showed that the only common gene in all extreme thermophiles sequenced at the time was a DNA reverse gyrase, lending further support to this enzyme's importance to thermophily (Forterre, 2002).

In extreme thermophiles, the thermal stress response involves the thermosome (a molecular chaperone also referred to as the rosettasome, or archeosome), a large HSP60-like protein complex originally discovered in *S. shibitae* (Trent *et al.*, 1990, 1991). In archaea, the thermosome is composed of multiple α and β subunits that form ring structures with either 8 or 9-fold symmetry (Kagawa *et al.*, 1995). A third γ subunit has been described in some extreme thermophiles that is down-regulated during heat shock (Archibald *et al.*, 1999, Kagawa *et al.*, 2003, Tachdjian & Kelly, 2006). The thermosome is a Group II chaperonin, conserved among archaea (Kagawa *et al.*, 1995), that shares structural and functional similarities to the bacterial Group I chaperonin GroEL complex (Trent *et al.*, 1991). In both heat shock complexes, cytosolic proteins are sequestered inside an internal cavity where they can refold. As demonstrated for the *Sa. solfataricus* thermosome, complexes are formed with different subunit compositions in a temperature-dependent manner, which supposedly have different substrate specificities (Chaston *et al.*, 2016). However, unlike the GroEL chaperonin of bacteria that has GroES to act as a lid for the complex, the thermosome does not have a separate subunit cap to close its internal cavity. Instead, the thermosome goes through conformational changes to close the apertures at each end (Schoehn *et al.*, 2000). These conformational changes are correlated with the binding and hydrolysis of ATP (Bigotti & Clarke, 2005). The apo-chaperonin is naturally in an open conformation and binding of ATP allows for the expansion of the thermosome entrance regions.

The hydrolysis of bound ATP causes a conformational change to enclose a denatured protein. Finally, the release of ADP and/or P_i liberates the folded protein and reopens the thermosome (Gutsche *et al.*, 2000, Gutsche *et al.*, 2000). The archaeal thermosomes are very similar in sequence to the cytosolic TCP1 eukaryotic chaperonins, the major difference being the number of subunits that compose each chaperonin's ring structures (Kagawa *et al.*, 1995).

In addition to the thermosome (~60 kDa subunits), thermoacidophiles also have HSP20 small heat shock proteins (sHSP), composed of 12 – 43 kDa subunits that also serve as chaperones during thermal stress response. HSP20 are common heat shock response proteins that are represented in every phylogenetic kingdom (Haslbeck *et al.*, 2005). These small proteins form larger structures of up to 50 subunits that can bind and stabilize proteins upon exposure to supra-optimal temperatures preventing protein aggregation (Wang *et al.*, 2010, Li *et al.*, 2012, Baes *et al.*, 2020). Similar to the thermosome, the sHSP can also refold denatured proteins.

If denatured or damaged proteins cannot be refolded via chaperones, they will be degraded by the proteasome. In eukaryotes, ubiquitin and the ubiquitin-related modifier-1 (Urm1) mark proteins for degradation by the proteasome. Likewise, in thermoacidophilic archaea, a Urm1-like molecule is covalently attached to proteins by an E1-like enzyme that is reliant on the hydrolysis of ATP. This urmylation designates a protein for degradation by the archaeal proteasome. The archaeal Urm1 protein is also degraded in the process, unlike the Urm1 signal in eukaryotic systems (Anjum *et al.*, 2015). The archaeal proteasome consists of two components: the 19S proteasome-activating nucleotidase (PAN) and the 20S proteolytic core particle. PANs are AAA^+ ATPase regulatory particles that unfold proteins in preparation for degradation by the core particle. The 20S core particle of the proteasome is a cylindrical stack of

four heptameric rings, which consists of two β rings flanked by two α rings. Within the core particle is an internal channel that contains the proteolytic site for protein degradation located within the two β rings. PAN is a homohexameric ring that associates with the outside α rings of the core particle. Comprehensive reviews of proteasomes and AAA⁺ ATPases are available (Bar-Nun & Glickman, 2012, Maupin-Furlow, 2012).

Another possible way thermophiles handle thermal stress is through compatible solutes that were originally described only as an osmoprotectant (Brown & Simpson, 1972). These organic molecules can accumulate intracellularly without interfering with cell metabolism to protect cytosolic components from other stressors, such as heat. For instance, *Mycobacterium smegmatis* mutants, unable to generate the compatible solute trehalose, exhibited sensitivities to elevated temperatures (Woodruff *et al.*, 2004). Though trehalose is present in all domains of life, it has been found in large quantities in several thermophilic archaea, such as *S. solfataricus* (Nicolaus *et al.*, 1988). On the other hand, the organic solutes di-myo-inositol-phosphate (DIP), di-mannosyl-di-myo-inositol-phosphate, di-glycerol-phosphate, mannosylglycerate and mannosylglyceramide are only observed in thermophilic organisms (reviewed in (Santos & Da Costa, 2002)). Also, cellular quantities of DIP have been shown to increase in response to increasing temperature in the thermophiles *Pyrococcus furiosus* (Martins & Santos, 1995) and *Thermotoga neapolitana* (Martins *et al.*, 1996). Additionally, di-mannosyl-di-myo-inositol-phosphate also increases in *T. neapolitana* due to thermal stress (Martins *et al.*, 1996). Although the exact mechanism of thermal protection by compatible solutes has not been determined, these compounds clearly have a part in the thermal stress response.

Some thermoacidophiles have pH optima near 0 (Schleper *et al.*, 1995), but most have cytosols with a near neutral pH, thereby resulting in a large pH differential across the membrane. Even *P. oshimae*, previously mentioned to have an internal pH of ~4.6, has a cytoplasmic pH several pH units above its environmental pH optima of < 1. Acidophiles (both thermophilic and mesophilic) must have ways to maintain their cytoplasm near neutral and still generate a proton motor force for energy generation. For instance, acidophiles have a reversed membrane potential, where the intracellular membrane is positively charged and the extracellular is negatively charged. This reverse membrane potential is driven by the active transport of potassium ions into the cell and prevents passive proton transport. Despite the large ΔpH across the membrane, acidophiles still generate ATP by coupling the influx of protons to the phosphorylation of ADP through a membrane-bound ATP synthase (Moll & Schäfer, 1988). However, this necessary influx of protons must be countered to maintain the intracellular pH of the cell. Proton pumps that are uniporters, symporters, and antiporters, are part of the energy generating respiratory chain that prevents the acidification of the intracellular environment by exporting internalized protons. These proton pumps are dependent on maintenance of the inverted membrane potential, which can be accomplished by pumping cations, such as potassium ions, into the cell (Schäfer, 1996). In fact, potassium ions have been shown to be a vital part of proton pumping in the respiration chain of *S. acidocaldarius* (Moll & Schäfer, 1988).

Alternatively, if protons do reach the cytoplasm, the cell must have ways to prevent the acidification of the cytosol. Intracellular molecules and proteins have buffering capacities that will neutralize internalized protons (Baker-Austin & Dopson, 2007). For example, basic amino acids, such as histidine, arginine, and lysine, can add to the buffering capacity of the cytoplasm.

As discussed in **Section 8.1.2**, members of the *Sulfolobales* have amino acid decarboxylases that may produce polyamines that are known to contribute to cytoplasmic buffering. In fact, reduction of amino acids in continuous cultures of *M. sedula* caused a decrease in internal pH and an increase in thermosome protein levels (Peeples & Kelly, 1995). Furthermore, protonated organic acids can permeate the cell and release a proton once inside the neutral cytoplasm. Heterotrophic acidophiles could possibly degrade organic acids without the production of a free proton to prevent the acidification of the cytoplasm (Baker-Austin & Dopson, 2007). As is the case for thermal stress, damaged proteins due to low pH can either be repaired by microbial chaperones or be marked for degradation by the proteasome.

7.1.1. Membrane Composition and Thermoacidophily

Thermoacidophiles maintain their intracellular pH by having membranes that are more impermeable to protons than neutralophiles (Konings *et al.*, 2002). The ether-linked lipids of archaeal membranes are more resistant to acid hydrolysis than the ester-linked lipids of bacterial membranes. Liposomes comprised of ether-linked lipids are more resistant to leakage due to thermal stress (Choquet *et al.*, 1994). Furthermore, tetraether lipids that span the microbial membrane make a monolayer that reduces the fluidity of the archaeal membrane, thereby decreasing its permeability to proton penetration and increasing heat tolerance. These tetraether lipids can also have multiple cyclopentyl rings that will further increase the packing of these lipids, making the membrane even less permeable to the diffusion of protons and more resistant to elevated temperatures. This has been demonstrated in liposomes derived from the membranes of *S. acidocaldarius* (Elferink *et al.*, 1994, Komatsu & Chong, 1998) and *P. oshimae* (van de

Vossenberg *et al.*, 1998). The most abundant tetraether lipids are the glycerol dibiphytanyl glycerol tetraethers (GDGTs) that can have up to 4 cyclopentyl rings on each chain. The number of ring structures within these tetraether lipids increased with increasing temperature in *Thermoplasma acidophilum* (Uda *et al.*, 2001), and with both decreasing pH and increasing temperature in *P. torridus* (Feyhl-Buska *et al.*, 2016) and *Acidilobus sulfurireducens* (Boyd *et al.*, 2011). These studies exemplify the importance of archaeal tetraether lipids with these cyclic moieties to survival in hot acid environments. However, only thermoacidophiles of the *Sulfolobales* have calditol, a cyclopentyl head group, ether linked to their GDGTs. Recently, the importance of this distinctive head group to acid resistance has been demonstrated in *S. acidocaldarius* as deletion of a calditol synthase resulted in a sensitivity to low pH (Zeng *et al.*, 2018).

It is clear that thermoacidophiles have several different mechanisms to persist in elevated temperatures and low pH, including the thermosome and proteasome to manage the impact of damaged proteins in the cytosol, membranes made up of tetraether lipids with a reverse potential, and proton pumps to maintain intracellular pH homeostasis (**Figure 12**).

7.2. Thermal Stress Response

While the importance of the thermosome, sHSPs, and proteasome to thermoacidophile heat tolerance is well known, the transcriptional thermal stress response of their associated genes is somewhat varied in the *Sulfolobales*. Although some sHSPs in other organisms are constitutively expressed, in *Sa. solfataricus* and *S. acidocaldarius* two sHSP are transcriptionally up-regulated during thermal stress (Tachdjian & Kelly, 2006, Li *et al.*, 2012, Baes *et al.*, 2020).

However, the expression of the α and β subunits of the thermosome are unchanged during thermal stress in *Sa. solfataricus* (Tachdjian & Kelly, 2006), while up-regulated in *Sa. shibatae* and *S. acidocaldarius* (Kagawa *et al.*, 2003, Baes *et al.*, 2020). Also, as previously mentioned, transcriptional expression of the γ subunit of the thermosome decreases during thermal stress in most studied species (Archibald *et al.*, 1999, Kagawa *et al.*, 2003, Tachdjian & Kelly, 2006), with the exception of *S. acidocaldarius* (Baes *et al.*, 2020). Although the proteasome plays a vital role in the heat shock response, the transcriptional expression of the 20S core components of the proteasome were unaffected by thermal stress in *Sa. solfataricus*, while the PAN regulatory component was significantly down-regulated under these conditions (Tachdjian & Kelly, 2006). However, post-transcriptional or -translational regulation, which has not yet been studied in detail, may also play a role in the thermosome's and proteasome's response to elevated temperatures.

The transcription of genes encoding components of the thermosome and proteasome are not significantly impacted by thermal stress, but this is not the case for much of the genome. In *Sa. solfataricus*, one-third of the genome, including sHSPs, was transcriptionally responsive to a 10°C temperature shift from 80°C to 90°C (Tachdjian & Kelly, 2006). Of note, many Type II Toxin-Antitoxin (TA) loci, which encode a ribonucleolytic Toxin and a corresponding Antitoxin protein, were also up-regulated. In particular, when the genes for one of these TA loci were deleted, the resulting *Sa. solfataricus* mutant became heat shock labile (Tachdjian & Kelly, 2006, Cooper *et al.*, 2009, Maezato *et al.*, 2011).

Thermoacidophiles can also use motility to seek out cooler conditions when thermal stress is encountered. Specifically, a temperature shift from 50°C to 80°C in *S. acidocaldarius*

resulted in an increase in swimming speed and run time. Also, when exposed to a temperature gradient, cells migrated away from the higher temperature to regions closer to their optimum temperature. However, a similar response was not seen when the pH was shifted from 2 to 4, indicating that this was a temperature specific stress response for the thermoacidophile (Lewus & Ford, 1999). Overall, thermoacidophiles mount a complex response to thermal stress that is varied among the members of the order *Sulfolobales*.

7.3. Metal Stress Response

Not only do thermoacidophiles have to combat the deleterious effects of life in hot acid, they also often encounter toxic heavy metals in their biotopes. In general, certain metal ions are essential for proper function of cells. Metals participate in many cellular processes, including as cofactors bound to proteins, as catalysts of redox reactions, and to transport electrons in the respiratory chain. In mesophilic organisms, the focus is on acquiring biologically important heavy metals that are scarce in their environment. But, in the case of acidophilic organisms, especially in mining environments, the aim is to avoid their influx of heavy metals into the cell or at least to reduce the effective intracellular concentration. Mesophilic bacteria do this by using active and passive systems to remove the metal ions from the cell, form complexes with these metals, or convert them into a less toxic form. It is also important to note that there are mechanisms associated with metabolic response to the reactive oxygen species (ROS) generated inside the cell after metal uptake (reviewed in (Lemire *et al.*, 2017)). Most of the studies on heavy metal resistance in *Crenarchaeota* have been done in *Sa. solfataricus* and *M. sedula*,

although many resistance mechanisms are still unknown, especially in species like *Sulfura metallicus*, which has unusually high metal resistance levels.

7.3.1. Uranium and Type II Toxin-Antitoxin Systems

Some extreme thermoacidophiles have adapted to toxic metals, such as uranium, through unusual resistance mechanisms. *M. sedula*, more so than a very similar species, *Metallosphaera prunae*, catalyzes the oxidization of U_3O_8 to soluble U(VI), mediated by Fe^{3+} generated from Fe^{2+} under chemolithoautotrophic conditions. However, the U(VI) so generated is toxic to the microbe (Mukherjee *et al.*, 2012). As an adaptation to uranium (U(VI) stress, *M. prunae* activates Type II TA systems to degrade cellular rRNA during uranium stress, thereby inducing a population-wide dormancy. This response is not seen in *M. sedula*. This resistance mechanism allows *M. prunae* to withstand the toxic effects of living in the uranium-rich environment from which it was initially isolated (Mukherjee *et al.*, 2012, Mukherjee *et al.*, 2017).

7.3.2. Active Transporters for Metal Resistance


Active transport efflux is one of the most common approaches used for cellular metal resistance and the Cop system, which is involved in copper (Cu) resistance, is the best studied example in archaea. It includes a transcriptional regulator (CopT), a metal-binding chaperone (CopM), and a Cu-transporting ATPase (CopA) (**Figure 13**) (Villafane *et al.*, 2011). The genes for this system form a cluster in the *Sa. solfataricus* genome (She *et al.*, 2001, Ettema *et al.*, 2003, Ettema *et al.*, 2006), which is also present in *M. sedula*, *S. acidocaldarius*, *Sulf. tokodaii* (Martínez-Bussenius *et al.*, 2017) and *Sulfura. metallicus* (Orell *et al.*, 2013, Martínez-Bussenius

et al., 2017). Furthermore, in *Sulfura. metallicus*, the cop cluster is duplicated in the genome, although one cluster has a lower transcriptional response to copper. This duplication likely relates to the high heavy metal resistance of this organism (Orell *et al.*, 2013).

Besides CopA, there is another copper-exporting ATPase, CopB, which contributes to Cu resistance and is present in some thermoacidophile genomes, including *Sa. solfataricus*. While CopA responds to intracellular Cu levels, CopB is constitutive (Völlmecke *et al.*, 2012). The Cop system is also responsive to cadmium (Cd), but apparently not to silver (Ag), as seen in *Sa. solfataricus* (Ettema *et al.*, 2006) and *Sulfura. metallicus* (Orell *et al.*, 2013). The *Sulfura. metallicus* Cop system also responds to chalcopyrite (CuFeS2) (Orell *et al.*, 2013) and likely reinforces its Cu resistance in mining environments.

7.3.3. Polyphosphate

Polyphosphates (PolyP) are ubiquitous molecules that play many cellular roles, including heavy metal resistance. These polymers are made of hundreds to thousands of inorganic orthophosphate (Pi) residues, linked by high energy bonds, similar to those in ATP. There are two major enzymes involved in metabolism of this molecule: polyphosphate kinase (PPK) and exopolyphosphatase (PPX). The first enlarges the PolyP chain, adding a Pi to the end in a reversible reaction, using ATP as substrate, while PPX catalyzes the reverse reaction, hydrolyzing PolyP starting from the terminal Pi residue (there are endo-polyphosphatases as well, that cleaves internal PolyP bonds) (Kornberg, 1999, Albi & Serrano, 2016):





In the *Crenarchaeota*, the enzyme that synthesizes PolyP has not been identified, since neither PPK1 nor PPK2 has been identified. Several archaea, especially in the *Sulfolobales*, possess a PPX enzyme (Martínez-Bussenius *et al.*, 2017, Wang *et al.*, 2019). Meanwhile, in *Euryarchaeotes*, a putative PPX was recently identified via bioinformatic analyses (Paula *et al.*, 2019). High concentrations of intracellular polyP are related to higher metal resistance in *Sulfolobales* species (Remonsellez *et al.*, 2006). Also, a *Sa. solfataricus* strain lacking PolyP due to overexpression of PPX, exhibited less copper resistance compared to the wild type strain (Soto *et al.*, 2019). Moreover, a spontaneous mutant with a functional PitA transporter, called *M. sedula* CuR1, was more resistant to copper and arsenic than the wild type strain that carries a truncated *pitA* (McCarthy *et al.*, 2014). PitA and Pho84 transporters are related to the uptake of Pi, and the exportation of Pi-Cu outside the cell (**Figure 13**). In *M. sedula*, Pho84-like transporters responded to the presence of Cu and are also proposed to be involved in Pi-Cu uptake (Rivero *et al.*, 2018). There are two ways in which PolyP helps directly to decrease intracellular effective concentration of metals. Since PolyP has negative charges, metal cations are attracted and complexed in the surface of the granule (**Figure 13**). On the other hand, the increase in metal concentration triggers PolyP degradation, and Cu associated with Pi is exported outside the cell by PitA or Pho84 transporters (**Figure 13**) (Keasling, 1997, Remonsellez *et al.*, 2006, Orell *et al.*, 2012, Grillo-Puertas *et al.*, 2014).

As mentioned, PolyP plays many roles in archaea and bacteria related to stress response and protein aggregation, the latter as an inorganic chaperone (Gray & Jakob, 2015). PolyP can also sequester metals, like Fe, preventing the Fenton reaction and generating more free radicals

(Gray & Jakob, 2015). Proteomic studies in the *Sa. solfataricus* PolyP (-) mutant revealed up-regulation of stress-related proteins when compared to the wild type strain (Soto *et al.*, 2019). Proteins, like peroxidases, were up-regulated, and other enzymes from various metabolic pathways exhibited changes that collectively reassembled metabolic oxidative stress response (Soto *et al.*, 2019).

7.3.4. Metal sequestration and transformation

Besides sequestration by PolyP granules, there are other molecules, such as DNA-binding proteins from starved cells (Dps) (**Figure 13**), that are proposed to bind to metals, thereby avoiding deleterious effects as seen in *Sa. solfataricus* (Wiedenheft *et al.*, 2005). *Sa. solfataricus* also contains a *mer* operon, required for Hg (II) reduction. The operon contains a regulator, MerR, that controls production of a mercuric reductase (MerA) along with other components (**Table 2**). MerR acts as both a repressor and a metal-responsive activator of mercury resistance genes by binding an operator sequence (*merO*) that also forms part of the operon (Schelert *et al.*, 2004, 2006).

7.3.5. Metal resistance and bioleaching

The use of mesoacidophiles and thermoacidophiles in biomining applications, that have reduced environmental impact, has driven interest in understanding heavy metal resistance in these organisms (Martínez-Bussenius *et al.*, 2017) (see **Section 9**). The relationship between metal resistance and lithoautotrophy was seen in *M. sedula* when *copA* was disrupted, which lowered copper resistance and consequently chalcopyrite bioleaching (Maezato *et al.*, 2012). In

contrast, a spontaneous mutant of *M. sedula*, strain CuR1, also showed increased bioleaching capability, associated with its higher Cu resistance (Maezato *et al.*, 2012). These examples illustrate that understanding metal resistance can lead to new strategies for bioleaching, where the *Sulfolobales* show great potential (see **Section 9**).

8. Metabolism

8.1. Heterotrophic metabolism

As mentioned in **Section 2**, microorganisms adapted to both high temperature and low pH, so called thermoacidophiles, have been identified mainly in the domain of Archaea and most of them belong to the two orders, the euryarchaeal *Thermoplasmatales*, including the genera *Thermoplasma*, *Picrophilus*, and the crenarchaeal *Sulfolobales* with the genera *Caldivirga*, *Thermocladium*, *Acidianus*, *Desulfurolobus*, *Metallosphaera*, *Sulfolobus*, *Saccharolobus* and *Sulfurococcus* (Bertoldo *et al.*, 2004, Zaparty & Siebers, 2011, Sakai & Kurosawa, 2018). Most of these organisms were isolated from solfataric fields or hot springs and are able to grow heterotrophically, using a variety of substrates as carbon and energy sources. Some have been also described as facultative heterotrophs that are also able to grow autotrophically by sulfur, metal or hydrogen oxidation. Among these species, the *Crenarchaeota* *Sa. solfataricus* and *S. acidocaldarius* have been most extensively analyzed with respect to the heterotrophic lifestyle and, therefore, represent archaeal model organisms for metabolic network reconstruction.

Sa. solfataricus shows high metabolic versatility and is able to utilize a broad spectrum of substrates, including monosaccharides (e.g. D-glucose, D-galactose, L-fucose, D-fructose, D/L-arabinose and D-xylose), disaccharides (e.g. cellobiose, maltose, sucrose, trehalose and lactose),

oligo- and polysaccharides (e.g., β -glucans, starch and dextrin), amino acids (e.g. glutamate), peptides and proteinaceous substrates (e.g. tryptone), and alcohols including aromatics (e.g. ethanol, phenol) (Quehenberger *et al.*, 2017, Schocke *et al.*, 2019). In contrast, *S. acidocaldarius* has a much narrower substrate spectrum; this could be attributed to its relatively smaller genome, which lacks numerous transport systems for substrate uptake. The metabolism of *S. acidocaldarius* is limited to a smaller range of substrates that include D-glucose, L-arabinose, D-xylose, sucrose, maltotriose, dextrin, starch, wheat bran, several fatty acids, and peptides and amino acids (Grogan, 1989, Wang *et al.*, 2019). Over the past several decades, the metabolic network, especially the central carbohydrate metabolism, in these two model organisms have been studied intensely, and is characterized by the presence of unusual and/or unique pathways and enzymes (Bräsen *et al.*, 2014, Quehenberger *et al.*, 2017). As such, we focus on these two model organisms to provide a prospective on heterotrophy in thermoacidophilic Archaea (see **Figure 14** for central metabolism overview).

8.1.1. Carbohydrates metabolism

Sugar catabolism: The two model organisms, *Sa. solfataricus* and *S. acidocaldarius*, can grow on a range of monosaccharides, disaccharides, and also polysaccharides (**Table 4**). A variety of polysaccharides-degrading enzymes have been studied in *Sa. solfataricus*, such as cellulase (Maurelli *et al.*, 2008), glucanase (Limauro *et al.*, 2001, Huang *et al.*, 2005, Girfoglio *et al.*, 2012), glucoamylase (Kim *et al.*, 2004), β -glycosidase (Noh & Oh, 2009), α -glucosidase (Rolfsmeier *et al.*, 1998, Wagner *et al.*, 2014), β -glucosidase (Moracci *et al.*, 1995, Shin *et al.*, 2013), α -amylase (Haseltine *et al.*, 1996, Wagner *et al.*, 2014), xylanase (Cannio *et al.*, 2004,

Maurelli *et al.*, 2008) and xylosidase (Moracci *et al.*, 2000). The degradation products, i.e. mono-, di- and/or oligosaccharides, are taken up into the cytoplasm primarily by various ATP binding cassette (ABC) transporters (Albers *et al.*, 2004, Choi *et al.*, 2013, Wagner *et al.*, 2017). The further processing of the monosaccharides follows a general scheme: the sugars are first oxidized to the corresponding lactone by sugar dehydrogenases, which in most cases belong to the medium chain dehydrogenase/reductase (MDR) superfamily (Lamble *et al.*, 2003, Brouns *et al.*, 2006, Nunn *et al.*, 2010, Haferkamp *et al.*, 2011, Wolf *et al.*, 2016, Reinhardt *et al.*, 2019), and in only some cases to the short chain dehydrogenase/reductase (SDR) superfamily (Yasutake *et al.*, 2007, Kim *et al.*, 2012). The sugar lactones are then hydrolyzed either enzymatically via lactonases or spontaneously in a non-enzymatic reaction favored by the extremely thermophilic conditions. The resulting sugar acids are subsequently dehydrated to the key intermediates, the 2-keto-3-deoxy sugar acids. These reactions are catalyzed by dehydratases mainly from the enolase superfamily. The further degradation then varies with respect to phosphorylation and (aldol) cleavage. In the modified branched ED pathway, the 2-keto-3-deoxy sugar acids, 2-keto-3-deoxy gluconate (KDG) and 2-keto-3-deoxy galactonate (KDGal) are either first phosphorylated by KDG kinase to KDPG/KDPGal and then cleaved to glyceraldehyde 3-phosphate (GAP) and pyruvate in the semi-phosphorylative (sp) ED branch, or they are directly cleaved by the same aldolase to yield glyceraldehyde (GA) and pyruvate in the non-phosphorylative (np) ED branch (Ahmed *et al.*, 2005, Lamble *et al.*, 2005). GAP is then directly oxidized to 3-phosphoglycerate (3PG) by a non-phosphorylating GAP dehydrogenase (GAPN), without coupling the oxidation to ATP generation via substrate level phosphorylation. 3PG is then further converted to a second molecule of pyruvate through the reaction sequence of the common lower Embden-Meyerhof-

Parnas (EMP) pathway. In the npED branch, GA is first oxidized in a ferredoxin-dependent manner to glycerate by GA: ferredoxin oxidoreductase and then phosphorylated to 2-phosphoglycerate (2PG) by means of 2-phosphoglycerate kinase. 2PG again enters the lower common shunt of the EMP. The 2-keto-3-deoxy-L-fuconate derived from the 6-deoxy hexose L-fucose is cleaved by the same aldolase to lactaldehyde, and pyruvate and the lactaldehyde is further oxidized in two consecutive steps to lactate and finally to a second molecule of pyruvate (Wolf *et al.*, 2016). Also, pentose degradation initially follows the same reaction sequence of oxidation and dehydration, finally leading to the corresponding 2-keto-3-deoxy acids, which are then further processed either in an aldolase dependent manner, called the Dahms pathway, or in an aldolase independent manner, referred to as the Weimberg pathway (Nunn *et al.*, 2010). The aldol cleavage in the Dahms pathway yields pyruvate and glycolaldehyde, which is then oxidized in two steps to glyoxylate, converted with acetyl-CoA to malate via malate synthase, and finally enters the TCA cycle as malate. In the Weimberg pathway, the 2-keto-3-deoxy pentanoates are converted via a second dehydration catalyzed by the 2-keto-3-deoxy xylonate dehydratase (KDXD) to 2-ketoglutarate semialdehyde (KGSA) and then oxidized through KGSA dehydrogenase (KGSADH) to 2-ketoglutarate entering the TCA cycle (Nunn *et al.*, 2010, Wagner *et al.*, 2017). While D-arabinose is only used as carbon source in *Sa. solfataricus* and degraded via the Weimberg pathway (Brouns *et al.*, 2006), L-arabinose and D-xylose are converted by both pathways to the same extent in *Sa. solfataricus* and *S. acidocaldarius* (Nunn *et al.*, 2010). However, deletion mutant analyses in *S. acidocaldarius* MW001 demonstrated that the Dahms pathway is dispensable, whereas the Weimberg pathway is essential for D-xylose degradation (Wagner *et al.*, 2017). In contrast to the other hexoses, another 6-deoxy sugar, i.e. L-

rhamnose, was also proposed to be degraded in an aldolase independent manner via the so called 2,4-di-keto pathway. The 2-keto-3-deoxy rhamnoate generated by the common initial reactions is then oxidized at C4 to 2,4-keto-3-deoxy-rhamnoate, which is cleaved by a hydrolase to yield lactate and pyruvate (Reinhardt *et al.*, 2019). The lactate is then subsequently oxidized to a second molecule of pyruvate.

An interesting feature of the sugar degradation routes in the *Sulfolobales* is the pronounced substrate promiscuity of the enzymes involved, especially in the upper part of the pathways catalyzing analogous reactions. This has first been described by Danson and co-workers, who found that the glucose dehydrogenase, the gluconate dehydratase and the KD(P)G aldolase from *Sa. solfataricus*, also accept the D-galactose derivatives as substrates (Lamble *et al.*, 2003, Lamble *et al.*, 2004, Lamble *et al.*, 2005). Subsequently, it was shown that the same sugar dehydrogenase is also responsible for the oxidation of the pentoses D- xylose and L-arabinose (Nunn *et al.*, 2010). However, D-arabinose and L-fucose demonstrated the opposite stereochemistry on C2-C4 compared to L-arabinose oxidized by another dehydrogenase, which also accepted L-rhamnose as a substrate (Brouns *et al.*, 2006, Wolf *et al.*, 2016, Reinhardt *et al.*, 2019). Interestingly, the dehydratase's conversion of L-rhamnonate is different from the conversion of D-arabonate and L-fuconate, indicating that the promiscuity of the dehydratases is less pronounced than that of the dehydrogenases (Reinhardt *et al.*, 2019). This has also been proposed for the gluconate/galactonate dehydratase, which presumably does not accept the pentose derivatives as substrates suggesting the presence of an alternative dehydratase, although this has not been confirmed (Nunn *et al.*, 2010). The KD(P)G aldolase, however, shows by far the most marked promiscuity, accepting 2-keto-3-deoxy acids derived from hexoses and

pentoses, i.e. D-glucose, D-galactose, D-xylose, L-arabinose, D-arabinose, and L-fucose, as well as the phosphorylated derivatives KDPG and KDPGal (Ahmed *et al.*, 2005, Lamble *et al.*, 2005, Nunn *et al.*, 2010, Wolf *et al.*, 2016). Thus, this enzyme plays a central role in sugar degradation via the branched ED pathway, comprising the npED and spED branch, as well as the Dahms pathway, both of which are involved in the breakdown of a wide variety of naturally occurring sugars in *Sulfolobales*.

The end products of the glycolytic pathways are pyruvate and the tricarboxylic acid (TCA) cycle intermediates, α -ketoglutarate and malate. As in all Archaea, the pyruvate produced is then oxidatively decarboxylated to acetyl-CoA by ferredoxin oxidoreductase (Kerscher *et al.*, 1982, Zhang *et al.*, 1996, Yan *et al.*, 2016). The resulting acetyl-CoA is then completely oxidized to CO₂ in the TCA cycle (Danson, 1988). Also, the oxidative decarboxylation of α -ketoglutarate is carried out by a ferredoxin-dependent oxidoreductase. NAD(P)⁺-dependent dehydrogenase complexes (i.e. pyruvate dehydrogenase complex), known from bacteria and eukaryotes, are not operative in the *Sulfolobales* or, for that matter, Archaea in general (Payne *et al.*, 2010).

Sugar anabolism: Sugar degradation in aerobic Archaea, including the *Sulfolobales*, proceeds via a modified branched ED pathway and is, at least in some ways, analogous to the Dahms and/or Weimberg pathways. Notably, for a complete EMP pathway, only a functional phosphofructokinase (PFK) is missing, which highlights the exclusively gluconeogenic function of the EMP in *Sulfolobales* (Bräsen *et al.*, 2014). In addition to the common lower shunt enzymes, gluconeogenesis is initiated by the phosphoenolpyruvate synthetase, bypassing the irreversible pyruvate kinase (PK) reaction (Haferkamp *et al.*, 2019), and also involves the classical glyceraldehyde-3-phosphate (GAPDH)/phosphoglycerate kinase (PGK) couple, which

bypasses the irreversible GAPN (Kouril *et al.*, 2013). Perhaps, the most striking difference to classical gluconeogenesis is the presence of the bifunctional fructose-bisphosphate aldolase/phosphatase (FBPA/ase), characterized among others from the *Sulfolobales* *M. sedula* and *Sa. solfataricus*, replacing the classical fructose-1,6-bisphosphate aldolase and F1,6BP phosphatase couple of the gluconeogenesis route in Bacteria and Eukarya (Say & Fuchs, 2010). This enzyme catalyzes the conversion of GAP and DHAP to F6P without liberating the F1,6BP intermediate.

All of these modifications, both in sugar catabolism and anabolism, have been discussed as a mechanism of metabolic thermodaptation, since the formation of extremely thermolabile triose phosphates, e.g. GAP, DHAP, and 1,3BPG, are avoided (Say & Fuchs, 2010, Kouril *et al.*, 2013, Figueiredo *et al.*, 2017, Zhang *et al.*, 2017). Also, the altered regulatory properties of the pathways and enzymes could contribute to thermodaptation, since the classical control points, e.g., HK, PFK and PK, are missing or changed in Archaea, including in the *Sulfolobales*. For example, the main glycolytic control point is the GAPN, instead of the sugar(phosphate) kinases being activated by glucose 1-phosphate (Ettema *et al.*, 2008, Kouril *et al.*, 2017). Also, the PKs so far characterized, including those from *Sulfolobus* spp., show divergent regulation compared to the enzymes from the classical bacterial and eukaryotic pathways (Haferkamp *et al.*, 2019, Johnsen *et al.*, 2019).

The phosphoglucose isomerase (PGI) interconverting F6P and G6P and the phosphoglucomutase (PGM) present in *Sulfolobales* represent the branch point to the synthesis of the storage compound glycogen and trehalose as the only compatible solutes in *Sulfolobus* spp. described so far (Martins *et al.*, 1997, Bräsen *et al.*, 2014). The additional presence of a

hexokinase suggests that these biosynthesis reactions can also be directly initiated from the substrate molecule glucose. However, the spED branch, particularly the KDG kinase as its key enzyme, may be involved in gluconeogenesis by providing triose phosphates for anabolic purposes by directing the flux from the upper branched ED to the hexose phosphate synthesis (Kouril *et al.*, 2013).

The pentose metabolism in the *Sulfolobales* and *Thermoplasmatales* are also characterized by a missing oxidative pentose phosphate pathway (OPPP). Furthermore, the non-oxidative pentose phosphate pathway (NOPPP) is only partially present in *Sulfolobales* lacking a ribulose-5-phosphate epimerase and a transaldolase, but is entirely present in the *Thermoplasmatales* (*Thermoplasma* and *Picrophilus*) (Bräsen *et al.*, 2014). However, the pentose precursor ribulose-5-phosphate, as in most Archaea, is provided by the ribulosemonophosphate (RuMP) pathway converting F6P to Ru5P and formaldehyde via D-arabinohexulose 6-phosphate. Ribose 5-phosphate (R5P) and erythrose 4-phosphate (E4P) are precursors of nucleotide and aromatic amino acid biosynthesis, respectively, and are then provided by the remaining reaction of the NOPPP. Thus, F6P is the main source of pentose/tetrose phosphates in most Archaea in general and, particularly, thermoacidophiles (Bräsen *et al.*, 2014).

8.1.2. Degradation of proteins and amino acids

Although thermoacidophilic Archaea are routinely grown on proteinaceous substrates, surprisingly little is known about the breakdown of amino acids in these organisms. It appears that certain *Sulfolobales* species grow well on these complex protein substrates, such as yeast

extract, caseinhydrolysate, NZ- amine, and on mixtures of amino acids, but hardly if at all on single amino acids as a sole carbon, energy and nitrogen source (Grogan, 1989, Stark *et al.*, 2017, Quehenberger *et al.*, 2019). The reason for this remains unclear so far. However, *Sa. solfataricus*, when growing on caseinhydrolysate, prefers certain amino acids like glutamate, methionine, leucine, phenylalanine, and isoleucine and, to a lesser extent, threonine, alanine, aspartate, glycine, and tyrosine (Stark *et al.*, 2017). Growth of *Sa. solfataricus* on glucose was stimulated most by glutamate and, to a lesser extent, by aspartate (Stark *et al.*, 2017).

Additionally, growth of *S. acidocaldarius* on glucose is enhanced by glutamate, followed by aspartate, arginine and lysine (Quehenberger *et al.*, 2019). For the breakdown of (poly)peptides though, many proteinase and peptidase encoding genes are present in the genomes of the *Sulfolobales*, but few have been studied in detail (Cannio *et al.*, 2010, Gogliettino *et al.*, 2010). Additionally, ABC transporters for di/oligopeptides have been identified in *Sa. solfataricus*, *Sulf. tokodaii* and *T. acidophilum* (Albers *et al.*, 2004).

There are generally three mechanisms known by which amino acid degradation is initiated: decarboxylation, transmination, and (oxidative) deamination. Decarboxylation leads to biogenic amines and there are some reports in *Sa. solfataricus* for amino acid decarboxylases, e.g. arginine decarboxylase, which might play a role in the biosynthesis of spermidine and putrescine (Giles & Graham, 2008, Esser *et al.*, 2013). However, the more common mechanism for amino acid breakdown in Archaea, which was intensively studied for *Thermococcales*, is the transamination with 2-oxoacids, mainly alpha-ketoglutarate, and/or the oxidative deamination using amino acid dehydrogenases, most importantly the glutamate dehydrogenase liberating ammonia and concomitantly reducing NAD(P)⁺ (Yokooji *et al.*, 2013, Awano *et al.*, 2014, Scott

et al., 2014). Both mechanisms ultimately lead to the formation of the corresponding 2-keto acids that are subsequently oxidatively decarboxylated by ferredoxin dependent 2-oxoacid: Fd oxidoreductases, which are also well known and characterized from *Sulfolobus* species, *Sulf. tokodaii* in particular (Kerscher *et al.*, 1982, Zhang *et al.*, 1996, Park *et al.*, 2006, Yan *et al.*, 2016). The reaction products are the corresponding CoA esters of the carboxylate backbone of the amino acids. In aerobes, these CoA esters are then completely oxidized to CO₂ by channeling these compounds into the central energy metabolic pathways via pyruvate, PEP, acetyl-CoA, and the TCA cycle intermediates. These pathways have also been identified in *Sa. solfataricus* by *in silico* metabolic reconstructions (Ulas *et al.*, 2012). In anaerobic archaea, most well studied in *Thermococcales*, the CoA esters play pivotal roles in fermentative energy generation by being the sole source of net ATP via substrate level phosphorylation. The key enzymes are the ADP-forming acyl-CoA synthetases coupling the CoA ester hydrolysis to the ATP formation from ADP and Pi (Awano *et al.*, 2014, Scott *et al.*, 2014, Weiße *et al.*, 2016). However, these ADP-forming acyl-CoA synthetases are also present in aerobic archaea, including the *Sulfolobales*. Specifically, during *Sa. solfataricus* growth on casein hydrolysate carboxylic acids, mainly isovalerate, are excreted into the medium (Stark *et al.*, 2017). Although such product formation appears unusual under aerobic conditions, it is well documented in bacterial model organisms, like *E. coli* and *Bacillus subtilis* (Bräsen & Schönheit, 2004), and this so-called “overflow metabolism” has also been observed in halophilic archaea (Bräsen & Schönheit, 2004). However, under acidophilic conditions, carboxylic acids at elevated concentrations act as uncouplers/protonophores that lead to the acidification of the cytosol and the breakdown of membrane gradients (Baker-Austin & Dopson, 2007). Moreover, the excretion of products

results in the loss of carbon, which may account for the less efficient growth of *Sa. solfataricus* on casein hydrolysate compared to glucose. Furthermore, in *Sa. solfataricus* pyroglutamate may form from glutamate at high temperature and low pH to inhibit growth of thermophilic archaea (Stark *et al.*, 2017). However, in *S. acidocaldarius*, pyroglutamate is not inhibitory and even serves as a carbon source, making *S. acidocaldarius* a better thermoacidophilic platform organism for applications with glutamate-containing media (Vetter *et al.*, 2019). Finally, it has not been determined whether the carboxylic acids formed during exponential growth can be reused by *Sulfolobus* species, as it has been shown for other aerobic organisms.

8.1.3. Degradation of lipids and fatty acids

In general, the mechanisms for fatty acid and, fatty acid-based, (phospho)lipid metabolism are not well understood in Archaea. However, *Sa. solfataricus* P1 can (partially) degrade corn oil as well as olive oil (Choi *et al.*, 2016), and *S. acidocaldarius* can cleave triacylglycerols (Zweerink *et al.*, 2017). Moreover, several extracellular esterases/lipases have been identified from *Sa. solfataricus*, *S. acidocaldarius* and *Sulf. tokodaii* (Suzuki *et al.*, 2004, Choi *et al.*, 2016, Zweerink *et al.*, 2017). A TetR-family transcription factor (FadR_{Sa}) plays a role in regulation of putative fatty acid metabolism related genes in *S. acidocaldarius* (**Table 2**) and growth of this organism on short-chain fatty acids, i.e. butyrate and hexanoate, as sole carbon sources has been demonstrated (Wang *et al.*, 2019).

Genomic analyses revealed that all the genes encoding homologs of the key enzymes involved in the bacterial-like β -oxidation are present in the genomes of several Archaea, including *Sa. solfataricus* and *S. acidocaldarius* (Dibrova *et al.*, 2014, Wang *et al.*, 2019),

implying that fatty acids could be degraded in these organisms through classical β -oxidation; but, it remains unproven. Nonetheless, *S. acidocaldarius* can grow on acetate (as also described for many other Archaea including methanogens, *Pyrobaculum*, halophilic Archaea) as the sole carbon and energy source. The glyoxylate shunt is operative under these conditions, demonstrating that C2 units can be assimilated by this reaction sequence (Uhrigshardt *et al.*, 2002).

As a product of lipid hydrolysis, glycerol can be utilized as a carbon and energy source by many bacteria. In Archaea, glycerol degradation has been examined in halophiles (Sherwood *et al.*, 2009, Rawls *et al.*, 2011, Williams *et al.*, 2017). Glycerol is taken up either by simple diffusion or glycerol transporters (Richey & Lin, 1972, Stroud *et al.*, 2003, Anderson *et al.*, 2011). The haloarchaea employ one of the bacterial-like mechanisms, first phosphorylating glycerol followed by sn-glycerol-3-phosphate oxidation to DHAP. So far, glycerol catabolism has not been studied in thermoacidophilic archaea, although there are some indications that *Sulfolobales* spp. do not utilize glycerol as carbon and energy source (Grogan, 1989). In contrast, genome-scale metabolic network reconstruction and modelling suggested glycerol as the most efficient carbon source for *Sa. solfataricus* (Ulas *et al.*, 2012).

8.1.4. Degradation of other substrates

In addition to the three major types of nutrients described above, other substrates also support heterotrophic growth of thermoacidophilic archaea. For instance, *Sa. solfataricus* grows on acetoin, citric acid, alcohols and phenol (Izzo *et al.*, 2005, Chong *et al.*, 2007, Wolf *et al.*, 2016). There are several alcohol dehydrogenases in *Sa. solfataricus*, allowing oxidation of

alcohols into aldehydes. The second step is then the oxidation to the corresponding carboxylic acids and subsequent activation of the CoA esters for further degradation (Chong *et al.*, 2007, Chong *et al.*, 2007). Phenol is degraded in *Sa. solfataricus* in a classic pathway, as reported in some bacteria, e.g. *Burkholderia pickettii* and *Pseudomonas stutzeri* OX1. It is first converted to catechol, which undergoes a series of ring cleavage reactions producing products that finally enter TCA cycle (Izzo *et al.*, 2005).

In conclusion, thermoacidophilic (facultative) heterotrophs, especially from the *Sulfolobales*, utilize a variety of substrates for cell growth (reported substrates with degradation pathways are concluded in **Table 4**). This metabolic versatility includes the potential for autotrophic growth, combined with the thermoacidophilic lifestyle, making them ideal candidates for the development as platform strains for the production of added-value compounds from renewable (waste) materials like lignocellulosics (see **Section 9**).

8.2. Autotrophy and Chemolithotrophy

8.2.1. Autotrophy

The inhospitable environments in which the *Sulfolobales* thrive often have a scarcity of organic carbon available. As a result, many of the species within this order rely on the autotrophic fixation of CO₂ to support growth. At present, six mechanisms for CO₂ fixation are known throughout the domains of life. Several of these mechanisms build carbon-carbon bonds by fixing CO₂ using oxygen-sensitive carboxylases, or in the case of the Calvin-Bassham-Benson (CBB) cycle, ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCo) (Hugler *et al.*, 2003). While RuBisCo is not oxygen-sensitive, it does have a highly detrimental side

reaction with oxygen, and the prevalence of oxygen-reducing terminal oxidases throughout the *Sulfolobales* (anaerobes included) seems to indicate that oxygen is essential for the organisms' ability to survive their extreme conditions. The autotrophic pathway in the *Sulfolobales* circumvents the dependence on RuBisCo by incorporating bicarbonate molecules rather than CO₂ (Gong *et al.*, 2019). This pathway, named the 3-hydropropionate/4-hydroxybutyrate (3-HP/4-HB) cycle after two of its prominent intermediates, generates one acetyl-CoA per cycle (Berg *et al.*, 2007). Initially identified in *M. sedula*, the 3-HP/4-HB pathway shares the enzymatic route from succinyl-CoA to acetyl-CoA with the dicarboxylate/4-hydroxybutyrate autotrophy cycle. However, the generation of succinyl-CoA eschews the traditional ferredoxin-powered pyruvate synthase and instead relies on the sequential addition of bicarbonate molecules and coenzyme-A moieties (Fuchs, 2011). One full rotation of the 3-HP/4-HB cycle costs the cell 4 ATP and 4 NADPH, making it only a moderately expensive route to carbon fixation (Gong *et al.*, 2019).

Since the discovery of the 3-HP/4-HB pathway, the enzymes responsible for each step in the pathway have been identified and characterized (Kockelkorn & Fuchs, 2009, Teufel *et al.*, 2009, Ramos-Vera *et al.*, 2011). *M. sedula* has served as the model organism for many of these efforts, and part of the pathway characterization has involved linking the enzymes of the 3-HP/4-HB pathway to open reading frames within *M. sedula*. This has proved to be challenging, considering the promiscuity of enzymes causing overlapping catalytic activities. In the case of 4-hydroxybutyrate-CoA synthetase, five candidate genes were identified based on bioinformatic analysis. However, three of these genes showed no activity for the desired reaction (Ramos-Vera *et al.*, 2011), while the remaining two did demonstrate this activity (albeit with orders of

magnitude of difference in V_{max} values). Further investigation into these two candidates determined that one was merely a promiscuous acyl-CoA synthetase rather than the genuine 4-hydroxybutyrate-CoA synthetase (Hawkins *et al.*, 2013).

As greater understanding of the precise enzymatic path of the 3-HP/4-HB cycle has developed, modeling of the pathway has become a point of interest. This effort has been greatly aided by the quantification of thermodynamic and kinetic parameters associated with each step in the pathway (Ramos-Vera *et al.*, 2011, Loder *et al.*, 2016). A kinetic model of the system in *M. sedula* revealed differing degrees of rotations the cycle can undergo, resulting in a different distribution of products including acetyl-CoA and succinyl-CoA (Loder *et al.*, 2016). This modeling supports previous exploration into the channeling of carbon into various biosynthetic pathways from the 3-HP/4-HB cycle. Based on this isotope distribution, it was determined that the majority of CO_2 taken up during autotrophy generates succinyl-CoA, representing a half-turn or one-and-a-half turns of the full cycle. Acetyl-CoA (one full turn) is generated during this process, but only enough for amino acids directly synthesized from acetyl-CoA. Otherwise, succinyl-CoA dominates as the product of this cycle (Estelmann *et al.*, 2011). Extension of 3-HP/4-HB cycle modeling was done to identify the effect of the pathway on carbon isotopes and demonstrated that the source of carbon for the cycle may not be extracellular bicarbonate. Instead, bicarbonate is formed intracellularly as CO_2 is taken up by the cell (Pearson *et al.*, 2019).

While this autotrophic cycle requires 16 steps to generate acetyl-CoA from two bicarbonate molecules, only 13 enzymes are involved in the pathway. This inconsistency points towards the unusual redundancy in enzyme function throughout this pathway and it manifests in

a variety of ways (**Figure 15**). For example, the two-step conversion of crotonyl-CoA to acetoacetyl-CoA is catalyzed by the crotonyl-CoA hydratase. The first step of this conversion, however, can also be catalyzed by 3-hydroxypropionyl-CoA dehydratase, which also serves a separate function of dehydrating 3-hydroxypropionyl-CoA into acryloyl-CoA. Other redundancies include the reduction of malonyl-CoA and succinyl-CoA by the aptly named malonyl-CoA/succinyl-CoA reductase and the catalysis of hydroxypropionate to 3-hydroxypropionyl-CoA by either 3-hydroxypropionyl-CoA synthetase or the promiscuous 4-hydroxybutyryl-CoA synthetase (Loder *et al.*, 2016). Notably, these overlaps in function reflect the structural similarity in the 3-hydroxypropionate half of the pathway and the 4-hydroxybutyrate half of the pathway. In fact, the two halves can almost be viewed as the same sequence of reactions, with enzymes acting on the same functional groups and the substrates varying only in length of the carbon chain.

The 3-HP/4-HB cycle is highly conserved within the *Sulfolobales*. In fact, with the exception of *Sulfo. acidophilus*, all genome-sequenced members of the order possess homologs to the characterized enzymes from *M. sedula* (Counts, 2020). In spite of this conservation, not all *Sulfolobales* appear to be capable of carbon fixation. A prime example of this is *S. acidocaldarius*, which at the time of its isolation was reported to grow chemolithoautotrophically in the presence of elemental sulfur (Brock *et al.*, 1972). More recently, it appears that the commonly used lab strain, *S. acidocaldarius* DSM 639, is a strict heterotroph (Zeldes *et al.*, 2019). This example may be explained through an incomplete transcriptional regulation pathway. Recently, a conserved transcriptional regulator, HhcR, was proposed to be a universal autotrophy regulator in the *Sulfolobales* (Leyn *et al.*, 2015) (**Table 2**).

Several genes involved in the 3-HP/4-HB cycle appear to be lacking the binding motif for this regulator in *S. acidocaldarius* (Zeldes *et al.*, 2019). Whether this lack of promoter binding regions is a result of strain domestication by extensive heterotrophic growth in the lab remains to be determined. Further complicating the network of autotrophy in the *Sulfolobales* is the existence of all genes necessary for the dicarboxylate/4-hydroxybutyrate in several chemolithoautotrophs, including *M. sedula*. However, these genes are transcribed at a low level even during autotrophic growth conditions (Berg *et al.*, 2010). One area of interest for this pathway is its application in engineered organisms. A partial pathway was engineered into *E. coli* to generate both 3-hydroxypropionate and 4-hydroxybutyrate as a means of generating a block copolymer product (Meng *et al.*, 2012). Parts of the cycle from *M. sedula* have been introduced into *Pyrococcus furiosus* in an effort to convert CO₂ into 3-hydroxypropionic acid (Keller *et al.*, 2013). Similarly, *E. coli* has again been host to 3-HP/4-HB enzymes in order to produce propionic acid and acrylic acid (Liu & Liu, 2016).

8.2.2. Chemolithotrophy at the Cell Surface

In order to power an autotrophic lifestyle, members of the *Sulfolobales* tend to rely on metals and inorganic compounds that prevail in their primordial environments. The oxidation of ferrous iron (Fe²⁺) is one such source of energy. Despite a very positive reduction potential (+0.77 V for Fe³⁺/Fe²⁺), chemolithoautotrophs are still able to leverage iron oxidation to drive the electron transport chain (ETC) (Amend & Shock, 2001). While this pathway has been studied most intensely in mesoacidophiles like *Acidithiobacillus ferrooxidans*, transcriptomic studies of several *Sulfolobales* members have shed light on enzymes responsible for thermoacidophilic iron

oxidation (Bathe & Norris, 2007, Auernik & Kelly, 2008, Kozubal *et al.*, 2011). *M. sedula* is a prolific iron oxidizer and has served as the organism of study to further elucidate the mechanism for iron oxidation. These results indicate control of iron oxidation by a locus known as the *fox* cluster. The cluster appears to be membrane-bound, with iron oxidation taking place at the surface of the cell membrane and funneling electrons into the ETC.

A variety of sulfur species also persist in the *Sulfolobales*' habitats. Anaerobic *Sulfolobales* reduce zero-valent sulfur to hydrogen sulfide (H_2S) in conjunction with oxidizing diatomic hydrogen. Much like iron oxidation, sulfur reduction takes place entirely at the surface of the cell membrane, where a membrane-bound sulfur reductase acts on zero-valent sulfur (**Figure 16 A**) (Laska *et al.*, 2003). The complex achieves this reduction by transferring electrons from reduced quinols (specifically *Sulfolobus* quinol). Quinone cycling links sulfur reduction to hydrogen oxidation, and it both regenerates the oxidized quinone pool and powers proton export (Kletzin *et al.*, 2004). The dominant form of zero-valent sulfur under acidic conditions is cyclooctasulfur, but experimental evidence suggests sulfane sulfur from polysulfide chains may be the substrate for sulfur reductase (Blumentals *et al.*, 1990).

8.2.3. Chemolithotrophy within the Cytoplasm

In contrast to iron oxidation and sulfur reduction, sulfur oxidation takes place largely in the cytoplasm, where it cycles through a number of different reduced inorganic compounds (RISCs) (**Figure 16 B**). The premier enzyme of sulfur oxidation in the *Sulfolobales* is the cytoplasmic sulfur oxygenase reductase (SOR). The 24-subunit homomeric enzyme disproportionates zero-valent sulfur into H_2S and sulfite (SO_3^{2-}) without the assistance of any

cofactors (Kletzin, 1989). An abiotic reaction of these two products generates thiosulfate ($S_2O_3^{2-}$) as a byproduct (Kletzin, 1992). These species are coupled to the ETC through a variety of membrane-bound oxidoreductases: sulfide:quinone oxidoreductase (SQR) for H_2S (Brito *et al.*, 2009), thiosulfate:quinone oxidoreductase (TQO) for $S_2O_3^{2-}$ (Muller *et al.*, 2004), and sulfite:acceptor oxidoreductase (SAOR) for SO_3^{2-} (Zimmermann *et al.*, 1999). In the case of SAOR, activity of the enzyme has been detected in *Acidianus ambivalens*, but the enzyme has not been linked to an open reading frame in any of the *Sulfolobales*. Caldariellaquinone is the primary acceptor of electrons for these oxidoreductases.

While the product of SQR is a polysulfide chain that can be recycled to SOR and the product of SAOR is fully oxidized sulfate (SO_4^{2-}), TQO generates tetrathionate ($S_4O_6^{2-}$) as a product (Muller *et al.*, 2004). Recent studies have investigated the possibility of tetrathionate acting as the substrate for a set of highly conserved genes in the *Sulfolobales*, the *hdr/dsr/tusA* locus. DsrE3A and TusA both appear to be sulfur trafficking proteins, which cleave the sulfur-sulfur bond of $S_4O_6^{2-}$ to regenerate $S_2O_3^{2-}$ and form an organic persulfide compound. The persulfide ultimately acts as the substrate for the membrane-bound heterodisulfide reductase (Hdr) complex, which generate SO_4^{2-} (Liu *et al.*, 2014). A tentative role of the Hdr complex is once again the reduction of quinones to conserve energy. However, recent studies in the dimethyl sulfide (DMS)-degrading *Hyphomicrobium denitrificans* have demonstrated the association of a lipoate-binding protein with the Hdr complex and the importance of the *hdr/dsr/tusA* locus for energy conservation from DMS (Koch & Dahl, 2018). Homologs of these binding proteins have been identified in some *Sulfolobales*, and the reduction potential of lipoate is sufficient to reduce NAD^+ , thereby conserving energy (Cao *et al.*, 2018). While the exact acceptor molecule of Hdr

has yet to be confirmed in the *Sulfolobales*, it seems clear that the complex and its traffickers play some role in conserving energy from tetrathionate. In addition, the complex provides a route to total oxidation of $\text{S}_2\text{O}_3^{2-}$ to SO_4^{2-} and may serve to detoxify the abiotic byproduct of SOR.

A final avenue to energy conservation is the phosphorylating pathway of cytoplasmic SO_3^{2-} oxidation. In this pathway sulfite is attached to AMP by adenyl-5'-phosphosulfate reductase (APSR) using an unknown electron acceptor. Adenylylsulfate:phosphate adenylyltransferase (APAT) then replaces the sulfite group with phosphate to generate ADP. Finally, two molecules of ADP are converted to ATP and AMP by adenylate kinase (AK), thereby generating ATP directly from SO_3^{2-} oxidation (Kappler & Dahl, 2001). Much like SAOR, activity of these enzymes has been detected in *A. ambivalens* cell extracts, but the enzymes have not been purified, characterized, or linked to a sequence in the genome (Zimmermann *et al.*, 1999).

Sulfur transport into the cytoplasm remains something of a mystery in the *Sulfolobales*. No transporter for zero-valent sulfur has been identified yet in any organism from this order, although a possible sulfate transporter in *Metallosphaera cuprina* has been identified in transcriptomic data (Jiang *et al.*, 2014). Evidence has been presented, however, for the passive diffusion of H_2S through the membrane (Mathai *et al.*, 2009). In addition, an extracellular tetrathionate hydrolase is expressed in the *Acidianus* spp., many of which are prolific sulfur oxidizers (Protze *et al.*, 2011). The reaction generates thiosulfate extracellularly, which may undergo abiotic reactions in the acidic environment to generate the appropriate sulfur species for transport across the membrane. Given the wide distribution of sulfur substrates used within the cytoplasm, it is difficult to say what RISC might act as the “starting point” of sulfur oxidation,

and understanding sulfur transport is still a key area of investigation for chemolithotrophy in the *Sulfolobales*.

8.2.4. Terminal Oxidases and the Electron Transport Chain (ETC)

The ETC in the *Sulfolobales* deviates from the traditional four-complex structure. While homologs to complexes I and II (NADH dehydrogenase and succinate dehydrogenase, respectively) have been identified (Lemos *et al.*, 2001, Melo *et al.*, 2004), complexes III and IV appear to be combined into single quinone oxidoreductase complexes. In fact, cytochrome c has yet to be found in any of the *Sulfolobales*. As a result, these quinone oxidoreductases are responsible for the reduction of molecular oxygen and pH homeostasis through proton pumping. A diverse array of terminal oxidases is present throughout the *Sulfolobales* and appear to relate to the mode of growth for each organism (**Table 5**). The DoxBCE complex is conserved throughout the *Sulfolobales*, including in the obligate anaerobe *Stygiolobus azoricus*, and it appears to sometimes co-purify with the DoxDA subunits of TQO (Purschke *et al.*, 1997). However, SoxABCDD'L and SoxEFGHIM are more varied in their distribution. SoxABCDD'L, which directly pumps protons out of the cell (Gleissner *et al.*, 1997), seems to be associated with aerobic growth. Meanwhile, SoxEFGHIM is absent from known chemolithoautotrophic organisms such as the *Acidianus* spp., and it is associated with heterotrophic growth (Lubben *et al.*, 1994). An additional quinol oxidase, SoxLN-CbsAB, has been identified in *A. ambivalens* and is highly similar to the complex III cytochrome *bc*₁ (Bandeiras *et al.*, 2009), but the electron acceptor of this complex has yet to be definitively established. There is a wide range of binding affinities and kinetic parameters for these structures (Schafer *et al.*, 2001), and it is possible that

the kinetics of quinol oxidation plays a key role in energy conservation from inorganic substrates in the *Sulfolobales*.

9. Potential and Current Uses of Thermoacidophiles in Biotechnological Applications

9.1. Challenges in establishing genetic systems in *Sulfolobales* species

A genetic toolbox is essential to study and understand the function of genes and proteins of an organism and it has to include three essential components: i) a strain, which is able to take up foreign DNA, ii) a vector system, allowing for the introduction of genetic material, and iii) a selection system for screening of mutated cells. Much effort has been put into the implementation of established systems from different prokaryotic organisms into the *Sulfolobales*. The main challenge of this objective is the natural growth conditions of this group. The high temperature and acidic environment lead to degradation of antibiotics and their resistance-mediating enzymes, which are mainly used as selectable markers in mesophilic organisms. There have been numerous attempts to adapt the systems to these difficult conditions, but either the attempt failed (Cammarano *et al.*, 1985, Grogan, 1989, Grogan, 1991, Aagaard *et al.*, 1994, Sanz *et al.*, 1994, Ruggero & Londei, 1996, Hjort & Bernander, 2001, Bini *et al.*, 2002, Reilly & Grogan, 2002) or positive results could not be reproduced (Aravalli & Garrett, 1997, Cannio *et al.*, 1998, Cannio *et al.*, 2001). Another more promising approach is the use of auxotrophic systems, where metabolically deficient strains are complemented with a functional gene, which becomes a selectable marker. Because these systems use the endogenous metabolic system for selection, they are not affected by the harsh environmental conditions. Therefore,

most genetic systems in the *Sulfolobales* make use of these auxotrophies for mutant selection, which are discussed in this section.

9.1.1. Transformation

The fundamental prerequisite for the establishment of a genetic tool is a reliable transformation protocol, consisting of a transformable strain and a way to introduce DNA into it. The first transformation in *Sulfolobales* was carried out by Schleper *et al.* in 1992 using electroporation, to test the infectivity of *S. shibatae* virus 1 (SSV1) (see **Section 3**) derived shuttle vector system in *Sa. solfataricus* P1 (Schleper *et al.*, 1992). This protocol has been transferred and improved over the years into various other related strains (Zillig *et al.*, 1993, Arnold *et al.*, 1999, Aucelli *et al.*, 2006). Transformation efficiency was later improved by altering the electroporation procedure as well as introducing a regeneration treatment to the cells after the electric shock (Kurosawa & Grogan, 2005, Albers & Driessen, 2007).

S. acidocaldarius expresses a restriction-modification enzyme, Sua I (Prangishvili *et al.*, 1985), cleaving GGCC sequences lacking N4-methylation on the first cytosine (Grogan, 2003). Therefore, the transformation protocol for *S. acidocaldarius* was significantly improved by methylating these sites on the plasmid DNA (Berkner *et al.*, 2007). Another possibility is to delete Sua1 and use the resulting strain as a host for further experiments (Suzuki & Kurosawa, 2016). While methylation of transformed plasmid DNA in *S. acidocaldarius* is mandatory, it is strain dependent in *Sa. islandicus* and *Sa. solfataricus*, with unmodified transformation possible for *Sa. solfataricus* P1 and *Sa. solfataricus* 98/2 and their derived strains (Stedman *et al.*, 1999,

Albers & Driesssen, 2007). In any case, electroporation has proven to be the most efficient means for the transformation of species used for genetic systems in the *Sulfolobales*.

9.1.2. Genetic stability

The most studied *Sulfolobales* species regarding genetic manipulation are *Sa. solfataricus*, *S. acidocaldarius* and *Sa. islandicus*. In contrast to the other two, *S. acidocaldarius* exhibits only a small number of insertion elements (Grogan *et al.*, 2001). These are highly mobile constructs resulting in an elevated mutation frequency, which is problematic for genetic studies, as whole sections of the genome can be inverted or rearranged within a couple of generations (Redder & Garrett, 2006). These elements are most abundant in *Sa. solfataricus* (Martusewitsch *et al.*, 2000), making it the most unstable genome. For *Sa. islandicus*, it depends on the strain, some of which have low mutation frequencies (Berkner & Lipps, 2008), making them preferable for studies of this species.

9.1.3. Cryptical and virus-based shuttle vectors

A broad spectrum of different viruses and plasmids were discovered for the *Sulfolobales* (reviewed by (Prangishvili *et al.*, 2001, Snyder *et al.*, 2003, Prangishvili & Garrett, 2004, Lipps, 2006), some of which were used to create the first generation of *Saccharolobus/Sulfolobus-E. coli* shuttle vector systems. The first derived genetic tool, used in different laboratories, was pMJ03 that consisted of the virus DNA of SSV1 and parts of the bacterial pUC18 vector (Jonuscheit *et al.*, 2003). An advantage of using SSV1 DNA is that no selectable markers are required, as the plasmid can transfect cultures independently. However, this is also the largest

disadvantage, since viral infection puts the cells under severe stress, causing other problems down the line (e.g. growth retardation or contamination of other close cultures). But with no other selection system available at that time, using virus DNA was the only reliable method for vector spreading.

Another approach was the use of naturally occurring cryptic plasmids as scaffolds for vector systems. The basis is the cryptic plasmids pRN1 and pRN2, which were extracted from *Sa. islandicus* (Keeling *et al.*, 1996, Keeling *et al.*, 1998), with pRN1 as the main plasmid backbone in *S. acidocaldarius* systems (Berkner *et al.*, 2007), and pRN2 in *Sa. islandicus* (Deng *et al.*, 2009). However, these systems require a selectable marker for propagation. As the development of these marker systems advanced, the existing system based on SSV1 was largely replaced by cryptic plasmids, which today form the basis for the most used systems in *S. acidocaldarius* and *Sa. islandicus* (Deng *et al.*, 2009, Berkner *et al.*, 2010, Wagner *et al.*, 2012).

9.1.4. LacS: first directed mutants in *Sulfolobus*

The first targeted mutation in *Sulfolobales* was performed by Blum and co-workers (Worthington *et al.*, 2003), based on a natural *lacS* deficient strain, *Sa. solfataricus* *lacS::IS1217*, that had an insertion in the endogenous *lacS* cluster making it unable to grow on lactose as a carbon source. Therefore, growth on lactose could be used as a selection after electroporation of linear fragments containing the *lacS* gene flanked by the region of the DNA where the *lacS* gene had to be inserted. Later, the strain *Sa. solfataricus* PBL2025 proved to be more efficient as the recipient strain. In this natural mutant, a 50 kB region spanning the locus SSO_3004 to SSO3050

were deleted, including the *lacS* gene (Schelert *et al.*, 2004). Improved transformation protocols and a 7-14 days adaptation time after the electroporation in minimal lactose medium enabled the more frequent use of this strain for genetic studies (Albers & Driessen, 2007). However, it only works for *Sa. solfataricus* and *Sa. islandicus*, since *S. acidocaldarius* is not able to grow on lactose as the sole carbon source. Therefore, other methods were sought for *S. acidocaldarius* on the basis of uracil auxotrophic strains and vector systems containing the *pyrEF* gene cassette as a selection marker.

9.1.5. Genetic system for *S. acidocaldarius*

The generation of markerless deletion mutants in archaea was first established in the *Euryarchaeota Haloferax volcanii* using uracil auxotrophy (Bitan-Banin *et al.*, 2003) and *Thermococcus kodakarensis* using uracil and tryptophan auxotrophic mutants (Sato *et al.*, 2005), as well as in the methanogens *Methanosaerina acetivorans* (Pritchett *et al.*, 2004) and *Methanococcus maripaludis* (Moore & Leigh, 2005). The genes *pyrE* and *pyrF* encode for the enzymes orotatephosphoribosyl transferase and orotidine-5'-monophosphate decarboxylase, respectively, which catalyze the last two steps of the uridine monophosphate synthesis pathway (Grogan & Gunsalus, 1993). Upon deletion of one of these two genes, cells lose the ability to grow without uracil supplementation. This deletion can be induced by exposure to the analog substrate 5'-fluoroorotic acid (5-FOA), which is metabolized to cytotoxic products, forcing mutations in the *pyrEF* cluster and generating auxotrophic colonies (Grogan, 1991, Kondo *et al.*, 1991). These generated deficient mutants were first used to test horizontal marker transfer and homologous recombination in *S. acidocaldarius* (Grogan, 1991, Kurosawa & Grogan, 2005).

S. acidocaldarius is capable of recombining linear DNA fragments into its genome via site specific interactions during rapid growth phases (Grogan & Stengel, 2008). The efficiency of the recombination rises proportionally with the length of the fragments used, with 10-30 nt as the minimum length (Kurosawa & Grogan, 2005). Additionally, attachment of short flanking sequences to a selectable marker (e.g. *pyrEF*) could lead to integration of the marker into a gene of interest (Sakofsky *et al.*, 2011), similar to a technique used in *S. cerevisiae* (Kelly *et al.*, 2001) or the one previously described to obtain *Sa. solfataricus lacS::IS1217* (Worthington *et al.*, 2003). Wagner *et al.* were able to generate a *pyrE* deficient mutant, called MW001, derived from *S. acidocaldarius* DSM639 using this approach (Wagner *et al.*, 2012). *S. acidocaldarius* MW001 contains a deletion of 322 bp (91-412 bp) in the *pyrE* gene and only grows in medium supplemented with uracil. In contrast to the typically used 5-FOA or UV light treatment (Grogan, 1991), this method ensures a low probability for additional mutations in the genome.

9.1.6. Generation of markerless deletion mutants in *S. acidocaldarius*

The general idea of this method is an integration of *pyrEF* into a deficient strain via homologous recombination of target sequences, which flank the marker cassette and can interact with the region around the gene of interest (GOI) (**Figure 17**). Positive clones can then be isolated in uracil free medium. The *pyrEF* sequence is derived from *Sa. solfataricus* to avoid homologous recombination between the *pyrEF* in the genome and plasmid. In addition to the auxotrophy, *lacS* from *Sa. solfataricus* was introduced into the vector system as a selectable marker allowing for standard blue/white selection with X-Gal staining, with positive clones exhibiting a blue color.

Following this, colonies are treated with 5-FOA and uracil, which imposes selective pressure, leading to a loop out of the *pyrEF* marker to avoid the formation of toxic byproducts. The successful marker deletion can again be additionally tested via blue/white staining, with successfully obtained mutants displaying a white coloring. Depending on the design of the experiment, the GOI can be deleted by two different approaches. The first is by cloning the upstream (US) and downstream (DS) regions around the GOI consecutively next to the *pyrEF* marker cassette. Single-crossover can then occur leading to an integration of the plasmid (**Figure 17 A, intermediate state**). Upon treatment with 5-FOA, single cross-over can occur again, now between either of the two US or DS regions, respectively, leading to either a deletion of the GOI or the regeneration of the wild type genotype (**Figure 17 A**) (Wagner *et al.*, 2009, Wagner *et al.*, 2012). In an alternative approach, a linearized plasmid or linear PCR fragment is used, which allows for a double-crossover. For the second crossover site, a part of the GOI sequence is used and cloned in front of *pyrEF* (**Figure 17, B**). As only the US is present in a merodiploid form, subsequent removal of *pyrEF* via loop out generates only mutated colonies (Wagner *et al.*, 2009, Wagner *et al.*, 2012).

This strategy is useful to test if a gene and its product have essential functions in the cell and, therefore, cannot be deleted. Removal of the *pyrEF* marker cassette by loop out allows for reuse of the marker, resulting in the possibility of multiple gene deletions in a single mutant (Meyer *et al.*, 2011, Henche *et al.*, 2012, Wagner *et al.*, 2012). The *S. acidocaldarius* MW001 strain has so far been used successfully in more than 100 studies.

9.1.7. Generation of markerless deletion mutants in *Sa. islandicus*

The starting point in *Sa. islandicus* was the generation of a *pyrEF* deletion mutant, *Sa. islandicus* E233 (She *et al.*, 2009), as a recipient strain for the generation of markerless deletion mutants (Deng *et al.*, 2009). Deng *et al.* were able to show that their genetic system worked by using the single-crossover (**Figure 17 A**) and double-crossover (**Figure 17 B**) strategies, as in *S. acidocaldarius*. Alternatively, the plasmid is introduced as a linear fragment again, with the US and DS region flanking the *pyrEF* marker on both sites, plus one additional US or DS site, leading to a double cross-over event. Substitution of the GOI with only *pyrEF* (**Figure 17 C**) leads to mutated cells after 5-FOA treatment (Deng *et al.*, 2009). However, the use of *pyrEF* was not successful in other *Sa. islandicus* strains, as the background growth on solid medium is always high (Zhang & Whitaker, 2012). Therefore, additional selectable markers had to be introduced to achieve higher selective pressure.

Zheng *et al.* showed that simvastatin, a thermostable antibiotic, inhibits the growth of *Sa. islandicus* (Zheng *et al.*, 2012). Shuttle vector systems were developed for *Sa. islandicus* 16.4 (Zhang & Whitaker, 2012) and *Sa. islandicus* REY15A (Zheng *et al.*, 2012), based on the resistance mediated by the overexpression of the 3-hydroxy-3-methylglutaryl coenzyme A reductase gene (*hmgA*). It became apparent, however, that simvastatin resistant cultures had fitness issues that could only be counteracted by enriching the mutants in liquid medium, resulting in a time-consuming protocol. Therefore, it was suggested to use the simvastatin selection only as a last resort.

By deleting the *argD* gene encoding for the arginine decarboxylase, Zhang *et al.* were able to generate an agmatine auxotrophic system (Zhang *et al.*, 2013). The methodology follows

the uracil auxotrophy strategy, as it is possible to recover growth by supplying agmatine or expressing *argD* in a vector system. With this positive selectable marker, the problems derived from using *pyrEF* could be solved for *Sa. islandicus* (Zhang *et al.*, 2013). In addition, a second counter selectable genetic marker was introduced. Through inactivation of a putative adenine phosphoribosyltransferase, resistance could be mediated to 6-methylpurine (6-MP), a purine analog (Zhang *et al.*, 2016). Like 5-FOA, the metabolism of 6-MP leads to toxic compounds, which forces the cell to mutate the gene of the catalyzing enzyme from the genome. This system has been successfully used to establish a transposon library, revealing the essential genome of *Sa. islandicus* (Zhang *et al.*, 2018). This counterselection was also later successfully implemented in *Thermococcus barophilus* (Birien *et al.*, 2018). **Table 6** summarizes the most frequently cited and applied methods of gene disruption and deletion.

9.1.8. Expression vectors: promoters and tags

The combination of the development of selectable markers and recipient strains stimulated efforts to create vectors for homologous protein expression, using different promoters and protein tags. Several attempts were made, but none were useful for high level expression of proteins or for the study of promoters using reporter genes (Aagaard *et al.*, 1996, Elferink *et al.*, 1996, Aravalli & Garrett, 1997, Cannio *et al.*, 1998). The first stable system for homologous expression and tagging of proteins was developed in *Sa. solfataricus* by Albers and co-workers (Jonuscheit *et al.*, 2003, Albers *et al.*, 2006). As previously mentioned, pMJ03 was designed using the SSV1 virus and pUC18 from *E. coli* (Jonuscheit *et al.*, 2003). In this work, the *pyrEF* complementation served as a selectable marker and the heat inducible promotor of the

chaperonin tf55 α gene were used. The reporter gene was *lacS*, that codes for a β -galactosidase, and the recipient strain was a double *pyrEF/lacS* *Sa. solfataricus* mutant. Under heat shock conditions (shift from 75°C to 88°C), an increase of greater than 10-fold gene expression was seen, measured by Northern blot analyses and activity assays of the enzyme. The promotor also had a strong basal expression. This vector was stable for 40-60 generations when cells were maintained in uracil, but propagation of transformants without selective pressure led to loss of the plasmid (Jonuscheit *et al.*, 2003).

Subsequently, this vector was modified. A sugar-inducible promoter (D-arabinose) was added instead of the tf55 α promotor along with cloning sites that allowed for the exchange of *lacS* for a gene of interest. Different tags for protein purification and detection, such as 6x-His or Strep, were also added, leading to the pSVA plasmids set (**Table 7**). The strength of the D-Ara promoter was tested using LacS as a reporter resulting in an increase in activity of 13-fold when D-arabinose was added to the medium. The amount of protein obtained was similar to that of the tf55 α promoter, but without the previously seen basal expression levels. The low basal expression can prevent adverse effects of high expression of proteins on growth before induction. While the His tag resulted in a 99% homogeneity rate in purification, Strep-tagged proteins co-eluted with a carboxylase from *Sa. solfataricus* when low yields of the recombinant protein were obtained (Albers *et al.*, 2006).

An advantage of pMJ03 and derived plasmids was that they were self-spreading, but the production of virus particles had an adverse effect on the transformed cells. A breakthrough came with the use of the plasmid pRN1 in *S. acidocaldarius* (Berkner *et al.*, 2007). It was stable in different *Sulfolobus* species and could be easily selected for in *S. acidocaldarius* using the

pyrEF marker cassette. Several promotors were tested including those from *tf55α dps*, *lacS*, *mal*, *gdhA* and *sac7d* (Berkner *et al.*, 2010). The promoter from the maltose binding protein (*mal*) showed low basal activity and increased expression in the presence of maltose or dextrin in the medium, leading to the plasmid pCmalLacS. The other promoters showed low (*dps*, *lacS*) or high constitutive expression (*gdhA* and *sac7d*) and were left aside. The promoter from *copMA* was also tested but exhibited even lower expression levels than the maltose promoter (Wagner *et al.*, 2012).

In a recent study, a D-xylose/L-arabinose promoter was tested and showed less basal activity than the widely used maltose inducible promoter (van der Kolk *et al.*, 2020). Furthermore, this promoter is also D-arabinose inducible, and can be used as a strong inducer similar to IPTG in *E. coli* since D-arabinose is not a growth substrate for *S. acidocaldarius* (van der Kolk *et al.*, 2020). This study led to the availability of several expression plasmids for *S. acidocaldarius*. Using FX cloning (Geertsma, 2013) for introduction of the desired gene, it is possible to choose between different promoters (D-xylose/L-arabinose /D-arabinose and maltose inducible) and several tags for proteins (StrepII, His, HA, His+Strep, etc.), either in the N-terminal or C-terminal region of the protein (**Table 7**) (van der Kolk *et al.*, 2020).

9.1.9. Mutants via CRISPR Cas and gene silencing

CRISPR systems and spacer acquisition are discussed in **Section 3**. CRISPR Cas systems are classified into three groups so far: type I, associated with Cas3 protein, type II, associated with Cas9, and type III, associated with Cas10. Type I and II recognizes DNA via PAM sequences in the protospacer in the target genome. Type III does not need a PAM sequence, but

instead a seed sequence, corresponding to a mismatch in the 5' end of the crRNA (van der Oost J, 2014).

In the *Sulfolobales*, the idea of using the proper CRISPR-Cas system was first explored in *Sa. islandicus*. Li et al. used the native type IA and IIIB systems of this organism to generate mutants for *lacS*, so the phenotype could be easily tested with X-gal (Li et al., 2016). The pSe-Rp plasmid was used as backbone and an artificial CRISPR array was introduced. The array consisted of two repeat sequences, flanking a spacer designed based on a protospacer sequence from the target gene. Also, the plasmid contained a donor region that does not exhibit DNA interference activity. Mutagenesis through deletion, insertion or point mutation is achieved via recombination of the donor region with the genomic DNA (Li et al., 2016).

The CRISPR RNA (crRNA) generated from the CRISPR array of the plasmid guides the native CRISPR-Cas system to self-targeting DNA from wild type cells, killing them but not the mutants, which accomplished recombination with the donor DNA. Using a plasmid containing the donor sequence and the CRISPR array led to better results than performing co-transformation using a CRISPR plasmid and a short DNA fragment as donor sequence (Li et al., 2016).

In *Sa. solfataricus*, on the other hand, type III CRISPR Cas was used to develop silencing via RNA interference (Zebec et al., 2016). *Sa. solfataricus* has two type IIIB CRISPR complexes, both targeting RNA, and one of them also DNA (Zhang et al., 2016). A plasmid containing a mini-CRISPR array with a spacer designed to target mRNA from β -galactosidase was used to demonstrate that 50% of gene silencing is possible to achieve using the native CRISPR Cas system from *Sa. solfataricus* (Zebec et al., 2014). In later work, silencing up to 90% could be accomplished using a CRISPR array containing five different spacers from the

same gene, in this case α -amylase (Zebec *et al.*, 2016). Lower levels of silencing were achieved using one to three spacers, resulting in 35% to 82% gene silencing. Since one of the type IIIB complexes also targets DNA, the protospacers in the genome were chosen to have a flanking region that matches with the 5' end handle of the crRNA. This inhibits DNA targeting since CRISPR-Cas system uses the repeat sequences to “protect itself” by recognizing the cells own genomic DNA (Manica & Schleper, 2013).

CRISPR has also been used to edit rod-shaped virus 2 (SIRV-2) from *Sa. islandicus*, using the archaeon as a host and its endogenous CRISPR system as machinery (Mayo-Muñoz *et al.*, 2018). This method was also used to investigate the core genome of the virus, generating knock out mutants, useful for probing the details of the infection process. In general, CRISPR Cas systems are powerful genetic tools, which have had a huge impact in the life science community (van der Oost J, 2014, Plagens *et al.*, 2015, Mougios *et al.*, 2016, Quehenberger *et al.*, 2017), and also show promise for studying and engineering thermoacidophiles.

9.2. The potential of thermoacidophiles as metabolic engineering platforms

As outlined in **Section 8**, representatives of the *Sulfolobales* have been examined for their autotrophic and heterotrophic life styles. They have enormous metabolic versatility that differs from species to species, with respect to their growth on a variety of carbon sources that include complex polymers (e.g. polysaccharides, proteins, lipids), monomers (e.g. carbohydrates, amino acids, fatty acids, alcohols, aldehydes), and CO₂. In addition, comprehensive biochemical and functional genomics data are available for members of the *Sulfolobales* (Bräsen *et al.*, 2014), including ¹³C NMR flux analysis for different carbon sources (Nunn *et al.*, 2010). Furthermore,

systems biology (www.sulfosys.com (Zaparty et al., 2009)), genome scale stoichiometric (FBA) (Ulas et al., 2012), and detailed kinetic models (e.g. for gluconeogenesis) (Kouril et al., 2013, Kouril et al., 2013, Kouril et al., 2017) have been established for one of the model *Sulfolobales*, *Sa. solfataricus*.

Critical for metabolic engineering is the availability of advanced genetic tools, which are described above for several species in the *Sulfolobales* and include in-frame markerless deletion mutants, ectopic integration of foreign DNA, and provides a homologous expression system (Wagner et al., 2014) (see **Section 8**). Thus, *S. acidocaldarius*, *Sa. islandicus*, and *Sa. solfataricus* are all a potential host “chassis” on which to build biosynthetic designs of increasing complexity, although *S. acidocaldarius*’s genetic stability might be an advantage. Another important criterion for metabolic engineering and application in biotechnology is ease of cultivation under aerobic conditions. Complex and minimal media have been described, e.g. for *S. acidocaldarius*, and high cell density cultivation has been established for *S. shibatae* and *S. acidocaldarius* (Quehenberger et al., 2017, Schocke et al., 2019, Quehenberger et al., 2020). The current genetic systems do require further improvement. In particular, the integration of larger gene clusters and the development of new regulatory strategies are needed to fully realize the biotechnological potential of these thermoacidophiles (Crosby et al., 2019).

9.2.1. Thermoacidophile Biotechnology

As mentioned above, by their very nature, thermoacidophiles are robust microorganisms that can handle industrial processing conditions well and therefore offer potential advantages as metabolic engineering hosts over more established but less extremophilic species, such as *E. coli*

and *S. cerevisiae* (Crosby *et al.*, 2019). Their cytosolic enzymes are adapted to high temperature and neutral pH, whereas their extracellular enzymes, such as amylases, cellulases and lipases, are also adapted to low pH. Thus, the properties of thermoacidophilic biocatalysts are consistent with process schemes used in lignocellulosic biomass pre-treatments, which are typically done at high temperatures and low pH. The utilization of thermoacidophiles and their enzymes offers certain benefits for industrial biotechnology (Turner *et al.*, 2007, Hess, 2008, Zeldes *et al.*, 2015, Straub *et al.*, 2018): at high temperature reaction rates increase and as does substrate accessibility for biopolymers such as starch and lignocellulosic carbohydrates, thereby enhancing biomass conversion. Since substrate solubility improves at higher temperatures, this enables mixing of otherwise viscous slurries. Furthermore, the energy input for cooling steps in bioreactors and thus production costs can be reduced. Under thermal conditions, volatile products can be removed through gas stripping and evaporation facilitating product recovery. As such, expensive distillation steps as well as inhibition by toxic products can be minimized, allowing for novel design “one-pot” strategies (Zeldes *et al.*, 2018). Particularly important, microbial contamination is negligible at high temperatures and low pH, so that the use of antibiotics and the need for pharmaceutical-like processing can be avoided (Marhuenda-Egea & Bonete, 2002, Champdore *et al.*, 2007, Quehenberger *et al.*, 2017, Cabrera & Blamey, 2018).

There are also some disadvantages and challenges to overcome to fully realize the biotechnological potential of thermoacidophiles as industrial microorganisms. As previously described, central metabolic pathways (e.g. for lipid or glycerol degradation) are still not well understood and further work is needed to unravel the metabolic complexity of promising representatives of the *Sulfolobales*. In particular, networks with regulation at the gene and

protein level require further basic research, and only a few transcriptional regulators have been investigated, and regulation via post-translational modification is also not well understood (See **Section 4**). Processes that truly exploit thermoacidophily remain to be developed, although advances in thermoacidophile genetic motivate such efforts.

9.3. Biomining Applications of Thermoacidophiles

The importance of acidophilic organisms in the breakdown of sulfidic ore has been known for many years. Acid mine drainage is a byproduct of acidophiles at work on pyritic mine waste and provides an environmental backdrop for studying the mechanism of biological oxidation (Schippers *et al.*, 1996). However, the same mechanism can be leveraged to extract base and precious metals from sulfidic ores through bioleaching operations. An important distinction to note is that bioleaching refers specifically to the dissolution of metals by bacteria and archaea, while biooxidation simply refers to the oxidation of metal and non-metal substrates by the same organisms.

At first glance, one might suspect that the acidophilic organisms directly attack the solid ore. However, closer inspection of the mechanism shows that the dissolution of both pyrite and other metal sulfides is an indirect result of biooxidation by acidophilic bacteria (**Figure 18**). The ore undergoes electrophilic attack in the presence of ferric ions (Fe^{3+}) and protons (Sand *et al.*, 2001). Other ions have also been proposed to facilitate this initial attack, including copper and silver (Hiroyoshi *et al.*, 2000, Zhao *et al.*, 2019). The role of biooxidation in the bioleaching process is then the regeneration of ferrous ions into ferric ions that can again attack the ore and the total oxidation of the freed sulfur to sulfate (Li *et al.*, 2013). Two distinct mechanisms for ore

dissolution underscore the importance of the distribution of sulfur in the system. In the presence of pyrite, thiosulfate is directly generated during the electrophilic attack (**Figure 18 A**). This initiates a cycle of abiotic sulfur reactions, in which greater than 80% of the sulfur product is converted to sulfate (Sand *et al.*, 2001). In contrast, the polysulfide mechanism applies to dissolution of metal sulfide ores, like chalcopyrite (**Figure 18 B**). Here, the sulfur product of the initial attack is hydrogen sulfide (H_2S), which then undergoes a series of abiotic chain elongation reactions to form polysulfides. Ultimately, these polysulfides cyclize to form the thermodynamically stable and water-insoluble S_8 ring (Steudel, 1996), which accounts for 90-99% of the final sulfur product depending on the metal sulfide species (Sand *et al.*, 2001). While pyrite is the dominant form of metal sulfide ores, mining streams generally contain a mixture of these various crystal structures (Neale *et al.*, 2009), and thus these mechanisms exist simultaneously.

A key issue associated with the polysulfide mechanism generating solid cyclic sulfur is the passivation of the ore's surface (Zhao *et al.*, 2019). Passivation is the formation of an inhibitory film on the surface of the ore that prevents further electrophilic attack (Klauber, 2008). In the case of chalcopyrite, a passivating layer of solid sulfur coats the ore and significantly slows the dissolution rate. Another component of this passivating effect is the formation of jarosites, complex ferric sulfate compounds, that is only an issue with long leaching times. However, the slowed dissolution rate caused by passivating sulfur may allow for the accumulation of jarosites, and both modes of inhibition are seen in chalcopyrite bioleaching (Klauber, 2008).

There are two primary formats for industrial-scale bioleaching. Heap leaching, as the name implies, is the open-air extraction of metals from large heaps of mine tailings. Leaching solution is percolated throughout the pile, with the metal-laden run-off collected for downstream processing (Schlitt, 2006). The capital investment of these operations is generally low and uses low-grade ore to maximize yield from mining operations. These heaps are often self-inoculated and contain a consortium of acidophilic organisms, including mesophilic *Acidithiobacillus* spp., moderately thermophilic *Sulfolobacillus* spp., and extremely thermophilic *Acidianus* spp. and *Metallosphaera* spp. (Pradhan *et al.*, 2008). Despite the open-air environment of the heaps, the highly exothermic oxidation of sulfur results in heap temperatures reaching as high as 81°C, which supports the growth of the *Sulfolobales* (Pradhan *et al.*, 2008).

The second format is a traditional bioreactor. While the capital cost of this setup is much higher than that of heap leaching, the more controlled environment and improved contact area dramatically reduces the leaching time. While heap operations run for months to years (Schlitt, 2006), bioreactor leaching has a retention time on the order of days (Neale *et al.*, 2009). These bioreactors are often still self-inoculating and contain a complex microbial landscape. Notably, the acidic conditions provide a natural barrier to contamination, and this barrier is further enhanced when working at the high temperatures of the *Sulfolobales*.

Bioleaching at the high temperatures required by the *Sulfolobales* offers distinct advantages. Given the large exotherm associated with sulfur oxidation, the reactor does not need to be heated to maintain the temperatures necessary for *Sulfolobales*' growth. In fact, cooling is necessary to maintain the constant temperatures (Neale *et al.*, 2009), and at higher temperatures less cooling is necessary. Furthermore, chalcopyrite dissolution kinetics are significantly

improved at higher temperatures (Watling, 2006). In a 10-day lab-scale bioleaching reactor, around 70% copper recovery was achieved with the mesophilic *Acidithiobacillus ferrooxidans* and roughly 85% copper recovery was achieved with the moderate thermophile *Sulfobacillus* (Mousavi *et al.*, 2005). At pilot scale, a thermoacidophile bioreactor containing *Acidianus*, *Metallosphaera*, and *Sulfolobus* spp. achieved 95% copper recovery (Neale *et al.*, 2009).

As understanding of the mechanisms for iron and sulfur oxidation improves, engineering an optimized bioleaching organism presents an intriguing possibility. In particular, this could overcome the obstacle of surface passivation caused by the accumulation of elemental sulfur and jarosites. Indeed, one means of controlling the passivation of ore is to lower the reduction potential of the reactor. This process controls the ratio of $\text{Fe}^{3+}/\text{Fe}^{2+}$ ions in order to limit the rate of elemental sulfur and jarosite formation. A variety of approaches to control the redox potential of the system have been explored, ranging from adding reagents to form new redox couples, controlling dissolved oxygen levels, and adjusting the microbe composition (Zhao *et al.*, 2019). This last option points towards the possibility of an optimized bioleaching organism that manages the redox potential through the relative rates of sulfur and iron oxidation. Efforts to generate an optimized strain of *M. sedula* through laboratory evolution had moderate success in this regard (McCarthy *et al.*, 2018). While natural evolution of the ability to breakdown chalcopyrite may provide novel insights, tailoring an organism to mitigate surface passivation may require a more controlled approach. As such, the genetic tools available to select members of the *Sulfolobales* provide a promising platform to expand their bioleaching capabilities.

10. Conclusions

The scientific and technological potential of the *Sulfolobales* has come a long way since their isolation more than a half century ago. During this time, molecular biology and genomics came of age, with all of the associated tools that can be brought to bear in understanding the microbiology of these thermoacidophiles. Here, the goal was to provide some historical perspective as well as to give an up-to-date overview of where the world of the *Sulfolobales* stands. Despite the length and breadth of this review, we have likely inadvertently left out important contributions to the field and thank all of those who have studied and reported on facets of these interesting microorganisms.

Acknowledgments

This work was supported by grants to R.M.K. from the U.S. Air Force Office of Sponsored Research (AFOSR) (FA9550-17-1-0268) and the U.S. National Science Foundation (CBET-1802939). J.A.C. and D.J.W. acknowledges support from an U.S. National Institutes of Health Biotechnology Traineeships (T32 GM008776-16, T32 GM133366-01). T.E.F.Q. was supported by the Deutsche Forschungsgemeinschaft (German Research Foundation) with an Emmy Noether grant (411069969). A.J. and J.B. were supported by the grant HotAcidFactory (031B0848C) from the BMBF (Federal Ministry of Education and Research, Germany). P.N. received funding from the VW Foundation on Momentum grant 94933. L.S. was supported by the Mercator foundation for support with a Mercur startup grant (Pr-2013-0010) and by the BMBF grant HotSysAPP, 031L0078A within the e:Bio2 funding initiative. L. Sch. was

supported by the BMBF grant HotAcidFACTORY, 031B0848A. B.S. and C.B. acknowledge funding from the VW foundation on the “Experiment?” LipideDivide grant 96725.

References

Aagaard C, Phan H, Trevisanato S & Garrett RA (1994) A spontaneous point mutation in the single 23S rRNA gene of the thermophilic archaeon *Sulfolobus acidocaldarius* confers multiple drug resistance. *J Bacteriol* **176**: 7744-7747.

Aagaard C, Leviev I, Aravalli RN, Forterre P, Prieur D & Garrett RA (1996) General vectors for archaeal hyperthermophiles: Strategies based on a mobile intron and a plasmid. *FEMS Microbiol Rev* **18**: 93-104.

Ahmed H, Ettema Thijs JG, Tjaden B, Geerling Ans CM, van der Oost J & Siebers B (2005) The semi-phosphorylative Entner–Doudoroff pathway in hyperthermophilic archaea: a re-evaluation. *Biochem J* **390**: 529-540.

Albers S-V & Driessens AJM (2007) Conditions for gene disruption by homologous recombination of exogenous DNA into the *Sulfolobus solfataricus* genome. *Archaea* **2**: 145-149.

Albers S-V & Pohlschröder M (2009) Diversity of archaeal type IV pilin-like structures. *Extremophiles* **13**: 403-410.

Albers S-V & Meyer BH (2011) The archaeal cell envelope. *Nat Rev Microbiol* **9**: 414-426.

Albers S-V, Szabó Z & Driessens AJM (2003) Archaeal homolog of bacterial type IV prepilin signal peptidases with broad substrate specificity. *J Bacteriol* **185**: 3918-3925.

Albers S-V, Koning SM, Konings WN & Driessens AJ (2004) Insights into ABC transport in archaea. *J Bioenerg Biomembr* **36**: 5-15.

Albers SV & Jarrell KF (2015) The archaeum: How Archaea swim. *Front Microbiol* **6**: 1-12.

Albers SV, Konings WN & Driessens AJ (1999) A unique short signal sequence in membrane-anchored proteins of Archaea. *Mol Microbiol* **31**: 1595-1596.

Albers SV, Koning SM, Konings WN & Driessens AJ (2004) Insights into ABC transport in archaea. *J Bioenerg Biomembr* **36**: 5-15.

Albers SV, Jonuscheit M, Dinkelaker S, Urich T, Kletzin A, Tampé R, Driessens AJM & Schleper C (2006) Production of recombinant and tagged proteins in the hyperthermophilic archaeon *Sulfolobus solfataricus*. *Appl Environ Microbiol* **72**: 102-111.

Albi T & Serrano A (2016) Inorganic polyphosphate in the microbial world. Emerging roles for a multifaceted biopolymer. *World J Microbiol Biotechnol* **32**: 27.

Amend JP & Shock EL (2001) Energetics of overall metabolic reactions of thermophilic and hyperthermophilic Archaea and Bacteria. *FEMS Microbiol Rev* **25**: 175-243.

Anderson I, Scheuner C, Goker M, Mavromatis K, Hooper SD, Porat I, Klenk HP, Ivanova N & Kyrpides N (2011) Novel insights into the diversity of catabolic metabolism from ten haloarchaeal genomes. *PLoS One* **6**: e20237.

Anjum RS, Bray SM, Blackwood JK, *et al.* (2015) Involvement of a eukaryotic-like ubiquitin-related modifier in the proteasome pathway of the archaeon *Sulfolobus acidocaldarius*. *Nat Commun* **6**: 8163.

Ao X, Li Y, Wang F, Feng M, Lin Y, Zhao S, Liang Y & Peng N (2013) The *Sulfolobus* initiator element is an important contributor to promoter strength. *J Bacteriol* **195**: 5216-5222.

Aravalli RN & Garrett RA (1997) Shuttle vectors for hyperthermophilic archaea. *Extremophiles* **1**: 183-191.

Aravind L (1999) DNA-binding proteins and evolution of transcription regulation in the archaea. *Nucl Acids Res* **27**: 4658-4670.

Archibald JM, Logsdon JM & Doolittle WF (1999) Recurrent paralogy in the evolution of archaeal chaperonins. *Curr Biology* **9**: 1053-S1056.

Arnold HP, Ziese U & Zillig W (2000) SNDV, a novel virus of the extremely thermophilic and acidophilic archaeon *Sulfolobus*. *Virology* **272**: 409-416.

Arnold HP, She Q, Phan H, Stedman K, Prangishvili D, Holz I, Kristjansson JK, Garrett R & Zillig W (1999) The genetic element pSSVx of the extremely thermophilic crenarchaeon *Sulfolobus* is a hybrid between a plasmid and a virus. *Mol Microbiol* **34**: 217-226.

Athukoralage JS, Rouillon C, Graham S, Grüsschow S & White MF (2018) Ring nucleases deactivate type III CRISPR ribonucleases by degrading cyclic oligoadenylate. *Nature* **562**: 277-280.

Athukoralage JS, Graham S, Rouillon C, Grüsschow S, Czekster CM & White MF (2020) The dynamic interplay of host and viral enzymes in type III CRISPR-mediated cyclic nucleotide signalling. *eLife* **9**: e55852.

Athukoralage JS, McMahon SA, Zhang C, Grüsschow S, Graham S, Krupovic M, Whitaker RJ, Gloster TM & White MF (2020) An anti-CRISPR viral ring nuclease subverts type III CRISPR immunity. *Nature* **577**: 572-575.

Atomi H, Matsumi R & Imanaka T (2004) Reverse gyrase is not a prerequisite for hyperthermophilic life. *J Bacteriol* **186**: 4829-4833.

Aucelli T, Contursi P, Girfoglio M, Rossi M & Cannio R (2006) A spreadable, non-integrative and high copy number shuttle vector for *Sulfolobus solfataricus* based on the genetic element pSSVx from *Sulfolobus islandicus*. *Nucl Acids Res* **34**: e114.

Auernik KS & Kelly RM (2008) Identification of components of electron transport chains in the extremely thermoacidophilic crenarchaeon *Metallosphaera sedula* through iron and sulfur compound oxidation transcriptomes. *Appl Environ Microbiol* **74**: 7723-7732.

Auernik KS, Maezato Y, Blum PH & Kelly RM (2008) The genome sequence of the metal-mobilizing, extremely thermoacidophilic archaeon *Metallosphaera sedula* provides insights into bioleaching-associated metabolism. *Appl Environ Microbiol* **74**: 682-692.

Awano T, Wilming A, Tomita H, Yokooji Y, Fukui T, Imanaka T & Atomi H (2014) Characterization of two members among the five ADP-Forming Acyl Coenzyme A (Acyl-CoA) Synthetases reveals the presence of a 2-(Imidazol-4-yl)Acetyl-CoA Synthetase in *Thermococcus kodakarensis*. *J Bacteriol* **196**: 140-147.

Baes R, Lemmens L, Mignon K, Carlier M & Peeters E (2020) Defining heat shock response for the thermoacidophilic model crenarchaeon *Sulfolobus acidocaldarius*. *Extremophiles*.

Baker-Austin C & Dopson M (2007) Life in acid: pH homeostasis in acidophiles. *Trends Microbiol* **15**: 165-171.

Bandeiras TM, Refojo PN, Todorovic S, Murgida DH, Hildebrandt P, Bauer C, Pereira MM, Kletzin A & Teixeira M (2009) The cytochrome ba complex from the thermoacidophilic crenarchaeote *Acidianus ambivalens* is an analog of bc(1) complexes. *Biochim Biophys Acta* **1787**: 37-45.

Bar-Nun S & Glickman MH (2012) Proteasomal AAA-ATPases: Structure and function. *Biochim Biophys Acta - Mol Cell Res* **1823**: 67-82.

Barrangou R & Horvath P (2017) A decade of discovery: CRISPR functions and applications. *Nat Microbiol* **2**: 17092.

Barrangou R, Fremaux C, Deveau H, Richards M, Boyaval P, Moineau S, Romero DA & Horvath P (2007) CRISPR provides acquired resistance against viruses in prokaryotes. *Science* **315**: 1709-1712.

Bathe S & Norris PR (2007) Ferrous iron- and sulfur-induced genes in *Sulfolobus metallicus*. *Appl Environ Microbiol* **73**: 2491-2497.

Bautista MA, Black JA, Youngblut ND & Whitaker RJ (2017) Differentiation and structure in *Sulfolobus islandicus* rod-shaped virus populations. *Viruses* **9**: 120.

Bell SD, Kosa PL, Sigler PB & Jackson SP (1999) Orientation of the transcription preinitiation complex in Archaea. *Proc Natl Acad Sci USA* **96**: 13662-13667.

Bell SD, Brinkman AB, Van Der Oost J & Jackson SP (2001) The archaeal TFIIE α homologue facilitates transcription initiation by enhancing TATA-box recognition. *EMBO Rep* **2**: 133-138.

Bell SD, Botting CH, Wardleworth BN, Jackson SP & White MF (2002) The interaction of Alba, a conserved archaeal chromatin protein, with Sir2 and its regulation by acetylation. *Science* **296**: 148-151.

Benelli D, Maone E & Londei P (2003) Two different mechanisms for ribosome/mRNA interaction in archaeal translation initiation. *Mol Microbiol* **50**: 635-643.

Berg IA, Kockelkorn D, Buckel W & Fuchs G (2007) A 3-hydroxypropionate/4-hydroxybutyrate autotrophic carbon dioxide assimilation pathway in Archaea. *Science* **318**: 1782-1786.

Berg IA, Ramos-Vera WH, Petri A, Huber H & Fuchs G (2010) Study of the distribution of autotrophic CO₂ fixation cycles in Crenarchaeota. *Microbiologyopen* **156**: 256-269.

Berg IA, Kockelkorn D, Ramos-Vera WH, Say RF, Zarzycki J, Hugler M, Alber BE & Fuchs G (2010) Autotrophic carbon fixation in archaea. *Nature Microbial Rev* **8**: 447-460.

Berkner S & Lipps G (2008) Mutation and reversion frequencies of different *Sulfolobus* species and strains. *Extremophiles* **12**: 263-270.

Berkner S, Grogan D, Albers S-V & Lipps G (2007) Small multicopy, non-integrative shuttle vectors based on the plasmid pRN1 for *Sulfolobus acidocaldarius* and *Sulfolobus solfataricus*, model organisms of the (cren-)archaea. *Nucl Acids Res* **35**: e88.

Berkner S, Włodkowski A, Albers S-V & Lipps G (2010) Inducible and constitutive promoters for genetic systems in *Sulfolobus acidocaldarius*. *Extremophiles* **14**: 249-259.

Bernander R & Poplawski A (1997) Cell cycle characteristics of thermophilic Archaea. *J Bacteriol* **179**: 4963-4969.

Bertoldo C, Dock C & Antranikian G (2004) Thermoacidophilic microorganisms and their novel biocatalysts. *Eng Life Sci* **4**: 521-532.

Bettstetter M, Peng X, Garrett RA & Prangishvili D (2003) AFV1, a novel virus infecting hyperthermophilic archaea of the genus *Acidianus*. *Virology* **315**: 68-79.

Bhoobalan-Chitty Y, Johansen TB, Di Cianni N & Peng X (2019) Inhibition of Type III CRISPR-Cas Immunity by an archaeal virus-encoded anti-CRISPR Protein. *Cell* **179**: 448-458.e411.

Bigotti MG & Clarke AR (2005) Cooperativity in the Thermosome. *J Mol Biol* **348**: 13-26.

Bini E, Dikshit V, Dirksen K, Drozda M & Blum P (2002) Stability of mRNA in the hyperthermophilic archaeon *Sulfolobus solfataricus*. *RNA* **8**: 1129-1136.

Birien T, Thiel A, Henneke G, Flament D, Moalic Y & Jebbar M (2018) Development of an effective 6-methylpurine counterselection marker for genetic manipulation in *Thermococcus barophilus*. *Genes* **9**: 77.

Bitan-Banin G, Ortenberg R & Mevarech M (2003) Development of a gene knockout system for the halophilic archaeon *Haloferax volcanii* by use of the *pyrE* gene. *J Bacteriol* **185**: 772-778.

Bize A, Karlsson EA, Ekefjärd K, Quax TEF, Pina M, Prevost M-C, Forterre P, Tenaillon O, Bernander R & Prangishvili D (2009) A unique virus release mechanism in the Archaea. *Proc Natl Acad Sci USA* **106**: 11306-11311.

Blombach F, Matelska D, Fouqueau T, Cackett G & Werner F (2019) Key concepts and challenges in archaeal transcription. *J Mol Biol* **431**: 4184-4201.

Blombach F, Salvadori E, Fouqueau T, *et al.* (2015) Archaeal TFE α / β is a hybrid of TFIIE and the RNA polymerase III subcomplex hRPC62/39. *eLife* **4**: e08378.

Blum H, Zillig W, Mallok S, Domdey H & Prangishvili D (2001) The genome of the archaeal virus SIRV1 has features in common with genomes of eukaryal viruses. *Virology* **281**: 6-9.

Blumentals, II, Itoh M, Olson GJ & Kelly RM (1990) Role of polysulfides in reduction of elemental sulfur by the hyperthermophilic archaeabacterium *Pyrococcus furiosus*. *Appl Environ Microbiol* **56**: 1255-1262.

Bolduc B, Shaughnessy DP, Wolf YI, Koonin EV, Roberto FF & Young M (2012) Identification of novel positive-strand RNA viruses by metagenomic analysis of archaea-dominated Yellowstone hot springs. *J Virology* **86**: 5562-5573.

Bolotin A, Quinquis B, Sorokin A & Ehrlich SD (2005) Clustered regularly interspaced short palindrome repeats (CRISPRs) have spacers of extrachromosomal origin. *Microbiologyopen* **151**: 2551-2561.

Bonneau R, Facciotti MT, Reiss DJ, *et al.* (2007) A predictive model for transcriptional control of physiology in a free living cell. *Cell* **131**: 1354-1365.

Borges AL, Davidson AR & Bondy-Denomy J (2017) The discovery, mechanisms, and evolutionary impact of anti-CRISPRs. *Annu Rev Virology* **4**: 37-59.

Boyd ES, Fecteau KM, Havig JR, Shock EL & Peters JW (2012) Modeling the habitat range of phototrophs in Yellowstone National Park: Toward the development of a comprehensive fitness landscape. *Front Microbiol* **3**: 221.

Boyd ES, Pearson A, Pi Y, Li W-J, Zhang YG, He L, Zhang CL & Geesey GG (2011) Temperature and pH controls on glycerol dibiphytanyl glycerol tetraether lipid composition in the hyperthermophilic crenarchaeon *Acidilobus sulfurireducens*. *Extremophiles* **15**: 59-65.

Boyd ES, Jackson RA, Encarnacion G, Zahn JA, Beard T, Leavitt WD, Pi Y, Zhang CL, Pearson A & Geesey GG (2007) Isolation, characterization, and ecology of sulfur-respiring crenarchaea inhabiting acid-sulfate-chloride-containing geothermal springs in Yellowstone National Park. *Appl Environ Microbiol* **73**: 6669-6677.

Brenneis M, Hering O, Lange C & Soppa J (2007) Experimental characterization of cis-acting elements important for translation and transcription in halophilic Archaea. *PLoS Genet* **3**: e229.

Brito JA, Sousa FL, Stelter M, Bandeiras TM, Vonrhein C, Teixeira M, Pereira MM & Archer M (2009) Structural and functional insights into sulfide:quinone oxidoreductase. *Biochemistry* **48**: 5613-5622.

Brito JA, Sousa FL, Stelter M, Bandeiras TM, Vonrhein C, Teixeira M, Pereira MM & Archer M (2009) Structural and functional insights into sulfide:quinone oxidoreductase. *Biochemistry* **48**: 5613-5622.

Brock TD, Brock KM, Belly RT & Weiss RL (1972) *Sulfolobus*: a new genus of sulfur-oxidizing bacteria living at low pH and high temperature. *Arch Mikrobiol* **84**: 54-68.

Brouns SJ, Jore MM, Lundgren M, Westra ER, Slijkhuis RJ, Snijders AP, Dickman MJ, Makarova KS, Koonin EV & Van Der Oost J (2008) Small CRISPR RNAs guide antiviral defense in prokaryotes. *Science* **321**: 960-964.

Brouns SJ, Walther J, Snijders APL, et al. (2006) Identification of the missing links in prokaryotic pentose oxidation pathways. *J Biol Chem* **281**: 27378-27388.

Brown AD & Simpson JR (1972) Water relations of sugar-tolerant yeasts: the role of intracellular polyols. *Journal of general microbiology* **72**: 589-591.

Brown PB & Wolfe GV (2006) Protist genetic diversity in the acidic hydrothermal environments of Lassen Volcanic National Park, USA. *J Eukaryot Microbiol* **53**: 420-431.

Brumfield SK, Ortmann AC, Ruigrok V, Suci P, Douglas T & Young MJ (2009) Particle assembly and ultrastructural features associated with replication of the lytic archaeal virus *Sulfolobus* turreted icosahedral virus. *J Virology* **83**: 5964-5970.

Bräsen C & Schönheit P (2004) Regulation of acetate and acetyl-CoA converting enzymes during growth on acetate and/or glucose in the halophilic archaeon *Haloarcula marismortui*. *FEMS Microbiol Lett* **241**: 21-26.

Bräsen C, Esser D, Rauch B & Siebers B (2014) Carbohydrate metabolism in Archaea: current insights into unusual enzymes and pathways and their regulation. *Microbiol Mol Biol Rev* **78**: 89-175.

Bulaev AG, Kanygina AV & Manolov AI (2017) Genome analysis of *Acidiplasma* sp. MBA-1, a polyextremophilic archaeon predominant in the microbial community of a bioleaching reactor. *Microbiologyopen* **86**: 89-95.

Cabrera MA & Blamey JM (2018) Biotechnological applications of archaeal enzymes from extreme environments. *Biol Res* **51**: 1-15.

Caillat C, Maity S, Miguet N, Roos WH & Weissenhorn W (2019) The role of VPS4 in ESCRT-III polymer remodeling. *Biochem Soc Trans* **47**: 441-448.

Cammarano P, Teichner A, Londei P, Acca M, Nicolaus B, Sanz JL & Amils R (1985) Insensitivity of archaebacterial ribosomes to protein synthesis inhibitors. Evolutionary implications. *EMBO J* **4**: 811-816.

Campbell KM, Kouris A, England W, Anderson RE, McCleskey RB, Nordstrom DK & Whitaker RJ (2017) *Sulfolobus islandicus* meta-populations in Yellowstone National Park hot springs. *Environ Microbiol* **19**: 2334-2347.

Cannio R, Contursi P, Rossi M & Bartolucci S (1998) An autonomously replicating transforming vector for *Sulfolobus solfataricus*. *J Bacteriol* **180**: 3237-3240.

Cannio R, Contursi P, Rossi M & Bartolucci S (2001) Thermoadaptation of a mesophilic hygromycin B phosphotransferase by directed evolution in hyperthermophilic Archaea: Selection of a stable genetic marker for DNA transfer into *Sulfolobus solfataricus*. *Extremophiles* **5**: 153-159.

Cannio R, Di Prizito N, Rossi M & Morana A (2004) A xylan-degrading strain of *Sulfolobus solfataricus*: Isolation and characterization of the xylanase activity. *Extremophiles* **8**: 117-124.

Cannio R, Catara G, Fiume I, Balestrieri M, Rossi M & Palmieri G (2010) Identification of a cell-bound extracellular protease overproduced by *Sulfolobus solfataricus* in peptide-rich media. *Protein & Peptide Lett* **17**: 78-85.

Cao J, Wang Q, Liu T, Peng N & Huang L (2018) Insights into the post-translational modifications of archaeal Sis10b (Alba): lysine-16 is methylated, not acetylated, and this does not regulate transcription or growth. *Mol Microbiol* **109**: 192-208.

Cao X, Koch T, Steffens L, Finkensieper J, Zigann R, Cronan JE & Dahl C (2018) Lipoate-binding proteins and specific lipoate-protein ligases in microbial sulfur oxidation reveal an atypical role for an old cofactor. *eLife* **7**.

Caspi Y & Dekker C (2018) Dividing the archaeal way: The ancient Cdv cell-division machinery. *Front Microbiol* **9**.

Champdore Md, Staiano M, Rossi M & D'Auria S (2007) Proteins from extremophiles as stable tools for advanced biotechnological applications of high social interest. *J R Soc Interface* **4**: 183-191.

Chaston JJ, Smits C, Aragao D, *et al.* (2016) Structural and functional insights into the evolution and stress adaptation of type II Chaperonins. *Structure* **24**: 364-374.

Chaudhury P, Quax TEF & Albers S-V (2018) Versatile cell surface structures of Archaea. *Mol Microbiol* **107**: 298-311.

Chen KY & Morris JC (1972) Kinetics of oxidation of aqueous sulfide by oxygen. *Environ Sci Technol* **6**: 529-537.

Chen L, Brügger K, Skovgaard M, *et al.* (2005) The genome of *Sulfolobus acidocaldarius*, a model organism of the *Crenarchaeota*. *J Bacteriol* **187**: 4992-4999.

Choi KH, Hwang S & Cha J (2013) Identification and characterization of MalA in the maltose/maltodextrin operon of *Sulfolobus acidocaldarius* DSM639. *J Bacteriol* **195**: 1789-1799.

Choi Y-H, Park Y-J, Yoon S-J & Lee H-B (2016) Purification and characterization of a new inducible thermostable extracellular lipolytic enzyme from the thermoacidophilic archaeon *Sulfolobus solfataricus* P1. *J Mol Catalysis B: Enzymatic* **124**: 11-19.

Chong PK, Burja AM, Radianingtyas H, Fazeli A & Wright PC (2007) Proteome and transcriptional analysis of ethanol-grown *Sulfolobus solfataricus* P2 reveals ADH2, a potential alcohol dehydrogenase. *J Proteome Res* **6**: 3985-3994.

Chong PK, Burja AM, Radianingtyas H, Fazeli A & Wright PC (2007) Translational and transcriptional analysis of *Sulfolobus solfataricus* P2 to provide insights into alcohol and ketone utilisation. *Proteomics* **7**: 424-435.

Choquet CG, Patel GB, Sprott GD & Beveridge TJ (1994) Stability of pressure-extruded liposomes made from archaeobacterial ether lipids. *Appl Microbiol Biotechnol* **42**: 375-384.

Clore AJ & Stedman KM (2007) The SSV1 viral integrase is not essential. *Virology* **361**: 103-111.

Cohen O, Doron S, Wurtzel O, Dar D, Edelheit S, Karunker I, Mick E & Sorek R (2016) Comparative transcriptomics across the prokaryotic tree of life. *Nucleic Acids Res* **44**: W46-W53.

Colman DR, Poudel S, Hamilton TL, Havig JR, Selensky MJ, Shock EL & Boyd ES (2018) Geobiological feedbacks and the evolution of thermoacidophiles. *ISME J* **12**: 225-236.

Cooper Charlotte R, Daugherty Amanda J, Tachdjian S, Blum Paul H & Kelly Robert M (2009) Role of *vapBC* toxin–antitoxin loci in the thermal stress response of *Sulfolobus solfataricus*. *Biochem Soc Trans* **37**: 123-126.

Coulson RMR, Touboul N & Ouzounis CA (2007) Lineage-specific partitions in archaeal transcription. *Archaea* **2**: 117-125.

Counts JA, Zeldes BM, Lee LL, Straub CT, Adams MWW & Kelly RM (2017) Physiological, metabolic and biotechnological features of extremely thermophilic microorganisms. *Wiley Interdiscip Rev Syst Biol Med* **9**.

Counts JA, D.J. Willard, and R.M. Kelly (2020) Life in hot acid: A genome-based reassessment of the archeal order *Sulfolobales*. *Environ Microbiol* **submitted**.

Couturier M & Lindås AC (2018) The DNA methylome of the hyperthermoacidophilic crenarchaeon *Sulfolobus acidocaldarius*. *Front Microbiol* **9**: 1-13.

Crosby JR, Laemthong T, Lewis AM, Straub CT, Adams MW & Kelly RM (2019) Extreme thermophiles as emerging metabolic engineering platforms. *Curr Opin Biotech* **59**: 55-64.

Cunningham Dueber EL, Corn JE, Bell SD & Berger JM (2007) Replication origin recognition and deformation by a heterodimeric archaeal Orc1 complex. *Science* **317**: 1210-1213.

Dahche H, Abdullah A, Potters MB & Kennelly PJ (2009) A PPM-family protein phosphatase from the thermoacidophile *Thermoplasma volcanium* hydrolyzes protein-bound phosphotyrosine. *Extremophiles* **13**: 371.

Danson MJ (1988) Archaebacteria: the comparative enzymology of their central metabolic pathways. *Adv Microbial Phys* **29**: 165-231.

Dar D, Prasse D, Schmitz RA & Sorek R (2016) Widespread formation of alternative 3' UTR isoforms via transcription termination in Archaea. *Nat Microbiol* **1**: 16143.

Darland G, Brock TD, Samsonoff W & Conti SF (1970) A thermophilic, acidophilic mycoplasma isolated from a coal refuse pile. *Science* **170**: 1416-1418.

Daum B, Quax TEF, Sachse M, *et al.* (2014) Self-assembly of the general membrane-remodeling protein PVAP into sevenfold virus-associated pyramids. *Proc Natl Acad Sci USA* **111**: 3829-3834.

Dellas N, Snyder JC, Bolduc B & Young MJ (2014) Archaeal viruses: Diversity, replication, and structure. *Annu Rev Virology* **1**: 399-426.

Deng L, Zhu H, Chen Z, Liang YX & She Q (2009) Unmarked gene deletion and host–vector system for the hyperthermophilic crenarchaeon *Sulfolobus islandicus*. *Extremophiles* **13**: 735.

Deng L, Garrett RA, Shah SA, Peng X & She Q (2013) A novel interference mechanism by a type IIIB CRISPR-Cmr module in *Sulfolobus*. *Mol Microbiol* **87**: 1088-1099.

Deng L, He F, Bhoobalan-Chitty Y, Martinez-Alvarez L, Guo Y & Peng X (2014) Unveiling cell surface and type IV secretion proteins responsible for archaeal ravidivirus entry. *J Virology* **88**: 10264-10268.

DeWerff SJ, Bautista MA, Pauly M, Zhang C & Whitaker RJ (2020) Killer Archaea: Virus-mediated antagonism to CRISPR-immune populations results in emergent virus-host mutualism. *mBio* **11**.

Di Fiore A, Fiorentino G, Vitale RM, Ronca R, Amodeo P, Pedone C, Bartolucci S & De Simone G (2009) Structural analysis of BldR from *Sulfolobus solfataricus* provides insights into the molecular basis of transcriptional activation in Archaea by MarR family proteins. *J Mol Biol* **388**: 559-569.

Dibrova DV, Galperin MY & Mulkidjanian AY (2014) Phylogenomic reconstruction of archaeal fatty acid metabolism. *Environ Microbiol* **16**: 907-918.

DiMaio F, Yu X, Rensen E, Krupovic M, Prangishvili D & Egelman EH (2015) A virus that infects a hyperthermophile encodes A-form DNA. *Science* **348**: 914-917.

Dionne I, Nookala RK, Jackson SP, Doherty AJ & Bell SD (2003) A heterotrimeric PCNA in the hyperthermophilic archaeon *Sulfolobus solfataricus*. *Mol Cell* **11**: 275-282.

Dionne I, Robinson N, McGeoch A, Marsh V, Reddish A & Bell S (2003) DNA replication in the hyperthermophilic archaeon *Sulfolobus solfataricus*. *31*: 674-676.

Dobro MJ, Samson RY, Yu Z, McCullough J, Ding HJ, Chong PLG, Bell SD & Jensen GJ (2013) Electron cryotomography of ESCRT assemblies and dividing *Sulfolobus* cells suggests that spiraling filaments are involved in membrane scission. *Mol Biol Cell* **24**: 2319-2327.

Dopson M (2016) Physiological and phylogenetic diversity of acidophilic Bacteria. *Acidophiles: Life in extremely acidic environments*, (Quatrini R & Barrie JD, eds.), p.^pp. 79-91. Caister Academic Press, Norfolk, UK.

Dorman CJ, Schumacher MA, Bush MJ, Brennan RG & Buttner MJ (2020) When is a transcription factor a NAP? *Curr Opin Microbiol* **55**: 26-33.

Doron S, Melamed S, Ofir G, Leavitt A, Lopatina A, Keren M, Amitai G & Sorek R (2018) Systematic discovery of antiphage defense systems in the microbial pangenome. *Science* **359**.

Drake JW (2009) Avoiding dangerous missense: thermophiles display especially low mutation rates. *PLoS Genet* **5**: e1000520.

Driessen RPC, Lin SN, Waterreus WJ, *et al.* (2016) Diverse architectural properties of Sso10a proteins: Evidence for a role in chromatin compaction and organization. *Sci Rep* **6**: 1-11.

Drozd M, Chandrashekhar K & Rajashekara G (2014) Polyphosphate-mediated modulation of *Campylobacter jejuni* biofilm growth and stability. *Virulence* **5**: 680-690.

Duan X & He Z-G (2011) Characterization of the specific interaction between archaeal FHA domain-containing protein and the promoter of a flagellar-like gene-cluster and its regulation by phosphorylation. *Biochem Biophys Res Commun* **407**: 242-247.

Duggin IG, McCallum SA & Bell SD (2008) Chromosome replication dynamics in the archaeon *Sulfolobus acidocaldarius*. *Proc Natl Acad Sci USA* **105**: 16737-16742.

Duggin IG, Dubarry N & Bell SD (2011) Replication termination and chromosome dimer resolution in the archaeon *Sulfolobus solfataricus*. *EMBO J* **30**: 145-153.

Elferink MG, Schleper C & Zillig W (1996) Transformation of the extremely thermoacidophilic archaeon *Sulfolobus solfataricus* via a self-spreading vector. *FEMS Microbiol Lett* **137**: 31-35.

Elferink MG, Albers SV, Konings WN & Driessen AJ (2001) Sugar transport in *Sulfolobus solfataricus* is mediated by two families of binding protein-dependent ABC transporters. *Mol Microbiol* **39**: 1494-1503.

Elferink MGL, de Wit JG, Driessen AJM & Konings WN (1994) Stability and proton-permeability of liposomes composed of archaeal tetraether lipids. *Biochim Biophys Acta - Biomembranes* **1193**: 247-254.

Ellen AF, Albers S-V & Driessens AJ (2010) Comparative study of the extracellular proteome of *Sulfolobus* species reveals limited secretion. *Extremophiles* **14**: 87.

Ellen AF, Albers S-V, Huibers W, *et al.* (2009) Proteomic analysis of secreted membrane vesicles of archaeal *Sulfolobus* species reveals the presence of endosome sorting complex components. *Extremophiles* **13**: 67-79.

England P, Wehenkel A, Martins S, Hoos S, André-Leroux G, Villarino A & Alzari PM (2009) The FHA-containing protein GarA acts as a phosphorylation-dependent molecular switch in mycobacterial signaling. *FEBS Lett* **583**: 301-307.

Esser D & Siebers B (2013) Atypical protein kinases of the RIO family in archaea. *Biochem Soc Trans* **41**: 399-404.

Esser D, Pham T, Reimann J, Albers S, Siebers B & Wright P (2012) Change of carbon source causes dramatic effects in the phospho-proteome of the archaeon *Sulfolobus solfataricus*. *J Proteome Res* **11**: 4823-4833.

Esser D, Kouril T, Talfournier F, Polkowska J, Schrader T, Brasen C & Siebers B (2013) Unraveling the function of paralogs of the aldehyde dehydrogenase super family from *Sulfolobus solfataricus*. *Extremophiles* **17**: 205-216.

Esser D, Hoffmann L, Pham TK, Bräsen C, Qiu W, Wright PC, Albers S-V & Siebers B (2016) Protein phosphorylation and its role in archaeal signal transduction. *FEMS Microbiol Rev* **40**: 625-647.

Estelmann S, Hugler M, Eisenreich W, Werner K, Berg IA, Ramos-Vera WH, Say RF, Kockelkorn D, Gad'on N & Fuchs G (2011) Labeling and enzyme studies of the central carbon metabolism in *Metallosphaera sedula*. *J Bacteriol* **193**: 1191-1200.

Ettema TJG, Huynen MA, de Vos WM & van der Oost J (2003) TRASH: a novel metal-binding domain predicted to be involved in heavy-metal sensing, trafficking and resistance. *Trends Biochem Sci* **28**: 170-173.

Ettema TJG, Ahmed H, Geerling ACM, van der Oost J & Siebers B (2008) The non-phosphorylating glyceraldehyde-3-phosphate dehydrogenase (GAPN) of *Sulfolobus solfataricus*: A key-enzyme of the semi-phosphorylative branch of the Entner–Doudoroff pathway. *Extremophiles* **12**: 75-88.

Ettema TJG, Brinkman AB, Lamers PP, Kornet NG, de Vos WM & van der Oost J (2006) Molecular characterization of a conserved archaeal copper resistance (cop) gene cluster and its copper-responsive regulator in *Sulfolobus solfataricus* P2. *Microbiologyopen* **152**: 1969-1979.

Facciotti MT, Reiss DJ, Pan M, *et al.* (2007) General transcription factor specified global gene regulation in Archaea. *Proc Natl Acad Sci USA* **104**: 4630-4635.

Farias ST & Bonato MC (2003) Preferred amino acids and thermostability. *Genetics and molecular research : GMR* **2**: 383-393.

Feng X, Sun M, Han W, Liang Yun X & She Q (2018) A transcriptional factor B paralog functions as an activator to DNA damage-responsive expression in Archaea. *Nucleic Acids Res* **46**: 7085-7096.

Feyhl-Buska J, Chen Y, Jia C, Wang J-X, Zhang CL & Boyd ES (2016) Influence of growth phase, pH, and temperature on the abundance and composition of tetraether lipids in the thermoacidophile *Picrophilus torridus*. *Front Microbiol* **7**.

Figueiredo AS, Kouril T, Esser D, Haferkamp P, Wieloch P, Schomburg D, Ruoff P, Siebers B & Schaber J (2017) Systems biology of the modified branched Entner-Doudoroff pathway in *Sulfolobus solfataricus*. *PLOS ONE* **12**: e0180331.

Fiorentino G, Del Giudice I, Bartolucci S, Durante L, Martino L & Del Vecchio P (2011) Identification and physicochemical characterization of BldR2 from *Sulfolobus solfataricus*, a novel archaeal member of the MarR transcription factor family. *Biochemistry* **50**: 6607-6621.

Forterre P (2002) A hot story from comparative genomics: reverse gyrase is the only hyperthermophile-specific protein. *Trends Genetics* **18**: 236-237.

Fouqueau T, Blombach F, Hartman R, Cheung ACM, Young MJ & Werner F (2017) The transcript cleavage factor paralogue TFS4 is a potent RNA polymerase inhibitor. *Nat Commun* **8**: 1-13.

Fröls S (2013) Archaeal biofilms: Widespread and complex. *Biochem Soc Trans* **41**: 393-398.

Fröls S, Gordon PM, Panlilio MA, Duggin IG, Bell SD, Sensen CW & Schleper C (2007) Response of the hyperthermophilic archaeon *Sulfolobus solfataricus* to UV damage. *J Bacteriol* **189**: 8708-8718.

Fröls S, Ajon M, Wagner M, Teichmann D, Zolghadr B, Folea M, Boekema EJ, Driessen AJM, Schleper C & Albers S-V (2008) UV-inducible cellular aggregation of the hyperthermophilic archaeon *Sulfolobus solfataricus* is mediated by pili formation. *Mol Microbiol* **70**: 938-952.

Fu C-y, Wang K, Gan L, Lanman J, Khayat R, Young MJ, Jensen GJ, Doerschuk PC & Johnson JE (2010) *In vivo* assembly of an archaeal virus studied with whole-cell electron cryotomography. *Structure* **18**: 1579-1586.

Fuchs G (2011) Alternative pathways of carbon dioxide fixation: Insights into the early evolution of life? *Annu Rev Microbiol* **65**: 631-658.

Fuchs T, Huber H, Teiner K, Burggraf S & Stetter KO (1995) *Metallosphaera prunae*, sp. nov., a novel metal-mobilizing, thermoacidophilic archaeum, isolated from a uranium mine in Germany. *Syst Appl Microbiol* **18**: 560-566.

Futterer O, Angelov A, Liesegang H, Gottschalk G, Schleper C, Schepers B, Dock C, Antranikian G & Liebl W (2004) Genome sequence of *Picrophilus torridus* and its implications for life around pH 0. *Proc Natl Acad Sci USA* **101**: 9091-9096.

Galperin MY, Makarova KS, Wolf YI & Koonin EV (2018) Phyletic distribution and lineage-specific domain architectures of archaeal two-component signal transduction systems. *J Bacteriol* **200**: e00681-00617.

Gambelli L, Meyer BH, McLaren M, Sanders K, Quax TEF, Gold VAM, Albers S-V & Daum B (2019) Architecture and modular assembly of *Sulfolobus* S-layers revealed by electron cryotomography. *Proc Natl Acad Sci USA* **116**: 25278-25286.

Gardner AF, Bell SD, White MF, Prangishvili D & Krupovic M (2014) Protein-Protein interactions leading to recruitment of the host DNA sliding clamp by the hyperthermophilic *Sulfolobus islandicus* rod-shaped Virus 2. *J Virology* **88**: 7105-7108.

Garrett RA, Shah SA, Vestergaard G, Deng L, Gudbergsdottir S, Kenchappa CS, Erdmann S & She Q (2011) CRISPR-based immune systems of the *Sulfolobales*: Complexity and diversity. *Biochem Soc Trans* **39**: 51-57.

Geertsma ER (2013) FX cloning: a versatile high-throughput cloning system for characterization of enzyme variants. *Enzyme Eng*, p.^pp. 133-148. Springer.

Giles TN & Graham DE (2008) Crenarchaeal Arginine Decarboxylase Evolved from an S-Adenosylmethionine Decarboxylase enzyme. *J Biol Chem* **283**: 25829-25838.

Girfoglio M, Rossi M & Cannio R (2012) Cellulose degradation by *Sulfolobus solfataricus* requires a cell-anchored endo-beta-1,4-glucanase. *J Bacteriol* **194**: 5091-5100.

Gleissner M, Kaiser U, Antonopoulos E & Schafer G (1997) The archaeal SoxABCD complex is a proton pump in *Sulfolobus acidocaldarius*. *J Biol Chem* **272**: 8417-8426.

Gigliettino M, Balestrieri M, Pocsfalvi G, Fiume I, Natale L, Rossi M & Palmieri G (2010) A highly selective oligopeptide binding protein from the archaeon *Sulfolobus solfataricus*. *J Bacteriol* **192**: 3123-3131.

Goldberg GW, Jiang W, Bikard D & Marraffini LA (2014) Conditional tolerance of temperate phages via transcription-dependent CRISPR-Cas targeting. *Nature* **514**: 633-637.

Golyshina OV, Yakimov MM, Lunsdorf H, Ferrer M, Nimtz M, Timmis KN, Wray V, Tindall BJ & Golyshin PN (2009) *Acidiplasma aeolicum* gen. nov., sp. nov., a euryarchaeon of the family *Ferroplasmaceae* isolated from a hydrothermal pool, and transfer of *Ferroplasma cupricumulans* to *Acidiplasma cupricumulans* comb. nov. *Int J Syst Evol Microbiol* **59**: 2815-2823.

Gomes-Filho JV & Randau L (2019) RNA stabilization in hyperthermophilic Archaea. *Ann NY Acad Sci* **1447**: 88-96.

Gong F, Zhu H, Zhou J, Zhao T, Xiao L, Zhang Y & Li Y (2019) Enhanced biological fixation of CO₂ using microorganisms. *An Economy Based on Carbon Dioxide and Water*, (Aresta M, Karimi IA & Kawi S, eds.), p.^pp. Springer, Cham.

Goto K, Tanimoto Y, Tamura T, Mochida K, Arai D, Asahara M, Suzuki M, Tanaka H & Inagaki K (2002) Identification of thermoacidophilic bacteria and a new *Alicyclobacillus* genomic species isolated from acidic environments in Japan. *Extremophiles* **6**: 333-340.

Gray MJ & Jakob U (2015) Oxidative stress protection by polyphosphate—new roles for an old player. *Curr Opin Microbiol* **24**: 1-6.

Greci MD & Bell SD (2020) Archaeal DNA replication. *Annu Rev Microbiol* **74**.

Grillo-Puertas M, Rintoul R, Villegas JM & Rapisarda VA (2012) Polyphosphate degradation in stationary phase triggers biofilm formation via LuxS quorum sensing system in *Escherichia coli*. *PLoS ONE* **7**.

Grillo-Puertas M, Schurig-Briccio LA, Rodríguez-Montelongo L, Rintoul MR & Rapisarda VA (2014) Copper tolerance mediated by polyphosphate degradation and low-affinity inorganic phosphate transport system in *Escherichia coli*. *BMC Microbiol* **14**: 72.

Grogan D, Palm P & Zillig W (1990) Isolate B12, which harbours a virus-like element, represents a new species of the archaeabacterial genus *Sulfolobus*, *Sulfolobus shibatae*, sp. nov. *Arch Microbiol* **154**: 594-599.

Grogan DW (1989) Phenotypic characterization of the archaeabacterial genus *Sulfolobus*: comparison of five wild-type strains. *J Bacteriol* **171**: 6710-6719.

Grogan DW (1991) Selectable mutant phenotypes of the extremely thermophilic archaeabacterium *Sulfolobus acidocaldarius*. *J Bacteriol* **173**: 7725-7727.

Grogan DW (1996) Organization and interactions of cell envelope proteins of the extreme thermoacidophile *Sulfolobus acidocaldarius*. *Can J Microbiol* **42**: 1163-1171.

Grogan DW (2003) Cytosine methylation by the SuaI restriction-modification system: implications for genetic fidelity in a hyperthermophilic archaeon. *J Bacteriol* **185**: 4657-4661.

Grogan DW & Gunsalus RP (1993) *Sulfolobus acidocaldarius* synthesizes UMP via a standard de novo pathway: results of biochemical-genetic study. *J Bacteriol* **175**: 1500-1507.

Grogan DW & Stengel KR (2008) Recombination of synthetic oligonucleotides with prokaryotic chromosomes: substrate requirements of the *Escherichia coli*/lambdaRed and *Sulfolobus acidocaldarius* recombination systems. *Mol Microbiol* **69**: 1255-1265.

Grogan DW, Carver GT & Drake JW (2001) Genetic fidelity under harsh conditions: Analysis of spontaneous mutation in the thermoacidophilic archaeon *Sulfolobus acidocaldarius*. *Proc Natl Acad Sci USA* **98**: 7928-7933.

Guan Z, Delago A, Nußbaum P, Meyer B, Albers S-V & Eichler J (2016) N-glycosylation in the thermoacidophilic archaeon *Sulfolobus acidocaldarius* involves a short dolichol pyrophosphate carrier. *FEBS Lett* **590**: 3168-3178.

Guo L, Ding J, Guo R, Hou Y, Wang DC & Huang L (2014) Biochemical and structural insights into RNA binding by Ssh10b, a member of the highly conserved Sac10b protein family in Archaea. *J Biol Chem* **289**: 1478-1490.

Guo L, Feng Y, Zhang Z, Yao H, Luo Y, Wang J & Huang L (2008) Biochemical and structural characterization of Cren7, a novel chromatin protein conserved among Crenarchaea. *Nucleic Acids Res* **36**: 1129-1137.

Gutsche I, Mihalache O & Baumeister W (2000) ATPase cycle of an archaeal chaperonin. *J Mol Biol* **300**: 187-196.

Gutsche I, Holzinger J, Rößle M, Heumann H, Baumeister W & May RP (2000) Conformational rearrangements of an archaeal chaperonin upon ATPase cycling. *Curr Biology* **10**: 405-408.

Haferkamp P, Tjaden B, Shen L, Bräsen C, Kouril T & Siebers B (2019) The carbon switch at the level of pyruvate and phosphoenolpyruvate in *Sulfolobus solfataricus* P2. *Front Microbiol* **10**.

Haferkamp P, Kutschki S, Treichel J, Hemeda H, Sewczyk K, Hoffmann D, Zaparty M & Siebers B (2011) An additional glucose dehydrogenase from *Sulfolobus solfataricus*: Fine-tuning of sugar degradation? *Biochem Soc Trans* **39**: 77-81.

Haile JD & Kennelly PJ (2011) The activity of an ancient atypical protein kinase is stimulated by ADP-ribose *in vitro*. *Arch Biochem Biophys* **511**: 56-63.

Hampton HG, Watson BNJ & Fineran PC (2020) The arms race between bacteria and their phage foes. *Nature* **577**: 327-336.

Han D & Krauss G (2009) Characterization of the endonuclease SSO2001 from *Sulfolobus solfataricus* P2. *FEBS Letters* **583**: 771-776.

Hanks SK (2003) Genomic analysis of the eukaryotic protein kinase superfamily: A perspective. *Genome Biology* **4**: 111.

Haring M, Rachel R, Peng X, Garrett RA & Prangishvili D (2005) Viral Diversity in hot springs of Pozzuoli, Italy, and characterization of a unique archaeal virus, *Acidianus* bottle-shaped virus, from a new family, the *Ampullaviridae*. *J Virology* **79**: 9904-9911.

Hartman R, Eilers BJ, Bollschweiler D, Munson-McGee JH, Engelhardt H, Young MJ & Lawrence CM (2019) The molecular mechanism of cellular attachment for an archaeal virus. *Structure* **27**: 1634-1646.e1633.

Haseltine C, Rolfsmeier M & Blum P (1996) The glucose effect and regulation of alpha-amylase synthesis in the hyperthermophilic archaeon *Sulfolobus solfataricus*. *J Bacteriol* **178**: 945-950.

Haslbeck M, Franzmann T, Weinfurtner D & Buchner J (2005) Some like it hot: the structure and function of small heat-shock proteins. *Nat Struct Mol Biol* **12**: 842-846.

Haurat MF, Figueiredo AS, Hoffmann L, Li L, Herr K, J. Wilson A, Beeby M, Schaber J & Albers SV (2017) ArnS, a kinase involved in starvation-induced archaellum expression. *Mol Microbiol* **103**: 181-194.

Hawkes RB, Franzmann PD, O'Hara G & Plumb JJ (2006) *Ferroplasma cupricumulans* sp. nov., a novel moderately thermophilic, acidophilic archaeon isolated from an industrial-scale chalcocite bioleach heap. *Extremophiles* **10**: 525-530.

Hawkins AS, Han Y, Bennett RK, Adams MW & Kelly RM (2013) Role of 4-hydroxybutyrate-CoA synthetase in the CO₂ fixation cycle in thermoacidophilic archaea. *J Biol Chem* **288**: 4012-4022.

He F, Bhoobalan-Chitty Y, Van LB, Kjeldsen AL, Dedola M, Makarova KS, Koonin EV, Brodersen DE & Peng X (2018) Anti-CRISPR proteins encoded by archaeal lytic viruses inhibit subtype I-D immunity. *Nat Microbiol* **3**: 461-469.

Held NL & Whitaker RJ (2009) Viral biogeography revealed by signatures in *Sulfolobus islandicus* genomes. *Environ Microbiol* **11**: 457-466.

Henche A-L, Koerdt A, Ghosh A & Albers S-V (2012) Influence of cell surface structures on crenarchaeal biofilm formation using a thermostable green fluorescent protein. *Environ Microbiol* **14**: 779-793.

Henche AL, Ghosh A, Yu X, Jeske T, Egelman E & Albers SV (2012) Structure and function of the adhesive type IV pilus of *Sulfolobus acidocaldarius*. *Environ Microbiol* **14**: 3188-3202.

Hess M (2008) Thermoacidophilic proteins for biofuel production. *Trends Microbiol* **16**: 414-419.

Hirata A, Klein BJ & Murakami KS (2008) The X-ray crystal structure of RNA polymerase from Archaea. *Nature* **451**: 851-854.

Hiroyoshi N, Miki H, Hirajima T & Tsunekawa M (2000) A model for ferrous-promoted chalcopyrite leaching. *Hydrometallurgy* **57**: 31-38.

Hjort K & Bernander R (2001) Cell cycle regulation in the hyperthermophilic crenarchaeon *Sulfolobus acidocaldarius*. *Mol Microbiol* **40**: 225-234.

Hobel CFV, Albers SV, Driessen AJM & Lupas AN (2008) The *Sulfolobus solfataricus* AAA protein Sso0909, a homologue of the eukaryotic ESCRT Vps4 ATPase. *Biochem Sci Trans* **36**: 94-98.

Hocher A, Rojec M, Swadling JB, Esin A & Warnecke T (2019) The DNA-binding protein HTa from *Thermoplasma acidophilum* is an archaeal histone analog. *eLife* **8**: e52542.

Hoffmann L, Schummer A, Reimann J, Haurat MF, Wilson AJ, Beeby M, Warscheid B & Albers SV (2017) Expanding the archaellum regulatory network—the eukaryotic protein kinases ArnC and ArnD influence motility of *Sulfolobus acidocaldarius*. *Microbiologyopen* **6**.

Hottiger T, Schmutz P & Wiemken A (1987) Heat-induced accumulation and futile cycling of trehalose in *Saccharomyces cerevisiae*. *J Bacteriol* **169**: 5518-5522.

Huang Q, Zhong Q, Mayaka J, Ni J & Shen Y (2017) Autophosphorylation and cross-phosphorylation of protein kinases from the Crenarchaeon *Sulfolobus islandicus*. *Front Microbiol* **8**: 2173.

Huang Q, Mayaka JBA, Zhong Q, Zhang C, Hou G, Ni J & Shen Y (2019) Phosphorylation of the archaeal Holliday junction resolvase Hjc inhibits its catalytic activity and facilitates DNA repair in *Sulfolobus islandicus* REY15A. *Front Microbiol* **10**: 1214.

Huang Y, Krauss G, Cottaz S, Driguez H & Lipps G (2005) A highly acid-stable and thermostable endo- β -glucanase from the thermoacidophilic archaeon *Sulfolobus solfataricus*. *Biochem J* **385**: 581-588.

Huber G & Stetter KO (1991) *Sulfolobus metallicus*, sp. nov., a novel strictly chemolithoautotrophic thermophilic archaeal species of metal-mobilizers. *Syst Appl Microbiol* **14**: 372-378.

Huber G, Spinnler C, Gambacorta A & Stetter KO (1989) *Metallosphaera sedula* gen, and sp. nov. represents a new genus of aerobic, metal-mobilizing, thermoacidophilic archaeabacteria. *Syst Appl Microbiol* **12**: 38-47.

Hugler M, Huber H, Stetter KO & Fuchs G (2003) Autotrophic CO₂ fixation pathways in archaea (Crenarchaeota). *Arch Microbiol* **179**: 160-173.

Hwang S & Maxwell KL (2019) Meet the Anti-CRISPRs: Widespread protein inhibitors of CRISPR-Cas systems. *CRISPR J* **2**: 23-30.

Häring M, Vestergaard G, Rachel R, Chen L, Garrett RA & Prangishvili D (2005) Virology: Independent virus development outside a host. *Nature* **436**: 1101-1102.

Inskeep W, Jay Z, Tringe S, Herrgard M & Rusch D (2013) The YNP Metagenome Project: Environmental parameters responsible for microbial distribution in the Yellowstone geothermal ecosystem. *Front Microbiol* **4**.

Iqbal J & Qureshi SA (2010) Selective depletion of *Sulfolobus solfataricus* transcription factor E under heat shock conditions. *J Bacteriol* **192**: 2887-2891.

Itoh T, Yoshikawa N & Takashina T (2007) *Thermogymnomonas acidicola* gen. nov., sp. nov., a novel thermoacidophilic, cell wall-less archaeon in the order *Thermoplasmatales*, isolated from a solfataric soil in Hakone, Japan. *Int J Syst Evol Microbiol* **57**: 2557-2561.

Itoh T, Suzuki K, Sanchez PC & Nakase T (1999) *Caldivirga maquilingensis* gen. nov., sp. nov., a new genus of rod-shaped crenarchaeote isolated from a hot spring in the Philippines. *Int J Syst Evol Microbiol* **49**: 1157-1163.

Itoh T, Suzuki K, Sanchez PC & Nakase T (2003) *Caldisphaera lagunensis* gen. nov., sp. nov., a novel thermoacidophilic crenarchaeote isolated from a hot spring at Mt Maquiling, Philippines. *Int J Syst Evol Microbiol* **53**: 1149-1154.

Itoh T, Miura T, Sakai HD, Kato S, Ohkuma M & Takashina T (2020) *Sulfuracidifex tepidarius* gen. nov., sp. nov. and transfer of *Sulfolobus metallicus* Huber and Stetter 1992 to the genus *Sulfuracidifex* as *Sulfuracidifex metallicus* comb. nov. *Int J Syst Evol Microbiol* **70**: 1837-1842.

Izzo V, Notomista E, Picardi A, Pennacchio F & Di Donato A (2005) The thermophilic archaeon *Sulfolobus solfataricus* is able to grow on phenol. *Res Microbiol* **156**: 677-689.

Jachlewski S, Jachlewski WD, Linne U, Bräsen C, Wingender J & Siebers B (2015) Isolation of extracellular polymeric substances from biofilms of the thermoacidophilic archaeon *Sulfolobus acidocaldarius*. *Front Bioeng Biotechnol* **3**: 123-123.

Jackson SA, McKenzie RE, Fagerlund RD, Kieper SN, Fineran PC & Brouns SJ (2017) CRISPR-Cas: Adapting to change. *Science* **356**.

Jarrell KF & Albers S-V (2012) The archaellum: an old motility structure with a new name. *Trends Microbiol* **20**: 307-312.

Jarrell KF, Ding Y, Meyer BH, Albers S-V, Kaminski L & Eichler J (2014) N-linked glycosylation in Archaea: a structural, functional, and genetic analysis. *Microbiol Molec Biol Rev* **78**: 304-341.

Jay ZJ, Beam JP, Dlakic M, Rusch DB, Kozubal MA & Inskeep WP (2018) *Marsarchaeota* are an aerobic archaeal lineage abundant in geothermal iron oxide microbial mats. *Nat Microbiol* **3**: 732-740.

Jiang CY, Liu LJ, Guo X, You XY, Liu SJ & Poetsch A (2014) Resolution of carbon metabolism and sulfur-oxidation pathways of *Metallosphaera cuprina* Ar-4 via comparative proteomics. *J Proteomics* **109**: 276-289.

Johnsen U, Reinhardt A, Landan G, Tria FDK, Turner JM, Davies C & Schönheit P (2019) New views on an old enzyme: allosteric regulation and evolution of archaeal pyruvate kinases. *FEBS J* **286**: 2471-2489.

Johnson DB, Okibe N & Roberto FF (2003) Novel thermo-acidophilic bacteria isolated from geothermal sites in Yellowstone National Park: Physiological and phylogenetic characteristics. *Arch Microbiol* **180**: 60-68.

Johnson T, Payne S, Grove R, McCarthy S, Oeltjen E, Mach C, Adamec J, Wilson MA, Cott KV & Blum P (2019) Methylation deficiency of chromatin proteins is a non-mutational and epigenetic-like trait in evolved lines of the archaeon *Sulfolobus solfataricus*. *J Biol Chem* **294**: 7821-7832.

Jonuscheit M, Martusewitsch E, Stedman KM & Schleper C (2003) A reporter gene system for the hyperthermophilic archaeon *Sulfolobus solfataricus* based on a selectable and integrative shuttle vector. *Mol Microbiol* **48**: 1241-1252.

Kagawa HK, Yaoi T, Brocchieri L, McMillan RA, Alton T & Trent JD (2003) The composition, structure and stability of a group II chaperonin are temperature regulated in a hyperthermophilic archaeon. *Mol Microbiol* **48**: 143-156.

Kagawa HK, Osipiuk J, Maltsev N, Overbeek R, Quaite-Randall E, Joachimiak A & Trent JD (1995) The 60 kDa heat shock proteins in the hyperthermophilic archaeon *Sulfolobus shibatae*. *J Mol Biol* **253**: 712-725.

Kalichuk V, Béhar G, Renodon-Cornière A, Danovski G, Obal G, Barbet J, Mouratou B & Pecorari F (2016) The archaeal "7 kDa DNA-binding" proteins: Extended characterization of an old gifted family. *Sci Rep* **6**: 1-10.

Kallioma-Sanford AK, Rodriguez-Castañeda FA, McLeod BN, Latorre-Roselló V, Smith JH, Reimann J, Albers SV & Barillà D (2012) Chromosome segregation in Archaea mediated by a hybrid DNA partition machine. *Proc Natl Acad Sci USA* **109**: 3754-3759.

Kamada K & Barillà D (2018) Combing chromosomal DNA mediated by the SMC complex: structure and mechanisms. *BioEssays* **40**: 1-8.

Kappler U & Dahl C (2001) Enzymology and molecular biology of prokaryotic sulfite oxidation. *FEMS Microbiol Lett* **203**: 1-9.

Karr EA, Isom CE, Trinh V & Peeters E (2017) Transcription factor-mediated gene regulation in Archaea. *RNA Metabolism and Gene Expression in Archaea*, Vol. 32 p.^pp. 27-69. Springer, Cham.

Kazlauskiene M, Kostiuk G, Venclovas Č, Tamulaitis G & Siksnys V (2017) A cyclic oligonucleotide signaling pathway in type III CRISPR-Cas systems. *Science* **357**: 605-609.

Keasling JD (1997) Regulation of intracellular toxic metals and other cations by hydrolysis of polyphosphate. *Ann NY Acad Sci* **829**: 242-249.

Keeling PJ, Klenk H-P, Singh RK, Schenk ME, Sensen CW, Zillig W & Doolittle WF (1998) *Sulfolobus islandicus* plasmids pRN1 and pRN2 share distant but common evolutionary ancestry. *Extremophiles* **2**: 391-393.

Keeling PJ, Klenk H-P, Singh RK, Feeley O, Schleper C, Zillig W, Doolittle WF & Sensen CW (1996) Complete nucleotide sequence of the *Sulfolobus islandicus* multicopy plasmid pRN1. *Plasmid* **35**: 141-144.

Keller MW, Schut GJ, Lipscomb GL, *et al.* (2013) Exploiting microbial hyperthermophilicity to produce an industrial chemical, using hydrogen and carbon dioxide. *Proc Natl Acad Sci USA* **110**: 5840-5845.

Kelley DS, Baross JA & Delaney JR (2002) Volcanoes, fluids, and life at mid-ocean ridge spreading centers. *Annu Rev Earth Planetary Sci* **30**: 385-491.

Kelly DE, Lamb DC & Kelly SL (2001) Genome-wide generation of yeast gene deletion strains. *Comp Funct Genomics* **2**: 236-242.

Kennelly P, Oxenrider K, Leng J, Cantwell J & Zhao N (1993) Identification of a serine/threonine-specific protein phosphatase from the archaeabacterium *Sulfolobus solfataricus*. *J Biol Chem* **268**: 6505-6510.

Kennelly PJ (2003) Archaeal protein kinases and protein phosphatases: insights from genomics and biochemistry. *Biochem J* **370**: 373-389.

Kennelly PJ (2014) Protein Ser/Thr/Tyr phosphorylation in the Archaea. *J Biol Chem* **289**: 9480-9487.

Kerscher L, Nowitzki S & Oesterhelt D (1982) Thermoacidophilic archaeabacteria contain bacterial-type ferredoxins acting as electron acceptors of 2-oxoacid:ferredoxin oxidoreductases. *Eur J Biochem* **128**: 223-230.

Kim D-j & Forst S (2001) Genomic analysis of the histidine kinase family in Bacteria and Archaea. *Microbiologyopen* **147**: 1197-1212.

Kim MS, Park JT, Kim YW, *et al.* (2004) Properties of a novel thermostable glucoamylase from the hyperthermophilic archaeon *Sulfolobus solfataricus* in relation to starch processing. *Appl Environ Microbiol* **70**: 3933-3940.

Kim SM, Paek KH & Lee SB (2012) Characterization of NADP+-specific l-rhamnose dehydrogenase from the thermoacidophilic archaeon *Thermoplasma acidophilum*. *Extremophiles* **16**: 447-454.

Klauber C (2008) A critical review of the surface chemistry of acidic ferric sulphate dissolution of chalcopyrite with regards to hindered dissolution. *Int J Miner Process* **86**: 1-17.

Kletzin A (1989) Coupled enzymatic production of sulfite, thiosulfate, and hydrogen sulfide from sulfur: purification and properties of a sulfur oxygenase reductase from the facultatively anaerobic archaeabacterium *Desulfurolobus ambivalens*. *J Bacteriol* **171**: 1638-1643.

Kletzin A (1992) Molecular characterization of the *sor* gene, which encodes the sulfur oxygenase/reductase of the thermoacidophilic archaeum *Desulfurolobus ambivalens*. *J Bacteriol* **174**: 5854-5859.

Kletzin A, Urich T, Muller F, Bandeiras TM & Gomes CM (2004) Dissimilatory oxidation and reduction of elemental sulfur in thermophilic archaea. *J Bioenerg Biomembr* **36**: 77-91.

Klingl A, Pickl C & Flechsler J (2019) Archaeal cell walls. *Sub-Cellular Biochem* **92**: 471-493.

Knüppel R, Christensen RH, Gray FC, Esser D, Strauß D, Medenbach J, Siebers B, MacNeill SA, LaRonde N & Ferreira-Cerca S (2018) Insights into the evolutionary conserved regulation of Rio ATPase activity. *Nucl Acids Res* **46**: 1441-1456.

Koch T & Dahl C (2018) A novel bacterial sulfur oxidation pathway provides a new link between the cycles of organic and inorganic sulfur compounds. *ISME J* **12**: 2479-2491.

Kockelkorn D & Fuchs G (2009) Malonic semialdehyde reductase, succinic semialdehyde reductase, and succinyl-coenzyme A reductase from *Metallosphaera sedula*: Enzymes of the autotrophic 3-hydroxypropionate/4-hydroxybutyrate cycle in *Sulfolobales*. *J Bacteriol* **191**: 6352-6362.

Koechler S, Farasin J, Cleiss-Arnold J & Arsène-Ploetze F (2015) Toxic metal resistance in biofilms: Diversity of microbial responses and their evolution. *Res Microbiol* **166**: 764-773.

Koerdt A, Gödeke J, Berger J, Thormann KM & Albers SV (2010) Crenarchaeal biofilm formation under extreme conditions. *PLOS ONE* **5**: e14104.

Koerdt A, Jachlewski S, Ghosh A, Wingender J, Siebers B & Albers SV (2012) Complementation of *Sulfolobus solfataricus* PBL2025 with an α -mannosidase: effects on surface attachment and biofilm formation. *Extremophiles* **16**: 115-125.

Koerdt A, Orell A, Pham TK, Mukherjee J, Włodkowski A, Karunakaran E, Biggs CA, Wright PC & Albers S-V (2011) Macromolecular fingerprinting of *Sulfolobus* species in biofilm: A transcriptomic and proteomic approach combined with spectroscopic analysis. *J Proteome Res* **10**: 4105-4119.

Komatsu H & Chong PL-G (1998) Low permeability of liposomal membranes composed of bipolar tetraether lipids from thermoacidophilic archaeabacterium *Sulfolobus acidocaldarius*. *Biochemistry* **37**: 107-115.

Kondo S, Yamagishi A & Oshima T (1991) Positive selection for uracil auxotrophs of the sulfur-dependent thermophilic archaeabacterium *Sulfolobus acidocaldarius* by use of 5-fluoroorotic acid. *J Bacteriol* **173**: 7698-7700.

Kondrat'eva TF, Pivovarova TA, Tsaplina IA, Fomchenko NV, Zhuravleva AE, Murav'ev MI, Melamud VS & Bulaev AG (2012) [Diversity of acidophilic chemolithotrophic microbial consortia in natural and anthropogenic ecosystems]. *Mikrobiologija* **81**: 3-27.

Konings WN, Albers S-V, Koning S & Driessen AJM (2002) The cell membrane plays a crucial role in survival of Bacteria and Archaea in extreme environments. *Antonie van Leeuwenhoek* **81**: 61-72.

Koonin EV, Makarova KS & Wolf YI (2017) Evolutionary genomics of defense systems in Archaea and Bacteria. *Ann Rev Microbiol* **71**: 233-261.

Korkhin Y, Unligil UM, Littlefield O, Nelson PJ, Stuart DI, Sigler PB, Bell SD & Abrescia NGA (2009) Evolution of complex RNA polymerases: The complete archaeal RNA polymerase structure. *PLoS Biol* **7**: e1000102.

Kornberg A, N.N. Rao and D. Ault-Riché (1999) Inorganic polyphosphate: A molecule of many functions. *Ann Rev Biochem* **68**: 89-125.

Kouril T, Eicher JJ, Siebers B & Snoep JL (2017) Phosphoglycerate kinase acts as a futile cycle at high temperature. *Microbiologyopen* **163**: 1604-1612.

Kouril T, Esser D, Kort J, Westerhoff HV, Siebers B & Snoep JL (2013) Intermediate instability at high temperature leads to low pathway efficiency for an *in vitro* reconstituted system of gluconeogenesis in *Sulfolobus solfataricus*. *FEBS J* **280**: 4666-4680.

Kouril T, Wieloch P, Reimann J, Wagner M, Zaparty M, Albers SV, Schomburg D, Ruoff P & Siebers B (2013) Unraveling the function of the two Entner-Doudoroff branches in the thermoacidophilic Crenarchaeon *Sulfolobus solfataricus* P2. *FEBS J* **280**: 1126-1138.

Kozubal MA, Dlakic M, Macur RE & Inskeep WP (2011) Terminal oxidase diversity and function in "*Metallosphaera yellowstonensis*": Gene expression and protein modeling suggest mechanisms of Fe(II) oxidation in the *Sulfolobales*. *Appl Environ Microbiol* **77**: 1844-1853.

Kozubal MA, Romine M, Jennings R, Jay ZJ, Tringe SG, Rusch DB, Beam JP, McCue LA & Inskeep WP (2013) *Geoarchaeota*: A new candidate phylum in the Archaea from high-temperature acidic iron mats in Yellowstone National Park. *ISME J* **7**: 622-634.

Kurosawa N & Grogan DW (2005) Homologous recombination of exogenous DNA with the *Sulfolobus acidocaldarius* genome: properties and uses. *FEMS Microbiol* **253**: 141-149.

Kurosawa N, Itoh YH, Iwai T, Sugai A, Uda I, Kimura N, Horiuchi T & Itoh T (1998) *Sulfurisphaera ohwakuensis* gen. nov., sp. nov., a novel extremely thermophilic acidophile of the order *Sulfolobales*. *Int J Syst Evol Microbiol* **48**: 451-456.

La Teana A, Benelli D, Londei P & Bläsi U (2013) Translation initiation in the crenarchaeon *Sulfolobus solfataricus*: Eukaryotic features but bacterial route. *Biochem Soc T* **41**: 350-355.

Lamble HJ, Heyer NI, Bull SD, Hough DW & Danson MJ (2003) Metabolic pathway promiscuity in the archaeon *Sulfolobus solfataricus* revealed by studies on Glucose Dehydrogenase and 2-Keto-3-deoxygluconate Aldolase. *J Biol Chem* **278**: 34066-34072.

Lamble HJ, Milburn CC, Taylor GL, Hough DW & Danson MJ (2004) Gluconate dehydratase from the promiscuous Entner-Doudoroff pathway in *Sulfolobus solfataricus*. *FEBS Lett* **576**: 133-136.

Lamble HJ, Theodossis A, Milburn CC, Taylor GL, Bull SD, Hough DW & Danson MJ (2005) Promiscuity in the part-phosphorylative Entner-Doudoroff pathway of the archaeon *Sulfolobus solfataricus*. *FEBS Lett* **579**: 6865-6869.

Laplagia C & Hartzell PL (1997) Stress-induced production of biofilm in the hyperthermophile *Archaeoglobus fulgidus*. *Appl Environ Microbiol* **63**: 3158-3163.

LaRonde NA (2014) The ancient microbial RIO kinases. *J Biol Chem* **289**: 9488-9492.

Laska S, Lottspeich F & Kletzin A (2003) Membrane-bound hydrogenase and sulfur reductase of the hyperthermophilic and acidophilic archaeon *Acidianus ambivalens*. *Microbiologyopen* **149**: 2357-2371.

Lassak K, Neiner T, Ghosh A, Klingl A, Wirth R & Albers SV (2012) Molecular analysis of the crenarchaeal flagellum. *Mol Microbiol* **83**: 110-124.

Laurens N, Driessens RP, Heller I, Vorselen D, Noom MC, Hol FJ, White MF, Dame RT & Wuite GJ (2012) Alba shapes the archaeal genome using a delicate balance of bridging and stiffening the DNA. *Nat Commun* **3**: 1-8.

Lechner J & Sumper M (1987) The primary structure of a prokaryotic glycoprotein. Cloning and sequencing of the cell surface glycoprotein gene of Halobacteria. *J Biol Chem* **262**: 9724-9729.

Lemire J, Alhasawi A, Appanna VP, Tharmalingam S & Appanna VD (2017) Metabolic defence against oxidative stress: the road less travelled so far. *J Appl Microbiol* **123**: 798-809.

Lemmens L, Maklad HR, Bervoets I & Peeters E (2019) Transcription regulators in Archaea: Homologies and differences with bacterial regulators. *J Mol Biol* **431**: 4132-4146.

Lemos RS, Gomes CM & Teixeira M (2001) *Acidianus ambivalens* Complex II typifies a novel family of succinate dehydrogenases. *Biochem Biophys Res Commun* **281**: 141-150.

Leng J, Cameron A, Buckel S & Kennelly PJ (1995) Isolation and cloning of a protein-serine/threonine phosphatase from an archaeon. *J Bacteriol* **177**: 6510-6517.

Leonard CJ, Aravind L & Koonin EV (1998) Novel families of putative protein kinases in Bacteria and Archaea: evolution of the “eukaryotic” protein kinase superfamily. *Genome Res* **8**: 1038-1047.

Lewis P & Ford RM (1999) Temperature-sensitive motility of *Sulfolobus acidocaldarius* influences population distribution in extreme environments. *J Bacteriol* **181**: 4020-4025.

Leyn SA, Rodionova IA, Li X & Rodionov DA (2015) Novel transcriptional regulons for autotrophic cycle genes in *Crenarchaeota*. *J Bacteriol* **197**: 2383-2391.

León-Sobrino C, Kot WP & Garrett RA (2016) Transcriptome changes in STSV2-infected *Sulfolobus islandicus* REY15A undergoing continuous CRISPR spacer acquisition. *Mol Microbiol* **99**: 719-728.

Li D-C, Yang F, Lu B, Chen D-F & Yang W-J (2012) Thermotolerance and molecular chaperone function of the small heat shock protein HSP20 from hyperthermophilic archaeon, *Sulfolobus solfataricus* P2. *Cell Stress Chaper* **17**: 103-108.

Li L, Banerjee A, Bischof LF, Maklad HR, Hoffmann L, Henche AL, Veliz F, Bildl W, Schulte U & Orell A (2017) Wing phosphorylation is a major functional determinant of the Lrs14-type biofilm and motility regulator AbfR1 in *Sulfolobus acidocaldarius*. *Mol Microbiol* **105**: 777-793.

Li L, Banerjee A, Bischof LF, et al. (2017) Wing phosphorylation is a major functional determinant of the Lrs14-type biofilm and motility regulator AbfR1 in *Sulfolobus acidocaldarius*. *Mol Microbiol* **105**: 777-793.

Li Y, Kawashima N, Li J, Chandra AP & Gerson AR (2013) A review of the structure, and fundamental mechanisms and kinetics of the leaching of chalcopyrite. *Adv Colloid Interface Sci* **197-198**: 1-32.

Li Y, Pan S, Zhang Y, Ren M, Feng M, Peng N, Chen L, Liang YX & She Q (2016) Harnessing Type I and Type III CRISPR-Cas systems for genome editing. *Nucleic Acids Res* **44**: e34-e34.

Lian H, Zeldes BM, Lipscomb GL, Hawkins AB, Han Y, Loder AJ, Nishiyama D, Adams MW & Kelly RM (2016) Ancillary contributions of heterologous biotin protein ligase and carbonic anhydrase for CO₂ incorporation into 3-hydroxypropionate by metabolically engineered *Pyrococcus furiosus*. *Biotechnol Bioeng* **113**: 2652-2660.

Lillestøl RK, Shah SA, Brügger K, Redder P, Phan H, Christiansen J & Garrett RA (2009) CRISPR families of the crenarchaeal genus *Sulfolobus*: Bidirectional transcription and dynamic properties. *Mol Microbiol* **72**: 259-272.

Limauro D, Cannio R, Fiorentino G, Rossi M & Bartolucci S (2001) Identification and molecular characterization of an endoglucanase gene, *celS*, from the extremely thermophilic archaeon *Sulfolobus solfataricus*. *Extremophiles* **5**: 213-219.

Lindas AC, Karlsson EA, Lindgren MT, Ettema TJ & Bernander R (2008) A unique cell division machinery in the Archaea. *Proc Natl Acad Sci USA* **105**: 18942-18946.

Lipps G (2006) Plasmids and viruses of the thermoacidophilic crenarchaeote *Sulfolobus*. *Extremophiles* **10**: 17-28.

Liu B, Ouyang S, Makarova KS, Xia Q, Zhu Y, Li Z, Guo L, Koonin EV, Liu ZJ & Huang L (2015) A primase subunit essential for efficient primer synthesis by an archaeal eukaryotic-type primase. *Nat Commun* **6**: 1-11.

Liu H, Wang K, Lindås AC & Peeters E (2016) The genome-scale DNA-binding profile of BarR, a β -alanine responsive transcription factor in the archaeon *Sulfolobus acidocaldarius*. *BMC Genomics* **17**: 569.

Liu H, Orell A, Maes D, van Wolferen M, Lindås A-C, Bernander R, Albers S-V, Charlier D & Peeters E (2014) BarR, an Lrp-type transcription factor in *Sulfolobus acidocaldarius*, regulates an aminotransferase gene in a β -alanine responsive manner. *Mol Microbiol* **92**: 625-639.

Liu LJ, Stockdreher Y, Koch T, et al. (2014) Thiosulfate transfer mediated by DsrE/TusA homologs from acidothermophilic sulfur-oxidizing archaeon *Metallosphaera cuprina*. *J Biol Chem* **289**: 26949-26959.

Liu Y, Ishino S, Ishino Y, Pehau-Arnaudet G, Krupovic M & Prangishvili D (2017) A novel type of polyhedral viruses infecting hyperthermophilic Archaea. *J Virology* **91**.

Liu Z & Liu T (2016) Production of acrylic acid and propionic acid by constructing a portion of the 3-hydroxypropionate/4-hydroxybutyrate cycle from *Metallosphaera sedula* in *Escherichia coli*. *J Ind Microbiol Biotechnol* **43**: 1659-1670.

Loder AJ, Han Y, Hawkins AB, Lian H, Lipscomb GL, Schut GJ, Keller MW, Adams MWW & Kelly RM (2016) Reaction kinetic analysis of the 3-hydroxypropionate/4-hydroxybutyrate CO₂ fixation cycle in extremely thermoacidophilic Archaea. *Metab Eng* **38**: 446-463.

Loomis WF, Shaulsky G & Wang N (1997) Histidine kinases in signal transduction pathways of eukaryotes. *J Cell Sci* **110**: 1141-1145.

Lower BH & Kennelly PJ (2002) The membrane-associated protein-serine/threonine kinase from *Sulfolobus solfataricus* is a glycoprotein. *J Bacteriol* **184**: 2614-2619.

Lower BH & Kennelly PJ (2003) Open reading frame sso2387 from the archaeon *Sulfolobus solfataricus* encodes a polypeptide with protein-serine kinase activity. *J Bacteriol* **185**: 3436-3445.

Lower BH, Bischoff KM & Kennelly PJ (2000) The archaeon *Sulfolobus solfataricus* contains a membrane-associated protein kinase activity that preferentially phosphorylates threonine residues *in vitro*. *J Bacteriol* **182**: 3452-3459.

Lower BH, Potters MB & Kennelly PJ (2004) A phosphoprotein from the archaeon *Sulfolobus solfataricus* with protein-serine/threonine kinase activity. *J Bacteriol* **186**: 463-472.

Lubben M, Castresana J & Warne A (1994) Terminal Oxidases of *Sulfolobus* - Genes and proteins. *Syst Appl Microbiol* **16**: 556-559.

Lundgren M, Andersson A, Chen L, Nilsson P & Bernander R (2004) Three replication origins in *Sulfolobus* species: Synchronous initiation of chromosome replication and asynchronous termination. *Proc Natl Acad Sci USA* **101**: 7046-7051.

Ma C, Pathak C, Lee SJ, Lee KY, Jang SB, Nam M, Im H, Yoon HJ & Lee BJ (2016) Alba from *Thermoplasma volcanium* belongs to α -NAT's: An insight into the structural aspects of Tv Alba and its acetylation by Tv Ard1. *Arch Biochem Biophys* **590**: 90-100.

Maezato Y, Johnson T, McCarthy S, Dana K & Blum P (2012) Metal resistance and lithoautotrophy in the extreme thermoacidophile *Metallosphaera sedula*. *J Bacteriol* **194**: 6856-6863.

Maezato Y, Daugherty A, Dana K, Soo E, Cooper C, Tachdjian S, Kelly RM & Blum P (2011) VapC6, a ribonucleolytic toxin regulates thermophilicity in the crenarchaeote *Sulfolobus solfataricus*. *RNA* **17**: 1381-1392.

Mai VQ, Chen X, Hong R & Huang L (1998) Small abundant DNA binding proteins from the thermoacidophilic archaeon *Sulfolobus shibatae* constrain negative DNA supercoils. *J Bacteriol* **180**: 2560-2563.

Makarova KS, Koonin EV & Albers S-V (2016) Diversity and evolution of Type IV pili systems in Archaea. *Front Microbiol* **7**: 667.

Makarova KS, Wolf YI, van der Oost J & Koonin EV (2009) Prokaryotic homologs of Argonaute proteins are predicted to function as key components of a novel system of defense against mobile genetic elements. *Biology Direct* **4**: 1-15.

Makarova KS, Grishin NV, Shabalina SA, Wolf YI & Koonin EV (2006) A putative RNA-interference-based immune system in prokaryotes: computational analysis of the predicted enzymatic machinery, functional analogies with eukaryotic RNAi, and hypothetical mechanisms of action. *Biology Direct* **1**: 7.

Makarova KS, Wolf YI, Iranzo J, Shmakov SA, Alkhnbashi OS, Brouns SJ, Charpentier E, Cheng D, Haft DH & Horvath P (2019) Evolutionary classification of CRISPR–Cas systems: a burst of class 2 and derived variants. *Nat Rev Microbiol* **1**-17.

Maklad HR, Gutierrez GJ, Esser D, Siebers B & Peeters E (2020) Phosphorylation of the acyl-CoA binding pocket of the FadR transcription regulator in *Sulfolobus acidocaldarius*. *Biochimie* **175**: 120-124.

Maklad HR, Gutierrez GJ, Esser D, Siebers B & Peeters E (2020) Phosphorylation of the acyl-CoA binding pocket of the FadR transcription regulator in *Sulfolobus acidocaldarius*. *Biochimie*.

Manica A & Schleper C (2013) CRISPR-mediated defense mechanisms in the hyperthermophilic archaeal genus *Sulfolobus*. *RNA Biology* **10**: 671-678.

Marhuenda-Egea FC & Bonete MaJ (2002) Extreme halophilic enzymes in organic solvents. *Curr Opin Biotech* **13**: 385-389.

Marinsek N, Barry ER, Makarova KS, Dionne I, Koonin EV & Bell SD (2006) GINS, a central nexus in the archaeal DNA replication fork. *EMBO Rep* **7**: 539-545.

Martin A, Yeats S, Janekovic D, Reiter WD, Aicher W & Zillig W (1984) SAV 1, a temperate uv-inducible DNA virus-like particle from the archaeabacterium *Sulfolobus acidocaldarius* isolate B12. *EMBO J* **3**: 2165-2168.

Martins LO & Santos H (1995) Accumulation of mannosylglycerate and di-myo-inositol-phosphate by *Pyrococcus furiosus* in response to salinity and temperature. *Appl Environ Microbiol* **61**: 3299-3303.

Martins LO, Carreto LS, Da Costa MS & Santos H (1996) New compatible solutes related to Di-myo-inositol-phosphate in members of the order *Thermotogales*. *J Bacteriol* **178**: 5644-5651.

Martins LO, Huber R, Huber H, Stetter KO, Da Costa MS & Santos H (1997) Organic solutes in hyperthermophilic Archaea. *Appl Environ Microbiol* **63**: 896-902.

Martusewitsch E, Sensen CW & Schleper C (2000) High spontaneous mutation rate in the hyperthermophilic archaeon *Sulfolobus solfataricus* is mediated by transposable elements. *J Bacteriol* **182**: 2574-2581.

Martínez-Alvarez L, Deng L & Peng X (2017) Formation of a viral replication focus in *Sulfolobus* cells infected by the ravidivirus *Sulfolobus islandicus* rod-shaped Virus 2. *J Virology* **91**.

Martínez-Bussenius C, Navarro CA & Jerez CA (2017) Microbial copper resistance: importance in biohydrometallurgy. *Microbial Biotechnol* **10**: 279-295.

Mathai JC, Missner A, Kugler P, Saparov SM, Zeidel ML, Lee JK & Pohl P (2009) No facilitator required for membrane transport of hydrogen sulfide. *Proc Natl Acad Sci USA* **106**: 16633-16638.

Maupin-Furlow J (2012) Proteasomes and protein conjugation across domains of life. *Nat Rev Microbiol* **10**: 100-111.

Maurelli L, Giovane A, Esposito A, Moracci M, Fiume I, Rossi M & Morana A (2008) Evidence that the xylanase activity from *Sulfolobus solfataricus* Oα is encoded by the endoglucanase precursor gene (sso1354) and characterization of the associated cellulase activity. *Extremophiles* **12**: 689-700.

Mayo-Muñoz D, He F, Jørgensen JB, Madsen PK, Bhoobalan-Chitty Y & Peng X (2018) Anti-CRISPR-based and CRISPR-based genome editing of *Sulfolobus islandicus* rod-shaped virus 2. *Viruses* **10**: 695.

McCarthy S, Ai C & Blum P (2018) Enhancement of *Metallosphaera sedula* bioleaching by targeted recombination and adaptive laboratory evolution. *Adv Appl Microbiol* **104**: 135-165.

McCarthy S, Ai C, Wheaton G, Tevratia R, Eckrich V, Kelly R & Blum P (2014) Role of an archaeal pita transporter in the copper and arsenic resistance of *Metallosphaera sedula*, an extreme thermoacidophile. *J Bacteriol* **196**: 3562-3570.

Meagher M, Epling LB & Enemark EJ (2019) DNA translocation mechanism of the MCM complex and implications for replication initiation. *Nat Commun* **10**: 1-13.

Melo AM, Bandeiras TM & Teixeira M (2004) New insights into type II NAD(P)H:quinone oxidoreductases. *Microbiol Mol Biol Rev* **68**: 603-616.

Meng DC, Shi ZY, Wu LP, Zhou Q, Wu Q, Chen JC & Chen GQ (2012) Production and characterization of poly(3-hydroxypropionate-co-4-hydroxybutyrate) with fully controllable structures by recombinant *Escherichia coli* containing an engineered pathway. *Metab Eng* **14**: 317-324.

Meyer BH & Albers S-V (2014) AglB, catalyzing the oligosaccharyl transferase step of the archaeal N-glycosylation process, is essential in the thermoacidophilic crenarchaeon *Sulfolobus acidocaldarius*. *Microbiologyopen* **3**: 531-543.

Meyer BH, Birich A & Albers S-V (2015) N-Glycosylation of the archaellum filament is not important for archaella assembly and motility, although N-Glycosylation is essential for motility in *Sulfolobus acidocaldarius*. *Biochimie* **118**: 294-301.

Meyer BH, Shams-Eldin H & Albers S-V (2017) AglH, a thermophilic UDP-N-acetylglucosamine-1-phosphate: dolichyl phosphate GlcNAc-1-phosphotransferase initiating protein N-glycosylation pathway in *Sulfolobus acidocaldarius*, is capable of complementing the eukaryal Alg7. *Extremophiles* **21**: 121-134.

Meyer BH, Peyfoon E, Dietrich C, Hitchen P, Panico M, Morris HR, Dell A & Albers S-V (2013) Agl16, a thermophilic glycosyltransferase mediating the last step of N-Glycan biosynthesis in the thermoacidophilic crenarchaeon *Sulfolobus acidocaldarius*. *J Bacteriol* **195**: 2177-2186.

Meyer BH, Zolghadr B, Peyfoon E, et al. (2011) Sulfoquinovose synthase - an important enzyme in the N-glycosylation pathway of *Sulfolobus acidocaldarius*. *Mol Microbiol* **82**: 1150-1163.

Mierzwa BE, Chiaruttini N, Redondo-Morata L, et al. (2017) Dynamic subunit turnover in ESCRT-III assemblies is regulated by Vps4 to mediate membrane remodelling during cytokinesis. *Nat Cell Biol* **19**: 787-798.

Mojica FJM, Díez-Villaseñor C, García-Martínez J & Soria E (2005) Intervening sequences of regularly spaced prokaryotic repeats derive from foreign genetic elements. *J Mol Evol* **60**: 174-182.

Moll R & Schäfer G (1988) Chemiosmotic H⁺ cycling across the plasma membrane of the thermoacidophilic archaebacterium *Sulfolobus acidocaldarius*. *FEBS Lett* **232**: 359-363.

Monroe N, Han H, Gonciarz MD, Eckert DM, Karren MA, Whitby FG, Sundquist WI & Hill CP (2014) The oligomeric state of the active Vps4 AAA ATPase. *J Mol Biol* **426**: 510-525.

Moore BC & Leigh JA (2005) Markerless mutagenesis in *Methanococcus maripaludis* demonstrates roles for alanine dehydrogenase, alanine racemase, and alanine permease. *J Bacteriol* **187**: 972-979.

Moracci M, Nucci R, Febbraio F, Vaccaro C, Vespa N, La Cara F & Rossi M (1995) Expression and extensive characterization of a beta-glycosidase from the extreme thermoacidophilic archaeon *Sulfolobus solfataricus* in *Escherichia coli*: Authenticity of the recombinant enzyme. *Enzyme Microb Technol* **17**: 992-997.

Moracci M, Cobucci Ponzano B, Trincone A, Fusco S, De Rosa M, van Der Oost J, Sensen CW, Charlebois RL & Rossi M (2000) Identification and molecular characterization of the first alpha-xylosidase from an archaeon. *J Biol Chem* **275**: 22082-22089.

Moriscot C, Gribaldo S, Jault JM, Krupovic M, Arnaud J, Jamin M, Schoehn G, Forterre P, Weissenhorn W & Renesto P (2011) Crenarchaeal CdvA forms double-helical filaments containing DNA and interacts with ESCRT-III-like CdvB. *PLoS ONE* **6**.

Mougiakos I, Bosma EF, de Vos WM, van Kranenburg R & van der Oost J (2016) Next generation prokaryotic engineering: the CRISPR-Cas toolkit. *Trends Biotechnol* **34**: 575-587.

Mousavi SM, Yaghmaei S, Vossoughi M, Jafari A & Hoseini SA (2005) Comparison of bioleaching ability of two native mesophilic and thermophilic bacteria on copper recovery from chalcopyrite concentrate in an airlift bioreactor. *Hydrometallurgy* **80**: 139-144.

Mukherjee A, Wheaton GH, Blum PH & Kelly R (2012) Uranium extremophily is an adaptive, rather than intrinsic, feature for extremely thermoacidophilic *Metallosphaera prunae*. *Proc Natl Acad Sci USA* **109**: 16702-16707.

Mukherjee A, Wheaton GH, Counts JA, Ijeomah B, Desai J & Kelly RM (2017) VapC toxins drive cellular dormancy under uranium stress for the extreme thermoacidophile *Metallosphaera prunae*. *Environ Microbiol* **19**: 2831-2842.

Muller FH, Bandeiras TM, Urich T, Teixeira M, Gomes CM & Kletzin A (2004) Coupling of the pathway of sulphur oxidation to dioxygen reduction: characterization of a novel membrane-bound thiosulphate:quinone oxidoreductase. *Mol Microbiol* **53**: 1147-1160.

Munson-Mcgee JH, Snyder JC & Young MJ (2018) Archaeal viruses from high-temperature environments. *Genes* **9**: 128.

Munson-Mcgee JH, Peng S, Dewerff S, Stepanauskas R, Whitaker RJ, Weitz JS & Young MJ (2018) A virus or more in (nearly) every cell: Ubiquitous networks of virus-host interactions in extreme environments. *ISME Journal* **12**: 1706-1714.

Muskhelishvili G, Palm P & Zillig W (1993) SSV1-encoded site-specific recombination system in *Sulfolobus shibatae*. *Mol & Gen Genetics* **237**: 334-342.

Märtens B, Manoharadas S, Hasenöhrl D, Manica A & Bläsi U (2013) Antisense regulation by transposon-derived RNAs in the hyperthermophilic archaeon *Sulfolobus solfataricus*. *EMBO Rep* **14**: 527-533.

Müller FH, Bandeiras TM, Urich T, Teixeira M, Gomes CM & Kletzin A (2004) Coupling of the pathway of sulphur oxidation to dioxygen reduction: characterization of a novel membrane-bound thiosulphate:quinone oxidoreductase. *Mol Microbiol* **53**: 1147-1160.

Neale JW, Robertson SW, Muller HH & Gericke M (2009) Integrated piloting of a thermophilic bioleaching process for the treatment of a low-grade nickel-copper sulphide concentrate. *J S Afr Inst Min Metall* **109**: 273-293.

Nguyen-Duc T, van Oeffelen L, Song N, Hassanzadeh-Ghassabeh G, Muyldermans S, Charlier D & Peeters E (2013) The genome-wide binding profile of the *Sulfolobus solfataricus* transcription factor Ss-LrpB shows binding events beyond direct transcription regulation. *BMC Genomics* **14**: 1-15.

Nicolaus B, Gambacorta A, Basso AL, Riccio R, De Rosa M & Grant WD (1988) Trehalose in Archaeabacteria. *Sys Appl Microbiol* **10**: 215-217.

Niewohner O, Garcia-Doval C, Rostøl JT, Berk C, Schwede F, Bigler L, Hall J, Marraffini LA & Jinek M (2017) Type III CRISPR-Cas systems produce cyclic oligoadenylate second messengers. *Nature* **548**: 543-548.

Niu Y, Xia Y, Wang S, et al. (2013) A prototypic lysine methyltransferase 4 from Archaea with degenerate sequence specificity methylates chromatin proteins Sul7d and Cren7 in different patterns. *J Biol Chem* **288**: 13728-13740.

Noh KH & Oh DK (2009) Production of the rare ginsenosides compound K, compound Y, and compound Mc by a thermostable beta-glycosidase from *Sulfolobus acidocaldarius*. *Biol Pharm Bull* **32**: 1830-1835.

Nordstrom KD, McCleskey BR & Ball JW (2009) Sulfur geochemistry of hydrothermal waters in Yellowstone National Park: IV Acid-sulfate waters. *Appl Geochem* **24**: 191-207.

Norris PR, Clark DA, Owen JP & Waterhouse S (1996) Characteristics of *Sulfovbacillus acidophilus* sp. nov. and other moderately thermophilic mineral-sulphide-oxidizing bacteria. *Microbiologyopen* **142**: 775-783.

Nunn CE, Johnsen U, Schonheit P, Fuhrer T, Sauer U, Hough DW & Danson MJ (2010) Metabolism of pentose sugars in the hyperthermophilic archaea *Sulfolobus solfataricus* and *Sulfolobus acidocaldarius*. *J Biol Chem* **285**: 33701-33709.

Nunn CEM, Johnsen U, Schönheit P, Fuhrer T, Sauer U, Hough DW & Danson MJ (2010) Metabolism of pentose sugars in the hyperthermophilic archaea *Sulfolobus solfataricus* and *Sulfolobus acidocaldarius*. *J Biol Chem* **285**: 33701-33709.

Obita T, Saksena S, Ghazi-Tabatabai S, Gill DJ, Perisic O, Emr SD & Williams RL (2007) Structural basis for selective recognition of ESCRT-III by the AAA ATPase Vps4. *Nature* **449**: 735-739.

Odling-Smee FJ, Laland KN & Feldman MW (1996) Niche Construction. *The American Naturalist* **147**: 641-648.

Ogino H, Ishino S, Kohda D & Ishino Y (2017) The RecJ2 protein in the thermophilic archaeon *Thermoplasma acidophilum* is a 3'-5' exonuclease that associates with a DNA replication complex. *J Biol Chem* **292**: 7921-7931.

Oke M, Kerou M, Liu H, Peng X, Garrett RA, Prangishvili D, Naismith JH & White MF (2011) A dimeric rep protein initiates replication of a linear archaeal virus genome: Implications for the rep mechanism and viral replication. *J Virology* **85**: 925-931.

Oke M, Carter LG, Johnson KA, et al. (2010) The scottish structural proteomics facility: Targets, methods and outputs. *J Struct Funct Genomics* **11**: 167-180.

Orell A, Remonsellez F, Arancibia R & Jerez CA (2013) Molecular characterization of copper and cadmium resistance determinants in the biomining thermoacidophilic archaeon *Sulfolobus metallicus*. *Archaea* **2013**: 289236.

Orell A, Navarro CA, Rivero M, Aguilar JS & Jerez CA (2012) Inorganic polyphosphates in extremophiles and their possible functions. *Extremophiles* **16**: 573-583.

Orell A, Tripp V, Aliaga-Tobar V, Albers SV, Maracaja-Coutinho V & Randau L (2018) A regulatory RNA is involved in RNA duplex formation and biofilm regulation in *Sulfolobus acidocaldarius*. *Nucleic Acids Res* **46**: 4794-4806.

Orell A, Peeters E, Vassen V, Jachlewski S, Schalles S, Siebers B & Albers S-V (2013) Lrs14 transcriptional regulators influence biofilm formation and cell motility of *Crenarchaea*. *ISME J* **7**: 1886-1898.

Ortmann AC, Brumfield SK, Walther J, McInerney K, Brouns SJ, van de Werken HJG, Bothner B, Douglas T, van de Oost J & Young MJ (2008) Transcriptome analysis of infection of the archaeon *Sulfolobus solfataricus* with *Sulfolobus* turreted icosahedral virus. *J Virology* **82**: 6874.

Palmieri G, Balestrieri M, Peter-Katalinić J, Pohlentz G, Rossi M, Fiume I & Pocsfalvi G (2013) Surface-exposed glycoproteins of hyperthermophilic *Sulfolobus solfataricus* P2 show a common N-glycosylation profile. *J Proteome Res* **12**: 2779-2790.

Papon N & Stock AM (2019) What do archaeal and eukaryotic histidine kinases sense? *F1000Research* **8**.

Park YJ, Yoo CB, Choi SY & Lee HB (2006) Purifications and characterizations of a ferredoxin and its related 2-oxoacid:ferredoxin oxidoreductase from the hyperthermophilic archaeon, *Sulfolobus solfataricus* P1. *J Biochem Mol Biol* **39**: 46-54.

Paula FS, Chin JP, Schnürer A, et al. (2019) The potential for polyphosphate metabolism in Archaea and anaerobic polyphosphate formation in *Methanosarcina mazei*. *Sci Rep* **9**: 17101.

Pauly MD, Bautista MA, Black JA & Whitaker RJ (2019) Diversified local CRISPR-Cas immunity to viruses of *Sulfolobus islandicus*. *Phil Trans Royal Soc B: Biol Sci* **374**.

Payne KAP, Hough DW & Danson MJ (2010) Discovery of a putative acetoin dehydrogenase complex in the hyperthermophilic archaeon *Sulfolobus solfataricus*. *FEBS Lett* **584**: 1231-1234.

Payne S, McCarthy S, Johnson T, North E & Blum P (2018) Nonmutational mechanism of inheritance in the Archaeon *Sulfolobus solfataricus*. *Proc Natl Acad Sci USA* **115**: 12271-12276.

Paytubi S & White MF (2009) The crenarchaeal DNA damage-inducible transcription factor B parologue TFB3 is a general activator of transcription. *Mol Microbiol* **72**: 1487-1499.

Pearson A, Hurley SJ, Elling FJ & Wilkes EB (2019) CO₂-dependent carbon isotope fractionation in Archaea, Part I: Modeling the 3HP/4HB pathway. *Geochim Cosmochim Acta* **261**: 368-382.

Peeples TL & Kelly RM (1995) Bioenergetic response of the extreme thermoacidophile *Metallosphaera sedula* to thermal and nutritional stresses. *Appl Environ Microbiol* **61**: 2314-2321.

Peeters E, Peixeiro N & Sezonov G (2013) Cis-regulatory logic in archaeal transcription. *Biochem Soc T* **41**: 326-331.

Peeters E, Driessens RP, Werner F & Dame RT (2015) The interplay between nucleoid organization and transcription in archaeal genomes. *Nat Rev Microbiol* **13**: 333-341.

Peng X, Blum H, She Q, Mallon S, Brügger K, Garrett RA, Zillig W & Prangishvili D (2001) Sequences and replication of genomes of the archaeal ravidiviruses SIRV1 and SIRV2: relationships to the archaeal lipothrixvirus SIFV and some eukaryal viruses. *Virology* **291**: 226-234.

Perez-Rueda E, Hernandez-Guerrero R, Martinez-Nuñez MA, Armenta-Medina D, Sanchez I & Ibarra JA (2018) Abundance, diversity and domain architecture variability in prokaryotic DNA-binding transcription factors. *PLoS ONE* **13**: e0195332.

Petrova OE & Sauer K (2012) Sticky situations: Key components that control bacterial surface attachment. *J Bacteriol* **194**: 2413 LP-2425.

Peyfoon E, Meyer B, Hitchen PG, Panico M, Morris HR, Haslam SM, Albers S-V & Dell A (2010) The S-layer glycoprotein of the crenarchaeote *Sulfolobus acidocaldarius* is glycosylated at multiple sites with chitobiose-linked N-glycans. *Archaea* **2010**.

Pietilä MK, Demina TA, Atanasova NS, Oksanen HM & Bamford DH (2014) Archaeal viruses and bacteriophages: comparisons and contrasts. *Trends Microbiol* **22**: 334-344.

Pina M, Bize A, Forterre P & Prangishvili D (2011) The archeoviruses. *FEMS Microbiol Rev* **35**: 1035-1054.

Pina M, Basta T, Quax TE, Joubert A, Baconnais S, Cortez D, Lambert S, Le Cam E, Bell SD & Forterre P (2014) Unique genome replication mechanism of the archaeal virus AFV 1. *Mol Microbiol* **92**: 1313-1325.

Plagens A, Richter H, Charpentier E & Randau L (2015) DNA and RNA interference mechanisms by CRISPR-Cas surveillance complexes. *FEMS Microbiol Rev* **39**: 442-463.

Plumb JJ, Haddad CM, Gibson JaE & Franzmann PD (2007) *Acidianus sulfidivorans* sp. nov., an extremely acidophilic, thermophilic archaeon isolated from a solfatara on Lihir Island, Papua New Guinea, and emendation of the genus description. *Int J Syst Evol Microbiol* **57**: 1418-1423.

Poranen MM, Daugelavičius R & Bamford DH (2002) Common principles in viral entry. *Ann Rev Microbiol* **53**: 521-538.

Pourcel C, Salvignol G & Vergnaud G (2005) CRISPR elements in *Yersinia pestis* acquire new repeats by preferential uptake of bacteriophage DNA, and provide additional tools for evolutionary studies. *Microbiologyopen* **151**: 653-663.

Pradhan N, Nathsarma KC, Rao KS, Sukla LB & Mishra BK (2008) Heap bioleaching of chalcopyrite: A review. *Minerals Eng* **21**: 355-365.

Prangishvili D (2015) Archaeal viruses: Living fossils of the ancient virosphere? *Ann NY Acad Sci* **1341**: 35-40.

Prangishvili D & Garrett RA (2004) Exceptionally diverse morphotypes and genomes of crenarchaeal hyperthermophilic viruses. *Biochem Soc Trans* **32**: 204-208.

Prangishvili D & Quax TE (2011) Exceptional virion release mechanism: One more surprise from archaeal viruses. *Curr Opin Microbiol* **14**: 315-320.

Prangishvili D, Stedman K & Zillig W (2001) Viruses of the extremely thermophilic archaeon *Sulfolobus*. *Trends Microbiol* **9**: 39-43.

Prangishvili D, Forterre P & Garrett RA (2006) Viruses of the Archaea: A unifying view. *Nat Rev Microbiol* **4**: 837-848.

Prangishvili D, Koonin EV & Krupovic M (2013) Genomics and biology of Ravidviruses, a model for the study of virus-host interactions in Archaea. *Biochem Soc Trans* **41**: 443-450.

Prangishvili D, Bamford DH, Forterre P, Iranzo J, Koonin EV & Krupovic M (2017) The enigmatic archaeal virosphere. *Nat Rev Microbiol* **15**: 724-739.

Prangishvili D, Arnold HP, Götz D, Ziese U, Holz I, Kristjansson JK & Zillig W (1999) A novel virus family, the Ravidviridae: Structure, virus-host interactions and genome variability of the *Sulfolobus* viruses SIRV1 and SIRV2. *Genetics* **152**: 1387-1396.

Prangishvili D, Vestergaard G, Häring M, Aramayo R, Basta T, Rachel R & Garrett RA (2006) Structural and genomic properties of the hyperthermophilic archaeal virus ATV with an extracellular stage of the reproductive cycle. *J Mol Biol* **359**: 1203-1216.

Prangishvili DA, Vashakidze RP, Chelidze MG & Gabriadze I (1985) A restriction endonuclease SuaI from the thermoacidophilic archaebacterium *Sulfolobus acidocaldarius*. *FEBS Lett* **192**: 57-60.

Pritchett MA, Zhang JK & Metcalf WW (2004) Development of a markerless genetic exchange method for *Methanoscincus acetivorans* C2A and its use in construction of new genetic tools for methanogenic archaea. *Appl Environ Microbiol* **70**: 1425-1433.

Prokofeva MI, Miroshnichenko ML, Kostrikina NA, Chernyh NA, Kuznetsiv BB, Tourova TP & Bonch-Osmolovskaya EA (2000) *Acidilobus aceticus* gen. nov., sp. nov., a novel anaerobic thermoacidophilic archaeon from continental hot vents in Kamchatka. *Int J Syst Evol Microbiol* **50**: 2001-2008.

Prokofeva MI, Kostrikina NA, Kolganova TV, Tourova TP, Lysenko AM, Lebedinsky AV & Bonch-Osmolovskaya EA (2009) Isolation of the anaerobic thermoacidophilic crenarchaeote *Acidilobus saccharovorans* sp. nov. and proposal of *Acidilobales* ord. nov., including *Acidilobaceae* fam. nov. and *Caldisphaeraceae* fam. nov. *Int J Syst Evol Microbiol* **59**: 3116-3122.

Protze J, Müller F, Lauber K, Naß B, Mentele R, Lottspeich F & Kletzin A (2011) An extracellular tetrathionate hydrolase from the thermoacidophilic archaeon *Acidianus ambivalens* with an activity optimum at pH 1. *Front Microbiol* **2**.

Prüschenk R, Baumeister W & Zillig W (1987) Surface structure variants in different species of *Sulfolobus*. *FEMS Microbiol Lett* **43**: 327-330.

Pulschen AA, Mutavchiev DR, Culley S, *et al.* (2020) Live imaging of a hyperthermophilic archaeon reveals distinct roles for two ESCRT-III homologs in ensuring a robust and symmetric division. *Curr Biol* **30**: 2852-2859 e2854.

Purschke WG, Schmidt CL, Petersen A & Schafer G (1997) The terminal quinol oxidase of the hyperthermophilic archaeon *Acidianus ambivalens* exhibits a novel subunit structure and gene organization. *J Bacteriol* **179**: 1344-1353.

Pérez-Rueda E & Janga SC (2010) Identification and genomic analysis of transcription factors in archaeal genomes exemplifies their functional architecture and evolutionary origin. *Mol Biol Evol* **27**: 1449-1459.

Quax TE, Krupovič M, Lucas S, Forterre P & Prangishvili D (2010) The *Sulfolobus* rod-shaped virus 2 encodes a prominent structural component of the unique virion release system in Archaea. *Virology* **404**: 1-4.

Quax TE, Lucas S, Reimann J, Pehau-Arnaudet G, Prevost M-C, Forterre P, Albers S-V & Prangishvili D (2011) Simple and elegant design of a virion egress structure in Archaea. *Proc Natl Acad Sci USA* **108**: 3354-3359.

Quax TEF & Daum B (2017) Structure and assembly mechanism of virus-associated pyramids. *Biophys Rev* **10**: 551-557.

Quax TEF, Voet M, Sismeiro O, *et al.* (2013) Massive activation of archaeal defense genes during viral infection. *J Virology* **87**: 8419-8428.

Quehenberger J, Pittenauer E, Allmaier G & Spadiut O (2020) The influence of the specific growth rate on the lipid composition of *Sulfolobus acidocaldarius*. *Extremophiles* **24**: 413-420.

Quehenberger J, Shen L, Albers S-V, Siebers B & Spadiut O (2017) *Sulfolobus* – A Potential key organism in future biotechnology. *Front Microbiol* **8**.

Quehenberger J, Albersmeier A, Glatzel H, Hackl M, Kalinowski J & Spadiut O (2019) A defined cultivation medium for *Sulfolobus acidocaldarius*. *J Biotechnol* **301**: 56-67.

Quemin ER, Lucas S, Daum B, Quax TE, Kühlbrandt W, Forterre P, Albers S-V, Prangishvili D & Krupovic M (2013) First insights into the entry process of hyperthermophilic archaeal viruses. *J Virology* **87**: 13379-13385.

Quemin ERJ & Quax TEF (2015) Archaeal viruses at the cell envelope: entry and egress. *Front Microbiol* **6**: 552-552.

Quemin ERJ, Chlanda P, Sachse M, Forterre P, Prangishvili D & Krupovic M (2016) Eukaryotic-like virus budding in Archaea. *mBio* **7**.

Qureshi SA, Bell SD & Jackson SP (1997) Factor requirements for transcription in the Archaeon *Sulfolobus shibatae*. *EMBO J* **16**: 2927-2936.

Ramos-Vera WH, Weiss M, Strittmatter E, Kockelkorn D & Fuchs G (2011) Identification of missing genes and enzymes for autotrophic carbon fixation in *Crenarchaeota*. *J Bacteriol* **193**: 1201-1211.

Rashid MH, Rumbaugh K, Passador L, Davies DG, Hamood AN, Iglewski BH & Kornberg A (2000) Polyphosphate kinase is essential for biofilm development, quorum sensing, and virulence of *Pseudomonas aeruginosa*. *Proc Natl Acad Sci USA* **97**: 9636-9641.

Rawls KS, Martin JH & Maupin-Furlow JA (2011) Activity and transcriptional regulation of bacterial protein-like Glycerol-3-Phosphate Dehydrogenase of the Haloarchaea in *Haloferax volcanii*. *J Bacteriol* **193**: 4469-4476.

Ray WK, Potters MB, Haile JD & Kennelly PJ (2015) Activation of SsoPK4, an Archaeal eIF2 α kinase homolog, by oxidized CoA. *Proteomes* **3**: 89-116.

Ray WK, Keith SM, DeSantis AM, Hunt JP, Larson TJ, Helm RF & Kennelly PJ (2005) A phosphohexamutase from the archaeon *Sulfolobus solfataricus* is covalently modified by phosphorylation on serine. *J Bacteriol* **187**: 4270-4275.

Redder P & Garrett RA (2006) Mutations and rearrangements in the genome of *Sulfolobus solfataricus* P2. *J Bacteriol* **188**: 4198-4206.

Reilly MS & Grogan DW (2002) Biological effects of DNA damage in the hyperthermophilic archaeon *Sulfolobus acidocaldarius*. *FEMS Microbiol Lett* **208**: 29-34.

Reimann J, Lassak K, Khadouma S, Ettema TJ, Yang N, Driessen AJ, Klingl A & Albers SV (2012) Regulation of archaella expression by the FHA and von Willebrand domain-containing proteins ArnA and ArnB in *Sulfolobus acidocaldarius*. *Mol Microbiol* **86**: 24-36.

Reimann J, Esser D, Orell A, Amman F, Pham TK, Noirel J, Lindås A-C, Bernander R, Wright PC & Siebers B (2013) Archaeal signal transduction: impact of protein phosphatase deletions on cell size, motility, and energy metabolism in *Sulfolobus acidocaldarius*. *Mol Cel Proteomics* **12**: 3908-3923.

Reina-Bueno M, Argandoña M, Nieto JJ, Hidalgo-García A, Iglesias-Guerra F, Delgado MJ & Vargas C (2012) Role of trehalose in heat and desiccation tolerance in the soil bacterium *Rhizobium etli*. *BMC Microbiol* **12**: 207.

Reinhardt A, Johnsen U & Schönheit P (2019) l-Rhamnose catabolism in Archaea. *Mol Microbiol* **111**: 1093-1108.

Remonsellez F, Orell A & Jerez CA (2006) Copper tolerance of the thermoacidophilic archaeon *Sulfolobus metallicus*: possible role of polyphosphate metabolism. *Microbiologyopen* **152**: 59-66.

Reno ML, Held NL, Fields CJ, Burke PV & Whitaker RJ (2009) Biogeography of the *Sulfolobus islandicus* pan-genome. *Proc Natl Acad Sci USA* **106**: 8605-8610.

Reysenbach A-L, Banta AB, Boone DR, Cary SC & Luther GW (2000) Microbial essentials at hydrothermal vents. *Nature* **404**: 835.

Reysenbach AL, Liu Y, Banta AB, Beveridge TJ, Kirshtein JD, Schouten S, Tivey MK, Von Damm KL & Voytek MA (2006) A ubiquitous thermoacidophilic archaeon from deep-sea hydrothermal vents. *Nature* **442**: 444-447.

Rice G, Tang L, Stedman K, Roberto F, Spuhler J, Gillitzer E, Johnson JE, Douglas T & Young M (2004) The structure of a thermophilic archaeal virus shows a double-stranded DNA viral capsid type that spans all domains of life. *Proc Natl Acad Sci USA* **101**: 7716-7720.

Richey D & Lin E (1972) Importance of facilitated diffusion for effective utilization of glycerol by *Escherichia coli*. *J Bacteriol* **112**: 784-790.

Rinker KD & Kelly RM (1996) Growth physiology of the hyperthermophilic archaeon *Thermococcus litoralis*: Development of a sulfur-free defined medium, characterization of an exopolysaccharide, and evidence of biofilm formation. *Appl Environ Microbiol* **62**: 4478-4485.

Risa GT, Hurtig F, Bray S, *et al.* (2019) Proteasome-mediated protein degradation resets the cell division cycle and triggers ESCRT-III-mediated cytokinesis in an archaeon. *bioRxiv* 774273-774273.

Rivero M, Torres-Paris C, Munoz R, Cabrera R, Navarro CA & Jerez CA (2018) Inorganic polyphosphate, exopolyphosphatase, and Pho84-like transporters may be involved in copper resistance in *Metallosphaera sedula* DSM 5348^T. *Archaea* **2018**: 5251061.

Robinson NP & Bell SD (2007) Extrachromosomal element capture and the evolution of multiple replication origins in archaeal chromosomes. *Proc Natl Acad Sci USA* **104**: 5806-5811.

Robinson NP, Dionne I, Lundgren M, Marsh VL, Bernander R & Bell SD (2004) Identification of two origins of replication in the single chromosome of the archaeon *Sulfolobus solfataricus*. *Cell* **116**: 25-38.

Rolfsmeier M, Haseltine C, Bini E, Clark A & Blum P (1998) Molecular characterization of the alpha-glucosidase gene (malA) from the hyperthermophilic archaeon *Sulfolobus solfataricus*. *J Bacteriol* **180**: 1287-1295.

Rostøl JT & Marraffini LA (2019) Non-specific degradation of transcripts promotes plasmid clearance during type III-A CRISPR–Cas immunity. *Nature Microbiol* **4**: 656-662.

Rothschild LJ & Mancinelli RL (2001) Life in extreme environments. *Nature* **409**: 1092-1101.

Rouillon C, Athukoralage JS, Graham S, Grüsschow S & White MF (2018) Control of cyclic oligoadenylate synthesis in a type III CRISPR system. *eLife* **7**.

Rowland EF, Bautista MA, Zhang C & Whitaker RJ (2020) Surface resistance to SSVs and SIRVs in pilin deletions of *Sulfolobus islandicus*. *Mol Microbiol* **113**: 718-727.

Ruggero D & Londei P (1996) Differential antibiotic sensitivity determined by the large ribosomal subunit in thermophilic archaea. *J Bacteriol* **178**: 3396-3398.

Sakai HD & Kurosawa N (2017) *Sulfodiicoccus acidiphilus* gen. nov., sp. nov., a sulfur-inhibited thermoacidophilic archaeon belonging to the order *Sulfolobales* isolated from a terrestrial acidic hot spring. *Int J Syst Evol Microbiol* **67**: 1880-1886.

Sakai HD & Kurosawa N (2018) *Saccharolobus caldissimus* gen. nov., sp. nov., a facultatively anaerobic iron-reducing hyperthermophilic archaeon isolated from an acidic terrestrial hot spring, and reclassification of *Sulfolobus solfataricus* as *Saccharolobus solfataricus* comb. nov. and *Sulfolobus shibatae* as *Saccharolobus shibatae* comb. nov. *Int J Syst Evol Microbiol* **68**: 1271-1278.

Sakai HD & Kurosawa N (2019) Complete genome sequence of the *Sulfodiicoccus acidiphilus* strain HS-1(T), the first crenarchaeon that lacks *polB3*, isolated from an acidic hot spring in Ohwaku-dani, Hakone, Japan. *BMC Res Notes* **12**: 444.

Sakofsky CJ, Runck LA & Grogan DW (2011) *Sulfolobus* mutants, generated via PCR products, which lack putative enzymes of UV photoproduct repair. *Archaea* **2011**: 864015.

Samson RY, Abeyrathne PD & Bell SD (2016) Mechanism of archaeal MCM helicase recruitment to DNA replication origins. *Mol Cell* **61**: 287-296.

Samson RY, Obita T, Freund SM, Williams RL & Bell SD (2008) A role for the ESCRT system in cell division in *Archaea*. *Science* **322**: 1710-1713.

Samson RY, Obita T, Hodgson B, Shaw MK, Chong PLG, Williams RL & Bell SD (2011) Molecular and structural basis of ESCRT-III recruitment to membranes during archaeal cell division. *Mol Cell* **41**: 186-196.

Samson RY, Xu Y, Gadelha C, *et al.* (2013) Specificity and function of archaeal DNA replication initiator proteins. *Cell Rep* **3**: 485-496.

Sand W, Gehrke T, Jozsa PG & Schippers A (2001) (Bio) chemistry of bacterial leaching - direct vs. indirect bioleaching. *Hydrometallurgy* **59**: 159-175.

Santos H & Da Costa MS (2002) Compatible solutes of organisms that live in hot saline environments. *Environ Microbiol* **4**: 501-509.

Sanz JL, Huber G, Huber H & Amils R (1994) Using protein synthesis inhibitors to establish the phylogenetic relationships of the *Sulfolobales* order. *J Mol Evol* **39**: 528-532.

Sato T, Fukui T, Atomi H & Imanaka T (2005) Improved and versatile transformation system allowing multiple genetic manipulations of the hyperthermophilic archaeon *Thermococcus kodakaraensis*. *Appl Environ Microbiol* **71**: 3889-3899.

Say RF & Fuchs G (2010) Fructose 1,6-bisphosphate aldolase/phosphatase may be an ancestral gluconeogenic enzyme. *Nature* **464**: 1077-1081.

Schafer G, Moll R & Schmidt CL (2001) Respiratory enzymes from *Sulfolobus acidocaldarius*. *Methods Enzymol* **331**: 369-410.

Schaller GE, Shiu S-H & Armitage JP (2011) Two-component systems and their co-option for eukaryotic signal transduction. *Curr Biology* **21**: R320-R330.

Scheele U, Erdmann S, Ungewickell EJ, Felisberto-Rodrigues C, Ortiz-Lombardía M & Garrett RA (2011) Chaperone role for proteins p618 and p892 in the extracellular tail development of *Acidianus* two-tailed virus. *J Virology* **85**: 4812-4821.

Schelert J, Drozda M, Dixit V, Dillman A & Blum P (2006) Regulation of mercury resistance in the Crenarchaeote *Sulfolobus solfataricus*. *J Bacteriol* **188**: 7141-7150.

Schelert J, Dixit V, Hoang V, Simbahan J, Drozda M & Blum P (2004) Occurrence and characterization of mercury resistance in the hyperthermophilic archaeon *Sulfolobus solfataricus* by use of gene disruption. *J Bacteriol* **186**: 427-437.

Schippers A, Jozsa P & Sand W (1996) Sulfur chemistry in bacterial leaching of pyrite. *Appl Environ Microbiol* **62**: 3424-3431.

Schleper C, Kubo K & Zillig W (1992) The particle SSV1 from the extremely thermophilic archaeon *Sulfolobus* is a virus: Demonstration of infectivity and of transfection with viral DNA. *Proc Natl Acad Sci USA* **89**: 7645-7649.

Schleper C, Puehler G, Holz I, Gambacorta A, Janekovic D, Santarius U, Klenk HP & Zillig W (1995) *Picrophilus* gen. nov., fam. nov.: a novel aerobic, heterotrophic, thermoacidophilic genus and family comprising archaea capable of growth around pH 0. *J Bacteriol* **177**: 7050-7059.

Schleper C, G. Puhler, H.-P. Klenk, and W. Zillig (1996) *Picrophilus oshimae* and *Picrophilus torridus* fam. nov., gen. nov., sp. nov., two species of hyperacidophilic, thermophilic, heterotrophic, aerobic archaea. *J Syst Evol Microbiol* **46**: 814-816.

Schlitt WJ (2006) Kennecott's million-ton test heap - The active leach program. *Miner Metall Proc* **23**: 1-16.

Schocke L, Bräsen C & Siebers B (2019) Thermoacidophilic *Sulfolobus* species as source for extremozymes and as novel archaeal platform organisms. *Curr Opin Biotechnol* **59**: 71-77.

Schoehn G, Quaite-Randall E, Jiménez JL, Joachimiak A & Saibil HR (2000) Three conformations of an archaeal chaperonin, TF55 from *Sulfolobus shibatae*. *J Mol Biol* **296**: 813-819.

Schult F, Le TN, Albersmeier A, Rauch B, Blumenkamp P, Van Der Does C, Goesmann A, Kalinowski J, Albers SV & Siebers B (2018) Effect of UV irradiation on *Sulfolobus acidocaldarius* and involvement of the general transcription factor TFB3 in the early UV response. *Nucleic Acids Res* **46**: 7179-7192.

Schäfer G (1996) Bioenergetics of the archaeabacterium *Sulfolobus*. *Biochim Biophys Acta - Bioenergetics* **1277**: 163-200.

Scott JW, Poole FL & Adams MW (2014) Characterization of ten heterotetrameric NDP-Dependent Acyl-CoA Synthetases of the hyperthermophilic archaeon *Pyrococcus furiosus*. *Archaea* **2014**: 176863.

Segerer A, Neuner A, Kristjansson JK & Stetter KO (1986) *Acidianus infernus* gen. nov., sp. nov., and *Acidianus brierleyi* Comb. nov. - facultatively aerobic, extremely acidophilic thermophilic sulfur-metabolizing archaeabacteria. *Int J Syst Evol Microbiol* **36**: 559-564.

Segerer A LT, Stetter KO. (1988) *Thermoplasma acidophilum* and *Thermoplasma volcanium* sp. nov. from solfatara fields. *Syst Appl Microbiol* **10**: 161-171.

Segerer AH, Trincone A, Gahrtz M & Stetter KO (1991) *Stygiolobus azoricus* gen. nov., sp. nov. represents a novel genus of anaerobic, extremely thermoacidophilic archaeabacteria of the order *Sulfolobales*. *Int J Syst Bacteriol* **41**: 495-501.

Serre MC, Letzelter C, Garel JR & Duguet M (2002) Cleavage properties of an archaeal site-specific recombinase, the SSV1 integrase. *J Biol Chem* **277**: 16758-16767.

She Q, Zhang C, Deng L, Peng N, Chen Z & Liang YX (2009) Genetic analyses in the hyperthermophilic archaeon *Sulfolobus islandicus*. *Biochem Soc Trans* **37**: 92-96.

She Q, Zhang C, Deng L, Peng N, Chen Z & Liang YX (2009) Genetic analyses in the hyperthermophilic archaeon *Sulfolobus islandicus*. *Biochem Soci Trans* **37**: 92-96.

She Q, Singh RK, Confalonieri F, et al. (2001) The complete genome of the crenarchaeon *Sulfolobus solfataricus* P2. *Proc Natl Acad Sci USA* **98**: 7835.

Sherwood KE, Cano DJ & Maupin-Furlow JA (2009) Glycerol-mediated repression of glucose metabolism and glycerol kinase as the sole route of glycerol catabolism in the haloarchaeon *Haloferax volcanii*. *J Bacteriol* **191**: 4307-4315.

Shi L, Potts M & Kennelly PJ (1998) The serine, threonine, and/or tyrosine-specific protein kinases and protein phosphatases of prokaryotic organisms: a family portrait. *FEMS Microbiol Rev* **22**: 229-253.

Shi X, Rao NN & Kornberg A (2004) Inorganic polyphosphate in *Bacillus cereus*: motility, biofilm formation, and sporulation. *Proc Natl Acad Sci USA* **101**: 17061-17065.

Shi Y (2009) Serine/threonine phosphatases: mechanism through structure. *Cell* **139**: 468-484.

Shin KC, Oh HJ, Kim BJ & Oh DK (2013) Complete conversion of major protopanaxadiol ginsenosides to compound K by the combined use of α -L-arabinofuranosidase and β -galactosidase from *Caldicellulosiruptor saccharolyticus* and β -glucosidase from *Sulfolobus acidocaldarius*. *J Biotechnol* **167**: 33-40.

Singer GAC & Hickey DA (2003) Thermophilic prokaryotes have characteristic patterns of codon usage, amino acid composition and nucleotide content. *Gene* **317**: 39-47.

Skórko R (1984) Protein phosphorylation in the archaebacterium *Sulfolobus acidocaldarius*. *Eur J Biochem* **145**: 617-622.

Smith SC, Kennelly PJ & Potts M (1997) Protein-tyrosine phosphorylation in the Archaea. *J Bacteriol* **179**: 2418-2420.

Snyder JC, Brumfield SK, Peng N, She Q & Young MJ (2011) *Sulfolobus* turreted icosahedral Virus c92 protein responsible for the formation of pyramid-like cellular lysis structures. *J Virology* **85**: 6287-6292.

Snyder JC, Stedman K, Rice G, Wiedenheft B, Spuhler J & Young MJ (2003) Viruses of hyperthermophilic Archaea. *Res Microbiol* **154**: 474-482.

Soto DF, Recalde A, Orell A, Albers S-V, Paradela A, Navarro CA & Jerez CA (2019) Global effect of the lack of inorganic polyphosphate in the extremophilic archaeon *Sulfolobus solfataricus*: A proteomic approach. *J Proteomics* **191**: 143-152.

Spudich JL & Stoeckenius W (1980) Light-regulated retinal-dependent reversible phosphorylation of *Halobacterium* proteins. *J Biol Chem* **255**: 5501-5503.

Stancik IA, Šestak MS, Ji B, Axelson-Fisk M, Franjevic D, Jers C, Domazet-Lošo T & Mijakovic I (2018) Serine/Threonine protein kinases from Bacteria, Archaea and Eukarya share a common evolutionary origin deeply rooted in the tree of life. *J Mol Biol* **430**: 27-32.

Stark H, Wolf J, Albersmeier A, Pham TK, Hofmann JD, Siebers B, Kalinowski J, Wright PC, Neumann-Schaal M & Schomburg D (2017) Oxidative Stickland reactions in an obligate aerobic organism - amino acid catabolism in the *Crenarchaeon Sulfolobus solfataricus*. *FEBS J* **284**: 2078-2095.

Stedman KM, Kosmicki NR & Diemer GS (2013) Codon usage frequency of RNA virus genomes from high-temperature acidic-environment metagenomes. *J Virology* **87**: 1919-1919.

Stedman KM, Schleper C, Rumpf E & Zillig W (1999) Genetic requirements for the function of the archaeal virus SSV1 in *Sulfolobus solfataricus*: construction and testing of viral shuttle vectors. *Genetics* **152**: 1397-1405.

Steudel R (1996) Mechanism for the formation of elemental sulfur from aqueous sulfide in chemical and microbiological desulfurization processes. *Ind Eng Chem Res* **35**: 7.

Straub CT, Counts JA, Nguyen DMN, *et al.* (2018) Biotechnology of extremely thermophilic archaea. *FEMS Microbiol Rev* **42**: 543-578.

Stroud RM, Miercke LJ, O'Connell J, Khademi S, Lee JK, Remis J, Harries W, Robles Y & Akhavan D (2003) Glycerol facilitator GlpF and the associated aquaporin family of channels. *Curr Opin Struct Biol* **13**: 424-431.

Sumper M, Berg E, Mengele R & Strobel I (1990) Primary structure and glycosylation of the S-layer protein of *Haloflexax volcanii*. *J Bacteriol* **172**: 7111-7118.

Suzuki S & Kurosawa N (2016) Disruption of the gene encoding restriction endonuclease SuAI and development of a host–vector system for the thermoacidophilic archaeon *Sulfolobus acidocaldarius*. *Extremophiles* **20**: 139-148.

Suzuki T, Iwasaki T, Uzawa T, Hara K, Nemoto N, Kon T, Ueki T, Yamagishi A & Oshima T (2002) *Sulfolobus tokodaii* sp. nov. (f. *Sulfolobus* sp. strain 7), a new member of the genus *Sulfolobus* isolated from Beppu Hot Springs, Japan. *Extremophiles* **6**: 39-44.

Suzuki Y, Miyamoto K & Ohta H (2004) A novel thermostable esterase from the thermoacidophilic archaeon *Sulfolobus tokodaii* strain 7. *FEMS Microbiol Lett* **236**: 97-102.

Swarts DC, Makarova K, Wang Y, Nakanishi K, Ketting RF, Koonin EV, Patel DJ & van der Oost J (2014) The evolutionary journey of Argonaute proteins. *Nat Struct & Mol Biol* **21**: 743-753.

Swarts DC, Jore MM, Westra ER, *et al.* (2014) DNA-guided DNA interference by a prokaryotic Argonaute. *Nature* **507**: 258-261.

Tachdjian S & Kelly RM (2006) Dynamic metabolic adjustments and genome plasticity are implicated in the heat shock response of the extremely thermoacidophilic archaeon *Sulfolobus solfataricus*. *J Bacteriol* **188**: 4553-4559.

Takayanagi S, Kawasaki H, Sugimori K, Yamada T, Sugai A, Ito T, Yamasato K & Shioda M (1996) *Sulfolobus hakonensis* sp. nov., a novel species of acidothermophilic archaeon. *Int J Syst Evol Microbiol* **46**: 377-382.

Takemata N, Samson RY & Bell SD (2019) Physical and functional compartmentalization of archaeal chromosomes. *Cell* **179**: 165-179.e118.

Tang TH, Polacek N, Zywicki M, Huber H, Brugger K, Garrett R, Bachellerie JP & Hüttenhofer A (2005) Identification of novel non-coding RNAs as potential antisense regulators in the archaeon *Sulfolobus solfataricus*. *Mol Microbiol* **55**: 469-481.

Tansey MR & Brock TD (1972) The upper temperature limit for Eukaryotic organisms. *Proc Nat Acad Sci USA* **69**: 2426-2428.

Taylor KA, Deatherage JF & Amos LA (1982) Structure of the S-layer of *Sulfolobus acidocaldarius*. *Nature* **299**: 840-842.

Tekaia F, Yeramian E & Dujon B (2002) Amino acid composition of genomes, lifestyles of organisms, and evolutionary trends: a global picture with correspondence analysis. *Gene* **297**: 51-60.

Teufel R, Kung JW, Kockelkorn D, Alber BE & Fuchs G (2009) 3-hydroxypropionyl-coenzyme A dehydratase and acryloyl-coenzyme A reductase, enzymes of the autotrophic 3-hydroxypropionate/4-hydroxybutyrate cycle in the *Sulfolobales*. *J Bacteriol* **191**: 4572-4581.

Trent JD, Osipiuk J & Pinkau T (1990) Acquired thermotolerance and heat shock in the extremely thermophilic archaebacterium *Sulfolobus* sp. strain B12. *J Bacteriol* **172**: 1478-1484.

Trent JD, Nimmessern E, Wall JS, Hartl FU & Horwich AL (1991) A molecular chaperone from a thermophilic archaebacterium is related to the eukaryotic protein t-complex polypeptide-1. *Nature* **354**: 490-493.

Tripepi M, Imam S & Pohlschröder M (2010) *Haloferax volcanii* flagella are required for motility but are not involved in PibD-dependent surface adhesion. *J Bacteriol* **192**: 3093-3102.

Tsai C-L, Tripp P, Sivabalasarma S, *et al.* (2020) The structure of the periplasmic FlaG-FlaF complex and its essential role for archaellar swimming motility. *Nature Microbiol* **5**: 216-225.

Tsuboi K, Sakai HD, Nur N, Stedman KM, Kurosawa N & Suwanto A (2018) *Sulfurisphaera javensis* sp. nov., a hyperthermophilic and acidophilic archaeon isolated from Indonesian hot spring, and reclassification of *Sulfolobus tokodaii* Suzuki et al. 2002 as *Sulfurisphaera tokodaii* comb. nov. *Int J Syst Evol Microbiol* **68**: 1907-1913.

Turner P, Mamo G & Karlsson EN (2007) Potential and utilization of thermophiles and thermostable enzymes in biorefining. *Microbial Cell Factories* **6**.

Uda I, Sugai A, Itoh YH & Itoh T (2001) Variation in molecular species of polar lipids from *Thermoplasma acidophilum* depends on growth temperature. *Lipids* **36**: 103-105.

Uhrigshardt H, Walden M, John H, Petersen A & Anemuller S (2002) Evidence for an operative glyoxylate cycle in the thermoacidophilic crenarchaeon *Sulfolobus acidocaldarius*. *FEBS Lett* **513**: 223-229.

Ulas T, Riemer SA, Zaparty M, Siebers B & Schomburg D (2012) Genome-scale reconstruction and analysis of the metabolic network in the hyperthermophilic archaeon *Sulfolobus solfataricus*. *PloS one* **7**: e43401.

Uldahl KB, Jensen SB, Bhoobalan-Chitty Y, Martínez-Álvarez L, Papathanasiou P & Peng X (2016) Life cycle characterization of *Sulfolobus monocaudavirus* 1, an extremophilic spindle-shaped virus with extracellular tail development. *J Virology* **90**: 5693-5699.

Urbieta MS, Rascovan N, Castro C, Revale S, Giaveno MA, Vazquez M & Donati ER (2014) Draft genome sequence of the novel thermoacidophilic archaeon *Acidianus copahuensis* strain ALE1, isolated from the Copahue volcanic area in Neuquén, Argentina. *Genome Announc* **2**: e00259-00214.

Valentine DL (2007) Adaptations to energy stress dictate the ecology and evolution of the Archaea. *Nat Rev Microbiol* **5**: 316-323.

van de Vossenberg JL, Driessen AJM, Zillig W & Konings WN (1998) Bioenergetics and cytoplasmic membrane stability of the extremely acidophilic, thermophilic archaeon *Picrophilus oshimae*. *Extremophiles* **2**: 67-74.

van der Kolk N, Wagner A, Wagner M, Waßmer B, Siebers B & Albers S-V (2020) Identification of XylR, the activator of arabinose/xylose inducible regulon in *Sulfolobus acidocaldarius* and its application for homologous protein expression. *Front Microbiol* **11**: 1066.

van der Oost J, Jackson RN & Wiedenheft B (2014) Unravelling the structural and mechanistic basis of CRISPR-Cas systems. *Nat Rev Microbiol* **12**: 479-492.

van Wolferen M, Orell A & Albers S-V (2018) Archaeal biofilm formation. *Nat Rev Microbiol* **16**: 699-713.

van Wolferen M, Shajahan A, Heinrich K, Brenzinger S, Black IM, Wagner A, Briegel A, Azadi P & Albers S-V (2020) Species-specific recognition of *Sulfolobales* mediated by UV-inducible pili and s-layer glycosylation patterns. *mBio* **11**: e03014-03019.

Vassart A, Van Wolferen M, Orell A, Hong Y, Peeters E, Albers S-V & Charlier D (2013) Sa-Lrp from *Sulfolobus acidocaldarius* is a versatile, glutamine-responsive, and architectural transcriptional regulator. *Microbiologyopen* **2**: 75-93.

Veith A, Klingl A, Zolghadr B, Lauber K, Mentele R, Lottspeich F, Rachel R, Albers S-V & Kletzin A (2009) *Acidianus*, *Sulfolobus* and *Metallosphaera* surface layers: structure, composition and gene expression. *Mol Microbiol* **73**: 58-72.

Vestergaard G, Shah SA, Bize A, Reitberger W, Reuter M, Phan H, Briegel A, Rachel R, Garrett RA & Prangishvili D (2008) *Stygiolobus* rod-shaped virus and the interplay of crenarchaeal rudiviruses with the CRISPR antiviral system. *J Bacteriol* **190**: 6837-6845.

Vetter AM, Helmecke J, Schomburg D & Neumann-Schaal M (2019) The Impact of pyroglutamate: *Sulfolobus acidocaldarius* has a growth advantage over *Saccharolobus solfataricus* in glutamate-containing media. *Archaea* **2019**: 3208051.

Villafane A, Voskoboinik Y, Ruhl I, Sannino D, Maezato Y, Blum P & Bini E (2011) CopR of *Sulfolobus solfataricus* represents a novel class of archaeal-specific copper-responsive activators of transcription. *Microbiologyopen* **157**: 2808-2817.

Vorontsov EA, Rensen E, Prangishvili D, Krupovic M & Chamot-Rooke J (2016) Abundant lysine methylation and N-terminal acetylation in *Sulfolobus islandicus* revealed by bottom-up and top-down proteomics. *Mol Cell Proteomics* **15**: 3388-3404.

Völlmecke C, Drees SL, Reimann J, Albers S-V & Lübben M (2012) The ATPases CopA and CopB both contribute to copper resistance of the thermoacidophilic archaeon *Sulfolobus solfataricus*. *Microbiologyopen* **158**: 1622-1633.

Wagner M, Berkner S, Ajon M, Driessen AJM, Lipps G & Albers S-V (2009) Expanding and understanding the genetic toolbox of the hyperthermophilic genus *Sulfolobus*. *Biochem Soc Trans* **37**: 97-101.

Wagner M, van Wolferen M, Wagner A, Lassak K, Meyer B, Reimann J & Albers S-V (2012) Versatile genetic tool box for the Crenarchaeote *Sulfolobus acidocaldarius*. *Front Microbiol* **3**.

Wagner M, Wagner A, Ma X, Kort JC, Ghosh A, Rauch B, Siebers B & Albers S-V (2014) Investigation of the *malE* promoter and MalR, a positive regulator of the maltose regulon, for an improved expression system in *Sulfolobus acidocaldarius*. *Appl Environ Microbiol* **80**: 1072-1081.

Wagner M, Shen L, Albersmeier A, van der Kolk N, Kim S, Cha J, Brasen C, Kalinowski J, Siebers B & Albers SV (2017) *Sulfolobus acidocaldarius* uptakes pentoses via a cut2-type ABC transporter and metabolizes them through the aldolase-independent Weimberg pathway. *Appl Environ Microbiol* **84**: 1273.

Wang B, Yang S, Zhang L & He Z-G (2010) Archaeal eukaryote-like serine/threonine protein kinase interacts with and phosphorylates a forkhead-associated-domain-containing protein. *J Bacteriol* **192**: 1956-1964.

Wang F, Liu Y, Su Z, Osinski T, de Oliveira GAP, Conway JF, Schouten S, Krupovic M, Prangishvili D & Egelman EH (2019) A packing for A-form DNA in an icosahedral virus. *Proc Natl Acad Sci* **116**: 22591-22597.

Wang F, Cvirkaitė-Krupovic V, Kreutzberger MA, Su Z, de Oliveira GA, Osinski T, Sherman N, DiMaio F, Wall JS & Prangishvili D (2019) An extensively glycosylated archaeal pilus survives extreme conditions. *Nat Microbiol* **4**: 1401-1410.

Wang H-C, Susko E & Roger AJ (2006) On the correlation between genomic G+C content and optimal growth temperature in prokaryotes: Data quality and confounding factors. *Biochem Biophys Res Comm* **342**: 681-684.

Wang K, Sybers D, Maklad HR, Lemmens L, Lewyllie C, Zhou X, Schult F, Bräsen C, Siebers B & Valegård K (2019) A TetR-family transcription factor regulates fatty acid metabolism in the archaeal model organism *Sulfolobus acidocaldarius*. *Nat Commun* **10**: 1-16.

Wang Y, Xu X, Wen Z, Li W, Yang B & Whiteley C (2010) Isolation, purification, and properties of a novel small heat shock protein from the hyperthermophile *Sulfolobus solfataricus*. *Appl Biochem Biotechnol* **162**: 476-485.

Ward L, Taylor MW, Power JF, Scott BJ, McDonald IR & Stott MB (2017) Microbial community dynamics in Inferno Crater Lake, a thermally fluctuating geothermal spring. *ISME J* **11**: 1158-1167.

Watling HR (2006) The bioleaching of sulphide minerals with emphasis on copper sulphides - A review. *Hydrometallurgy* **84**: 81-108.

Weinberger AD, Wolf YI, Lobkovsky AE, Gilmore MS & Koonin EV (2012) Viral diversity threshold for adaptive immunity in prokaryotes. *mBio* **3**: e00456-00412.

Weiße RH-J, Faust A, Schmidt M, Schönheit P & Scheidig AJ (2016) Structure of NDP-forming Acetyl-CoA synthetase ACD1 reveals a large rearrangement for phosphoryl transfer. *Proc Natl Acad Sci USA* **113**: E519-E528.

Wiedenheft B, Mosolff J, Willits D, Yeager M, Dryden KA, Young M & Douglas T (2005) An archaeal antioxidant: Characterization of a Dps-like protein from *Sulfolobus solfataricus*. *Proc Natl Acad Sci USA* **102**: 10551-10556.

Williams TJ, Allen M, Tschitschko B & Cavigchioli R (2017) Glycerol metabolism of *Haloarchaea*. *Environ Microbiol* **19**: 864-877.

Willkomm S, Oellig CA, Zander A, Restle T, Keegan R, Grohmann D & Schneider S (2017) Structural and mechanistic insights into an archaeal DNA-guided Argonaute protein. *Nat Microbiol* **2**: 17035-17035.

Wolf J, Stark H, Fafenrot K, et al. (2016) A systems biology approach reveals major metabolic changes in the thermoacidophilic archaeon *Sulfolobus solfataricus* in response to the carbon source L-fucose versus D-glucose. *Mol Microbiol* **102**: 882-908.

Woodruff PJ, Carlson BL, Siridechadilok B, Pratt MR, Senaratne RH, Mougous JD, Riley LW, Williams SJ & Bertozzi CR (2004) Trehalose 1s required for growth of *Mycobacterium smegmatis*. *J Biol Chem* **279**: 28835-28843.

Worthington P, Hoang V, Perez-Pomares F & Blum P (2003) Targeted disruption of the α -amylase gene in the hyperthermophilic archaeon *Sulfolobus solfataricus*. *J Bacteriol* **185**: 482-488.

Wurtzel O, Sapra R, Chen F, Zhu Y, Simmons BA & Sorek R (2010) A single-base resolution map of an archaeal transcriptome. *Genome Res* **20**: 133-141.

Xu N, Li Y, Zhao YT, Guo L, Fang YY, Zhao JH, Wang XJ, Huang L & Guo HS (2012) Identification and characterization of small RNAs in the hyperthermophilic archaeon *Sulfolobus solfataricus*. *PLoS ONE* **7**: e35306.

Xu Y, Gristwood T, Hodgson B, Trinidad JC, Albers SV & Bell SD (2016) Archaeal orthologs of Cdc45 and GINS form a stable complex that stimulates the helicase activity of MCM. *Proc Natl Acad Sci USA* **113**: 13390-13395.

Yan Z, Maruyama A, Arakawa T, Fushinobu S & Wakagi T (2016) Crystal structures of archaeal 2-oxoacid:ferredoxin oxidoreductases from *Sulfolobus tokodaii*. *Sci Rep* **6**: 33061.

Yang N & Driessen AJM (2014) Deletion of *cdvB* paralogous genes of *Sulfolobus acidocaldarius* impairs cell division. *Extremophiles* **18**: 331-339.

Yasutake Y, Nishiya Y, Tamura N & Tamura T (2007) Structural insights into unique substrate selectivity of *Thermoplasma acidophilum* D-aldohexose dehydrogenase. *J Mol Biol* **367**: 1034-1046.

Yokooji Y, Sato T, Fujiwara S, Imanaka T & Atomi H (2013) Genetic examination of initial amino acid oxidation and glutamate catabolism in the hyperthermophilic archaeon *Thermococcus kodakarensis*. *J Bacteriol* **195**: 1940-1948.

Zago MA, Dennis PP & Omer AD (2005) The expanding world of small RNAs in the hyperthermophilic archaeon *Sulfolobus solfataricus*. *Mol Microbiol* **55**: 1812-1828.

Zaparty M & Siebers B (2011) Physiology, metabolism, and enzymology of thermoacidophiles. *Extremophiles Handbook*, (Horikoshi K, ed.) p.^pp. 601-639. Springer Japan, Tokyo.

Zaparty M, Esser D, Gertig S, et al. (2009) "Hot standards" for the thermoacidophilic archaeon *Sulfolobus solfataricus*. *Extremophiles* **14**: 119-142.

Zaramela LS, Vêncio RZN, Ten-Caten F, Baliga NS & Koide T (2014) Transcription start site associated RNAs (TSSaRNAs) are ubiquitous in all domains of life. *PLoS ONE* **9**: e107680.

Zebec Z, Zink IA, Kerou M & Schleper C (2016) Efficient CRISPR-Mediated post-transcriptional gene silencing in a hyperthermophilic archaeon using multiplexed crRNA expression. *G3* **6**: 3161-3168.

Zebec Z, Manica A, Zhang J, White MF & Schleper C (2014) CRISPR-mediated targeted mRNA degradation in the archaeon *Sulfolobus solfataricus*. *Nucleic Acids Res* **42**: 5280-5288.

Zeldes BM, Straub CT, Otten JK, Adams MW & Kelly RM (2018) A synthetic enzymatic pathway for extremely thermophilic acetone production based on the unexpectedly thermostable acetoacetate decarboxylase from *Clostridium acetobutylicum*. *Biotechnol Bioeng* **115**: 2951-2961.

Zeldes BM, Keller MW, Loder AJ, Straub CT, Adams MW & Kelly RM (2015) Extremely thermophilic microorganisms as metabolic engineering platforms for production of fuels and industrial chemicals. *Front Microbiol* **6**: 1209.

Zeldes BM, Loder AJ, Counts JA, Haque M, Widney KA, Keller LM, Albers S-V & Kelly RM (2019) Determinants of sulphur chemolithoautotrophy in the extremely thermoacidophilic *Sulfolobales*. *Environ Microbiol* **21**: 3696-3710.

Zeldovich KB, Berezovsky IN & Shakhnovich EI (2007) Protein and DNA sequence determinants of thermophilic adaptation. *PLoS Comput Biol* **3**: e5-e5.

Zeng Z, Liu X-L, Wei JH, Summons RE & Welander PV (2018) Calditol-linked membrane lipids are required for acid tolerance in *Sulfolobus acidocaldarius*. *Proc Natl Acad Sci* **115**: 12932-12937.

Zhang C & Whitaker RJ (2012) A broadly applicable gene knockout system for the thermoacidophilic archaeon *Sulfolobus islandicus* based on simvastatin selection. *Microbiologyopen* **158**: 1513-1522.

Zhang C, Cooper TE, Krause DJ & Whitaker RJ (2013) Augmenting the genetic toolbox for *Sulfolobus islandicus* with a stringent positive selectable marker for agmatine prototrophy. *Appl Environ Microbiol* **79**: 5539-5549.

Zhang C, She Q, Bi H & Whitaker RJ (2016) The *apt*/6-methylpurine counterselection system and its applications in genetic studies of the hyperthermophilic archaeon *Sulfolobus islandicus*. *Appl Environ Microbiol* **82**: 3070-3081.

Zhang C, Phillips APR, Wipfler RL, Olsen GJ & Whitaker RJ (2018) The essential genome of the crenarchaeal model *Sulfolobus islandicus*. *Nat Commun* **9**: 1-11.

Zhang C, Wipfler RL, Li Y, Wang Z, Hallett EN & Whitaker RJ (2019) Cell structure changes in the hyperthermophilic crenarchaeon *Sulfolobus islandicus* lacking the S-layer. *mBio* **10**.

Zhang C, Tian B, Li S, Ao X, Dalgaard K, Gökce S, Liang Y & She Q (2013) Genetic manipulation in *Sulfolobus islandicus* and functional analysis of DNA repair genes. *Biochem Soc Trans* **41**: 405-410.

Zhang J & White MF (2013) Hot and crispy: CRISPR-Cas systems in the hyperthermophile *Sulfolobus solfataricus*. *Biochem Soc Trans* **41**: 1422-1426.

Zhang J, Graham S, Tello A, Liu H & White MF (2016) Multiple nucleic acid cleavage modes in divergent type III CRISPR systems. *Nucleic Acids Res* **44**: 1789-1799.

Zhang Q, Iwasaki T, Wakagi T & Oshima T (1996) 2-oxoacid:ferredoxin oxidoreductase from the thermoacidophilic archaeon, *Sulfolobus* sp. strain 7. *J Biochem* **120**: 587-599.

Zhang R, Neu TR, Blanchard V, Vera M & Sand W (2019) Biofilm dynamics and EPS production of a thermoacidophilic bioleaching archaeon. *New Biotechnology* **51**: 21-30.

Zhang R, Neu TR, Zhang Y, Bellenberg S, Kuhlicke U, Li Q, Sand W & Vera M (2015) Visualization and analysis of EPS glycoconjugates of the thermoacidophilic archaeon *Sulfolobus metallicus*. *Appl Microbiol Biotechnol* **99**: 7343-7356.

Zhang RY, Neu TR, Bellenberg S, Kuhlicke U, Sand W & Vera M (2015) Use of lectins to *in situ* visualize glycoconjugates of extracellular polymeric substances in acidophilic archaeal biofilms. *Microbial Biotechnol* **8**: 448-461.

Zhang Y, Kouril T, Snoep JL, Siebers B, Barberis M & Westerhoff HV (2017) The peculiar glycolytic pathway in hyperthermophilic archaea: Understanding its whims by experimentation *in silico*. *Int J Mol Sci* **18**: 876.

Zhang Z, Zhao M, Chen Y, Wang L, Liu Q, Dong Y, Gong Y & Huang L (2019) Architectural roles of Cren7 in folding crenarchaeal chromatin filament. *Mol Microbiol* **111**: 556-569.

Zhang Z, Zhan Z, Wang B, Chen Y, Chen X, Wan C, Fu Y & Huang L (2020) Archaeal chromatin proteins Cren7 and Sul7d compact DNA by bending and bridging. *mBio* **11**: 1-15.

Zhao HB, Zhang YS, Zhang X, Qian L, Sun ML, Yang Y, Zhang YS, Wang J, Kim H & Qiu GZ (2019) The dissolution and passivation mechanism of chalcopyrite in bioleaching: An overview. *Miner Eng* **136**: 140-154.

Zheng T, Huang Q, Zhang C, Ni J, She Q & Shen Y (2012) Development of a simvastatin selection marker for a hyperthermophilic acidophile, *Sulfolobus islandicus*. *Appl Environ Microbiol* **78**: 568-574.

Zhu Y, Klompe SE, Vlot M, van der Oost J & Staals RHJ (2018) Shooting the messenger: RNA-targetting CRISPR-Cas systems. *Biosci Rep* **38**: BSR20170788.

Zillig W, Stetter KO & Janekovic D (1979) DNA-dependent RNA polymerase from the archaeabacterium *Sulfolobus acidocaldarius*. *Eur J Biochem* **96**: 597-604.

Zillig W, Stetter KO, Wunderl S, Schulz W, Priess H & Scholz I (1980) The *Sulfolobus*-“Caldariella” group: Taxonomy on the basis of the structure of DNA-dependent RNA polymerases. *Arch Microbiol* **125**: 259-269.

Zillig W, Yeats S, Holz I, Böck A, Rettenberger M, Groppe F & Simon G (1986) *Desulfurolobus ambivalens*, gen. nov., sp. nov., an autotrophic archaebacterium facultatively oxidizing or reducing sulfur. *Syst Appl Microbiol* **8**: 197-203.

Zillig W, Kletzin A, Schleper C, Holz I, Janečková D, Hain J, Lanzendörfer M & Kristjánsson JK (1993) Screening for *Sulfolobales*, their plasmids and their viruses in Icelandic Solfataras. *Syst Appl Microbiol* **16**: 609-628.

Zimmermann P, Laska S & Kletzin A (1999) Two modes of sulfite oxidation in the extremely thermophilic and acidophilic archaeon *Acidianus ambivalens*. *Arch Microbiol* **172**: 76-82.

Zolghadr B, Kling A, Koerdt A, Driessen AJM, Rachel R & Albers SV (2010) Appendage-mediated surface adherence of *Sulfolobus solfataricus*. *J Bacteriol* **192**: 104-110.

Zweerink S, Kallnik V, Ninck S, *et al.* (2017) Activity-based protein profiling as a robust method for enzyme identification and screening in extremophilic Archaea. *Nature Commun* **8**: 15352.

Zähringer U, Moll H, Hettmann T, Knirel YA & Schäfer G (2000) Cytochrome b558/566 from the archaeon *Sulfolobus acidocaldarius* has a unique Asn-linked highly branched hexasaccharide chain containing 6-sulfoquinovose. *Eur J Biochem* **267**: 4144-4149.

Figure Legends

Figure 1.1 Timeline of thermoacidophile isolations and major events.

Timeline contains the organism's name at the time of the associated event. The following are the current classifications: *Sulfolobus brierleyi* (f. *Acidianus brierleyi*), *Saccharolobus solfataricus* (f. *Sulfolobus solfataricus*), *Acidianus ambivalens* (f. *Desulfurolobus ambivalens*), *Saccharolobus shibatae* (f. *Sulfolobus shibatae*) *Sulfuracidifex metallicus* (f. *Sulfolobus metallicus*), *Metallosphaera hakonensis* (f. *Sulfolobus hakonensis*), *Sulfurisphaera tokadaii* (f. *Sulfolobus tokodaii*), *Saccharolobus islandicus* (f. *Sulfolobus islandicus*).

Figure 2.1 16S Phylogeny tree of thermoacidophilic organisms.

Figure 3.1 Thermoacidophilic archaeal viruses and their infection mechanisms.

(A) Schematic representation of virion morphologies of viruses infecting thermoacidophilic archaea, as described in the text. **(B)** Segmented tomographic volume of a SIRV2 virion (red) attached to a surface filament of *Sa. islandicus* (green) with help of the three terminal virion fibers (blue). Inset depicts a magnification of the interaction between the tail fibers and the surface structure. Scale bars, 500 nm. **(C)** Volume segmentations of electron microscopy tomograms showing SSV1 maturation and release by budding. Scale bar, 50 nm. **(D)** Transmission electron micrograph of a thin section of a SIRV2 infected *Sa. islandicus* cell displaying several pyramidal egress structures. Scale bar, 100 nm. **(E)** Transmission electron micrographs of an isolated pyramidal egress structure in open conformation isolated after SIRV2

infection of *Sa.islandicus*. Scale bar, 100 nm Adapted from (Bize *et al.*, 2009, Quax *et al.*, 2011, Quemin *et al.*, 2013, Quemin *et al.*, 2016).

Figure 4.1 Main principles in genome organization and genetic information processing in the *Sulfolobales*. Conceptual schemes representing the major elements and principles of macro-level organization of the genomic DNA (adapted from (Takemata *et al.*, 2019) **(A)**, micro-level organization of the genomic DNA (partially adapted from (Peeters *et al.*, 2015) **(B)**, initiation of replication **(C)** and initiation of transcription **(D)**.

Figure 5.1 Archaeal cell surface structures involved in planktonic and biofilm growth. **Upper image:** Electronic microscopy image from a *S. acidocaldarius* cell where archaellum and pilus can be seen. **Lower image:** Schematic model with all proposed cell surface appendages in the *Sulfolobales*: the Aap pili (archaeal adhesive pili), the Ups pili (UV induced pili), the archaellum and the threads. Also depicted is the S-layer and its proteins: SlaA and SlaB.

Figure 5.2 Confocal Laser Microscopy images from static biofilm from *S. acidocaldarius* in days 3 to 7 of growth. Cells (DNA) stained with 4',6-diamidino-2-phenylindole (DAPI; blue); extracellular glucose and mannose residues stained with fluorescently labeled concanavalin A (conA; green); and N- acetyl-d- glucosamine residues stained with fluorescently labeled lectin IB4 (yellow). Scale bars: 20 μ m.

Figure 5.3 Model of the *Sulfolobus* S-layer. (A) The *Sulfolobus* S-layer consists of the 2 protein subunits: SlaA dimers (red, orange, yellow) form the outer S-layer canopy. Each SlaA

protein is predicted to be rich in β -strands. The SlaA dimer has a boomerang-like shape, the angle of which determines the S-layer unit cell size. SlaB trimers (gray) form the membrane anchors of the S-layer. Each SlaB is predicted to consist of an N-terminal transmembrane domain (TMD), a coiled-coil domain (CC), and 2–3 C-terminal β -sandwich domains (β). SlaA and SlaB proteins are highly glycosylated (green). (B) Electron microscopy image from negatively stained isolated S-layer from *S. acidocaldarius*. (C) SDS/PAGE showing surface layer proteins. M, marker; 1, washed once; 2, washed twice; 3, washed 3 times in detergent. (D) Subtomogram average of fully assembled S-layer. (E) Subtomogram average of SlaB-depleted S-layer. (F) Difference map (pink) overlaid with the complete S-layer visualizes location of SlaB. (Scale bars, and C–E, 20 nm) (Figure adapted from Gambeli et al. 2019)

Figure 5.4. Glycosylation. (A) Comparison of the N-glycan trees of three different *Sulfolobus/Saccharolobus* species. (B) The current understanding of the N-glycosylation pathway in *S. acidocaldarius*. The N-glycan biosynthesis is initiated by adding nucleotide-activated monosaccharides sequentially to the lipid carrier DolP on the cytoplasmic side of the membrane. The fully assembled DolPP-linked N-glycan (hexasaccharide) is translocated across the membrane and then transferred by AlgB on the specific N-glycosylation sequons in secreted proteins. Suagr code is shown in (A).

Figure 5.5: Schematic model of the cell division process in *Sulfolobus acidocaldarius*. (I) CdvA (red) is the first protein of the *S. acidocaldarius* cell division machinery that arrives at the future site of cell division, before DNA (light blue) segregation starts. (II) During nucleoid

condensation (blue) CdVB (light green) forms a ring-like structure at midcell that is anchored to the membrane by CdVA. **(III)** CdVB provides a scaffold for CdVB1 and B2 (green) that are positioned at the cell center in ring-like structures. Additionally, CdVC (yellow), a homolog of the hexameric ATPase Vps4, localizes at the septum while nucleoid segregation and initial membrane invagination starts. **(IV)** After nucleoid segregation, the CdVB-ring undergoes proteasomal (purple) degradation. **(V)** Upon CdVB removal CdVB1 and B2 constrict, leading to the final division of the cell. **(VI)** Directly after fission, the new born cells have an oval shape that rapidly changes to the typical coccoid shape of *S. acidocaldarius* cells. CdVA and CdVC are organized in a ring-like structure as well. However, for a better overview of the model, only the organization of both proteins at mid-cell was indicated.

Figure 6.1 Major mechanisms of thermoacidophily. 1) Thermoacidophiles have an inverted membrane potential with a positive charge on the inside of the cellular membrane and a negative charge on the outside to prevent the acidification of the cytoplasm by the passive diffusion of protons. 2) The inverted membrane potential is maintained by transporting cations such as K^+ into the cytoplasm. 3) Cyclopentyl ring moieties on tetraether lipids increase packing of the tetraether lipids decreasing the permeability of the membrane by protons and increasing cellular heat stability. 4) Tetraether lipids make a monolayer that is less permeable to protons and more heat stable than diether lipids. 5) Proton pumps export protons from the cytoplasm to prevent the acidification of the cytoplasm. 6) Heat damaged or protonated proteins can either be degraded via the proteasome or properly refolded by the thermosome.

Figure 6.2 Model of heavy metal resistance in *Sulfolobales*. 1) CopA and CopB export Cu outside the cell with ATP consumption. CopM is a metal chaperone that forms part of the Cop system that also includes a transcriptional factor called CopT (not shown). 2) PolyP can sequester cations via its negatively charged surface. 3) PolyP can also be degraded by PPX into inorganic phosphate to be exported outside the cell along with cations via PitA or Pho84 transporters. 4) Some proteins also act to sequester metal ions, for example Dps. The mechanism for the influx of metals is still unknown.

Figure 7.1: Predicted protein kinase and protein phosphatase homologs in the four different *Sulfolobales* species *S. acidocaldarius*, *Sa. islandicus*, *Sa. solfataricus* and *Sulfuri. tokodaii*. Depicted are the different canonical and non-canonical Hanks type protein kinases and protein phosphatases with their correspondent domain structure. (Lower *et al.*, 2000, Lower & Kennelly, 2002, Lower & Kennelly, 2003, Lower *et al.*, 2004, Haile & Kennelly, 2011, Esser & Siebers, 2013, Ray *et al.*, 2015, Esser *et al.*, 2016, Hoffmann *et al.*, 2017, Huang *et al.*, 2017).

Figure 7.2: Representation of known protein phosphorylation target enzymes in different *Sulfolobales* strains. (*Sa. solfataricus* in pink, *Sa. islandicus* in orange, *S. acidocaldarius* in blue and *Sulfuri. tokodaii* in grey) with their physiological function. From upper left to right: The FadR transcriptional regulator represses transcription of the fatty acid gene cluster and dissociates from the DNA upon binding to acyl-CoA. Phosphorylation of FadR by the ePK ArnC (Saci_1196) prevents acyl-CoA binding and thus hinders transcription of the gene cluster (Maklad *et al.*, 2020). The archaellum regulatory network consists of the gene

cluster *arlBXGFHIJ* (*flaBXGFHIJ*), which encodes the motility structure, the archaellum, and is under the control of two promoters, one upstream of *arlB* (*flaB*) being induced under starvation and one weak promoter upstream of *arlX* (*flaX*). The two negative regulators ArnA (Saci_1210) and ArnB (Saci_1211) were shown to be phosphorylated by the ePKs ArnC and ArnD (Saci_1694) and dephosphorylated by the PP PP2A (Saci_0884). Deletion of the PP2A led to a hypermotile phenotype suggesting a negative influence on the gene cluster (Reimann *et al.*, 2012, Hoffmann *et al.*, 2017). The DNA binding protein AbfR1 (Saci_0446) is a positive regulator of the *arlB* (*flaB*) promoter (Orell *et al.*, 2013). Phosphorylation of AbfR1 inhibits DNA binding and thus regulates biofilm formation and motility (Li *et al.*, 2017). The FHA domain containing protein ST0829 was shown to interact and be phosphorylated by the ePK ST1565 indicating a role in transcription regulation (Duan & He, 2011). The Holliday Junction Resolvase (Hjc) (SiRe_1431) is phosphorylated by the aPK SiRe_0171 facilitating DNA repair (Huang *et al.*, 2019). The phosphohexomutase (SSO0207) exhibited a decreased V_{max} value after being phosphorylated (Ray *et al.*, 2005). The Rio kinases (Saci_0796 and Saci_0965) were shown to play a role in the ribosome maturation of the small subunit (SSU) (Knüppel *et al.*, 2018).

Figure 8.1 Overview of the central metabolism in *Sulfolobales*. Dashed arrows indicate pathways, which have not yet been experimentally demonstrated. Abbreviations: F6P, fructose 6-phosphate; DHAP, dihydroxyacetone phosphate; GAP, glyceraldehyde 3-phosphate; D-KDG, 2-keto-3-deoxy-D-gluconate; D-KDGal, 2-keto-3-deoxy-D-galactonate; D-KDA, 2-keto-3-deoxy-D-arabinoate; L-KDA, 2-keto-3-deoxy-L-arabinoate; D-KDX, 2-keto-3-deoxy-D-

xylonate; AA, amino acid; ED, Entner-Doudoroff pathway; EMP, Embden-Meyerhof-Parnas pathway; RuMP, reversed ribulose monophosphate pathway; TCA, Tricarboxylic acid cycle; 3HP/4HB, 3-hydroxypropionate/4-hydroxybutyrate cycle; ABC, ATP binding cassette transporters; RC, respiratory chain.

Figure 8.2 The enzymatic pathway of the 3-hydroxypropionate/4-hydroxybutyrate (3-HP/4-HB) cycle as characterized in *Metallosphaera sedula*. Enzyme names are contained within the yellow oval with arrows indicating reactions for which they have known catalytic activity. Enzymes in pink have shown activity for only a single reaction in the 3-HP/4-HB cycle; enzymes in green exhibit activity on multiple steps in the cycle.

Figure 8.3 Current knowledge of the mechanism of sulfur oxidation and reduction in the *Sulfolobales*. A) Sulfur reduction and B) sulfur oxidation. Solid arrows indicate involvement in a reaction; dotted arrows represent transport of species; dashed lines indicate that the function is suspected but has not been demonstrated experimentally in the *Sulfolobales*. Enzyme colors indicate general grouping of function: coupled to electron transport chain (blue), involved in transporting or trafficking sulfur species (yellow), transformation of sulfur species with no energy conservation (orange), transformation of sulfur species directly coupled to energy-conserving biomolecules (green). Abbreviations: sulfur reductase (Sre), hydrogenase (Hyn), heterodisulfide reductase (Hdr), tetrathionate hydrolase (TetH), sulfide:quinone oxidoreductase (SQO), sulfite:acceptor oxidoreductase (SAOR), thiosulfate:quinone oxidoreductase (TQO), sulfur oxygenase reductase (SOR), adenosine-5'-phosphosulfate

reductase (APSR), adenosine-5'-phosphosulfate (APS), adenylylsulfate:phosphate adenylyltransferase (APAT), ATP sulfurylase (ATPS), adenylate kinase (AK).

Figure 9.1 Mechanisms for the generation of markerless deletion mutants. A)

Plasmid integration occurs via single-crossover, resulting in a merodiploidal form. After counterselection with 5-FOA, the *pyrEF* marker cassette is looped out, either with or without the GOI, resulting in a theoretical ration of one to one in mutated and wild type cells. B) and C) Double-crossover is feasible by introducing a linearized vector. Depending on the experimental design, either parts of the GOI (B) or an upstream (US) region (C) are introduced for recombination. Counterselection with 5-FOA produces marker-free deletion mutants. D) A plasmid containing a CRISPR array as well as a repair fragment with homologous sequences to the GOI are introduced into a recipient strain. Upon induction, crRNA is transcribed and forms a ribonucleoprotein complex with the endogenous Cas protein, scanning the genomic DNA for the spacer sequence and cutting it. Only colonies which conducted recombination with the repair fragment survive. GOI, gene of interest; US, upstream; DS, downstream, *pyrEF*, *pyrEF* marker cassette; crRNA, CRISPR RNA; crRNP, ribonucleoprotein complex consisting of crRNA and Cas protein.

Figure 9.2 Mechanisms of sulfidic ore dissolution. A) thiosulfate mechanism B) and polysulfide mechanism. Green dashed arrows indicate biological steps; solid arrows indicate spontaneous abiotic reactions; blue dashed-dotted arrows indicate an overall transformation involving multiple reaction steps; yellow dashed-dotted arrows represent phase transition.

Bold boxes around a species indicates that this is the dominant sulfur product of the dissolution process.

References:

Bize A, Karlsson EA, Ekefjärd K, Quax TEF, Pina M, Prevost M-C, Forterre P, Tenaillon O, Bernander R & Prangishvili D (2009) A unique virus release mechanism in the Archaea. *Proc Natl Acad Sci USA* **106**: 11306-11311.

Duan X & He Z-G (2011) Characterization of the specific interaction between archaeal FHA domain-containing protein and the promoter of a flagellar-like gene-cluster and its regulation by phosphorylation. *Biochem Biophys Res Commun* **407**: 242-247.

Esser D & Siebers B (2013) Atypical protein kinases of the RIO family in archaea. *Biochem Soc Trans* **41**: 399-404.

Esser D, Hoffmann L, Pham TK, Bräsen C, Qiu W, Wright PC, Albers S-V & Siebers B (2016) Protein phosphorylation and its role in archaeal signal transduction. *FEMS Microbiol Rev* **40**: 625-647.

Haile JD & Kennelly PJ (2011) The activity of an ancient atypical protein kinase is stimulated by ADP-ribose *in vitro*. *Arc Biochem Biophys* **511**: 56-63.

Hoffmann L, Schummer A, Reimann J, Haurat MF, Wilson AJ, Beeby M, Warscheid B & Albers SV (2017) Expanding the archaeal regulatory network—the eukaryotic protein kinases ArnC and ArnD influence motility of *Sulfolobus acidocaldarius*. *MicrobiologyOpen* **6**.

Huang Q, Zhong Q, Mayaka J, Ni J & Shen Y (2017) Autophosphorylation and cross-phosphorylation of protein kinases from the crenarchaeon *Sulfolobus islandicus*. *Front Microbiol* **8**: 2173.

Huang Q, Mayaka JBA, Zhong Q, Zhang C, Hou G, Ni J & Shen Y (2019) Phosphorylation of the archaeal Holliday junction resolvase Hjc inhibits its catalytic activity and facilitates DNA repair in *Sulfolobus islandicus* REY15A. *Front Microbiol* **10**: 1214.

Knüppel R, Christensen RH, Gray FC, Esser D, Strauß D, Medenbach J, Siebers B, MacNeill SA, LaRonde N & Ferreira-Cerca S (2018) Insights into the evolutionary conserved regulation of Rio ATPase activity. *Nucleic Acids Res* **46**: 1441-1456.

Li L, Banerjee A, Bischof LF, Maklad HR, Hoffmann L, Henche AL, Veliz F, Bildl W, Schulte U & Orell A (2017) Wing phosphorylation is a major functional determinant of the Lrs14-type biofilm and motility regulator AbfR1 in *Sulfolobus acidocaldarius*. *Mol Microbiol* **105**: 777-793.

Lower BH & Kennelly PJ (2002) The membrane-associated protein-serine/threonine kinase from *Sulfolobus solfataricus* is a glycoprotein. *J Bacteriol* **184**: 2614-2619.

Lower BH & Kennelly PJ (2003) Open reading frame sso2387 from the archaeon *Sulfolobus solfataricus* encodes a polypeptide with protein-serine kinase activity. *J Bacteriol* **185**: 3436-3445.

Lower BH, Bischoff KM & Kennelly PJ (2000) The archaeon *Sulfolobus solfataricus* contains a membrane-associated protein kinase activity that preferentially phosphorylates threonine residues *in vitro*. *J Bacteriol* **182**: 3452-3459.

Lower BH, Potters MB & Kennelly PJ (2004) A phosphoprotein from the archaeon *Sulfolobus solfataricus* with protein-serine/threonine kinase activity. *J Bacteriol* **186**: 463-472.

Maklad HR, Gutierrez GJ, Esser D, Siebers B & Peeters E (2020) Phosphorylation of the acyl-CoA binding pocket of the FadR transcription regulator in *Sulfolobus acidocaldarius*. *Biochimie*.

Orell A, Peeters E, Vassen V, Jachlewski S, Schalles S, Siebers B & Albers S-V (2013) Lrs14 transcriptional regulators influence biofilm formation and cell motility of *Crenarchaea*. *ISME J* **7**: 1886-1898.

Peeters E, Driessen RP, Werner F & Dame RT (2015) The interplay between nucleoid organization and transcription in archaeal genomes. *Nat Rev Microbiol* **13**: 333-341.

Quax TE, Lucas S, Reimann J, Pehau-Arnaudet G, Prevost M-C, Forterre P, Albers S-V & Prangishvili D (2011) Simple and elegant design of a virion egress structure in Archaea. *Proc Natl Acad Sci* **108**: 3354-3359.

Quemin ER, Lucas S, Daum B, Quax TE, Kühlbrandt W, Forterre P, Albers S-V, Prangishvili D & Krupovic M (2013) First insights into the entry process of hyperthermophilic archaeal viruses. *J Virol* **87**: 13379-13385.

Quemin ERJ, Chlanda P, Sachse M, Forterre P, Prangishvili D & Krupovic M (2016) Eukaryotic-like virus budding in archaea. *mBio* **7**.

Ray WK, Potters MB, Haile JD & Kennelly PJ (2015) Activation of SsoPK4, an archaeal eIF2 α kinase homolog, by oxidized CoA. *Proteomes* **3**: 89-116.

Ray WK, Keith SM, DeSantis AM, Hunt JP, Larson TJ, Helm RF & Kennelly PJ (2005) A phosphohexomutase from the archaeon *Sulfolobus solfataricus* is covalently modified by phosphorylation on serine. *J Bacteriol* **187**: 4270-4275.

Reimann J, Lassak K, Khadouma S, Ettema TJ, Yang N, Driessen AJ, Klingl A & Albers SV (2012) Regulation of archaella expression by the FHA and von Willebrand domain-containing proteins ArnA and ArnB in *Sulfolobus acidocaldarius*. *Mol Microbiol* **86**: 24-36.

Takemata N, Samson RY & Bell SD (2019) Physical and functional compartmentalization of archaeal chromosomes. *Cell* **179**: 165-179.e118.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



286x190mm (150 x 150 DPI)




Figure 1.1 Timeline of thermoacidophile isolations and major events. Timeline contains the organism's names at the time of the associated event. The following are the current classifications: *Sulfolobus brierleyi* (f. *Acidianus brierleyi*), *Saccharolobus solfataricus* (f. *Sulfolobus solfataricus*), *Acidianus ambivalens* (f. *Desulfurolobus ambivalens*), *Saccharolobus shibatae* (f. *Sulfolobus shibatae*) *Sulfuracidifex metallicus* (f. *Sulfolobus metallicus*), *Metallosphaera hakonensis* (f. *Sulfolobus hakonensis*), *Sulfurisphaera tokodaii* (f. *Sulfolobus tokodaii*), *Saccharolobus islandicus* (f. *Sulfolobus islandicus*).

254x190mm (300 x 300 DPI)




Figure 2.1 16s Phylogeny tree of thermoacidophilic organisms.

99x190mm (300 x 300 DPI)




Figure 3.1 Thermoacidophilic archaeal viruses and their infection mechanisms. (A) Schematic representation of virion morphologies of viruses infecting thermoacidophilic archaea as described in the text. (B) Segmented tomographic volume of a SIRV2 virion (red) attached to a surface filament of *Sa. islandicus* (green) with help of the three terminal virion fibers (blue). Inset depicts a magnification of the interaction between the tail fibers and the surface structure. Scale bars, 500 nm. (C) Volume segmentations of Electron microscopy tomograms showing SSV1 maturation and release by budding. Scale bar, 50 nm. (D) Transmission electron micrograph of a thin section of a SIRV2 infected *Sa. islandicus* cell displaying several pyramidal egress structures. Scale bar, 100 nm. (E) Transmission electron micrographs of an isolated pyramidal egress structure in open conformation isolated after SIRV2 infection of *Sa. islandicus*. Scale bar, 100 nm. Adapted from (Bize et al., 2009, Quax et al., 2011, Quemin et al., 2013, Quemin et al., 2016).

107x193mm (300 x 300 DPI)




Figure 4.1 Main principles in genome organization and genetic information processing in the *Sulfobolales*. Conceptual schemes representing the major elements and principles of macro-level organization of the genomic DNA (adapted from Takemata et al., 2019) (A), micro-level organization of the genomic DNA (partially adapted from Peeters et al., 2015) (B), initiation of replication (C) and initiation of transcription (D).




Figure 5.1 Archaeal cell surface structures involved in planktonic and biofilm growth. Upper
image: Electronic microscopy image from a *S. acidocaldarius* cell where archaealum and pilus can be seen.
Lower image: Schematic model with all proposed cell surface appendages in the *Sulfolobales*: the Aap pili (archaeal adhesive pili), the Ups pili (UV induced pili), the archaealum and the threads. Also depicted is the S-layer and its proteins: SlaA and SlaB.

162x181mm (300 x 300 DPI)




Figure 5.2 Confocal Laser Microscopy images from static biofilm from *S. acidocaldarius* in days 3 to 7 of growth. Cells (DNA) stained with 4',6-diamidino-2-phenylindole (DAPI; blue); extracellular glucose and mannose residues stained with fluorescently labeled concanavalin A (conA; green); and N- acetyl-d-glucosamine residues stained with fluorescently labeled lectin IB4 (yellow). Scale bars: 20 μ m.





Figure 5.3 Model of the *Sulfolobus* S-layer. (A) The *Sulfolobus* S-layer consists of the 2 protein subunits: SlaA dimers (red, orange, yellow) form the outer S-layer canopy. Each SlaA protein is predicted to be rich in β-strands. The SlaA dimer has a boomerang-like shape, the angle of which determines the S-layer unit cell size. SlaB trimers (gray) form the membrane anchors of the S-layer. Each SlaB is predicted to consist of an N-terminal transmembrane domain (TMD), a coiled-coil domain (CC), and 2–3 C-terminal β-sandwich domains (β). SlaA and SlaB proteins are highly glycosylated (green). (B) Electron microscopy image from negatively stained isolated S-layer from *S. acidocaldarius*. (C) SDS/PAGE. M, marker; 1, washed once; 2, washed twice; 3, washed 3 times in detergent. (D) Subtomogram average of fully assembled S-layer. (E) Subtomogram average of SlaB-depleted S-layer. (F) Difference map (pink) overlaid with the complete S-layer visualizes location of SlaB. (Scale bars, and C–E, 20 nm) (Figure adapted from Gambeli et al. 2019)

183x138mm (300 x 300 DPI)

A



B




Figure 5.4. Glycosylation. (A) Comparison of the N-glycan trees of three different Sulfolobus/Saccharolobus species. (B) The current understanding of the N-glycosylation pathway in *S. acidocaldarius*. The N-glycan biosynthesis is initiated by adding nucleotide-activated monosaccharides sequentially to the lipid carrier DolP on the cytoplasmic side of the membrane. The fully assembled DolPP-linked N-glycan (hexasaccharide) is translocated across the membrane and then transferred by AlgB on the specific N-glycosylation sequons in secreted proteins. Suagr code is shown in (A).




Figure 5.5: Schematic model of the cell division process in *Sulfolobus acidocaldarius*. (I) CdvA (red) is the first protein of the *S. acidocaldarius* cell division machinery that arrives at the future site of cell division, before DNA (light blue) segregation starts. (II) During nucleoid condensation (blue) CdvB (light green) forms a ring-like structure at midcell that is anchored to the membrane by CdvA. (III) CdvB provides a scaffold for CdvB1 and B2 (green) that are positioned at the cell center in ring-like structures. Additionally, CdvC (yellow), a homolog of the hexameric ATPase Vps4, localizes at the septum while nucleoid segregation and initial membrane invagination starts. (IV) After nucleoid segregation, the CdvB-ring undergoes proteasomal (purple) degradation. (V) Upon CdvB removal CdvB1 and B2 constrict, leading to the final division of the cell. (VI) Directly after fission the new born cells have an oval shape that rapidly changes to the typical coccoid shape of *S. acidocaldarius* cells. CdvA and CdvC are organized in a ring-like structure as well, however, for a better overview in the model the organization of both proteins at midcell was only indicated.




Figure 6.1 Major mechanisms of thermoacidophily. **1)** Thermoacidophiles have an inverted membrane potential with a positive charge on the inside of the cellular membrane and a negative charge on the outside to prevent the acidification of the cytoplasm by the passive diffusion of protons. **2)** The inverted membrane potential is maintained by transporting cations such as K^+ into the cytoplasm. **3)** Cyclopentyl ring moieties on tetraether lipids increase packing of the tetraether lipids decreasing the permeability of the membrane by protons and increasing cellular heat stability. **4)** Tetraether lipids make a monolayer that is less permeable to protons and more heat stable than diether lipids. **5)** Proton pumps export protons from the cytoplasm to prevent the acidification of the cytoplasm. **6)** Heat damaged or protonated proteins can either be degraded via the proteasome or properly refolded by the thermosome.

307x173mm (150 x 150 DPI)




Figure 6.2 Model of heavy metal resistance in *Sulfolobales*. **1)** CopA and CopB export Cu outside the cell with ATP consumption. CopM is a metal chaperone that forms part of the Cop system, which also includes a transcriptional factor called CopT (not shown). **2)** PolyP can sequestrate cations in its negative surface, **3)** but it can also be degraded by PPX into inorganic phosphate to be exported outside the cell along with cations via PitA or Pho84 transporters. **4)** Some proteins also act sequestering metal ions, as for example Dps. The mechanism for which metals enter the cell is still unknown.




Figure 7.1: Predicted protein kinase and protein phosphatase homologs in the four different *Sulfolobales* species *S. acidocaldarius*, *Sa. islandicus*, *Sa. solfataricus* and *Sulfuri. tokodaii*.

Depicted are the different canonical and non-canonical Hanks type protein kinases and protein phosphatases with their correspondent domain structure. (Lower et al., 2000, Lower & Kennelly, 2002, Lower & Kennelly, 2003, Lower et al., 2004, Haile & Kennelly, 2011, Esser & Siebers, 2013, Ray et al., 2015, Esser et al., 2016, Hoffmann et al., 2017, Huang et al., 2017).

195x154mm (300 x 300 DPI)




Figure 7.2: Representation of known target enzymes in different *Sulfolobales* strains. (*Sa. solfataricus* in pink, *Sa. islandicus* in orange, *S. acidocaldarius* in blue and *Sulfurisphaera tokodaii* in grey) with their physiological function. From upper left to right: The FadR transcriptional regulator represses transcription of the fatty acid gene cluster and dissociates from the DNA upon binding to acyl-CoA. Phosphorylation of FadR by the ePK ArnC (Saci_1196) prevents acyl-CoA binding and thus hinders transcription of the gene cluster (Maklad et al., 2020). The archaellum regulatory network consists of the gene cluster arlBXGFHIJ (flaBXGFHIJ), which encodes the motility structure, the archaellum, and is under the control of two promoters, one upstream of *arlB* (flaB) being induced under starvation and one weak promoter upstream of *arlX* (flaX). The two negative regulators ArnA (Saci_1210) and ArnB (Saci_1211) were shown to be phosphorylated by the ePKs ArnC and ArnD (Saci_1694) and dephosphorylated by the PP2A (Saci_0884). Deletion of the PP2A led to a hypermotile phenotype suggesting a negative influence on the gene cluster (Reimann et al., 2012, Hoffmann et al., 2017). The DNA binding protein AbfR1 (Saci_0446) is a positive regulator of the *arlB* (flaB) promoter (Orell et al., 2013). Phosphorylation of AbfR1 inhibits DNA binding and thus regulates biofilm formation and motility (Li et al., 2017). The FHA domain containing protein ST0829 was shown to interact and be phosphorylated by the ePK ST1565 indicating a role in transcription regulation (Duan & He, 2011). The Holliday Junction Resolvase (Hjc) (SiRe_1431) is phosphorylated by the aPK SiRe_0171 facilitating DNA repair (Huang et al., 2019). The phosphohexomutase (SSO0207) exhibited a decreased V_{max} value after being phosphorylated (Ray et al., 2005). The Rio kinases (Saci_0796 and Saci_0965) were shown to play a role in the ribosome maturation of the small subunit (SSU) (Knüppel et al., 2018).

178x152mm (150 x 150 DPI)




Figure 8.1 An overview of the central metabolism in Sulfolobales. Dashed arrows indicate pathways, which have not yet been experimentally demonstrated. Abbreviations: F6P, fructose 6-phosphate; DHAP, dihydroxyacetone phosphate; GAP, glyceraldehyde 3-phosphate; D-KDG, 2-keto-3-deoxy-D-gluconate; D-KDGal, 2-keto-3-deoxy-D-galactonate; D-KDA, 2-keto-3-deoxy-D-arabinonate; L-KDA, 2-keto-3-deoxy-L-arabinonate; D-KDX, 2-keto-3-deoxy-D-xylonate; AA, amino acid; ED, Entner-Doudoroff pathway; EMP, Embden-Meyerhof-Parnas pathway; RuMP, reversed ribulose monophosphate pathway; TCA, Tricarboxylic acid cycle; 3HP/4HB, 3-hydroxypropionate/4-hydroxybutyrate cycle; ABC, ATP binding cassette transporters; RC, respiratory chain.




Figure 8.2 The enzymatic pathway of the 3-hydroxypropionate/4-hydroxybutyrate (3-HP/4-HB) cycle as characterized in *Metallosphaera sedula*. Enzyme names are contained within the yellow oval with arrows indicating reactions for which they have known catalytic activity. Enzymes in pink have shown activity for only a single reaction in the 3-HP/4-HB cycle; enzymes in green exhibit activity on multiple steps in the cycle.

338x190mm (150 x 150 DPI)




Figure 8.3 Current knowledge of the mechanism of sulfur oxidation and reduction in the Sulfolobales. A) Sulfur reduction and B) sulfur oxidation. Solid arrows indicate involvement in a reaction; dotted arrows represent transport of species; dashed lines indicate that the function is suspected but has not been demonstrated experimentally in the *Sulfolobales*. Enzyme colors indicate general grouping of function: coupled to electron transport chain (blue), involved in transporting or trafficking sulfur species (yellow), transformation of sulfur species with no energy conservation (orange), transformation of sulfur species directly coupled to energy-conserving biomolecules (green). Abbreviations: sulfur reductase (Sre), hydrogenase (Hyn), heterodisulfide reductase (Hdr), tetrathionate hydrolase (TetH), sulfide:quinone oxidoreductase (SQO), sulfite:acceptor oxidoreductase (SAOR), thiosulfate:quinone oxidoreductase (TQO), sulfur oxygenase reductase (SOR), adenosine-5'-phosphosulfate reductase (APSR), adenosine-5'-phosphosulfate (APS), adenylylsulfate:phosphate adenylyltransferase (APAT), ATP sulfurylase (ATPS), adenylate kinase (AK).

340x193mm (150 x 150 DPI)




Figure 9.1 Different mechanisms for the generation of markerless deletion mutants. A) Plasmid integration occurs via single-crossover, resulting in a merodiploidal form. After counterselection with 5-FOA, the *pyrEF* marker cassette is looped out, either with or without the GOI, resulting in a theoretical ratio of one to one in mutated and wild type cells. **B)** and **C)** Double-crossover is feasible by introducing a linearized vector. Depending on the experimental design, either parts of the GOI (**B**) or an upstream (US) region (**C**) are introduced for recombination. Counterselection with 5-FOA produces marker-free deletion mutants. **D)** A plasmid containing a CRISPR array as well as a repair fragment with homologous sequences to the GOI are introduced into a recipient strain. Upon induction, crRNA is transcribed and forms a ribonucleoprotein complex with the endogenous Cas protein, scanning the genomic DNA for the spacer sequence and cutting it. Only colonies which conducted recombination with the repair fragment survive. GOI, gene of interest; US, upstream; DS, downstream, *pyrEF*, *pyrEF* marker cassette; crRNA, CRISPR RNA; crRNP, ribonucleoprotein complex consisting of crRNA and Cas protein.




Figure 9.2 Mechanisms of sulfidic ore dissolution. A) Thiosulfate mechanism **B)** and polysulfide mechanism. Green dashed arrows indicate biological steps; solid arrows indicate spontaneous abiotic reactions; blue dashed-dotted arrows indicate an overall transformation involving multiple reaction steps; yellow dashed-dotted arrows represent phase transition. Bold boxes around a species indicates that this is the dominant sulfur product of the dissolution process.

339x191mm (150 x 150 DPI)

Figure Legends

Figure 1.1 Timeline of thermoacidophile isolations and major events.

Timeline contains the organism's names at the time of the associated event. The following are the current classifications: *Sulfolobus brierleyi* (f. *Acidianus brierleyi*), *Saccharolobus solfataricus* (f. *Sulfolobus solfataricus*), *Acidianus ambivalens* (f. *Desulfurolobus ambivalens*), *Saccharolobus shibatae* (f. *Sulfolobus shibatae*) *Sulfuracidifex metallicus* (f. *Sulfolobus metallicus*), *Metallosphaera hakonensis* (f. *Sulfolobus hakonensis*), *Sulfurisphaera tokadaii* (f. *Sulfolobus tokodaii*), *Saccharolobus islandicus* (f. *Sulfolobus islandicus*).

Figure 2.1 16S Phylogeny tree of thermoacidophilic organisms.

Figure 3.1 Thermoacidophilic archaeal viruses and their infection mechanisms.

(A) Schematic representation of virion morphologies of viruses infecting thermoacidophilic archaea as described in the text. **(B)** Segmented tomographic volume of a SIRV2 virion (red) attached to a surface filament of *Sa. islandicus* (green) with help of the three terminal virion fibers (blue). Inset depicts a magnification of the interaction between the tail fibers and the surface structure. Scale bars, 500 nm. **(C)** Volume segmentations of Electron microscopy tomograms showing SSV1 maturation and release by budding. Scale bar, 50 nm. **(D)** Transmission electron micrograph of a thin section of a SIRV2 infected *Sa. islandicus* cell displaying several pyramidal egress structures. Scale bar, 100 nm. **(E)** Transmission electron micrographs of an isolated pyramidal egress structure in open conformation isolated after SIRV2 infection of *Sa. islandicus*. Scale bar, 100 nm Adapted from (Bize *et al.*, 2009, Quax *et al.*, 2011, Quemin *et al.*, 2013, Quemin *et al.*, 2016).

Figure 4.1 Main principles in genome organization and genetic information

processing in the *Sulfolobales*. Conceptual schemes representing the major elements and principles of macro-level organization of the genomic DNA (adapted from (Takemata *et al.*, 2019) (A), micro-level organization of the genomic DNA (partially adapted from (Peeters *et al.*, 2015) (B), initiation of replication (C) and initiation of transcription (D).

Figure 5.1 Archaeal cell surface structures involved in planktonic and biofilm

growth. Upper image: Electronic microscopy image from a *S. acidocaldarius* cell where archaellum and pilus can be seen. **Lower image:** Schematic model with all proposed cell surface appendages in the *Sulfolobales*: the Aap pili (archaeal adhesive pili), the Ups pili (UV induced pili), the archaellum and the threads. Also depicted is the S-layer and its proteins: SlaA and SlaB.

Figure 5.2 Confocal Laser Microscopy images from static biofilm from

***S. acidocaldarius* in days 3 to 7 of growth.** Cells (DNA) stained with 4',6-diamidino-2-phenylindole (DAPI; blue); extracellular glucose and mannose residues stained with fluorescently labeled concanavalin A (conA; green); and N- acetyl-d- glucosamine residues stained with fluorescently labeled lectin IB4 (yellow). Scale bars: 20 μ m.

Figure 5.3 Model of the *Sulfolobus* S-layer. (A) The *Sulfolobus* S-layer consists of the 2

protein subunits: SlaA dimers (red, orange, yellow) form the outer S-layer canopy. Each SlaA protein is predicted to be rich in β -strands. The SlaA dimer has a boomerang-like shape, the angle of which determines the S-layer unit cell size. SlaB trimers (gray) form the membrane anchors of the S-layer. Each SlaB is predicted to consist of an N-terminal transmembrane

1
2
3 domain (TMD), a coiled-coil domain (CC), and 2–3 C-terminal β -sandwich domains (β). SlaA
4 and SlaB proteins are highly glycosylated (green). (B) Electron microscopy image from
5 negatively stained isolated S-layer from *S. acidocaldarius*. (C) SDS/PAGE. M, marker; 1,
6 washed once; 2, washed twice; 3, washed 3 times in detergent. (D) Subtomogram average of
7 fully assembled S-layer. (E) Subtomogram average of SlaB-depleted S-layer. (F) Difference map
8 (pink) overlaid with the complete S-layer visualizes location of SlaB. (Scale bars, and C–E, 20
9 nm) (Figure adapted from Gambeli et al. 2019)

10
11
12
13
14
15
16
17
18
19
20
21
22 **Figure 5.4. Glycosylation.** (A) Comparison of the N-glycan trees of three different
23 *Sulfolobus/Saccharolobus* species. (B) The current understanding of the N-glycosylation
24 pathway in *S. acidocaldarius*. The N-glycan biosynthesis is initiated by adding nucleotide-
25 activated monosaccharides sequentially to the lipid carrier DolP on the cytoplasmic side of the
26 membrane. The fully assembled DolPP-linked N-glycan (hexasaccharide) is translocated across
27 the membrane and then transferred by AlgB on the specific N-glycosylation sequons in secreted
28 proteins. Suagr code is shown in (A).

29
30
31
32
33
34
35
36
37
38
39
40 **Figure 5.5: Schematic model of the cell division process in *Sulfolobus acidocaldarius*.**
41
42 (I) CdvA (red) is the first protein of the *S. acidocaldarius* cell division machinery that arrives at
43 the future site of cell division, before DNA (light blue) segregation starts. (II) During nucleoid
44 condensation (blue) CdvB (light green) forms a ring-like structure at midcell that is anchored to
45 the membrane by CdvA. (III) CdvB provides a scaffold for CdvB1 and B2 (green) that are
46 positioned at the cell center in ring-like structures. Additionally, CdvC (yellow), a homolog of
47 the hexameric ATPase Vps4, localizes at the septum while nucleoid segregation and initial
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 membrane invagination starts. **(IV)** After nucleoid segregation, the CdvB-ring undergoes
4 proteasomal (purple) degradation. **(V)** Upon CdvB removal CdvB1 and B2 constrict, leading to
5 the final division of the cell. **(VI)** Directly after fission the new born cells have an oval shape that
6 rapidly changes to the typical coccoid shape of *S. acidocaldarius* cells. CdvA and CdvC are
7 organized in a ring-like structure as well, however, for a better overview in the model the
8 organization of both proteins at midcell was only indicated.
9
10
11
12
13
14
15
16
17
18

19 **Figure 6.1 Major mechanisms of thermoacidophily.** **1)** Thermoacidophiles have an
20 inverted membrane potential with a positive charge on the inside of the cellular membrane and a
21 negative charge on the outside to prevent the acidification of the cytoplasm by the passive
22 diffusion of protons. **2)** The inverted membrane potential is maintained by transporting cations
23 such as K^+ into the cytoplasm. **3)** Cyclopentyl ring moieties on tetraether lipids increase packing
24 of the tetraether lipids decreasing the permeability of the membrane by protons and increasing
25 cellular heat stability. **4)** Tetraether lipids make a monolayer that is less permeable to protons and
26 more heat stable than diether lipids. **5)** Proton pumps export protons from the cytoplasm to
27 prevent the acidification of the cytoplasm. **6)** Heat damaged or protonated proteins can either be
28 degraded via the proteasome or properly refolded by the thermosome.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 **Figure 6.2 Model of heavy metal resistance in *Sulfolobales*.** **1)** CopA and CopB export
46 Cu outside the cell with ATP consumption. CopM is a metal chaperone that forms part of the
47 Cop system, which also includes a transcriptional factor called CopT (not shown). **2)** PolyP can
48 sequestrate cations in its negative surface, **3)** but it can also be degraded by PPX into inorganic
49 phosphate to be exported outside the cell along with cations via PitA or Pho84 transporters. **4)**
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Some proteins also act sequestering metal ions, as for example Dps. The mechanism for which
4 metals enter the cell is still unknown.
5
6
7
8
9

10 **Figure 7.1: Predicted protein kinase and protein phosphatase homologs in the four**
11 **different *Sulfolobales* species *S. acidocaldarius*, *Sa. islandicus*, *Sa. solfataricus* and *Sulfuri-***
12 ***tokodaii*.** Depicted are the different canonical and non-canonical Hanks type protein kinases and
13 protein phosphatases with their correspondent domain structure. (Lower *et al.*, 2000, Lower &
14 Kennelly, 2002, Lower & Kennelly, 2003, Lower *et al.*, 2004, Haile & Kennelly, 2011, Esser &
15 Siebers, 2013, Ray *et al.*, 2015, Esser *et al.*, 2016, Hoffmann *et al.*, 2017, Huang *et al.*, 2017).
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 7.2: Representation of known target enzymes in different *Sulfolobales* strains. (*Sa. solfataricus* in pink, *Sa. islandicus* in orange, *S. acidocaldarius* in blue and *Sulfuri-*
tokodaii in grey) with their physiological function. From upper left to right: The FadR
transcriptional regulator represses transcription of the fatty acid gene cluster and dissociates from
the DNA upon binding to acyl-CoA. Phosphorylation of FadR by the ePK ArnC (Saci_1196)
prevents acyl-CoA binding and thus hinders transcription of the gene cluster (Maklad *et al.*,
2020). The archaellum regulatory network consists of the gene cluster *arlBXGFHIJ*
(*flaBXGFHIJ*), which encodes the motility structure, the archaellum, and is under the control of
two promoters, one upstream of *arlB* (*flaB*) being induced under starvation and one weak
promoter upstream of *arlX* (*flaX*). The two negative regulators ArnA (Saci_1210) and ArnB
(Saci_1211) were shown to be phosphorylated by the ePKs ArnC and ArnD (Saci_1694) and
dephosphorylated by the PP PP2A (Saci_0884). Deletion of the PP2A led to a hypermotile
phenotype suggesting a negative influence on the gene cluster (Reimann *et al.*, 2012, Hoffmann

et al., 2017). The DNA binding protein AbfR1 (Saci_0446) is a positive regulator of the *arlB* (*flaB*) promoter (Orell et al., 2013). Phosphorylation of AbfR1 inhibits DNA binding and thus regulates biofilm formation and motility (Li et al., 2017). The FHA domain containing protein ST0829 was shown to interact and be phosphorylated by the ePK ST1565 indicating a role in transcription regulation (Duan & He, 2011). The Holliday Junction Resolvase (Hjc) (SiRe_1431) is phosphorylated by the aPK SiRe_0171 facilitating DNA repair (Huang et al., 2019). The phosphohexomutase (SSO0207) exhibited a decreased Vmax value after being phosphorylated (Ray et al., 2005). The Rio kinases (Saci_0796 and Saci_0965) were shown to play a role in the ribosome maturation of the small subunit (SSU) (Knüppel et al., 2018).

Figure 8.1 An overview of the central metabolism in *Sulfolobales*. Dashed arrows indicate pathways, which have not yet been experimentally demonstrated. Abbreviations: F6P, fructose 6-phosphate; DHAP, dihydroxyacetone phosphate; GAP, glyceraldehyde 3-phosphate; D-KDG, 2-keto-3-deoxy-D-gluconate; D-KDGal, 2-keto-3-deoxy-D-galactonate; D-KDA, 2-keto-3-deoxy-D-arabinoate; L-KDA, 2-keto-3-deoxy-L-arabinoate; D-KDX, 2-keto-3-deoxy-D-xylose; AA, amino acid; ED, Entner-Doudoroff pathway; EMP, Embden-Meyerhof-Parnas pathway; RuMP, reversed ribulose monophosphate pathway; TCA, Tricarboxylic acid cycle; 3HP/4HB, 3-hydroxypropionate/4-hydroxybutyrate cycle; ABC, ATP binding cassette transporters; RC, respiratory chain.

Figure 8.2 The enzymatic pathway of the 3-hydroxypropionate/4-hydroxybutyrate (3-HP/4-HB) cycle as characterized in *Metallosphaera sedula*. Enzyme names are contained within the yellow oval with arrows indicating reactions for which they have known catalytic

1
2
3 activity. Enzymes in pink have shown activity for only a single reaction in the 3-HP/4-HB cycle;
4 enzymes in green exhibit activity on multiple steps in the cycle.
5
6
7
8
9

10 **Figure 8.3 Current knowledge of the mechanism of sulfur oxidation and reduction**
11 in the *Sulfolobales*. **A)** Sulfur reduction and **B)** sulfur oxidation. Solid arrows indicate
12 involvement in a reaction; dotted arrows represent transport of species; dashed lines indicate that
13 the function is suspected but has not been demonstrated experimentally in the *Sulfolobales*.
14 Enzyme colors indicate general grouping of function: coupled to electron transport chain (blue),
15 involved in transporting or trafficking sulfur species (yellow), transformation of sulfur species
16 with no energy conservation (orange), transformation of sulfur species directly coupled to
17 energy-conserving biomolecules (green). Abbreviations: sulfur reductase (Sre), hydrogenase
18 (Hyn), heterodisulfide reductase (Hdr), tetrathionate hydrolase (TetH), sulfide:quinone
19 oxidoreductase (SQO), sulfite:acceptor oxidoreductase (SAOR), thiosulfate:quinone
20 oxidoreductase (TQO), sulfur oxygenase reductase (SOR), adenosine-5'-phosphosulfate
21 reductase (APSR), adenosine-5'-phosphosulfate (APS), adenylylsulfate:phosphate
22 adenylyltransferase (APAT), ATP sulfurylase (ATPS), adenylate kinase (AK).
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 9.1 Different mechanisms for the generation of markerless deletion mutants.

A) Plasmid integration occurs via single-crossover, resulting in a merodiploidal form. After
counterselection with 5-FOA, the *pyrEF* marker cassette is looped out, either with or without the
GOI, resulting in a theoretical ration of one to one in mutated and wild type cells. **B)** and **C)**
Double-crossover is feasible by introducing a linearized vector. Depending on the experimental
design, either parts of the GOI (**B**) or an upstream (US) region (**C**) are introduced for
recombination. Counterselection with 5-FOA produces marker-free deletion mutants. **D)** A

1
2
3 plasmid containing a CRISPR array as well as a repair fragment with homologous sequences to
4 the GOI are introduced into a recipient strain. Upon induction, crRNA is transcribed and forms a
5 ribonucleoprotein complex with the endogenous Cas protein, scanning the genomic DNA for the
6 spacer sequence and cutting it. Only colonies which conducted recombination with the repair
7 fragment survive. GOI, gene of interest; US, upstream; DS, downstream, *pyrEF*, *pyrEF* marker
8 cassette; crRNA, CRISPR RNA; crRNP, ribonucleoprotein complex consisting of crRNA and
9 Cas protein.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 9.2 Mechanisms of sulfidic ore dissolution. A) thiosulfate mechanism B) and
polysulfide mechanism. Green dashed arrows indicate biological steps; solid arrows indicate
spontaneous abiotic reactions; blue dashed-dotted arrows indicate an overall transformation
involving multiple reaction steps; yellow dashed-dotted arrows representative phase transition.
Bold boxes around a species indicates that this is the dominant sulfur product of the dissolution
process.

References:

Bize A, Karlsson EA, Ekefjärd K, Quax TEF, Pina M, Prevost M-C, Forterre P, Tenaillon O, Bernander R & Prangishvili D (2009) A unique virus release mechanism in the Archaea. *Proc Natl Acad Sci USA* **106**: 11306-11311.

Duan X & He Z-G (2011) Characterization of the specific interaction between archaeal FHA domain-containing protein and the promoter of a flagellar-like gene-cluster and its regulation by phosphorylation. *Biochem Biophys Res Commun* **407**: 242-247.

Esser D & Siebers B (2013) Atypical protein kinases of the RIO family in archaea. *Biochem Soc Trans* **41**: 399-404.

Esser D, Hoffmann L, Pham TK, Bräsen C, Qiu W, Wright PC, Albers S-V & Siebers B (2016) Protein phosphorylation and its role in archaeal signal transduction. *FEMS Microbiol Rev* **40**: 625-647.

Haile JD & Kennelly PJ (2011) The activity of an ancient atypical protein kinase is stimulated by ADP-ribose *in vitro*. *Arc Biochem Biophys* **511**: 56-63.

Hoffmann L, Schummer A, Reimann J, Haurat MF, Wilson AJ, Beeby M, Warscheid B & Albers SV (2017) Expanding the archaellum regulatory network—the eukaryotic protein kinases ArnC and ArnD influence motility of *Sulfolobus acidocaldarius*. *MicrobiologyOpen* **6**.

Huang Q, Zhong Q, Mayaka J, Ni J & Shen Y (2017) Autophosphorylation and cross-phosphorylation of protein kinases from the crenarchaeon *Sulfolobus islandicus*. *Front Microbiol* **8**: 2173.

Huang Q, Mayaka JBA, Zhong Q, Zhang C, Hou G, Ni J & Shen Y (2019) Phosphorylation of the archaeal Holliday junction resolvase Hjc inhibits its catalytic activity and facilitates DNA repair in *Sulfolobus islandicus* REY15A. *Front Microbiol* **10**: 1214.

Knüppel R, Christensen RH, Gray FC, Esser D, Strauß D, Medenbach J, Siebers B, MacNeill SA, LaRonde N & Ferreira-Cerca S (2018) Insights into the evolutionary conserved regulation of Rio ATPase activity. *Nucleic Acids Res* **46**: 1441-1456.

Li L, Banerjee A, Bischof LF, Maklad HR, Hoffmann L, Henche AL, Veliz F, Bildl W, Schulte U & Orell A (2017) Wing phosphorylation is a major functional determinant of the Lrs14-type biofilm and motility regulator AbfR1 in *Sulfolobus acidocaldarius*. *Mol Microbiol* **105**: 777-793.

Lower BH & Kennelly PJ (2002) The membrane-associated protein-serine/threonine kinase from *Sulfolobus solfataricus* is a glycoprotein. *J Bacteriol* **184**: 2614-2619.

Lower BH & Kennelly PJ (2003) Open reading frame sso2387 from the archaeon *Sulfolobus solfataricus* encodes a polypeptide with protein-serine kinase activity. *J Bacteriol* **185**: 3436-3445.

1
2
3 Lower BH, Bischoff KM & Kennelly PJ (2000) The archaeon *Sulfolobus solfataricus* contains a
4 membrane-associated protein kinase activity that preferentially phosphorylates threonine
5 residues *in vitro*. *J Bacteriol* **182**: 3452-3459.
6
7 Lower BH, Potters MB & Kennelly PJ (2004) A phosphoprotein from the archaeon *Sulfolobus*
8 *solfataricus* with protein-serine/threonine kinase activity. *J Bacteriol* **186**: 463-472.
9
10 Maklad HR, Gutierrez GJ, Esser D, Siebers B & Peeters E (2020) Phosphorylation of the acyl-
11 CoA binding pocket of the FadR transcription regulator in *Sulfolobus acidocaldarius*. *Biochimie*.
12
13 Orell A, Peeters E, Vassen V, Jachlewski S, Schalles S, Siebers B & Albers S-V (2013) Lrs14
14 transcriptional regulators influence biofilm formation and cell motility of *Crenarchaea*. *ISME J*
15 7: 1886-1898.
16
17 Peeters E, Driessen RP, Werner F & Dame RT (2015) The interplay between nucleoid
18 organization and transcription in archaeal genomes. *Nat Rev Microbiol* **13**: 333-341.
19
20 Quax TE, Lucas S, Reimann J, Pehau-Arnaudet G, Prevost M-C, Forterre P, Albers S-V &
21 Prangishvili D (2011) Simple and elegant design of a virion egress structure in Archaea. *Proc
22 Natl Acad Sci* **108**: 3354-3359.
23
24 Quemin ER, Lucas S, Daum B, Quax TE, Kühlbrandt W, Forterre P, Albers S-V, Prangishvili D
25 & Krupovic M (2013) First insights into the entry process of hyperthermophilic archaeal viruses.
26 *J Virol* **87**: 13379-13385.
27
28 Quemin ERJ, Chlanda P, Sachse M, Forterre P, Prangishvili D & Krupovic M (2016)
29 Eukaryotic-like virus budding in archaea. *mBio* **7**.
30
31 Ray WK, Potters MB, Haile JD & Kennelly PJ (2015) Activation of SsoPK4, an archaeal eIF2 α
32 kinase homolog, by oxidized CoA. *Proteomes* **3**: 89-116.
33
34 Ray WK, Keith SM, DeSantis AM, Hunt JP, Larson TJ, Helm RF & Kennelly PJ (2005) A
35 phosphohexamutase from the archaeon *Sulfolobus solfataricus* is covalently modified by
36 phosphorylation on serine. *J Bacteriol* **187**: 4270-4275.
37
38 Reimann J, Lassak K, Khadouma S, Ettema TJ, Yang N, Driessen AJ, Klingl A & Albers SV
39 (2012) Regulation of archaella expression by the FHA and von Willebrand domain-containing
40 proteins ArnA and ArnB in *Sulfolobus acidocaldarius*. *Mol Microbiol* **86**: 24-36.
41
42 Takemata N, Samson RY & Bell SD (2019) Physical and functional compartmentalization of
43 archaeal chromosomes. *Cell* **179**: 165-179.e118.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2.1 Thermoacidophile Organisms

Kingdom	Phylum/Division	Genus/Species	T _{opt} (°C)	pH _{opt}	Isolation Site (Locale, Country)	Reference
Eukarya	Rhodophyta	<i>Galdieria</i> <i>Sulphuraria</i> (Merola)	45	2-3	Solfatara (Pozzuoli, Campania, Italy)	(Merola, 1982)
	Proteobacteria	<i>Acidithiobacillus</i> (<i>A. caldus</i>)	25-45 (45)	2.0-4.0 (2.0- 2.5)	Coal spoil enrichment (Belfast, Northern Ireland, United Kingdom)	(Dopson, 2016)
	<i>Nitrospirae</i>	<i>Leptospirillum</i> (<i>L. ferriphilum</i>)	30-43 (30- 37)	1.4-3.0 (1.4- 1.8)	Bioleaching Tank (South Africa)	(Coram & Rawlings, 2002)
		<i>Sulfobacillus</i> <i>thermosulfidooxidans</i>	50-55	1.7-2.4	Copper-zinc-pyrite ore (Nikolaev Mine East Kazakhstan, Kazakhstan)	(Bogdanova <i>et al.</i> , 2006)
		<i>Sulfobacillus sibericus</i>	55	2.0-2.5	Nezhdaninskoe ore deposit East Siberia, Republic of Sakha, Russian Federation)	(Melamud <i>et al.</i> , 2003)
	<i>Aquificae</i>	<i>Alicyclobacillus</i> (<i>A. acidocaldarius</i>)	35-65 (60- 65)	1.5-4.5 (3.0- 4.0)	Hot spring (Yellowstone NP, Wyoming, USA)	(Darland & Brock, 1971) (Karavaiko <i>et al.</i> , 2005)
		<i>Hydrogenobaculum</i> <i>acidophilum</i>	65	3.0-4.0	Solfatara (Tsumagoi, Gunma, Japan)	(Shima & Suzuki, 1993)

1	Kingdom	Phylum/Division	Genus/Species	T _{opt} (°C)	pH _{opt}	Isolation Site (Locale, Country)	Reference
2	Archaea	Candidate <i>Geoarchaeota</i>	Uncultured	60-78	3.5	Norris Geyser Basin (Yellowstone, Wyoming, USA)	(Kozubal <i>et al.</i> , 2013)
3		Candidate <i>Marsarchaeota</i>	Uncultured	50-80	3.0-3.5	Thermal Springs (Yellowstone, Wyoming, USA)	(Jay <i>et al.</i> , 2018)
4		<i>Euryarchaea</i>	<i>Aciduliprofundum boonei</i>	70	4.2-4.8	Deep sea vents (Mariner, Lau Basin, near Tonga)	(Reysenbach <i>et al.</i> , 2006)
5			<i>Acidiplasma cupricumulans</i>	53.6	1.0-1.2	Mineral bioleaching heap (Undisclosed, Myanmar)	(Hawkes <i>et al.</i> , 2006)
6			<i>Acidiplasma aeolicum</i>	45	1.4-1.6	Hydrothermal pool (Vulcano Island, Messina, Italy)	(Golyshina <i>et al.</i> , 2009)
7			<i>Picrophilus</i> (<i>P. torridus</i> / <i>P. oshimae</i>)	60	0.7	Solfatara (Hokkaido, Japan)	(Schleper, 1996)
8			<i>Thermoplasma volcanium</i>	60	2.0	Solfatara (Vulcano Island, Messina, Italy)	(Segerer A, 1988)
9			<i>Thermoplasma acidophilum</i>	59	1.0-2.0	Coal refuse pile (Friar Tuck Mine, Indiana, USA)	(Darland <i>et al.</i> , 1970)
10			<i>Thermogymnomononas acidicola</i>	60	3.0	Solfatara (Ohwaku-dani, Hakone, Japan)	(Itoh <i>et al.</i> , 2007)
11		<i>Crenarchaea</i> (<i>Non-Sulfolobales</i>)	<i>Acidilobus aceticus</i>	85	3.9	Thermal spring (Moutnovski, Kamchatka, Russia)	(Prokofeva <i>et al.</i> , 2000)
12			<i>Acidilobus saccharovorans</i>	80-85	3.5-4.0	Thermal spring (Uzon Caldera, Kamchatka, Russia)	(Prokofeva <i>et al.</i> , 2009)
13			<i>Caldisphaera laguensis</i>	70-75	3.5-4.0	Hot spring (Mt. Maquiling, Laguna, Philippines)	(Itoh <i>et al.</i> , 2003)
14			<i>Caldivirga maquilingensis</i>	85	3.7-4.2	Hot spring (Mt. Maquiling, Laguna, Philippines)	(Itoh <i>et al.</i> , 1999)

1	Archaea <i>Crenarchaea</i> <i>(Sulfolobales)</i>	<i>Acidianus ambivalens</i>	81	2.5	Solfatara (Leihnukur, Iceland)	(Zillig <i>et al.</i> , 1986)
2		<i>Acidianus brierleyi</i>	70	1.5-2.0	Thermal spring drainage (Yellowstone, Wyoming, USA)	(Brierley & Brierley, 1973, Segerer <i>et al.</i> , 1986)
3		<i>Acidianus infernus</i>	90	2.0	Mud pot (Naples, Campania, Italy)	(Segerer <i>et al.</i> , 1986)
4		<i>Acidianus sulfidivorans</i>	74	0.8-1.4	Solfatara (Lihir Island, Papua New Guinea)	(Plumb <i>et al.</i> , 2007)
5		<i>Metallosphaera cuprina</i>	65	3.5	Thermal spring (Tengchong, Yunnan, China)	(Liu <i>et al.</i> , 2011)
6		<i>Metallosphaera hakonensis</i>	70	3.0	Thermal spring (Ohwaku-dani, Hakone, Japan)	(Takayanagi <i>et al.</i> , 1996)
7		<i>Metallosphaera prunae</i>	75	2.5	Uranium slag heap (Ronneburg, Hesse, Germany)	(Fuchs <i>et al.</i> , 1995)
8		<i>Metallosphaera sedula</i>	75	2.5	Thermal pool (Naples, Campania, Italy)	(Huber <i>et al.</i> , 1989)
9		<i>Saccharolobus caldissimus</i>	85	3.0	Thermal spring (Ohwaku-dani, Hakone, Japan)	(Sakai & Kurosawa, 2018)
10		<i>Saccharolobus shibatae</i>	81	3.0	Mud pot (Kyushu, Japan)	(Grogan <i>et al.</i> , 1990)
11		<i>Saccharolobus solfataricus</i>	87	4.5	Thermal spring (Agnano, Campania, Italy)	(Zillig <i>et al.</i> , 1980)
12		<i>Stygiolobus azoricus</i>	80	2.5-3.0	Solfatara (São Miguel Island, Azores, Portugal)	(Segerer <i>et al.</i> , 1991)
13		<i>Sulfodiicoccus acidiphilus</i>	65-70	3.0-3.5	Solfatara (Ohwaku-dani, Hakone, Japan)	(Sakai & Kurosawa, 2017)
14		<i>Sulfolobus acidocaldarius</i>	70-75	2.0-3.0	Thermal spring (Yellowstone, Wyoming, USA)	(Brock <i>et al.</i> , 1972)
15		<i>Sulfuracidifex tepidarius</i>	65	3.5	Solfatara (Ohwaku-dani, Hakone, Japan)	(Itoh <i>et al.</i> , 2020)
16		<i>Sulfuracidifex metallicus</i>	65	2.0-3.0	Solfatara (Krafla, Iceland)	
17		<i>Sulfurisphaera ohwakuensis</i>	84	2.0	Thermal spring (Ohwaku, Hakone, Japan)	(Kurosawa <i>et al.</i> , 1998)
18		<i>Sulfurisphaera tokodaii</i>	80	2.5-3.0	Hot spring (Beppu, Kyushu, Japan)	(Suzuki <i>et al.</i> , 2002)
19		<i>Sulfurisphaera javensis</i>	80-85	2.5-4.0	Thermal spring (Java, Indonesia)	(Tsuboi <i>et al.</i> , 2018)

1
2 References:
3
4
56 Bogdanova TI, Tsaplina IA, Kondrat'eva TF, Duda VI, Suzina NE, Melamud VS, Tourova TP & Karavaiko GI (2006) *Sulfobacillus*
7 *thermotolerans* sp. nov., a thermotolerant, chemolithotrophic bacterium. *Int J Syst Evol Microbiol* **56**: 1039-1042.
89 Brierley CL & Brierley JA (1973) A chemoautotrophic and thermophilic microorganism isolated from an acid hot spring. *Can J*
10 *Microbiol* **19**: 183-188.
1112 Brock TD, Brock KM, Belly RT & Weiss RL (1972) *Sulfolobus*: a new genus of sulfur-oxidizing bacteria living at low pH and high
13 temperature. *Arch Microbiol* **84**: 54-68.
1415 Coram NJ & Rawlings DE (2002) Molecular relationship between two groups of the genus *Leptospirillum* and the finding that
16 *Leptospirillum ferriphilum* sp. nov. dominates South African commercial biooxidation tanks that operate at 40 C. *Appl Environ*
17 *Microbiol* **68**: 838-845.
1820 Darland G & Brock TD (1971) *Bacillus acidocaldarius* sp. nov., an acidophilic thermophilic spor-forming bacterium. *J Gen Microbiol*
21 **67**: 9-15.
2224 Darland G, Brock TD, Samsonoff W & Conti SF (1970) A thermophilic, acidophilic mycoplasma isolated from a coal refuse pile. *Science*
25 **170**: 1416-1418.
2627 Dopson M (2016) Physiological and phylogenetic diversity of acidophilic Bacteria. *Acidophiles: Life in extremely acidic*
28 *environments*, (Quatrini R & Barrie JD, eds.), p.^pp. 79-91. Caister Academic Press, Norfolk, UK.
2931 Fuchs T, Huber H, Teiner K, Burggraf S & Stetter KO (1995) *Metallosphaera prunae*, sp. nov., a novel metal-mobilizing,
32 thermoacidophilic archaeum, isolated from a uranium mine in Germany. *Syst App Microbiol* **18**: 560-566.
3334 Golyshina OV, Yakimov MM, Lunsdorf H, Ferrer M, Nimtz M, Timmis KN, Wray V, Tindall BJ & Golyshin PN (2009) *Acidiplasma*
35 *aeolicum* gen. nov., sp. nov., a euryarchaeon of the family *Ferroplasmaceae* isolated from a hydrothermal pool, and transfer of
36 *Ferroplasma cupricumulans* to *Acidiplasma cupricumulans* comb. nov. *Int J Syst Evol Microbiol* **59**: 2815-2823.
3739 Grogan D, Palm P & Zillig W (1990) Isolate B12, which harbours a virus-like element, represents a new species of the archaeabacterial
40 genus *Sulfolobus*, *Sulfolobus shibatae*, sp. nov. *Arch Microbiol* **154**: 594-599.
41

1 Hawkes RB, Franzmann PD, O'Hara G & Plumb JJ (2006) *Ferroplasma cupricumulans* sp. nov., a novel moderately thermophilic,
2 acidophilic archaeon isolated from an industrial-scale chalcocite bioleach heap. *Extremophiles* **10**: 525-530.

3

4 Huber G, Spinnler C, Gambacorta A & Stetter KO (1989) *Metallosphaera sedula* gen. and sp. nov. represents a new genus of aerobic,
5 metal-mobilizing, thermoacidophilic archaeabacteria. *Syst Appl Microbiol* **12**: 38-47.

6

7 Itoh T, Yoshikawa N & Takashina T (2007) *Thermogymnomonas acidicola* gen. nov., sp. nov., a novel thermoacidophilic, cell wall-less
8 archaeon in the order *Thermoplasmatales*, isolated from a solfataric soil in Hakone, Japan. *Int J Syst Evol Microbiol* **57**: 2557-2561.

9

10 Itoh T, Suzuki K, Sanchez PC & Nakase T (1999) *Caldivirga maquilingensis* gen. nov., sp. nov., a new genus of rod-shaped
11 crenarchaeote isolated from a hot spring in the Philippines. *Int J Syst Evol Microbiol* **49**: 1157-1163.

12

13 Itoh T, Suzuki K, Sanchez PC & Nakase T (2003) *Caldisphaera lagunensis* gen. nov., sp. nov., a novel thermoacidophilic crenarchaeote
14 isolated from a hot spring at Mt Maquiling, Philippines. *Int J Syst Evol Microbiol* **53**: 1149-1154.

15

16 Itoh T, Miura T, Sakai HD, Kato S, Ohkuma M & Takashina T (2020) *Sulfuracidifex tepidarius* gen. nov., sp. nov. and transfer of
17 *Sulfolobus metallicus* Huber and Stetter 1992 to the genus *Sulfuracidifex* as *Sulfuracidifex metallicus* comb. nov. *Int J Syst Evol
18 Microbiol* **70**: 1837-1842.

19

20 Jay ZJ, Beam JP, Dlakic M, Rusch DB, Kozubal MA & Inskeep WP (2018) *Marsarchaeota* are an aerobic archaeal lineage abundant in
21 geothermal iron oxide microbial mats. *Nat Microbiol* **3**: 732-740.

22

23 Karavaiko GI, Bogdanova TI, Tourova TP, Kondrat'eva TF, Tsaplina IA, Egorova MA, Krasil'nikova EN & Zakharchuk LM (2005)
24 Reclassification of '*Sulfobacillus thermosulfidooxidans* subsp. *thermotolerans*' strain K1 as *Alicyclobacillus tolerans* sp. nov. and
25 *Sulfobacillus disulfidooxidans* Dufresne et al. 1996 as *Alicyclobacillus disulfidooxidans* comb. nov., and emended description of the
26 genus *Alicyclobacillus*. *Int J Syst Evol Microbiol* **55**: 941-947.

27

28 Kozubal MA, Romine M, Jennings R, Jay ZJ, Tringe SG, Rusch DB, Beam JP, McCue LA & Inskeep WP (2013) *Geoarchaeota*: A new
29 candidate phylum in the Archaea from high-temperature acidic iron mats in Yellowstone National Park. *ISME J* **7**: 622-634.

30

31

32 Kurosawa N, Itoh YH, Iwai T, Sugai A, Uda I, Kimura N, Horiuchi T & Itoh T (1998) *Sulfurisphaera ohwakuensis* gen. nov., sp. nov., a
33 novel extremely thermophilic acidophile of the order *Sulfolobales*. *Int J Syst Evol Microbiol* **48**: 451-456.

34

35 Liu L-J, You X-Y, Guo X, Liu S-J & Jiang C-Y (2011) *Metallosphaera cuprina* sp. nov., an acidothermophilic, metal-mobilizing archaeon.
36 *Int J Syst Evol* **61**: 2395-2400.

37

38

39

40

41

42

43

44

45

46

47

1 Melamud VS, Pivovarova TA, Tourova TP, Kolganova TV, Osipov GA, Lysenko AM, Kondrat'eva TF & Karavaiko GI (2003) *Sulfovbacillus*
2 *sibericus* sp. nov., a new moderately thermophilic bacterium. *MicrobiologyOpen* **72**: 605-612.

3

4 Merola A, Castaldo, R., De Luca, P., Gambardella, R., Musacchio, A., Taddei, R. (1982) Revision of *Cyanidium caldarium*. Three species
5 of acidophilis agae. *G Bot Lt* **116**: 189-195.

6

7 Plumb JJ, Haddad CM, Gibson JaE & Franzmann PD (2007) *Acidianus sulfidivorans* sp. nov., an extremely acidophilic, thermophilic
8 archaeon isolated from a solfatara on Lihir Island, Papua New Guinea, and emendation of the genus description. *Int J Syst Evol*
9 *Microbiol* **57**: 1418-1423.

10

11

12 Prokofeva MI, Miroshnichenko ML, Kostrikina NA, Chernyh NA, Kuznetsiv BB, Tourova TP & Bonch-Osmolovskaya EA (2000)
13 *Acidilobus aceticus* gen. nov., sp. nov., a novel anaerobic thermoacidophilic archaeon from continental hot vents in Kamchatka. *Int J*
14 *Syst Evol Microbiol* **50**: 2001-2008.

15

16

17 Prokofeva MI, Kostrikina NA, Kolganova TV, Tourova TP, Lysenko AM, Lebedinsky AV & Bonch-Osmolovskaya EA (2009) Isolation of
18 the anaerobic thermoacidophilic crenarchaeote *Acidilobus saccharovorans* sp. nov. and proposal of *Acidilobales* ord. nov., including
19 *Acidilobaceae* fam. nov. and *Caldisphaeraceae* fam. nov. *Int J Syst Evol Microbiol* **59**: 3116-3122.

20

21

22 Reysenbach A-L, Liu Y, Banta AB, Beveridge TJ, Kirshtein JD, Schouten S, Tivey MK, Von Damm KL & Voytek MA (2006) A ubiquitous
23 thermoacidophilic archaeon from deep-sea hydrothermal vents. *Nature* **442**: 444-447.

24

25 Sakai HD & Kurosawa N (2017) *Sulfodiicoccus acidiphilus* gen. nov., sp. nov., a sulfur-inhibited thermoacidophilic archaeon belonging
26 to the order *Sulfolobales* isolated from a terrestrial acidic hot spring. *Int J Syst Evol Microbiol* **67**: 1880-1886.

27

28 Sakai HD & Kurosawa N (2018) *Saccharolobus caldissimus* gen. nov., sp. nov., a facultatively anaerobic iron-reducing
29 hyperthermophilic archaeon isolated from an acidic terrestrial hot spring, and reclassification of *Sulfolobus solfataricus* as
30 *Saccharolobus solfataricus* comb. nov. and *Sulfolobus shibatae* as *Saccharolobus shibatae* comb. nov. *Int J Syst Evol Microbiol* **68**:
31 1271-1278.

32

33

34 Schleper C, G. Puhler, H.-P. Klenk, and W. Zillig (1996) *Picrophilus oshimae* and *Picrophilus torridus* fam. nov., gen. nov., sp. nov., two
35 species of hyperacidophilic, thermophilic, heterotrophic, aerobic archaea. *J Syst Evol Microbiol* **46**: 814-816.

36

37

38 Segerer A, Neuner A, Kristjansson JK & Stetter KO (1986) *Acidianus infernus* gen. nov., sp. nov., and *Acidianus brierleyi* comb. nov.:
39 facultatively aerobic, extremely acidophilic thermophilic sulfur-metabolizing archaeabacteria. *Int J Syst Evol* **36**: 559-564.

40

41

42

43

44

45

46

47

1 Segerer A LT, Stetter KO. (1988) *Thermoplasma acidophilum* and *Thermoplasma volcanium* sp. nov. from solfatara fields. *Syst Appl*
2 *Microbiol* **10**: 161-171.

3

4 Segerer AH, Trincone A, Gahrtz M & Stetter KO (1991) *Stygiolobus azoricus* gen. nov., sp. nov. represents a novel genus of anaerobic,
5 extremely thermoacidophilic archaeabacteria of the order Sulfolobales. *Int J Syst Bacteriol* **41**: 495-501.

6

7 Shima S & Suzuki K-I (1993) *Hydrogenobacter acidophilus* sp. nov., a thermoacidophilic, aerobic, hydrogen-oxidizing bacterium
8 requiring elemental sulfur for growth. *Int J Syst Evol Microbiol* **43**: 703-708.

9

10 Suzuki T, Iwasaki T, Uzawa T, Hara K, Nemoto N, Kon T, Ueki T, Yamagishi A & Oshima T (2002) *Sulfolobus tokodaii* sp. nov.(f.
11 Sulfolobus sp. strain 7), a new member of the genus *Sulfolobus* isolated from Beppu Hot Springs, Japan. *Extremophiles* **6**: 39-44.

12

13

14 Takayanagi S, Kawasaki H, Sugimori K, Yamada T, Sugai A, Ito T, Yamasato K & Shioda M (1996) *Sulfolobus hakonensis* sp. nov., a
15 novel species of acidothermophilic archaeon. *Int J Syst Evol* **46**: 377-382.

16

17 Tsuboi K, Sakai HD, Nur N, Stedman KM, Kurosawa N & Suwanto A (2018) *Sulfurisphaera javensis* sp. nov., a hyperthermophilic and
18 acidophilic archaeon isolated from Indonesian hot spring, and reclassification of *Sulfolobus tokodaii* Suzuki et al. 2002 as
19 *Sulfurisphaera tokodaii* comb. nov. *Int J Syst Evol Microbiol* **68**: 1907-1913.

20

21

22 Zillig W, Stetter KO, Wunderl S, Schulz W, Priess H & Scholz I (1980) The *Sulfolobus*-“Caldariella” group: taxonomy on the basis of the
23 structure of DNA-dependent RNA polymerases. *Arch Microbiol* **125**: 259-269.

24

25

26 Zillig W, Yeats S, Holz I, Böck A, Rettenberger M, Groppe F & Simon G (1986) *Desulfurolobus ambivalens*, gen. nov., sp. nov., an
27 autotrophic archaeabacterium facultatively oxidizing or reducing sulfur. *Syst Appl Microbiol* **8**: 197-203.

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

Table 4.1. Overview of characterized transcription regulators in thermoacidophilic archaea and their viruses. N.A. = not applicable (unknown based on published information). ¹Hypothesized; ²UNIPROT number; ³Multifunctional protein with enzymatic and transcription regulatory domains; ⁴Genbank accession number; ⁵Viral regulators are often difficult to classify into a family because of a lack of homology.

Name	Family	Microbial or viral species	Gene number	Physiological role	Regulatory action	Reference
C68	AbrB	<i>S. islandicus</i> plasmid-virus pSSVx	ORFC68	(CRISPR-mediated) virus-host interactions ¹	Activation ¹	(Contursi <i>et al.</i> , 2011)
MerR	ArsR	<i>S. solfataricus</i>	SSO2688	Mercury resistance	Repression	(Schelert <i>et al.</i> , 2006)
IdeR	DtxR	<i>T. acidophilum</i>	TA0872	Iron uptake and homeostasis	Repression	(Yeo <i>et al.</i> , 2012, Yeo <i>et al.</i> , 2014)
ArnA	FHA	<i>S. acidocaldarius</i>	<i>Saci_1210</i>	Motility	Repression	(Duan & He, 2011, Reimann <i>et al.</i> , 2012)
		<i>S. tokodaii</i>	ST0829			
YtrA	GntR	<i>S. acidocaldarius</i>	<i>Saci_1851</i>	Expression of membrane proteins	Repression	(Lemmens <i>et al.</i> , 2019)
BarR	Lrp	<i>S. acidocaldarius</i>	<i>Saci_2136</i>	β-alanine metabolism	Activation	(Liu <i>et al.</i> , 2014)
		<i>S. tokodaii</i>	ST1115			
Lrp	Lrp	<i>S. acidocaldarius</i>	<i>Saci_1588</i>	Global regulation of metabolism and physiology	Dual	(Enoru-Eta <i>et al.</i> , 2000, Vassart <i>et al.</i> , 2013)
		<i>S. solfataricus</i>	SSO0606			
LysM	Lrp	<i>S. acidocaldarius</i>	<i>Saci_0752</i>	Amino acid transport and metabolism	Activation	(Brinkman <i>et al.</i> , 2002, Song <i>et al.</i> , 2013)
		<i>S. solfataricus</i>	SSO0157			
LrpB	Lrp	<i>S. solfataricus</i>	SSO2131	Regulation of pyruvate ferredoxin oxidoreductase and permeases	Dual	(Peeters <i>et al.</i> , 2009, Peeters <i>et al.</i> , 2013)
AbfR1	Lrs14	<i>S. acidocaldarius</i>	<i>Saci_0446</i>	Biofilm formation and motility	Dual	(Orell <i>et al.</i> , 2013, Li <i>et al.</i> , 2017)
Sta1	Lrs14	<i>S. solfataricus</i>	SSO0048	Regulation of SIRV1 viral gene expression	Activation	(Kessler <i>et al.</i> , 2006)
Lrs14	Lrs14	<i>S. solfataricus</i>	SSO1101	N.A.	Repression	(Bell & Jackson, 2000)
Csa3a	MarR	<i>S. islandicus</i>	<i>SiRe_0764</i>	CRISPR spacer acquisition	Activation	(Liu <i>et al.</i> , 2015)
BldR	MarR	<i>S. solfataricus</i>	SSO1352	Detoxification of aromatic compounds	Activation	(Fiorentino <i>et al.</i> , 2007)
BldR2	MarR	<i>S. solfataricus</i>	SSO1082	Stress response to aromatic compounds	N.A.	(Fiorentino <i>et al.</i> , 2011)
N.A.	MarR	<i>S. tokodaii</i>	ST1710	N.A.	N.A.	(Kumarevel <i>et al.</i> , 2009)

MLPT v	MarR	<i>T. volcanium</i>	BAB59904 ₂	N.A.	N.A.	(Liu <i>et al.</i> , 2010)
RbkR	MarR ²	<i>T. acidophilum</i>	<i>TaI064</i>	Riboflavin biosynthesis	Activation ¹	(Rodionova <i>et al.</i> , 2017)
		<i>M. yellowstonensis</i>	EHP68448. 1 ⁴			
FadR	TetR	<i>S. acidocaldarius</i>	<i>Saci_1107</i>	Fatty acid and lipid metabolism	Repression	(Wang <i>et al.</i> , 2019)
HhcR	TrmB	<i>M. yellowstonensis</i>	H2C8P4 ⁴	Autotrophic metabolism	N.A.	(Leyn <i>et al.</i> , 2015)
MalR	TrmB	<i>S. acidocaldarius</i>	<i>Saci_1161</i>	Maltose transport and metabolism	Activation	(Wagner <i>et al.</i> , 2014)
ArnB	vWA	<i>S. acidocaldarius</i>	<i>Saci_1211</i>	Motility	Repression	(Reimann <i>et al.</i> , 2012)
XylR	N.A.	<i>S. acidocaldarius</i>	<i>Saci_2116</i>	Arabinose/xylose transport and metabolism	Activation	(van der Kolk <i>et al.</i> , 2020)
ArnR	N.A.	<i>S. acidocaldarius</i>	<i>Saci_1180</i>	Motility; type IV pili surface structures	Activation	(Lassak <i>et al.</i> , 2013, Bischof <i>et al.</i> , 2019)
ArnR1	N.A.	<i>S. acidocaldarius</i>	<i>Saci_1171</i>	Motility; type IV pili surface structures	Activation	(Lassak <i>et al.</i> , 2013, Bischof <i>et al.</i> , 2019)
CopR/ CopT	N.A.	<i>S. solfataricus</i>	SSO2652	Copper homeostasis	Repression	(Ettema <i>et al.</i> , 2006, Villafane <i>et al.</i> , 2009)
Fur	N.A.	<i>T. volcanium</i>	TVN0292	Oxidative stress	N.A.	(Minoshima <i>et al.</i> , 2014)
SvtR	N.A. ⁵	<i>S. islandicus</i> rod-shaped virus 1 (SIRV1)	ORF56b	Viral development	Repression	(Guillière <i>et al.</i> , 2009)
RIP	N.A. ⁵	<i>Acidianus</i> two-tailed virus (ATV)	ORF145	Global regulation of host transcription	Repression	(Sheppard <i>et al.</i> , 2016)
Stf76	N.A. ⁵	<i>S. islandicus</i> plasmid-virus pSSVx	ORF76	N.A.	N.A.	(Contursi <i>et al.</i> , 2014)
F55	N.A. ⁵	<i>Sulfolobus</i> spindle-shaped virus 1 (SSV1)	T _{lys}	Viral lysogeny and UV induction	Repression	(Fusco <i>et al.</i> , 2015)

References:

Bell SD & Jackson SP (2000) Mechanism of autoregulation by an archaeal transcriptional repressor. *J Biol Chem* **275**: 31624-31629.

Bischof LF, Haurat MF & Albers SV (2019) Two membrane-bound transcription factors regulate expression of various type-IV-pili surface structures in *Sulfolobus acidocaldarius*. *PeerJ* **7**: e6459.

Brinkman AB, Bell SD, Lebbink RJ, De Vos WM & Der Van Oost J (2002) The *Sulfolobus solfataricus* Lrp-like protein LysM regulates lysine biosynthesis in response to lysine availability. *J Biol Chem* **277**: 29537-29549.

Contursi P, D'Ambrosio K, Pirone L, Pedone E, Aucelli T, She Q, De Simone G & Bartolucci S (2011) C68 from the *Sulfolobus islandicus* plasmid-virus pSSVx is a novel member of the AbrB-like transcription factor family. *Biochem J* **435**: 157-166.

Contursi P, Farina B, Pirone L, Fusco S, Russo L, Bartolucci S, Fattorusso R & Pedone E (2014) Structural and functional studies of Stf76 from the *Sulfolobus islandicus* plasmid-virus pSSVx: A novel peculiar member of the winged helix-turn-helix transcription factor family. *Nucleic Acids Res* **42**: 5993-6011.

Duan X & He Z-G (2011) Characterization of the specific interaction between archaeal FHA domain-containing protein and the promoter of a flagellar-like gene-cluster and its regulation by phosphorylation. *Biochem Biophys Res Commun* **407**: 242-247.

Enoru-Eta J, Gigot D, Thia-Toong TL, Glansdorff N & Charlier D (2000) Purification and characterization of Sa-Lrp, a DNA-binding protein from the extreme thermoacidophilic archaeon *Sulfolobus acidocaldarius* homologous to the bacterial global transcriptional regulator Lrp. *J Bacteriol* **182**: 3661-3672.

Ettema TJG, Brinkman AB, Lamers PP, Kornet NG, de Vos WM & van der Oost J (2006) Molecular characterization of a conserved archaeal copper resistance (cop) gene cluster and its copper-responsive regulator in *Sulfolobus solfataricus* P2. *Microbiology* **152**: 1969-1979.

Fiorentino G, Ronca R, Cannio R, Rossi M & Bartolucci S (2007) MarR-like transcriptional regulator involved in detoxification of aromatic compounds in *Sulfolobus solfataricus*. *J Bacteriol* **189**: 7351-7360.

Fiorentino G, Del Giudice I, Bartolucci S, Durante L, Martino L & Del Vecchio P (2011) Identification and physicochemical characterization of BldR2 from *Sulfolobus solfataricus*, a novel archaeal member of the MarR transcription factor family. *Biochemistry* **50**: 6607-6621.

Fusco S, She Q, Fiorentino G, Bartolucci S & Contursi P (2015) Unravelling the Role of the F55 Regulator in the Transition from Lysogeny to UV Induction of *Sulfolobus* Spindle-Shaped Virus 1. *J Virol* **89**: 6453-6461.

Guillière F, Peixeiro N, Kessler A, Raynal B, Desnoues N, Keller J, Delepierre M, Prangishvili D, Sezonov G & Guijarro JI (2009) Structure, function, and targets of the transcriptional regulator SvtR from the hyperthermophilic archaeal virus SIRV1. *J Biol Chem* **284**: 22222-22237.

1
2
3 Kessler A, Sezonov G, Guijarro JI, Desnoues N, Rose T, Delepierre M, Bell SD &
4 Prangishvili D (2006) A novel archaeal regulatory protein, Sta1, activates transcription from
5 viral promoters. *Nucleic Acids Res* **34**: 4837-4845.
6

7
8 Kumarevel T, Tanaka T, Umehara T & Yokoyama S (2009) ST1710-DNA complex crystal
9 structure reveals the DNA binding mechanism of the MarR family of regulators. *Nucleic
10 Acids Res* **37**: 4723-4735.
11

12 Lassak K, Peeters E, Wróbel S & Albers SV (2013) The one-component system ArnR: A
13 membrane-bound activator of the crenarchaeal archaellum. *Mol Microbiol* **88**: 125-139.
14

15 Lemmens L, Tilleman L, De Koning E, Valegård K, Lindås AC, Van Nieuwerburgh F, Maes
16 D & Peeters E (2019) YtrASa, a GntR-family transcription factor, represses two genetic loci
17 encoding membrane proteins in *Sulfolobus acidocaldarius*. *Front Microbiol* **10**: 2084.
18

19 Leyn SA, Rodionova IA, Li X & Rodionov DA (2015) Novel transcriptional regulons for
20 autotrophic cycle genes in crenarchaeota. *J Bacteriol* **197**: 2383-2391.
21

22 Li L, Banerjee A, Bischof LF, Maklad HR, Hoffmann L, Henche AL, Veliz F, Bildl W,
23 Schulte U & Orell A (2017) Wing phosphorylation is a major functional determinant of the
24 Lrs14-type biofilm and motility regulator AbfR1 in *Sulfolobus acidocaldarius*. *Mol Microbiol*
25 **105**: 777-793.
26

27 Liu H, Orell A, Maes D, van Wolferen M, Lindås A-C, Bernander R, Albers S-V, Charlier D
28 & Peeters E (2014) BarR, an Lrp-type transcription factor in *Sulfolobus acidocaldarius*,
29 regulates an aminotransferase gene in a β-alanine responsive manner. *Mol Microbiol* **92**: 625-
30 639.
31

32 Liu T, Li Y, Wang X, Ye Q, Li H, Liang Y, She Q & Peng N (2015) Transcriptional
33 regulator-mediated activation of adaptation genes triggers CRISPR *de novo* spacer
34 acquisition. *Nucleic Acids Res* **43**: 1044-1055.
35

36 Liu Z, Walton TA & Rees DC (2010) A reported archaeal mechanosensitive channel is a
37 structural homolog of MarR-like transcriptional regulators. *Protein Sci* **19**: 808-814.
38

39 Minoshima H, Ikeda Y, Fujii M, Satoh M, Ishikawa T, Itoh Y, Kawashima-Ohya Y, Tomobe
40 K, Omata Y & Kawashima T (2014) Specificity of Fur binding to the oxidative stress
41 response gene promoter in the facultative anaerobic archaeon *Thermoplasma volcanium*. *Biol
42 Pharm Bull* **37**: 481-485.
43

44 Orell A, Peeters E, Vassen V, Jachlewski S, Schalles S, Siebers B & Albers S-V (2013) Lrs14
45 transcriptional regulators influence biofilm formation and cell motility of *Crenarchaea*. *ISME
46 J* **7**: 1886-1898.
47

48 Peeters E, Peixeiro N & Sezonov G (2013) Cis-regulatory logic in archaeal transcription.
49 *Biochem Soc T* **41**: 326-331.
50

51 Peeters E, Albers SV, Vassart A, Driessens AJM & Charlier D (2009) Ss-LrpB, a
52 transcriptional regulator from *Sulfolobus solfataricus*, regulates a gene cluster with a pyruvate
53 ferredoxin oxidoreductase-encoding operon and permease genes. *Mol Microbiol* **71**: 972-988.
54

1
2
3 Reimann J, Lassak K, Khadouma S, Ettema TJ, Yang N, Driessen AJ, Klingl A & Albers SV
4 (2012) Regulation of archaella expression by the FHA and von Willebrand domain-containing
5 proteins ArnA and ArnB in *Sulfolobus acidocaldarius*. *Mol Microbiol* **86**: 24-36.
6
7 Rodionova IA, Vetting MW, Li X, Almo SC, Osterman AL & Rodionov DA (2017) A novel
8 bifunctional transcriptional regulator of riboflavin metabolism in Archaea. *Nucleic Acids Res*
9 **45**: 3785-3799.
10
11 Schelert J, Drozda M, Dixit V, Dillman A & Blum P (2006) Regulation of mercury resistance
12 in the crenarchaeote *Sulfolobus solfataricus*. *J Bacteriol* **188**: 7141-7150.
13
14 Sheppard C, Blombach F, Belsom A, *et al.* (2016) Repression of RNA polymerase by the
15 archaeo-viral regulator ORF145/RIP. *Nat Commun* **7**: 1-13.
16
17 Song N, Duc TN, Van Oeffelen L, Muyldermans S, Peeters E & Charlier D (2013) Expanded
18 target and cofactor repertoire for the transcriptional activator LysM from *Sulfolobus*. *Nucleic*
19 *Acids Res* **41**: 2932-2949.
20
21 van der Kolk N, Wagner A, Wagner M, Waßmer B, Siebers B & Albers S-V (2020)
22 Identification of XylR, the activator of arabinose/xylose inducible regulon in *Sulfolobus*
23 *acidocaldarius* and its application for homologous protein expression. *Front Microbiol* **11**:
24 1066.
25
26 Vassart A, Van Wolferen M, Orell A, Hong Y, Peeters E, Albers S-V & Charlier D (2013)
27 Sa-Lrp from *Sulfolobus acidocaldarius* is a versatile, glutamine-responsive, and architectural
28 transcriptional regulator. *Microbiologyopen* **2**: 75-93.
29
30 Villafane AA, Voskoboinik Y, Cuebas M, Ruhl I & Bini E (2009) Response to excess copper
31 in the hyperthermophile *Sulfolobus solfataricus* strain 98/2. *Biochem Biophys Res Commun*
32 **385**: 67-71.
33
34 Wagner M, Wagner A, Ma X, Kort JC, Ghosh A, Rauch B, Siebers B & Albers S-V (2014)
35 Investigation of the *malE* promoter and MalR, a positive regulator of the maltose regulon, for
36 an improved expression system in *Sulfolobus acidocaldarius*. *Appl Environ Microbiol* **80**:
37 1072-1081.
38
39 Wang K, Sybers D, Maklad HR, Lemmens L, Lewyllie C, Zhou X, Schult F, Bräsen C,
40 Siebers B & Valegård K (2019) A TetR-family transcription factor regulates fatty acid
41 metabolism in the archaeal model organism *Sulfolobus acidocaldarius*. *Nat Commun* **10**: 1-
42 16.
43
44 Yeo HK, Park YW & Lee JY (2014) Structural analysis and insight into metal-ion activation
45 of the iron-dependent regulator from *Thermoplasma acidophilum*. *Acta Crystallogr D* **70**:
46 1281-1288.
47
48 Yeo HK, Kang J, Park YW, Sung JS & Lee JY (2012) Crystallization and preliminary X-ray
49 diffraction analysis of the metalloregulatory protein DtxR from *Thermoplasma acidophilum*.
50 *Acta Crystallogr F* **68**: 172-174.
51
52
53
54
55
56
57
58
59
60

Table 5.1 Biofilms of the *Sulfolobales*

Organism	Adhesion	Biofilm morphology	EPS components	Dispersion	References
<i>Sa. solfataricus</i>	Archaealla and Ups pili	Carpet-like structure, low density of cells (20-30 mm thick)	Glucose, galactose, mannose and N-acetyl-glucosamine residues	Unknown	(Koerdt <i>et al.</i> , 2010, Koerdt <i>et al.</i> , 2012)
<i>S. acidocaldarius</i>	Aap and Ups pili	Dense biofilm with tower-like structures (25-35 mm thick)	Glucose, galactose, mannose and N-acetyl-glucosamine residues, eDNA and proteins	Depends on archaealla	(Koerdt <i>et al.</i> , 2010, Henche <i>et al.</i> , 2012, Koerdt <i>et al.</i> , 2012)
<i>Sulfuri. tokodaii</i>	Unknown	Carpet-like structure with towers (25-35 mm thick)	Glucose, galactose, mannose and N-acetyl-glucosamine residues	Unknown	(Koerdt <i>et al.</i> , 2010)
<i>Sa. metallicus</i>	Unknown	Micro and macro colonies on elemental sulfur	Galactose, mannose and N-acetyl-glucosamine residues, eDNA and proteins	Unknown	(Zhang <i>et al.</i> , 2015)

References:

Henche AL, Koerdt A, Ghosh A & Albers SV (2012) Influence of cell surface structures on crenarchaeal biofilm formation using a thermostable green fluorescent protein. *Environmental microbiology* **14**: 779-793.

Koerdt A, Gödeke J, Berger J, Thormann KM & Albers SV (2010) Crenarchaeal biofilm formation under extreme conditions. *PLoS ONE* **5**: e14104.

Koerdt A, Jachlewski S, Ghosh A, Wingender J, Siebers B & Albers SV (2012) Complementation of *Sulfolobus solfataricus* PBL2025 with an α -mannosidase: effects on surface attachment and biofilm formation. *Extremophiles* **16**: 115-125.

Zhang R, Neu TR, Zhang Y, Bellenberg S, Kuhlicke U, Li Q, Sand W & Vera M (2015) Visualization and analysis of EPS glycoconjugates of the thermoacidophilic archaeon *Sulfolobus metallicus*. *Appl Microbiol Biotechnol* **99**: 7343-7356.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 8.1. An overview of the reported substrates and degradation pathways in *Sa. solfataricus* and *S. acidocaldarius*.

Substrate	Organism	Reference
Carbohydrates	Cellulose	<i>Sa. solfataricus</i> (Girfoglio <i>et al.</i> , 2012)
	Xylan	<i>Sa. solfataricus</i> (Cannio <i>et al.</i> , 2004)
	Glucose	<i>Sa. solfataricus</i> , <i>S. acidocaldarius</i> (De Rosa <i>et al.</i> , 1984, Ahmed <i>et al.</i> , 2005)
	Galactose	<i>Sa. solfataricus</i> (Lamble <i>et al.</i> , 2003)
	D-arabinose	<i>Sa. solfataricus</i> (Brouns <i>et al.</i> , 2006)
	L-arabinose	<i>Sa. solfataricus</i> , <i>S. acidocaldarius</i> (Nunn <i>et al.</i> , 2010, Wagner <i>et al.</i> , 2017)
	D-xylose	<i>Sa. solfataricus</i> , <i>S. acidocaldarius</i> (Nunn <i>et al.</i> , 2010, Wagner <i>et al.</i> , 2017)
Proteins/peptides/ amino acids	Proteins/peptides	<i>Sa. solfataricus</i> , <i>S. acidocaldarius</i> (Lin & Tang, 1990, Gogliettino <i>et al.</i> , 2014)
	Glutamate, methionine, leucine, phenylalanine, isoleucine, threonine, alanine, asparagine, glycine, tyrosine and serine	<i>Sa. solfataricus</i> (Stark <i>et al.</i> , 2017)
Lipids/fatty acids	Olive oil, corn oil, p-nitrophenyl (PNP)-butyrate, PNP- caprylate, PNP- palmitate	<i>Sa. solfataricus</i> (Choi <i>et al.</i> , 2016)
	Tributyrin, tricaproin	<i>S. acidocaldarius</i> (Zweerink <i>et al.</i> , 2017)
	Butyrate, hexanoate	<i>S. acidocaldarius</i> (Wang <i>et al.</i> , 2019)
Other substrates	Ethanol	<i>Sa. solfataricus</i> (Chong <i>et al.</i> , 2007)
	Pechnol	<i>Sa. solfataricus</i> (Izzo <i>et al.</i> , 2005)

References:

Ahmed H, Ettema Thijs JG, Tjaden B, Geerling Ans CM, van der Oost J & Siebers B (2005) The semi-phosphorylative Entner–Doudoroff pathway in hyperthermophilic archaea: a re-evaluation. *Biochem J* **390**: 529-540.

Brouns SJ, Walther J, Snijders APL, *et al.* (2006) Identification of the missing links in prokaryotic pentose oxidation pathways. *J Biol Chem* **281**: 27378-27388.

Cannio R, Di Prizito N, Rossi M & Morana A (2004) A xylan-degrading strain of *Sulfolobus solfataricus*: Isolation and characterization of the xylanase activity. *Extremophiles : life under extreme conditions* **8**: 117-124.

Choi Y-H, Park Y-J, Yoon S-J & Lee H-B (2016) Purification and characterization of a new inducible thermostable extracellular lipolytic enzyme from the thermoacidophilic archaeon *Sulfolobus solfataricus* P1. *J Mol Catalysis B: Enzymatic* **124**: 11-19.

Chong PK, Burja AM, Radianingtyas H, Fazeli A & Wright PC (2007) Proteome and transcriptional analysis of ethanol-grown *Sulfolobus solfataricus* P2 reveals ADH2, a potential alcohol dehydrogenase. *J Proteome Res* **6**: 3985-3994.

De Rosa M, Gambacorta A, Nicolaus B, Giardina P, Poerio E & Buonocore V (1984) Glucose metabolism in the extreme thermoacidophilic archaebacterium *Sulfolobus solfataricus*. *Biochem J* **224**: 407-414.

Girfoglio M, Rossi M & Cannio R (2012) Cellulose degradation by *Sulfolobus solfataricus* requires a cell-anchored endo-beta-1-4-glucanase. *Journal of bacteriology* **194**: 5091-5100.

Gigliettino M, Riccio A, Cocca E, Rossi M, Palmieri G & Balestrieri M (2014) A new pepstatin-insensitive thermopsin-like protease overproduced in peptide-rich cultures of *Sulfolobus solfataricus*. *Int J Mol Sci* **15**: 3204-3219.

Izzo V, Notomista E, Picardi A, Pennacchio F & Di Donato A (2005) The thermophilic archaeon *Sulfolobus solfataricus* is able to grow on phenol. *Res Microbiol* **156**: 677-689.

Lamble HJ, Heyer NI, Bull SD, Hough DW & Danson MJ (2003) Metabolic pathway promiscuity in the archaeon *Sulfolobus solfataricus* revealed by studies on glucose dehydrogenase and 2-keto-3-deoxygluconate aldolase. *J Biol Chem* **278**: 34066-34072.

Lin X-l & Tang J (1990) Purification, characterization, and gene cloning of thermopsin, a thermostable acid protease from *Sulfolobus acidocaldarius*. *J Biol Chem* **265**: 1490-1495.

Nunn CE, Johnsen U, Schonheit P, Fuhrer T, Sauer U, Hough DW & Danson MJ (2010) Metabolism of pentose sugars in the hyperthermophilic archaea *Sulfolobus solfataricus* and *Sulfolobus acidocaldarius*. *J Biol Chem* **285**: 33701-33709.

1
2
3 Stark H, Wolf J, Albersmeier A, Pham TK, Hofmann JD, Siebers B, Kalinowski J,
4 Wright PC, Neumann-Schaal M & Schomburg D (2017) Oxidative Stickland
5 reactions in an obligate aerobic organism–amino acid catabolism in the Crenarchaeon
6 *Sulfolobus solfataricus*. *FEBS J* **284**: 2078-2095.
7
8

9 Wagner M, Shen L, Albersmeier A, van der Kolk N, Kim S, Cha J, Brasen C,
10 Kalinowski J, Siebers B & Albers SV (2017) *Sulfolobus acidocaldarius* uptakes
11 pentoses via a cut2-type ABC transporter and metabolizes them through the aldolase-
12 independent Weimberg pathway. *Applied and environmental microbiology* **84**: 1273.
13
14

15 Wang L, Liu Q, Wu X, Huang Y, Wise MJ, Liu Z, Wang W, Hu J & Wang C (2019)
16 Bioinformatics analysis of metabolism pathways of archaeal energy reserves. *Sci Rep*
17 **9**: 1034.
18
19

20 Wolf J, Stark H, Fafenrot K, *et al.* (2016) A systems biology approach reveals major
21 metabolic changes in the thermoacidophilic archaeon *Sulfolobus solfataricus* in
22 response to the carbon source L-fucose versus D-glucose. *Mol Microbiol* **102**: 882-
23 908.
24
25

26 Zweerink S, Kallnik V, Ninck S, *et al.* (2017) Activity-based protein profiling as a
27 robust method for enzyme identification and screening in extremophilic Archaea.
28 *Nature Commun* **8**: 15352.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 8-2: Distribution of Oxidase Complexes in the *Sulfolobales*

Organism	SoxABCDD'L	SoxEFGHIM	DoxBCE	SoxLN-CbsAB
<i>Acidianus ambivalens</i>			X	X
<i>Acidianus brierleyi</i>	X		X	X
<i>Metallosphaera cuprina</i>	X	X	X	X
<i>Metallosphaera sedula</i>	X	X	X	X
<i>Saccharolobus islandicus</i>	X	X	X	X
<i>Saccharolobus solfataricus</i>	X	X	X	
<i>Stygiolobus azoricus</i>			X	
<i>Sulfodiicoccus acidophilus</i>		X	X	X
<i>Sulfolobus acidocaldarius</i>	X	X	X	X
<i>Sulfuracidifex metallicus</i>	X		X	X
<i>Sulfurisphaera tokodaii</i>	X	X	X	X

1
2
3 **Table 9.1 Most used knock out systems in *Sulfolobales***
4

5 6 7 8 <i>Sulfolobus acidocaldarius</i>	Markerless deletion mutants via crossover based on <i>pyrEF/5-FOA</i> counterselection (Wagner <i>et al.</i> , 2012)
9 10 11 12 <i>Saccharolobus solfataricus</i>	Gene disruption via permanent insertion of <i>lacS</i> reporter gene via homologous recombination (Albers & Driessen, 2007)
13 14 15 16 17 <i>Saccharolobus islandicus</i>	Markerless deletion mutants via crossover based on <i>pyrEF/5-FOA</i> counterselection improved with <i>argD</i> selection (Zhang <i>et al.</i> , 2013)
18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60	Addition of <i>apt/6-MP</i> counterselection (Zhang <i>et al.</i> , 2016)
	CRISPR-based gene knock out (Li <i>et al.</i> , 2016)

References:

Albers S-V & Driessen AJM (2007) Conditions for gene disruption by homologous recombination of exogenous DNA into the *Sulfolobus solfataricus* genome. *Archaea* **2**: 145-149.

Li Y, Pan S, Zhang Y, Ren M, Feng M, Peng N, Chen L, Liang YX & She Q (2016) Harnessing Type I and Type III CRISPR-Cas systems for genome editing. *Nucleic Acids Res* **44**: e34-e34.

Wagner M, van Wolferen M, Wagner A, Lassak K, Meyer B, Reimann J & Albers S-V (2012) Versatile genetic tool box for the Crenarchaeote *Sulfolobus acidocaldarius*. *Front Microbiol* **3**.

Zhang C, Cooper TE, Krause DJ & Whitaker RJ (2013) Augmenting the genetic toolbox for *Sulfolobus islandicus* with a stringent positive selectable marker for agmatine prototrophy. *Appl Environ Microbiol* **79**: 5539-5549.

Zhang C, She Q, Bi H & Whitaker RJ (2016) The *apt/6-methylpurine* counterselection system and its applications in genetic studies of the hyperthermophilic archaeon *Sulfolobus islandicus*. *Appl Environ Microbiol* **82**: 3070-3081.

Table 9.2 Most used expression vectors in *Sulfolobales*

Organism	Name	Expression vectors				Reference
		Promoter/ Inducer	Selection marker	Feature		
<i>S. acidocaldarius</i>	pCmalLacS	<i>mal</i> maltose	<i>pyrEF</i> , <i>lacS</i> , <i>amp</i>	-	(Berkner <i>et al.</i>)	
	pSVAaraFX	<i>araS</i> arabinose				
	pSVAmalFX	<i>mal</i> maltose	<i>pyrEF</i> , <i>lacS</i> , <i>amp</i>	HA, 6xHis, StrepII, Twin-Strep, 10xHis + StrepII tags at the C and N terminus	(van der Kolk <i>et al.</i>)	
	pSVAxylFX	<i>xylR</i> xylose				
<i>Sa. solfataricus</i>	pSVA	<i>araS</i> arabinose	<i>pyrEF</i> , <i>lacS</i> , <i>amp</i>	6xHis, 10x His, StrepII tags	(Albers <i>et al.</i>)	
<i>Sa. islandicus</i>	pSeSD	<i>ara-SD</i> arabinose	<i>pyrEF</i> , <i>lacS</i> , <i>amp</i>	6xHis tag, 2 protease sites for tag removal	(Peng <i>et al.</i>)	

References:

Albers SV, Jonascheit M, Dinkelaker S, Urich T, Kletzin A, Tampé R, Driessen AJM & Schleper C (2006) Production of recombinant and tagged proteins in the hyperthermophilic archaeon *Sulfolobus solfataricus*. *Appl Environ Microbiol* **72**: 102-111.

Berkner S, Włodkowska A, Albers S-V & Lipps G (2010) Inducible and constitutive promoters for genetic systems in *Sulfolobus acidocaldarius*. *Extremophiles* **14**: 249-259.

Peng N, Deng L, Mei Y, Jiang D, Hu Y, Awayez M, Liang Y & She Q (2012) A synthetic arabinose-inducible promoter confers high levels of recombinant protein expression in hyperthermophilic archaeon *Sulfolobus islandicus*. *Appl Environ Microbiol* **78**: 5630-5637.

van der Kolk N, Wagner A, Wagner M, Waßmer B, Siebers B & Albers S-V (2020) Identification of XylR, the activator of arabinose/xylose inducible regulon in *Sulfolobus acidocaldarius* and its application for homologous protein expression. *Front Microbiol* **11**: 1066.