

1 Local and non-local controls on seasonal variations in water
2 availability and use by riparian trees along a hydroclimatic gradient
3

4 Christopher I. Sargeant¹, Michael Bliss Singer^{*2,3,4}
5

6 ¹School of Earth and Environmental Sciences, University of St Andrews, St Andrews, KY16
7 9AL, UK

8 ²School of Earth and Environmental Sciences, Cardiff University, Main Building, Park Place,
9 Cardiff, CF10 3AT, UK

10 ³Water Research Institute, Cardiff University, The Sir Martin Francis Building, Museum
11 Avenue Cardiff, CF10 3AX, UK

12 ⁴Earth Research Institute, University of California, Santa Barbara, 6832 Ellison Hall, Santa
13 Barbara, CA 93106, USA
14

15 *Corresponding Author: Michael Bliss Singer

16 **Email:** bliss@eri.ucsb.edu
17
18

19 **Author Contributions:** CIS and MBS designed research and wrote the paper. CIS performed
20 research and analyzed data.
21

22 **Competing Interest Statement:** The authors declare no conflicts of interest.
23

24 **Keywords:** Riparian vegetation, Groundwater, Streamflow, Rhône, France
25
26
27

Abstract

As global climate change continues to impact regional water cycles, we may expect further shifts in water availability to forests that create challenges for certain species and biomes. Lowland deciduous riparian forests are particularly vulnerable because tree species cannot migrate out of the stream corridor, and they rely on root zone water availability that is controlled by variations in both local climate conditions (e.g., precipitation, evaporation, and infiltration) and non-local hydroclimatic forcing (e.g., streamflow, snowmelt, recharge). To determine how the seasonal water source usage of riparian trees is controlled by local versus non-local variability in hydroclimatic regime, we reconstructed the seasonal oxygen isotope ($\delta^{18}\text{O}$) signature of water used by two riparian tree species with contrasting rooting depths, comprising ~800 $\delta^{18}\text{O}$ tree-ring cellulose measurements from 12 tree-level decadal time-series at sub-annual resolution (6 samples per year), along a strong hydroclimatic gradient within the Rhône River basin, SE France. These results were evaluated alongside $\delta^{18}\text{O}$ measurements made from potential endmember water sources and independent hydroclimatic metrics to characterize the seasonal evolution of both potential water availability at distinct rooting depths and tree water source use, and to investigate the generalized riparian tree response to seasonal variations in local versus non-local hydroclimatic forcing over a decade. We show: 1) distinct seasonal water use between species, based on differential access to groundwater; 2) substantial source switching in both species based on evolving water availability; and 3) riparian trees are more dependent on locally controlled soil moisture with distance downstream, creating increased vulnerability to locally increasing temperatures. We also find that deeply rooted trees in lowland riparian floodplains are potentially vulnerable to climate change because of their large dependence on water supply from mountains. This effect is more pronounced downstream, where seasonal water table decline may lead to loss of required water for deeply rooted trees.

Introduction

Despite a growing consensus that forests across the globe are becoming increasingly vulnerable to water stress (1-5), insufficient information exists on the availability of specific water sources and the usage of this water by vegetation throughout the growing season across

hydroclimatic gradients. Knowledge of water availability to plants and their use of this water in favorable hydrologic years versus drought periods will improve understanding of the proportional balance of groundwater versus infiltrated precipitation. It will particularly inform us about the evolution of key climatically controlled water sources for plant species rooted at different depths, and the corresponding vegetation responses to hydroclimatic forcing, whether locally or non-locally expressed. This knowledge gap is compounded by a lack of historical (i.e., decadal-multidecadal) baseline data, from which spatiotemporal trends of water source usage can be evaluated to predict and mitigate deleterious forest responses to shifting hydroclimate regimes under global climate change (1, 6, 7). Lowland environments are of great concern because trees are already subjected to high levels of water stress, yet adaptation strategies such as upslope species migration are impossible because riparian trees in Mediterranean climates are adapted to lowlands and exist in isolated patches (8-12). A new predictive understanding of the variation in water source availability and usage in response to local and non-local hydroclimatic forcing could support management interventions that prevent or mitigate large-scale forest diebacks, such as those recently attributed to increasing temperatures, precipitation variability, and root-zone moisture deficits (1, 7, 13-17).

Trees in riparian zones commonly utilize either phreatic (saturated) zone moisture or vadose (unsaturated) zone moisture sources, depending on their rooting architecture (18-21). These water sources are comprised of an unknown mixture of regional groundwater flow, streamflow support of the shallow alluvial aquifer, and locally infiltrated precipitation (22-26). Within seasonally dry climates (e.g., Mediterranean biomes), the cycles of the hydroclimate and vegetative growth may be out of phase, such that root zone soil moisture becomes limited during the summer period, when evaporative demand is highest and floodplain water tables are depressed due to declining streamflow. Small shifts in vegetative water stress resulting from changes in water availability could thus have important implications for tree growth and health, potentially driving shifts in forest community composition and function, ecosystem services, and carbon sequestration (27-31). Climatic changes have already begun to significantly influence the water cycle in many seasonally dry regions, affecting both the input/output fluxes of water and the storage of water in soils, streams, and groundwater aquifers (23, 32-34). Source water

availability within riparian zones is also potentially affected by anthropogenic activities (e.g., dams, river training, groundwater abstraction), which are particularly common in Mediterranean systems, disrupting natural hydrological regimes and thereby impact water storage at different rooting depths (22, 35-39). The increasing likelihood of water cycle shifts in Mediterranean climate systems suggests the need for more detailed investigation of systematic patterns and timing of source water usage across known gradients of floodplain water availability.

One can conceive of water availability to vegetation in large river floodplains as a balance between (Figure 1): 1) local climate, where shallow soil moisture is driven by the stochastic difference between precipitation inputs and evapotranspiration losses (40, 41), and 2) non-local climate, where water table elevation and corresponding capillary rise is driven by regional groundwater recharge and streamflow regimes that support the shallow phreatic aquifer (42, 43). We refer to these as local and non-local hydroclimatic controls on water availability, respectively, noting that there may be marked differences in the magnitude, frequency, and timing of changes to the local and non-local water cycle that create differences in riparian zone water availability at any downstream location. Climate projections in Mediterranean-type ecosystems indicate that temperature and precipitation regimes will become progressively warmer and drier with less mountain snowpack (9, 44, 45). These trends have the potential to alter the spatiotemporal availability of water within floodplains at different points along a river network, and thus to the overlying forest ecosystem (11, 46). Nevertheless, our understanding of how hydroclimate is expressed in terms of seasonal tree water source availability is limited, which restricts our ability to adequately manage for the future consequences of climate change to Mediterranean riparian forests.

Isotopic ratios of oxygen ($\delta^{18}\text{O}$) extracted from xylem samples, annual growth rings, and potential endmember waters have been particularly helpful in identifying and disentangling contributions from different source waters, where deeper water is usually associated with lighter isotopic values and soil water that has undergone evaporative losses becomes isotopically heavier (23-25, 47). They have enabled improved understanding of the evolution of seasonally available water sources at critical periods of growth, and to identify the dominant water source

usage over an entire season of growth (48, 49). Recent developments in the study of $\delta^{18}\text{O}$ from tree ring cellulose have enabled the analysis of water source variability at sub-annual resolution in the same reference frame as the potential contributing endmember sources for co-occurring riparian tree species that typically use different water sources (48, 50). This sub-annual information about water source usage is critical for predicting how riparian forests may respond to climatic changes that affect either local or non-local controls on water availability. Where there are substantial isotopic differences between river/phreatic water and precipitation/vadose zone water, $\delta^{18}\text{O}$ signatures support a relative comparison of source water uptake by riparian trees from shallow versus deeper floodplain water reservoirs, which represent local versus non-local hydroclimatic controls, respectively.

This paper aims to elucidate the dynamic behavior of hydroclimatic controls on subsurface hydrology through analysis of sub-annual patterns of tree water source use. We use seasonally resolved dendro-isotopic analyses of $\delta^{18}\text{O}$ extracted from tree ring cellulose, representing signatures of source waters (Supplementary Material), to generalize local and non-local hydroclimatic expressions of floodplain water availability. We conduct this study within three riparian forest sites spanning a strong hydroclimatic gradient along the Rhône River in SE France. These sites span a range of 3.5°C and 140 mm in growing season temperature and precipitation, respectively, over the period 2000-2010 from Peage de Rousillon - 'Upstream' to Donzere-Mondragon - 'Midstream' to Mas Thibert - 'Downstream' (Figure 1b). Along this hydroclimatic gradient, historical temperatures have already increased >2°C between 1970 and 2015, yet precipitation trends are equivocal (Figure S1). Each site contains common, and co-occurring Mediterranean tree species, *Fraxinus excelsior* and *Populus nigra/alba* (Figure 2a). *Fraxinus* and *Populus* have been shown to exhibit contrasting rooting depths, where the former is restricted to soils above the gravel layer (~ less than 1 m deep) and the latter has most of its root mass within gravels deeper than 1 m (31, 49). The investigation utilizes an 11-year dataset of sub-annual in tree-ring cellulose ($\delta^{18}\text{O}_{\text{cell}}$) (n = 792) from individual, co-located *Fraxinus*-*Populus* pairs from both near channel (NC) and interior floodplain (IF) locations within each study plot along the Rhône hydroclimatic gradient (Figure 2a). We then employ an inverse biomechanistic model (50) to determine the $\delta^{18}\text{O}$ of source water ($\delta^{18}\text{O}_{\text{msw}}$) utilized by each tree

during the formation of each respective sub-annual $\delta^{18}O_{cell}$ sample (Supplementary Material, Figure S2).

Our broader goal was to investigate how potential water variability along the hydroclimatic gradient affects water source usage by riparian trees. To accomplish this, we statistically compared distributions of $\delta^{18}O$ via Student's t and Mann-Whitney tests to identify site differences (and similarities) in potential water sources to riparian trees rooted at different positions. We then compared distributions of $\delta^{18}O_{msw}$ between tree species and along the hydroclimatic gradient via the Kolmogorov-Smirnov test, and we interpreted the median values in the context of $\delta^{18}O$ for these potential water sources - separately analyzed river, phreatic, vadose, non-growing season (NGS) precipitation, growing season (GS) precipitation (Supplementary Material). We then use seasonal $\delta^{18}O_{msw}$ data for all growth years to develop characteristic patterns of seasonal water use for each tree and species along the gradient.

Finally, we evaluate these characteristic seasonal $\delta^{18}O_{msw}$ time series for each site against stream stage records (representing non-local controls on water table elevation) and a drought index (representing local controls on soil moisture availability as a function of the cumulative water balance over an antecedent period). Specifically, we assessed non-local controls on water availability in the shallow alluvial aquifer based on characteristic seasonal variations in upper percentiles of local stream stage within the Rhône River at each site, which integrates water production over the entire upstream drainage area. And we quantified water availability in the vadose zone based on the Standardized Precipitation and Evaporation Index-SPEI (51), a drought index that reflects the local balance over previous periods between input precipitation and evapotranspiration that affect soil moisture. SPEI is calculated based on potential evapotranspiration from atmospheric variables rather than actual evapotranspiration. Therefore, a positive value of SPEI over the growing season suggests that soil moisture should be available to plants at some point in the growing season based on the time integral of rainfall minus PET over the antecedent months. Thus, we connect the characteristic seasonal plant water usage for these differentially rooted tree species along the hydroclimatic gradient to the

local versus non-local controls on water availability in the rooting zones of the floodplain at different positions within the river basin.

Results

Modeled Source Water Use from Dendro-Isotopes and Potential Endmember Sources

Generally, we find systematically higher values of $\delta^{18}O_{msw}$ with downstream distance along the climate gradient (ensemble medians of -11‰, -9‰, -7‰), consistent with the downstream warming climate (Figures 2, 3a-c, Figure S1, Table S1). However, there is substantial variability in $\delta^{18}O_{msw}$ between species, rooting locations, and sites (Figure S3a, Table S1). For example, at NC locations, *Fraxinus* $\delta^{18}O_{msw}$ values are on average isotopically heavier than those of *Populus*, consistent with shallower water sources (Figure 3, Table S1). We also find that some individual trees use a large range of water sources, evidenced by a broad distribution with a muted peak (e.g., *IF Populus* at Midstream, Figure 3b). Furthermore, there was notable within-site statistical similarity in $\delta^{18}O_{msw}$ distributions between *Populus* trees at both within-site rooting positions for both Upstream and Downstream sites, as well as between *Fraxinus* trees at Midstream (Figure 3a-c).

In terms of endmember water sources, the $\delta^{18}O$ values in river samples along the climate gradient were statistically similar (mean value of -10.5‰, Figure 3d-f; Figure S3), suggesting a well-mixed pool of non-locally derived water in the Rhône. In contrast, $\delta^{18}O$ in locally controlled GS and NGS precipitation (ppt) vary along the hydroclimatic gradient, becoming isotopically heavier from north to south, e.g., -5‰ to -3‰ for GS ppt (and relatively heavier in the growing season). This downstream heavier trend is partially mirrored in the vadose zone samples (rising from -10‰ to -8‰, north to south), though the differences are not statistically significant. Additionally, we find that the vadose zone moisture at all sites appears to derive from the lower half of NGS ppt inputs, associated with winter precipitation under colder conditions (Figure 3d-f). At Upstream and Downstream sites, there is statistical similarity between the average values for river water and phreatic water (Figure S3b), reflecting a tight connection between the Rhône River and its alluvial floodplain aquifer, compared with Midstream, where an isotopically heavier groundwater source from the Massif Central contributes to the alluvial aquifer (48). *Populus*

trees at each site and at all but one rooting location (Upstream *IF*, where both species exhibit isotopically light $\delta^{18}O_{msw}$) tend to use isotopically lighter water than *Fraxinus* trees (Figure 3, Table S1). Generally, the lighter value of $\delta^{18}O_{msw}$ used by *Populus* trees corresponds more closely to non-locally controlled phreatic water sourced from the Rhône alluvial aquifer than it does with locally controlled vadose zone water, yet the substantial variability over the decadal time period indicates hydroclimatically controlled seasonal variations in water sources for both *Populus* and *Fraxinus* trees (Figure 3, Table S2).

Seasonal Patterns of Water Use

The subannual $\delta^{18}O_{msw}$ values averaged over all years (Figure 4a-l) indicate the seasonal evolution of source water $\delta^{18}O$ utilized by each tree at each rooting position and site along the hydroclimatic gradient, based on more than a decade of sub-annual samples. There are notable patterns that emerge. For example, *Populus* trees at all three sites and rooting positions begin their seasonal growth using relatively isotopically heavy water, after which their water use becomes progressively lighter (non-locally controlled) until the last stages of latewood development, when isotopic enrichment toward heavier, locally controlled $\delta^{18}O_{msw}$ is evident (Figures 3,4a-c). This suggests that *Populus* trees undergo seasonal water source switching from shallow (locally controlled) to deep (non-locally controlled) and back to shallow rooting depths (Fig3d-f). In contrast, *Fraxinus* trees do not exhibit systematic patterns across rooting positions and sites, although there appears to be consistent seasonal water usage between years for *Fraxinus* within each site. For example, *Fraxinus* at Upstream begin seasonal growth using the lightest $\delta^{18}O_{msw}$ (reflecting locally controlled NGS ppt, Figure 3d), which then rises and plateaus during earlywood formation (reflective of local evaporative enrichment), before rising again such that the isotopically heaviest water is used at the end of the growing season (LW3, Figure 4a,c).

We recognize one clear consistency across all trees at all sites: a dip in $\delta^{18}O_{msw}$ for initial latewood growth (LG1 or LG2) followed by an increase in $\delta^{18}O_{msw}$ at the end of the growing season for all sites (LG3). This result suggests systematic source switching during latewood growth to a relatively shallow water source that has undergone significant evaporative

enrichment (heavier isotopic value). This pattern is most dramatically observable for the *NC Populus* at Downstream (Figure 4j). The spread of annual $\delta^{18}\text{O}_{msw}$ values is largest at Upstream, and lowest for Downstream (Figures 3, S3). This result suggests more interannual consistency in water use with distance downstream along the hydroclimatic gradient, where trees are more constrained in their use of variable moisture pools.

Climatic Controls on Seasonal Water Availability

The variations in $\delta^{18}\text{O}_{msw}$ between tree species within and between sites along the hydroclimatic gradient in SE France support further analysis of the local versus non-local hydroclimatic controls on water availability (e.g., via river stage and SPEI), which affect which water sources are used at certain periods of the growing season. The results of these analyses are presented in Figure 5, along with the mean seasonal patterns of $\delta^{18}\text{O}_{msw}$. The Rhône River stage data show an evolving pattern with distance downstream. Due to contributions of snowmelt runoff from the Alps, stage (h) is generally highest at the onset of earlywood growth (EW1) at all sites (Figure 5d-f), and then declines more with distance downstream (0.3 - 0.7 m for h_{95} , from north to south, for example), before slightly rebounding at the end of latewood growth (LW3). The main differences in h percentiles between sites is when the low point occurs and to what level it rebounds. For example, the trough in h at Upstream occurs during earlywood production, while it occurs during the latewood growth period for Midstream and Downstream. This is consistent with the expansion of drainage area with distance downstream that delays the annual flood wave in the Rhône (52), particularly that which is derived from French Alps snowmelt runoff (Downstream lies downstream of the Rhône's confluence with the Alps-draining Durance River). Additionally, the seasonal variation in h increases with distance downstream, reaching a range of >0.7 m for all three h percentiles at Downstream. High values of h relative to the average generally drive the filling of the shallow alluvial aquifer under the floodplain, which drains more slowly than it fills, leaving behind high non-locally controlled water availability for deeper rooting species (53). In general, the stage values at Upstream and Midstream are ~5-6 m below the surface elevation at tree rooting positions, although h is only 1-2 m below the surface at Downstream (Figure 5d-f), suggesting greater access of aquifer water to trees at this site.

267

268 The SPEI values at each site show a similar pattern for 1 to 4 months of antecedent water
269 balance (Figure 5g-i). For example, the 1-month SPEI rises to its peak at the end of earlywood
270 formation (EW3), and then essentially flatlines for the rest of the growing season, whereas the
271 2-month SPEI peaks at LW1, the 3-month index peaks at LW2, and the 4-month SPEI peaks
272 at LW3. This pattern suggests that the accumulated soil moisture throughout the growing
273 season is most available to trees in the final stages of both earlywood and latewood growth.
274 The SPEI values with longer antecedence reflect soil moisture surplus derived from locally
275 controlled NGS precipitation (which will be exhausted earlier), while the 1-month antecedence
276 SPEI, for example, mainly reflects contributions from GS precipitation (also locally controlled).

277

278 ***Hydroclimatic Patterns in Seasonal Water Use***

279 In the context of seasonal changes in h and SPEI, the characteristic patterns of seasonal water
280 use for each tree indicate differences in the availability of deep versus shallow moisture sources
281 throughout the growing season and corresponding source switching by both species to maintain
282 growth throughout the season. We summarize the water availability results by these metrics
283 separately for the initiation of growth, the middle of the growing season, and the end of growth.

284

285 At growth initiation (EW1) Rhône stream stage (h) is highest for all sites, yet this does not
286 necessarily correspond to the use of non-locally controlled phreatic water in early stages of
287 growth because Mediterranean trees will often only access this water source once the water
288 table declines, leaving behind isotopically light capillary water in deeper soil layers (54, 55).
289 Thus, the starting values of $\delta^{18}O_{msw}$ values vary dramatically between sites (Figure 5a-c),
290 depending instead on the isotopic signature of readily available water in shallow soil layers from
291 locally controlled NGS ppt, the volume of which is apparently higher upstream and declines
292 downstream (Figure 5g-i). The $>5\text{‰}$ range in starting $\delta^{18}O_{msw}$ values at Upstream suggest
293 access from deep versus shallow root for *Populus* and *Fraxinus* trees, respectively, whereas
294 all trees at the Midstream and the Downstream sites appear to use similar water to begin growth
295 (except for the NC *Fraxinus*, Figure 5a-c).

296

In the middle stages of growth, characteristic use of isotopically light (relatively) water for all trees suggests availability and use of water from deeper rooting depths. Lighter $\delta^{18}\text{O}_{msw}$ in mid-growth are associated with non-locally controlled phreatic or capillary water supported by the passage of Rhône River snowmelt runoff wave and/or the accumulation of deeper soil moisture by locally controlled NGS precipitation several months prior to growth (Figures 2d-f,4). The trough of $\delta^{18}\text{O}_{msw}$ occurs around LW1, and here there are similarities in water use signatures for all trees at all sites, except the *NC Fraxinus* at Midstream and both *Fraxinus* trees at Downstream (Figure 5a-c), which are apparently rooted more shallowly such that they access a different (more enriched) moisture pool that reflects lower antecedent moisture (56) (Figure 5g-i).

In the final stage of growth, there is consistent isotopic enrichment in locally controlled soil moisture, which is evident in the $\delta^{18}\text{O}_{msw}$ data for each of the Rhône sites, particularly for the more shallowly rooted *Fraxinus* (Figure 5a-c). In fact, the seasonal patterns of water use suggest that both riparian tree species are adapted to take advantage of soil moisture at the end of the growing season, even if the isotopic value of this enriched moisture pool varies substantially year to year (Figure S4). At this stage, water may become very limited in both deep and shallow soil layers, such that trees are using decreasing pools of locally controlled soil moisture that may be briefly replenished by summer rains (GS ppt), which are subjected to high evaporation losses, producing high variability in end-of-season $\delta^{18}\text{O}_{msw}$. It is also clearest from the LW3 $\delta^{18}\text{O}_{msw}$ values that the hydroclimate gradient exerts a strong control on source water availability and isotopic enrichment.

Discussion

Seasonal water source use variations throughout the growing period are an expression of adaptive plant responses to annual variations in water availability in phreatic or vadose zone stores. Thus, characteristic patterns of seasonal water usage may develop under a stationary climate, where climatic expression within hydrology regionally (non-locally) and locally are predictable. Clearly, there will be favorable and unfavorable years in terms of water availability in these phreatic versus vadose zone stores, but water source switching between shallow and

deep roots would support healthy growth if both hydrologic stores are not negatively affected at the same time (e.g., a regional drought affecting upland snowpack and lowland rainfall). The analysis presented here shows a seasonal cycle of water availability and use by two co-located tree species with contrasting rooting depths that reflects the influence of: 1) non-local climatic controls on phreatic zone (and deeper soil layers) water availability and use in the middle of the growing season, especially by the more deeply rooted *Populus*; 2) accumulation of soil moisture from locally controlled NGS precipitation primarily benefitting *Fraxinus*; 3) contributions from locally controlled GS precipitation to shallow zone water availability; and 4) locally high temperatures resulting in soil drying within shallow layers at the end of the growing season (Figures 2,4). The local and non-local hydroclimatic controls on riparian water availability and tree water use are summarized schematically in Figure 1. This figure shows that non-local controls on water availability are dominated by snowmelt from the mountains and their contribution to streamflow selectively affects deeply rooting species, whereas the local balance between precipitation and evapotranspiration affects water availability to shallowly rooting species in riparian forests.

The climate is becoming warmer in SE France (a $\sim 2^{\circ}\text{C}$ increase in GS temperature between 1970 and 2015, Figure S1) and in many seasonally dry environments around the world, where riparian trees are adapted to varying water availability by switching water sources between deeper and shallower rooting zones to support their growth and strong hydroclimatic gradients (23, 48, 57, 58). Under nonstationary climatic forcing, the local and non-local water balance may become perturbed. Upland precipitation may shift in phase from snow to rain, leading to an increase in winter flooding but also to a smaller spring snowmelt pulse. Additionally, warmer upland spring temperatures may cause earlier/faster melting of remaining snowpack, shifting the timing/magnitude of the snowmelt flood pulse (Figure 5d-f). These non-local effects may be exacerbated at riparian floodplain sites by locally controlled warmer lowland summer temperatures in the absence of precipitation changes (Figure S1), which may result in higher soil evaporation, leading to faster drying, and/or to rapid exhaustion of soil moisture at deeper depths in the profile. Furthermore, a non-local shift to more regional rainfall and less snowfall would support to a higher contribution of locally controlled NGS precipitation to summer soil

moisture (41-43). All these changes would affect the availability of moisture in both phreatic and vadose zone storage at key periods of tree growth (42, 43), but it is challenging to anticipate where and by how much.

The increasing vulnerability of riparian forests to drought stress suggests there is a necessity to improve the understanding of seasonal water source availability to trees. For example, systematic differences in co-located trees of different species suggest different water use strategies that may play out over seasonal timescales. Considering the Rhône hydroclimatic gradient, we show an increasing influence of (locally controlled) vadose zone water availability with distance downstream towards warmer climates, especially for *Fraxinus*. This suggests that within warmer, drier environments in the southern Rhône basin, source water availability to trees is responsive to local drought conditions, or even individual very dry seasons, leading to strong vulnerabilities in *Fraxinus* because their rooting architecture gives them fewer options for source switching. In contrast, *Populus*, a phreatophyte with a deeper rooting architecture, is less susceptible to negative effects of soil moisture drying, especially where the water table is quite shallow (at Midstream, Figure 5f). Downstream experiences the highest availability of Rhône-sourced phreatic water of any site, supporting uptake and growth of deeply rooted trees especially over the latewood period. However, the high spring values of h at all three sites likely support an elevated water table and/or capillary water in deeper soil layers after water table recession, potentially leading to high water availability of isotopically light water to deep roots during latewood growth (Figure 5a-c).

Along the entire hydroclimatic gradient, *Populus* trees consistently use shallow phreatic water to support mid-season growth (Figure 5a-c), suggesting that their key water source is non-locally controlled. This means that *Populus* and *Fraxinus* trees along the Rhône are responsive and vulnerable to distinct aspects of the hydroclimatic system (non-local controls on the shallow alluvial aquifer for *Populus* and local controls on soil moisture for *Fraxinus*, Figure 1). We suspect this disconnect in controls on water availability to co-located riparian trees with distinct rooting depths is a common feature of many lowland river floodplains. Under stationary climatic conditions there should be (on average) sufficient floodplain water for both species, and each

could manage a season of water limitation arising from an unfavorable hydrologic year via below-normal growth (31). The progression of the growing season also brings a progressive isotopic enrichment due to evaporation and higher temperatures during precipitation events and increased evaporation, leading to higher values of $\delta^{18}\text{O}$ in soil moisture at LW3 than in EW3. In general, deeper water sources may be exhausted by August due to water table decline and depletion of hyporheic bank storage (48, 53), especially for the drier (most downstream sites), requiring trees of both species to access this enriched soil moisture to complete their annual rings at the end of the growing season. However, given that riparian trees are transpiring throughout the growing season (59), the true soil moisture availability will generally decline throughout the season, especially because GS precipitation is generally low in Mediterranean environments. This poses risks to both species in years of unfavorable streamflow, when they will compete for a dramatically limited water source.

A nonstationary climate, on the other hand, may pose species-specific challenges, depending on whether it affects the water balance locally or non-locally. For example, if climatic changes affect upland snowpack/snowmelt, it is likely to limit phreatic zone water availability (non-locally controlled), especially at locations most distant from the mountains, such that phreatophyte species (e.g., *Populus*) may lose their buffer against locally warmer and drier conditions, and therefore become highly sensitive to soil moisture deficits (60). In contrast, if local growing season temperatures increase (which is already happening along the entire Rhône, Figure S1) and/or local precipitation declines (more equivocal), soil moisture deficits will selectively impact shallowly rooted species (e.g., *Fraxinus*). It is also possible for both climatic changes (local and non-local) to occur simultaneously, negatively affecting both phreatic zone and vadose water availability (drought conditions) in lowland riparian forests. If this should occur, the inter-species competition for water will favor the survival of the more drought-tolerant species, *Fraxinus* (56), which has made more investment in shallow roots, and it would have important consequences for riparian forest composition along this and other river systems.

Since water deficits have been identified as the major drivers of forest die-back (1), alterations to forest species composition (e.g., hydroclimatic changes may favor the expansion of a

particular species) may reduce biodiversity and ecosystem services. We have shown here that climate change may affect locally versus non-locally controlled water availability, leading to correspondingly different responses in *Fraxinus* and *Populus* trees along the Rhône hydroclimatic gradient. Ecologically, these spatially restricted lowland riparian forests have no adaptive escape hatch to ameliorate the effect of such changes (8-12). These forest patches are ecologically isolated with few opportunities to expand their range or even recruit on new floodplain surfaces, given their adaptation to lowlands and anthropogenic pressures in lowland floodplains (61-63). Thus, lowland riparian forests, especially in Mediterranean climate regimes are 'sitting ducks' for climate change, due to their dependence on locally and non-locally controlled root zone water availability (Figure 1) that will undergo changes with atmospheric warming and alterations to precipitation regimes. We suggest that characterization of seasonal water source availability and tree water use enables identification of water supply issues to riparian forests, and therefore allows for potential intervention to climate change at regional (water management) and local (forest management) scales.

Acknowledgments

Support for this work came from the U.S. National Science Foundation (BCS-1660490, EAR-1700555), the U.S. Department of Defense's Strategic Environmental Research Development Program (RC18-1006). It was also partly funded by Labex DRIIHM (ANR-11-LABX-0010) as part of the Observatoire Hommes-Milieux Vallée du Rhône (OHM VR) and performed within the framework of the EUR H2O'Lyon (ANR-17-EURE-0018) of Université de Lyon (UdL) through the "Investissements d'Avenir" program operated by the French National Research Agency (ANR). The lab work was done based on an in-kind Natural Environmental Research Council Facility Grant via the British Geological Society's Stable Isotope Facility (Melanie Leng). We also thank various collaborators in France who assisted with access and equipment for field work. These include: Hervé Piégay, Bernard Pont, Jérémie Riquier, and Kristell Michel.

References


1. C. D. Allen *et al.*, A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *For. Ecol. Manage.* **259**, 660-684 (2010).
2. C. D. Allen, D. D. Breshears, N. G. McDowell, On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere* **6**, art129 (2015).
3. N. G. McDowell, C. D. Allen, Darcy's law predicts widespread forest mortality under climate warming. *Nature Clim. Change* **5**, 669-672 (2015).
4. H. D. Adams *et al.*, A multi-species synthesis of physiological mechanisms in drought-induced tree mortality. *Nature Ecology & Evolution* 10.1038/s41559-017-0248-x (2017).
5. J. S. Clark *et al.*, The impacts of increasing drought on forest dynamics, structure, and biodiversity in the United States. *Global Change Biol.* **22**, 2329-2352 (2016).
6. D. D. Breshears *et al.*, Tree die-off in response to global change-type drought: mortality insights from a decade of plant water potential measurements. *Front. Ecol. Environ.* **7**, 185-189 (2009).
7. D. D. Breshears *et al.*, Regional vegetation die-off in response to global-change-type drought. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 15144-15148 (2005).
8. F. Giorgi, Climate change hot-spots. *Geophys. Res. Lett.* **33**, 4 (2006).
9. F. Giorgi, P. Lionello, Climate change projections for the Mediterranean region. *Global Planet. Change* **63**, 90-104 (2008).
10. J. C. Stella, J. Riddle, H. Piégay, M. Gagnage, M.-L. Trémélo, Climate and local geomorphic interactions drive patterns of riparian forest decline along a Mediterranean Basin river. *Geomorphology* **202**, 101-114 (2013).
11. S. J. Capon *et al.*, Riparian Ecosystems in the 21st Century: Hotspots for Climate Change Adaptation? *Ecosystems* **16**, 359-381 (2013).

- 472 12. S. R. Loarie *et al.*, The velocity of climate change. *Nature* **462**, 1052-1055
473 (2009).
- 474 13. G. P. Asner *et al.*, Progressive forest canopy water loss during the 2012–
475 2015 California drought. *Proceedings of the National Academy of Sciences*
476 **113**, E249-E255 (2016).
- 477 14. W. R. L. Anderegg, L. D. L. Anderegg, K. L. Kerr, A. T. Trugman, Widespread
478 drought-induced tree mortality at dry range edges indicates that climate
479 stress exceeds species' compensating mechanisms. *Global Change Biol.* **25**,
480 3793-3802 (2019).
- 481 15. W. R. Anderegg, J. M. Kane, L. D. Anderegg, Consequences of widespread
482 tree mortality triggered by drought and temperature stress. *Nature*
483 *Climate Change* **3**, 30-36 (2013).
- 484 16. A. Park Williams *et al.*, Temperature as a potent driver of regional forest
485 drought stress and tree mortality. *Nature Climate Change* **3**, 292-297
486 (2013).
- 487 17. P. J. van Mantgem *et al.*, Widespread Increase of Tree Mortality Rates in
488 the Western United States. *Science* **323**, 521-524 (2009).
- 489 18. R. B. Jackson *et al.*, A global analysis of root distributions for terrestrial
490 biomes. *Oecologia* **108**, 389-411 (1996).
- 491 19. S. Rood, S. Bigelow, A. Hall, Root architecture of riparian trees: river cut-
492 banks provide natural hydraulic excavation, revealing that cottonwoods
493 are facultative phreatophytes. *Trees* **25**, 907-917 (2011).
- 494 20. Y. Yang, R. J. Donohue, T. R. McVicar, Global estimation of effective plant
495 rooting depth: Implications for hydrological modeling. *Water Resour. Res.*
496 **52**, 8260-8276 (2016).
- 497 21. Y. Fan, G. Miguez-Macho, E. G. Jobbágy, R. B. Jackson, C. Otero-Casal,
498 Hydrologic regulation of plant rooting depth. *Proceedings of the National*
499 *Academy of Sciences* **114**, 10572-10577 (2017).
- 500 22. C. A. Williams, D. J. Cooper, Mechanisms of riparian cottonwood decline
501 along regulated rivers. *Ecosystems* **8**, 382-395 (2005).
- 502 23. M. B. Singer *et al.*, Floodplain ecohydrology: Climatic, anthropogenic, and
503 local physical controls on partitioning of water sources to riparian trees.
504 *Water Resour. Res.* **50**, 4490-4513 (2014).
- 505 24. T. E. Dawson, J. R. Ehleringer, Streamside trees that do not use stream
506 water. *Nature* **350**, 335-337 (1991).
- 507 25. J. Renee Brooks, H. R. Barnard, R. Coulombe, J. J. McDonnell, Ecohydrologic
508 separation of water between trees and streams in a Mediterranean
509 climate. *Nature Geosci* **3**, 100-104 (2010).
- 510 26. G. Skiadaresis, J. Schwarz, K. Stahl, J. Bauhus, Groundwater extraction
511 reduces tree vitality, growth and xylem hydraulic capacity in *Quercus*
512 *robur* during and after drought events. *Scientific Reports* **11**, 5149 (2021).
- 513 27. N. Bréda, R. Huc, A. Granier, E. Dreyer, Temperate forest trees and stands
514 under severe drought: a review of ecophysiological responses, adaptation
515 processes and long-term consequences. *Ann. For. Sci.* **63**, 625-644 (2006).
- 516 28. S. J. Lite, J. C. Stromberg, Surface water and ground-water thresholds for
517 maintaining *Populus-Salix* forests, San Pedro River, Arizona. *Biol. Conserv.*
518 **125**, 153-167 (2005).
- 519 29. J. C. Stromberg *et al.*, Altered stream-flow regimes and invasive plant
520 species: the *Tamarix* case. *Global Ecol. Biogeogr.* **16**, 381-393 (2007).

- 521 30. D. Sarris, D. Christodoulakis, C. Körner, Impact of recent climatic change
522 on growth of low elevation eastern Mediterranean forest trees. *Clim.*
523 *Change* **106**, 203-223 (2011).
- 524 31. M. B. Singer *et al.*, Contrasting water-uptake and growth responses to
525 drought in co-occurring riparian tree species. *Ecohydrology* **6**, 402-412
526 (2013).
- 527 32. M. B. Singer, K. Michaelides, Deciphering the expression of climate change
528 within the Lower Colorado River basin by stochastic simulation of
529 convective rainfall. *Environmental Research Letters* **12** (2017).
- 530 33. D. Petra, Vulnerability to the impact of climate change on renewable
531 groundwater resources: a global-scale assessment. *Environmental*
532 *Research Letters* **4**, 035006 (2009).
- 533 34. J. R. Gremer, J. B. Bradford, S. M. Munson, M. C. Duniway, Desert grassland
534 responses to climate and soil moisture suggest divergent vulnerabilities
535 across the southwestern United States. *Global Change Biol.* **21**, 4049-4062
536 (2015).
- 537 35. C. Nilsson, K. Berggren, Alterations of Riparian Ecosystems Caused by
538 River Regulation. *Bioscience* **50**, 783-792 (2000).
- 539 36. M. B. Singer, The influence of major dams on hydrology through the
540 drainage network of the Sacramento Valley, California. *River Res. Appl.* **23**,
541 55-72 (2007).
- 542 37. M. B. Singer, R. Aalto, Floodplain development in an engineered setting
543 *Earth Surface Processes and Landforms* **34**, 291-304 (2009).
- 544 38. M. B. Singer, T. Dunne, An empirical-stochastic, event-based model for
545 simulating inflow from a tributary network: Theoretical framework and
546 application to the Sacramento River basin, California. *Water Resour. Res.*
547 **40**, W07506, doi:07510.01029/02003WR002725 (2004).
- 548 39. M. C. Rains, J. E. Mount, E. W. Larsen, Simulated changes in shallow
549 groundwater and vegetation distributions under different reservoir
550 operations scenarios. *Ecol. Appl.* **14**, 192-207 (2004).
- 551 40. I. Rodriguez-Iturbe, A. Porporato, F. Laio, L. Ridolfi, Plants in water-
552 controlled ecosystems: active role in hydrologic processes and response
553 to water stress: I. Scope and general outline. *Advances in Water Resources*
554 **24**, 695-705 (2001).
- 555 41. A. Porporato, P. D'Odorico, F. Laio, L. Ridolfi, I. Rodriguez-Iturbe,
556 Ecohydrology of water-controlled ecosystems. *Advances in Water*
557 *Resources* **25**, 1335-1348 (2002).
- 558 42. F. Laio, S. Tamea, L. Ridolfi, P. D'Odorico, I. Rodriguez-Iturbe,
559 Ecohydrology of groundwater-dependent ecosystems: 1. Stochastic water
560 table dynamics. *Water Resour. Res.* **45**, n/a-n/a (2009).
- 561 43. S. Tamea, F. Laio, L. Ridolfi, P. D'Odorico, I. Rodriguez-Iturbe,
562 Ecohydrology of groundwater-dependent ecosystems: 2. Stochastic soil
563 moisture dynamics. *Water Resour. Res.* **45**, n/a-n/a (2009).
- 564 44. J. I. López-Moreno *et al.*, Different sensitivities of snowpacks to warming
565 in Mediterranean climate mountain areas. *Environmental Research Letters*
566 **12**, 074006 (2017).
- 567 45. S. D. Polade, A. Gershunov, D. R. Cayan, M. D. Dettinger, D. W. Pierce,
568 Precipitation in a warming world: Assessing projected hydro-climate

- changes in California and other Mediterranean climate regions. *Scientific Reports* **7**, 10783 (2017).
46. S. B. Rood *et al.*, Declining summer flows of Rocky Mountain rivers: Changing seasonal hydrology and probable impacts on floodplain forests. *Journal of Hydrology* **349**, 397-410 (2008).
 47. C. Gazis, X. Feng, A stable isotope study of soil water: evidence for mixing and preferential flow paths. *Geoderma* **119**, 97-111 (2004).
 48. C. I. Sargeant, M. B. Singer, Sub-annual variability in historical water source use by Mediterranean riparian trees. *Ecohydrology* **9**, 1328-1345 (2016).
 49. J. M. Sánchez-Pérez, E. Lucot, T. Bariac, M. Trémoières, Water uptake by trees in a riparian hardwood forest (Rhine floodplain, France). *Hydrological Processes* **22**, 366-375 (2008).
 50. C. I. Sargeant, M. B. Singer, C. Vallet-Coulomb, Identification of Source-Water Oxygen Isotopes in Trees Toolkit (ISO-Tool) for Deciphering Historical Water Use by Forest Trees. *Water Resour. Res.* **55**, 10954-10975 (2019).
 51. S. M. Vicente-Serrano, S. Beguería, J. I. López-Moreno, A Multiscalar Drought Index Sensitive to Global Warming: The Standardized Precipitation Evapotranspiration Index. *J. Clim.* **23**, 1696-1718 (2010).
 52. J. M. Olivier *et al.*, "The Rhône River Basin" in *The Rivers of Europe*. (Academic Press, 2009), chap. 7, pp. 247-295.
 53. C. M. Evans, D. G. Dritschel, M. B. Singer, Modeling Subsurface Hydrology in Floodplains. *Water Resour. Res.* **54**, 1428-1459 (2018).
 54. C. Evans, M. B. Singer, D. G. Dritschel, A subsurface hydrology model that traces oxygen isotopes. *Eos Transactions AGU* **97**, EP52B-03. (2016).
 55. D. E. Busch, N. L. Ingraham, S. D. Smith, Water Uptake in Woody Riparian Phreatophytes of the Southwestern United States: A Stable Isotope Study. *Ecol. Appl.* **2**, 450-459 (1992).
 56. G. Marigo, J. P. Peltier, J. Girel, G. Pautou, Success in the demographic expansion of *Fraxinus excelsior* L. *Trees-Struct. Funct.* **15**, 1-13 (2000).
 57. L. B. Flanagan, T. E. Orchard, T. N. Tremel, S. B. Rood, Using stable isotopes to quantify water sources for trees and shrubs in a riparian cottonwood ecosystem in flood and drought years. *Hydrological Processes* **33**, 3070-3083 (2019).
 58. K. A. Snyder, D. G. Williams, Water sources used by riparian trees varies among stream types on the San Pedro River, Arizona. *Agricultural and Forest Meteorology* **105**, 227-240 (2000).
 59. M. Mayes *et al.*, Climate sensitivity of water use by riparian woodlands at landscape scales. *Hydrological Processes* **34**, 4884-4903 (2020).
 60. N. M. Amlin, S. B. Rood, Drought stress and recovery of riparian cottonwoods due to water table alteration along Willow Creek, Alberta. *Trees-Struct. Funct.* **17**, 351-358 (2003).
 61. M. B. Singer, R. Aalto, L. A. James, Status of the lower Sacramento Valley flood-control system within the context of its natural geomorphic setting. *Nat. Hazards Rev.* **9**, 104-115 (2008).
 62. J.-P. Bravard, P. Gaydou, "Historical Development and Integrated Management of the Rhône River Floodplain, from the Alps to the Camargue Delta, France" in *Geomorphic Approaches to Integrated*

- 618 Floodplain Management of Lowland Fluvial Systems in North America and
 619 Europe, P. F. Hudson, H. Middelkoop, Eds. (Springer New York, 2015),
 620 10.1007/978-1-4939-2380-9_12 chap. 12, pp. 289-320.
 621 63. J. Stella, P. Rodríguez-González, S. Dufour, J. Bendix, Riparian vegetation
 622 research in Mediterranean-climate regions: common patterns, ecological
 623 processes, and considerations for management. *Hydrobiologia* **719**, 291-
 624 315 (2013).



625
 626 **Figure 1.** Schematic of local and non-local hydroclimatic controls affecting water availability to
 627 riparian trees and their use of available water.




Figure 2. Map of the study region and field sites (a) with mean growing season (May-September) temperature and total precipitation for each site (b). Numbered boxes correspond to the nearest climate stations for each field site: 1 = Bron, 2 = Montélimar, 3 = Arles Tour du Valat. Aerial imagery from GoogleEarth ©2017, CNES / Airbus, Maxar Technologies. Imagery dates: Upstream -22/08/2016, Midstream-07/9/2015, Downstream-26/03/2016.




Figure 3. (a-c) Kernel density plots of all sub-annual $\delta^{18}\text{O}_{\text{msw}}$ values calculated for each near-channel (NC) and interior floodplain (IF) *Fraxinus* and *Populus* at each of the study sites. (d-f) Potential endmember water source $\delta^{18}\text{O}$ characterization for each site. The non-growing season and growing season precipitation $\delta^{18}\text{O}$ are denoted as NGS ppt and GS ppt, respectively.





Figure 4. Modeled source water $\delta^{18}\text{O}$ ($\delta^{18}\text{O}_{msw}$) for the sub-annual $\delta^{18}\text{O}_{cell}$ samples obtained from each tree for the 2000-2010 period at study sites, Upstream (a-d), Midstream (e-h), and Downstream (j-l). Blue and red lines correspond to the sub-annual $\delta^{18}\text{O}_{msw}$ means for *Fraxinus* and *Populus*, respectively. Grey shading represents ± 1 SD.

647
648



649

650 **Figure 5.** Mean sub-annual $\delta^{18}O_{msw}$ for sampled trees at each site and floodplain location (a-
651 c), corresponding growing season river stage percentiles (d-f) and mean seasonal SPEI series
652 for 1-6-month timescales (g-i).

653