



RNAAS RESEARCH NOTES OF THE AAS



IOP

OPEN ACCESS

ZTF-identified HW Virginis Systems

Josie Enenstein¹, Kevin Burdge¹, and Thomas Prince¹

Published April 2021 ¥ © 2021. The Author(s). Published by the American Astronomical Society.

[Research Notes of the AAS, Volume 5, Number 4](#)

Citation Josie Enenstein *et al* 2021 *Res. Notes AAS* 5 90

626 Total downloads

Share this article

--	--	--	--

--

Article information

Author e-mails

josieenenstein@gmail.com

josieenenstein@gmail.com

Author affiliations

¹ Division of Physics Mathematics and Astronomy, California Institute of Technology, Pasadena, CA 91125, USA; josieenenstein@gmail.com

ORCID iDs

Josie Enenstein <https://orcid.org/0000-0002-4084-9098>

Kevin Burdge <https://orcid.org/0000-0002-7226-836X>

Thomas Prince <https://orcid.org/0000-0002-8850-3627>

Dates

Received January 2021

Revised March 2021

Accepted March 2021

Published April 2021

DOI

<https://doi.org/10.3847/2515-5172/abf4d3>

Keywords

[Common envelope binary stars](#); [Light curves](#); [Light curve classification](#); [Eclipsing binary stars](#); [Common envelope evolution](#); [Interacting binary stars](#); [O subdwarf stars](#); [Automated telescopes](#); [Optical astronomy](#); [B subdwarf stars](#); [Surveys](#); [Cataclysmic variable stars](#)

- ☐ Journal RSS
- ☐ Sign up for new issue notifications
- ☐ Create citation alert

Abstract

We present 46 HW Virginis Post-Common Envelope Binary Systems identified by the Zwicky Transient Facility, 26 of which appear to be new discoveries. Through the application of the Box Least-Squares Algorithm to ZTF data, we searched a data set of 17,473 systems for periodicity and phase-folded all of the ZTF lightcurves of the objects at their best period. Through visual analysis of these phase-folded lightcurves, we identified the 46 HW Vir Systems.

Export citation and abstract

- [BibTeX](#)
- [RIS](#)

Previous article in issue

Next article in issue

☐ Related links

- [NASA ADS Record](#) ☐
- [About Related Links](#)



Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

1. Introduction

HW Virginis (HW Vir) systems are Post Common Envelope Binaries (Schaffenroth et al. [2019](#)) consisting of a hot subluminescent dwarf star (Heber [2016](#)) and an M dwarf companion star (Ivanova et al. [2013](#)). All data from this project were taken on the Samuel Oschin Schmidt 48 inch telescope of the Zwicky Transient Facility (ZTF) at Palomar Observatory in Southern California (Bellm et al. [2019](#)) (Graham et al. [2019](#)). This survey has a decl. limit of -30° , a 47 deg^2 field of view, and has a 5σ limiting magnitude at about 20.6 for *r*-band and 20.8 in *g*-band (Burdge et al. [2020](#)). Using data from the Gaia hot subdwarf catalog (Geier et al. [2019](#)) and cross-matching it with ZTF data, a list of objects observed by ZTF was created. Through periodicity analysis, we identify 46 HW Vir candidates. This collection of objects can be used as training data for supervised machine learning methods to help create an automated system capable of identifying HW Vir lightcurves.


2. Data Analysis

In order to extract the sample of lightcurves, we cross-matched the catalog of 39,800 hot subdwarfs reported in Geier et al. ([2019](#)) with the ZTF DR3 archival lightcurve database, and analyzed all objects with more than 50 5σ detections in individual ZTF-*g* and ZTF-*r* exposures, yielding 17,473 candidates. Due to its speed, we used a graphics processing unit implementation of the BLS algorithm (Kovács et al. [2002](#)) to determine the best period for each object, and visually inspected all 4572 of the phase-folded lightcurves in the sample with best periods falling between 1 hr and 10 days in order to identify the HW Vir candidates.

3. Results

We include a machine-readable table containing our data from our 46 HW Vir systems. We crossmatched our data with the EREBOS survey, which contains 161 HW Vir, (Schaffenroth et al. [2019](#)) to find that 26 of our HW Vir appear to be newly

discovered systems. Our galactic coordinate plot of these systems' locations shows the 46 HW Vir to be generally concentrated between galactic longitudes of -20° and 20° (see Figure 1). The median orbital period of the systems identified in this work is roughly 0.16 days (3.8 hr), which is quite short. Objects with periods this short will interact, and many will evolve into mass-transferring cataclysmic variables.



ZTF is a project led by PI S. R. Kulkarni at Caltech, and includes IPAC; WIS, Israel; OKC, Sweden; JSI/UMd, USA; UW, USA; DESY, Germany; NRC, Taiwan; UW Milwaukee, USA; and LANL USA. ZTF acknowledges the generous support of the NSF under AST MSIP Grant No 1440341. We thank the Caltech Summer Research Connections program for high-school students, under which Josie Enenstein did her research work. This work is based on observations obtained with the Samuel Oschin 48 inch Telescope at the Palomar Observatory.

[Show References](#)



[Journals](#) [Books](#) [About IOPscience](#) [Contact us](#) [Developing countries access](#) [IOP Publishing open access policy](#)

© Copyright 2021 IOP Publishing [Terms & conditions](#) [Disclaimer](#) [Privacy & cookie policy](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies.