



Initial Data Rigidity Results

Michael Eichmair¹ , Gregory J. Galloway², Abraão Mendes³

¹ Faculty of Mathematics, University of Vienna, Vienna, Austria. E-mail: michael.eichmair@univie.ac.at

² Department of Mathematics, University of Miami, Coral Gables, FL, USA.

E-mail: galloway@math.miami.edu

³ Instituto de Matemática, Universidade Federal de Alagoas, Maceió, AL, Brazil.

E-mail: abraao.mendes@im.ufal.br

Received: 2 October 2020 / Accepted: 3 February 2021

© The Author(s) 2021

Abstract: We prove several rigidity results related to the spacetime positive mass theorem. A key step is to show that certain marginally outer trapped surfaces are weakly outermost. As a special case, our results include a rigidity result for Riemannian manifolds with a lower bound on their scalar curvature.

1. Introduction

In this paper we establish several rigidity results for initial data sets that are motivated by the spacetime positive mass theorem.

An initial data set (M, g, K) consists of a connected Riemannian manifold (M, g) and a symmetric $(0, 2)$ -tensor field K . In addition, we assume that M is oriented throughout this paper.

Let (M, g, K) be an initial data set.

The *local energy density* μ and the *local current density* J of (M, g, K) are given by

$$\mu = \frac{1}{2} \left(R - |K|^2 + (\operatorname{tr} K)^2 \right) \quad \text{and} \quad J = \operatorname{div} (K - (\operatorname{tr} K) g).$$

Here, R is the scalar curvature of (M, g) . The initial data set is said to satisfy the *dominant energy condition* (DEC for short) if

$$\mu \geq |J|.$$

Let $\Sigma \subset M$ be a two-sided hypersurface with unit normal ν and

$$H = \operatorname{div}_{\Sigma} \nu$$

be the associated mean curvature. The *future outgoing null expansion scalar* θ^+ and *past outgoing null expansion scalar* θ^- of Σ are the quantities

$$\theta^+ = H + \text{tr}_\Sigma(K) \quad \text{and} \quad \theta^- = H - \text{tr}_\Sigma(K).$$

The hypersurface Σ is *outer trapped* if $\theta^+ < 0$, *weakly outer trapped* if $\theta^+ \leq 0$, and *marginally outer trapped* if $\theta^+ = 0$. In the latter case, we refer to Σ as a *marginally outer trapped surface* (MOTS for short). Unless stated otherwise, we require that MOTS are closed, i.e. compact and without boundary.

We also consider the quantities

$$\chi^+ = A + K|_\Sigma \quad \text{and} \quad \chi^- = A - K|_\Sigma$$

where A is the second fundamental form of Σ . Our sign convention is such that $H = \text{tr}_\Sigma A$ and thus $\theta^\pm = \text{tr}_\Sigma \chi^\pm$.

Initial data sets arise naturally in general relativity. Let $\bar{M}$ be a spacelike hypersurface in a spacetime, i.e. a time-oriented Lorentzian manifold, $(\bar{M}, \bar{g})$. Let g be the Riemannian metric induced on M and K be the second fundamental form with respect to the future-pointing unit normal u of M in $\bar{M}$. Then (M, g, K) is an initial data set. In this setting, χ^+ and χ^- are the *null second fundamental forms* with respect to the *null normal fields*

$$\ell^+ = v + u|_\Sigma \quad \text{and} \quad \ell^- = v - u|_\Sigma$$

of Σ viewed as a surface in $\bar{M}$. Note that $\theta^\pm = \text{div}_\Sigma \ell^\pm$.

An initial data set (M, g, K) is said to be *time-symmetric* or *Riemannian* if $K = 0$. In this case, the DEC is the requirement that the scalar curvature of (M, g) be non-negative. Moreover, Σ is a MOTS if and only if it is a minimal surface in (M, g) . Quite generally, MOTS share many properties with minimal surfaces, which they generalize; cf. e.g. the survey article [3].

The following version of the spacetime positive mass theorem has been obtained by L.-H. Huang, D. A. Lee, R. Schoen, and the first-named author in [11].

Theorem 1.1 [11]. *Let (M, g, K) be an n -dimensional asymptotically flat initial data set with ADM energy-momentum vector (E, P) . Assume that $3 \leq n \leq 7$. If the dominant energy condition $\mu \geq |J|$ is satisfied, then $E \geq |P|$.*

We refer to [11] for the definition of the energy-momentum vector. The case of equality $E = |P|$ has recently been characterized by L.-H. Huang and D. Lee [17].

In [19], J. Lohkamp has presented a different proof of Theorem 1.1 for all $n \geq 3$. His method is by reduction to and proof of the following result: Let (M, g, K) be an initial data set that is isometric to Euclidean space, with $K = 0$, outside some bounded open set $U \subset M$. Then one cannot have $\mu > |J|$ on U ; see [19, Theorem 2]. In particular, if (M, g, K) satisfies the DEC, there must be a point in U where $\mu = |J|$. The goal of our first result is to show that a much stronger conclusion holds when $3 \leq n \leq 7$.

Under the assumption of Lohkamp's result stated above, one obtains by obvious inclusion and identification a compact initial data set $(\tilde{M}, \tilde{g}, \tilde{K})$ with boundary $\partial\tilde{M} = \Sigma_1 \cup \Sigma_2$ where Σ_1 and Σ_2 are flat $(n-1)$ -tori in $(\tilde{M}, \tilde{g})$ that both are totally geodesic in the spacetime sense, i.e. $\chi^\pm = 0$ with respect to either choice of unit normal. In particular, both are MOTS. With this compactification (also used by Lohkamp) in mind, we state our first main rigidity result.

Theorem 1.2. *Let (M, g, K) be an n -dimensional, $3 \leq n \leq 7$, compact-with-boundary initial data set. Suppose that (M, g, K) satisfies the DEC, $\mu \geq |J|$. Suppose also that the boundary can be expressed as a disjoint union $\partial M = \Sigma_0 \cup S$ of non-empty unions of components such that the following conditions hold:*

- (1) $\theta^+ \leq 0$ on Σ_0 with respect to the normal that points into M .
- (2) $\theta^+ \geq 0$ on S with respect to the normal that points out of M .
- (3) M satisfies the homotopy condition with respect to Σ_0 .
- (4) Σ_0 satisfies the cohomology condition.

Then, the following hold:

- (i) $M \cong [0, \ell] \times \Sigma_0$ for some $\ell > 0$.
Let $\Sigma_t \cong \{t\} \times \Sigma_0$ with unit normal v_t in direction of the foliation.
- (ii) $\chi^+ = 0$ on Σ_t for every $t \in [0, \ell]$.
- (iii) Σ_t is a flat torus with respect to the induced metric for every $t \in [0, \ell]$.
- (iv) $\mu + J(v_t) = 0$ on Σ_t for every $t \in [0, \ell]$. In particular, $\mu = |J|$ on M .

The definitions of the cohomology condition and the homotopy condition are given in Sect. 3. The cohomology condition ensures that Σ_0 does not admit a metric of positive scalar curvature. The homotopy condition holds, for example, if M has almost product topology $M \cong ([0, 1] \times \Sigma_0) \# N$ where N is a closed manifold. It implies that Σ_0 is connected. A priori, we allow S to have multiple components.

The assumptions of Theorem 1.2 are satisfied in the compactified picture (after Lohkamp) described above. Note that Theorem 1.2 provides a relatively simple proof of Theorem 2 in [19] in dimensions $3 \leq n \leq 7$. In conjunction with Corollary 2.11 in [19], this leads to an alternative proof of Theorem 1.1 stated above. Conversely, note that under the assumptions of Theorem 1.2, the proof of Theorem 2 in [19] implies that $\mu = |J|$.

Theorem 1.2 is a global version of the local rigidity result for MOTS obtained in [13] and stated here as Theorem 2.1. We apply Lemma 3.2 to ensure that the weakly outermost condition (see Sect. 2) of this local rigidity result holds in our setting.

Imposing a convexity condition on the spacetime second fundamental form K , we are able to prove the following, stronger rigidity result. Note also that the boundary conditions are different from those in Theorem 1.2.

Theorem 1.3. *Let (M, g, K) be an n -dimensional, $3 \leq n \leq 7$, compact-with-boundary initial data set. Suppose that (M, g, K) satisfies the DEC, $\mu \geq |J|$. Suppose also that the boundary can be expressed as a disjoint union $\partial M = \Sigma_0 \cup S$ of non-empty unions of components such that the following conditions hold:*

- (1) $\theta^+ \leq 0$ on Σ_0 with respect to the normal that points into M .
- (2) $\theta^- \geq 2(n-1)\epsilon$ on S with respect to the normal that points out of M , where $\epsilon = 0$ or $\epsilon = 1$.
- (3) M satisfies the homotopy condition with respect to Σ_0 .
- (4) Σ_0 satisfies the cohomology condition.
- (5) $K + \epsilon g$ is $(n-1)$ -convex.

Then, the following hold:

- (i) (Σ_0, g_0) is a flat torus, where g_0 is the induced metric on Σ_0 .
- (ii) (M, g) is isometric to $([0, \ell] \times \Sigma_0, dt^2 + e^{2\epsilon t} g_0)$ for some $\ell > 0$.
- (iii) $K = (1 - \epsilon) a dt^2 - \epsilon g$ on M , where $a \in C^\infty(M)$ depends only on $t \in [0, \ell]$.
- (iv) $\mu = 0$ and $J = 0$ on M .

The definition of $(n - 1)$ -convexity is recalled in Sect. 2. Note that $K + \epsilon g$ is $(n - 1)$ -convex in the special case where it is positive semi-definite. The case $\epsilon = 1$ is relevant in the asymptotically hyperbolic or asymptotically hyperboloidal setting. Theorem 1.3 contains the following Riemannian result.

Corollary 1.4. *Let (M, g) be an n -dimensional, $3 \leq n \leq 7$, connected, oriented, compact-with-boundary Riemannian manifold. Suppose that the scalar curvature of (M, g) satisfies $R \geq -n(n - 1)\epsilon$, where $\epsilon = 0$ or $\epsilon = 1$. Suppose also that the boundary can be expressed as a disjoint union $\partial M = \Sigma_0 \cup S$ of non-empty unions of components such that the following conditions hold:*

- (1) *The mean curvature of Σ_0 in (M, g) with respect to the normal that points into M satisfies $H \leq (n - 1)\epsilon$.*
- (2) *The mean curvature of S in (M, g) with respect to the normal that points out of M satisfies $H \geq (n - 1)\epsilon$.*
- (3) *M satisfies the homotopy condition with respect to Σ_0 .*
- (4) *Σ_0 satisfies the cohomology condition.*

Then (Σ_0, g_0) is a flat torus, where g_0 is the induced metric on Σ_0 . Moreover, (M, g) is isometric to $([0, \ell] \times \Sigma_0, dt^2 + e^{2\epsilon t} g_0)$ for some $\ell > 0$.

Note the similarity of Corollary 1.4 with the rigidity result Theorem 1 in [8] in the Ricci curvature setting, due to C. B. Croke and B. Kleiner. Theorem 1.1 in [2] follows from Corollary 1.4 in the special case where $\epsilon = 1$. Corollary 1.4 should also be compared to the results of H. C. Jang and the second-named author in [14], which require an *outermost* condition. An alternative proof of Corollary 1.4 may be given using area minimization when $\epsilon = 0$ and minimization of the so-called brane action when $\epsilon = 1$. The MOTS methodology presented in this paper gives a synthetic way of treating both cases, and much more, simultaneously.

We review some background material on MOTS in Sect. 2. In Sect. 3, we establish criteria to verify the weakly outermost condition for MOTS. In Sect. 4, we give a proof of Theorem 1.2. In Sect. 5, we give the proofs of Theorem 1.3 and Corollary 1.4 and also consider some additional results. Finally, in Sect. 6, we show how to embed the initial data set in Theorem 1.3 into a quotient of Minkowski space.

2. Preliminaries

We recall several results for MOTS that are needed in this paper.

Let (M, g, K) be an initial data set and $\Sigma \subset M$ be a closed MOTS with unit normal ν .

Let $\{\Sigma_t\}_{|t| < \epsilon}$ be a variation of Σ , where

$$\Sigma_t = \{\exp_x(t \phi(x) \nu(x)) : x \in \Sigma\}$$

for some $\phi \in C^\infty(M)$. We may view the expansion scalar θ^+ of these hypersurfaces as a parameter-dependent function on Σ . We recall from e.g. [4, p. 861] or [3, p. 20] that

$$\left. \frac{d}{dt} \right|_{t=0} \theta^+(t, \cdot) = L \phi \tag{1}$$

where

$$L \phi = -\Delta \phi + 2 \langle X, \nabla \phi \rangle + \left(Q - |X|^2 + \operatorname{div}(X) \right) \phi$$

and

$$Q = \frac{1}{2} R_\Sigma - \frac{1}{2} |\chi^+|^2 - \mu - J(v).$$

Here, Δ is the non-positive definite Laplace-Beltrami operator, ∇ the gradient, div the divergence, and R_Σ the scalar curvature of $(\Sigma, \langle \cdot, \cdot \rangle)$. Moreover, X is the tangent field of Σ that is dual to the form $K(v, \cdot)$.

If there is a $\phi \in C^\infty(\Sigma)$ with $\phi > 0$ and

$$L\phi \geq 0,$$

then Σ is called a *stable MOTS*; cf. [4, p. 868]. We refer in passing to related notions of stability for MOTS with boundary and their applications; see e.g. [15, p. 3], [12, Sect. 2], or [1, Sect. 5].

Assume now that Σ is a boundary in M . More precisely, assume that v points towards a top-dimensional submanifold $M_+ \subset M$ such that $\partial M_+ = \Sigma \cup S$ where S is a union of components of ∂M . We think of M_+ as the region outside of Σ . Then Σ is called an *outermost MOTS* if there is no closed embedded surface in M_+ with $\theta^+ \leq 0$ that is homologous to and different from Σ . If there is no such surface with $\theta^+ < 0$, then Σ is called *weakly outermost*.

We now state the local rigidity result for MOTS from [13] mentioned in the introduction.

Theorem 2.1 ([13], Theorem 3.1). *Let (M, g, K) be an n -dimensional, $n \geq 3$, initial data set that satisfies the DEC, $\mu \geq |J|$. Suppose that $\Sigma \subset M$ is a connected weakly outermost MOTS that does not support a metric of positive scalar curvature. There is a neighborhood $U \subset M$ of Σ with $U \cap M_+ \cong [0, \delta) \times \Sigma$ such that the following hold:*

- (i) $\Sigma_0 = \Sigma$ where $\Sigma_t \cong \{t\} \times \Sigma$ for every $t \in [0, \delta)$.
- (ii) $\chi^+ = 0$ on Σ_t for every $t \in [0, \delta)$ with respect to the unit normal v_t in direction of the foliation. In particular, Σ_t is a MOTS for every $t \in [0, \delta)$.
- (iii) The metric induced on Σ_t by g is Ricci-flat for every $t \in [0, \delta)$.
- (iv) $\mu + J(v_t) = 0$ on Σ_t for every $t \in [0, \delta)$. In particular, $\mu = |J|$ on $U \cap M_+$.

Theorem 2.1 implies the following result related to the topology of apparent horizons.

Corollary 2.2. *Let (M, g, K) be an n -dimensional, $n \geq 3$, initial data set that satisfies the DEC, $\mu \geq |J|$. Suppose that $\Sigma \subset M$ is an outermost MOTS in (M, g, K) . Then Σ admits a metric of positive scalar curvature.*

We will apply the following existence result for MOTS. It was obtained by L. Andersson and J. Metzger [5] in dimension $n = 3$ and then, using different techniques, by the first-named author [9, 10] in dimensions $3 \leq n \leq 7$. The approaches in [5, 9, 10] are all based on an idea of R. Schoen to construct MOTS between suitably trapped hypersurfaces by forcing a blow up of the Jang equation. See also [3] for a survey of these existence results.

Theorem 2.3 [5, 9, 10]. *Let (M, g, K) be an n -dimensional, $3 \leq n \leq 7$, compact-with-boundary initial data set. Suppose that the boundary can be expressed as a disjoint union $\partial M = \Sigma_{\text{in}} \cup \Sigma_{\text{out}}$ where $\Sigma_{\text{in}}, \Sigma_{\text{out}}$ are non-empty unions of components of ∂M with $\theta^+ \leq 0$ on Σ_{in} with respect to the normal pointing into M and with $\theta^+ > 0$ on Σ_{out} with respect to the normal pointing out of M . Then there is an outermost MOTS in (M, g, K) that is homologous to Σ_{out} .*

Some of our results require a convexity condition on the spacetime second fundamental form K of the initial data set (M, g, K) . We say that a symmetric $(0, 2)$ -tensor field P is $(n - 1)$ -convex if, at every point, the sum of the smallest $(n - 1)$ eigenvalues of P with respect to g is non-negative. In particular, if P is $(n - 1)$ -convex, then $\text{tr}_\Sigma P \geq 0$ for every hypersurface $\Sigma \subset M$. This convexity condition has been used by the third-named author in [20] in a related context.

3. The Weakly Outermost Condition

In this section, we assume that (M, g, K) is a compact initial data set. While we do not assume a priori that M is topologically a product, we require a more general condition of a similar flavor. Let Σ_0 be a union of components of ∂M . We say that M satisfies the *homotopy condition* with respect to Σ_0 provided that there exists a continuous map $\rho : M \rightarrow \Sigma_0$ such that $\rho \circ i : \Sigma_0 \rightarrow \Sigma_0$ is homotopic to id_{Σ_0} where $i : \Sigma_0 \rightarrow M$ is the inclusion map. Since M is connected by assumption, this condition implies that Σ_0 is connected. Note that this homotopy condition is satisfied if there is a retraction of M onto Σ_0 .

An orientable, closed manifold N of dimension m is said to satisfy the *cohomology condition* if there are classes $\omega_1, \dots, \omega_m \in H^1(N, \mathbb{Z})$ whose cup product

$$\omega_1 \smile \dots \smile \omega_m \in H^m(N, \mathbb{Z})$$

is non-zero. Such a manifold N has a component that does not admit a metric of positive scalar curvature; see [22, Theorem 5.2] and the discussion of [18, Theorem 2.28]. Note that every manifold diffeomorphic to $\mathbb{T}^m$ or, more generally, to $\mathbb{T}^m \# Q$ with Q oriented and closed satisfies the cohomology condition; cf. [22, Theorem 5.1].

Lemma 3.1. *Let M be an orientable, compact n -dimensional, $n \geq 3$, manifold with boundary. Let Σ_0 be a union of components of ∂M . Suppose that M satisfies the homotopy condition with respect to Σ_0 and that Σ_0 satisfies the cohomology condition. Then every closed, embedded hypersurface $\Sigma \subset M$ homologous to Σ_0 satisfies the cohomology condition.*

In particular, by the preceding discussion, Σ does not support a metric of positive scalar curvature.

Proof. Let $\rho : M \rightarrow \Sigma_0$ be a continuous map such that $\rho \circ i \simeq \text{id}_{\Sigma_0}$, where $i : \Sigma_0 \rightarrow M$ is the inclusion map. Let $\omega_1, \dots, \omega_{n-1} \in H^1(\Sigma_0, \mathbb{Z})$ be classes with $\omega_1 \smile \dots \smile \omega_{n-1} \neq 0$. Let $j : \Sigma \rightarrow M$ be the inclusion map. The map $\sigma = \rho \circ j : \Sigma \rightarrow \Sigma_0$ induces a map $H_{n-1}(\Sigma, \mathbb{Z}) \rightarrow H_{n-1}(\Sigma_0, \mathbb{Z})$. Since Σ and Σ_0 are homologous,

$$\sigma_*[\Sigma] = \rho_*(j_*[\Sigma]) = \rho_*(i_*[\Sigma_0]) = (\text{id}_{\Sigma_0})_*[\Sigma_0] = [\Sigma_0].$$

Using this, we can conclude the proof arguing as in [18, p. 45]. Note that

$$\sigma_*([\Sigma] \frown (\sigma^* \omega_1 \smile \dots \smile \sigma^* \omega_{n-1})) = [\Sigma_0] \frown (\omega_1 \smile \dots \smile \omega_{n-1}) \neq 0.$$

In particular, $\sigma^* \omega_1 \smile \dots \smile \sigma^* \omega_{n-1} \neq 0$. □

We combine the previous lemma with Corollary 2.2 and Theorem 2.3 to show that the weakly outermost condition follows from seemingly weaker assumptions.

Lemma 3.2. *Let (M, g, K) be an n -dimensional, $3 \leq n \leq 7$, compact-with-boundary initial data set. Suppose that (M, g, K) satisfies the DEC, $\mu \geq |J|$. Suppose also that the boundary can be expressed as a disjoint union $\partial M = \Sigma_0 \cup S$ of non-empty unions of components such that the following conditions hold:*

- (1) $\theta^+ \leq 0$ on Σ_0 with respect to the normal that points into M .
- (2) $\theta^+ \geq 0$ on S with respect to the normal that points out of M .
- (3) M satisfies the homotopy condition with respect to Σ_0 .
- (4) Σ_0 satisfies the cohomology condition.

Then Σ_0 is a weakly outermost MOTS in (M, g, K) .

In the proof of this result below, we compute expansion scalars with respect to different spacetime second fundamental forms. For clarity, we indicate by a subscript which spacetime second fundamental form is used in the computation.

Proof. First, we show that Σ_0 is a MOTS.

Suppose that θ_K^+ is not identically zero on Σ_0 . It follows from [5, Lemma 5.2] that there is a hypersurface $\Sigma \subset M$ obtained as a small perturbation of Σ_0 into M such that $\theta_K^+ < 0$ on Σ with respect to the normal pointing away from Σ_0 . Let W be the connected, compact region bounded by Σ and S in M . Observe that $\theta_{-K}^+ \leq 0$ on S with respect to the normal that points into W and $\theta_{-K}^+ > 0$ on Σ with respect to the normal that points out of W . Note that the initial data set $(W, g, -K)$ satisfies the DEC. Applying Theorem 2.3 to this initial data set, we obtain an outermost MOTS $\tilde{\Sigma}$ in $(W, g, -K)$ that is homologous to and disjoint from Σ . Clearly, $\tilde{\Sigma}$ is homologous to Σ_0 . By Lemma 3.1, $\tilde{\Sigma}$ does not support a metric of positive scalar curvature. This contradicts Corollary 2.2 applied to the initial data set $(W, g, -K)$. Thus Σ_0 is a MOTS in (M, g, K) .

Next, we show that Σ_0 is a weakly outermost MOTS.

Suppose, by contradiction, that Σ_0 is not weakly outermost. Then there is a hypersurface $\Sigma \subset M$ homologous to Σ_0 such that $\theta_K^+ < 0$ on Σ with respect to the normal that points away from Σ_0 . We may assume that each component of Σ is homologically non-trivial in M . Let W be the compact region in M bounded by Σ and S . Assume first that W is connected. Applying Theorem 2.3, we obtain an outermost MOTS $\tilde{\Sigma}$ in $(W, g, -K)$ homologous to Σ . As before, by Lemma 3.1, $\tilde{\Sigma}$ does not support a metric of positive scalar curvature. This contradicts Corollary 2.2. In the general case, we apply this argument separately to each component of W . At least one of the outermost MOTS obtained in this way does not carry a metric of positive scalar curvature. Again, this contradicts Corollary 2.2. $\square$

4. Proof of Theorem 1.2

Theorem 1.2 may be viewed as a global version of Theorem 2.1. We emphasize that Theorem 1.2 does not require the *weakly outermost* assumption.

We start with the following observation.

Lemma 4.1. *Under the assumptions of Theorem 1.2, there is a neighborhood of Σ_0 in M diffeomorphic to $[0, \delta) \times \Sigma_0$ such that the leaves $\Sigma_t \cong \{t\} \times \Sigma_0$ satisfy properties (ii)-(iv) of the conclusion of Theorem 1.2.*

Proof. According to Lemma 3.2, Σ_0 is a weakly outermost MOTS in (M, g, K) . We may apply Theorem 2.1. We claim that the foliation from Theorem 2.1 has the asserted properties. It only remains to show that each Σ_t is isometric to a flat torus. To see

this, note the estimate $b_1(\Sigma_t) \geq n$ for the first Betti number of Σ_t . This follows from the cohomology condition, Poincaré duality, and the fact that $H^n(M, \mathbb{Z})$ is torsion free. Conversely, by a classical result of Bochner, see e.g. [21, p. 208], it holds that $b_1(\Sigma_t) \leq n$ with equality if and only if Σ_t is isometric to a flat torus. $\square$

Proof of Theorem 1.2. We use ν to denote the unit normal field of the foliation $\{\Sigma_t\}_{t \in [0, \delta]}$ from Lemma 4.1. Note that the divergence of ν evaluated on Σ_t is equal to the mean curvature of Σ_t . Since every leaf Σ_t is a MOTS, we see that the divergence of ν is bounded. By the divergence theorem,

$$\text{vol}(\Sigma_t) = \text{vol}(\Sigma_0) + \int_{U_t} \text{div}(\nu)$$

where $U_t \cong [0, t] \times \Sigma_0$ is the collar between Σ_0 and Σ_t . This argument shows that $\text{vol}(\Sigma_t)$ is bounded independently of $t \in [0, \delta]$.

Note that the second fundamental form of each Σ_t is bounded independently as well, since the null second fundamental form of each Σ_t vanishes.

To proceed, we briefly recall a standard fact. For convenience of exposition, we extend (M, g) across its boundary to a homogeneously regular manifold. Given $C > 0$, there is a small constant $r > 0$ with the following property. Let $\Sigma \subset M$ be a closed and two-sided surface whose second fundamental form is bounded by C . Let $p \in M$ be such that $\Sigma \cap B_r(p) \neq \emptyset$. Then Σ contains the graph of a function $f : \{y \in \mathbb{R}^{n-1} : |y| < r\} \rightarrow \mathbb{R}$ with $|f(0)| < r$, $|Df| \leq 1$, and $|D^2 f| \leq 2C$, where an appropriately rotated geodesic coordinate system with center at p is used to identify $B_{2r}(p)$ with the Euclidean ball $\{x \in \mathbb{R}^n : |x| < 2r\}$. In fact, it is possible to choose the geodesic coordinate system so that $\Sigma \cap B_r(p)$ is covered by such graphs.

It follows from these facts that the leaves $\{\Sigma_t\}_{t \in [0, \delta]}$ have a smooth *immersed* limit Σ_δ as $t \nearrow \delta$. We use an idea of L. Andersson and J. Metzger [5] to show that Σ_δ is embedded. For if not, we can find for every $\eta > 0$ a leaf Σ_t and $p \in M$ such that $\Sigma_t \cap B_r(p)$ contains the graphs of two functions f_1, f_2 with the properties stated above and such that $|f_1(0) - f_2(0)| < \eta$. In fact, we can arrange for the layer between these graphs to lie to the outside of Σ_t . Arguing exactly as in Sect. 6 of [5], if $\eta > 0$ is sufficiently small, it is possible to glue in a neck to connect Σ_t across this layer to obtain a surface with non-positive expansion and negative expansion around the neck. By flowing this surface outward at the speed of its expansion as in Lemma 5.2 of [5], one obtains a surface homologous to Σ_t with everywhere negative expansion. This contradicts the fact that Σ_0 is weakly outermost.

It is easy to see from the proof of Theorem 2.1 that $\{\Sigma_t\}_{t \in [0, \delta]}$ is a foliation. Recall that M and $\Sigma_\delta \cong \Sigma_0$ are connected. By the strong maximum principle as in e.g. [6, Proposition 3.1] or [5, Proposition 2.4], we have that $\Sigma_\delta = S$ if $\Sigma_\delta \cap S \neq \emptyset$. Note that the assumptions of the theorem continue to hold if we replace Σ_0 by Σ_δ and M by the complement of U in M . The result now follows by a continuity argument. $\square$

Example 4.2. The following example shows that there is still a fair amount of flexibility in the initial data sets covered by Theorem 1.2.

Let $\mathbb{R}_1^3$ be Minkowski space with standard coordinates t, x, y, z . Consider the box $\mathcal{B} = \{(x, y, z) : 0 \leq x \leq 1, 0 \leq y \leq 1, 0 \leq z \leq 1\}$ in the $t = 0$ slice. Let $f : \mathcal{B} \rightarrow \mathbb{R}$ be a smooth function that vanishes near the boundary of $\mathcal{B}$ and whose graph is spacelike in $\mathbb{R}_1^3$. We identify opposite sides in the x and the y coordinate to obtain an initial data set (M, g, K) with $M \cong \mathbb{T}^2 \times [0, 1]$. Let Σ_0 be the torus corresponding to $t = z = 0$. Note that (M, g, K) satisfies the conditions of Theorem 1.2. The foliation in the conclusion

of Theorem 1.2 arises from intersecting the graph of f with the null hypersurfaces $\mathcal{H}_c : t = z + c$. This can be understood using the following standard argument; see e.g. [7, Appendix A]. The hypersurfaces $\mathcal{H}_c$ are totally geodesic null hypersurfaces, i.e. each has vanishing null second fundamental form with respect to any null vector field K_c tangent to $\mathcal{H}_c$. Since K_c is orthogonal to every spacelike cross section, it follows that all these cross sections have vanishing null second fundamental form. In particular, they are MOTS. Moreover, again because $\mathcal{H}_c$ is totally geodesic, the induced metric on every spacelike cross section is invariant under the flow generated by K_c . It follows that any two such cross sections are isometric.

5. Proof of Theorem 1.3 and Further Consequences

As in the proof of Lemma 3.2, we will compute several quantities with respect to different spacetime second fundamental forms. We indicate by subscript the second fundamental form that is used.

Theorem 1.3 follows from the local rigidity result below.

Lemma 5.1. *Assumptions as in Theorem 1.3. Then (Σ_0, g_0) is a flat torus, where g_0 is the metric on Σ_0 induced by g . Moreover, there is a neighborhood U of Σ_0 in M such that the following hold:*

- (i) (U, g) is isometric to $([0, \delta) \times \Sigma_0, dt^2 + e^{2\epsilon t} g_0)$, for some $\delta > 0$.
- (ii) $K = (1 - \epsilon) a dt^2 - \epsilon g$ on U , where a depends only on $t \in [0, \delta)$.
- (iii) $\mu = 0$ and $J = 0$ on U .

Proof. By assumption,

$$\theta_K^- = H - \text{tr}_S K \geq 2(n-1)\epsilon$$

on S , where H is the mean curvature of S with respect to the normal pointing out of M . Using also the assumption that $K + \epsilon g$ is $(n-1)$ -convex, we obtain

$$H \geq \text{tr}_S K + 2(n-1)\epsilon = \text{tr}_S K + (n-1)\epsilon + (n-1)\epsilon \geq (n-1)\epsilon.$$

Therefore,

$$\theta_K^+ = H + \text{tr}_S K \geq (n-1)\epsilon + \text{tr}_S K \geq 0$$

on S . It follows from Lemma 3.2 that Σ_0 is a weakly outermost MOTS in (M, g, K) . By Theorem 2.1, Σ_0 has a neighborhood $U_1 \cong [0, \delta_1) \times \Sigma_0$ in M such that the following hold:

- We have that

$$g = \phi_1^2 ds^2 + g_1(s)$$

on U_1 , where $g_1(s)$ is the metric on $\Sigma_1(s) \cong \{s\} \times \Sigma_0$ induced by g .

- Every leaf $\Sigma_1(s)$ is a MOTS. In fact,

$$0 = \chi_K^+(s) = A_1(s) + K|_{\Sigma_1(s)},$$

where $A_1(s)$ is the second fundamental form of $\Sigma_1(s)$ in M computed with respect to the unit normal $\nu_1(s)$ in direction of the foliation.

– We have that

$$\mu_K + J_K(v_1(s)) = 0.$$

Consider now the initial data set (M, g, P) , where

$$P = -K - 2\epsilon g.$$

Note that (M, g, P) satisfies the DEC. In fact,

$$\mu_P - |J_P| = \mu_K - |J_K| + 2(n-1)(\text{tr } K + n\epsilon)\epsilon \geq 0$$

where we have used the assumption that $K + \epsilon g$ is $(n-1)$ -convex.

By assumption, $\theta_K^+ = H + \text{tr}_{\Sigma_0} K \leq 0$ on Σ_0 , where H is the mean curvature of Σ_0 with respect to the normal that points into M . It follows that

$$H \leq -\text{tr}_{\Sigma_0} K \leq (n-1)\epsilon$$

on Σ_0 and thus

$$\theta_P^+ = H + \text{tr}_{\Sigma_0} P = H - \text{tr}_{\Sigma_0} K - 2(n-1)\epsilon \leq 0.$$

Also,

$$\theta_P^+ = \theta_K^- - 2\epsilon(n-1) \geq 0$$

on S . By Lemma 3.2, Σ_0 is a weakly outermost MOTS in (M, g, P) . It follows from Theorem 2.1 that there is a neighborhood U_2 of Σ_0 in M diffeomorphic to $[0, \delta_2) \times \Sigma_0$ for some $\delta_2 > 0$, such that the following hold:

– We have that

$$g = \phi_2^2 dt^2 + g_2(t)$$

on U_2 , where $g_2(t)$ is the metric on $\Sigma_2(t) \cong \{t\} \times \Sigma_0$ induced by g .

– Every leaf $\Sigma_2(t)$ is a MOTS. In fact,

$$0 = \chi_P^+(t) = A_2(t) + P|_{\Sigma_2(t)},$$

where $A_2(t)$ is the second fundamental form of $\Sigma_2(t)$ in M computed with respect to the unit normal $v_2(t)$ in direction of the foliation.

– $(\Sigma_2(t), g_2(t))$ is Ricci flat.

– We have that

$$\mu_P + J_P(v_2(t)) = 0.$$

Decreasing $\delta_2 > 0$, if necessary, we may assume that $U_2 \subset U_1$. Fix $t \in (0, \delta_2)$ and note that $\Sigma_1(s) \cap \Sigma_2(t) \neq \emptyset$ for some $s \in (0, \delta_1)$, since $\Sigma_2(t) \subset U_2 \subset U_1$ and $\Sigma_1(0) \cap \Sigma_2(t) = \Sigma_0 \cap \Sigma_2(t) = \emptyset$. Let

$$s_0 = s_0(t) = \inf\{s \in (0, \delta_1) : \Sigma_1(s) \cap \Sigma_2(t) \neq \emptyset\}$$

and note that $\Sigma_1(s_0) \cap \Sigma_2(t) \neq \emptyset$. In particular, $s_0 > 0$. Also, $\Sigma_1(s) \cap \Sigma_2(t) = \emptyset$ for all $s \in [0, s_0)$. This means that $\Sigma_2(t)$ is contained in the region outside of $\Sigma_1(s_0)$.

The mean curvature of $\Sigma_1(s_0)$ is given by

$$H_1(s_0) = \text{tr } \chi_K^+(s_0) - \text{tr}_{\Sigma_1(s_0)} K = -\text{tr}_{\Sigma_1(s_0)} K \leq (n-1)\epsilon.$$

For the mean curvature of $\Sigma_2(t)$, we have the estimate

$$H_2(t) = \text{tr } \chi_P^+(t) - \text{tr}_{\Sigma_2(t)} P = \text{tr}_{\Sigma_2(t)} K + 2(n-1)\epsilon \geq (n-1)\epsilon.$$

In particular,

$$H_1(s_0) \leq H_2(t)$$

so that

$$\Sigma_1(s_0) = \Sigma_2(t)$$

by the maximum principle.

We see that the foliations $\{\Sigma_1(s)\}_{s \in [0, \delta_1]}$ and $\{\Sigma_2(t)\}_{t \in [0, \delta_2]}$ are the same after reparametrization. Below, we will denote this foliation of a neighborhood U of Σ_0 in M by $\{\Sigma(t)\}_{t \in [0, \delta]}$. Note that $\chi_K^+ = 0$ and $\chi_P^+ = 0$ on each leaf $\Sigma(t)$. Let $\nu(t)$ be the unit normal of $\Sigma(t)$ in direction of the foliation, $g(t)$ the induced metric, $A(t)$ the second fundamental form with respect to $\nu(t)$, and ϕ the lapse function of the foliation. By (1), we have that

$$0 = \frac{\partial \theta_K^+}{\partial t} = -\Delta \phi + 2\langle X_K, \nabla \phi \rangle + \left(\text{div } X_K - |X_K|^2 \right) \phi$$

where

$$Q_K = \frac{1}{2} R_{\Sigma(t)} - (\mu_K + J_K(\nu(t))) - \frac{1}{2} |\chi_K^+|^2$$

vanishes. Arranging terms as in [16, (2.9)], we obtain that

$$\text{div}(X_K - \nabla \ln \phi) - |X_K - \nabla \ln \phi|^2 = 0.$$

Integrating both sides of this equation over $\Sigma(t)$ and applying the divergence theorem, we obtain that

$$X_K = \nabla \ln \phi$$

on $\Sigma(t)$. By the same argument, we find

$$X_P = \nabla \ln \phi.$$

From the definition of P ,

$$X_P = -X_K \quad \mu_P = \mu_K + 2(n-1)(\text{tr } K + n\epsilon) \quad J_P = -J_K.$$

Thus,

$$\mu_K = -J_K(\nu(t)) = J_P(\nu(t)) = -\mu_P = -\mu_K - 2(n-1)(\text{tr } K + n\epsilon)\epsilon$$

and

$$\nabla \ln \phi = X_K = -X_P = -\nabla \ln \phi.$$

It follows that

$$|J_K| \leq \mu_K = -(n-1)(\operatorname{tr} K + n\epsilon)\epsilon \leq 0 \quad \text{and} \quad \nabla \ln \phi = 0.$$

From this, we conclude that

$$\mu_K = 0 \quad J_K = 0 \quad (\operatorname{tr} K + n\epsilon)\epsilon = 0$$

on U . Moreover, the lapse function ϕ is constant on $\Sigma(t)$ for every $t \in [0, \delta)$.

Using that

$$0 = \chi_K^+ = A(t) + K|_{\Sigma(t)} \quad \text{and} \quad 0 = \chi_P^+ = A(t) - K|_{\Sigma(t)} - 2\epsilon g(t),$$

we obtain

$$A(t) = \epsilon g(t) = -K|_{\Sigma(t)}$$

and thus

$$g = dt^2 + e^{2\epsilon t} g_0.$$

Using also that $K(v(t), \cdot)|_{\Sigma(t)} = 0$ since $X_K = 0$, we see that

$$K = a dt^2 - \epsilon g(t)$$

on U . If $\epsilon = 1$, we have $\operatorname{tr} K = -n$. Thus $a = -1$ and hence $K = -g$ on U . If $\epsilon = 0$, we use that $d(\operatorname{tr} K) = \operatorname{div} K$ (since $J_K = 0$) to see that a is constant on every leaf $\Sigma(t)$. Finally, the same argument as in the proof of Lemma 4.1 shows that (Σ_0, g_0) is a flat torus. This completes the proof. $\square$

Proof of Theorem 1.3. Let

$$\ell = \sup\{\delta : \text{the conclusion of Lemma 5.1 holds with this value of } \delta > 0\}.$$

Note that $\ell < \infty$ since M is compact. Reasoning the embeddedness of the final sheet as in the proof of Theorem 1.2, we see that (M, g) is isometric to the warped product $([0, \ell] \times \Sigma_0, dt^2 + e^{2\epsilon t} g_0)$. Moreover, we see that (iii) and (iv) of Theorem 1.3 hold. $\square$

Proof of Corollary 1.4. Let $K = -\epsilon g$ and note that (M, g, K) satisfies the DEC. In fact, a straightforward calculation gives that

$$\mu_K = \frac{1}{2} (R + n(n-1)\epsilon) \geq 0 \quad \text{and} \quad J_K = 0.$$

The expansion θ_K^+ of Σ_0 in (M, g, K) computed with respect to the normal that points into M satisfies

$$\theta_K^+ = H + \operatorname{tr}_{\Sigma_0} K = H - (n-1)\epsilon \leq 0.$$

The expansion θ_K^- of S in (M, g, K) computed with respect to the normal that points out of M satisfies

$$\theta_K^- = H - \operatorname{tr}_S K = H + (n-1)\epsilon \geq 2(n-1)\epsilon.$$

Thus Theorem 1.3 applies to (M, g, K) and gives the assertion. $\square$

The next theorem establishes a rigidity result under the boundary conditions of Theorem 1.2, assuming a volume minimizing condition on Σ_0 .

Theorem 5.2. *Let (M, g, K) be an n -dimensional, $3 \leq n \leq 7$, compact-with-boundary initial data set. Suppose that (M, g, K) satisfies the DEC, $\mu \geq |J|$. Suppose also that the boundary can be expressed as a disjoint union $\partial M = \Sigma_0 \cup S$ of non-empty unions of components such that the following conditions hold:*

- (1) $\theta^+ \leq 0$ on Σ_0 with respect to the normal that points into M .
- (2) $\theta^+ \geq 0$ on S with respect to the normal that points out of M .
- (3) M satisfies the homotopy condition with respect to Σ_0 .
- (4) Σ_0 satisfies the cohomology condition.
- (5) K is $(n - 1)$ -convex.
- (6) Σ_0 is volume minimizing in (M, g) .

Then, the following hold:

- (i) (Σ_0, g_0) is a flat torus, where g_0 is the metric on Σ_0 induced by g .
- (ii) (M, g) is isometric to $([0, \ell] \times \Sigma_0, dt^2 + g_0)$, for some $\ell > 0$.
- (iii) $K = a dt^2$ on M , where a depends only on $t \in [0, \ell]$.
- (iv) $\mu = 0$ and $J = 0$ on M .

As shown in the following example, Theorem 5.2 fails to hold if one drops either the $(n - 1)$ -convexity assumption or the volume minimizing assumption.

Example 5.3. Let (Σ_0, g_0) be the square flat $(n - 1)$ -torus. Let (M, g) be the cylinder $([0, \ell] \times \Sigma_0, dt^2 + e^{2\epsilon t} g_0)$ and $K = -\epsilon g$, where $\epsilon = -1$ or $\epsilon = 1$. The second fundamental form of $\Sigma_t = \{t\} \times \Sigma_0$ in (M, g) with respect to the normal in direction of increasing values of t is given by $A(t) = \epsilon e^{2\epsilon t} g_0$. Then (M, g, K) satisfies all the assumptions of Theorem 5.2 except for the volume minimizing assumption in the case where $\epsilon = -1$ and the $(n - 1)$ -convexity assumption in the case where $\epsilon = 1$.

Proof of Theorem 5.2. It follows from Lemma 3.2 that Σ_0 is weakly outermost. Then, by Theorem 2.1, there exists a neighborhood U of Σ_0 in M diffeomorphic to $[0, \delta) \times \Sigma_0$ for some $\delta > 0$, such that:

- We have that

$$g = \phi^2 dt^2 + g(t)$$

on U , where $g(t)$ is the metric on $\Sigma(t) \cong \{t\} \times \Sigma_0$ induced by g .

- Every leaf $\Sigma(t)$ is a MOTS. In fact,

$$\chi_K^+(t) = A(t) + K|_{\Sigma(t)} = 0,$$

where $A(t)$ is the second fundamental form of $\Sigma(t)$ in (M, g) .

- $(\Sigma(t), g(t))$ is Ricci flat.
- We have that

$$\mu_K + J_K(v(t)) = 0,$$

where $v(t)$ is the unit normal field on $\Sigma(t)$ in direction of increasing values of t .

Since $\text{tr}_{\Sigma(t)} K \geq 0$, we have

$$H(t) \leq H(t) + \text{tr}_{\Sigma(t)} K = \text{tr } \chi_K^+(t) = 0$$

where $H(t)$ is the mean curvature of $\Sigma(t)$ in (M, g) . The first variation formula for the volume of $(\Sigma(t), g(t))$ gives

$$\frac{d}{dt} \text{Vol}(\Sigma(t), g(t)) = \int_{\Sigma(t)} \phi H(t) d \text{vol}_{g(t)} \leq 0.$$

In particular,

$$\text{Vol}(\Sigma(t), g(t)) \leq \text{Vol}(\Sigma_0, g_0) \quad (2)$$

for every $t \in [0, \delta)$. Since Σ_0 is volume minimizing by assumption, we obtain

$$\text{Vol}(\Sigma(t), g(t)) = \text{Vol}(\Sigma_0, g_0)$$

for all $t \in [0, \delta)$. Then, by (2), we have $H(t) = 0$, which implies $\text{tr}_{\Sigma(t)} K = 0$, for each $t \in [0, \delta)$. Therefore $\theta_K^+ = \theta_K^- = 0$ on $\Sigma(t)$, for each $t \in [0, \delta)$.

As in the proof of Lemma 5.1, the first variation of θ_K^+ gives that $X_K = \nabla \ln \phi$ on $\Sigma(t)$. On the other hand, the first variation of $\theta_K^- = \theta_{-K}^+$ gives that $X_{-K} = \nabla \ln \phi$ on Σ_t . Proceeding as in the proof of Lemma 5.1, we obtain the following *local rigidity*:

- (Σ_0, g_0) is a flat torus, where g_0 is the metric on Σ_0 induced by g .
- (U, g) is isometric to $([0, \delta) \times \Sigma_0, dt^2 + g_0)$.
- $K = a dt^2$ on U , where a depends only on $t \in [0, \delta)$.
- $\mu_K = 0$ and $J_K = 0$ on U .

Observe that $\Sigma(t)$ is also volume minimizing in (M, g) . The assertion follows from this local rigidity as in the proof of Theorem 1.3. $\square$

6. Embedding of the Initial Data into a Quotient of Minkowski Space

In this section we show how, under the assumptions of Theorem 1.3, (M, g) can be isometrically embedded into a quotient of the Minkowski spacetime in a such way that K is exactly its second fundamental form. This, together with Theorem 1.3, characterizes the geometry—both intrinsic and extrinsic—of the initial data set (M, g, K) under natural conditions. The same holds under the assumptions of Theorem 5.2.

Theorem 6.1. *Assumptions as in Theorem 1.3 or Theorem 5.2. There is an isometric embedding of (M, g) into a quotient of Minkowski space in a such way that K is its second fundamental form.*

Consider the Minkowski spacetime $\mathbb{R}_1^n$ of dimension $n + 1$, i.e. $\mathbb{R} \times \mathbb{R} \times \mathbb{R}^{n-1}$ with the Lorentzian metric

$$g_M = -dt^2 + dr^2 + d\mathbf{x}^2$$

where $d\mathbf{x}^2$ is the standard Euclidean metric on $\mathbb{R}^{n-1}$.

Given a smooth function $t : \mathbb{R} \rightarrow \mathbb{R}$, define $r : \mathbb{R} \rightarrow \mathbb{R}$ by

$$r(s) = \int_0^s \sqrt{1 + t'(\sigma)^2} d\sigma.$$

Consider the spacelike hypersurface

$$N_0 = \{(t(s), r(s), x) : s \in \mathbb{R}, x \in \mathbb{R}^{n-1}\} \subset \mathbb{R}_1^n.$$

Note that (N_0, h_0) is isometric to $(\mathbb{R} \times \mathbb{R}^{n-1}, ds^2 + d\mathbf{x}^2)$ where h_0 is the metric induced by g_M . Straightforward calculations show that the second fundamental form of N in $\mathbb{R}_1^n$ with respect to $\frac{\partial}{\partial s}$ is given by $P = b ds^2$, where $b : \mathbb{R} \rightarrow \mathbb{R}$ is the function

$$b = \frac{t''}{\sqrt{1+t'^2}}.$$

Now, consider the hyperbolic space $\mathbb{H}^n$ of dimension n , that is, the n -manifold $\mathbb{R}_+ \times \mathbb{R}^{n-1}$ endowed with the metric

$$g_H = \frac{1}{x_0^2} (dx_0^2 + d\mathbf{x}^2).$$

Using the change of variables x_0 to $s = -\ln x_0$, we may write

$$g_H = ds^2 + e^{2s} d\mathbf{x}^2.$$

Thus $(N_1, h_1) = (\mathbb{R} \times \mathbb{R}^{n-1}, ds^2 + e^{2s} d\mathbf{x}^2)$ is isometric to hyperbolic space.

Consider the hypersurface

$$H = \{(t, r, x) \in \mathbb{R} \times \mathbb{R} \times \mathbb{R}^{n-1} : -t^2 + r^2 + |x|^2 = -1, t < 0\},$$

where $|x|^2 = x_1^2 + \dots + x_{n-1}^2$. Recall that H with the metric induced by g_M is isometric to $\mathbb{H}^n$ and that the second fundamental form of H in $\mathbb{R}_1^n$ with respect to the future directed unit normal is given by $-g_M|_H$.

Proof of Theorem 6.1. From Theorem 1.3 and Theorem 5.2, we know that (M, g) is isometric to $([0, \ell] \times \Sigma_0, ds^2 + e^{2\epsilon t} g_0)$, where (Σ_0, g_0) is a flat torus and $K = (1 - \epsilon)a dt^2 - \epsilon g$ for some function $a : [0, \ell] \rightarrow \mathbb{R}$. Therefore, (M, g) is isometric to a quotient of $([0, \ell] \times \mathbb{R}^{n-1}, ds^2 + e^{2\epsilon t} d\mathbf{x}^2)$. In the case where $\epsilon = 0$, we can take $t : [0, \ell] \rightarrow \mathbb{R}$ to be the solution of

$$\frac{t''}{\sqrt{1+t'^2}} = a$$

with initial condition $t(0) = 0$ and $t'(0) = 0$. Therefore, identifying (N_0, h_0) with $(\mathbb{R} \times \mathbb{R}^{n-1}, ds^2 + d\mathbf{x}^2)$, it follows from the above remarks that we can embed (M, g) into a quotient of $\mathbb{R}_1^n$ in a such way that the second fundamental form of M is given by $P = a ds^2 = K$. In the case where $\epsilon = 1$, it suffices to identify (N_1, h_1) with (H, h) , where h is the metric on H induced by g_M . $\square$

Acknowledgements. Michael Eichmair is supported by the START-Project Y963-N35 of the Austrian Science Fund. Gregory J. Galloway acknowledges the support of NSF Grant DMS-1710808. Abraão Mendes is grateful to the University of Miami where much of his work on this project was carried out. He was supported in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001. The authors would like to thank Dan A. Lee and Pengzi Miao for useful discussions on the topic of this paper.

Funding Open access funding provided by University of Vienna.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- Alaee, A., Lesourd, M., Yau, S.-T.: Stable Surfaces and Free Boundary Marginally Outer Trapped Surfaces (preprint). [arXiv:2009.07933](https://arxiv.org/abs/2009.07933) (2020)
- Andersson, L., Cai, M.-L., Galloway, G.J.: Rigidity and positivity of mass for asymptotically hyperbolic manifolds. *Ann. Henri Poincaré* **9**(1), 1–33 (2008)
- Andersson, L., Eichmair, M., Metzger, J.: Jang's equation and its applications to marginally trapped surfaces, Complex analysis and dynamical systems IV. Part 2, *Contemp. Math.*, vol. 554, Amer. Math. Soc., Providence, RI, pp. 13–45 (2011)
- Andersson, L., Mars, M., Simon, W.: Stability of marginally outer trapped surfaces and existence of marginally outer trapped tubes. *Adv. Theor. Math. Phys.* **12**(4), 853–888 (2008)
- Andersson, L., Metzger, J.: The area of horizons and the trapped region. *Commun. Math. Phys.* **290**(3), 941–972 (2009)
- Ashtekar, A., Galloway, G.J.: Some uniqueness results for dynamical horizons. *Adv. Theor. Math. Phys.* **9**(1), 1–30 (2005)
- Chruściel, P.T., Delay, E., Galloway, G.J., Howard, R.: Regularity of horizons and the area theorem. *Ann. Henri Poincaré* **2**(1), 109–178 (2001)
- Croke, C.B., Kleiner, B.: A warped product splitting theorem. *Duke Math. J.* **67**(3), 571–574 (1992)
- Eichmair, M.: The Plateau problem for marginally outer trapped surfaces. *J. Differ. Geom.* **83**(3), 551–583 (2009)
- Eichmair, M.: Existence, regularity, and properties of generalized apparent horizons. *Commun. Math. Phys.* **294**(3), 745–760 (2010)
- Eichmair, M., Huang, L.-H., Lee, D.A., Schoen, R.: The spacetime positive mass theorem in dimensions less than eight. *J. Eur. Math. Soc. (JEMS)* **18**(1), 83–121 (2016)
- Eichmair, M., Metzger, J.: Jenkins–Serrin-type results for the Jang equation. *J. Differ. Geom.* **102**(2), 207–242 (2016)
- Galloway, G.J.: Rigidity of outermost MOTS: the initial data version. *Gen. Relativity Gravit.* **50**(3), Art. 32, 7 (2018)
- Galloway, G.J., Jang, H.C.: Some scalar curvature warped product splitting theorems. *Proc. Am. Math. Soc.* **148**(6), 2617–2629 (2020)
- Galloway, G.J., Ó Murchadha, N.: Some remarks on the size of bodies and black holes. *Class. Quantum Gravity* **25**(10), 105009, 9 (2008)
- Galloway, G.J., Schoen, R.: A generalization of Hawking's black hole topology theorem to higher dimensions. *Commun. Math. Phys.* **266**(2), 571–576 (2006)
- Huang, L.-H., Lee, D.A.: Equality in the spacetime positive mass theorem. *Commun. Math. Phys.* **376**(3), 2379–2407 (2020)
- Lee, D.A.: *Geometric Relativity*, Graduate Studies in Mathematics, vol. 201. American Mathematical Society, Providence (2019)
- Lohkamp, J.: The Higher Dimensional Positive Mass Theorem II, preprint. [arXiv:1612.07505](https://arxiv.org/abs/1612.07505) (2016)
- Mendes, A.: Rigidity of marginally outer trapped (hyper)surfaces with negative σ -constant. *Trans. Am. Math. Soc.* **372**(8), 5851–5868 (2019)
- Petersen, P.: *Riemannian Geometry*, second ed. Graduate Texts in Mathematics, vol. 171. Springer, New York (2006)
- Schoen, R., Yau, S.-T.: Positive Scalar Curvature and Minimal Hypersurface Singularities, preprint. [arXiv:1704.05490](https://arxiv.org/abs/1704.05490) (2017)