




Contents lists available at ScienceDirect

Advances in Mathematics

www.elsevier.com/locate/aim



When are multidegrees positive?

Federico Castillo^a, Yairon Cid-Ruiz^a, Binglin Li^b,
Jonathan Montaña^{c,*}¹, Naizhen Zhang^d

^a Max Planck Institute for Mathematics in the Sciences, Inselstraße 22, 04103 Leipzig, Germany

^b Department of Statistics, 310 Herty Drive, University of Georgia, Athens, GA 30602, USA

^c Department of Mathematical Sciences, New Mexico State University, PO Box 30001, Las Cruces, NM 88003-8001, USA

^d Institut für Differentialgeometrie, Leibniz Universität Hannover, Welfengarten 1, 30167 Hannover, Germany



ARTICLE INFO

Article history:

Received 3 June 2020

Received in revised form 29 July 2020

Accepted 6 August 2020

Available online 26 August 2020

Communicated by Ezra Miller

MSC:

primary 14C17, 13H15

secondary 52B40, 13A30

Keywords:

Positivity

Multidegrees

Mixed multiplicities

Multiprojective scheme

ABSTRACT

Let $\mathbb{k}$ be an arbitrary field, $\mathbb{P} = \mathbb{P}_{\mathbb{k}}^{m_1} \times_{\mathbb{k}} \cdots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_p}$ be a multiprojective space over $\mathbb{k}$, and $X \subseteq \mathbb{P}$ be a closed subscheme of $\mathbb{P}$. We provide necessary and sufficient conditions for the positivity of the multidegrees of X . As a consequence of our methods, we show that when X is irreducible, the support of multidegrees forms a discrete algebraic polymatroid. In algebraic terms, we characterize the positivity of the mixed multiplicities of a standard multigraded algebra over an Artinian local ring, and we apply this to the positivity of mixed multiplicities of ideals. Furthermore, we use our results to recover several results

* Corresponding author.

E-mail addresses: Federico.Castillo@mis.mpg.de (F. Castillo), cidruiz@mis.mpg.de (Y. Cid-Ruiz), binglinligeometry@uga.edu (B. Li), jmon@nmsu.edu (J. Montaña), naizhen.zhang@math.uni-hannover.de (N. Zhang).

URL: <https://ycid.github.io> (Y. Cid-Ruiz).

¹ The fourth author is supported by National Science Foundation Grant DMS #2001645.

Polymatroids
Hilbert polynomial

in the literature in the context of combinatorial algebraic
geometry.

© 2020 Elsevier Inc. All rights reserved.

1. Introduction

Let $\mathbb{k}$ be an arbitrary field, $\mathbb{P} = \mathbb{P}_{\mathbb{k}}^{m_1} \times_{\mathbb{k}} \cdots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_p}$ be a multiprojective space over $\mathbb{k}$, and $X \subseteq \mathbb{P}$ be a closed subscheme of $\mathbb{P}$. The *multidegrees* of X are fundamental invariants that describe algebraic and geometric properties of X . For each $\mathbf{n} = (n_1, \dots, n_p) \in \mathbb{N}^p$ with $n_1 + \cdots + n_p = \dim(X)$ one can define the *multidegree of X of type $\mathbf{n}$ with respect to $\mathbb{P}$* , denoted by $\deg_{\mathbb{P}}^{\mathbf{n}}(X)$, in different ways (see Definition 2.7, Remark 2.8 and Remark 2.9). In classical geometrical terms, when $\mathbb{k}$ is algebraically closed, $\deg_{\mathbb{P}}^{\mathbf{n}}(X)$ equals the number of points (counting multiplicity) in the intersection of X with the product $L_1 \times_{\mathbb{k}} \cdots \times_{\mathbb{k}} L_p \subset \mathbb{P}$, where $L_i \subset \mathbb{P}_{\mathbb{k}}^{m_i}$ is a general linear subspace of dimension $m_i - n_i$ for each $1 \leq i \leq p$.

The study of multidegrees goes back to pioneering work by van der Waerden [60]. From a more algebraic point of view, multidegrees receive the name of *mixed multiplicities* (see Definition 2.7). More recent papers where the notion of multidegree (or mixed multiplicity) is studied are, e.g., [1,9,11,23,33,36,41,42,57].

The main goal of this paper is to answer the following fundamental question considered by Trung [57] and by Huh [25] in the case $p = 2$.

- For $\mathbf{n} \in \mathbb{N}^p$ with $n_1 + \cdots + n_p = \dim(X)$, when do we have that $\deg_{\mathbb{P}}^{\mathbf{n}}(X) > 0$?

Our main result says that the positivity of $\deg_{\mathbb{P}}^{\mathbf{n}}(X)$ is determined by the dimensions of the images of the natural projections from $\mathbb{P}$ restricted to the irreducible components of X . First, we set a basic notation: for each $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq \{1, \dots, p\}$, let $\Pi_{\mathfrak{J}}$ be the natural projection

$$\Pi_{\mathfrak{J}} : \mathbb{P} = \mathbb{P}_{\mathbb{k}}^{m_1} \times_{\mathbb{k}} \cdots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_p} \rightarrow \mathbb{P}_{\mathbb{k}}^{m_{j_1}} \times_{\mathbb{k}} \cdots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_{j_k}}.$$

The following is the main theorem of this article. Here, we give necessary and sufficient conditions for the positivity of multidegrees.

Theorem A (Theorem 3.12, Corollary 3.13). *Let $\mathbb{k}$ be an arbitrary field, $\mathbb{P} = \mathbb{P}_{\mathbb{k}}^{m_1} \times_{\mathbb{k}} \cdots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_p}$ be a multiprojective space over $\mathbb{k}$, and $X \subseteq \mathbb{P}$ be a closed subscheme of $\mathbb{P}$. Let $\mathbf{n} = (n_1, \dots, n_p) \in \mathbb{N}^p$ be such that $n_1 + \cdots + n_p = \dim(X)$. Then, $\deg_{\mathbb{P}}^{\mathbf{n}}(X) > 0$ if and only if there is an irreducible component $Y \subseteq X$ of X that satisfies the following two conditions:*

- $\dim(Y) = \dim(X)$.
- For each $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq \{1, \dots, p\}$ the inequality

$$n_{j_1} + \cdots + n_{j_k} \leq \dim(\Pi_{\mathfrak{J}}(Y))$$

holds.

When $\mathbb{k}$ is the field of complex numbers Theorem A is essentially covered by the geometric results in [34, Theorems 2.14, 2.19],² however their methods do not extend to arbitrary fields. Here we follow an algebraic approach that allows us to prove the result for all fields, and hence a general version for algebras over Artinian local rings (see Theorem B). The main idea in the proof of Theorem A is the study of the dimensions of the images of the natural projections after cutting by a general hyperplane (see Theorem 3.7).

We note that if $p = 2$ and X is arithmetically Cohen-Macaulay, the conclusion of Theorem A in the irreducible case also holds for X (see [57, Corollary 2.8]). In Example 5.2 we show that this is not necessarily true for $p > 2$.

If X is irreducible, then the function $r : 2^{\{1, \dots, p\}} \rightarrow \mathbb{Z}$ defined by $r(\mathfrak{J}) := \dim(\Pi_{\mathfrak{J}}(Y))$ is a submodular function, i.e., $r(\mathfrak{J}_1 \cap \mathfrak{J}_2) + r(\mathfrak{J}_1 \cup \mathfrak{J}_2) \leq r(\mathfrak{J}_1) + r(\mathfrak{J}_2)$ for any two subsets $\mathfrak{J}_1, \mathfrak{J}_2 \subseteq \{1, \dots, p\}$, as proved in Proposition 5.1 (see also Definition 2.16). By the Submodular Theorem (see, e.g., [7, Theorem 3.11] or [44, Appendix B]) and the inequalities of Theorem A, the points $\mathbf{n} \in \mathbb{N}^p$ for which $\deg_{\mathbb{P}}^{\mathbf{n}}(X) > 0$ are the lattice points of a *generalized permutohedron*. Defined by A. Postnikov in [48] generalized permutohedra are polytopes obtained by deforming usual permutohedra. In recent years this family of polytopes has been studied in relation to other fields such as probability, combinatorics, and representation theory (see [44, 45, 49]).

In a more algebraic flavor, we state the translation of Theorem A to the mixed multiplicities of a standard multigraded algebra over an Artinian local ring (see Definition 2.13).

Theorem B (Corollary 3.14). *Let A be an Artinian local ring and R be a finitely generated standard $\mathbb{N}^p$ -graded A -algebra. For each $1 \leq j \leq p$, let $\mathfrak{m}_j \subset R$ be the ideal generated by the elements of degree $\mathbf{e}_j$, where $\mathbf{e}_j \in \mathbb{N}^p$ denotes the j -th elementary vector. Let $\mathfrak{N} = \mathfrak{m}_1 \cap \dots \cap \mathfrak{m}_p \subset R$. Let $\mathbf{n} = (n_1, \dots, n_p) \in \mathbb{N}^p$ be such that $n_1 + \dots + n_p = \dim(R/(0 :_R \mathfrak{N}^\infty)) - p$. Then, $e(\mathbf{n}; R) > 0$ if and only if there is a minimal prime ideal $\mathfrak{P} \in \text{Min}(0 :_R \mathfrak{N}^\infty)$ of $(0 :_R \mathfrak{N}^\infty)$ that satisfies the following two conditions:*

- (a) $\dim(R/\mathfrak{P}) = \dim(R/(0 :_R \mathfrak{N}^\infty))$.
- (b) For each $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq \{1, \dots, p\}$ the inequality

$$n_{j_1} + \dots + n_{j_k} \leq \dim\left(\frac{R}{\mathfrak{P} + \sum_{j \notin \mathfrak{J}} \mathfrak{m}_j}\right) - k$$

holds.

² In Remark 3.15 we briefly discuss how (over the complex numbers) Theorem A can be obtained by using the results in [34, §2.2].

For a given finite set of ideals in a Noetherian local ring, such that one of them is zero-dimensional, we can define their mixed multiplicities by considering a certain associated standard multigraded algebra (see [58] for more information). These multiplicities have a long history of interconnecting problems from commutative algebra, algebraic geometry, and combinatorics, with applications to the topics of Milnor numbers, mixed volumes, and integral dependence (see, e.g., [25,27,55,58]). As a direct consequence of Theorem B we are able to give a characterization for the positivity of mixed multiplicities of ideals (see Corollary 4.3). In another related result, we focus on homogeneous ideals generated in one degree; this case is of particular importance due to its relation with rational maps between projective varieties. In this setting, we provide more explicit conditions for positivity in terms of the analytic spread of products of these ideals (see Theorem 4.4).

Going back to the setting of Theorem A, we switch our attention to the following discrete set

$$\text{MSupp}_{\mathbb{P}}(X) = \{\mathbf{n} \in \mathbb{N}^p \mid \deg_{\mathbb{P}}^{\mathbf{n}}(X) > 0\},$$

which we call the *support of X with respect to P*. When X is irreducible, we show that $\text{MSupp}_{\mathbb{P}}(X)$ is a (*discrete*) *polymatroid* (see §2.3, Proposition 5.1). The latter result was included in an earlier version of this paper when $\mathbb{k}$ is algebraically closed, and an alternative proof is given by Brändén and Huh in [3, Corollary 4.7] using the theory of Lorentzian polynomials. An advantage of our approach is that we can describe the corresponding rank submodular functions of the polymatroids, a fact that we exploit in the applications of Section 6. Additionally, our results are valid when X is just irreducible and not necessarily geometrically irreducible over $\mathbb{k}$ (i.e., we do not need to assume that $X \times_{\mathbb{k}} \bar{\mathbb{k}}$ is irreducible for an algebraic closure $\bar{\mathbb{k}}$ of $\mathbb{k}$); it should be noticed that this generality is not covered by the statements in [3] and [34].

Discrete polymatroids [24] have also been studied under the name of M-convex sets [46]. Polymatroids can also be described as the integer points in a generalized permutohedron [48], so they are closely related to submodular functions, which are well studied in optimization, see [38] and [52, Part IV] for comprehensive surveys on submodular functions, their applications, and their history. There are two distinguishable types of polymatroids, linear and algebraic polymatroids, whose main properties are inherited by their representation in terms of other algebraic structures. Theorem A allows us to define another type of polymatroids, that we call *Chow polymatroids*, and which interestingly lies in between the other two. In the following theorem we summarize our main results in this direction.

Theorem C (Theorem 5.5). *Over an arbitrary field $\mathbb{k}$, we have the following inclusions of families of polymatroids*

$$\left(\text{Linear polymatroids} \right) \subseteq \left(\text{Chow polymatroids} \right) \subseteq \left(\text{Algebraic polymatroids} \right).$$

Moreover, when $\mathbb{k}$ is a field of characteristic zero, the three families coincide.

If $\mathbb{k}$ has positive characteristic, then these types of polymatroids do not agree. In fact, there exist examples of polymatroids which are algebraic over any field of positive characteristic but never linear (see Remark 5.7).

Theorem A can be applied to particular examples of varieties coming from combinatorial algebraic geometry. In §6.1 we do so to matrix Schubert varieties; in this case the multidegrees are the coefficients of Schubert polynomials, thus our results allow us to give an alternative proof to a recent conjecture regarding the support of these polynomials (see Theorem 6.3). In §6.2 and §6.3 we study certain embeddings of flag varieties and of the moduli space $\overline{\mathcal{M}}_{0,p+3}$, respectively (see Proposition 6.7 and Proposition 6.8). In §6.4 we recover a well-known characterization for the positivity of mixed volumes of convex bodies (see Theorem 6.9).

We now outline the contents of the article. In Section 2 we set up the notation used throughout the document. We also include key preliminary definitions and results, paying special attention to the connection between mixed multiplicities of standard multigraded graded algebras and multidegrees of their corresponding schemes. Section 3 is devoted to the proof of Theorem A and Theorem B. Our results for mixed multiplicities of ideals are included in Section 4. In Section 5 we relate our results to the theory of polymatroids. In particular, we show the proof of Theorem C. We finish the paper with Section 6 where the applications to combinatorial algebraic geometry are presented.

We conclude the Introduction with an illustrative example. The following example is constructed following the same ideas in Proposition 5.4.


Example 1.1. Consider the polynomial ring $S = \mathbb{k}[v_1, v_2, v_3][w_1, w_2, w_3]$ with the $\mathbb{N}^3$ -grading $\deg(v_i) = (0, 0, 0)$, $\deg(w_i) = \mathbf{e}_i$ for $1 \leq i \leq 3$. Let T be the $\mathbb{N}^3$ -graded polynomial ring $T = \mathbb{k}[x_0, \dots, x_3][y_0, \dots, y_3][z_0, \dots, z_3]$ where $\deg(x_i) = \mathbf{e}_1$, $\deg(y_i) = \mathbf{e}_2$ and $\deg(z_i) = \mathbf{e}_3$. Consider the $\mathbb{N}^3$ -graded $\mathbb{k}$ -algebra homomorphism

$$\begin{aligned} \varphi = T \rightarrow S, \quad & \begin{array}{llll} x_0 \mapsto w_1, & x_1 \mapsto v_1 w_1, & x_2 \mapsto v_1 w_1, & x_3 \mapsto v_1 w_1, \\ y_0 \mapsto w_2, & y_1 \mapsto v_1 w_2, & y_2 \mapsto v_2 w_2, & y_3 \mapsto (v_1 + v_2) w_2, \\ z_0 \mapsto w_3, & z_1 \mapsto v_1 w_3, & z_2 \mapsto v_2 w_3, & z_3 \mapsto v_3 w_3. \end{array} \end{aligned}$$

Note that $\mathfrak{P} = \text{Ker}(\varphi) \subset T$ is an $\mathbb{N}^3$ -graded prime ideal. Let $Y \subset \mathbb{P} = \mathbb{P}_{\mathbb{k}}^3 \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^3 \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^3$ be the closed subscheme corresponding to $\mathfrak{P}$. In this case, one can easily compute the dimension of the projections $\Pi_{\mathfrak{J}}(Y)$ for each $\mathfrak{J} \subseteq \{1, 2, 3\}$, and so Theorem A implies that $\text{MSupp}_{\mathbb{P}}(Y)$ is given by all $\mathbf{n} = (n_1, \dots, n_3) \in \mathbb{N}^3$ satisfying the following conditions:

$$\begin{aligned} n_1 + n_2 + n_3 &= 3 = \dim(Y), \\ n_1 + n_2 &\leq 2 = \dim(\Pi_{\{1,2\}}(Y)), \quad n_1 + n_3 \leq 3 = \dim(\Pi_{\{1,3\}}(Y)), \\ n_2 + n_3 &\leq 3 = \dim(\Pi_{\{2,3\}}(Y)), \\ n_1 &\leq 1 = \dim(\Pi_{\{1\}}(Y)), \quad n_2 \leq 2 = \dim(\Pi_{\{2\}}(Y)), \quad n_3 \leq 3 = \dim(\Pi_{\{3\}}(Y)). \end{aligned}$$

Hence $\text{MSupp}_{\mathbb{P}}(Y) = \{(0, 0, 3), (0, 1, 2), (0, 2, 1), (1, 0, 2), (1, 1, 1)\} \subset \mathbb{N}^3$. This set can also be represented graphically as follows:



Additionally, by using `Macaulay2` [21] we can compute that its *multidegree polynomial* (see Definition 2.10) is equal to:

$$\deg_{\mathbb{P}}(Y; t_1, t_2, t_3) = t_1^3 t_2^3 + t_1^3 t_2^2 t_3 + t_1^3 t_2 t_3^2 + t_1^2 t_2^3 t_3 + t_1^2 t_2^2 t_3^2.$$

We note that here we are following the convention that $\text{MSupp}_{\mathbb{P}}(Y)$ is given by the complementary degrees of the polynomial $\deg_{\mathbb{P}}(Y; t_1, t_2, t_3)$; for instance, the term $t_1^3 t_2^3$ corresponds to the point $(3, 3, 3) - (3, 3, 0) = (0, 0, 3) \in \text{MSupp}_{\mathbb{P}}(Y)$.

2. Notation and preliminaries

In this section, we set up the notation that is used throughout the paper. We also present some preliminary results needed in the proofs of our main theorems.

Let $p \geq 1$ be a positive integer. If $\mathbf{n} = (n_1, \dots, n_p), \mathbf{m} = (m_1, \dots, m_p) \in \mathbb{Z}^p$ are two multi-indexes, we write $\mathbf{n} \geq \mathbf{m}$ whenever $n_i \geq m_i$ for all $1 \leq i \leq p$, and $\mathbf{n} > \mathbf{m}$ whenever $n_j > m_j$ for all $1 \leq j \leq p$. For each $1 \leq i \leq p$, let $\mathbf{e}_i \in \mathbb{N}^p$ be the i -th elementary vector $\mathbf{e}_i = (0, \dots, 1, \dots, 0)$. Let $\mathbf{0} \in \mathbb{N}^p$ and $\mathbf{1} \in \mathbb{N}^p$ be the vectors $\mathbf{0} = (0, \dots, 0)$ and $\mathbf{1} = (1, \dots, 1)$ of p copies of 0 and 1, respectively. For any $\mathbf{n} = (n_1, \dots, n_p) \in \mathbb{Z}^p$, we define its weight as $|\mathbf{n}| := n_1 + \dots + n_p$. Let $[p]$ denote the set $[p] := \{1, \dots, p\}$.

For clarity of exposition we first introduce the main concepts in the theory of multidegrees over an arbitrary field. Later, we also work over Artinian local rings; we highlight important details in this more general setting in §2.2.

2.1. The case over a field

We begin by introducing a general setup for Theorem A and its preparatory results.

Setup 2.1. Let k be an arbitrary field. Let R be a finitely generated standard $\mathbb{N}^p$ -graded algebra over k , that is, $[R]_0 = k$ and R is finitely generated over k by elements of degree $\mathbf{e}_i$ with $1 \leq i \leq p$. For each subset $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq [p] = \{1, \dots, p\}$ denote by $R_{(\mathfrak{J})}$ the standard $\mathbb{N}^k$ -graded k -algebra given by

$$R_{(\mathfrak{J})} := \bigoplus_{\substack{i_1 \geq 0, \dots, i_p \geq 0 \\ i_j = 0 \text{ if } j \notin \mathfrak{J}}} [R]_{(i_1, \dots, i_p)};$$

for instance, for each $1 \leq j \leq p$, $R_{(j)}$ denotes the standard $\mathbb{N}$ -graded $\mathbb{k}$ -algebra $R_{(j)} := \bigoplus_{k \geq 0} [R]_{k \cdot e_j}$. For each $1 \leq j \leq p$, let $\mathfrak{m}_j \subset R$ be the ideal $\mathfrak{m}_j := ([R]_{e_j})$. Let $\mathfrak{N} \subset R$ be the multigraded irrelevant ideal $\mathfrak{N} := \mathfrak{m}_1 \cap \dots \cap \mathfrak{m}_p$. For each $\mathfrak{J} \subseteq [p]$, let $\mathfrak{N}_{\mathfrak{J}} \subset R_{(\mathfrak{J})}$ be the corresponding multigraded irrelevant ideal $\mathfrak{N}_{\mathfrak{J}} := \left(\bigcap_{j \in \mathfrak{J}} \mathfrak{m}_j \right) \cap R_{(\mathfrak{J})}$. Let X be the multiprojective scheme $X := \text{MultiProj}(R)$ (see Definition 2.2 below) and $X_{\mathfrak{J}}$ be the multiprojective scheme $X_{\mathfrak{J}} := \text{MultiProj}(R_{(\mathfrak{J})})$ for each $\mathfrak{J} \subseteq [p]$. To avoid trivial situations, we always assume that $X \neq \emptyset$.

Definition 2.2. The multiprojective scheme $\text{MultiProj}(R)$ is given by $\text{MultiProj}(R) := \{\mathfrak{P} \in \text{Spec}(R) \mid \mathfrak{P} \text{ is } \mathbb{N}^p\text{-graded and } \mathfrak{P} \not\supseteq \mathfrak{N}\}$, and its scheme structure is obtained by using multi-homogeneous localizations (see, e.g., [28, §1]).

The inclusion $R_{(\mathfrak{J})} \hookrightarrow R$ induces the natural projection

$$\Pi_{\mathfrak{J}} : X \rightarrow X_{\mathfrak{J}}, \quad \mathfrak{P} \in X \mapsto \mathfrak{P} \cap R_{(\mathfrak{J})} \in X_{\mathfrak{J}}.$$

We embed X as a closed subscheme of a multiprojective space $\mathbb{P} := \mathbb{P}_{\mathbb{k}}^{m_1} \times_{\mathbb{k}} \dots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_p}$. Then, for each $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq [p]$, $\Pi_{\mathfrak{J}} : X \rightarrow X_{\mathfrak{J}}$ corresponds with the restriction to X and to $X_{\mathfrak{J}}$ of the natural projection

$$\Pi_{\mathfrak{J}} : \mathbb{P} \rightarrow \mathbb{P}_{\mathbb{k}}^{m_{j_1}} \times_{\mathbb{k}} \dots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_{j_k}},$$

and $X_{\mathfrak{J}}$ becomes a closed subscheme of $\mathbb{P}_{\mathbb{k}}^{m_{j_1}} \times_{\mathbb{k}} \dots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_{j_k}}$.

For any multi-homogeneous element $x \in R$, the closed subscheme $\text{MultiProj}(R/xR) \subseteq X$ is denoted by $X \cap V(x)$.

Notation 2.3. From now on, $\mathfrak{J} = \{j_1, \dots, j_k\}$ denotes a subset of $[p]$. Set $r := \dim(X)$ and $r(\mathfrak{J}) := \dim(\Pi_{\mathfrak{J}}(X))$ for each $\mathfrak{J} \subseteq [p]$. For a singleton set $\{i\} \subseteq [p]$, $r(\{i\})$ and $\Pi_{\{i\}}$ are simply denoted by $r(i)$ and Π_i , respectively.

Note that the image of $\Pi_{\mathfrak{J}} : X \rightarrow X_{\mathfrak{J}}$ can be described by the following isomorphism

$$\Pi_{\mathfrak{J}}(X) \cong \text{MultiProj} \left(\frac{R_{(\mathfrak{J})}}{R_{(\mathfrak{J})} \cap (0 :_R \mathfrak{N}^\infty)} \right). \quad (1)$$

Remark 2.4. Since $R = R_{(\mathfrak{J})} \oplus \left(\sum_{j \notin \mathfrak{J}} \mathfrak{m}_j \right)$, we obtain a natural isomorphism $R_{(\mathfrak{J})} \cong \frac{R}{\sum_{j \notin \mathfrak{J}} \mathfrak{m}_j}$ of $\mathbb{N}^k$ -graded $\mathbb{k}$ -algebras where $k = |\mathfrak{J}|$.

We now provide some preparatory results.

Lemma 2.5. Under Setup 2.1, the following statements hold:

- (i) $r = \dim(X) = \dim(R/(0 :_R \mathfrak{N}^\infty)) - p$.
- (ii) There is an isomorphism

$$\Pi_{\mathfrak{J}}(X) \cong \text{MultiProj} \left(R / \left((0 :_R \mathfrak{N}^\infty) + \sum_{j \notin \mathfrak{J}} \mathfrak{m}_j \right) \right). \quad (2)$$

- (iii) If $(0 :_R \mathfrak{N}^\infty) = 0$, then $\Pi_{\mathfrak{J}}(X) \cong \text{MultiProj}(R_{(\mathfrak{J})}) = X_{\mathfrak{J}}$.
- (iv) If $(0 :_R \mathfrak{N}^\infty) = 0$, then $(0 :_{R_{(\mathfrak{J})}} \mathfrak{N}_{\mathfrak{J}}^\infty) = 0$.

Proof. (i) This formula follows from [28, Lemma 1.2] (also, see [10, Corollary 3.5]).

(ii) From the natural maps $R_{(\mathfrak{J})} \hookrightarrow R \twoheadrightarrow R/(0 :_R \mathfrak{N}^\infty)$, we obtain a natural isomorphism

$$R_{(\mathfrak{J})} / (R_{(\mathfrak{J})} \cap (0 :_R \mathfrak{N}^\infty)) \xrightarrow{\cong} (R/(0 :_R \mathfrak{N}^\infty))_{(\mathfrak{J})}.$$

By using Remark 2.4 it follows that $(R/(0 :_R \mathfrak{N}^\infty))_{(\mathfrak{J})} \cong R / \left((0 :_R \mathfrak{N}^\infty) + \sum_{j \notin \mathfrak{J}} \mathfrak{m}_j \right)$. Therefore, the claimed isomorphism is obtained from (1).

- (iii) It follows directly from part (ii) and Remark 2.4.
- (iv) This part is clear. $\square$

Let $P_R(\mathbf{t}) = P_R(t_1, \dots, t_p) \in \mathbb{Q}[\mathbf{t}] = \mathbb{Q}[t_1, \dots, t_p]$ be the *Hilbert polynomial* of R (see, e.g., [23, Theorem 4.1], [9, Theorem 3.4]). Then, the degree of P_R is equal to r and

$$P_R(\mathbf{v}) = \dim_{\mathbb{k}}([R]_{\mathbf{v}})$$

for all $\mathbf{v} \in \mathbb{N}^p$ such that $\mathbf{v} \gg \mathbf{0}$. Furthermore, if we write

$$P_R(\mathbf{t}) = \sum_{n_1, \dots, n_p \geq 0} e(n_1, \dots, n_p) \binom{t_1 + n_1}{n_1} \cdots \binom{t_p + n_p}{n_p}, \quad (3)$$

then $0 \leq e(n_1, \dots, n_r) \in \mathbb{Z}$ for all $n_1 + \cdots + n_p = r$.

Remark 2.6. The following are basic properties of Hilbert polynomials.

- (i) Since $\mathfrak{N}^k(0 :_R \mathfrak{N}^\infty) = 0$ for $k \gg 0$ we have $\dim_{\mathbb{k}}([R]_{\mathbf{v}}) = \dim_{\mathbb{k}}([R/(0 :_R \mathfrak{N}^\infty)]_{\mathbf{v}})$ for $\mathbf{v} \gg \mathbf{0}$. Thus,

$$P_R(\mathbf{t}) = P_{R/(0 :_R \mathfrak{N}^\infty)}(\mathbf{t}).$$

- (ii) Let $\mathbb{L}$ be a field extension of $\mathbb{k}$. Then, $\mathbf{R} \otimes_{\mathbb{k}} \mathbb{L}$ is a finitely generated standard $\mathbb{N}^p$ -graded $\mathbb{L}$ -algebra and $\dim_{\mathbb{L}} ([\mathbf{R} \otimes_{\mathbb{k}} \mathbb{L}]_{\mathbf{v}}) = \dim_{\mathbb{k}} ([\mathbf{R}]_{\mathbf{v}})$ for all $\mathbf{v} \in \mathbb{N}^p$. Thus,

$$P_{\mathbf{R} \otimes_{\mathbb{k}} \mathbb{L}}(\mathbf{t}) = P_{\mathbf{R}}(\mathbf{t}).$$

In particular, one can always assume $\mathbb{k}$ is an infinite field (for instance, we can substitute $\mathbb{k}$ by a purely transcendental field extension $\mathbb{k}(\xi)$).

Under the notation of (3) we define the following invariants.

Definition 2.7. Let $\mathbf{n} = (n_1, \dots, n_p) \in \mathbb{N}^p$ with $|\mathbf{n}| = r$. Then:

- (i) $e(\mathbf{n}, \mathbf{R}) := e(n_1, \dots, n_p)$ is the *mixed multiplicity of $\mathbf{R}$ of type $\mathbf{n}$* .
(ii) $\deg_{\mathbb{P}}^{\mathbf{n}}(X) := e(\mathbf{n}, \mathbf{R})$ is the *multidegree of $X = \text{MultiProj}(\mathbf{R})$ of type $\mathbf{n}$ with respect to $\mathbb{P}$* .

As stated in the Introduction, in classical geometrical terms, when $\mathbb{k}$ is algebraically closed, $\deg_{\mathbb{P}}^{\mathbf{n}}(X)$ is also equal to the number of points (counting multiplicity) in the intersection of X with the product $L_1 \times_{\mathbb{k}} \cdots \times_{\mathbb{k}} L_p \subset \mathbb{P}$, where $L_i \subseteq \mathbb{P}_{\mathbb{k}}^{m_i}$ is a general linear subspace of dimension $m_i - n_i$ for each $1 \leq i \leq p$ (see [60], [9, Theorem 4.7]).

The multidegrees of X can be defined easily in terms of Chow rings and in terms of Hilbert series.

Remark 2.8. The Chow ring of $\mathbb{P} = \mathbb{P}_{\mathbb{k}}^{m_1} \times_{\mathbb{k}} \cdots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_p}$ is given by

$$A^*(\mathbb{P}) = \frac{\mathbb{Z}[H_1, \dots, H_p]}{\left(H_1^{m_1+1}, \dots, H_p^{m_p+1}\right)}$$

where H_i represents the class of the inverse image of a hyperplane of $\mathbb{P}_{\mathbb{k}}^{m_i}$ under the natural projection $\Pi_i : \mathbb{P} \rightarrow \mathbb{P}_{\mathbb{k}}^{m_i}$. Then, the class of the cycle associated to X coincides with

$$[X] = \sum_{\substack{0 \leq n_i \leq m_i \\ |\mathbf{n}|=r}} \deg_{\mathbb{P}}^{\mathbf{n}}(X) H_1^{m_1-n_1} \cdots H_p^{m_p-n_p} \in A^*(\mathbb{P}).$$

Remark 2.9. By considering the Hilbert series $\text{Hilb}_{\mathbf{R}}(t_1, \dots, t_p) := \sum_{\mathbf{v} \in \mathbb{N}^p} \dim_{\mathbb{k}} ([\mathbf{R}]_{\mathbf{v}}) t_1^{v_1} \cdots t_p^{v_p}$ of $\mathbf{R}$, one can analogously define the notions of mixed multiplicities and multidegrees (see [43, §8.5], [9, Theorem A]). Here we quickly derive this analogous definition because we shall use it in §6.1. Let $S = \mathbb{k}[x_{1,0}, x_{1,1}, \dots, x_{1,m_1}] \otimes_{\mathbb{k}} \cdots \otimes_{\mathbb{k}} \mathbb{k}[x_{p,0}, x_{p,1}, \dots, x_{p,m_p}]$ be the multigraded polynomial ring corresponding with $\mathbb{P} = \mathbb{P}_{\mathbb{k}}^{m_1} \times_{\mathbb{k}} \cdots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_p}$, that is $\mathbb{P} = \text{MultiProj}(S)$. By considering an S -free resolution of $\mathbf{R}$, we can write

$$\text{Hilb}_R(\mathbf{t}) = \mathcal{K}(R; \mathbf{t}) / (\mathbf{1} - \mathbf{t})^{m+1} = \mathcal{K}(R; \mathbf{t}_1, \dots, \mathbf{t}_p) / \prod_{i=1}^p (1 - \mathbf{t}_i)^{m_i+1},$$

where $\mathcal{K}(R; \mathbf{t})$ is called the *K-polynomial* of R (see [43, Definition 8.21]). Let $\mathcal{C}(R; \mathbf{t}) \in \mathbb{Z}[\mathbf{t}_1, \dots, \mathbf{t}_p]$ be the sum of all the terms in $\mathcal{K}(R; \mathbf{1} - \mathbf{t})$ of total degree equal to $\dim(S) - \dim(R)$ (see [43, Definition 8.45]). Then, if $(0 :_R \mathfrak{N}^\infty) = 0$, we obtain the equality

$$\mathcal{C}(R; \mathbf{t}) = \sum_{\substack{0 \leq n_i \leq m_i \\ |\mathbf{n}|=r}} \deg_{\mathbb{P}}^{\mathbf{n}}(X) \mathbf{t}_1^{m_1-n_1} \dots \mathbf{t}_p^{m_p-n_p}.$$

Proof. From [9, Theorem A(I)] we have $\text{Hilb}_R(\mathbf{t}) = \sum_{|\mathbf{k}|=\dim(R)} Q_{\mathbf{k}}(\mathbf{t}) / (\mathbf{1} - \mathbf{t})^{\mathbf{k}}$ where $Q_{\mathbf{k}}(\mathbf{t}) \in \mathbb{Z}[\mathbf{t}]$. The assumption $(0 :_R \mathfrak{N}^\infty) = 0$ gives that $\dim(R) = r + p$ (see Lemma 2.5(i)). Hence, by using [9, Theorem A(II,III)] we obtain that $\deg_{\mathbb{P}}^{\mathbf{n}}(X) = e(\mathbf{n}; R) = Q_{\mathbf{n}+1}(\mathbf{1})$ for all $|\mathbf{n}| = r$. Also, the assumption $(0 :_R \mathfrak{N}^\infty) = 0$ and [9, Theorem 2.8(ii)] imply that $Q_{\mathbf{k}}(\mathbf{1}) = 0$ when $|\mathbf{k}| = \dim(R)$ and $\mathbf{k}_i = 0$ for some $1 \leq i \leq p$.

After writing $\text{Hilb}_R(\mathbf{t}) = \sum_{|\mathbf{k}|=\dim(R)} Q_{\mathbf{k}}(\mathbf{t})(\mathbf{1} - \mathbf{t})^{m+1-\mathbf{k}} / (\mathbf{1} - \mathbf{t})^{m+1}$, we obtain the equality

$$\mathcal{K}(R; \mathbf{t}) = \sum_{|\mathbf{k}|=\dim(R)} Q_{\mathbf{k}}(\mathbf{t})(\mathbf{1} - \mathbf{t})^{m+1-\mathbf{k}}.$$

Making the substitution $\mathbf{t}_i \mapsto (1 - \mathbf{t}_i)$ and choosing the terms of total degree $\dim(S) - \dim(R) = \sum_{i=1}^p m_i - r$, it follows that $\mathcal{C}(R; \mathbf{t}) = \sum_{|\mathbf{k}|=\dim(R)} Q_{\mathbf{k}}(\mathbf{1}) \mathbf{t}^{m+1-\mathbf{k}} = \sum_{|\mathbf{n}|=r} Q_{\mathbf{n}+1}(\mathbf{1}) \mathbf{t}^{m-\mathbf{n}}$. So, the result is clear. $\square$

Although in the proofs of Theorem A and Theorem B we do not exploit the fact that multidegrees can be defined as in Remark 2.8, we do encode the multidegrees in a homogeneous polynomial that mimics the cycle associated to X in the Chow ring $A^*(\mathbb{P})$. The following objects are the main focus of this paper.

Definition 2.10. Let $X \subseteq \mathbb{P} = \mathbb{P}_{\mathbb{k}}^{m_1} \times_{\mathbb{k}} \dots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_p}$ be a closed subscheme with $r = \dim(X)$. We denote the *multidegree polynomial of X with respect to $\mathbb{P}$* as the homogeneous polynomial

$$\deg_{\mathbb{P}}(X; \mathbf{t}_1, \dots, \mathbf{t}_p) := \sum_{\substack{0 \leq n_i \leq m_i \\ |\mathbf{n}|=r}} \deg_{\mathbb{P}}^{\mathbf{n}}(X) \mathbf{t}_1^{m_1-n_1} \dots \mathbf{t}_p^{m_p-n_p} \in \mathbb{N}[\mathbf{t}_1, \dots, \mathbf{t}_p]$$

of degree $m_1 + \dots + m_p - r$. We say that the *support of X with respect to $\mathbb{P}$* is given by

$$\text{MSupp}_{\mathbb{P}}(X) := \{\mathbf{n} \in \mathbb{N}^p \mid \deg_{\mathbb{P}}^{\mathbf{n}}(X) > 0\}.$$

Remark 2.11. Note that under the assumption $(0 :_R \mathfrak{N}^\infty) = 0$ we obtain the equality $\deg_{\mathbb{P}}(X; \mathbf{t}) = \mathcal{C}(R; \mathbf{t})$.

2.2. The case over an Artinian local ring

In this subsection, we show how the mixed multiplicities are defined for a standard multigraded algebra over an Artinian local ring.

Setup 2.12. Keep the notations and assumptions introduced in Setup 2.1 and now substitute the field $\mathbb{k}$ by an Artinian local ring A .

In this setting, the notion of mixed multiplicities is defined essentially in the same way as in Definition 2.7.

Definition 2.13. Let $P_R(\mathbf{t}) = P_R(t_1, \dots, t_p) \in \mathbb{Q}[\mathbf{t}] = \mathbb{Q}[t_1, \dots, t_p]$ be the *Hilbert polynomial* of R (see, e.g., [23, Theorem 4.1], [9, Theorem 3.4]). Then, as before, the degree of P_R is equal to $\dim(R/(0 :_R \mathfrak{N}^\infty)) - p$ and

$$P_R(\mathbf{v}) = \text{length}_A([R]_{\mathbf{v}})$$

for all $\mathbf{v} \in \mathbb{N}^p$ such that $\mathbf{v} \gg \mathbf{0}$. If we write $P_R(\mathbf{t}) = \sum_{\mathbf{n}_1, \dots, \mathbf{n}_p \geq 0} e(\mathbf{n}_1, \dots, \mathbf{n}_p) \binom{t_1 + \mathbf{n}_1}{\mathbf{n}_1} \cdots \binom{t_p + \mathbf{n}_p}{\mathbf{n}_p}$, then $0 \leq e(\mathbf{n}_1, \dots, \mathbf{n}_p) \in \mathbb{Z}$ for all $\mathbf{n}_1 + \cdots + \mathbf{n}_p = \dim(R/(0 :_R \mathfrak{N}^\infty)) - p$. For each $\mathbf{n} = (\mathbf{n}_1, \dots, \mathbf{n}_p) \in \mathbb{N}^p$ with $|\mathbf{n}| = \dim(R/(0 :_R \mathfrak{N}^\infty)) - p$, we set that $e(\mathbf{n}, R) := e(\mathbf{n}_1, \dots, \mathbf{n}_p)$ is the *mixed multiplicity of R of type $\mathbf{n}$* .

2.3. Polymatroids

In this subsection we include some relevant information about polymatroids.

Definition 2.14. Let E be a finite set and r a function $r : 2^E \rightarrow \mathbb{Z}_{\geq 0}$ satisfying the following two properties: (i) it is *non-decreasing*, i.e., $r(\mathfrak{T}_1) \leq r(\mathfrak{T}_2)$ if $\mathfrak{T}_1 \subseteq \mathfrak{T}_2 \subseteq E$, and (ii) it is *submodular*, i.e., $r(\mathfrak{T}_1 \cap \mathfrak{T}_2) + r(\mathfrak{T}_1 \cup \mathfrak{T}_2) \leq r(\mathfrak{T}_1) + r(\mathfrak{T}_2)$ if $\mathfrak{T}_1, \mathfrak{T}_2 \subseteq E$. The function r is called a *rank function on E* . We usually let $E = [p]$.

A (discrete) *polymatroid* $\mathcal{P}$ on $[p]$ with rank function r is a collection of points in $\mathbb{N}^p$ of the following form

$$\mathcal{P} = \left\{ \mathbf{x} = (x_1, \dots, x_p) \in \mathbb{N}^p \mid \sum_{j \in \mathfrak{J}} x_j \leq r(\mathfrak{J}), \forall \mathfrak{J} \subseteq [p], \sum_{i \in [p]} x_i = r([p]) \right\}.$$

By definition, a polymatroid consists of the integer points of a polytope (the convex hull of $\mathcal{P}$), we call that polytope a *base polymatroid polytope*. We note that a polymatroid is completely determined by its rank function.

Remark 2.15. If the rank function of $\mathcal{P}$ satisfies $r(\{i\}) \leq 1$ for every $i \in [p]$, then $\mathcal{P}$ is called a *matroid*. In other words, matroids are discrete polymatroids where every integer point is an element of $\{0, 1\}^p$. A general reference for matroids is [47].

In the following definition we consider the standard notions of linear and algebraic matroids (see [47, Chapter 6]) and adapt them to the polymatroid case.

Definition 2.16. Let $\mathcal{P}$ be a polymatroid.

- We say $\mathcal{P}$ is *linear* over a field $\mathbb{k}$ if there exists a $\mathbb{k}$ -vector space V and subspaces $V_i, i \in [p]$ such that for every $\mathfrak{J} \subseteq [p]$ we have $r(\mathfrak{J}) = \dim_{\mathbb{k}} \left(\sum_{j \in \mathfrak{J}} V_j \right)$ [47, Proposition 1.1.1]. The vector space V together with the subspaces V_i for $1 \leq i \leq p$, are a *linear representation* of $\mathcal{P}$.
- We say $\mathcal{P}$ is *algebraic* over a field $\mathbb{k}$ if there exists a field extension $\mathbb{k} \hookrightarrow \mathbb{L}$ and intermediate field extensions $\mathbb{L}_i, i \in [p]$ such that for every $\mathfrak{J} \subseteq [p]$ we have $r(\mathfrak{J}) = \text{trdeg}_{\mathbb{k}} \left(\bigwedge_{j \in \mathfrak{J}} \mathbb{L}_j \right)$, where $\bigwedge_{j \in \mathfrak{J}} \mathbb{L}_j$ is the *compositum* of the subfields, i.e., the smallest subfield in $\mathbb{L}$ containing all of them [47, Theorem 6.7.1]. The field $\mathbb{L}$ together with the subfields $\mathbb{L}_i$ for $1 \leq i \leq p$, are an *algebraic representation* of $\mathcal{P}$.

3. A characterization for the positivity of multidegrees

In this section, we focus on characterizing the positivity of multidegrees and our main goal is to prove Theorem A and Theorem B. Throughout this section we continue using the same notations and assumptions of Section 2.

We begin with the following result that relates the Hilbert polynomial $P_R(\mathbf{t}) \in \mathbb{Q}[\mathbf{t}]$ of R with the dimensions $r(\mathfrak{J}) = \dim(\Pi_{\mathfrak{J}}(X))$ of the schemes $\Pi_{\mathfrak{J}}(X)$. It extends [57, Theorem 1.7] to a multigraded setting.

Proposition 3.1. Assume Setup 2.1. For each $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq [p]$, let $\deg(P_R; \mathfrak{J})$ be the degree of the Hilbert polynomial P_R in the variables $t_{j_1}, \dots, t_{j_k}$. Then, for every such $\mathfrak{J} = \{j_1, \dots, j_k\}$ we have that

$$\deg(P_R; \mathfrak{J}) = r(\mathfrak{J}).$$

Proof. We may assume that $(0 :_R \mathfrak{N}^\infty) = 0$ and $\mathbb{k}$ is an infinite field by Remark 2.6. Fix $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq [p]$ and let $w \in \mathbb{N}$ be such that $\dim_{\mathbb{k}}([R]_{\mathbf{n}}) = P_R(\mathbf{n})$ for every $\mathbf{n} = (n_1, \dots, n_p) \geq w\mathbf{1}$. Let $(d_1, \dots, d_k)$ be such that $\delta := \deg(P_R; \mathfrak{J}) = d_1 + \dots + d_k$ and $t_{j_1}^{d_1} \cdots t_{j_k}^{d_k}$ divides a term of P_R .

Let q be a polynomial in the variables $\{t_i \mid i \notin \mathfrak{J}\}$ such that $P_R - q \cdot t_{j_1}^{d_1} \cdots t_{j_k}^{d_k}$ has no term divisible by $t_{j_1}^{d_1} \cdots t_{j_k}^{d_k}$. Let $\mathbf{s} = (s_i \mid i \notin \mathfrak{J}) \in \mathbb{N}^{p-|\mathfrak{J}|}$ be a vector of integers such that $\mathbf{s} \geq w\mathbf{1}$ and $q(\mathbf{s}) \neq 0$. Thus, if one evaluates $t_i = s_i$ in P_R for every $i \notin \mathfrak{J}$ one obtains a polynomial Q on the variables $t_{j_1}, \dots, t_{j_k}$ of degree δ . On the other hand, by [9, Theorem 3.4], for $n_{j_1}, \dots, n_{j_k} \geq w$ this polynomial Q coincides with the Hilbert polynomial of the $R_{(\mathfrak{J})}$ -module generated by $[R]_{\mathbf{s}'}$, where $s'_i = s_i$ if $i \notin \mathfrak{J}$ and $s'_i = 0$ otherwise. Call this module M .

Since $(0 :_{\mathbb{R}} \mathfrak{N}^\infty) = 0$, for every $1 \leq i \leq p$ we have $\text{grade}(\mathfrak{m}_i) \geq 1$, and then there exist elements $y_i \in [\mathbb{R}]_{\mathfrak{e}_i}$ which are non-zero-divisors (see, e.g., [6, Lemma 1.5.12]). From the fact that $y_1^{s'_1} \cdots y_p^{s'_p} \in M$, it follows that $\text{Ann}_{\mathbb{R}_{(\mathfrak{J})}}(M) = 0$. Therefore, $\delta = \dim(\text{Supp}(M) \cap X_{(\mathfrak{J})}) = r(\mathfrak{J})$, by [9, Theorem 3.4], finishing the proof. $\square$

In the following remark we gather some basic relations for the radicals of certain ideals.

Remark 3.2. (i) Let $I, J, K \subset \mathbb{R}$ be ideals. If $J \subset \sqrt{K}$, then $I + J \subset \sqrt{I + K}$. In particular, if $\sqrt{J} = \sqrt{K}$, then $\sqrt{I + J} = \sqrt{I + K}$.

(ii) For any element $x \in \mathfrak{m}_1$, since $(x :_{\mathbb{R}} \mathfrak{m}_1^\infty) \mathfrak{m}_1^k \subset (x)$ for some $k > 0$, it follows that $\sqrt{(x)} = \sqrt{(x :_{\mathbb{R}} \mathfrak{m}_1^\infty) \mathfrak{m}_1^k} = \sqrt{(x :_{\mathbb{R}} \mathfrak{m}_1^\infty) \cap \mathfrak{m}_1}$.

If $\mathbb{k}$ is an infinite field, then for each $1 \leq i \leq p$ we say that a property **P** is satisfied by a *general element* in the $\mathbb{k}$ -vector space $[\mathbb{R}]_{\mathfrak{e}_i}$, if there exists a dense open subset $\mathcal{U}$ of $[\mathbb{R}]_{\mathfrak{e}_i}$ with the Zariski topology such that every element in $\mathcal{U}$ satisfies the property **P**.

The following three technical lemmas are important steps for the proof of Theorem 3.7.

Lemma 3.3. Assume Setup 2.1 with $\mathbb{k}$ being an infinite field. Suppose that $\mathbb{R}$ is a domain. Let $x \in [\mathbb{R}]_{\mathfrak{e}_1}$ be a general element. Then, we have the equality $\sqrt{(x :_{\mathbb{R}} \mathfrak{N}^\infty)} = \sqrt{(x :_{\mathbb{R}} \mathfrak{m}_1^\infty)}$.

Proof. Since $(0 :_{\mathbb{R}} \mathfrak{N}^\infty) = 0$, we have that $\text{ht}(\mathfrak{m}_j) \geq 1$ for every $1 \leq j \leq p$. Consider the following finite set of prime ideals

$$\mathfrak{S} = \{\mathfrak{P} \in \text{Spec}(\mathbb{R}) \mid \mathfrak{P} \in \text{Min}(\mathfrak{m}_j) \text{ for some } 2 \leq j \leq p \text{ and } \mathfrak{P} \not\supseteq \mathfrak{m}_1\}.$$

By using the Prime Avoidance Lemma and the fact that $\mathbb{k}$ is infinite, for a general element $x \in [\mathbb{R}]_{\mathfrak{e}_1}$ we have that $x \notin \bigcup_{\mathfrak{P} \in \mathfrak{S}} \mathfrak{P}$. If $\mathfrak{P} \in \text{Min}(x :_{\mathbb{R}} \mathfrak{m}_1^\infty)$, then $\text{ht}(\mathfrak{P}) \leq 1$ by Krull's Principal Ideal Theorem, and so we would have that $\mathfrak{P} \in \mathfrak{S}$ whenever $\mathfrak{P} \not\supseteq \mathfrak{m}_1$ and $\mathfrak{P} \supseteq \mathfrak{m}_j$ for some $2 \leq j \leq p$. Therefore, for any $\mathfrak{P} \in \text{Spec}(\mathbb{R})$ and a general element $x \in [\mathbb{R}]_{\mathfrak{e}_1}$, if $\mathfrak{P} \in \text{Min}(x :_{\mathbb{R}} \mathfrak{m}_1^\infty)$ we get $\mathfrak{P} \supseteq (x :_{\mathbb{R}} \mathfrak{N}^\infty) = (x :_{\mathbb{R}} (\mathfrak{m}_1 \cap \mathfrak{m}_2 \cap \cdots \cap \mathfrak{m}_p)^\infty)$; so, $\sqrt{(x :_{\mathbb{R}} \mathfrak{N}^\infty)} = \sqrt{(x :_{\mathbb{R}} \mathfrak{m}_1^\infty)}$. $\square$

The lemma below is necessary for some reduction arguments in Theorem 3.7.

Lemma 3.4. Assume Setup 2.1 with $\mathbb{k}$ being an infinite field. Suppose that $\mathbb{R}$ is a domain. Let $x \in [\mathbb{R}]_{\mathfrak{e}_1}$ be a general element and set $Z = X \cap V(x) = \text{MultiProj}(\mathbb{R}/x\mathbb{R})$. Then, for each $\mathfrak{J} = \{1, j_2, \dots, j_k\} \subseteq [p]$, the following statements hold:

- (i) $\dim(\Pi_{\mathfrak{J}}(Z)) = \dim(X_{\mathfrak{J}} \cap V(x))$, where $X_{\mathfrak{J}} \cap V(x) = \text{MultiProj}(\mathbb{R}_{(\mathfrak{J})}/x\mathbb{R}_{(\mathfrak{J})})$.
- (ii) $\dim(\Pi_{\mathfrak{L}}(Z)) = \dim(\Pi'_{\mathfrak{L}}(X_{\mathfrak{J}} \cap V(x)))$, where $\mathfrak{L} = \mathfrak{J} \setminus \{1\}$ and $\Pi'_{\mathfrak{L}}$ denotes the natural projection $\Pi'_{\mathfrak{L}} : \mathbb{P}_{\mathbb{k}}^{m_1} \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_{j_2}} \times_{\mathbb{k}} \cdots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_{j_k}} \rightarrow \mathbb{P}_{\mathbb{k}}^{m_{j_2}} \times_{\mathbb{k}} \cdots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_{j_k}}$.

Proof. For notational purposes, let $\mathfrak{b}_{\mathfrak{J}} := \mathfrak{m}_1 \cap \mathbf{R}_{(\mathfrak{J})}$.

(i) From (2) we have that $\Pi_{\mathfrak{J}}(\mathbf{Z}) \cong \text{MultiProj}(\mathbf{R}/((\mathfrak{x} :_{\mathbf{R}} \mathfrak{N}^{\infty}) + \sum_{j \notin \mathfrak{J}} \mathfrak{m}_j))$. Since we are assuming $1 \in \mathfrak{J}$, from Remark 2.4 we obtain the natural isomorphism

$$\mathbf{R}_{(\mathfrak{J})} / \left(\mathfrak{x} :_{\mathbf{R}_{(\mathfrak{J})}} \mathfrak{b}_{\mathfrak{J}}^{\infty} \right) \xrightarrow{\cong} \mathbf{R} / \left((\mathfrak{x} :_{\mathbf{R}} \mathfrak{m}_1^{\infty}) + \sum_{j \notin \mathfrak{J}} \mathfrak{m}_j \right);$$

indeed, for $\ell \geq 0$ and $\mathfrak{y} \in \mathbf{R}_{(\mathfrak{J})}$, one notices that $\mathfrak{b}_{\mathfrak{J}}^{\ell} \cdot \mathfrak{y} \in \mathfrak{x} \mathbf{R}_{(\mathfrak{J})}$ if and only if $\mathfrak{m}_1^{\ell} \cdot \mathfrak{y} \in \mathfrak{x} \mathbf{R}$.

By Lemma 3.3 and Remark 3.2(i) we have $\sqrt{(\mathfrak{x} :_{\mathbf{R}} \mathfrak{m}_1^{\infty}) + \sum_{j \notin \mathfrak{J}} \mathfrak{m}_j} = \sqrt{(\mathfrak{x} :_{\mathbf{R}} \mathfrak{N}^{\infty}) + \sum_{j \notin \mathfrak{J}} \mathfrak{m}_j}$, and by applying Lemma 3.3 to the ring $\mathbf{R}_{(\mathfrak{J})}$ we obtain $\sqrt{(\mathfrak{x} :_{\mathbf{R}_{(\mathfrak{J})}} \mathfrak{b}_{\mathfrak{J}}^{\infty})} = \sqrt{(\mathfrak{x} :_{\mathbf{R}_{(\mathfrak{J})}} \mathfrak{N}_{\mathfrak{J}}^{\infty})}$. It follows that

$$\mathbf{R}_{(\mathfrak{J})} / \sqrt{(\mathfrak{x} :_{\mathbf{R}_{(\mathfrak{J})}} \mathfrak{N}_{\mathfrak{J}}^{\infty})} \cong \mathbf{R} / \sqrt{(\mathfrak{x} :_{\mathbf{R}} \mathfrak{N}^{\infty}) + \sum_{j \notin \mathfrak{J}} \mathfrak{m}_j}, \quad (4)$$

which gives the result.

(ii) By using (2) we obtain that $\Pi_{\mathfrak{L}}(\mathbf{Z}) \cong \text{MultiProj}(\mathbf{R}/((\mathfrak{x} :_{\mathbf{R}} \mathfrak{N}^{\infty}) + \sum_{j \notin \mathfrak{J}} \mathfrak{m}_j + \mathfrak{m}_1))$ and that $\Pi'_{\mathfrak{L}}(\mathbf{X}_{\mathfrak{J}} \cap \mathbf{V}(\mathfrak{x})) \cong \text{MultiProj}(\mathbf{R}_{(\mathfrak{J})} / ((\mathfrak{x} :_{\mathbf{R}_{(\mathfrak{J})}} \mathfrak{N}_{\mathfrak{J}}^{\infty}) + \mathfrak{b}_{\mathfrak{J}}))$. Since the isomorphism in (4) can be extended to

$$\mathbf{R}_{(\mathfrak{J})} / \left(\sqrt{(\mathfrak{x} :_{\mathbf{R}_{(\mathfrak{J})}} \mathfrak{N}_{\mathfrak{J}}^{\infty})} + \mathfrak{b}_{\mathfrak{J}} \right) \cong \mathbf{R} / \left(\sqrt{(\mathfrak{x} :_{\mathbf{R}} \mathfrak{N}^{\infty}) + \sum_{j \notin \mathfrak{J}} \mathfrak{m}_j} + \mathfrak{m}_1 \right),$$

the result follows from Remark 3.2(i). $\square$

We continue with the next auxiliary lemma that allows us to simplify the proof of Theorem 3.7.

Lemma 3.5. Assume Setup 2.1 with $\mathbb{k}$ being an infinite field. Suppose that $\mathbf{R}$ is a domain and $r(1) \geq 1$. Let $\mathfrak{x} \in [\mathbf{R}]_{e_1}$ be a general element and set $\mathbf{Z} = \mathbf{X} \cap \mathbf{V}(\mathfrak{x}) = \text{MultiProj}(\mathbf{R}/\mathfrak{x}\mathbf{R})$. Then, the following statements hold:

- (i) If $1 \in \mathfrak{J} \subseteq [p]$, then $\dim(\Pi_{\mathfrak{J}}(\mathbf{Z})) = r(\mathfrak{J}) - 1$; in particular, $\dim(\mathbf{Z}) = r - 1$.
- (ii) If $1 \notin \mathfrak{J}$ and $r(\mathfrak{K}) > r(\mathfrak{J})$, where $\mathfrak{K} = \{1\} \cup \mathfrak{J} \subseteq [p]$, then $\dim(\Pi_{\mathfrak{J}}(\mathbf{Z})) = r(\mathfrak{J})$.

Proof. (i) First, from Lemma 3.4(i) it suffices to compute $\dim(\mathbf{X}_{\mathfrak{J}} \cap \mathbf{V}(\mathfrak{x}))$, where $\mathbf{X}_{\mathfrak{J}} \cap \mathbf{V}(\mathfrak{x}) = \text{MultiProj}(\mathbf{R}_{(\mathfrak{J})}/\mathfrak{x}\mathbf{R}_{(\mathfrak{J})})$. For $\mathfrak{J} = \{1, j_2, \dots, j_k\} \subseteq [p]$, note that $\Pi_1(\mathbf{X}) \cong \Pi'_1(\mathbf{X}_{\mathfrak{J}})$, where Π'_1 denotes the natural projection $\Pi'_1 : \mathbb{P}_{\mathbb{k}}^{m_1} \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_{j_2}} \times_{\mathbb{k}} \dots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_{j_k}} \rightarrow \mathbb{P}_{\mathbb{k}}^{m_1}$. Therefore, neither the assumption nor the conclusion changes if we substitute $\mathbf{R}$ by $\mathbf{R}_{(\mathfrak{J})}$ and $\mathbf{X}$ by $\mathbf{X}_{\mathfrak{J}}$, and we do so.

From the short exact sequence

$$0 \rightarrow R(-\mathbf{e}_1) \xrightarrow{x} R \rightarrow R/xR \rightarrow 0,$$

we obtain $P_{R/xR}(\mathbf{t}) = P_R(\mathbf{t}) - P_R(\mathbf{t} - \mathbf{e}_1)$. By using Proposition 3.1, $\deg(P_R; \mathbf{t}_1) = r(1) \geq 1$ and so P_R is non-constant as a univariate polynomial in the variable $\mathbf{t}_1$. Thus, $P_{R/xR}(\mathbf{t}) \neq 0$ which implies that $(x :_R \mathfrak{N}^\infty)$ is a proper ideal. So, Krull's Principal Ideal Theorem yields that $\text{ht}(x :_R \mathfrak{N}^\infty) = 1$ and that

$$\dim(Z) = \dim(R/(x :_R \mathfrak{N}^\infty)) - p = \dim(R) - 1 - p = (r + p) - 1 - p = r - 1.$$

(ii) By using Lemma 3.4(ii), we can substitute R by $R_{(\mathfrak{K})}$ and X by $X_{\mathfrak{K}}$, and we do so. So, we may assume that $\mathfrak{K} = [p]$ and $\mathfrak{J} = \{2, \dots, p\}$. From (2) we get the isomorphism

$$\Pi_{\mathfrak{J}}(Z) \cong \text{MultiProj}\left(R / ((x :_R \mathfrak{N}^\infty) + \mathfrak{m}_1)\right). \quad (5)$$

The equality

$$\sqrt{(x :_R \mathfrak{N}^\infty) + \mathfrak{m}_1} = \sqrt{(x :_R \mathfrak{m}_1^\infty) + \mathfrak{m}_1} \quad (6)$$

follows from Lemma 3.3 and Remark 3.2(i). The assumption yields that

$$\text{ht}(\mathfrak{m}_1) = \dim(R) - \dim(R_{(\mathfrak{J})}) = (r + p) - (r(\mathfrak{J}) + p - 1) \geq 2,$$

then as a consequence Krull's Principal Ideal Theorem it follows that $\sqrt{(x)} = \sqrt{(x :_R \mathfrak{m}_1^\infty)}$; therefore, Remark 3.2(i) implies that $\sqrt{(x :_R \mathfrak{m}_1^\infty) + \mathfrak{m}_1} = \sqrt{\mathfrak{m}_1}$. By summing up, we obtain the equalities $\dim(R / ((x :_R \mathfrak{N}^\infty) + \mathfrak{m}_1)) = \dim(R/\mathfrak{m}_1) = r(\mathfrak{J}) + p - 1$, and so the result follows. $\square$

The next important theorem computes the dimension of the image of the projections $\Pi_{\mathfrak{J}}$ after cutting with a general hyperplane under certain conditions. For the proof of this result, we need the following version of Grothendieck's Connectedness Theorem. For that, we recall the definitions

$$c(R) := \min \{ \dim(R/\mathfrak{a}) \mid \mathfrak{a} \subset R \text{ is an ideal and } \text{Spec}(R) \setminus V(\mathfrak{a}) \text{ is disconnected} \},$$

$$\text{sdim}(R) := \min \{ \dim(R/\mathfrak{P}) \mid \mathfrak{P} \in \text{Min}(R) \} \text{ and}$$

$$\text{ara}(\mathfrak{a}) := \min \{ n \mid \sqrt{(\mathfrak{a}_1, \dots, \mathfrak{a}_n)} = \sqrt{\mathfrak{a}} \text{ and } \mathfrak{a}_i \in R \}$$

for any ideal $\mathfrak{a} \subset R$.

Lemma 3.6 ([5, Proposition 2.1], [57, Lemma 2.6]). *For two proper homogeneous ideals $\mathfrak{a}, \mathfrak{b} \subset R$, if $\min\{\dim(R/\mathfrak{a}), \dim(R/\mathfrak{b})\} > \dim(R/(\mathfrak{a} + \mathfrak{b}))$, then*

$$\dim(R/(\mathfrak{a} + \mathfrak{b})) \geq \min\{c(R), \text{sdim}(R) - 1\} - \text{ara}(\mathfrak{a} \cap \mathfrak{b}).$$

We are now ready to present the following theorem.

Theorem 3.7. *Assume Setup 2.1 with $\mathbb{k}$ being an infinite field. Suppose that R is a domain and $r(1) \geq 1$. Let $x \in [R]_{e_1}$ be a general element and set $Z = X \cap V(x) = \text{MultiProj}(R/xR)$. Then, for each $\mathfrak{J} \subseteq [p]$ we have that*

$$\dim(\Pi_{\mathfrak{J}}(Z)) = \min \left\{ r(\mathfrak{J}), r(\mathfrak{J} \cup \{1\}) - 1 \right\}.$$

Proof. For each $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq \mathfrak{K} = \{h_1, \dots, h_\ell\} \subseteq [p]$ we have that $\Pi_{\mathfrak{J}}(Z) = \Pi'_{\mathfrak{J}}(\Pi_{\mathfrak{K}}(Z))$ where $\Pi'_{\mathfrak{J}}$ denotes the natural projection $\Pi'_{\mathfrak{J}} : \mathbb{P}_{\mathbb{k}}^{m_{h_1}} \times_{\mathbb{k}} \dots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_{h_\ell}} \rightarrow \mathbb{P}_{\mathbb{k}}^{m_{j_1}} \times_{\mathbb{k}} \dots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_{j_k}}$. So, from Lemma 3.5(i) it follows that the inequality “ $\leq$ ” holds in the desired equality.

Due to Lemma 3.5, in order to show the reversed inequality “ $\geq$ ”, it suffices to show that $\dim(\Pi_{\mathfrak{J}}(Z)) \geq r(\mathfrak{J}) - 1$ when $1 \notin \mathfrak{J}$ and $r(\mathfrak{K}) = r(\mathfrak{J})$, where $\mathfrak{K} = \{1\} \cup \mathfrak{J} \subseteq [p]$. By using Lemma 3.4(ii), we assume may that $\mathfrak{K} = [p]$ and $\mathfrak{J} = \{2, \dots, p\}$. From (5) and (6), the proof would be complete if we prove the inequality $\dim(R/((x :_R m_1^\infty) + m_1)) \geq (r_{\mathfrak{J}} - 1) + (p - 1) = r + p - 2$.

By using Lemma 3.3 and Lemma 3.5(i) we obtain that

$$\dim(R/(x :_R m_1^\infty)) = \dim(R/(x :_R \mathfrak{N}^\infty)) = (r - 1) + p = r + p - 1,$$

and since $r(\mathfrak{J}) = r$, we have

$$\dim(R/m_1) = \dim(R_{(\mathfrak{J})}) = r(\mathfrak{J}) + (p - 1) = r + (p - 1) = r + p - 1.$$

Moreover, (6) and Lemma 3.5(i) yield that

$$\dim(R/((x :_R m_1^\infty) + m_1)) = \dim(R/((x :_R \mathfrak{N}^\infty) + m_1)) \leq (r - 1) + (p - 1) = r + p - 2.$$

Since $x \in m_1$, Remark 3.2(ii) gives that $\text{ara}((x :_R m_1^\infty) \cap m_1) = \text{ara}((x)) = 1$. As R is a domain, $c(R) = \text{sdim}(R) = r + p$. Therefore, from Lemma 3.6 we obtain that

$$\dim(R/((x :_R m_1^\infty) + m_1)) \geq \min\{r + p, (r + p) - 1\} - 1 = r + p - 2.$$

So, the proof is complete. $\square$

Notation 3.8. Let $\{x_0, \dots, x_s\}$ be a basis of the $\mathbb{k}$ -vector space $[R]_{e_1}$. Consider a purely transcendental field extension $\mathbb{L} := \mathbb{k}(z_0, \dots, z_s)$ of $\mathbb{k}$, and set $R_{\mathbb{L}} := R \otimes_{\mathbb{k}} \mathbb{L}$ and $X_{\mathbb{L}} := X \otimes_{\mathbb{k}} \mathbb{L} = \text{MultiProj}(R_{\mathbb{L}}) \subseteq \mathbb{P} \otimes_{\mathbb{k}} \mathbb{L} = \mathbb{P}_{\mathbb{L}}^{m_1} \times_{\mathbb{L}} \dots \times_{\mathbb{L}} \mathbb{P}_{\mathbb{L}}^{m_p}$. We say that $z := z_0 x_0 + \dots + z_s x_s \in [R_{\mathbb{L}}]_{e_1}$ is the *generic element* of $[R_{\mathbb{L}}]_{e_1}$.

In the following remark we explain that field extensions as in Notation 3.8 preserve the domain assumption.

Remark 3.9. Suppose that R is a domain and consider a purely transcendental field extension $\mathbb{k}(\xi)$. Then, $R \otimes_{\mathbb{k}} \mathbb{k}(\xi)$ is also a domain; indeed, one can see that $R \otimes_{\mathbb{k}} \mathbb{k}(\xi)$ is a subring of the field of fractions $\text{Quot}(R[\xi])$ of the polynomial ring $R[\xi]$. So, when R is a domain one can extend $\mathbb{k}$ to an infinite field without losing the assumption of R being a domain.

The lemma below shows that the Hilbert function modulo a generic element coincides with the one modulo a general element.

Lemma 3.10. Assume Notation 3.8 with $\mathbb{k}$ being an infinite field. Let $x \in [R]_{e_1}$ be a general element, then

$$\dim_{\mathbb{k}}([R/xR]_{\mathbf{v}}) = \dim_{\mathbb{L}}([R_{\mathbb{L}}/zR_{\mathbb{L}}]_{\mathbf{v}})$$

for all $\mathbf{v} \in \mathbb{N}^p$.

Proof. Let T be the polynomial ring $T = \mathbb{k}[z_0, \dots, z_s]$ and consider the finitely generated T -algebra given by $S = (R \otimes_{\mathbb{k}} T) / \mathfrak{w}(R \otimes_{\mathbb{k}} T)$ where $\mathfrak{w} = z_0x_0 + \dots + z_sx_s \in R \otimes_{\mathbb{k}} T$. From the Grothendieck's Generic Freeness Lemma (see, e.g., [39, Theorem 24.1], [13, Theorem 14.4]) there exists an element $0 \neq \mathfrak{a} \in T$ such that $S_{\mathfrak{a}}$ is a free $T_{\mathfrak{a}}$ -module. Hence, for any $\mathfrak{p} \in \text{Spec}(T)$ inside the dense open subset $D(\mathfrak{a}) \subset \text{Spec}(T)$, if $k(\mathfrak{p})$ denotes the residue field $k(\mathfrak{p}) = T_{\mathfrak{p}}/\mathfrak{p}T_{\mathfrak{p}}$ of $T_{\mathfrak{p}}$, one has that

$$\dim_{k(\mathfrak{p})}([S_{\mathfrak{a}} \otimes_{T_{\mathfrak{a}}} k(\mathfrak{p})]_{\mathbf{v}}) = \dim_{\text{Quot}(T)}([S_{\mathfrak{a}} \otimes_{T_{\mathfrak{a}}} \text{Quot}(T)]_{\mathbf{v}}) = \dim_{\mathbb{L}}([R_{\mathbb{L}}/zR_{\mathbb{L}}]_{\mathbf{v}})$$

for all $\mathbf{v} \in \mathbb{N}^p$. Note that for any $\beta = (\beta_0, \dots, \beta_s) \in \mathbb{k}^{s+1}$ with $\mathfrak{p}_{\beta} = (z_0 - \beta_0, \dots, z_s - \beta_s) \in D(\mathfrak{a})$ one has the isomorphisms

$$S_{\mathfrak{a}} \otimes_{T_{\mathfrak{a}}} k(\mathfrak{p}_{\beta}) \cong \frac{R \otimes_{\mathbb{k}} T}{(z_0x_0 + \dots + z_sx_s, z_0 - \beta_0, \dots, z_s - \beta_s)} \cong R/(\beta_0x_0 + \dots + \beta_sx_s)R.$$

So, the result follows. $\square$

We now obtain Theorem A when X is an irreducible scheme.

Remark 3.11. We first provide a couple of general words regarding the proof of Theorem 3.12 below and where the irreducibility assumption comes into play. The proof is achieved by iteratively cutting with generic hyperplanes (following Notation 3.8) to arrive to a zero-dimensional situation, and the main constraint is to control the dimension of the image of all the possible projections after cutting with a general hyperplane (see (7)). Our main tool to control those dimensions is Theorem 3.7, where it is needed to assume that R is a domain. When X is irreducible, by just taking the reduced scheme structure $X_{\text{red}} = \text{MultiProj}(R/\sqrt{0})$ we can easily reduce to the case where R is a domain. To maintain the irreducibility assumption during the inductive process, we use a

“generic” version of Bertini’s Theorem as presented in [16, Proposition 1.5.10]. It should be noted that the usual versions of Bertini’s Theorem for irreducibility require X to be geometrically irreducible and that the dimension of the image of certain morphism is bigger or equal than two (see [32, Theoreme 6.10, Corollaire 6.11]). Finally, Lemma 3.10 is used to relate the process of cutting with a *generic* hyperplane with the one of cutting with a *general* hyperplane.

Theorem 3.12. *Assume Setup 2.1. Suppose that X is irreducible. Let $\mathbf{n} = (n_1, \dots, n_p) \in \mathbb{N}^p$ such that $|\mathbf{n}| = r$. Then, $\deg_{\mathbb{P}}^{\mathbf{n}}(X) > 0$ if and only if for each $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq [p]$ the inequality $n_{j_1} + \dots + n_{j_k} \leq r(\mathfrak{J})$ holds.*

Proof. From Proposition 3.1 it is clear that the inequalities $n_{j_1} + \dots + n_{j_k} \leq r(\mathfrak{J})$ are a necessary condition for $\deg_{\mathbb{P}}^{\mathbf{n}}(X) = e(\mathbf{n}; R) > 0$. Therefore, it suffices to show that they are also sufficient.

Assume that $n_{j_1} + \dots + n_{j_k} \leq r(\mathfrak{J})$ for every $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq [p]$. We may also assume that $(0 :_R \mathfrak{N}^\infty) = 0$ by Remark 2.6(i). Hence, the condition of X being irreducible implies that $\sqrt{0} \subset R$ is a prime ideal. Since the associativity formula for mixed multiplicities (see, e.g., [9, Lemma 2.7]) yields that

$$e(\mathbf{n}; R) = \text{length}_{R_{\sqrt{0}}} (R_{\sqrt{0}}) \cdot e(\mathbf{n}; R/\sqrt{0}),$$

we can assume that R is a domain, and we do so. In addition, by Remark 2.6(ii), Proposition 3.1, and Remark 3.9 we may also assume that $\mathbb{k}$ is an infinite field.

We proceed by induction on r . If $r = 0$, then [9, Theorem 3.10] implies $e(\mathbf{0}; R) > 0$.

Suppose now that $r \geq 1$. Without any loss of generality, perhaps after changing the grading, we can assume that $n_1 \geq 1$. Let $\mathbb{L}$, $R_{\mathbb{L}}$, $X_{\mathbb{L}}$ and z be defined as in Notation 3.8. Let $x \in [R]_{e_1}$ be a general element. Set $S = R/xR$, $Z = X \cap V(x) = \text{MultiProj}(S)$, $T = R_{\mathbb{L}}/zR_{\mathbb{L}}$, $W = X_{\mathbb{L}} \cap V(z) = \text{MultiProj}(T)$ and $\mathbf{n}' = \mathbf{n} - \mathbf{e}_1$. Then, [9, Lemma 3.9] and Lemma 3.10 yield that $e(\mathbf{n}; R) = e(\mathbf{n}'; S) = e(\mathbf{n}'; T)$. From [16, Proposition 1.5.10] we obtain that W is also an irreducible scheme. By the assumed inequalities and because $n_1 \geq 1$ we have that for each $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq [p]$ the following inequality holds

$$n'_{j_1} + \dots + n'_{j_k} \leq \min \left\{ r(\mathfrak{J}), r(\mathfrak{J} \cup \{1\}) - 1 \right\}, \quad (7)$$

and the latter is equal to $\dim(\Pi_{\mathfrak{J}}(Z))$ by Theorem 3.7. Moreover, by Lemma 3.10 and Proposition 3.1, we also have $\dim(\Pi_{\mathfrak{J}}(W)) = \dim(\Pi_{\mathfrak{J}}(Z))$; here, by an abuse of notation $\Pi_{\mathfrak{J}}(W)$ denotes the image of the natural projection $\Pi_{\mathfrak{J}} : \mathbb{P} \otimes_{\mathbb{k}} \mathbb{L} \rightarrow \mathbb{P}_{\mathbb{L}}^{m_{j_1}} \times_{\mathbb{L}} \dots \times_{\mathbb{L}} \mathbb{P}_{\mathbb{L}}^{m_{j_k}}$ restricted to W .

Finally, by using the inductive hypothesis applied to the irreducible scheme W , we obtain that $e(\mathbf{n}; R) = e(\mathbf{n}'; T) > 0$, and so the result follows. $\square$

Now we are ready to show the general version of Theorem A.

Corollary 3.13. Assume Setup 2.1. Let $\mathbf{n} = (n_1, \dots, n_p) \in \mathbb{N}^p$ such that $|\mathbf{n}| = \dim(X)$. Then, $\deg_{\mathbb{P}}^{\mathbf{n}}(X) > 0$ if and only if there is an irreducible component $Y \subseteq X$ of X that satisfies the following two conditions:

- (a) $\dim(Y) = \dim(X)$.
- (b) For each $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq [p]$ the inequality $n_{j_1} + \dots + n_{j_k} \leq \dim(\Pi_{\mathfrak{J}}(Y))$ holds.

Proof. We may assume that $(0 :_{\mathbb{R}} \mathfrak{N}^{\infty}) = 0$ by Remark 2.6(i). By the associativity formula for mixed multiplicities (see, e.g., [9, Lemma 2.7]) we get the equation

$$\deg_{\mathbb{P}}^{\mathbf{n}}(X) = e(\mathbf{n}; \mathbb{R}) = \sum_{\substack{\mathfrak{P} \in \text{Min}(\mathbb{R}) \\ \dim(\mathbb{R}/\mathfrak{P}) = r+p}} \text{length}_{\mathbb{R}_{\mathfrak{P}}}(\mathbb{R}_{\mathfrak{P}}) \cdot e(\mathbf{n}; \mathbb{R}/\mathfrak{P}).$$

Thus, $e(\mathbf{n}; \mathbb{R}) > 0$ if and only if $e(\mathbf{n}; \mathbb{R}/\mathfrak{P}) > 0$ for some minimal prime $\mathfrak{P} \in \text{Min}(\mathbb{R})$ of maximal dimension. So, the result is clear from Theorem 3.12. $\square$

Below we have a proof for Theorem B.

Corollary 3.14. Assume Setup 2.12. Let $\mathbf{n} = (n_1, \dots, n_p) \in \mathbb{N}^p$ such that $\dim(\mathbb{R}/(0 :_{\mathbb{R}} \mathfrak{N}^{\infty})) - p = |\mathbf{n}|$. Then, $e(\mathbf{n}; \mathbb{R}) > 0$ if and only if there is a minimal prime ideal $\mathfrak{P} \in \text{Min}(0 :_{\mathbb{R}} \mathfrak{N}^{\infty})$ of $(0 :_{\mathbb{R}} \mathfrak{N}^{\infty})$ that satisfies the following two conditions:

- (a) $\dim(\mathbb{R}/\mathfrak{P}) = \dim(\mathbb{R}/(0 :_{\mathbb{R}} \mathfrak{N}^{\infty}))$.
- (b) For each $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq [p]$ the inequality $n_{j_1} + \dots + n_{j_k} \leq \dim\left(\frac{\mathbb{R}}{\mathfrak{P} + \sum_{j \notin \mathfrak{J}} \mathfrak{m}_j}\right) - k$ holds.

Proof. As in Corollary 3.13, after assuming that $(0 :_{\mathbb{R}} \mathfrak{N}^{\infty}) = 0$ and using the associativity formula for mixed multiplicities, we obtain that $e(\mathbf{n}; \mathbb{R}) > 0$ if and only if $e(\mathbf{n}; \mathbb{R}/\mathfrak{P}) > 0$ for some minimal prime $\mathfrak{P} \in \text{Min}(\mathbb{R})$ of maximal dimension. Note that, for each $\mathfrak{P} \in \text{Min}(\mathbb{R})$, $\mathbb{R}/\mathfrak{P}$ is naturally a finitely generated standard $\mathbb{N}^p$ -graded algebra over a field. So, the result follows by using Theorem 3.12. $\square$

Finally, for the sake of completeness, we provide a brief discussion on how Theorem 3.12 can be recovered (over the complex number) from the related results of [34, §2.2].

Remark 3.15. Assume $\mathbb{k} = \mathbb{C}$. For the closed subscheme $X \subset \mathbb{P} = \mathbb{P}_{\mathbb{k}}^{m_1} \times_{\mathbb{k}} \dots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_p}$, let L_i be the pullback of $\mathcal{O}_{\mathbb{P}_{\mathbb{k}}^{m_i}}(1)$ to X . Take E_i to be $|L_i|$. Following the notation in [34, §2.2], for each $\emptyset \neq \mathfrak{J} \subseteq [p]$, denote by

$$\Phi_{\mathfrak{J}} : X \rightarrow \mathbb{P}(E_{\mathfrak{J}}^{\vee})$$

the Kodaira map corresponding with the linear system $E_{\mathfrak{J}}$. Let $\tau_{\mathfrak{J}}$ be the dimension of the closure of the image of $\Phi_{\mathfrak{J}}$ ([34, Definition 2.12]). Consequently, it is easy to check that $\dim(\Pi_{\mathfrak{J}}(X)) = \tau_{\mathfrak{J}}$. Thus, [34, Theorems 2.14, 2.19] translate into the following statement: $\dim(\Pi_{\mathfrak{J}}(X)) \geq |\mathfrak{J}|$ if and only if for general hyperplanes $H_j \in |\Pi_j^* \mathcal{O}_{\mathbb{P}_{\mathbb{k}}^{m_i}}(1)|$ ($j \in \mathfrak{J}$), $X \cap (\bigcap_{j \in \mathfrak{J}} H_j) \neq \emptyset$. The latter is equivalent to the condition $[X] \cdot \prod_{j \in \mathfrak{J}} [H_j] \neq 0$ on intersection of classes. Theorem 3.12 (over the complex numbers) eventually follows from applying this statement finitely many times to relevant index subsets $\mathfrak{J}$.

4. Positivity of the mixed multiplicities of ideals

In this section, we characterize the positivity of the mixed multiplicities of ideals. The results obtained here are a consequence of applying Theorem B to a certain multigraded algebra. For the particular case of ideals generated in one degree in graded domains we obtain a neat characterization in Theorem 4.4.

Throughout this section we use the following setup.

Setup 4.1. Let R be a Noetherian local ring with maximal ideal $\mathfrak{m} \subset R$ (or a finitely generated standard graded algebra over a field $\mathbb{k}$ with graded irrelevant ideal $\mathfrak{m} \subset R$).

Let $J_0 \subset R$ be an $\mathfrak{m}$ -primary ideal and $J_1, \dots, J_p \subset R$ be arbitrary ideals (homogeneous in the graded case). The *multi-Rees algebra* of the ideals $J_0, J_1, \dots, J_p$ is given by

$$\mathcal{R}(J_0, \dots, J_p) := R[J_0 t_0, \dots, J_p t_p] = \bigoplus_{i_0 \geq 0, \dots, i_p \geq 0} J_0^{i_0} \cdots J_p^{i_p} t_0^{i_0} \cdots t_p^{i_p} \subset R[t_0, \dots, t_p],$$

where $t_0, \dots, t_p$ are new variables. Note that $\mathcal{R}(J_0, \dots, J_p)$ is naturally a standard $\mathbb{N}^{p+1}$ -graded algebra and that, for $0 \leq k \leq p$, the ideal $\mathfrak{m}_k$ generated by elements of degree $\mathbf{e}_k$ is given by

$$\mathfrak{m}_k := J_k t_k \mathcal{R}(J_0, \dots, J_p) \subset \mathcal{R}(J_0, \dots, J_p).$$

Let $\mathfrak{N} := \mathfrak{m}_0 \cap \cdots \cap \mathfrak{m}_p$ be the corresponding multigraded irrelevant ideal. Since J_0 is $\mathfrak{m}$ -primary, we obtain that

$$T(J_0 \mid J_1, \dots, J_p) := \mathcal{R}(J_0, \dots, J_p) \otimes_R R/J_0 = \bigoplus_{i_0 \geq 0, i_1 \geq 0, \dots, i_p \geq 0} J_0^{i_0} J_1^{i_1} \cdots J_p^{i_p} / J_0^{i_0+1} J_1^{i_1} \cdots J_p^{i_p}$$

is a finitely generated standard $\mathbb{N}^{p+1}$ -graded algebra over the Artinian local ring R/J_0 . For simplicity of notation, throughout this section we fix $\mathcal{R} := \mathcal{R}(J_0, \dots, J_p)$ and $T := T(J_0 \mid J_1, \dots, J_p)$. Let r be the integer $r := \dim(\text{MultiProj}(T))$, which coincides with the degree of the Hilbert polynomial $P_T(\mathbf{u}_1, \dots, \mathbf{u}_{p+1})$ of the $\mathbb{N}^{p+1}$ -graded R/J_0 -algebra T . From [58, Theorem 1.2(a)] we have the equality $r = \dim(R/(0 :_R (J_1 \cdots J_p)^\infty)) - 1$.

Definition 4.2. Under the above notations, for each $\mathbf{n} \in \mathbb{N}^{p+1}$ with $|\mathbf{n}| = r$, we say that

$$e_{\mathbf{n}}(J_0 \mid J_1, \dots, J_p) := e(\mathbf{n}; T)$$

is the *mixed multiplicity* of $J_0, J_1, \dots, J_p$ of type $\mathbf{n}$.

The main focus in this section is to characterize when $e_{\mathbf{n}}(J_0 \mid J_1, \dots, J_p) > 0$. As a direct consequence of Theorem B we get the following general criterion for the positivity $e_{\mathbf{n}}(J_0 \mid J_1, \dots, J_p)$.

Corollary 4.3. Assume Setup 4.1 and the notations above. Let $\mathbf{n} = (n_0, n_1, \dots, n_p) \in \mathbb{N}^{p+1}$ such that $|\mathbf{n}| = r$. Then, $e_{\mathbf{n}}(J_0 \mid J_1, \dots, J_p) > 0$ if and only if there is a minimal prime ideal $\mathfrak{P} \in \text{Min}(0 :_T \mathfrak{N}^\infty)$ of $(0 :_T \mathfrak{N}^\infty)$ that satisfies the following two conditions:

- (a) $\dim(T/\mathfrak{P}) = \dim(T/(0 :_T \mathfrak{N}^\infty))$.
- (b) For each $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq \{0\} \cup [p]$ the inequality, $n_{j_1} + \dots + n_{j_k} \leq \dim\left(\frac{T}{\mathfrak{P} + \sum_{j \notin \mathfrak{J}} m_j T}\right) - k$ holds.

We now focus on the case where R is a graded $\mathbb{k}$ -domain and each ideal J_i is generated in one degree. In this case, our characterization depends on the *analytic spread* of certain ideals; recall that the analytic spread of an ideal $I \subset R$ is given by $\ell(I) := \dim(\mathcal{R}(I)/\mathfrak{m}\mathcal{R}(I))$.

Theorem 4.4. Let R be a finitely generated standard graded domain over a field $\mathbb{k}$ with graded irrelevant ideal $\mathfrak{m} \subset R$. Let $J_0 \subset R$ be an $\mathfrak{m}$ -primary ideal and $J_1, \dots, J_p \subset R$ be arbitrary ideals. Suppose that, for each $0 \leq i \leq p$, J_i is generated by homogeneous elements of the same degree $d_i > 0$. Let $\mathbf{n} = (n_0, n_1, \dots, n_p) \in \mathbb{N}^{p+1}$ such that $|\mathbf{n}| = \dim(R) - 1$. Then, $e_{\mathbf{n}}(J_0 \mid J_1, \dots, J_p) > 0$ if and only if for each $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq [p]$ the inequality

$$n_{j_1} + \dots + n_{j_k} \leq \ell(J_{j_1} \cdots J_{j_k}) - 1$$

holds.

Proof. First, note that $r = \dim(R) - 1$.

Since J_0 is $\mathfrak{m}$ -primary, the kernel of the canonical map $T \twoheadrightarrow T' := \mathcal{R} \otimes_R R/\mathfrak{m}$ is nilpotent. Therefore, the conditions (a), (b) in Corollary 4.3 are satisfied for T if and only if they are satisfied for T' .

Consider the $\mathbb{N}^{p+1}$ -graded domain given by

$$\mathcal{F} := \bigoplus_{i_0 \geq 0, \dots, i_p \geq 0} \left[J_0^{i_0} \right]_{i_0 d_0} \left[J_1^{i_1} \right]_{i_1 d_1} \cdots \left[J_p^{i_p} \right]_{i_p d_p} t_0^{i_0} t_1^{i_1} \cdots t_p^{i_p} \subset \mathcal{R}.$$

Since $[J_0^{i_0}]_{i_0 d_0} [J_1^{i_1}]_{i_1 d_1} \cdots [J_p^{i_p}]_{i_p d_p} \cong J_0^{i_0} J_1^{i_1} \cdots J_p^{i_p} \otimes_{\mathbf{R}} \mathbf{R}/\mathbf{m}$, we have the isomorphism $T' \cong \mathcal{F}$ and so T' is a domain.

For any $\mathfrak{K} = \{h_1, \dots, h_s\} \subseteq \{0\} \cup [p]$, since we have the natural isomorphism $\mathcal{R}/\sum_{h \notin \mathfrak{K}} \mathbf{m}_h \cong \mathcal{R}(J_{h_1}, \dots, J_{h_s}) = \mathcal{R}[J_{h_1} t_{h_1}, \dots, J_{h_s} t_{h_s}]$, it follows that

$$\dim(T'/\sum_{h \notin \mathfrak{K}} \mathbf{m}_h T') = \dim(\mathcal{R}(J_{h_1}, \dots, J_{h_s}) \otimes_{\mathbf{R}} \mathbf{R}/\mathbf{m}).$$

After using the Segre embedding we get the isomorphism $\text{MultiProj}(\mathcal{R}(J_{h_1}, \dots, J_{h_s}) \otimes_{\mathbf{R}} \mathbf{R}/\mathbf{m}) \cong \text{Proj}(\mathcal{R}(J_{h_1} \cdots J_{h_s}) \otimes_{\mathbf{R}} \mathbf{R}/\mathbf{m})$ and, accordingly, from Lemma 2.5(i) we have

$$\dim(\mathcal{R}(J_{h_1}, \dots, J_{h_s}) \otimes_{\mathbf{R}} \mathbf{R}/\mathbf{m}) = \dim(\mathcal{R}(J_{h_1} \cdots J_{h_s}) \otimes_{\mathbf{R}} \mathbf{R}/\mathbf{m}) + s - 1 = \ell(J_{h_1} \cdots J_{h_s}) + s - 1,$$

(also, see [2, Corollary 3.10]).

So, $e_{\mathbf{n}}(J_0 | J_1, \dots, J_p) > 0$ if and only if for each $\mathfrak{K} = \{h_1, \dots, h_s\} \subseteq \{0\} \cup [p]$ the inequality $n_{h_1} + \cdots + n_{h_s} \leq \dim(T'/\sum_{h \notin \mathfrak{K}} \mathbf{m}_h T') - s = \ell(J_{h_1} \cdots J_{h_s}) - 1$ holds.

For any $\mathfrak{K} = \{0, h_2, \dots, h_s\} \subseteq \{0\} \cup [p]$, as J_0 is $\mathbf{m}$ -primary, from [27, Theorem 5.1.4, Proposition 5.1.6] we obtain

$$\begin{aligned} \dim(T'/\sum_{h \notin \mathfrak{K}} \mathbf{m}_h T') &= \dim(\mathcal{R}(J_0, J_{h_2}, \dots, J_{h_s}) \otimes_{\mathbf{R}} \mathbf{R}/J_0) \\ &= \dim\left(\text{gr}_{J_0 \mathcal{R}(J_{h_2}, \dots, J_{h_s})}(\mathcal{R}(J_{h_2}, \dots, J_{h_s}))\right) \\ &= \dim(\mathbf{R}) + s - 1. \end{aligned}$$

Therefore, we only need check the inequalities corresponding to the subsets $\mathfrak{J} \subseteq [p]$, and so the result follows. $\square$

Remark 4.5. Note that in Theorem 4.4 the conditions for the positivity of $e_{\mathbf{n}}(J_0 | J_1, \dots, J_p)$ do not involve the $\mathbf{m}$ -primary ideal J_0 (see [58, Corollary 1.8(a)]).

Remark 4.6. We note that if in Theorem 4.4 we have $\ell(J_i) = \dim(\mathbf{R})$ for every $1 \leq i \leq p$, then by [29, Lemma 4.7] for each $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq [p]$ we also have $\ell(J_{j_1} \cdots J_{j_k}) = \dim(\mathbf{R})$. Therefore, by Theorem 4.4 it follows that $e_{\mathbf{n}}(J_0 | J_1, \dots, J_p) > 0$ for every $\mathbf{n} \in \mathbb{N}^{p+1}$ such that $|\mathbf{n}| = \dim(\mathbf{R}) - 1$.

5. Polymatroids

We recall that Theorem A implies that $\text{MSupp}_{\mathbb{P}}(X)$ (see Definition 2.10) is the set of integer points in a polytope when X is irreducible. In this section we explore properties of these discrete sets.

Following standard notations, we say that X is a *variety* over $\mathbb{k}$ if X is a reduced and irreducible separated scheme of finite type over $\mathbb{k}$ (see, e.g., [Tag 020C, 56]). In the

following two results we connect the theory of polymatroids (see §2.3) with $\text{MSupp}_{\mathbb{P}}(X)$ when X is a variety.

Proposition 5.1. *Let $X \subseteq \mathbb{P} = \mathbb{P}_{\mathbb{k}}^{m_1} \times_{\mathbb{k}} \cdots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_p}$ be a multiprojective variety over an arbitrary field $\mathbb{k}$. Then $\text{MSupp}_{\mathbb{P}}(X)$ is a discrete algebraic polymatroid over $\mathbb{k}$.*

Proof. We consider the associated $\mathbb{N}^p$ -graded $\mathbb{k}$ -domain R . Let ξ be the generic point of X and set $\mathbb{L} := \mathcal{O}_{X, \xi}$. For each $\mathfrak{J} \subseteq [p]$, let $X_{\mathfrak{J}} = \Pi_{\mathfrak{J}}(X) = \text{MultiProj}(R_{(\mathfrak{J})})$ and $\xi_{\mathfrak{J}}$ be the generic point of $X_{\mathfrak{J}}$, and notice that

$$\mathcal{O}_{X_{\mathfrak{J}}, \xi_{\mathfrak{J}}} = \{f/g \mid f, g \in R_{(\mathfrak{J})}, g \neq 0, \deg(f) = \deg(g)\} \subseteq \mathbb{L}.$$

For $1 \leq i \leq p$, let $\mathbb{L}_i := \mathcal{O}_{X_i, \xi_i}$. Then, for each $\mathfrak{J} \subseteq [p]$, we have that $\mathcal{O}_{X_{\mathfrak{J}}, \xi_{\mathfrak{J}}} = \bigwedge_{j \in \mathfrak{J}} \mathbb{L}_j$ and that $\dim(X_{\mathfrak{J}}) = \text{trdeg}_{\mathbb{k}}(\mathcal{O}_{X_{\mathfrak{J}}, \xi_{\mathfrak{J}}})$ (see, e.g., [20, Theorem 5.22], [22, Exercise II.3.20]). Finally, the result follows from Theorem A. $\square$

In [57, Corollary 2.8] it is shown that the conclusion of Theorem 3.12 holds if $p = 2$ and X is arithmetically Cohen-Macaulay. The following example shows that this result does not always hold for $p > 2$.

Example 5.2. Let $S = \mathbb{k}[x_1, \dots, x_{12}, y_1, \dots, y_{12}]$ be a polynomial rings with an $\mathbb{N}^{12}$ -grading induced by $\deg(x_i) = \deg(y_i) = \mathbf{e}_i$ for $1 \leq i \leq 12$. Let Δ be the simplicial complex given by the boundary of the icosahedron. We note that Δ is a Cohen-Macaulay complex (because it is a triangulation of the sphere S^2 [53, Corollary II.4.4]), but it is not a (poly)matroid (see Remark 2.15) since not every restriction is pure [53, Proposition III.3.1].

Let $J_{\Delta} = \{(x_{i_1} + y_{i_1}) \cdots (x_{i_k} + y_{i_k}) \mid \{i_1, \dots, i_k\} \notin \Delta\}$ and $X_{\Delta} = \text{MultiProj}(S/J_{\Delta}) \subset \mathbb{P} = \mathbb{P}_{\mathbb{k}}^1 \times \cdots \times \mathbb{P}_{\mathbb{k}}^1$. The definition of J_{Δ} is a modification on the definition of I_{Δ} , the Stanley-Reisner ideal of Δ with monomials in the variables $\{x_1, \dots, x_{12}\}$ [43, Chapter 1], [53, Chapter II]. It can be easily verified that I_{Δ} is the initial ideal of J_{Δ} with respect to any elimination order with $\{x_1, \dots, x_{12}\} \geq \{y_1, \dots, y_{12}\}$. Since the ideal J_{Δ} is obtained from I_{Δ} by a linear change of variables, we have a similar primary decomposition as [43, Theorem 1.7], so no component is supported on any coordinate subspace and thus J_{Δ} is saturated with respect to the irrelevant ideal of S . By [24, Corollary 3.3.5], X_{Δ} is arithmetically Cohen-Macaulay. Moreover, since Hilbert functions are preserved by Gröbner degenerations, the multidegree of X_{Δ} coincides with $\mathcal{C}(S/I_{\Delta}; \mathbf{t})$ (see Remark 2.9). Thus, $\text{MSupp}_{\mathbb{P}}(X_{\Delta})$ consists of all the incidence vectors of the facets of Δ [43, Theorem 1.7] and then it is not a polymatroid.

With Proposition 5.1 in hand, we can introduce the following class of polymatroids.

Definition 5.3. A polymatroid $\mathcal{P}$ is *Chow* over a field $\mathbb{k}$ if there exists a variety $X \subseteq \mathbb{P} = \mathbb{P}_{\mathbb{k}}^{m_1} \times_{\mathbb{k}} \cdots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_p}$ such that $\text{MSupp}_{\mathbb{P}}(X) = \mathcal{P}$.

The following statement follows as an easy corollary of the main result in [37], when $\mathbb{k}$ is an infinite field. Here we give a simple direct argument for an arbitrary field $\mathbb{k}$.

Proposition 5.4. *A linear polymatroid over an arbitrary field $\mathbb{k}$ is Chow over the same field.*

Proof. Let V be a $\mathbb{k}$ -vector space and $V_1, \dots, V_p$ be arbitrary subspaces. Let S be the polynomial ring $S = \text{Sym}(V) = \mathbb{k}[x_1, \dots, x_q]$, where $q = \dim_{\mathbb{k}}(V)$. By using the isomorphism $[S]_1 \cong V$, we identify each V_i with a $\mathbb{k}$ -subspace U_i of $[S]_1$. For each $1 \leq i \leq p$, let $\{x_{i,1}, x_{i,2}, \dots, x_{i,r_i}\} \subset [S]_1$ be a basis of the $\mathbb{k}$ -vector space U_i . Let T be the $\mathbb{N}^p$ -graded polynomial ring

$$T := \mathbb{k}[y_{i,j} \mid 1 \leq i \leq p, 0 \leq j \leq r_i, \deg(y_{i,j}) = \mathbf{e}_i].$$

Induce an $\mathbb{N}^p$ -grading on $S[t_1, \dots, t_p]$ given by $\deg(t_i) = \mathbf{e}_i$ and $\deg(x_j) = \mathbf{0}$. Consider the $\mathbb{N}^p$ -graded $\mathbb{k}$ -algebra homomorphism

$$\varphi = T \rightarrow S[t_1, \dots, t_p], \quad \begin{array}{ll} y_{i,0} \mapsto t_i & \text{for } 1 \leq i \leq p \\ y_{i,j} \mapsto x_{i,j} t_i & \text{for } 1 \leq i \leq p, 1 \leq j \leq r_i. \end{array}$$

Note that $\mathfrak{P} := \text{Ker}(\varphi)$ is an $\mathbb{N}^p$ -graded prime ideal. Set $R := T/\mathfrak{P}$ and $X := \text{MultiProj}(R)$. By construction, for each $\mathfrak{J} \subseteq [p]$, we obtain the isomorphism

$$R_{(\mathfrak{J})} \cong \mathbb{k}[x_{i,j} t_i \mid i \in \mathfrak{J}, 1 \leq j \leq r_i] [t_i \mid i \in \mathfrak{J}] \subset S[t_1, \dots, t_p];$$

thus, it is clear that

$$\begin{aligned} \dim(R_{(\mathfrak{J})}) &= \text{trdeg}_{\mathbb{k}}(\mathbb{k}[x_{i,j} t_i \mid i \in \mathfrak{J}, 1 \leq j \leq r_i] [t_i \mid i \in \mathfrak{J}]) \\ &= \text{trdeg}_{\mathbb{k}}(\mathbb{k}[x_{i,j} \mid i \in \mathfrak{J}, 1 \leq j \leq r_i] [t_i \mid i \in \mathfrak{J}]) \\ &= \dim_{\mathbb{k}}\left(\sum_{i \in \mathfrak{J}} U_i\right) + |\mathfrak{J}| = \dim_{\mathbb{k}}\left(\sum_{i \in \mathfrak{J}} V_i\right) + |\mathfrak{J}|. \end{aligned}$$

Therefore, Lemma 2.5 yields that $\dim(\Pi_{\mathfrak{J}}(X)) = \dim_{\mathbb{k}}(\sum_{i \in \mathfrak{J}} V_i)$, and so the result follows from Theorem A. $\square$

The following is the main theorem of this section. Here we summarize the results presented above to show that the class of Chow polymatroids lies in between the ones introduced in Definition 2.16.

Theorem 5.5. *Over an arbitrary field $\mathbb{k}$, we have the following inclusions of families of polymatroids*

$$(\text{Linear polymatroids}) \subseteq (\text{Chow polymatroids}) \subseteq (\text{Algebraic polymatroids}).$$

Moreover, when $\mathbb{k}$ is a field of characteristic zero, the three families coincide.

Proof. The first inclusion follows from Proposition 5.4; the second from Proposition 5.1. In the characteristic zero case linear and algebraic polymatroids coincide by [30, Corollary, Page 166] (also, see Remark 5.6). $\square$

Remark 5.6. The result mentioned above from [30] is stated for matroids but the arguments go unchanged for polymatroids.

Remark 5.7. Over finite fields there are algebraic matroids that are not linear. An example is the Non-Pappus matroid described in [47, Page 517], it is algebraic over any field of positive characteristic but not linear over any field.

Classifying linear polymatroid rank functions is a difficult problem. For linear matroids over a field of characteristic zero, the poetically titled “*The missing axiom of matroid theory is lost forever*” [59] together with a recent addition [40] shows that there is no finite list of axioms that characterize which rank functions are linear. For fields of positive characteristic, Rota conjectured in 1971 that for each field there is a list of finite restrictions. A proof of Rota’s conjecture has been announced by Geelen, Gerards, and Whittle, but expected to be several hundred of pages long. Little is known about the algebraic case. In [31] there is an example of a matroid that is not algebraic over any field: the Vamos matroid V_8 [47, Page 511]. For these reasons we do not expect further characterizations of Chow polymatroids.

We finish this section with the following question.

Question 5.8. Are all algebraic polymatroids Chow?

6. Applications

In this section we relate our results to several objects from combinatorial algebraic geometry.

6.1. Schubert polynomials

Let $\mathfrak{S}_p$ be the symmetric group on the set $[p]$. For every $i \in [p-1]$ we have the transposition $s_i := (i, i+1) \in \mathfrak{S}_p$. Recall that the set $S = \langle s_i, 1 \leq i < p \rangle$ generates $\mathfrak{S}_p$. The *length* $l(\pi)$ of a permutation π is the least amount of elements in S needed to obtain π . Alternatively, the length is equal to the number of *inversions*, i.e., $l(\pi) = \{(i, j) \in [p] \times [p] : i < j, \pi(i) > \pi(j)\}$. The permutation $\pi_0 = (p, p-1, \dots, 2, 1)$ (in one line notation) is the longest permutation, it has length $\frac{p(p-1)}{2}$.

Definition 6.1. The *Schubert polynomials* $\mathfrak{S}_\pi \in \mathbb{Z}[t_1, \dots, t_p]$ are defined recursively in the following way. First we define $\mathfrak{S}_{\pi_0} := \prod_i t_i^{p-i}$, and for any permutation π and transposition s_i with $l(s_i\pi) < l(\pi)$ we let

$$\mathfrak{S}_{s_i\pi} = \frac{\mathfrak{S}_\pi - s_i\mathfrak{S}_\pi}{t_i - t_{i+1}},$$

where $\mathfrak{S}_p$ acts on $\mathbb{Z}[t_1, \dots, t_p]$ by permutation of variables. For more information see [18, Chapter 10].

Next we define *matrix Schubert varieties* following [43, Chapter 15]. Let $\mathbb{k}$ be an algebraic closed field and $M_p(\mathbb{k})$ be the $\mathbb{k}$ -vector space of $p \times p$ matrices with entries in $\mathbb{k}$. As an affine variety we define its coordinate ring as $R_p := \mathbb{k}[x_{ij} : (i, j) \in [p] \times [p]]$. Furthermore we consider an $\mathbb{N}^p$ -grading on R_p by letting $\deg(x_{ij}) = \mathbf{e}_i$.

Definition 6.2. Let π be a permutation matrix. The matrix Schubert variety $\overline{X}_\pi \subset M_p(\mathbb{k})$ is the subvariety

$$\overline{X}_\pi = \{Z \in M_p(\mathbb{k}) \mid \text{rank}(Z_{m \times n}) \leq \text{rank}(\pi_{m \times n}) \text{ for all } m, n\},$$

where $Z_{m \times n}$ is the restriction to the first m rows and n columns. This is an irreducible variety and the prime ideal $I(\overline{X}_\pi)$ is multihomogeneous [43, Theorem 15.31]. By [43, Theorem 15.40], the Schubert polynomial $\mathfrak{S}_\pi$ equals the multidegree polynomial of the variety corresponding to the ideal $I(\overline{X}_\pi)$ (see Definition 2.10).

Following [45] we say a polynomial $f = \sum_{\mathbf{n}} c_{\mathbf{n}} t^{\mathbf{n}} \in \mathbb{Z}[t_1, \dots, t_p]$ have the Saturated Newton Polytope property (SNP for short) if $\text{supp}(f) := \{\mathbf{n} \in \mathbb{N}^p \mid c_{\mathbf{n}} > 0\} = \text{ConvexHull}\{\mathbf{n} \in \mathbb{N}^p \mid c_{\mathbf{n}} > 0\} \cap \mathbb{N}^p$, in other words, if the support of f consists of the integer points of a polytope. In [45, Conjecture 5.5] it was conjectured that the Schubert polynomials have SNP property and they even conjectured a set of defining inequalities for the Newton polytope in [45, Conjecture 5.13]. A. Fink, K. Mészáros, and A. St. Dizier confirmed the full conjecture in [15]. As noted by the authors of [26] the combination of Proposition 5.1 (they use the equivalent [3, Corollary 10.2]) and [43, Theorem 15.40] (which is also included in [26, Theorem 6]) is enough to give an alternative proof to [45, Conjecture 5.5].

Theorem 6.3. For any permutation π , the Schubert polynomial $\mathfrak{S}_\pi$ has SNP and its Newton polytope is a polymatroid polytope.

The Newton polytope of a polynomial f is by definition the convex hull of the exponents in the support of f , however in by our convention in Definition 2.10 MSupp consists of the complementary exponents. This does not change the conclusion that the resulting polytope is a polymatroid polytope.




Fig. 1. Example of a diagram in $[5] \times [5]$.

Codimensions of projections. We now use Theorem A to give a combinatorial interpretation for the codimensions of the natural projections of matrix Schubert varieties. First we need some terminology.

A *diagram* D is a subset of a $p \times p$ grid whose boxes are indexed by the set $[p] \times [p]$. The authors of [45] define a function $\theta_D : 2^{[p]} \mapsto \mathbb{Z}$ as follows: for a subset $\mathfrak{J} \subseteq [p]$ and $c \in [p]$, we construct a word $W_D^c(\mathfrak{J})$ by reading the column c of $[p] \times [p]$ from top to bottom and recording

- (if $(r, c) \notin D$ and $r \in \mathfrak{J}$,
-) if $(r, c) \in D$ and $r \notin \mathfrak{J}$,
- $\star$ if $(r, c) \in D$ and $r \in \mathfrak{J}$;

let $\theta_D^c(\mathfrak{J}) = \# \text{ paired "()" in } W_D^c(\mathfrak{J}) + \# \star \text{ in } W_D^c(\mathfrak{J})$, and finally $\theta_D(\mathfrak{J}) = \sum_{i=1}^p \theta_D^i(\mathfrak{J})$.

Example 6.4. For example, let D be the diagram depicted in Fig. 1 and $\mathfrak{J} = \{2, 3\}$, then $\theta_D(\mathfrak{J}) = 3$.

For any $\pi \in \mathfrak{S}_p$ we can define its *Rothe diagram* as

$$D_\pi := \{(i, j) \mid 1 \leq i, j \leq n, \pi(i) > j \text{ and } \pi^{-1}(j) > i\} \subset [p] \times [p].$$

For example when $\pi = 42531$ then D_π is the diagram of Fig. 1.

Theorem 6.5. Let $\pi \in \mathfrak{S}_p$, then for any $\mathfrak{J} \subseteq [p]$ the projection $\Pi_{\mathfrak{J}}(\overline{X_\pi})$ onto the rows indexed by $\mathfrak{J}$ has codimension $\theta_{D_\pi}([p]) - \theta_{D_\pi}(\mathfrak{J}')$, where $\mathfrak{J}' = [p] \setminus \mathfrak{J}$ is the complement of $\mathfrak{J}$.

Proof. In [15, Theorem 10] the authors give a proof of [45, Conjecture 5.13], which in our setup (recall the indexing in Definition 2.10) states that $\text{MSupp}(\overline{X_\pi})$ is equal to

$$\left\{ \mathbf{n} \in \mathbb{N}^p \mid \sum_{j \in \mathfrak{J}} ((p-1) - n_j) \leq \theta_{D_\pi}(\mathfrak{J}), \forall \mathfrak{J} \subsetneq [p], \sum_{j \in [p]} ((p-1) - n_j) = \theta_{D_\pi}([p]) \right\}.$$

The first inequalities can be rewritten as $(p-1)|\mathfrak{J}| - \theta_{D_\pi}(\mathfrak{J}) \leq \sum_{j \in \mathfrak{J}} n_j$, and combining them with $(p-1)p - \theta_{D_\pi}([p]) = \sum_{j \in [p]} n_j$ we obtain

$$\sum_{j \in \mathfrak{J}'} n_j \leq (p-1)|\mathfrak{J}'| - \left(\theta_{D_\pi}([p]) - \theta_{D_\pi}(\mathfrak{J}') \right).$$

By Theorem A we must have that $\text{codim}(\Pi_{\mathfrak{J}}(\overline{X_\pi})) = \theta_{D_\pi}([p]) - \theta_{D_\pi}(\mathfrak{J}')$ for every $\mathfrak{J} \subseteq [p]$, as we wanted to show. $\square$

Remark 6.6. Notice that $\theta_{D_\pi}([p])$ counts the total number of boxes in D_π , which is equal to the length of π (see [43, Definition 15.13]). So the case $\mathfrak{J} = [p]$ of Theorem 6.5 above is equivalent to the well-known fact that the codimension of a matrix Schubert variety is equal to the length of the permutation (see [43, Theorem 15.31]).

6.2. Flag varieties

We now focus on a multiprojective embedding of flag varieties. We first review some terminology. For more information the reader is referred to [18] or [4].

In this subsection we work over an algebraically closed field $\mathbb{k}$. Consider the *complete flag variety* $\text{Fl}(V)$ of a $\mathbb{k}$ -vector space V of dimension $p+1$. This variety parametrizes complete flags, i.e., sequences $V_\bullet := (V_0, \dots, V_{p+1})$ such that $\{0\} = V_0 \subset V_1 \subset V_2 \subset \dots \subset V_p \subset V_{p+1} = V$, and each V_i is a linear subspace of V of dimension i . One can embed this variety in a product of Grassmannians $\text{Fl}(V) \hookrightarrow \text{Gr}(1, V) \times \text{Gr}(2, V) \times \dots \times \text{Gr}(p, V)$ as the subvariety cut out by incidence relations.

Furthermore, each Grassmannian can be embedded in a projective space via the Plücker embedding $\iota_i : \text{Gr}(i, V) \rightarrow \mathbb{P}_{\mathbb{k}}^{m_i}$ for $1 \leq i \leq p$. By considering the product of these maps, we obtain a multiprojective embedding of $\iota : \text{Fl}(V) \hookrightarrow \mathbb{P}_{\mathbb{k}}^{m_1} \times_{\mathbb{k}} \dots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_p}$. For convenience we also call ι the *Plücker embedding*. The proposition below computes the corresponding multidegree support.

Proposition 6.7. *Let V be a $\mathbb{k}$ -vector space of dimension $p+1$ and let X be the image of the Plücker embedding $\iota : \text{Fl}(V) \hookrightarrow \mathbb{P} = \mathbb{P}_{\mathbb{k}}^{m_1} \times_{\mathbb{k}} \dots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_p}$, then*

$$\text{MSupp}_{\mathbb{P}}(X) = \left\{ \mathbf{n} \in \mathbb{N}^p \mid 1 \leq n_k \leq \sum_{j=1}^k (-j) - \sum_{i=1}^{k-1} n_i, \forall k \in [p], \sum_{j=1}^p n_j = \binom{p+1}{2} \right\}; \quad (8)$$

Proof. We need to compute the dimension of $\Pi_{\mathfrak{J}}(X)$ for each $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq [p]$. The key observation is that $\Pi_{\mathfrak{J}}(X)$ is isomorphic to the *partial flag variety* $\text{Fl}_{\mathfrak{J}}(V)$: it parametrizes flags $W_\bullet := \{0\} = V_0 \subset V_2 \subset V_3 \subset \dots \subset V_k \subset V_{k+1} = V$, where $\dim V_k = j_k$. Hence

$$\dim(\Pi_{\mathfrak{J}}(X)) = \dim(\text{Fl}_{\mathfrak{J}}(V)) = \sum_{1 \leq i < j \leq k+1} d_i d_j = \mathcal{S}(\mathfrak{J});$$

here, for each $\mathfrak{J} = \{j_1, \dots, j_k\} \subseteq [p]$, we set $\mathcal{S}(\mathfrak{J}) := \sum_{1 \leq i < j \leq k+1} d_i d_j$, where $d_i := j_i - j_{i-1}$ and by convention $j_0 := 0$, $j_{k+1} := p + 1$. For a proof of the second equality see [4, §1.2]. From Theorem A it follows that

$$\text{MSupp}_{\mathbb{P}}(X) = \left\{ \mathbf{n} \in \mathbb{N}^p \mid \sum_{j \in \mathfrak{J}} n_j \leq \mathcal{S}(\mathfrak{J}), \forall \mathfrak{J} \subseteq [p], \sum_{j=1}^p n_j = \binom{p+1}{2} \right\}. \quad (9)$$

It can be checked that the description in (9) coincides with the one in (8). $\square$

The pullbacks of the classes H_i from $\mathbb{P}_{\mathbb{k}}^{m_1} \times_{\mathbb{k}} \dots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^{m_p}$ to $\text{Fl}(V)$ are called the Schubert divisors, so Proposition 6.7 amounts to a criterion for which powers of these classes intersect. These intersections are called Grassmannian Schubert problems in [50]. In [50, Theorem 1.2] K. Purbhoo and F. Sottile give a stronger statement by providing an explicit combinatorial formula using *filtered tableau* to compute the exact intersection numbers.

6.3. A multiprojective embedding of $\overline{M}_{0,p+3}$

The moduli space $\overline{M}_{0,p+3}$ parametrizes rational stable curves with $p + 3$ marked points. Here we apply our methods to an embedding considered in [8]. The starting point is the closed embedding $\Psi_p : \overline{M}_{0,p+3} \rightarrow \overline{M}_{0,p+2} \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^p$ constructed by S. Keel and J. Tevelev in [35, Corollary 2.7]. By iterating this construction we obtain an embedding $\overline{M}_{0,p+3} \hookrightarrow \mathbb{P}_{\mathbb{k}}^1 \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^2 \times_{\mathbb{k}} \dots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^p$ (see [8, Corollary 3.2]). In [8], R. Cavalieri, M. Gillespie, and L. Monin computed the corresponding multidegree which turns out to be related to parking functions. As an easy consequence of our Theorem A, we can compute its support.

Proposition 6.8. *Let X be the image of $\overline{M}_{0,p+3} \hookrightarrow \mathbb{P} = \mathbb{P}_{\mathbb{k}}^1 \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^2 \times_{\mathbb{k}} \dots \times_{\mathbb{k}} \mathbb{P}_{\mathbb{k}}^p$, then*

$$\text{MSupp}_{\mathbb{P}}(X) = \left\{ \mathbf{n} \in \mathbb{N}^p \mid \sum_{i=1}^k n_i \leq k, \forall 1 \leq k \leq p-1, \sum_{i=1}^p n_i = p \right\}. \quad (10)$$

Proof. First, as explained in [8, §3] we have $\dim(\Pi_{[p]}(X)) = \dim(\Pi_{\{p\}}(X)) = p$. Also, by construction $\Pi_{[p-1]}(X) \cong \overline{M}_{0,p+2}$, and thus $\dim(\Pi_{[i-1]}(X)) = i-1$ for every $2 \leq i \leq p$. So, by induction one gets $\dim(\Pi_{\{i\}}(X)) = i$ for all $1 \leq i \leq p$.

To use Theorem A, we must compute $\dim(\Pi_{\mathfrak{J}}(X))$ for all $\mathfrak{J} \subseteq [p]$. Let $m := \max\{j \mid j \in \mathfrak{J}\}$, then as explained above we have $\dim(\Pi_{[m]}(X)) = \dim(\Pi_{\{m\}}(X)) = m$ and so we must have $\dim(\Pi_{\mathfrak{J}}(X)) = m$. By Theorem A we obtain that $\text{MSupp}_{\mathbb{P}}(X)$ is equal to

$$\{\mathbf{n} \in \mathbb{N}^p \mid \sum_{i \in \mathfrak{J}} n_i \leq \max\{j \mid j \in \mathfrak{J}\}, \forall \mathfrak{J} \subseteq [p], \sum_{i=1}^p n_i = p\},$$

but it is straightforward to check that the inequalities in (10) are enough to describe the same set. $\square$

The cardinality of $\text{MSupp}_{\mathbb{P}}(X)$ is equal to the Catalan number C_n (see [54, Exercise 86]). For a comprehensible survey on Catalan numbers see [54].

6.4. Mixed volumes

In this subsection we assume $\mathbb{k}$ is an algebraically closed field. We begin by reviewing the definition of mixed volumes of convex bodies, as a general reference see [14, Chapter IV]. Let $\mathbf{K} = (K_1, \dots, K_p)$ be a p -tuple of convex bodies in $\mathbb{R}^d$. The volume polynomial $v(\mathbf{K}) \in \mathbb{Z}[w_1, \dots, w_p]$ is defined as

$$v(K_1, \dots, K_p) := \text{Vol}_d(w_1 K_1 + \dots + w_p K_p).$$

This is a homogeneous polynomial of degree d . If the coefficients of $v(\mathbf{K})$ are written as $\binom{d}{\mathbf{n}} V(\mathbf{K}; \mathbf{n}) w^{\mathbf{n}}$, then the numbers $V(\mathbf{K}; \mathbf{n})$ are called the *mixed volumes* of $\mathbf{K}$. A natural question to ask is: when are mixed volumes positive? The relation between mixed volumes and toric varieties (see (11) below) together with Theorem A allows us to give another proof of a classical theorem formulated on the non-vanishing of mixed volumes [51, Theorem 5.1.8].

Theorem 6.9. *Let $\mathbf{K} = (K_1, \dots, K_p)$ be a p -tuple of convex bodies in $\mathbb{R}^d$. Then, $V(\mathbf{K}; \mathbf{n}) > 0$ if and only if $\sum_{i=1}^p n_i = d$ and $\sum_{i \in \mathfrak{J}} n_i \leq \dim(\sum_{i \in \mathfrak{J}} K_i)$ for every subset $\mathfrak{J} \subseteq [p]$.*

We first indicate how to reduce to the case of polytopes. The basic idea is that convex bodies can be approximated by polytopes in the Hausdorff metric [51, Section 1.8]. However, the condition for positivity as stated in Theorem 6.9 is a priori not stable under limits. To fix this we invoke an equivalent condition more suitable for the limiting argument.

Lemma 6.10. *It suffices to show Theorem 6.9 for polytopes.*

Proof. This follows from two facts. The first is that mixed volumes $V(\mathbf{K}; \mathbf{n})$ are continuous [51, Theorem 5.1.7] and monotonous [51, Equation 5.25] on each entry. The second fact is that for a given sequence $\mathbf{K} = (K_1, \dots, K_p) \subset (\mathbb{R}^d)^p$ of convex bodies, by [51, Lemma 5.1.9] the following conditions are equivalent:

- (1) $\sum_{i=1}^p n_i = d$ and $\sum_{i \in \mathfrak{J}} n_i \leq \dim(\sum_{i \in \mathfrak{J}} K_i)$ for every subset $\mathfrak{J} \subseteq [p]$.
- (2) There exist line segments $S_{i,1}, S_{i,2}, \dots, S_{i,n_i} \subseteq K_i$, for every i , such that $\{S_{i,j}\}_{1 \leq i \leq p, 1 \leq j \leq n_i}$ has segments in d linearly independent directions.

We now assume the statement of Theorem 6.9 is true when each K_i is a polytope and show that it follows in the case where each K_i is an arbitrary convex body.

If $V(\mathbf{K}; \mathbf{n}) > 0$ then by continuity we can find polytopes $\mathbf{P} = (P_1, \dots, P_p)$ with $V(\mathbf{P}; \mathbf{n}) > 0$ and $P_i \subseteq K_i$ for each $i \in [p]$. By assumption, the sequence $\mathbf{P}$ satisfies condition (2) above and hence so does the sequence $\mathbf{K}$.

Conversely, suppose condition (2) holds for $\mathbf{K}$, then by continuity we can find polytopes $\mathbf{P} = (P_1, \dots, P_p)$ with $P_i \subseteq K_i$ arbitrarily close so that (2) holds for $\mathbf{P}$ too. Due to the assumption, we have $V(\mathbf{P}; \mathbf{n}) > 0$ and thus $V(\mathbf{K}; \mathbf{n}) \geq V(\mathbf{P}; \mathbf{n}) > 0$ by monotonicity. $\square$

To finish the proof of Theorem 6.9 we need some preliminary results about toric varieties and lattice polytopes. As an initial step we recall some facts about basepoint free divisors; a general reference is [12, Section 6]. Let Σ be a fan and let P be a lattice polytope whose normal fan coarsens Σ . Then, P induces a basepoint free divisor D_P in the toric variety Y_Σ [12, Proposition 6.2.5]. Here, being basepoint free means that the complete linear series $|D_P|$ induces a morphism $\phi_P : Y_\Sigma \rightarrow \mathbb{P}_k^{m_i}$ for some $m_i \in \mathbb{N}$ such that $\phi^*(H) = D_P \in A^*(Y_\Sigma)$, where H is the class of a hyperplane in the projective space $\mathbb{P}_k^{m_i}$.

Lemma 6.11. *Let $K_1, \dots, K_p$ be lattice polytopes and let $K := K_1 + \dots + K_p$ be their Minkowski sum. Let Y be the toric variety associated to Σ , the normal fan of K , then for each $\mathfrak{J} \subseteq [p]$ we have a map $\phi_{\mathfrak{J}} : Y \rightarrow \prod_{j \in \mathfrak{J}} \mathbb{P}_k^{m_j}$ such that $\dim(\phi_{\mathfrak{J}}(Y)) = \dim(\sum_{i \in \mathfrak{J}} K_i)$.*

Proof. The fan Σ is the common refinement of the normal fans of $K_1, \dots, K_p$ [61, Proposition 7.12], so each K_i induces a basepoint free divisor D_i on $Y = Y_\Sigma$ and thus also a map $\phi_i : Y \rightarrow \mathbb{P}_k^{m_i}$. By the universal property of fiber products these maps induce a canonical map $\phi_{\mathfrak{J}} : Y \rightarrow \prod_{j \in \mathfrak{J}} \mathbb{P}_k^{m_j}$ for each $\mathfrak{J} \subseteq [p]$. It remains to compute the dimensions of the images of these maps.

By composing with the Segre embedding $\varphi_{\mathfrak{J}} : \prod_{j \in \mathfrak{J}} \mathbb{P}_k^{m_j} \rightarrow \mathbb{P}_k^m$ we obtain a map $\varphi_{\mathfrak{J}} \circ \phi_{\mathfrak{J}} : Y \rightarrow \mathbb{P}_k^m$. Let H be the class of a hyperplane in $A^*(\mathbb{P}_k^m)$, we have that $\varphi_{\mathfrak{J}}^*(H) = \sum_{i \in \mathfrak{J}} H_i \in A^*(\prod_{j \in \mathfrak{J}} \mathbb{P}_k^{m_j})$ where each H_i is the pullback of a hyperplane in the i -th factor (see, e.g., [22, Exercise 5.11]). Then $(\varphi_{\mathfrak{J}} \circ \phi_{\mathfrak{J}})^*(H) = \sum_{i \in \mathfrak{J}} D_i \in A^*(Y)$. This means that the morphism $\varphi_{\mathfrak{J}} \circ \phi_{\mathfrak{J}}$ corresponds to the complete linear series $|D_{K_{\mathfrak{J}}}|$ where $K_{\mathfrak{J}} = \sum_{j \in \mathfrak{J}} K_j$, hence the image has dimension $\dim(K_{\mathfrak{J}}) = \dim(\sum_{j \in \mathfrak{J}} K_j)$ [12, Theorem 6.1.22]. $\square$

Lemma 6.12. *In the setup of Lemma 6.11, if $\mathfrak{J} = [p]$ then after scaling each polytope if necessary, $\phi = \phi_{\mathfrak{J}}$ is an embedding.*

Proof. By construction the normal fan of $K_{[p]} = \sum_{i=1}^p K_i$ is Σ , so the corresponding divisor is ample [12, Theorem 6.1.14]. By replacing the list of polytopes by large enough scalings we obtain a very ample divisor, hence an embedding. $\square$

Proof of Theorem 6.9. By using Lemma 6.10, we can assume that each K_i is a polytope. Additionally, we can reduce to the case where each K_i is a lattice polytope since any polytope can be approximated by lattice polytopes (see [17, Page 120]). Let $K = K_1 + \cdots + K_p$ and let Y be the toric projective variety associated to the normal fan of K . Each lattice polytope K_i induces a basepoint free divisor D_i on Y . As explained in [17, Eq. (2), Page 116], the fundamental connection between mixed volumes and intersection products is given by the following equation

$$V(K; \mathbf{n}) = (D_1^{n_1} \cdots D_p^{n_p}) / d!, \quad (11)$$

where the numerator is the intersection product of the divisors in Y . Notice that positivity of mixed volumes is unchanged by scaling so whenever needed we can scale each polytope.

By Lemma 6.12 we have an embedding $\phi : Y \rightarrow \prod_{i=1}^p \mathbb{P}_k^{m_i}$ such that the pullback of each $H_i \in A^*(\prod_{i=1}^p \mathbb{P}_k^{m_i})$ is D_i . By using the projection formula [19, Proposition 2.5(c)], we can consider the product $(H_1^{n_1} \cdots H_p^{n_p}) / d!$ instead of the one in (11). From the fact that Y is irreducible we are now in the Setup 2.1. So, the result follows by Remark 2.8, Theorem A, and Lemma 6.11. $\square$

Acknowledgments

We thank the reviewer for his/her suggestions for the improvement of this work. We would like to thank Chris Eur, Maria Gillespie, June Huh, David Speyer, and Mauricio Velasco for useful conversations. We are also grateful with Frank Sottile for useful comments on an earlier version (in particular for the simplification of the statement of Proposition 6.7). Special thanks to Brian Osserman for many insightful conversations and encouragements. The computer algebra system Macaulay2 [21] was of great help to compute several examples in the preparation of this paper.

References

- [1] P.B. Bhattacharya, The Hilbert function of two ideals, *Proc. Camb. Philos. Soc.* 53 (1957) 568–575.
- [2] C. Bivià-Ausina, J. Montaña, Analytic spread and integral closure of decomposable modules, *arXiv preprint arXiv:2001.08313*, 2020.
- [3] P. Brändén, J. Huh, Lorentzian polynomials, *arXiv:1902.03719*, 2020.
- [4] M. Brion, Lectures on the Geometry of Flag Varieties, *Topics in Cohomological Studies of Algebraic Varieties*, Trends Math., Birkhäuser, Basel, 2005, pp. 33–85.
- [5] M. Brodmann, J. Rung, Local cohomology and the connectedness dimension in algebraic varieties, *Comment. Math. Helv.* 61 (3) (1986) 481–490.
- [6] W. Bruns, J. Herzog, *Cohen-Macaulay Rings*, 2nd ed., Cambridge Studies in Advanced Mathematics, Cambridge University Press, 1998.
- [7] F. Castillo, F. Liu, Deformation cones of nested braid fans, *Int. Math. Res. Not.* (20 2006) 20, rnaa090.
- [8] R. Cavalieri, M. Gillespie, L. Monin, Projective embeddings of $\overline{M}_{0,n}$ and parking functions, *arXiv preprint arXiv:1912.12343*, 2020.
- [9] Y. Cid-Ruiz, Mixed multiplicities and projective degrees of rational maps, *J. Algebra* (2020), *arXiv preprint arXiv:2001.00547*, in press.

- [10] Y. Cid-Ruiz, A. Simis, Degree of rational maps and specialization, *Int. Math. Res. Not.* (2020) rnaa183, <https://doi.org/10.1093/imrn/rnaa183>.
- [11] A. Conca, E. De Negri, E. Gorla, Cartwright-Sturmfels ideals associated to graphs and linear spaces, *J. Algebraic Comb.* 2 (3) (2018) 231–257.
- [12] D.A. Cox, J.B. Little, H.K. Schenck, *Toric Varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011.
- [13] D. Eisenbud, *Commutative Algebra with a View Towards Algebraic Geometry*, Graduate Texts in Mathematics, vol. 150, Springer-Verlag, 1995.
- [14] G. Ewald, *Combinatorial Convexity and Algebraic Geometry*, Graduate Texts in Mathematics, vol. 168, Springer-Verlag, New York, 1996.
- [15] A. Fink, K. Mészáros, A.St. Dizier, Schubert polynomials as integer point transforms of generalized permutahedra, *Adv. Math.* 332 (2018) 465–475.
- [16] H. Flenner, L. O’Carroll, W. Vogel, *Joins and Intersections*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 1999.
- [17] W. Fulton, *Introduction to Toric Varieties: The William H. Roever Lectures in Geometry*, Annals of Mathematics Studies, vol. 131, Princeton University Press, Princeton, NJ, 1993.
- [18] W. Fulton, *Young Tableaux: With Applications to Representation Theory and Geometry*, London Mathematical Society Student Texts, vol. 35, Cambridge University Press, Cambridge, 1997.
- [19] W. Fulton, *Intersection theory*, second, in: *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge (Results in Mathematics and Related Areas. 3rd Series)*, in: *A Series of Modern Surveys in Mathematics*, vol. 2, Springer-Verlag, Berlin, 1998.
- [20] U. Görtz, T. Wedhorn, *Algebraic Geometry I: Schemes with Examples and Exercises*, Advanced Lectures in Mathematics, Vieweg + Teubner, Wiesbaden, 2010.
- [21] D.R. Grayson, M.E. Stillman, Macaulay2, a software system for research in algebraic geometry, Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [22] R. Hartshorne, *Algebraic Geometry*, Graduate Texts in Mathematics, vol. 52, Springer-Verlag, New York-Heidelberg, 1977.
- [23] M. Herrmann, E. Hyry, J. Ribbe, Z. Tang, Reduction numbers and multiplicities of multigraded structures, *J. Algebra* 197 (2) (1997) 311–341.
- [24] J. Herzog, T. Hibi, *Monomial Ideals*, Graduate Texts in Mathematics, vol. 260, Springer-Verlag, London, Ltd., London, 2011.
- [25] J. Huh, Milnor numbers of projective hypersurfaces and the chromatic polynomial of graphs, *J. Am. Math. Soc.* 25 (3) (2012) 907–927.
- [26] J. Huh, J.P. Matherne, K. Mészáros, A.St. Dizier, Logarithmic concavity of Schur and related polynomials, 2019.
- [27] C. Huneke, I. Swanson, *Integral Closure of Ideals, Rings, and Modules*, vol. 13, Cambridge University Press, 2006.
- [28] E. Hyry, The diagonal subring and the Cohen-Macaulay property of a multigraded ring, *Trans. Am. Math. Soc.* 351 (6) (1999) 2213–2232.
- [29] E. Hyry, Cohen-Macaulay multi-Rees algebras, *Compos. Math.* 130 (3) (2002) 319–343.
- [30] A.W. Ingleton, Representation of matroids, in: *Combinatorial Mathematics and Its Applications*, Proc. Conf., Oxford, 1969, Academic Press, London, 1971, pp. 149–167.
- [31] A.W. Ingleton, R.A. Main, Non-algebraic matroids exist, *Bull. Lond. Math. Soc.* 7 (1975) 144–146.
- [32] J.-P. Jouanolou, *Théorèmes de Bertini et applications*, Université Louis Pasteur, Département de Mathématique, Institut de Recherche Mathématique Avancée, Strasbourg, 1979.
- [33] D. Katz, S. Mandal, J.K. Verma, Hilbert functions of bigraded algebras, in: *Commutative Algebra*, Trieste, 1992, World Sci. Publ., River Edge, NJ, 1994, pp. 291–302.
- [34] K. Kaveh, A.G. Khovanskii, Complete intersections in spherical varieties, *Sel. Math. New Ser.* 22 (4) (2016) 2099–2141.
- [35] S. Keel, J. Tevelev, Equations for $\overline{M}_{0,n}$, *Int. J. Math.* 20 (9) (2009) 1159–1184.
- [36] A. Knutson, E. Miller, Gröbner geometry of Schubert polynomials, *Ann. Math.* (2) 161 (3) (2005) 1245–1318.
- [37] B. Li, Images of rational maps of projective spaces, *Int. Math. Res. Not.* 13 (2018) 4190–4228.
- [38] L. Lovász, *Submodular Functions and Convexity*, Mathematical Programming: The State of the Art (Bonn, 1982) Springer, Berlin, 1983, pp. 235–257.
- [39] H. Matsumura, *Commutative Ring Theory*, 1st ed., Cambridge Studies in Advanced Mathematics, vol. 8, Cambridge University Press, 1989.
- [40] D. Mayhew, M. Newman, G. Whittle, Yes, the ‘missing axiom’ of matroid theory is lost forever, *Trans. Am. Math. Soc.* 370 (8) (2018) 5907–5929.

- [41] M. Michałek, B. Sturmfels, C. Uhler, P. Zwiernik, Exponential varieties, *Proc. Lond. Math. Soc.* (3) 112 (1) (2016) 27–56.
- [42] M. Michałek, L. Monin, J. Wiśniewski, Maximum likelihood degree and space of orbits of a C^* action, arXiv preprint, arXiv:2004.07735, 2020.
- [43] E. Miller, B. Sturmfels, *Combinatorial Commutative Algebra*, Graduate Texts in Mathematics, vol. 227, Springer-Verlag, New York, 2005.
- [44] F. Mohammadi, C. Uhler, C. Wang, J. Yu, Generalized permutohedra from probabilistic graphical models, *SIAM J. Discrete Math.* 32 (1) (2018) 64–93.
- [45] C. Monical, N. Tokcan, A. Yong, Newton polytopes in algebraic combinatorics, *Sel. Math. New Ser.* 25 (5) (2019) 66.
- [46] K. Murota, *Discrete Convex Analysis*, SIAM Monographs on Discrete Mathematics and Applications, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2003.
- [47] J.G. Oxley, *Matroid Theory*, Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1992.
- [48] A. Postnikov, Permutohedra, associahedra, and beyond, *Int. Math. Res. Not.* 6 (2009) 1026–1106.
- [49] A. Postnikov, V. Reiner, L. Williams, Faces of generalized permutohedra, *Doc. Math.* 13 (2008) 207–273.
- [50] K. Purbhoo, F. Sottile, A Littlewood-Richardson rule for Grassmannian permutations, *Proc. Am. Math. Soc.* 137 (6) (2009) 1875–1882.
- [51] R. Schneider, *Convex Bodies: The Brunn-Minkowski Theory*, Encyclopedia of Mathematics and Its Applications, vol. 151, Cambridge University Press, Cambridge, 2014.
- [52] A. Schrijver, Matroids, trees, stable sets, Chapters 39–69, in: *Combinatorial Optimization. Polyhedra and Efficiency*, Vol. B, in: *Algorithms and Combinatorics*, vol. 24, Springer-Verlag, Berlin, 2003.
- [53] R.P. Stanley, *Combinatorics and Commutative Algebra*, 2, Birkhäuser, Boston, 1996.
- [54] R.P. Stanley, *Catalan Numbers*, Cambridge University Press, New York, 2015.
- [55] B. Teissier, Cycles évanescents, sections planes et conditions de Whitney, *Astérisque* 7 (8) (1973) 285–362.
- [56] The Stacks project authors, the stacks project, 2019.
- [57] N.V. Trung, Positivity of mixed multiplicities, *Math. Ann.* 319 (1) (2001) 33–63.
- [58] N.V. Trung, J. Verma, Mixed multiplicities of ideals versus mixed volumes of polytopes, *Trans. Am. Math. Soc.* 359 (10) (2007) 4711–4727.
- [59] P. Vámos, The missing axiom of matroid theory is lost forever, *J. Lond. Math. Soc.* (2) 18 (3) (1978) 403–408.
- [60] B.L. Van der Waerden, On Hilbert’s Function, Series of Composition of Ideals and a Generalization of the Theorem of Bezout, *Proc. Roy. Acad. Amsterdam*, vol. 31, 1929, pp. 749–770.
- [61] G.M. Ziegler, *Lectures on Polytopes*, Graduate Texts in Mathematics, vol. 152, Springer-Verlag, New York, 1995.