

Open r -Spin Theory I: Foundations

Alexandr Buryak^{1,2,*}, Emily Clader³ and Ran J. Tessler⁴

¹Faculty of Mathematics, National Research University Higher School of Economics, 6 Usacheva str., Moscow, 119048, Russian Federation,

²Novosibirsk State University, 1 Pirogova str., Novosibirsk, 630090,

Russian Federation, ³San Francisco State University, San Francisco, CA

94132-1722, USA, and ⁴Incumbent of the Lilian and George Lytle Career Development Chair: Department of Mathematics, Weizmann Institute of Science, POB 26, Rehovot 7610001, Israel

*Correspondence to be sent to: e-mail: buryaksh@gmail.com

We lay the foundation for a version of r -spin theory in genus zero for Riemann surfaces with boundary. In particular, we define the notion of r -spin disks, their moduli space, and the Witten bundle; we show that the moduli space is a compact smooth orientable orbifold with corners, and we prove that the Witten bundle is canonically relatively oriented relative to the moduli space. In the sequel to this paper, we use these constructions to define open r -spin intersection theory and relate it to the Gelfand–Dickey hierarchy, thus providing an analog of Witten's r -spin conjecture in the open setting.

1 Introduction

On a smooth marked curve $(C; z_1, \dots, z_n)$, an *r -spin structure* is a line bundle S together with an isomorphism

$$S^{\otimes r} \cong \omega_C \left(- \sum_{i=1}^n a_i [z_i] \right),$$

Received March 13, 2020; Revised October 1, 2020; Accepted November 6, 2020
Communicated by Prof. Jake Solomon

where $a_i \in \{0, 1, \dots, r-1\}$. There is a natural compactification $\overline{\mathcal{M}}_{g, \{a_1, \dots, a_n\}}^{1/r}$ of the moduli space of r -spin structures on smooth curves, and this space admits a virtual fundamental class c_W known as *Witten's class*. In genus zero, Witten's class is defined by

$$c_W := e(R^1\pi_*\mathcal{S})^\vee, \quad (1.1)$$

where $\pi : \mathcal{C} \rightarrow \overline{\mathcal{M}}_{0, \{a_1, \dots, a_n\}}^{1/r}$ is the universal curve and \mathcal{S} is the universal r -spin structure. In higher genus, on the other hand, $R^1\pi_*\mathcal{S}$ may not be a vector bundle, and there are several (all much more intricate) versions of the definition of Witten's class [6, 7, 13, 21, 25].

Given any of these definitions, one defines the *closed r -spin intersection numbers* by

$$\langle \tau_{d_1}^{a_1} \cdots \tau_{d_n}^{a_n} \rangle_g^{\overline{\mathcal{M}}_{g, \{a_1, \dots, a_n\}}^{1/r}} := r^{1-g} \int_{\overline{\mathcal{M}}_{g, \{a_1, \dots, a_n\}}^{1/r}} c_W \cap \psi_1^{d_1} \cdots \psi_n^{d_n}, \quad (1.2)$$

where $\psi_1, \dots, \psi_n \in H^2(\overline{\mathcal{M}}_{g, \{a_1, \dots, a_n\}}^{1/r})$ are the 1st Chern classes of the cotangent line bundles at the n marked points. This theory has received a great deal of attention in recent years; for example, it led to a proof of a conjecturally complete set of tautological relations on $\overline{\mathcal{M}}_{g, n}$ [23], and it is a special case of Fan–Jarvis–Ruan–Witten theory [13] as well as the gauged linear sigma model [14]. For our purposes, perhaps the most interesting feature of r -spin theory was proven by Faber–Shadrin–Zvonkine [12]: after a simple change of variables, the generating function of the closed r -spin intersection numbers becomes a tau function of the r th Gelfand–Dickey hierarchy. This statement generalizes Witten's celebrated conjecture (proven by Kontsevich) regarding the generating function of ψ -integrals on $\overline{\mathcal{M}}_{g, n}$.

A different direction in which the intersection theory of $\overline{\mathcal{M}}_{g, n}$ can be generalized is the consideration of Riemann surfaces with boundary. This work was undertaken by Pandharipande, Solomon, and the 3rd author in [24], in which a moduli space $\overline{\mathcal{M}}_{0, k, l}$ was constructed that parameterizes tuples $(\Sigma; x_1, \dots, x_k; z_1, \dots, z_l)$ where Σ is a stable disk, $x_i \in \partial\Sigma$ are boundary marked points, and $z_j \in \Sigma \setminus \partial\Sigma$ are internal marked points. The authors also constructed intersection numbers on $\overline{\mathcal{M}}_{0, k, l}$ that can be viewed as integrals of ψ -classes at the internal marked points. This construction was extended in [28] to all genera, yielding a generating function F^o of open intersection numbers. In order to define the extension, Solomon and the 3rd author introduced *graded*

2-spin structures and proved that the moduli space of 2-spin surfaces with boundary is canonically oriented. The open analog of Witten's conjecture was proved by the 1st and 3rd authors in [2, 5, 29], relating $\exp(F^0)$ to the wave function of the KdV hierarchy.

Combining r -spin theory with open theory, one might hope to define *open r -spin theory* and generalize Witten's conjecture to this setting. In order to do so even in genus zero, though, one first must define an appropriate open r -spin moduli space $\overline{\mathcal{M}}_{0,k,\{a_1,\dots,a_l\}}^{1/r}$ and an open analog of Witten's bundle $(R^1\pi_*\mathcal{S})^\vee$. We carry out the construction of such a moduli space in this paper, referring to its objects as graded r -spin disks.

1.1 Moduli space of graded r -spin disks

Graded r -spin disks are defined, roughly, as follows. Let C be an orbifold curve equipped with an involution $\phi : C \rightarrow C$ that realizes the coarse underlying curve $|C|$ as a union of two Riemann surfaces Σ and $\overline{\Sigma}$ (where $\overline{\Sigma}$ is obtained from Σ by reversing the complex structure) glued along their common boundary:

$$|C| = \Sigma \cup_{\partial \Sigma} \overline{\Sigma}.$$

Let $z_1, \dots, z_l \in \Sigma \setminus \partial \Sigma$ be a collection of internal marked points, let $\bar{z}_i := \phi(z_i) \in \overline{\Sigma}$ be their conjugates, and let $x_1, \dots, x_k \in \partial \Sigma$ be a collection of boundary marked points. On such a curve, a *graded r -spin structure* with twists a_1, \dots, a_l is an orbifold line bundle S on C together with an isomorphism

$$|S|^{\otimes r} \cong \omega_{|C|} \otimes \mathcal{O} \left(- \sum_{i=1}^l a_i [z_i] - \sum_{i=1}^l a_i [\bar{z}_i] - \sum_{j=1}^k (r-2) [x_j] \right)$$

on the coarse underlying curve $|C|$, an anti-holomorphic involution $\tilde{\phi} : S \rightarrow S$ lifting ϕ , and a certain orientation of $(S|_{\partial \Sigma \setminus \{x_j\}})^{\tilde{\phi}}$, which we refer to as a *grading*. In what follows, we prove that there exists a moduli space $\overline{\mathcal{M}}_{0,k,\{a_1,\dots,a_l\}}^{1/r}$ of graded r -spin disks with twists a_1, \dots, a_l , and that this moduli space is a compact, orientable, effective, smooth orbifold with corners.

On $\overline{\mathcal{M}}_{0,k,\{a_1,\dots,a_l\}}^{1/r}$, there is an *open Witten bundle*, which is a real vector bundle, defined roughly as

$$\mathcal{W} := (R^0\pi_*(\mathcal{S}^\vee \otimes \omega_\pi))_+ = (R^1\pi_*\mathcal{S})_-^\vee,$$

where “+” denotes the space of $\tilde{\phi}$ -invariant sections and “−” the space of $\tilde{\phi}$ -anti-invariant sections. There are also cotangent line bundles $\mathbb{L}_1, \dots, \mathbb{L}_l$ at the internal marked points. We define these bundles carefully below and explore their behavior under forgetful morphisms and restriction to boundary strata.

It is straightforward to show that the cotangent line bundles have canonical complex orientations. The open Witten bundle, on the other hand, is a real vector bundle and hence it is not clear that it is orientable at all. One of the main results of this paper is that not only is \mathcal{W} orientable, but it carries a *canonical relative orientation* relative to the moduli space; the grading plays a central role in the construction of this canonical orientation. We also analyze the behavior of the canonical orientation under restriction to boundary strata.

1.2 Companion works

This paper lays the foundations for the sequel [4], in which we use the construction of $\overline{\mathcal{M}}_{0,k, \{a_1, \dots, a_l\}}^{1/r}$ and its associated bundles to define genus-zero open r -spin intersection numbers and prove an open r -spin version of Witten’s conjecture. In particular, in [4], we calculate all the genus-zero open r -spin intersection numbers and prove an explicit relationship between their generating function and the genus-zero part of the Gelfand–Dickey wave function. In addition to verifying the generalization of Witten’s conjecture in genus zero, this leads to a conjecture for higher-genus open intersection numbers.

The content of [4] also illuminates an intriguing connection between open r -spin theory and an extension of closed r -spin theory, in which one allows a single marked point with twist -1 . We define this “closed extended r -spin theory” carefully in the companion paper [3] to this work, and in [4], we make the correspondence between the two theories precise.

1.3 Plan of the paper and the main ideas

The structure of the current paper is as follows. In Section 2, we define r -spin disks and introduce the notion of a grading, and in Section 3, we describe the moduli space of graded r -disks, its orbifold structure, and its orientation. Section 4 contains the definition of the cotangent line bundles \mathbb{L}_i and the open Witten bundle \mathcal{W} , as well as an investigation of the behavior of these bundles under certain important morphisms.

In Section 5, we establish the canonical relative orientation of \mathcal{W} and analyze its behavior under the relevant morphisms. This section is the main technical contribution of this paper, and in particular, it is where the importance of the grading is manifested;

without the grading, as we discuss in Remark 2.8, the Witten bundle may not be canonically oriented relative to the moduli space. However, equipped with the grading, for any collection of internal twists and a specific number of boundary twists, we are able to produce in Construction 5.6 an explicit basis for the fiber of the Witten bundle. The canonical relative orientation of the Witten bundle for any collection of twists arises from there by analyzing the behavior of this basis under degenerations of the underlying surface.

2 Graded r -Spin Disks

We denote by $[n]$ the set $\{1, 2, \dots, n\}$ and by \mathbb{N} the set of natural numbers $\mathbb{Z}_{\geq 1}$. Throughout what follows, a *marking* of a set A is a function

$$m : A \rightarrow 2^{\mathbb{N}}$$

such that, for all distinct $a, a' \in A$, we have $m(a) \cap m(a') = \emptyset$. A marking is *strict* if \emptyset is not in its image.

Given a marking, we identify elements of $A \setminus m^{-1}(\emptyset)$ with their images in $2^{\mathbb{N}}$, and if the image is a singleton, we identify it with an element of \mathbb{N} . Such functions are used in what follows to label the marked points on a curve; the possibility of marking some points by \emptyset or with a set is desired to handle marked points that arise via normalization of a nodal curve.

2.1 Smooth r -spin surfaces

Recall that an *orbifold Riemann surface* is a smooth, proper, possibly disconnected, one-dimensional Deligne–Mumford stack over \mathbb{C} . We sometimes refer to such a surface as *closed*, to distinguish it from the Riemann surfaces considered below that may have boundary.

A (smooth) *marked orbifold Riemann surface with boundary* is a tuple

$$(C, \phi, \Sigma, \{z_i\}_{i \in I}, \{x_j\}_{j \in B}, m^I, m^B),$$

in which

- (i) C is a (closed) orbifold Riemann surface;
- (ii) $\phi : C \rightarrow C$ is an anti-holomorphic involution (*conjugation*) that realizes the coarse underlying Riemann surface $|C|$ topologically as two Riemann

surfaces Σ and $\bar{\Sigma}$ (where $\bar{\Sigma}$ is obtained from Σ by reversing the complex structure) glued along their common boundary $\partial\Sigma = \partial\bar{\Sigma} = \text{Fix}(|\phi|)$:

$$|C| = \Sigma \cup_{\partial\Sigma} \bar{\Sigma};$$

- (iii) $z_i \in C$ is a collection of distinct points (the *internal marked points*) labeled by the set I , whose images in $|C|$ lie in $\Sigma \setminus \partial\Sigma$, with *conjugate marked points* $\bar{z}_i := \phi(z_i)$;
- (iv) $x_j \in \text{Fix}(\phi)$ is a collection of distinct points (the *boundary marked points*) labeled by the set B , whose images in $|C|$ lie in $\partial\Sigma$;
- (v) the only nontrivial isotropy of C occurs at the (internal, conjugate, and boundary) marked points;
- (vi) $m^I : I \rightarrow 2^{\mathbb{N}}$ and $m^B : B \rightarrow 2^{\mathbb{N}}$ are maps such that, for any connected component C' of C with marked points labeled by $I' \subseteq I$ and $B' \subseteq B$, the restrictions $m^I|_{I'}$ and $m^B|_{B'}$ are markings, and whenever $C' \cap \phi(C') \neq \emptyset$, the marking $m^I|_{I'}$ is strict.

A marked orbifold Riemann surface C is *stable* if each genus-zero connected component has at least three marked points (including conjugate marked points) and each genus-one connected component has at least one marked point.

We observe that the choice of a preferred half $\Sigma \subseteq |C|$, which is a part of the data of an orbifold Riemann surface with boundary, endows $\partial\Sigma$ with a canonical orientation. In what follows, we typically suppress ϕ from the notation and write \bar{x} for $\phi(x)$ when x lies in Σ .

An *isomorphism* of marked orbifold Riemann surfaces with boundary

$$(C_1, \phi_1, \Sigma_1, \{z_{1,i}\}_{i \in I}, \{x_{1,j}\}_{j \in B}, m_1^{I_1}, m_1^{B_1}) \cong (C_2, \phi_2, \Sigma_2, \{z_{2,i}\}_{i \in I_2}, \{x_{2,j}\}_{j \in B_2}, m_2^{I_2}, m_2^{B_2})$$

consists of an isomorphism $s : C_1 \rightarrow C_2$ and bijections $f^I : I_1 \rightarrow I_2$, $f^B : B_1 \rightarrow B_2$ such that

- (i) $s \circ \phi_1 = \phi_2 \circ s$,
- (ii) $s(\Sigma_1) = \Sigma_2$,
- (iii) $s(z_{1,j}) = z_{2,f^B(j)}$ for all $j \in B_1$ and $s(x_{1,i}) = x_{2,f^I(i)}$ for all $i \in I_1$,
- (iv) $m_1^{I_1} = m_2^{I_2} \circ f^I$ and $m_1^{B_1} = m_2^{B_2} \circ f^B$.

Let C be a marked orbifold Riemann surface with boundary in which every marked point has isotropy group $\mathbb{Z}/r\mathbb{Z}$. We denote

$$\omega_{C,\log} := \rho^* \omega_{|C|,\log} = \omega_C \left(\sum_{i \in I} [z_i] + \sum_{i \in I} [\bar{z}_i] + \sum_{j \in B} [x_j] \right), \quad (2.1)$$

where $\rho : C \rightarrow |C|$ is the morphism to the coarse underlying Riemann surface and $[z_i]$, $[\bar{z}_i]$, and $[x_j]$ denote the degree-1/r orbifold divisors of the marked points. An r -spin structure on C is an orbifold line bundle L together with an isomorphism

$$\tau : L^{\otimes r} \cong \omega_{C,\log} \quad (2.2)$$

and an anti-holomorphic involution $\tilde{\phi} : L \rightarrow L$ lifting ϕ such that $\tilde{\phi}^{\otimes r}$ agrees under τ with the involution on $\omega_{C,\log}$ induced by ϕ . We denote by $\text{mult}_p(L)$ the multiplicity of L at a point p , which is defined as the integer $m \in \{0, 1, \dots, r-1\}$ such that the local structure of the total space of L near p is $[\mathbb{C}^2/(\mathbb{Z}/r\mathbb{Z})]$ with the action

$$\zeta \cdot (x, v) = (\zeta x, \zeta^m v)$$

by the canonical generator ζ of the isotropy group.

Associated to an r -spin structure L on C , there is a unique twisted r -spin structure S on C , defined as the complex line bundle

$$S := L \otimes \mathcal{O} \left(- \sum_{i \mid \text{mult}_{z_i}(L)=0} r[z_i] - \sum_{i \mid \text{mult}_{\bar{z}_i}(L)=0} r[\bar{z}_i] - \sum_{j \mid \text{mult}_{x_j}(L)=0} r[x_j] \right). \quad (2.3)$$

This bundle satisfies

$$S^{\otimes r} \cong \rho^* \left(\omega_{|C|,\log} \otimes \mathcal{O} \left(- \sum_{i \mid \text{mult}_{z_i}(L)=0} r[z_i] - \sum_{i \mid \text{mult}_{\bar{z}_i}(L)=0} r[\bar{z}_i] - \sum_{j \mid \text{mult}_{x_j}(L)=0} r[x_j] \right) \right), \quad (2.4)$$

in which, on the right-hand side, $[z_i]$, $[\bar{z}_i]$, and $[x_j]$ now denote the corresponding divisors on $|C|$. It follows that the coarse underlying bundle $|S| := \rho_* S$ satisfies

$$|S|^{\otimes r} \cong \omega_{|C|} \otimes \mathcal{O} \left(- \sum_{i \in I} a_i [z_i] - \sum_{i \in I} a_i [\bar{z}_i] - \sum_{j \in B} b_j [x_j] \right) \quad (2.5)$$

with $a_i, b_j \in \{0, 1, \dots, r-1\}$ defined by the requirement that

$$a_i \equiv \text{mult}_{z_i}(L) - 1 \pmod{r}, \quad b_j \equiv \text{mult}_{x_j}(L) - 1 \pmod{r}.$$

We refer to the numbers a_i and b_j as internal and boundary *twists*. In fact, pushforward under ρ defines an equivalence of categories between bundles S satisfying (2.4) and bundles $|S|$ satisfying (2.5) for some choice of a_i and b_j ; hence, in particular, the data of $|S|$ and its twists are equivalent to the data of an r -spin structure on C . See the appendix of [10] for a more detailed discussion of this equivalence.

Let

$$J := S^\vee \otimes \omega_C, \quad (2.6)$$

which inherits an involution that we also denote by $\tilde{\phi}$. Using the fact that

$$\omega_C = \rho^* \omega_{|C|} \otimes \mathcal{O} \left(\sum_{i \in I} (r-1)[z_i] + \sum_{i \in I} (r-1)[\bar{z}_i] + \sum_{j \in B} (r-1)[x_j] \right),$$

which follows directly from (2.1), one shows easily that

$$J = \rho^* \left(|S|^\vee \otimes \omega_{|C|} \right) \otimes \mathcal{O} \left(\sum_{i \in I} \text{mult}_{z_i}(J)[z_i] + \sum_{i \in I} \text{mult}_{\bar{z}_i}(J)[\bar{z}_i] + \sum_{j \in B} \text{mult}_{x_j}(J)[x_j] \right),$$

and hence

$$|J| = |S|^\vee \otimes \omega_{|C|}. \quad (2.7)$$

In particular, since $|J|$ is a bundle on a non-orbifold curve and hence has integral degree, the following observation is immediate.

Observations 2.1. The twists a_i, b_j for a twisted r -spin structure on a smooth marked orbifold Riemann surface with boundary satisfy the following congruence condition:

$$e := \frac{2 \sum a_i + \sum b_j + (g-1)(r-2)}{r} \in \mathbb{Z}, \quad (2.8)$$

where g is the genus of C . (In case C is disconnected, we define $g = \sum_i g(C_i) - l + 1$, where C_1, \dots, C_l are the connected components of C .)

All of the above also works in the more familiar setting of closed marked orbifold Riemann surfaces $(C, \{z_i\}_{i \in I}, m^I)$. However, in this case, it is important in what follows to allow the possibility of limited -1 twists; see Observation 4.1. Thus, we define a *closed twisted r -spin structure* as a closed marked orbifold Riemann surface equipped with an orbifold line bundle S satisfying

$$S^{\otimes r} \cong \rho^* \left(\omega_{|C|, \log} \otimes \mathcal{O} \left(- \sum_{i \in I_0} r[z_i] \right) \right),$$

where $I_0 \subseteq I$ is a subset of the marked points such that $\text{mult}_{z_i}(S) = 0$ for all $i \in I_0$ and each connected component of C contains at most one marked such point z_i with $i \notin I_0$. In this case, one has

$$|S|^{\otimes r} \cong \omega_{|C|} \otimes \mathcal{O} \left(- \sum_{i \in I} a_i [z_i] \right)$$

with $a_i \in \{-1, 0, \dots, r-1\}$ and $a_i = -1$ for at most one marked point z_i in each connected component of C . Analogously to Observation 2.1, we have the following.

Observations 2.2. The twists a_i for a closed twisted r -spin structure satisfy

$$\frac{\sum a_i + (g-1)(r-2)}{r} \in \mathbb{Z}. \quad (2.9)$$

An *isomorphism* of r -spin structures consists of an isomorphism of marked orbifold Riemann surfaces with boundary, as defined above, together with an isomorphism $\tilde{s} : s^* L_2 \cong L_1$ of the spin bundles such that

- (i) \tilde{s} commutes with the involutions, that is, $\tilde{s} \circ \tilde{\phi}_1 = \tilde{\phi}_2 \circ \tilde{s}$;
- (ii) the diagram

$$\begin{array}{ccc} s^* L_2^{\otimes r} & \xrightarrow{\tau_1} & s^* \omega_{C_2, \log} \\ \tilde{s}^{\otimes r} \downarrow & & \downarrow \\ L_1^{\otimes r} & \xrightarrow{\tau_2} & \omega_{C_1, \log} \end{array}$$

commutes, where the right-hand vertical arrow is induced by s .

Similarly, an isomorphism of twisted r -spin structures is an isomorphism $\tilde{s} : s^*S_2 \cong S_1$, satisfying the analogs of (i) and (ii).

2.2 Gradings on smooth r -spin disks

From here forward, we restrict to Riemann surfaces with boundary in which each connected component has genus zero, which we refer to as *disks*. The definitions extend to higher genus, as well, but they are not needed for the current work. We denote by $(C, \phi, \Sigma, \{z_i\}_{i \in I}, \{x_j\}_{j \in B}, m^I, m^B)$ a smooth marked orbifold disk. Note that we allow the case $\partial\Sigma = \emptyset$, and that if a smooth marked orbifold disk is connected (meaning that C is connected), then $\partial\Sigma \neq \emptyset$.

Let $A \subseteq \partial\Sigma \setminus \{x_j\}_{j \in B}$ be an open subset (where we identify the boundary marked points with their images in $|C|$), and let $s \in \Gamma(A, \omega_{|C|})^\phi$ be a section fixed under the fiberwise involution on $\omega_{|C|}|_A$ induced by ϕ . We call s *positive* if, for any $p \in A$ and any tangent vector $v \in T_p(\partial\Sigma)$ in the direction of orientation, we have $\langle s(p), v \rangle > 0$. A similar notion of positivity applies to ϕ -fixed sections of any tensor power of $\omega_{|C|}$ over A .

Definition 2.3. Given a twisted r -spin structure S , a *lifting of S over A* is a continuous, $\tilde{\phi}$ -invariant section

$$v \in \Gamma\left(A, |S|^{\tilde{\phi}}\right)$$

such that the image of $v^{\otimes r}$ under the map on sections induced by the injection

$$|S|^{\otimes r} \rightarrow \omega_{|C|} \tag{2.10}$$

is positive. A *lifting of J over A* is a $\tilde{\phi}$ -invariant section

$$w \in \Gamma\left(A, |J|^{\tilde{\phi}}\right)$$

for which there exists a lifting v of S over A with $\langle w, v \rangle \in \Gamma(A, \omega_{|C|})$ positive on A , where $\langle -, - \rangle$ is induced by the natural pairing between $|S|^\vee$ and $|S|$. We consider two liftings v and v' (of either S or J) *equivalent* if $v = cv'$ for a continuous function $c : A \rightarrow \mathbb{R}^+$. We write $[v]$ for the equivalence class of v .

Observe that a twisted r -spin structure admits a lifting of S over A precisely if it admits a lifting of J over A . Moreover, there is a bijection between equivalence classes of

liftings of S and of J , in which $[v]$ corresponds to $[w]$ if $\langle w, v \rangle$ is everywhere positive for all representatives v and w of $[v]$ and $[w]$. If a twisted r -spin structure admits a lifting (of either S or J) over all of $\partial\Sigma \setminus \{x_j\}_{j \in B}$, we call it *compatible*.

We now define the notion of a lifting “alternating” at a boundary marked point.

Definition 2.4. Let w be a lifting of J over $\partial\Sigma \setminus \{x_i\}_{i \in B}$. A boundary marked point x_j is said to be *illegal* if there exists a lifting $w' \in [w]$ that can be continuously extended to x_j without vanishing. If x_j is not illegal, we say that it is *legal* and that w *alternates* at x_j . (One can define legality in the exact same manner for liftings of S , and it is straightforward to see that x_j is legal for the class $[w]$ of liftings of J precisely if it is legal for the corresponding class $[v]$ of liftings of S .)

It is immediate that legality is well defined under equivalence of v . Furthermore, legality and compatibility are closely related to the twists.

Proposition 2.5.

- (1) When r is odd, any twisted r -spin structure is compatible, and there is a unique equivalence class of liftings.
- (2) When r is even, the boundary twists b_j in a compatible twisted r -spin structure must be even. Whenever the boundary twists are even, either the r -spin structure is compatible or it becomes compatible after replacing $\tilde{\phi}$ by $\xi \circ \tilde{\phi} \circ \xi^{-1}$ for ξ an r th root of -1 , which yields an isomorphic r -spin structure.
- (3) Suppose r is odd and v is a lifting over a punctured neighborhood of a boundary marked point x_j . Then x_j is legal if and only if its twist is odd.
- (4) Suppose r is even. If a lifting over $\partial\Sigma \setminus \{x_j\}_{j \in B}$ alternates precisely at a subset $D \subseteq \{x_j\}_{j \in B}$, then

$$\frac{2 \sum a_i + \sum b_j + 2}{r} \equiv |D| \pmod{2}. \quad (2.11)$$

If (2.11) holds, then there exist exactly two liftings (up to equivalence) that alternate precisely at $D \subseteq \{x_j\}_{j \in B}$, one of which is the negative of the other.

Proof. We begin by choosing trivializations

$$\omega_{|C|}^\phi \Big|_{\partial\Sigma} \cong \partial\Sigma \times \mathbb{R} \quad (2.12)$$

and

$$\omega_{|C|} \left(- \sum_{i \in I} a_i[z_i] - \sum_{i \in I} a_i[\bar{z}_i] - \sum_{j \in B} b_j[x_j] \right)^\phi \Big|_{\partial \Sigma \setminus \{x_j\}_{j \in B}} \cong (\partial \Sigma \setminus \{x_j\}_{j \in B}) \times \mathbb{R}$$

such that a section of either of these bundles is positive precisely if its image lies in the positive ray $\mathbb{R}^+ \subseteq \mathbb{R}$ in each fiber. Let I_j be the connected component of $\partial \Sigma \setminus \{x_j\}_{j \in B}$ defined by the property that x_j is the left endpoint of the closure of I_j with respect to the orientation of $\partial \Sigma$. Since the r th tensor power of a $\tilde{\phi}$ -invariant section of $|S|$ is ϕ -invariant, one can see that on each I_j , either there is a section $v_j \in \Gamma(I_j, |S|^{\tilde{\phi}})$ with $v_j^{\otimes r}$ mapping to 1 under the composition of (2.10) and (2.12), or there is a section v_j with $v_j^{\otimes r}$ mapping to -1 .

Now, for the 1st item, suppose r is odd. Then, by replacing v_j with $-v_j$ if necessary, we find for each I_j a section $v_j \in \Gamma(I_j, |S|^{\tilde{\phi}})$ such that $v_j^{\otimes r}$ maps to 1 under the composition of (2.10) and (2.12). Thus, there is always a lifting.

If r is even, then if there is a section $v_j \in \Gamma(I_j, |S|^{\tilde{\phi}})$ with $v_j^{\otimes r}$ mapping to the constant section $\epsilon_j = \pm 1$, then $(-v_j)^{\otimes r}$ maps to $(-1)^r \epsilon_j = \epsilon_j$, and there is no real section that maps to $-\epsilon_j$. We see that when r is even, the structure is compatible precisely if $\epsilon_j = 1$ for all j .

Suppose the elements of B are enumerated cyclically so that x_j follows x_{j-1} in the cyclic order of $j \in \{1, \dots, |B|\}$ around the boundary. Choose a local coordinate x on $\partial \Sigma$ centered around a boundary marked point x_j . Then the map on local sections induced by (2.10) is multiplication by the local section x^{b_j} of $\mathcal{O}(b_j[x_j])$. The section

$$x^{-b_j} \in \Gamma \left(U \setminus \{x_j\}, \omega_{|C|} \left(- \sum_{i \in I} a_i[z_i] - \sum_{i \in I} a_i[\bar{z}_i] - \sum_{j \in B} b_j[x_j] \right)^\phi \right),$$

defined in a small punctured neighborhood $U \setminus \{x_j\}$ without any other markings, extends to all of U as a nowhere-vanishing local section that we denote by s . In addition, there exists a section $v \in \Gamma(|S|^{\tilde{\phi}}|_{U \cap I_j})$ with $v^{\otimes r}$ mapping to $\epsilon_j s$. Since s does not vanish, v can be extended to all of U without vanishing in such a way that $v^{\otimes r}$ maps to s . Given that x^{-b_j} changes sign at x_j precisely when b_j is odd, we see that

$$\epsilon_j = (-1)^{b_j} \epsilon_{j-1}.$$

Thus, if each ϵ_j equals 1, then all b_j must be even. In case all b_j are even, it could still be the case that $\epsilon_j = -1$ for each j , but then replacing $\tilde{\phi}$ by $\xi \circ \tilde{\phi} \circ \xi^{-1}$ reverses the notion of positivity and hence ensures that the spin structure is compatible. The 2nd item follows.

For the 3rd item, by the considerations of the previous two items, for r odd there is a unique r th root

$$v(x) = x^{-b_j/r}.$$

Its r th power $v^{\otimes r}$ does not change sign after crossing x_j (and hence gives rise to a lifting) precisely if b_j is even.

The last item is a consequence of the real zero count for a section of $|J|$. Using (2.7), we have

$$\deg(|J|) = \frac{2 \sum a_i + \sum b_i + 2 - 2r}{r} \equiv \frac{2 \sum a_i + \sum b_i + 2}{r} \pmod{2}.$$

Viewing the degree as the number of zeroes minus the number of poles of a meromorphic section, one sees also that

$$\deg(|J|) \equiv \deg(|J|_{|\partial\Sigma}) \pmod{2},$$

since we may choose a $\tilde{\phi}$ -invariant meromorphic section, and for such sections non-real zeroes and poles come in conjugate pairs.

The number of zeroes minus the number of poles of a $\tilde{\phi}$ -invariant meromorphic section of $|J|_{|\partial\Sigma}$ is even precisely if the real subbundle $|J|^{\tilde{\phi}}$ on $\partial\Sigma$ is orientable. The orientability of $|J|^{\tilde{\phi}} \rightarrow \partial\Sigma$, on the other hand, can be deduced from the number of legal marked points on $\partial\Sigma$: a section w , as in Definition 2.4, gives a trivialization of $|J|^{\tilde{\phi}}_{|\partial\Sigma}$ away from boundary marked points, and the transition functions between these trivializations are sign-reversing exactly at the legal marked points. Equation (2.11) follows. The same considerations also allow us, assuming (2.11) holds, to construct a lifting that alternates precisely at the points of D . The equivalence class of such a lifting is determined by choosing a lifting at any unmarked boundary points, and there are exactly two such choices. ■

The last paragraph of the above proof also yields the following statement.

Corollary 2.6. The bundle $|J|^{\tilde{\phi}}$ on $\partial\Sigma$ is orientable if and only if the number of legal marked points on $\partial\Sigma$ is even.

If S is a compatible twisted spin structure on a smooth marked orbifold disk, then we define a *grading* of S as a lifting w for J over $\partial\Sigma \setminus \{x_i\}_{i \in B}$ that alternates at every boundary marked point. We apply the same notion of equivalence as above to the grading w .

Given this, we can now define our main objects of study, in the case where C is smooth.

Definition 2.7. A *graded r -spin structure* on a smooth marked orbifold disk is a compatible twisted spin structure S in which all boundary twists are $r - 2$, together with an equivalence class $[w]$ of gradings.

Remark 2.8. The key reason for introducing the grading is that it induces a notion of positivity—that is, a choice of a preferred real ray in $J_q^{\hat{\phi}}$ at any unmarked boundary point q . Without this choice, for even r , the Witten bundle may not be orientable relative to the moduli space. Indeed, the grading cancels the automorphism of the spin bundle obtained by scaling its fibers by -1 . This automorphism lifts to the Witten bundle as scaling by -1 again. Thus, if the rank of the Witten bundle is odd, this automorphism is orientation reversing, and in particular the Witten bundle is neither orientable nor relatively orientable. However, as we will show in Theorem 5.1, the Witten bundle on the moduli space of graded r -spin structures indeed admits a canonical relative orientation.

The positivity induced by the grading will also play a crucial role in the sequel, [4], in the definition of boundary conditions for the open r -spin intersection numbers.

For that definition we also need to restrict to boundary twists of $r - 2$. In the closed r -spin cohomological field theory, twists of $r - 2$ correspond to point constraints, which are those that couple nontrivially with the unit. In the open setting the role of boundary twist $r - 2$ appears in the construction of the intersection numbers, specifically in [4, Proposition 3.20]. In that work, we use the combinatorics of Witten bundle ranks that arise from these specific twists in our construction of canonical boundary conditions for the bundles $\bigoplus \mathbb{L}_i^{\oplus d_i} \oplus \mathcal{W}$. These boundary conditions are necessary in order to ensure that the Euler class of such a bundle has a well-defined integral over a moduli space with boundary.

Remark 2.9. Graded 2-spin structures were first introduced and explored by Solomon and the 3rd author in [27], in which an equivalent definition of graded structures is also described, purely in terms of the preferred half-surface Σ considered as an orbifold Riemann surface with corners; a similar definition can be given for the graded r -spin

case as well. The moduli space of graded 2-spin surfaces is described in [28], where the authors prove that the higher-genus open intersection theory of [24] is defined. A combinatorial description of graded 2-spin structures in terms of Kasteleyn orientations of triangulations appears in [29], and it is used to prove a combinatorial formula for open intersection numbers.

A smooth marked orbifold disk together with a twisted spin structure and a grading is called a *smooth graded r -spin disk*. An *isomorphism* of graded r -spin disks consists of an isomorphism of twisted spin structures, as defined above, such that the image of the equivalence class of gradings $[w_1]$ on C_1 under the map on sections induced by s and \tilde{s} is the equivalence class $[w_2]$.

If $\text{Fix}(\phi) = \emptyset$, the above notion of grading is vacuous, but we require an additional datum.

Definition 2.10. A *smooth graded r -spin sphere* is a smooth r -spin sphere together with a choice of a distinguished marking z_i , referred to as the *anchor*, such that

- (1) $m^I(z_i) = \emptyset$ and z_i is the only marked point marked \emptyset ;
- (2) if there is a marked point with twist -1 , then it must be the anchor;
- (3) if the twist a_i of the anchor is $r-1$, we have a map $\tau' : (|S| \otimes \mathcal{O}([z_i]))^{\otimes r} \big|_{z_i} \rightarrow \mathbb{C}$ defined as the composition

$$(|S| \otimes \mathcal{O}([z_i]))^{\otimes r} \big|_{z_i} \rightarrow \omega_{|C|}([z_i]) \big|_{z_i} \cong \mathbb{C},$$

where the 2nd identification is the residue map. In this case, we also fix an involution $\tilde{\phi}$ on the fiber $(|S| \otimes \mathcal{O}([z_i]))_{z_i}$ and require it to satisfy two properties: first,

$$\tau'(\tilde{\phi}(v)^{\otimes r}) = -\overline{\tau'(v^{\otimes r})}$$

for all $v \in (|S| \otimes \mathcal{O}([z_i]))_{z_i}$, where $w \mapsto \overline{w}$ is the standard conjugation; and second,

$$\left\{ \tau'(v^{\otimes r}) \mid v \in (|S| \otimes \mathcal{O}([z_i]))_{z_i}^{\tilde{\phi}} \right\} \supseteq i\mathbb{R}_+,$$

where i is the root of -1 in the upper half plane. Finally, we pick a *positive direction* on $(|S| \otimes \mathcal{O}([z_i]))_{z_i}^{\tilde{\phi}}$, meaning a connected component V of $(|S| \otimes \mathcal{O}([z_i]))_{z_i}^{\tilde{\phi}} \setminus \{0\}$ such that $\tau'(v^{\otimes r}) \in i\mathbb{R}_+$ for any $v \in V$.

Since

$$|J|_{z_i} \otimes (|S| \otimes \mathcal{O}([z_i]))_{z_i} \cong \omega_{|C|}([z_i])|_{z_i},$$

the involution $\tilde{\phi}$ induces a conjugation on the fiber $|J|_{z_i}$, also denoted by $\tilde{\phi}$, by the requirement that

$$\langle \tilde{\phi}(w), \tilde{\phi}(v) \rangle = -\overline{\langle w, v \rangle}$$

under the identification of $\omega_{|C|}([z_i])_{z_i}$ with \mathbb{C} . Similarly, a positive direction is defined on $|J|_{z_i}^{\tilde{\phi}}$ by the requirement that $w \in |J|_{z_i}^{\tilde{\phi}}$ is positive if, for any positive $v \in (|S| \otimes \mathcal{O}([z_i]))_{z_i}^{\tilde{\phi}}$, one has

$$\langle w, v \rangle \in i\mathbb{R}_+.$$

An *isomorphism* of graded r -spin spheres is an isomorphism of r -spin spheres that preserves the anchor and, in the case where the anchor has twist $r - 1$, also preserves the involution $\tilde{\phi}$ and the positive direction.

Remark 2.11. The anchor is an auxiliary tool that is useful when sphere components arise from normalization. The anchor of a sphere component should be thought of as the half node at which the component met a disk component, or met a simple path of sphere components connecting it to a disk component, or met a contracted boundary node. This situation is relevant in what follows when the anchor has twist -1 or $r - 1$.

An anchor of twist -1 appears when there is an internal Ramond node. In this case, as we shall see below, the Witten bundle does not decompose as a direct sum. Still, by choosing a preferred half edge (the anchor) we can obtain a slightly weaker decomposition, Proposition 4.7, which will be crucial for the inductive construction of the intersection theory in the sequel. An anchor of twist $r - 1$, on the other hand, appears when a graded r -spin disk without boundary markings degenerates to a sphere, with the boundary contracting to a point. In this case, the global parity constraint induced by the grading determines the twist of the contracted boundary node to be $r - 1$, and the residue map described above is the limiting notion of grading under this degeneration.

We observe the following parity constraint, which follows from Observation 2.1 when r is odd and from item 4 of Proposition 2.5 when r is even.

Observations 2.12. For any graded spin disk, we have

$$e \equiv |B| - 1 \pmod{2}, \quad (2.13)$$

where e is defined by equation (2.8) with $g = 0$ and $b_j = r - 2$ for all j .

In both the closed (sphere) theory and the open (disk) theory, one has an existence and a uniqueness result.

Proposition 2.13. If $(C, \{z_1, \dots, z_l\})$ is a marked sphere and $a_1, \dots, a_l \in \{-1, 0, \dots, r-1\}$ are twists satisfying (2.9) such that at most one a_i equals -1 , then there exists a unique twisted r -spin structure on C with twists a_1, \dots, a_l . The only automorphisms are given by scaling S fiberwise by an r th root of unity. For any choice of z_i , there exists a unique graded r -spin structure on (C, S) with anchor z_i , and it has no automorphisms if the anchor has twist $r-1$.

Suppose $(C, \phi, \Sigma, \{z_1, \dots, z_l\}, \{x_1, \dots, x_k\})$ is a connected marked disk and $a_1, \dots, a_l \in \{0, 1, \dots, r-1\}$, $b_1, \dots, b_k \in \{0, 1, \dots, r-2\}$ are such that (2.8) holds. If r is even, let $D \subseteq [k]$ be an arbitrary set for which (2.11) holds, and if r is odd, let $D := \{i \in [k] \mid 2 \nmid b_i\}$. Then C admits a unique r -spin structure with a lifting such that the internal twists are given by the integers a_i , the boundary twists are given by the integers b_j , and D is the set of legal boundary marked points. The r -spin structure has no automorphisms that preserve the lifting.

In particular, when all b_j are equal to $r-2$ and (2.8) and (2.13) hold, there exists a unique graded r -spin structure on this disk with internal twists given by the integers a_i , and this graded structure has no automorphisms.

Proof. The 1st part is well known; see, for example, the appendix of [10]. The statement regarding the grading in the closed setting is clear, except for the automorphism claim in case the anchor has twist $r-1$. In this case, when r is odd, scaling by a root of unity is not compatible with the involution $\tilde{\phi}$ in the fiber of the anchor. When r is even, scaling by -1 is compatible with the involution but not with the choice of positive direction.

Now, fix a connected marked disk and twists a_1, \dots, a_l such that (2.8) and (2.11) hold. Note that the integrality requirement for the closed case (Observation 2.2) is

satisfied on the closed marked Riemann surface

$$(C; \{z_1, \dots, z_l, \bar{z}_1, \dots, \bar{z}_l, x_1, \dots, x_k\}),$$

so there exists a twisted r -spin structure S on C .

To define a twisted r -spin structure in the sense of surfaces with boundary, we must produce an involution

$$\tilde{\phi} : |S| \rightarrow |S|$$

lifting the involution $|\phi|$ on $|C|$. In order to do so, choose a boundary point $p \in \partial\Sigma \setminus \{x_1, \dots, x_k\}$, and choose a vector $v \in |S|_p$ such that the image of $v^{\otimes r}$ in $\omega_{|C|}|_p$, under the injection (2.10), is positive. For any other point $q \in \Sigma$ that is not a marked point, and any $u \in |S|_q$, we define $\tilde{\phi}(q, u) = (\phi(q), \tilde{\phi}(u))$ as follows. Let $\tilde{\gamma}$ be a path in the total space of

$$|S|_{|\Sigma \setminus \{z_i\}_{i \in [l]}, \{x_j\}_{j \in [k]}}$$

with $\tilde{\gamma}(0) = (p, v)$ and $\tilde{\gamma}(1) = (q, u)$, and let γ be the image of $\tilde{\gamma}^{\otimes r}$ image in the total space of the bundle $\omega_{|C|}|_{\Sigma \setminus \{z_i\}_{i \in [l]}, \{x_j\}_{j \in [k]}}$. There is a unique path in the total space of $|S|_{|\phi(\Sigma) \setminus \{\tilde{z}_i\}_{i \in [l]}, \{x_j\}_{j \in [k]}}$, denoted by $\tilde{\phi}(\tilde{\gamma})$ and starting at (p, v) , whose r th power maps to $\phi(\gamma)$. Define $\tilde{\phi}(q, u)$ to be its endpoint.

It is easy to see that the above definition is independent of choices, and the conjugation extends uniquely, up to isomorphism, to the marked points. So far, we have defined a twisted r -spin structure on the marked disk, and what remains is to define the grading. For this, one can take as the lifting any smooth section v of $|\tilde{\phi}|_{\partial\Sigma \setminus \{x_j\}_{j \in [k]}}$ that alternates at each boundary marking; such a section exists because the parity constraint (2.11) is satisfied. The lifting determines a grading w .

The absence of automorphisms, when r is odd, is due to the involution: fiberwise multiplication by an r th root of unity does not commute with $\tilde{\phi}$, so it is not an automorphism. When r is even, multiplication by -1 does commute with the involution, but it does not preserve the grading w , so it is again not an automorphism of the graded structure. ■

2.3 Stable graded r -spin disks

Thus far, we have considered only graded r -spin structures on smooth curves. In order to compactify the moduli space of such objects, however, we also need to allow the curves C to be nodal.

A *nodal marked orbifold Riemann surface with boundary* is a tuple

$$(C, \phi, \Sigma, \{z_i\}_{i \in I}, \{x_j\}_{j \in B}, m^I, m^B),$$

defined exactly as in Section 2.1 except that C is a nodal, possibly disconnected, orbifold Riemann surface (curve) as in [1], and $\text{Fix}(|\phi|)$ is locally homeomorphic near every point either to an open subset of \mathbb{R} , to the union of the coordinate axes

$$\{xy = 0\} \subseteq \mathbb{R}^2,$$

or to a single point. We recall from [1] that C is only allowed to have a nontrivial isotropy at marked points and nodes, each of which has isotropy group $\mathbb{Z}/r\mathbb{Z}$, and all nodes are required to be balanced in the sense that, in the local picture $\{xy = 0\} \subseteq \mathbb{C}^2$ at the node, the action of the distinguished generator ζ of the isotropy group is given by

$$(x, y) \mapsto (\zeta x, \zeta^{-1} y).$$

The nodes in a Riemann surface with boundary can be divided into three types:

- (1) Internal nodes, which are nodes in the interior Σ (together with a conjugate node in $\overline{\Sigma}$);
- (2) Boundary nodes, which are nodes in $\partial\Sigma$, around which $\partial\Sigma$ is locally homeomorphic to the union of the coordinate axes

$$\{xy = 0\} \subseteq \mathbb{R}^2;$$

- (3) Contracted boundary nodes, which are nodes arising in the limiting case where one component of $\text{Fix}(|\phi|)$ is a single point.

The three types of nodes are illustrated in Figure 1.

A nodal Riemann surface is *stable* if every connected component of its normalization is stable.




Fig. 1. The closed curve $|C|$, with the open disk Σ shaded.

An *r-spin structure* on a nodal marked orbifold Riemann surface with boundary is defined, exactly as in the smooth case, as a complex line bundle L on C with an isomorphism τ as in (2.2), together with an involution $\tilde{\phi} : L \rightarrow L$ lifting ϕ that is compatible with τ . There is an associated *twisted r-spin structure* S , defined by

$$S := L \otimes \mathcal{O} \left(- \sum_{i \in I_0} r[z_i] - \sum_{i \in I_0} r[\bar{z}_i] - \sum_{j \mid \text{mult}_{x_j}(L)=0} r[x_j] \right), \quad (2.14)$$

where $I_0 \subseteq I$ is a subset of the marked points such that $\text{mult}_{z_i}(L) = 0$ for all $i \in I_0$, and such that the connected components of C not meeting the ϕ -fixed locus contain at most one multiplicity-zero marked point z_i with $i \notin I_0$, whereas the connected components meeting the ϕ -fixed locus do not contain any multiplicity-zero z_i with $i \notin I_0$. The bundle S satisfies

$$S^{\otimes r} \cong \rho^* \left(\omega_{|C|, \log} \otimes \mathcal{O} \left(- \sum_{i \in I_0} r[z_i] - \sum_{i \in I_0} r[\bar{z}_i] - \sum_{j \mid \text{mult}_{x_j}(L)=0} r[x_j] \right) \right), \quad (2.15)$$

and we again define $J = S^\vee \otimes \omega_C$, as in (2.6).

Remark 2.14. Closed orbifold Riemann surfaces have additional “ghost” automorphisms in the presence of nodes, which play a role in our calculation of the automorphism groups of stable graded *r*-spin disks below. Specifically, in the local picture of a node as $\{xy = 0\} \subseteq \mathbb{C}^2$, there is one ghost automorphism of the form

$$(x, y) \mapsto (\xi x, y) \quad (2.16)$$

for each r th root of unity ξ . These act trivially on the coarse underlying curve $|C|$, but they act nontrivially on the orbifold C and induce a nontrivial action on orbifold line bundles. Indeed, let q be a node of C with branches p and p' (given locally by $y = 0$ and $x = 0$, respectively), and let m be the multiplicity of S at p (where multiplicity at a branch of a node is defined analogously to multiplicity at a marked point). Then, if g is a ghost automorphism given by (2.16), the lift $\tilde{g} : g^*S \cong S$ multiplies the fiber over p by ξ^m , or in other words, changes the gluing of the fibers of S over p and p' by a factor of ξ^m ; see [8, Proposition 2.5.3].

From here on, we restrict again to the case where each connected component has genus zero. Let $n : \widehat{C} \rightarrow C$ be the normalization morphism. Then $n^*L \rightarrow \widehat{C}$ is an r -spin structure, but n^*S is not a twisted spin structure, in general, since it may not satisfy the requisite conditions on the subset I_0 . Still, there is a canonical way to associate to S a twisted spin structure on \widehat{C} , by setting

$$\widehat{S} := n^*S \otimes \mathcal{O} \left(- \sum_{q \in \mathcal{R}} r[q] \right). \quad (2.17)$$

Here, $[q]$ is the degree- $1/r$ orbifold divisor of a point q , and \mathcal{R} is the subset of the half-nodes $q \in \widehat{C}$ (thought of as marked points of \widehat{C}) with $\text{mult}_q(n^*S) = 0$ that satisfy one of the following:

- (1) $n(q)$ is a boundary node;
- (2) $n(q)$ is a contracted boundary node;
- (3) $n(q)$ is an internal node of C that belongs to a connected component not containing any marked point of twist -1 and not meeting the ϕ -fixed locus;
- (4) $n(q)$ is an internal node of C that belongs to a connected component containing a marked point of twist -1 , and if one normalizes C only at $n(q)$, then the half node corresponding to q is in the same connected component as the marked point of twist -1 ;
- (5) $n(q)$ is an internal node of C that belongs to a connected component meeting the ϕ -fixed locus, and if one normalizes C only at $n(q)$, then the half node corresponding to q is in the connected component meeting the ϕ -fixed locus.

Thus, for each irreducible component C_l of \widehat{C} , if $\{z_i\}_{i \in I_l}$, $\{\bar{z}_i\}_{i \in I_l}$, and $\{x_j\}_{j \in B_l}$ are the marked points lying in C_l and $\{p_k\}_{k \in N_l}$ are the branches of nodes in C_l , we have the equation

$$\left(|\widehat{S}|_{|C_l|} \right)^{\otimes r} \cong \omega_{|C_l|} \otimes \mathcal{O} \left(- \sum_{i \in I_l} a_i[z_i] - \sum_{i \in I_l} a_i[\bar{z}_i] - \sum_{j \in B_l} b_j[x_j] - \sum_{h \in N_l} c_h[p_h] \right) \quad (2.18)$$

with

$$a_i, c_h \in \{-1, 0, \dots, r-1\}, \quad b_j \in \{0, \dots, r-1\}.$$

The numbers a_i , b_j , and c_h are the *twists* of the corresponding internal marked points, boundary marked points, and half nodes.

Note that if p and p' are the two branches of a node, then we have

$$c_p + c_{p'} \equiv r-2 \pmod{r}.$$

If $c_p \equiv -1 \pmod{r}$ (and hence $c_{p'} \equiv -1 \pmod{r}$, as well), we say that the node is *Ramond*. Otherwise, we have $c_p + c_{p'} = r-2$, and we say that the node is *Neveu–Schwarz*.

As long as C does not have a contracted boundary node, the notion of *lifting* can be defined as before, as a continuous, $\tilde{\phi}$ -invariant section of $|S|$ on (an open subset of) the complement of the special points in $\partial\Sigma$ whose r th power is positive, and we say that a twisted r -spin structure is *compatible* if it admits a lifting over the entire complement A of the special points in $\partial\Sigma$. Two liftings v and v' are *equivalent* if there is a continuous function $c : A \rightarrow \mathbb{R}^+$ such that $v = cv'$. We similarly define liftings and equivalence for J , and the same correspondence between equivalence classes of liftings for S and J holds. If w is a lifting of J over A , then it induces a lifting of

$$\widehat{J} := \omega_{\widehat{C}} \otimes \widehat{S}^\vee$$

over the complement of the special points in the boundary $\widehat{\partial\Sigma}$ of the normalization. We say that w *alternates* at a marked point or half-node q , and that q is *legal*, if the induced lifting on \widehat{J} alternates at q . Otherwise, the lifting does not alternate, and the point is said to be *illegal*. A lifting w of J over A that alternates at all boundary marked points is a *grading* if, in addition, one of the two half nodes of every Neveu–Schwarz boundary node is legal and the other is illegal.

Observations 2.15. In a compatible r -spin structure for even r , all boundary half nodes have twists of even parity, by item 2 of Proposition 2.5. In particular, in this case, there are no Ramond boundary nodes, since the twist of such a node is odd. When r is odd, Ramond boundary nodes may exist, but both of their half nodes are necessary illegal. Indeed, since the twist of a Ramond boundary halfnode is $r-1$ (by the definition of \widehat{S} and the twists in (2.17) and (2.18)), which is even, the illegality of Ramond boundary half nodes follows from item 3 of Proposition 2.5.

If C has a contracted boundary node, on the other hand, then in any connected component with such a node, the complement of the special points in $\partial\Sigma$ is empty, so we must adapt the definition of a lifting. Given that each connected component of C has genus zero, there can only be one contracted boundary node in each connected component. Restrict to one such component, and let q be the contracted boundary node. Assume, additionally, that q is Ramond. Recall that the fiber $\omega_{|C|}|_q$ is canonically identified with \mathbb{C} via the residue, and the involution ϕ is sent, under this identification, to the involution $z \mapsto -\bar{z}$, whose fixed points are the purely imaginary numbers (The residue of a conjugation-invariant form ζ can be calculated as $\frac{1}{2\pi i} \oint_L \zeta$, where $L \subseteq \Sigma$ is a small loop surrounding q whose orientation is such that q is to the left of L . Applying conjugation and using the invariance of ζ shows that the residue is imaginary.). We define a *lifting* as a $\tilde{\phi}$ -invariant element

$$v \in \Gamma(\{q\}, |S|) = |S| |_q$$

such that the image of $v^{\otimes r}$ under the map

$$|S|^{\otimes r}|_q \rightarrow \omega_{|C|}|_q$$

is positive imaginary, meaning that it lies in $i\mathbb{R}_+$. In this case, we call the twisted r -spin structure *compatible* if the contracted boundary node is Ramond and a lifting exists. Two liftings are equivalent if, at the contracted boundary node, they differ by multiplication by a positive number. There always exists a $\tilde{\phi}$ -invariant $w \in |J| |_q$ such that $\langle v, w \rangle$ is positive imaginary, and we refer to this w as a *grading*; this is the limiting case of the notion of grading for smooth curves.

With or without a contracted boundary node, we now have the following definition:

Definition 2.16. A *stable genus-zero graded r -spin surface* is a nodal marked orbifold Riemann surface with boundary whose coarse underlying surface $(|C|, \{z_i\}, \{\bar{z}_i\}, \{x_j\})$ is a stable Riemann surface in which each connected component has genus zero, together with

- (1) a compatible twisted r -spin structure S in which all boundary marked points have twist $r - 2$ and all contracted boundary nodes are Ramond;
- (2) an equivalence class of gradings;

(3) a choice of one distinguished special point (called the *anchor* and marked \emptyset) in each connected component C' of C that is either disjoint from the set $\text{Fix}(\phi)$ or meets the set $\text{Fix}(\phi)$ in a single contracted boundary node. If either a contracted boundary node or marked point of twist -1 exists, we require the anchor to be this point; if not, the anchor is simply required to be a marked point. We also require that the collection of anchors is ϕ -invariant, so that it descends to Σ . Finally, if the twist of an anchor z_i is $r - 1$, we fix an involution $\tilde{\phi}$ on the fiber $(|S| \otimes \mathcal{O}([z_i]))_{z_i}$ and an orientation of the $\tilde{\phi}$ -fixed subspace, as in Definition 2.10.

The internal and boundary marked points are required to satisfy the same properties as in the smooth case, and in particular, an anchor, if it is a marked point, is the only marked point in its connected component that is marked \emptyset .

We conclude this subsection with an existence and uniqueness result analogous to Proposition 2.13. Here, we refer to irreducible components of C that do not meet the preimage of $\text{Fix}(|\phi|)$, or that meet the preimage of $\text{Fix}(|\phi|)$ in a single contracted boundary node, as *sphere components*, and we refer to the other irreducible components as *disk components*.

Proposition 2.17. Suppose that $(C, \phi, \Sigma, \{z_1, \dots, z_l\}, \{x_1, \dots, x_k\})$ is a connected stable marked disk and $a_1, \dots, a_l \in \{0, 1, \dots, r-1\}$ are such that (2.8) and (2.13) hold with $|B| = k$ and $b_j = r-2$ for all j . Then there exists a unique graded r -spin structure on this stable disk with internal twists given by the integers a_i .

The order of the automorphism group of the graded r -spin structure is r^n where $2n$ is the number of internal nodes of C .

Proof. The proof of the existence and uniqueness of the graded structure closely mirrors the proof of Proposition 2.13. In particular, there exists a unique twisted r -spin structure on the closed genus-zero surface C with the given twists at all internal, conjugate, and boundary marked points, so what remains is to produce the involution $\tilde{\phi} : S \rightarrow S$ and the sections v and w .

First, suppose that there is no contracted boundary node. Then the involution $\tilde{\phi}$ can be defined component by component. Namely, starting from some boundary point p , we first construct the involution on a single component using the argument of Proposition 2.13. At each node, there is an identification of the fibers of S on the two half nodes, so when we encounter a boundary half node, the involution on one side

induces the involution on the other side, and we use the other half node as the basepoint for the construction of the involution in its component. An analogous treatment works for conjugate internal nodes. The resulting involution is unique up to an isomorphism.

The lifting v and the grading w can be defined componentwise, uniquely up to isomorphism (as we prove below), using either the parity of the twist (in the case where r is odd) or the parity constraint (2.11) (in the case where r is even) to determine whether it alternates at each halfnode. Ramond boundary nodes appear only when r is odd, and then there is no choice in the lifting. For Neveu–Schwarz boundary nodes, there are two choices of lifting when r is even, but they are equivalent via the isomorphism scaling the fibers of S by -1 .

To prove that there is a unique choice of grading satisfying the requisite condition at a Neveu–Schwarz boundary node q , let p and p' be the two branches with respective twists c_p and $c_{p'}$. When r is odd, the fact that w alternates at exactly one branch is immediate from item 3 of Proposition 2.5, since exactly one of c_p and $c_{p'}$ is odd. When r is even, assume for simplicity that the nodal Riemann surface C consists of two sphere components joined at the node q . Then, adding the parity constraints (2.11) on the two components, one obtains

$$\frac{4 + 2 \sum a_i + \sum b_i + c_p + c_{p'}}{r} \equiv k + \delta_p^{\text{alt}} + \delta_{p'}^{\text{alt}} \pmod{2},$$

where k is the number of boundary marked points, and δ_p^{alt} is defined to be 1 if w alternates at p and 0 otherwise. Combining this with the parity constraint (2.11) on the entire curve, one finds

$$\frac{2 + c_p + c_{p'}}{r} \equiv \delta_p^{\text{alt}} + \delta_{p'}^{\text{alt}} \pmod{2},$$

so we indeed see that if $c_p + c_{p'} = r - 2$, then w alternates at exactly one branch.

This completes the proof of the existence of the graded r -spin structure in the case where there is no contracted boundary node. When there is a contracted boundary node, it must be Ramond. To see this, apply constraint (2.8) to see that the internal twists satisfy

$$-2 - 2 \sum a_i \equiv 0 \pmod{r}. \quad (2.19)$$

On a single component of C , on the other hand, the degree of the restriction of S is

$$\frac{-2 - \sum a_i - c_p}{r},$$

so we must have

$$-2 - \sum a_i - c_p \equiv 0 \pmod{r}. \quad (2.20)$$

Combining (2.19) and (2.20), we find that $2(c_p + 1) \equiv 0 \pmod{r}$. When r is odd, this is sufficient to conclude that $c_p \equiv r - 1 \pmod{r}$. When r is even, we apply (2.13) to find

$$\frac{2 + 2 \sum a_i}{r} \equiv 0 \pmod{2}.$$

Therefore, we have

$$\frac{1 + \sum a_i}{r} \in \mathbb{Z},$$

so $1 + \sum a_i \equiv 0 \pmod{r}$. Combining this with (2.20) again shows that $c_p \equiv r - 1 \pmod{r}$.

When there is a contracted boundary node q , the involution $\tilde{\phi}$ can be defined first in the fiber of q , by choosing a lifting of the conjugation on $\omega_{\log}|_q$. There are r such choices; when r is odd, they are all equivalent, while when r is even, there are $r/2$ equivalent choices that make the structure compatible, but in either case, we choose one. We then extend the conjugation to the other components of C , if there are any, component by component, using the argument in the beginning of this proof. Since the contracted boundary node is Ramond, we can now choose the grading as above. We observe that, again, there is no choice when r is odd and there are two equivalent choices when r is even.

Finally, we compute the order of the automorphism group. The closed case is known (see, e.g., [18, Proposition 1.18] or [11, Section 2.3]): an r -spin structure on a closed, genus-zero stable curve with N nodes has r^{N+1} automorphisms. Namely, each node contributes r ghost automorphisms of the curve, which can each be lifted to the spin bundle, as in Remark 2.14, and each of the resulting automorphisms of the spin structure can be composed with a global fiberwise scaling by an r th root of unity.

Let us now consider which of these automorphisms respect the graded structure. By compatibility with the involution, a ghost automorphism at an internal node determines the ghost automorphism at its conjugate node but is otherwise unconstrained.

Nontrivial ghost automorphisms at boundary and contracted boundary nodes, on the other hand, cannot respect both the conjugation and the graded structure. Indeed, at a boundary node, (2.16) may constitute an automorphism only if ξ is real. For odd r , this implies that $\xi = 1$. When r is even, $\xi = -1$ is also a possibility; however, in this case, by Remark 2.14, this ghost automorphism acts on the spin bundle by changing the gluing of the fibers at the node by a factor of ξ^m , in which m is the multiplicity at one half node. By Observation 2.15, the twists at the half nodes are even, so m is odd. It follows that $\xi^m = (-1)^m = -1$, and hence the ghost automorphism does not respect the grading. At a contracted boundary node, we consider the automorphisms in local coordinates, in which they have the form

$$(x, y) \mapsto (\xi x, \xi y),$$

where ξ and ζ are r th roots of unity. Such an automorphism is compatible with the conjugation, which locally has the form $(x, y) \mapsto (\bar{y}, \bar{x})$, only if $\zeta = \bar{\xi} = \xi^{-1}$. But in this case, given the orbifold structure at the node, it is the identity automorphism.

It follows, then, that there are r^n contributing ghost automorphisms, where $2n$ is the number of internal nodes. Since we have already argued in Proposition 2.13 that the fiberwise scalings on disk components do not respect the graded structure, there are no further automorphisms. ■

3 Moduli Space of Graded r -Spin Disks

Henceforth, we usually denote an r -spin disk with a lifting simply by Σ , the preferred half, suppressing most of the notation.

3.1 Stable graded r -spin graphs

It is useful to encode some of the combinatorial data of graded r -spin disks in a decorated dual graph.

Definition 3.1. A *genus-zero pre-stable dual graph* is a tuple

$$\Gamma = (V, H, \sigma_0, \sim, H^{CB}, m),$$

in which

- (i) V is a finite set (the *vertices*) equipped with a decomposition $V = V^O \sqcup V^C$ into *open* and *closed vertices*;

- (ii) H is a finite set (the *half-edges*) equipped with a decomposition $H = H^B \cup H^I$ into *boundary* and *internal half edges*;
- (iii) $\sigma_0 : H \rightarrow V$ is a function, viewed as associating to each half edge the vertex from which it emanates;
- (iv) \sim is an equivalence relation on H , which decomposes as a pair of equivalence relations \sim_B on H^B and \sim_I on H^I . The equivalence classes are required to be of size 1 or 2, and those of size 1 are referred to as *tails*. We denote by $T^B \subseteq H^B$ and $T^I \subseteq H^I$ the sets of equivalence classes of size 1 in H^B and H^I , respectively;
- (v) H^{CB} is a subset of T^I , the *contracted boundary tails*;
- (vi) m is a function given by

$$m = m^B \sqcup m^I : T^B \sqcup (T^I \setminus H^{CB}) \rightarrow 2^{\mathbb{N}},$$

where m^B and m^I (the *boundary* and *internal markings*) satisfy the definition of a marking when restricted to any connected component of Γ , and where m^I is strict on connected components with an open vertex or a contracted boundary tail.

Note that (V, H, σ_0) defines a graph and that we do not require this graph to be connected; denote its set of connected components by $\text{Conn}(\Gamma) = \{\Lambda_i\}$. We require the above data to satisfy the following conditions:

- (1) For each boundary half-edge $h \in H^B$, we have $\sigma_0(h) \in V^O$;
- (2) For each Λ_i , we have $h^1(\Lambda_i) = 0$;
- (3) Each Λ_i contains at most one half edge in H^{CB} , and if Λ_i contains such a half edge, then all vertices of Λ_i are closed;
- (4) For each Λ_i , the sub-graph formed by its open vertices (if any exist) and their incident boundary edges is connected.

Conditions (2), (3), and (4) guarantee that for any nodal graded r -spin surface with dual graph Γ , each connected component of the closed surface C has genus zero.

We refer to elements of T^B as *boundary tails* and to elements of $T^I \setminus H^{CB}$ as *internal tails*, and we denote $T := T^I \sqcup T^B$. Note that \sim induces a fixed-point-free involution on $H \setminus T$, which we denote by σ_1 . Write

$$E^B := (H^B \setminus T^B) / \sim_B, \quad E^I := (H^I \setminus T^I) / \sim_I$$

and refer to these as *boundary edges* and *internal edges*. The set of *edges* is $E := E^B \sqcup E^I$. Denote by σ_0^B the restriction of σ_0 to H^B , and similarly for σ_0^I .

For each vertex v , set $k(v) := |(\sigma_0^B)^{-1}(v)|$ and $l(v) := |(\sigma_0^I)^{-1}(v)|$. We say that an open vertex $v \in V^O$ is *stable* if $k(v) + 2l(v) > 2$, and we say that a closed vertex $v \in V^C$ is *stable* if $l(v) > 2$. A graph is *stable* if all of its vertices are stable, and it is *closed* if $V^O = \emptyset$. A graph is *smooth* if there are no edges or contracted boundary tails.

Definition 3.2. An *isomorphism* between two genus-zero pre-stable dual graphs

$$\Gamma = (V, H, \sigma_0, \sim, H^{CB}, m) \text{ and } \Gamma' = (V', H', \sigma'_0, \sim', H'^{CB}, m')$$

is a pair $f = (f^V, f^H)$, where $f^V : V \rightarrow V'$ and $f^H : H \rightarrow H'$ are bijections satisfying

- (1) $h_1 \sim h_2$ if and only if $f(h_1) \sim' f(h_2)$,
- (2) $f^V \circ \sigma_0 = \sigma'_0 \circ f^H$,
- (3) $m = m' \circ f^H$,
- (4) $f(H^{CB}) = H'^{CB}$.

We denote by $\text{Aut}(\Gamma)$ the group of automorphisms of Γ .

Pre-stable dual graphs encode the discrete data of a marked orbifold Riemann surface with boundary. In order to encode the additional data of a twisted spin structure and a lifting, we must add further decorations.

Definition 3.3. A *genus-zero twisted r -spin dual graph with a lifting* is a genus-zero pre-stable dual graph Γ as above, together with maps

$$\text{tw} : H \rightarrow \{-1, 0, 1, \dots, r-1\}$$

(the *twist*) and

$$\text{alt} : H^B \rightarrow \mathbb{Z}/2\mathbb{Z}$$

and a subset $T^* \subseteq T^I$ (the *anchors*), satisfying the following conditions:

- (i) Any connected component of Γ that is not stable consists either of (a) a single open vertex with a single internal tail, or (b) a single closed vertex with exactly two tails, one of which is in H^{CB} and the other of which is in H^I .

- (ii) Every closed connected component contains exactly one tail in T^* . All contracted boundary tails and all tails t with $\text{tw}(t) = -1$ belong to T^* . Open connected components have no tails in T^* . Any element of $T^* \setminus H^{CB}$ is marked by \emptyset and is the only tail marked by \emptyset in its connected component.
- (iii) For any vertex v , the total number of incident half-edges h with $h \in T^*$ or $\text{tw}(h) = -1$ is at most one.
- (iv) For any contracted boundary tail $t \in H^{CB}$, we have $\text{tw}(t) = r - 1$.
- (v) For any open vertex $v \in V^O$,

$$2 \sum_{h \in (\sigma_0^I)^{-1}(v)} \text{tw}(h) + \sum_{h \in (\sigma_0^B)^{-1}(v)} \text{tw}(h) \equiv r - 2 \pmod{r}$$

and

$$\frac{2 \sum_{h \in (\sigma_0^I)^{-1}(v)} \text{tw}(h) + \sum_{h \in (\sigma_0^B)^{-1}(v)} \text{tw}(h) + 2}{r} \equiv \sum_{h \in (\sigma_0^B)^{-1}(v)} \text{alt}(h) \pmod{2}.$$

- (vi) For any closed vertex $v \in V^C$,

$$\sum_{h \in \sigma_0^{-1}(v)} \text{tw}(h) \equiv r - 2 \pmod{r}.$$

- (vii) For any half-edge $h \in H \setminus T$, we have

$$\text{tw}(h) + \text{tw}(\sigma_1(h)) \equiv r - 2 \pmod{r},$$

and at most one of $\text{tw}(h)$ and $\text{tw}(\sigma_1(h))$ equals -1 . No boundary half-edge h satisfies $\text{tw}(h) = -1$. In case $h \in H^I \setminus T^I$ satisfies $\text{tw}(h) \equiv -1 \pmod{r}$, then $\text{tw}(h) = r - 1$ precisely if, after detaching the edge, h belongs to the connected component containing an anchor $t^* \in T^*$ (if h is in a closed connected component of Γ) or an open vertex $v \in V^O$ (if h is in an open connected component of Γ).

- (viii) For any boundary half-edge $h \in H^B \setminus T^B$, if $\text{tw}(h) \neq r - 1$ we have

$$\text{alt}(h) + \text{alt}(\sigma_1(h)) = 1,$$

and if $\text{tw}(h) = r - 1$ then $\text{alt}(h) = \text{alt}(\sigma_1(h)) = 0$.

(ix) If r is odd, then for any $h \in H^B$,

$$\text{alt}(h) \equiv \text{tw}(h) \pmod{2},$$

and if r is even, then for any $h \in H^B$,

$$\text{tw}(h) \equiv 0 \pmod{2}.$$

Boundary half-edges h with $\text{alt}(h) = 1$ are called *legal*, and those with $\text{alt}(h) = 0$ are called *illegal*. Half-edges h with $\text{tw}(h) \in \{-1, r-1\}$ are called *Ramond*, and those with $\text{tw}(h) \in \{0, \dots, r-2\}$ are called *Neveu–Schwarz*. An edge is called *Ramond* if one (hence both) of its half edges is Ramond, and Neveu–Schwarz otherwise.

We say that a genus-zero twisted r -spin dual graph with a lifting is *stable* if the underlying dual graph is stable, in the sense specified above. An *isomorphism* between genus-zero twisted r -spin dual graphs with liftings consists of an isomorphism in the sense of Definition 3.3 that respects tw , alt , and T^* . Analogously to Definition 2.16, we define a *genus-zero graded r -spin graph* to be a genus-zero twisted r -spin dual graph with a lifting such that every boundary tail $h \in H^B$ has

$$\text{tw}(h) = r-2, \quad \text{alt}(h) = 1.$$

Any stable graded r -spin disk Σ induces a stable genus-zero graded r -spin graph $\Gamma(\Sigma)$. Namely, the correspondence associates an irreducible component of Σ to a vertex, a marked point to a tail, a node to an edge, and a contracted boundary node to a contracted boundary tail. The twist is represented by tw and the alternation by alt , and the anchors correspond to elements of T^* .

3.2 Moduli of stable graded r -spin disks

In the situation without boundary, there is a well-studied moduli space $\overline{\mathcal{M}}_{g,n}^{1/r}$ of stable Riemann surfaces with r -spin structure, which is known to be a smooth Deligne–Mumford stack with projective coarse moduli, for which the forgetful map to $\overline{\mathcal{M}}_{g,n}$ is finite (see [8] or, in the setting of a slightly different compactification, [16]). This moduli space admits a decomposition into open and closed substacks,

$$\overline{\mathcal{M}}_{g,n}^{1/r} = \bigsqcup_{\vec{a}=(a_1, \dots, a_n)} \overline{\mathcal{M}}_{g, \vec{a}}^{1/r}, \quad (3.1)$$

where $a_i \in \{0, 1, \dots, r-1\}$ for each i and $\overline{\mathcal{M}}_{g, \vec{a}}^{1/r}$ denotes the substack of r -spin structures with twist a_i at the i th marked point. In genus zero, the situation is even simpler: according to Proposition 2.13, for any choice of \vec{a} such that (2.9) holds, the moduli space $\overline{\mathcal{M}}_{0, \vec{a}}^{1/r}$ has coarse moduli isomorphic to $\overline{\mathcal{M}}_{0, n}$ and generic additional isotropy $\mathbb{Z}/r\mathbb{Z}$. The isomorphism of coarse moduli is given by the smooth map For_{spin} that forgets the spin structure.

When there is no boundary, it is also straightforward to add the information of a grading. Indeed, the moduli space of graded r -spin spheres for which the anchor does not have twist $r-1$ is canonically isomorphic to $\overline{\mathcal{M}}_{0, n}^{1/r}$, whereas the moduli space of graded r -spin spheres with twist \vec{a} and anchor twisted $r-1$ is canonically an r -to-1 cover of the moduli space $\overline{\mathcal{M}}_{0, \vec{a}'}^{1/r}$ on which \vec{a}' agrees with \vec{a} except that the anchor has twist -1 .

To generalize the construction of the moduli space to the open setting, we first note that in [24] the moduli space $\overline{\mathcal{M}}_{0, k, l}$ of connected stable marked disks with boundary marked points marked by $\{1, \dots, k\}$ and internal marked points marked by $\{1, \dots, l\}$ was considered. It is a smooth orientable manifold with corners in the sense of [19], and its dimension is

$$\dim_{\mathbb{R}}(\overline{\mathcal{M}}_{0, k, l}) = k + 2l - 3.$$

Let $\overline{\mathcal{M}}_{0, k, l}^{1/r}$ denote the set of isomorphism classes of connected stable graded r -spin disks, with boundary and internal marked points as above. There is a set-theoretic decomposition analogous to (3.1),

$$\overline{\mathcal{M}}_{0, k, l}^{1/r} = \bigsqcup_{\vec{a}} \overline{\mathcal{M}}_{0, k, \vec{a}}^{1/r},$$

in which $\overline{\mathcal{M}}_{0, k, \vec{a}}^{1/r} \subseteq \overline{\mathcal{M}}_{0, k, l}^{1/r}$ consists of those disks for which the i th internal marked point has twist a_i . By Proposition 2.17, whenever $\overline{\mathcal{M}}_{0, k, \vec{a}}^{1/r} \neq \emptyset$, there is a bijection given by the forgetful map

$$\text{For}_{\text{spin}} : \overline{\mathcal{M}}_{0, k, \vec{a}}^{1/r} \rightarrow \overline{\mathcal{M}}_{0, k, l},$$

and we use these to give $\overline{\mathcal{M}}_{0, k, l}^{1/r}$ the structure of a manifold with corners.

This describes the coarse underlying space of $\overline{\mathcal{M}}_{0,k,l}^{1/r}$. The main theorem of this section is that it can also be given an orbifold-with-corners structure in the sense of [22, Section 3].

Theorem 3.4. The moduli space $\overline{\mathcal{M}}_{0,k,l}^{1/r}$ of connected stable graded r -spin disks with boundary marked points marked by $\{1, \dots, k\}$ and internal marked points marked by $\{1, \dots, l\}$ is a compact smooth orientable orbifold with corners of real dimension $k+2l-3$. Its universal bundle admits a universal grading.

We split the proof of Theorem 3.4 into three parts. Lemma 3.6 shows that the moduli space is a compact smooth orbifold with corners, Lemma 3.8 proves the existence of the universal grading, and Proposition 3.13 together with Observation 3.10 proves the orientability.

Remark 3.5. In fact, for all $r \geq 2$, the moduli space $\overline{\mathcal{M}}_{g,k,l}^{1/r}$ of connected stable graded genus g marked r -spin surfaces with boundary is a compact smooth orbifold with corners of real dimension $k+2l+3g-3$, and its universal bundle admits a universal grading. For $r=2$, this is proven in [28], and the moduli space is moreover shown in [29] to be canonically oriented. The general r case will be proven elsewhere; in this case, the moduli space may not be orientable.

Lemma 3.6. $\overline{\mathcal{M}}_{0,k,l}^{1/r}$ has the structure of a compact smooth orbifold with corners.

Proof. We describe a procedure that defines an orbifold-with-corners structure on $\overline{\mathcal{M}}_{0,k,l}^{1/r}$, analogous to the procedure performed in [22, Section 2]. To define the procedure, we make reference to the following sequence:

$$\overline{\mathcal{M}}_{0,k,l}^{1/r} \xrightarrow{(5)} \widehat{\mathcal{M}}_{0,k,l}^{1/r} \xrightarrow{(4)} \widetilde{\mathcal{M}}_{0,k,l}^{1/r} \xrightarrow{(3)} \overline{\mathcal{M}}_{0,k,l}^{1/r, \mathbb{Z}_2} \xrightarrow{(2)} \overline{\mathcal{M}}_{0,k+2l}^{1/r, \mathbb{Z}_2} \xrightarrow{(1)} \overline{\mathcal{M}}_{0,k+2l}^{'1/r}. \quad (3.2)$$

The moduli spaces and maps appearing in (3.2) are defined as they appear in what follows.

Step 1: First, $\overline{\mathcal{M}}_{0,k+2l}^{'1/r}$ is the suborbifold of $\overline{\mathcal{M}}_{0,k+2l}^{1/r}$ given by the condition that the 1st k markings have twist $r-2$ and that the integer defined in (2.9) is of the same parity as $k+1$. Inside this space, $\overline{\mathcal{M}}_{0,k+2l}^{1/r, \mathbb{Z}_2}$ is the fixed locus of the involution defined by

$$(C; w_1, \dots, w_{k+2l}, S) \mapsto (\overline{C}; w_1, \dots, w_k, w_{k+l+1}, \dots, w_{k+2l}, w_{k+1}, \dots, w_{k+l}, \overline{S}),$$

where \bar{C} and \bar{S} are the same as C and S but with the conjugate complex structure (more details on the fixed point functor on stacks can be found in [26]). As the fixed locus of an anti-holomorphic involution, $\overline{\mathcal{M}}_{0,k+2l}^{1/r, \mathbb{Z}_2}$ has the structure of a real orbifold. It parameterizes isomorphism types of marked spin spheres with a real structure (an involution $\tilde{\phi}$ covering the conjugation ϕ on C) and the prescribed twists, and it maps to $\overline{\mathcal{M}}_{0,k+2l}^{1/r}$ (in general, it is not a sub-orbifold, since some isotropy is lost), so it inherits a universal curve via pullback.

Remark 3.7. Let us digress to discuss the isotropy of $\overline{\mathcal{M}}_{0,k+2l}^{1/r, \mathbb{Z}_2}$, especially near nodal strata (Without spin structure, the nodal strata of the real moduli space $\overline{\mathcal{M}}_{0,k,l}$ are discussed in [20, Section 3], and in the closed case with spin structure, the nodal strata of $\overline{\mathcal{M}}_{0,l}^{1/r}$ are discussed in [9, Section 4]). The generic point of $\overline{\mathcal{M}}_{0,k+2l}^{1/r, \mathbb{Z}_2}$ has isotropy coming from scaling the fibers of the spin bundle by real r th roots of unity. When r is odd, there are no such roots and hence no generic isotropy, while when r is even, there is generic $\mathbb{Z}/2\mathbb{Z}$ isotropy. Nodal strata have additional $\mathbb{Z}/r\mathbb{Z}$ isotropy for each internal Neveu–Schwarz node, coming from the ghost automorphisms.

For boundary nodes, there is a difference in behavior for r odd or even. When r is odd, boundary nodes also contribute no further isotropy. Furthermore, if $U \times [-1, 1]$ is a neighborhood in the moduli space of a curve with a single boundary node, such that $U \times \{0\}$ is the intersection with the nodal stratum and (u, t) for $t \neq 0$ corresponds to a smooth real sphere, then the passage from $t < 0$ to $t > 0$ geometrically corresponds to flipping one of the two disk components and defining the involution on the spin bundle in the unique possible way. Thus, in this case, the behavior near the node is exactly like in the real, non-spin case, and the spin moduli continues to be a trivial degree-one cover of the non-spin moduli generically. When r is even, on the other hand, boundary nodes contribute additional $\mathbb{Z}/2\mathbb{Z}$ isotropy, coming from ghost automorphisms of the form $(x, y) \rightarrow (-x, y)$; see the discussion in the proof of Proposition 2.17. On the moduli level, the picture is that a neighborhood of a nodal curve with a single boundary node looks locally like $U \times [-1, 1]/(\mathbb{Z}/2\mathbb{Z})$, where $U \times \{0\}$ is the nodal locus and the generator of $\mathbb{Z}/2\mathbb{Z}$ takes (u, t) to $(u, -t)$.

Contracted boundary nodes add no additional isotropy, as we saw in the end of the proof of Proposition 2.17. On the moduli level, again let $U \times [-1, 1]$ be a neighborhood in the moduli space of a curve with a contracted boundary, such that $U \times \{0\}$ is the intersection with the nodal stratum and (u, t) for $t > 0$ corresponds to a smooth real sphere on which the conjugation has nonempty fixed locus. Then (u, t) for $t < 0$

corresponds to a real sphere on which the conjugation has no fixed points, so after taking the quotient by the conjugation, the result is a marked real projective plane. ■

Step 2: Returning to our discussion of (3.2), the next step is to cut $\overline{\mathcal{M}}_{0,k+2l}^{1/r, \mathbb{Z}_2}$ along the real simple normal crossings divisor consisting of curves with at least one real node, via the “real hyperplane blow-up”, as in [22], yielding an orbifold with corners $\widetilde{\mathcal{M}}_{0,k,l}^{1/r, \mathbb{Z}_2}$. We direct the reader to [22, Section 3.3] for more details, but the idea is the following.

Near the real divisor consisting of curves with a contracted boundary node—or, when r is odd, near the real divisor consisting of curves with a boundary node (a real node that is not an isolated fixed point of the conjugation)—the real blow-up is the standard cutting procedure that can be defined without a spin structure. In the notation of Remark 3.7, the real blow-up corresponds to the natural quotient map

$$U \times [-1, 0] \sqcup U \times [0, 1] \rightarrow U \times [-1, 0] \cup U \times [0, 1] = U \times [-1, 1].$$

When r is even, the blow-up near the real divisor consisting of curves with a boundary node is a topologically trivial operation, but nontrivial on the orbifold level. In particular, in local charts, the real blow-up, when r is even, is equivalent to blowing up before taking the extra $\mathbb{Z}/2\mathbb{Z}$ quotient mentioned in Remark 3.7, and then taking the quotient, so it kills the additional $\mathbb{Z}/2\mathbb{Z}$ isotropy on nodal strata. In other words, the blow-up is locally the map

$$(U \times [-1, 0] \sqcup U \times [0, 1]) / (\mathbb{Z}/2\mathbb{Z}) \rightarrow (U \times [-1, 0] \cup U \times [0, 1]) / (\mathbb{Z}/2\mathbb{Z}) = U \times [-1, 1] / (\mathbb{Z}/2\mathbb{Z}),$$

where the generator of $\mathbb{Z}/2\mathbb{Z}$ takes $(u, t) \in U \times [-1, 0] \sqcup U \times [0, 1]$ to $(u, -t)$.

Step 3: Consider the subset of $\widetilde{\mathcal{M}}_{0,k,l}^{1/r, \mathbb{Z}_2}$ whose generic point is a smooth marked real spin sphere with nonempty real locus. Then $\widetilde{\mathcal{M}}_{0,k,l}^{1/r}$ is the disconnected 2-to-1 cover of this subset given by the choice of a distinguished connected disk component of $C \setminus C^\phi$. Equivalently, in the generic (smooth) situation, we first restrict to the connected components of $\widetilde{\mathcal{M}}_{0,k,l}^{1/r, \mathbb{Z}_2}$ consisting of real spheres on which the conjugation has nonempty fixed locus, and then we choose an orientation for C^ϕ . It is important to note, however, that this choice can be uniquely continuously extended to points in the boundary of $\widetilde{\mathcal{M}}_{0,k,l}^{1/r}$.

Step 4: Inside $\widetilde{\mathcal{M}}_{0,k,l}^{1/r}$, we denote by $\widehat{\mathcal{M}}_{0,k,l}^{1/r}$ the union of connected components such that the marked points w_{k+1}, \dots, w_{k+l} lie in the distinguished stable disk and, for even r , the spin structure is compatible.

Step 5: Finally, $\overline{\mathcal{M}}_{0,k,l}^{1/r}$ is the cover of $\widehat{\mathcal{M}}_{0,k,l}^{1/r}$ given by a choice of grading. When r is odd, this is the identity, while when r is even, it is a 2-to-1 cover given by forgetting the global $\mathbb{Z}/2\mathbb{Z}$ isotropy.

As a topological space, $\overline{\mathcal{M}}_{0,k,l}^{1/r}$ is indeed the same space defined previously, but now it has the additional structure of an orbifold with corners. The proof that $\widehat{\mathcal{M}}_{0,k,l}^{1/r}$ is an orbifold with corners is identical to the proof of the analogous Theorem 2 in [22]. The space $\overline{\mathcal{M}}_{0,k,l}^{1/r}$ then inherits the orbifold-with-corners structure from $\widehat{\mathcal{M}}_{0,k,l}^{1/r}$. It is moreover compact since compactness is preserved at every step.

Over $\overline{\mathcal{M}}_{0,k,l}^{1/r}$, there is a universal curve whose fibers are compatible stable spin disks. The content of Lemma 3.8 below is that one may construct a graded structure, in a continuous way, on the fibers.

Lemma 3.8. One can continuously choose a grading for the fibers of the universal curve of $\overline{\mathcal{M}}_{0,k,l}^{1/r}$. This choice is unique when r is odd, while for even r it is unique up to a global change of grading in each connected component of the moduli space.

Proof. We first prove the lemma for the universal curve over

$$\text{Int}(\mathcal{M}_{0,k,l}^{1/r}) := \overline{\mathcal{M}}_{0,k,\{a_1, \dots, a_l\}}^{\frac{1}{r}} \setminus \partial \overline{\mathcal{M}}_{0,k,\{a_1, \dots, a_l\}}^{\frac{1}{r}}.$$

The statement is clear when r is odd, and its uniqueness up to a possible global change of grading in each component is immediate when r is even, given the existence. We hence prove the existence for even r .

The fact that this choice can be made locally in a continuous way is straightforward. The obstruction to making such a choice global in $\text{Int}(\mathcal{M}_{0,k,l}^{1/r})$ is the possible existence of a loop

$$\gamma : \partial \bar{\Delta} \rightarrow \text{Int}(\overline{\mathcal{M}}_{0,k,\{a_1, \dots, a_l\}}^{\frac{1}{r}}),$$

where $\bar{\Delta}$ is the closed unit disk, along which a continuous choice of grading alternates.

Since the strata corresponding to disks with an internal “bubble” are of codimension two, it may be assumed that $\gamma(\partial \bar{\Delta}) \subseteq \mathcal{M}_{0,k,\vec{a}}^{1/r}$. We first show that γ may be extended to $\hat{\gamma} : \bar{\Delta} \rightarrow \overline{\mathcal{M}}_{0,k,\vec{a}}^{1/r}$ by verifying that $\pi_1(\overline{\mathcal{M}}_{0,k,\vec{a}})$ is trivial. This is true when $l = 0$, as every connected component of $\overline{\mathcal{M}}_{0,k,0}$ is the contractible associahedron. By considering the forgetful map $\overline{\mathcal{M}}_{0,k,1} \rightarrow \overline{\mathcal{M}}_{0,k,0}$, whose fiber is contractible, it is easy to see that $\overline{\mathcal{M}}_{0,k,1}$ is also contractible. For $l > 1$, it is enough to consider an arbitrary

$\gamma : \partial \bar{\Delta} \rightarrow \mathcal{M}_{0,k,l}$ and to show that it can be extended to the disk. By working in the unit disk model where z_1 is mapped to the origin and x_1 to 1, one may write $\gamma(\theta)$ as

$$(1, x_2(\theta), \dots, x_k(\theta), 0, z_2(\theta), \dots, z_l(\theta)).$$

An extension $\hat{\gamma}$ may be written as

$$\begin{aligned} \hat{\gamma}(r, \theta) = (1, e^{r \log x_2(\theta) + (1-r) \frac{2\pi i}{k}}, e^{r \log x_3(\theta) + (1-r) \frac{4\pi i}{k}}, \dots, \\ e^{r \log x_k(\theta) + (1-r) \frac{2(k-1)\pi i}{k}}, 0, r^{l-1} z_2(\theta), r^{l-2} z_3(\theta), \dots, r z_l(\theta)), \end{aligned}$$

where we define the logarithm by excluding the positive real ray.

As $\bar{\Delta}$ is contractible, standard homotopy arguments now show that we can uniquely extend the grading from an arbitrary grading at $\gamma(1)$ to a grading for all points of $\hat{\gamma}(\bar{\Delta})$. Restricting to $\partial \bar{\Delta}$, the grading defines a grading for the points of γ . Thus, the grading does not alternate along γ , and therefore it can be defined globally.

Given a grading for the fibers of the universal curve over the interior of the moduli space, we extend it to fibers over the boundary by continuity. If Σ_t is a family of smooth graded r -spin surfaces converging to Σ_0 , then the gradings of Σ_t determine, by continuity, a compatible lifting on Σ_0 away from special points. An argument as in Proposition 2.17 shows that this lifting is in fact a grading, and it is independent of the family Σ_t .

Suppose $\Sigma \in \partial \bar{\mathcal{M}}_{0,k,l}^{1/r}$ has a contracted boundary node, which in particular forces that $k = 0$. Using the same argument as in Proposition 2.17 for the contracted boundary case, we observe that the contracted boundary node must be Ramond. The limit of the grading in the smooth case, at the boundary stratum consisting of surfaces with a contracted boundary node, is precisely a grading in the sense of contracted boundary nodes defined above. ■

Remark 3.9. It is interesting to note that even for nodal spin disks with Neveu-Schwarz nodes, the choice of grading cannot be performed independently for different components, if it is required to be continuous. It is the real blow-up stage in the construction of the orbifold with corners that fixes this choice, up to a global change of grading in each connected component of the moduli space.

The above results can be carried out in greater generality. First, if the images of the markings are any sets B and I , one can clearly define the space $\bar{\mathcal{M}}_{0,B,I}^{1/r}$ in the same way

as above. Furthermore, associated to each connected stable genus-zero twisted graded r -spin dual graph Γ , there is a closed suborbifold with corners $\overline{\mathcal{M}}_{\Gamma}^{1/r} \subseteq \overline{\mathcal{M}}_{0,B,I}^{1/r}$ whose general point is a graded r -spin disk with dual graph Γ . We also allow for the possibility that Γ is disconnected, in which case $\overline{\mathcal{M}}_{\Gamma}^{1/r}$ is defined as the product of the moduli spaces $\overline{\mathcal{M}}_{\Gamma_i}^{1/r}$ associated to its connected components. Inside $\overline{\mathcal{M}}_{\Gamma}^{1/r}$, we define $\mathcal{M}_{\Gamma}^{1/r}$ as the open suborbifold consisting of graded r -spin disks whose dual graph is precisely Γ .

There are forgetful maps between the moduli spaces, but we note that marked points can only be forgotten if their twist is zero (otherwise (2.5) is not preserved), and boundary marked points can only be forgotten if they are in addition illegal (otherwise the grading does not descend to the moduli space with fewer marked points). We define

$$\text{For}_{B',I'} : \overline{\mathcal{M}}_{\Gamma}^{1/r} \rightarrow \overline{\mathcal{M}}_{\Gamma'}^{1/r} \quad (3.3)$$

for $B', I' \subseteq \mathbb{Z}$ by forgetting all twist-zero internal marked points marked by I' and all twist-zero illegal boundary marked points marked by B' . This process may create unstable components; we repeatedly contract them. If the process ends with some unstable components, we remove them. We denote by $\text{for}_{B',I'}(\Gamma)$ the graph Γ' resulting from this procedure.

3.3 Orientation of $\overline{\mathcal{M}}_{0,k,\{a_1 \dots, a_l\}}^{\frac{1}{r}}$

In the following section, we describe a natural orientation on the spaces $\overline{\mathcal{M}}_{0,k,\{a_1 \dots, a_l\}}^{\frac{1}{r}}$, thereby completing the proof of Theorem 3.4. The ideas presented here are not new; in particular, they are similar to those presented in [24, Section 2.5] and are closely related to the earlier discussion in [15, Section 2.1.2]. First, we reduce the question of orientability to a simpler setting.

Observations 3.10. We claim that the moduli space $\overline{\mathcal{M}}_{0,k,\{a_1 \dots, a_l\}}^{\frac{1}{r}}$ is orientable exactly if $\mathcal{M}_{0,k,l}$ is orientable and that an orientation on $\overline{\mathcal{M}}_{0,k,l}$ induces one on $\overline{\mathcal{M}}_{0,k,\{a_1 \dots, a_l\}}^{\frac{1}{r}}$ by pullback under the map

$$\text{For}_{\text{spin}} : \overline{\mathcal{M}}_{0,k,\{a_1 \dots, a_l\}}^{\frac{1}{r}} \rightarrow \overline{\mathcal{M}}_{0,k,l}$$

that forgets the graded spin structure.

To see this, note that the map $\text{For}_{\text{spin}} : \mathcal{M}_{0,k,\{a_1 \dots, a_l\}}^{\frac{1}{r}} \rightarrow \mathcal{M}_{0,k,l}$ on the open moduli spaces is a diffeomorphism on the coarse underlying level, which means that the open

locus $\mathcal{M}_{0,k,\{a_1,\dots,a_l\}}^{\frac{1}{r}}$ is indeed orientable precisely if $\mathcal{M}_{0,k,l}$ is orientable. To pass from the open locus to the full moduli space, one can construct $\overline{\mathcal{M}}_{0,k,\{a_1,\dots,a_l\}}^{\frac{1}{r}}$ (respectively, $\overline{\mathcal{M}}_{0,k,l}$) from $\mathcal{M}_{0,k,\{a_1,\dots,a_l\}}^{\frac{1}{r}}$ (respectively, $\mathcal{M}_{0,k,l}$) in two stages. First, add loci parameterizing disks without boundary nodes or contracted boundary components; such loci are of real codimension two and hence do not affect orientability. Then, add the boundary of the moduli space; this contains strata of real codimension one, but the fact that they lie in the boundary means that they do not affect orientability. This proves the claim.

We henceforth discuss orientations on $\mathcal{M}_{0,k,l}$, but, in light of Observation 3.10, all statements carry over to the spin case. Furthermore, orientations can be studied on a connected component at a time, and the connected components of both $\overline{\mathcal{M}}_{0,B,\vec{a}}^{1/r}$ and $\overline{\mathcal{M}}_{0,B,\vec{a}}$ are indexed by cyclic orders of B .

Notation 3.11. If $\pi : [|B|] \rightarrow B$ is an order of B , we denote its induced cyclic order by $\hat{\pi}$, and we denote by $\overline{\mathcal{M}}_{0,B,\vec{a}}^{1/r, \hat{\pi}}$ (respectively, $\overline{\mathcal{M}}_{0,B,\vec{a}}^{\hat{\pi}}$) the connected component of $\overline{\mathcal{M}}_{0,B,\vec{a}}^{1/r}$ (respectively, $\overline{\mathcal{M}}_{0,B,\vec{a}}$) that parameterizes disks for which the cyclic order of boundary markings taken along the boundary of the disk, with its canonical orientation, is $\hat{\pi}$. We denote by $\mathcal{M}_{0,k,l}^{\text{main}}, \overline{\mathcal{M}}_{0,k,l}^{\text{main}} \subseteq \overline{\mathcal{M}}_{0,k,l}$ the subspaces where the induced cyclic order on the boundary marked points is the cyclic order induced from the standard order π^{std} on $[k]$.

We denote by $\text{Ord}(B)$ the set of all orders of B , by $\text{Cyc}(B)$ the set of cyclic orders, and by S_B the group of permutations of B . Note that S_B acts both on $\text{Ord}(B)$, by composition, and on $\overline{\mathcal{M}}_{0,B,\vec{a}}^{1/r}$, by permuting markings.

Definition 3.12. Let $\{\tilde{o}^\pi = \tilde{o}_{0,B,I}^\pi\}$ be a family of orientations, where B runs over all sets of size k , I runs over all sets of size l , π runs over all orders of B , and $\tilde{o}_{0,B,I}^\pi$ is an orientation for $\overline{\mathcal{M}}_{0,B,I}^{\hat{\pi}}$. We say such a family is *covariant* if, whenever $f^B : B \rightarrow B'$ and $f^I : I \rightarrow I'$ are bijections and $F : \overline{\mathcal{M}}_{0,B,I} \rightarrow \overline{\mathcal{M}}_{0,B',I'}$ is the induced map, we have $\tilde{o}_{0,B,I}^\pi = F^* \tilde{o}_{0,B',I'}^{\pi \circ f^B}$. A family $\{\tilde{o}^\pi = \tilde{o}_{0,B,\{\vec{a}\}}^\pi\}$ of orientations of $\overline{\mathcal{M}}_{0,B,\{\vec{a}_i\}_{i \in I}}^{1/r, \hat{\pi}}$ is covariant if it is the pullback of a covariant family of orientations for $\overline{\mathcal{M}}_{0,B,I}^{\hat{\pi}}$.

The fiber of the forgetful map $\text{For}_{k+1} : \mathcal{M}_{0,k,l+1} \rightarrow \mathcal{M}_{0,k,l}$ is a punctured disk with a canonical complex orientation. For $k \geq 1$, the fiber of the forgetful map $\mathcal{M}_{0,k+1,l} \rightarrow \mathcal{M}_{0,k,l}$ is a union of open intervals, so it is canonically oriented as the boundary of an oriented disk (as above). Denote this orientation by $o_{\text{For}_{k+1}^{-1}(\Sigma)}$.

Proposition 3.13. Suppose $k + 2l \geq 3$. Then there exists a unique covariant family of orientations $\tilde{o}_{0,B,I}^\pi$ for the spaces $\overline{\mathcal{M}}_{0,B,I}^{\hat{\pi}}$ with the following properties:

- (1) In the zero-dimensional case where $k = l = 1$, the orientation is positive, while when $k = 3$ and $l = 0$, the orientations are negative.
- (2) Fix an integer h , and let $\pi \in \text{Ord}(B)$ and $g \in S_B$ be such that g sends $x_{\pi(i)}$ to $x_{\pi(i+h)}$ cyclically. Then g preserves the orientation of $\overline{\mathcal{M}}_{0,B,I}^{\hat{\pi}}$ if and only if $h(|B| - 1)$ is even (this holds for any orientation of the moduli space of disks).
- (3) The orientation $\tilde{o}_{0,k,l+1}^\pi$ agrees with the orientation induced from $\tilde{o}_{0,k,l}^\pi$ by the fibration $\overline{\mathcal{M}}_{0,k,l+1} \rightarrow \overline{\mathcal{M}}_{0,k,l}$ and the complex orientation on the fiber.
- (4) On $\overline{\mathcal{M}}_{0,k+1,l}^{\text{main}}$, we have $\tilde{o}_{0,k+1,l}^{\pi^{\text{std}}} = o_{\text{For}_{k+1}^{-1}(\Sigma)} \otimes \text{For}_{k+1}^* \tilde{o}_{0,k,l}^{\pi^{\text{std}}}$.

Remark 3.14. For k odd, the orientations described here differ from those of [24] by $(-1)^{\frac{k-1}{2}}$. This choice is more natural from the point of view of integrable hierarchies.

Proof of Proposition 3.13. If orientations $\tilde{o}_{0,k,l}^\pi$ exist, then properties (1)–(4) imply that they are unique. It remains to check existence.

For property (2) to hold, we must show that permutations of labels that map the component $\overline{\mathcal{M}}_{0,k,l}^{\text{main}}$ to itself affect the orientation according to their sign. This can be checked with respect to any orientation. Write

$$U := \left\{ (z, w) \mid \begin{array}{ll} z = (z_1, \dots, z_k) \in (S^1)^k, & z_i \neq z_j, \quad i \neq j \\ w = (w_1, \dots, w_l) \in (\text{int}D^2)^l, & w_i \neq w_j, \quad i \neq j \end{array} \right\}.$$

Denote by $U^{\text{main}} \subseteq U$ the subset where the cyclic order of z_1, \dots, z_k on $S^1 = \partial D^2$ (with respect to the orientation induced from the complex orientation of D^2) agrees with the standard order of $[k]$. Then

$$\mathcal{M}_{0,k,l}^{\text{main}} = U^{\text{main}} / \text{PSL}_2(\mathbb{R}).$$

When k is odd, cyclic permutations of the boundary labels preserve the orientation of U^{main} and thus also $\mathcal{M}_{0,k,l}^{\text{main}}$ and $\overline{\mathcal{M}}_{0,k,l}^{\text{main}}$. When k is even, a cyclic permutation of boundary labels that moves each boundary label by h multiplies the orientation by the sign $(-1)^h$. Renaming internal markings is a complex map that preserves orientations trivially, and similarly, arbitrary permutations of the interior labels preserve the orientation of $\overline{\mathcal{M}}_{0,k,l}^{\text{main}}$.

A direct calculation shows that the orientation on $\overline{\mathcal{M}}_{0,3,1}$ induced by property (3) from $\tilde{\mathfrak{o}}_{0,3,0}^\pi$ agree with the orientation induced by property (4) from $\tilde{\mathfrak{o}}_{0,1,1}^\pi$. Thus, the required $\tilde{\mathfrak{o}}_{0,3,1}^\pi$ exists. Existence of $\tilde{\mathfrak{o}}_{0,k,l}^\pi$ satisfying properties (3) and (4) for other k and l follows from the commutativity of the diagram of forgetful maps

$$\begin{array}{ccc} \overline{\mathcal{M}}_{0,k+1,l+1}^{\text{main}} & \longrightarrow & \overline{\mathcal{M}}_{0,k+1,l}^{\text{main}} \\ \downarrow & & \downarrow \\ \overline{\mathcal{M}}_{0,k,l+1}^{\text{main}} & \longrightarrow & \overline{\mathcal{M}}_{0,k,l}^{\text{main}}. \end{array}$$

Covariance, at this point, gives a unique way to extend the orientations to other connected components and to moduli spaces for different B, I . \blacksquare

Notation 3.15. Denote by $\tilde{\mathfrak{o}}_{0,B,\{a_i\}_{i \in I}}^\pi$ the orientation on $\overline{\mathcal{M}}_{0,B,\{a_i\}_{i \in I}}^{1/r, \hat{\pi}}$ defined as $\text{For}_{\text{spin}}^* \tilde{\mathfrak{o}}_{0,B,I}^\pi$.

For the next lemma we need the following facts. Let Γ be an r -spin dual graph with a lifting that corresponds to a stratum of $\overline{\mathcal{M}}_{0,B,\{a_i\}_{i \in I}}$. Assume furthermore it consists of two vertices v_1 and v_2 connected by a single edge e , with boundary labels B and internal labels I . We have

$$i_\Gamma^* \det(T\overline{\mathcal{M}}_{0,B,\{a_i\}_{i \in I}}^{1/r}) \cong \det(N) \otimes \det(T\overline{\mathcal{M}}_{\text{detach}_e(\Gamma)}^{1/r}), \quad (3.4)$$

where N is the normal bundle of $\overline{\mathcal{M}}_\Gamma^{1/r}$ relative to $\overline{\mathcal{M}}_{0,B,\{a_i\}_{i \in I}}^{1/r}$; this can be identified with the tensor product of the tangent line bundles at the half nodes on the moduli spaces $\overline{\mathcal{M}}_{v_1}^{1/r}$ and $\overline{\mathcal{M}}_{v_2}^{1/r}$ associated to the vertices (To be more precise, the isomorphism (3.4) holds only on the coarse level, as the actions of the isotropy groups of the moduli space on the fibers are not the same; see Remarks 4.5 and 4.8. However, these actions are clearly orientation preserving, so this issue does not affect our orientation analysis, and will therefore be ignored below.). These two tangent lines are canonically oriented complex lines when e is internal and are canonically oriented real lines when e is boundary. In both cases, N carries a canonical orientation, in the 1st case it is the complex orientation. In the 2nd case $\overline{\mathcal{M}}_\Gamma^{1/r}$ is a real codimension 1 boundary stratum, hence N is a real one-dimensional bundle. Nonzero elements in the fibers correspond to either inward-pointing vectors, vectors that point to the interior of $\overline{\mathcal{M}}_{0,B,\{a_i\}_{i \in I}}^{1/r}$, or outward-pointing ones. The canonical orientation of N is the one that makes the outward directions positive.

Lemma 3.16. The orientations $\tilde{o}_{0,B,\{a_i\}_{i \in I}}^\pi$ satisfy the following two properties:

(1) Write $I = I_1 \sqcup I_2$, and take $B = B_1 \sqcup B_2$. Let Γ be the graph with two open vertices, v_1 and v_2 , connected by an edge e , where the vertex v_i has internal tails labeled by I_i and k_i boundary tails labeled by B_i , and let h_i denote the half edges of v_i . Let $\hat{\pi}$ be a cyclic ordering of B in which all tails of v_1 are consecutive and all tails of v_2 are consecutive. Denote by π the unique ordering of B such that, for any $\Sigma \in \mathcal{M}_\Gamma^{\hat{\pi}}$ with normalization $\Sigma_1 \sqcup \Sigma_2$ (where Σ_i corresponds to v_i), we have

- under π , the marked points of Σ_1 appear before those of Σ_2 ;
- when π is restricted to the points of Σ_i , it agrees with the order of the points on $\partial \Sigma_i$ with its natural orientation, starting after the node.

Let π_1 be the restriction of π to the points of Σ_1 , but adding the half-node x_{h_1} in the end, and let π_2 be the restriction to points of Σ_2 , but adding the half-node x_{h_2} in the beginning. By (3.4) we have

$$\det(T\overline{\mathcal{M}}_{0,B,\{a_i\}_{i \in I}}^{\hat{\pi}})|_{\mathcal{M}_\Gamma^{\hat{\pi}}} = \det(N) \otimes \det(T\mathcal{M}_\Gamma^{\hat{\pi}}),$$

where N is the normal bundle with the canonical orientation o_N given by the outward-pointing normal. Then

$$\tilde{o}^\pi|_{\mathcal{M}_\Gamma^{\hat{\pi}}} = (-1)^{(|B_1|-1)|B_2|} o_N \otimes (\tilde{o}_{0,B_1 \cup \{h_1\}, I_1}^{\pi_1} \boxtimes \tilde{o}_{0,\{h_2\} \cup B_2, I_2}^{\pi_2}).$$

(2) Let Γ be a graph with two vertices, an open vertex v^o and a closed vertex v^c . By (3.4) we have

$$\det(T\mathcal{M}_\Gamma) = \det(N) \boxtimes \det(T\mathcal{M}_{v^c}) \boxtimes \det(T\mathcal{M}_{v^o}),$$

where N is again the normal bundle. Then, for any order π ,

$$\tilde{o}^\pi|_{\mathcal{M}_\Gamma} = o_N \otimes (\tilde{o}_{v^o}^\pi \boxtimes \tilde{o}_{v^c}), \quad (3.5)$$

where o_N and \tilde{o}_{v^c} are the canonical complex orientations.

Proof. We prove the 1st item (which is analogous to Lemma 3.16 in [24]) by induction on the dimension $\overline{\mathcal{M}}_{0,B,I}^{1/r}$. By Observation 3.10, it is enough to prove the analogous claim for $\overline{\mathcal{M}}_{0,k,l}$. The proof of the 2nd item uses exactly the same arguments, so it is omitted.

Covariance shows that it is enough to prove the claim when $B = [k]$ and π is the order $1, \dots, k$. The base cases where the moduli space has dimension one or two can be checked by hand. Suppose, then, that the desired statement holds for all moduli spaces of dimension less than n , where $n \geq 3$. After possibly interchanging the roles of v_1 and v_2 , we can assume that a tail of v_2 can be forgotten without affecting stability; here, we use item (2) of Proposition 3.13 and the fact that $\dim_{\mathbb{R}} \mathcal{M}_{v_i} \equiv |B_i| \pmod{2}$ to see that the interchanging affects the equation with the correct sign. Let Γ' be the stable graph obtained by removing a tail of v_2 . If this tail is a boundary tail, assume it is labeled k . Write v'_2 for the resulting vertex of Γ' and π' for the restriction of π to $[k-1]$.

Consider first the case where the forgotten tail corresponds to a boundary point x_k . With the notation of Proposition 3.13, we have $\tilde{o}_{0,k,l}^{\pi} = o_{\text{For}_k^{-1}(\Sigma)} \otimes \text{For}_k^* \tilde{o}_{0,k,l}^{\pi'}$, so

$$\tilde{o}^{\pi}|_{\mathcal{M}_{\Gamma}} = o_{\text{For}_k^{-1}(\Sigma)} \otimes \text{For}_k^* \tilde{o}^{\pi'}|_{\mathcal{M}_{\Gamma'}}. \quad (3.6)$$

By induction, we have

$$\tilde{o}^{\pi'}|_{\mathcal{M}_{\Gamma'}} = (-1)^{(|B_1|-1)(|B_2|-1)} o_N \otimes (\tilde{o}_{v_1}^{\pi_1} \boxtimes \tilde{o}_{v'_2}^{\pi_2}). \quad (3.7)$$

We can identify $\text{For}_k^{-1}(N|_{\mathcal{M}_{\Gamma'}}) \cong N_{\mathcal{M}_{\Gamma'}}$, and the identification preserves natural orientations. Finally,

$$\tilde{o}_{v'_2}^{\pi_2} = o_{\text{For}_k^{-1}(\Sigma)} \otimes \text{For}_k^* \tilde{o}_{v'_2}^{\pi'}. \quad (3.8)$$

Putting equations (3.6), (3.7), and (3.8) together, and recalling that $\dim_{\mathbb{R}} \mathcal{M}_{v_1} \equiv |B_1| \pmod{2}$, we obtain the result.

If the forgotten tail corresponds to an internal marked point labeled i , then

$$\tilde{o}_{0,k,l}^{\pi} = o_{\text{For}_i^{-1}(\Sigma)} \otimes \text{For}_i^* \tilde{o}_{0,k,l-1}^{\pi} \Rightarrow \tilde{o}^{\pi}|_{\mathcal{M}_{\Gamma}} = o_{\text{For}_i^{-1}(\Sigma)} \otimes \text{For}_i^* \tilde{o}^{\pi}|_{\mathcal{M}_{\Gamma'}}, \quad (3.9)$$

where we abuse notation somewhat by using For_i to denote the map forgetting the i th internal marked point. By induction, we have

$$\tilde{o}^{\pi'}|_{\mathcal{M}_{\Gamma'}} = (-1)^{(|B_1|-1)|B_2|} o_N \otimes \tilde{o}_{v_1}^{\pi_1} \boxtimes \tilde{o}_{v'_2}^{\pi_2}. \quad (3.10)$$

Observe that $\text{For}_i^{-1}(N|_{\mathcal{M}_{\Gamma'}}) \cong N_{\mathcal{M}_{\Gamma'}}$, and the equation preserves natural orientations. Finally,

$$\tilde{\mathfrak{o}}_{V_2}^{\pi_2} = o_{\text{For}_i^{-1}(\Sigma)} \otimes \text{For}_i^* \tilde{\mathfrak{o}}_{V_2}^{\pi'_2}. \quad (3.11)$$

Putting equations (3.9), (3.10), and (3.11) together, and noting that $\dim_{\mathbb{R}} \text{For}_i^{-1}(\Sigma) = 2$, we obtain the result. \blacksquare

4 Associated Vector Bundles

4.1 Witten bundle

In closed genus-zero r -spin theory, the virtual fundamental class is defined using the *Witten bundle* $(R^1\pi_* \mathcal{S})^\vee$, where $\pi : \mathcal{C} \rightarrow \overline{\mathcal{M}}_{0,n}^{1/r}$ is the universal family and $\mathcal{S} \rightarrow \mathcal{C}$ is the universal twisted r -spin bundle. This is an orbifold vector bundle with the fibers

$$H^1(\mathcal{C}, \mathcal{S})^\vee \cong H^0(\mathcal{C}, J),$$

of complex rank $\frac{\sum a_i - (r-2)}{r}$.

Observations 4.1. As observed in [17] and studied in detail in [3], the Witten bundle is a bundle as long as at most one marked point has twist -1 and all other twists are non-negative, since this ensures that $R^0\pi_* \mathcal{S} = 0$.

We now define an open analog of the Witten bundle. Denote by $\pi : \mathcal{C} \rightarrow \overline{\mathcal{M}}_{0,k+2l}^{1/r, \mathbb{Z}_2}$ the universal curve over the moduli space of real spin spheres defined above, and by $\mathcal{S} \rightarrow \mathcal{C}$ the universal spin bundle. Then $R^1\pi_* \mathcal{S}$ is a vector bundle, since (2.5) implies that spin structures have negative degree and hence $R^0\pi_* \mathcal{S} = 0$. There are universal involutions

$$\phi : \mathcal{C} \rightarrow \mathcal{C} \text{ and } \tilde{\phi} : \mathcal{S} \rightarrow \mathcal{S},$$

which induce an involution on $R^1\pi_* \mathcal{S}$. Let

$$\mathcal{W}^{\text{pre}} := (R^0\pi_* \mathcal{J})_+ = (R^1\pi_* \mathcal{S})_-^\vee$$

be the vector bundle of $\tilde{\phi}$ -invariant sections of J , where $\mathcal{J} := \mathcal{S}^\vee \otimes \omega_\pi$; the 2nd equality uses Serre duality, under which invariant sections become anti-invariant. From here,

the *open Witten bundle*, which is a real vector bundle, is defined as

$$\mathcal{W} := \varpi^* \mathcal{W}^{\text{pre}},$$

where $\varpi : \overline{\mathcal{M}}_{0,k,l}^{1/r} \rightarrow \overline{\mathcal{M}}_{0,k+2l}^{1/r, \mathbb{Z}_2}$ is the composition of the morphisms defined above.

Remark 4.2. We pull back \mathcal{W} from the moduli space $\overline{\mathcal{M}}_{0,k+2l}^{1/r, \mathbb{Z}_2}$ of spheres in order to avoid the need to define derived pushforward in the orbifold-with-corners context. To avoid cluttering the notation in what follows, however, we often write $\mathcal{W} = (R^0\pi_* \mathcal{J})_+$ even on $\overline{\mathcal{M}}_{0,k,l}^{1/r}$. Whenever we write such expressions, they should be understood as pulled back under ϖ . In addition, we sometimes write $\mathcal{W} = (R^0\pi_* \mathcal{J})_+$ on $\overline{\mathcal{M}}_{0,n}^{1/r}$, where no involution is needed; in this case, the subscript should be ignored and the equation is to be interpreted as $\mathcal{W} = R^0\pi_* \mathcal{J}$.

Remark 4.3. Due to the canonical isomorphism

$$H^0(J) = H^0(|J|), \quad (4.1)$$

the fibers of the Witten bundle can equivalently be viewed as sections of $|J|$ on $|C|$. Furthermore, if C is a graded r -spin disk and $p \in C$ is a non-orbifold point, then the fiber J_p is identified with the fiber $|J|_{\rho(p)}$ over the image point $\rho(p) \in |C|$. In particular, if $s \in H^0(J)$ is an element of the fiber of the Witten bundle over C corresponding under (4.1) to $\rho_*(s) \in H^0(|J|)$, then the evaluation of s at p agrees under the identification $J_p = |J|_{\rho(p)}$ with the evaluation of $\rho_*(s)$ at $\rho(p)$. Because of these observations, we view the fibers of the Witten bundle interchangeably as $H^0(J)$ or as $H^0(|J|)$ in what follows.

The real rank of \mathcal{W} is

$$\frac{2 \sum a_i + \sum b_j - (r - 2)}{r}, \quad (4.2)$$

which is the number e defined in (2.8) after setting $g = 0$. Indeed, a standard Riemann–Roch calculation shows that (4.2) is the complex rank of $R^0\pi_* \mathcal{J}$, and taking involution-invariant parts reduces the real rank by half. Furthermore, in the notation of (3.3), there is the canonical isomorphism

$$\mathcal{W} = \text{For}_{B', I'}^* \mathcal{W} \quad (4.3)$$

for any subsets $B', I' \subseteq \mathbb{Z}$. This identification involves replacing a graded r -spin disk C by a partially coarsened (and possibly stabilized) disk C' . However, in the same vein as Remark 4.3, if $s \in H^0(J)$ is an element of the fiber of \mathcal{W} over C that corresponds under (4.3) to an element $\text{For}_{B', I'}^*(s)$ in the fiber of $\text{For}_{B', I'}^* \mathcal{W}$ over C' , and if $p \in C$ is a non-special point whose component is not stabilized when passing to C' , then the evaluation of s at p coincides under (4.3) with the evaluation of $\text{For}_{B', I'}^*(s)$ at the image of p .

4.2 Decomposition properties of the Witten bundle

The open Witten bundle, like its closed analog, satisfies a decomposition property along nodes. In order to state the property, we must define a normalization morphism on the moduli spaces, which can be described by a “detaching” operation on graphs.

Definition 4.4. Let Γ be a genus-zero graded r -spin graph, and let e be an edge of Γ with half-edges h and h' . Then the *detaching* of Γ at e is the disconnected graph

$$\text{detach}_e(\Gamma) = (V', H', \sigma'_0, \sim', H'^{CB}, m'),$$

defined to agree with Γ except that $h \not\sim' h'$. We keep alt and tw the same, and we extend the marking and the anchor as follows. If e is a boundary edge, set $m'(h) = m'(h') = \emptyset$. If e is an internal edge, then exactly one of the components of $\text{detach}_e(\Gamma)$ is closed and unanchored; suppose, without loss of generality, that this is the component containing h . Then we set h to be the anchor of its component, and we set $m'(h) = \emptyset$ and $m'(h')$ to be the union of the markings of the internal tails $h'' \neq h$ in the same component as h .

If $t \in H^{CB}$ is a contracted boundary tail, then the *detaching* of Γ at t is the graph $\text{detach}_t(\Gamma)$ defined to agree with Γ except that $t \in (T')^I \setminus (H')^{CB}$. We keep alt and tw the same and leave t as the anchor.

Note that the new internal and boundary markings still satisfy the requirements of Definitions 3.1 and 3.3. In particular, since there is a canonical identification of $E(\Gamma) \setminus \{e\}$ with the edges of $\text{detach}_e(\Gamma)$ and of $E(\Gamma)$ with the edges of $\text{detach}_t(\Gamma)$, one can also iterate the detaching process. For any subset $N \subseteq E(\Gamma) \cup H^{CB}(\Gamma)$, we denote by $\text{detach}_N(\Gamma)$ the graph obtained by performing detach_f for each element $f \in N$; the result is independent of the order in which the detachings are performed. When we write $\text{detach}(\Gamma)$ without any subscript, we mean $\text{detach}_{E(\Gamma)}(\Gamma)$.

Let Γ be a stable r -spin dual graph, and let $\widehat{\Gamma} = \text{detach}_N(\Gamma)$ for some set $N \subseteq E(\Gamma) \cup H^{CB}(\Gamma)$ of edges and contracted boundary tails. Unlike the moduli space of curves,

the r -spin moduli space does not always have a gluing map $\overline{\mathcal{M}}_{\widehat{\Gamma}}^{1/r} \rightarrow \overline{\mathcal{M}}_{0,k,l}^{1/r}$, because there is no canonical way to glue the fibers of the spin bundle at the internal nodes. Instead, we consider the following diagram of morphisms:

$$\overline{\mathcal{M}}_{\widehat{\Gamma}}^{1/r} \xleftarrow{q} \overline{\mathcal{M}}_{\widehat{\Gamma}} \times_{\overline{\mathcal{M}}_{\Gamma}} \overline{\mathcal{M}}_{\Gamma}^{1/r} \xrightarrow{\mu} \overline{\mathcal{M}}_{\Gamma}^{1/r} \xrightarrow{i_{\Gamma}} \overline{\mathcal{M}}_{0,k,l}^{1/r}. \quad (4.4)$$

Here, $\overline{\mathcal{M}}_{\Gamma}$, $\overline{\mathcal{M}}_{\widehat{\Gamma}}$ are the moduli spaces of marked disks with dual graphs $\Gamma, \widehat{\Gamma}$, respectively. The morphism q is defined by sending the spin structure S to \widehat{S} as in (2.17).

The map μ is an isomorphism, though we distinguish between its domain and codomain because they have different universal objects. While the map q is in general not an isomorphism (see Remark 4.5 below), it has degree one. This fact is known in the closed case (see, e.g., [11]); the key point is that each element of $\overline{\mathcal{M}}_{\widehat{\Gamma}}^{1/r}$ has a single geometric preimage under q , and in both the domain and codomain of q , the order of the automorphism group is r^c , where c is the number of components. By the analysis of automorphisms in Proposition 2.17, the same argument applies in the open case.

Remark 4.5. The reason that q is not, in general, an isomorphism is that it does not induce an isomorphism on automorphism groups. Automorphisms on a normalized curve are given by separate fiberwise scalings on each sphere component; on a nodal curve, however, scalings by ζ and η at opposite branches of a node only glue to give a global automorphism if one can act by a ghost automorphism of C to make the scalings at the two branches agree—in other words (by Remark 2.14), if there exists $\xi \in \mathbb{Z}/r\mathbb{Z}$ for which $\zeta \cdot \xi^m = \eta$, where m is the multiplicity of S at the ζ -branch of the node. Thus, q is an isomorphism only if each internal node satisfies $\gcd(r, m) = 1$, so that such ξ exists. (Boundary nodes need not satisfy any condition, since fiberwise scaling on a disk component is not an automorphism of the graded spin structure.)

Notation 4.6. For Γ and N as above, we denote the map $q \circ \mu^{-1} : \overline{\mathcal{M}}_{\Gamma}^{1/r} \rightarrow \overline{\mathcal{M}}_{\widehat{\Gamma}}^{1/r}$ by Detach_N . When $N = \{e\}$ is a singleton, we denote this map by Detach_e .

There are two natural universal curves over the fiber product $\overline{\mathcal{M}}_{\widehat{\Gamma}} \times_{\overline{\mathcal{M}}_{\Gamma}} \overline{\mathcal{M}}_{\Gamma}^{1/r}$. We define \mathcal{C}_{Γ} by the fiber diagram

$$\begin{array}{ccc} \mathcal{C}_{\Gamma} & \longrightarrow & \mathcal{C} \\ \pi \downarrow & & \downarrow \\ \overline{\mathcal{M}}_{\widehat{\Gamma}} \times_{\overline{\mathcal{M}}_{\Gamma}} \overline{\mathcal{M}}_{\Gamma}^{1/r} & \xrightarrow{i_{\Gamma} \circ \mu} & \overline{\mathcal{M}}_{0,k,l}^{1/r} \end{array}$$

and $\mathcal{C}_{\widehat{\Gamma}}$ by the fiber diagram

$$\begin{array}{ccc} \widehat{\mathcal{C}} & \xleftarrow{\quad} & \mathcal{C}_{\widehat{\Gamma}} \\ \downarrow & & \downarrow \widehat{\pi} \\ \overline{\mathcal{M}}_{\widehat{\Gamma}}^{1/r} & \xleftarrow{q} & \overline{\mathcal{M}}_{\widehat{\Gamma}} \times_{\overline{\mathcal{M}}_{\Gamma}} \overline{\mathcal{M}}_{\Gamma}^{1/r}, \end{array}$$

in which $\widehat{\mathcal{C}}$ is the universal curve over $\overline{\mathcal{M}}_{\widehat{\Gamma}}^{1/r}$. There are universal bundles \mathcal{S} and $\widehat{\mathcal{S}}$ on these two universal curves, and they are related by the universal normalization morphism

$$n : \mathcal{C}_{\widehat{\Gamma}} \rightarrow \mathcal{C}_{\Gamma}.$$

We can now state the decomposition properties of the Witten bundle. We state the properties in the case where $N = \{e\}$ for a single edge e , but all can be readily generalized to the setting where more than one edge is detached.

Proposition 4.7. Let Γ be a stable genus-zero twisted r -spin dual graph with a lifting. Suppose that Γ has a single edge e , so the general point of $\overline{\mathcal{M}}_{\Gamma}^{1/r}$ is a stable r -spin disk with two components C_1 and C_2 meeting at a node p . Let $\widehat{\Gamma} = \text{detach}_e(\Gamma)$. Let \mathcal{W} and $\widehat{\mathcal{W}}$ denote the Witten bundles on $\overline{\mathcal{M}}_{0,k,l}^{1/r}$ and $\overline{\mathcal{M}}_{\widehat{\Gamma}}^{1/r}$, respectively.

Then, topologically, the Witten bundle decomposes as follows along the node p :

- (i) If e is a Neveu–Schwarz edge, then $\mu^* i_{\Gamma}^* \mathcal{W} = q^* \widehat{\mathcal{W}}$.
- (ii) If e is a Ramond boundary edge, then there is an exact sequence

$$0 \rightarrow \mu^* i_{\Gamma}^* \mathcal{W} \rightarrow q^* \widehat{\mathcal{W}} \rightarrow \mathcal{T}_+ \rightarrow 0, \quad (4.5)$$

where \mathcal{T}_+ is a trivial real line bundle.

- (iii) If e is a Ramond internal edge connecting two closed vertices, write $q^* \widehat{\mathcal{W}} = \widehat{\mathcal{W}}_1 \boxplus \widehat{\mathcal{W}}_2$, in which $\widehat{\mathcal{W}}_1$ is the Witten bundle on the component containing the anchor of Γ (defined via $\widehat{\mathcal{S}}|_{C_1}$) and $\widehat{\mathcal{W}}_2$ is the Witten bundle on the other component. Then there is an exact sequence

$$0 \rightarrow \widehat{\mathcal{W}}_2 \rightarrow \mu^* i_{\Gamma}^* \mathcal{W} \rightarrow \widehat{\mathcal{W}}_1 \rightarrow 0. \quad (4.6)$$

Furthermore, if $\widehat{\Gamma}'$ is defined to agree with $\widehat{\Gamma}$ except that the twist at each Ramond tail is $r-1$, and $q' : \overline{\mathcal{M}}_{\widehat{\Gamma}'} \times_{\overline{\mathcal{M}}_{\Gamma}} \overline{\mathcal{M}}_{\Gamma}^{1/r} \rightarrow \overline{\mathcal{M}}_{\widehat{\Gamma}'}^{1/r}$ is defined analogously

to q , then there is an exact sequence

$$0 \rightarrow \mu^* i_{\Gamma}^* \mathcal{W} \rightarrow (q')^* \widehat{\mathcal{W}}' \rightarrow \mathcal{T} \rightarrow 0, \quad (4.7)$$

where $\widehat{\mathcal{W}}'$ is the Witten bundle on $\overline{\mathcal{M}}_{\widehat{\Gamma}'}^{1/r}$ and \mathcal{T} is a line bundle whose r th power is trivial.

- (iv) If e is a Ramond internal edge connecting an open vertex to a closed vertex, write $q^* \widehat{\mathcal{W}} = \widehat{\mathcal{W}}_1 \boxplus \widehat{\mathcal{W}}_2$, in which $\widehat{\mathcal{W}}_1$ is the Witten bundle on the disk component (defined via $\widehat{\mathcal{S}}|_{\mathcal{C}_1}$) and $\widehat{\mathcal{W}}_2$ is the Witten bundle on the sphere component. Then the exact sequences (4.6) and (4.7) both hold.
- (v) Suppose that Γ has a single vertex, no edges, and a contracted boundary tail t , and let $\widehat{\Gamma} = \text{detach}_t(\Gamma)$. If \mathcal{W} and $\widehat{\mathcal{W}}$ denote the Witten bundles on $\overline{\mathcal{M}}_{0,k,l}^{1/r}$ and $\overline{\mathcal{M}}_{\widehat{\Gamma}}^{1/r}$, respectively, then the sequence (4.5) also holds in this case.

Remark 4.8. We say that the Witten bundle decomposes “topologically” as above to emphasize that, while the coarse underlying Witten bundle behaves as above, the action of the isotropy groups of the moduli space on the fibers may not respect these identifications. (This is only relevant for internal nodes, since boundary and contracted boundary nodes do not contribute isotropy to the moduli space.) For example, in the case of a Neveu–Schwarz internal node, an element of the fiber of $q^* \widehat{\mathcal{W}}$ is acted on by independently scaling the sections of J on the two components, while Remark 4.5 shows that this is not in general possible for elements of the fiber of $\mu^* i_{\Gamma}^* \mathcal{W}$.

Proof of Proposition 4.7. First, let us fix some notation. Letting $\mathcal{J} = \mathcal{S}^{\vee} \otimes \omega_{\pi}$ and decomposing $\mathcal{C}_{\widehat{\Gamma}}$ into components \mathcal{C}_1 and \mathcal{C}_2 , we define:

$$\pi_i = \widehat{\pi}|_{\mathcal{C}_i}, \quad \mathcal{S}_i = n^* \mathcal{S}|_{\mathcal{C}_i}, \quad \widehat{\mathcal{S}}_i = \widehat{\mathcal{S}}|_{\mathcal{C}_i}, \quad \widehat{\mathcal{J}}_i = \widehat{\mathcal{S}}_i^{\vee} \otimes \omega_{\pi_i}$$

for $i = 1, 2$. We view $\mu^* i_{\Gamma}^* \mathcal{W} = (R^0 \pi_* \mathcal{J})_+$ and $q^* \widehat{\mathcal{W}} = (R^0 \pi_{1*} \widehat{\mathcal{J}}_1 \oplus R^0 \pi_{2*} \widehat{\mathcal{J}}_2)_+$.

Suppose that e is Neveu–Schwarz. Then the normalization exact sequence yields

$$0 \rightarrow \mathcal{S} \rightarrow n_* n^* \mathcal{S} \rightarrow \mathcal{S}|_{\Delta_p} \rightarrow 0,$$

where $\Delta_p \subseteq \mathcal{C}_{\Gamma}$ is the orbifold divisor corresponding to the node p . Since the twist of every tail of Γ is non-negative, except at most one tail that may have twist -1 , we have

$R^0\pi_{1*}\mathcal{S}_1 = R^0\pi_{2*}\mathcal{S}_2 = 0$, and we obtain

$$0 \rightarrow \sigma_p^*\mathcal{S} \rightarrow R^1\pi_*\mathcal{S} \rightarrow R^1\pi_{1*}\mathcal{S}_1 \oplus R^1\pi_{2*}\mathcal{S}_2 \rightarrow 0. \quad (4.8)$$

The assumption that e is Neveu–Schwarz implies both that $\widehat{\mathcal{S}}_i = \mathcal{S}_i$ for $i = 1, 2$ and that $\sigma_p^*\mathcal{S} = 0$, since sections of an orbifold line bundle necessarily vanish at Neveu–Schwarz points. Thus, dualizing and taking involution-invariant parts of (4.8) shows that the fibers of $\mu^*i_\Gamma^*\mathcal{W}$ and $q^*\widehat{\mathcal{W}}$ are canonically identified.

Suppose now that e is a Ramond boundary edge. Then

$$\widehat{\mathcal{S}}_i = \mathcal{S}_i \otimes \mathcal{O}(-r\Delta_{i,p}),$$

where $\Delta_{i,p} \subseteq \mathcal{C}_i$ is the orbifold divisor corresponding to the half node in \mathcal{C}_i . The normalization exact sequence for \mathcal{J} yields

$$0 \rightarrow R^0\pi_*\mathcal{J} \rightarrow R^0\widehat{\pi}_*(n^*\mathcal{J}) \rightarrow \sigma_p^*\mathcal{J} \rightarrow 0. \quad (4.9)$$

Now, passing to coarse underlying bundles (which does not affect cohomology), the middle term can be re-written:

$$|n^*\mathcal{J}|_{\mathcal{C}_i} = |n|^* \left(|\mathcal{S}|^\vee \otimes \omega_{|\pi|} \right) \Big|_{|\mathcal{C}_i|} = |\mathcal{S}_i|^\vee \otimes \mathcal{O}(\Delta_{i,p}) \otimes \omega_{|\pi|} = |\widehat{\mathcal{J}}_i|.$$

Thus, the sequence (4.9) can be re-expressed as

$$0 \rightarrow R^0\pi_*\mathcal{J} \rightarrow R^0\pi_{1*}\widehat{\mathcal{J}}_1 \oplus R^0\pi_{2*}\widehat{\mathcal{J}}_2 \rightarrow \sigma_p^*\mathcal{J} \rightarrow 0. \quad (4.10)$$

Because e is Ramond, the bundle $\mathcal{T} := (\sigma_p^*\mathcal{J})$ has trivial r th power, using that the restriction of both ω_π and $\omega_{\pi,\log}$ to the locus of nodes is trivial. Taking involution-invariant parts in (4.10) yields (4.5), where the real line \mathcal{T}_+ is trivialized using the grading.

Next, suppose that e is a Ramond internal edge connecting two closed vertices. Denote the two vertices of Γ by v_1 and v_2 , and let v_1 be the vertex supporting the anchor. Then $R^0\pi_{2*}\mathcal{S}_2 = 0$, and if v_1 contains at least one tail of positive twist, then $R^0\pi_{1*}\mathcal{S}_1 = 0$, as well. In this case, we still have the sequence (4.8). We also have

$$0 \rightarrow \sigma_p^*\mathcal{S} \rightarrow R^1\pi_{1*}\widehat{\mathcal{S}}_1 \rightarrow R^1\pi_{1*}\mathcal{S}_1 \rightarrow 0. \quad (4.11)$$

The sequences (4.8) and (4.11) fit together into a commutative diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \sigma_p^* \mathcal{S} & \longrightarrow & R^1 \pi_* \widehat{\mathcal{S}}_1 & \longrightarrow & R^1 \pi_* \mathcal{S}_1 & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & \sigma_p^* \mathcal{S} & \longrightarrow & R^1 \pi_* \mathcal{S} & \longrightarrow & R^1 \pi_* \mathcal{S}_1 \oplus R^1 \pi_* \mathcal{S}_2 & \longrightarrow & 0,
 \end{array}$$

in which the middle vertical arrow can be constructed by dualizing the sequence (4.10). By the Snake Lemma the cokernel of the middle vertical arrow is $R^1 \pi_{2*} \mathcal{S}_2 = R^1 \pi_{2*} \widehat{\mathcal{S}}_2$, so

$$0 \rightarrow R^1 \pi_{1*} \widehat{\mathcal{S}}_1 \rightarrow R^1 \pi_* \mathcal{S} \rightarrow R^1 \pi_{2*} \widehat{\mathcal{S}}_2 \rightarrow 0.$$

Dualizing and taking involution-invariant parts prove (4.6).

Still assuming that e is a Ramond internal edge connecting two closed vertices, suppose that every tail of v_1 except the anchor has twist zero. In this case the anchor must have twist -1 . Then $R^0 \pi_{1*} \mathcal{S}_1$ is one-dimensional, and in the normalization exact sequence

$$0 \rightarrow R^0 \pi_{1*} \mathcal{S}_1 \oplus R^1 \pi_{2*} \mathcal{S}_2 \rightarrow \sigma_p^* \mathcal{S} \rightarrow R^1 \pi_* \mathcal{S} \rightarrow R^1 \pi_{1*} \mathcal{S}_1 \oplus R^1 \pi_{2*},$$

the 1st map is an isomorphism. Hence, $R^1 \pi_* \mathcal{S} \cong R^1 \pi_{1*} \mathcal{S}_1 \oplus R^1 \pi_{2*} \mathcal{S}_2$. A similar argument shows that the 1st map in

$$0 \rightarrow R^0 \pi_{1*} \mathcal{S}_1 \rightarrow \sigma_p^* \mathcal{S}_1 \rightarrow R^0 \pi_{1*} \widehat{\mathcal{S}}_1 \rightarrow R^0 \pi_{1*} \mathcal{S}_1 \rightarrow 0$$

is an isomorphism, so we also have $R^1 \pi_{1*} \widehat{\mathcal{S}}_1 \cong R^1 \pi_{1*} \mathcal{S}_1$. Since $\widehat{\mathcal{S}}_2 = \mathcal{S}_2$ by construction, we conclude that $R^1 \pi_* \mathcal{S} \cong R^1 \pi_{1*} \widehat{\mathcal{S}}_1 \oplus R^1 \pi_{2*} \widehat{\mathcal{S}}_2$, which implies that (4.6) holds (and, in fact, splits) in this case. The proof of (4.7) is identical to the proof of (4.5) above.

If e is a Ramond internal edge connecting an open vertex to a closed vertex, then the proof that (4.6) holds is identical to the proof for an edge joining two closed vertices (with $R^0 \pi_{1*} \mathcal{S}_1 = R^0 \pi_{2*} \mathcal{S}_2 = 0$), and the proof that (4.7) holds is exactly as in (4.5). Finally, in the situation where Γ has a single vertex, no edges, and a contracted boundary tail, the exact sequence (4.5) still holds, by the same proof. ■

One further observation will be needed later.

Observations 4.9. Suppose $I' \subseteq [l]$ is such that $\sum_{i \in I'} a_i \leq r - 1$. Let Γ be a graph with exactly two vertices, a closed vertex v^c containing exactly the internal tails labeled by

I' and an open vertex v^o , connected by a single edge. Then $\overline{\mathcal{M}}_{\widehat{\Gamma}}^{1/r} = \overline{\mathcal{M}}_{v^c} \times \overline{\mathcal{M}}_{v^o}$. One can compute that the twist at the node is $r - 2 - \sum_{i \in I'} a_i$ and the Witten bundle \mathcal{W}_{v^c} on $\overline{\mathcal{M}}_{v^c}$ has rank zero, so Proposition 4.7 implies that

$$\mu^* i_{\Gamma}^* \mathcal{W}_{\Gamma} = q^* \pi_2^* \mathcal{W}_{v^o},$$

where π_2 is the projection to the 2nd factor.

4.3 Cotangent line bundles

For each $i \in I$, a cotangent line bundle \mathbb{L}_i is defined on the moduli space of stable marked disks as the line bundle whose fiber over $(C, \phi, \Sigma, \{z_i\}, \{x_j\}, m^I, m^B)$ is the cotangent line $T_{z_i}^* \Sigma$. Alternatively, \mathbb{L}_i is the pullback via the doubling map of the usual line bundle \mathbb{L}_i on $\overline{\mathcal{M}}_{0,k+2l}$. We define cotangent line bundles \mathbb{L}_i on $\overline{\mathcal{M}}_{0,k,l}^{1/r}$ by the pullback under the morphism forgetting the spin structure, and for any graph Γ , we let \mathbb{L}_i^{Γ} be the pullback of \mathbb{L}_i to $\overline{\mathcal{M}}_{\Gamma}^{1/r}$.

We mention a few important properties of these bundles in the observation below. The proofs are identical to the proofs of the analogous statements in [24, Section 3.5].

Observations 4.10.

- (i) If e is an edge of Γ , Γ_1 and Γ_2 are the two connected components of $\text{detach}_e(\Gamma)$, and i is a marking of an internal tail of Γ_1 , then

$$\mathbb{L}_i^{\Gamma} = \text{Proj}_1^* \mathbb{L}_i^{\Gamma_1},$$

where $\text{Proj}_1 : \overline{\mathcal{M}}_{\Gamma}^{1/r} \rightarrow \overline{\mathcal{M}}_{\Gamma_1}^{1/r}$ is the projection.

- (ii) If $B' \subseteq \mathbb{Z} \setminus \{i\}$, $I' \subseteq \mathbb{Z}$, and $\Gamma' = \text{for}_{B', I'}(\Gamma)$, then there exists a canonical morphism

$$t_{\Gamma, B', I'} : \text{For}_{B', I'}^* \mathbb{L}_i^{\Gamma'} \rightarrow \mathbb{L}_i^{\Gamma},$$

which is an isomorphism away from strata where the component containing z_i is contracted by the forgetful map, and which vanishes identically on the remaining strata.

- (iii) \mathbb{L}_i^{Γ} is canonically oriented as a complex line bundle.

5 Orientation: Constructions and Properties

5.1 Relative orientation of the Witten bundle

The open Witten bundle does not carry a canonical complex orientation. Nevertheless, it is canonically relatively oriented, relative to the moduli space of stable graded r -spin disks. Before stating this precisely, we require some notation.

Consider a set B of boundary markings, a set I of internal markings, and a set of twists $\{a_i\}_{i \in I}$. Given decompositions $I = I_1 \sqcup I_2$ and $B = B_1 \sqcup B_2$ such that

$$|B_1| - 1 = \sum_{i \in I_1} a_i \pmod{r}, \quad |B_2| = \sum_{i \in I_2} a_i \pmod{r} \quad (5.1)$$

and hence

$$\frac{2 \sum_{i \in I_2} a_i + |B_2|(r-2)}{r} \equiv |B_2| \pmod{2},$$

let $\Gamma = \Gamma_{I_1, I_2; B_1, B_2}$ be the graph with two open vertices v_1 and v_2 joined by a boundary edge e , in which v_i contains the internal tails marked by I_i and the boundary tails marked by B_i . If h_i is the half edge of e incident to v_i , then a straightforward computation shows that

$$\text{tw}(h_1) = \text{alt}(h_1) = 0.$$

Let us denote by $\mathcal{B}\Gamma$ the graph with vertices v'_1 and v_2 obtained by detaching e and forgetting h_1 ; this is a very special case of the notion of the “base” of a graph Γ , defined in general in the sequel. Let $\overline{\mathcal{M}}_{v'_1}^{1/r}$, $\overline{\mathcal{M}}_{v_1}^{1/r}$, and $\overline{\mathcal{M}}_{v_2}^{1/r}$ be the moduli spaces of stable graded r -spin disks corresponding to the vertices v'_1 , v_1 , and v_2 , thought of as r -spin graphs.

By the discussion in Section 3.2, the maps q and μ are isomorphisms in the case of a single boundary edge, which implies that

$$\overline{\mathcal{M}}_{\Gamma}^{1/r} \cong \overline{\mathcal{M}}_{v'_1}^{1/r} \times \overline{\mathcal{M}}_{v_2}^{1/r}.$$

Composing with the forgetful map forgetting the twist-zero tail h_1 on v_1 , we obtain a map

$$F_{\Gamma} : \overline{\mathcal{M}}_{\Gamma}^{1/r} \rightarrow \overline{\mathcal{M}}_{\mathcal{B}\Gamma}^{1/r} \cong \overline{\mathcal{M}}_{v'_1}^{1/r} \times \overline{\mathcal{M}}_{v_2}^{1/r}.$$

By Proposition 4.7 and the fact that the Witten class pulls back under the forgetful map,

$$i_\Gamma^* \mathcal{W} = F_\Gamma^* \mathcal{W}_{\mathcal{B}\Gamma} = \mathcal{W}_{v'_1} \boxplus \mathcal{W}_{v_2}. \quad (5.2)$$

In this situation, we also have

$$i_\Gamma^* \det(T\overline{\mathcal{M}}_{0,B,\{a_i\}_{i \in I}}^{1/r}) \cong \det(N) \otimes \det(f_\Gamma) \otimes F_\Gamma^* \det(T\overline{\mathcal{M}}_{\mathcal{B}\Gamma}^{1/r}), \quad (5.3)$$

where N is again the normal bundle of $\overline{\mathcal{M}}_\Gamma^{1/r}$, and f_Γ is the fiber of F_Γ (which can be identified with the fiber of the map on $\overline{\mathcal{M}}_{v_1}$ that forgets the marked point x_{h_1} associated to h_1). Note that f_Γ also carries a canonical orientation, as a subset of the boundary of a disk.

The main theorem of this section is the following.

Theorem 5.1. The morphism $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{0,B,\{a_i\}_{i \in I}}^{1/r}$ is canonically relatively oriented. More precisely, for any sets B and I of boundary and internal markings, and any set $\{a_i\}_{i \in I}$ of the internal twists, there exists a distinguished orientation $o_{0,B,\{a_i\}_{i \in I}}$ of $T\overline{\mathcal{M}}_{0,B,\{a_i\}_{i \in I}}^{1/r} \oplus \mathcal{W}$. These orientations satisfy the following properties:

- (i) If $f^B : B \rightarrow B'$ and $f^I : I \rightarrow I'$ are bijections such that f^I preserves twists, and if

$$F : \overline{\mathcal{M}}_{0,B,\{a_i\}_{i \in I}}^{1/r} \rightarrow \overline{\mathcal{M}}_{0,B',\{a_i\}_{i \in I'}}^{1/r}$$

is the induced map, then $o_{0,B,\{a_i\}_{i \in I}} = F^* o_{0,B',\{a_i\}_{i \in I'}}$. In particular the canonical orientation is invariant to renumbering marked points.

- (ii) If $I = I_1 \sqcup I_2$ and $B = B_1 \sqcup B_2$ satisfy (5.1) and $\Gamma = \Gamma_{I_1, I_2; B_1, B_2}$, then under the isomorphisms (5.2) and (5.3), we have

$$i_\Gamma^* o_{0,B,\{a_i\}_{i \in I}} = o_N \otimes o_{h_1} \otimes F_\Gamma^*(o_{0,B_1,I_1} \boxtimes o_{\{h_2\} \cup B_2, I_2}),$$

in which o_N and o_{h_1} are the canonical orientations on N and f_Γ , respectively (Since the rank of the Witten bundle and the dimension of the moduli are of the same parity, the canonical orientations represent *even variables*, this means, in particular, that $i_\Gamma^* o_{0,B,\{a_i\}_{i \in I}} = o_N \otimes o_{h_1} \otimes F_\Gamma^*(o_{\{h_2\} \cup B_2, I_2} \boxtimes o_{0,B_1,I_1})$ also holds.).

(iii) If Γ is a graph with one open vertex v^o and one closed vertex v^c , then under the identification

$$i_\Gamma^* \det(T\overline{\mathcal{M}}_{0,B,\{a_i\}_{i \in I}}^{1/r}) = \det(N) \otimes (\det(T\overline{\mathcal{M}}_{v^o}) \boxtimes \det(T\overline{\mathcal{M}}_{v^c})),$$

we have

$$i_\Gamma^* o_{0,B,\{a_i\}_{i \in I}} = o_N \otimes (o_{v^o} \boxtimes o_{v^c}), \quad (5.4)$$

where o_N is canonically defined as above and o_{v^o}, o_{v^c} are the orientations on the total space associated to the two vertices, o_{v^c} is the standard complex one.

Remark 5.2. Note that in the last item we have used that if $0 \rightarrow A \xrightarrow{f} B \rightarrow C \rightarrow 0$ is an exact sequence of vector bundles, then there is a canonical isomorphism

$$\det(A) \otimes \det(C) \cong \det(B).$$

In case e is Neveu–Schwarz, the Witten bundle decomposes as a direct sum by Proposition 4.7, but when e is Ramond, we use (4.6) in order to write (5.4). In both cases, the decomposition may not respect the isotropy group actions, but these actions are orientation preserving as they are induced from scaling the closed Witten bundle summand by roots of unity.

The proof of Theorem 5.1 is the content of the remainder of this section. Equipped with this theorem, we will have a canonical relative orientation o_Γ on $\mathcal{W} \rightarrow \overline{\mathcal{M}}_\Gamma^{1/r}$ whenever Γ is a connected smooth graded graph. Furthermore, we can extend the definition to the case where Γ is a smooth graded graph, but not necessarily connected, by putting

$$o_\Gamma = \boxtimes_{\Lambda \in \text{Conn}(\Gamma)} o_\Lambda,$$

in which we use that the Witten bundle \mathcal{W} on $\overline{\mathcal{M}}_\Gamma^{1/r}$ is canonically isomorphic to the direct sum of the Witten bundles on the moduli spaces $\overline{\mathcal{M}}_\Lambda^{1/r}$ associated to $\Lambda \in \text{Conn}(\Gamma)$.

The key point in the proof of Theorem 5.1 is Construction 5.6, which defines an explicit frame for the Witten bundle of tuples $(\{a_i\}_{i \in I}, k)$, for $k = k(a_1, \dots, a_l)$. By

studying the properties of these frames and using compatibility relations we are able to construct orientations in full generality and analyze their properties.

Remark 5.3. Theorem 5.1 determines a unique family of orientations, up to the following ambiguity: if $o_{0,B,\{a_i\}_{i \in I}}$ is a family satisfying the conditions of the theorem, then $(-1)^{|B|-1} o_{0,B,\{a_i\}_{i \in I}}$ also satisfies these conditions; see Lemma 5.17 and Remark 5.18 below.

5.2 Extending the internal twists

Throughout this section, we write $\mathcal{W}_{0,k,\vec{a}}$ for the Witten bundle on $\overline{\mathcal{M}}_{0,k,\vec{a}}^{1/r}$ for any tuple $\vec{a} = \{a_1, \dots, a_l\}$.

Thus far, we have assumed that in the open r -spin moduli space $\overline{\mathcal{M}}_{0,k,\{a_1, \dots, a_l\}}^1$, the internal twists satisfy $a_i \in \{-1, 0, 1, \dots, r-1\}$. However, when defining orientations, it is useful to allow more general twists. In fact, there exists a smooth orientable orbifold with corners $\overline{\mathcal{M}}_{0,k,\vec{a}}^{1/r}$ for *any* tuple of non-negative integers $\vec{a} = \{a_1, \dots, a_l\}$, parameterizing stable nodal marked orbifold Riemann surfaces with boundary together with an orbifold line bundle S , an isomorphism

$$|S|^{\otimes r} \cong \omega_{|C|} \otimes \mathcal{O} \left(- \sum_{i \in I} a_i [z_i] - \sum_{i \in I} a_i [\bar{z}_i] - \sum_{j \in B} (r-2) [x_j] \right),$$

and a grading. Moreover, there is a Witten bundle on this moduli space, defined as before.

The relationship between the Witten bundle with twist a_i and twist $a_i + r$ was observed by Jarvis–Kimura–Vaintrob [18] in the closed case, and the same is true in the open theory.

Lemma 5.4. Let $\vec{a} = \{a_1, \dots, a_l\}$ be a tuple of non-negative integers, and let $\vec{a}' = \{a_1 + r, a_2, \dots, a_l\}$. Then there is an isomorphism

$$\kappa : \overline{\mathcal{M}}_{0,k,\vec{a}'}^{1/r} \rightarrow \overline{\mathcal{M}}_{0,k,\vec{a}}^{1/r},$$

and the Witten bundles on $\overline{\mathcal{M}}_{0,k,\vec{a}'}^{1/r}$ and $\overline{\mathcal{M}}_{0,k,\vec{a}}^{1/r}$ are related by a short exact sequence

$$0 \rightarrow \tilde{\mathbb{L}} \rightarrow \mathcal{W}_{0,k,\vec{a}'} \rightarrow \kappa^* \mathcal{W}_{0,k,\vec{a}} \rightarrow 0,$$

in which $\tilde{\mathbb{L}}$ is an orbifold line bundle satisfying

$$\tilde{\mathbb{L}}^{\otimes r} \cong (\mathbb{L}_1^\vee)^{\otimes(a_1+1)}.$$

Proof. The isomorphism κ is given by sending the twisted r -spin structure S' to

$$S := S' \otimes \mathcal{O}(r[z_1] + r[\bar{z}_1]),$$

where $[z_1]$ and $[\bar{z}_1]$ denote the orbifold divisors of the 1st internal marked point and its conjugate, so that

$$|S| = |S'| \otimes \mathcal{O}([z_1] + [\bar{z}_1])$$

on $|C|$. If S' denotes the universal twisted r -spin structure on the universal curve \mathcal{C} over $\overline{\mathcal{M}}_{0,k,\vec{a}'}^{1/r}$ and $S := S' \otimes \mathcal{O}(\Delta_{z_1} + \Delta_{\bar{z}_1})$ for the divisors $\Delta_{z_1}, \Delta_{\bar{z}_1} \subseteq \mathcal{C}$ corresponding to z_1 and \bar{z}_1 , then the short exact sequence

$$0 \rightarrow |S'| \rightarrow |S| \rightarrow |S|_{\Delta_{z_1} + \Delta_{\bar{z}_1}} \rightarrow 0$$

implies

$$0 \rightarrow R^0\pi_*\left(|S|_{\Delta_{z_1} + \Delta_{\bar{z}_1}}\right) \rightarrow R^1\pi_*|S'| \rightarrow R^1\pi_*|S| \rightarrow 0.$$

Taking $\tilde{\phi}$ -invariants part yields

$$0 \rightarrow \sigma_1^*|S| \rightarrow \left(R^1\pi_*|S'|\right)_+ \rightarrow \left(R^1\pi_*|S|\right)_+ \rightarrow 0,$$

where σ_1 is the section of $|\mathcal{C}|$ associated to the 1st internal marked point. The 1st bundle in this sequence satisfies

$$(\sigma_1^*|S|)^{\otimes r} \cong \sigma_1^*\left(\omega_{|\pi|, \log}\left(-\sum_{i=1}^l (a_i + 1)\Delta_{z_i}\right)\right) \cong \mathbb{L}_1^{\otimes(a_1+1)},$$

since $\sigma_1^*\omega_{|\pi|, \log}$ is trivial. Dualizing thus proves the claim. ■

5.3 Orientability of the Witten bundle

Before we show how to choose canonical orientations, we first confirm that the Witten bundles are orientable. For this, recall the action of S_B on $\overline{\mathcal{M}}_{0,B,\tilde{a}}^{1/r}$ by permuting markings, and note that this action lifts canonically to $\mathcal{W}_{0,k,\tilde{a}}$.

Lemma 5.5. The bundles $\mathcal{W}_{0,k,\{a_1,\dots,a_l\}}$ are orientable.

Moreover, suppose $g \in S_B$ preserves the component $\overline{\mathcal{M}}_{0,B,\tilde{a}}^{1/r,\hat{\pi}}$, meaning that there exists an integer h such that

$$g(\pi(i)) = \pi(i + h),$$

interpreted cyclically. Then, for any orientation \mathfrak{o} of $\mathcal{W}_{0,k,\{a_1,\dots,a_l\}} \rightarrow \overline{\mathcal{M}}_{0,B,\tilde{a}}^{1/r,\hat{\pi}}$, the permutation g acts with sign ε^h , where $\varepsilon = \varepsilon_k = (-1)^{k-1}$. The action of $g \in S_l$ by permuting the internal markings, which also lifts canonically to $\mathcal{W}_{0,k,\{a_1,\dots,a_l\}}$, is orientation preserving.

Proof. Suppose, first, that $l \geq 1$ and $k - 1 \geq \sum a_i$. Observe that

$$k - 1 \equiv \sum a_i \pmod{r}.$$

There exists a (usually non-unique) tuple $(\tilde{a}_1, \dots, \tilde{a}_l)$ with

$$\tilde{a}_i \geq 0, \quad \tilde{a}_i \equiv a_i \pmod{r}, \quad \sum \tilde{a}_i = k - 1.$$

By Lemma 5.4, the bundles $\mathcal{W} := \mathcal{W}_{0,k,\{a_1,\dots,a_l\}}$ and $\tilde{\mathcal{W}} := \mathcal{W}_{0,k,\{\tilde{a}_1,\dots,\tilde{a}_l\}}$ differ by a bundle E that is a direct sum of complex line bundles. Thus, one has

$$\det(E) \otimes \det(\mathcal{W}) \cong \det(\tilde{\mathcal{W}}),$$

so, since E is canonically oriented, orienting \mathcal{W} is equivalent to orienting $\tilde{\mathcal{W}}$. Note that S_B and S_l act with the trivial sign on E . We can thus reduce to the case where $k - 1 = \sum a_i$.

In this case, the Witten bundle is of real rank $e = \sum a_i = k - 1$. Since the boundary of the base space does not affect the orientability of a bundle, $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{0,k,\{a_1,\dots,a_l\}}^{\frac{1}{r}}$ is orientable precisely if $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{0,k,\{a_1,\dots,a_l\}}^{\frac{1}{r}} \setminus \partial \mathcal{M}_{0,k,\{a_1,\dots,a_l\}}^{\frac{1}{r}}$ is orientable. Furthermore, this is the case exactly if \mathcal{W} is orientable on the moduli space $\mathcal{M}_{0,k,\{a_1,\dots,a_l\}}^{\frac{1}{r}}$ of smooth graded r -spin disks, since compact strata of real codimension two do not affect orientability. It is therefore enough to prove that the bundle \mathcal{W} is orientable

over each connected component $\mathcal{M}_{0,k,\{a_1,\dots,a_l\}}^{\frac{1}{r},\hat{\pi}}$ of $\mathcal{M}_{0,k,\{a_1,\dots,a_l\}}^{\frac{1}{r}}$ associated to an order $\pi : [k] \rightarrow [k]$ with induced cyclic order $\hat{\pi}$.

The following gives explicit sections of $\mathcal{W} \rightarrow \mathcal{M}_{0,k,\{a_1,\dots,a_l\}}^{\frac{1}{r},\hat{\pi}}$ that form a basis in any fiber.

Construction/Notation 5.6. Let Σ be a smooth graded r -spin disk. Identifying Σ with the upper half plane, let

$$\xi_{ij} = \xi_{ij}^\pi = \frac{(x_{\pi(j)} - x_{\pi(i)})dw}{(w - x_{\pi(i)})(w - x_{\pi(j)})}, \quad i, j \in [k], \quad \xi_j = \frac{\sqrt{-1}(\bar{z}_j - z_j)dw}{(w - z_j)(w - \bar{z}_j)}, \quad j \in [l],$$

where $\sqrt{-1}$ is the root in the upper half plane. Define sections

$$(\sigma_j)_\Sigma = (\sigma_j^\pi)_\Sigma = (\sigma_{0,k,\{a_1,\dots,a_l\};j}^\pi)_\Sigma = \left((-1)^{r+1} \prod_{i \in [l]} \xi_i^{a_i} \prod_{i \in [k]} \xi_{i(i+1)}^{-1} \xi_{1(1+j)}^r \right)^{\frac{1}{r}} \in H^0(\Sigma, |J|),$$

for $j \in [k-1]$, where $k+1$ is taken to be 1. More precisely, inside the parentheses in the above formula, a global section of

$$\omega_{|C|}^{\otimes r-1} \otimes \mathcal{O} \left(\sum_{i=1}^l a_i [z_i] + \sum_{i=1}^l a_i [\bar{z}_i] + \sum_{j=1}^k (r-2) [x_j] \right) \cong |J|^{\otimes r}$$

is written. This section is ϕ -invariant and positive on the arc from x_1 to x_{1+j} , with respect to the canonical orientation. One can construct an r th root in the sense of a global section σ_j of $|J|$ whose r th power is mapped to the expression in parentheses. Indeed, this can be done locally away from special points, and the order of zeroes and poles at special points guarantee that the construction extends globally and univalently. When r is odd, a real r th root is unique. When r is even, there are two real r th roots, one of which agrees with the grading on the arc from x_1 to x_{1+j} and the other of which agrees on the complementary arc; we choose σ_j to be the one that agrees on the arc from x_1 to x_{1+j} . ■

We claim that, for any $\Sigma \in \mathcal{M}_{0,k,\{a_1,\dots,a_l\}}^{\frac{1}{r},\hat{\pi}}$, the sections $(\sigma_j^\pi)_\Sigma$ for $j \in [k-1]$ form a basis of the fiber of \mathcal{W} over Σ . To see this, observe first that the forms ξ_j and ξ_{ij} are $\text{PSL}_2(\mathbb{R})$ invariant and conjugation invariant. It is immediate that for all j , σ_j is nowhere vanishing on $\mathcal{M}_{0,k,\{a_1,\dots,a_l\}}^{\frac{1}{r},\hat{\pi}}$. As the number of those sections is $k-1$, it is enough to show

that they are linearly independent. But this is clear, since $(\sigma_j)_\Sigma$ has poles only at $x_{\pi(1)}$ and $x_{\pi(1+j)}$, and by calculating coefficients of poles (e.g., at $x_{\pi(i)}$ for $i \in \{2, \dots, k\}$), we see that if

$$\sum c_j (\sigma_j)_\Sigma = 0,$$

then $c_j = 0$ for all $j \in [k-1]$. Thus, $\mathcal{W} \rightarrow \mathcal{M}_{0,k,\{a_1, \dots, a_l\}}^{\frac{1}{r}, \hat{r}}$ is orientable.

Remark 5.7. Let $(\Sigma, S, [w])$ be a graded spin disk as above and $(\Sigma', S', [w']) = (\Sigma, S, [-w])$ the same disk with opposite grading. The two disks represent the same moduli point, and we would expect that the orientation described above will be the same, whether we calculate it at the fiber of Σ or of Σ' . As a sanity check we shall verify it directly. Recall that the isomorphism α between $(\Sigma, S, [w])$ and $(\Sigma', S', [w'])$ is realized as the identity on the surface and acts by negation on the bundle and grading. The induced map on the fibers of the Witten bundles $H^0(J) \cong H^0(J')$ is given by sending a section $s' \in H^0(J')$ to $\alpha^*(s') = -s'$. Since the grading is also flipped, it is evident that $\alpha^*(\sigma'_j) = \sigma_j$, where σ'_j is defined via Construction 5.6 applied to Σ' . Thus, the isomorphism α takes the constructed orientation of the Witten fiber at Σ' to that of the Witten bundle at Σ , as expected (In higher genus, changing the grading of one boundary is not an isomorphism. In that case, the orientation may change. For example, in [28], which deals with the all-genus $r = 2$ case, it is proven that changing the grading of a boundary with m boundary marking changes the canonical orientation of the moduli space by $(-1)^{m-1}$; see also [29, Section 6.2].).

The next case is $l = 0$. Recall from equation (3.3) that $\mathcal{W}_{0,k,\{0\}}$ is the pullback of $\mathcal{W}_{0,k,\emptyset}$ by the forgetful map $\overline{\mathcal{M}}_{0,k,\{0\}} \rightarrow \overline{\mathcal{M}}_{0,k,\emptyset}$. The fiber of this forgetful map is an open disk, and in particular, it is contractible. Thus, $\mathcal{W}_{0,k,\emptyset}$ is orientable exactly if $\mathcal{W}_{0,k,\{0\}}$ is orientable, and the latter is orientable by the 1st case.

The last case is when $l \geq 1$ and $k-1 < \sum a_i$. Let $k' = 1 + \sum a_i$, so that $\mathcal{W}_{0,k',\{a_1, \dots, a_l\}}$ is orientable by the 1st case. Consider the boundary stratum $\overline{\mathcal{M}}_\Gamma^{1/r}$ defined by a graph Γ with two open vertices v_1 and v_2 , in which v_1 has $k' - k$ boundary tails and no internal tails, while v_2 has the rest. Note that the half-edge h_1 of v_1 has $\text{tw}(h_1) = \text{alt}(h_1) = 0$. By Proposition 4.7 and equation (3.3), we have an identification

$$i_\Gamma^* \mathcal{W}_{0,k',\{a_1, \dots, a_l\}} \cong \mathcal{W}_{0,k'-k+1,\emptyset} \boxplus \mathcal{W}_{0,k,\{a_1, \dots, a_l\}}.$$

Hence,

$$\det(\mathcal{W}_{0,k',\{a_1,\dots,a_l\}})|_{\overline{\mathcal{M}}_\Gamma} \simeq \det(\mathcal{W}_{0,k'-k+1,\emptyset}) \boxtimes \det(\mathcal{W}_{0,k,\{a_1,\dots,a_l\}}).$$

As $\mathcal{W}_{0,k',\{a_1,\dots,a_l\}}$ and $\mathcal{W}_{0,k'-k+1,\emptyset}$ are orientable by the 1st two steps, $\mathcal{W}_{0,k,\{a_1,\dots,a_l\}}$ is also orientable.

Regarding the signs of actions of S_B and S_I , consider first the case where $l = 1$. In this situation, $k \equiv 1 + a_1 \pmod{r}$. As noted above, the bundles \mathcal{W}_{0,k,a_1} and $\mathcal{W}_{0,k,\{k-1\}}$ differ by a direct sum of complex line bundles by Lemma 5.4, and this decomposition is invariant under S_B . The claim in this case follows, then, if we prove it for the bundle $\mathcal{W}_{0,k,\{k-1\}}$. Moreover, it suffices to prove the lemma for $h = -1$, so we assume this in what follows.

Observe that, since the bundle is orientable and all components of the moduli space are isomorphic, the claim can be verified for any specific component by fixing any orientation \mathfrak{o} for \mathcal{W} over this component and comparing $\mathfrak{o}_{g \cdot \Sigma}$ to $g^* \mathfrak{o}_\Sigma$, where $g \cdot \Sigma$ is obtained from Σ by renaming markings according to g and Σ is an arbitrary point in the specified component.

Using these comments, fix a generic smooth Σ and write $\Sigma' = g \cdot \Sigma$. Let \mathfrak{o}^π be the orientation determined by the ordered basis $(\sigma_1^\pi, \dots, \sigma_{k-1}^\pi)$. Comparing $\mathfrak{o}_{g \cdot \Sigma}^\pi$ with $g^* \mathfrak{o}_\Sigma^\pi$ is equivalent to calculating the ratio of \mathfrak{o}^π to $\mathfrak{o}^{g \cdot \pi}$ at the same point of the moduli space. To this end, first observe that the residues of $(\sigma_j^\pi)_\Sigma$ at $x_{\pi(1)}$ and $x_{\pi(1+j)}$ are real and of opposite sign. The sign of the residue at $x_{\pi(1)}$ is independent of j , and moreover, it equals the sign of the residue of $(\sigma_j^{\pi'})_\Sigma$ at $x_{\pi'(1)}$ for $\pi' = g \cdot \pi$. Thus, the coordinate change between the ordered bases $(e_1, \dots, e_{k-1}) = (\sigma_1^\pi, \dots, \sigma_{k-1}^\pi)_{g \cdot \Sigma}$, $(e'_1, \dots, e'_{k-1}) = (\sigma_1^{\pi'}, \dots, \sigma_{k-1}^{\pi'})_{g \cdot \Sigma}$ is given by

$$e'_1 = -\alpha_1 e_1 + \beta_1 e_2, \dots, e'_{k-2} = -\alpha_{k-2} e_1 + \beta_{k-2} e_{k-1}, e'_{k-1} = -\alpha_{k-1} e_1,$$

where α_i, β_i are positive numbers depending on Σ . Hence, the induced sign on the orientation is equal to the sign of $\prod (-\alpha_i)$, which is $(-1)^{k-1}$.

The case $l = 0$ follows by the same argument as in the proof of orientability above: we first add an internal point with twist zero, and then we reduce to the case $l = 1$.

Finally, for $l > 1$, we compare the two orientations at a generic point of \mathcal{M}_Γ , where Γ is a graph with two vertices, a closed vertex v_1 with all internal tails and an open vertex v_2 with only boundary tails. By Proposition 4.7, we have $\mu^* \mathcal{W}|_{\mathcal{M}_\Gamma} \cong q^* (\mathcal{W}_{v_1} \boxplus$

\mathcal{W}_{v_2}), and the decomposition is invariant under the actions of S_B and S_I . Now, μ^* and q^* are complex maps and hence preserve signs. Furthermore, \mathcal{W}_{v_1} is canonically oriented as a complex bundle, $g \in S_B$ acts on it trivially, and $g \in S_I$ (which preserves twists) acts on it as a complex isomorphism, hence preserves sign. Thus, the sign induced by g on the orientation of $\mathcal{W}_{0,k,\{a_1,\dots,a_l\}}$ (or, equivalently, on the orientation of $\mathcal{W}|_{\mathcal{M}_\Gamma}$) is the sign it induces on the orientation of \mathcal{W}_{v_2} . It is 1 for $g \in S_I$, and it equals $(-1)^{k-1}$ for $g \in S_B$ by the 1st case.

5.4 Choosing canonical orientations

Notation 5.8. Denote by $\Gamma_{0,k,\{a_1,\dots,a_l\}}$ the connected graded smooth graph with a single vertex v , boundary tails marked by $[k]$, and internal tails marked $[l]$, such that the i th internal tail has twist a_i .

Throughout what follows, we continue with the notation from the proof of Lemma 5.5.

Definition 5.9. Fix a set of twists \vec{a} of size l and an integer $k \geq 0$. Let $\{\sigma^\pi\}$ be a family of orientations, one for each $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{0,B,\vec{a}}^{\hat{\pi},1/r}$, in which π runs over all orderings of B and B runs over all sets of size k . We say that the family $\{\sigma^\pi\}$ is *covariant* if, whenever $f^B : B \rightarrow B'$ and $f^I : I \rightarrow I'$ are bijections such that f^I preserves twists, we have $\sigma_{0,B,\vec{a}}^\pi = F^* \sigma_{0,B',\vec{a}}^{f^B \circ \pi}$, where $F : (\mathcal{W} \rightarrow \overline{\mathcal{M}}_{0,B,\{a_i\}_{i \in I}}^{1/r}) \rightarrow (\mathcal{W} \rightarrow \overline{\mathcal{M}}_{0,B',\{a_i\}_{i \in I'}}^{1/r})$ is the induced map.

Remark 5.10. By the 2nd part of Lemma 5.5, whenever $\{\sigma^\pi\}$ is a covariant family and $g \in S_B$ cyclically satisfies $g(\pi(i)) = \pi(i+h)$ for some integer h , we have the equality $\sigma^\pi = \varepsilon^h \sigma^{g \cdot \pi}$, where $\varepsilon = \varepsilon_k = (-1)^{k-1}$. Thus, associated to any multiset (\vec{a}, k) are exactly two covariant orientations, determined by choosing σ^π for a single π to be any orientation of the Witten bundle of $\overline{\mathcal{M}}_{0,k,\vec{a}}^{1/r,\hat{\pi}}$, and then extending covariantly.

Definition 5.11. Suppose $a_i \leq r-1$ for each i and $k-1 = \sum_{i \in [l]} a_i$. When $k > 1$, for an order π of the boundary marked points, denote by $\sigma_{0,k,\vec{a}}^\pi$ the orientation of $\mathcal{W}_{0,k,\vec{a}}^{1/r,\hat{\pi}}$ defined using the ordered basis $\sigma^\pi = (\sigma_1^\pi, \dots, \sigma_{k-1}^\pi)$ on the smooth locus. For any I of size l and B of size $k = 1 + \sum_{i \in I} a_i$, this uniquely defines a covariant family of orientations $\{\sigma_{0,B,\{a_i\}_{i \in I}}^\pi\}_\pi$. When $k = 1$ and hence all $a_i = 0$, the Witten bundle is zero-dimensional; in this case, define its orientations $\{\sigma_{0,\{b\},\{0\}_{i \in I}}\}$ to be the positive orientations.

In case $l = 0$, \vec{a} is the empty vector, and $k = r+1$, Remark 5.10 implies that there are exactly two covariant families, induced by the two choices of orientation of

the Witten bundle on the space $\overline{\mathcal{M}}_{0,B,\emptyset}^{\frac{1}{r}}$ with $|B| = r + 1$. Fix one of these orientations, to be chosen later, and denote it by σ_{r+1} .

Now, let $l, k \geq 0$, and fix $a_1, \dots, a_l \in \{0, \dots, r - 1\}$ with $\overline{\mathcal{M}}_{0,k,\{a_1, \dots, a_l\}}^{\frac{1}{r}} \neq \emptyset$. Suppose we have a covariant family of orientations with twists $\{a_1, \dots, a_l\}$ and B of size k . We now show how to induce, given σ_{r+1} as above, a covariant family of orientations with the same twists but for B' of size $k' := k \pm r$, assuming k' is non-negative.

Suppose, first, that $k' = k - r$. Let B' be a set of size $k - r$, and let π' be an order of B' . Let B be a set of size k containing B' , and let π be an order of B that extends π' , such that the 1st $k - r$ element of B with respect to π are those of B' .

Let Γ be a graph with two open vertices and no closed vertices, for which there is one vertex v_0 that contains all the internal tails, all the boundary tails marked by B' , and one half-edge h_0 . The other vertex, v_1 , contains the remaining r boundary tails marked by $B \setminus B'$ and the half-edge $\sigma_1 h_0$. By construction, $\text{tw}(h_0) = \text{alt}(h_0) = 0$. Observe that in $\mathcal{M}_{\Gamma}^{1/r, \hat{\pi}}$, the boundary marked points on the component corresponding to v_0 are cyclically ordered by $\hat{\pi}'$. The Witten bundle \mathcal{W}_{Γ} on $\overline{\mathcal{M}}_{\Gamma}^{1/r}$ can be identified with $\mathcal{W}_{v_0} \boxplus \mathcal{W}_{v_1}$, so we have an identification

$$\det(\mathcal{W}_{\Gamma}) = \det(\mathcal{W}_{v_0}) \boxtimes \det(\mathcal{W}_{v_1}).$$

Define an orientation $\text{ind}_{\Gamma, \pi \rightarrow \pi'} \sigma^{\pi}$ on $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{0, B', \{a_i\}_{i \in [l]}}^{\frac{1}{r}}$ as the unique orientation satisfying

$$\sigma^{\pi}|_{\mathcal{W}_{\overline{\mathcal{M}}_{\Gamma}^{1/r, \hat{\pi}}}} = (\text{ind}_{\Gamma, \pi \rightarrow \pi'} \sigma^{\pi}) \boxtimes \sigma_{r+1}^{\pi_1},$$

where π_1 is the order on the half edges of v_1 induced from π , starting from $\sigma_1 h_0$.

This procedure defines $\sigma^{\pi'}$ uniquely from σ^{π} . The construction is easily seen to be independent of choices and yields a covariant family. Moreover, inverting the steps allows us to define σ^{π} uniquely from $\sigma^{\pi'}$, and if the latter comes from a covariant family, so will the former. Thus, the case $k' = k + r$ is also treated.

Therefore, given twists $\{a_i\}_{i \in I}$, we can uniquely define a covariant family of orientations σ^{π} for any set B such that $\overline{\mathcal{M}}_{0, B, \vec{a}}^{1/r} \neq \emptyset$ by inducing the orientations iteratively, starting from the covariant family of Definition 5.11 (the case $k = \sum a_i + 1$) or, when $I = \emptyset$, from σ_{r+1} .

Definition 5.12. Given \mathfrak{o}_{r+1} and the orientations of Definition 5.11, define $\{\mathfrak{o}^\pi\}$ as the unique covariant family of orientations induced by the above procedure from the family $\{\mathfrak{o}_{0,B,\{a_i\}_{i \in I}}^\pi\}_\pi$, defined when $k - 1 = \sum a_i$.

Observations 5.13. If $I \subseteq I'$ and $a_i = 0$ for all $i \in I'$, then $\mathfrak{o}_{0,B,\{a_i\}_{i \in I'}}^\pi = \text{For}_{I' \setminus I}^* \mathfrak{o}_{0,B,\{a_i\}_{i \in I}}^\pi$.

Indeed, the case $|B| - 1 = \sum a_i$ follows from the definitions, while the general case follows easily by noting that the forgetful map behaves well with respect to the induction procedure.

5.5 Properties of the orientations

The family of orientations in Definition 5.12 respects the decomposition properties of the Witten bundle.

To make this precise, we first state two lemmas, whose proofs are postponed to the next section. We first fix decompositions $I = I_1 \sqcup I_2$, $B = B_1 \sqcup B_2$, and $k = k_1 + k_2$ such that

$$k_1 - 1 \equiv \sum_{i \in I_1} a_i \pmod{r}, \quad k_2 \equiv \sum_{i \in I_2} a_i \pmod{r}.$$

Let Γ be a graph with two open vertices, v_1 and v_2 , connected by an edge e , in which the vertex v_i has internal tails labeled by I_i and k_i boundary tails labeled by B_i . If h_i are the half edges of v_i , then a simple calculation shows that

$$\text{tw}(h_1) = \text{alt}(h_1) = 0.$$

Let π be an order in which the elements of B_1 are consecutive and come before the elements of B_2 . Consider $\Sigma \in \mathcal{M}_\Gamma^{\hat{\pi}}$, where the normalization of Σ has components Σ_1 and Σ_2 corresponding to v_1 and v_2 , respectively. Let π_1 be the restriction of π to the points of Σ_1 , and let π_2 be the restriction to points of Σ_2 but with the half-node x_{h_2} added as the 1st point.

On $\mathcal{M}_\Gamma^{\hat{\pi}}$, the bundle \mathcal{W}_Γ is again identified with $\mathcal{W}_{v_1} \boxplus \mathcal{W}_{v_2}$, so we have the identification

$$\det(\mathcal{W}_\Gamma) = \det(\mathcal{W}_{v_1}) \boxtimes \det(\mathcal{W}_{v_2}), \quad (5.5)$$

and this is respected by the orientations in Definition 5.12. That is, we have the following statement.

Lemma 5.14. For Σ as above, the orientation σ^π agrees with $\sigma_{0,B_1,I_1}^{\pi_1} \boxtimes \sigma_{0,(h_2) \cup B_2,I_2}^{\pi_2}$ under the isomorphism (5.5).

Similarly, the family of orientations in Definition 5.12 satisfies a decomposition property along internal nodes. To specify this, write

$$m^c(\{a_i\}_{i \in I}) = \frac{\sum a_i - (r - 2) + ((r - 2 - \sum a_i) \text{ (mod } r)')}{r},$$

where $x \text{ (mod } r)'$ is the unique element of $\{-1, 0, \dots, r - 2\}$ congruent to x modulo r .

Now, let Γ be a graph with two vertices, an open vertex v^o and a closed vertex v^c . By Proposition 4.7 and Remark 5.2, the Witten bundle \mathcal{W}_Γ on $\overline{\mathcal{M}}_\Gamma^{1/r}$ satisfies

$$\det(\mathcal{W}_\Gamma) \cong (q \circ \mu^{-1})^* \left(\det(\mathcal{W}_{v^o}) \boxtimes \det(\mathcal{W}_{v^c}) \right), \quad (5.6)$$

and we have the following.

Lemma 5.15. There exists $\delta = \pm 1$, depending only on the choice of σ_{r+1} , such that for any Γ, v^o , and v^c as above and any order π , the orientation $\sigma^\pi|_{\mathcal{M}_\Gamma^{\hat{\pi}}}$ agrees with $\delta^{m^c(\{a_i\}_{i \in I^c})} \sigma_{v^o}^\pi \boxtimes \sigma_{v^c}$ under the isomorphism (5.6), where σ_{v^c} is the canonical complex orientation and I^c are the labels of v^c . Moreover, changing σ_{r+1} to the opposite orientation changes δ to $-\delta$.

Definition 5.16. Define σ_{r+1} to be the unique covariant family of orientations of $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{0,r+1,\emptyset}^{1/r}$ for which the $\delta = 1$ in Lemma 5.15. This induces orientations $\sigma_{0,B,\vec{a}}^\pi$ on the Witten bundle for all B and \vec{a} by Definition 5.12.

We can now complete the proof of Theorem 5.1.

Proof of Theorem 5.1. Let $\tilde{\sigma}_{0,B,\vec{a}}^\pi$ be the orientations on the moduli spaces $\overline{\mathcal{M}}_{0,B,\vec{a}}^{1/r}$, described explicitly in Notation 3.15, and let $\sigma_{0,B,\vec{a}}^\pi$ be the orientations on the Witten

bundle described in Definition 5.16. By Lemma 5.14 and Proposition 3.13, the relative orientation

$$o_{0,B,\{a_i\}_{i \in I}} = \tilde{o}_{0,B,\{a_i\}_{i \in I}}^\pi \otimes o_{0,B,\{a_i\}_{i \in I}}^\pi \quad (5.7)$$

for the Witten bundle on $\overline{\mathcal{M}}_{0,B,\{a_i\}_{i \in I}}^{\frac{1}{r},\hat{\pi}}$ is independent of the choice of π . The same argument, together with Remark 5.10, shows the invariance property.

For the 2nd item, first note that, under the notation of Lemma 5.14 and using item (2) of Proposition 3.13, we may write

$$o_{v_1}^{\pi'_1} = o_{h_1} \boxtimes o_{v'_1}^{\pi_1},$$

where π'_1 is the extension of π_1 defined by writing h_1 as the last element, and the pullback of $o_{v'_1}^{\pi_1}$ is with respect to the map that forgets the half-node n_{h_1} . Note also that

$$\dim_{\mathbb{R}} \overline{\mathcal{M}}_{v'_1}^{1/r} \equiv |B_1| - 1 \pmod{2}, \quad \text{rank}_{\mathbb{R}} \mathcal{W}_{v_2} \equiv |B_2| \pmod{2}.$$

From here, the 2nd item is a consequence of Lemma 5.14 and Lemma 3.16, where the sign $(-1)^{(|B_1|-1)|B_2|}$ disappears when commuting $\tilde{o}_{v'_2}^{\pi_2}$ with $o_{v'_1}^{\pi_1}$.

The last item is a direct consequence of Lemmas 5.15 and 3.16. ■

Lemma 5.17. The properties of Theorem 5.1 characterize precisely two families of orientations: the family $\{o_{0,B,\{a_i\}_{i \in I}}\}_{B,\{a_i\}_{i \in I}}$ and the family $\{(-1)^{|B|-1} o_{0,B,\{a_i\}_{i \in I}}\}_{B,\{a_i\}_{i \in I}}$.

Proof. Suppose that $\{o'_{0,B,\{a_i\}_{i \in I}}\}_{B,\{a_i\}_{i \in I}}$ is a different family of orientations satisfying the requirements of Theorem 5.1. Let $\delta_{B,\vec{a}_i} \in \{\pm 1\}$ be the ratio of $o'_{0,B,\{a_i\}_{i \in I}}$ to $o_{0,B,\{a_i\}_{i \in I}}$. Then item (i) shows that $\delta_{B,\vec{a}_i} = \delta_{|B|,\vec{a}_i}$. Item (iii) shows that $\delta_{|B|,\vec{a}_i} = \delta_{|B|,\{\sum a_i \pmod{r}\}}$. Since $\sum a_i = |B| - 1 \pmod{r}$, we denote $\delta_{|B|,\{\sum a_i \pmod{r}\}}$ by $\delta_{|B|}$. Finally, item (ii) shows that $\delta_{a+b} = \delta_{a+1}\delta_b$. Thus, $\delta_1 = 0$ and $\delta_k = \delta_2^{k-1}$, where $\delta_2 \in \pm 1$. The claim follows. ■

Remark 5.18. This ambiguity from Lemma 5.17 is killed by specifying the orientation of Witten's bundle for a single real one-dimensional moduli space: the additional requirement is that if we orient $\overline{\mathcal{M}}_{0,2,\{1\}}^{\hat{\pi}}$ for $\pi = (1, 2)$ by \tilde{o}^π , then the bundle $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{0,2,\{1\}}^{\hat{\pi}}$ is oriented so that sections that are positive with respect to the grading on the arc from x_1 to x_2 .

5.6 Proof of Lemmas 5.14 and 5.15

We now return to the proofs of the two lemmas from the previous section.

Proof of Lemma 5.14. We first treat the case

$$k_1 - 1 = \sum_{i \in I_1} a_i, \quad k_2 = \sum_{i \in I_2} a_i.$$

For convenience, assume that $B = [k]$, that $B_1 = [k_1]$, and that π is the standard order. We show that if $j < k_1$, then up to rescaling by a positive function, the j th basis element of σ^π converges to $\sigma_{0,k_1,\{a_i\}_{i \in I_1};j}^{\pi_1}$ as one approaches a point in $\mathcal{M}_\Gamma^{1/r,\hat{\pi}}$, and that if $j \geq k_1$, it converges to $\sigma_{0,1+k_2,\{a_i\}_{i \in I_2};j-k_1+1}^{\pi_1}$. This verification implies the lemma.

We assume that $k_1, k_2 > 0$, since otherwise, the result is straightforward. Recall that a vector $u \in (\mathcal{W}_{0,k,\vec{a}})_\Sigma$ for $\Sigma \in \mathcal{M}_\Gamma^{1/r,\hat{\pi}}$ can be written as $u = u_1 \boxplus u_2$ with $u_i \in (\mathcal{W}_{V_i})_{\Sigma_i}$. To calculate a coordinate expression for u_i , let $\{\Sigma_t\}_{t \in (0,1/2)}$ be a path in $\mathcal{M}_{0,k,\{a_1,\dots,a_l\}}^{\frac{1}{r}}$ such that $\lim_{t \rightarrow 0} \Sigma_t = \Sigma$. One can model $(\Sigma_t)_{t > 0}$ on the upper half plane, preserving the complex orientation, such that all markings of Σ_{3-i} tend to 0 as $t \rightarrow 0$ but all markings of Σ_i tend to finite, nonzero limits. The resulting marked upper half plane is a model for Σ_i in which x_{h_i} is mapped to the origin. If the vectors $u_t \in \mathcal{W}_{\Sigma_t}$ converge to u , then their expressions in the coordinates induced from the upper half-plane model converge to the coordinate expression for u_i . Moreover, as $t \rightarrow 0$, the ratio between any two markings going to zero is bounded away from zero, since otherwise, $\Sigma \in \overline{\mathcal{M}}_\Gamma^{1/r} \setminus \mathcal{M}_\Gamma^{1/r}$.

Suppose, first, that $j \geq k_1$, so that $1+j$ is a marked point of Σ_2 . Recall that

$$\sigma_j^\pi = \left((-1)^{r+1} \prod_{i \in [l]} \xi_i^{a_i} \prod_{i \in [k]} \xi_{i(i+1)}^{-1} \xi_{1(1+j)}^r \right)^{\frac{1}{r}}$$

and we have

$$\sigma_{1+j-k_1}^{\pi_2} = \left((-1)^{r+1} \prod_{i \in I_2} \xi_i^{a_i} \prod_{i \in [k_2+1]} (\xi_{i(i+1)}^{\pi_2})^{-1} (\xi_{1(2+j-k_1)}^{\pi_2})^r \right)^{\frac{1}{r}},$$

where $x_{\pi_2(1)} = x_{h_2}$. Then, by the discussion above,

$$\left(\frac{\prod_{h \in [k_1-1]} (x_{\pi(h+1)} - x_{\pi(h)})}{\prod_{h \in I_1} (\sqrt{-1}(\bar{z}_h - z_h))^{a_h}} \right)^{\frac{1}{r}} (\sigma_j^\pi)_{\Sigma_t} \rightarrow \sigma_j^{\pi_2}.$$

The function in the above by which we multiply $(\sigma_j^\pi)_{\Sigma_t}$ is positive. Moreover, for t close enough to zero, the ratios between any two factors $x_{\pi(h+1)} - x_{\pi(h)}$ and $\bar{z}_h - z_h$ are bounded, hence they tend to zero uniformly. In fact, since $\sum_{h \in I_1} a_h = k_1 - 1$, the expression

$$\frac{\prod_{h \in [k_1-1]} (x_{\pi(h+1)} - x_{\pi(h)})}{\prod_{h \in I_1} (\sqrt{-1}(\bar{z}_h - z_h))^{a_h}}$$

has a nonzero limit. Calculating the same expression but in a gauge fixing for which the points of Σ_2 approach zero as $t \rightarrow 0$, while those of Σ_1 have distinct finite images, the same argument shows that the limiting section, when projected to \mathcal{W}_{V_1} , has zero of order $\frac{1}{r}$.

Suppose, now, that $j < k_1$, so that $1+j$ is a marked point of Σ_1 . Recall that

$$\sigma_j^{\pi_1} = \left((-1)^{r+1} \prod_{i \in I_1} \xi_i^{a_i} \prod_{i \in [k_1]} (\xi_{i(i+1)}^{\pi_1})^{-1} (\xi_{1(1+j)}^{\pi_1})^r \right)^{\frac{1}{r}}.$$

Write x_{h_1} for the half node of Σ_1 . Again, we work with the upper half-plane model for Σ_1 , where x_{h_1} is mapped to the origin and the orientation of the image of $\partial \Sigma_1$ agrees with the standard real orientation of \mathbb{R} . Consider again a path of smooth surfaces Σ_t , where $t \in (0, \frac{1}{2})$, that converges to Σ . Choose $i_j \in I_j$ and write

$$C_{i_1 i_2} = -\frac{(\bar{z}_{i_1} - z_{i_1})(\bar{z}_{i_2} - z_{i_2})}{(z_{i_1} - z_{i_2})(\bar{z}_{i_1} - \bar{z}_{i_2})}.$$

This is a well-defined, positive function on $\mathcal{M}_{0,k, \{a_1, \dots, a_l\}}^{\frac{1}{r}}$, so it has a positive r th root; furthermore, it vanishes at \mathcal{M}_Γ . The same considerations as above reveal that the limit of $\frac{\sigma_j^\pi(\Sigma_t)}{C_{i_1 i_2}^{1/r}}$, when projected on \mathcal{W}_{V_1} , is nonzero, but its projection on \mathcal{W}_{V_2} vanishes to order $1 - 1/r$. Moreover, the projection of the limit on \mathcal{W}_{V_1} has the zero profile of $\sigma_j^{\pi_1}$, hence they agree up to multiplication by a real function, by degree reasons and involution invariance. This function is positive, since both sections are positive on the arc from x_1 to x_{1+j} .

We now turn to the general case. Write

$$m_1 = \frac{k_1 - 1 - \sum_{i \in I_1} a_i}{r}, \quad m_2 = \frac{k_2 - \sum_{i \in I_2} a_i}{r}.$$

The proof is by induction on $m = |m_1| + |m_2|$. The case $m = 0$ has been treated above. Note that, perhaps by applying Observation 5.13, we may assume I_1 and I_2 are nonempty. Suppose that the claim has been proven for $m - 1$, and assume $m_2 < 0$.

Let $B'_2 \supseteq B_2$ be a set of size $k'_2 = |B_2| + r$, and write $B' = B_1 \cup B'_2$. Consider the graph Γ' obtained by attaching to Γ an open vertex, connected to v_2 by an edge whose half-edge h in v_2 has $\text{tw}(h) = \text{alt}(h) = 0$, and that has r boundary tails labeled by $B' \setminus B$. Let π' be an order on B' extending π , such that the last elements are those of $B \setminus B'$ and are ordered so that tails belonging to the same open vertex are labeled consecutively. Let Γ_2 be the component of v_2 in $\text{detach}_e(\Gamma')$. By the construction of the induced orientation, we have

$$\mathfrak{o}^{\pi'}|_{\overline{\mathcal{M}}_{\Gamma'}^{\hat{\pi}'}} = \mathfrak{o}^{\pi}|_{\overline{\mathcal{M}}_{\Gamma}^{\hat{\pi}}} \boxtimes \mathfrak{o}'_{r+1}, \quad (5.8)$$

where \mathfrak{o}'_{r+1} is the orientation for the Witten bundle of the new component with respect to the order induced from π' after putting the node as the 1st element. On the other hand, by the induction assumption, we have

$$\mathfrak{o}^{\pi'}|_{\overline{\mathcal{M}}_{\Gamma'}^{\hat{\pi}'}} = \mathfrak{o}_{0, B_1, I_1}^{\pi_1} \boxtimes \mathfrak{o}_{0, \{h_2\} \cup B'_2, I_2}^{\pi'_2}, \quad (5.9)$$

where π'_2 is the restriction of π_2 to B'_2 . By the construction of the induced orientation, this time with respect to $\overline{\mathcal{M}}_{0, \{h_2\} \cup B'_2, \{a_i\}_{i \in I_2}}^{1/r, \pi'_2}$, we have

$$\mathfrak{o}^{\pi'_2}|_{\overline{\mathcal{M}}_{\Gamma_2}^{\hat{\pi}'_2}} = \mathfrak{o}^{\pi_2} \boxtimes \mathfrak{o}'_{r+1}. \quad (5.10)$$

Putting these observations together, we see that

$$\mathfrak{o}^{\pi}|_{\overline{\mathcal{M}}_{\Gamma}^{\hat{\pi}}} = \mathfrak{o}_{0, B_1, I_1}^{\pi_1} \boxtimes \mathfrak{o}_{0, \{h_2\} \cup B_2, I_2}^{\pi_2}. \quad (5.11)$$

The case $m_2 > 0$ is treated similarly to the above, so we omit it. The remaining case is $m_2 = 0$. In this case, $m_1 \neq 0$, and the proof is similar, so we merely remark on the changes. First, we work with the graph Γ' obtained from Γ by attaching a new vertex with r boundary tails, no internal tails, and one legal half edge of twist $r - 2$ to v_1 . We choose the order π' so that boundary tails of v_1 come first and those of v_2 come last. If we let e_1 be the edge of v_2 and e_2 the edge of the new vertex, then we compare the orientation expressions for $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{\Gamma'}^{\pi'}$ obtained in two ways. First, we induce orientation

from $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{d_{e_1, e_2}}^{\hat{\pi}'} \Gamma'$ to $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{d_{e_1}}^{\hat{\pi}'} \Gamma'$ via Ind, and then to $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{\Gamma'}^{\hat{\pi}'}$, to obtain (using the above notation)

$$\mathfrak{o}^{\pi'}|_{\overline{\mathcal{M}}_{\Gamma'}^{\hat{\pi}'}} = \epsilon_1 (-1)^{rk_2} \mathfrak{o}^{\pi_1} \boxtimes \mathfrak{o}^{\pi_2} \boxtimes \mathfrak{o}'_{r+1}.$$

Here, ϵ_1 is the sign appearing from inducing the orientation from $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{d_{e_1}}^{\hat{\pi}'} \Gamma'$ to $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{\Gamma'}^{\hat{\pi}'}$ and $(-1)^{rk_2}$ comes from the fact that, in π' , the tails of the new vertex are not last, so we have to perform a shift in π' in order to calculate the sign induced using Ind. Using Lemma 5.5, we see that the total added sign is $(-1)^{rk_2}$. Next, we induce orientation from $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{d_{e_1, e_2}}^{\hat{\pi}'} \Gamma'$ to $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{d_{e_2}}^{\hat{\pi}'} \Gamma'$, and then, via Ind, to $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{\Gamma'}^{\hat{\pi}'}$. The result is

$$\mathfrak{o}^{\pi'}|_{\overline{\mathcal{M}}_{\Gamma'}^{\hat{\pi}'}} = \epsilon_2 \mathfrak{o}^{\pi_1} \boxtimes \mathfrak{o}'_{r+1} \boxtimes \mathfrak{o}^{\pi_2} = \epsilon_2 (-1)^{rk_2} \mathfrak{o}^{\pi_1} \boxtimes \mathfrak{o}^{\pi_2} \boxtimes \mathfrak{o}'_{r+1},$$

in which ϵ_2 is the sign that appears from inducing the orientation to $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{d_{e_2}}^{\hat{\pi}'} \Gamma'$ and the sign $(-1)^{rk_2}$ comes from changing the order of \mathfrak{o}^{π_2} and \mathfrak{o}'_{r+1} . One of ϵ_1 or ϵ_2 is determined by induction to be 1, so the other is also 1. The case $m_2 = 0$ is thus proven. \blacksquare

Proof of Lemma 5.15. Write

$$\mathfrak{o}^{\pi}|_{\overline{\mathcal{M}}_{\Gamma}^{\hat{\pi}}} = \epsilon_{\{a_i\}_{i \in I^C}, \{a_i\}_{i \in I^O}, B}^{\pi} \mathfrak{o}_{V^O}^{\pi} \boxtimes \mathfrak{o}_{V^C},$$

where I^O are the labels in the open part and I^C in the closed part. By covariance, $\epsilon_{\{a_i\}_{i \in I^C}, \{a_i\}_{i \in I^O}, B}^{\pi}$ is independent of π and can be written as $\epsilon_{\{a_i\}_{i \in I^C}, \{a_i\}_{i \in I^O}, |B|}$. We prove the lemma by showing the following:

- (1) $\epsilon_{\{a_i\}_{i \in I^C}, \{a_i\}_{i \in I^O}, B} = \epsilon_{\{a_i\}_{i \in I^C}}$, meaning that it depends only on $\{a_i\}_{i \in I^C}$.
- (2) If $I^C = I_1 \cup I_2$ where neither I_1 nor I_2 is empty and $|I_1| \geq 2$, then

$$\epsilon_{\{a_i\}_{i \in I^C}} = \epsilon_{\{a_i\}_{i \in I_1}} \epsilon_{\{a_i\}_{i \in I_2} \cup \{\sum_{i \in I_1} a_i \bmod r\}}.$$

Thus, ϵ is fully determined by its value on pairs of elements.

- (3) If $a + b < r$, then $\epsilon_{\{a, b\}} = 1$.
- (4) $\epsilon_{\{a, b + c \pmod r\}} \epsilon_{\{b, c\}} = \epsilon_{\{a + b \pmod r, c\}} \epsilon_{\{a, b\}}$, and when $a < r - 1$ but $a + b \geq r$

$$\epsilon_{\{a, b\}} = \epsilon_{\{1, a + b - r\}} \epsilon_{\{a, b\}} = \epsilon_{\{1 + a, b\}} \epsilon_{\{1, a\}} = \epsilon_{\{1 + a, b\}},$$

so that for $r \leq a + b \leq 2r - 2$, $\epsilon_{\{a_i, b\}} = \delta \in \{\pm 1\}$ is a constant.

(5) $\epsilon_{\{a_i\}_{i \in I^C}} = \delta^{m^c(\{a_i\}_{i \in I^C})}$.

For the 1st item, it is enough to show that $\epsilon_{\{a_i\}_{i \in I^C}, \{a_i\}_{i \in I^O}, B} = \epsilon_{\{a_i\}_{i \in I^C}, \{a_i\}_{i \in I^O}, B'}$ where $B \subseteq B'$ and $I^O \subseteq I'^O$. To show this, consider the graph Γ' consisting of two open vertices, v^o and v' , and a closed vertex, where the open vertices are connected by an edge e' whose half-edge h in v^o has $\text{tw}(h) = \text{alt}(h) = 0$, and the component of v^o in $\text{detach}_{e'}(\Gamma')$ is Γ . Denote the internal edge of Γ (and of Γ') by e . Let π' be an order extending π to B' such that the 1st elements are those of B . By abuse of notation, denote also by π' the restriction of π to $B' \setminus B$. We calculate the orientation of $\mathcal{W}|_{\mathcal{M}_{\Gamma'}^{\pi'}}$ in two ways. First, by the definition of ϵ applied to the moduli space for the graph $d_{e'}\Gamma$ obtained by smoothing the edge e' of Γ' , we have

$$\mathfrak{o}^{\pi'}|_{\mathcal{M}_{d_{e'}\Gamma'}^{\pi'}} = \epsilon_{\{a_i\}_{i \in I^C}, \{a_i\}_{i \in I^O}, B'} \mathfrak{o}_{0, B', \{a_i\}_{i \in I'^O} \cup \{\sum_{i \in I^C} a_i \pmod{r}\}}^{\pi'} \boxtimes \mathfrak{o}_{v^c}.$$

Applying Lemma 5.14 to $\overline{\mathcal{M}}_{0, B', \{a_i\}_{i \in I'^O} \cup \{\sum_{i \in I^C} a_i \pmod{r}\}}^{\frac{1}{r}}$ gives

$$\mathfrak{o}^{\pi'}|_{\mathcal{M}_{\Gamma'}^{\pi'}} = \epsilon_{\{a_i\}_{i \in I^C}, \{a_i\}_{i \in I^O}, B'} \mathfrak{o}_{v^o}^{\pi} \boxtimes \mathfrak{o}_{v'}^{\pi'} \boxtimes \mathfrak{o}_{v^c}.$$

On the other hand, by Lemma 5.14 applied to $\overline{\mathcal{M}}_{0, B', \{a_i\}_{i \in I^C \cup I'^O}}^{\frac{1}{r}}$, we have

$$\mathfrak{o}^{\pi'}|_{\mathcal{M}_{d_{e'}\Gamma'}^{\pi'}} = \mathfrak{o}_{0, B, \{a_i\}_{i \in I^C \cup I^O}}^{\pi} \boxtimes \mathfrak{o}_{v'}^{\pi'}.$$

The claim now follows since, by the definition of ϵ , we have

$$\mathfrak{o}^{\pi'}|_{\mathcal{M}_{\Gamma'}^{\pi'}} = \epsilon_{\{a_i\}_{i \in I^C}, \{a_i\}_{i \in I^O}, B} \mathfrak{o}_{v^o}^{\pi} \boxtimes \mathfrak{o}_{v^c} \boxtimes \mathfrak{o}_{v'}^{\pi'}.$$

For the 2nd item, first note that if Λ is a connected, closed r -spin graph consisting of two vertices v_1 and v_2 and an edge between them, then

$$\mathfrak{o}^{\pi}|_{\mathcal{M}_{\Lambda}} = \mathfrak{o}_{v_1} \boxtimes \mathfrak{o}_{v_2},$$

where all the orientations in the equation are the canonical complex orientations. Consider a graph Γ' with two internal closed vertices v_1 and v_2 and an open vertex

v^o , where v^o is connected to v_2 by e_2 and v_2 is connected to v_1 by e_1 . We again calculate the orientation of $\mathcal{W} \rightarrow \mathcal{M}_{\Gamma'}$ in two ways. First, by the definition of ϵ ,

$$\mathfrak{o}^\pi|_{\mathcal{M}_{de_1\Gamma}^{\hat{\pi}}} = \epsilon_{\{a_i\}_{i \in I^C}} \mathfrak{o}_{v^o}^\pi \boxtimes \mathfrak{o}_{v^c},$$

where v^c is the closed vertex of $d_{e_1}\Gamma$ and I^C are the labels of its tails. Then

$$\mathfrak{o}^\pi|_{\mathcal{M}_{\Gamma}^{\hat{\pi}}} = \epsilon_{\{a_i\}_{i \in I^C}} \mathfrak{o}_{v^o}^\pi \boxtimes \mathfrak{o}_{v_1} \boxtimes \mathfrak{o}_{v_2}.$$

On the other hand, by the definition of ϵ ,

$$\mathfrak{o}^\pi|_{\mathcal{M}_{de_2\Gamma}^{\hat{\pi}}} = \epsilon_{\{a_i\}_{i \in I_1}} \boxtimes \mathfrak{o}_{v^o}^\pi \boxtimes \mathfrak{o}_{v_1},$$

where v^o is the open vertex of $d_{e_2}\Gamma$ and I_i are the labels of the tails on v_i . Again by the definition of ϵ ,

$$\mathfrak{o}^\pi|_{\mathcal{M}_{\Gamma}^{\hat{\pi}}} = \epsilon_{\{a_i\}_{i \in I_1}} \epsilon_{\{a_i\}_{i \in I_2} \cup \{\sum_{i \in I_1} a_i \pmod r\}} \mathfrak{o}_{v^o}^\pi \boxtimes \mathfrak{o}_{v_1} \boxtimes \mathfrak{o}_{v_2},$$

as claimed.

For the 3rd item, we use Observation 4.9. Let $I = [l]$ and $B = [k]$ and assume that $k - 1 = \sum_{i \in [l]} a_i$. Suppose $I' \subseteq [l]$ is such that $\sum_{i \in I'} a_i \leq r - 1$, and let Γ be the graph with a closed vertex v^c containing exactly the tails labeled I' and an internal edge to the open vertex v^o with internal tails labeled by $[l] \setminus I'$ and boundary tails labeled $[k]$. We claim, in this situation, that the two orientations $\mathfrak{o}^\pi|_{\mathcal{M}_{\Gamma}^{\hat{\pi}}}$ and $\mathfrak{o}_{0,k,\{a_i\}_{i \in [l] \setminus I'} \cup \{\sum a_i\}_{i \in I'}}^\pi$ on $\mathcal{W} \rightarrow \overline{\mathcal{M}}_{\Gamma}^\pi$ agree (where for convenience we omit the pullback maps from the notation).

To prove this claim, write $a = \sum_{i \in I'} a_i$, and let $\sigma_j = \sigma_{0,k,\{a_i\}_{i \in [l]};j}^\pi$ and $\sigma'_j = \sigma_{0,k,\{a_i\}_{i \in [l] \setminus I'} \cup \{a\};j}^\pi$. Then it is a direct computation to verify that, when $\Sigma' \rightarrow \Sigma \in \mathcal{M}_{\Gamma}^\pi$, the section $(\sigma_j)_{\Sigma'}$ converges to a section $(\hat{\sigma}_j)_{\Sigma}$ of \mathcal{W}_{Σ} that is the pullback of σ'_j . Indeed, denoting by z_{h^o} the half node in the disk component of Σ , the projection to \mathcal{W}_{v^o} of the limit of $(\sigma_j)_{\Sigma'}$ is

$$\left(\left(\frac{\sqrt{-1}(\bar{z}_{h^o} - z_{h^o})dw}{(w - z_{h^o})(w - \bar{z}_{h^o})} \right)^a \prod_{h \in [l] \setminus I'} \xi_h^{a_h/r} \prod_{h \in [k]} \xi_{h(h+1)}^{-1} \xi_{1(1+j)}^r \right)^{\frac{1}{r}},$$

as claimed. Thanks to the 1st item, this claim implies the 3rd item.

The 4th item is a direct consequence of the 2nd item, by partitioning the set $\{a, b, c\}$ in two different ways. Its 2nd part uses the 3rd item, applied twice to $\{1, a, b\}$.

The last item follows by induction on $|I^C| \geq 2$. The 3rd and 4th items serve as the base case. Suppose, then, that we have shown the claim for $|I^C| = n$, and let $|I^C| = n + 1$. Write $I^C = I_1 \cup I_2$ with $I_2 = \{a, b\}$, and write $c = a + b \pmod{r}$. Then

$$\epsilon_{\{a_i\}_{i \in I^C}} = \epsilon_{\{a_i\}_{i \in I_1} \cup \{c\}} \epsilon_{a, b} = (-1)^p,$$

where the power p is given by

$$p = \frac{c + \sum a_i - (r - 2) + (r - 2 - c - \sum_{i \in I_1} a_i \pmod{r})'}{r} + \frac{a + b - (r - 2) + (r - 2 - a - b) \pmod{r}'}{r}.$$

The induction now follows from the definition of m^c and the two equations

$$r - 2 - c - \sum_{i \in I_1} a_i \pmod{r}' = r - 2 - \sum_{i \in I^C} a_i \pmod{r}',$$

$$c + (r - 2 - a - b) \pmod{r}' = r - 2.$$

The “moreover” statement of the lemma is straightforward from the definitions. Indeed, when $k = 1 + \sum a_i$, the orientation is defined without σ_{r+1} , so it does not change when σ_{r+1} changes. For general k' , the construction of orientations in Definition 5.12 uses the map $\text{ind}_{k \rightarrow k'}$, and it is immediate from the definition of $\text{ind}_{k \rightarrow k'}$ that changing σ_{r+1} changes the orientation by a factor of $(-1)^{\frac{k' - \sum_{i \in I} a_i - 1}{r}}$. The claim now follows from noting that

$$\frac{\sum_{i \in I} a_i - k' + 1}{r} = m^c(\{a_i\}_{i \in I^C}) + \frac{\{\sum_{i \in I^C} a_i \pmod{r}\} + \sum_{i \in I^0} a_i - k' + 1}{r}.$$

■

Funding

The work was supported by the Mathematical Center in Akademgorodok under agreement No. 075-2019-1675 with the Ministry of Science and Higher Education of the Russian Federation (to A.B.); the National Science Foundational Division of Mathematical Sciences, (1810969 to E.C.); the Center for New Scientists of Weizmann Institute (to R.T.); Dr Max Rössler (to R.T.); the Walter Haefner Foundation (to R.T.); the Eidgenössische Technische Hochschule Zürich Foundation (to R.T.); and the Israel Science Foundation (grant 335/19 to R.T.).

Acknowledgments

The authors would like to thank R. Pandharipande, D. Ross, J. Solomon, and E. Witten for interesting discussions related to open r -spin theory. The authors are grateful to J. Guéré for enlightening discussions concerning the Ramond sector and to A. Netser Zernik for his enriching explanations about orbifolds with corners.

References

- [1] Abramovich, D. and A. Vistoli. "Compactifying the space of stable maps." *J. Am. Math. Soc.* 15 (2002): 27–76.
- [2] Buryak, A. "Open intersection numbers and the wave function of the KdV hierarchy." *Mosc. Math. J.* 16 (2016): 27–44.
- [3] Buryak, A., E. Clader, and R. J. Tessler. "Closed extended r -spin theory and the Gelfand–Dickey wave function." *J. Geom. Phys.* 137 (2019): 132–53.
- [4] Buryak, A., E. Clader, and R. J. Tessler. "Open r -spin theory II: the analogue of Witten's conjecture for r -spin disks." (2018): preprint arXiv:1809.02536v4.
- [5] Buryak, A. and R. J. Tessler. "Matrix models and a proof of the open analog of Witten's conjecture." *Comm. Math. Phys.* 353 (2017): 1299–328.
- [6] Chang, H.-L., J. Li, and W.-P. Li. "Witten's top Chern class via cosection localization." *Invent. Math.* 200 (2015): 1015–63.
- [7] Chiodo, A. "The Witten top Chern class via K-theory." *J. Algebraic Geom.* 15 (2006): 681–707.
- [8] Chiodo, A. "Stable twisted curves and their r -spin structures." *Ann. Inst. Fourier (Grenoble)* 58 (2008): 1635–89.
- [9] Chiodo, A. "Towards Global Mirror Symmetry." *Lecture Notes From 2011 Summer School on Moduli of Curves and Gromov–Witten Theory*. University Joseph Fourier (Grenoble 1), 2011.
- [10] Chiodo, A. and Y. Ruan. "Landau–Ginzburg/Calabi–Yau correspondence for quintic threefolds via symplectic transformations." *Invent. Math.* 182 (2010): 117–65.
- [11] Chiodo, A. and D. Zvonkine. "Twisted r -spin potential and Givental's quantization." *Adv. Theor. Math. Phys.* 13 (2009): 1335–69.
- [12] Faber, C., S. Shadrin, and D. Zvonkine. "Tautological relations and the r -spin Witten conjecture." *Ann. Sci. Éc. Norm. Supér. (4)* 43 (2010): 621–58.
- [13] Fan, H., T. J. Jarvis, and Y. Ruan. "The Witten equation, mirror symmetry, and quantum singularity theory." *Ann. Math. (2)* 178 (2013): 1–106.
- [14] Fan, H., T. J. Jarvis, and Y. Ruan. "A mathematical theory of the gauged linear sigma model." *Geom. Topol.* 22 (2018): 235–303.
- [15] Fukaya, K., Y.-G. Oh, H. Ohta, and K. Ono. "Lagrangian Intersection Floer Theory: Anomaly and Obstruction." Part I. Volume 46.1 of *AMS/IP Studies in Advanced Mathematics*. Providence, RI: American Mathematical Society. Somerville, MA: International Press, 2009.
- [16] Jarvis, T. J. "Geometry of the moduli of higher spin curves." *Int. J. Math.* 11 (2000): 637–63.

- [17] Jarvis, T. J., T. Kimura, and A. Vaintrob. "Gravitational Descendants and the Moduli Space of Higher Spin Curves." *Advances in Algebraic Geometry Motivated by Physics (Lowell, MA, 2000)*, vol. 276 of Contemp. Math. 167–77. Providence, RI: Amer. Math. Soc., 2001.
- [18] Jarvis, T. J., T. Kimura, and A. Vaintrob. "Moduli spaces of higher spin curves and integrable hierarchies." *Compos. Math.* 126 (2001): 157–212.
- [19] Joyce, D. "On Manifolds With Corners." *Advances in Geometric Analysis*, vol. 21 of Adv. Lect. Math. (ALM). 225–58. Somerville, MA: Int. Press, 2012.
- [20] Liu, C.-C. M. "Moduli of J -holomorphic curves with Lagrangian boundary conditions and open Gromov–Witten invariants for an S^1 -equivariant pair." (2020): preprint arXiv:math/0210257.
- [21] Mochizuki, T. "The virtual class of the moduli stack of stable r -spin curves." *Comm. Math. Phys.* 264 (2006): 1–40.
- [22] Netser Zernik, A. "Moduli of open stable maps to a homogeneous space." (2017): preprint arXiv:1709.07402.
- [23] Pandharipande, R., A. Pixton, and D. Zvonkine. "Relations on $\overline{\mathcal{M}}_{g,n}$ via 3-spin structures." *J. Am. Math. Soc.* 28 (2015): 279–309.
- [24] Pandharipande, R., J. P. Solomon, and R. J. Tessler. "Intersection theory on moduli of disks, open KdV and Virasoro." (2014): preprint arXiv:1409.2191.
- [25] Polishchuk, A. and A. Vaintrob. "Algebraic Construction of Witten's Top Chern Class." *Advances in Algebraic Geometry Motivated by Physics (Lowell, MA, 2000)*. vol. 276 of Contemp. Math. 229–49. Providence, RI: Amer. Math. Soc., 2001.
- [26] Romagny, M. "Group actions on stacks and applications." *Michigan Math. J.* 53 (2005): 209–36.
- [27] Solomon, J. P. and R. J. Tessler. "Graded Riemann surfaces with boundary and the open Arf invariant." (2020): In preparation.
- [28] Solomon, J. P. and R. J. Tessler. "Intersection theory on the moduli space of graded Riemann surfaces with boundary." (2020): In preparation.
- [29] Tessler, R. J. "The combinatorial formula for open gravitational descendants." (2015): preprint arXiv:1507.04951.