

RESEARCH



# Problems in arithmetic topology

Claudio Gómez-González\* and Jesse Wolfson

\*Correspondence:  
claudijg@uci.edu  
Department of Mathematics,  
University of California-Irvine,  
Irvine, CA, USA  
JW is supported in part by NSF  
Grant Nos. DMS-1811846 and  
DMS-1944862. Support for the  
workshop was provided by PIMS  
and by NSF Grant No.  
DMS-1856737

## Abstract

We present a list of problems in arithmetic topology posed at the June 2019 PIMS/NSF workshop on “Arithmetic Topology.” Three problem sessions were hosted during the workshop in which participants proposed open questions to the audience and engaged in shared discussions from their own perspectives as working mathematicians across various fields of study. Participants were explicitly asked to provide problems of various levels of difficulty, with the goal of capturing a cross section of exciting challenges in the field that could help guide future activity. The problems, together with references and brief discussions when appropriate, are collected below into three categories: (1) topological analogues of arithmetic phenomena, (2) point counts, stability phenomena and the Grothendieck ring, and (3) tools, methods and examples.

Three problem sessions were hosted during the workshop in which participants proposed open questions to the audience and engaged in shared discussions from their own perspectives as working mathematicians across various fields of study. Participants were explicitly asked to provide problems of various levels of difficulty, with the goal of capturing a cross section of exciting challenges in the field that could help guide future activity. The problems, together with references and brief discussions when appropriate, are collected below into three categories: (1) topological analogues of arithmetic phenomena, (2) point counts, stability phenomena and the Grothendieck ring, and (3) tools, methods and examples.

## 1 Topological analogues

### 1.1 Craig Westerland

This problem concerns the homology of random topological objects and the Cohen–Lenstra distribution, originally introduced to heuristically understand experimental observations about the class groups of number fields [18]. The Cohen–Lenstra distribution on finite (abelian) groups posits that the probability of each group  $G$  being in the support of the distribution is inversely proportional to  $|G|^n |\text{Aut}(G)|^m$  for some  $n, m \geq 0$ . Since its first use, it has appeared throughout number theory and in various models of random integral matrix cokernels; see, for example, [15, 27, 58, 86].

Working one prime  $p$  at a time, one can compute the  $p$ -power torsion and torsion-free part of the integral homology of a given chain complex by tensoring with the  $p$ -adic integers  $\mathbb{Z}_p$  before taking homology. If one further specifies the total homological rank  $r$

of such complexes, the Cohen–Lenstra distribution specifies a discrete measure on the resulting module  $M = \mathbb{Z}_p^r \times N$  (for  $N$  an abelian  $p$ -group). We can then ask the following question:

**Conjecture 1** Let  $X_\bullet$  be a random simplicial complex. Does the torsion  $H_*(X_\bullet)_{tors}$  have a Cohen–Lenstra distribution?

This question was originally posed by Matt Kahle, Frank Lutz, Andrew Newman and Kyle Parsons, along with experimental evidence supporting the conjecture, in [50]. Westerland raised this standing problem and also discussed an analogue for random manifolds where one can introduce a torsion linking pairing

$$\begin{aligned} H^i(X; \mathbb{Z})_{tors} \times H^{\dim X - i - 1}(X; \mathbb{Z})_{tors} &\xrightarrow{\cup} H^{\dim X - 1}(X; \mathbb{Z})_{tors} \\ &\xrightarrow{\delta} H^{\dim X}(X; \mathbb{Q}/\mathbb{Z}) \cong \mathbb{Q}/\mathbb{Z}. \end{aligned}$$

Delaunay [21] introduced heuristics in the context of a bilinear pairing

$$\mu : G \times G \rightarrow \mathbb{Q}/\mathbb{Z},$$

originally in the symplectic context of Tate–Shafarevitch groups but later expanded by others (e.g., [8, 85]) to additional phenomena, where  $\text{Aut}(G)$  is replaced with  $\text{Aut}^\mu(G)$  of automorphisms preserving the pairing.

**Question 2** Are there Delaunay heuristics in the random manifold case?

Andy Putman mentioned useful models for random 3-manifolds (e.g., Dunfield–Thurston manifolds via random classes in the mapping class group  $\text{Mod}_g$ ) that might help approach this problem.

### 1.2 Daniel Litt

Motivated by the Frobenius action on the pro- $p$  Galois group of a curve over  $\mathbb{F}_p$ , consider a random automorphism  $\varphi$  of a free pro- $p$  group  $F$ , and construct a random Weil group

$$W_\varphi := F \rtimes \langle \varphi \rangle.$$

What is the representation theory of  $W_\varphi$ ? Do the predictions from geometric Langlands theory hold? For example:

**Problem 3** Given an irreducible continuous representation  $\rho : W_\varphi \rightarrow \text{GL}_n(\mathbb{F}_p((t)))$ , does the restriction  $\rho|_F$  always have finite image?

For  $F$  the geometric étale fundamental group of a smooth projective curve  $X$  over  $\mathbb{F}_q$ , the analogous problem was posed by de Jong [20] and proven by Gaitsgory [33]. Similarly, one could ask:

**Problem 4** Let  $L/\mathbb{Q}_p$  be a finite extension, and let  $\rho : W_\varphi \rightarrow \text{GL}_n(L)$  be a continuous irreducible representation. For almost every  $\varphi$  (i.e., with probability 1), does  $H_{cts}^1(F; \rho \otimes \rho^\vee)^\varphi = 0$ ?

Viewing  $H_{cts}^1(F; \rho \otimes \rho^\vee)^\varphi$  as the tangent space to  $\rho$  in the character variety of  $F$ , we can understand the previous problem as a “cohomological rigidity” statement. Like the first problem, this problem is a theorem for  $F$  the geometric étale fundamental group of a smooth projective curve  $X$  over  $\mathbb{F}_q$ : indeed, a theorem of L. Lafforgue [59] implies that

$\rho$  is pure of some weight, so  $\rho \otimes \rho^\vee$  is pure of weight 0. Deligne's bounds [22] therefore imply that  $H_{cts}^1(F; \rho \otimes \rho^\vee)$  is pure of weight 1, and hence the invariants of Frobenius are trivial.

More generally, both of the above problems arise from trying to understand in what sense the action of Frobenius on the geometric étale fundamental group of a smooth projective curve over  $\mathbb{F}_q$  is “special”? Affirmative answers to both of the questions above would suggest that it isn't!

## 2 Point counting, stability phenomena and the Grothendieck ring

### 2.1 Akshay Venkatesh

In many interesting number theory problems (e.g., counting number fields), one has not only a main term in the asymptotic count, but a secondary term or more. For example, the number of cubic fields of discriminant up to  $X$  is

$$aX + bX^{5/6} + \text{lower order terms.}$$

We have very little understanding of these lower-order terms, and they are not just of theoretical interest: when one tries to verify the conjectures numerically, one finds that the secondary terms are dominant in the computational range.

So the question, following the framework of the Weil conjectures, is:

**Problem 5** What is the topological meaning of secondary terms appearing in asymptotic counts in number theory?

They do not correspond to stable cohomology classes (these are the main terms), but to some kind of slightly weaker structure, which is still much better behaved than cohomology near middle degree.

### 2.2 Zachary Himes

The recent work of Galatius–Kupers–Randal-Williams [34, 35] develops and applies a homology theory based on  $E_k$ -algebras, originally introduced to study  $k$ -fold loop spaces [64], with the goal of showing results “beyond homological stability” which they call *secondary stability*. For example, in [35],  $E_k$ -homology is used to measure the failure of homological stability itself as a stable phenomenon in the context of mapping class groups, and in [73], this perspective is used to explain the homological stability phenomena which underlie the breakthrough work of Ellenberg–Venkatesh–Westerland [27].

**Question 6** Building off [27], can one prove Hurwitz spaces have stability (for more general groups than the dihedral groups)? Secondary stability, or higher-order stability? What can one say about the homology of Hurwitz spaces after restricting to subsets of connected components?

### 2.3 Daniel Litt

**Problem 7** Fix  $n, d$ , and  $q$ . How many supersingular hypersurfaces  $X \subset \mathbb{P}^n$  exist of degree  $d$  over  $\mathbb{F}_q$ ? In particular, does such a hypersurface exist for every degree, dimension, and characteristic? Moreover, the set of such varieties are the  $\mathbb{F}_q$ -points of some supersingular locus  $V$ —what is the class  $[V]$  in the Grothendieck ring?

A great deal is known about the supersingular locus in particular cases. See, for example, the work of Oort and Li [63] on moduli spaces of supersingular abelian varieties, especially on the case of principally polarized abelian varieties  $S_{g,1}$ .

#### 2.4 Aaron Landesman

Cohen–Lenstra heuristics can be used to describe distributions of class groups of certain global fields. In many ways, the Selmer groups of elliptic curves behave similarly to class groups. As such, Landesman provided two conjectures—a number-theoretic version and a topological version—relating topological techniques to the average size of Selmer groups of elliptic curves over function fields. We describe the latter here, framed as a question of homological stability, which would likely imply the number-theoretic conjecture. For more details, see [60].

**Definition 8** The *moduli space of Weierstrass models* is given by

$$\mathcal{W}_d := \{(a(S, T), b(S, T)) \in \mathbb{C}[S, T]^2 \mid \text{homogeneous degree } 4d \text{ and } 6d, \text{ respectively}\} \subseteq \mathbb{C}^{10d+2}.$$

A point  $W = (a(S, T), b(S, T)) \in \mathcal{W}_d$  gives rise to an elliptic surface  $S_W$  via

$$S_W := \{([S : T], [X : Y : Z]) \in \mathbb{CP}^1 \times \mathbb{CP}^2 \mid Y^2Z = X^3 + a(S, T)XZ^2 + b(S, T)Z^3\},$$

together with the natural projection map  $\varphi_W : S_W \rightarrow \mathbb{CP}^1$ . The *universal family of Weierstrass models*,  $\mathcal{UW}_d$ , is the space parameterizing models  $W$  and a point  $p$  on the corresponding elliptic surface  $S_W$ . That is,

$$\mathcal{UW}_d := \{(W, p) \in \mathcal{W}_d \times (\mathbb{CP}^1 \times \mathbb{CP}^2) \mid p \in S_W\}.$$

There are natural projection maps

$$\mathcal{UW}_d \xrightarrow{f} \mathbb{CP}^1 \times \mathcal{W}_d \xrightarrow{g} \mathcal{W}_d.$$

The Selmer space is constructed as  $\text{Sel}_n^d(\mathbb{C}) := R^1g_*(R^1f_*(\mu_n))$ .

**Question 9** For fixed  $n$ , do  $\text{Sel}_n^d(\mathbb{C})$  satisfy homological stability as  $d \rightarrow \infty$ ?

In particular, Landesman provided a more precise prediction:

**Conjecture 10** There are constants  $A$  and  $B$  depending on  $n$  so that

$$\dim H_i(\text{Sel}_n^d(\mathbb{C}); \mathbb{Q}) = \dim H_i(\text{Sel}_n^{d+1}(\mathbb{C}); \mathbb{Q})$$

whenever  $d \geq Ai + B$ .

#### 2.5 Will Sawin

Continuing the paradigm put forward by Vakil–Wood [82], Sawin proposed further comparison of three types of convergence for families of varieties:

- (1) Homological stability, i.e., isomorphisms between low-degree singular cohomology groups (equivalently, high-degree compactly-supported cohomology groups), ideally in some way respecting arithmetic structures such as Galois representations, Hodge structures, etc.

- (2) Point counts, in particular after normalizing by a factor of  $q^{-\dim}$ .
- (3) Convergence in the Grothendieck ring under the dimension filtration; see, for example, [82].

In particular, Sawin suggested fixing a degree  $d$  and considering a sequence  $H_n$  of smooth degree  $d$  hypersurfaces in  $\mathbb{P}^n$  as follows:

**Question 11** Does  $\lim_{n \rightarrow \infty} \frac{[H_n] - [\mathbb{P}^{n-1}]}{[\mathbb{A}^{n-1}]}$  exist? Is the limit zero?

This convergence makes sense for many objects that can be obtained via maps out of the Grothendieck group (i.e., Hodge structures), so a negative result would show the Grothendieck ring is richer than these. In the setting of homological stability, this question reduces to the Lefschetz hyperplane theorem; with respect to counting points over  $\mathbb{F}_q$ , the limit can be controlled when  $\sqrt{q}$  is larger than  $d - 1$  using Lefschetz together with the Weil conjectures. While any progress toward answering this question is interesting in its own right, there is also an application to proving Grothendieck ring analogues of the Browning–Sawin circle method [13].

## 2.6 Ronno Das

In 1849, Cayley and Salmon [14] showed that every smooth cubic surface over an algebraically closed field contains exactly 27 lines. If we instead let  $X$  be a smooth cubic surface over  $k = \mathbb{R}$  or  $\mathbb{F}_q$ , for example, the number of lines contained in  $X$  and defined over  $k$  can be strictly less than 27. In fact, as shown by Das [19], the average number of lines on a cubic surface over  $\mathbb{F}_q$  is exactly 1 for all but finitely many characteristics.

More generally, as suggested in correspondence with Ravi Vakil, one can consider lines on del Pezzo surfaces of degree  $1 \leq d \leq 9$ , where  $d = 3$  is the case of cubic surfaces. See for example [7] and [75] for more details.

**Problem 12** What is the average number of  $\mathbb{F}_q$ -lines of a degree 1 del Pezzo surface over  $\mathbb{F}_q$ ?

The average number of lines on a del Pezzo surface defined over  $\mathbb{F}_q$  is 1 for degree  $d \geq 3$ , except when  $d = 7$  where the average is 2. Moreover, the  $d = 2$  case is closely related to Bergvall’s computations [6].

**Problem 13** Is there a uniform proof for all degrees  $1 \leq d \leq 9$ ?

## 2.7 Jesse Kass

When  $q$  is odd or when  $k = \mathbb{R}$ , Segre [77] observed that a line  $\ell$  on a cubic surface  $X$  comes equipped with a distinguished involution  $L$  that is necessarily *hyperbolic* or *elliptic* as an element of  $\mathrm{PGL}_2$ , determined by whether or not the fixed points of  $L$  are defined over  $k$ .

**Definition 14** A line  $\ell$  on a cubic surface  $X$ , both defined either over  $\mathbb{R}$  or  $\mathbb{F}_q$  for  $q$  odd, is said to be *hyperbolic* (respectively, *elliptic*) if the distinguished involution  $L$  (see [77]) is hyperbolic (respectively, elliptic).

See the papers of Finashin–Kharlamov [30], Okonek–Teleman [69], and Kass–Wickelgren [54] on how this distinction gives rise to restrictions for the number of lines on cubic surfaces defined over  $\mathbb{R}$  and  $\mathbb{F}_q$  with odd characteristic.

**Problem 15** Let  $X$  be a cubic surface over  $\mathbb{F}_q$ , a finite field of odd characteristic, and with 27 lines defined over  $\mathbb{F}_q$ . Let  $H$  and  $E$  be the number of hyperbolic and elliptic lines, respectively.

- (1) What pairs  $(H, E)$  can appear?
- (2) Statistically, how often do each appear?
- (3) What subgraphs of the intersection graph of lines can be given by hyperbolic and elliptic lines?

### 2.8 Isabel Vogt

**Problem 16** Let  $X$  be a smooth cubic hypersurface defined over  $\mathbb{F}_q$  and fix  $n \leq q + 1$  points from  $X$ . Does there exist an algebraic map  $f : \mathbb{P}_{\mathbb{F}_q}^1 \rightarrow X$  such that the  $n$  points are contained in the image? If so, what degree is necessary?

As proved by Kollar [57], the answer to the first question is yes when  $q \geq 8$ , but the degree of  $f$  grows rapidly. In general, for  $X$  a fixed hypersurface with  $g : Z \rightarrow X$  a map on some finite subscheme  $Z$  of  $\mathbb{P}^1$ , one can consider the space

$$\text{Mor}_d(\mathbb{P}^1, X; g) := \{f \in \text{Mor}_d(\mathbb{P}^1, X) : f|_Z = g\}$$

of maps extending  $g$  to all of  $\mathbb{P}^1$ . The preceding question becomes that of the  $\mathbb{F}_q$ -points of  $\text{Mor}_d(\mathbb{P}^1, X; g)$ . In this spirit, Vogt offered the following general problem:

**Problem 17** Asking these questions over  $\mathbb{C}$ , how does the topology of the associated spaces vary as  $d$  grows?

As a related example, fixing a degree  $d$  holomorphic function  $g : \mathbb{CP}^{m-1} \rightarrow \mathbb{CP}^n$ , Mostovoy [65, 67] considered the space

$$\text{Mor}_d(\mathbb{CP}^m, \mathbb{CP}^n; g) := \{f : \mathbb{CP}^m \rightarrow \mathbb{CP}^n | f \circ i = g\}$$

for some fixed embedding  $i : \mathbb{CP}^{m-1} \hookrightarrow \mathbb{CP}^m$ , and exhibited a map

$$\text{Mor}_d(\mathbb{CP}^m, \mathbb{CP}^n; g) \rightarrow \Omega^{2m} \mathbb{CP}^n$$

that induces isomorphisms in homology up to degree  $d(2n - 2m + 1) - 1$ . Since the latter space has a homotopy type naturally invariant of  $d$ , this result (and subsequent generalizations into toric varieties [66]) gives a homological stability result in the spirit of Problem 17.

### 2.9 Vlad Matei

**Problem 18** Consider the variety of monic square-free polynomials,

$$\text{Poly}_n(k) := \{(a_{n-1}, \dots, a_0) \in k^n \mid t^n + a_{n-1}t^{n-1} + \dots + a_0 \text{ is square-free}\}.$$

Fixing  $n$  and  $k = \mathbb{F}_q$ , what is the distribution of the  $a_i \in \mathbb{F}_q$  for  $i \leq n - 2$ ?

In the  $q \gg 1$  regime, the coefficients are well-known to be uniformly distributed. On the other hand, fixing  $q$ , one can see immediately that  $a_{n-1}$  is uniformly distributed simply by

elementary substitutions. The question of fixing some of the coefficients is of particular interest to computer scientists; see, for example, [39] for a history of the problem. Matei outlined two distinct ways to study the aforementioned problem:

The first is by dualizing—considering the discriminant hypersurface  $\Delta \subset \mathbb{A}^n$  and picking out  $\mathbb{F}_q$ -points by their coordinates. These, in turn, can be thought of as hyperplane sections of  $\Delta$ . The downside of this approach is that we would need an explicit formula for  $\Delta$  in terms of the coefficients, which becomes computationally infeasible even in small degrees.

The second approach is by considering hypersurfaces determined by the symmetric sums of the roots  $\{z_1, \dots, z_n\}$ . Writing  $\sigma_k$  for the  $k$ th symmetric sum, one could consider hypersurfaces of the form  $\sigma_k = a \in \mathbb{F}_q$  on the open locus of distinct roots. In this case, the objective is to understand the  $S_n$ -action on these hypersurfaces and the associated cohomology groups, in order to count  $\mathbb{F}_q$  points on the quotient.

### 2.10 Oishee Banerjee

While many of the results concerning the cohomology of moduli spaces of polynomials concentrate on data encoded in their zeroes, i.e., configuration spaces on  $\mathbb{C}$ , one can also proceed by studying ramification. Such work is in the spirit of long-standing open problems concerning the topology of Hurwitz spaces.

Indeed, the irreducibility of the Hurwitz space is a classical result proved in [16], where the topology of its subvarieties corresponding to specific ramification loci is almost completely unknown. Banerjee [4] studied the stable cohomology of these Hurwitz spaces satisfying certain conditions. In addition, her work shows that the étale cohomology does not stabilize in positive characteristic, which is in contrast to comparable stability results (see [27, 29].)

**Problem 19** Fix a prime power  $q$  and integer  $n$  with  $\text{char}(\mathbb{F}_q) > n + 1$ . For which  $f(t) \in \text{Poly}_n(\mathbb{F}_q)$  does the anti-derivative

$$F(t) := \int_0^t f(x) dx$$

give a simply-branched mapping  $F : \mathbb{A}^1 \rightarrow \mathbb{A}^1$ ? Alternatively, how many simply-branched maps  $F : \mathbb{A}^1 \rightarrow \mathbb{A}^1$  over  $\mathbb{F}_q$  of degree  $n + 1$  have  $\frac{dF}{dt} \in \text{Poly}_n(\mathbb{F}_q)$ ?

### 2.11 Patricia Hersh

This question is motivated by the task of moving from studying the configuration space of  $n$  distinct points on a manifold, either entirely labeled or unlabeled, to instead looking at situations where the  $n$  points are partitioned into groups of points that cannot be distinguished from each other but can be distinguished from the other groups of points.

If  $A$  is a hyperplane arrangement, one can count the number of  $\mathbb{F}_q$  points in the complement  $X$  of  $A$  using the Möbius function  $\mu_A$  of the intersection poset  $L_A$  (see, for example, [80]). In short:

$$\#X(\mathbb{F}_q) = \sum_{y \in L_A} \mu(\hat{0}, y) q^{\dim A - \text{rk } y}. \quad (2.1)$$

Hersh–Kleinberg introduced a multiplicative deformation of the Möbius function,  $\mu'$ , defined somewhat similarly to  $\mu_A$  but (1) replacing the recursion for  $\mu_A$  by a recursion that

forces  $\mu'$  to be multiplicative, and (2) replacing the intersection poset  $L_A$  in the counting formula above by the multiset partition poset ordered by refinement. In particular,  $\mu'$  records multiplicities of incidences and thus behaves more elegantly in various contexts (see [47] for details).

**Question 20** *Replacing  $\mu$  with  $\mu'$  in (2.1) and replacing  $L_A$  by the multiset partition poset, does the resulting formula have arithmetic meaning?*

Additional references are [36] and [46], the former remarked by Melanie Wood.

### 2.12 Joseph Gunther

**Question 21** *Let  $\mathcal{C}(a, b)$  be moduli space of smooth complex curves in  $\mathbb{P}^1 \times \mathbb{P}^1$  of bi-degree  $(a, b)$ . Is there a homological stability for  $\mathcal{C}(a, b)$  as  $b \rightarrow \infty$  with  $a$  fixed?*

This problem is solved over  $\mathbb{F}_q$  by [28] and a motivic treatment is found in [9].

### 2.13 Vlad Matei

In 1984, Kani–Rosen [52] studied relations between idempotents in the algebra of rational endomorphisms of a fixed abelian variety. In particular, if a group  $G$  covered by subgroups  $H_1, \dots, H_r$  acts on a smooth variety  $X$  over  $\mathbb{Q}$ , we obtain idempotents  $\epsilon_{H_i}$  given by

$$\epsilon_{H_i} := \frac{1}{H_i} \sum_{h \in H_i} h$$

in the group algebra  $\mathbb{Q}[G]$  that come with linear relations of the form

$$\sum_{i=1}^r a_i \epsilon_{H_i} = 0.$$

**Question 22** *Do these transfer to relations in the Grothendieck ring? That is, are there  $a_i \in \mathbb{Z}$  such that*

$$\sum_{i=1}^r a_i [X/H_i] = 0?$$

Matei observed that this holds for the action of  $G = S_3$  on affine space  $\mathbb{A}^3$ , but Daniel Litt commented that he expects this to be false mod  $p$  generically.

## 3 Developing tools, methods, and new examples

### 3.1 Will Sawin

Sawin's talk outlined joint work with Tim Browning on a geometric version of the Hardy–Littlewood circle method, used to compute the compactly supported cohomology of the space of rational curves on a smooth hypersurface (see [13]).

**Question 23** *Can we apply these methods to other cases, such as to maps*

- (1) *from higher genus curves?*
- (2) *into projective varieties?*
- (3) *into complete intersections?*
- (4) *with local conditions?*
- (5) *with Hodge structure modules?*

### 3.2 Michael Fried

For  $n \geq 4$ , two *moduli* spaces associated with  $\mathbb{P}_z^1$  appear often:

The moduli of ordered (distinct) points  $\mathcal{M}_{0,n} := ((\mathbb{P}_z^1)^n \setminus \Delta_n) / \mathrm{PGL}_2(\mathbb{C})$ , and

The moduli of unordered (distinct) points  $J_n := (\mathbb{P}^n \setminus D_n) / \mathrm{PGL}_2(\mathbb{C})$

where  $\Delta_n$  is the “fat diagonal,” i.e., the locus of points with repetitions, and  $D_n$  is the discriminant (treating the homogeneous coordinates in  $\mathbb{P}^n$  as the coefficients of a degree  $n$  polynomial in one variable). For  $n = 4$ , these are the (open) classical lines  $\mathbb{P}_\lambda^1 \setminus \{0, 1, \infty\}$  and  $\mathbb{P}_j^1 \setminus \{\infty\}$ . The fundamental group of  $\mathcal{M}_{0,n}$  mod outer automorphisms,  $\pi_n$ , is the pure (spherical) braid group on  $n$  strands mod its center  $\mathbb{Z}/2\mathbb{Z}$ . Note that  $\pi_4$  is a free group on two generators,  $F_2$ .

The Grothendieck–Teichmüller group is a conjectured description (cf. [24, 41, 49]) of the absolute Galois group  $G_{\mathbb{Q}}$  given by its action on the profinite fundamental group(oids) of  $\mathcal{M}_{0,4}$  and  $\mathcal{M}_{0,5}$ . Even if correct, Grothendieck’s conjecture is not useful without a way to name elements in the action of  $G_{\mathbb{Q}}$  (cf. [49, Question 2, p. 6]).

Ihara replaced  $\pi_n$  by  $\pi_n^{nil}$ , its nilpotent completion, as a way to name the elements in their action; note that this will not capture all of  $G_{\mathbb{Q}}$ . He, along with Anderson, documented aspects of this for explicit collections of covers (e.g., [1, 2]). Foremost was his identification of the second commutator quotient of  $G_{\mathbb{Q}}$  (the 2-step nilpotent quotient), using the 3-, 4-, and 5-cycle relations developed by Drinfeld (and apparently known to Grothendieck). This generalized the Kronecker–Weber description to the abelian quotient of  $G_{\mathbb{Q}}$  as the (Galois group over  $\mathbb{Q}$  of the) cyclotomic closure of  $\mathbb{Q}$  (see [48]).

**Problem 24** Go beyond the nilpotent quotient  $\pi_n^{nil}$  to give a naming scheme for elements in  $G_{\mathbb{Q}}$ .

We propose that a moduli-theoretic approach is possible using reduced Hurwitz space covers (see [31] and [3, 32]). These covers of  $J_n$ —defined by finite groups and conjugacy classes within these group, and parameterizing branched covers of  $\mathbb{P}^1$  with fixed branching data and monodromy group—pull back to unramified covers of  $\mathcal{M}_{0,n}$  whose components and their fields of definition are controlled by an explicit braid group action. For example, for  $n = 4$ , components, genuses and cusps of these covers of  $\mathcal{M}_{0,4} = \mathbb{P}_\lambda^1 \setminus \{0, 1, \infty\}$  are efficiently computable. We propose that these Hurwitz moduli spaces give a way to identify the  $G_{\mathbb{Q}}$  action. Using versions of Deligne’s tangential base points [23] based in  $J_n$  rather than  $\mathcal{M}_{0,n}$  (significant even for  $n = 4$ ) thereby gives tests relating Drinfeld’s relations to the fields of definition of components of Hurwitz space.

**Special case:** even the case when the Hurwitz space pullbacks to  $\mathcal{M}_{0,n}$  are nilpotent covers should be doable and significant (e.g., in shedding new light on Ihara’s results).

### 3.3 Jesse Kass

The following was discussed in the context of developing an enriched discriminant or conductor using  $\mathbb{A}^1$ -homotopy theory, in the context of the mini-course by Kirsten Wickelgren and work by Marc Levine; see, for example, [61].

Consider a generically separable non-constant Galois cover  $f : Y \rightarrow X$  of smooth projective curves with Galois group  $G$ . Associated with  $df$ , there is a certain Euler class, the *enriched different*, which is related to the “enriched Riemann–Hurwitz formula” due to

Levine. In classical theory, the different is related to the discriminant and the conductor; see, for example, [68] for an overview.

**Question 25** *Is there an enriched conductor associated with  $y \in Y$  and  $x = f(y)$ ?*

**Definition 26** The classical conductor is defined via the Artin character (see [40]). In this context, where  $G_y := \text{Stab}_G(y)$ , we have an *enriched Artin character*

$$G_y \rightarrow \text{GW}(k)$$

$\sigma \mapsto$  local Lefschetz trace of  $\sigma$ ,

where  $\text{GW}(k)$  is the Grothendieck–Witt group associated with the field  $k$ .

**Question 27** *Is the enriched Artin character the character of some map?*

More generally, the task here is to continue enriching Serre’s book [40].

### 3.4 Daniel Litt

Following the work of Marc Levine, Jesse Kass, Kirsten Wickelgren, and others (see, for example, [54, 55, 62]), Daniel Litt asked about creating enriched Gromov–Witten invariants.

**Question 28** *Can we develop enriched Gromov–Witten invariants in the non-enumerative case? Can we relax the conditions, e.g., for non-smooth varieties?*

Kass remarked that there are computations over the real numbers, for example [37, 78, 84], which might give us some hint on how to define them.

### 3.5 Hannah Knight

In 1979, based on intuition from Morse theory, Segal [76] proved that the map

$$\text{Rat}_d^n(\mathbb{C}) \hookrightarrow \Omega^2 \mathbb{CP}^n$$

is a homotopy equivalence through dimension  $(2n - 1)d$ , where  $\Omega^2 \mathbb{CP}^n$  is the space of based continuous maps  $\mathbb{CP}^1 \rightarrow \mathbb{CP}^n$  with the compact-open topology. Part of his proof included exhibiting homological stability via (non-algebraic) maps

$$\text{Rat}_d^n(\mathbb{C}) \hookrightarrow \text{Rat}_{d+1}^n(\mathbb{C}).$$

This work inspired many generalizations in the 40 years since, for example extending the domain to genus  $g \geq 1$  curves and the target to Grassmannians or toric varieties; see, for example, [12, 17, 42, 43, 51, 56]. Jun-Yong Park remarked that his joint work with Changho-Han and Hunter Spink [44, 72] showed stability for maps from  $\mathbb{P}^1$  to weighted projective spaces  $\mathbb{P}(a, b)$ . Kirsten Wickelgren commented that the Chow ring of stable maps from  $\mathbb{P}^1 \rightarrow \mathbb{P}^n$  is known to stabilize in the degree, due to Pandharipande [70].

Knight asked what was known about extensions of Segal’s results to the case

$$\text{Hol}_d(\mathbb{CP}^m, \mathbb{CP}^n) = \{(f_0, \dots, f_n) : \text{each } f_i \in \mathbb{C}[x_0, \dots, x_m]_d \text{ with no common root}\},$$

where  $1 < m \leq n$  and  $d \geq 1$ . Claudio Gómez-González remarked that this work had been carried out, first by Mostovoy [65, 67] and, more generally, by Mostovoy–Munguia–Villanueva [66]. Further, [38] calculates the stable homology as  $d \rightarrow \infty$ , and shows that both the unstable rational homology and the corresponding point counts seem to be

independent of  $d$  (except for dimension reasons, for the latter). Benson Farb added that in general, little is known about spaces of algebraic maps when the (complex) dimension of the domain is greater than one (except when those spaces are finite). This motivates:

**Question 29** *Can we develop more general methods to study the topology of spaces of algebraic maps  $X \rightarrow Y$  between varieties, especially when  $X$  has dimension greater than 1?*

**Question 30** *Can we replace homology/cohomology with the Chow ring in more general settings?*

### 3.6 Jesse Kass

**Question 31** *What are the standing obstructions to computing cohomologies of discriminants? Are there problems that can serve as benchmarks for extending the existing techniques?*

Orsola Tommasi commented that she uses Vassiliev's method (see [81, 83]). Many papers cited here use some variant of Vassiliev's spectral sequence; Mostovoy [67], for example, use a truncated version of the spectral sequence when the limiting terms become too cumbersome to compute.

### 3.7 Jun-Yong Park

Trace formulas give a concrete way to work with arithmetic topology. However, moduli functors that we want to study are often represented in the category of stacks rather than schemes, which are harder to work with. Kai Behrend [5] established a trace formula for algebraic stacks, which has extended this arithmetic topology bridge to many moduli stacks. For example, Han and Park [44, 45, 71] computed the  $\ell$ -adic cohomology of the Hom stack  $\text{Hom}_n(\mathbb{P}^1, \mathbb{P}(\lambda_0, \dots, \lambda_N))$  and its class in the Grothendieck ring, with connections to fine moduli stacks of elliptic and hyperelliptic fibrations over  $\mathbb{P}^1$ .

**Question 32** *Can we develop a robust and general theory of point counting to understand moduli stacks arithmetically?*

Wei Ho remarked on important connections to the work of Ellenberg–Satriano–Zureick–Brown [26], bringing together many arithmetic problems about counting points, such as the Batyrev–Manin conjecture, the Malle conjecture, and more via, among other things, establishing a theory of heights on stacks. See, for example, the paper by Boggess and Sankar [10], which discusses stacky versions of these conjectures in the context of the Ellenberg–Satriano–Zureick–Brown framework.

### 3.8 Inna Zakharevich

Consider the Grothendieck ring of varieties over a field  $k$ ,  $K_0(\text{Var}_k)$ , equipped with its dimension filtration, where the  $n$ th graded piece of the filtration is given by those elements that can be represented as sums of varieties of dimension at most  $n$ , i.e., the image of the natural map

$$K_0(\text{Var}_k^{\leq n}) := \mathbb{Z}[X \mid \dim X \leq n] / ([X] = [Y] + [X - Y]) \rightarrow K_0(\text{Var}_k). \quad (3.1)$$

The quotients  $K_0(\text{Var}_k^{\leq n})/K_0(\text{Var}_k^{\leq n-1})$  have a simple presentation as the free abelian group on the birational isomorphism classes of  $n$ -dimensional varieties.

However, as shown by Karzhemanov [53] in 2014 and Borisov [11] in 2015, the map (3.1) is non-injective when  $k \subseteq \mathbb{C}$ . The failure of injectivity is measured by a spectral sequence whose differentials compute the differences

$$[X - U] - [X - V],$$

where  $U$  and  $V$  are isomorphic open subsets identified by a birational automorphism of  $X$ ; for details, see [87]. This motivates the following problem:

**Problem 33** Fix a field  $k$ , variety  $X$ , and a birational automorphism  $f : X \dashrightarrow X$ . Let  $U, V \subseteq X$  be open subsets such that  $f : U \rightarrow V$  is an isomorphism. Give an explicit example such that the complements  $X - U$  and  $X - V$ , which will be equal in the Grothendieck ring, are not piecewise isomorphic.

As remarked above, such an example must exist for  $k \subseteq \mathbb{C}$ , though none is known; it is unknown if examples exist over other fields.

### 3.9 Zinovy Reichstein

**Problem 34** Let  $G$  be a finite group and fix subgroups  $H_1, \dots, H_r \leq G$ , not necessarily distinct. Moreover, fix  $d > 0$  and faithful  $d$ -dimensional representations  $V_i$  of each of the  $H_i$ . Is there a  $d$ -dimensional irreducible variety  $X$  over  $\mathbb{C}$  with a  $G$  action such that there are  $r$  distinct points  $x_1, \dots, x_r$  with  $x_i$  fixed by  $H_i$  and isotropy representation given by  $V_i$ ?

Reichstein remarked that this is analogous in spirit to the Chinese Remainder Theorem and can be done if each  $H_i$  is abelian; see [74, Theorem 8.6] for details. Moreover, when  $r = 1$  and  $H = G$ , one can simply take  $X = V$ , while for  $r = 2$  and  $H_1 = H_2 = G$ , it is an unpublished result of A. Kresch. In the special case, where  $H_1, \dots, H_r$  are the non-conjugate Sylow subgroups of  $G$  (i.e., a list of Sylow subgroups for the different primes dividing  $|G|$ ), this problem arose in the context of work on essential dimension of finite groups (see [Conjecture 11.5] [25]).

### 3.10 Hunter Spink

**Problem 35** Let  $R$  be a ring with  $m_1, \dots, m_k \in R$ . Assume that for any subset  $A \subset \{1, \dots, k\}$ , we have

$$\sum_{i \in A} m_i \in R^\times.$$

Also set  $a_i = m_1 x_1^i + \dots + m_k x_k^i \in R[x_1, \dots, x_k]$ . What is the smallest  $n$  such that there exist monic  $f(z), g(z)$  with coefficients in  $R[a_1, \dots, a_n]$  giving

$$\frac{f(z)}{g(z)} = \frac{m_1}{z - x_1} + \dots + \frac{m_k}{z - x_k}?$$

For context on this problem, see the paper by Spink and Dennis Tseng [79] on incidence strata of affine varieties that also outlines their conjecture of  $n = 2^k + 1$  to problem 35.

### Acknowledgements

We thank all of the people who posed problems, both for their original suggestions and for their help clarifying and refining our write-ups of them. We thank Christin Bibby, Aaron Landesman, Lior Silberman, and Bena Tshishiku for transcribing the problem sessions in real time. We thank Benson Farb for helpful comments. We thank all the participants

who commented and asked questions during the problem sessions. We thank the anonymous referee for helpful comments. Last, we thank Alejandro Adem, Craig Westerland and Melanie Wood for co-organizing the workshop, and we thank the PIMS staff for their hospitality and support in organizing the workshop.

Received: 7 January 2021 Accepted: 22 March 2021 Published online: 12 April 2021

## References

1. Anderson, G., Ihara, Y.: Pro- $\ell$  branched coverings of  $\mathbb{P}^1$  and higher circular  $\ell$ -units. *Ann. Math.* **128**, 271–293 (1988)
2. Anderson, G., Ihara, Y.: Pro- $\ell$  branched coverings of  $\mathbb{P}^1$  and higher circular  $\ell$ -units. II. *Int. J. Math.* **1**(2), 119–148 (1990)
3. Bailey, P., Fried, M., Hurwitz monodromy, spin separation and higher levels of a modular tower., Arithmetic fundamental groups and noncommutative algebra (Berkeley, CA, : Proc. Sympos. Pure Math., vol. 70. AMS **2002**, 79–220 (1999)
4. Banerjee, O.: Cohomology of the space of polynomial maps on  $\mathbb{A}^1$  with prescribed ramification. *Adv. Math.* **359**, 106881 (2020)
5. Behrend, K.: The Lefschetz trace formula for algebraic stacks. *Invent. Math.* **112**(1), 127–149 (1993)
6. Bergvall, O.: Equivariant cohomology of moduli spaces of genus three curves with level two structure. *Geom. Dedicata* **202**(1), 165–191 (2019)
7. Bergvall, O., Gounelas, F.: Cohomology of moduli spaces of del Pezzo surfaces, arXiv preprint [arXiv:1904.10249](https://arxiv.org/abs/1904.10249) (2019)
8. Bhargava, M., Kane, D., Lenstra, H., Poonen, B., Rains, E.: Modeling the distribution of ranks, Selmer groups, and Shafarevich-Tate groups of elliptic curves. *Camb. J. Math.* **3**(3), 275–321 (2015)
9. Bilu, M., Howe, S.: Motivic Euler products in motivic statistics, [arXiv:1910.05207](https://arxiv.org/abs/1910.05207), 2019
10. Boggess, B., Sankar, S.: Counting elliptic curves with a rational  $n$ -isogeny for small  $n$ , [arXiv:2009.05223](https://arxiv.org/abs/2009.05223)
11. Borisov, L.: Class of the affine line is a zero divisor in the Grothendieck ring. *J. Algeb. Geom.* **27**(2), 203–209 (2018)
12. Boyer, C.P., Hurtubise, J.C., Milgram, R.J.: Stability theorems for spaces of rational curves. *Int. J. Math.* **12**(2), 223–262 (2001)
13. Browning, T., Sawin, W.: A geometric version of the circle method. *Ann. Math.* **191**(3), 893–948 (2020)
14. Cayley, A.: *On the triple tangent planes of surfaces of the third order*, Cambridge Library Collection - Mathematics, vol. 1, pp. 445–456, Cambridge University Press, 2009
15. Clancy, J., Kaplan, N., Leake, T., Payne, S., Wood, M.M.: On a Cohen-Lenstra heuristic for Jacobians of random graphs. *J. Algeb. Combin.* **42**(3), 701–723 (2015)
16. Clebsch, A.: *Zur theorie der Riemann'schen fläche*, *Mathematische Annalen* **6**, 216–230
17. Cohen, F.R., Cohen, R.L., Mann, B.M., Milgram, R.J.: The topology of rational functions and divisors of surfaces. *Acta Math.* **166**, 163–221 (1991)
18. H. Cohen, H. and Lenstra, *Heuristics on class groups of number fields*, Number Theory Noordwijkerhout 1983, Springer, 1984, pp. 33–62
19. Das, R.: The space of cubic surfaces equipped with a line. *Math. Z.* (2020)
20. de Jong, A.J.: A conjecture on arithmetic fundamental groups. *Israel J. Math.* **121**, 61–64 (2001)
21. Delaunay, C.: Heuristics on Tate-Shafarevitch groups of elliptic curves defined over  $\mathbb{Q}$ . *Exp. Math.* **10**(2), 191–196 (2001)
22. Deligne, P.: *La conjecture de Weil. II*, *Publ. Math. IHÉS* **52** (1980), no. 5522, 137–252
23. Deligne, P.: *Le groupe fondamental de la droite projective moins trois points*, Galois groups over  $\mathbb{Q}$ , vol. 16, Springer, 1989, pp. 79–297
24. Drinfeld, V.: *On quasi-triangular quasi-Hopf algebras and some group closely associated with  $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$* , *Algebra i Analiz* **2** (1990), no. 4, 149–181
25. Duncan, A., Reichstein, Z.: Pseudo-reflection groups and essential dimension. *J. Lond. Math. Soc.* **90**(3), 879–902 (2014)
26. Ellenberg, J., Satriano, M., Zureick-Brown, D.: *Heights on stacks and a generalized Batyrev–Manin–Malle conjecture*, in preparation
27. Ellenberg, J., Venkatesh, A., Westerland, C.: Homological stability for Hurwitz spaces and the Cohen-Lenstra conjecture over function fields. *Ann. Math.* **183**, (2009)
28. Erman, D., Wood, M.M.: Semiample Bertini theorems over finite fields. *Duke Math. J.* **164**(1), 1–38 (2015)
29. Farb, B., Wolfson, J.: Étale homological stability and arithmetic statistics. *Quart. J. Math.* **69**, (2015)
30. Finashin, V., Kharlamov, S.: Abundance of real lines on real projective hypersurfaces. *Int. Math. Res. Notices* **16**, 3639–3646 (2012)
31. Fried, M.: Moduli relations between  $\ell$ -adic representations and the regular inverse Galois problem. *Grad. J. Math.* **5**(1), 38–75 (2020)
32. Fried, M., Volklein, H.: The inverse Galois problem and rational points on moduli spaces. *Mathematische Annalen* **290**(4), 771–800 (1991)
33. Gaitsgory, D.: On de Jong's conjecture. *Israel J. Math.* **157**, 155–191 (2007)
34. Galatius, S., Kupers, A., Randal-Williams, O.: *Cellular  $E_k$ -algebras*, [arXiv:1805.07184](https://arxiv.org/abs/1805.07184), 2018
35. Galatius, S., Kupers, A., Randal-Williams, O.:  $E_2$ -cells and mapping class groups. *Publ. Math. IHÉS* **130**, 1–61 (2019)
36. Garton, D.: Random matrices, the Cohen-Lenstra heuristics, and roots of unity. *Algeb. Number Theory* **9**(1), 149–171 (2015)
37. Georgieva, P., Zinger, A.: Real Gromov-Witten theory in all genera and real enumerative geometry: Construction. *Ann. Math.* **188**(3), 685–752 (2018)
38. Gómez-González, C.: *Spaces of non-degenerate maps between complex projective spaces*, *Res Math Sci* **7** (2020), no. 26
39. Granger, R.: On the enumeration of irreducible polynomials over  $GF(q)$  with prescribed coefficients. *Finite Fields Appl.* **57**, 156–229 (2019)
40. Greenberg, M.J., Serre, J.P.: *Local Fields*. Graduate Texts in Mathematics. Springer, New York (2013)

41. Grothendieck, A.: *Esquisse d'un programme*, Geometric Galois actions, vol. 242, Cambridge University Press, 1997, pp. 5–48
42. Guest, M.A.: The topology of the space of rational curves on a toric variety. *Acta Math.* **174**(1), 119–145 (1995)
43. Guest, M.A., Kozłowski, A., Yamaguchi, K.: Spaces of polynomials with roots of bounded multiplicity. *Fund. Math.* **161**(1–2), 93–117 (1999)
44. Han, C., Park, J.-Y.: Arithmetic of the moduli of semistable elliptic surfaces. *Math. Ann.* **375**(3–4), 1745–1760 (2019)
45. Han, C., Park, J.-Y.: *Enumerating algebraic curves and abelian varieties over global function fields with lower order terms*, [arXiv:2002.00563](https://arxiv.org/abs/2002.00563), 2020
46. Hersh, P.: Chain decomposition and the flag f-vector. *J. Combin. Theory Ser. A* **103**(1), 27–52 (2003)
47. Hersh, P., Kleinberg, R.: *A multiplicative deformation of the mobius function for the poset of partitions of a multiset*, *Communicating Mathematics, Contemp. Math.*, vol. 479, AMS, 01 2009, pp. 113–118
48. Ihara, Y.: What's next after  $\mathbb{Q}(\mu_{\ell^\infty})$ ?, *Algebraic analysis, geometry, and number theory*. JHU Press **153–159**, (1989)
49. Ihara, Y.: *Braids, Galois groups and some arithmetic functions*, *Proceedings of 1990 ICM* (1990)
50. Kahle, M., Lutz, F., Newman, A., Parsons, K.: Cohen-Lenstra heuristics for torsion in homology of random complexes. *Exp. Math.* **29**(3), 357–359 (2020)
51. Kallel, S., Milgram, R.J.: The geometry of the space of holomorphic maps from a Riemann surface to a complex projective space. *J. Differ. Geom.* **47**(2), 321–375 (1997)
52. Kani, E., Rosen, M.: Idempotent relations and factors of Jacobians. *Mathematische Annalen* **284**(2), 307–328 (1989)
53. Karzhemanov, I.: On the cut-and-paste property of algebraic varieties. *Bull. Sci. Math.* **166**, 102920 (2021)
54. Kass, J., Wickelgren, K.: *An arithmetic count of the lines on a smooth cubic surface*, [arXiv:1708.01175](https://arxiv.org/abs/1708.01175)
55. Kass, J., Wickelgren, K.: The class of Eisenbud-Khimshiashvili-Levine is the local  $\mathbf{A}^1$ -brouwer degree. *Duke Math. J.* **168**(3), 429–469 (2019)
56. Kirwan, F.: On spaces of maps from Riemann surfaces to Grassmannians and applications to the cohomology of moduli of vector bundles. *Ark. Mat.* **24**(1–2), 221–275 (1985)
57. Kollar, J., Derenthal, U.: *Looking for rational curves on cubic hypersurfaces*, [arXiv:0710.5516](https://arxiv.org/abs/0710.5516), 2007
58. Koplewitz, S.: Sandpile groups and the coeulerian property for random directed graphs. *Adv. Appl. Math.* **90**, 145–159 (2017)
59. Lafforgue, L.: Chtoucas de Drinfeld et correspondance de Langlands. *Invent. Math.* **147**, 1–241 (2002)
60. Landesman, A.: *The geometric average size of Selmer groups over function fields*, [arXiv:1811.00966](https://arxiv.org/abs/1811.00966), 2018
61. Levine, M.: *Aspects of enumerative geometry with quadratic forms*, [arXiv:1703.03049](https://arxiv.org/abs/1703.03049), 2017
62. Levine, M.: *Toward an algebraic theory of Welschinger invariants*, [arXiv:1808.02238](https://arxiv.org/abs/1808.02238), 2018
63. Li, K.Z., Oort, F.: *Moduli of Spersingular Abelian Varieties*. Lecture Notes in Mathematics. Springer, Berlin (2006)
64. May, J.P.: *The Geometry of Iterated Loop Spaces*. Lecture Notes in Mathematics. Springer, Berlin (2006)
65. Mostovoy, J.: Spaces of rational maps and the Stone-Weierstrass theorem. *Topology* **45**(2), 281–293 (2006)
66. Mostovoy, J., Munguia-Villanueva, E.: Spaces of morphisms from a projective space to a toric variety. *Rev. Colombiana Mat.* **48**(1), 41–53 (2014)
67. Mostovoy, Jacob: Truncated simplicial resolutions and spaces of rational maps. *Quart. J. Math.* **63**, (2012)
68. Neukirch, J., Schappacher, N.: *Algebraic number theory*. Grundlehren der mathematischen Wissenschaften, Springer, Berlin Heidelberg (2013)
69. Okonek, C., Spindler, A.: Intrinsic signs and lower bounds in real algebraic geometry. *J. Reine Angew. Math.* **688**, 219–241 (2014)
70. Pandharipande, R.: *The Chow ring of the Hilbert scheme of rational normal curves*, [arXiv:alg-geom/9607025](https://arxiv.org/abs/alg-geom/9607025), 1996
71. Park, J.-Y.:  $\ell$ -adic étale cohomology of the moduli of elliptic and hyperelliptic fibrations, [arXiv:1812.11694](https://arxiv.org/abs/1812.11694), 2018
72. Park, J.-Y., Spink, H.: *Motive of the moduli stack of rational curves on a weighted projective stack*, *Research in the Mathematical Sciences* **8** (2021), no. 1, #1
73. Randall-Williams, O.: *Homology of Hurwitz spaces and the Cohen–Lenstra heuristics for function fields (after Ellenberg, Venkatesh, and Westerland)*, Séminaire Bourbaki, vol. 71, 2019
74. Reichstein, Z., Youssin, B.: Splitting fields of G-varieties. *Pac. J. Math.* **200**(1), 207–249 (2001)
75. Schicho, J.: *Elementary theory of del Pezzo surfaces*, *Computational Methods for Algebraic Spline Surfaces* (Berlin, Heidelberg), pp. 77–94. Springer, Berlin Heidelberg (2005)
76. Segal, G.: The topology of spaces of rational functions. *Acta Math.* **143**, 39–72 (1979)
77. Segre, B.: The non-singular cubic surfaces: A new method of investigation with special reference to questions of reality. Clarendon Press, Oxford (1942)
78. Solomon, J.P., Tukachinsky, S.B.: *Point-like bounding chains in open Gromov–Witten theory*, [arXiv:1608.02495](https://arxiv.org/abs/1608.02495), 2016
79. Spink, H., Tseng, D.: *Incidence strata of affine varieties with complex multiplicities*, [arXiv:1903.05011](https://arxiv.org/abs/1903.05011), 2019
80. Stanley, R. P.: *An introduction to hyperplane arrangements*, *Geometric Combinatorics* (Bernd Sturmfels, Ezra Miller, and Victor Reiner, eds.), American Mathematical Society, 2007, pp. 389–496
81. Tommasi, O.: Stable cohomology of spaces of non-singular hypersurfaces. *Adv. Math.* **265**, 428–440 (2014)
82. Vakil, R., Wood, M.M.: Discriminants in the Grothendieck ring. *Duke Math. J.* **164**(6), 1139–1185 (2015)
83. Vassiliev, V.A.: *Complements of discriminants of smooth maps: Topology and applications*, *Translations of mathematical monographs*, American Mathematical Society, 1994
84. Welschinger, J.-Y.: Invariants of real symplectic 4-manifolds and lower bounds in real enumerative geometry. *Invent. Math.* **162**(1), 195–234 (2005)
85. Wood, M.M.: The distribution of sandpile groups of random graphs. *J. Amer. Math. Soc.* **30**(4), 915–958 (2017)
86. Wood, M.M.: Random integral matrices and the Cohen-Lenstra heuristics. *Amer. J. Math.* **141**(2), 383–398 (2019)
87. Zakharevich, I.: The annihilator of the Lefschetz motive. *Duke Math. J.* **166**(11), 1989–2022 (2017)

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.