



Potential dispersal of tardigrades by birds through endozoochory: evidence from Sub-Antarctic White-bellied Seedsnipe (*Attagis malouinus*)

Michael W. Robertson¹ · Nicholas J. Russo² · Sandra J. McInnes³ · Bernard Goffinet² · Jaime E. Jiménez^{1,4}

Received: 28 December 2019 / Revised: 25 April 2020 / Accepted: 8 May 2020
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Abstract

Tardigrades are potentially dispersed by birds, but the extent of the interactions between birds and tardigrades is virtually unknown. We discovered nine tardigrades within feces of White-bellied Seedsnipe (*Attagis malouinus*) collected from high Andean tundra on Navarino Island, Chile. Eight of the tardigrade specimens began moving once rehydrated. Two specimens belonged to the genus *Adropion* (Hypsibiidae), one to the *Macrobiotus* (Macrobiotidae), and five could not be identified. A ninth specimen was a species of *Isohypsibius* in an embryonic egg state. These tardigrades could have passed through the avian digestive tract after incidental ingestion or burrowed into the feces post-defecation to feed on microorganisms and undigested plant matter present in the feces. To our knowledge, this is the first discovery of tardigrades in bird feces and may have implications for tardigrade distributions if birds transport tardigrades endogenously.

Keywords Bird feces · Cape Horn region · Herbivory

Introduction

Tardigrades are microscopic metazoans that inhabit terrestrial and aquatic (freshwater and marine) environments throughout the world (Ramazzotti and Maucci 1983; Schill 2018). The phylum Tardigrada is ubiquitous, cosmopolitan, and currently consists of *ca.* 1200 species (Guidetti and Bertolani 2005; Degma and Guidetti 2007; Vicente et al. 2013; Degma et al. 2009–2019). Some tardigrades can enter a cryptobiotic tun state upon desiccation, which allows them to survive various environmental stressors. While in this

protective form of suspended animation, they are thought to disperse passively by wind over great distances (Kristensen 1987; Bertolani et al. 1990; Pugh and McInnes 1998).

Tardigrades occur in bird nests, suggesting that birds may disperse tardigrades by carrying them in their feathers or transporting them altogether with collected material for their nests, e.g., mosses or lichens (Kaczmarek et al. 2018; Mogle et al. 2018). Here, we present the discovery of tardigrades inhabiting the droppings of White-bellied Seedsnipe (*Attagis malouinus* Boddaert, 1783) and propose two possible explanations for their presence: ingestion and endogenous dispersal or post-defecation colonization.

Michael W. Robertson and Nicholas J. Russo have contributed equally.

✉ Michael W. Robertson
MichaelRobertson2@my.unt.edu

¹ Department of Biological Sciences, University of North Texas, Denton, TX, USA

² Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT, USA

³ British Antarctic Survey, Natural Environment Research Council, Madingley Road, Cambridge, UK

⁴ Advanced Environmental Research Institute, University of North Texas, Denton, USA

Materials and methods

This study was conducted during a three-month period from January to March 2018. We collected 26 White-bellied Seedsnipe droppings opportunistically from Andean alpine habitats along the Dientes de Navarino trail on Navarino Island, Chile. Samples were air dried and kept undisturbed at room temperature until examination. The dried external crust surface of the droppings was removed with a razor blade prior to observation to exclude airborne contaminants. Droppings were sliced into 1 mm-wide discs and diluted with filtered

spring water for examination under a dissecting microscope. Live tardigrades recovered from the feces were placed in filtered spring water and unresponsive tardigrades were placed in 90% ethanol. Each tardigrade was mounted on a glass slide using de Faure's mounting media and observed under differential interference contrast at 400 × (using an Olympus BX20 microscope). Most individuals were identified to genus or species (Ramazzotti and Maucci 1983; Degma et al. 2009–2019). Permanent mounts of the specimens were retained by SJM at the British Antarctic Survey.

Results and discussion

Tardigrades have previously been recorded on Navarino Island by Richters (1908) and Rahm (1931, 1932) who reported four taxa *Echiniscus testudo* (Doyère, 1840), *Pseudechiniscus suillus* (Ehrenberg, 1853), *Mesobiotus furciger* (Murray, 1907), and *Macrobiotus hufelandi* (C.A.S. Schultze, 1833), all now considered species complexes. We found nine tardigrade specimens in seedsnipe feces, representing at least four species from three genera. One of the recovered tardigrades was unresponsive after hydration and was presumed dead. All tardigrades were recovered from the interior of two different White-bellied Seedsnipe droppings collected from the same location above tree line (Navarino Island; 579 m a.s.l., 55° 01' 17" S 67° 43' 07" W). We confirmed two distinct species of the genus *Adropion* (Hypsibiidae). The first *Adropion* specimen had three placoids, a microplacoid, and a septulum reminiscent of *Adropion greveni* (Dastych 1984) or *A. onorei* (Pilato et al. 2002), both recorded from South America (Kaczmarek et al. 2015). The differences included the two anterior macroplacoids being of equal size, while in both *A. greveni* and *A. onorei* the second macroplacoid was the shortest; a much larger septulum and larger claws than either *Adropion greveni* or *A. onorei*; and lacked the distinct spines or crenulation at the claw base. The second *Adropion* had three macroplacoids, but no microplacoid or septulum, robust claws with an obvious accessory spine on the inner claws, no bars, lunules, or teeth at the base of the claws, and did not match any *Adropion* recorded from South America. These two specimens of *Adropion* may represent new species, but more material should be collected to confirm their distinction and variation before a formal taxonomic proposal is made. One specimen is believed to belong to the genus *Macrobiotus* (Macrobiotidae). This genus includes several species complexes that require careful taxonomic analysis, including examining egg morphology, to properly differentiate the species. Without eggs, we were unable to identify the species group, which could belong to the *hufelandi*, *pallari* or *echinogenitus* groups. The last specimen, an *Isohypsibius*

sp., was an individual observed in an embryonic egg stage and therefore, the species could not be determined. The remaining five tardigrades were unidentifiable due to degradation while traveling between laboratories.

All but one of the tardigrades discovered regained vigor upon hydration, suggesting that they were in a cryptobiotic state due to desiccation (Welnicz et al. 2011). We suggest two possible reasons for the presence of tardigrades in droppings of White-bellied Seedsnipe: (1) incidental ingestion and endogenous dispersal or (2) post-defecation colonization.


White-bellied Seedsnipes belong to an endemic group of plovers (Charadriiformes: Thinocoridae) of Antarctic origin. They are associated with harsh high elevation and open landscapes of the southern Andes during the breeding season and migrate to lowland grasslands during the winter months (Ibarguchi 2011). Seedsnipes are found from 39° S (Bío-Bío) to 55° 57' S (Cape Horn Island, eBird 2020), are specialized herbivores that resemble Ptarmigans (*Lagopus*) from the Northern Hemisphere. While little is known of their diet, Fjeldså and Kirwan (2020) concur with our observations (JEJ) that they consume crowberry (*Empetrum rubrum*) and grasses near mosses and lichens where tardigrades are present.

If ingested as adults or as eggs, the tardigrades could survive the passage through the birds' digestive tract (Fig. 1). Alternatively, the tardigrades could have burrowed into the already-voided droppings in search of microorganisms or undigested plant tissue as a food source. Indeed, we found numerous live microorganisms in White-bellied Seedsnipe feces during this research. In either case, as the droppings dried, the sunbaked and wind-blown outer crust of the feces may have prevented escape and forced the tardigrades to enter a cryptobiotic state where they lay dormant until their resurrection in the Petri dish where conditions were once again favorable.

In previous studies, tardigrades were found occurring in feces of land snails in Puerto Rico (Fox and García 1962; Fox 1966). In the case of the land snail species *Bulimulus exilis* (Gmelin, 1791), Fox and García (1962) and Fox (1966) believed the most likely explanation for the presence of live tardigrades in feces was that they were incidentally ingested by the snails and passed live through the animals' digestive system. This is the closest link to the work we present here and suggests the potential for tardigrades to move through another organism's digestive tract unscathed.

To our knowledge, our observations are the first of tardigrades inhabiting avian feces. In future studies, the examination of freshly defecated droppings may yield evidence to support the ingestion hypothesis. Experiments with captive birds could be used to determine whether tardigrades survive passage through the avian digestive tract

Fig. 1 Illustration depicting an example of the possible mechanism of endozoochorous dispersal of tardigrades by White-bellied Seedsnipe (*Attagis malouinus*). **a** Unseen tardigrades present in the environment are ingested by a bird during foraging. **b** Tardigrades pass through the digestive system from incidental consumption and are encapsulated within the feces. **c** Tardigrades could be present in various developmental stages: as adults, free laid eggs, or eggs present in the molted cuticle. **d** Ornamented tardigrade egg



(Hillerman et al. 1953). If tardigrades pass through the avian digestive system and remain viable post-defecation, this would add to the growing evidence that birds play a role as long-distance dispersers of tardigrades.

Acknowledgements This study was funded by the National Science Foundation (NSF) International Research Experiences for Students (IRES) 1658651 “Cross-cutting interdisciplinary research and integration of ecology and biocultural conservation in the world’s southernmost forests.” The authors would like to thank the Sub-Antarctic Biocultural Conservation Program for lodging during the experiment, Tamara Contador and the Wankara Lab for providing their laboratory space and equipment, Peter Convey for transferring the tardigrades to the UK, and Jorge Talamantes for providing the illustration of Fig. 1. The authors would also like to thank Roy Mackenzie and Jose A. Sanchez-Ruiz for their input and critique. A special thanks to Łukasz Kaczmarek and Harry Meyer for providing peer review for the publication.

Compliance with ethical standards

Conflicts of interest The authors declare that they have no conflict of interest.

References

- Bertolani R, Rebecchi L, Beccaccioli G (1990) Dispersal of *Ramazzottius* and other tardigrades in relation to type of reproduction. *Invertebr Reprod Dev* 18:153–157. <https://doi.org/10.1080/07924259.1990.9672137>
- Dastych H (1984) The Tardigrada from Antarctica with description of several new species. *Acta Zool Cracov* 27:377–436
- Degma P, Guidetti R (2007) Notes to the current checklist of Tardigrada. *Zootaxa* 1579:41–53. <https://doi.org/10.11646/zootaxa.1579.1.2>
- Degma P, Bertolani R, Guidetti R (2009–2019) Actual checklist of Tardigrada species. https://doi.org/10.25431/11380_1178608
- Doyère PLN (1840) Memoire sur les Tardigrades. *Ann des Sci Nat Zool Paris Ser* 2(14):269–362
- Ehrenberg CG (1853) Diagnoses novarum formarum. *Verhandlungen der Königlich Preussische Akademie der Wissenschaften zu Berlin* 8:526–533
- eBird. Checklist: <https://birdsoftheworld.org/bow/species/whbsee2/cur/introduction>. eBird: An online database of bird distribution and abundance. eBird, Ithaca, New York. <https://www.ebird.org>. Accessed 22 Apr 2020
- Fjeldså J, Kirwan GM (2020) White-bellied Seedsnipe (*Attagis malouinus*). In: del Hoyo J, Elliott A, Sargatal J, Christie DA,

- de Juana E (eds) Birds of the world. Ver. 1. Cornell Lab of Ornithology, Ithaca
- Fox I (1966) On the significance of tardigrades in the digestive tract of snails. In: Proceedings of the first international congress of parasitology, pp 1064–1065
- Fox I, García-Moll I (1962) *Echiniscus molluscorum*, new tardigrade from the feces of the land snail, *Bulimulus exilis* (Gmelin) in Puerto Rico (Tardigrada: Scutichiniscidae). J Parasitol 48:177. <https://doi.org/10.2307/3275559>
- Guidetti R, Bertolani RB (2005) Tardigrade taxonomy: an updated check list of the taxa and a list of characters for their identification. Zootaxa 845:1. <https://doi.org/10.11646/zootaxa.845.1.1>
- Gmelin JF (1791) Caroli a Linnaei Systema Naturae per Regna Tria Naturae, Ed. 13. Tome 1(6). GE Beer, Lipsiae [Leipzig], pp 3289
- Hillerman JP, Kratzer FH, Wilson WO (1953) Food passage through chickens and turkeys and some regulating factors. Poultry Sci 32:332–335. <https://doi.org/10.3382/ps.0320332>
- Ibarguchi, G (2011) Biogeography and diversification of the Andean seedsnipes (Thinocoridae): an Antarctic avian lineage? Dissertation, Queen's University, Ontario, Canada. pp 1–379
- Kaczmarek Ł, Michalczyk Ł, McInnes SJ (2015) Annotated zoogeography of non-marine Tardigrada. Part II: South America. Zootaxa 3923:1–107. <https://doi.org/10.11646/zootaxa.3923.1.1>
- Kaczmarek Ł, Parnikoza I, Gawlak M, Esefeld J, Peter HU, Kozeretska I, Roszkowska M (2018) Tardigrades from *Larus dominicanus* Lichtenstein, 1823 nests on the Argentine Islands (maritime Antarctic). Polar Biol 41:283–301. <https://doi.org/10.1007/s00300-017-2190-4>
- Kristensen RM (1987) Generic revision of the Echiniscidae (Heterotardigrada), with a discussion of the origin of the family. In: Bertolani R (ed) Biology of tardigrades. Selected symposia and sonographs U.Z.I., Vol 1. Mucchi, Modena, pp 261–335
- Mogle MJ, Kimball SA, Miller WR, McKown RD (2018) Evidence of avian-mediated long distance dispersal in American tardigrades. PeerJ 6:e5035. <https://doi.org/10.7717/peerj.5035>
- Murray J (1907) Encystment of Tardigrada. Trans R Soc Edinb 45:837–854
- Pilato G, Binda MG, Napolitano A, Moncada E (2002) Remarks on some species of tardigrades from South America with the description of two new species. J Nat Hist 38:1081–1086. <https://doi.org/10.1080/0022293031000071541>
- Pugh PJA, McInnes SJ (1998) The origin of Arctic terrestrial and freshwater tardigrades. Polar Biol 19:177–182. <https://doi.org/10.1007/s003000050232>
- Rahm G (1931) Tardigrada of the South of America (esp. Chile). Rev Chil Hist Nat Pura y Aplicada 35:118–141
- Rahm G (1932) Freilebende Nematoden, Rotatorien und Tardigraden aus Südamerika (besonders aus Chile). Zool Anz 98:94–128
- Ramazzotti G, Maucci W (1983) Il Phylum Tardigrada. Mem dell'Ist Ital di Idrobiol Pallanza 41:1–1012
- Richters F (1908) Moosbewohner. In: Wissenschaftliche Ergebnisse der Schwedischen Südpolar-Expedition, 1901–1903 unter Leitung von Dr. Otto Nordenskjöld. Lithographisches Institut des Generalstabs, Stockholm. Vol. VI, No. 2. pp 1–17
- Schill RO (ed) (2018) Water bears: The biology of tardigrades. In: Zoological monographs. <https://doi.org/10.1007/978-3-319-95702-9>
- Schultze CAS (1833) Macrobiotus Hufelandii animal e crustaceorum classe novum, reviviscendi post diuturnam asphixiam et ariditatem potens. Apud Carolum Curths, Berolini, p 7
- Vicente F, Fontoura P, Cesari M, Rebecchi L, Guidetti R, Serrano A, Bertolani R (2013) Integrative taxonomy allows the identification of synonymous species and the erection of a new genus of Echiniscidae (Tardigrada, Heterotardigrada). Zootaxa 3613:557–572. <https://doi.org/10.11646/zootaxa.3613.6.3>
- Welnicz W, Grohme MA, Kaczmarek Ł, Schill RO, Frohme M (2011) Anhydrobiosis in tardigrades—the last decade. J Insect Physiol 57:577–583. <https://doi.org/10.1016/j.jinsphys.2011.03.019>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.