

# High-quality assemblies for three invasive social wasps from the *Vespula* genus

1 Thomas W.R. Harrop<sup>\*,1</sup>, Joseph Guhlin<sup>\*,1</sup>, Gemma M. McLaughlin<sup>\*</sup>, Elizabeth Permina<sup>\*</sup>,  
 2 Peter Stockwell<sup>†</sup>, Josh Gilligan<sup>\*</sup>, Marissa F. Le Lec<sup>\*</sup>, Monica A. M. Gruber<sup>‡</sup>, Oliver  
 3 Quinn<sup>‡</sup>, Mackenzie Lovegrove<sup>\*</sup>, Elizabeth J. Duncan<sup>§</sup>, Emily J. Remnant<sup>\*\*</sup>,  
 4 Jens Van Eeckhoven<sup>§</sup>, Brittany Graham<sup>§</sup>, Rosemary A. Knapp<sup>§</sup>, Kyle W. Langford<sup>††</sup>, Zev  
 5 Kronenberg<sup>††</sup>, Maximilian O. Press<sup>††</sup>, Stephen M. Eacker<sup>††</sup>, Erin E. Wilson-Rankin<sup>‡‡</sup>,  
 6 Jessica Purcell<sup>‡‡</sup>, Philip J. Lester<sup>‡</sup>, Peter K. Dearden<sup>\*</sup>

7 1 These authors contributed equally to the work.

8 \* Genomics Aotearoa and Department of Biochemistry, University of Otago, Dunedin  
 9 9054, Aotearoa-New Zealand.

10 † Pathology Department, University of Otago, Dunedin, Aotearoa-New Zealand.

11 ‡ Centre for Biodiversity and Restoration Ecology, School of Biological Sciences, Victoria  
 12 University of Wellington, PO Box 600, Wellington, New Zealand.

13 § School of Biology, Faculty of Biological Sciences, University of Leeds, Leeds LS2 9JT,  
 14 United Kingdom.

15 \*\* School of Life and Environmental Sciences, University of Sydney, Sydney NSW 2006,  
 16 Australia.

17 †† Phase Genomics Inc., 1617 8th Ave N, Seattle, WA 98109, USA.

18 ‡‡ Department of Entomology, University of California, Riverside, CA 92521, USA.

## 19 Accession numbers:

- 20 • Raw reads: [PRJNA643352](#)
- 21 • Genomes and annotations: The version of the assemblies and annotations described
- 22 in this paper is hosted on Zenodo under DOI [10.5281/zenodo.4001020](#). The
- 23 genomes and annotations have also been uploaded to GenBank under accessions
- 24 JACSDY000000000 (*Vespula pensylvanica*), JACSDZ000000000 (*Vespula*

25 *germanica*) and JACSEA0000000000 (*Vespula vulgaris*). The annotations used in  
26 this paper and the annotations hosted by GenBank differ because records were  
27 removed to pass NCBI validation.

28 **Running head:** Genomes of invasive social wasps

29 **Keywords:** *Vespula germanica*, *Vespula pensylvanica*, *Vespula vulgaris*,  
30 Hymenoptera, social insects, genomes

31 **Correspondence:**

32 Peter K. Dearden

33 PO Box 56

34 Dunedin 9054

35 New Zealand

36 +64 3 479 7863

37 peter.dearden@otago.ac.nz

## Abstract

38 Social wasps of the genus *Vespula* have spread to nearly all landmasses worldwide and  
39 have become significant pests in their introduced ranges, affecting economies and  
40 biodiversity. Comprehensive genome assemblies and annotations for these species are  
41 required to develop the next generation of control strategies and monitor existing  
42 chemical control. We sequenced and annotated the genomes of the common wasp  
43 (*Vespula vulgaris*), German wasp (*Vespula germanica*), and the western yellowjacket  
44 (*Vespula pensylvanica*). Our chromosome-level *Vespula* assemblies each contain 176–  
45 179 Mb of total sequence assembled into 25 scaffolds, with 10–200 unanchored  
46 scaffolds, and 16,566–18,948 genes. We annotated gene sets relevant to the applied  
47 management of invasive wasp populations, including genes associated with  
48 spermatogenesis and development, pesticide resistance, olfactory receptors, immunity  
49 and venom. These genomes provide evidence for active DNA methylation in Vespidae  
50 and tandem duplications of venom genes. Our genomic resources will contribute to the  
51 development of next-generation control strategies, and monitoring potential resistance  
52 to chemical control.

## Introduction

Social wasps (Hymenoptera: Vespidae) are remarkable because their highly eusocial lifestyle appears to have evolved independently of other eusocial Hymenoptera (Piekarski *et al.* 2018; Hines *et al.* 2007; Peters *et al.* 2017). The eusocial lifestyle is characterized by overlapping generations of adults living together, cooperative care of offspring and reproductive division of labor (Michener and Brothers 1974). Along with their foraging flexibility and predatory ability, eusociality appears to play a major role in the ecological success of social wasps (Grimaldi and Engel 2005; Lester and Beggs 2019).

Vespid wasps can be effective pollinators of plants including ivy or orchid species (Cheng *et al.* 2009; Jacobs *et al.* 2010). They are generalist predators and effect biological control of some pest species (Donovan 2003). Their ecological success can also be problematic. Invasive colonies of common wasps (*Vespula vulgaris*) can contain up to 230,000 workers, while nests of the western yellowjacket (*Vespula pensylvanica*) containing up to half a million individuals have been observed (Lester and Beggs 2019). Colonies are smaller in the native ranges (Archer and Turner 2014 p. 61). In New Zealand's native beech forests, Vespid wasp populations can reach up to 40 nests per hectare and have a biomass similar to, or greater than, the combined biomasses of birds and mammals (Thomas *et al.* 1990; Lester *et al.* 2017). These invasive Vespid populations (Figure 1; Lester and Beggs 2019) have major impacts on ecosystems because of their large colony sizes, reproductive capacity and flexible predation.

Current control methods for vespid wasps are limited, with pesticides containing Fipronil being the most common and widespread chemical control method (Edwards *et al.* 2017). The use of this neurotoxic pesticide over large areas, and over consecutive years, may select for resistance. Fipronil resistance has been observed in other economically important insect pests (Matsumura *et al.* 2008; reviewed in Feyereisen *et al.* 2015). Next-generation pest control technologies, including gene drives, have been proposed as part of an alternative solution for controlling or eradicating invasive social wasps (Dearden *et al.* 2018). Targets for genetic modification include developmental genes associated with wasp fitness or fecundity (Lester and Beggs 2019; Dearden *et al.* 2018). Gene drives that have immune system targets have been proposed and developed

in the laboratory for other pests such as mosquitoes (*e.g.* Gantz *et al.* 2015; Dong *et al.* 2018). Detailed knowledge of *Vespula* genomes is required to ensure that targeted control of Vespids does not affect other beneficial Hymenoptera, such as honeybees or biocontrol agents.

Five genomes from the *Polistes* genus of the Vespidae subclade are available in the NCBI Genome database. Genomes for *Polistes canadensis* and *Polistes dominula* were assembled from short reads (Standage *et al.* 2016; Patalano *et al.* 2015). Genomes for *Polistes dorsalis*, *Polistes fuscatus* and *Polistes metricus*, based on Pacific Biosciences single-molecule sequencing, were recently deposited in the NCBI database (Miller *et al.* 2020). The three long-read genomes have improved contiguity compared to the short-read genomes produced for *P. dominula* and *P. canadensis* (Standage *et al.* 2016; Patalano *et al.* 2015; Miller *et al.* 2020). Here, we report chromosome-level assemblies for three *Vespula* species that are invasive pests in their introduced ranges (Figure 1; Lester and Beggs 2019). We undertake manual and computational annotation and phylogenomic analyses, with emphasis on sets of genes relevant to chemical control, olfaction, venom and DNA methylation. These resources will be useful for understanding the biology of vespid wasps, developing next-generation control strategies, and monitoring resistance to current chemical control.

**Figure 1:** *Vespula* species are major invasive pests. A–C. Examples of (A) *Vespula germanica*, (B) *Vespula pensylvanica* (Jon Sullivan, public domain), and (C) *Vespula vulgaris* (Sid Mosdell, CC BY 2.0). D. Native and invaded ranges of the sequenced *Vespula* species. E. Rooted phylogenetic species tree of *Vespula* species and other sequenced Hymenoptera, with *Drosophila melanogaster* as an outspecies, derived from 415 orthogroups using OrthoFinder.

## Materials and Methods

**Genome assembly and scaffolding:** *Vespula pensylvanica* samples were collected in Volcano, Hawaii in August, 2017 (lat 19.43, lon -155.21). The *V. pensylvanica* assembly was produced by Dovetail Genomics (<https://dovetailgenomics.com/>), starting with an Illumina library generated from a single haploid male to produce 115.5 Gb of 150bp paired-end reads. After trimming adaptors and removing bases or truncating reads with low quality scores, the draft genome was assembled with Meraculous 2.2.6 using a kmer size of 55, which produced the best fit with a constrained heterozygous model (Chapman *et al.* 2011, 2016). Chromosome-scale scaffolds were generated using Chicago and Dovetail Hi-C data from diploid females, and implemented using Dovetail's proprietary HiRise software.

We collected *V. vulgaris* samples in May 2015, from Richmond Hill Forest Park, New Zealand (lat -41.3292, lon 173.4637). *V. germanica* samples were collected in March 2018, from Lincoln, New Zealand (lat -43.6384, lon 172.4784). We used libraries generated from a single, haploid male to produce 17.1 Gb of 150b paired-end reads and 27.3 Gb of 125b paired-end reads, respectively. After trimming adaptor sequences, removing contaminants and verifying pairing with BBMap 38.00 (Bushnell 2014), we assembled draft genomes with Meraculous 2.2.6 (Chapman *et al.* 2011, 2016). We repeated assembly with a range of parameters, and used BUSCO analysis (Simão *et al.* 2015), assembly size and contiguity statistics to choose the best set of parameters for each dataset. Code for running and assessing the assemblies is hosted at <https://github.com/tomharrop/vger-illumina> and <https://github.com/tomharrop/vvul-illumina>. Draft genomes were scaffolded by Phase Genomics using Hi-C data generated from pools of 20 larvae. Chromatin conformation capture data was generated using Phase Genomics Proximo Hi-C 2.0 Kit, which is a commercially available version of the Hi-C protocol (Lieberman-Aiden *et al.* 2009). The Phase Genomics Proximo Hi-C genome scaffolding platform was used to create chromosome-scale scaffolds from the corrected assembly as described Bickhart *et al.* (2017).

**Genome curation:** Command-line arguments and scripts can be found at [https://github.com/jguhlin/vespula\\_paper](https://github.com/jguhlin/vespula_paper). Assembled genomes were cleared of

contamination by removing contigs which had BLAST taxonomy results which did not include *Polistes*, *Vespula*, or the word “wasp”. Contigs without BLAST results were kept if they contained predicted genes found in a Hymenoptera orthogroup from an initial gene prediction. Remaining contigs were only kept if they fell within 2 standard deviations of mean GC% of our kept contigs. The largest 25 chromosomes of *V. pensylvanica* were renamed to chromosomes and ordered according to size. Our *V. vulgaris* and *V. germanica* assemblies were aligned to the *V. pensylvanica* genome using D-GENIES, which inserts contigs into syntenic locations, flanked by 100 Ns, to assign chromosome names to those most related to those in *V. pensylvanica* (Cabanettes and Klopp 2018). Scaffolds were numbered with four digits in order of size.

**Repeat Masking:** Repeats were identified using RepeatModeler 2.0.1 ([github.com/ Dfam-consortium/ RepeatModeler](https://github.com/Dfam-consortium/RepeatModeler)) and RepeatMasker 4.0.9 ([repeatmasker.org/ RMDownload.html](https://repeatmasker.org/RMDownload.html)) via the funannotate pipeline (Palmer and Stajich 2019).

**RNA sequencing:** *Vespula vulgaris* queens, workers, and larvae were sampled from mature nests in the native range of Belgium and the introduced range in New Zealand and total RNA transcriptome data was generated as described by Gruber *et al.* (2019).

**Gene Prediction:** We performed iterative gene prediction using the Funannotate pipeline v1.6.0, manual annotation, and extrinsic protein evidence (Palmer and Stajich 2019). For *V. vulgaris* we used RNA-seq reads (described in the previous section) from *V. vulgaris* queens, workers, and larvae as additional evidence. The reads were trimmed with sickle ([github.com/ najoshi/ sickle](https://github.com/najoshi/sickle)) and aligned to our assembly using STAR in two-pass mode (Dobin *et al.* 2012). Gene predictions were performed on the assembly using funannotate predict with the RNA alignments and extrinsic protein of all *Vespula* proteins from NCBI, *Apis mellifera*, *Nasonia vitripennis*, and the UniProt SWISS-PROT database (Boutet *et al.* 2007; Pruitt *et al.* 2005; The Honeybee Genome Sequencing Consortium 2006; Werren *et al.* 2010).

Initial predictions were evaluated with GeneValidator with UniProt SWISS-PROT database as the high-quality targets. Genes whose protein predictions scored > 90 were used to train Augustus via the optimize\_augustus.pl script (Drăgan *et al.* 2016). The



prediction step of funannotate was then re-run as before using the trained vvulg  
AUGUSTUS species definition. This allowed retention of high-quality gene predictions  
from the target species to be used as a training set for AUGUSTUS gene prediction, a  
component of the funannotate pipeline. This process was repeated for *V. pensylvanica*  
and *V. germanica* using *V. vulgaris* species definition as the initial AUGUSTUS species  
in the first iteration of gene prediction, generating a species-specific configuration in the  
following round. This two-step gene prediction with validation and training using high-  
confidence gene calls between the first and second round allowed for the creation of  
species-specific AUGUSTUS models.

**Manual Curation:** Genes were manually curated in WebApollo (Lee *et al.* 2013).  
These manual annotations took precedence over intersecting computational gene  
predictions. Manual annotation was performed on *V. vulgaris* and lifted over to *V.*  
*germanica* and *V. pensylvanica* where possible.

**Gene Family Specific Predictions:** Gene-family specific predictions were enhanced  
using AUGUSTUS-PPX for the LGIC and Olfactory families (Keller *et al.* 2011). Protein  
sequences of interest from external sources were clustered based on bitscore using  
BLAST+ and MCL (Dongen 2000). Clusters of protein sequences were converted to  
protein profiles via AUGUSTUS tool msa2prfl.pl Assemblies were searched with  
fastBlockSearch and gene prediction was performed on matched regions with an  
additional flanking sequence of 1kbp. These predictions took precedence over  
intersecting computationally predicted genes.

**Annotation:** Further annotation was performed with funannotate using InterProScan  
5.32-71.0 (Mitchell *et al.* 2019). Genes were renamed using custom scripts. Protein  
predictions were compared with GeneValidator to both our Hymenoptera + *Drosophila*  
Protein Set and UniProt-SwissProt to generate GV scores and statistics. Proteins were  
compared with the publicly available genomes from Hymenoptera base using  
OrthoFinder (Elsik *et al.* 2016; Emms and Kelly 2019).

**Methylation analysis:** Nucleotide and dinucleotide content of gene body sequences  
were calculated using a custom perl script. CpG[o/e] was calculated as the number of  
CpG dinucleotides divided by the number of C nucleotides times the number of G

190 nucleotides. The number of components in CpG[o/e] distributions was estimated in R  
191 (R Core Team 2015) using mclust model-based clustering (Scrucca *et al.* 2016). The best  
192 fitting model was identified among several non-nested models using Bayesian  
193 information criteria (BIC).

194 **Data availability:** Raw sequence data are hosted in the NCBI Sequence Read Archive  
195 under accession PRJNA643352. Assembled genomes are available on GenBank under  
196 accessions JACSDY0000000000 (*Vespula pensylvanica*), JACSDZ0000000000 (*Vespula*  
197 *germanica*) and JACSEA0000000000 (*Vespula vulgaris*).

## Results and discussion

**Genome assemblies and annotation:** We used a combination of short-read Illumina sequencing and Hi-C scaffolding to assemble draft genomes for *V. germanica*, *V. pensylvanica*, and *V. vulgaris*. The genomes each contain 176–179 Mb of total sequence assembled into 25 superscaffolds (Figure 2A; Supplementary table 1; N<sub>50</sub> length 8.30–8.53 Mb), which likely represent the 25 chromosomes observed in *Vespula* karyotypes (Hoshiba *et al.* 1989). Each draft genome also contains 10–200 unanchored scaffolds (N<sub>50</sub> lengths 1.77–2.28 kb; Supplementary table 2). These genomes are similar in size to the genomes of the closely related European paper wasp, *Polistes dominula* (Standage *et al.* 2016), and the red paper wasp, *Polistes canadensis* (Patalano *et al.* 2015). However, the contiguity in our *Vespula* assemblies is higher than Illumina-based assemblies of other Vespidae, and comparable to the latest-generation *Apis mellifera* assembly (Table 1). We ordered and named scaffolds in the *Vespula* assemblies based on scaffold length in *V. pensylvanica*. The three genomes are highly syntenic, with evidence of some structural rearrangements (Figure 2B; Supplementary Figure 1). Repeat masking masked 17.86%, 18.75% and 18.71% of the *V. vulgaris*, *V. pensylvanica*, and *V. germanica* genomes, respectively. We predicted 16,751, 17,854, and 19,142 genes for *V. vulgaris*, *V. germanica*, and *V. pensylvanica*, respectively. We found between 92.2% and 96.0% of expected single-copy orthologs using BUSCO with the Hymenoptera lineage dataset (Simão *et al.* 2015). The contiguity of our *Vespula* assemblies and completeness of our annotations indicates that the combination of short-read sequencing and Hi-C scaffolding on haploid material is an effective strategy for assembling high-quality hymenopteran genomes.

**Figure 2:** A. Assembled chromosome lengths of the three *Vespula* species. Chromosomes were named in order of scaffold length in *V. pensylvanica*. B. Synteny between selected chromosomes of *Vespula* species. Chromosome 1 has minor translocations and inversions but is syntenic overall. Chromosome 2 has a larger inversion between *V. pensylvanica* and *V. germanica* that is not present in *V. vulgaris*. Chromosome 5 has a large translocation and inversion between *V. pensylvanica* and *V. germanica*. Chromosome lengths are scaled to chromosome 1. C. Shared orthogroups between the three *Vespula* species. D. The pan and core genomes of Hymenoptera. The core genome (the set of orthogroups present in all genomes sampled) decreases as more genomes are sampled, to a final core genome size of 3,092

orthogroups when all 26 hymenopteran genomes are compared. The pan genome (the total set of orthogroups present in one or more of the sampled genomes) does not continue to grow as more genomes are sampled, indicating a closed hymenopteran pan genome. We predicted more orthogroups (13,141–14,022 orthogroups) in *Vespula* species than in non-*Vespula* Hymenoptera (median 9,193 orthogroups), which resulted in a larger pan genome in comparisons that included *Vespula* species. We only analysed genes that were assigned to an orthogroup, and we plotted a random subset of 1,000 comparisons for genome numbers that resulted in more than 1000 comparisons.

221 To predict divergence time of *Vespula* species from *Polistes*, we reconstructed a  
222 phylogeny using other hymenopteran genomes. Based on a published estimate for  
223 Hymenoptera (Peters *et al.* 2017), the divergence time of Vespid wasps from their last  
224 common ancestor with *P. dominula* is estimated to be 51 million years (95% CI: 34–71  
225 million years). Mitochondrial genomes suggest a divergence time of 75 mya (Huang *et*  
226 *al.* 2019). Scaling our ultrametric phylogenetic tree to the former estimate places  
227 separation of *V. vulgaris* from *V. pensylvanica* and *V. germanica* at ~6 mya, and  
228 separation of *V. pensylvanica* from *V. germanica* at 4.5 mya (Supplementary Figure 2).

229 We manually curated 361 gene models in *V. vulgaris* and used these curations to  
230 improve automated prediction steps for the other two species. During manual curation,  
231 we focused on a range of gene sets relevant to the evolution and applied management of  
232 invasive *Vespula* spp., including olfactory receptors, pesticide resistance, immunity and  
233 viruses, venom, and spermatogenesis and development.

234 To investigate relationships between *Vespula* genes, we clustered predicted proteins  
235 into orthogroups with predicted proteins from the Hymenoptera Genome Database,  
236 using *Drosophila melanogaster* as the outgroup (Hoskins *et al.* 2015; Elsik *et al.* 2016;  
237 Emms and Kelly 2019). Each orthogroup contains a set of genes putatively descended  
238 from a single gene in the last common ancestor of the species represented in the  
239 orthogroup. Between 82.6% and 88.4% of our predicted *Vespula* proteins belonged to  
240 orthogroups. *V. vulgaris* shares 12,560 and 12,084 orthogroups with *V. pensylvanica*  
241 and *V. germanica*, respectively, and *V. pensylvanica* shares 13,209 orthogroups with *V.*  
242 *germanica* (Figure 2C). Orthogroups including other hymenopteran species allowed us  
243 to predict the core- and pan-genomes for Hymenoptera (Figure 2D). This analysis  
244 suggests that Hymenoptera have a closed pan-genome, because as we include more

245 genomes the rate of discovery of new orthogroups decreases. We also observed more  
246 orthogroups in our Vespidae genomes than in other Hymenoptera, which could indicate  
247 over-prediction resulting from our annotation.

248 93.7–94.5% of predicted orthogroups in *P. canadensis* and 96.0–96.9% in *P. dominula*  
249 clustered into orthogroups that contained a *Vespula* gene. *Vespula* genes from our  
250 annotations were present in 58.9–62.8% of all orthogroups across Hymenoptera and *D.*  
251 *melanogaster*, compared to 40.7% and 41.2% for *P. canadensis* and *P. dominula*,  
252 respectively (Table 2). Other hymenopteran species had a member in 37.8–44.5% of  
253 orthogroups, indicating gene families may be missing in predictions amongst other  
254 species. Most of the genes we annotated were in shared orthogroups, with 100 genes  
255 (0.1–0.4% per species) in species-specific orthogroups. Other hymenopteran genomes  
256 had 0–3.3% (mean 1.7%) of genes in species-specific orthogroups. Our annotation  
257 results suggest that Vespidae wasps have more genes than other Hymenoptera and/or  
258 gene annotation in other Hymenoptera is incomplete. This could be resolved by re-  
259 annotation of other hymenopteran genes using a comparative approach.

**Table 1.** Comparison of Vespidae and honeybee genome assemblies.

1. Apart from our *Vespula* assemblies, repeat content was counted from lowercase nucleotides in published assemblies.

| Species                     | Sequencing Strategy                 | Total Sequence | Largest    | Scaffolds/contigs | $N_{50}$ length | Ns         | Gaps   | Repeats <sup>1</sup> (%) |
|-----------------------------|-------------------------------------|----------------|------------|-------------------|-----------------|------------|--------|--------------------------|
| <i>Vespula vulgaris</i>     | Illumina + Hi-C                     | 176,275,134    | 19,426,332 | 35                | 8,304,510       | 4,147,610  | 49,679 | 17.12                    |
| <i>Vespula germanica</i>    | Illumina + Hi-C                     | 178,312,246    | 19,524,135 | 133               | 8,396,154       | 1,783,864  | 18,963 | 18.89                    |
| <i>Vespula pensylvanica</i> | Illumina + Hi-C                     | 179,379,562    | 19,704,315 | 225               | 8,532,720       | 444,951    | 4,987  | 19.15                    |
| <i>Polistes canadensis</i>  | Illumina                            | 211,202,212    | 3,185,661  | 3,836             | 521,566         | 14,106,256 | 15,755 | 41.60                    |
| <i>Polistes dominula</i>    | Illumina                            | 208,026,220    | 7,126,315  | 1,483             | 1,625,592       | 7,426,626  | 14,286 | 44.50                    |
| <i>Polistes fuscatus</i>    | PacBio Sequel + Illumina + Dovetail | 219,116,742    | 19,629,704 | 187               | 9,116,088       | 4,436,170  | 1,204  | 41.79                    |
| <i>Polistes metricus</i>    | PacBio Sequel + Illumina            | 219,838,961    | 15,979,625 | 216               | 4,634,047       | 1,082,619  | 459    | 44.48                    |

|                              |                                        |             |            |       |            |            |        |       |
|------------------------------|----------------------------------------|-------------|------------|-------|------------|------------|--------|-------|
| <i>Polistes dorsalis</i>     | 10x Genomics                           | 209,288,276 | 20,305,868 | 5,129 | 5,372,633  | 4,771,923  | 8,816  | 42.35 |
| <i>Apis mellifera</i> 4.5    | Sanger + SOLiD + 454                   | 250,287,000 | 29,893,408 | 5,321 | 13,219,345 | 21,165,099 | 12,690 | 5.28  |
| <i>Apis mellifera</i> HAv3.1 | PacBio + 10x Chromium + BioNano + Hi-C | 225,250,884 | 27,754,200 | 177   | 13,619,445 | 1,313,614  | 51     | 44.63 |

**Table 2.** Gene content and orthogroup representation for selected hymenopteran genomes

| Species                     | Genes  | Orthogroups represented | Orthogroups represented (%) | Species-specific orthogroups | Genes in species-specific orthogroups (%) |
|-----------------------------|--------|-------------------------|-----------------------------|------------------------------|-------------------------------------------|
| <i>Vespula vulgaris</i>     | 16,751 | 13,141                  | 58.9%                       | 5                            | 0.1%                                      |
| <i>Vespula germanica</i>    | 17,854 | 13,739                  | 61.6%                       | 23                           | 0.4%                                      |
| <i>Vespula pensylvanica</i> | 19,142 | 14,022                  | 62.8%                       | 11                           | 0.1%                                      |
| <i>Polistes canadensis</i>  | 10,518 | 9,086                   | 40.7%                       | 12                           | 0.2%                                      |
| <i>Polistes dominula</i>    | 11,069 | 9,193                   | 41.2%                       | 27                           | 0.5%                                      |
| <i>Apis mellifera</i>       | 14,064 | 8,949                   | 40.1%                       | 35                           | 0.6%                                      |
| <i>Nasonia vitripennis</i>  | 14,647 | 8,901                   | 39.9%                       | 616                          | 11.1%                                     |

**Evidence for active DNA methylation:** DNA methylation has been functionally linked to caste specification in honeybees and ants and division of labour in honeybees (Herb *et al.* 2012; Kucharski *et al.* 2008; Bonasio *et al.* 2012; Guan *et al.* 2013). DNA methylation may be integral for aspects of eusociality (reviewed by Li-Byarlay *et al.* 2013), although recent studies have found no consistent link (Bewick *et al.* 2017; Glastad *et al.* 2017). In mice, Dnmt3 enzymes catalyse *de novo* DNA methylation (reviewed by Goll and Bestor 2005). The genomes of *P. canadensis* and *P. dominula* do not encode a *Dnmt3* homolog (Standage *et al.* 2016; Patalano *et al.* 2015; Bewick *et al.* 2017). While this manuscript was under revision, three new *Polistes* genomes based on Pacific Biosciences single-molecule sequencing were deposited in the NCBI database



270 (Miller *et al.* 2020). Genome annotations have not been reported for the three long-read  
271 *Polistes* genomes. We aligned all five *Polistes* genomes and our *V. germanica* and *V.*  
272 *pensylvanica* genomes against Chr11 of the *V. vulgaris* genome, which encodes *Dnmt3*.  
273 None of the five *Polistes* genomes have a region that aligns to the *V. vulgaris* *Dnmt3*  
274 locus (Figure 3A). All three *Vespula* genomes encode an ortholog of *Dnmt3*, indicating  
275 the lack of this gene in *Polistes* is due to gene loss following the divergence from the  
276 *Vespula* lineage approximately 50 million years ago (Peters *et al.* 2017). *Vespula*  
277 genomes all contain a single ortholog of *Dnmt1* and have evidence of active DNA  
278 methylation (Figure 3B; Supplementary Figure 3).

279

**Figure 3:** A. Whole-genome alignments of five *Polistes* genomes and genomes of *V. germanica* and *V. pensylvanica* against Chr11 of the *V. vulgaris* genome, showing the *Dnmt3* locus (Vvulg11g01820) and the surrounding 20 kb. None of the *Polistes* assemblies have regions that are homologous to the *Dnmt3* locus, and annotations for *P. canadensis* and *P. dominula* do not include a *Dnmt3* gene (Standage *et al.* 2016; Patalano *et al.* 2015; Bewick *et al.* 2017). All three *Vespula* genomes encode an ortholog of *Dnmt3*. B. Frequency histograms of CpG[o/e] observed in coding sequences of *Drosophila melanogaster*, *Apis mellifera*, *Polistes canadensis*, *Polistes dominula*, *Vespula germanica*, *Vespula vulgaris* and *Vespula pensylvanica*. The y-axis depicts the number of genes (Frequency) with CpG[o/e] values given on the x-axis. The distribution of CpG[o/e] in Vespid wasps is a trimodal distribution, with a low-CpG[o/e] peak consistent with the presence of historical DNA methylation in all five Vespid species.

280 **Gene content and duplications:** We found more olfactory receptor (OR) genes in  
281 *Vespula* genomes than in the genomes of *P. dominula* and *P. canadensis*. We predicted  
282 120 OR genes in *V. vulgaris*, 133 in *V. germanica*, and 102 in *V. pensylvanica*.  
283 Annotations for *P. dominula* and *P. canadensis* contain 94 and 72 OR genes,  
284 respectively (Supplementary table 3). In contrast, honeybee and *Nasonia* genomes  
285 encode more OR genes (170 and 301, respectively; Robertson and Wanner 2006;  
286 Robertson *et al.* 2010). *Vespula* OR genes cluster into 28 orthogroups. The co-receptor  
287 *Orco* is present in all genomes and forms a stable orthogroup (orthogroup 7148; Larsson  
288 *et al.* 2004; Jones *et al.* 2005). In contrast, there are significant expansions of particular  
289 OR orthogroups in the Vespid wasps, and these differ from the groups expanded in  
290 *Nasonia* and honeybee (Robertson *et al.* 2010; Robertson and Wanner 2006).  
291 Orthogroup members are arranged in expanded tandem arrays on chromosome 3

(orthogroup 51), chromosome 13 (orthogroup 2434), and chromosome 25 (orthogroup 232) of *Vespula* genomes. In these clusters, numbers of genes vary between species, implying that duplications and deletions are recent and ongoing. Variation in olfactory receptors between wasp species, and between these wasps and other Hymenoptera may indicate species-specific olfactory biology. These may be key to understanding the social behavior and pheromone signalling systems present within these species.

Some of the genes encoding venom components also have variable copy numbers in *Vespula* genomes. The major allergens in *Vespula* venoms are phospholipase A1, hyaluronidase, and antigen 5 (Biló *et al.* 2005; Kolarich *et al.* 2007). Phospholipase A1 is found in three tandem copies in the *P. dominula* and *V. germanica* genomes (chromosome 9 in *Vespula*), and one copy in each of the *V. pensylvanica* and *V. vulgaris* genomes. The phylogenetic placement of these duplicates in *Polistes* and *V. germanica* (Figure 4) implies that these are independent amplifications. The hyaluronidase gene is duplicated in our three Vespid genomes, but not *P. dominula*. These are tandem duplications that appear to have been present in the last common ancestor of the three *Vespula* species. *P. dominula*, *V. germanica* and *V. pensylvanica* also have two Antigen 5 genes, but these duplications appear ancient before the common ancestor of Hymenoptera (Figure 4). In *Vespula* species, one copy is on chromosome 6 and one is on chromosome 7. In *V. vulgaris*, the chromosome 6 copy is absent. Duplication of venom genes in Vespids is no surprise, given the importance of venom to their life cycle.

**Figure 4:** Ultrametric trees of orthogroups encoding Vespid wasp venom components in insect genomes. A. Venom hyaluronidase phylogram showing duplication of this gene in Vespid genomes but not *Polistes*. B. Venom phospholipase phylogram indicating independent duplications in *Polistes* and *V. germanica* from a single gene ancestor in related Hymenoptera. C. Antigen 5 phylogram, indicating two genes encode this venom component in Vespid wasps and *Polistes*, one of which is missing from the *V. vulgaris* genome.

Primary chemical control of *Vespula* populations is through the use of baits containing a low concentration of the phenylpyrazole insecticide, Fipronil (Lester and Beggs 2019; Edwards *et al.* 2017). We used targeted prediction to identify ligand-gated ion channel

(LGIC), olfactory receptor, and spermatogenesis genes using Augustus protein profiles (Keller *et al.* 2011). Our annotation of the Fipronil target site, the GABA receptor *Resistant to dieldrin (Rdl)*, did not suggest the presence of the classical Ala301 mutation that confers high resistance. *Vespula* LGIC receptors are highly conserved, with one-to-one orthology in *Apis* and *Bombus* (Supplementary Figure 4). This suggests that any chemicals targeting *Vespula* LGICs will also affect bees, as is the case with Fipronil.

**Conclusions:** We have produced chromosome-level genome assemblies for three invasive social wasps in the *Vespula* genus. Our approach of short-read sequencing and Hi-C scaffolding using haploid material allowed us to produce assemblies that exceed the genome quality targets suggested by the i5k insect genome sequencing initiative (scaffold  $N_{50}$  length > 300 kb; Richards and Murali 2015). Using manual curation and computational prediction, we have identified genes that may encode specific biology suitable for targeting with next-generation control technologies, and genes that may be affected by selection by current chemical controls.

These are the first three genomes from this branch of the Aculeata subclade, which will be useful in phylogenetic comparisons of the remarkable life history characteristics of Hymenoptera. In particular, these genomes will be valuable for understanding the evolution of eusociality, which has appeared twice in Vespid wasps independently of other Hymenoptera (Piekarski *et al.* 2018; Hines *et al.* 2007; Peters *et al.* 2017). Comparing the genomes of Vespid wasps, which are highly eusocial, with the closely related paper wasps, which are primitively eusocial, may also help our understanding of how evolution elaborates mechanisms of colonial living.

Vespid wasps are significant invasive pests in many parts of the world. These genomes will be of major importance for applied management of *Vespula*, in programs using both chemical control methods and for next-generation applications. Our assemblies will provide species-specific targets for novel control methods, such as RNA interference, gene drives and the deployment of damaging viruses. The genomic resources we have developed will also be essential for monitoring the effects of next-generation control methods and measuring genomic variation across natural populations.

**Acknowledgements:** The authors would like to thank P.M. Dearden for editing the manuscript. New Zealand Genomics Ltd for generation of the Common wasp genome data. The New Zealand National eScience Infrastructure (NeSI) for computational support. D. Hart and D.J. Champion for I.T. support and B.P. Dearden for critical discussions in the formulation of this work.

**Funding:** This project was supported by Genomics Aotearoa (to PKD) and the Biological Heritage National Science Challenge (to P.J.L.) both funded by the Ministry of Business Innovation and Employment (Hikina Whakatutuki), Government of New Zealand, as well as a US National Science Foundation Grant and UC Riverside Seed Grant to JP and EWR; Dovetail Genomics Matching Funds Grant to JP.

**Availability of data and materials:** Raw reads are hosted in the NCBI Sequence Read Archive under accession PRJNA643352. The version of the assemblies and annotations described in this paper is hosted on Zenodo under DOI [10.5281/zenodo.4001020](https://doi.org/10.5281/zenodo.4001020). The genomes and annotations have also been uploaded to GenBank under accessions JACSDY000000000 (*Vespula pensylvanica*), JACSDZ000000000 (*Vespula germanica*) and JACSEA000000000 (*Vespula vulgaris*). The annotations used in this paper and the annotations hosted by GenBank differ because records were removed to pass NCBI validation.

**Authors' contributions:** P.J.L. and P.K.D. conceived and designed the project. All authors aided in obtaining and analysing the genomic data. P.J.L., P.K.D., T.W.R.H., J.G., E.J.R. and E.J.D. wrote the manuscript draft, and all authors participated in the revision of the final version. All authors read and approved the final manuscript.

**Ethics approval and consent to participate:** Common wasps (*V. vulgaris*) were collected under the permit National Authorisation Number 38337-RES from the Department of Conservation in New Zealand. Samples of other wasps were collected from private land where no permit was required. No other ethical approval was required.

**Consent for publication:** Not applicable.

374 **Competing interests:** The authors declare that they have no competing interests.

## Literature Cited

Archer, M. E., and J. Turner, 2014 *Handbooks for the identification of British insects. Vol. 6, Pt. 6: The vespoid wasps (Tiphidae, Mutillidae, Sapygidae, Scoliidae and Vespidae) of the British Isles*. Field Studies Council, Telford.

Bewick, A. J., K. J. Vogel, A. J. Moore, and R. J. Schmitz, 2017 Evolution of DNA Methylation across Insects. *Mol Biol Evol* 34: 654–665. doi: [10.1093/molbev/msw264](https://doi.org/10.1093/molbev/msw264).

Bickhart, D. M., B. D. Rosen, S. Koren, B. L. Sayre, and A. R. Hastie *et al.*, 2017 Single-molecule sequencing and chromatin conformation capture enable de novo reference assembly of the domestic goat genome. *Nat Genet* 49: 643–650. doi: [10.1038/ng.3802](https://doi.org/10.1038/ng.3802).

Biló, B. M., F. Rueff, H. Mosbech, F. Bonifazi, and J. N. G. Oude- Elberink, 2005 Diagnosis of Hymenoptera venom allergy. *Allergy* 60: 1339–1349. doi: [10.1111/j.1398-9995.2005.00963.x](https://doi.org/10.1111/j.1398-9995.2005.00963.x).

Bonasio, R., Q. Li, J. Lian, N. Mutti, and L. Jin *et al.*, 2012 Genome-wide and Caste-Specific DNA Methylomes of the Ants *Camponotus floridanus* and *Harpegnathos saltator*. *Current Biology* 22: 1755–1764. doi: [10.1016/j.cub.2012.07.042](https://doi.org/10.1016/j.cub.2012.07.042).

Boutet, E., D. Lieberherr, M. Tognolli, M. Schneider, and A. Bairoch, 2007 UniProtKB/Swiss-Prot, pp. 89–112 in *Plant Bioinformatics: Methods and Protocols*, edited by D. Edwards. *Methods in Molecular Biology*<sup>TM</sup>, Humana Press, Totowa, NJ. doi: [10.1007/978-1-59745-535-0\\_4](https://doi.org/10.1007/978-1-59745-535-0_4).

Bushnell, B., 2014 BBMap: A Fast, Accurate, Splice-Aware Aligner: Lawrence Berkeley National Lab. (LBNL), Berkeley, CA (United States) LBNL-7065E.

Cabanettes, F., and C. Klopp, 2018 D-GENIES: dot plot large genomes in an interactive, efficient and simple way. *PeerJ* 6: e4958. doi: [10.7717/peerj.4958](https://doi.org/10.7717/peerj.4958).

Chapman, J. A., I. Y. Ho, E. Goltsman, and D. S. Rokhsar, 2016 Meraculous2: fast accurate short-read assembly of large polymorphic genomes. arXiv:1608.01031 [cs, q-bio].

Chapman, J. A., I. Ho, S. Sunkara, S. Luo, and G. P. Schroth *et al.*, 2011 Meraculous: De Novo Genome Assembly with Short Paired-End Reads. *PLOS ONE* 6: e23501. doi: [10.1371/journal.pone.0023501](https://doi.org/10.1371/journal.pone.0023501).

Cheng, J., J. Shi, F.-Z. Shangguan, A. Dafni, and Z.-H. Deng *et al.*, 2009 The pollination of a self-incompatible, food-mimic orchid, *Coelogyne fimbriata* (Orchidaceae), by female *Vespula* wasps. *Ann Bot* 104: 565–571. doi: [10.1093/aob/mcp029](https://doi.org/10.1093/aob/mcp029).

Dearden, P. K., N. J. Gemmell, O. R. Mercier, P. J. Lester, and M. J. Scott *et al.*, 2018 The potential for the use of gene drives for pest control in New Zealand: a perspective. *Journal of the Royal Society of New Zealand* 48: 225–244. doi: [10.1080/03036758.2017.1385030](https://doi.org/10.1080/03036758.2017.1385030).

- Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, and C. Zaleski *et al.*, 2012 STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* (Oxford, England) 29: 15–21. doi: [10.1093/bioinformatics/bts635](https://doi.org/10.1093/bioinformatics/bts635).
- Dongen, S. van, 2000 Graph clustering by flow simulation: University of Utrecht.
- Dong, Y., M. L. Simoes, E. Marois, and G. Dimopoulos, 2018 CRISPR/Cas9-mediated gene knockout of *Anopheles gambiae* FREP1 suppresses malaria parasite infection. *PLoS Pathog* 14: e1006898. doi: [10.1371/journal.ppat.1006898](https://doi.org/10.1371/journal.ppat.1006898).
- Donovan, B. J., 2003 Potential manageable exploitation of social wasps, *Vespula* spp. (Hymenoptera: Vespidae), as generalist predators of insect pests. *International Journal of Pest Management* 49: 281–285. doi: [10.1080/0967087031000123698](https://doi.org/10.1080/0967087031000123698).
- Drăgan, M.-A., I. Moghul, A. Priyam, C. Bustos, and Y. Wurm, 2016 GeneValidator: identify problems with protein-coding gene predictions. *Bioinformatics* 32: 1559–1561. doi: [10.1093/bioinformatics/btw015](https://doi.org/10.1093/bioinformatics/btw015).
- Edwards, E., R. Toft, N. Joice, and I. Westbrooke, 2017 The efficacy of Vespex® wasp bait to control *Vespula* species (Hymenoptera: Vespidae) in New Zealand. *International Journal of Pest Management* 63: 266–272. doi: [10.1080/09670874.2017.1308581](https://doi.org/10.1080/09670874.2017.1308581).
- Elsik, C. G., A. Tayal, C. M. Diesh, D. R. Unni, and M. L. Emery *et al.*, 2016 Hymenoptera Genome Database: integrating genome annotations in HymenopteraMine. *Nucleic Acids Res* 44: D793–D800. doi: [10.1093/nar/gkv1208](https://doi.org/10.1093/nar/gkv1208).
- Emms, D. M., and S. Kelly, 2019 OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biology* 20: 238. doi: [10.1186/s13059-019-1832-y](https://doi.org/10.1186/s13059-019-1832-y).
- Feyereisen, R., W. Dermauw, and T. Van Leeuwen, 2015 Genotype to phenotype, the molecular and physiological dimensions of resistance in arthropods. *Pesticide Biochemistry and Physiology* 121: 61–77. doi: [10.1016/j.pestbp.2015.01.004](https://doi.org/10.1016/j.pestbp.2015.01.004).
- Gantz, V. M., N. Jasinskiene, O. Tatarenkova, A. Fazekas, and V. M. Macias *et al.*, 2015 Highly efficient Cas9-mediated gene drive for population modification of the malaria vector mosquito *Anopheles stephensi*. *PNAS* 112: E6736–E6743. doi: [10.1073/pnas.1521077112](https://doi.org/10.1073/pnas.1521077112).
- Glastad, K. M., S. V. Arsenault, K. L. Vertacnik, S. M. Geib, and S. Kay *et al.*, 2017 Variation in DNA Methylation Is Not Consistently Reflected by Sociality in Hymenoptera. *Genome Biol Evol* 9: 1687–1698. doi: [10.1093/gbe/evx128](https://doi.org/10.1093/gbe/evx128).
- Goll, M. G., and T. H. Bestor, 2005 Eukaryotic cytosine methyltransferases. *Annu. Rev. Biochem.* 74: 481–514. doi: [10.1146/annurev.biochem.74.010904.153721](https://doi.org/10.1146/annurev.biochem.74.010904.153721).
- Grimaldi, D. A., and M. S. Engel, 2005 *Evolution of the insects*. Cambridge University Press, Cambridge [U.K.] ; New York.
- Gruber, M. A. M., O. Quinn, J. W. Baty, J. Dobelmann, and J. Haywood *et al.*, 2019 Fitness and microbial networks of the common wasp, *Vespula vulgaris* (Hymenoptera:

Vespidae), in its native and introduced ranges. *Ecological Entomology* 44: 512–523. doi: [10.1111/een.12732](https://doi.org/10.1111/een.12732).

Guan, C., A. B. Barron, X. J. He, Z. L. Wang, and W. Y. Yan *et al.*, 2013 A Comparison of Digital Gene Expression Profiling and Methyl DNA Immunoprecipitation as Methods for Gene Discovery in Honeybee (*Apis mellifera*) Behavioural Genomic Analyses. *PLOS ONE* 8: e73628. doi: [10.1371/journal.pone.0073628](https://doi.org/10.1371/journal.pone.0073628).

Herb, B. R., F. Wolschin, K. D. Hansen, M. J. Aryee, and B. Langmead *et al.*, 2012 Reversible switching between epigenetic states in honeybee behavioral subcastes. *Nature Neuroscience* 15: 1371–1373. doi: [10.1038/nn.3218](https://doi.org/10.1038/nn.3218).

Hines, H. M., J. H. Hunt, T. K. O'Connor, J. J. Gillespie, and S. A. Cameron, 2007 Multigene phylogeny reveals eusociality evolved twice in vespid wasps. *PNAS* 104: 3295–3299. doi: [10.1073/pnas.0610140104](https://doi.org/10.1073/pnas.0610140104).

Hoshiba, H., M. Matsuura, and H. T. Imai, 1989 Karyotype evolution in the social wasps. *Jpn J Genet.*, 遺伝学雑誌 64: 209–222. doi: [10.1266/jjg.64.209](https://doi.org/10.1266/jjg.64.209).

Hoskins, R. A., J. W. Carlson, K. H. Wan, S. Park, and I. Mendez *et al.*, 2015 The Release 6 reference sequence of the *Drosophila melanogaster* genome. *Genome Res.* 25: 445–458. doi: [10.1101/gr.185579.114](https://doi.org/10.1101/gr.185579.114).

Huang, P., J. M. Carpenter, B. Chen, and T.-J. Li, 2019 The first divergence time estimation of the subfamily Stenogastrinae (Hymenoptera: Vespidae) based on mitochondrial phylogenomics. *International Journal of Biological Macromolecules* 137: 767–773. doi: [10.1016/j.ijbiomac.2019.06.239](https://doi.org/10.1016/j.ijbiomac.2019.06.239).

Jacobs, J. H., S. J. Clark, I. Denholm, D. Goulson, and C. Stoate *et al.*, 2010 Pollinator effectiveness and fruit set in common ivy, *Hedera helix* (Araliaceae). *Arthropod-Plant Interactions* 4: 19–28. doi: [10.1007/s11829-009-9080-9](https://doi.org/10.1007/s11829-009-9080-9).

Jones, W. D., T.-A. T. Nguyen, B. Kloss, K. J. Lee, and L. B. Vosshall, 2005 Functional conservation of an insect odorant receptor gene across 250 million years of evolution. *Current Biology* 15: R119–R121. doi: [10.1016/j.cub.2005.02.007](https://doi.org/10.1016/j.cub.2005.02.007).

Keller, O., M. Kollmar, M. Stanke, and S. Waack, 2011 A novel hybrid gene prediction method employing protein multiple sequence alignments. *Bioinformatics* 27: 757–763. doi: [10.1093/bioinformatics/btr010](https://doi.org/10.1093/bioinformatics/btr010).

Kolarich, D., A. Loos, R. Léonard, L. Mach, and G. Marzban *et al.*, 2007 A proteomic study of the major allergens from yellow jacket venoms. *Proteomics* 7: 1615–1623. doi: [10.1002/pmic.200600800](https://doi.org/10.1002/pmic.200600800).

Kucharski, R., J. Maleszka, S. Foret, and R. Maleszka, 2008 Nutritional Control of Reproductive Status in Honeybees via DNA Methylation. *Science* 319: 1827–1830. doi: [10.1126/science.1153069](https://doi.org/10.1126/science.1153069).



Larsson, M. C., A. I. Domingos, W. D. Jones, M. E. Chiappe, and H. Amrein *et al.*, 2004 Or83b Encodes a Broadly Expressed Odorant Receptor Essential for *Drosophila* Olfaction. *Neuron* 43: 703–714. doi: [10.1016/j.neuron.2004.08.019](https://doi.org/10.1016/j.neuron.2004.08.019).

Lee, E., G. A. Helt, J. T. Reese, M. C. Munoz-Torres, and C. P. Childers *et al.*, 2013 Web Apollo: a web-based genomic annotation editing platform. *Genome Biology* 14: R93. doi: [10.1186/gb-2013-14-8-r93](https://doi.org/10.1186/gb-2013-14-8-r93).

Lester, P. J., and J. R. Beggs, 2019 Invasion Success and Management Strategies for Social *Vespula* Wasps. *Annual Review of Entomology* 64: 51–71. doi: [10.1146/annurev-ento-011118-111812](https://doi.org/10.1146/annurev-ento-011118-111812).

Lester, P. J., J. Haywood, M. E. Archer, and C. R. Shortall, 2017 The long-term population dynamics of common wasps in their native and invaded range. *Journal of Animal Ecology* 86: 337–347. doi: [10.1111/1365-2656.12622](https://doi.org/10.1111/1365-2656.12622).

Li-Byarlay, H., Y. Li, H. Stroud, S. Feng, and T. C. Newman *et al.*, 2013 RNA interference knockdown of *DNA methyl-transferase 3* affects gene alternative splicing in the honey bee. *PNAS* 110: 12750–12755. doi: [10.1073/pnas.1310735110](https://doi.org/10.1073/pnas.1310735110).

Lieberman-Aiden, E., N. L. van Berkum, L. Williams, M. Imakaev, and T. Ragoczy *et al.*, 2009 Comprehensive mapping of long range interactions reveals folding principles of the human genome. *Science* 326: 289–293. doi: [10.1126/science.1181369](https://doi.org/10.1126/science.1181369).

Matsumura, M., H. Takeuchi, M. Satoh, S. Sanada-Morimura, and A. Otuka *et al.*, 2008 Species-specific insecticide resistance to imidacloprid and fipronil in the rice planthoppers *Nilaparvata lugens* and *Sogatella furcifera* in East and South-east Asia. *Pest Manag. Sci.* 64: 1115–1121. doi: [10.1002/ps.1641](https://doi.org/10.1002/ps.1641).

Michener, C. D., and D. J. Brothers, 1974 Were workers of eusocial hymenoptera initially altruistic or oppressed? *Proc Natl Acad Sci U S A* 71: 671–4. doi: [10.1073/pnas.71.3.671](https://doi.org/10.1073/pnas.71.3.671).

Miller, S. E., A. W. Legan, M. T. Henshaw, K. L. Ostevik, and K. Samuk *et al.*, 2020 Evolutionary dynamics of recent selection on cognitive abilities. *PNAS* 117: 3045–3052. doi: [10.1073/pnas.1918592117](https://doi.org/10.1073/pnas.1918592117).

Mitchell, A. L., T. K. Attwood, P. C. Babbitt, M. Blum, and P. Bork *et al.*, 2019 InterPro in 2019: improving coverage, classification and access to protein sequence annotations. *Nucleic Acids Res* 47: D351–D360. doi: [10.1093/nar/gky1100](https://doi.org/10.1093/nar/gky1100).

Palmer, J., and J. Stajich, 2019 *nextgenusfs/funannotate: funannotate v1.6.0*. Zenodo. doi: [10.5281/zenodo.3354704](https://doi.org/10.5281/zenodo.3354704).

Patalano, S., A. Vlasova, C. Wyatt, P. Ewels, and F. Camara *et al.*, 2015 Molecular signatures of plastic phenotypes in two eusocial insect species with simple societies. *Proc Natl Acad Sci USA* 112: 13970–13975. doi: [10.1073/pnas.1515937112](https://doi.org/10.1073/pnas.1515937112).

Peters, R. S., L. Krogmann, C. Mayer, A. Donath, and S. Gunkel *et al.*, 2017 Evolutionary History of the Hymenoptera. *Current Biology* 27: 1013–1018. doi: [10.1016/j.cub.2017.01.027](https://doi.org/10.1016/j.cub.2017.01.027).

Piekarski, P. K., J. M. Carpenter, A. R. Lemmon, E. Moriarty Lemmon, and B. J. Sharanowski, 2018 Phylogenomic evidence overturns current conceptions of social evolution in wasps (Vespidae). *Mol Biol Evol* 35: 2097–2109. doi: [10.1093/molbev/msy124](https://doi.org/10.1093/molbev/msy124).

Pruitt, K. D., T. Tatusova, and D. R. Maglott, 2005 NCBI Reference Sequence (RefSeq): a curated non-redundant sequence database of genomes, transcripts and proteins. *Nucleic Acids Res* 33: D501–D504. doi: [10.1093/nar/gki025](https://doi.org/10.1093/nar/gki025).

R Core Team, 2015 R: A Language and Environment for Statistical Computing.

Richards, S., and S. C. Murali, 2015 Best practices in insect genome sequencing: what works and what doesn't. *Current Opinion in Insect Science* 7: 1–7. doi: [10.1016/j.cois.2015.02.013](https://doi.org/10.1016/j.cois.2015.02.013).

Robertson, H. M., J. Gadau, and K. W. Wanner, 2010 The insect chemoreceptor superfamily of the parasitoid jewel wasp *Nasonia vitripennis*. *Insect Molecular Biology* 19: 121–136. doi: [10.1111/j.1365-2583.2009.00979.x](https://doi.org/10.1111/j.1365-2583.2009.00979.x).

Robertson, H. M., and K. W. Wanner, 2006 The chemoreceptor superfamily in the honey bee, *Apis mellifera*: Expansion of the odorant, but not gustatory, receptor family. *Genome Res.* 16: 1395–1403. doi: [10.1101/gr.5057506](https://doi.org/10.1101/gr.5057506).

Scrucca, L., M. Fop, T. B. Murphy, and A. E. Raftery, 2016 mclust 5: Clustering, Classification and Density Estimation Using Gaussian Finite Mixture Models. *R J* 8: 289–317.



Simão, F. A., R. M. Waterhouse, P. Ioannidis, E. V. Kriventseva, and E. M. Zdobnov, 2015 BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31: 3210–3212. doi: [10.1093/bioinformatics/btv351](https://doi.org/10.1093/bioinformatics/btv351).





Standage, D. S., A. J. Berens, K. M. Glastad, A. J. Severin, and V. P. Brendel *et al.*, 2016 Genome, transcriptome and methylome sequencing of a primitively eusocial wasp reveal a greatly reduced DNA methylation system in a social insect. *Molecular Ecology* 25: 1769–1784. doi: [10.1111/mec.13578](https://doi.org/10.1111/mec.13578).



The Honeybee Genome Sequencing Consortium, 2006 Insights into social insects from the genome of the honeybee *Apis mellifera*. *Nature* 443: 931–949. doi: [10.1038/nature05260](https://doi.org/10.1038/nature05260).

Thomas, C. D., H. Moller, G. M. Plunkett, and R. J. Harris, 1990 The prevalence of introduced *Vespula vulgaris* wasps in a New Zealand beech forest community. *New Zealand journal of ecology* 13: 63–72.

Werren, J. H., S. Richards, C. A. Desjardins, O. Niehuis, and J. Gadau *et al.*, 2010  
Functional and Evolutionary Insights from the Genomes of Three Parasitoid *Nasonia*  
Species. *Science* 327: 343–348. doi: [10.1126/science.1178028](https://doi.org/10.1126/science.1178028).











Vespula-specific  
duplications




A) Venom hyaluronidase

Independent  
duplications  
in *Polistes*  
and *germanica*



B) Venom phospholipase

Antigen 5 clade 1



C) Venom Antigen 5