

Mathematical Models and Methods in Applied Sciences
(2021)
© World Scientific Publishing Company
DOI: 10.1142/S0218202521500196



The Stationary Boussinesq problem under singular forcing

Alejandro Allendes* and Enrique Otárola†

*Departamento de Matemática,
Universidad Técnica Federico Santa María,
Valparaíso, Chile*

**alejandro.allendes@usm.cl*

†enrique.otarola@usm.cl

Abner J. Salgado

*Department of Mathematics,
University of Tennessee,
Knoxville, TN 37996, USA
asalgad1@utk.edu*

Received 15 May 2020

Revised 13 December 2020

Accepted 22 January 2021

Published

Communicated by L. Beirao da Veiga

In Lipschitz two- and three-dimensional domains, we study the existence for the so-called Boussinesq model of thermally driven convection under singular forcing. By singular we mean that the heat source is allowed to belong to $H^{-1}(\varpi, \Omega)$, where ϖ is a weight in the Muckenhoupt class A_2 that is regular near the boundary. We propose a finite element scheme and, under the assumption that the domain is convex and $\varpi^{-1} \in A_1$, show its convergence. In the case that the thermal diffusion and viscosity are constants, we propose an *a posteriori* error estimator and show its reliability. We also explore efficiency estimates.

Keywords: Boussinesq problem; Navier–Stokes equations; singular sources; Muckenhoupt weights; weighted estimates; finite element approximation; *a posteriori* error estimates.

AMS Subject Classification 2020: 35Q35, 35Q30, 35R06, 76Dxx, 65N15, 65N30, 65N50

1. Introduction

The purpose of this paper is to study existence, uniqueness, and approximation results for the so-called Boussinesq model of thermally driven convection. While this problem has been considered before in different contexts and there are such results already available in the literature,^{3,10,15,21,33,41–43} our main source of novelty and originality here is that we allow the heat source to be singular, say a

†Corresponding author.

2 *A. Allendes, E. Otárola & A. J. Salgado*

Dirac measure concentrated in a lower dimensional object so that the problem cannot be understood with the usual energy setting; as it was done, for instance, in Refs. 10, 21, 33, 42 and 43. Let us make this discussion precise. Let $\Omega \subset \mathbb{R}^d$, with $d \in \{2, 3\}$, be an open and bounded domain with Lipschitz boundary $\partial\Omega$. We are interested in existence, uniqueness, and approximation of solutions to the following system of partial differential equations (PDEs) in strong form:

$$\begin{cases} -\operatorname{div}(\nu(\mathbf{T})\nabla \mathbf{u}) + (\mathbf{u} \cdot \nabla)\mathbf{u} + \nabla \mathbf{p} = \mathbf{g}\mathbf{T}, & \text{in } \Omega, \\ \operatorname{div} \mathbf{u} = 0, & \text{in } \Omega, \\ -\operatorname{div}(\kappa(\mathbf{T})\nabla \mathbf{T}) + \operatorname{div}(\mathbf{u}\mathbf{T}) = \mathcal{H}, & \text{in } \Omega, \\ \mathbf{u} = \mathbf{0}, \quad \mathbf{T} = 0, & \text{on } \partial\Omega. \end{cases} \quad (1.1)$$

The unknowns are the velocity $\mathbf{u}$, pressure $\mathbf{p}$, and temperature $\mathbf{T}$ of the fluid, respectively. The data are the viscosity coefficient ν , gravity $\mathbf{g}$, the thermal diffusivity coefficient κ , and the externally applied heat source $\mathcal{H}$. Our main source of interest is the case of a rough $\mathcal{H}$, so that standard energy arguments do not apply to obtain suitable estimates. We will make precise assumptions quantifying this below.

Our presentation will be organized as follows. We collect background information, and the main assumptions under which we shall operate in Sec. 2, where we also introduce a notion of solution for (1.1); see Definition 2.2. Existence of solutions is presented in Sec. 3. The numerical analysis of problem (1.1) begins in Sec. 4, where we introduce a finite-element-like numerical scheme, show that it always has solutions, and that these converge. Section 5 continues the numerical analysis by introducing an *a posteriori* error estimator for our problem and showing its reliability. We also explore efficiency estimates. Finally, a series of numerical experiments are presented in Sec. 6. We show the performance of the devised error estimator within an adaptive loop and explore our model beyond what our theory can handle.

2. Notation and Main Assumptions

Throughout this paper, $d \in \{2, 3\}$ and $\Omega \subset \mathbb{R}^d$ is an open and bounded domain with Lipschitz boundary $\partial\Omega$. If $\mathcal{W}$ and $\mathcal{Z}$ are Banach function spaces, we write $\mathcal{W} \hookrightarrow \mathcal{Z}$ to denote that $\mathcal{W}$ is continuously embedded in $\mathcal{Z}$. We denote by $\mathcal{W}'$ and $\|\cdot\|_{\mathcal{W}}$ the dual and the norm of $\mathcal{W}$, respectively.

For $E \subset \Omega$ open and $f : E \rightarrow \mathbb{R}$, we set

$$\int_E f \, dx = \frac{1}{|E|} \int_E f \, dx, \quad |E| = \int_E 1 \, dx.$$

Given $p \in (1, \infty)$, we denote by p' its Hölder conjugate, i.e. the real number such that $1/p + 1/p' = 1$. By $a \lesssim b$ we mean $a \leq Cb$, with a constant C that neither depends on a , b , nor the discretization parameters. The value of C might change at each occurrence. If the particular value of a constant is of relevance, then we will assign it a name. For instance, throughout our paper, by $C_{4 \rightarrow 2}$ we shall denote the best constant in the embedding $\mathbf{H}_0^1(\Omega) \hookrightarrow \mathbf{L}^4(\Omega)$.

2.1. Weighted function spaces and their embeddings

A weight is a locally integrable and nonnegative function defined on $\mathbb{R}^d$. If ϖ is a weight and $p \in [1, \infty)$, we say that ϖ belongs to the so-called Muckenhoupt class A_p if^{18,35,45}

$$[\varpi]_{A_p} := \sup_B \left(\int_B \varpi \, dx \right) \left(\int_B \varpi^{1/(1-p)} \, dx \right)^{p-1} < \infty, \quad p \in (1, \infty),$$

$$[\varpi]_{A_1} := \sup_B \left(\int_B \varpi \, dx \right) \sup_{x \in B} \frac{1}{\varpi(x)} < \infty, \quad p = 1,$$

where the supremum is taken over all balls B in $\mathbb{R}^d$. In addition, $A_\infty := \bigcup_{p \geq 1} A_p$. We call $[\varpi]_{A_p}$, for $p \in [1, \infty)$, the Muckenhoupt characteristic of ϖ .

Distances to lower dimensional objects are prototypical examples of Muckenhoupt weights. In particular, if $\mathcal{K} \subset \Omega$ is a smooth compact submanifold of dimension $k \in \{0, 1, \dots, d-1\}$ then, owing to Lemma 2.3 item (vi) in Ref. 22, we have that

$$d_{\mathcal{K}}^\alpha(x) = \text{dist}(x, \mathcal{K})^\alpha$$

belongs to the class A_p provided $\alpha \in (-(d-k), (d-k)(p-1))$. This allows us to identify three particular cases:

- (i) Let $d > 1$ and $z \in \Omega$, then the weight $d_z^\alpha \in A_2$ if and only if $\alpha \in (-d, d)$.
- (ii) Let $d \geq 2$ and $\gamma \subset \Omega$ be a smooth closed curve without self-intersections. We have that $d_\gamma^\alpha \in A_2$ if and only if $\alpha \in (-(d-1), d-1)$.
- (iii) Finally, if $d = 3$ and $\Gamma \subset \Omega$ is a smooth surface without boundary, then $d_\Gamma^\alpha \in A_2$ if and only if $\alpha \in (-1, 1)$.

Since the aforementioned lower dimensional objects are strictly contained in Ω , there is a neighborhood of $\partial\Omega$ where the weight has no degeneracies or singularities. In fact, it is continuous and strictly positive. Inspired by Definition 2.5 in Ref. 22, this observation motivates us to define a restricted class of Muckenhoupt weights.

Definition 2.1. (Class $A_p(D)$) Let $D \subset \mathbb{R}^d$ be a Lipschitz domain. For $p \in (1, \infty)$ we say that $\varpi \in A_p$ belongs to $A_p(D)$ if there is an open set $\mathcal{G} \subset D$, and positive constants $\varepsilon > 0$ and $\varpi_l > 0$, such that:

$$\{x \in \Omega : \text{dist}(x, \partial D) < \varepsilon\} \subset \mathcal{G}, \quad \varpi \in C(\bar{\mathcal{G}}), \quad \varpi_l \leq \varpi(x) \quad \forall x \in \bar{\mathcal{G}}.$$

Remark 2.1. ($d_z^\alpha \in A_2(\Omega)$ and $d_z^{-\alpha} \in A_1$) Let $z \in \Omega$ and $\alpha \in (d-2, d)$. Define $d_z(x) = |x - z|$. Then, we have that the weight d_z^α is such that $d_z^\alpha \in A_2(\Omega)$ and $d_z^{-\alpha} \in A_1$.

From the A_p -condition and Hölder's inequality follows that an A_p -weight satisfies the so-called *strong doubling property*; see Proposition 1.2.7 in Ref. 45: Let

4 A. Allendes, E. Otárola & A. J. Salgado

$\varpi \in A_p$ with $p \in (1, \infty)$ and $E \subset \mathbb{R}^d$ be a measurable subset of a ball $B \subset \mathbb{R}^d$. Then,

$$\varpi(B) \leq C_{p,\varpi} \left(\frac{|B|}{|E|} \right)^p \varpi(E). \quad (2.1)$$

The following embedding results will be of importance in our analysis.

Proposition 2.1. (Weighted embedding) *Let $p \in (1, \infty)$ and $\varpi \in A_p$. There is $\delta > 0$ such that if*

$$k \in \left[1, \frac{d}{d-1} + \delta \right],$$

then $W_0^{1,p}(\varpi, \Omega) \hookrightarrow L^{kp}(\varpi, \Omega)$. If, in addition, $\mathbf{S}_{\text{sing}}(\varpi) \Subset \Omega$ then, the embedding is compact for $1 \leq k \leq d/(d-1)$. Here, $\mathbf{S}_{\text{sing}}(\varpi)$ denotes the set of singularities defined in Sec. 4.1 of Ref. 28.

Proof. Theorem 1.3 in Ref. 20 guarantees that the embedding is continuous. The compactness of the embedding follows from Theorem 4.12 in Ref. 28. $\square$

Proposition 2.2. (Embedding with different metrics) *Let $1 \leq p \leq q < \infty$, $\varpi \in A_p$, and $\rho \in A_q$. If the pair (ρ, ϖ) satisfies the compatibility condition*

$$\frac{r}{R} \left(\frac{\rho(B_r)}{\rho(B_R)} \right)^{1/q} \left(\frac{\varpi(B_R)}{\varpi(B_r)} \right)^{1/p} \lesssim 1, \quad 0 < r \leq R,$$

then, we have that, $W_0^{1,p}(\varpi, \Omega) \hookrightarrow L^q(\rho, \Omega)$.

Proof. See Theorem 6.1 in Ref. 36. $\square$

Proposition 2.3. (Boundedness) *Let $d \in \{2, 3\}$ and $\varpi \in A_2$. For every $\mathbf{g} \in \mathbf{L}^\infty(\Omega)$, $\theta \in H_0^1(\varpi, \Omega)$, and $\mathbf{v} \in \mathbf{H}_0^1(\Omega)$, we have that*

$$\left| \int_{\Omega} \theta \mathbf{g} \cdot \mathbf{v} \, dx \right| \leq C_{e,1} \|\mathbf{g}\|_{\mathbf{L}^\infty(\Omega)} \|\nabla \theta\|_{\mathbf{L}^2(\varpi, \Omega)} \|\nabla \mathbf{v}\|_{\mathbf{L}^2(\Omega)}. \quad (2.2)$$

If, in addition, the weight satisfies $\varpi^{-1} \in A_1$, then provided $r \in H_0^1(\varpi^{-1}, \Omega)$, we have

$$\left| \int_{\Omega} \theta \mathbf{v} \cdot \nabla r \, dx \right| \leq C_{e,2} \|\nabla \mathbf{v}\|_{\mathbf{L}^2(\Omega)} \|\nabla \theta\|_{\mathbf{L}^2(\varpi, \Omega)} \|\nabla r\|_{\mathbf{L}^2(\varpi^{-1}, \Omega)}. \quad (2.3)$$

In both estimates, the constants depend only on Ω and ϖ .

Proof. Since the weight $\varpi \in A_2$, we have that it satisfies the strong doubling property (2.1) with $p = 2$. Thus, for $0 < r \leq R$ and $q \leq d/(d-1)$, we have

$$\frac{r}{R} \left(\frac{|B_r|}{|B_R|} \right)^{1/q} \left(\frac{\varpi(B_R)}{\varpi(B_r)} \right)^{1/2} \lesssim \left(\frac{r}{R} \right)^{1+d/q} \frac{|B_R|}{|B_r|} \lesssim 1.$$

We thus invoke Proposition 2.2 with $p = 2$ and $\rho = 1$ to conclude that $H_0^1(\varpi, \Omega) \hookrightarrow L^q(\Omega)$ provided $q \leq d/(d-1)$. Consequently, for $q \leq d/(d-1)$, we have

$$\left| \int_{\Omega} \theta \mathbf{g} \cdot \mathbf{v} \, dx \right| \leq \|\mathbf{g}\|_{\mathbf{L}^\infty(\Omega)} \|\theta\|_{L^q(\Omega)} \|\mathbf{v}\|_{\mathbf{L}^{q'}(\Omega)} \lesssim \|\mathbf{g}\|_{\mathbf{L}^\infty(\Omega)} \|\nabla \theta\|_{L^2(\varpi, \Omega)} \|\mathbf{v}\|_{\mathbf{L}^{q'}(\Omega)}.$$

Notice that $q' \geq d$. Choosing $q = d/(d-1)$ and utilizing a standard Sobolev embedding yields estimate (2.2).

We now prove inequality (2.3). To accomplish this task, we first notice that $\varpi^{-1} \in A_1$ implies $\varpi \in L^\infty(\Omega)$. Second, since Proposition 2.1 guarantees the existence of $\epsilon > 0$ such that $H_0^1(\varpi, \Omega) \hookrightarrow L^{\mu+\epsilon}(\varpi, \Omega)$, for $\mu = 2d/(d-1)$, we can conclude the existence of $\ell \geq 2d/(d-1)$ and $m \leq 2d$ such that $m^{-1} + \ell^{-1} = 1/2$ and

$$\begin{aligned} \left| \int_{\Omega} \theta \mathbf{v} \cdot \nabla r \, dx \right| &\leq \|\varpi\|_{L^\infty(\Omega)}^{1/m} \|\mathbf{v}\|_{\mathbf{L}^m(\Omega)} \|\theta\|_{L^\ell(\varpi, \Omega)} \|\nabla r\|_{\mathbf{L}^2(\varpi^{-1}, \Omega)} \\ &\lesssim \|\varpi\|_{L^\infty(\Omega)}^{1/m} \|\nabla \mathbf{v}\|_{\mathbf{L}^2(\Omega)} \|\nabla \theta\|_{L^2(\varpi, \Omega)} \|\nabla r\|_{\mathbf{L}^2(\varpi^{-1}, \Omega)}, \end{aligned} \quad (2.4)$$

where we have also used a standard, unweighted, Sobolev embedding to handle the term involving $\mathbf{v}$. This yields (2.3) and concludes the proof. $\square$

2.2. Main assumptions and definition of solution

Having described the functional setting that we shall adopt and some of its more relevant properties, we can precisely state the assumptions under which we shall operate.

- *Domain:* Let $d \in \{2, 3\}$. We assume that Ω is a bounded domain in $\mathbb{R}^d$ with Lipschitz boundary $\partial\Omega$. When dealing with discretization, we shall further assume that Ω is a polytope.
- *Gravity:* The gravity is a constant vector $\mathbf{g} \in \mathbb{R}^d$. We set $g = |\mathbf{g}|$.
- *Viscosity:* The viscosity is a function $\nu \in C^{0,1}(\mathbb{R})$ that is strictly positive and bounded, i.e. there are positive constants ν_- and ν_+ such that $\nu_- \leq \nu_+$ and

$$\nu_- \leq \nu(t) \leq \nu_+ \quad \forall t \in \mathbb{R}.$$

- *Thermal diffusivity:* The thermal coefficient is a function $\kappa \in C^{0,1}(\mathbb{R})$ that is, moreover, strictly positive and bounded, i.e. there are positive constants κ_- and κ_+ such that $\kappa_- \leq \kappa_+$ and

$$\kappa_- \leq \kappa(t) \leq \kappa_+ \quad \forall t \in \mathbb{R}.$$

To quantify the oscillation of the thermal diffusivity we shall introduce

$$\Lambda(\kappa) := \frac{\kappa_-}{\kappa_+} \in (0, 1].$$

- *Weight:* We assume that we have a weight $\varpi \in A_2(\Omega)$ such that $\varpi^{-1} \in A_1$ and $\mathbf{S}_{\text{sing}}(\varpi) \Subset \Omega$. A canonical example of this scenario is given in Remark 2.1; Example 4.4 in Ref. 28 shows that $|\mathbf{S}_{\text{sing}}(\varpi)| = 0$.

6 *A. Allendes, E. Otárola & A. J. Salgado*

- *Heat source:* We allow the heat source to be singular, and we quantify this by assuming that it belongs to the dual of a weighted space. Namely, we assume that $\mathcal{H} \in H^{-1}(\varpi, \Omega) := H_0^1(\varpi^{-1}, \Omega)'$.

With these assumptions at hand we can define our notion of solution.

Definition 2.2. (Weak solution) We say that the triple $(\mathbf{T}, \mathbf{u}, \mathbf{p}) \in H_0^1(\varpi, \Omega) \times \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$ is a weak solution to (1.1) if

$$\begin{cases} \int_{\Omega} (\nu(\mathbf{T}) \nabla \mathbf{u} : \nabla \mathbf{v} + (\mathbf{u} \cdot \nabla) \mathbf{u} \cdot \mathbf{v} - \mathbf{p} \operatorname{div} \mathbf{v} - \mathbf{T} \mathbf{g} \cdot \mathbf{v}) \, dx = 0, \\ \int_{\Omega} q \operatorname{div} \mathbf{u} \, dx = 0, \\ \int_{\Omega} (\kappa(\mathbf{T}) \nabla \mathbf{T} \cdot \nabla r - \mathbf{T} \mathbf{u} \cdot \nabla r) \, dx = \langle \mathcal{H}, r \rangle, \end{cases} \quad (2.5)$$

for all $\mathbf{v} \in \mathbf{H}_0^1(\Omega)$, $q \in L_0^2(\Omega)$, and $r \in H_0^1(\varpi^{-1}, \Omega)$. Here, $\langle \cdot, \cdot \rangle$ denotes the duality pairing between $H_0^1(\varpi^{-1}, \Omega)$ and its dual $H^{-1}(\varpi, \Omega)$.

We immediately comment that, owing to our assumptions on data and definition of solution, all terms in this definition are meaningful; see Proposition 2.3.

3. Existence of Solutions

The main goal in this section is to show that problem (1.1) has, under the assumptions stated in Sec. 2.2, a solution in the sense of Definition 2.2. We proceed in several steps.

3.1. The Navier–Stokes equation with prescribed temperature

We begin by making a simple observation. Given $\theta \in H_0^1(\varpi, \Omega)$, let us consider the following problem: Find $(\mathbf{u}, p) \in \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$ such that, for all $(\mathbf{v}, q) \in \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$,

$$\begin{cases} \int_{\Omega} (\nu(\theta) \nabla \mathbf{u} : \nabla \mathbf{v} + (\mathbf{u} \cdot \nabla) \mathbf{u} \cdot \mathbf{v} - p \operatorname{div} \mathbf{v}) \, dx = \int_{\Omega} \theta \mathbf{g} \cdot \mathbf{v} \, dx \\ \int_{\Omega} q \operatorname{div} \mathbf{u} \, dx = 0. \end{cases} \quad (3.1)$$

We present the following result regarding existence and uniqueness of solutions to (3.1). We will make use of the fact that, on Lipschitz domains, the divergence operator is surjective from $\mathbf{H}_0^1(\Omega)$ to $L_0^2(\Omega)$.^{11,12} This implies that there is a constant B , that depends only on d and Ω , for which

$$B \|q\|_{L^2(\Omega)} \leq \sup_{\mathbf{v} \in \mathbf{H}_0^1(\Omega)} \frac{\int_{\Omega} \operatorname{div} \mathbf{v} q \, dx}{\|\nabla \mathbf{v}\|_{\mathbf{L}^2(\Omega)}} \quad \forall q \in L_0^2(\Omega). \quad (3.2)$$

Theorem 3.1. (Existence and uniqueness) *For every $\theta \in H_0^1(\varpi, \Omega)$ problem (3.1) has at least one solution. In addition, if $C_P C_{e,1} g \|\nabla \theta\|_{\mathbf{L}^2(\varpi, \Omega)} < \nu_-^2$, where C_P is a constant that depends only on Ω and d , then this solution is unique and satisfies*

$$\begin{aligned} \|\nabla \mathbf{u}\|_{\mathbf{L}^2(\Omega)} &\leq \frac{C_{e,1} g}{\nu_-} \|\nabla \theta\|_{\mathbf{L}^2(\varpi, \Omega)}, \\ \|p\|_{L^2(\Omega)} &\leq \frac{C_{e,1} g}{B} \|\nabla \theta\|_{\mathbf{L}^2(\varpi, \Omega)} \left(1 + \frac{\nu_+}{\nu_-} + \frac{C_{4 \rightarrow 2}^2 C_{e,1} g}{\nu_-^2} \|\nabla \theta\|_{\mathbf{L}^2(\varpi, \Omega)} \right), \end{aligned}$$

where $C_{4 \rightarrow 2}$ is the best constant in the embedding $\mathbf{H}_0^1(\Omega) \hookrightarrow \mathbf{L}^4(\Omega)$ and B is the constant appearing in estimate (3.2).

Proof. Since $\theta \in H_0^1(\varpi, \Omega)$, the function $\bar{\nu}(x) := \nu(\theta(x))$ is bounded, measurable, and strictly positive. Define the functional

$$\mathcal{F}_\theta : \mathbf{v} \mapsto \int_{\Omega} \theta \mathbf{g} \cdot \mathbf{v} \, dx.$$

Owing to Proposition 2.3 we have that $\mathcal{F}_\theta \in \mathbf{H}^{-1}(\Omega)$. In addition, Proposition 2.3 also shows that

$$\|\mathcal{F}_\theta\|_{\mathbf{H}^{-1}(\Omega)} \leq C_{e,1} \|g\|_{\mathbf{L}^\infty(\Omega)} \|\nabla \theta\|_{\mathbf{L}^2(\varpi, \Omega)} = C_{e,1} g \|\nabla \theta\|_{\mathbf{L}^2(\varpi, \Omega)}.$$

Thus, the standard theory of existence and uniqueness under small data (or large viscosity) for the Navier–Stokes equation applies; see Chap. II, §1 in Ref. 44.

We now obtain the claimed estimates. The bound on $\mathbf{u}$ follows from standard energy arguments and the bound on $\mathcal{F}_\theta$. The bound on p is obtained from (3.2). Indeed, using the first equation in (3.1), inequality (3.2) yields

$$B \|p\|_{L^2(\Omega)} \leq \nu_+ \|\nabla \mathbf{u}\|_{\mathbf{L}^2(\Omega)} + C_{4 \rightarrow 2}^2 \|\nabla \mathbf{u}\|_{\mathbf{L}^2(\Omega)}^2 + \|\mathcal{F}_\theta\|_{\mathbf{H}^{-1}(\Omega)}.$$

Substitute the obtained bounds in terms of $\nabla \theta$ to conclude. $\square$

3.2. The stationary heat equation with convection

Here, we study the existence of solutions to a stationary heat equation with convection and under singular forcing. Namely, given $\varkappa \in L^\infty(\Omega)$ with $0 < \varkappa_- \leq \varkappa \leq \varkappa_+$, $\mathbf{u} \in \mathbf{H}_0^1(\Omega)$, and $\mathcal{H} \in H^{-1}(\varpi, \Omega)$, we consider the following stationary heat equation: Find $T \in H_0^1(\varpi, \Omega)$ such that

$$\int_{\Omega} (\varkappa \nabla T \cdot \nabla r - T \mathbf{u} \cdot \nabla r) \, dx = \langle \mathcal{H}, r \rangle \quad \forall r \in H_0^1(\varpi^{-1}, \Omega). \quad (3.3)$$

As a first step, we state a well-posedness result for the case $\mathbf{u} = \mathbf{0}$.

Proposition 3.1. (Well-posedness for $\mathbf{u} = \mathbf{0}$) *There is a constant Λ_0 , depending only on Ω and ϖ , such that, if $\Lambda(\varkappa) \geq \Lambda_0$, problem (3.3) with $\mathbf{u} = \mathbf{0}$ is well-posed.*

8 A. Allendes, E. Otárola & A. J. Salgado

This, in particular, implies that

$$\|\nabla T\|_{\mathbf{L}^2(\varpi, \Omega)} \leq C_{\varkappa} \sup_{r \in H_0^1(\varpi^{-1}, \Omega)} \frac{\int_{\Omega} \varkappa \nabla T \cdot \nabla r \, dx}{\|\nabla r\|_{\mathbf{L}^2(\varpi^{-1}, \Omega)}} \quad \forall T \in H_0^1(\varpi, \Omega). \quad (3.4)$$

The constant $C_{\varkappa}$ depends only on Λ_0 , $\Lambda(\varkappa)$, Ω , d , and ϖ .

Proof. See Theorem 12 in Ref. 40. $\square$

We now study the case with nonzero convection.

Proposition 3.2. (Well-posedness for $\mathbf{u} \neq \mathbf{0}$) Assume that $\Lambda(\varkappa) \geq \Lambda_0$, where Λ_0 is defined in Proposition 3.1. If

$$C_{\varkappa} C_{e,2} \|\nabla \mathbf{u}\|_{\mathbf{L}^2(\Omega)} \leq q < 1, \quad (3.5)$$

then problem (3.3) is well-posed. This, in particular, implies that the solution T of problem (3.3) satisfies the estimate

$$\|\nabla T\|_{\mathbf{L}^2(\varpi, \Omega)} \leq C_H(q) \|\mathcal{H}\|_{H^{-1}(\varpi, \Omega)}, \quad C_H(q) = \frac{C_{\varkappa}}{1-q}.$$

Proof. Let us introduce the linear map $\mathcal{A} : H_0^1(\varpi, \Omega) \rightarrow H^{-1}(\varpi, \Omega)$ via

$$\langle \mathcal{A}T, r \rangle := \int_{\Omega} \varkappa \nabla T \cdot \nabla r \, dx, \quad \forall T \in H_0^1(\varpi, \Omega), \quad \forall r \in H_0^1(\varpi^{-1}, \Omega).$$

Clearly, $\mathcal{A}$ is a bounded linear operator and, moreover, owing to the inf-sup estimate (3.4), $\mathcal{A}$ is invertible with $\|\mathcal{A}^{-1}\|_{\mathcal{L}(H^{-1}(\varpi, \Omega), H_0^1(\varpi, \Omega))} \leq C_{\varkappa}$.

Given $\mathbf{u} \in \mathbf{H}_0^1(\Omega)$, we introduce the map $\mathcal{B}_{\mathbf{u}} : H_0^1(\varpi, \Omega) \rightarrow H^{-1}(\varpi, \Omega)$ defined by

$$\langle \mathcal{B}_{\mathbf{u}}T, r \rangle = - \int_{\Omega} T \mathbf{u} \cdot \nabla r \, dx, \quad \forall T \in H_0^1(\varpi, \Omega), \quad \forall r \in H_0^1(\varpi^{-1}, \Omega).$$

Estimate (2.3) shows that $\mathcal{B}_{\mathbf{u}}$ is a bounded linear map which satisfies the estimate

$$\|\mathcal{B}_{\mathbf{u}}\|_{\mathcal{L}(H_0^1(\varpi, \Omega), H^{-1}(\varpi, \Omega))} \leq C_{e,2} \|\nabla \mathbf{u}\|_{\mathbf{L}^2(\Omega)}.$$

Since it will be needed later, we now show that $\mathcal{B}_{\mathbf{u}}$ is compact. Let $\{T_n\}_{n \geq 0}$ be a bounded sequence in $H_0^1(\varpi, \Omega)$. Since Proposition 2.1 guarantees that, for $k \leq d/(d-1)$, the embedding $H_0^1(\varpi, \Omega) \hookrightarrow L^{2k}(\varpi, \Omega)$ is compact, we conclude the existence of a subsequence $\{T_{n_j}\}_{j \geq 0}$ of $\{T_n\}_{n \geq 0}$ such that $T_{n_j} \rightharpoonup T^*$ in $H_0^1(\varpi, \Omega)$ and $T_{n_j} \rightarrow T^*$ in $L^{2d/(d-1)}(\varpi, \Omega)$ as $j \uparrow \infty$. Thus, estimate (2.4) yields

$$\|\mathcal{B}_{\mathbf{u}}T_{n_j} - \mathcal{B}_{\mathbf{u}}T^*\|_{H^{-1}(\varpi, \Omega)} \lesssim \|T_{n_j} - T^*\|_{L^{2d/(d-1)}(\varpi, \Omega)} \|\nabla \mathbf{u}\|_{\mathbf{L}^2(\Omega)} \rightarrow 0, \quad j \uparrow \infty.$$

This shows that $\{\mathcal{B}_{\mathbf{u}}T_{n_j}\}_{j \geq 0}$ converges in $H^{-1}(\varpi, \Omega)$ and thus that $\mathcal{B}_{\mathbf{u}}$ is compact. We notice that, in view of (2.4), the map $\mathcal{B}_{\mathbf{u}}$ is well-defined on the space $L^{2d/(d-1)}(\varpi, \Omega)$.

With this notation, we have that problem (3.3) can be written as

$$(\mathcal{A} + \mathcal{B}_{\mathbf{u}})T = \mathcal{H} \iff (I + \mathcal{A}^{-1}\mathcal{B}_{\mathbf{u}})T = \mathcal{A}^{-1}\mathcal{H}$$

in $H^{-1}(\varpi, \Omega)$. Since $\mathcal{A}^{-1}\mathcal{B}_{\mathbf{u}}$ is continuous, assumption (3.5) implies that this problem has a unique solution, because

$$\|\mathcal{A}^{-1}\mathcal{B}_{\mathbf{u}}\|_{\mathcal{L}(H_0^1(\varpi, \Omega))} \leq C_{\varkappa}C_{e,2}\|\nabla\mathbf{u}\|_{\mathbf{L}^2(\Omega)} \leq q < 1.$$

Moreover, we have the estimate

$$\|\nabla T\|_{\mathbf{L}^2(\varpi, \Omega)} \leq \frac{\|\mathcal{A}^{-1}\|_{\mathcal{L}(H^{-1}(\varpi, \Omega), H_0^1(\varpi, \Omega))}}{1 - \|\mathcal{A}^{-1}\mathcal{B}_{\mathbf{u}}\|_{\mathcal{L}(H_0^1(\varpi, \Omega))}} \|\mathcal{H}\|_{H^{-1}(\varpi, \Omega)} \leq \frac{C_{\varkappa}}{1 - q} \|\mathcal{H}\|_{H^{-1}(\varpi, \Omega)}.$$

Notice that $C_H(q) = C_{\varkappa}/(1 - q)$ depends only on q , Λ_0 , $\Lambda(\varkappa)$, Ω , d , and ϖ . $\square$

3.3. Existence of solutions

Having studied each one of the subproblems separately, we proceed to show existence of solutions to (2.5) via a fixed point argument. To accomplish this task, we define the map $\mathfrak{F} : H_0^1(\varpi, \Omega) \times \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega) \rightarrow H_0^1(\varpi, \Omega) \times \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$ by $\mathfrak{F}(\theta, \mathbf{u}, p) := (\mathbf{T}, \mathbf{u}, p)$, where $(\mathbf{T}, \mathbf{u}, p)$ solves

$$\int_{\Omega} (\nu(\theta)\nabla\mathbf{u} : \nabla\mathbf{v} + (\mathbf{u} \cdot \nabla\mathbf{u}) \cdot \mathbf{v} - p \operatorname{div} \mathbf{v}) \, dx = \int_{\Omega} \theta \mathbf{g} \cdot \mathbf{v} \, dx \quad \forall \mathbf{v} \in \mathbf{H}_0^1(\Omega), \quad (3.6)$$

$$\int_{\Omega} q \operatorname{div} \mathbf{u} \, dx = 0 \quad \forall q \in L_0^2(\Omega), \quad (3.7)$$

$$\int_{\Omega} (\kappa(\theta)\nabla\mathbf{T} \cdot \nabla r - \mathbf{T}\mathbf{u} \cdot \nabla r) \, dx = \langle \mathcal{H}, r \rangle \quad \forall r \in H_0^1(\varpi^{-1}, \Omega). \quad (3.8)$$

Note that the definition of $(\mathbf{T}, \mathbf{u}, p)$ implies solving a stationary Navier–Stokes equation with prescribed temperature θ and a stationary heat equation with convection $\mathbf{u}$. The following result shows that the map $\mathfrak{F}$ is well-defined. To concisely state it we define

$$\begin{cases} \mathfrak{B}_T = \{\theta \in H_0^1(\varpi, \Omega) : \|\nabla\theta\|_{\mathbf{L}^2(\varpi, \Omega)} \leq S\}, \\ S = \frac{\nu_-}{gC_{e,1}} \min \left\{ \frac{\nu_-}{C_P}, \frac{1}{2C_{\varkappa}C_{e,2}} \right\}, \end{cases} \quad (3.9)$$

$$\begin{cases} \mathfrak{B}_{\mathbf{u}} = \{\mathbf{u} \in \mathbf{H}_0^1(\Omega) : \|\nabla\mathbf{u}\|_{\mathbf{L}^2(\Omega)} \leq G\}, \\ G = \frac{1}{2C_{\varkappa}C_{e,2}}, \end{cases} \quad (3.10)$$

$$\begin{cases} \mathfrak{B}_p = \{p \in L_0^2(\Omega) : \|p\|_{L^2(\Omega)} \leq P\}, \\ P = \frac{C_{e,1}g}{B} S \left(1 + \frac{\nu_+}{\nu_-} + \frac{C_{4 \rightarrow 2}^2 C_{e,1}g}{\nu_-^2} S \right) \end{cases} \quad (3.11)$$

and $\mathfrak{B} = \mathfrak{B}_T \times \mathfrak{B}_{\mathbf{u}} \times \mathfrak{B}_p$.

Proposition 3.3. ($\mathfrak{F}$ is well-defined) *Assume that $\Lambda(\kappa) \geq \Lambda_0$, where Λ_0 is defined in Proposition 3.1. If the heat source $\mathcal{H} \in H^{-1}(\varpi, \Omega)$ satisfies the estimate*

$$\|\mathcal{H}\|_{H^{-1}(\varpi, \Omega)} \leq \frac{S}{C_H(1/2)} = \frac{S}{2C_\kappa}, \quad (3.12)$$

then $\mathfrak{F}$ is well-defined on $\mathfrak{B}$. In addition, we have $\mathfrak{F}(\mathfrak{B}) \subset \mathfrak{B}$.

Proof. Let $\theta \in \mathfrak{B}_T$. Invoke Theorem 3.1 to conclude the existence of a unique $(\mathbf{u}, \mathbf{p}) \in \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$ that solves (3.6) and (3.7). Moreover, $\mathbf{u}$ and $\mathbf{p}$ satisfy the estimates

$$\begin{aligned} \|\nabla \mathbf{u}\|_{\mathbf{L}^2(\Omega)} &\leq \frac{C_{e,1}g}{\nu_-} \|\nabla \theta\|_{\mathbf{L}^2(\varpi, \Omega)} \leq \frac{C_{e,1}g}{\nu_-} \frac{\nu_-}{gC_{e,1}} \frac{1}{2C_\kappa C_{e,2}} = G, \\ \|\mathbf{p}\|_{L^2(\Omega)} &\leq \frac{C_{e,1}g}{B} \|\nabla \theta\|_{\mathbf{L}^2(\varpi, \Omega)} \left(1 + \frac{\nu_+}{\nu_-} + \frac{C_{4 \rightarrow 2}^2 C_{e,1}g}{\nu_-^2} \|\nabla \theta\|_{\mathbf{L}^2(\varpi, \Omega)} \right) \leq P. \end{aligned}$$

Consequently, $\mathbf{u} \in \mathfrak{B}_\mathbf{u}$ and $\mathbf{p} \in \mathfrak{B}_p$. Now, since $\Lambda(\kappa) \geq \Lambda_0$ and $\mathbf{u} \in \mathfrak{B}_\mathbf{u}$, we invoke Proposition 3.2, with $q = 1/2$, to conclude that there exists a unique $\mathbf{T}$ that solves (3.8). Moreover, the condition on $\mathcal{H}$ guarantees that

$$\|\nabla \mathbf{T}\|_{\mathbf{L}^2(\varpi, \Omega)} \leq C_H(1/2) \|\mathcal{H}\|_{H^{-1}(\varpi, \Omega)} \leq S,$$

which implies that $\mathbf{T} \in \mathfrak{B}_T$. We have thus proved all the statements. $\square$

As a last preparatory step we show that the mapping $\mathfrak{F}$ is weakly continuous.

Lemma 3.1. (Weak continuity) *The mapping $\mathfrak{F} : \mathfrak{B} \rightarrow \mathfrak{B}$ is weakly continuous.*

Proof. Let the sequence $\{(\theta_n, \mathbf{u}_n, p_n)\}_{n \geq 0} \subset \mathfrak{B}$ be such that $(\theta_n, \mathbf{u}_n, p_n) \rightharpoonup (\theta, \mathbf{u}, p)$ in $H_0^1(\varpi, \Omega) \times \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$. As the set $\mathfrak{B}$ is closed and convex, it is weakly closed. Therefore, $(\theta, \mathbf{u}, p) \in \mathfrak{B}$. Set $(\mathbf{T}_n, \mathbf{u}_n, p_n) = \mathfrak{F}(\theta_n, \mathbf{u}_n, p_n)$ and $(\mathbf{T}, \mathbf{u}, p) = \mathfrak{F}(\theta, \mathbf{u}, p)$. We must show that $(\mathbf{T}_n, \mathbf{u}_n, p_n) \rightharpoonup (\mathbf{T}, \mathbf{u}, p)$.

Owing to the reverse Hölder inequality, Theorem 7.4 in Ref. 18, we have that, for some $\epsilon > 0$, the embedding $H_0^1(\varpi, \Omega) \hookrightarrow W^{1,1+\epsilon}(\Omega)$ is continuous. Since $W^{1,1+\epsilon}(\Omega)$ is compactly embedded in $L^{1+\epsilon}(\Omega)$, we obtain that $\theta_n \rightarrow \theta$ in $L^{1+\epsilon}(\Omega)$. The continuity of κ implies then that $\kappa(\theta_n) \rightarrow \kappa(\theta)$ almost everywhere in Ω ; see Theorem 7 in Ref. 9. Now, since $\{(\mathbf{T}_n, \mathbf{u}_n, p_n)\}_{n \geq 0} \subset \mathfrak{B}$ is bounded, we can extract a weakly convergent subsequence $\{(\mathbf{T}_{n_k}, \mathbf{u}_{n_k}, p_{n_k})\}_{k \geq 0}$ such that $(\mathbf{T}_{n_k}, \mathbf{u}_{n_k}, p_{n_k}) \rightharpoonup (\tilde{\mathbf{T}}, \tilde{\mathbf{u}}, \tilde{p})$ in $H_0^1(\varpi, \Omega) \times \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$ as $k \uparrow \infty$. The previous discussion shows that, for every $r \in H_0^1(\varpi^{-1}, \Omega)$, we have

$$\int_{\Omega} \kappa(\theta_{n_k}) \nabla \mathbf{T}_{n_k} \cdot \nabla r \, dx \rightarrow \int_{\Omega} \kappa(\theta) \nabla \tilde{\mathbf{T}} \cdot \nabla r \, dx, \quad k \uparrow \infty.$$

Similar arguments for the remaining terms that comprise the definition of $\mathfrak{F}$ show that, in the limit, we must have $(\tilde{T}, \tilde{u}, \tilde{p}) = \mathfrak{F}(\theta, \mathbf{u}, p)$. Consequently, $(\tilde{T}, \tilde{u}, \tilde{p}) = (T, u, p)$. Since problem (3.6)–(3.8) admits a unique solution, any convergent subsequence converges to the same limit, which implies that the whole sequence must do so to (T, u, p) , which established the desired weak continuity of $\mathfrak{F}$. $\square$

We now proceed to obtain existence via a fixed point argument.

Theorem 3.2. (Existence) *Assume that $\Lambda(\kappa) \geq \Lambda_0$, where Λ_0 is defined in Proposition 3.1. If the heat source $\mathcal{H} \in H^{-1}(\varpi, \Omega)$ satisfies (3.12), then there is a $(T, u, p) \in H_0^1(\varpi, \Omega) \times \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$ that solves (1.1) in the sense of Definition 2.2. Moreover, we have that $(T, u, p) \in \mathfrak{B}$.*

Proof. We wish to invoke the Leray–Schauder fixed point theorem, Theorem 8.8 in Ref. 16, for the map $\mathfrak{F}$ over $\mathfrak{B} = \mathfrak{B}_T \times \mathfrak{B}_u \times \mathfrak{B}_p$, where $\mathfrak{B}_T$, $\mathfrak{B}_u$, and $\mathfrak{B}_p$ are defined in (3.9), (3.10), and (3.11), respectively. Notice that $\mathfrak{B}$ is nonempty, closed, bounded, and convex. Since Proposition 3.3 already showed that $\mathfrak{F}(\mathfrak{B}) \subset \mathfrak{B}$, it remains to show the compactness of $\mathfrak{F}$. In other words, we must improve on Lemma 3.1 by showing the weak–strong continuity of $\mathfrak{F}$. To accomplish this task, let $\{(\theta_n, \mathbf{u}_n, p_n)\}_{n \geq 0} \subset \mathfrak{B}$ be such that $(\theta_n, \mathbf{u}_n, p_n) \rightharpoonup (\theta, \mathbf{u}, p) \in \mathfrak{B}$, in $H_0^1(\varpi, \Omega) \times \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$, as $n \uparrow \infty$. We already know, via Lemma 3.1, that $(T_n, u_n, p_n) = \mathfrak{F}(\theta_n, \mathbf{u}_n, p_n) \rightharpoonup \mathfrak{F}(\theta, \mathbf{u}, p) = (T, u, p)$, in $H_0^1(\varpi, \Omega) \times \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$.

Let $r \in H_0^1(\varpi^{-1}, \Omega)$. Invoke the problems that (T_n, u_n) and (T, u) satisfy and observe that the difference $\mathbf{e}_{T,n} := T - T_n$ verifies the relation

$$\int_{\Omega} (\kappa(\theta_n) \nabla \mathbf{e}_{T,n} - \mathbf{e}_{T,n} \mathbf{u}) \cdot \nabla r \, dx = \int_{\Omega} (T_n(\mathbf{u} - \mathbf{u}_n) + (\kappa(\theta_n) - \kappa(\theta)) \nabla T) \cdot \nabla r \, dx,$$

i.e. $\mathbf{e}_{T,n}$ is the solution to a heat equation with convection; the problem that was studied in Sec. 3.2. Let us denote the functional on the right-hand side of this expression by $\mathcal{H}_n$. Since $\mathbf{u} \in \mathfrak{B}_u$ and $\Lambda(\kappa) \geq \Lambda_0$, we can invoke Proposition 3.2 to conclude that

$$\|\nabla \mathbf{e}_{T,n}\|_{\mathbf{L}^2(\varpi, \Omega)} \leq C_H(1/2) \|\mathcal{H}_n\|_{H^{-1}(\varpi, \Omega)}.$$

The arguments that led to (2.4) show the existence of $m < 2d$ and $\ell > 2d/(d-1)$ such that

$$\sup_{r \in H_0^1(\varpi^{-1}, \Omega)} \frac{\int_{\Omega} T_n(\mathbf{u} - \mathbf{u}_n) \cdot \nabla r \, dx}{\|\nabla r\|_{\mathbf{L}^2(\varpi^{-1}, \Omega)}} \lesssim \|\nabla T_n\|_{\mathbf{L}^2(\varpi, \Omega)} \|\mathbf{u} - \mathbf{u}_n\|_{\mathbf{L}^m(\Omega)} \rightarrow 0,$$

where we have also used the compact embedding $H_0^1(\Omega) \hookrightarrow L^m(\Omega)$. For the second term we observe that, since κ is continuous, and hence bounded, we have that $(\kappa(\theta_n) - \kappa(\theta)) \nabla T \rightarrow \mathbf{0}$ in $\mathbf{L}^2(\varpi, \Omega)$. In conclusion, $T_n \rightarrow T$ in $H_0^1(\varpi, \Omega)$.

12 *A. Allendes, E. Otárola & A. J. Salgado*

An argument similar to that used to obtain uniqueness for the stationary Navier–Stokes equation, cf. Chap. II, §1 in Ref. 44, shows that $\mathbf{u}_n \rightarrow \mathbf{u}$ in $\mathbf{H}_0^1(\Omega)$. Finally, (3.2) implies that $\mathbf{p}_n \rightarrow \mathbf{p}$ in $L_0^2(\Omega)$.

The theorem is thus proved. $\square$

4. Discretization and Convergence

Let us now study a finite-element-like scheme to approximate the solution of (1.1). To that effect, we assume that we have at hand, for each $h > 0$, finite-dimensional spaces $W_h \subset H_0^1(\varpi, \Omega) \cap H_0^1(\varpi^{-1}, \Omega)$, $\mathbf{X}_h \subset \mathbf{H}_0^1(\Omega)$, and $M_h \subset L_0^2(\Omega)$ that are dense in the limit. Moreover, we assume that the pair $(\mathbf{X}_h, M_h)$ is compatible, in the sense that there is a constant $\beta > 0$ such that, for all $h > 0$,

$$\beta \|q_h\|_{L^2(\Omega)} \leq \sup_{\mathbf{v}_h \in \mathbf{X}_h} \frac{\int_{\Omega} \operatorname{div} \mathbf{v}_h q_h \, dx}{\|\nabla \mathbf{v}_h\|_{\mathbf{L}^2(\Omega)}} \quad \forall q_h \in M_h. \quad (4.1)$$

We also assume that the $H_0^1(\Omega)$ projection onto W_h is $H_0^1(\varpi^{\pm 1}, \Omega)$ stable. In other words, there is a constant $\gamma > 0$ such that, for all $h > 0$,

$$\gamma \|\nabla r_h\|_{\mathbf{L}^2(\varpi^{\pm 1}, \Omega)} \leq \sup_{\theta_h \in W_h} \frac{\int_{\Omega} \nabla r_h \cdot \nabla \theta_h \, dx}{\|\nabla \theta_h\|_{\mathbf{L}^2(\varpi^{\mp 1}, \Omega)}} \quad \forall r_h \in W_h. \quad (4.2)$$

Finally, we assume that there is an interpolation operator $\pi_W : H_0^1(\varpi^{-1}, \Omega) \rightarrow W_h$ which is stable and has suitable approximation properties: For all $r \in H_0^1(\varpi^{-1}, \Omega)$, we have

$$\|\nabla \pi_W r\|_{\mathbf{L}^2(\varpi^{-1}, \Omega)} \lesssim \|\nabla r\|_{\mathbf{L}^2(\varpi^{-1}, \Omega)}, \quad \|\nabla(\pi_W r - r)\|_{\mathbf{L}^2(\varpi^{-1}, \Omega)} \xrightarrow{h \rightarrow 0} 0. \quad (4.3)$$

Examples of triples verifying our assumptions are plentiful within the finite element literature. Pairs that satisfy (4.1) can be found, for instance, in Refs. 14, 19 and 26. In addition, Theorem 3.2 and Corollary 3.4 in Ref. 17 show that if Ω is convex, and W_h consists of continuous functions that are piecewise polynomials of degree $k \geq 1$ over a quasiuniform mesh of Ω of size h , then our assumptions on the weight ϖ guarantee that (4.2) holds. Finally, in this setting, Ref. 36 constructs interpolants that satisfy (4.3).

As in the continuous case, we will say that a triple $(\mathbf{T}_h, \mathbf{u}_h, \mathbf{p}_h) \in W_h \times \mathbf{X}_h \times M_h$ is a discrete solution to (1.1) if

$$\left\{ \begin{array}{l} \int_{\Omega} \left(\nu(\mathbf{T}_h) \nabla \mathbf{u}_h : \nabla \mathbf{v}_h + (\mathbf{u}_h \cdot \nabla \mathbf{u}_h) \cdot \mathbf{v}_h + \frac{1}{2} \operatorname{div} \mathbf{u}_h \mathbf{u}_h \cdot \mathbf{v}_h \right. \\ \quad \left. - \mathbf{p}_h \operatorname{div} \mathbf{v}_h - \mathbf{T}_h \mathbf{g} \cdot \mathbf{v}_h \right) dx = 0, \\ \int_{\Omega} q_h \operatorname{div} \mathbf{u}_h \, dx = 0, \\ \int_{\Omega} (\kappa(\mathbf{T}_h) \nabla \mathbf{T}_h \cdot \nabla r_h - \mathbf{T}_h \mathbf{u}_h \cdot \nabla r_h) \, dx = \langle \mathcal{H}, r_h \rangle, \end{array} \right. \quad (4.4)$$

for all $\mathbf{v}_h \in \mathbf{X}_h$, $q_h \in M_h$, and $r_h \in W_h$. Our main objective here will be to show that, under similar assumptions to Theorem 3.2, problem (4.4) always has a solution and that, as $h \rightarrow 0$, these solutions weakly converge, up to subsequences, to a solution of (1.1) in the sense of Definition 2.2.

4.1. A discrete stationary heat equation with variable coefficient

As a first step to achieve our goals we must prove a discrete version of Proposition 3.1. The proof of the following result is, essentially, an adaption of Proposition 8.6.2 in Ref. 13.

Proposition 4.1. (Weighted stability) *If $\varkappa \in L^\infty(\Omega)$ is such that $0 < \varkappa_- \leq \varkappa \leq \varkappa_+$ and*

$$\Lambda(\varkappa) \geq \Lambda_1 := \max \left\{ \Lambda_0, 1 - \gamma \left(1 - \frac{1}{2\varkappa_+} \right) \right\}, \quad (4.5)$$

then, for all $h > 0$, we have that

$$\frac{\gamma}{2} \|\nabla r_h\|_{\mathbf{L}^2(\varpi^{\pm 1}, \Omega)} \leq \sup_{\theta_h \in W_h} \frac{\int_{\Omega} \varkappa \nabla r_h \cdot \nabla \theta_h \, dx}{\|\nabla \theta_h\|_{\mathbf{L}^2(\varpi^{\mp 1}, \Omega)}} \quad \forall r_h \in W_h, \quad (4.6)$$

where $\gamma > 0$ is the constant appearing in estimate (4.2).

Proof. As mentioned above, the proof essentially follows the perturbation argument developed in Proposition 8.6.2 of Ref. 13. Let us define the bilinear form

$$B_{\varkappa} : H_0^1(\varpi^{\pm 1}, \Omega) \times H_0^1(\varpi^{\mp 1}, \Omega) \rightarrow \mathbb{R}, \quad B_{\varkappa}(r, \theta) := \int_{\Omega} \left(1 - \frac{\varkappa}{\varkappa_+} \right) \nabla r \cdot \nabla \theta \, dx.$$

Note that, for $r \in H_0^1(\varpi^{\pm 1}, \Omega)$ and $\theta \in H_0^1(\varpi^{\mp 1}, \Omega)$,

$$|B_{\varkappa}(r, \theta)| \leq (1 - \Lambda(\varkappa)) \|\nabla r\|_{\mathbf{L}^2(\varpi^{\pm 1}, \Omega)} \|\nabla \theta\|_{\mathbf{L}^2(\varpi^{\mp 1}, \Omega)},$$

and that

$$\int_{\Omega} \nabla r \cdot \nabla \theta \, dx = B_{\varkappa}(r, \theta) + \frac{1}{\varkappa_+} \int_{\Omega} \varkappa \nabla r \cdot \nabla \theta \, dx.$$

Thus, owing to (4.2) we have that, for any $h > 0$ and any $r_h \in W_h$,

$$(\gamma + \Lambda(\varkappa) - 1) \|\nabla r_h\|_{\mathbf{L}^2(\varpi^{\pm 1}, \Omega)} \leq \frac{1}{\varkappa_+} \sup_{\theta_h \in W_h} \frac{\int_{\Omega} \varkappa \nabla r_h \cdot \nabla \theta_h \, dx}{\|\nabla \theta_h\|_{\mathbf{L}^2(\varpi^{\mp 1}, \Omega)}}.$$

The restriction on $\Lambda(\varkappa)$ allows us to conclude. $\square$

With the previous result at hand, we can show that a discrete version of (3.3) always has a solution for sufficiently small convection and that, more importantly, the discrete solutions are uniformly bounded with respect to h . Given $\varkappa \in L^\infty(\Omega)$

14 *A. Allendes, E. Otárola & A. J. Salgado*

with $0 < \kappa_- \leq \kappa \leq \kappa_+$, $\mathbf{u} \in \mathbf{H}_0^1(\Omega)$, and $\mathcal{H} \in H^{-1}(\varpi, \Omega)$ we consider the following problem: Find $T_h \in W_h$ such that

$$\int_{\Omega} (\kappa \nabla T_h \cdot \nabla r_h - T_h \mathbf{u} \cdot \nabla r_h) \, dx = \langle \mathcal{H}, r_h \rangle \quad \forall r_h \in W_h. \quad (4.7)$$

Corollary 4.1. (Well-posedness) *Assume that $\Lambda(\kappa) \geq \Lambda_1$, where Λ_1 is defined in (4.5), and that $\mathbf{u} \in \mathbf{H}_0^1(\Omega)$ satisfies*

$$\frac{2C_{e,2}}{\gamma} \|\nabla \mathbf{u}\|_{\mathbf{L}^2(\Omega)} \leq q < 1.$$

Then, for every $h > 0$, problem (4.7) has a unique solution. Moreover, T_h satisfies

$$\|\nabla T_h\|_{\mathbf{L}^2(\varpi, \Omega)} \leq \frac{2}{\gamma(1-q)} \|\mathcal{H}\|_{H^{-1}(\varpi, \Omega)}.$$

Proof. Repeat *verbatim* the proof of Proposition 3.2 replacing (3.4) by (4.6) and C_{κ} by $2/\gamma$. $\square$

4.2. Existence and stability

Having studied a discrete diffusion equation with variable coefficient on weighted spaces, we can proceed and show that, under similar assumptions to Theorem 3.2, our discrete problem (4.4), always has solutions and that, moreover, these are uniformly bounded with respect to $h > 0$. This will be the first step to show, via a compactness argument, the convergence of discrete solutions to a solution of (1.1), in the sense of Definition 2.2.

We proceed via a fixed point argument. We define, for each $h > 0$, the map

$$\begin{aligned} \mathfrak{F}_h : W_h \times \mathbf{X}_h \times M_h &\rightarrow W_h \times \mathbf{X}_h \times M_h, \quad (\theta_h, \mathbf{u}_h, p_h) \mapsto \mathfrak{F}_h(\theta_h, \mathbf{u}_h, p_h) \\ &= (\mathsf{T}_h, \mathbf{u}_h, \mathbf{p}_h) \end{aligned}$$

by the following procedure: Let the pair $(\mathbf{u}_h, \mathbf{p}_h) \in \mathbf{X}_h \times M_h$ be such that, for all $(\mathbf{v}_h, q_h) \in \mathbf{X}_h \times M_h$,

$$\begin{cases} \int_{\Omega} \left(\nu(\theta_h) \nabla \mathbf{u}_h : \nabla \mathbf{v}_h + (\mathbf{u}_h \cdot \nabla) \mathbf{u}_h \cdot \mathbf{v}_h + \frac{1}{2} \operatorname{div} \mathbf{u}_h \mathbf{u}_h \cdot \mathbf{v}_h \right) \, dx \\ \quad - \int_{\Omega} \mathbf{p}_h \operatorname{div} \mathbf{v}_h \, dx = \int_{\Omega} \theta_h \mathbf{g} \cdot \mathbf{v}_h \, dx, \\ \int_{\Omega} q_h \operatorname{div} \mathbf{u}_h \, dx = 0, \end{cases} \quad (4.8)$$

and $\mathsf{T}_h \in W_h$ be defined as the solution to

$$\int_{\Omega} (\kappa(\theta_h) \nabla \mathsf{T}_h \cdot \nabla r_h - \mathsf{T}_h \mathbf{u}_h \cdot \nabla r_h) \, dx = \langle \mathcal{H}, r_h \rangle \quad \forall r_h \in W_h. \quad (4.9)$$

For reasons similar to the continuous case, this map is well-defined, if we restrict it to a ball of appropriate size. To quantify that, we introduce

$$\begin{cases} \mathfrak{B}_T^h = \{\theta_h \in W_h : \|\nabla \theta_h\|_{\mathbf{L}^2(\varpi, \Omega)} \leq \tilde{S}\}, \\ \tilde{S} = \frac{\nu_-}{gC_{e,1}} \min \left\{ \frac{\nu_-}{C_P}, \frac{\gamma}{4C_{e,2}} \right\}, \\ \mathfrak{B}_u^h = \{\mathbf{u}_h \in \mathbf{X}_h : \|\nabla \mathbf{u}_h\|_{\mathbf{L}^2(\Omega)} \leq \tilde{G}\}, \\ \tilde{G} = \frac{\gamma}{4C_{e,2}}, \\ \mathfrak{B}_p^h = \{p_h \in M_h : \|p_h\|_{L^2(\Omega)} \leq \tilde{P}\}, \\ \tilde{P} = \frac{C_{e,1}g}{\beta} \tilde{S} \left(1 + \frac{\nu_+}{\nu_-} + \frac{3}{2} \frac{C_{4 \rightarrow 2}^2 C_{e,1}g}{\nu_-^2} \tilde{S} \right), \end{cases}$$

and $\mathfrak{B}^h = \mathfrak{B}_T^h \times \mathfrak{B}_u^h \times \mathfrak{B}_p^h$. Notice that neither $\tilde{S}$, $\tilde{G}$, nor $\tilde{P}$ are dependent on the parameter $h > 0$.

Proposition 4.2. ($\mathfrak{F}_h$ is well-defined) *Assume that $\Lambda(\kappa) \geq \Lambda_1$, where Λ_1 is defined in Proposition 4.1. If the heat source $\mathcal{H} \in H^{-1}(\varpi, \Omega)$ satisfies the estimate*

$$\|\mathcal{H}\|_{H^{-1}(\varpi, \Omega)} \leq \frac{\gamma \tilde{S}}{4},$$

then, for every $h > 0$, the mapping $\mathfrak{F}_h$ is well-defined on $\mathfrak{B}^h$. Moreover, $\mathfrak{F}_h(\mathfrak{B}^h) \subset \mathfrak{B}^h$.

Proof. The proof essentially repeats that of Proposition 3.3. For this reason, we only sketch it. Let $(\theta_h, \mathbf{u}_h, p_h) \in \mathfrak{B}^h$. Since we have skew symmetrized the convective term, we know that problem (4.8) always has a solution which satisfies

$$\|\nabla \mathbf{u}_h\|_{\mathbf{L}^2(\Omega)} \leq \frac{1}{\nu_-} \|\theta_h \mathbf{g}\|_{\mathbf{H}^{-1}(\Omega)} \leq \frac{C_{e,1}g}{\nu_-} \|\nabla \theta_h\|_{\mathbf{L}^2(\varpi, \Omega)} \leq \tilde{G},$$

$$\beta \|\mathbf{p}_h\|_{L^2(\Omega)} \leq \nu_+ \|\nabla \mathbf{u}_h\|_{\mathbf{L}^2(\Omega)} + \frac{3}{2} C_{4 \rightarrow 2}^2 \|\nabla \mathbf{u}_h\|_{\mathbf{L}^2(\Omega)}^2 + \frac{3}{2} \|\theta_h \mathbf{g}\|_{\mathbf{H}^{-1}(\Omega)} \leq \beta \tilde{P},$$

where we used (2.2) and the fact that $\theta_h \in \mathfrak{B}_T^h$, to obtain the estimate for $\mathbf{u}_h$, and the inf-sup condition (4.1) for the estimate of $\mathbf{p}_h$. In addition, we observe that the conditions on the data for this solution to be unique are met; see Theorem IV.3.1 in Ref. 25. Corollary 4.1, with $q = 1/2$, then implies that problem (4.9) has a unique solution, which satisfies

$$\|\nabla \mathbf{T}_h\|_{\mathbf{L}^2(\varpi)} \leq \frac{4}{\gamma} \|\mathcal{H}\|_{H^{-1}(\varpi)} \leq \tilde{S},$$

where we used the assumption on $\mathcal{H}$. Thus, as we intended to show, $\mathfrak{F}_h$ is well-defined on $\mathfrak{B}^h$ and $\mathfrak{F}_h(\mathfrak{B}^h) \subset \mathfrak{B}^h$. $\square$

We conclude by showing existence of solutions, via a fixed point argument.

Theorem 4.1. (Existence) *Assume that $\Lambda(\kappa) \geq \Lambda_1$, where Λ_1 is defined in Proposition 4.1. If the heat source $\mathcal{H} \in H^{-1}(\varpi, \Omega)$ satisfies the estimate*

$$\|\mathcal{H}\|_{H^{-1}(\varpi, \Omega)} \leq \frac{\gamma \tilde{S}}{4},$$

then, for every $h > 0$, there is a triple $(\mathbf{T}_h, \mathbf{u}_h, \mathbf{p}_h) \in \mathfrak{B}^h \subset W_h \times \mathbf{X}_h \times M_h$ that solves (4.4).

Proof. Since we are now in finite dimensions, we will apply Brouwer's fixed point theorem; see Theorem 3.2 in Ref. 16. For that, we only need to verify the continuity of $\mathfrak{F}_h$. This is achieved by repeating *verbatim* the proof of Lemma 3.1 and using that we are in finite dimensions to pass from weak to strong convergence. $\square$

4.3. Convergence

The results of the previous section show that, provided the heat source $\mathcal{H}$ and the oscillation of κ are not too large, then for every $h > 0$ problem (4.4) has a solution, and that this family of solutions remains bounded uniformly in $h > 0$. We can then pass to a (not relabeled) weakly convergent subsequence $(\mathbf{T}_h, \mathbf{u}_h, \mathbf{p}_h) \rightharpoonup (\mathbf{T}, \mathbf{u}, \mathbf{p})$. We will show here that this limit must be a solution to (1.1), in the sense of Definition 2.2.

We begin with some notation. We define

$$\begin{aligned} \hat{\mathfrak{B}}_T &= \{\theta \in H_0^1(\varpi, \Omega) : \|\nabla \theta\|_{L^2(\varpi, \Omega)} \leq \hat{S}\}, \quad \hat{\mathfrak{B}}_T^h = \hat{\mathfrak{B}}_T \cap W_h, \\ \hat{S} &= \min\{S, \tilde{S}\}, \quad \hat{\mathfrak{B}}_{\mathbf{u}} = \{\mathbf{u} \in \mathbf{H}_0^1(\Omega) : \|\nabla \mathbf{u}\|_{L^2(\Omega)} \leq \hat{G}\}, \\ \hat{\mathfrak{B}}_{\mathbf{u}}^h &= \hat{\mathfrak{B}}_{\mathbf{u}} \cap \mathbf{X}_h, \quad \hat{G} = \min\{G, \tilde{G}\}, \quad \hat{\mathfrak{B}}_p = \{p \in L_0^2(\Omega) : \|p\|_{L^2(\Omega)} \leq \hat{P}\}, \\ \hat{\mathfrak{B}}_p^h &= \hat{\mathfrak{B}}_p \cap M_h, \quad \hat{P} = \min\{P, \tilde{P}\}, \\ \hat{\mathfrak{B}} &= \hat{\mathfrak{B}}_T \times \hat{\mathfrak{B}}_{\mathbf{u}} \times \hat{\mathfrak{B}}_p, \quad \text{and} \quad H = \min\left\{\frac{\gamma \tilde{S}}{4}, \frac{S}{2C_\kappa}\right\}. \end{aligned}$$

Theorem 4.2. (Convergence) *Assume that $\Lambda(\kappa) \geq \Lambda_1$, where Λ_1 is defined in Proposition 4.1. If the heat source $\mathcal{H} \in H^{-1}(\varpi, \Omega)$ satisfies*

$$\|\mathcal{H}\|_{H^{-1}(\varpi)} \leq H,$$

then the family $\{(\mathbf{T}_h, \mathbf{u}_h, \mathbf{p}_h) \in \mathfrak{B}^h\}_{h>0}$ of solutions to (4.4) converges weakly (up to subsequences) to an element of $\mathfrak{B}$. Moreover, this limit is a solution to (1.1) in the sense of Definition 2.2.

Proof. The assumptions guarantee that we can invoke Theorem 4.1 to ascertain the existence of $(\mathbf{T}_h, \mathbf{u}_h, \mathbf{p}_h) \in \hat{\mathfrak{B}}_T^h \times \hat{\mathfrak{B}}_{\mathbf{u}}^h \times \hat{\mathfrak{B}}_p^h$ that solve (4.4). Moreover, since $\hat{\mathfrak{B}}_T^h \times \hat{\mathfrak{B}}_{\mathbf{u}}^h \times \hat{\mathfrak{B}}_p^h \subset \hat{\mathfrak{B}}$, this family of solutions remains in a bounded set, and we can extract weakly convergent subsequences which for simplicity of notation we do not

relabel. Let us denote this limit by $(\mathbf{T}, \mathbf{u}, \mathbf{p}) \in \hat{\mathfrak{B}}$, and show that it is a solution to (2.5).

Observe now that:

- The compact embedding $H_0^1(\varpi, \Omega) \hookrightarrow L^{1+\epsilon}(\Omega)$ implies, by continuity of ν and κ , that we have $\nu(\mathbf{T}_h) \rightarrow \nu(\mathbf{T})$ and $\kappa(\mathbf{T}_h) \rightarrow \kappa(\mathbf{T})$ almost everywhere in Ω .
- Let $q \in C_0^\infty(\Omega)$ with zero average, and $q_h \in M_h$ be its L^2 -projection onto M_h . Then we have that

$$\left| \int_{\Omega} q \operatorname{div} \mathbf{u} \, dx \right| \leq \left| \int_{\Omega} (q - q_h) \operatorname{div} \mathbf{u}_h \, dx \right| + \left| \int_{\Omega} q \operatorname{div} (\mathbf{u} - \mathbf{u}_h) \, dx \right| \rightarrow 0,$$

where we used that $\mathbf{u}_h$ is discretely solenoidal, the strong convergence $q_h \rightarrow q$ in $L_0^2(\Omega)$ and the weak convergence $\operatorname{div} \mathbf{u}_h \rightharpoonup \operatorname{div} \mathbf{u}$. In conclusion $\mathbf{u}$ is solenoidal.

- We now show that the pair $(\mathbf{T}, \mathbf{u})$ satisfies the heat equation with convection. Let now $r \in H_0^1(\varpi^{-1}, \Omega)$ be arbitrary and introduce $r_h = \pi_W r \in W_h$, where the operator π_W is the one that satisfies (4.3). The almost everywhere convergence of $\kappa(\mathbf{T}_h)$ then implies

$$\begin{aligned} \int_{\Omega} \kappa(\mathbf{T}_h) \nabla \mathbf{T}_h \cdot \nabla r_h \, dx &= \int_{\Omega} \kappa(\mathbf{T}_h) \nabla \mathbf{T}_h \cdot \nabla r \, dx + \int_{\Omega} \kappa(\mathbf{T}_h) \nabla \mathbf{T}_h \cdot \nabla (r_h - r) \, dx \\ &\rightarrow \int_{\Omega} \kappa(\mathbf{T}) \nabla \mathbf{T} \cdot \nabla r \, dx. \end{aligned}$$

Finally, let $k > 2d/(d-1)$ such that we have the compact embedding $H^1(\Omega) \hookrightarrow L^m(\Omega)$, with $k^{-1} + m^{-1} = 1/2$, i.e. $m < 2d$. Consequently, $\mathbf{u}_h \rightarrow \mathbf{u}$ in $\mathbf{L}^m(\Omega)$. This is sufficient to assert that

$$\int_{\Omega} \mathbf{T}_h \mathbf{u}_h \cdot \nabla r \, dx \rightarrow \int_{\Omega} \mathbf{T} \mathbf{u} \cdot \nabla r \, dx.$$

- It remains to deal with the momentum equation, but most of the terms are standard here and have been treated in several other works, see for instance Refs. 23, 27 37, 38 and 39. The only somewhat nonstandard term is

$$\int_{\Omega} \mathbf{T}_h \mathbf{g} \cdot \mathbf{v}_h \, dx,$$

but the estimates of Proposition 2.3 can be used to assert convergence of this term as well.

In conclusion, the limit is a solution and the theorem is proved. $\square$

5. A Posteriori Error Estimates

In this section, we design and analyze an *a posteriori* error estimator for the finite-dimensional approximation (4.4) of problem (2.5). To be able to do so, in addition to the assumptions stated in Sec. 2.2, we shall require that:

- The viscosity ν and the thermal diffusivity κ are independent of the temperature, i.e. they are *positive constants*.

In addition, to be able to develop an explicit *a posteriori* error estimator and show its reliability, we must be more specific in the structure of the discrete spaces we are dealing with. For this reason, in addition to the assumptions of Sec. 4, we assume that, for $h > 0$, the spaces W_h , $\mathbf{X}_h$, and M_h are constructed using finite elements over a conforming and shape regular mesh $\mathcal{T}_h = \{K\}$ of $\bar{\Omega}$. In this setting, however, the parameter h does not bear the meaning of a mesh size. Rather, it can be thought of as $h = 1/k$, where $k \in \mathbb{N}_0$ is the index set in a sequence of refinements of an original partition $\mathcal{T}_0$. For definiteness, we select the pair of discrete velocity/pressure spaces $(\mathbf{X}_h, M_h)$ from the following (popular) options:

- (1) The lowest order Taylor Hood element,^{19,30,46} which is defined by

$$\mathbf{X}_h = \{\mathbf{v}_h \in \mathbf{C}(\bar{\Omega}) : \forall K \in \mathcal{T}_h, \mathbf{v}_h|_K \in \mathbb{P}_2(K)^d\} \cap \mathbf{H}_0^1(\Omega), \quad (5.1)$$

$$M_h = \{q_h \in L_0^2(\Omega) \cap C(\bar{\Omega}) : \forall K \in \mathcal{T}_h, q_h|_K \in \mathbb{P}_1(K)\}. \quad (5.2)$$

- (2) The mini element, which is considered in Refs. 7 and 19 and is defined by

$$\mathbf{X}_h = \{\mathbf{v}_h \in \mathbf{C}(\bar{\Omega}) : \forall K \in \mathcal{T}_h, \mathbf{v}_h|_K \in [\mathbb{P}_1(K) \oplus \mathbb{B}(K)]^d\} \cap \mathbf{H}_0^1(\Omega), \quad (5.3)$$

$$M_h = \{q_h \in L_0^2(\Omega) \cap C(\bar{\Omega}) : \forall K \in \mathcal{T}_h, q_h|_K \in \mathbb{P}_1(K)\}, \quad (5.4)$$

where $\mathbb{B}(K)$ denotes the space spanned by local bubble functions.

Both pairs satisfy the compatibility condition (4.1) and are such that $\mathbf{X}_h \subset \mathbf{H}_0^1(\Omega)$ and $M_h \subset L_0^2(\Omega)$. We will set the finite element space W_h as

$$W_h = \{w_h \in C(\bar{\Omega}) : \forall K \in \mathcal{T}_h, w_h|_K \in \mathbb{P}_k(K)\} \cap H_0^1(\Omega),$$

where $k = 2$ when the Taylor–Hood element (5.1)–(5.2) is used to approximate the velocity and pressure variables and $k = 1$ for when the mini element (5.3)–(5.4) is considered. Notice that, for any $\varpi \in A_2$, $W_h \subset W_0^{1,\infty}(\Omega) \subset H_0^1(\varpi^{\pm 1}, \Omega)$.

We begin our analysis by introducing some preliminary notions. We define the temperature error $\mathbf{e}_T$, the velocity error $\mathbf{e}_u$, and the pressure error $\mathbf{e}_p$ as follows:

$$\mathbf{e}_T := T - T_h \in H_0^1(\varpi, \Omega), \quad \mathbf{e}_u := \mathbf{u} - \mathbf{u}_h \in \mathbf{H}_0^1(\Omega), \quad \mathbf{e}_p := p - p_h \in L_0^2(\Omega).$$

We also define, for an open set $D \subset \Omega$, the following norms on the space $H_0^1(\varpi, D) \times \mathbf{H}_0^1(D) \times L^2(D)$:

$$\|(w, \mathbf{w}, s)\|_D^2 := \nu \|\nabla \mathbf{w}\|_{\mathbf{L}^2(D)}^2 + \|\operatorname{div} \mathbf{w}\|_{L^2(D)}^2 + \|s\|_{L^2(D)}^2 + \|\nabla w\|_{\mathbf{L}^2(\varpi, D)}^2,$$

$$\|(w, \mathbf{w}, s)\|_D^2 := \nu \|\nabla \mathbf{w}\|_{\mathbf{L}^2(D)}^2 + \|s\|_{L^2(D)}^2 + \|\nabla w\|_{\mathbf{L}^2(\varpi, D)}^2.$$

5.1. Ritz projection

To perform a reliability analysis for the devised *a posteriori* error estimator we shall introduce a so-called Ritz projection (φ, Φ, ψ) of the residuals. This projection

is defined as the solution to the following problem: Find $(\varphi, \Phi, \psi) \in H_0^1(\varpi, \Omega) \times \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$ such that

$$\int_{\Omega} \nabla \varphi \cdot \nabla r \, dx = \Upsilon(r), \quad \forall r \in H_0^1(\varpi^{-1}, \Omega), \quad (5.5)$$

$$\nu \int_{\Omega} \nabla \Phi : \nabla \mathbf{v} \, dx = \Xi(\mathbf{v}), \quad \forall \mathbf{v} \in \mathbf{H}_0^1(\Omega), \quad (5.6)$$

$$\int_{\Omega} \psi q \, dx = \Sigma(q), \quad \forall q \in L_0^2(\Omega), \quad (5.7)$$

where the functionals $\Upsilon \in H^{-1}(\varpi, \Omega)$, $\Xi \in \mathbf{H}^{-1}(\Omega)$, and $\Sigma \in L_0^2(\Omega)$ are defined, respectively, by

$$\begin{aligned} \Upsilon(r) &:= \int_{\Omega} (\kappa \nabla \mathbf{e}_{\mathbf{T}} \cdot \nabla r - \mathbf{T} \mathbf{e}_{\mathbf{u}} \cdot \nabla r - \mathbf{e}_{\mathbf{T}} \mathbf{u}_h \cdot \nabla r) \, dx, \\ \Xi(\mathbf{v}) &:= \int_{\Omega} \left(\nu \nabla \mathbf{e}_{\mathbf{u}} : \nabla \mathbf{v} - \mathbf{e}_{\mathbf{p}} \operatorname{div} \mathbf{v} \right. \\ &\quad \left. + (\mathbf{u} \cdot \nabla) \mathbf{e}_{\mathbf{u}} \cdot \mathbf{v} + (\mathbf{e}_{\mathbf{u}} \cdot \nabla) \mathbf{u}_h \cdot \mathbf{v} + \frac{1}{2} \operatorname{div} \mathbf{e}_{\mathbf{u}} \mathbf{u}_h \cdot \mathbf{v} - \mathbf{e}_{\mathbf{T}} \mathbf{g} \cdot \mathbf{v} \right) \, dx, \\ \Sigma(q) &:= - \int_{\Omega} q \operatorname{div} \mathbf{e}_{\mathbf{u}} \, dx. \end{aligned} \quad (5.8)$$

The following result yields the well-posedness of problem (5.5)–(5.7). Recall that, by $C_{4 \rightarrow 2}$ we denote the best constant in the Sobolev embedding $\mathbf{H}_0^1(\Omega) \hookrightarrow \mathbf{L}^4(\Omega)$. To shorten notation we define

$$\mathfrak{B}(\mathbf{u}, \mathbf{u}_h) := \frac{C_{4 \rightarrow 2}^2}{\nu} \left(\|\nabla \mathbf{u}\|_{\mathbf{L}^2(\Omega)} + \frac{3}{2} \|\nabla \mathbf{u}_h\|_{\mathbf{L}^2(\Omega)} \right), \quad \mathfrak{A}(\mathbf{u}, \mathbf{u}_h) := 1 + \mathfrak{B}(\mathbf{u}, \mathbf{u}_h).$$

Proposition 5.1. (Ritz projection) *Problem (5.5)–(5.7) admits a unique solution $(\varphi, \Phi, \psi) \in H_0^1(\varpi, \Omega) \times \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$. In addition, we have the estimate*

$$\begin{aligned} \|(\varphi, \Phi, \psi)\|_{\Omega}^2 &\leq \left[3\mathfrak{A}(\mathbf{u}, \mathbf{u}_h)^2 + 2 \frac{C_1^2 C_{e,2}^2}{\nu} \|\nabla \mathbf{T}\|_{\mathbf{L}^2(\varpi, \Omega)}^2 \right] \nu \|\nabla \mathbf{e}_{\mathbf{u}}\|_{\mathbf{L}^2(\Omega)}^2 \\ &\quad + \|\operatorname{div} \mathbf{e}_{\mathbf{u}}\|_{L^2(\Omega)}^2 + \frac{3}{\nu} \|\mathbf{e}_{\mathbf{p}}\|_{L^2(\Omega)}^2 \\ &\quad + \left[\frac{3g^2 C_{e,1}^2}{\nu} + 2C_1^2 (\kappa + C_{e,2} \|\nabla \mathbf{u}_h\|_{\mathbf{L}^2(\Omega)})^2 \right] \|\nabla \mathbf{e}_{\mathbf{T}}\|_{\mathbf{L}^2(\varpi, \Omega)}^2, \end{aligned} \quad (5.9)$$

where C_1 denotes the constant in the inf-sup estimate (3.4) with $\varkappa = 1$.

Proof. Since $\Xi \in \mathbf{H}^{-1}(\Omega)$, the Lax–Milgram lemma immediately yields the existence of a unique $\Phi \in \mathbf{H}_0^1(\Omega)$ that solves problem (5.6). In addition, estimate (2.2)

20 *A. Allendes, E. Otárola & A. J. Salgado*

and the standard Sobolev embedding $\mathbf{H}_0^1(\Omega) \hookrightarrow \mathbf{L}^4(\Omega)$ yield

$$\begin{aligned} \sqrt{\nu} \|\nabla \Phi\|_{\mathbf{L}^2(\Omega)} &\leq \mathfrak{A}(\mathbf{u}, \mathbf{u}_h) \sqrt{\nu} \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)} \\ &\quad + \frac{1}{\sqrt{\nu}} \|\mathbf{e}_p\|_{L^2(\Omega)} + \frac{gC_{e,1}}{\sqrt{\nu}} \|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(\varpi, \Omega)}. \end{aligned} \quad (5.10)$$

On the other hand, since $\mathbf{e}_u \in \mathbf{H}_0^1(\Omega)$, similar arguments reveal the existence and uniqueness of $\psi \in L_0^2(\Omega)$ that solves problem (5.7) together with the bound

$$\|\psi\|_{L^2(\Omega)} \leq \|\operatorname{div} \mathbf{e}_u\|_{L^2(\Omega)}. \quad (5.11)$$

Next, we invoke the inf-sup condition (3.4) for the variational form of the Dirichlet Laplace operator on weighted spaces to conclude that there exists a unique $\varphi \in H_0^1(\varpi, \Omega)$ that solves (5.5). In addition, we have that φ satisfies the estimate

$$\begin{aligned} \|\nabla \varphi\|_{\mathbf{L}^2(\varpi, \Omega)} &\leq C_1 [\kappa \|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(\varpi, \Omega)} + C_{e,2} \|\nabla \mathbf{T}\|_{\mathbf{L}^2(\varpi, \Omega)} \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)} \\ &\quad + C_{e,2} \|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(\varpi, \Omega)} \|\nabla \mathbf{u}_h\|_{\mathbf{L}^2(\Omega)}], \end{aligned} \quad (5.12)$$

where C_1 denotes the constant in the inf-sup estimate (3.4) with $\varkappa = 1$.

The desired estimate (5.9) thus follows from collecting estimates (5.10), (5.11), and (5.12). This concludes the proof. $\square$

5.2. An upper bound for the error

We now prove that the energy norm of the error can be bounded in terms of the energy norm of the Ritz projection, which in turn will allow us to provide a computable upper bound for the error.

Theorem 5.1. (Upper bound for the error) *Assume that the solutions to (2.5) and (4.4) are such that the following inequalities hold:*

$$\begin{aligned} \max \left\{ C_\kappa C_{e,2}, \frac{3C_{4 \rightarrow 2}^2}{2\nu} \right\} \|\nabla \mathbf{u}_h\|_{\mathbf{L}^2(\Omega)} &\leq \frac{1}{16}, \\ \frac{C_{4 \rightarrow 2}^2}{\nu} \|\nabla \mathbf{u}\|_{\mathbf{L}^2(\Omega)} &\leq \frac{1}{16}, \\ \frac{gC_\kappa C_{e,1} C_{e,2}}{\nu} \|\nabla \mathbf{T}\|_{\mathbf{L}^2(\varpi, \Omega)} &\leq \frac{15}{16^2}. \end{aligned} \quad (5.13)$$

Then, we have that

$$|||(\mathbf{e}_T, \mathbf{e}_u, \mathbf{e}_p)|||_\Omega \lesssim |||(\varphi, \Phi, \psi)|||_\Omega,$$

where the hidden constant is independent of $(\mathbf{e}_T, \mathbf{e}_u, \mathbf{e}_p)$ and (φ, Φ, ψ) but depends on ν and the constant involved in the inf-sup condition (3.2).

Proof. We divide the proof in six steps.

Step 1. We first bound $\|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(\varpi, \Omega)}$. Owing to Proposition 3.1 we have that there is a positive constant C_κ such that the following inf-sup condition holds:

$$\|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(\varpi, \Omega)} \leq C_\kappa \sup_{r \in H_0^1(\varpi^{-1}, \Omega)} \frac{\int_\Omega \kappa \nabla \mathbf{e}_T \cdot \nabla r \, dx}{\|\nabla r\|_{\mathbf{L}^2(\varpi^{-1}, \Omega)}}. \quad (5.14)$$

To estimate the right-hand side of (5.14) we rewrite Eq. (5.5) as

$$\int_\Omega \kappa \nabla \mathbf{e}_T \cdot \nabla r \, dx = \int_\Omega (\mathbf{T} \mathbf{e}_u \cdot \nabla r + \mathbf{e}_T \mathbf{u}_h \cdot \nabla r + \nabla \varphi \cdot \nabla r) \, dx \quad \forall r \in H_0^1(\varpi^{-1}, \Omega).$$

Utilize the inf-sup condition (5.14) and estimate (2.3), twice, to obtain

$$\begin{aligned} \|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(\varpi, \Omega)} &\leq C_\kappa (C_{e,2} \|\nabla \mathbf{T}\|_{\mathbf{L}^2(\varpi, \Omega)} \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)} \\ &\quad + C_{e,2} \|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(\varpi, \Omega)} \|\nabla \mathbf{u}_h\|_{\mathbf{L}^2(\Omega)} + \|\nabla \varphi\|_{\mathbf{L}^2(\varpi, \Omega)}), \end{aligned}$$

which, upon utilizing the first estimate in (5.13), yields the bound

$$\frac{15}{16} \|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(\varpi, \Omega)} \leq C_\kappa (C_{e,2} \|\nabla \mathbf{T}\|_{\mathbf{L}^2(\varpi, \Omega)} \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)} + \|\nabla \varphi\|_{\mathbf{L}^2(\varpi, \Omega)}). \quad (5.15)$$

Step 2. We now control $\|\mathbf{e}_p\|_{L^2(\Omega)}$. To do this, we rewrite Eq. (5.6) as follows:

$$\begin{aligned} \int_\Omega \mathbf{e}_p \operatorname{div} \mathbf{v} \, dx &= \int_\Omega (\nu \nabla \mathbf{e}_u : \nabla \mathbf{v} + (\mathbf{u} \cdot \nabla) \mathbf{e}_u \cdot \mathbf{v} + (\mathbf{e}_u \cdot \nabla) \mathbf{u}_h \cdot \mathbf{v} \\ &\quad + \frac{1}{2} \operatorname{div} \mathbf{e}_u \mathbf{u}_h \cdot \mathbf{v} - \mathbf{e}_T \mathbf{g} \cdot \mathbf{v} - \nu \nabla \Phi : \nabla \mathbf{v}) \, dx =: \mathfrak{X}(\mathbf{v}) \quad \forall \mathbf{v} \in \mathbf{H}_0^1(\Omega). \end{aligned}$$

The standard inf-sup condition for the divergence (3.2) thus yields the estimate

$$B \|\mathbf{e}_p\|_{L^2(\Omega)} \leq \|\mathfrak{X}\|_{\mathbf{H}^{-1}(\Omega)}.$$

Moreover, Sobolev embeddings and estimate (2.2) imply that

$$\begin{aligned} \|\mathfrak{X}\|_{\mathbf{H}^{-1}(\Omega)} &\leq \nu (\|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)} + \|\nabla \Phi\|_{\mathbf{L}^2(\Omega)}) \\ &\quad + \mathfrak{B}(\mathbf{u}, \mathbf{u}_h) \nu \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)} + C_{e,1} g \|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(\varpi, \Omega)}. \end{aligned}$$

To conclude this step, we invoke (5.15) to arrive at

$$\begin{aligned} B \|\mathbf{e}_p\|_{L^2(\Omega)} &\leq \nu (\|\nabla \Phi\|_{\mathbf{L}^2(\Omega)} + \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)}) + \frac{16 C_\kappa C_{e,1} g}{15} \|\nabla \varphi\|_{\mathbf{L}^2(\varpi, \Omega)} \\ &\quad + \left[\mathfrak{B}(\mathbf{u}, \mathbf{u}_h) + \frac{16 C_\kappa C_{e,1} C_{e,2} g}{15 \nu} \|\nabla \mathbf{T}\|_{\mathbf{L}^2(\varpi, \Omega)} \right] \nu \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)}. \quad (5.16) \end{aligned}$$

Step 3. Set $\mathbf{v} = \mathbf{e}_u$ in (5.6) and $q = -\mathbf{e}_p$ in (5.7). Adding the obtained relations, and using the skew symmetry of convection when the first argument is solenoidal, we see that

$$\begin{aligned} \nu \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)}^2 &\leq \nu \|\nabla \Phi\|_{\mathbf{L}^2(\Omega)} \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)} + \|\psi\|_{L^2(\Omega)} \|\mathbf{e}_p\|_{L^2(\Omega)} \\ &\quad + \frac{3 C_{4 \rightarrow 2}^2}{2 \nu} \|\nabla \mathbf{u}_h\|_{\mathbf{L}^2(\Omega)} \nu \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)}^2 + C_{e,1} g \|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(\varpi, \Omega)} \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)}. \end{aligned}$$

22 *A. Allendes, E. Otárola & A. J. Salgado*

Estimate (5.15) and Young's inequality imply that

$$\begin{aligned} \nu(1 - \varepsilon_1) \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)}^2 &\leq \nu \|\nabla \Phi\|_{\mathbf{L}^2(\Omega)} \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)} + \|\mathbf{e}_p\|_{L^2(\Omega)} \|\psi\|_{L^2(\Omega)} \\ &\quad + \left[\frac{3C_{4 \rightarrow 2}^2}{2\nu} \|\nabla \mathbf{u}_h\|_{\mathbf{L}^2(\Omega)} + \frac{16C_\kappa C_{e,1} C_{e,2} g}{15\nu} \|\nabla \mathbf{T}\|_{\mathbf{L}^2(\varpi, \Omega)} \right] \\ &\quad \times \nu \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)}^2 + C_{\varepsilon_1} \left(\frac{16C_\kappa C_{e,1} C_{e,2} g}{15\sqrt{\nu}} \right)^2 \|\nabla \varphi\|_{\mathbf{L}^2(\varpi, \Omega)}^2. \end{aligned}$$

The pressure error estimate (5.16), and repeated applications of Young's inequality imply that, for some $\varepsilon \in (0, 1)$

$$\begin{aligned} (1 - \varepsilon) \nu \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)}^2 &\leq C_{\varepsilon, B} \nu \|\nabla \Phi\|_{\mathbf{L}^2(\Omega)}^2 + C_{\varepsilon, B, \nu} \|\psi\|_{L^2(\Omega)}^2 + C_{\varepsilon, B, \nu} \|\nabla \varphi\|_{\mathbf{L}^2(\varpi, \Omega)}^2 \\ &\quad + \left[\frac{3C_{4 \rightarrow 2}^2}{2\nu} \|\nabla \mathbf{u}_h\|_{\mathbf{L}^2(\Omega)} + \frac{16C_\kappa C_{e,1} C_{e,2} g}{15\nu} \|\nabla \mathbf{T}\|_{\mathbf{L}^2(\varpi, \Omega)} \right] \\ &\quad \times \nu \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)}^2 \\ &\quad + \frac{1}{2} \left[\mathfrak{B}(\mathbf{u}, \mathbf{u}_h) + \frac{16C_\kappa C_{e,1} C_{e,2} g}{15\nu} \|\nabla \mathbf{T}\|_{\mathbf{L}^2(\varpi, \Omega)} \right]^2 \nu \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)}^2. \end{aligned}$$

Now, estimates (5.13) imply that we can choose $\varepsilon \in (0, 1)$ so that

$$\nu \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)}^2 \lesssim \nu \|\nabla \Phi\|_{\mathbf{L}^2(\Omega)}^2 + \|\psi\|_{L^2(\Omega)}^2 + \|\nabla \varphi\|_{\mathbf{L}^2(\varpi, \Omega)}^2,$$

where the hidden constant depends on ε , B , and ν .

Step 5. The previous estimate combined with estimates (5.15) and (5.16) yields

$$\begin{aligned} &\|\nabla \mathbf{e}_\mathbf{T}\|_{\mathbf{L}^2(\varpi, \Omega)}^2 + \nu \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(\Omega)}^2 + \|\mathbf{e}_p\|_{L^2(\Omega)}^2 \\ &\lesssim \|\nabla \varphi\|_{\mathbf{L}^2(\varpi, \Omega)}^2 + \nu \|\nabla \Phi\|_{\mathbf{L}^2(\Omega)}^2 + \|\psi\|_{L^2(\Omega)}^2. \end{aligned}$$

Step 6. Conclude with the obvious observation that, since $\operatorname{div} \mathbf{e}_u \in L_0^2(\Omega)$, (5.7) implies

$$\|\operatorname{div} \mathbf{e}_u\|_{L^2(\Omega)} \leq \|\psi\|_{L^2(\Omega)}.$$

The claimed upper bound for the error has been obtained, and the theorem has been proved. $\square$

5.3. A residual-type error estimator

In this section, we design an *a posteriori* error estimator for the finite-dimensional approximation (4.4) of problem (2.5). To be able to do so, we will assume that the singular forcing $\mathcal{H}$ has a particular structure, that is:

- The singular forcing term $\mathcal{H}$ has the form $\mathcal{H} = \mathfrak{h} \delta_z$, where δ_z corresponds to the Dirac delta supported at the interior point $z \in \Omega$ and $\mathfrak{h} \in \mathbb{R}$.

To handle such a singular forcing term, we introduce the weight $\mathbf{d}_z^\alpha$, where $\mathbf{d}_z(x) := |x - z|$ and $\alpha \in (d - 2, d)$. We must immediately notice the following two important properties: First, owing to Remark 2.1, we have that the weight $\mathbf{d}_z^\alpha$ is such that $\mathbf{d}_z^\alpha \in A_2(\Omega)$ and $\mathbf{d}_z^{-\alpha} \in A_1$. Second, $\delta_z \in H^{-1}(\mathbf{d}_z^\alpha, \Omega)$; see Lemma 7.1.3 in Ref. 32 and Remark 21.18 in Ref. 29. Simply put, all the assumptions we have made so far apply to this particular choice of $\mathcal{H}$ and weight.

5.3.1. Notation

Before presenting and analyzing our *a posteriori* error estimator we first need to introduce and set some notation. We recall that $\mathcal{T}_h = \{K\}$ is a conforming and shape regular partition of $\bar{\Omega}$ into closed simplices K with size $h_K = \text{diam}(K) \approx |K|^{1/d}$. We denote by $\mathcal{S}$ the set of *internal* $(d-1)$ -dimensional inter-element boundaries S of $\mathcal{T}_h$. For $S \in \mathcal{S}$, we indicate by h_S the diameter of S . If $K \in \mathcal{T}_h$, we define $\mathcal{S}_K$ as the subset of $\mathcal{S}$ that contains the sides of K . For $S \in \mathcal{S}$, we set $\mathcal{N}_S = \{K^+, K^-\}$, where $K^+, K^- \in \mathcal{T}_h$ are such that $S = K^+ \cap K^-$. For $K \in \mathcal{T}_h$, we define the following *stars* or *patches* associated with the element K

$$\mathcal{N}_K := \{K' \in \mathcal{T}_h : \mathcal{S}_K \cap \mathcal{S}_{K'} \neq \emptyset\}, \quad \mathcal{S}_K := \{K' \in \mathcal{T}_h : K \cap K' \neq \emptyset\}. \quad (5.17)$$

In an abuse of notation, below we denote by $\mathcal{N}_K$ and $\mathcal{S}_K$ either the sets themselves, or the union of its elements.

5.3.2. A posteriori error estimator

We define an error estimator that can be decomposed as the sum of two contributions: a contribution related to the discretization of the stationary Navier–Stokes equations and another one associated to the discretization of the stationary heat equation with convection and singular forcing.

To present the contribution related to the stationary Navier–Stokes equations, we define, for an element $K \in \mathcal{T}_h$ and an internal side $S \in \mathcal{S}$, the *element residual* $\mathcal{R}_K$ and the *interelement residual* $\mathcal{J}_S$ as

$$\begin{aligned} \mathcal{R}_K &:= \left(\nu \Delta \mathbf{u}_h - (\mathbf{u}_h \cdot \nabla) \mathbf{u}_h - \frac{1}{2} \text{div} \mathbf{u}_h \mathbf{u}_h - \nabla \mathbf{p}_h + \mathbb{T}_h \mathbf{g} \right) |_K, \\ \mathcal{J}_S &:= \llbracket (\nu \nabla \mathbf{u}_h - \mathbf{p}_h \mathbf{I}) \cdot \mathbf{n} \rrbracket, \end{aligned} \quad (5.18)$$

where $(\mathbb{T}_h, \mathbf{u}_h, \mathbf{p}_h)$ denotes a solution to the discrete problem (4.4) and $\mathbf{I} \in \mathbb{R}^{d \times d}$ denotes the identity matrix. For a discrete tensor-valued function $\mathbf{v}_h$, we denote by $\llbracket \mathbf{v}_h \cdot \mathbf{n} \rrbracket$ the jump, which is defined, on the internal side $S \in \mathcal{S}$ shared by the distinct elements $K^+, K^- \in \mathcal{N}_S$, by

$$\llbracket \mathbf{v}_h \cdot \mathbf{n} \rrbracket = \mathbf{v}_h|_{K^+} \cdot \boldsymbol{\nu}^+ + \mathbf{v}_h|_{K^-} \cdot \mathbf{n}^-.$$

Here $\mathbf{n}^+, \mathbf{n}^-$ are unit normals on S pointing toward K^+, K^- , respectively. With $\mathcal{R}_K$ and $\mathcal{J}_S$ at hand, we define, for $K \in \mathcal{T}_h$, the *element indicator*

$$\mathcal{E}_K^2 := h_K^2 \|\mathcal{R}_K\|_{\mathbf{L}^2(K)}^2 + \|\text{div} \mathbf{u}_h\|_{L^2(K)}^2 + h_K \|\mathcal{J}_S\|_{\mathbf{L}^2(\partial K \setminus \partial \Omega)}^2. \quad (5.19)$$

We now introduce the contribution associated to the stationary heat equation with convection. To accomplish this task, we define, for $K \in \mathcal{T}_h$ and an internal side $S \in \mathcal{S}$, the *element residual* $\mathfrak{R}_K$ and the *interelement residual* $\mathfrak{J}_S$ as

$$\begin{aligned}\mathfrak{R}_K &:= (\kappa \Delta \mathbf{T}_h - \mathbf{u}_h \cdot \nabla \mathbf{T}_h - \operatorname{div} \mathbf{u}_h \mathbf{T}_h)|_K, \\ \mathfrak{J}_S &:= \llbracket (\kappa \nabla \mathbf{T}_h - \mathbf{T}_h \mathbf{u}_h) \cdot \mathbf{n} \rrbracket.\end{aligned}\tag{5.20}$$

With $\mathfrak{R}_K$ and $\mathfrak{J}_S$ at hand, we define, for $K \in \mathcal{T}_h$ and $\alpha \in (d-2, d)$,

$$\begin{aligned}\mathfrak{E}_K^2 &:= h_K^2 D_K^\alpha \|\mathfrak{R}_K\|_{L^2(K)}^2 + h_K D_K^\alpha \|\mathfrak{J}_S\|_{L^2(\partial K \setminus \partial \Omega)}^2 \\ &\quad + |\mathfrak{h}| h_K^{\alpha+2-d} \#(\{z\} \cap K),\end{aligned}\tag{5.21}$$

where $D_K := \max_{x \in K} |x - z|$. For a set E , by $\#(E)$ we mean its cardinality. Thus, $\#(\{z\} \cap K)$ equals one if $z \in K$ and zero otherwise. Here, we must recall that we consider our elements K to be closed sets.

With all these ingredients at hand, we define the local error indicator $\mathcal{E}_K^2 := \mathcal{E}_K^2 + \mathfrak{E}_K^2$ and the *a posteriori* error estimators

$$\mathcal{E}_h := \left[\sum_{K \in \mathcal{T}_h} \mathcal{E}_K^2 \right]^{\frac{1}{2}}, \quad \mathfrak{E}_h := \left[\sum_{K \in \mathcal{T}_h} \mathfrak{E}_K^2 \right]^{\frac{1}{2}}, \quad \mathcal{E}_h^2 := \left[\sum_{K \in \mathcal{T}_h} \mathcal{E}_K^2 \right]^{\frac{1}{2}}.\tag{5.22}$$

5.4. Reliability estimates

We present the following global reliability estimate for the estimator $\mathcal{E}_h$.

Theorem 5.2. (Global reliability) *Let $(\mathbf{T}, \mathbf{u}, \mathbf{p}) \in H_0^1(\mathbf{d}_z^\alpha, \Omega) \times \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$ be a solution to (2.5) and $(\mathbf{T}_h, \mathbf{u}_h, \mathbf{p}_h) \in W_h \times \mathbf{X}_h \times M_h$ be its finite element approximation obtained as solution to (4.4). Let $\alpha \in (d-2, d)$. In the framework of Theorem 5.1, we have the following a posteriori error estimate:*

$$\|(\mathbf{e}_\mathbf{T}, \mathbf{e}_\mathbf{u}, \mathbf{e}_\mathbf{p})\|_\Omega \lesssim \mathcal{E}_h,\tag{5.23}$$

where the hidden constant is independent of the continuous and discrete solutions, the size of the elements in the mesh $\mathcal{T}_h$, and $\#\mathcal{T}_h$.

Proof. We proceed in several steps.

Step 1. Invoke Theorem 5.1, and the obvious bound $\|\operatorname{div} \Phi\|_{L^2(\Omega)} \leq \|\nabla \Phi\|_{\mathbf{L}^2(\Omega)}$ to arrive at the estimate

$$\|(\mathbf{e}_\mathbf{T}, \mathbf{e}_\mathbf{u}, \mathbf{e}_\mathbf{p})\|_\Omega^2 \lesssim \|\nabla \varphi\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, \Omega)}^2 + \|\nabla \Phi\|_{\mathbf{L}^2(\Omega)}^2 + \|\psi\|_{L^2(\Omega)}^2.$$

It thus suffices to bound $\|\nabla \Phi\|_{\mathbf{L}^2(\Omega)}$, $\|\psi\|_{L^2(\Omega)}$, and $\|\nabla \varphi\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, \Omega)}$.

Step 2. We control $\|\nabla \Phi\|_{\mathbf{L}^2(\Omega)}$. To accomplish this task, we invoke Eq. (5.6), the fact that $(\mathbf{T}, \mathbf{u}, \mathbf{p})$ solves problem (2.5), and an integration by parts formula to conclude that, for every $\mathbf{v} \in \mathbf{H}_0^1(\Omega)$, we have

$$\mathbf{I} := \int_\Omega \nu \nabla \Phi : \nabla \mathbf{v} \, dx = \sum_{K \in \mathcal{T}_h} \int_K \mathcal{R}_K \cdot \mathbf{v} \, dx + \sum_{S \in \mathcal{S}} \int_S \mathcal{J}_S \cdot \mathbf{v} \, ds.\tag{5.24}$$

Denote by I_h the Clément interpolation operator. We utilize the first equation of problem (4.4) with $\mathbf{v}_h = I_h \mathbf{v}$ and an integration by parts formula, again, to arrive at

$$\mathbf{I} = \sum_{K \in \mathcal{T}_h} \int_K \mathcal{R}_K \cdot (\mathbf{v} - I_h \mathbf{v}) \, dx + \sum_{S \in \mathcal{S}} \int_S \mathcal{J}_S \cdot (\mathbf{v} - I_h \mathbf{v}) \, ds.$$

We now invoke standard approximation properties for the interpolation operator I_h and a scaled trace inequality to conclude that

$$|\mathbf{I}| \lesssim \sum_{K \in \mathcal{T}_h} \left(h_K \|\mathcal{R}_K\|_{\mathbf{L}^2(K)} \|\nabla \mathbf{v}\|_{\mathbf{L}^2(S_K)} + \sum_{S \in \mathcal{S}_K} h_K^{\frac{1}{2}} \|\mathcal{J}_S\|_{\mathbf{L}^2(S)} \|\nabla \mathbf{v}\|_{\mathbf{L}^2(S_K)} \right).$$

Set $\mathbf{v} = \Phi$, use the Cauchy–Schwarz inequality in $\mathbb{R}^{\#\mathcal{T}_h}$ and the finite overlapping property of stars to obtain

$$\|\nabla \Phi\|_{\mathbf{L}^2(\Omega)} \lesssim \left(\sum_{K \in \mathcal{T}_h} h_K^2 \|\mathcal{R}_K\|_{\mathbf{L}^2(K)}^2 + h_K \|\mathcal{J}_S\|_{\mathbf{L}^2(\partial K \setminus \partial \Omega)}^2 \right)^{\frac{1}{2}}. \quad (5.25)$$

Step 3. In this step we bound $\|\psi\|_{L^2(\Omega)}$. Set $q = \psi$ in (5.7). This yields

$$\begin{aligned} \|\psi\|_{L^2(\Omega)}^2 &= \int_{\Omega} \psi \operatorname{div} \mathbf{u}_h \, dx \leq \sum_{K \in \mathcal{T}_h} \|\psi\|_{L^2(K)} \|\operatorname{div} \mathbf{u}_h\|_{L^2(K)} \\ &\leq \left(\sum_{K \in \mathcal{T}_h} \|\operatorname{div} \mathbf{u}_h\|_{L^2(K)}^2 \right)^{\frac{1}{2}} \|\psi\|_{L^2(\Omega)}. \end{aligned} \quad (5.26)$$

Notice that we have used that $\int_{\Omega} q \operatorname{div} \mathbf{u} \, dx = 0$ for every $q \in L_0^2(\Omega)$.

Step 4. Estimates (5.25) and (5.26) immediately yield that

$$(\|\nabla \Phi\|_{\mathbf{L}^2(\Omega)}^2 + \|\psi\|_{L^2(\Omega)}^2)^{\frac{1}{2}} \lesssim \left(\sum_{K \in \mathcal{T}_h} \mathcal{E}_K^2 \right)^{\frac{1}{2}} = \mathcal{E}_h. \quad (5.27)$$

Step 5. Our goal now is to bound $\|\nabla \varphi\|_{\mathbf{L}^2(\mathbf{d}_z^{\alpha}, \Omega)}$. To accomplish this task, we invoke the problem that φ solves, i.e. problem (5.5), and the fact that $(\mathbf{T}, \mathbf{u}, \mathbf{p})$ solves (2.5). These arguments, combined with an integration by parts formula, yield

$$\mathbf{II} := \int_{\Omega} \nabla \varphi \cdot \nabla r \, dx = \langle \mathbf{h} \delta_z, r \rangle + \sum_{K \in \mathcal{T}_h} \int_K \mathfrak{R}_K r \, dx + \sum_{S \in \mathcal{S}} \int_S \mathfrak{J}_S r \, ds,$$

for every $r \in H_0^1(\mathbf{d}_z^{-\alpha}, \Omega)$. We recall that $\mathfrak{R}_K$ and $\mathfrak{J}_S$ are defined in (5.20). Invoke the discrete problem (4.4) and an integration by parts formula, again, to arrive at

$$\mathbf{II} = \langle \mathbf{h} \delta_z, r - \pi_W r \rangle + \sum_{K \in \mathcal{T}_h} \int_K \mathfrak{R}_K (r - \pi_W r) \, dx + \sum_{S \in \mathcal{S}} \int_S \mathfrak{J}_S (r - \pi_W r) \, ds,$$

where we recall that π_W denotes the quasi-interpolation operator onto W_h , constructed in Ref. 36, that satisfies (4.3). We control $\langle \mathbf{h} \delta_z, r - \pi_W r \rangle$ on the basis of

26 *A. Allendes, E. Otárola & A. J. Salgado*

Theorem 4.7 in Ref. 1 and stability and interpolation estimates for π_W derived in Ref. 36. In fact, let K be such that $z \in K$, then

$$\begin{aligned} |\langle \mathfrak{h} \delta_z, r - \pi_W r \rangle| &\lesssim |\mathfrak{h}| h_K^{\frac{\alpha}{2} - \frac{d}{2}} \|r - \pi_W r\|_{L^2(d_z^{-\alpha}, K)} \\ &\quad + |\mathfrak{h}| h_K^{\frac{\alpha}{2} + 1 - \frac{d}{2}} \|\nabla(r - \pi_W r)\|_{L^2(d_z^{-\alpha}, K)} \\ &\lesssim |\mathfrak{h}| h_K^{\frac{\alpha}{2} + 1 - \frac{d}{2}} \|\nabla r\|_{L^2(d_z^{-\alpha}, S_K)}. \end{aligned}$$

Notice now that

$$\begin{aligned} \int_K \Re_K(r - \pi_W r) dx &\leq \|\Re_K\|_{L^2(K)} \|r - \pi_W r\|_{L^2(K)} \\ &\lesssim h_K D_K^{\frac{\alpha}{2}} \|\Re_K\|_{L^2(K)} \|\nabla r\|_{L^2(d_z^{-\alpha}, S_K)}, \end{aligned}$$

where we have used Proposition 4 in Ref. 4. Similar arguments yield, upon using Proposition 5 in Ref. 4, the control of the jump term. We can thus invoke the inf-sup condition (3.4) to arrive at the estimate

$$\|\nabla \varphi\|_{L^2(d_z^\alpha, \Omega)} \lesssim \sup_{r \in H_0^1(d_z^{-\alpha}, \Omega)} \frac{\int_\Omega \nabla \varphi \cdot \nabla r}{\|\nabla r\|_{L^2(d_z^{-\alpha}, \Omega)}} \lesssim \mathfrak{E}_h. \quad (5.28)$$

Step 6. Collecting the estimates (5.27) and (5.28) we obtain the reliability bound (5.23). This concludes the proof. $\square$

5.5. Efficiency estimates

In this section, we analyze efficiency properties for the local error indicator $\mathcal{E}_K$ on the basis of standard bubble function arguments.⁴⁷ Before proceeding with such analysis, we introduce the following notation: For an edge, triangle or tetrahedron G , let $\mathcal{V}(G)$ be the set of vertices of G . With this notation at hand, we introduce the following standard element and edge bubble functions. Let $K \in \mathcal{T}_h$ and $S \in \mathcal{S}$. We define

$$\Upsilon_K := (d+1)^{d+1} \prod_{v \in \mathcal{V}(K)} \lambda_{v|K}, \quad \Upsilon_S := d^d \prod_{v \in \mathcal{V}(S)} \lambda_{v|K} \quad \text{with } K \in \mathcal{N}_S. \quad (5.29)$$

In these formulas, by $\lambda_{v|K}$, we denote the barycentric coordinate function associated to $v \in \mathcal{V}(K)$.

We will also make use of the following bubble functions, whose construction we owe to Ref. 1. Given $K \in \mathcal{T}_h$, we introduce Ψ_K , which satisfies $0 \leq \Psi_K \leq 1$,

$$\Psi_K(z) = 0, \quad |K| \lesssim \int_K \Psi_K dx, \quad \|\nabla \Psi_K\|_{L^\infty(R_K)} \lesssim h_K^{-1}, \quad (5.30)$$

and there exists a simplex $K^* \subset K$ such that $R_K := \text{supp}(\Psi_K) \subset K^*$. Notice that, since Ψ_K satisfies (5.30), we have that, for every $m \in \mathbb{N}$

$$\|\theta\|_{L^2(R_K)} \lesssim \|\Psi_K^{\frac{1}{2}} \theta\|_{L^2(R_K)} \quad \forall \theta \in \mathbb{P}_m(R_K), \quad (5.31)$$

where the hidden constant depends on m , but does not depend on θ or K . Given $S \in \mathcal{S}$, we also introduce an edge bubble function Ψ_S , which satisfies $0 \leq \Psi_S \leq 1$,

$$\Psi_S(z) = 0, \quad |S| \lesssim \int_S \Psi_S \, ds, \quad \|\nabla \Psi_S\|_{\mathbf{L}^\infty(R_S)} \lesssim h_S^{-1}, \quad (5.32)$$

and $R_S := \text{supp}(\Psi_S)$ is such that, if $\mathcal{N}_S = \{K, K'\}$, there are simplices $K_* \subset K$ and $K'_* \subset K'$ such that $R_S \subset K_* \cup K'_* \subset K \cup K'$.

The following identities are essential to perform the upcoming efficiency analysis. Invoke (5.24), (5.6), and (5.8) to arrive at

$$\begin{aligned} \sum_{K \in \mathcal{T}_h} \int_K \mathcal{R}_K \cdot \mathbf{v} \, dx + \sum_{S \in \mathcal{S}} \int_S \mathcal{J}_S \cdot \mathbf{v} \, ds &= \int_\Omega \nu \nabla \Phi : \nabla \mathbf{v} \, dx = \int_\Omega \left(\nu \nabla \mathbf{e}_u : \nabla \mathbf{v} \right. \\ &\quad \left. - \mathbf{e}_p \operatorname{div} \mathbf{v} + (\mathbf{u} \cdot \nabla) \mathbf{e}_u \cdot \mathbf{v} + (\mathbf{e}_u \cdot \nabla) \mathbf{u}_h \cdot \mathbf{v} + \frac{1}{2} \operatorname{div} \mathbf{e}_u \mathbf{u}_h \cdot \mathbf{v} - \mathbf{e}_T \mathbf{g} \cdot \mathbf{v} \right) dx \end{aligned} \quad (5.33)$$

for all $\mathbf{v} \in \mathbf{H}_0^1(\Omega)$. We recall the reader that $\mathcal{R}_K$ and $\mathcal{J}_S$ are defined in (5.18). Similarly, for every $r \in H_0^1(\mathbf{d}_z^{-\alpha}, \Omega)$, we have

$$\begin{aligned} \sum_{K \in \mathcal{T}_h} \int_K \mathcal{R}_K r \, dx + \sum_{S \in \mathcal{S}} \int_S \mathcal{J}_S r \, ds + \langle \mathfrak{h} \delta_z, r \rangle &= \int_\Omega \nabla \varphi \cdot \nabla r \, dx \\ &= \int_\Omega (\kappa \nabla \mathbf{e}_T \cdot \nabla r - \mathbf{T} \mathbf{e}_u \cdot \nabla r - \mathbf{e}_T \mathbf{u}_h \cdot \nabla r) \, dx. \end{aligned} \quad (5.34)$$

In order to be in position to prove an efficiency result, we must obtain higher integrability of the velocity in our problem in the case $d = 3$. The following result provides this.

Proposition 5.2. (Higher integrability) *Let $d = 3$, $\alpha \in (1, 3)$, and $(\mathbf{T}, \mathbf{u}, \mathbf{p}) \in H_0^1(\mathbf{d}_z^\alpha, \Omega) \times \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$ denote the solution to (2.5) in the sense of Definition 2.2 with κ and ν constant, and $\mathcal{H} = \mathfrak{h} \delta_z$, with $z \in \Omega$. Then we have that*

$$\frac{1}{\nu} \|\nabla \mathbf{u}\|_{\mathbf{L}^3(\Omega)} \leq C,$$

where C denotes a positive constant.

Proof. We recall that we are assuming that the domain Ω is at least Lipschitz. Write the momentum component of the stationary Navier–Stokes equations as

$$-\nu \Delta \mathbf{u} + \nabla \mathbf{p} = \mathbf{g} \mathbf{T} - (\mathbf{u} \cdot \nabla) \mathbf{u}.$$

In three dimensions, we have that $\nabla \mathbf{u} \in \mathbf{L}^3(\Omega)$, and since $\alpha \in (1, 3)$ that $\mathbf{g} \mathbf{T} \in \mathbf{L}^2(\Omega)$. Owing then to Corollary 1.7 in Ref. 34 (with $\alpha = -1$ and $q = 2$) the result follows. $\square$

Our efficiency result is as follows.

Theorem 5.3. (Efficiency) *Let $(\mathbf{T}, \mathbf{u}, \mathbf{p}) \in H_0^1(\mathbf{d}_z^\alpha, \Omega) \times \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$ be a solution to (2.5) and $(\mathbf{T}_h, \mathbf{u}_h, \mathbf{p}_h) \in W_h \times \mathbf{X}_h \times M_h$ be its finite element approximation obtained as solution to (4.4). Let $\alpha \in (d-2, d)$. In the framework of Theorem 5.1, we have the following efficiency estimate:*

$$\begin{aligned} \mathcal{E}_K^2 &\lesssim \sum_{K' \in \mathcal{S}_K} (\|\mathbf{e}_\mathbf{T}\|_{L^2(\mathbf{d}_z^\alpha, K')}^2 + \nu^2 \|\mathbf{e}_\mathbf{u}\|_{\mathbf{H}^1(K')}^2 + \|\mathbf{e}_\mathbf{p}\|_{L^2(K')}^2) \\ &\quad + \|\operatorname{div} \mathbf{e}_\mathbf{u}\|_{L^2(K)}^2, \end{aligned} \quad (5.35)$$

where the hidden constant is independent of the continuous and discrete solutions, the size of the elements in the mesh $\mathcal{T}_h$, and $\#\mathcal{T}_h$.

Proof. We examine each of the contributions of $\mathcal{E}_K$ separately so the proof involves several steps.

Step 1. Let $K \in \mathcal{T}_h$. We bound the term $h_K^2 \|\mathcal{R}_K\|_{\mathbf{L}^2(K)}^2$ in (5.19). To accomplish this task, we first utilize estimate (5.31) and write

$$\|\mathcal{R}_K\|_{\mathbf{L}^2(K)}^2 = \int_K |\mathcal{R}_K|^2 dx \lesssim \int_K |\mathcal{R}_K|^2 \Psi_K dx = \int_K \mathcal{R}_K \cdot \boldsymbol{\Theta}_K dx, \quad (5.36)$$

where $\boldsymbol{\Theta}_K := \Psi_K \mathcal{R}_K$. Set $\mathbf{v} = \boldsymbol{\Theta}_K$ as a test function on identity (5.33) and utilize that $\operatorname{supp} \boldsymbol{\Theta}_K \subset K^* \subset K$. We proceed differently according to the spatial dimension. If $d = 2$, then we have

$$\begin{aligned} \int_K \mathcal{R}_K \cdot \boldsymbol{\Theta}_K dx &\leq (\nu \|\nabla \mathbf{e}_\mathbf{u}\|_{\mathbf{L}^2(K)} + \|\mathbf{e}_\mathbf{p}\|_{L^2(K)}) \|\nabla \boldsymbol{\Theta}_K\|_{\mathbf{L}^2(K)} \\ &\quad + \|\mathbf{g}\|_{\mathbf{L}^\infty(K)} \|\mathbf{e}_\mathbf{T}\|_{L^2(\mathbf{d}_z^\alpha, K)} \|\boldsymbol{\Theta}_K\|_{\mathbf{L}^2(\mathbf{d}_z^{-\alpha}, K)} + \left(\|\mathbf{u}\|_{\mathbf{L}^2(K)} \|\nabla \mathbf{e}_\mathbf{u}\|_{\mathbf{L}^2(K)} \right. \\ &\quad \left. + \|\mathbf{e}_\mathbf{u}\|_{\mathbf{L}^2(K)} \|\nabla \mathbf{u}_h\|_{\mathbf{L}^2(K)} + \frac{1}{2} \|\nabla \mathbf{e}_\mathbf{u}\|_{\mathbf{L}^2(K)} \|\mathbf{u}_h\|_{\mathbf{L}^2(K)} \right) \|\boldsymbol{\Theta}_K\|_{\mathbf{L}^\infty(K)}. \end{aligned} \quad (5.37)$$

Now, observe that estimate (5.6) in Ref. 1 yields $\|\boldsymbol{\Theta}_K\|_{\mathbf{L}^2(\mathbf{d}_z^{-\alpha}, K)} \lesssim D_K^{-\alpha/2} \|\mathcal{R}_K\|_{\mathbf{L}^2(K)}$ while basic arguments reveal that $\|\nabla \boldsymbol{\Theta}_K\|_{\mathbf{L}^2(K)} \lesssim h_K^{-1} \|\mathcal{R}_K\|_{\mathbf{L}^2(K)}$ and $\|\boldsymbol{\Theta}_K\|_{\mathbf{L}^\infty(K)} \lesssim h_K^{-1} \|\mathcal{R}_K\|_{\mathbf{L}^2(K)}$. With these estimates at hand, we utilize (5.36) and (5.13) to obtain

$$h_K^2 \|\mathcal{R}_K\|_{\mathbf{L}^2(K)}^2 \lesssim \|\mathbf{e}_\mathbf{T}\|_{L^2(\mathbf{d}_z^\alpha, K)}^2 + \nu^2 \|\mathbf{e}_\mathbf{u}\|_{\mathbf{H}^1(K)}^2 + \|\mathbf{e}_\mathbf{p}\|_{L^2(K)}^2. \quad (5.38)$$

Observe that $D_K^{-\frac{\alpha}{2}} = (\max_{x \in K} |x - z|)^{-\frac{\alpha}{2}} \lesssim h_K^{-\frac{\alpha}{2}} \lesssim h_K^{-1}$, because $h_K \lesssim D_K$ and $\alpha \in (0, 2)$; see Sec. 5.2 in Ref. 1 for details. If $d = 3$, we proceed as follows:

$$\begin{aligned} \int_K \mathcal{R}_K \cdot \boldsymbol{\Theta}_K dx &\leq (\nu \|\nabla \mathbf{e}_\mathbf{u}\|_{\mathbf{L}^2(K)} + \|\mathbf{e}_\mathbf{p}\|_{L^2(K)}) \|\nabla \boldsymbol{\Theta}_K\|_{\mathbf{L}^2(K)} \\ &\quad + \|\mathbf{g}\|_{\mathbf{L}^\infty(K)} \|\mathbf{e}_\mathbf{T}\|_{L^2(\mathbf{d}_z^\alpha, K)} \|\boldsymbol{\Theta}_K\|_{\mathbf{L}^2(\mathbf{d}_z^{-\alpha}, K)} \end{aligned}$$

$$\begin{aligned}
& + \left[\|u\|_{\mathbf{L}^3(K)} \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(K)} + \|\mathbf{e}_u\|_{\mathbf{L}^2(K)} \|\nabla u\|_{\mathbf{L}^3(K)} \right. \\
& \left. + \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(K)} \left(\|\mathbf{e}_u\|_{\mathbf{L}^3(K)} + \frac{1}{2} \|u_h\|_{\mathbf{L}^3(K)} \right) \right] \|\Theta_K\|_{\mathbf{L}^6(K)}.
\end{aligned}$$

We thus invoke $\|\Theta_K\|_{\mathbf{L}^6(K)} \lesssim h_K^{-1} \|\Theta_K\|_{\mathbf{L}^2(K)}$ and similar arguments to the ones used to treat the two-dimensional case to conclude that (5.38) holds also for $d = 3$.

Step 2. Let $K \in \mathcal{T}$ and $S \in \mathcal{S}_K$. We bound $h_K \|\mathcal{J}_S\|_{\mathbf{L}^2(S)}^2$. Define $\Lambda_S := \Psi_S \mathcal{J}_S$, where $\mathcal{J}_S$ and Ψ_S are as in (5.18) and (5.32), respectively. Basic properties of Ψ_S yield

$$\|\mathcal{J}_S\|_{\mathbf{L}^2(S)}^2 = \int_S |\mathcal{J}_S|^2 \, ds \lesssim \int_S |\mathcal{J}_S|^2 \Psi_S \, ds = \int_S \mathcal{J}_S \cdot \Lambda_S \, ds. \quad (5.39)$$

Notice that $\text{supp } \Lambda_S \subset K_* \cup K'_* \subset \mathcal{N}_S$. Setting $\mathbf{v} = \Lambda_S$ in (5.33) yields

$$\begin{aligned}
\int_S \mathcal{J}_S \cdot \Lambda_S \, ds &= \sum_{K' \in \mathcal{N}_S} \int_{K'} (\nu \nabla \mathbf{e}_u : \nabla \Lambda_S - \mathbf{e}_p \text{div } \Lambda_S + (\mathbf{u} \cdot \nabla) \mathbf{e}_u \cdot \Lambda_S \\
&\quad + (\mathbf{e}_u \cdot \nabla) u_h \cdot \Lambda_S + \frac{1}{2} \text{div } \mathbf{e}_u u_h \cdot \Lambda_S - \mathbf{e}_T \mathbf{g} \cdot \Lambda_S - \mathcal{R}_{K'} \cdot \Lambda_S) \, dx.
\end{aligned}$$

On the other hand, by shape regularity, we have that

$$\begin{aligned}
\|\Lambda_S\|_{\mathbf{L}^2(K')} &\approx |K'|^{\frac{1}{2}} |S|^{-\frac{1}{2}} \|\Lambda_S\|_{\mathbf{L}^2(S)} \approx h_{K'}^{\frac{1}{2}} \|\Lambda_S\|_{\mathbf{L}^2(S)} \approx h_{K'}^{\frac{1}{2}} \|\mathcal{J}_S\|_{\mathbf{L}^2(S)}, \\
\|\nabla \Lambda_S\|_{\mathbf{L}^2(K')} &\lesssim h_{K'}^{-1} \|\Lambda_S\|_{\mathbf{L}^2(K')} \approx h_{K'}^{-\frac{1}{2}} \|\mathcal{J}_S\|_{\mathbf{L}^2(S)}.
\end{aligned}$$

In view of (5.39), the estimates for Λ_S previously stated, and the arguments that yield estimate (5.38) for $\|\mathcal{R}_K\|_{\mathbf{L}^2(K')}$, we are capable of obtaining that

$$h_K \|\mathcal{J}_S\|_{\mathbf{L}^2(S)}^2 \lesssim \sum_{K' \in \mathcal{N}_S} (\|\mathbf{e}_T\|_{L^2(d_z^\alpha, K')}^2 + \nu^2 \|\mathbf{e}_u\|_{\mathbf{H}^1(K')}^2 + \|\mathbf{e}_p\|_{L^2(K')}^2).$$

Step 3. We now bound the residual term associated with the incompressibility constraint. Since $\text{div } \mathbf{u} = 0$, for any $K \in \mathcal{T}$, we have

$$\|\text{div } u_h\|_{L^2(K)}^2 = \|\text{div } \mathbf{e}_u\|_{L^2(K)}^2.$$

Finally, combining all the previous results (5.35) follows. $\square$

Theorem 5.4. (Efficiency) *Let $(\mathbf{T}, \mathbf{u}, p) \in H_0^1(d_z^\alpha, \Omega) \times \mathbf{H}_0^1(\Omega) \times L_0^2(\Omega)$ be a solution to (2.5) and $(\mathbf{T}_h, u_h, p_h) \in W_h \times \mathbf{X}_h \times M_h$ be its finite element approximation obtained as solution to (4.4). Let $\alpha \in (d-2, d)$. In the framework of Theorem 5.1, we have the following efficiency estimate:*

$$\begin{aligned}
\mathfrak{E}_K^2 &\lesssim \sum_{K' \in \mathcal{N}_S} ((\nu^2 + \kappa^2) \|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(d_z^\alpha, K')}^2 + \nu^2 h_K^{-2} \|\mathbf{e}_T\|_{L^2(d_z^\alpha, K')}^2 \\
&\quad + \nu^2 \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(K')}^2 + \nu^2 h_K^{-2} \|\mathbf{e}_u\|_{\mathbf{L}^2(K')}^2),
\end{aligned} \quad (5.40)$$

30 *A. Allendes, E. Otárola & A. J. Salgado*

where the hidden constant is independent of the continuous and discrete solutions, the size of the elements in the mesh $\mathcal{T}_h$, and $\#\mathcal{T}_h$.

Proof. We proceed in several steps.

Step 1. Let $K \in \mathcal{T}_h$. We bound $h_K^2 D_K^\alpha \|\mathfrak{R}_K\|_{L^2(K)}^2$ in (5.21). Define $\phi_K := \Psi_K^2 \mathfrak{R}_K$, where $\mathfrak{R}_K$ and Ψ_K are as in (5.20) and (5.30), respectively. Invoke (5.31) to arrive at

$$\|\mathfrak{R}_K\|_{L^2(K)}^2 \lesssim \int_K \mathfrak{R}_K \phi_K \, dx. \quad (5.41)$$

Set $r = \phi_K \in H_0^1(d_z^{-\alpha}, \Omega)$ as a test function in (5.34). Utilize that $R_K = \text{supp } \phi_K \subset K^* \subset K$ and the property $\phi_K(z) = 0$ to obtain

$$\begin{aligned} \int_K \mathfrak{R}_K \phi_K \, dx &= \int_K \kappa \nabla \mathbf{e}_T \cdot (2\Psi_K \nabla \Psi_K \mathfrak{R}_K + \Psi_K^2 \nabla \mathfrak{R}_K) \, dx \\ &\quad - \int_K \mathbf{T}(\Psi_K \mathbf{e}_u) \cdot (2\nabla \Psi_K \mathfrak{R}_K + \Psi_K \nabla \mathfrak{R}_K) \, dx \\ &\quad - \int_K (\Psi_K \mathbf{e}_T) \mathbf{u}_h \cdot (2\nabla \Psi_K \mathfrak{R}_K + \Psi_K \nabla \mathfrak{R}_K) \, dx =: \text{I} + \text{II} + \text{III}. \end{aligned}$$

We first observe that Proposition 8 in Ref. 4 yields

$$\|2\nabla \Psi_K \mathfrak{R}_K + \Psi_K \nabla \mathfrak{R}_K\|_{L^2(d_z^{-\alpha}, K)} \lesssim h_K^{-1} D_K^{-\alpha/2} \|\mathfrak{R}_K\|_{L^2(K)}. \quad (5.42)$$

We now proceed differently according to the spatial dimension. If $d = 2$, we utilize (2.4) and (5.42) and estimate II as follows:

$$\begin{aligned} \text{II} &\lesssim \|\mathbf{T}\|_{L^4(d_z^\alpha, K)} \|\Psi_K \mathbf{e}_u\|_{L^4(K)} h_K^{-1} D_K^{-\alpha/2} \|\mathfrak{R}_K\|_{L^2(K)} \\ &\lesssim \|\mathbf{T}\|_{L^4(d_z^\alpha, \Omega)} h_K^{1/2} \|\nabla(\Psi_K \mathbf{e}_u)\|_{L^2(K)} h_K^{-1} D_K^{-\alpha/2} \|\mathfrak{R}_K\|_{L^2(K)} \\ &\lesssim \nu h_K^{1/2} (h_K^{-1} \|\mathbf{e}_u\|_{L^2(K)} + \|\nabla \mathbf{e}_u\|_{L^2(K)}) h_K^{-1} D_K^{-\alpha/2} \|\mathfrak{R}_K\|_{L^2(K)}, \end{aligned} \quad (5.43)$$

where we have also used Lemma II.3.2 in Ref. 24 and the smallness assumption (5.13). We now control the term III. To accomplish this task, we utilize (2.4) and (5.42) to obtain

$$\begin{aligned} \text{III} &\lesssim \|\Psi_K \mathbf{e}_T\|_{L^4(d_z^\alpha, K)} \|\mathbf{u}_h\|_{L^4(K)} h_K^{-1} D_K^{-\alpha/2} \|\mathfrak{R}_K\|_{L^2(K)} \\ &\lesssim \|\nabla(\Psi_K \mathbf{e}_T)\|_{L^2(d_z^\alpha, K)} \|\mathbf{u}_h\|_{L^4(\Omega)} h_K^{-1} D_K^{-\alpha/2} \|\mathfrak{R}_K\|_{L^2(K)} \\ &\lesssim \nu (h_K^{-1} \|\mathbf{e}_T\|_{L^2(d_z^\alpha, K)} + \|\nabla \mathbf{e}_T\|_{L^2(d_z^\alpha, K)}) h_K^{-1} D_K^{-\alpha/2} \|\mathfrak{R}_K\|_{L^2(K)}. \end{aligned} \quad (5.44)$$

Here, we have also used $\|\Psi_K \mathbf{e}_T\|_{L^4(d_z^\alpha, K)} \lesssim \|\nabla(\Psi_K \mathbf{e}_T)\|_{L^2(d_z^\alpha, K)}$, which follows from Theorem 1.2 in Ref. 20. A collection of the estimates obtained for II and III

yields

$$\begin{aligned} \int_K \Re_K \phi_K \, dx &\leq [\kappa \|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, K)} + \nu h_K^{1/2} (h_K^{-1} \|\mathbf{e}_u\|_{\mathbf{L}^2(K)} + \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(K)}) \\ &\quad + \nu (h_K^{-1} \|\mathbf{e}_T\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, K)} + \|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, K)})] h_K^{-1} D_K^{-\alpha/2} \|\Re_K\|_{L^2(K)}. \end{aligned}$$

Consequently,

$$\begin{aligned} h_K^2 D_K^\alpha \|\Re_K\|_{\mathbf{L}^2(K)}^2 &\lesssim (\kappa^2 + \nu^2) \|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, K)}^2 + \nu^2 h_K \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(K)}^2 \\ &\quad \times \nu^2 h_K^{-2} \|\mathbf{e}_T\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, K)}^2 + \nu^2 h_K^{-1} \|\mathbf{e}_u\|_{\mathbf{L}^2(K)}^2. \end{aligned} \quad (5.45)$$

Similar arguments yield the following bound for the three-dimensional case:

$$\begin{aligned} \text{II} &\lesssim \|\mathbf{T}\|_{\mathbf{L}^3(\mathbf{d}_z^\alpha, K)} \|\Psi_K \mathbf{e}_u\|_{\mathbf{L}^6(K)} h_K^{-1} D_K^{-\alpha/2} \|\Re_K\|_{L^2(K)} \\ &\lesssim \nu (h_K^{-1} \|\mathbf{e}_u\|_{\mathbf{L}^2(K)} + \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(K)}) h_K^{-1} D_K^{-\alpha/2} \|\Re_K\|_{L^2(K)}, \end{aligned} \quad (5.46)$$

upon utilizing estimate II.3.7 in Ref. 24, and

$$\begin{aligned} \text{III} &\lesssim \|\Psi_K \mathbf{e}_T\|_{L^3(\mathbf{d}_z^\alpha, K)} \|\mathbf{u}_h\|_{\mathbf{L}^6(K)} h_K^{-1} D_K^{-\alpha/2} \|\Re_K\|_{L^2(K)} \\ &\lesssim \nu h_K^{1/6} (h_K^{-1} \|\mathbf{e}_T\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, K)} + \|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, K)}) h_K^{-1} D_K^{-\alpha/2} \|\Re_K\|_{L^2(K)}. \end{aligned} \quad (5.47)$$

Here, we have used the estimate $\|\Psi_K \mathbf{e}_T\|_{L^3(\mathbf{d}_z^\alpha, K)} \lesssim h_K^{1/6} \|\nabla(\Psi_K \mathbf{e}_T)\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, K)}$, which follows from Theorem 1.2 in Ref. 20. Consequently,

$$\begin{aligned} h_K^2 D_K^\alpha \|\Re_K\|_{\mathbf{L}^2(K)}^2 &\lesssim (\kappa^2 + \nu^2 h_K^{1/3}) \|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, K)}^2 + \nu^2 \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(K)}^2 \\ &\quad \times \nu^2 h_K^{-5/3} \|\mathbf{e}_T\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, K)}^2 + \nu^2 h_K^{-2} \|\mathbf{e}_u\|_{\mathbf{L}^2(K)}^2. \end{aligned} \quad (5.48)$$

Step 2. Let $K \in \mathcal{T}_h$ and $S \in \mathcal{S}_K$. The bound of $h_K D_K^\alpha \|\mathfrak{J}_S\|_{L^2(S)}^2$ follows similar arguments as the ones developed in Step 2 upon utilizing (5.45) and (5.48). In fact, we have, in two and three dimensions,

$$\begin{aligned} h_K D_K^\alpha \|\mathfrak{J}_S\|_{L^2(S)}^2 &\lesssim \sum_{K' \in \mathcal{N}_S} ((\nu^2 + \kappa^2) \|\nabla \mathbf{e}_T\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, K')}^2 + \nu^2 h_K^{-2} \|\mathbf{e}_T\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, K')}^2 \\ &\quad + \nu^2 \|\nabla \mathbf{e}_u\|_{\mathbf{L}^2(K')}^2 + \nu^2 h_K^{-2} \|\mathbf{e}_u\|_{\mathbf{L}^2(K')}^2). \end{aligned}$$

Step 3. We now control the term $|\mathfrak{h}| h_K^{\alpha+2-d} \#(\{z\} \cap K)$ in (5.21). Let $K \in \mathcal{T}_h$, and notice first that, if $T \cap \{z\} = \emptyset$, then estimate (5.40), follows from the estimates derived in the previous steps. If, on the other hand, $K \cap \{z\} = \{z\}$, then we must obtain a bound for the term $|\mathfrak{h}| h_K^{\alpha+2-d}$. To do so we follow the arguments developed in the proof of Theorem 5.3 in Ref. 1 that yield the existence of a smooth η such that

$$\eta(z) = 1, \quad \|\eta\|_{L^\infty(\Omega)} = 1, \quad \|\nabla \eta\|_{L^\infty(\Omega)} = h_K^{-1}, \quad \Omega_\eta := \text{supp}(\eta) \subset \mathcal{S}_K,$$

32 *A. Allendes, E. Otárola & A. J. Salgado*

where $\mathcal{S}_K$ is defined in (5.17). With this function at hand, define $r_\eta := \mathfrak{h}\eta^2 \in W_0^{1,\infty}(\Omega) \subset H_0^1(\mathbf{d}_z^{-\alpha}, \Omega)$ and notice that

$$\begin{aligned} |\mathfrak{h}|^2 &= \langle \mathfrak{h}\delta_z, r_\eta \rangle = \int_{\Omega} (\kappa \nabla \mathbf{T} \cdot \nabla r_\eta - \mathbf{T} \mathbf{u} \cdot \nabla r_\eta) \, dx \\ &= \int_{\Omega} \nabla \varphi \cdot \nabla r_\eta \, dx + \int_{\Omega} (\kappa \nabla \mathbf{T}_h \cdot \nabla r_\eta - \mathbf{T}_h \mathbf{u}_h \cdot \nabla r_\eta) \, dx, \end{aligned}$$

where we have used Eq. (5.5). We thus apply similar arguments to the ones used in the previous steps, integration by parts, and basic estimates to arrive, in two dimensions, at

$$\begin{aligned} |\mathfrak{h}|^2 &\lesssim \kappa \|\nabla \mathbf{e}_\mathbf{T}\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, \mathcal{S}_K)} \|\nabla r_\eta\|_{\mathbf{L}^2(\mathbf{d}_z^{-\alpha}, \mathcal{S}_K)} \\ &\quad + \nu h_K^{1/2} (h_K^{-1} \|\mathbf{e}_\mathbf{u}\|_{\mathbf{L}^2(\mathcal{S}_K)} + \|\nabla \mathbf{e}_\mathbf{u}\|_{\mathbf{L}^2(\mathcal{S}_K)}) \|\mathfrak{h} \nabla \eta\|_{\mathbf{L}^2(\mathbf{d}_z^{-\alpha}, \mathcal{S}_K)} \\ &\quad + \nu (h_K^{-1} \|\mathbf{e}_\mathbf{T}\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, \mathcal{S}_K)} + \|\nabla \mathbf{e}_\mathbf{T}\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, \mathcal{S}_K)}) \|\mathfrak{h} \nabla \eta\|_{\mathbf{L}^2(\mathbf{d}_z^{-\alpha}, \mathcal{S}_K)} \\ &\quad + \sum_{K' \in \mathcal{T}_h: K' \subset \mathcal{S}_K} \left(\|\mathfrak{R}_K\|_{\mathbf{L}^2(K')} \|r_\eta\|_{L^2(K')} \right. \\ &\quad \left. + \sum_{S \in \mathcal{S}_{K'}: S \not\subset \partial \mathcal{S}_K} \|\mathfrak{J}_S\|_{L^2(S)} \|r_\eta\|_{L^2(S)} \right), \end{aligned}$$

where we have also used the smallness assumption (5.13). Using the shape regularity of the mesh, in conjunction with the fact that, since $z \in K$, $h_K \approx D_K$, the bounds

$$\|\nabla \eta\|_{\mathbf{L}^2(\mathbf{d}_z^{-\alpha}, \mathcal{S}_K)} \lesssim h_K^{\frac{d-2}{2} - \frac{\alpha}{2}}, \quad \|\eta\|_{L^2(\mathcal{S}_K)} \lesssim h_K^{\frac{d}{2}}, \quad \|\eta\|_{L^2(S)} \lesssim h_K^{\frac{d-1}{2}},$$

allow us to conclude that

$$\begin{aligned} |\mathfrak{h}| &\lesssim h_K^{\frac{d-2}{2} - \frac{\alpha}{2}} ((\kappa + \nu) \|\nabla \mathbf{e}_\mathbf{T}\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, \mathcal{S}_K)} + \nu h_K^{1/2} \|\nabla \mathbf{e}_\mathbf{u}\|_{\mathbf{L}^2(\mathcal{S}_K)} \\ &\quad \times \nu h_K^{-1/2} \|\mathbf{e}_\mathbf{u}\|_{\mathbf{L}^2(\mathcal{S}_K)} + \nu h_K^{-1} \|\mathbf{e}_\mathbf{T}\|_{\mathbf{L}^2(\mathbf{d}_z^\alpha, \mathcal{S}_K)}) \\ &\quad + \sum_{K' \in \mathcal{T}_h: K' \subset \mathcal{S}_K} h_{K'}^{\frac{d-2}{2} - \frac{\alpha}{2}} \left(h_{K'} D_{K'}^{\frac{\alpha}{2}} \|\mathfrak{R}_K\|_{\mathbf{L}^2(K')} \right. \\ &\quad \left. + \sum_{S \in \mathcal{S}_{K'}: S \not\subset \partial \mathcal{S}_K} h_{K'}^{\frac{1}{2}} D_{K'}^{\frac{\alpha}{2}} \|\mathfrak{J}_S\|_{L^2(S)} \right). \end{aligned}$$

The estimate in three dimensions being similar.

Finally, combining all the previous results (5.35) follows. $\square$

6. Numerical Experiments

In this section, we conduct a series of numerical examples that illustrate the performance of the *a posteriori* error estimator we have devised and analyzed in Sec. 5. The examples have been carried out with the help of a code that we implemented using C++. All matrices have been assembled exactly and global linear systems were solved using the multifrontal massively parallel sparse direct solver (MUMPS).^{5,6} The *element* and *interelement* residuals are computed with the help of quadrature formulas which are exact for all the expressions involved. To visualize finite element approximations we have used the open-source application ParaView.^{2,8}

For a given partition $\mathcal{T}_h$, we solve (4.4) with the discrete spaces $\mathbf{X}_h$, M_h , and W_h given by (5.1), (5.2), and the space of continuous piecewise polynomial functions of degree two, respectively. To be precise, to adaptively solve the nonlinear system (4.4) we proceed as in Algorithm 1. We comment that, in Algorithm 2, for an initial partition $\mathcal{T}_0$, the initial guesses $\mathbf{T}_h^0 \in W_h$ and $(\mathbf{u}_h^0, \mathbf{p}_h^0) \in \mathbf{X}_h \times W_h$ are obtained as the respective solutions to the following problems:

$$\int_{\Omega} \kappa \nabla \mathbf{T}_h^0 \cdot \nabla r_h \, dx = \langle \mathcal{H}, r_h \rangle \quad \forall r_h \in W_h$$

and

$$\int_{\Omega} \nu \nabla \mathbf{u}_h^0 : \nabla \mathbf{v}_h - \int_{\Omega} \mathbf{p}_h^0 \operatorname{div} \mathbf{v}_h \, dx = \int_{\Omega} \mathbf{T}_h^0 \mathbf{g} \cdot \mathbf{v}_h \, dx, \quad \int_{\Omega} q_h \operatorname{div} \mathbf{u}_h^0 \, dx = 0,$$

for all $\mathbf{v}_h \in \mathbf{X}_h$ and $q_h \in M_h$, respectively. Once the discrete solution is obtained, we compute, for all $K \in \mathcal{T}_h$, the local *a posteriori* error indicators $\mathcal{E}_K$, defined in Sec. 5.3.2, to derive the adaptive mesh refinement procedure described in Algorithm 1. A sequence of adaptively refined meshes is thus generated from an initial mesh.

Finally, we denote the total number of degrees of freedom by $\text{Ndof} = \dim(W_h) + \dim(\mathbf{X}_h) + \dim(M_h)$.

Algorithm 1: Adaptive Algorithm.

Input: Initial mesh $\mathcal{T}_0$, interior point $z \in \Omega$, and parameters ν , κ , $\mathbf{h}$, and $\alpha \in (0, 2)$;

1: Solve the discrete problem (4.4) by using **Algorithm 2**;

2: For each $K \in \mathcal{T}_h$ compute the local error indicators $\mathcal{E}_K$ defined in Sec. 5.3.2;

3: Mark an element $K \in \mathcal{T}_h$ for refinement if

$$\mathcal{E}_K > \frac{1}{2} \max_{K' \in \mathcal{T}_h} \mathcal{E}_{K'};$$

4: From step **3**, construct a new mesh, using a longest edge bisection algorithm.³¹ Set $i \leftarrow i + 1$, and go to step **1**.

Algorithm 2: Fixed-Point Algorithm.

Input: Initial guess $(\mathbf{T}_h^0, \mathbf{u}_h^0, \mathbf{p}_h^0) \in W_h \times \mathbf{X}_h \times M_h$ and $\text{tol} = 10^{-8}$.

1: For $i \geq 0$, find $(\mathbf{u}_h^{i+1}, p_h^{i+1}) \in \mathbf{X}_h \times M_h$ such that

$$\begin{aligned} \int_{\Omega} (\nu \nabla \mathbf{u}_h^{i+1} : \nabla \mathbf{v}_h + (\mathbf{u}_h^i \cdot \nabla) \mathbf{u}_h^{i+1} \cdot \mathbf{v}_h + \frac{1}{2} \operatorname{div} \mathbf{u}_h^i \mathbf{u}_h^{i+1} \cdot \mathbf{v}_h) dx \\ - \int_{\Omega} p_h^{i+1} \operatorname{div} \mathbf{v}_h dx = \int_{\Omega} \mathbf{T}_h^i \mathbf{g} \cdot \mathbf{v}_h dx, \quad \forall \mathbf{v}_h \in \mathbf{X}_h, \\ \int_{\Omega} \operatorname{div} \mathbf{u}_h^{i+1} q_h dx, \quad \forall q_h \in M_h. \end{aligned}$$

Then, $\mathbf{T}_h^{i+1} \in W_h$ is found as the solution of

$$\int_{\Omega} (\kappa \nabla \mathbf{T}_h^{i+1} \cdot \nabla r_h - \mathbf{T}_h^{i+1} \mathbf{u}_h^{i+1} \cdot \nabla r_h) dx = \langle \mathbf{h} \delta_z, r_h \rangle \quad \forall r_h \in W_h.$$

2: If $\|(\mathbf{T}_h^{i+1}, \mathbf{u}_h^{i+1}, p_h^{i+1}) - (\mathbf{T}_h^i, \mathbf{u}_h^i, p_h^i)\|_2 > \text{tol}$, set $i \leftarrow i + 1$, and go to step **1**.

Otherwise, **return** $(\mathbf{T}_h, \mathbf{u}_h, p_h) = (\mathbf{T}_h^{i+1}, \mathbf{u}_h^{i+1}, p_h^{i+1})$.

We now explore the performance of the devised *a posteriori* error estimator in two problems with homogeneous Dirichlet boundary conditions on convex and non-convex domains. In all the numerical experiments, we have considered $\nu = \kappa = 1$, $\mathbf{g} = [1, 0]^\top$, $\mathbf{h} = 1$, $z = (0.5, 0.5)$, and different values for the exponent of the Muckenhoupt weight: $\alpha \in \{0.1, 0.5, 1.0, 1.5, 1.9\}$. We let

- (i) $\Omega = (0, 1)^2$, for example 1, and
- (ii) $\Omega = (-1, 1)^2 \setminus [0, 1) \times [-1, 0)$, for example 2.

In Figs. 1 and 3 we present, within the setting of examples 1 and 2, respectively, experimental rates of convergence for the error estimator $\mathcal{E}_h$. We also present the initial meshes used in the adaptive algorithm. We observe that optimal experimental rates of convergence are attained for all the values of the parameter α that we have considered. We also observe that a better value of the estimator, at a fixed mesh, can be obtained for values of α closer to two. We notice that, when α is small, after a certain number of adaptive iterations, there are elements K around z such that $|K| \approx 10^{-16}$. This makes impossible more computations within the adaptive procedure.

In Figs. 2 and 4 we present, for examples 1 and 2, respectively, a series of meshes obtained after 30 adaptive iterations. We observe that most of the refinement is concentrated around the singular source point. For the case of example 2, and after 30 adaptive refinements, the adaptive loop also concentrates the refinement around the reentrant corner when $\alpha \in [1, 2)$.

Finally, in Fig. 5 we present, for the setting of example 2 with $\alpha = 1.5$, $|\mathbf{u}_h|$, its associated streamlines, the pressure p_h , and the temperature $\mathbf{T}_h$ over a mesh




Fig. 1. Example 1: Experimental rates of convergence for the error estimator $\mathcal{E}_h$ considering $\alpha \in \{0.1, 0.5, 1.0, 1.5, 1.9\}$ (left) and the initial mesh used in the adaptive algorithm (right).




Fig. 2. Example 1: Meshes obtained after 30 iterations of our adaptive loop for (a) $\alpha = 0.1$ (232 elements and 121 vertices); (b) $\alpha = 0.5$ (232 elements and 121 vertices); (c) $\alpha = 1.0$ (392 elements and 209 vertices); (d) $\alpha = 1.5$ (592 elements and 309 vertices); and (e) $\alpha = 1.9$ (1056 elements and 553 vertices).




Fig. 3. Example 2: Experimental rates of convergence for the error estimator $\mathcal{E}_h$ considering $\alpha \in \{0.1, 0.5, 1.0, 1.5, 1.9\}$ (left) and the initial mesh used in the adaptive algorithm, which contains 96 elements and 65 vertices (right).




Fig. 4. Example 2: Meshes obtained after 30 iterations of our adaptive loop for (a) $\alpha = 0.1$ (328 elements and 181 vertices); (b) $\alpha = 0.5$ (328 elements and 181 vertices); (c) $\alpha = 1.0$ (539 elements and 291 vertices); (d) $\alpha = 1.5$ (847 elements and 450 vertices); and (e) $\alpha = 1.9$ (1380 elements and 728 vertices).




Fig. 5. Example 2: Finite element approximations of $|u_h|$, combined with its streamlines (left), pressure p_h (center), and temperature T_h (right) over a mesh containing 16105 elements and 8178 vertices obtained after 65 adaptive refinements ($\alpha = 1.5$).

containing 16105 elements and 8178 vertices; the latter being obtained after 65 iterations of our adaptive loop.

Acknowledgments

AA has been partially supported by CONICYT through FONDECYT project 1210729. EO has been partially supported by CONICYT through FONDECYT project 11180193. AJS has been partially supported by NSF grant DMS-1720213.

References

1. J. P. Agnelli, E. M. Garau and P. Morin, *A posteriori* error estimates for elliptic problems with Dirac measure terms in weighted spaces, *ESAIM Math. Model. Numer. Anal.* **48** (2014) 1557–1581.
2. J. P. Ahrens, B. Geveci and C. C. W. Law, ParaView: An end-user tool for large-data visualization, in *Visualization Handbook* (Elsevier, 2005).
3. A. Allendes, C. Naranjo and E. Otárola, Stabilized finite element approximations for a generalized Boussinesq problem: *A posteriori* error analysis, *Comput. Methods Appl. Mech. Eng.* **361** (2020) 112703.
4. A. Allendes, E. Otárola and A. J. Salgado, *A posteriori* error estimates for the Stokes problem with singular sources, *Comput. Methods Appl. Mech. Eng.* **345** (2019) 1007–1032.
5. P. R. Amestoy, I. S. Duff, J.-Y. L'Excellent and J. Koster, A fully asynchronous multifrontal solver using distributed dynamic scheduling, *SIAM J. Matrix Anal. Appl.* **23** (2001) 15–41 (electronic).
6. P. R. Amestoy, A. Guermouche, J.-Y. L'Excellent and S. Pralet, Hybrid scheduling for the parallel solution of linear systems, *Parallel Comput.* **32** (2006) 136–156.
7. D. N. Arnold, F. Brezzi and M. Fortin, A stable finite element for the Stokes equations, *Calcolo* **21** (1984) 337–344 (1985).
8. U. Ayachit, *The ParaView Guide: A Parallel Visualization Application* (Kitware, Inc., 2015).
9. R. G. Bartle and J. T. Joichi, The preservation of convergence of measurable functions under composition, *Proc. Amer. Math. Soc.* **12** (1961) 122–126.

38 *A. Allendes, E. Otárola & A. J. Salgado*

10. C. Bernardi, B. Métivet and B. Pernaud-Thomas, Couplage des équations de Navier–Stokes et de la chaleur: le modèle et son approximation par éléments finis, *RAIRO Modél. Math. Anal. Numér.* **29** (1995) 871–921.
11. M. Bogovskiĭ, Solution of the first boundary value problem for an equation of continuity of an incompressible medium, *Dokl. Akad. Nauk SSSR* **248** (1979) 1037–1040.
12. M. Bogovskiĭ, Solutions of some problems of vector analysis, associated with the operators div and grad, in *Theory of Cubature Formulas and the Application of Functional Analysis to Problems of Mathematical Physics*, Trudy Sem. S. L. Soboleva, No. 1, Vol. 1980 (Akad. Nauk SSSR Sibirsk. Otdel., Inst. Mat., Novosibirsk, 1980), pp. 5–40, 149.
13. S. C. Brenner and L. R. Scott, *The Mathematical Theory of Finite Element Methods*, 3rd edn., Texts in Applied Mathematics, Vol. 15 (Springer, 2008).
14. P. G. Ciarlet, *The Finite Element Method for Elliptic Problems* (SIAM, 2002).
15. E. Colmenares, G. N. Gatica and R. Oyarzúa, Analysis of an augmented mixed-primal formulation for the stationary Boussinesq problem, *Numer. Methods Partial Differential Equations* **32** (2016) 445–478.
16. K. Deimling, *Nonlinear Functional Analysis* (Springer-Verlag, 1985).
17. I. Drelichman, R. G. Durán and I. Ojea, A weighted setting for the numerical approximation of the Poisson problem with singular sources, *SIAM J. Numer. Anal.* **58** (2020) 590–606.
18. J. Duoandikoetxea, *Fourier Analysis*, Graduate Studies in Mathematics, Vol. 29 (Amer. Math. Soc., 2001), translated and revised from the 1995 Spanish original by David Cruz-Urbe.
19. A. Ern and J.-L. Guermond, *Theory and Practice of Finite Elements*, *Applied Mathematical Sciences*, Vol. 159 (Springer-Verlag, 2004).
20. E. B. Fabes, C. E. Kenig and R. P. Serapioni, The local regularity of solutions of degenerate elliptic equations, *Comm. Partial Differential Equations* **7** (1982) 77–116.
21. M. Farhloul, S. Nicaise and L. Paquet, A mixed formulation of Boussinesq equations: Analysis of nonsingular solutions, *Math. Comput.* **69** (2000) 965–986.
22. R. Farwig and H. Sohr, Weighted L^q -theory for the Stokes resolvent in exterior domains, *J. Math. Soc. Japan* **49** (1997) 251–288.
23. X. Feng and S. Wise, Analysis of a Darcy–Cahn–Hilliard diffuse interface model for the Hele–Shaw flow and its fully discrete finite element approximation, *SIAM J. Numer. Anal.* **50** (2012) 1320–1343.
24. G. P. Galdi, *An Introduction to the Mathematical Theory of the Navier–Stokes Equations: Steady-State Problems*, 2nd edn., Springer Monographs in Mathematics (Springer, 2011).
25. V. Girault and P.-A. Raviart, *Finite Element Approximation of the Navier–Stokes equations*, Lecture Notes in Mathematics, Vol. 749 (Springer-Verlag, 1979).
26. V. Girault and P.-A. Raviart, *Finite Element Methods for Navier–Stokes Equations: Theory and Algorithms*, Springer Series in Computational Mathematics, Vol. 5 (Springer-Verlag, 1986).
27. G. Grün, On convergent schemes for diffuse interface models for two-phase flow of incompressible fluids with general mass densities, *SIAM J. Numer. Anal.* **51** (2013) 3036–3061.
28. D. D. Haroske and L. Skrzypczak, Entropy and approximation numbers of embeddings of function spaces with Muckenhoupt weights, II. General weights, *Ann. Acad. Sci. Fenn. Math.* **36** (2011) 111–138.

29. J. Heinonen, T. Kilpeläinen and O. Martio, *Nonlinear Potential Theory of Degenerate Elliptic Equations* (Dover Publications, Inc., 2006), unabridged republication of the 1993 original.
30. P. Hood and C. Taylor, Navier–Stokes equations using mixed interpolation, *Finite Element Methods in Flow Problems* (1974) 121–132.
31. I. Kossaczky, A recursive approach to local mesh refinement in two and three dimensions, *J. Comput. Appl. Math.* **55** (1994) 275–288.
32. V. Kozlov, V. Maz’ya and J. Rossmann, *Elliptic Boundary Value Problems in Domains with Point Singularities* (Amer. Math. Soc., 1997).
33. S. A. Lorca and J. L. Boldrini, Stationary solutions for generalized Boussinesq models, *J. Differential Equations* **124** (1996) 389–406.
34. M. Mitrea and M. Wright, Boundary value problems for the Stokes system in arbitrary Lipschitz domains, *Astérisque* **344** (2012) viii+241.
35. B. Muckenhoupt, Weighted norm inequalities for the Hardy maximal function, *Trans. Amer. Math. Soc.* **165** (1972) 207–226.
36. R. H. Nochetto, E. Otárola and A. J. Salgado, Piecewise polynomial interpolation in Muckenhoupt weighted Sobolev spaces and applications, *Numer. Math.* **132** (2016) 85–130.
37. R. H. Nochetto, A. J. Salgado and I. Tomas, A diffuse interface model for two-phase ferrofluid flows, *Comput. Methods Appl. Mech. Eng.* **309** (2016) 497–531.
38. R. H. Nochetto, A. J. Salgado and I. Tomas, The equations of ferrohydrodynamics: Modeling and numerical methods, *Math. Models Methods Appl. Sci.* **26** (2016) 2393–2449.
39. R. H. Nochetto, A. J. Salgado and S. W. Walker, A diffuse interface model for electrowetting with moving contact lines, *Math. Models Methods Appl. Sci.* **24** (2014) 67–111.
40. E. Otárola and A. J. Salgado, The Poisson and Stokes problems on weighted spaces in Lipschitz domains and under singular forcing, *J. Math. Anal. Appl.* **471** (2019) 599–612.
41. R. Oyarzúa, T. Qin and D. Schötzau, An exactly divergence-free finite element method for a generalized Boussinesq problem, *IMA J. Numer. Anal.* **34** (2014) 1104–1135.
42. C. E. Pérez, J.-M. Thomas, S. Blancher and R. Creff, The steady Navier–Stokes/energy system with temperature-dependent viscosity. I. Analysis of the continuous problem, *Internat. J. Numer. Methods Fluids* **56** (2008) 63–89.
43. C. E. Pérez, J.-M. Thomas, S. Blancher and R. Creff, The steady Navier–Stokes/energy system with temperature-dependent viscosity. II. The discrete problem and numerical experiments, *Internat. J. Numer. Methods Fluids* **56** (2008) 91–114.
44. R. Temam, *Navier–Stokes Equations: Theory and Numerical Analysis* (AMS Chelsea Publishing, 2001), reprint of the 1984 edition.
45. B. O. Turesson, *Nonlinear Potential Theory and Weighted Sobolev Spaces, Lecture Notes in Mathematics*, Vol. 1736 (Springer-Verlag, 2000).
46. R. Verfürth, *A posteriori* error estimators for the Stokes equations, *Numer. Math.* **55** (1989) 309–325.
47. R. Verfürth, *A Posteriori Error Estimation Techniques for Finite Element Methods*, Numerical Mathematics and Scientific Computation (Oxford Univ. Press, 2013).