

Throwing a Sofa Through the Window

Dan Halperin 

Tel Aviv University, School of Computer Science, Israel

Micha Sharir 

Tel Aviv University, School of Computer Science, Israel

Itay Yehuda 

Tel Aviv University, School of Computer Science, Israel

Abstract

We study several variants of the problem of moving a convex polytope K , with n edges, in three dimensions through a flat rectangular (and sometimes more general) window. Specifically:

- (i) We study variants where the motion is restricted to translations only, discuss situations where such a motion can be reduced to sliding (translation in a fixed direction), and present efficient algorithms for those variants, which run in time close to $O(n^{8/3})$.
- (ii) We consider the case of a *gate* (an unbounded window with two parallel infinite edges), and show that K can pass through such a window, by any collision-free rigid motion, iff it can slide through it, an observation that leads to an efficient algorithm for this variant too.
- (iii) We consider arbitrary compact convex windows, and show that if K can pass through such a window W (by any motion) then K can slide through a slab of width equal to the diameter of W .
- (iv) We show that if a purely translational motion for K through a rectangular window W exists, then K can also slide through W keeping the same orientation as in the translational motion. For a given fixed orientation of K we can determine in linear time whether K can translate (and hence slide) through W keeping the given orientation, and if so plan the motion, also in linear time.
- (v) We give an example of a polytope that cannot pass through a certain window by translations only, but can do so when rotations are allowed.
- (vi) We study the case of a circular window W , and show that, for the regular tetrahedron K of edge length 1, there are two thresholds $1 > \delta_1 \approx 0.901388 > \delta_2 \approx 0.895611$, such that (a) K can slide through W if the diameter d of W is ≥ 1 , (b) K cannot slide through W but can pass through it by a purely translational motion when $\delta_1 \leq d < 1$, (c) K cannot pass through W by a purely translational motion but can do it when rotations are allowed when $\delta_2 \leq d < \delta_1$, and (d) K cannot pass through W at all when $d < \delta_2$.
- (vii) Finally, we explore the general setup, where we want to plan a general motion (with all six degrees of freedom) for K through a rectangular window W , and present an efficient algorithm for this problem, with running time close to $O(n^4)$.

2012 ACM Subject Classification Theory of computation $\rightarrow$ Computational geometry

Keywords and phrases Motion planning, Convex polytopes in 3D

Digital Object Identifier 10.4230/LIPIcs.SoCG.2021.41

Related Version *Full Version:* <https://arxiv.org/abs/2102.04262>

Funding Work on this paper by DH and IY has been supported in part by the Israel Science Foundation (grant no. 1736/19), by NSF/US-Israel-BSF (grant no. 2019754), by the Israel Ministry of Science and Technology (grant no. 103129), by the Blavatnik Computer Science Research Fund, and by a grant from Yandex. Work by MS and IY has been supported in part by Grant 260/18 from the Israel Science Foundation. Work by MS has also been supported by Grant G-1367-407.6/2016 from the German-Israeli Foundation for Scientific Research and Development, and by the Blavatnik Research Fund in Computer Science at Tel Aviv University.



© Dan Halperin, Micha Sharir, and Itay Yehuda;
licensed under Creative Commons License CC-BY 4.0


37th International Symposium on Computational Geometry (SoCG 2021).

Editors: Kevin Buchin and Éric Colin de Verdière; Article No. 41; pp. 41:1–41:16

Leibniz International Proceedings in Informatics




LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany



Acknowledgements We deeply thank Pankaj Agarwal and Boris Aronov for useful interactions concerning this work. In particular, Boris has suggested an alternative proof of the key topological property in the analysis in Section 4, and Pankaj has been instrumental in discussions concerning the range searching problem in Section 2. We are also grateful to Lior Hadassi for interaction involving the topological property presented in Section 4, and to Eytan Tirosh for help with the alternative proof of Lemma 1.

1 Introduction

Let K be a convex polytope (a “sofa”) in $\mathbb{R}^3$ with n edges, and let W be a rectangular window, placed in the xy -plane in the axis-parallel position $[0, a] \times [0, b]$, where a and b are the respective width and height of W . We assume that the complement of W in the xy -plane is a solid wall that K must avoid. The problem is to determine whether K can be moved, in a collision-free manner, from any position that is fully contained in the upper halfspace $z > 0$, through W , to any position that is fully contained in the lower halfspace $z < 0$, and, if so, to plan such a motion (see Figure 1).



■ **Figure 1** Moving a convex polytope K through a window W .

A continuous motion of a rigid body in three dimensions has six degrees of freedom, three of translation and three of rotation, and in the general form of the problem, studied in Section 8, we allow all six degrees. On the way, we will study simpler versions where only restricted types of motion are allowed, such as purely translational motion (that has only three degrees of freedom), a translational motion in a fixed direction, which we refer to as *sliding* (one degree of freedom), or a translational motion combined with rotations around the vertical axis only (four degrees of freedom), etc. Some of our main results show that, in certain favorable situations, the existence of a general collision-free motion of K through W implies the existence of a restricted motion of one of these types. This allows us to solve the problem in a significantly more efficient manner.

In terms of the *free configuration space* $\mathcal{F}$ of K , all the placements of K that are fully contained in the upper (resp., lower) halfspace are free, and form a connected subset $\mathcal{F}^+$ (resp., $\mathcal{F}^-$) of $\mathcal{F}$. Our problem, in general, is to determine whether both $\mathcal{F}^+$ and $\mathcal{F}^-$ are contained in the same connected component of $\mathcal{F}$. This interpretation applies to the general setup, with six degrees of freedom, as well as to any other subclass of motions, with fewer degrees of freedom.

Motion planning is an intensively studied problem in computational geometry and robotics. A systematic and general way to describe the free space $\mathcal{F}$ is by using *constraint surfaces*, namely surfaces describing all the configurations where some feature on the boundary of the moving object (K in our case) touches a feature on the boundary of the free workspace (W in our case); see, e.g., [12, 18, 19]. These surfaces partition the configuration space into cells, each of which is either fully contained in $\mathcal{F}$ or fully contained in the forbidden portion of the

configuration space. This representation is based on the arrangement $\mathcal{A}$ of the constraint surfaces, whose number in our case is $O(n)$, one surface for each feature (edge or vertex) of W and each feature (edge, vertex or face) of K . Each cell of $\mathcal{A}$ is fully contained either in $\mathcal{F}$ or in its complement. Hence the complexity of $\mathcal{F}$ is $O(n^d)$, where d is the number of degrees of freedom (namely, the dimension of the configuration space) [12]. To exploit this representation, we construct and transform it into a so-called discrete *connectivity graph*, which can be searched for the existence of a motion of the desired kind.

One common way of doing this is to further decompose the arrangement into subcells of constant complexity, using *vertical decomposition* [4]. Such constructions are easily implementable for motion planning with two degrees of freedom [8], but become significantly more involved for problems with three or more degrees of freedom. This has led to the development of alternative methods, such as sampling-based techniques (see [5, Chapter 7] and [10]), the best known of which are PRM [16] and RRT [17], which have dozens of variants. While extremely successful in solving practical problems, they trade-off the completeness of the arrangement approach with efficiency, and may fail when the setting contains tight passages [21, 22], a situation that can arise in the problems that we study in this paper.

Toussaint [25] studied movable separability of sets, where he collected a variety of tight-setting motion planning problems, similar in nature to the problems studied here. These problems are interesting theoretically (see, e.g., [24] for such a problem and its intriguing solution), but also from an applied perspective, since motion in tight settings often arises in manufacturing processes such as *assembly planning* [11] or *casting and molding* [3].

It was in Toussaint's review [25] that we encountered the problem of "throwing" a polytope through a window. Although Toussaint's paper was published 35 years ago, we are not aware of any previous progress on this specific problem. We remark that the word *sofa* in the title of the paper is borrowed from the classical two-dimensional *moving sofa* problem (see, e.g., [6, 9]), which is to find the shape of largest area that can be moved through a corner in an L-shaped corridor whose legs have width 1.

Our results. We first consider, in Section 2, sliding motions (translations in a fixed direction) of K . We characterize situations in which such a sliding motion exists, and present efficient algorithms, with runtime close to $O(n^{8/3})$, for finding such a motion when one exists.

We next consider, in Section 3, the case where W is an unbounded slab, enclosed between two parallel unbounded lines (we call it a *gate*). We show that if K can pass through such a gate W , by any collision-free rigid motion, it can also slide through W , making the general motion planning problem through a gate particularly easy to solve.

In Section 4, we consider arbitrary compact convex windows, and show that if K can move through such a window W , by an arbitrary collision-free motion, then K can slide through a gate of width equal to the diameter of W , and this holds in any sliding direction. This requires nontrivial topological arguments, presented in Section 4 and in the full version of the paper [14].

We then consider, in Section 5, purely translational motions of K through a rectangular window W , and prove that the existence of such a purely-translational collision-free motion implies the existence of a collision-free sliding motion keeping the same orientation as in the translational motion. For a given fixed orientation of K we can determine in linear time whether K can translate (and hence slide) through W keeping the given orientation, and if so plan the motion, also in linear time. We also give a near-linear time algorithm for planning the motion of K through an arbitrary (flat) polygonal window of fixed (constant) complexity.

In Section 6, we show that rotations are sometimes needed, by giving an example of a convex polytope K (actually a tetrahedron) that can move through a square window W by a collision-free motion that includes rotation (only around an axis orthogonal to W), but there is no purely translational motion of K through W .

In Section 7, we consider the problem of passing through a circular window W , and show that, for the regular tetrahedron K of edge length 1, there are two thresholds $1 > \delta_1 \approx 0.901388 > \delta_2 \approx 0.895611$, such that (i) K can slide through the window W if the diameter d of W is ≥ 1 , (ii) K cannot slide through W but can pass through it by a purely translational motion when $\delta_1 \leq d < 1$, (iii) K cannot pass through W by a purely translational motion but can do it with rotations when $\delta_2 \leq d < \delta_1$, and (iv) K cannot pass through W at all when $d < \delta_2$.

We finally consider, in Section 8, the general problem, with all six degrees of freedom. We present an efficient algorithm, which runs in time close to $O(n^4)$, for constructing the free configuration space, from which one can construct, within a comparable time bound, a valid motion through W if one exists.

2 Translation in a fixed direction

In this section we address the case in which the movement is purely translational in a single fixed direction. Such a motion, to which we refer as a *sliding motion*, has only one degree of freedom. In the most restricted version (which is very easy to solve), we are given a fixed orientation of K at a fixed initial placement, and also the direction of motion. In this section we study a more general setting, in which we seek values for these parameters – orientation, initial placement, and direction of motion, for which such a sliding motion of K through W is possible (or determine that no such motion is possible).

In Section 2.1 we observe that if a sliding motion for K exists, then K can also slide in a direction orthogonal to the plane of the window. Using this and other structural properties of the problem, we transform the problem at hand into a certain range searching problem. We present an efficient novel solution to the latter problem, which yields an algorithm for solving our original problem, whose running time is close to $O(n^{8/3})$.

2.1 The existence of an orthogonal sliding motion


For the most general version of the sliding motion, in which none of the parameters (orientation, initial placement, and direction of motion) is prespecified, we have:

► **Lemma 1.** *If K can slide through W from some starting placement in some direction, then K can slide through W , possibly from some other starting placement (and at another orientation), by translating it in the negative z -direction.*

Proof. Let K_0 be the starting placement of K and let $\vec{v}$ be the direction of motion through W , for which the resulting sliding motion is collision-free. Form the infinite prism $\Pi_0 := \bigcup_{\lambda \in \mathbb{R}} (K_0 + \lambda \vec{v})$ that K_0 spans in direction $\vec{v}$. The premise of the lemma implies that the intersection of Π_0 with the xy -plane is contained in W .

Let W_0 be the orthogonal projection of W onto some plane orthogonal to $\vec{v}$. Note that W_0 is a parallelogram, and that, by construction, K_0 can pass through W_0 when translated in direction $\vec{v}$. By an old result, reviewed and proved by Debrunner and Mani-Levitska [7], it follows that, when mapped rigidly into the xy -plane, W_0 (the “shadow” of W in direction $\vec{v}$) can be placed fully within W (see Figure 2).¹

¹ Curiously, as shown in [7], this property, of containing your shadows, may fail in higher dimensions.



■ **Figure 2** The projection of W (green) can be located in a congruent copy of W (blue).

Now rotate and translate $\mathbb{R}^3$ so that $\vec{v}$ becomes the (negative) z -direction, and the image of W_0 is fully contained in (the former, untransformed copy of) W . Then the image of K under this transformation can be moved vertically down through W , in a collision-free manner, as asserted. ◀

Debrunner and Mani-Levitska's proof is involved, and applies to an arbitrary planar convex shape (showing that it contains its projection in any direction). In the full version of the paper [14] we give a simple alternative proof for the case of a rectangle.

2.2 Finding a sliding motion

The following discussion is with respect to a fixed initial placement K_0 of K . For a given direction $\vec{v}$, the *projected silhouette* of K_0 in direction $\vec{v}$ is the boundary of the convex polygon obtained by the projection of K_0 in direction $\vec{v}$, within the image plane $h_{\vec{v}}$ (which is orthogonal to $\vec{v}$). The silhouette itself is the cyclic sequence of vertices and edges of K , whose projections form the projected silhouette.² The silhouette and its projection do not change combinatorially, that is, when represented as a cyclic sequence of vertices and edges of K (or of their projections), as long as $\vec{v}$ is not parallel to any face of K_0 . We thus form the set of the $O(n)$ great circles on $\mathbb{S}^2$ that are parallel to the faces of K_0 , and construct their arrangement $\mathcal{A}_0$ on $\mathbb{S}^2$, which is also known as the *aspect graph* of K_0 [20]. In each face φ of $\mathcal{A}_0$ the combinatorial structure of the silhouette is fixed, but the projected silhouette varies continuously as $\vec{v}$ moves in φ .

A *view* of K_0 is a pair $(\vec{v}, \theta)$, where $\vec{v}$ is a direction, and θ is the angle of rotation of the projected silhouette within $h_{\vec{v}}$ (translations within that plane are ignored). The space of views is thus three-dimensional. A fixed view $(\vec{v}, \theta)$ fixes the uppermost, leftmost, bottommost and rightmost vertices w_t, w_l, w_b and w_r of the projected silhouette. The view is valid if

$$x_{w_r} - x_{w_l} \leq a \quad \text{and} \quad y_{w_t} - y_{w_b} \leq b \quad (1)$$

(in the coordinate frame of $h_{\vec{v}}$ when rotated by θ).

When $\vec{v}$ is fixed and θ varies, we get $O(n)$ quadruples (w_t, w_l, w_b, w_r) of the projected silhouette. The view is valid if the inequalities in (1) (which depend on $(\vec{v}, \theta)$) have a solution for one such quadruple, which lies in the appropriate portion of the view space (in which w_t, w_l, w_b, w_r are indeed the four extreme vertices). The existence of a valid view is equivalent to the existence of a sliding motion of K_0 through W , after suitably shifting K_0 and rotating it around $\vec{v}$ by θ . We give more details in the full version [14], where we show that the total number of quadruples of vertices is $O(n^3)$, from which we obtain an algorithm for finding a valid view, with near-cubic running time.

² The silhouette is indeed such a cycle of vertices and edges of ∂K for generic directions $\vec{v}$. When $\vec{v}$ is parallel to a face f of K_0 , the entire f is part of the silhouette.

2.3 An improved algorithm

We next present an improved, albeit more involved algorithm that solves the problem of finding a sliding motion of K , if one exists, in time $O(n^{8/3}\text{polylog}(n))$. The problem of finding a direction $\vec{v}$ in which we can slide K through W is equivalent to the problem of finding a placement of W in some plane h , such that the projected silhouette of K on h is contained in W , which in turn is equivalent to verifying that all the projected vertices of K fit into that placement of W .

An equivalent way of checking for the latter characterization is to look for two unit vectors x and y in $\mathbb{R}^3$ (which will be the directions of the axes of W in the desired placement; note that h is spanned by x and y) that satisfy:

- (i) x and y are perpendicular to each other.
- (ii) For every segment e connecting two vertices of K we have $\langle x, e \rangle \leq a$.
- (iii) For every segment e connecting two vertices of K we have $\langle y, e \rangle \leq b$.

(Note that since we go over all ordered pairs of vertices of K in (ii), (iii), we actually require that $|\langle x, e \rangle| \leq a$ and $|\langle y, e \rangle| \leq b$ for each such segment e .) Every inequality in (ii) defines a halfspace that has to contain x . We intersect those $O(n^2)$ halfspaces, to obtain a convex polytope Q of complexity $O(n^2)$, and intersect Q with the unit sphere $\mathbb{S}^2$ to obtain the admissible region A of the vectors x that satisfy (ii), in $O(n^2 \log n)$ time. The region A is bounded by circular arcs (not necessarily arcs of great circles), which meet at the *vertices* of A . We apply the same procedure for y using the suitable collection of halfspaces in (iii), and obtain the admissible region B for the vectors y that satisfy (iii), also in $O(n^2 \log n)$ time.

To satisfy (i) too, we need to check whether there exist an orthogonal pair of vectors $x \in A$, $y \in B$. We use the following lemma, whose proof is given in the full version [14].


► **Lemma 2.** *Let S_A denote the set of all vertices of A , and let T_A denote the set of the points that are closest locally to the north pole of $\mathbb{S}^2$ along each circular arc of ∂A . (If the direction of the north pole of $\mathbb{S}^2$ is generic, T_A is finite and $|S_A \cup T_A| = O(n^2)$.) Define similarly the sets S_B, T_B . If there exist an orthogonal pair $(x, y) \in A \times B$ then there exist such an orthogonal pair so that either $x \in S_A \cup T_A$ or $y \in S_B \cup T_B$.*

We iterate over the points of $S_A \cup T_A$. For each such point v let C_v be the great circle orthogonal to v , and let $\mathcal{C}$ denote the collection of these $O(n^2)$ great circles. We face the problem of determining whether any great circle in $\mathcal{C}$ crosses B , which is essentially the problem of determining whether any great circle in $\mathcal{C}$ crosses an arc of ∂B . By centrally projecting the setup onto two parallel planes, the problem is further reduced to the problem where we have a set of $M = O(n^2)$ lines and a set of $N = O(n^2)$ arcs of conic sections in the plane, and the goal is to determine whether there is a line-arc intersection. The important role of Lemma 2 is that it enables us to reduce the quest for an orthogonal pair (x, y) into a finite number ($O(n^2)$) of range-searching queries, by asserting that if an orthogonal pair exists, then there exists an orthogonal pair where at least one of x, y is from a prescribed discrete set of points.

In the full version [14] we give full details of our solution. Here is a brief sketch. With no loss of generality, assume that each input arc is x -monotone and convex. Then a rightward-oriented line ℓ intersects an arc γ if and only if one of the following conditions holds:

- (i) The two endpoints of γ lie on different sides of ℓ . See Figure 3(i).
- (ii) The two endpoints of γ lie to the left of ℓ , γ has a tangent that is parallel to ℓ (that is, the slope a of ℓ lies between the slopes of the tangents to γ at its endpoints), and ℓ lies to the left of the tangent to γ with slope a . See Figure 3(ii).

Using carefully designed multi-level range-searching data structures, we can test whether some pair of an input line and an input arc satisfy (i) or (ii), in time $O(M^{2/3}N^{2/3}\text{polylog}(M+N)) = O(n^{8/3}\text{polylog}(n))$.³ Hence, within the same time bound, we can find a valid view of K , and thus a collision-free sliding motion of K through W , if one exists. That is, we have:



■ **Figure 3** A line ℓ intersecting a convex x -monotone elliptic arc γ : (i) The two endpoints of γ lie on different sides of ℓ . (ii) The two endpoints lie to the left of ℓ and ℓ lies to the left of the parallel tangent to the arc. (iii) The two endpoints lie to the right of ℓ (and then there is no intersection). (iv) The two endpoints lie to the left of ℓ but γ has no tangent parallel to ℓ (and then there is no intersection).

► **Theorem 3.** *Given K and W as above, we can determine whether K can slide through W in a collision-free manner, and, if so, find such a sliding motion, in time $O(n^{8/3}\text{polylog}(n))$.*

3 Unbounded windows

In this section we consider the variant in which W is an infinite slab in the xy -plane, bounded by, say, two vertical lines $x = 0$ and $x = a$. We refer to such a window as a *gate*. We show:


► **Theorem 4.** *Let K be a convex polytope that can be moved by some collision-free rigid motion through a gate W . Then there exists a sliding collision-free motion of K through W .*

We can therefore apply the machinery of Theorem 3, and conclude that we can determine whether K can be moved through W by a collision-free motion, in time $O(n^{8/3}\text{polylog}(n))$.

The full details of the proof of the theorem are given in the full version [14]. Here is brief sketch. By projecting the moving polytope K and W onto the xz -plane, W projects to the interval $g := [0, a] \times \{0\}$, and K projects to a time-varying convex polygon that starts from a placement that lies in the upper halfplane $z > 0$ and reaches a placement that lies in the lower halfplane $z < 0$. For technical reasons, we approximate K by a smooth convex body, and reduce the problem to the case where K is smooth and convex.


³ We are not aware of any published result that solves this problem efficiently. A different solution, with a similar performance bound, was improvised and sketched to us by Pankaj Agarwal, and we thank him deeply for the useful interaction concerning this problem.

At any time t during the motion, the projected planar region $\pi(K(t))$ (where $K(t)$ is the placement of K at time t) meets g at some interval $I(t)$ (we ignore the prefix and suffix of the motion where $K(t)$ does not yet meet, or no longer meets W). We consider the two tangents to $\pi(K(t))$ at the endpoints $\kappa^-(t)$, $\kappa^+(t)$ of $I(t)$, and note that, at the beginning of the motion, the wedge that these tangents form and that contains $K(t)$ points upwards, and at the end of the motion it points downwards; see Figure 4.



■ **Figure 4** Moving the projection of K through g . Left: At the beginning of the crossing of g , the tangents $\tau^-(t)$ and $\tau^+(t)$ “open up” (with respect to their sides that contain $K(t)$). Right: At the end of the crossing, they “open down”.

Since the tangents vary continuously (because $K(t)$ is always smooth), there must be a time t_0 at which the two tangents are parallel to each other, and thus span a slab σ (in the xz -plane) whose width is clearly $\leq a$. See Figure 5. The Cartesian product of σ and the y -axis yields a slab σ^* in $\mathbb{R}^3$, whose cross-section with the xy -plane is contained in W . This in turn implies that K can slide through W , and completes the proof. $\square$



■ **Figure 5** The critical instance t_0 where the tangents at $\kappa^-(t_0)$ and at $\kappa^+(t_0)$ become (anti-) parallel.

4 From passing through an arbitrary convex window to sliding through a gate

In this section we prove a similar yet different property of a convex polytope passing through an arbitrary compact planar convex window, not necessarily rectangular.

► **Theorem 5.** *Let W be an arbitrary compact convex region in the xy -plane. Let K be a convex polytope that can be moved by some collision-free motion (possibly full rigid motion, with six degrees of freedom) through W , and let d be the diameter of W (the maximum distance between any pair of points in W). Let h be an arbitrary plane, and let K_h be the*

orthogonal projection of K on h . Then K_h can be rigidly placed between two parallel lines at distance d . That is, for any fixed direction $\vec{v}$, K can slide, from its (arbitrary) initial placement, in direction $\vec{v}$ through a gate of width d , in a plane perpendicular to $\vec{v}$.

We provide two different topology-based proofs of Theorem 5, both presented in full detail in the full version [14]. We sketch here one of these proofs. But first here is an interesting corollary of the theorem, also proved in [14].

► **Corollary 6.** *If K can be moved through a rectangular window W of dimensions $a \times b$ by some collision-free motion, then K can slide through a rectangle of dimensions $\min(a, b) \times \sqrt{a^2 + b^2}$.*

Sketch of a proof of Theorem 5. Similar to the previous section, we first prove the theorem for smooth strongly convex compact bodies, and then extend the result to polytopes (see [14]). Consider the motion of K , now assumed to be a smooth, strongly convex, and compact body, during the time interval $[0, 1]$. Assume that at $t = 0$ (resp., at time $t = 1$), K lies fully above (resp., below) the xy -plane.

Fix some direction $\vec{v}$, and let $C = C(\vec{v})$ denote the silhouette of K when viewed in direction $\vec{v}$. Let h be some plane orthogonal to $\vec{v}$, and let π_h denote the orthogonal projection onto h . Parameterize a point $u \in C$ by the orientation θ of the tangent at $\pi_h(u)$ to $K_h := \pi_h(K)$, which is well defined since K is smooth and strongly convex, and let $\gamma = \gamma_h$ be the inverse of π_h , that is, $\gamma(\theta)$ is the unique point $u \in C$ such that $\pi_h(u) = \theta$. Since K is assumed to be strongly convex, K_h is also strongly convex, and γ is a well-defined and continuous function on $\mathbb{S}^1$. We extend γ to a bivariate function $\gamma^* : \mathbb{S}^1 \times [0, 1] \mapsto \mathbb{R}^3$, so that $\gamma^*(\theta, t)$ is the position (in the ambient 3-space) of $\gamma(\theta)$ at time t during the motion of K .

Let $\delta : \mathbb{S}^1 \times [0, 1] \mapsto \mathbb{R}$ be the function $\delta(\theta, t) = z(\gamma^*(\theta, t))$, namely, the z -coordinate of the corresponding point $\gamma(\theta)$ of C at time t . Note that at time $t = 0$ (resp., $t = 1$), δ is positive (resp., negative) at each θ , since K lies fully above (resp., below) the xy -plane at that time. Put $M := \max_{\theta \in \mathbb{S}^1} \delta(\theta, 0)$ and $m := \min_{\theta \in \mathbb{S}^1} \delta(\theta, 1)$, so $M > 0$ and $m < 0$.

The functions $\delta_0(\theta) = \delta(\theta, 0)$ and $\delta_1(\theta) = \delta(\theta, 1)$ are defined and continuous on $\mathbb{S}^1$, and we extend each of them to the closed unit disk $\mathcal{B}^1$ bounded by $\mathbb{S}^1$, in polar coordinates, which, for technical reasons, we write in reverse order as (θ, r) , by


$$\begin{aligned}\delta_0^*(\theta, r) &= r\delta_0(\theta) + (1-r)M \\ \delta_1^*(\theta, r) &= r\delta_1(\theta) + (1-r)m.\end{aligned}$$

It is easily checked that these extensions are well defined and continuous over $\mathcal{B}^1$. Moreover, $\delta_0^*(\theta, r) > 0$ and $\delta_1^*(\theta, r) < 0$ for every θ .

We now take our function δ , which is so far defined on the side surface S of the cylinder $\mathbb{S}^1 \times [0, 1]$, and extend it to the entire boundary $S^* := S \cup B_0 \cup B_1$ of the cylinder, so that δ coincides with δ_0^* on the base B_0 of the cylinder at $t = 0$, and with δ_1^* on the base B_1 at $t = 1$. Clearly, the extended δ is well defined and continuous over S^* .

To simplify the forthcoming analysis, we identify S^* with the unit sphere $\mathbb{S}^2$, which we parameterize by (θ, z) , where $\theta \in \mathbb{S}^1$ is the horizontal orientation of the point on $\mathbb{S}^2$ and z is its z -coordinate (so θ is not well defined at the north and south poles of $\mathbb{S}^2$). We use the simple homeomorphism f that maps a point $(\theta, t) \in S$ to $(\theta, t - 1/2) \in \mathbb{S}^2$, maps a point $(\theta, r) \in B_0$ to $(\theta, -1 + r/2) \in \mathbb{S}^2$, and maps a point $(\theta, r) \in B_1$ to $(\theta, 1 - r/2) \in \mathbb{S}^2$. See Figure 6 for an illustration.

Define a function G from $\mathbb{S}^2$ to $\mathbb{R}^2$ by $G(\theta, z) = (\delta(\theta, z), \delta(\theta + \pi, z))$, for $(\theta, z) \in \mathbb{S}^2$. Our goal is to show that $G(\mathbb{S}^2)$ contains the origin. Note that, by construction, $G(f(B_0))$ is fully contained in the positive quadrant $Q_1 := \{(x, y) \mid x, y > 0\}$, and $G(f(B_1))$ is fully contained



■ **Figure 6** Identifying S^* with the unit sphere $\mathbb{S}^2$. B_0 is shown in green, B_1 in orange, and S in light blue. In (i) S^* is depicted, in (ii) an intermediate snapshot of the deformation is shown, for visual convenience, and in (iii) the final unit ball is shown, divided into the three parts that correspond to B_0 , S and B_1 .

in the negative quadrant $Q_3 := \{(x, y) \mid x, y < 0\}$. Thus, if $G(\mathbb{S}^2)$ contains the origin then so does $G(f(S))$. Once this property is established, it provides us with a pair (θ, z) such that $\delta(\theta, z) = \delta(\theta + \pi, z) = 0$, which means that there are two antipodal points $u, v \in C$ that pass through W simultaneously. Therefore their distance must be at most the diameter of W , and hence also the distance between the parallel tangent planes through them, which is a slab parallel to $\vec{v}$ of width at most d that contains K , as asserted.

Assume to the contrary that $G(\mathbb{S}^2)$ does not contain the origin. Then we can normalize G to the function

$$H(\theta, z) := \frac{G(\theta, z)}{\|G(\theta, z)\|}, \quad \text{for } (\theta, z) \in \mathbb{S}^2,$$

which maps $\mathbb{S}^2$ continuously to the unit circle $\mathbb{S}^1$. The function G , and thus also the function H , are symmetric with respect to the line $y = x$ in $\mathbb{R}^2$, meaning that

$$\begin{aligned} G(\theta + \pi, z) &= \Sigma(G(\theta, z)), & \text{for } (\theta, z) \in \mathbb{S}^2, & \text{ and thus also} \\ H(\theta + \pi, z) &= \Sigma(H(\theta, z)), & \text{for } (\theta, z) \in \mathbb{S}^2, \end{aligned}$$

where Σ is the reflection about $y = x$, that is, $\Sigma(x, y) = (y, x)$.

We now use the property that the real line is a *covering space* of $\mathbb{S}^1$, in the specific (and easily verified) sense that the continuous map $p : \mathbb{R} \mapsto \mathbb{S}^1$, given by $p(x) = e^{2\pi i x}$, for $x \in \mathbb{R}$, is surjective, and, for each $w \in \mathbb{S}^1$, there exists an open neighborhood U of w such that $p^{-1}(U)$ is the disjoint union of open sets in $\mathbb{R}$, each of which is mapped homeomorphically to U by p . The map p is called the *covering map*.

A well known property of covering spaces is the *lifting property* (reviewed, e.g., in [1]; see also [15]), a special case of which asserts, in the specific context used here, that, if φ is any continuous map from $\mathbb{S}^2$ to $\mathbb{S}^1$ then φ can be *lifted* to a map $\psi : \mathbb{S}^2 \mapsto \mathbb{R}$, so that $p \circ \psi = \varphi$.

Applying the lifting property to the function H , we get a continuous mapping $T : \mathbb{S}^2 \mapsto \mathbb{R}$, such that $p \circ T = H$, so we have the property that

$$p(T(\theta + \pi, z)) = \Sigma(p(T(\theta, z))), \quad \text{for } (\theta, z) \in \mathbb{S}^2.$$

As is easily checked, we have $\Sigma(e^{iy}) = e^{i(\pi/2 - y)}$, and therefore, for a point $x \in \mathbb{R}$, we have

$$\Sigma(p(x)) = \Sigma(e^{2\pi i x}) = e^{\pi i/2 - 2\pi i x} = p(1/4 - x), \quad \text{so}$$

$$p(T(\theta + \pi, z)) = p(1/4 - T(\theta, z)), \quad \text{for } (\theta, z) \in \mathbb{S}^2.$$

This in turn implies, by the definition of p , that

$$T(\theta + \pi, z) = 1/4 + k_{\theta, z} - T(\theta, z),$$

for some integer $k_{\theta, z}$. However, since T is continuous, there must be a single integer k such that $k_{\theta, z} \equiv k$ for all θ and z . That is, we have

$$T(\theta + \pi, z) + T(\theta, z) = 1/4 + k, \quad \text{for all } (\theta, z) \in \mathbb{S}^2. \quad (2)$$

By an easy application of the mean-value theorem (which is also a special case of the Borsuk-Ulam theorem in dimension 1), there exist θ_0 and θ_1 such that, recalling that the value $z = -1/2$ (resp., $z = 1/2$) corresponds to points on the lower (resp., upper) circle bounding S ,

$$\begin{aligned} T(\theta_0 + \pi, -1/2) &= T(\theta_0, -1/2) \\ T(\theta_1 + \pi, 1/2) &= T(\theta_1, 1/2). \end{aligned}$$

Substituting in (2), we get

$$T(\theta_0, -1/2) = T(\theta_1, 1/2) = 1/8 + k/2.$$

However, by construction, $H(\theta_0, -1/2)$ lies in the first quadrant Q_1 , and $H(\theta_1, 1/2)$ lies in the third quadrant Q_3 . Hence we have $T(\theta_0, -1/2) \in (0, 1/4) + \mathbb{Z}$ and $T(\theta_1, 1/2) \in (1/2, 3/4) + \mathbb{Z}$, but $1/8 + k/2$ can belong to only one of these sets (depending on whether k is even or odd). This contradiction shows that $G(\mathbb{S}^2)$, and thus also $G(f(S))$, contains the origin, as asserted.

So far the proof was for a smooth strongly convex compact bodies. The extension to the case of a convex polytope K is done exactly as in the analysis in the preceding section (which is spelled out in the full version [14]). $\blacktriangleleft$

5 Purely translational motions

In this section we show that purely translational motions of K through a rectangular window W are not more powerful than sliding. Specifically, we have the following theorem, which is, in a sense, a strengthening of Lemma 1. The proofs of the theorems in this section are given in [14].

► **Theorem 7.** *If K can be moved through a rectangular window W by a purely translational collision-free motion in some fixed orientation Θ , then K can be moved through W , possibly from some other starting position, by sliding while keeping the same orientation Θ .*

We also address a more restricted case where we are given a prescribed orientation Θ and we wish to find a purely translational motion for K with this orientation. We denote the polytope K at orientation Θ (ignoring translations) by K^Θ . We obtain the following results.

► **Theorem 8.** *Given an orientation Θ , we can determine whether a translational motion for K^Θ through the rectangular W exists, and if so find a sliding motion for K^Θ through W , in $O(n)$ time.*

► **Theorem 9.** *Let W be an arbitrary (not necessarily convex) polygonal window with k edges, lying in the xy -plane. Given a prescribed orientation Θ of K , we can determine whether a translational motion for K^Θ through W exists, and, if so, find such a motion, in $O(nk \log k \log kn)$ randomized expected time.*


6 Rotations are needed

So far we have considered versions of the problem in which we were able to show that the existence of an arbitrary collision-free motion of K through W implies that K can also slide through W (or, in one instance, through another window related to W). However, perhaps not very surprisingly, this is not the case in general. We show in this and the following section that in general rotations are needed to obtain a collision-free motion of the polytope through the window.

► **Lemma 10.** *Let W be a square window with side length $\sqrt{5}$. Let $A = (0,0,0)$, $B = (1,3,0)$, $C = (1,0,h)$, $D = (0,3,h)$ be four points, where $h \gg 1$ is a sufficiently large parameter, and let K be the tetrahedron $ABCD$ (see Figure 7). Then*

1. *K cannot pass through W by any purely translational collision-free motion (for sufficiently large $h \gg 1$).*
2. *K can pass through W by a collision-free motion with only two degrees of freedom: translating in the z -direction combined with rotation around a z -vertical axis (for any value of $h > 0$).*

The proof is given in the full version [14].



■ **Figure 7** The tetrahedron $K = ABCD$ of Lemma 10.

7 The case of a circular window

In this section we study the case where W is a circular window. There are (at least) three possible types of motion of K through W : sliding, purely translational motion, and general motion with all six degrees of freedom. In this section we show that these types are not equivalent, as spelled out in the following theorem.

► **Theorem 11.** *Let K be the regular tetrahedron of side length 1. Then there exist two threshold parameters $1 > \delta_1 \approx 0.901388 > \delta_2 \approx 0.895611$, so that, denoting by d the diameter of W , we have:*

- (i) *K can slide through W if $d \geq 1$.*
- (ii) *K cannot slide through W , but can pass through W by a purely translational motion, if $\delta_1 \leq d < 1$.*
- (iii) *K cannot pass through W by a purely translational motion, but can pass through W by a general motion, if $\delta_2 \leq d < \delta_1$.*
- (iv) *K cannot pass through W at all if $d < \delta_2$.*

Proof sketch; the full proof is in [14].

- (i) If $d \geq 1$ then, as is easy to show, K can slide through W , because K can be enclosed in a cylinder of diameter 1, whose axis is orthogonal to two opposite edges of K .
- (ii) We need to show that, for this range of d , (a) K cannot slide through W at any fixed orientation and direction of sliding, but (b) K can pass through W by a collision-free purely translational motion.

For (a), a suitable adaptation of Lemma 1 shows that if K can slide through W in some direction then it can also slide in the vertical z -direction. It therefore suffices to show that K cannot be contained in a cylinder of diameter smaller than 1. A proof of this fact can be found in [2]. For the sake of completeness, we reproduce the proof in the full version [14], where we also provide the full details of the proof of (b).

- (iii) We construct a simple five-step motion of K through W , or rather of W through K . The details are given in the full version [14], and are illustrated in Figure 8.




Figure 8 Moving W around K . (i) The initial configuration. (ii) Translating W until it contains the triangle BUV . (iii) Rotating W around UV until it contains the rectangle $U'V'U'V$. (iv) Translating W until it contains the symmetric rectangle, with edge lengths swapped. The remainder of the motion is a fully symmetric reversal of the first two steps.

For this five-step motion to work, there has to exist a cross section through B (BUV in Figure 8(ii)) that can be enclosed by a disc of diameter d (and we show, in the full version [14], that this ensures that the entire motion is collision-free). A numerical calculation shows that the cross section through a vertex with the smallest enclosing disc is such that the diameter of this disc is $\delta_2 \approx 0.895611$. This establishes (iii).

- (iv) When $d < \delta_2$ no motion is possible because no cross section through a vertex can be enclosed by a disc of diameter d , as just argued. ◀

8 Planning general rigid collision-free motion of a convex polytope through a rectangular window

Finally we deal with the general case, in which the motion of K has all six degrees of freedom. By standard (and general) arguments in algorithmic motion planning the free configuration space for this problem has complexity $O(n^6)$, and it can be computed in $O(n^{8+\epsilon})$ time [13], from which we can easily extract a solution path, when one exists, within the same time bound. We show here that we can exploit the special structure of the problem at hand to find a solution, or detect and notify that none exists, in time close to $O(n^4)$. We sketch below the main ideas; the full details are given in the full version [14].

If there is a solution path for K to move through W with all six degrees of freedom, then there is also a canonical solution path where at all times at which K intersects the plane of W (namely the xy -plane), K touches the bottom and left edges of W with two edges e_b

and e_ℓ (possibly with the closure of these edges, namely with vertices of K , and possibly with more than one edge touching a side of W). During this motion, for every point on the path define e_t to be the edge of K whose intersection with the xy -plane has the largest y -coordinate, and e_r to be the edge of K whose intersection with the xy -plane has the largest x -coordinate.

We now split the canonical solution path into maximal open segments, along which the open edges e_b, e_ℓ, e_t , and e_r are fixed. We construct a collection of four-dimensional subspaces of the full-dimensional configuration space, one for each such quadruple e_b, e_ℓ, e_t, e_r of four edges, consisting of those free placements that have those four edges as the extreme edges in the x - and y -directions within the xy -plane. This can be done in total $O(n^4)$ time since each of these subspaces has constant descriptive complexity.

The major remaining problem is to economically detect the free connections among these $O(n^4)$ subspaces. The efficiency of our approach relies on the following lemma (proved in [14]), which asserts that the total number of certain quintuplets of edges of K that encode these connections is only $O(n^4)$, rather than $O(n^5)$, and that they can be computed efficiently:

► **Lemma 12.** *The maximum number of quintuplets $(e_r, e_t, e_\ell, e_b, e_\xi)$, where e_r, e_t, e_ℓ, e_b are as defined above, and e_ξ is another edge of K whose intersection with the xy -plane h_{xy} has the same x - (respectively, y -) coordinate as the intersection with h_{xy} of e_ℓ or e_r (respectively, e_b or e_t), is $O(n^4)$. All these quintuplets can be computed in $O(n^3 \lambda_q(n) \log n)$ time⁴ for some small constant q .*

This in turn leads to the following summary result.

► **Theorem 13.** *Given a convex polytope K with n edges and a rectangular window W , we can construct a collision-free motion of K through W , if one exists, or determine that no such motion exists, in time $O(n^3 \lambda_q(n) \log n)$, for some small constant q . The algorithm requires $O(n^4)$ storage.*

The proofs and the algorithm are given in the full version [14].

9 Conclusion and further research

In this paper we have studied a variety of problems concerning collision-free motion of a convex polytope through a planar window, under several kinds of allowed motion – sliding (translating in a fixed direction), purely translational motion, and general motion. We have presented several structural properties and characterizations of such motions, and obtained efficient algorithms for several special cases, as well as for the general case.

There are several open problems and directions for further research. One such direction is to show that the near-quartic upper bound, established in Section 8, on the cost of the general motion planning problem for K and a rectangular window W is almost tight in the worst case, in the specific sense of establishing a lower bound $\Omega(n^4)$ on the worst-case complexity of the resulting free configuration space, a property that we conjecture to hold, and have in fact an initial plan for establishing this bound.

In addition, in Section 6, we presented an example in which a rotation is needed to pass a polytope through a rectangular window. However, in this construction we only used rotation about the line perpendicular to the plane that contains the window. This suggests

⁴ $\lambda_q(n)$ is a near-linear function related to Davenport-Schinzel sequences [23].

the conjecture that every convex polytope that can pass through a rectangular window W can also pass through W by a motion consisting of arbitrary translations and rotations only about the line perpendicular to the plane of W . The results of Section 7 show that for circular windows this claim is false in general, but the status of the conjecture is still open for a rectangular window.

It is also not clear what can be said about the motion of a general non-convex polytope through a rectangular, general convex, or even non-convex window. There are several variants of this question, depending on the type of motion that we allow, both in terms of structural properties of the motion, and of the efficiency of algorithms for performing it.

References

- 1 https://en.wikipedia.org/wiki/Covering_space.
- 2 <https://math.stackexchange.com/questions/1364880/smallest-cylinder-into-which-a-regular-tetrahedron-can-fit>.
- 3 P. Bose, D. Halperin, and S. Shamaï. On the separation of a polyhedron from its single-part mold. In *13th IEEE Conference on Automation Science and Engineering*, pages 61–66, 2017.
- 4 B. Chazelle, H. Edelsbrunner, L. J. Guibas, and M. Sharir. A singly exponential stratification scheme for real semi-algebraic varieties and its applications. *Theor. Comput. Sci.*, 84(1):77–105, 1991.
- 5 H. Choset, K. M. Lynch, S. Hutchinson, G. Kantor, W. Burgard, L. E. Kavraki, and S. Thrun. *Principles of Robot Motion: Theory, Algorithms, and Implementation*. MIT Press, 2005.
- 6 H. T. Croft, K. J. Falconer, and R. K. Guy. *Unsolved Problems in Geometry*. Problem Books in Mathematics. Springer Verlag, Heidelberg, 1991.
- 7 H. E. Debrunner and P. Mani-Levitska. Can you cover your shadows? *Discrete Comput. Geom.*, 1:45–58, 1986.
- 8 E. Fogel, D. Halperin, and R. Wein. *CGAL Arrangements and their Applications – A Step-by-Step Guide*, volume 7 of *Geometry and Computing*. Springer, 2012.
- 9 P. Gibbs. A computational study of sofas and cars. *Computer Science*, 2:1–5, 2014.
- 10 D. Halperin, L. Kavraki, and K. Solovey. Robotics. In *Handbook of Discrete and Computational Geometry*, chapter 51, pages 1343–1376. Chapman & Hall/CRC, 3 edition, 2018.
- 11 D. Halperin, J.-C. Latombe, and R. H. Wilson. A general framework for assembly planning: The motion space approach. *Algorithmica*, 3-4:577–601, 2000.
- 12 D. Halperin, O. Salzman, and M. Sharir. Algorithmic motion planning. In *Handbook of Discrete and Computational Geometry*, chapter 50, pages 1311–1342. Chapman & Hall/CRC, 3 edition, 2018.
- 13 D. Halperin and M. Sharir. Arrangements. In *Handbook of Discrete and Computational Geometry*, chapter 28, pages 723–762. Chapman & Hall/CRC, 3 edition, 2018.
- 14 D. Halperin, M. Sharir, and I. Yehuda. Throwing a sofa through the window. *CoRR*, abs/2102.04262, 2021. [arXiv:2102.04262](https://arxiv.org/abs/2102.04262).
- 15 A. Hatcher. *Algebraic Topology*. Cambridge University Press, Cambridge, 2002.
- 16 L. E. Kavraki, P. Šestka, J.-C. Latombe, and M. H. Overmars. Probabilistic roadmaps for path planning in high dimensional configuration spaces. *IEEE Transactions on Robotics*, 12(4):566–580, 1996.
- 17 J. J. Kuffner and S. M. LaValle. RRT-Connect: An efficient approach to single-query path planning. In *IEEE International Conference on Robotics and Automation (ICRA)*, pages 995–1001, 2000.
- 18 J.-C. Latombe. *Robot Motion Planning*, volume 124 of *The Kluwer international series in engineering and computer science*. Kluwer, 1991.
- 19 S. M. LaValle. *Planning Algorithms*. Cambridge University Press, 2006.
- 20 W. H. Plantinga and C. R. Dyer. Visibility, occlusion, and the aspect graph. *Int. J. Computer Vision*, 5:137–160, 1990.

41:16 Throwing a Sofa Through the Window

- 21 O. Salzman, M. Hemmer, and D. Halperin. On the power of manifold samples in exploring configuration spaces and the dimensionality of narrow passages. *IEEE Trans Autom. Sci. Eng.*, 12(2):529–538, 2015.
- 22 O. Salzman, M. Hemmer, B. Raveh, and D. Halperin. Motion planning via manifold samples. *Algorithmica*, 67(4):547–565, 2013.
- 23 M. Sharir and P. K. Agarwal. *Davenport–Schinzel Sequences and Their Geometric Applications*. Cambridge University Press, New York, 1995.
- 24 J. Snoeyink and J. Stolfi. Objects that cannot be taken apart with two hands. *Discrete Comput. Geom.*, 12:367–384, 1994.
- 25 G. Toussaint. Movable separability of sets. *Comput. Geom.*, 2:335–375, 1985.