

RESEARCH



Everywhere local solubility for hypersurfaces in products of projective spaces

Tom Fisher¹, Wei Ho^{2*} and Jennifer Park³

*Correspondence:
weihuo@umich.edu

²Department of Mathematics,
University of Michigan, Ann
Arbor, MI 48109, USA

Full list of author information is
available at the end of the article

Abstract

We prove that a positive proportion of hypersurfaces in products of projective spaces over \mathbb{Q} are everywhere locally soluble, for almost all multidegrees and dimensions, as a generalization of a theorem of Poonen and Voloch [25]. We also study the specific case of genus 1 curves in $\mathbb{P}^1 \times \mathbb{P}^1$ defined over \mathbb{Q} , represented as bidegree (2, 2)-forms, and show that the proportion of everywhere locally soluble such curves is approximately 87.4%. As in the case of plane cubics [2], the proportion of these curves in $\mathbb{P}^1 \times \mathbb{P}^1$ soluble over \mathbb{Q}_p is a rational function of p for each finite prime p . Finally, we include some experimental data on the Hasse principle for these curves.

1 Introduction

Let V be a variety defined over \mathbb{Q} . The study of rational points on V often involves determining the local points $V(\mathbb{Q}_v)$ for completions \mathbb{Q}_v of \mathbb{Q} . We say that V is *(globally) soluble* if the set $V(\mathbb{Q})$ of rational points is nonempty, and V is *everywhere locally soluble* if the set $V(\mathbb{Q}_v)$ is nonempty for all places $v \leq \infty$ of \mathbb{Q} .

Poonen and Voloch [25] show that a positive proportion of all hypersurfaces in \mathbb{P}^n of fixed degree d are everywhere locally soluble for $n, d \geq 2$ and $(n, d) \neq (2, 2)$. In particular, they prove that this proportion (as a limit) is exactly the product $c = \prod_v c_v > 0$ of local factors c_v , where c_v is the proportion of hypersurfaces that have a \mathbb{Q}_v -point. In [1], Bright, Browning, and Loughran generalized this theorem to other families of varieties. Poonen and Voloch also conjecture that the proportion of globally soluble hypersurfaces is c when $2 \leq d \leq n$ (with $c = 0$ when $(n, d) = (2, 2)$), and is 0 when $d > n + 1$. This conjecture implies that the Hasse principle is true for 100% and for 0% of the everywhere locally soluble hypersurfaces in the two cases, respectively. Browning, Le Boudec, and Sawin [10] have recently proved this conjecture for $2 \leq d \leq n$ and $(n, d) \neq (3, 3)$, i.e., that the Hasse principle is satisfied 100% of the time for Fano hypersurfaces of degree d and dimension at least 3 in \mathbb{P}^n .

However, Poonen and Voloch [25] do not make any conjectures about local versus global solubility in the boundary (Calabi-Yau) case where $d = n + 1$; this case is perhaps

the most interesting and the most mysterious. The simplest of these cases, that of plane cubics ($d = 3$ and $n = 2$), has been studied by Bhargava, Cremona, and the first author [2], in which the authors explicitly compute the proportion c of everywhere locally soluble plane cubics; the same authors [3] also study the proportion of everywhere locally soluble hyperelliptic curves. In addition, Bhargava [9] shows that a positive proportion of plane cubics fail the Hasse principle and a positive proportion satisfy the Hasse principle.

In this paper, we are interested in answering the analogous questions and computing the explicit proportion of everywhere locally soluble hypersurfaces defined in *products* of projective spaces. We also study a specific family of genus one curves over \mathbb{Q} (analogous to the plane cubics studied in [2]) and determine the proportion of such curves that are everywhere locally soluble. In forthcoming work, Bhargava and the second author show that the Hasse principle fails for a positive proportion of curves in this family [7]. In this paper, we include some data on the success and failure of the Hasse principle for randomly selected sets of these curves with bounded coefficients.

In Sect. 2, we begin by proving the analogue of the theorem of Poonen and Voloch for hypersurfaces in products of projective spaces:

Theorem 1.1 *Let $d_1, \dots, d_k, n_1, \dots, n_k$ be positive integers. The proportion of multidegree (d_1, \dots, d_k) hypersurfaces over \mathbb{Q} in $\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k}$ that are everywhere locally soluble tends to a real number $c > 0$, where c is the product $\prod_v c_v$ over all places v and c_v is the proportion of such multidegree homogeneous polynomials with a nontrivial zero over \mathbb{Q}_v , as long as one of the following holds:*

- (i) $k = 1$ and $n_1 \geq 2$ but not $(n_1, d_1) = (2, 2)$;
- (ii) $k \geq 2$.

For the cases with $k = 1$ not covered by (i), it may be shown that the proportion of everywhere locally soluble hypersurfaces is 0.

We also predict the success or failure of the Hasse principle for families of hypersurfaces in products of projective spaces in many cases, depending on the multidegree of the hypersurfaces:

Conjecture 1.2 For any positive integers d_i, n_i for $i \in \{1, \dots, k\}$, consider the family of multidegree (d_1, \dots, d_k) -hypersurfaces in $\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k}$ over \mathbb{Q} . Then

- (i) If $d_i > n_i + 1$ for all $i = 1, \dots, k$, then 100% of everywhere locally soluble (d_1, \dots, d_k) -hypersurfaces fail the Hasse principle.
- (ii) If $d_i < n_i + 1$ for some $i = 1, \dots, k$, then 100% of everywhere locally soluble (d_1, \dots, d_k) -hypersurfaces satisfy the Hasse principle. With these conditions, we have that the proportion of soluble hypersurfaces equals the product of local factors $c = \prod_v c_v$.

Note that the numerical conditions in Conjecture 1.2(ii) are significantly more general than just requiring the hypersurface to be Fano, which is equivalent to having $d_i < n_i + 1$ for *all* i (not just some i). Still, as in the original Poonen–Voloch heuristic, there are some remaining boundary cases, namely when $d_i \geq n_i + 1$ for all i with equality for at least one i . In this paper, we compute the explicit probability of everywhere local solubility for the simplest such case with $k > 1$, namely the family of bidegree $(2, 2)$ curves in $\mathbb{P}^1 \times \mathbb{P}^1$.

Explicitly, these are the curves defined by the bihomogeneous bidegree (2, 2)-polynomials of the form

$$\begin{aligned} F(X_0, X_1, Y_0, Y_1) = & a_{00}X_0^2Y_0^2 + a_{01}X_0^2Y_0Y_1 + a_{02}X_0^2Y_1^2 \\ & + a_{10}X_0X_1Y_0^2 + a_{11}X_0X_1Y_0Y_1 + a_{12}X_0X_1Y_1^2 \\ & + a_{20}X_1^2Y_0^2 + a_{21}X_1^2Y_0Y_1 + a_{22}X_1^2Y_1^2 \end{aligned} \quad (1.3)$$

where $a_{ij} \in \mathbb{Q}$ for $i, j \in \{0, 1, 2\}$. Smooth curves of this form have genus one. We show:

Theorem 1.4 *For a finite prime p , the probability a bidegree (2, 2)-form with coefficients in \mathbb{Z}_p is soluble over \mathbb{Q}_p is*

$$\rho(p) = 1 - \frac{p(p-1)(p^2-1)f(p)}{8(p^8-1)(p^9-1)}$$

where $f(p) = 4p^{11} - 4p^{10} + 4p^9 - p^8 + 5p^7 - 2p^6 + 5p^5 - p^4 + 2p^3 - 2p^2 + 6p - 2$.

The probability a (2, 2)-form is soluble over the reals (when we choose the coefficients uniformly at random in $[-B, \dots, B]$ with B large) appears to be about 96.46%, based on a Monte Carlo simulation. Note that the probability of solubility over \mathbb{R} is greater than $7/8$, since for insolubility $a_{00}, a_{02}, a_{20}, a_{22}$ must all have the same sign.

Theorem 1.4 gives $\prod_{p < \infty} \rho(p) \approx 0.90592$. Combining this with the estimate of the previous paragraph, we have

$$\prod_{p \leq \infty} \rho(p) \approx 0.8739,$$

thereby giving the following theorem:

Theorem 1.5 *The probability that a random (2, 2)-form with coefficients in \mathbb{Q} is p -locally soluble for all $p < \infty$ is approximately 90.592%; assuming the truth of the Monte Carlo experiment, the probability that a random (2, 2)-form with coefficients in \mathbb{Q} is everywhere locally soluble is approximately 87.39%.*

A heuristic similar to that in [9] suggests that of the (2, 2)-forms that are everywhere locally soluble, exactly $1/4$ should be globally soluble. In Sect. 6, we report on an experiment to test this prediction numerically.

2 Generalization of Poonen–Voloch’s theorem and conjecture

In this section, we generalize the main result and conjectures of [25] to hypersurfaces in products of projective spaces. The main idea of the proof of the theorem is identical to that of [25] but relies on a more general combinatorial inequality. We work over the field \mathbb{Q} for this section.

Fix an integer $k \geq 1$ and positive integers n_i, d_i for $1 \leq i \leq k$. We consider multidegree (d_1, \dots, d_k) hypersurfaces in the product space $\mathcal{P} := \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k}$. In particular, let $\mathbb{Z}[\{x_{ij}\}]_{\mathbf{d}}$ denote the set of multihomogeneous polynomials in $\mathbb{Z}[\{x_{ij}\} : 1 \leq i \leq k, 0 \leq j \leq n_i]$ of multidegree $\mathbf{d} = (d_1, \dots, d_k)$. There are $m_i := \binom{n_i+d_i}{d_i}$ monomials of degree d_i in $(n_i + 1)$ variables, so the total number of monomials of multidegree \mathbf{d} in $\mathbb{Z}[\{x_{ij}\}]_{\mathbf{d}}$ is $m = \prod_{i=1}^k m_i$.

Define the *height* $h(f)$ of $f \in \mathbb{Z}[\{x_{ij}\}]_{\mathbf{d}}$ to be the maximum of the absolute values of the coefficients of f . Let $M_{\mathbb{Q}}$ be the set of places of \mathbb{Q} . We define the following counts for

$H > 0$ and $v \in M_{\mathbb{Q}}$:

$$\begin{aligned} N_{\text{tot}}(H) &:= \#\{f \in \mathbb{Z}[\{x_{ij}\}]_{\mathbf{d}} : h(f) \leq H\} = (2\lfloor H \rfloor + 1)^m \\ N(H) &:= \#\{f \in \mathbb{Z}[\{x_{ij}\}]_{\mathbf{d}} : h(f) \leq H \text{ and } \exists \mathbf{x} = \{x_{ij}\} \in \prod_{i=1}^k \mathbb{Z}^{n_i+1} \setminus \{0\} \text{ with } f(\mathbf{x}) = 0\} \\ N_v(H) &:= \#\{f \in \mathbb{Z}[\{x_{ij}\}]_{\mathbf{d}} : h(f) \leq H \text{ and } \exists \mathbf{x} = \{x_{ij}\} \in \prod_{i=1}^k \mathbb{Q}_v^{n_i+1} \setminus \{0\} \text{ with } f(\mathbf{x}) = 0\} \\ N_{\text{loc}}(H) &:= \#\{f \in \mathbb{Z}[\{x_{ij}\}]_{\mathbf{d}} : h(f) \leq H \text{ and } \forall v \in M_{\mathbb{Q}}, \exists \mathbf{x} \in \prod_{i=1}^k \mathbb{Q}_v^{n_i+1} \setminus \{0\} \text{ with } f(\mathbf{x}) = 0\}. \end{aligned}$$

In other words, $N_{\text{tot}}(H)$ is the total number of multidegree \mathbf{d} polynomials in \mathcal{P} of height at most H , and $N(H)$, $N_v(H)$, and $N_{\text{loc}}(H)$ are the number of such polynomials that are globally soluble, soluble over \mathbb{Q}_v , and everywhere locally soluble, respectively. The limits

$$\lim_{H \rightarrow \infty} \frac{N(H)}{N_{\text{tot}}(H)} \quad \text{and} \quad \lim_{H \rightarrow \infty} \frac{N_{\text{loc}}(H)}{N_{\text{tot}}(H)},$$

if they exist, will be called the proportion of globally soluble and everywhere locally soluble multidegree \mathbf{d} hypersurfaces, respectively. Note that for any place v , the local proportion $\lim_{H \rightarrow \infty} N_v(H)/N_{\text{tot}}(H)$ exists. Indeed, as explained in Remark 2.3 of [25], after normalizing the Haar measure (or Lebesgue measure for $v = \infty$) on the space \mathbb{Z}_v^m of the multihomogeneous polynomials of degree \mathbf{d} in the variables $\{x_{ij} : 1 \leq i \leq k, 0 \leq j \leq n_i\}$, we see that this local proportion is the measure of the v -adically closed subset of \mathbb{Z}_v^m corresponding to the multihomogeneous polynomials with a nontrivial zero over \mathbb{Q}_v . We restate Theorem 1.1 using this language, now ignoring some of the trivial cases where some $d_i = 1$:

Theorem 2.1 *Let $c_v = \lim_{H \rightarrow \infty} N_v(H)/N_{\text{tot}}(H)$. In the limit $H \rightarrow \infty$, the proportion $N_{\text{loc}}(H)/N_{\text{tot}}(H)$ tends to $c := \prod_v c_v > 0$, if one of the following conditions holds:*

- (i) $k = 1$ and $n_1, d_1 \geq 2$ but not $(n_1, d_1) = (2, 2)$;
- (ii) $k = 2$, and if $n_1 = n_2 = 1$, then $d_1, d_2 \geq 2$;
- (iii) $k \geq 3$.

Proof If f is absolutely irreducible modulo p , then for sufficiently large p , the Lang-Weil estimate guarantees a smooth point on the hypersurface $f = 0$ modulo p , which may be lifted to a \mathbb{Q}_p -point by Hensel's lemma. By Lemmas 20 and 21 of [24], it suffices to show that the space of reducible polynomials is of codimension at least 2 in the space of all polynomials in $\mathbb{Z}[\{x_{ij}\}]_{\mathbf{d}}$. This follows from Lemma 2.2 below, which shows that the product of the projective spaces of multidegree (r_1, \dots, r_k) polynomials and of multidegree $(d_1 - r_1, \dots, d_k - r_k)$ polynomials has dimension at most $m - 3$, where $m - 1$ is the dimension of the projective space of multidegree (d_1, \dots, d_k) polynomials. \square

Lemma 2.2 *Fix a positive integer k . Given positive integers n_i, d_i, r_i for $1 \leq i \leq k$ such that $0 \leq r_i \leq d_i$ for all i but $0 < \sum_i r_i < \sum_i d_i$, we have*

$$\prod_{i=1}^k \binom{n_i + r_i}{n_i} + \prod_{i=1}^k \binom{n_i + d_i - r_i}{n_i} < \prod_{i=1}^k \binom{n_i + d_i}{n_i} \quad (2.3)$$

if one of the following conditions holds:

- (i) $k = 1$ and $n_1, d_1 \geq 2$ but not $(n_1, d_1) = (2, 2)$;
- (ii) $k = 2$, and if $n_1 = n_2 = 1$, then $d_1, d_2 \geq 2$;
- (iii) $k \geq 3$.

Proof We define

$$S = \{(A_1, \dots, A_k) : A_i \subset \{1, 2, \dots, n_i + d_i\} \text{ with } |A_i| = n_i\}$$

and subsets

$$S_1 = \{(A_1, \dots, A_k) \in S : A_i \cap \{1, \dots, r_i\} = \emptyset \text{ for all } 1 \leq i \leq k\},$$

$$S_2 = \{(A_1, \dots, A_k) \in S : A_i \cap \{r_i + 1, \dots, d_i\} = \emptyset \text{ for all } 1 \leq i \leq k\}.$$

Then $|S_1 \cap S_2| = 1$. To prove $|S_1| + |S_2| < |S|$ it suffices to show that $|S \setminus (S_1 \cup S_2)| \geq 2$.

- (i) If $k = 1$ then by choosing A_1 with

$$|A_1 \cap \{1, \dots, r_1\}| = 1 \quad \text{and} \quad |A_1 \cap \{r_1 + 1, \dots, d_1\}| = 1$$

(and so $|A_1 \cap \{d_1 + 1, \dots, n_1 + d_1\}| = n_1 - 2$) we have $|S \setminus (S_1 \cup S_2)| \geq r_1(d_1 - r_1) \binom{n_1}{2}$, which is at least 2 under the stated hypotheses.

- (ii) If $k = 2$ then by choosing (A_1, A_2) with

$$\begin{aligned} |A_1 \cap \{1, \dots, r_1\}| &= \delta, & |A_1 \cap \{r_1 + 1, \dots, d_1\}| &= 1 - \delta, \\ |A_2 \cap \{1, \dots, r_2\}| &= 1 - \delta, & |A_2 \cap \{r_2 + 1, \dots, d_2\}| &= \delta, \end{aligned}$$

for $\delta = 0, 1$, we have $|S \setminus (S_1 \cup S_2)| \geq (r_1(d_2 - r_2) + r_2(d_1 - r_1))n_1n_2$, which is at least 2 under the stated hypotheses.

- (iii) If $k \geq 3$ then $|S \setminus (S_1 \cup S_2)| \geq \sum_{i \neq j} r_i(d_j - r_j) \geq 2$. \square

We also discuss the analogue of the conjecture of Poonen–Voloch in our setting:

Conjecture 2.4 For any positive integers d_i, n_i for $i \in \{1, \dots, k\}$, consider the family of multidegree (d_1, \dots, d_k) -hypersurfaces in $\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k}$ over \mathbb{Q} . Then

- (i) If $d_i > n_i + 1$ for all $i = 1, \dots, k$, then 100% of everywhere locally soluble (d_1, \dots, d_k) -hypersurfaces fail the Hasse principle.
- (ii) If $d_i < n_i + 1$ for some $i = 1, \dots, k$, then 100% of everywhere locally soluble (d_1, \dots, d_k) -hypersurfaces satisfy the Hasse principle.

Combining Conjecture 2.4(ii) with Theorem 2.1 implies that the proportion of globally soluble hypersurfaces in \mathcal{P} is a product of local densities for many cases (the last part of Conjecture 1.2): If $d_i < n_i + 1$ for some $i = 1, \dots, k$, we have that

$$\lim_{H \rightarrow \infty} \frac{N(H)}{N_{\text{tot}}(H)} = c = \prod_v c_v. \quad (2.5)$$

Remark 2.6 If $d_i < n_i + 1$ for some i , and yet the hypotheses of Theorem 2.1 are not satisfied then either some $d_i = 1$ or $(k, n_1, d_1) = (1, 2, 2)$. In the first case both sides of (2.5) are 1. In the second case concerning plane conics, see, e.g., Theorem 2 of [4], which states that $c_p = 1 - \frac{p}{2(p+1)^2}$ for p a prime; and immediately implies the result of Serre [28] that both sides of (2.5) are 0 in this case.

2.1 Motivation for Conjecture 2.4(i)

Fix a point $a = (a_1, \dots, a_k) \in \mathbb{Z}^{n_1+1} \times \dots \times \mathbb{Z}^{n_k+1}$ with each $a_i \neq 0$ and having coprime coordinates, and consider the set of multidegree (d_1, \dots, d_k) -polynomials vanishing on a . The set of these polynomials form a hyperplane in \mathbb{Z}^m , and to count the multidegree (d_1, \dots, d_k) polynomials up to height H vanishing on a is to count the set of integral points contained in this hyperplane, whose coordinates are bounded by H . As in [25], the number of integral points of height at most H in this hyperplane is given by

$$\frac{c(a)H^{m-1}}{\phi(a)} + O(H^{m-2}),$$

where $\phi(a)$ denotes the covolume of the lattice of integer points on the hyperplane of polynomials vanishing on a , and $c(a)$ denotes the $(m-1)$ -dimensional volume of the part of the hyperplane inside $[-1, 1]^m$.

Ignoring the error term, we get that

$$N(H) \leq H^{m-1} \sum_a \frac{c(a)}{\phi(a)},$$

where the sum ranges over $a \in \mathbb{Z}^{n_1+1} \times \dots \times \mathbb{Z}^{n_k+1}$, excluding the zero vectors in each of the components. Since $c(a)$ is bounded by definition, it remains to understand the convergence of $\sum_a \frac{1}{\phi(a)}$. Lemma 3.1 of [25] shows that $\phi(a)$ equals the Euclidean norm of the vector b formed by the monomials of degree (d_1, \dots, d_k) in the coordinates of (a_1, \dots, a_k) (the coprime condition in that lemma is not needed). Let $\phi_i(a_i)$ denote the Euclidean norm of the vector formed by plugging in the coordinates of a_i into each of the degree d_i monomials in $n_i + 1$ variables. Because $\phi(a) = \prod_{i=1}^k \phi_i(a_i)$, we have

$$\begin{aligned} & \sum_{a \in (\mathbb{Z}^{n_1+1} \setminus \{0\}) \times \dots \times (\mathbb{Z}^{n_k+1} \setminus \{0\})} \frac{1}{\phi(a)} \\ &= \left(\sum_{a_1 \in \mathbb{Z}^{n_1+1} \setminus \{0\}} \frac{1}{\phi_1(a_1)} \right) \dots \dots \left(\sum_{a_k \in \mathbb{Z}^{n_k+1} \setminus \{0\}} \frac{1}{\phi_k(a_k)} \right). \end{aligned} \quad (2.7)$$

We claim that each of the sums $\sum_{a_i \in \mathbb{Z}^{n_i+1} \setminus \{0\}} \frac{1}{\phi_i(a_i)}$ converges.

We decompose $\mathbb{Z}^{n_i+1} \setminus \{0\} = T_{n_i+1} \cup T_{n_i} \cup \dots \cup T_1$, where T_j consists of the vectors in \mathbb{Z}^{n_i+1} with exactly j nonzero coordinates. Let $a_i = (a_{i0}, \dots, a_{in_i})$. Then by applying the AM-GM inequality to the definition of $\phi_i(a_i)$, we have

$$\phi_i(a_i) \geq m_i^{1/2} \left| \prod_{\ell=0}^{n_i} a_{i\ell} \right|^{\frac{d_i}{n_i+1}}. \quad (2.8)$$

In fact, for $a_i \in T_j$, the product over ℓ on the right side of (2.8) may be taken to be the product of only the j nonzero $a_{i\ell}$ if we replace m_i with $m_{ij} := \binom{j+d_i}{d_i}$ and $n_i + 1$ by j . We thus have

$$\begin{aligned} \sum_{a \in \mathbb{Z}^{n_i+1} \setminus \{0\}} \frac{1}{\phi_i(a_i)} &= \sum_{j=1}^{n_i+1} \sum_{a_i \in T_j} \frac{1}{\phi_i(a_i)} \\ &\leq \sum_{j=1}^{n_i+1} \sum_{a_i \in T_j} m_{ij}^{-1/2} \left| \prod_{\ell: a_{i\ell} \neq 0} a_{i\ell} \right|^{-\frac{d_i}{j}}. \end{aligned}$$

The above sum converges if $d_i > j$ for all $j = 1, \dots, n_i + 1$, so it converges if $d_i > n_i + 1$. Thus, the original quantity (2.7) converges if $d_i > n_i + 1$ for all $i = 1, \dots, k$.

This heuristic therefore predicts $N(H) = O(H^{m-1})$. Combining this with the estimate $N_{\text{tot}}(H) \sim (2H)^m$ gives Conjecture 2.4(i).

2.2 Motivation for Conjecture 2.4(ii)

We may restrict ourselves to only consider polynomials f for which $f = 0$ defines a smooth geometrically integral hypersurface X in \mathcal{P} (see also [25, Remark 2.1]). We use a well known conjecture of Colliot-Thélène (see, e.g., [13]):

Conjecture 2.9 (Colliot-Thélène) Let X be a smooth proper geometrically integral variety over a number field. If X is (geometrically) rationally connected, then the Brauer-Manin obstruction to the Hasse principle for X is the only obstruction.

In particular, we show in Proposition 2.10 that if X is a multidegree (d_1, \dots, d_k) -hypersurface in $\mathcal{P} = \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k}$ with $d_i < n_i + 1$ for some i , then X is rationally connected. We will show in Proposition 2.11 below that the Brauer-Manin obstruction is vacuous for smooth complete intersections (of dimension at least 3) in \mathcal{P} over number fields. Thus, conditional on Conjecture 2.9, we see that the Hasse principle is satisfied for these X of dimension ≥ 3 .

In almost all cases with $d_i < n_i + 1$ for some i where X has dimension 1 or 2, either some $d_i = 1$ (and so X has a rational point) or X is a quadric hypersurface (and so is known to satisfy the Hasse principle). One nontrivial case is handled in the proof of [25, Proposition 3.4], which shows, under Conjecture 2.9, that the Hasse principle holds for a density 1 set of cubic surfaces in \mathbb{P}^2 . The last remaining nontrivial case is that of bidegree $(2, d_2)$ -surfaces in $\mathbb{P}^2 \times \mathbb{P}^1$ with $d_2 \geq 2$. Here, it is possible for the Hasse principle to not be satisfied (e.g., the famous example of Iskovskih [21] where $d_2 = 4$), but as we explain in Lemma 2.13 below, Conjecture 2.9 still implies that, for any fixed $d_2 \geq 2$, the Hasse principle holds for 100% of these hypersurfaces. The argument, which relies on the generic member of this family having no Brauer-Manin obstruction by [20, Theorem 2.6], is analogous to that for cubic surfaces.

Proposition 2.10 *Fix an integer $k \geq 1$ and positive integers d_i, n_i for $i \in \{1, \dots, k\}$. Let X be a smooth integral multidegree (d_1, \dots, d_k) -hypersurface in $\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k}$ over \mathbb{C} . Then X is rationally connected if and only if $d_i < n_i + 1$ for some i .*

Proof One direction is easy: if $d_i > n_i$ for all i , then $H^0(X, \omega_X)$ is nonzero, hence X is not rationally connected. For the reverse direction, without loss of generality, suppose $d_1 < n_1 + 1$. Then consider the projection $\pi : X \rightarrow \mathbb{P}^{n_2} \times \dots \times \mathbb{P}^{n_k}$ (note that this codomain is a point if $k = 1$). The fibers of this projection π are degree d_1 hypersurfaces in \mathbb{P}^{n_1} ; the general ones are smooth and, by assumption, Fano and thus rationally connected [12, 22].

We apply a theorem of Graber–Harris–Starr [18], which states that if $X \rightarrow Y$ is a dominant morphism of complex varieties where the general fiber and Y are both rationally connected, then X is rationally connected as well. Here the projection map π is surjective by a dimension count, and the codomain $\mathbb{P}^{n_2} \times \dots \times \mathbb{P}^{n_k}$ is rationally connected, so X itself is rationally connected. \square

In the following proposition, the notation $\text{Br } Y$ refers to the cohomological Brauer group $H^2_{\text{ét}}(Y, \mathbb{G}_m)$ for a scheme Y .

Proposition 2.11 *Let K be a number field. Let X be a smooth complete intersection in a product $\mathcal{P} = \mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$ satisfying $\dim X \geq 3$. Then the natural map $\text{Br } K \rightarrow \text{Br } X$ is an isomorphism, hence the Brauer-Manin obstruction for X is vacuous.*

Proof We need only slightly modify the proof in [25, Appendix A and B] for the same result where X is a smooth complete intersection in a single projective space \mathbb{P}^n (see also [27, Proposition 2.6] for a smooth complete intersection in the product of two projective spaces). We summarize the argument here. We use the low degree exact sequence from the Leray spectral sequence

$$0 \rightarrow \text{Pic } X \rightarrow (\text{Pic } X_{\bar{K}})^{G_K} \rightarrow \text{Br } K \rightarrow \ker(\text{Br } X \rightarrow \text{Br } X_{\bar{K}}) \rightarrow H^1(K, \text{Pic } X_{\bar{K}}) \quad (2.12)$$

where $G_K = \text{Gal}(\bar{K}/K)$. We need to check that $\text{Pic } X \rightarrow (\text{Pic } X_{\bar{K}})^{G_K}$ is an isomorphism, $H^1(K, \text{Pic } X_{\bar{K}}) = 0$, and $\text{Br } X_{\bar{K}} = 0$.

The restriction map $\text{Pic } \mathcal{P}_{\bar{K}} \rightarrow \text{Pic } X_{\bar{K}}$ is an isomorphism since X is a smooth complete intersection of dimension at least 3 [19, Corollary 3.3]. Then since $\text{Pic } \mathcal{P} \rightarrow \text{Pic } \mathcal{P}_{\bar{K}}$ is an isomorphism, we find that the injections

$$\text{Pic } X \hookrightarrow (\text{Pic } X_{\bar{K}})^{G_K} \hookrightarrow \text{Pic } X_{\bar{K}}$$

are isomorphisms and $H^1(K, \text{Pic } X_{\bar{K}}) = 0$.

To show that $\text{Br } X_{\bar{K}} = 0$ here, we use the Kummer sequence to obtain, for any prime ℓ ,

$$\begin{array}{ccccccc} 0 & \longrightarrow & (\text{Pic } \mathcal{P}_{\bar{K}})/\ell & \longrightarrow & H^2(\mathcal{P}_{\bar{K}}, \mathbb{Z}/\ell\mathbb{Z}) & & \\ & & \downarrow & & \downarrow \psi & & \\ 0 & \longrightarrow & (\text{Pic } X_{\bar{K}})/\ell & \longrightarrow & H^2(X_{\bar{K}}, \mathbb{Z}/\ell\mathbb{Z}) & \longrightarrow & (\text{Br } X_{\bar{K}})[\ell] \longrightarrow 0. \end{array}$$

The top horizontal injection is in fact an isomorphism, since both groups are rank k over $\mathbb{Z}/\ell\mathbb{Z}$ (using the Künneth formula to compute $H^2(\mathcal{P}, \mathbb{Z}/\ell\mathbb{Z})$). By a version of Weak Lefschetz (see, e.g., [25, Corollary B.5] with $V = \mathcal{P}$), the vertical map ψ is an isomorphism, so we have $(\text{Br } X_{\bar{K}})[\ell] = 0$ for all ℓ . Since $\text{Br } X_{\bar{K}}$ is torsion, it is in fact 0.

Thus, from (2.12), the map $\text{Br } K \rightarrow \text{Br } X$ is an isomorphism, so no elements of $\text{Br } X$ obstruct rational points on X . \square

Lemma 2.13 *Assume Conjecture 2.9. Fix an integer $d \geq 2$ and consider the family of bidegree $(2, d)$ hypersurfaces in $\mathbb{P}^2 \times \mathbb{P}^1$. Then 100% of everywhere locally soluble such hypersurfaces satisfy the Hasse principle.*

Proof These surfaces S are conic bundles over \mathbb{P}^1 , say given by equations of the form $f(x_1, x_2, x_3, y_1, y_2) = 0$, where x_1, x_2, x_3 and y_1, y_2 are coordinates for \mathbb{P}^2 and \mathbb{P}^1 , respectively. We may also represent f as a symmetric 3×3 matrix $M(y_1, y_2)$ of degree d forms in y_1, y_2 , i.e., such that $f(x, y) = \mathbf{x}^T M \mathbf{x}$. Let Z be the singular locus in \mathbb{P}^1 of the fibration $S \rightarrow \mathbb{P}^1$; explicitly, Z is defined by the determinant of the matrix $M(y_1, y_2)$, which has degree $3d$ in y_1, y_2 , and degree 3 in the coefficients $a_1, \dots, a_{6(d+1)}$ of f .

For the generic member of this family over $\mathbb{Q}(a_1, \dots, a_{6(d+1)})$, we claim that Z is irreducible. This follows, e.g., from specialization, since for a prime p , the determinant of the

3×3 matrix

$$\begin{pmatrix} 0 & y_2^d & y_1^d \\ y_2^d & y_1^d & 0 \\ y_1^d & 0 & py_2^d \end{pmatrix}$$

is irreducible by Eisenstein's criterion. Thus, the singular locus is irreducible for the generic surface over $\mathbb{Q}(a_1, \dots, a_{6(d+1)})$, and by Hilbert irreducibility, the same is true for a density 1 set of such surfaces over \mathbb{Q} . Finally, if Z is irreducible, Theorem 2.6 of [20] immediately implies that there is no Brauer-Manin obstruction for the surface S , so by Conjecture 2.9, the Hasse principle holds. \square

3 Counting polynomials over finite fields

Our strategy for proving Theorems 1.4 and 1.5 can be viewed as an extension of Hensel's lemma. If the reduction of the $(2, 2)$ -form mod p has a smooth \mathbb{F}_p -point, then Hensel's lemma implies that this can be lifted to a \mathbb{Q}_p -point; conversely, if the reduction of the $(2, 2)$ -form has no \mathbb{F}_p -points, then clearly it has no \mathbb{Q}_p -points. The p -adic solubility of many $(2, 2)$ -forms can be determined in this way, so it is crucial to understand $(2, 2)$ -forms over finite fields. We work over a finite field \mathbb{F}_q , where q is a power of a prime p , for this section, since it is no extra work to do so, though we will only need these results over \mathbb{F}_p for the later sections.

3.1 Preliminaries

Many $(2, 2)$ -forms modulo p contain binary quadratic forms as one of its factors, so we collect some relevant calculations here. These are easy to check.

Lemma 3.1 *Of the q^2 monic quadratic polynomials $f \in \mathbb{F}_q[X]$,*

- $q(q-1)/2$ have two distinct roots in \mathbb{F}_q ;
- $(q^2-q)/2$ have conjugate roots in \mathbb{F}_{q^2} ;
- q have a double root in \mathbb{F}_q .

Lemma 3.2 *Of the q^3 binary quadratic forms $f \in \mathbb{F}_q[X, Y]$,*

- $(q-1)(q+1)q/2$ have two distinct roots in \mathbb{F}_q ;
- $(q-1)(q^2-q)/2$ have two conjugate roots in \mathbb{F}_{q^2} ;
- $(q-1)(q+1)$ have a double root in \mathbb{F}_q ;
- 1 is the zero form.

3.2 Reducible $(2, 2)$ -forms

Now we look at the bihomogeneous polynomials of bidegree (d_1, d_2) in $\mathbb{F}_q[X_0, X_1; Y_0, Y_1]$ for $0 \leq d_0, d_1 \leq 2$, starting with the irreducible (d_1, d_2) -forms with $(d_1, d_2) \neq (2, 2)$. These correspond to (d_1, d_2) -curves in $\mathbb{P}^1 \times \mathbb{P}^1$, where we assume that (X_0, X_1) corresponds to the coordinates in the first factor of \mathbb{P}^1 and (Y_0, Y_1) corresponds to the coordinates in the second \mathbb{P}^1 .

Lemma 3.3 *The number of irreducible bihomogeneous polynomials in $\mathbb{F}_q[X_0, X_1; Y_0, Y_1]$ (with bidegrees as indicated) are as follows:*

| Bidegree | Number of forms up to scaling by \mathbb{F}_q^\times |
|----------|--|
| (1,0) | $m_{10} = q + 1$ |
| (2,0) | $m_{20} = (q^2 - q)/2$ |
| (1,1) | $m_{11} = q^3 - q$ |
| (2,1) | $m_{21} = q^5 - q^3$ |

Proof. There are $q^2 - 1$ nonzero (1, 0)-forms, and so $q + 1$ up to scaling. The number of irreducible (2, 0)-forms is taken from Lemma 3.2, taking into account scaling. The coefficients of a (1, 1)-form may naturally be arranged as a 2 by 2 matrix, and the form is irreducible if and only if this matrix is nonsingular. Therefore each irreducible (1, 1)-form defines the graph of a Möbius map, and so $m_{11} = |\mathrm{PGL}_2(\mathbb{F}_q)| = q^3 - q$. Finally we compute

$$m_{21} = (q^6 - 1)/(q - 1) - m_{10}m_{11} - m_{10}m_{20} - m_{10}^2(m_{10} + 1)/2 = q^5 - q^3.$$

We now consider all the different ways in which a (2, 2)-form can factor. We use the notation $(a_1, b_1)^{e_1} \cdots (a_r, b_r)^{e_r}$ to denote the bidegrees of the irreducible factors, with multiplicity. For example, the *factorization type* $(1, 0)^2(0, 1)(0, 1)$ indicates that the (2, 2)-form factors as a product $F_{10}^2 F_{01} G_{01}$, where F_{10} , F_{01} , and G_{01} are irreducible polynomials in $\mathbb{F}_q[X_0, X_1; Y_0, Y_1]$ with bidegrees (1, 0), (0, 1), and (0, 1), respectively, and that F_{01} is not an \mathbb{F}_q^\times -multiple of G_{01} .

Lemma 3.4 *The number of reducible (2, 2)-forms over \mathbb{F}_q with each factorization type are as follows. Moreover, the curve $C \subset \mathbb{P}^1 \times \mathbb{P}^1$ defined by such a form either always has a smooth \mathbb{F}_q -point, or never has a smooth \mathbb{F}_q -point, as indicated in the right hand column.*

| Factorization type | Number of forms up to scaling by \mathbb{F}_q^\times | Smooth point? |
|--|--|---------------|
| (1,1)(1,1) | $\binom{m_{11}}{2} = (q^3 - q)(q^3 - q - 1)/2$ | yes |
| (2,1)(0,1) or (1,2)(1,0) | $2m_{21}m_{10} = 2q^3(q + 1)^2(q - 1)$ | yes |
| (1,1)(1,0)(0,1) | $m_{11}m_{10}^2 = q(q + 1)^3(q - 1)$ | yes |
| (1,0)(1,0)(0,1)(0,1) | $\binom{m_{10}}{2}^2 = q^2(q + 1)^2/4$ | yes |
| (2,0)(0,1)(0,1) or (0,2)(1,0)(1,0) | $2m_{20}\binom{m_{10}}{2} = q^2(q + 1)(q - 1)/2$ | yes |
| (2,0)(0,2) | $m_{20}^2 = q^2(q - 1)^2/4$ | no |
| (1, 0)^2(0, 1)(0, 1) or (0, 1)^2(1, 0)(1, 0) | $2m_{10}\binom{m_{10}}{2} = q(q + 1)^2$ | yes |
| (2, 0)(0, 1)^2 or (0, 2)(1, 0)^2 | $2m_{20}m_{10} = q(q + 1)(q - 1)$ | no |
| (1, 1)^2 | $m_{11} = q(q + 1)(q - 1)$ | no |
| (1, 0)^2(0, 1)^2 | $m_{10}^2 = (q + 1)^2$ | no |

Proof The counts all follow from Lemma 3.3 as indicated. In each of the cases listed as having smooth \mathbb{F}_q -points, there is an irreducible factor of multiplicity 1 with bidegree (1, 0), (0, 1), (1, 1), (2, 1) or (1, 2). Each such factor defines a smooth curve of genus 0 which, by projection to one of the factors, is isomorphic to \mathbb{P}^1 . Moreover a case-by-case analysis shows that this curve meets the other components of C in at most 2 points. Since $\#\mathbb{P}^1(\mathbb{F}_q) = q + 1 > 2$ this shows that C has a smooth \mathbb{F}_q -point. In the remaining cases each irreducible factor is either repeated or has bidegree (2, 0) or (0, 2). So in these cases there are no smooth \mathbb{F}_q -points. \square

3.3 Irreducible (2, 2)-forms

We now consider the irreducible (2, 2)-forms over \mathbb{F}_q . We distinguish between those that are absolutely irreducible (i.e., do not factor over \mathbb{F}_q) and those that factor over \mathbb{F}_{q^2} as the product of a bidegree (1, 1)-form and its conjugate. In the latter case, we say the form has factorization type $(1, 1)\overline{(1, 1)}$.

Lemma 3.5 *Let $C \subset \mathbb{P}^1 \times \mathbb{P}^1$ be a curve defined by an absolutely irreducible (2, 2)-form $F \in \mathbb{F}_q[X_0, X_1; Y_0, Y_1]$. Then C has a smooth \mathbb{F}_q -point.*

Proof If C is smooth then it has genus 1, and the lemma follows by the Hasse-Weil bound. If C is singular, then it has geometric genus 0. The normalization is a smooth genus 0 curve, in fact \mathbb{P}^1 itself (e.g., by the Hasse-Weil bound), and thus has $(q + 1)$ \mathbb{F}_q -points. Since the preimage of the singular point is at most length 2, the curve C must have at least one smooth \mathbb{F}_q -point. \square

Remark 3.6 The numbers of irreducible (2, 2)-forms are as follows:

| Factorization type | Number of forms up to scaling by \mathbb{F}_q^\times |
|-------------------------------------|--|
| smooth | $q^4(q + 1)^2(q - 1)^2$ |
| absolutely irreducible yet singular | $q^3(q + 1)^2(q - 1)^2$ |
| $(1, 1)\overline{(1, 1)}$ | $(q^3 - q)(q^3 + q - 1)/2$ |

In the case $(1, 1)\overline{(1, 1)}$ the $(1, 1)$ -forms are irreducible over \mathbb{F}_{q^2} , since the original (2, 2)-form was irreducible over \mathbb{F}_q . Each therefore defines the graph of a Möbius map, and so the count in the last row is $(\# \text{PGL}_2(\mathbb{F}_{q^2}) - \# \text{PGL}_2(\mathbb{F}_q))/2$. We omit the details of the other two counts since these are not needed for the proof of Theorem 1.4. However, as a check on our calculations, we note that these counts, together with those in Lemma 3.4, do indeed add up to $(q^9 - 1)/(q - 1)$.

If F has factorization type $(1, 1)\overline{(1, 1)}$ then C is geometrically the union of two rational curves. We subdivide into cases according as these meet in

- (i) a pair of points defined over \mathbb{F}_q ,
- (ii) a pair of conjugate points defined over \mathbb{F}_{q^2} ,
- (iii) a single point defined over \mathbb{F}_q .

Lemma 3.7 *The number of (2, 2)-forms over \mathbb{F}_q (up to scaling by \mathbb{F}_q^\times) in cases (i), (ii) and (iii) above are, respectively,*

$$\begin{aligned} n_{11} &= q^3(q + 1)^2(q - 1)/4, \\ n_{12} &= q^2(q + 1)(q - 1)^2(q - 2)/4, \\ n_{13} &= q(q + 1)^2(q - 1)^2/2. \end{aligned}$$

Proof (i) The singular points can be any pair of points in $\mathbb{P}^1(\mathbb{F}_q) \times \mathbb{P}^1(\mathbb{F}_q)$ that remain distinct under both projection maps. This last condition comes from the fact, noted in Remark 3.6, that each (1, 1)-form defines the graph of a Möbius map. We make a change of coordinates to move the singular points to $((0 : 1), (0 : 1))$ and $((1 : 0), (1 : 0))$. By hypothesis F factors as the product of two (1, 1)-forms

over \mathbb{F}_{q^2} , and these are now linear combinations of X_0Y_1 and X_1Y_0 . Therefore $F = f(X_0Y_1, X_1Y_0)$ for some irreducible binary quadratic form f . We compute n_{11} as the product of the $(q^2 - q)/2$ choices for f (up to scaling), and the $q^2(q + 1)^2/2$ choices for the (unordered) pair of singular points.

- (ii) We write $\mathbb{F}_{q^2} = \mathbb{F}_q(\alpha)$ where $\alpha^2 + r\alpha + s = 0$ for some $r, s \in \mathbb{F}_q$. As noted in (i), the singular points remain distinct under both projection maps. After a change of coordinates, defined over \mathbb{F}_q , we may therefore assume that the singular points are $((\alpha : 1), (\alpha : 1))$ and its Galois conjugate. Then $F = f(X_0Y_0 + rX_0Y_1 + sX_1Y_1, X_0Y_1 - X_1Y_0)$ for some irreducible binary quadratic form f . There are $(q^2 - q)/2$ choices for f (up to scaling), but one of these gives $F = (X_0^2 + rX_0X_1 + sX_1^2)(Y_0^2 + rY_0Y_1 + sY_1^2)$. Therefore n_{12} is the product of $(q + 1)(q - 2)/2$ and the $(q^2 - q)^2/2$ choices for the singular points.
- (iii) We make a change of coordinates to move the singular point to $((0 : 1), (0 : 1))$. Then F is the product of $\alpha X_0Y_0 + \beta X_0Y_1 + \gamma X_1Y_0$, and its Galois conjugate, for some $\alpha, \beta, \gamma \in \mathbb{F}_{q^2}$ with $\beta, \gamma \neq 0$. If β and γ are a basis for \mathbb{F}_{q^2} as an \mathbb{F}_q -vector space, then by substitutions of the form $X_1 \leftarrow X_1 + \lambda X_0$ and $Y_1 \leftarrow Y_1 + \mu Y_0$ with $\lambda, \mu \in \mathbb{F}_q$ we may reduce to the case $\alpha = 0$. But then the two components also meet at $((1 : 0), (1 : 0))$, which cannot happen in case (iii). Therefore $F = f(X_0Y_0, X_0Y_1 + cX_1Y_0)$ for some irreducible binary quadratic form f and constant $c \in \mathbb{F}_q^\times$. We compute n_{13} as the product of the $(q^2 - q)/2$ choices for f (up to scaling), the $q - 1$ choices for c , and the $(q + 1)^2$ choices for the singular point.

As a final check, we note that $n_{11} + n_{12} + n_{13} = (q^3 - q)(q^3 + q - 1)/2$. \square

4 Local solubility for bidegree (2, 2)-forms

Fix a prime p , and consider the space of all (2, 2)-forms $F \in \mathbb{Z}_p[X_0, X_1; Y_0, Y_1]$ with its natural product Haar measure when viewed as a copy of \mathbb{Z}_p^9 . In this section, we determine the density of \mathbb{Q}_p -soluble forms in this space.

In order to determine the solubility of a given form F , it will often suffice to look at its reduction mod p , denoted $\bar{F} \in \mathbb{F}_p[X_0, X_1; Y_0, Y_1]$, and look for a smooth \mathbb{F}_p -point on the curve \bar{C} defined by \bar{F} , so that we may apply Hensel's lemma. As seen in the tables of Sect. 3 (where we now take $q = p$), it is easy to see that most of the factorization types for \bar{F} have a smooth \mathbb{F}_p -point on \bar{C} . According to the results of Sect. 3, only five cases require further consideration, which we analyze in Sect. 4.3.

4.1 Preliminaries

Let $\nu(a) = \nu_p(a)$ denote the p -adic valuation of $a \in \mathbb{Q}_p$.

We will often keep track of the valuations of the coefficients of the (2, 2)-form (1.3) as a 3×3 table:

$$\begin{array}{ccc} \nu(a_{00}) & \nu(a_{01}) & \nu(a_{02}) \\ \nu(a_{10}) & \nu(a_{11}) & \nu(a_{12}) \\ \nu(a_{20}) & \nu(a_{21}) & \nu(a_{22}), \end{array} \tag{4.1}$$

where each entry is the valuation of the coefficient of the corresponding monomial term in

$$\begin{array}{ccc} X_0^2Y_0^2 & X_0^2Y_0Y_1 & X_0^2Y_1^2 \\ X_0X_1Y_0^2 & X_0X_1Y_0Y_1 & X_0X_1Y_1^2 \\ X_1^2Y_0^2 & X_1^2Y_0Y_1 & X_1^2Y_1^2. \end{array} \tag{4.2}$$

4.2 A useful lemma

Before analyzing the cases where Hensel's lemma does not directly apply, we note that the following lemma will be used several times in the next section, and is, in some sense, typical of the arguments we use. In fact, we use it once in the analysis of Case 3 and twice in the analysis of Case 5 (in Lemmas 4.11 and 4.15).

Lemma 4.3 *Let $a_{00}, a_{01}, a_{02}, a_{10}, a_{11}, a_{20}$ be any fixed elements of \mathbb{Z}_p satisfying*

$$\nu(a_{00}) \geq 2, \quad \nu(a_{01}) \geq 2, \quad \nu(a_{02}) = 1,$$

$$\nu(a_{10}) \geq 1, \quad \nu(a_{11}) \geq 1,$$

$$\nu(a_{20}) = 0.$$

Let

$$\mathcal{S} := \left\{ \sum_{i,j=0}^2 a_{ij} X_0^{2-i} X_1^i Y_0^{2-j} Y_1^j : a_{12} \in p\mathbb{Z}_p, a_{21} \in \mathbb{Z}_p, a_{22} \in \mathbb{Z}_p \right\}.$$

Then the proportion of the polynomials in \mathcal{S} that have \mathbb{Q}_p -solutions for which $p \nmid X_1 Y_1$ is $1/2$.

Proof Let $\sigma(a_{00}, a_{01}, a_{02}, a_{10}, a_{11}, a_{20})$ be the desired probability (that a polynomial in \mathcal{S} has a \mathbb{Q}_p -solution with $p \nmid X_1 Y_1$), and let $\tau(a_{00}, a_{01}, a_{02}, a_{10}, a_{11}, a_{20})$ be the corresponding probability when \mathcal{S} is replaced by its subset

$$\mathcal{T} := \left\{ \sum_{i,j=0}^2 a_{ij} X_0^{2-i} X_1^i Y_0^{2-j} Y_1^j : a_{12}, a_{21}, a_{22} \in p\mathbb{Z}_p \right\}.$$

If $F \in \mathcal{S}$, then $F(0, 1; Y_0, 1)$ reduces mod p to a quadratic polynomial in Y_0 , i.e., the coefficient of Y_0^2 is nonzero mod p . Furthermore, all such quadratic polynomials occur with equal probability. By Lemma 3.1, this quadratic splits into distinct factors over \mathbb{F}_p with probability $\frac{1}{2}(1 - \frac{1}{p})$. In this case, the point $((0 : 1), (\alpha : 1))$, where α is one of the roots of the quadratic, is a smooth \mathbb{F}_p -point, so by Hensel's lemma, the curve defined by $F = 0$ has a \mathbb{Q}_p -point of the form $((0 : 1), (\tilde{\alpha} : 1))$ for some lift $\tilde{\alpha} \in \mathbb{Z}_p$. If the quadratic is instead irreducible, as happens with probability $\frac{1}{2}(1 - \frac{1}{p})$, then there are no \mathbb{Q}_p -points with $p \nmid X_1$. It remains to consider the case $F \equiv a_{20} X_1^2 (Y_0 - c Y_1)^2 \pmod{p}$, for some $0 \leq c \leq p - 1$. Transforming F by the substitution $Y_0 \leftarrow Y_0 + c Y_1$ we find

$$\sigma \begin{pmatrix} a_{00} & a_{01} & a_{02} \\ a_{10} & a_{11} & \\ a_{20} & & \end{pmatrix} = \frac{1}{2} \left(1 - \frac{1}{p} \right) + \frac{1}{p^2} \sum_{c=0}^{p-1} \tau \begin{pmatrix} a_{00} & 2ca_{00} + a_{01} & c^2 a_{00} + ca_{01} + a_{02} \\ a_{10} & 2ca_{10} + a_{11} & \\ a_{20} & & \end{pmatrix}. \quad (4.4)$$

We note in particular that the arguments of τ satisfy the conditions in the statement of the lemma.

If $F \in \mathcal{T}$ then $F \equiv a_{20} X_1^2 Y_0^2 \pmod{p}$. For a solution with $p \nmid X_1$ we need $p \mid Y_0$. This suggests making the substitution $Y_0 \leftarrow p Y_0$. Dividing through by p , and then swapping the X 's and Y 's we find

$$\tau \begin{pmatrix} a_{00} & a_{01} & a_{02} \\ a_{10} & a_{11} & \\ a_{20} & & \end{pmatrix} = \sigma \begin{pmatrix} pa_{00} & pa_{10} & pa_{20} \\ a_{01} & a_{11} & \\ p^{-1}a_{02} & & \end{pmatrix}. \quad (4.5)$$

Using (4.4) and (4.5) to solve for σ and τ , we find that $\sigma = \tau = 1/2$. \square

4.3 The cases without smooth \mathbb{F}_p -points

As remarked before, these are the factorization types of the $(2, 2)$ -form F over \mathbb{F}_p which do not immediately yield a smooth \mathbb{F}_p -point in the reduction of the corresponding curve.

| Case number | Factorization type | Number of forms up to scaling by \mathbb{F}_p^\times |
|-------------|--------------------------------------|--|
| 1 | $(1, 1)(1, 1)$ | $n_1 := (p^3 - p)(p^3 + p - 1)/2$ |
| 2 | $(2, 0)(0, 2)$ | $n_2 := p^2(p - 1)^2/4$ |
| 3 | $(2, 0)(0, 1)^2$ or $(0, 2)(1, 0)^2$ | $n_3 := p(p + 1)(p - 1)$ |
| 4 | $(1, 1)^2$ | $n_4 := p(p + 1)(p - 1)$ |
| 5 | $(1, 0)^2(0, 1)^2$ | $n_5 := (p + 1)^2$ |

Let $n_0 = (p^9 - 1)/(p - 1) - (n_1 + n_2 + n_3 + n_4 + n_5)$ be the number of forms lying in none of the 5 cases, and let ξ_i be the probability of solubility in case i . Then the overall probability of solubility is

$$\rho = \frac{n_0 + n_1\xi_1 + n_2\xi_2 + n_3\xi_3 + n_4\xi_4 + n_5\xi_5}{(p^9 - 1)/(p - 1)}.$$

In this section we compute ξ_1, \dots, ξ_5 and hence obtain the final answer stated in Theorem 1.4. In the context of computing ξ_5 , it is helpful to make the following definition for ξ'_i , and to compute the ξ'_i alongside the ξ_i :

Definition 4.6 For $1 \leq i \leq 4$, we let ξ'_i be the probability of solubility given the following conditions: we are in case i , the point $((0 : 1), (0 : 1))$ is a singular point on the reduction mod p , and $\nu(a_{22}) \geq 2$. We write ξ'_{1j} for $1 \leq j \leq 3$ for the same probability in cases 1(i), 1(ii), and 1(iii), as defined below. We define ξ'_5 in the same way, except that we require that the singular point $((0 : 1), (0 : 1))$ is *not* the point where the two lines meet.

We compute the values of ξ_i and ξ'_i in the next sections.

4.3.1 Case 1

In this case the reduction of our curve mod p is geometrically the union of two rational curves. We subdivide into the cases (i), (ii), (iii) as defined immediately before Lemma 3.7, and note that $n_1 = n_{11} + n_{12} + n_{13}$. Writing $\xi_{11}, \xi_{12}, \xi_{13}$ for the probabilities of solubility in cases 1(i), 1(ii) and 1(iii), respectively, we have

$$\xi_1 = (n_{11}\xi_{11} + n_{12}\xi_{12} + n_{13}\xi_{13})/n_1.$$

Case 1(i)

In this case, the two components meet at a pair of points defined over \mathbb{F}_p . We begin by computing ξ_{11} . As in the proof of Lemma 3.7(i) we may assume that $\bar{F} = f(X_0Y_1, X_1Y_0)$ for some irreducible binary quadratic form f . The only \mathbb{F}_p -points on the reduction are the singular points $((0 : 1), (0 : 1))$ and $((1 : 0), (1 : 0))$. We must decide if they lift to \mathbb{Q}_p -points. Let α be the probability that the singular point $((0 : 1), (0 : 1))$ lifts.

Since $((0 : 1), (0 : 1))$ is a singular point on the curve defined by \bar{F} , we deduce that all the valuations $\nu(a_{12}), \nu(a_{21}), \nu(a_{22})$ are ≥ 1 . If $\nu(a_{22}) = 1$, then the singular point $((0 : 1), (0 : 1))$ does not lift. Otherwise (with probability $1/p$), we have $\nu(a_{22}) \geq 2$. Then

the valuations of the coefficients of F satisfy

$$\begin{array}{lll} \geq 1 & \geq 1 & = 0 \\ \geq 1 & \geq 0 & \geq 1 \\ = 0 & \geq 1 & \geq 2 \end{array}$$

where the equalities follow from f being irreducible. Making the substitutions $X_0 \leftarrow pX_0$, $Y_0 \leftarrow pY_0$ and dividing through by p^2 , we obtain a $(2, 2)$ -form $G(X_0, X_1, Y_0, Y_1)$ whose coefficients b_{ij} have valuations satisfying

$$\begin{array}{lll} \geq 3 & \geq 2 & = 0 \\ \geq 2 & \geq 0 & \geq 0 \\ = 0 & \geq 0 & \geq 0. \end{array}$$

We now investigate whether $\bar{G} \in \mathbb{F}_p[X_0, X_1; Y_0, Y_1]$ is absolutely irreducible. Define a ternary quadratic form $Q(X_0, Y_0, Z_0)$ by $\bar{G}(X_0, 1, Y_0, 1) = Q(X_0, Y_0, 1)$, so that the zero-set of Q in $\mathbb{A}_{X_0, Y_0}^2 \subset \mathbb{P}^2$ coincides with the zero-set of \bar{G} in $\mathbb{A}_{X_0}^1 \times \mathbb{A}_{Y_0}^1 \subset \mathbb{P}^1 \times \mathbb{P}^1$. Then the curve defined by Q (and thus, the curve defined by \bar{G}) is geometrically irreducible if and only if the discriminant of Q is nonzero, equivalently

$$b_{22} \operatorname{disc}(f) - f(b_{21}, -b_{12}) \not\equiv 0 \pmod{p}.$$

(This argument still works in characteristic 2 provided that the formula for the discriminant of a ternary quadratic form is scaled by appropriate powers of 2.)

If the curve defined by \bar{G} is geometrically irreducible, then the argument in Lemma 3.5 shows that it has a smooth \mathbb{F}_p -point. Otherwise (with probability $1/p$), the reduction mod p is geometrically the union of two rational curves meeting at $((1 : 0), (1 : 0))$ and an \mathbb{F}_p -point of the form $((\lambda : 1), (\mu : 1))$. The two rational curves are not defined over \mathbb{F}_p , since the binary quadratic form f is irreducible. We make the substitutions $X_0 \leftarrow X_0 + \lambda X_1$ and $Y_0 \leftarrow Y_0 + \mu Y_1$ to move the second point of intersection to $((0 : 1), (0 : 1))$, and start over again considering whether this singular point lifts. The probability that it lifts is again α . We thus obtain the recursive formula

$$\alpha = \frac{1}{p} \left(\left(1 - \frac{1}{p} \right) + \frac{1}{p} \alpha \right),$$

and so $\alpha = 1/(p+1)$.

We are interested in the probability that at least one of the singular points lifts. Since these events depend on different coefficients of the $(2, 2)$ -form they are independent. Therefore

$$\xi_{11} = 1 - \left(1 - \frac{1}{p+1} \right)^2 = \frac{2p+1}{(p+1)^2}.$$

A small modification of this argument (as required by Definition 4.6) gives

$$\xi'_{11} = 1 - \left(1 - \frac{1}{p+1} \right) \left(1 - \frac{p}{p+1} \right) = \frac{p^2 + p + 1}{(p+1)^2}.$$

Case 1(ii)

The two components meet at a pair of conjugate points defined over \mathbb{F}_{p^2} . There are no \mathbb{F}_p -points on the reduction. Therefore $\xi_{12} = 0$ and ξ'_{12} is not defined.

Case 1(iii)

The two components meet at a single point defined over \mathbb{F}_p . As in the proof of Lemma 3.7(iii) we may assume that $\bar{F} = f(X_0 Y_0, X_0 Y_1 + X_1 Y_0)$ for some irreducible binary quadratic form f . The only \mathbb{F}_p -point on the reduction is the singular point $((0 : 1), (0 : 1))$. If this lifts to a \mathbb{Q}_p -point then we must have $\nu(a_{22}) \geq 2$. Since this is exactly the condition in Definition 4.6 it follows that $\xi_{13} = (1/p)\xi'_{13}$.

We show in Sect. 5, using results from [3], that

$$\xi_{13} = \frac{2p^{10} + 3p^9 - p^5 + 2p^4 - 2p^2 - 3p - 1}{2(p+1)^2(p^9 - 1)}.$$

As noted in the last paragraph, we have $\xi'_{13} = p\xi_{13}$.

4.3.2 Case 2

This is the case $(2, 0)(0, 2)$. There are no \mathbb{F}_p -points on the reduction. Therefore $\xi_2 = 0$ and ξ'_2 is not defined.

4.3.3 Case 3

This is the case $(2, 0)(0, 1)^2$ or $(0, 2)(1, 0)^2$. We may assume without loss of generality that $\bar{F} = f(X_0, X_1)Y_0^2$ for some irreducible binary quadratic form f . The coefficients satisfy

$$\begin{array}{ccc} = 0 & \geq 1 & \geq 1 \\ \geq 0 & \geq 1 & \geq 1 \\ = 0 & \geq 1 & \geq 1 \end{array}$$

where the equalities follow from f being irreducible. Making the substitution $Y_0 \leftarrow pY_0$ and dividing through by p gives

$$\begin{array}{ccc} = 1 & \geq 1 & \geq 0 \\ \geq 1 & \geq 1 & \geq 0 \\ = 1 & \geq 1 & \geq 0 \end{array}$$

The reduction mod p is now $g(X_0, X_1)Y_1^2$ for some binary quadratic form g . If g is irreducible, splits, or has repeated roots, then the probability of solubility is 0, 1, or $1/2$, respectively. In the last of these cases, we are using Lemma 4.3: more specifically, we assume the double root is at $(X_0 : X_1) = (0 : 1)$, make the substitution $X_0 \leftarrow pX_0$, divide through by p , and then apply the lemma. Note that the lemma applies as $\nu(a_{20}) = 0$, and there are no solutions with $p \mid X_1 Y_1$ in view of the substitutions we made to reach this situation.

If g is identically zero, then we divide through by p , to obtain a $(2, 2)$ -form H satisfying the *line condition*, by which we mean that $H(X_0, X_1, 1, 0) \pmod{p}$ is an irreducible binary quadratic form. Writing δ_{line} for the probability of solubility in this case, we have

$$\xi_3 = \frac{1}{p^3} \left(\frac{p^3 - p}{2} \cdot 1 + \frac{p(p-1)^2}{2} \cdot 0 + (p^2 - 1) \cdot \frac{1}{2} + \delta_{\text{line}} \right) \quad (4.7)$$

and

$$\xi'_3 = \frac{1}{p^2} \left(p(p-1) \cdot 1 + (p-1) \cdot \frac{1}{2} + \delta_{\text{line}} \right). \quad (4.8)$$

Lemma 4.9 *There are (up to scaling by \mathbb{F}_p^\times) exactly $p^7(p-1)/2$ forms over \mathbb{F}_p satisfying the line condition. The numbers of these in Cases 1 to 3 are*

$$\begin{aligned} r_{11} &= p^3(p+1)(p-1)^2/4 \\ r_{12} &= p^2(p+1)(p-1)^2(p-2)/4 \\ r_{13} &= p^2(p+1)(p-1)^2/2 \\ r_2 &= p^2(p-1)^2/4 \\ r_3 &= p^2(p-1)/2 \end{aligned}$$

There are none in Cases 4 and 5.

Proof It is easy to check that forms in Cases 1(ii) and 2 always satisfy the line condition, and those in Cases 4 and 5 never satisfy the line condition. By double counting pairs consisting of (2, 2)-forms and (0, 1)-forms (both up to scalars) that meet in a pair of conjugate points over \mathbb{F}_{p^2} , we find that $(p+1)r_{11} = (p-1)r_{13}$ and $(p+1)r_2 = pr_{13}$. In Case 3 we must count the forms $f(X_0, X_1)g(Y_0, Y_1)^2$ where f is an irreducible binary quadratic form, and g is a linear form with $g(1, 0) \neq 0$. We find that r_3 is the product of the $(p^2-p)/2$ choices for f and the p choices for g . \square

Let $r_0 = p^7(p-1)/2 - (r_{11} + r_{12} + r_{13} + r_2 + r_3)$ be the number of forms satisfying the line condition not in Cases 1 to 3. Then

$$\delta_{\text{line}} = \frac{r_0 + r_{11}\xi_{11} + r_{12}\xi_{12} + r_{13}\xi_{13} + r_2\xi_2 + r_3\xi_3}{p^7(p-1)/2} \quad (4.10)$$

Using the values of $\xi_{11}, \xi_{12}, \xi_{13}$ and ξ_2 already computed, we can now solve (4.7) and (4.10) for ξ_3 and δ_{line} . We then use (4.8) to compute ξ'_3 . We find that

$$\xi_3 = \frac{p^{10} + 2p^9 + p^6 - 2p^5 + 2p^3 + p^2 - 3p - 2}{2(p+1)(p^9-1)}$$

and

$$\xi'_3 = \frac{2p^{10} + p^9 + p^7 - 2p^6 + 2p^4 + p^3 - 2p^2 - 2p - 1}{2(p+1)(p^9-1)}.$$

4.3.4 Case 4

This is the case $(1, 1)^2$. By a change of coordinates we may assume

$$F \equiv (X_0 Y_1 - X_1 Y_0)^2 \pmod{p}.$$

We show in Sect. 5, using results from [3], that

$$\xi_4 = \frac{5p^{10} + 8p^9 + p^8 - p^7 + 2p^6 - 3p^5 + 4p^3 - 10p - 6}{8(p+1)(p^9-1)},$$

and

$$\xi'_4 = \frac{4p^{10} + 3p^9 - p^7 + 2p^6 - 2p^5 + 2p^3 - p^2 - 5p - 2}{4(p+1)(p^9-1)}.$$

4.3.5 Case 5

This is the case $(1, 0)^2(0, 1)^2$. By a change of coordinates we may assume

$$F \equiv X_0^2 Y_0^2 \pmod{p}.$$

The coefficients of F have valuations satisfying

$$\begin{array}{lll} = 0 & \geq 1 & \geq 1 \\ \geq 1 & \geq 1 & \geq 1 \\ \geq 1 & \geq 1 & \geq 1. \end{array}$$

Let Q and Q' be the binary quadratic forms over \mathbb{F}_p determined by the last row and column, i.e.,

$$\begin{aligned} Q(Y_0, Y_1) &= \frac{1}{p}F(0, 1; Y_0, Y_1) \pmod{p} \\ Q'(X_0, X_1) &= \frac{1}{p}F(X_0, X_1; 0, 1) \pmod{p} \end{aligned}$$

Note that these forms have the same last coefficient $c \in \mathbb{F}_p$ as they share one entry in the coefficient matrix corresponding to $X_1^2 Y_1^2$. Writing ξ_{51} and ξ_{52} for the probabilities of solubility in the cases $c \neq 0$ and $c = 0$, respectively, we have

$$\xi_5 = (1 - 1/p)\xi_{51} + (1/p)\xi_{52}.$$

Lemma 4.11 *We have $\xi_{51} = 3/4$.*

Proof The coefficients satisfy

$$\begin{array}{lll} = 0 & \geq 1 & \geq 1 \\ \geq 1 & \geq 1 & \geq 1 \\ \geq 1 & \geq 1 & = 1. \end{array}$$

The reduction mod p is the union of two double lines, meeting at a single point. Any \mathbb{Q}_p -point has $p \mid X_0$ or $p \mid Y_0$, but not both since $v(a_{22}) = 1$. In other words, any \mathbb{Q}_p -point must reduce to lie on exactly one of the lines.

To investigate whether there are solutions with $p \mid X_0$ we make the substitution $X_0 \leftarrow pX_0$ and divide by p to get

$$\begin{array}{lll} = 1 & \geq 2 & \geq 2 \\ \geq 1 & \geq 1 & \geq 1 \\ \geq 0 & \geq 0 & = 0. \end{array}$$

We then apply Lemma 4.3 (with $Y_0 \leftrightarrow Y_1$). The probability of a solution with $p \mid X_0$ and the probability of a solution with $p \mid Y_0$ are each $1/2$. The lemma also implies that these two events are independent of each other, so the probability of insolubility of these polynomials is $1/4$. Hence, the probability of solubility is $3/4$. \square

Definition 4.12 Let δ_1 and δ_2 be the probabilities of solubility in the cases

$$\begin{array}{llllll} \geq 1 & \geq 1 & \geq 0 & \geq 1 & \geq 1 & = 0 \\ \geq 1 & \geq 0 & \geq 0 & \geq 1 & \geq 0 & \geq 0 \\ = 0 & \geq 0 & \geq 0 & = 0 & \geq 0 & \geq 0 \end{array}$$

(The subscript is the number of equalities in the matrix.) Let δ_1^* and δ_2^* be the probabilities when we change the top left ≥ 1 to $= 1$. Let ε_1 and ε_2 be the probabilities when we change the top left ≥ 1 to ≥ 2 .

Clearly we have

$$\begin{aligned} \delta_1 &= (1 - 1/p)\delta_1^* + (1/p)\varepsilon_1 & (4.13) \\ \delta_2 &= (1 - 1/p)\delta_2^* + (1/p)\varepsilon_2 \end{aligned}$$

Lemma 4.14 *We have*

$$\xi_{52} = \left(1 - \frac{1}{p^2}\right) + \frac{1}{p^2} \left(\frac{1}{p} \left(1 - \frac{1}{p}\right)^2 \delta_2^* + 2 \frac{1}{p} \left(1 - \frac{1}{p}\right) \delta_1^* + \left(\frac{1}{p}\right)^2 \varepsilon_1 \right)$$

Proof If at least one of the forms Q and Q' has distinct roots in \mathbb{F}_p then the $(2, 2)$ -form is soluble over \mathbb{Q}_p . This happens with probability $1 - 1/p^2$. Otherwise (with probability $1/p^2$) the coefficients satisfy

$$\begin{array}{lll} = 0 & \geq 1 & \geq 1 \\ \geq 1 & \geq 1 & \geq 2 \\ \geq 1 & \geq 2 & \geq 2 \end{array}$$

(The bottom right ≥ 2 comes from the assumption that $c = 0$, and the two adjacent ≥ 2 entries arise from assuming that neither Q nor Q' has distinct roots in \mathbb{F}_p .) We split into 3 cases:

- (i) Suppose $\nu(a_{02}) = \nu(a_{20}) = 1$. If $\nu(a_{22}) = 2$ then the $(2, 2)$ -form is insoluble over \mathbb{Q}_p . Otherwise, we find by substituting $X_0 \leftarrow pX_0$, $Y_0 \leftarrow pY_0$, and dividing through by p^3 , that the probability of solubility is δ_2^* .
- (ii) Suppose $\nu(a_{02}) = 1$ and $\nu(a_{20}) \geq 2$. We find by substituting $X_0 \leftarrow pX_0$, and dividing through by p^2 , that the probability of solubility is δ_1^* . The case where $\nu(a_{02}) \geq 2$ and $\nu(a_{20}) = 1$ works in exactly the same way via the substitution $Y_0 \leftarrow pY_0$.
- (iii) Suppose $\nu(a_{02}) \geq 2$ and $\nu(a_{20}) \geq 2$. Via either of the substitutions in (ii), the probability of solubility is ε_1 .

Combining these gives the desired expression for ξ_{52} . \square

Lemma 4.15 *We have*

$$\xi'_5 = \left(1 - \frac{1}{p}\right) + \frac{1}{p} \left(1 - \frac{1}{p}\right) \frac{3}{4} + \left(\frac{1}{p}\right)^2 \left(1 - \frac{1}{p}\right) + \left(\frac{1}{p}\right)^3 \delta_1.$$

Proof According to Definition 4.6, we may suppose the coefficients of F satisfy

$$\begin{array}{lll} = 0 & \geq 1 & \geq 2 \\ \geq 1 & \geq 1 & \geq 1 \\ \geq 1 & \geq 1 & \geq 1 \end{array}$$

If $\nu(a_{12}) = 1$, then Q' has distinct roots in \mathbb{F}_p and so F is soluble over \mathbb{Q}_p . Otherwise (with probability $1/p$), we have $\nu(a_{12}) \geq 2$.

If $\nu(a_{22}) = 1$, then by an argument similar to Lemma 4.11, the probability of solubility is $3/4$. (The solutions with $p \mid X_0$ are analysed in exactly the same way as before, whereas to analyse those with $p \mid Y_0$ we substitute $Y_0 \leftarrow pY_0$ and then $X_1 \leftarrow pX_1$.)

Otherwise (with probability $1/p$), we have $\nu(a_{22}) \geq 2$. If $\nu(a_{21}) = 1$, then Q has distinct roots in \mathbb{F}_p and so F is soluble over \mathbb{Q}_p . Otherwise (with probability $1/p$), we have $\nu(a_{21}) \geq 2$. We find by substituting $Y_0 \leftarrow pY_0$ and then dividing through by p^2 that the probability of solubility is δ_1 . \square

Lemma 4.16 *There are p^5 possibilities for \bar{F} (up to scaling by \mathbb{F}_p^\times) satisfying the conditions in the definition of δ_1 . The number of these in Cases 1 to 5 are*

$$s_{11} = p^3(p-1)/2,$$

$$s_{12} = 0,$$

$$s_{13} = p(p-1)^2/2,$$

$$s_2 = 0,$$

$$s_3 = p(p-1)/2,$$

$$s_4 = p(p-1)$$

$$s_5 = p.$$

In Cases 1(i), 1(iii), and 4, these forms also satisfy the conditions in the definition of δ_2 . In Cases 3 and 5, they do not.

Proof The conditions in the definition of δ_1 are that $\bar{F} = 0$ is singular at $((1:0), (1:0))$ but does not contain the line $Y_1 = 0$. We have $(p+1)^2 s_{11} = 2n_{11}$ and $(p+1)^2 s_{13} = n_{13}$. In Cases 1(ii) and 2, there are no \mathbb{F}_p -points so $s_{12} = s_2 = 0$. In Cases 3, 4, and 5, we count the forms $X_1^2 f(Y_0, Y_1)$, $(\alpha X_1 Y_0 + \beta X_0 Y_1 + \gamma X_1 Y_1)^2$, and $X_1^2 g(Y_0, Y_1)^2$ where f is an irreducible binary quadratic form, $\alpha, \beta, \gamma \in \mathbb{F}_p^\times$ with $\alpha\beta \neq 0$, and g is a linear form with $g(1, 0) \neq 0$. Finally, it is only in Cases 3 and 5 that the reduction mod p contains the line $X_1 = 0$. \square

For the final computation of ξ_5 , let $s_0 = p^5 - (s_{11} + s_{13} + s_3 + s_4 + s_5)$ and $t_0 = p^4(p-1) - (s_{11} + s_{13} + s_4)$ so that

$$\delta_1 = \frac{s_0 + s_{11}\xi_{11} + s_{13}\xi_{13} + s_3\xi_3 + s_4\xi_4 + s_5\xi_5}{p^5}$$

$$\delta_2 = \frac{t_0 + s_{11}\xi_{11} + s_{13}\xi_{13} + s_4\xi_4}{p^4(p-1)}.$$

Replacing each ξ by ξ' (see Definition 4.6) we have

$$\varepsilon_1 = \frac{s_0 + s_{11}\xi'_{11} + s_{13}\xi'_{13} + s_3\xi'_3 + s_4\xi'_4 + s_5\xi'_5}{p^5}$$

$$\varepsilon_2 = \frac{t_0 + s_{11}\xi'_{11} + s_{13}\xi'_{13} + s_4\xi'_4}{p^4(p-1)}.$$

Putting together all the equations derived in this section, together with the previously computed ξ 's, we are now able to solve for ξ_5 . We find $\xi_5 = f(p)/g(p)$ where

$$f(p) = 6p^{18} + 8p^{17} + 2p^{16} - 8p^{15} + 16p^{14} - 12p^{13} - 4p^{12} + 3p^{11} + 9p^{10} - 35p^9 + 8p^8 - 11p^7 + 3p^6 - p^5 + 8p^4 - 6p^3 - 4p^2 + 10p + 8,$$

$$g(p) = 8(p+1)(p^9 - 1)(p^8 - 1).$$

5 Relation to binary quartics

We compute some of the probabilities required in Sect. 4 by reducing them to probabilities already computed in [3]. The basic idea is that a $(2, 2)$ -form determines a binary quartic form, by writing the $(2, 2)$ -form as a binary quadratic form in Y_0, Y_1 and taking the discriminant. However, since we also want results in the case $p = 2$, we will in fact work with generalised binary quartics, defined as follows.

Definition 5.1 A *generalised binary quartic* (G_2, G_4) is a pair of binary forms of degrees 2 and 4. A generalised binary quartic (G_2, G_4) is *soluble* over a field K if for some $X_0, X_1, Z \in K$ with $(X_0, X_1) \neq (0, 0)$ we have $Z^2 + G_2(X_0, X_1)Z = G_4(X_0, X_1)$.

We write $\mathbb{Z}_p[X_0, X_1] = \bigoplus_d \mathbb{Z}_p[X_0, X_1]_d$ and $\mathbb{Z}_p[X_0, X_1; Y_0, Y_1] = \bigoplus_{d,e} \mathbb{Z}_p[X_0, X_1; Y_0, Y_1]_{de}$ for the gradings of these rings by degree d and by bidegree (d, e) , respectively.

Lemma 5.2 Let $\ell, a \in \mathbb{F}_p$ such that $Z^2 + \ell Z - a$ is irreducible over \mathbb{F}_p , and let

$$\begin{aligned} \mathcal{S} &:= \{(G_2, G_4) \in \mathbb{Z}_p[X_0, X_1]_2 \times \mathbb{Z}_p[X_0, X_1]_4 : G_2 \equiv \ell X_0^2 \pmod{p} \text{ and } G_4 \equiv a X_0^4 \pmod{p}\}, \\ \mathcal{T} &:= \{(G_2, G_4) \in \mathbb{Z}_p[X_0, X_1]_2 \times \mathbb{Z}_p[X_0, X_1]_4 : G_2 \equiv 0 \pmod{p} \text{ and } G_4 \equiv 0 \pmod{p}\}, \end{aligned}$$

and $\mathcal{T}^* := \{(G_2, G_4) \in \mathcal{T} : G_4(0, 1) \not\equiv 0 \pmod{p^2}\}$. Then the proportions of generalised binary quartics in \mathcal{S} , \mathcal{T} and \mathcal{T}^* that are soluble over \mathbb{Q}_p are, respectively,

$$\begin{aligned} \sigma &= \frac{2p^{10} + 3p^9 - p^5 + 2p^4 - 2p^2 - 3p - 1}{2(p+1)^2(p^9 - 1)}, \\ \tau &= \frac{5p^{10} + 8p^9 + p^8 - p^7 + 2p^6 - 3p^5 + 4p^3 - 10p - 6}{8(p+1)(p^9 - 1)}, \end{aligned}$$

and

$$\tau^* = \frac{5p^{10} + 5p^9 - p^7 + 3p^6 - 4p^5 + 4p^3 - 8p - 4}{8(p+1)(p^9 - 1)}.$$

Proof These probabilities were computed in [3]. The probability σ was computed following Corollary 18, where it was denoted λ . The probabilities τ and τ^* were computed in Sects. 2.3.6 and 2.3.5, where they were denoted σ_4 and σ'_4 . \square

We define a map

$$\begin{aligned} \Phi : \quad \mathbb{Z}_p[X_0, X_1; Y_0, Y_1]_{22} &\rightarrow \mathbb{Z}_p[X_0, X_1]_2 \times \mathbb{Z}_p[X_0, X_1]_4 \\ F_0 Y_0^2 + F_1 Y_0 Y_1 + F_2 Y_1^2 &\mapsto (F_1, -F_0 F_2). \end{aligned}$$

It is easy to check that a $(2, 2)$ -form F is soluble over \mathbb{Q}_p if and only if the generalised binary quartic $\Phi(F)$ is soluble over \mathbb{Q}_p . We write Φ_p for the corresponding map on forms with coefficients in \mathbb{F}_p . The cases (i) and (ii) in the following lemma relate to Cases 1(iii) and 4 in Sect. 4.3.

Lemma 5.3 Let $\bar{F} \in \mathbb{F}_p[X_0, X_1; Y_0, Y_1]_{22}$ take one of the following forms:

- (i) $\bar{F} = f(X_0 Y_0, X_0 Y_1 + X_1 Y_0)$ where f is an irreducible binary quadratic form,
- (ii) $\bar{F} = (X_0 Y_1 - X_1 Y_0)^2$.

Then Φ restricts to a measure-preserving map

$$\begin{aligned} \{F \in \mathbb{Z}_p[X_0, X_1; Y_0, Y_1]_{22} : F \equiv \bar{F} \pmod{p}\} \\ \rightarrow \{G \in \mathbb{Z}_p[X_0, X_1]_2 \times \mathbb{Z}_p[X_0, X_1]_4 : G \equiv \Phi_p(\bar{F}) \pmod{p}\}. \end{aligned}$$

Proof The proof comes down to showing that the derivative of Φ_p at \bar{F} is a surjective linear map $\mathbb{F}_p^9 \rightarrow \mathbb{F}_p^8$. In cases (i) and (ii), this linear map is given by

$$\begin{aligned} F_0 Y_0^2 + F_1 Y_0 Y_1 + F_2 Y_1^2 &\mapsto (F_1, -f(0, 1)X_0^2 F_0 - f(X_0, X_1)F_2), \text{ and} \\ F_0 Y_0^2 + F_1 Y_0 Y_1 + F_2 Y_1^2 &\mapsto (F_1, -X_0^2 F_0 - X_1^2 F_2), \end{aligned}$$

respectively, which are both surjective. \square

Case 1(iii).

We use these lemmas to compute ξ_{13} , that is, the probability of solubility where $F \bmod p$ is of the form indicated in Lemma 5.3(i). Let f have coefficients a, b, c . Then $\Phi_p(\bar{F})$ is the generalised binary quartic with equation

$$(Z + cX_0X_1)^2 + bX_0^2(Z + cX_0X_1) + aX_0^4 = 0.$$

From this we see that $\xi_{13} = \sigma$ as defined in Lemma 5.2.

Case 4.

We use these lemmas to compute ξ_4 , that is, the probability of solubility where $F \bmod p$ is of the form indicated in Lemma 5.3(ii). Since $\Phi_p(\bar{F})$ is identically zero, we see that $\xi_4 = \tau$ as defined in Lemma 5.2. To compute ξ'_4 we must consider $(2, 2)$ -forms F that additionally satisfy $\nu(a_{22}) \geq 2$. Under the measure-preserving map in Lemma 5.3(ii) these are mapped to $\mathcal{T} \setminus \mathcal{T}^*$. Therefore $\tau = (1 - 1/p)\tau^* + (1/p)\xi'_4$ and so

$$\xi'_4 = p\tau - (p - 1)\tau^* = \frac{4p^{10} + 3p^9 - p^7 + 2p^6 - 2p^5 + 2p^3 - p^2 - 5p - 2}{4(p + 1)(p^9 - 1)}.$$

Remark 5.4 The same approach could be used to compute ξ_{11} , and indeed our answer agrees with [3, Lemma 15].

6 Connections to the Hasse principle

In Theorem 1.5, we determined that the proportion of $(2, 2)$ -forms that are everywhere locally soluble is $c \approx 0.8739$. As we explain further in Sects. 6.1 and 6.2, a heuristic similar to [9, Conjectures 6 and 7] predicts that the proportion of everywhere locally soluble $(2, 2)$ -forms that are globally soluble is $\frac{1}{4}$, i.e., in the notation of Sect. 2, that $\lim_{H \rightarrow \infty} \frac{N(H)}{N_{\text{loc}}(H)} = \frac{1}{4}$ and thus $\lim_{H \rightarrow \infty} \frac{N(H)}{N_{\text{tot}}(H)} = \frac{1}{4}c$.

In this section we report on some experiments to test this conjecture numerically. A similar study in the case of plane cubics was made in [15]. With the one exception noted below, all computations were performed using Magma [5] and the data may be found at [14].

6.1 Experiments and results

For each $H \in \{10, 30, 100, 300, 1000\}$, we chose 1000 $(2, 2)$ -forms (i.e., polynomials of the form (1.3)), with coefficients chosen uniformly at random from $[-H, H] \cap \mathbb{Z}$. The numbers of these that were soluble or everywhere locally soluble (ELS) were as follows:

The second column gives our initial estimate for the number of soluble $(2, 2)$ -forms out of the 1000. The lower bound was obtained by searching for rational solutions, with the assistance of 4-descent in Magma. The upper bound was obtained by computing the Cassels-Tate pairing on the 2-Selmer group of the Jacobian. For the improved estimates in the third column we used a range of methods, described more fully below, that are conditional on standard conjectures and sometimes were only practical for H sufficiently small.

| H | initial range | | improved range | | # ELS |
|------|---------------|-----------|----------------|-----------|-------|
| | # soluble | # soluble | # soluble | # soluble | |
| 10 | [753, 755] | | 753 | | 885 |
| 30 | [640, 652] | | 642 | | 885 |
| 100 | [536, 582] | | 549 | | 875 |
| 300 | [378, 502] | | [432, 433] | | 867 |
| 1000 | [275, 464] | | [357, 464] | | 879 |

For the first 4 experiments, we also give the breakdown of these totals by the rank of the Jacobian elliptic curve E/\mathbb{Q} . The annotations + and – in the case $H = 300$ indicate that we should add or subtract one if the remaining form whose solubility has not yet been decided turns out to be soluble.

| | rank $E(\mathbb{Q})$ | singular | 0 | 1 | 2 | 3 | 4 | 5 | 6 | Total |
|-----------|----------------------|----------|---|------|-----|------|-----|----|---|-------|
| $H = 10$ | # ELS | 5 | 0 | 86 | 344 | 313 | 116 | 21 | 0 | 885 |
| | # soluble | 5 | 0 | 16 | 285 | 310 | 116 | 21 | 0 | 753 |
| $H = 30$ | # ELS | 0 | 0 | 122 | 310 | 291 | 129 | 29 | 4 | 885 |
| | # soluble | 0 | 0 | 1 | 208 | 274 | 126 | 29 | 4 | 642 |
| $H = 100$ | # ELS | 0 | 0 | 171 | 321 | 257 | 96 | 25 | 5 | 875 |
| | # soluble | 0 | 0 | 0 | 205 | 221 | 93 | 25 | 5 | 549 |
| $H = 300$ | # ELS | 0 | 0 | 230– | 373 | 187+ | 58 | 19 | 0 | 867 |
| | # soluble | 0 | 0 | 0 | 210 | 151+ | 52 | 19 | 0 | 432+ |

Although we can see from our first table that the proportion of everywhere locally soluble forms that are globally soluble is decreasing with H , this hardly amounts to strong evidence that the limit is $1/4$. However, the prediction of $1/4$ arises since, in the limit, it is expected that (i) 50% of the Jacobians have rank 1 and 50% have rank 2 (by, e.g., the Minimalist Conjecture), and (ii) the proportions of forms that are soluble in these two cases are 0 and $1/2$, respectively (as explained below). Our second table therefore provides much stronger evidence for the conjecture, and indeed we see that the convergence in (ii) is happening much faster than that in (i).

6.2 The marked point and heuristics

As we saw in Sect. 5, if $F \in \mathbb{Z}[X_0, X_1; Y_0, Y_1]$ is a $(2, 2)$ -form, then it determines a pair of binary quartics. These binary quartics have the same discriminant, which is accordingly called the discriminant of F . We should expect a randomly chosen $(2, 2)$ -form to have nonzero discriminant (and hence define a smooth curve). This was true in all our experiments, except for 5 cases with $H = 10$, which were all in any case soluble. From now on we assume that the discriminant is nonzero, and write E for the Jacobian of $C = \{F = 0\} \subset \mathbb{P}^1 \times \mathbb{P}^1$. Since the discriminant is a degree 12 polynomial in the coefficients of F , the conductor and discriminant of E each have size about H^{12} .

There are two maps $C \rightarrow \mathbb{P}^1$, given by projection to each factor, and the difference of fibres is a nonzero marked point $P_0 \in E(\mathbb{Q})$. There is an explicit formula for P_0 (see [6, Section 6.1.2] or [17, Lemma 2.1]) in terms of the coefficients of F . As might be predicted from this formula, we found in our experiments that P_0 had canonical height at most

$\log(2H^2)$. The torsion subgroup of $E(\mathbb{Q})$ was trivial in all but 6 cases with $H = 10$, when it had order 2. In only one of these cases was P_0 a torsion point.

We should expect that for a randomly chosen $(2, 2)$ -form, the associated binary quartics should not have any rational roots (i.e., linear factors). This was true in all but 69, 9 and 1 of our examples with $H = 10, 30$ and 100 . We should also expect that $P_0 \notin 2E(\mathbb{Q})$, and this was true in all but 6 cases when $H = 10$, and 2 cases when $H = 30$. If $P_0 \notin 2E(\mathbb{Q})$ and the associated binary quartics do not have any rational roots, then if $\text{rank } E(\mathbb{Q}) = 1$, the $(2, 2)$ -form is not soluble. We thus expect that the $(2, 2)$ -form is not soluble in general if $\text{rank } E(\mathbb{Q}) = 1$.

We now explain why half of the forms with rank 2 Jacobian are expected to be soluble. For an elliptic curve E of rank 2, we want to estimate the proportion of elements in $\text{Sel}^2(E/\mathbb{Q})$ that are in the image of $E(\mathbb{Q})/2E(\mathbb{Q})$. The average size of $\text{Sel}^2(E/\mathbb{Q})$ is 6 in this family [8], but two of the 2-Selmer group elements correspond to $(2, 2)$ -forms for which one of the associated binary quartics has a rational root, which should only happen 0% of the time when we order by height. Now the size of $E(\mathbb{Q})/2E(\mathbb{Q})$ is 4 for the 100% of elliptic curves E for which there is no 2-torsion, but we also subtract the same 2 elements corresponding to the $(2, 2)$ -forms for which one of the associated binary quartics has a rational root. We thus predict that $(4 - 2)/(6 - 2) = 1/2$ of the forms are soluble.

Remark 6.1 Although one of the best methods for finding generators of large height on an elliptic curve E/\mathbb{Q} is to use Heegner points, this only works for curves of rank 1. Since our elliptic curves all come with a known point of infinite order, this method was of no use to us. We instead relied almost exclusively on descent methods.

6.3 The initial estimates: computing ranks of elliptic curves

The curve defined by a $(2, 2)$ -form is isomorphic to the curve defined by either of the associated binary quartics. Our interest is therefore in deciding the solubility of the genus one curves associated to binary quartics.

Let C/\mathbb{Q} be a genus one curve defined by a binary quartic, and let E/\mathbb{Q} be its Jacobian. If C is everywhere locally soluble, then it defines a class $[C]$ in the 2-Selmer group $\text{Sel}^2(E/\mathbb{Q})$. Moreover $C(\mathbb{Q}) \neq \emptyset$ if and only if $[C] \in \text{im}(\delta)$ where δ is the connecting map in the Kummer exact sequence

$$0 \rightarrow E(\mathbb{Q})/2E(\mathbb{Q}) \xrightarrow{\delta} \text{Sel}^2(E/\mathbb{Q}) \rightarrow \text{III}(E/\mathbb{Q})[2] \rightarrow 0.$$

Given a point $P \in E(\mathbb{Q})$ the Magma function `GenusOneModel(2, P)` computes a binary quartic representing $\delta(P)$. In conjunction with the function `IsEquivalent` for testing equivalence of binary quartics, this gives a convenient way of reducing the problem of deciding whether $C(\mathbb{Q}) \neq \emptyset$ to that of finding generators for $E(\mathbb{Q})$.

An initial upper bound for the rank of $E(\mathbb{Q})$ is obtained by 2-descent, that is, by computing the 2-Selmer group $\text{Sel}^2(E/\mathbb{Q})$. This upper bound can sometimes be improved by computing the Cassels-Tate pairing. Let S_n be the image of the natural map $\text{Sel}^{2^n}(E/\mathbb{Q}) \rightarrow \text{Sel}^2(E/\mathbb{Q})$. If $\xi, \eta \in S_n$, say with $\xi' \mapsto \xi$ and $\eta' \mapsto \eta$, then there is an alternating pairing

$$\langle , \rangle_n : S_n \times S_n \rightarrow \mathbb{F}_2; \quad (\xi, \eta) \mapsto \langle \xi', \eta \rangle_{\text{CT}} = \langle \xi, \eta' \rangle_{\text{CT}} \quad (6.2)$$

whose kernel is S_{n+1} . We note the inclusions of \mathbb{F}_2 -vector spaces

$$\text{im}(\delta) \subset \dots \subset S_3 \subset S_2 \subset S_1 = \text{Sel}^2(E/\mathbb{Q}).$$

The function `CasselsStatePairing` in Magma, written by S. Donnelly, computes the pairings

$$\langle , \rangle_{\text{CT}} : \text{Sel}^2(E/\mathbb{Q}) \times \text{Sel}^2(E/\mathbb{Q}) \rightarrow \mathbb{F}_2, \quad (6.3)$$

and

$$\langle , \rangle_{\text{CT}} : \text{Sel}^2(E/\mathbb{Q}) \times \text{Sel}^4(E/\mathbb{Q}) \rightarrow \mathbb{F}_2, \quad (6.4)$$

taking as input either a pair of binary quartics, or a binary quartic and a quadric intersection. (A variant of his method for computing (6.3) is described in [16].) We may thus compute the pairing (6.2) for $n = 1$ and $n = 2$.

A lower bound for the rank of $E(\mathbb{Q})$ may be obtained by searching for points either directly on E , or better on one of its 4-coverings as computed using `FourDescent` in Magma. In this way we obtained generators for a subgroup $\Gamma \subset E(\mathbb{Q})$ of known points. In all cases where it is possible that $\text{rank } \Gamma < \text{rank } E(\mathbb{Q})$ we searched up to height 10^{10} on the 4-coverings.

Our initial (unconditional) estimate on the number of $(2, 2)$ -forms that are soluble was obtained as follows. First, if the curve C defined by our $(2, 2)$ -form is not everywhere locally soluble, then it is obviously not soluble. Otherwise C (or more precisely one of the associated binary quartics) determines a class $[C] \in \text{Sel}^2(E/\mathbb{Q})$. If $[C] \in \delta(\Gamma)$ then we know that $C(\mathbb{Q}) \neq \emptyset$, and indeed from the generators for Γ we may compute an explicit solution. Otherwise we look for $[D] \in \text{Sel}^2(E/\mathbb{Q})$ with $\langle [C], [D] \rangle_{\text{CT}} \neq 0$. If we succeed in finding a binary quartic D with these properties, then it is a witness to the fact that $C(\mathbb{Q}) = \emptyset$.

6.4 The improved estimates

Both the improved estimates, and the second table (giving the breakdown by rank of the Jacobian) are conditional on the following two standard conjectures.

- Parity conjecture: This is the parity part of the Birch–Swinnerton-Dyer conjecture, i.e., the Mordell-Weil rank of an elliptic curve E/\mathbb{Q} is even or odd according as its root number $w(E/\mathbb{Q})$ is $+1$ or -1 .
- Generalised Riemann Hypothesis (GRH) : This is needed for the class number calculations for 2-descent (but would be easy to remove for small H) and for the computation of analytic rank bounds.

For ease of exposition, we will assume (as is the case in all examples of interest) that $E(\mathbb{Q})$ has trivial torsion subgroup. We write $r_{2^n} = \dim_{\mathbb{F}_2} S_n$ for the upper bound on the rank of $E(\mathbb{Q})$ obtained by 2^n -descent. Thus we have

$$\text{rank } \Gamma \leq \text{rank } E(\mathbb{Q}) \leq \dots \leq r_8 \leq r_4 \leq r_2.$$

It is a theorem, originally due to Monsky [23], that $w(E/\mathbb{Q}) = (-1)^{r_2}$.

We improve our lower bounds on the number of forms that are soluble by using the parity conjecture. Indeed, if $\text{rank } \Gamma = r_{2^n} - 1$ for some n , then we may conclude by the parity conjecture that $\text{im}(\delta) = S_n$. We mainly used this idea with $n = 1$, when the conclusion is that binary quartics with Jacobian E satisfy the Hasse principle, but also used it with $n = 2$ in four examples with $H = 1000$.

In the two examples where we used the parity conjecture in the case $H = 30$, we were also able to find the missing generators using `EightDescent` in Magma [29]. In

particular on the curve $\{F = 0\} \subset \mathbb{P}^1 \times \mathbb{P}^1$ where

$$\begin{aligned} F = & 27X_0^2Y_0^2 - X_0X_1Y_0^2 + 17X_1^2Y_0^2 - 27X_0^2Y_0Y_1 + 15X_0X_1Y_0Y_1 + 9X_1^2Y_0Y_1 \\ & - 25X_0^2Y_1^2 - 12X_0X_1Y_1^2 - 13X_1^2Y_1^2, \end{aligned}$$

we found the solution

$$\begin{aligned} X_0 &= 5998800628516423107297133082973646629266881508307007941326966876023, \\ X_1 &= 342294900150114936634770190317380320064921533929615189995360150770683, \\ Y_0 &= 246468494594162038245191010835877699291643209107952263886240062422805, \\ Y_1 &= -206172926328604047309514129427033995615708844556361901784128916991449. \end{aligned}$$

This maps to a point on the Jacobian of canonical height $\widehat{h} \approx 644.736$, which is well beyond the range that could be found by 4-descent. Unfortunately it was not practical to run `EightDescent` in the experiments with $H = 100, 300$ and 1000 , and so our improved lower bounds in those cases remain conditional on the parity conjecture.

The main method we used to improve the upper bounds on the number of forms that are soluble was to compute the Cassels-Tate pairing (6.4). In the experiments with $H = 10, 30, 100$ and 300 , we were left with 0, 1, 2 and 5 examples where $r_2 = r_4 = r_8 = 3$, yet (despite searching on all 4-coverings up to height bound 10^{10}) we could only find one generator. The elliptic curves in question are recorded in the following table.

| | |
|--|----------------|
| $y^2 = x^3 - 385216x - 118546643$ | $\Delta = 2.0$ |
| $y^2 + xy = x^3 - x^2 - 21940631x - 10062163381$ | $\Delta = 2.6$ |
| $y^2 + xy = x^3 - x^2 - 130106786x - 418444299752$ | $\Delta = 2.5$ |
| $y^2 = x^3 + x^2 - 674939767x + 9768411280745$ | $\Delta = 3.5$ |
| $y^2 + xy + y = x^3 + 1365438724x + 1088450102306$ | $\Delta = 3.0$ |
| $y^2 + xy = x^3 + x^2 + 13646956x + 36868880351052$ | $\Delta = 3.6$ |
| $y^2 + y = x^3 + 463718380x - 1653282652263$ | $\Delta = 3.9$ |
| $y^2 + xy = x^3 + x^2 - 6811523942x + 180704627470189$ | |

In all but the last of these examples, we were able to prove that the rank is 1 by using Sage [26] to compute an upper bound on the analytic rank. The parameter Δ we used for this calculation (see [11]) is recorded in the right hand column. In the last example we obtained no rank bound better than 3, despite taking $\Delta = 4.0$.

Unfortunately it was not practical to compute the pairing (6.4) in the experiment with $H = 1000$. So we are left with a large number of unresolved cases. Writing $t = \text{rank } \Gamma$ for the number of generators known, there were 90 cases with $(r_2, r_4, t) = (3, 3, 1)$, 14 cases with $(r_2, r_4, t) = (4, 4, 2)$, and one each with $(r_2, r_4, t) = (4, 4, 1), (5, 3, 1)$ and $(5, 5, 3)$.

Acknowledgements

Much of this work was completed during a workshop organized by Alexander Betts, Tim Dokchitser, Vladimir Dokchitser and Celine Maistret, a trimester organized by David Harari, Emmanuel Peyre, and Alexei Skorobogatov, and a workshop organized by Michael Stoll; we thank the organizers as well as Baskerville Hall, Institut Henri Poincaré, and Franken-Akademie Schloss Schney, respectively, for their hospitality during those periods. We also thank Bhargav Bhatt, John Cremona, David Harari, Max Lieblich, Daniel Loughran, Bjorn Poonen, and the anonymous referee for helpful

conversations and comments. WH was partially supported by NSF grants DMS-1701437 and DMS-1844763 and the Sloan Foundation. JP was partially supported by NSF grant DMS-1902199.

Author details

¹DPMMS, Centre for Mathematical Sciences, University of Cambridge, Wilberforce Road, Cambridge CB3 0WB, UK,

²Department of Mathematics, University of Michigan, Ann Arbor, MI 48109, USA, ³Department of Mathematics, The Ohio State University, Columbus, OH 43210, USA.

Received: 26 June 2020 Accepted: 12 October 2020 Published online: 5 January 2021

References

- Bright, M.J., Browning, T.D., Loughran, D.: Failures of weak approximation in families. *Compos. Math.* **152**(7), 1435–1475 (2016). MR3530447
- Bhargava, M., Cremona, J.E., Fisher, T.A.: The proportion of plane cubic curves over \mathbb{Q} that everywhere locally have a point. *Int. J. Number Theory* **12**(4), 1077–1092 (2016)
- Bhargava, M., Cremona, J.E., Fisher, T.A.: The proportion of genus one curves over \mathbb{Q} defined by a binary quartic that everywhere locally have a point, (2020), to appear in *International J. Number Theory*, arXiv:2004.12085
- Bhargava, M., Cremona, J.E., Fisher, T.A., Jones, N.G., Keating, J.P.: What is the probability that a random integral quadratic form in n variables has an integral zero? *Int. Math. Res. Not. IMRN* **12**, 3828–3848 (2016). MR3544620
- Bosma, W., Cannon, J., Playoust, C.: The Magma algebra system. I. The user language, 1997, pp. 235–265. Computational algebra and number theory (London, 1993). MR1484478
- Bhargava, M., Ho, W.: Coregular spaces and genus one curves. *Camb. J. Math.* **4**(1), 1–119 (2016). MR3472915
- Bhargava, M., Ho, W.: The Hasse principle for some genus one curves in $\mathbb{P}^1 \times \mathbb{P}^1$, (2020). In preparation
- Bhargava, M., Ho, W.: On average sizes of Selmer groups and ranks in families of elliptic curves having marked points, (2020). Preprint
- Bhargava, M.: A positive proportion of plane cubics fail the Hasse principle, (2014). arXiv:1402.1131
- Browning, T., Le Boudec, P., Sawin, W.: The Hasse principle for random Fano hypersurfaces, (2020). arXiv:2006.02356
- Bober, J.W.: Conditionally bounding analytic ranks of elliptic curves, ANTS X—Proceedings of the Tenth Algorithmic Number Theory Symposium, pp. 135–144 (2013). MR3207411
- Campana, F.: Connexité rationnelle des variétés de Fano. *Ann. Sci. École Norm. Sup. (4)* **25**(5), 539–545 (1992). MR1191735
- Collot-Thélène, J.-L.: Points rationnels sur les fibrations, Higher dimensional varieties and rational points (Budapest, 2001), pp. 171–221 (2003). MR2011747
- Fisher, T.A., Ho, W., Park, J.: Data for “Everywhere local solubility for hypersurfaces in products of projective spaces”, (2019). Available at <https://www.dpmms.cam.ac.uk/~taf1000/papers/probs22.html>
- Fisher, T.A.: The proportion of plane cubic curves with a rational point, (2015). Oberwolfach report, Explicit methods in number theory
- Fisher, T.A.: On binary quartics and the Cassels-Tate pairing, (2016). <http://www.dpmms.cam.ac.uk/~taf1000/papers/bq-ctp.html>
- Fisher, T.A., Radičević, L.: Some minimisation algorithms in arithmetic invariant theory. *J. Théor. Nombres Bordeaux* **30**(3), 801–828 (2018). MR3938627
- Graber, T., Harris, J., Starr, J.: Families of rationally connected varieties. *J. Amer. Math. Soc.* **16**(1), 57–67 (2003). MR1937199
- Hartshorne, R.: Ample subvarieties of algebraic varieties. Lecture Notes in Mathematics, vol. 156. Springer, Berlin (1970)
- Iskovskih, V.A.: Rational surfaces with a pencil of rational curves. *Mat. Sb. (N.S.)* **74**(116), 608–638 (1967). MR0220734
- Iskovskih, V.A.: A counterexample to the Hasse principle for systems of two quadratic forms in five variables. *Mat. Zametki* **10**, 253–257 (1971). MR0286743
- Kollar, J., Miyaoka, Y., Mori, S.: Rational connectedness and boundedness of Fano manifolds. *J. Differ. Geom.* **36**(3), 765–779 (1992). MR1189503
- Monsky, P.: Generalizing the Birch-Stephens theorem. I. Modular curves. *Math. Z.* **221**(3), 415–420 (1996). MR1381589
- Poonen, B., Stoll, M.: The Cassels-Tate pairing on polarized abelian varieties. *Ann. Math. (2)* **150**(3), 1109–1149 (1999). MR1740984
- Poonen, B., Voloch, J.F.: Random Diophantine equations, Arithmetic of higher-dimensional algebraic varieties (Palo Alto, CA, 2002), (2004), pp. 175–184. With appendices by Jean-Louis Colliot-Thélène and Nicholas M. Katz. MR2029869
- The Sage Developers, Sagemath, the Sage Mathematics Software System (Version 8.7), (2019). <https://www.sagemath.org>
- Schindler, D.: Manin’s conjecture for certain bimprojective hypersurfaces. *J. Reine Angew. Math.* **714**, 209–250 (2016). MR3491888
- Serre, J.-P.: Spécialisation des éléments de $\text{Br}_2(\mathbf{Q}(T_1, \dots, T_n))$. *C. R. Acad. Sci. Paris Sér. I Math.* **311**(7), 397–402 (1990). MR1075658
- Stamminger, S.: Explicit 8-descent on elliptic curves, PhD thesis, Bremen (2005)

Publisher’s Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.