

REVIEW

Nutrient and stress tolerance traits linked to fungal responses to global change: Four case studies

Kathleen K. Treseder¹*, Charlotte J. Alster¹, Linh Anh Cat², Morgan E. Gorris³, Alexander L. Kuhn¹, Karissa G. Lovero¹, Frank Hagedorn⁴, Jennifer F. Kerekes⁵, Theresa A. McHugh⁶, and Emily F. Solly⁷

In this case study analysis, we identified fungal traits that were associated with the responses of taxa to 4 global change factors: elevated CO₂, warming and drying, increased precipitation, and nitrogen (N) enrichment. We developed a trait-based framework predicting that as global change increases limitation of a given nutrient, fungal taxa with traits that target that nutrient will represent a larger proportion of the community (and vice versa). In addition, we expected that warming and drying and N enrichment would generate environmental stress for fungi and may select for stress tolerance traits. We tested the framework by analyzing fungal community data from previously published field manipulations and linking taxa to functional gene traits from the MycoCosm Fungal Portal. Altogether, fungal genera tended to respond similarly to 3 elements of global change: increased precipitation, N enrichment, and warming and drying. The genera that proliferated under these changes also tended to possess functional genes for stress tolerance, which suggests that these global changes—even increases in precipitation—could have caused environmental stress that selected for certain taxa. In addition, these genera did not exhibit a strong capacity for C breakdown or P acquisition, so soil C turnover may slow down or remain unchanged following shifts in fungal community composition under global change. Since we did not find strong evidence that changes in nutrient limitation select for taxa with traits that target the more limiting nutrient, we revised our trait-based framework. The new framework sorts fungal taxa into Stress Tolerating versus C and P Targeting groups, with the global change elements of increased precipitation, warming and drying, and N enrichment selecting for the stress tolerators.

Keywords: Elevated CO₂, Precipitation, Warming, Drying, Nitrogen enrichment, Nitrogen fertilization, Limiting nutrients, Synthesis, Fungal traits, Growth form, Extracellular enzymes, Functional genes, Tissue phosphorus concentration

Introduction

Recent decades have witnessed a discovery stage in fungal ecology, powered in part by advances in high throughput DNA sequencing (Lindahl et al., 2013; Jansson and

Hofmockel, 2020). We have learned that fungal communities frequently shift under human-induced global change factors such as elevated CO2, warming, drought, increased precipitation, and nitrogen (N) enrichment (reviewed in Allison and Martiny, 2008; Castro et al., 2010; Pickles et al., 2012; Treseder et al., 2012; Mohan et al., 2014; Classen et al., 2015; Jansson and Hofmockel, 2020). These changes have now been well-documented in field studies (e.g., Hagedorn et al., 2013; Kerekes et al., 2013; McHugh and Schwartz, 2016; Treseder et al., 2016). Nevertheless, we have a poor understanding of why specific fungal taxa respond the way they do (Martiny et al., 2015). For example, why does the abundance of a given taxon increase under elevated CO2 (or another global change factor) while another decreases? If we understand underlying mechanisms for these shifts, we may better predict fungal contributions to carbon and nutrient cycling under global change.

Global changes could select for (or against) fungal taxa with certain physiological or genetic traits (e.g., Romero-

Department of Ecology and Evolutionary Biology, University of California Irvine, Irvine, CA, USA

²Cabrillo National Monument, National Park Service, San Diego, CA, USA

³Los Alamos National Lab, Los Alamos, NM, USA

⁴Swiss Federal Institute for Forest, Snow and Landscape Research, Birmensdorf, Switzerland

⁵Center for Science and Math Education, San Francisco State University, San Francisco, CA, USA

⁶Department of Life and Environmental Sciences, Carroll College, Helena, MT, USA

⁷Department of Environmental Systems Science, ETH Zürich, Zürich, Switzerland

^{*}Corresponding author: Email: treseder@uci.edu

	C limitation	P limitation	N limitation	Stress
Elevated CO ₂	_	+	+	-
Warming & drying	+	+	+	+
Increased precipitation	_	-	-	-
N enrichment	+	+	-	+
Mechanisms:	C acquisition	P acquisition	N acquisition	Stress tolerance
Functional genes:	Glycoside hydrolases	Phosphatase P transporters	Chitinase N transporters Lignin peroxidase	β 1,3 glucan synthase Cold-induced RNA helicase Heat shock proteins Trehalase

Figure 1. Hypothesized effects of global changes on nutrient limitation of fungal growth and activity. Mechanisms that might underlie fungal responses to nutrient limitation or stress, and related functional genes, are also included. See Table S1 for a complete listing of functional genes. DOI: https://doi.org/10.1525/elementa.2020.00144.f1

Olivares et al., 2021). Here, we focus on traits that enable fungi to acquire and use carbon (C), N, and phosphorus (P) from the environment. Fungi require these nutrients to grow and reproduce (Sinsabaugh et al., 2008; Sinsabaugh et al., 2009; Strickland and Rousk, 2010; Mouginot et al., 2014). Yet, global changes can alter the relative availability of C, N, and P (Vitousek, 2004; Chapin et al., 2011). For instance, elevated CO₂ may alleviate C limitation for fungi by boosting availability of plant C and exacerbate N or P limitation (Chagnon et al., 2013; Johnson et al., 2013). Accordingly, fungi with N- or P-acquisition traits may proliferate under elevated CO2, while those that invest in C-acquisition traits may decline or remain unchanged (Velicer and Lenski, 1999; Treseder, 2005; Malik et al., 2019). Other global change factors could also alter the relative abundance of C versus N or P, each in their own way.

Global change can also cause environmental stress for fungi, apart from changes in limiting nutrients (Mohan et al., 2014; Lenoir et al., 2016; Morrison et al., 2018; Malik et al., 2019; Garcia et al., 2020). For example, warming (and the drying often associated with it) in arid ecosystems could generate drought stress (Schimel et al., 2007; Manzoni et al., 2012; Homyak et al., 2018). In addition, N enrichment can lead to soil acidification, changes in osmotic potential, or the formation of toxic N-containing compounds such as melanoidins (Broadbent, 1965; Baath et al., 1981; Vitousek et al., 1997; Treseder, 2008; Liu and Greaver, 2010; Lu et al., 2011). Stress tolerance traits may be advantageous under these circumstances.

We created a framework to predict responses of fungal communities to elevated CO₂, warming, increased precipitation, and N enrichment and then tested the framework on 1 case study for each global change element (Table S1; Hagedorn et al., 2013; Kerekes et al., 2013; McHugh and Schwartz, 2016; Treseder et al., 2016). We acquired data on nutrient and stress tolerance—related traits of the fungal taxa by assessing functional genes in published fungal genomes housed in the U.S. Department of Energy's MycoCosm Fungal Portal (Grigoriev et al., 2014). Then, we used phylogenetic independent contrasts to test for relationships between the global change responses and those functional genes. If fungal taxa respond to global changes depending on nutrient or stress tolerance—related traits, these responses could help us identify broadly applicable mechanisms underlying community shifts. This approach could allow us to use a trait-based framework to predict fungal community shifts in response to future environmental conditions.

Trait-based framework

We based our approach on the idea that fungal taxa should invest more toward acquiring nutrients that limit their growth and reproduction (Read, 1991). Liebig's Law of the Minimum suggests that an organism's activity is constrained by the scarcest nutrient that it requires, even if other required nutrients are readily available (Liebig, 1843). Consequently, fungal taxa that preferentially target the limiting nutrients may displace groups that focus on other, nonlimiting nutrients. The displacement could be particularly strong if evolutionary or physiological tradeoffs apply-if resources invested in acquiring a nonlimiting nutrient cannot be invested in acquiring the limiting nutrient (Velicer and Lenski, 1999; Allison et al., 2010b; Treseder et al., 2011; Malik et al., 2019). Altogether, this framework suggests that the relative availability of nutrients selects for or against individual fungal taxa.

Global changes may alter the relative availability of C, N, or P (Figure 1). Elevated CO₂ can increase C fluxes from plants to litter and soil, potentially decreasing N and P availability under certain vegetation types (Staddon et al., 1999; Norby et al., 2001; de Graaff et al., 2006; Hungate et al., 2009; Dieleman et al., 2010; Norby and Zak, 2011). Accordingly, fungi that decompose litter and soil organic matter may become more N- and P-limited and less C-limited. In addition, mycorrhizal fungi may have an advantage under elevated CO2 since C allocation from host plants to the fungi can increase (Diaz, 1996; Compant et al., 2010). Nitrogen fertilization or anthropogenic N deposition can enrich N supplies to fungi, alleviating N limitation and exacerbating C- and P-limitation (Treseder and Allen, 2002; Johnson et al., 2013; Treseder et al., 2018). Increased precipitation may improve the availability of all 3 nutrients because the water can dissolve and carry the nutrients to the fungi and stimulate decomposer activity (Schimel and Bennett, 2004; Chapin et al., 2011; Manzoni et al., 2012; Manzoni et al., 2014). Warming can stimulate microbial activity when water is not limiting but can lead to higher water evaporation rates and soil drying in some ecosystems (Rebetez and Dobbertin, 2004; Allison and Treseder, 2008; Allison and Treseder, 2011). Consequently, warming could enhance C and N availability in moist ecosystems (Solly et al., 2017a; Solly et al., 2017b), while it could reduce availability of C, N, and P if warming induces moisture limitation (Allison and Treseder, 2011; Parker and Schimel, 2011). We expect that fungal taxa will respond to these shifts in nutrient limitations depending on their nutrient-related traits.

Stress tolerance traits could also influence the distribution of fungal taxa (Malik et al., 2019). Certain global change factors tend to reduce fungal abundance, potentially because the changes create environmental stresses for fungi. Fungal biomass generally declines as N enrichment increases (Treseder, 2008). Warming experiments have varied widely in the extent to which fungal abundance responds, and reductions are more common when warming is accompanied by drying (Chen et al., 2015; Romero-Olivares et al., 2017b; Gao and Yan, 2019). Elevated CO2 tends to increase mycorrhizal fungal biomass (Treseder, 2004; Dong et al., 2018), so it may alleviate environmental stress for those fungi. Altogether, we predicted that N enrichment and warming (if accompanied by drying) may select for taxa with stress tolerance traits, while elevated CO2 and increased precipitation may select against them (Figure 1).

Methods

Selected traits

Response traits

The extent to which a given fungal group becomes more or less abundant in response to global change can be considered a trait (sensu Lavorel and Garnier, 2002). We quantified response traits as the increase or decrease in relative abundance of a given fungal group in response to a global change factor. For example, if a given fungal group represented 30% of the fungal community under ambient CO₂ controls, but rose to 40% under experimental CO₂ enrichment, its elevated CO₂ response was +10%.

In this analysis, positive response traits will indicate proliferation under global change, while negative response traits indicate a reduction.

Functional gene traits

To test our trait-based framework, we focused on functional genes for C, N, and P acquisition and environmental stress tolerance. Certainly, possession of a gene does not confirm that the gene is expressed or translated (Pradet-Balade et al., 2001; Wilmes and Bond, 2006; Myrold et al., 2014). Instead, we consider gene possession an indication that the fungal group has the genetic capacity for that trait. Moreover, gene frequencies or copy numbers can influence the extent to which a trait is expressed (Zhou et al., 2011).

We selected functional genes encoding extracellular enzymes that break down organically bound C for C acquisition (Lynd et al., 2002; Edwards et al., 2008; Martínez et al., 2009), chitin or lignin complexes for N acquisition (Bending and Read, 1997; Caimey and Burke, 1998; Talbot et al., 2012), and C-bound phosphate for P acquisition (Sinsabaugh, 1994). In addition, amino acid permeases, ammonium transporters, nitrate transporters, and phosphate transporters can improve uptake rates of N or P into hyphae (Versaw and Metzenberg, 1995; Nehls et al., 1999; Mitsuzawa, 2006; Slot et al., 2007). For stress tolerance, β-1,3 glucan synthase allows fungi to incorporate this carbohydrate into their cell walls to prevent desiccation and freezing damage (Bowman and Free, 2006; Latgé, 2007). Furthermore, cold-induced RNA helicase and heat shock proteins help improve cold and heat tolerance, respectively (Schade et al., 2004; Owttrim, 2006; Tiwari et al., 2015). Moreover, trehalase produces trehalose, a compatible solute that protects fungi from water loss, freezing damage, and heat shock (Wiemken, 1990; Estruch, 2000). We determined functional gene frequencies (# copies per 10,000 genes) encoding each of these traits by examining 692 published whole fungal genomes representing 111 genera (Table 1; Figure S1). We used search terms (Table 1) in MycoCosm to identify existing gene annotations (last accessed February 24, 2021; Grigoriev et al., 2014).

Study selection

For each of the 4 global change factors, we selected 1 previously published study based on predetermined selection criteria (Table S1). First, we used field-based studies, so that findings could be applicable to an ecosystem setting. Second, we chose studies in which the global change factor was experimentally manipulated in comparison to a control, so we could calculate the response trait based on relative abundance in the experimental versus control treatments. Third, the relative abundance data for each fungal taxon needed to be accessible, either via published databases or directly from authors. Fourth, we focused on studies that characterized fungal communities at the species to genus level to allow greatest flexibility in taxonomic rank analysis. This latter criterion restricted us to studies that sequenced the highly variable ITS1 or ITS2 region. In contrast, we placed

Table 1. Functional genes and their search terms in MycoCosm (Grigoriev et al., 2014). DOI: https://doi.org/10.1525/elementa.2020.00144.t1

Functional Gene	Glycoside Hydrolase (GH) Activity	Search Term
Amino acid permease	n.a.	IPR004762
Ammonium transporter	n.a.	IPR001905
β-1,3 glucan synthase	n.a.	GO:0000148
Chitinase (GH 18)	Chitinase	IPR001223
Crystalline cellulase (AA 9, formerly GH 61)	n.a.	IPR005103
Fungal lignin peroxidase	n.a.	IPR001621
GH 1	β-glucosidase and β-galactosidase	IPR001360
GH 7	Cleave β -1,4 glycosidic bonds in cellulose/ β -1,4-glucans	IPR001722
GH 9	Cellulases	IPR001701
GH 12	Endoglucanase	IPR002594
GH 13	Substrates containing α-glucoside linkages	GH13
GH 15	Hydrolyze the non-reducing end residues of α -glucosides	IPR011613
GH 28	Polygalacturonase	IPR00743
GH 31	α-glucosidase	IPR000322
GH 32	Invertase	IPR001362
GH 76	Endo-acting α-mannanases	IPR005198
GH 81	Endo-β-1,3-glucanase	IPR005200
GH 85	Endo-β-N-acetylglucosaminidase	IPR005201
GH 92	Exo-acting α-mannosidases	IPR012939
Heat shock protein	n.a.	IPR002068
Nitrate transporter	n.a.	IPR004737
Phosphatase	n.a.	IPR000560
Phosphate transporter	n.a.	HMMPFAM: PF01384
Cold-induced RNA helicase	n.a.	IPR014014
Trehalase	n.a.	GO:0005991

n.a. = not applicable.

no restrictions on ecosystem type, geography, or sequencing platform (e.g., Illumina or 454). If samples were taken at multiple time points within the same plots, we used the last samplings to allow the longest exposure to the global change treatment. Upon study selection, we contacted the lead authors of the studies and invited them to collaborate on the analysis.

Fungal taxa identification

For most of the studies, we used taxonomic names provided by the investigators unless those taxonomies had been revised after publication. In those cases, we updated taxonomic names to match current names from Index Fungorum (2021, accessed February 24, 2021). For the N enrichment study, Kerekes et al. (2013) had not identified taxa to genus or species. For that study, we used UNITE (Nilsson et al., 2018) to match representative sequences to these taxonomic ranks where possible. Across

studies, between 18.2% and 71.2% of taxa were identified to at least the genus level (Table S1).

Experimental designs

Increased precipitation

We calculated responses to increased precipitation using fungal community composition from samples taken at the final sampling (i.e., 30 days after onset of treatments) of the McHugh and Schwartz (2016) study. They collected soil from the top 5 cm of 4 pairs of control and watered plots.

Nitrogen enrichment

Kerekes et al. (2013) characterized fungal taxa in the litter layer in 4 control and four N-fertilized plots.

Warming and drying

Warming and drying responses of taxa were taken directly from supplementary information in Treseder et al. (2016). They sampled fungi in the litter horizon of 5 control and 5 warmed plots.

Elevated CO₂

We calculated elevated CO₂ responses using data supplied by Hagedom et al. (2013). They had characterized fungal community composition in 10 blocks of control and elevated CO₂ plots. Each block encompassed 4 plots: a control plot and elevated CO2 plot, each centered on a Larix decidua tree; and a control plot and elevated CO₂ plot, each centered on a Pinus uncinata tree. There were 40 plots in total. From each of these plots, they collected soil from 3 horizons: litter layer, F horizon, and H horizon, after 9 plant growing seasons of CO2 enrichment. The soils were nutrient poor and acidic Ranker or Podzols with a 10to 20-cm thick organic layer. We calculated mean relative abundances across all soil horizons and both tree species. By combining soil horizons, we obtained a more generalizable pattern of fungal responses than if we had analyzed each horizon separately. On the other hand, it limited our ability to discern horizon-specific responses.

Trait compilation

We compiled the responses of taxa to elevated CO2, warming and drying, increased precipitation, and N enrichment with functional gene frequencies (Table S2). Specifically, for each response and functional gene trait, we calculated genus-level means (e.g., average response of all taxa within Boletus). We chose to take genus-level means instead of species-level means for 2 reasons. First, many taxa were identifiable to genus but not species. Second, the majority of species with response traits were detected in just 1 experiment or did not have whole genomes available in MycoCosm. By taking genus-level means, we were able to compare traits across more taxa. We lost species-level resolution but gained statistical power. Many of the functional gene traits we examined tend to vary most at the order to subphylum levels (Treseder and Lennon, 2015), so we likely captured much of the trait variation with genuslevel means.

Relationships between traits

We used phylogenetic independent contrasts to examine relationships between each pairwise combination of response and functional gene traits. Since the global change experiments had sequenced the ITS1 or ITS2 regions, which are highly variable, it was not feasible to construct an accurate phylogeny from these sequences. Instead, we used the fungal phylogeny of Choi and Kim (2017), which was developed from 235 whole genome sequences of fungal strains. We downloaded the phylogeny from Github (2020, accessed March 12, 2020). Of the 111 genera in our dataset, 41 were represented in the phylogeny. For each of the remaining genera, we used the closest taxon in the tree (Table S3). Sixty-one of the genera were assigned to a taxon of the same family or order in the phylogeny. We pruned the tree to remove all taxa not represented in at least 1 case study (Figure S1).

For the phylogenetic independent contrasts, we used the *aotf* function in Phylocom version 4.2 (Webb et al., 2008). This function determined the difference in the values of a specific trait between daughter clades of each node in the phylogenetic tree. It then took a series of correlations of the contrasts between each pairwise combination of traits, minimizing errors along the x and y axes. We used the default parameters for this function, so more recently diverged clades were weighted more in the correlations. Where the correlations included an outlier with large leverage, we ranked the contrasts. We focused on pairwise relationships with r values greater than 0.5 or less than -0.5, which corresponded to an unadjusted P < 0.00001 for correlations between functional gene traits. The aotf function also identified nodes at which each response trait diverged significantly (Figure S1).

To view relationships between all traits included in the case study analysis, we performed a nonmetric multidimensional scaling analysis using the Kruskal Method with a square (similarities) model and 2-dimensional output (SPSS, 2017). The input was the matrix of r values from the phylogenetic independent contrasts (Table S4).

Relationships between traits

Our results were unexpected. The fungal traits sorted into 3 distinct groups based on their relationships to one another (Figure 2). Responses to N enrichment tended to be positively related to responses to warming and drying as well as increased precipitation response. Collectively, these response traits clustered with most of the stress tolerance traits: β-1,3 glucan synthase, trehalase, and cold-induced RNA helicase (Group A; Figure 2). Thus, we will refer to this group of traits as the "Stress Tolerating Group." A second cluster of traits featured C-targeting enzymes and phosphate transporters (Group B; Figure Specifically, crystalline cellulase was positively related to phosphate transporters and GH Families 12 and 81 (endoglucanases). In turn, GH Family 12 was positively related to GH Family 7 (cellulases and β-1,4-glucanases). This group, which we will refer to as "C and P Targeting Group," was distinct from the group that included global change response traits. Altogether, none of our predictions were upheld. In fact, the clustering of increased precipitation response with stress tolerance traits was opposite to its predicted relationship.

Stress Tolerating Group

Why did genera that increased under climate change also tend to possess stress tolerance traits? Bijlsma and Loeschcke (2005) noted that environmental stress is a function of both the *stressor* (e.g., increased precipitation) and the *stressed* (e.g., resident fungal taxa). In other words, environmental conditions that are suboptimal for most fungi could be optimal for local fungi that have adapted to them. For example, although rainfall could alleviate drought stress for many fungi, it might generate environmental stress in fungi adapted to arid ecosystems (Van Gestel et al., 1993; Fierer et al., 2003; Schimel et al., 2007). Perhaps increased precipitation, N enrichment, and warming and drying induced stress because they exposed




Figure 2. Relationships between traits. Traits are arrayed based on a 2-dimensional nonmetric multidimensional scaling analysis. Traits located closer together tend to be more closely related. Each symbol represents 1 trait. Lines connect traits with phylogenetic independent contrast r values greater than 0.5 (cyan) or less than -0.5 (yellow). Line thickness is proportional to |r|. A given trait was assigned to a group if it was related (at |r| > 0.5) to at least one other trait within the group. "Group A" is highlighted because it includes global change responses. Correlation graphs for each linked pair of traits are below (Group A: Figures 3, 4, and S2; Group B: Figure 5; Group C: Figure S3). An interactive version of this figure can be accessed at Polinode (2021). DOI: https://doi.org/10.1525/elementa.2020.00144.f2

fungi to environmental conditions they did not typically experience.

This may have been the case in the increased precipitation manipulation we analyzed. As the climate warms, precipitation regimes change with it (Intergovernmental Panel on Climate Change [IPCC], 2014). On average, precipitation rates are expected to increase globally (IPCC, 2014). In the increased precipitation experiment by McHugh and Schwartz (2016), month-long water additions in an Arizonan semiarid grassland yielded faster soil respiration rates, larger inorganic N pools, and significant shifts in the fungal community. Previous research has indicated that fungal community composition frequently shifts with water availability (Castro et al., 2010; Hawkes et al., 2011; Maestre et al., 2015; Matulich et al., 2015; McHugh and Schwartz, 2015; Gao et al., 2016; He et al., 2017; Zhao et al., 2017; Jansson and Hofmockel, 2020), although not everywhere (Barnard

et al., 2013; Jumpponen and Jones, 2014; Zhang et al., 2016a).

In this experiment, water additions may have increased fungal stress. Fungal genera whose relative abundance increased following water additions also dedicated larger proportions of their genomes to trehalase and β-1,3 glucan synthase genes (Figure 3a and b). Counterintuitively, water additions to chronically dry soils can be physiologically challenging to microbes (Schimel et al., 2007). As soils dry, microbes tend to accumulate osmolytes to reduce water loss (Harris, 1981). When soils are rewetted, the osmolytes could drive water into their cells, raise internal pressures, and potentially burst the cells (Kieft et al., 1987). Strong cell walls can help offset this risk (Kieft et al., 1987), so reinforcement with β-1,3 glucan might be advantageous in this case. In addition, microbes can quickly remove osmolytes by transforming or excreting them (Wood et al., 2001). It is unclear why fungal genera




Figure 3. Relationships between fungal responses to increased precipitation versus select functional genes in the Stress Tolerating Group. Symbols represent phylogenetic nodes (Figure S1). Contrasts are differences in trait values between daughter clades of that node. For functional genes, units are number of copies per 10,000 genes. Symbol size is proportional to the node's assigned weight in the analysis. Lines are best fit. *P* values are unadjusted for multiple comparisons. Increased precipitation responses are calculated from McHugh and Schwartz (2016), and gene frequencies are from MycoCosm (Grigoriev et al., 2014). DOI: https://doi.org/10.1525/elementa.2020.00144.f3

with the capacity to construct trehalose tended to perform better in the McHugh and Schwartz (2016) study, given that osmolytes like trehalose can be disadvantageous during rewetting. Perhaps because trehalose production can be reversed upon rewetting, it allowed fungi to tolerate drought stress while allowing them the flexibility to acclimate quickly to changing water availability (Harris, 1981). Gene frequencies for ammonium transporters and cellulases (GH Family 9) were also positively related to responses to increased precipitation (Figure 3c and d), possibly because these stress tolerance traits require N for enzyme construction and C for trehalose production.

Trehalase genes coincided with β -1,3 glucan synthase genes and another stress tolerance trait: cold-induced RNA helicase genes (**Figure 4a and b**). In an earlier examination of 157 whole fungal genomes, Treseder and Lennon (2015) had noted that genes for trehalase, β -1,3 glucan synthase, and cold-induced RNA helicases are positively related to one another across the fungal tree of life. Our current analysis of 692 fungal genomes reinforced that observation.

These 3 traits may form part of a multifaceted "environmental stress response," which has previously been defined for ascomycetes (Gasch and Werner-Washburne, 2002; Gasch, 2007). In an environmental stress response, hundreds of stress tolerance-related genes are expressed when fungi are exposed to a stressor such as heat shock. After the environmental stress response has commenced, the fungus is then tolerant of other stresses it has not been previously exposed to, such as extreme pH (Gasch, 2007; Gasch and Werner-Washburne, 2002). At the same time, other genes, including those involving carbohydrate metabolism, are repressed, ostensibly to divert resources to stress tolerance traits (Gasch, 2002; Zakrzewska et al., 2011). This type of trade-off may also have been apparent in the negative relationship we observed between cold-induced RNA helicase and α-glucosidase (GH Family 13) genes (Figure 4c). If trehalase, β-1,3 glucan synthase, and cold-induced RNA helicases were indeed part of a larger environmental stress response, then it is possible that an unidentified stress response trait-




Figure 4. Relationships between functional genes in the Stress Tolerant Group. Symbols represent phylogenetic nodes (Figure S1). Contrasts are differences in trait values between daughter clades. Units are number of copies per 10,000 genes. Symbol size is proportional to the node's assigned weight in the analysis. Lines are best fit. *P* values are unadjusted for multiple comparisons. Data are from MycoCosm (Grigoriev et al., 2014). DOI: https://doi.org/10.1525/elementa.2020.00144.f4

other than trehalase—was driving tolerance for increased precipitation.

Responses to N enrichment and warming and drying also belonged to the Stress Tolerating Group. Specifically, fungal genera that responded positively to N enrichment in the Kerekes et al. (2013) tropical forest experiment also tended to respond positively to increased precipitation and warming and drying (Figure S2a and b). In this site, N additions had reduced soil pH from 5.1 to 4.9, which may have contributed to fungal stress (Corre et al., 2010). Nitrogen enrichment shifts fungal community composition (Johnson, 1993; Egerton-Warburton and Allen, 2000; Allison et al., 2008; Allison et al., 2010a; Entwistle et al., 2013; Amend et al., 2016; Morrison et al., 2016; Jia et al., 2017; Chen et al., 2018) more often than not (Porras-Alfaro et al., 2011; Cassman et al., 2016; McHugh et al., 2017). In a temperate forest in the northeastern United States, N fertilization favors stress-tolerant fungi and those with higher gene frequencies of ammonium transporters and amino acid permeases (Morrison et al., 2018; Romero-Olivares et al., 2021). Our finding that responses to N enrichment in the tropical forest tended to cluster with stress tolerance and ammonium transporter genes was consistent with those from the temperate forest.

With respect to warming and drying in the boreal forest study, departures from ambient temperature or moisture (or both) might have been stressful for local fungi. In the boreal forest experiment, warming increases evapotranspiration and dries the topsoil by about 22% (Allison and Treseder, 2008). As a result, fungal abundance declines (Allison and Treseder, 2008). After 8 years, the fungal community in surface litter had shifted significantly (Treseder et al., 2016; Romero-Olivares et al., 2017a). Moreover, decomposition rates of the litter slowed, especially for cellulose (Romero-Olivares et al., 2017a). Metatranscriptomic profiles of the fungal community in this experiment indicate that warming and drying induces the expression of cell maintenance genes while repressing the expression of GH genes, potentially owing to trade-offs between stress tolerance and decomposition (Romero-Olivares et al., 2019). In addition, fungi belonging to known stress-tolerant taxa displayed higher gene transcription rates in the warming and drying treatment compared to controls (Romero-Olivares et al., 2019). These findings mirror those reported from the Hubbard Brook Experimental Forest in the northeastern United States, where warming combined with intensified freeze/thaw cycles selected for stress-tolerant genes in fungi and bacteria (Garcia et al., 2020).

Carbon and Phosphorus Targeting Group

The C and P Targeting Group of traits was centered on crystalline cellulose genes (Auxiliary Activity Family 9; Levasseur et al., 2013; **Figure 2**). This enzyme family releases cellulose chains from highly crystalline or cross-linked structures, which tend to be recalcitrant (Harris et al., 2010; Langston et al., 2011). Fungal genera with relatively strong capacities for crystalline cellulase production tended to be capable of glucan breakdown as well (**Figure 5a and c**, GH Families 12 and 81). Moreover, gene frequencies for endoglucanase production (GH Family 12) were positively related to those for GH Family 7, which breaks β -1,4 glycosidic bonds in cellulose and β -1,4-glucan (**Figure 5d**; Shoemaker et al., 1983; Teeri et al., 1983; Ilmen et al., 1997). This suite of carbohydrate-targeting




Figure 5. Relationships between functional genes in the C and P Targeting Group. Symbols represent phylogenetic nodes (Figure S1). Contrasts are differences in trait values between daughter clades. Units are number of copies per 10,000 genes. Symbol size is proportional to the node's assigned weight in the analysis. Lines are best fit. *P* values are unadjusted for multiple comparisons. Data are from MycoCosm (Grigoriev et al., 2014). DOI: https://doi.org/10.1525/elementa.2020.00144.f5

genes may form a decomposer lifestyle in fungi, distinct from a stress tolerance lifestyle. Previous studies have noted that C-targeting extracellular enzymes tend to coincide within fungal taxa (e.g., Eastwood et al., 2011; Floudas et al., 2012; Riley et al., 2014; Treseder and Lennon, 2015).

Phosphate transporters were also linked to this trait group. Specifically, gene frequencies of phosphate transporters were positively related to those of crystalline cellulase (Figure 5b). Fungi use P to build DNA, RNA, adenosine triphosphate, and phospholipids (Elser et al., 1996). Fungi might invest in P uptake to meet stoichiometric demands associated with increased C use (Sinsabaugh et al., 2008; Sinsabaugh et al., 2009; Sinsabaugh et al., 2016).

Unlinked trait: Responses to elevated CO₂

Several traits were not strongly linked to any others, so they did not belong to any group here. Responses to elevated CO₂ in alpine treeline was one example. Unlike the other response traits, it did not cluster with the Stress Tolerating Group. The numbers of sequences and identified taxa from this experiment were comparable to the other experiments (Table S1), so lack of statistical power is not a likely explanation. In this study, an increase in CO₂ concentrations from approximately 370 ppm (ambient) to 563–600 ppm (elevated) yields no significant change in

fine root biomass or soil fungal biomass (Hagedom et al., 2013). Likewise, the soil fungal community does not shift significantly at the operational taxonomic unit (i.e., species) level (Hagedorn et al., 2013). In fact, soil bacterial communities may have been more sensitive to elevated CO₂ than are fungi, given that bacterial composition shifts marginally significantly (Hagedorn et al., 2013). In this ecosystem, temperature limitation of plants may constrain their responses to elevated CO₂ (Dawes et al., 2011), which in turn may minimize effects on belowground C supplies and progressive nutrient limitation. Essentially, fungi in this system may have experienced little change in environmental conditions under elevated CO₂.

This case study may not be representative of other elevated CO₂ studies. Elsewhere, mycorrhizal fungi frequently become more abundant (Treseder, 2004; Dong et al., 2018). Likewise, fungal community composition often shifts (Weber et al., 2011), sometimes owing to increases in fine root production (Lipson et al., 2014). These CO₂ effects in other ecosystems might be more readily linked to stress tolerance, resource acquisition, or other lifestyles.

Revisiting the trait-based framework

We revised our trait-based framework to center links between stress tolerance traits and responses to increased




Figure 6. Revised trait-based framework relating global changes to the relative abundance of fungal taxa and their functional gene traits. DOI: https://doi.org/10.1525/elementa.2020.00144.f6

precipitation, N enrichment, and warming and drying (Figure 6). We suggest that taxa with strong cell walls (from β -1,3 glucose), osmolytes (e.g., trehalose), and cold tolerance (via cold-induced RNA helicase) may have an advantage under these conditions. The advantage may be conferred by these specific stress tolerance traits or by a larger environmental stress response with which they co-occur (Gasch and Werner-Washburne, 2002; Gasch, 2007).

In contrast, C- and P-acquisition traits may be disadvantageous under these circumstances. If stress-tolerating fungi compete for space or resources with C and P acquiring fungi, the latter group may become less prevalent. Furthermore, GH 13 genes (α-glucosidases) were negatively related to cold-induced RNA helicase genes (Figure 4c), which implies that investment in C acquisition might trade off with stress tolerance. On the other hand, if competition or trade-offs are not significant, the C and P acquiring taxa may not necessarily decline in absolute abundance upon exposure to increased precipitation, N enrichment, or warming and drying. Nevertheless, fungal species are known to compete with one another (Boddy, 2000; Fukami et al., 2007; Kolesidis et al., 2019). It would be worth investigating whether stress tolerators compete with decomposers under natural settings (Cray et al., 2013; Ho et al., 2017; Malik et al., 2019).

Another question to explore is the timescale during which stress tolerance might be advantageous under global change. If stress is induced when fungi are exposed to unfamiliar environmental conditions, will the stress decline once fungi acclimate or adapt? Here, the increased precipitation experiment was relatively brief—the water additions lasted for 1 month (McHugh and Schwartz, 2016). By comparison, N enrichment had been maintained for 9 years at the time of sampling (Kerekes et al., 2013). Moreover, the warming and drying experiment had been ongoing for 8 years. In laboratory settings, fungi can adapt or acclimate to new conditions within hours to months

(Dettman et al., 2008; DeAngelis et al., 2010; Romero-Olivares et al., 2015; Zhang et al., 2016b). However, we might expect these mechanisms to operate more slowly in the field, where conditions can be less optimal, and stress levels are less extreme or abrupt (Leuzinger et al., 2011; Gao et al., 2020). In addition, direct, immediate effects of the global change itself need not be the only sources of stress for fungi. Indirect effects via shifts in the plant community, changes in soil texture, disturbance, or other intermediate processes could sustain stress over a longer time.

The revised trait-based framework suggests potential consequences of global changes for fungal contributions to nutrient cycling (sensu Lavorel and Garnier, 2002; Chagnon et al., 2013; Treseder and Lennon, 2015; Ho et al., 2017; Malik et al., 2019). In ecosystems exposed to increased precipitation, N enrichment, or warming and drying, we might expect that β-1,3 glucan incorporation into fungal cell walls will increase owing to its contributions to stress tolerance. B-1,3 glucan forms cross-linkages with chitin (Cabib, 2009), forming a recalcitrant C complex that may remain in the soil after the fungi die (Klis, 1994; Treseder and Lennon, 2015). This process could augment soil C storage. Conversely, as C and P targeting fungi use the products of extracellular cellulases, they essentially convert a portion of the cellulose-C to CO2. If competition with stress-tolerating fungi reduce the abundance of C and P targeting fungi, then the fungal community may produce less CO2 via this process. If competition is not significant, however, this may not be a consequence. Next generation trait-based models can use these trait linkages to predict ecosystem dynamics under global change (Follows et al., 2007; Allison, 2012; Allison, 2014; Wieder et al., 2015).

Conclusions

In summary, we found that stress tolerance traits were positively related to the responses of fungal genera to 3

types of global change: increased precipitation, N enrichment, and warming and drying. Capacity for C and P acquisition seemed less critical, as functional genes for these traits were less strongly linked to global change responses. Although we had expected that ability to capture limiting nutrients under global change would influence responses of fungal genera, we found little evidence to support this notion. We developed a revised trait-based framework predicting that increased precipitation, N enrichment, and warming and drying will generate stress in fungi and select for taxa with stress tolerance traits. Stress-tolerating fungi may outcompete C and P targeting fungi if space or resources are limiting. Essentially, elements of global change-even those that might otherwise seem beneficial to fungi like increased precipitation-may generate environmental stress and select for stress-tolerating

Data accessibility statement

The following datasets were generated:

Trait database, uploaded as Table S2 of this publication.

Supplemental files

The supplemental files for this article can be found as follows:

Figures S1– S3. PDF Table S1. PDF Tables S2–S4. xlsx

Acknowledgments

We thank all authors who contributed to the studies synthesized here: A. Amend, T. Bruns, J. Choi, A. Ekblad, B. Frey, I. T. Handa, M. Hartmann, S. Hättenschwiler, D. Hiltbrunner, M. Kaspari, S-H. Kim, M. R. Maltz, Y. Marusenko, A. Miltner, R. Nilsson, A. Romero-Olivares, E. Schwartz, B. Stevenson, and K. Streit. Thanks also to L. Tedersoo and an anonymous reviewer for helpful feedback.

Funding

This work was funded by U.S. NSF DEB-1912525 and the U.S. Department of Energy Office of Biological and Environmental Research, Grant Number: DE-SC0020382. Funding was also provided by the Velux Foundation (Grant 371) and the Swiss State Secretary for Education, Research and Innovation (COST Action ES1203).

Competing interests

The authors declare no competing interests.

Author contributions

Performed data assimilation and analysis and developed the conceptual framework: KKT, CJA, LAC, MEG, ALK, KGL. Helped write the manuscript, with KKT leading: All authors.

Provided data and intellectual contributions: FH, JFK, TAM, EFS.

References

- Allison, SD. 2012. A trait-based approach for modelling microbial litter decomposition. *Ecology Letters* 15: 1058–1070.
- Allison, SD. 2014. Modeling adaptation of carbon use efficiency in microbial communities. Frontiers in Microbiology 5(571). DOI: http://dx.doi.org/10. 3389/fmicb.2014.00571.
- Allison, SD, Czimczik, CI, Treseder, KK. 2008. Microbial activity and soil respiration under nitrogen addition in Alaskan boreal forest. Global Change Biology 14: 1156–1168.
- Allison, SD, Gartner, TB, Mack, MC, McGuire, K, Treseder, K. 2010a. Nitrogen alters carbon dynamics during early succession in boreal forest. *Soil Biology and Biochemistry* 42(7): 1157–1164. DOI: http://dx.doi.org/10.1016/j.soilbio.2010.03.026.
- Allison, SD, Martiny, JBH. 2008. Resistance, resilience, and redundancy in microbial communities. Proceedings of the National Academy of Sciences of the United States of America 105: 11512–11519. DOI: http://dx.doi.org/10.1073/pnas.0801925105.
- Allison, SD, Treseder, KK. 2008. Warming and drying suppress microbial activity and carbon cycling in boreal forest soils. Global Change Biology 14(12): 2898–2909.
- Allison, SD, Treseder, KK. 2011. Climate change feed-backs to microbial decomposition in boreal soils. Fungal Ecology 4(6): 362–374. DOI: http://dx.doi.org/10.1016/j.funeco.2011.01.003.
- Allison, SD, Weintraub, MN, Gartner, TB, Waldrop, MP. 2010b. Evolutionary-economic principles as regulators of soil enzyme production and ecosystem function, in Shukla, GC, Varma, A eds., Soil enzymology. Heidelberg, Germany: Springer: 245–258.
- Amend, AS, Martiny, AC, Allison, SD, Berlemont, R, Goulden, ML, Lu, Y, Treseder, KK, Weihe, C, Martiny, JB. 2016. Microbial response to simulated global change is phylogenetically conserved and linked with functional potential. *The* ISME Journal 10(1): 109–118. DOI: http://dx.doi.org/10.1038/ ismej.2015.96.
- Baath, E, Lundgren, B, Soderstrom, B. 1981. Effects of nitrogen fertilization on the activity and biomass of fungi and bacteria in a podzolic soil. Zentralblatt Fur Bakteriologie Mikrobiologie Und Hygiene I Abteilung Originale C-Allgemeine Angewandte Und Okologische Mikrobiologie 2(1): 90–98.
- Barnard, RL, Osborne, CA, Firestone, MK. 2013. Responses of soil bacterial and fungal communities to extreme desiccation and rewetting. *The ISME Journal* 7(11): 2229–2241. DOI: http://dx.doi.org/10.1038/ismej.2013.104.
- Bending, GD, Read, DJ. 1997. Lignin and soluble phenolic degradation by ectomycorrhizal and ericoid mycorrhizal fungi. Mycological Research 101: 1348–1354. DOI: http://dx.doi.org/10.1017/s0953756297004140.
- Bijlsma, R, Loeschcke, V. 2005. Environmental stress, adaptation and evolution: an overview. Journal of

- Evolutionary Biology **18**(4): 744–749. DOI: http://dx.doi.org/10.1111/j.1420-9101.2005.00962.x.
- Boddy, L. 2000. Interspecific combative interactions between wood-decaying basidiomycetes. FEMS Microbiology Ecology 31(3): 185–194. DOI: http:// dx.doi.org/10.1111/j.1574-6941.2000.tb00683.x.
- Bowman, SM, Free, SJ. 2006. The structure and synthesis of the fungal cell wall. *Bioessays* **28**(8): 799–808. DOI: http://dx.doi.org/10.1002/bies.20441.
- Broadbent, FE. 1965. Effects of fertilizer nitrogen on the release of soil nitrogen. Soil Science Society of America Proceedings 29: 692–696.
- Cabib, E. 2009. Two novel techniques for determination of polysaccharide cross-links show that Crh1p and Crh2p attach chitin to both beta(1–6)- and beta (1–3)glucan in the Saccharomyces cerevisiae cell wall. Eukaryotic Cell 8(11): 1626–1636. DOI: http://dx.doi.org/10.1128/ec.00228-09.
- Cairney, JWG, Burke, RM. 1998. Extracellular enzyme activities of the ericoid mycorrhizal endophyte Hymenoscyphus ericae (Read) Korf & Kernan: their likely roles in decomposition of dead plant tissue in soil. Plant and Soil 205(2): 181–192. DOI: http://dx.doi.org/10.1023/a:1004376731209.
- Cassman, NA, Leite, MFA, Pan, Y, de Hollander, M, van Veen, JA, Kuramae, EE. 2016. Plant and soil fungal but not soil bacterial communities are linked in long-term fertilized grassland. Scientific Reports 6(1): 23680. DOI: http://dx.doi.org/10.1038/ srep23680.
- Castro, HF, Classen, AT, Austin, EE, Norby, RJ, Schadt, CW. 2010. Soil microbial community responses to multiple experimental climate change drivers. Applied and Environmental Microbiology 76(4): 999–1007. DOI: http://dx.doi.org/10.1128/aem. 02874-09.
- Chagnon, PL, Bradley, RL, Maherali, H, Klironomos, JN. 2013. A trait-based framework to understand life history of mycorrhizal fungi. *Trends in Plant Science* 18(9): 484–491. DOI: http://dx.doi.org/10.1016/j. tplants.2013.05.001.
- Chapin, FS, Matson, PA, Vitousek, PM, Chapin, MC. 2011. *Principles of terrestrial ecosystem ecology.* 2nd ed. New York, NY: Springer.
- Chen, J, Luo, YQ, Xia, JY, Jiang, LF, Zhou, XH, Lu, M, Liang, J, Shi, Z, Shelton, S, Cao, J. 2015. Stronger warming effects on microbial abundances in colder regions. *Scientific Reports* 5: 10. DOI: http://dx.doi. org/10.1038/srep18032.
- Chen, W, Xu, R, Wu, Y, Chen, J, Zhang, Y, Hu, T, Yuan, X, Zhou, L, Tan, T, Fan, J. 2018. Plant diversity is coupled with beta not alpha diversity of soil fungal communities following N enrichment in a semi-arid grassland. Soil Biology and Biochemistry 116: 388–398. DOI: http://dx.doi.org/10.1016/j.soilbio. 2017.10.039.
- Choi, J. 2020. Available at https://github.com/jaejinchoi/ FFP. Accessed 12 March 2020.
- Choi, J, Kim, S-H. 2017. A genome Tree of Life for the Fungi kingdom. Proceedings of the National

- Academy of Sciences 114(35): 9391–9396. DOI: http://dx.doi.org/10.1073/pnas.1711939114.
- Classen, AT, Sundqvist, MK, Henning, JA, Newman, GS, Moore, JAM, Cregger, MA, Moorhead, LC, Patterson, CM. 2015. Direct and indirect effects of climate change on soil microbial and soil microbial-plant interactions: What lies ahead? *Ecosphere* 6(8): art130. DOI: http://dx.doi.org/10.1890/es15-00217.1.
- Compant, S, van der Heijden, MGA, Sessitsch, A. 2010. Climate change effects on beneficial plantmicroorganism interactions. *FEMS Microbiology Ecology* **73**(2): 197–214. DOI: http://dx.doi.org/10. 1111/j.1574-6941.2010.00900.x.
- Corre, MD, Veldkamp, E, Arnold, J, Wright, SJ. 2010. Impact of elevated N input on soil N cycling and losses in old-growth lowland and montane forests in Panama. *Ecology* 91(6): 1715–1729. DOI: http:// dx.doi.org/10.1890/09-0274.1.
- Cray, JA, Bell, ANW, Bhaganna, P, Mswaka, AY, Timson, DJ, Hallsworth, JE. 2013. The biology of habitat dominance; can microbes behave as weeds? *Microbial Biotechnology* 6(5): 453–492. DOI: http://dx.doi.org/10.1111/1751-7915.12027.
- Dawes, MA, Hagedorn, F, Zumbrunn, T, Handa, IT, Hättenschwiler, S, Wipf, S, Rixen, C. 2011. Growth and community responses of alpine dwarf shrubs to in situ CO₂ enrichment and soil warming. New Phytologist 191(3): 806–818. DOI: http://dx.doi.org/10.1111/j.1469-8137.2011.03722.x.
- DeAngelis, KM, Silver, WL, Thompson, AW, Firestone, MK. 2010. Microbial communities acclimate to recurring changes in soil redox potential status. Environmental Microbiology 12(12): 3137–3149. DOI: http://dx.doi.org/10.1111/j.1462-2920.2010. 02286.x.
- de Graaff, MA, van Groenigen, KJ, Six, J, Hungate, B, van Kessel, C. 2006. Interactions between plant growth and soil nutrient cycling under elevated CO₂: A meta-analysis. Global Change Biology 12(11): 2077–2091. DOI: http://dx.doi.org/10.1111/j.1365-2486.2006.01240.x.
- Dettman, J, Anderson, J, Kohn, L. 2008. Divergent adaptation promotes reproductive isolation among experimental populations of the filamentous fungus Neurospora. BMC Evolutionary Biology 8(1): 35.
- Diaz, S. 1996. Effects of elevated [CO₂] at the community level mediated by root symbionts. *Plant and Soil* 187(2): 309–320.
- Dieleman, WIJ, Luyssaert, S, Rey, A, De Angelis, P, Barton, CVM, Broadmeadow, MS, Broadmeadow, SB, Chigwerewe, KS, Crookshanks, M, Dufrêne, E, Jarvis, PG. 2010. Soil N modulates soil C cycling in CO₂-fumigated tree stands: A meta-analysis. *Plant Cell and Environment* 33(12): 2001–2011. DOI: http://dx.doi.org/10.1111/j.1365-3040.2010. 02201.x.
- Dong, YL, Wang, ZY, Sun, H, Yang, WC, Xu, H. 2018. The response patterns of arbuscular mycorrhizal and

- ectomycorrhizal symbionts under elevated CO₂: A meta-analysis. *Frontiers in Microbiology* **9**: 14. DOI: http://dx.doi.org/10.3389/fmicb.2018.01248.
- Eastwood, DC, Floudas, D, Binder, M, Majcherczyk, A, Schneider, P, Aerts, A, Asiegbu, FO, Baker, SE, Barry, K, Bendiksby, M, Blumentritt, M. 2011. The plant cell wall-decomposing machinery underlies the functional diversity of forest fungi. *Science* 333(6043): 762–765. DOI: http://dx.doi.org/10. 1126/science.1205411.
- Edwards, IP, Upchurch, RA, Zak, DR. 2008. Isolation of fungal cellobiohydrolase I genes from sporocarps and forest soils by PCR. Applied and Environmental Microbiology 74(11): 3481–3489. DOI: http://dx. doi.org/10.1128/aem.02893-07.
- Egerton-Warburton, LM, Allen, EB. 2000. Shifts in arbuscular mycorrhizal communities along an anthropogenic nitrogen deposition gradient. *Ecological* Applications 10: 484–496.
- Elser, JJ, Dobberfuhl, DR, MacKay, NA, Schampel, JH. 1996. Organism size, life history, and N: P stoichiometry: Toward a unified view of cellular and ecosystem processes. *Bioscience* 46(9): 674–685.
- Entwistle, EM, Zak, DR, Edwards, IP. 2013. Long-term experimental nitrogen deposition alters the composition of the active fungal community in the forest floor. Soil Science Society of America Journal 77(5): 1648–1658. DOI: http://dx.doi.org/10.2136/ sssaj2013.05.0179.
- Estruch, F. 2000. Stress-controlled transcription factors, stress-induced genes and stress tolerance in budding yeast. FEMS Microbiology Reviews 24(4): 469–486. DOI: http://dx.doi.org/10.1016/S0168-6445(00) 00035-8.
- Fierer, N, Schimel, JP, Holden, PA. 2003. Influence of drying–rewetting frequency on soil bacterial community structure. *Microbial Ecology* 45(1): 63–71. DOI: http://dx.doi.org/10.1007/s00248-002-1007-2.
- Floudas, D, Binder, M, Riley, R, Barry, K, Blanchette, RA, Henrissat, B, Martínez, AT, Otillar, R, Spatafora, JW, Yadav, JS, Aerts, A. 2012. The Paleozoic origin of enzymatic lignin decomposition reconstructed from 31 fungal genomes. *Science* 336(6089): 1715–1719. DOI: http://dx.doi.org/10. 1126/science.1221748.
- Follows, MJ, Dutkiewicz, S, Grant, S, Chisholm, SW. 2007. Emergent biogeography of microbial communities in a model ocean. *Science* 315(5820): 1843– 1846. DOI: http://dx.doi.org/10.1126/science. 1138544.
- Fukami, T, Beaumont, HJE, Zhang, X-X, Rainey, PB. 2007. Immigration history controls diversification in experimental adaptive radiation. *Nature* 446(7134): 436–439. DOI: http://dx.doi.org/10. 1038/nature05629.
- Gao, C, Kim, Y-C, Zheng, Y, Yang, W, Chen, L, Ji, NN, Wan, SQ, Guo, LD. 2016. Increased precipitation, rather than warming, exerts a strong influence on arbuscular mycorrhizal fungal community in

- a semiarid steppe ecosystem. *Botany* **94**(6): 459–469. DOI: http://dx.doi.org/10.1139/cjb-2015-0210.
- Gao, D, Bai, E, Li, M, Zhao, C, Yu, K, Hagedorn, F. 2020. Responses of soil nitrogen and phosphorus cycling to drying and rewetting cycles: A meta-analysis. *Soil Biology and Biochemistry* 148: 107896. DOI: http://dx.doi.org/10.1016/j.soilbio.2020.107896.
- Gao, WL, Yan, DH. 2019. Warming suppresses microbial biomass but enhances N recycling. Soil Biology and Biochemistry 131: 111–118. DOI: http://dx.doi.org/ 10.1016/j.soilbio.2019.01.002.
- Garcia, MO, Templer, PH, Sorensen, PO, Sanders-DeMott, R, Groffman, PM, Bhatnagar, JM. 2020. Soil microbes trade-off biogeochemical cycling for stress tolerance traits in response to year-round climate change. Frontiers in Microbiology 11(616). DOI: http://dx.doi.org/10.3389/fmicb.2020.00616.
- Gasch, AP. 2002. The environmental stress response: a common yeast response to environmental stresses, in Hohmann, S, Mager, P eds., Yeast stress responses. Berlin, Germany: Springer-Verlag: 11–70.
- Gasch, AP. 2007. Comparative genomics of the environmental stress response in ascomycete fungi. *Yeast* 24(11): 961–976. DOI: http://dx.doi.org/10.1002/yea.1512.
- Gasch, AP, Werner-Washburne, M. 2002. The genomics of yeast responses to environmental stress and starvation. Functional & Integrative Genomics 2(4–5): 181–92. DOI: http://dx.doi.org/10.1007/s10142-002-0058-2.
- Grigoriev, IV, Nikitin, R, Haridas, S, Kuo, A, Ohm, R, Otillar, R, Riley, R, Salamov, A, Zhao, X, Korzeniewski, F, Smirnova, T. 2014. MycoCosm portal: Gearing up for 1000 fungal genomes. *Nucleic Acids Research* 42(1): D699–704.
- Hagedorn, F, Hiltbrunner, D, Streit, K, Ekblad, A, Lindahl, B, Miltner, A, Frey, B, Handa, IT, Hättenschwiler, S. 2013. Nine years of CO₂ enrichment at the alpine treeline stimulates soil respiration but does not alter soil microbial communities. *Soil Biology and Biochemistry* 57: 390–400. DOI: http://dx.doi.org/10.1016/j.soilbio.2012.10.001.
- Harris, PV, Welner, D, McFarland, KC, Re, E, Poulsen, JCN, Brown, K, Salbo, R, Ding, H, Vlasenko, E, Merino, S, Xu, F. 2010. Stimulation of lignocellulosic biomass hydrolysis by proteins of Glycoside Hydrolase Family 61: Structure and function of a large, enigmatic family. *Biochemistry* 49(15): 3305–3316. DOI: http://dx.doi.org/10.1021/bi100009p.
- Harris, RF. 1981. Effect of water potential on microbial growth and activity, in Parr, JF, Gardner, WR, Elliott, LF eds., Water potential relations in soil microbiology. Madison, WI: American Society of Agronomy: 23–95.
- Hawkes, CV, Kivlin, SN, Rocca, JD, Huguet, V, Thomsen, MA, Suttle, KB. 2011. Fungal community responses to precipitation. Global Change Biology

- **17**(4): 1637–1645. DOI: http://dx.doi.org/10.1111/j.1365-2486.2010.02327.x.
- He, D, Shen, W, Eberwein, J, Zhao, Q, Ren, L, Wu, QL. 2017. Diversity and co-occurrence network of soil fungi are more responsive than those of bacteria to shifts in precipitation seasonality in a subtropical forest. *Soil Biology and Biochemistry* 115: 499–510. DOI: http://dx.doi.org/10.1016/j.soilbio.2017.09. 023.
- Ho, A, Di Lonardo, DP, Bodelier, PLE. 2017. Revisiting life strategy concepts in environmental microbial ecology. FEMS Microbiol Ecol 93(3): 14. DOI: http://dx.doi.org/10.1093/femsec/fix006.
- Homyak, PM, Allison, SD, Huxman, TE, Goulden, ML, Treseder, KK. 2018. Effects of drought manipulation on soil nitrogen cycling: A meta-analysis. *Jour*nal of Geophysical Research: Biogeosciences 122: 3260–3272. DOI: http://dx.doi.org/10.1002/ 2017JG004146.
- Hungate, BA, van Groenigen, KJ, Six, J, Jastrow, JD, Luo, YQ, De GRAAFF, MA, van Kessel, C, Osenberg, CW. 2009. Assessing the effect of elevated carbon dioxide on soil carbon: A comparison of four meta-analyses. Global Change Biology 15(8): 2020– 2034. DOI: http://dx.doi.org/10.1111/j.1365-2486. 2009.01866.x.
- Ilmen, M, Saloheimo, A, Onnela, ML, Penttila, ME. 1997. Regulation of cellulase gene expression in the filamentous fungus *Trichoderma reesei*. Applied and Environmental Microbiology 63(4): 1298–1306.
- IndexFungorum. 2021. Available at http://www. indexfungorum.org/. Accessed 24 February 2021.
- Intergovernmental Panel on Climate Change. 2014.

 Climate Change 2013: The Physical Science Basis:

 Working Group I Contribution to the Fifth Assessment
 Report of the International Panel on Climate Change.

 London, UK: Cambridge University Press.
- Jansson, JK, Hofmockel, KS. 2020. Soil microbiomes and climate change. *Nature Reviews Microbiology* 18(1): 35–46. DOI: http://dx.doi.org/10.1038/ s41579-019-0265-7.
- Jia, M, Liu, C, Li, Y, Xu, S, Han, G, Huang, J, Jin, B, Zou, Y, Zhang, G. 2017. Response of fungal composition and diversity to simulated nitrogen deposition and manipulation of precipitation in soils of an Inner Mongolia desert steppe of northern China. Canadian Journal of Soil Science 97(4): 613–625.
- Johnson, NC. 1993. Can fertilization of soil select less mutualistic mycorrhizae? *Ecological Applications* 3(4): 749–757.
- Johnson, NC, Angelard, C, Sanders, IR, Kiers, ET. 2013. Predicting community and ecosystem outcomes of mycorrhizal responses to global change. *Ecology Letters* 16: 140–153. DOI: http://dx.doi.org/10.1111/ele.12085.
- Jumpponen, A, Jones, KL. 2014. Tallgrass prairie soil fungal communities are resilient to climate change. Fungal Ecology 10: 44–57. DOI: http://dx.doi.org/ 10.1016/j.funeco.2013.11.003.

- Kerekes, JF, Kaspari, M, Stevenson, B, Nilsson, RH, Hartmann, M, Amend, A, Bruns, TD. 2013. Nutrient enrichment increased species richness of leaf litter fungal assemblages in a tropical forest. *Molecular Ecology* 22(10): 2827–2838. DOI: http://dx.doi. org/10.1111/mec.12259.
- Kieft, TL, Soroker, E, Firestone, MK. 1987. Microbial biomass response to a rapid increase in water potential when dry soil is wetted. Soil Biology and Biochemistry 19(2): 119–126. DOI: http://dx.doi.org/ 10.1016/0038-0717(87)90070-8.
- Klis, FM. 1994. Cell wall assembly in yeast. *Yeast* 10(7): 851–869. DOI: http://dx.doi.org/10.1002/yea. 320100702.
- Kolesidis, DA, Boddy, L, Eastwood, DC, Yuan, C, Fowler, MS. 2019. Predicting fungal community dynamics driven by competition for space. *Fungal Ecology* 41: 13–22. DOI: http://dx.doi.org/10.1016/j. funeco.2019.04.003.
- Langston, JA, Shaghasi, T, Abbate, E, Xu, F, Vlasenko, E, Sweeney, MD. 2011. Oxidoreductive cellulose depolymerization by the enzymes cellobiose dehydrogenase and glycoside hydrolase 61. Applied and Environmental Microbiology 77 (19): 7007–7015. DOI: http://dx.doi.org/10.1128/aem.05815-11.
- Latgé, J-P. 2007. The cell wall: a carbohydrate armour for the fungal cell. *Molecular Microbiology* 66(2): 279– 290. DOI: http://dx.doi.org/10.1111/j.1365-2958. 2007.05872.x.
- Lavorel, S, Garnier, E. 2002. Predicting changes in community composition and ecosystem functioning from plant traits: Revisiting the Holy Grail. Functional Ecology 16(5): 545–556.
- Lenoir, I, Fontaine, J, Lounès-Hadj Sahraoui, A. 2016. Arbuscular mycorrhizal fungal responses to abiotic stresses: A review. *Phytochemistry* 123: 4–15. DOI: http://dx.doi.org/10.1016/j.phytochem.2016.01. 002.
- Leuzinger, S, Luo, Y, Beier, C, Dieleman, W, Vicca, S, Körner, C. 2011. Do global change experiments overestimate impacts on terrestrial ecosystems? Trends in Ecology & Evolution 26(5): 236–241. DOI: http://dx.doi.org/10.1016/j.tree.2011.02.011.
- Levasseur, A, Drula, E, Lombard, V, Coutinho, PM, Henrissat, B. 2013. Expansion of the enzymatic repertoire of the CAZy database to integrate auxiliary redox enzymes. *Biotechnology for Biofuels* 6: 41. DOI: http://dx.doi.org/10.1186/1754-6834-6-41.
- Liebig, JV. 1843. Chemistry in its applications to agriculture and physiology. 3rd English ed. London, UK: Taylor Walton.
- Lindahl, BD, Nilsson, RH, Tedersoo, L, Abarenkov, K, Carlsen, T, Kjøller, R, Kõljalg, U, Pennanen, T, Rosendahl, S, Stenlid, J, Kauserud, H. 2013. Fungal community analysis by high-throughput sequencing of amplified markers - a user's guide. New Phytologist 199(1): 288–299. DOI: http://dx. doi.org/10.1111/nph.12243.
- Lipson, DA, Kuske, CR, Gallegos-Graves, LV, Oechel, WC. 2014. Elevated atmospheric CO₂ stimulates soil

- fungal diversity through increased fine root production in a semiarid shrubland ecosystem. *Global Change Biology* **20**(8): 2555–2565. DOI: http://dx.doi.org/10.1111/gcb.12609.
- Liu, L, Greaver, TL. 2010. A global perspective on belowground carbon dynamics under nitrogen enrichment. *Ecology Letters* 13(7): 819–828. DOI: http:// dx.doi.org/10.1111/j.1461-0248.2010.01482.x.
- Lu, M, Yang, Y, Luo, Y, Fang, C, Zhou, X, Chen J, Yang, X, Li, B. 2011. Responses of ecosystem nitrogen cycle to nitrogen addition: a meta-analysis. *New Phytologist* 189(4): 1040–1050. DOI: http://dx.doi.org/10. 1111/j.1469-8137.2010.03563.x.
- Lynd, LR, Weimer, PJ, van Zyl, WH, Pretorius, IS. 2002. Microbial cellulose utilization: Fundamentals and biotechnology. Microbiology and Molecular Biology Reviews 66(3): 506–577.
- Maestre, FT, Delgado-Baquerizo, M, Jeffries, TC, Eldridge, DJ, Ochoa, V, Gozalo, B, Quero, JL, Garcia-Gomez, M, Gallardo, A, Ulrich, W, Bowker, MA. 2015. Increasing aridity reduces soil microbial diversity and abundance in global drylands. Proceedings of the National Academy of Sciences 112(51): 15684–15689. DOI: http://dx.doi.org/10.1073/pnas.1516684112.
- Malik, AA, Martiny, JBH, Brodie, EL, Martiny, AC, Treseder, KK, Allison, SD. 2019. Defining trait-based microbial strategies with consequences for soil carbon cycling under climate change. The ISME Journal. DOI: http://dx.doi.org/10.1038/s41396-019-0510-0.
- Manzoni, S, Schaeffer, SM, Katul, G, Porporato, A, Schimel, JP. 2014. A theoretical analysis of microbial eco-physiological and diffusion limitations to carbon cycling in drying soils. Soil Biology and Biochemistry 73: 69–83. DOI: http://dx.doi.org/10. 1016/j.soilbio.2014.02.008.
- Manzoni, S, Schimel, JP, Porporato, A. 2012. Responses of soil microbial communities to water stress: results from a meta-analysis. *Ecology* 93(4): 930–938. DOI: http://dx.doi.org/10.1890/11-0026.1.
- Martínez, ÁT, Ruiz-Dueñas, FJ, Martínez, MJ, del Río, JC, Gutiérrez, A. 2009. Enzymatic delignification of plant cell wall: from nature to mill. *Current Opinion in Biotechnology* 20(3): 348–357. DOI: http://dx.doi.org/10.1016/j.copbio.2009.05.002.
- Martiny, JBH, Jones, SE, Lennon, JT, Martiny, AC. 2015. Microbiomes in light of traits: A phylogenetic perspective. *Science* 350(6261). DOI: http://dx.doi.org/ 10.1126/science.aac9323.
- Matulich, KL, Weihe, C, Allison, SD, Amend, AS, Berlemont, R, Goulden, ML, Kimball, S, Martiny, AC, Martiny, JB. 2015. Temporal variation overshadows the response of leaf litter microbial communities to simulated global change. *The ISME Journal* 9(11): 2477–2489. DOI: http://dx.doi.org/10.1038/ismej. 2015.58.
- McHugh, TA, Morrissey, EM, Mueller, RC, Gallegos-Graves, LV, Kuske, CR, Reed, SC. 2017. Bacterial, fungal, and plant communities exhibit no biomass

- or compositional response to two years of simulated nitrogen deposition in a semiarid grassland. *Environmental Microbiology* **19**(4): 1600–1611. DOI: http://dx.doi.org/10.1111/1462-2920.13678.
- McHugh, TA, Schwartz, E. 2015. Changes in plant community composition and reduced precipitation have limited effects on the structure of soil bacterial and fungal communities present in a semiarid grassland. *Plant and Soil* 388(1): 175–186. DOI: http://dx.doi.org/10.1007/s11104-014-2269-4.
- McHugh, TA, Schwartz, E. 2016. A watering manipulation in a semiarid grassland induced changes in fungal but not bacterial community composition. *Pedobiologia* **59**(3): 121–127. DOI: http://dx.doi.org/10.1016/j.pedobi.2016.04.003.
- Mitsuzawa, H. 2006. Ammonium transporter genes in the fission yeast *Schizosaccharomyces pombe*: Role in ammonium uptake and a morphological transition. *Genes to Cells* 11(10): 1183–1195. DOI: http:// dx.doi.org/10.1111/j.1365-2443.2006.01014.x.
- Mohan, JE, Cowden, CC, Baas, P, Dawadi, A, Frankson, PT, Helmick, K, Hughes, E, Khan, S, Lang, A, Machmuller, M, Taylor, M. 2014. Mycorrhizal fungi mediation of terrestrial ecosystem responses to global change: mini-review. Fungal Ecology 10: 3– 19. DOI: http://dx.doi.org/10.1016/j.funeco.2014. 01.005.
- Morrison, EW, Frey, SD, Sadowsky, JJ, van Diepen, LTA, Thomas, WK, Pringle, A. 2016. Chronic nitrogen additions fundamentally restructure the soil fungal community in a temperate forest. *Fungal Ecology* 23: 48–57. DOI: http://dx.doi.org/10.1016/j. funeco.2016.05.011.
- Morrison, EW, Pringle, A, van Diepen, LTA, Frey, SD. 2018. Simulated nitrogen deposition favors stresstolerant fungi with low potential for decomposition. Soil Biology and Biochemistry 125: 75–85. DOI: http://dx.doi.org/10.1016/j.soilbio.2018.06.027.
- Mouginot, C, Kawamura, R, Matulich, KL, Berlemont, R, Allison, SD, Amend, AS, Martiny, AC. 2014. Elemental stoichiometry of Fungi and Bacteria strains from grassland leaf litter. Soil Biology and Biochemistry 76: 278–285. DOI: http://dx.doi.org/ 10.1016/j.soilbio.2014.05.011.
- Myrold, DD, Zeglin, LH, Jansson, JK. 2014. The potential of metagenomic approaches for understanding soil microbial processes. Soil Science Society of America Journal 78(1): 3–10. DOI: http://dx.doi.org/10. 2136/sssaj2013.07.0287dgs.
- Nehls, U, Kleber, R, Wiese, J, Hampp, R. 1999. Isolation and characterization of a general amino acid permease from the ectomycorrhizal fungus *Amanita mus*caria. New Phytologist 144(2): 343–349.
- Nilsson, RH, Larsson, K-H, Taylor, AFS, Bengtsson-Palme, J, Jeppesen, TS, Schigel, D, Kennedy, P, Picard, K, Glöckner, FO, Tedersoo, L, Saar, I. 2018. The UNITE database for molecular identification of fungi: handling dark taxa and parallel taxonomic classifications. *Nucleic Acids Research*. DOI: http://dx.doi.org/10.1093/nar/gky1022.

- Norby, RJ, Cotrufo, MF, Ineson, P, O'Neill, EG, Canadell, JG. 2001. Elevated CO₂, litter chemistry, and decomposition: A synthesis. *Oecologia* **127**(2): 153–165.
- Norby, RJ, Zak, DR. 2011. Ecological lessons from Free-Air CO₂ Enrichment (FACE) experiments, in Futuyma, DJ, Shaffer, HB, Simberloff, D eds., Annual review of ecology, evolution, and systematics, Vol. 42. Palo Alto, CA: Annual Reviews: 181–203.
- Owttrim, GW. 2006. RNA helicases and abiotic stress. Nucleic Acids Research 34(11): 3220–3230. DOI: http://dx.doi.org/10.1093/nar/gkl408.
- Parker, SS, Schimel, JP. 2011. Soil nitrogen availability and transformations differ between the summer and the growing season in a California grassland. *Applied Soil Ecology* 48(2): 185–192. DOI: http:// dx.doi.org/10.1016/j.apsoil.2011.03.007.
- Pickles, BJ, Egger, KN, Massicotte, HB, Green, DS. 2012. Ectomycorrhizas and climate change. Fungal Ecology 5(1): 73–84. DOI: http://dx.doi.org/10. 1016/j.funeco.2011.08.009.
- Polinode. 2021. Available at https://app.polinode.com/ networks/explore/6045b7a2e63e6c0011ee77f7/ 604a6ddbe95bf60011849440. Accessed 25 June 2021.
- Porras-Alfaro, A, Herrera, J, Natvig, DO, Lipinski, K, Sinsabaugh, RL. 2011. Diversity and distribution of soil fungal communities in a semiarid grassland. *Mycologia* 103(1): 10–21. DOI: http://dx.doi.org/ 10.3852/09-297.
- Pradet-Balade, B, Boulmé, F, Beug, H, Müllner, EW, Garcia-Sanz, JA. 2001. Translation control: Bridging the gap between genomics and proteomics? *Trends* in *Biochemical Sciences* 26(4): 225–229.
- Read, DJ. 1991. Mycorrhizas in ecosystems Nature's response to the "Law of the minimum," in Hawksworth DL ed., Frontiers in mycology. Regensburg, Germany: CAB International: 101–130.
- Rebetez, M, Dobbertin, M. 2004. Climate change may already threaten Scots pine stands in the Swiss Alps. Theoretical and Applied Climatology 79(1): 1–9. DOI: http://dx.doi.org/10.1007/s00704-004-0058-3.
- Riley, R, Salamov, AA, Brown, DW, Nagy, LG, Floudas, D, Held, BW, Levasseur, A, Lombard, V, Morin, E, Otillar, R, Lindquist, EA. 2014. Extensive sampling of basidiomycete genomes demonstrates inadequacy of the white-rot/brown-rot paradigm for wood decay fungi. *Proceedings of the National Academy of Sciences* 111: 9923–9928. DOI: http://dx.doi.org/10.1073/pnas.1400592111.
- Romero-Olivares, AL, Allison, SD, Treseder, KK. 2017a. Decomposition of recalcitrant carbon under experimental warming in boreal forest. *PLoS ONE* 12(6): e0179674. DOI: http://dx.doi.org/10.1371/journal.pone.0179674.
- Romero-Olivares, AL, Allison, SD, Treseder, KK. 2017b. Soil microbes and their response to experimental warming over time: A meta-analysis of field studies. *Soil Biology and Biochemistry* **107**: 32–40. DOI: http://dx.doi.org/10.1016/j.soilbio.2016.12.026.

- Romero-Olivares, AL, Meléndrez-Carballo, G, Lago-Lestón, A, Treseder, KK. 2019. Soil metatranscriptomes under long-term experimental warming and drying: Fungi allocate resources to cell metabolic maintenance rather than decay. Frontiers in Microbiology 10(1914). DOI: http://dx.doi.org/10.3389/ fmicb.2019.01914.
- Romero-Olivares, AL, Morrison, EW, Pringle, A, Frey, SD. 2021. Linking genes to traits in fungi. *Microbial Ecology*. DOI: http://dx.doi.org/10.1007/s00248-021-01687-x.
- Romero-Olivares, AL, Taylor, J, Treseder, KK. 2015. Neurospora discreta as a model to assess adaptation of soil fungi to warming. BMC Evolutionary Biology 15(1): 198.
- Schade, B, Jansen, G, Whiteway, M, Entian, KD, Thomas, DY. 2004. Cold adaptation in budding yeast. Molecular Biology of the Cell 15(12): 5492– 5502. DOI: http://dx.doi.org/10.1091/mbc.E04-03-0167.
- Schimel, J, Balser, TC, Wallenstein, M. 2007. Microbial stress-response physiology and its implications for ecosystem function. *Ecology* 88(6): 1386–1394.
- Schimel, JP, Bennett, J. 2004. Nitrogen mineralization: Challenges of a changing paradigm. *Ecology* 85(3): 591–602.
- Shoemaker, S, Schweickart, V, Ladner, M, Gelfand, D, Kwok, S, Myambo, KA, Innis, M. 1983. Molecular clonin of exo-cellobiohydrolase I from *Trichoderma* reesi strain L27. Nature Biotechnology 1(8): 691–696. DOI: http://dx.doi.org/10.1038/nbt1083-691.
- Sinsabaugh, RL. 1994. Enzymatic analysis of microbial pattern and process. *Biology and Fertility of Soils* 17(1): 69–74.
- Sinsabaugh, RL, Hill, BH, Follstad Shah, JJ. 2009. Ecoenzymatic stoichiometry of microbial organic nutrient acquisition in soil and sediment. *Nature* 462(7274): 795–798. DOI: http://dx.doi.org/10. 1038/nature08632.
- Sinsabaugh, RL, Lauber, CL, Weintraub, MN, Ahmed, B, Allison, SD, Crenshaw, C, Contosta, AR, Cusack, D, Frey, S, Gallo, ME, Gartner, TB. 2008. Stoichiometry of soil enzyme activity at global scale. *Ecology Letters* 11(11): 1252–1264. DOI: http://dx. doi.org/10.1111/j.1461-0248.2008.01245.x.
- Sinsabaugh, RL, Turner, BL, Talbot, JM, Waring, BG, Powers, JS, Kuske, CR, Moorhead, DL, Follstad Shah, JJ. 2016. Stoichiometry of microbial carbon use efficiency in soils. *Ecological Monographs* 86(2): 172–189. DOI: http://dx.doi.org/10.1890/15-2110.1.
- Slot, JC, Hallstrom, KN, Matheny, PB, Hibbett, DS. 2007. Diversification of NRT2 and the origin of its fungal homolog. *Molecular Biology and Evolution* 24(8): 1731–1743. DOI: http://dx.doi.org/10. 1093/molbev/msm098.
- Solly, EF, Djukic, I, Moiseev, PA, Andreyashkina, NI, Devi, NM, Göransson, H, Mazepa, VS, Shiyatov, SG, Trubina, MR, Schweingruber, FH, Wilmking, M. 2017a. Treeline advances and associated shifts in

- the ground vegetation alter fine root dynamics and mycelia production in the South and Polar Urals. *Oecologia* **183**(2): 571–586. DOI: http://dx.doi. org/10.1007/s00442-016-3785-0.
- Solly, EF, Lindahl, BD, Dawes, MA, Peter, M, Souza, RC, Rixen, C, Hagedorn, F. 2017b. Experimental soil warming shifts the fungal community composition at the alpine treeline. *New Phytologist* 215(2): 766– 778. DOI: http://dx.doi.org/10.1111/nph.14603.
- SPSS. 2017. Systat 13.2. Systat Software, Inc.
- Staddon PL, Fitter AH, Graves JD. 1999. Effect of elevated atmospheric CO₂ on mycorrhizal colonization, external mycorrhizal hyphal production and phosphorus inflow in Plantago lanceolata and Trifolium repens in association with the arbuscular mycorrhizal fungus Glomus mosseae. Global Change Biology 5(3): 347–358. DOI: http://dx.doi.org/10.1046/j. 1365-2486.1999.00230.x.
- Strickland, MS, Rousk, J. 2010. Considering fungal: bacterial dominance in soils Methods, controls, and ecosystem implications. Soil Biology and Biochemistry 42(9): 1385–1395.
- Talbot, JM, Yelle, DJ, Nowick, J, Treseder, KK. 2012. Litter decay rates are determined by lignin chemistry. *Biogeochemistry* **108**(1–3): 279–295. DOI: http://dx.doi.org/10.1007/s10533-011-9599-6.
- Teeri, T, Salovuori, I, Knowles, J. 1983. The molecular cloning of the major cellulase gene from *Trichoder-ma reesi*. *Nature Biotechnology* 1(8): 696–699. DOI: http://dx.doi.org/10.1038/nbt1083-696.
- Tiwari, S, Thakur, R, Shankar, J. 2015. Role of heat-shock proteins in cellular function and in the biology of fungi. *Biotechnology Research International* 2015: 132635–132635. DOI: http://dx.doi.org/10. 1155/2015/132635.
- Treseder, KK. 2004. A meta-analysis of mycorrhizal responses to nitrogen, phosphorus, and atmospheric CO₂ in field studies. New Phytologist 164(2): 347–355.
- Treseder, KK. 2005. Nutrient acquisition strategies of fungi and their relation to elevated atmospheric CO₂, in Dighton, J, Oudemans, P, White, J eds., *The* fungal community. 3rd ed. Boca Raton, FL: Marcel Dekker: 713–731.
- Treseder, KK. 2008. Nitrogen additions and microbial biomass: A meta-analysis of ecosystem studies. Ecology Letters 11(10): 1111–1120.
- Treseder, KK, Allen, EB, Egerton-Warburton, LM, Hart, MM, Klironomos, JN, Maherali, H, Tedersoo, L. 2018. Arbuscular mycorrhizal fungi as mediators of ecosystem responses to nitrogen deposition: A trait-based predictive framework. *Journal of Ecology* 106(2): 480–489. DOI: http://dx.doi.org/10.1111/ 1365-2745.12919.
- Treseder, KK, Allen, MF. 2002. Direct nitrogen and phosphorus limitation of arbuscular mycorrhizal fungi: A model and field test. New Phytologist 155(3): 507–515
- Treseder, KK, Balser, TC, Bradford, MA, Brodie EL, Dubinsky EA, Eviner, VT, Hofmockel, KS, Lennon,

- JT, Levine, UY, MacGregor, BJ, Pett-Ridge, J. 2012. Integrating microbial ecology into ecosystem models: Challenges and priorities. *Biogeochemistry* **109**: 7–18.
- Treseder, KK, Kivlin, SN, Hawkes, CV. 2011. Evolutionary trade-offs among decomposers determine responses to nitrogen enrichment. *Ecology Letters* 14: 933–938. DOI: http://dx.doi.org/10.1111/j. 1461-0248.2011.01650.x.
- Treseder, KK, Lennon, JT. 2015. Fungal traits that drive ecosystem dynamics on land. Microbiology and Molecular Biology Reviews 79: 243–262.
- Treseder, KK, Marusenko, Y, Romero-Olivares, AL, Maltz, MR. 2016. Experimental warming alters potential function of the fungal community in boreal forest. Global Change Biology 22(10): 3395—3404. DOI: http://dx.doi.org/10.1111/gcb.13238.
- Van Gestel, M, Merckx, R, Vlassak, K. 1993. Microbial biomass responses to soil drying and rewetting: The fate of fast- and slow-growing microorganisms in soils from different climates. Soil Biology and Biochemistry 25(1): 109–123. DOI: http://dx.doi.org/ 10.1016/0038-0717(93)90249-B.
- Velicer, GJ, Lenski, RE. 1999. Evolutionary trade-offs under conditions of resource abundance and scarcity: Experiments with bacteria. *Ecology* 80(4): 1168–1179.
- Versaw, WK, Metzenberg, RL. 1995. Repressible cationphosphate symporters in *Neurospora crassa*. *Proceedings of the National Academy of Sciences* 92(9): 3884–3887. DOI: http://dx.doi.org/10.1073/pnas. 92.9.3884.
- Vitousek, PM. 2004. Nutrient cycling and limitation: Hawai'i as a model system. Oxford, UK: Princeton University Press (Princeton Environmental Institute series).
- Vitousek, PM, Aber, JD, Howarth, RW, Likens, GE, Matson, PA, Schindler, DW, Schlesinger, WH, Tilman, DG. 1997. Human alteration of the global nitrogen cycle: Sources and consequences. *Ecological Applications* 7(3): 737–750. DOI: http://dx.doi.org/ 10.2307/2269431.
- Webb, CO, Ackerly, DD, Kembel, SW. 2008. Phylocom: Software for the analysis of phylogenetic community structure and trait evolution. *Bioinformatics* 24(18): 2098–100.
- Weber, CF, Zak, DR, Hungate, BA, Jackson, RB, Vilgalys, R, Evans, RD, Schadt, CW, Megonigal, JP, Kuske, CR. 2011. Responses of soil cellulolytic fungal communities to elevated atmospheric CO₂ are complex and variable across five ecosystems. *Environmental Microbiology* 13(10): 2778–2793. DOI: http://dx.doi.org/10.1111/j.1462-2920.2011. 02548.x.
- Wieder, WR, Grandy, AS, Kallenbach, CM, Taylor, PG, Bonan, GB. 2015. Representing life in the Earth system with soil microbial functional traits in the MIMICS model. Geoscientific Model Development 8(6): 1789–1808. DOI: http://dx.doi.org/10.5194/ gmd-8-1789-2015.

- Wiemken, A. 1990. Trehalose in yeast, stress protectant rather than reserve carbohydrate. Antonie Van Leeuwenhoek International Journal of General and Molecular Microbiology 58(3): 209–217. DOI: http://dx. doi.org/10.1007/bf00548935.
- Wilmes, P, Bond, PL. 2006. Metaproteomics: Studying functional gene expression in microbial ecosystems. *Trends in Microbiology* 14(2): 92–97. DOI: http://dx. doi.org/10.1016/j.tim.2005.12.006.
- Wood, JM, Bremer, E, Csonka, LN, Kraemer, R, Poolman, B, van der Heide, T, Smith, LT. 2001. Osmosensing and osmoregulatory compatible solute accumulation by bacteria. Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology 130(3): 437–460. DOI: http://dx.doi.org/10.1016/S1095-6433(01)00442-1.
- Zakrzewska, A, van Eikenhorst, G, Burggraaff, JEC, Vis, DJ, Hoefsloot H, Delneri, D, Oliver, SG, Brul, S, Smits, GJ. 2011. Genome-wide analysis of yeast stress survival and tolerance acquisition to analyze the central trade-off between growth rate and cellular robustness. *Molecular Biology of the Cell* 22(22): 4435–4446. DOI: http://dx.doi.org/10.1091/mbc. E10-08-0721.

- Zhang, K, Shi, Y, Jing, X, He, J-S, Sun, R, Yang, Y, Shade, A, Chu, H. 2016a. Effects of short-term warming and altered precipitation on soil microbial communities in alpine grassland of the Tibetan Plateau. Frontiers in Microbiology 7(1032). DOI: http://dx.doi.org/10. 3389/fmicb.2016.01032.
- Zhang, S, Zhang, X, Chang, C, Yuan, Z, Wang, T, Zhao, Y, Yang, X, Zhang, Y, La, G, Wu, K, Zhang, Z. 2016b. Improvement of tolerance to lead by filamentous fungus *Pleurotus ostreatus* HAU-2 and its oxidative responses. *Chemosphere* 150: 33–39. DOI: http://dx.doi.org/10.1016/j.chemosphere.2016.02.003.
- Zhao, Q, Jian, S, Nunan, N, Maestre, FT, Tedersoo, L, He, J, Wei, H, Tan, X, Shen, W. 2017. Altered precipitation seasonality impacts the dominant fungal but rare bacterial taxa in subtropical forest soils. *Biology and Fertility of Soils* 53(2): 231–245. DOI: http://dx.doi.org/10.1007/s00374-016-1171-z.
- Zhou, J, Lemos, B, Dopman, EB, Hartl, DL. 2011. Copy number variation: The balance between gene dosage and expression in *Drosophila* melanogaster. *Genome Biology and Evolution* 3: 1014–1024. DOI: http://dx.doi.org/10.1093/gbe/evr023.

How to cite this article: Treseder, KK, Alster, CJ, Anh Cat, L, Gorris, ME, Kuhn, AL, Lovero, KG, Hagedorn, F, Kerekes, JF, McHugh, TA, Solly, EF. 2021. Nutrient and stress tolerance traits linked to fungal responses to global change: Four case studies. *Elementa: Science of the Anthropocene* 9(1). DOI: https://doi.org/10.1525/elementa.2020.00144

Domain Editor-in-Chief: Detlev Helmig, Boulder AIR LLC, Boulder, CO, USA

Associate Editor: Daniel Liptzin, Soil Health Institute, Morrisville, NC, USA

Knowledge Domain: Atmospheric Science

Published: August 6, 2021 Accepted: July 16, 2021 Submitted: September 24, 2020

Copyright: © 2021 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See http://creativecommons.org/licenses/by/4.0/.

