MNRAS **505**, 4874–4889 (2021) Advance Access publication 2021 May 31

Evolution of Wolf–Rayet stars as black hole progenitors

E. R. Higgins,^{1*} A. A. C. Sander¹⁰,¹ J. S. Vink¹⁰ and R. Hirschi^{2,3}

¹Armagh Observatory and Planetarium, College Hill, Armagh BT61 9DG, N. Ireland

²Astrophysics Group, Keele University, Keele, Staffordshire ST5 5BG, UK

³Kavli Institute for the Physics and Mathematice of the Universe (WPI), University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa 277-8583, Japan

Accepted 2021 May 23. Received 2021 May 20; in original form 2021 March 26

ABSTRACT

Evolved Wolf–Rayet stars form a key aspect of massive star evolution, and their strong outflows determine their final fates. In this study, we calculate grids of stellar models for a wide range of initial masses at five metallicities (ranging from solar down to just 2 per cent solar). We compare a recent hydrodynamically consistent wind prescription with two earlier frequently used wind recipes in stellar evolution and population synthesis modelling, and we present the ranges of maximum final masses at core He-exhaustion for each wind prescription and metallicity Z. Our model grids reveal qualitative differences in mass-loss behaviour of the wind prescriptions in terms of 'convergence'. Using the prescription from Nugis & Lamers the maximum stellar black hole is found to converge to a value of 20–30 M_{\odot}, independent of host metallicity; however, when utilizing the new physically motivated prescription from Sander & Vink there is no convergence to a maximum black hole mass value. The final mass is simply larger for larger initial He-star mass, which implies that the upper black hole limit for He-stars below the pair-instability (Z_{PI}) to be as high as 50 per cent Z_{\odot}, corresponding to the host metallicity of the Large Magellanic Cloud. Moreover, while the Nugis & Lamers prescription would not predict any black holes above the approx 130 M_{\odot} pair-instability limit, with Sander & Vink winds included, we demonstrate a potential channel for very massive helium stars to form such massive black holes at \sim 2 per cent Z_{\odot} or below.

Key words: stars: black holes - stars: evolution - stars: massive - stars: mass-loss - stars: Wolf-Rayet.

1 INTRODUCTION

The most massive stars are central to many fields of Astrophysics, driving the evolution of galaxies, recycling fusion material, as well as enriching their surroundings through strong stellar winds. While they become increasingly rare with increasing mass, they are also the progenitors of the heaviest stellar mass black holes (BHs). Recently, the detection of merging BHs via gravitational waves (GWs) has queried the spectrum of BH masses for different galaxies, with massive BHs initially detected at 30–40 M_{\odot} by Abbott et al. (2016). More recently, the detection of an 85 M_{\odot} and 66 M_{\odot} BH merger in GW 190521 witnessed an interesting challenge for stellar evolution and GW theorists. The mass of each BH in this event appears to lie in the so-called 'pair instability' (PI) mass gap, where the star is thought to rip itself apart in a violent explosion or through pulsations in a 'pulsational pair-instability' (PPI) supernova. As such, Abbott et al. (2020) suggested this extreme GW event to be the product of second generation BHs. Stellar evolution studies such as Vink et al. (2020) and Farrell et al. (2021) have, however, hypothesized alternative solutions for creating first generation heavy BHs such as in GW 190521 by maintaining a low core mass and retaining the large H envelope, which might even be possible for metallicities up to 0.1 Z_{\odot} (Vink et al. 2020).

The various evolutionary paths a massive star may take on route to becoming a black hole include stars that are stripped of their envelope at the end of core H-burning towards a Helium (He) star. For a sufficiently high mass-loss rate (Sander, Vink & Hamann 2020; Shenar et al. 2020), such stars will appear as classical Wolf– Rayet (WR) stars. The star may have been stripped through binary interactions (e.g. Podsiadlowski, Joss & Hsu 1992) or as a result of strong stellar winds in the case of single massive stars (e.g. Yoon et al. 2012; Georgy et al. 2013). Massive He stars and WR stars are considered to be the final evolutionary stage before forming a BH, and as a result, WR mass loss is the key in establishing the BH progenitor mass range.

In spite of this, the levels of He star mass-loss rates have been debated in particular with regards to uncertainties in their mass- and metallicity-dependence. For lower-mass objects, which will likely not appear as WR stars (Vink 2017; Sander & Vink 2020), there is essentially an absence of observations already at solar metallicity ($Z = Z_{\odot}$) between hot subdwarfs (e.g. Schootemeijer et al. 2018; Wang, Gies & Peters 2018) and WR stars (e.g. Hamann et al. 2019; Sander et al. 2019) with the notable exception of HD 45166 (Groh, Oliveira & Steiner 2008). At low Z, the situation gets even worse due to the lack of observed WR stars below the regime of the Small Magellanic Cloud (SMC; $Z \approx 0.2 Z_{\odot}$). Given the need to estimate the mass loss of He stars in stellar evolution, population synthesis, or supernova statistics (e.g. Eldridge & Stanway 2009; Woosley, Sukhbold & Janka 2020), there is a high demand for theoretically

^{*} E-mail: erin.higgins@armagh.ac.uk

determined mass-loss rates of massive stars, in particular for He stars and WRs.

Besides directly determining the upper-mass limit of BHs, the mass-loss dependent evolution of He stars as a function of metallicity also has consequences for the fraction of type IIb and Ibc stripped supernovae (SNe) as stronger winds could remove more material from the outer layers of a star, thereby altering the parameters of the eventual core collapse event (e.g. Gilkis et al. 2019). All this of course has major implications for the field of GW astronomy and detections of future BH mergers.

Until 2005, those stellar evolution models that included a metallicity dependence of WR stars usually relied on the *total* metallicity (e.g. Nugis & Lamers 2000) that is dominated by self-enriched surface material (nitrogen in the case of WN stars, carbon for WC stars). In a pilot study of late-type WN and WC stars, Vink & de Koter (2005) showed that it is not the CNO abundances but the *initial* iron (Fe) abundance that is the key in setting the mass-loss metallicity dependence, enabling Eldridge & Vink (2006) to model the observed drop in the WC/WN ratio towards lower host galaxy metallicity, and to predict heavier BHs in lower Z host galaxies.

This initial Z, Fe-dependent mass-loss of WR stars was also included in the GRB progenitor evolution by Yoon, Langer & Norman (2006) as well as in the maximum BH population study of Belczynski et al. (2010), which was subsequently utilized to infer the low host Z of the first gravitational wave source GW 150914 (Abbott et al. 2016; Vink 2019). It is thus clear that in order to correctly predict the BH mass distribution with Z, we need to properly understand the intricacies of WR mass loss as a function of Z, appreciating there is no requirement for this dependence to be a simple power law. In fact, Sander & Vink (2020) recently showed a more complex behaviour, and it is our aim in the current paper to explore the implications of this new WR mass-loss formulations for the BH mass function with Z. We defer a more detailed study of WR evolution in terms of WN, WC, WO sub-types to the future.

In Section 2, we present our method of evolving He stars and outline our model grid. Mass-loss rates of He stars are discussed in Section 3, including the implementation of three comparative prescriptions in our evolution models. Results from our Standard Grid of models are presented in Section 4, while comparable results for our Alternative Grid of models are outlined in Section 5. A discussion of the upper mass limit of black holes in terms of PI is shown in Section 6, with conclusions outlined in Section 7.

2 STELLAR EVOLUTION MODELS

We utilized the stellar evolution code MESA (version 8845; Paxton et al. 2011, 2013, 2015) in calculating detailed models of massive stars, focusing on the core He-burning phase of evolution. We provide a range of initial helium Zero-Age-Main-Sequence (He-ZAMS) masses and metallicities. The range of initial masses comprises 20, 40, 60, 80, 100, 120, and 200 M_{\odot} with a comparable range of alternative models provided in Tables A3 and A4. The initial composition is based on tables provided by Grevesse & Sauval (1998) with Z_{\odot} = 0.014, and Y = 0.266. We also employed OPAL opacity tables from Rogers & Nayfonov (2002). We calculate models for five solar-scaled metallicities: Z_{\odot}, 0.5 Z_{\odot} [Large Magellanic Cloud (LMC)], 0.2 Z_{\odot} (SMC), 0.1 Z_{\bigcirc}, and 0.02 Z_{\bigcirc}, providing a range of low metallicity environments that may host the heaviest stellar mass black holes.

We adopt the standard mixing-length theory of convection by Böhm-Vitense (1958) with $\alpha_{MLT} = 1.5$, implementing the Ledoux criterion, allowing for semiconvective mixing which we include

here with an efficiency parameter of $\alpha_{sc} = 100$, favouring blueward evolution (see Schootemeijer et al. 2019; Higgins & Vink 2020). We consider additional mixing by convective core overshooting on the main sequence, but do not include overshooting of the Helium core.

2.1 Pre-Helium evolution

To create a grid of He star models, we evolve H-burning models towards WR stars via extreme mixing, which promotes bluewards evolution due to additional H dredged into the core. Rather than inducing rapid rotation, we employ an artificially large increase of the convective core by exponential overshooting in our Standard Grid of models presented in this work. We include core convective overshooting above the H-burning core with a diffusive method for values of f_{ov} up to 0.9. In principle this extreme mixing could be achieved in Nature by various paths, including strong winds, rapid rotation, and/or binary evolution. In order to aid evolution towards a He star stage without numerical complexities in the expanding stellar envelope we included the option of MLT⁺⁺ in MESA which increases the transport of convective energy in low-density envelopes.

Rotation was included with angular momentum transport and chemical mixing coefficients from Heger, Langer & Woosley (2000). In our Standard Grid of He star models, for all masses and metallicities the initial rotation rate is set to 20 per cent critical, whereas for our Alternative Grid (see Section 5) of quasi-chemical homogeneously evolved (QCHE) models we implement higher values of 60 per cent, with 40 per cent at the highest masses >120 M_{\odot} . While increased mixing by rotation promotes evolution towards the He main sequence, the core He-burning models have sufficiently spun down due to angular momentum loss by main sequence stellar winds such that the varied initial rotation rates are all reduced to \leq 150 km s⁻¹.

For our Standard Grid of He stars, the initial H-burning ZAMS models implement zero mass loss in order to create He star models that remain massive enough on the He ZAMS to explore the full range of initial masses at all Z. The standard mass-loss recipe enabled as part of the 'Dutch' wind routine by default in MESA (e.g. Paxton et al. 2015) would implement Vink, de Koter & Lamers (2001) rates during H-burning, followed by that of Nugis & Lamers (2000) which sets the mass-loss rate for hot stars with less than 40 per cent surface H. encompassing the WR regime. Note that the default implementation of the Nugis & Lamers (2000) recipe in MESA (used in this study) includes a Z-dependency in terms of the current metallicity. This leads to almost no Z-scaling for the NL00 mass-loss rates due to self-enrichment. Other studies (e.g. Yusof et al. 2013) consider the initial metallicity for the Z-dependency of Nugis & Lamers (2000; following Eldridge & Vink 2006), in line with the Z_{Fe} -dependence from Vink & de Koter (2005) that leads to a stronger Z-dependence of the Nugis & Lamers (2000) rates.

In this study, we do not discuss any explicit impact of different wind mass loss treatments before the He ZAMS, but instead focus on the impact of stellar winds after the stars have become He stars (cf. Section 3). Fig. 1 illustrates the structure evolution of a 40 M_{\odot} helium star at Z_{\odot} . The green hatched lines represent the convective core, with the helium abundance shown in blue by the colour bar. The model shown in Fig. 1 implements mass loss from Sander & Vink (2020), and gives a final mass of 17.8 M_{\odot} at core He-exhaustion.

2.2 Helium star evolution

In all cases our models evolve from core H exhaustion towards the He ZAMS position at log $T_{\rm eff} \sim 5.0$ and continue their evolution as a



Figure 1. Structure evolution (Kippenhahn) diagram of a 40 M_{\odot} Helium star at Z_{\odot} with mass-loss rates from Sander & Vink (2020). The colour bar denotes the Helium abundance and the green hatched region illustrates the convective core size.

Helium star until core He exhaustion where the evolution is stopped. Our models do not evolve to cooler $T_{\rm eff}$ towards RSG evolution after core H-burning given that the scope of this study is to test He-rich WR stars from the He ZAMS to He exhaustion. As outlined above, we do not initiate our models on the He ZAMS. Artificially altering the Helium abundance does not account for the by-products created during core H-burning (¹⁴N) and would not provide a comparable structure to evolved WR stars. Therefore, we provide two comparable methods of producing pure He stars from the H ZAMS to the He ZAMS.

We present our Standard Grid of models in Section 4 that omits stellar winds during the core H-burning stage of evolution in order to probe the entire mass range for Helium stars on the He ZAMS at all Z. We provide an Alternative Grid in Section 5 that does include stellar winds during core H-burning, highlighting an Alternative method of creating He star models through the addition of rapid rotation. For both grids of models we focus on the core He-burning stage of evolution where we test three wind recipes for WR stars. As a result, we do not probe the evolution during core H-burning but rather compare the differing methods of reaching the He ZAMS. We find that the conclusions are interchangeable, irrespective of which method of fully mixing H-burning stars towards pure He stars is implemented. The benefit of our Standard Grid is the ability to test the evolution of 50–200 M_{\odot} He stars at $Z_{\odot},$ whereas the Alternative Grid approach limits the derived He star mass range due to the inclusion of mass loss, providing insights into the possible realistic initial masses of He stars. This is severe for high metallicities and would limit our studies to He ZAMS masses of about $\sim 20 \ M_{\odot}$ at Z_{\odot} .

3 WOLF-RAYET MASS-LOSS RATES

In contrast to OB-type stars, the theory for classical WR winds is still in its infancy (e.g. Nugis & Lamers 2002; Gräfener & Hamann 2005; Gräfener et al. 2017; Grassitelli et al. 2018; Sander et al. 2020). Consequently, empirically determined mass-loss recipes are used in

stellar evolution models. The recently derived formulae for massive He stars winds by Sander & Vink (2020) mark the first theoretically rooted recipes in this field. In this study, we present a comparison of the most utilized recipes for WR mass loss alongside the new prescription from Sander & Vink (2020).

Based on empirical mass-loss rates obtained from radio fluxes with a non-trivial clumping correction (Nugis, Crowther & Willis 1998) and emission lines, Nugis & Lamers (2000) derived relations for the mass-loss rates of WN and WC stars including a combined recipe of the form $\dot{M}(L, Y, Z)$. The latter is commonly applied in stellar evolution codes including MESA (Paxton et al. 2013), GENEC (Ekström et al. 2012), FRANEC (Chieffi & Limongi 2013), STAREVOL (Martins & Palacios 2017), or PARSEC (Chen et al. 2015).

Another widely used recipe, commonly applied in population synthesis modelling (e.g. Belczynski et al. 2010), is a combination of a clumping-corrected empirical mass-loss relation of Hamann, Koesterke & Wessolowski (1995) with the iron-metallicity [Z_{Fe}dependent mass-loss scaling of Vink & de Koter (2005)]. Based on the tailored atmosphere analysis of a WN sample, Hamann et al. (1995) derived a recipe of the form $\dot{M}(L)$ that can be extended to the form $\dot{M}(X, L)$ when accounting for a fit of the hydrogen content with the mass loss. Although various later studies with more elaborated models were made (e.g. Crowther et al. 2002; Sander et al. 2014; Hainich et al. 2015), it is - with certain adjustments discussed below - still employed in most population synthesis models. Given the fact that the work of Hamann et al. (1995) was performed before the introduction of clumping and iron-line blanketing in model atmospheres - two major additions that considerably affected the derived stellar and wind parameters - the original values were actually overestimating the mass loss of the WR stars. Moreover, (Vink & de Koter 2005) obtained a clear metallicity-dependence of WR winds, which has to be taken into account. At solar metallicity, Yoon & Langer (2005) showed that scaling down the results from Hamann et al. (1995) by a factor of 10 would align them more with the recipe form Nugis & Lamers (2000). This scaling was then combined in Yoon et al. (2006) with the Z_{Fe} -scaling from Vink & de Koter (2005).

With these mentioned adjustments, both widely used recipes essentially give values of \dot{M} that are in the regime of observed WR stars in the Milky Way. However, this does not imply that their scalings are necessarily physically meaningful. When employing current stellar atmosphere models with prescribed wind velocity fields, there is considerable scatter in the obtained results (e.g. Tramper et al. 2015; Hamann et al. 2019; Shenar et al. 2019) with regressions yielding very different *L*-dependencies from those in Nugis & Lamers (2000). Mitigating these was the effort of Tramper, Sana & de Koter (2016), though the introduced dependencies are still empirically and not physically motivated, thus making extrapolations beyond the regime of observed objects highly questionable (Vink 2017).

More recently, Yoon (2017) compared the mass-loss rates of various WR types (e.g. WNE, WC, WO) derived from stellar atmosphere calculations (see Hamann, Gräfener & Liermann 2006; Sander, Hamann & Todt 2012; Hainich et al. 2014, 2015; Tramper et al. 2015) with theoretical wind recipes as implemented in stellar evolution codes. Yoon (2017) explored a range of metallicity, surface helium, and luminosity dependencies on WR mass-loss rates, correlated with populations of WNE stars and WC/WO stars from stellar atmosphere models, and the implications for populations of SNe progenitors. Finding a steep dependence on initial metallicity of WNE stars compared to WC/WO stars, Yoon (2017) combined the relation of Hainich et al. (2014) and Tramper et al. (2016) with

4877

the earlier results from Nugis & Lamers (2000) and an additional scaling factor motivated by the lower clumping used in Hamann et al. (2006). Beside this additional scaling, the resulting recipe essentially switched from Nugis & Lamers (2000) to Tramper et al. (2016) at Y = 0.9, thereby resulting in higher mass-loss rates for WC/WO stars.

The recipe suggested by Yoon (2017) was created to yield evolutionary tracks surpassing the observed WR luminosity ranges in the Milky Way and the LMC - in particular when making use of a scaling factor larger than unity. As such, it is neither directly based on empirically determined wind parameters, nor on underlying theoretical considerations. Its parameter dependence results from the underlying incorporated recipes, making it not well suited for extrapolations. The recipe by Yoon (2017) reflects the empirical finding that observed WC stars have higher mass-loss rates than WN stars with the same luminosity. However, this does not imply that a WN and WC star with the same L also have the same mass. In fact, Sander et al. (2020) could show with hydrodynamically consistent atmosphere modelling, that the mass loss of WC stars is lower than the one for WN stars with the same luminosity and mass (in the same host environment). Thus, the empirical result of higher massloss rates for WCs compared to WNs of the same luminosity and the same initial metallicity implies that these WC stars likely have a lower current mass than WN stars. In light of the results obtained by Sander et al. (2020), the increase of \dot{M} at the onset of the WC stage in a mass-loss recipe is questionable. In this effect, the Yoon (2017) recipe is similar to Nugis & Lamers (2000), despite the difference in individual coefficients. We thus refrain from testing a larger range of recipes in this work and only compare the new theoretical relation with the two most widely used recipes outlined in the following Section 3.1.

3.1 WR mass-loss prescriptions

In this paper, we study how much the two mainly used recipes are affecting the resulting final masses (cf. Section 4.2) of He stars and how these results change when using a physically motivated recipe based on hydrodynamically consistent stellar atmospheres. For the latter, we take the $\dot{M}(L)$ -recipe provided by Sander & Vink (2020; hereafter SV20).

$$\dot{M}_{\rm SV20} = \dot{M}_{10} \left(\log \frac{L}{L_0} \right)^{\alpha} \left(\frac{L}{10 \cdot L_0} \right)^{3/4} \tag{1}$$

with coefficients \dot{M}_{10} , L_0 and α provided for a range of metallicities in table 1. In this work, we are studying the imprint of mass loss on the He main sequence, where our stars essentially follow a clear M-L relation (cf. Section 3.3). Thus, we can implement the mass-loss recipe from Sander & Vink (2020) in the $\dot{M}(L, Z_{init})$ -form, i.e. as a function depending only on luminosity L and initial metallicity Z_{init} . This is further justified as stars that have lost all of their hydrogen layers will spend most of their remaining lifetime on the He main sequence. However, we stress that the treatment used in this work does not account for other evolutionary stages, e.g. stars beyond central He burning or stars that have retained part of their hydrogen envelope, as their M-L relation can be significantly different and the change of their temperature and chemical abundances would have to be taken into account. In this work, we use a dedicated set of input parameters given in Table 1 for the free parameters in equation (1). We apply these values in a MESA subroutine where we switch on this mass-loss recipe during the core He-burning phase. There have been minor updates in SV20 to the specified coefficients outlined in Table 1 for the finalized set of atmosphere models presented in SV20.

Table 1. Coefficients at various Z for $\dot{M}(L)$ by SV20. (Minor revisions have been made to the coefficients in the resulting formulae of Sander & Vink (2020); however, while we have tested the updated relations, the results are indistinguishable.).

$Z_{\rm init}/Z_{\odot}$	α	$\log L_0$	$\log \dot{M}_{10}$
1.0	1.301	5.043	-4.075
).5	1.327	5.335	- 3.812
).2	1.299	5.668	- 3.523
).1	1.165	5.906	- 3.331
0.02	0.938	6.478	-2.941

However, we tested multiple calculations to verify that these minor differences are indistinguishable in their model results. For future applications, we suggest to use the metallicity fit relations provided in SV20 that are also implemented in the publicly available PYTHON script.¹

In MESA, the standard mass-loss recipes are accumulated in the so-called 'Dutch' wind scheme, which employs Vink et al. (2001) for hot and H-rich stars, de Jager, Nieuwenhuijzen & van der Hucht (1988) for cool stars, and the aforementioned Nugis & Lamers (2000; hereafter NL00) for hot, H-depleted stars. NL00's implementation in MESA has the form

$$\log \dot{M}_{\rm NL00} = -11.00 + 1.29 \log L + 1.73 \log Y + 0.47 \log Z_{\rm cur}$$
(2)

where it should be noted that Z_{cur} here refers to the *total* current Z, i.e. it is enhanced due to self-enrichment of elements such as carbon, which is no longer considered to be the physically dominant component (Vink & de Koter 2005).

The final set of comparison models implements mass-loss rates as suggested by Yoon et al. (2006) where the recipe from Hamann et al. (1995) has been reduced by a factor of 10 in order to account for clumping. The resulting wind recipe implemented in stellar population models, often simply referred to as 'Hamann 95' (hereafter H95+, where the '+' represents the adjustments for clumping and Fe-scaling), thus reads

$$\log \dot{M}_{\rm H95+} = -12.95 + 1.5 \log L - 2.85 X + 0.85 \log (Z_{\rm init}/Z_{\odot}).$$
(3)

In summary, we evolve stellar models toward the He-ZAMS and probe the core He-burning evolution with three wind recipes:

(i) SV20: Physically motivated mass-loss description resulting from detailed sets of dynamically consistent atmosphere models by Sander & Vink (2020).

(ii) NL00: Standard recipe implemented in stellar evolution codes from Nugis & Lamers (2000), although accounting for total Z (self-enriched) rather than initial Fe-scaled Z_{init} .

(iii) H95+: Yoon et al. (2006) implementation as often used in Population Synthesis modelling. Rates have been extracted from Hamann et al. (1995), reduced by a factor 10 to account for clumping, and including the Fe-dependent scaling from Vink & de Koter (2005).

3.2 Comparison of wind recipes

The key points to consider regarding the impact of different massloss recipes are the absolute mass-loss rates and their Z-dependence including in particular the actual implementation for the latter. While

¹The script is available at https://armagh.space/asander



Figure 2. Mass-loss rate as a function of luminosity at the start of the He ZAMS with Z_{\odot} for each wind prescription considered in this work SV20 (red), H95+ (green), and NL00 (blue).

mass-loss recipes are written in the form of assigning a mass-loss rate to a given abundance, it is in reality the resulting opacity that drives the mass loss. Thus, depending on the ionization stage, not every element available at the surface will actually provide significant line opacity to drive a stellar wind. The works of Vink & de Koter (2005) and Gräfener & Hamann (2005) could show that decisive line opacities in the winds of classical WR stars stem from iron and not from other elements such as CNO despite their large abundances. Performing detailed locally consistent calculations, Sander et al. (2020) recently demonstrated that due to the deep launching of the winds, the ionization stage of many elements at the sonic point is simply too high to provide considerable line opacities, while the iron-group M-shell ions can still contribute to the total line opacity there. While the CNO abundance is increased due to self-enrichment, the abundance of the iron group elements are confined to their initial value, at least when neglecting external influences such as stellar mergers or the pollution due to the infall of a planetary companion. Consequently, the Z-scalings derived by Vink & de Koter (2005) and Sander & Vink (2020) should be performed with the initial Z (or simply $Z_{Fe}/Z_{Fe, \odot}$, i.e. the iron mass fraction relative to the solar value), not the current surface abundances. This is not the case in the recipe by Nugis & Lamers (2000) and while it was adopted by Yoon et al. (2006), this greater insight has not been universally adopted in the literature.

In the recent mass-loss recipe by Sander & Vink (2020), the Z (i.e. Z_{init}) dependence is physical, and shown to be more complex than that of a simple power law. This is illustrated in Figs 2 and 3 where we plot the $\dot{M}(L)$ -behaviour of the different recipes at the He ZAMS for solar and SMC metallicity. While both NL00 as well as H95+ use power-law scalings, the SV20 curves reflect the finding of Sander & Vink (2020), highlighting when a star has a sufficient *L/M*-ratio to reach WR-type winds, which is particularly noticeable in Fig. 3 depicting the SMC-like case. However, once the regime of dense WR winds is fully reached, the *Z*-dependence in SV20 is flatter than any other current \dot{M} -recipe with its mass loss changing only by a factor of two for an order of magnitude change in metallicity.



Figure 3. Mass-loss rate as a function of luminosity at the start of the He ZAMS with Z_{SMC} for each wind prescription as described in Fig. 2.

Focusing on the resulting range of BH masses, Woosley et al. (2020) investigated the effect of He star mass-loss when employing different recipes. While they did not calculate detailed structure models for each mass-loss recipe, the overall loss of mass during the He-burning stage could be approximated sufficiently by using previous work (Woosley 2019) yielding the luminosity as a function of mass and lifetime. The initial findings from Sander et al. (2020) - in particular the strong deviation from a power law in $\dot{M}(L/M)$ - were already taken into account in Woosley et al. (2020), but the underlying model sequence was small and the models were not tailored to represent (L, M)-combinations along He ZAMS as later done in Sander & Vink (2020). We avoid these obstacles (e.g. the limited ability to extrapolate the results from Sander et al. 2020) in our current work by employing the newer calculations from SV20. The imprint on the resulting BH masses between Woosley et al. (2020) and our work can be quite large, in particular for higher masses, where SV20 predict much higher mass-loss rates than what would be inferred from extrapolating the Sander et al. (2020) relation. This discrepancy is rooted in the underlying polynomial fit for \dot{M} in Sander et al. (2020), which eventually yields a local maximum in \dot{M} . Higher mass calculations in SV20 could instead show that such a maximum does not exist and a different mathematical description had to be employed for \dot{M} to avoid such an incorrect 'asymptotic behaviour'.

3.3 The mass-luminosity relation

In this study, we provide stellar evolution models with mass-loss rates based on Sander & Vink (2020). The mass-loss recipe in Sander & Vink (2020) was derived from a set of atmosphere models assuming an M-L relation for chemically homogeneous helium stars by Gräfener et al. (2011). To test whether this M-L relation is actually representative for our structure models on the He ZAMS, we compare our results of (L, M)-tuples with the relation from Gräfener et al. (2011) in the Appendix (see Fig. A1). The comparison highlights an excellent overall agreement between the M-L relation for He stars



Figure 4. Structure evolution of a 40 M_{\odot} star with each mass-loss recipe (solid line with starred markers represent SV20, dash-dotted lines with triangle markers represent NL00, and dashed lines with circle markers employ H95+ rates). Left: Models with Z_{\odot} . Right: Z_{SMC} models. The purple lines show the total mass (M_{T}), blue lines outline the convective core mass (M_{conv}), and red lines show the final CO core mass (M_{CO}) at He exhaustion. The *x*-axis time-scale illustrates the evolution from the He-ZAMS until core He-exhaustion.

with our models (blue triangles) and that of Gräfener et al. (2011; black solid line) for a broad range of masses and metallicities.

4 RESULTS FROM THE STANDARD GRID

For our Standard Grid of helium WR stars, we have calculated a wide range of models with initial masses 20–200 M_{\odot} , at Z_{\odot} , 0.5 Z_{\odot} , 0.2 Z_{\odot} , 0.1 Z_{\odot} , 0.02 Z_{\odot} (see Tables A1 and A2). We evolve H-burning models towards becoming He stars via extreme artificial convective mixing as outlined in Section 2. These He-ZAMS models have less than 1 per cent surface H at all Z and for all initial masses by the He-ZAMS. We now compare the evolution during the He-main sequence, and ultimately the final masses which may be utilized as a proxy for BH progenitor masses since the final evolutionary stages post-He burning are extremely short.

The final mass at core He-exhaustion changes drastically with decreasing Z for the different wind recipes. At Z_{\odot} , the highest final masses are provided by models with SV20 winds, with models implementing NL00 showing the lowest final masses. The situation is comparable at SMC-like Z, where SV20 models show the highest final masses, closely followed by those with H95+ winds, but NL00 models remain the lowest mass models at He-exhaustion. At the lowest Z, we find that H95+ models have higher final masses than models which include SV20 rates. This is due to the steeper L-dependencies of H95+, compared to the theoretical recipe by SV20.

4.1 Structure and abundances

We compare the structure evolution of a 40 M_{\odot} model, as in Fig. 1 for SV20, with the subsequent two recipes in Fig. 4, for Z_{\odot} (left) and Z_{SMC} (right). This figure highlights the change in final mass and core mass with Z and mass-loss rate. The total mass is shown in purple with the solid lines representing SV20 models, dash-dotted lines illustrating models with NL00, and dashed lines showing H95+ models. Convective core masses are also provided in blue, with final CO core masses shown in red. We show here that at Z_{\odot} (left), the final mass is relatively similar for all recipes, with a variation of 2–3 M_{\odot} . However, the situation shifts at lower Z, where even at Z_{SMC} (right) the change in final mass can be 8–10 M_{\odot} . The drop in final mass can be seen for the NL00 model (dash–dotted) compared with those of SV20 and H95+ which have a much more shallow decline in total mass during core He-burning.

Another interesting consequence of the different mass-loss treatments is the resulting abundance profile at He-exhaustion. Fig. 5 shows the chemical abundance profile as a function of the mass coordinate throughout the star (from core to surface) for the same models as in Fig. 4. ⁴He is represented by red lines, ¹²C is shown by blue lines, and ¹⁶O in purple. Once again, the clear trend in mass loss between the three different recipes is clearly visible. In particular, one can notice the strong stripping of outer layers with the NL00 formula even at Z_{SMC} (right-hand panel). Both at Z_{\odot} as well as at Z_{SMC} the dominant surface abundance at He exhaustion is carbon with the resulting value being almost independent of the mass-loss recipe and metallicity. However, this finding does not hold for other mass ranges. For our 60 M_{\odot} models, the stripping is stronger and oxygen can become the most abundant element at the surface. On the contrary, the stripping is weaker for the 20 M_{\odot} case, where WR-type mass-loss even breaks down completely at Z_{SMC} and we are left with a He-dominated atmosphere at all metallicities except Z_{\odot} .

Until core-He exhaustion, we never see a maximum of $X_C \approx 0.5$ at the surface as a re-occurring pattern in all of our abundance profiles. When progressing inside, the carbon fraction is getting larger while the He fraction is shrinking. After X_C reaches a maximum, it declines further inwards and oxygen becomes the dominant element with relative mass fractions of around 0.8 with only minor spreads due to mass and mass-loss treatment. Consequently, we would not expect to observe WC and WO stars with $X_C > 0.5$ at their surfaces. In their empirical analyses of WC and WO stars, both Sander et al. (2012) and Tramper et al. (2015) get fractions of up to 0.62. This



Figure 5. Final abundance profile of a 40 M_{\odot} star at core He-exhaustion with each mass-loss recipe (solid lines represent SV20, dashed lines represent NL00, and dash-dotted lines show H95+ models). Left: Models with Z_{\odot} . Right: Z_{SMC} models. The red lines show ⁴He abundances, blue lines illustrate ¹²C, and purple lines show ¹⁶O. The *x*-axis illustrates the mass co-ordinate from core to surface at core He-exhaustion.

might very well be within the uncertainties of the present studies, but a more in-depth look in future studies with both structure and evolution models could yield important constraints on the parameters and precise evolutionary stage of known WC and WO stars.

For our 40 M_{\odot} models, the second most abundant element at the surface at Z_{\odot} is oxygen with mass fractions between 25 per cent and ~ 40 per cent. From both the temperatures as well as the surface abundance, we expect such stars to have a WO-type spectrum. Neglecting the solution resulting from the overestimation of mass loss by NL00, the situation at Z_{SMC} is quite different: The oxygen abundance is lower here and potentially even below the remaining amount of helium at the surface. While these fractions could probably still result in a WO-type spectrum, the different O/He-ratios could potentially provide an interesting albeit indirect metallicity constraint for observed WO stars. The current observational constraints (Tramper et al. 2015; Shenar et al. 2016) do not contradict this scenario, but they also do not provide clear evidence. This is not surprising as the O/He-ratio at the surface also depends on the mass regime with higher mass models also showing larger oxygen fractions at the surface. Thus, a certain ratio could either be reached by a more massive star at lower metallicity or a less massive star at higher metallicity. Still, if additional constraints are available which could break this degeneracy, the surface abundances could potentially provide important indirect diagnostics to otherwise inaccessible pieces of information.

The overall evolution of our model sample in the Hertzsprung– Russell diagram (HRD) is highlighted in Figs 6 and 7 where we show representative models for 20, 40, and 60 M_{\odot} at Z_{\odot} (left) and Z_{SMC} (right), with mass-loss recipes from SV20 (red solid lines), NL00 (blue dash–dotted lines), and H95+ (green dashed lines). Additional representative models for Z_{\odot} and Z_{SMC} are shown in Figs A2 and A3 with initial masses of 40, 80, and 120 M_{\odot}. The drop in luminosity seen in the blue tracks (NL00 mass loss) reflects the stronger winds applied in this recipe compared to the red and green tracks. At low *Z*, models that implement the new SV20 mass loss may reach the 'breakdown regime' at lower masses as the luminosity there is not sufficient to support the optically thick winds. This is evident in the qualitatively different shape of the 20 M_{\odot} -track at Z_{SMC} . Without an optically thick wind, the model loses less than a solar mass throughout the whole He-burning lifetime. Consequentially, the star hardly changes its position in the HRD until the core contracts and this leads to an increase in the surface luminosity.

4.2 Final masses at core Helium exhaustion

The final mass at the end of core He-burning provides a much needed proxy for BH progenitor masses as the remaining nuclear burning phases are exponentially shorter than for H or He-burning. We probe the variety of final masses for our grid of WR models as a function of decreasing Z to establish the upper stellar BH mass limit, before considering additional reductions due to (P)PI. By further comparing the most commonly implemented stellar wind prescriptions for hot, helium stars, we also provide a direct contrast which considers the ramifications of implementing each recipe in stellar evolution.

In particular, when studying a wide range of initial masses across all Z, we find that the maximum final mass for models with NL00 winds is ~ 30 M_☉, with the exception of the highest initial mass model at {1/50th Z_☉. This means that with the standard 'Dutch' wind recipe implemented in MESA (also included in many other stellar evolution codes) one would need to consider very massive Helium stars (i.e. above 200 M_☉) in order to reach a final compact object mass above 30 M_☉, or limit heavy BH progenitors to essentially primordial Z. In other words, we find that the high mass-loss rates for He stars even at low Z due to the – physically incorrect – self-enrichment of the NL00 recipe has a tremendous impact on the evolutionary channels for GW progenitors as it prevents the formation of heavier black holes from He stars. In contrast, the H95+ prescription from Yoon et al. (2006) provides a similar range of final masses to that of SV20.

Fig. 8 depicts the range of final masses at different metallicities Z, highlighting the upper mass limits for each wind recipe. The





Figure 6. Hertzsprung–Russell diagram of 20–60 M_{\odot} He-burning stars evolving from the He-ZAMS as WRs, calculated at Z_{\odot} . Red solid tracks represent models that employ SV20 mass loss, those with NL00 rates are shown in blue dash–dotted lines, and green dashed lines illustrate models that include H95+ mass loss.

solid red line highlights the maximum final mass for SV20 models at all Z, with the same shown for NL00 models in the dashed blue line, and for H95+ models in the dash-dotted green line. The coloured symbols represent the various metallicities, with triangles representing SV20 models, H95+ models with circles, and stars for models implementing NL00 rates. At Z_{\odot} , we find that the most massive final mass is provided by SV20 models, with much lower final masses from both NL00 and H95+. However, at lower Z ($\sim Z_{SMC}$), the situation begins to shift towards similar final masses with H95+ and SV20 due to the different Z-dependence of both recipes. On the other hand, NL00 models remain below 35 M_{\odot} at all Z above $0.02 Z_{\odot}$. For models at (and below) 0.1 Z_{\odot} , the maximum

Figure 7. Same as Fig. 6, but for models calculated at Z_{SMC} . At this metallicity, the SV20 description predicts no strong wind mass loss for the 20 M_{\odot} model, thus yielding a completely different behaviour in the HRD.

final masses are now provided by H95+, while the SV20 models remain much more massive than those of NL00.

Similarly, we provide the range of final masses as a function of initial He ZAMS mass in Fig. 9. SV20 models are shown in solid lines, with NL00 in dash-dotted lines, and H95+ in dashed lines. We illustrate that for decreasing Z, the final mass of each set of models increases due to Z-dependent winds. The dash-dotted lines of NL00 do not inherently increase as in the SV20 or H95+ models due to the stronger mass loss caused by self-enrichment during core He-burning. Once again we notice that the NL00 recipe yields fundamentally different behaviour for the final masses at lower Z than SV20 and H95+.



Figure 8. Final mass at core He-exhaustion as a function of Z for each mass-loss recipe. The most massive final masses for each recipe are shown by the solid line in red for SV20, NL00 with a dashed blue line, and H95+ with the dash-dotted green line. Coloured markers denote the varied Z for each set of models, with triangles denoting SV20 models, circles for H95+ models, and finally stars for NL00 models.

So far, we did not take PI effects into account, but simply illustrated the consequences of extrapolating empirical mass-loss formulae without a deeper physical motivation. We now highlight the additional limitations due to PI in Figs 10, 12, and 13, where each model's final CO core mass is shown as a function of initial He mass (SV20 shown in red, NL00 in blue, and H95+ in green). These figures demonstrate that some high mass models in our grid enter the PPI regime where the CO core mass is above the estimated 40 M_{\odot} limit for stripped stars (Woosley 2017; Farmer et al. 2019). We further discuss the implication for the resulting BH mass limit in Section 6.

5 RESULTS FROM QCHE ALTERNATIVE GRID

As an alternative approach to our standard He star grid, we have calculated another grid of models incorporating a different evolutionary path towards He stars. This 'Alternative Grid' consists of rotating models with 60 per cent critical rotation which evolve towards becoming He stars during the MS by increased mixing. Since the critical rotation rate scales with increased mass, models that are greater than 150 M_{\odot} have a reduced rotation rate of 40 per cent critical. Stellar winds are included during the MS, and as a result the stars spin-down towards $\sim 150~{\rm km\,s^{-1}}$ by the onset of core He-burning such that models remain fully mixed but are no longer rapidly rotating. Our Alternative Grid mimics a channel of QCHE, which has been suggested for some WR stars (e.g. Martins et al. 2009; Hainich et al. 2015).

In our Alternative Grid, the mass loss during core H-burning only uses the (Vink et al. 2001) description and does not depend on the He star wind recipe. To ensure this, we removed the common, but physically questionable switch to a WR-wind recipe based on the hydrogen surface abundance. Since we account for main-sequence mass loss in this method, the masses of the models on the He ZAMS are lower than the initial masses of the models. Although our treatment of core H-burning can only act as a proxy for the actual processes happening in this stage, the overall loss of masses provides us with some insights about the different range of masses to be expected on the He ZAMS and the restrictions for the final masses





Figure 9. Final mass as a function of initial He ZAMS mass for each Z and mass-loss recipe (SV20 in solid lines, H95+ in dashed lines, NL00 in dash-dotted lines).



Figure 10. Final CO core mass as a function of He ZAMS mass for each mass-loss recipe (SV20 in red triangles, NL00 in blue stars, H95+ in green circles). The shaded regions correspond to mass limits where above a certain mass, larger CO core masses lead to PPI supernovae and PI supernovae respectively.

of black hole progenitors. Although we start with initial masses up to 170 M_{\odot} in our Alternative Grid, we only reach $\sim 20 M_{\odot}$ on the He ZAMS at Z_{\odot} with final masses of black hole progenitors on the order of $\sim 10 M_{\odot}$. In our stellar evolution models for $0.2 Z_{\odot}$ (SMC), the highest ZAMS masses yield a He star of $\approx 50 M_{\odot}$ with a final mass of 32 M_{\odot} when applying the SV20 and H95+ wind recipes, while the NL00 recipe leads to a much lower $M_{\rm f}$ of 21.5 M_{\odot} . This effectively means a shift by about 10 M_{\odot} in the range of reasonably expected BH masses at SMC metallicity. The comparison between all three wind recipes alters in the lowest metallicity environments due to the



Figure 11. Maximum final mass for the Alternative Grid as a function of log Z/Z_{\odot} for all wind recipes, where the solid red line illustrates the maximum mass for the Sander & Vink (2020) models. Red triangles correspond to models calculated with SV20 winds, green triangles with H95+, and blue triangles with NL00. Implications of the Alternative Grid on the maximum BH mass across *Z* with prior evolution taken into consideration.

 $\dot{M}(Z)$ relations, with the highest resulting masses obtained in H95+ followed by SV20 and NL00, the latter two being separated by a large margin. For 2 per cent Z_{\odot} , H95+ models provide the highest masses of 120 M_{\odot} by core helium exhaustion, while models with the SV20 recipe have final masses up to 100 M_{\odot} for the same initial H ZAMS mass of 170 M_o. Comparatively, models with the NL00 prescription result in a much lower $M_{
m f} \sim 28~{
m M}_{\odot}$ leading to a change in final mass of $\Delta M \approx 92 M_{\odot}$ compared to that of H95+. The trend in final masses deduced from our alternative method is comparable to the results from our standard method described in Section 4, suggesting robust results irrespective of which method is used in creating He star models. The Alternative Grid effectively leads to a finer spacing of initial He masses at higher metallicities, giving us some additional insights. As previously seen in Section 4.2 for the Standard Grid, we see that models which implement NL00 yield final masses no greater than 30 M_{\odot} for $Z \ge 2$ per cent Z_{\odot}.

Our findings are summarized in Fig. 11, where we depict the maximum final mass formed from our alternative models as a function of the host metallicity, indicating the results using the SV20 recipe with a red line. In all cases, the highest final masses correspond to models with the highest initial He masses.

6 DISCUSSION ON PI

The upper mass limit of stellar mass black holes below the PI gap is a key issue for theorists that has multiple uncertainties from massive star evolution, and scarce observational constraints. As the last evolutionary stage before core collapse with a time scale where winds can remove a considerable amount of mass from the star, the mass loss of He stars provides a crucial ingredient that could determine the upper BH mass limit. If indeed, the extent of these WR winds is responsible for setting the upper BH mass limit, then each recipe must be tested for convergence to the point where a maximum



Figure 12. Final CO masses at core He-exhaustion as a function of initial metallicity for each wind recipe (SV20 in red triangles, NL00 in blue stars, H95+ in green circles). The maximum final CO core mass (CO_{max}) models, for each metallicity, which include mass-loss rates from SV20 are represented by solid red lines, H95+ in dash–dotted green lines, and NL00 in dashed blue lines. The grey shaded region highlights the region where PPI supernovae occur, with CO core masses above 40 M_{\odot} .

final mass is reached irrespective of increasing initial mass. In this work, we tested the range of masses produced at core He-exhaustion (see Fig. 8) finding a critical mass limit for all Z above 2 per cent Z_{\odot} , i.e. irrespective of initial mass, with the NL00 recipe. Interestingly, this is not the case for the SV20 and H95+ models, which seem to have increased final masses with increased initial masses at all Z. This means that if WR winds were most closely represented by the NL00 prescription, then the upper BH mass limit would indeed be set by WR mass loss.

Since there is no convergence on the upper mass limit by SV20 and H95+ models, such that we may have much higher final masses if extremely high initial masses are invoked, we also must consider other dependencies. Fig. 9 highlights the correlation of initial He ZAMS mass with the final BH mass, particularly for the SV20 and H95+ models (solid and dashed lines, respectively), suggesting that the upper BH mass limit could be set by the maximum initial He ZAMS mass at each Z. When calculating our grid of alternative models, we found that due to strong mass loss on the MS, models with $\sim Z_{\odot}$ have reduced He ZAMS masses compared to even Z_{SMC} models. We excluded this effect of the prior evolution on core Heburning in our Standard Grid of models by excluding H-burning mass loss, in order to probe the full WR mass range across all Z. Still, it is important to note that the MS evolution is key in setting the initial mass of He and WR stars - directly impacting the final mass range. If the initial ZAMS mass is not high enough, and mass loss is strong enough on the MS, then our final BH mass limit will be dominated by the prior evolution on the MS. We do not explore the complete picture of pre-WR evolution in this work, as this in itself contains many uncertainties (such as treatment of internal mixing, proximity to the Eddington limit at the highest initial masses, and the extent of MS mass loss). However, we can illustrate the importance of the MS evolution through results from our Alternative Grid (Section 5),



Figure 13. Final CO masses at core He-exhaustion as a function of initial metallicity for each wind recipe. SV20 are represented by solid red lines, H95+ in dash-dotted green lines, and NL00 in dashed blue lines. The maximum final mass (M_f) models, for each metallicity, which include mass-loss rates from SV20, are represented by solid light red lines, H95+ in dash-dotted light green lines and NL00 in dashed light blue lines. The grey shaded region highlights the region where PPI pair-instability supernovae occur, with CO core masses above 40 M_{\odot}.

where we include the QCHE method as a proxy for evolutionary channels towards WRs. Ultimately a combination of mixing and mass loss will provide various channels towards He stars, and our results show that this can lead to lowered initial He masses, and as a result lower final masses at high $Z (\sim 0.2-1 Z_{\odot})$.

If we consider a wide range of initial He ZAMS masses, but with the realization that WR winds may not set the upper BH mass limit (as hinted by the recent SV20 results), and the initial mass may drive the maximum final mass up to extremely high final masses, PI would introduce an additional cut-off for the resulting BH masses. Stars above a certain CO core mass do not collapse to form BHs at the end of their lives – except for a range of very massive stars with M_{He} above ~133 M_{\odot} (Heger & Woosley 2002) – but rather rip themselves apart in PI supernovae leaving no remnant behind.

In the lower mass range of this regime PPI occurs where large eruptions from pulsations lead to the loss of a significant amount of mass. This leads to a so-called PI gap in the resulting BH mass range. This region could be located at varying mass ranges for different Z, directly impacted by the mass loss history of the stars during H and He-burning (see Fig. 12). Studies by Woosley (2017) and Farmer et al. (2019) propose that the mass range above which PPI occurs is driven by the CO core mass. In fact, stars which have a CO core above $\sim 40 \text{ M}_{\odot}$ are predicted to enter the PPI gap, (Woosley 2019; Woosley et al. 2020). In order to have increased final masses which do not enter the PPI regime, stars should have small convective cores. Vink et al. (2020) shows that by maintaining a small core with low convective core overshooting ($f_{ov} = 0.01$), and retaining a large H envelope, the effective PI gap can be much smaller than commonly anticipated as even massive BHs up to about 90 Mo, such as the one seen in the GW event GW190521, may be formed.

In this study, we endeavour to calculate grids of WR models through various methods, mainly by increased internal mixing

through convection (Standard Grid) and rotation (Alternative Grid). Higher internal mixing leads to increased convective core masses, and as a result will push our WR models towards PPI at lower masses, since the core masses in our models are relatively high, compared to models with standard/lowered mixing (e.g. $\alpha_{ov} \sim 0.1$ -0.5, or < 40 per cent critical rotation). Fig. 13 highlights the small difference between the CO core mass and total final mass (e.g. see change from red line to pink line, or light blue line to blue line) for all Z and WR mass-loss prescriptions. At Z_{\odot} and Z_{LMC} , most of our calculations predict BHs of varying mass depending on the wind prescription. This situation alters at lower Z, where even at SMC, more PPISN are predicted, and for lower initial masses - from 200 M_{\odot} down to 100 M_{\odot} for SV20 and H95+ models. At 2–10 per cent Z_{\odot} , PI already occurs for 60 M_{\odot} He-star models and higher, again with rates from SV20 and H95+. NL00 models only enter PI at 2 per cent Z_{\odot} with initial He masses ${\geq}60~M_{\odot}.$

The critical upper Z limit for the occurrence of PI, Z_{PI} , also sets the limit of maximum BH masses and seems to be in the 10– 50 per cent Z_{\odot} region. The higher Z models are set by the initial He mass, dominated by prior evolution on the MS. At the lowest Z (2–10 per cent Z_{\odot}) all three recipes predict PPISN, but only SV20 and H95+ models predict PPISN at $0.02 < Z/Z_{\odot} < 0.5$, giving a much wider Z range. Interestingly, the observed Z_{PI} could be lower than the derived Z_{PI} due to prior evolution limiting the range of He star masses. Based on our calculations approximating prior evolution, we would expect to have the observed Z_{PI} at $Z \le 0.1 Z_{\odot}$. These results will have consequences for observations of PPISN/PISN and the spectrum of BH masses at varied Z.

7 CONCLUSIONS

In this study we have presented two grids of stellar evolution models that incorporate the mass-loss description from Sander & Vink (2020) based on dynamically consistent atmosphere calculations of massive helium star winds. Our evolution models span a wide range of initial masses and metallicities, focusing on the core He-burning phase. As a comparison we have calculated models with standard hot, helium star winds as implemented in many stellar evolution codes from Nugis & Lamers (2000), as well as models that include Yoon et al. (2006) rates commonly implemented in population synthesis codes (e.g. Belczynski et al. 2010). The change in final mass as a function of wind recipe highlights the importance of implementing proper descriptions of wind mass loss in stellar evolution codes. By applying the first theoretically derived mass-loss description for classical WR stars (SV20), we included a qualitative and quantitative improvement to stellar evolution models of He stars that also provides an insight into the range of black hole masses at various metallicities. This qualitative difference in the treatment of WR winds can be observed in the behaviour of the upper BH mass limit at each Z. Our results uncover the convergence of a maximum BH mass with the NL00 prescription, whereas there is no convergence on the final mass for models implementing the SV20 description. As a result, our grids illustrate that if there is no convergence based on wind recipes then the prior evolution or PI is responsible for limiting the upper BH mass.

At solar metallicity, the change in the final mass due to implementing the SV20 description is generally limited. None the less, the SV20 description allows for higher final masses at Z_{\odot} than other descriptions do. Due to the more complex shape of the description by SV20, this is not true at the lowest Z-range, where the H95+ description would allow slightly higher final masses than SV20. At all Z, models that implement the NL00 recipe have final masses below \sim 30 $M_{\odot},$ except for the highest masses at 0.02 Z_{\odot} shown in Section 4. This means that a 40 M_{\odot} BH could not be formed from a WR star using the standard 'Dutch' wind recipe as included in MESA, unless an extremely low Z is enforced. In contrast, we find that with the SV20 mass loss a 40 M_{\odot} BH may be formed in the range 0.1–0.2 Z_{\odot} , with final masses up to 56 M_{\odot} at 0.1 Z_{\odot} . While we did not continue our calculations after encountering PI, we do find interesting limitations. Our results highlight that the NL00 wind prescription leads to no stellar BHs formed above the second BH mass gap, unlike with SV20. Extremely low metallicity environments such as I Zw 18 ($\sim 0.02~Z_{\odot})$ provide an insight into the early Universe and likely host some of the heaviest stellar mass black holes that may be detected as gravitational wave sources (Vink et al. 2020). Calculations with the SV20 description show final masses of up to 140 M_{\odot} at 0.02 $Z_{\odot},$ hinting that there might be a channel for very massive He stars to form black holes above the second BH mass gap, though the most massive BHs may form from H-rich stars.

We have analysed the dependencies on the critical mass limit of BHs below the PI gap for a range of Z. Our calculations show that only the standard, self-enriched NL00 mass-loss rates really yield a limit due to WR mass loss, in this case at 20–30 M_{\odot} , while the new physically-motivated SV20 rates, and H95+ models do not converge towards such a critical mass limit. Thus, the initial mass of the He star is the determining factor. This value is both influenced by the prior evolution on the MS (including the H ZAMS mass) and the effects of mass loss before reaching the He ZAMS. Eventually, PPI effectively avoids the formation of BHs above a certain CO core mass limit. Interestingly, the upper BH mass for each Z can also be used as a proxy for the Z_{PI}, where PI supernovae may be observed. With the NL00 mass loss prescription, only models at (and below) 2 per cent Z_{\odot} eventually enter the PI regime. This Z_{PI} limit is increased by models with the SV20 and H95+ descriptions, reaching this PI regime already at 10–50 per cent Z_{\odot} , depending on what we consider as a reasonable upper limit for the He star mass. Our results have consequences for the observations of (P)PISNe, which could already occur at considerably higher metallicities than previously assumed, and the mass spectrum of observed BHs as a function of Z.

ACKNOWLEDGEMENTS

We would like to thank the MESA developers for making their stellar evolution code publicly available, and the anonymous referee for their constructive comments. JSV and ERH are supported by STFC funding under grant number ST/R000565/1 in the context of the BRIDGCE UK Network. AACS is Öpik Research Fellow at Armagh Observatory & Planetarium. This article is based upon work from the 'ChETEC' COST Action (CA16117), supported by COST (European Cooperation in Science and Technology)'. RH acknowledges support from the IReNA AccelNet Network of Networks, supported by the National Science Foundation under Grant No. OISE-1927130 and from the World Premier International Research Centre Initiative (WPI Initiative), MEXT, Japan. The authors would like to thank Sarah Lightfoot for preliminary test models during her Ogden Trust Summer internship at Keele in 2019.

DATA AVAILABILITY

The data underlying this article will be shared on reasonable request to the corresponding author.

REFERENCES

- Abbott B. P. et al., 2016, Phys. Rev. Lett., 116, 061102
- Abbott R. et al., 2020, Phys. Rev. Lett., 125, 101102
- Belczynski K., Dominik M., Bulik T., O'Shaughnessy R., Fryer C., Holz D. E., 2010, ApJ, 715, L138
- Böhm-Vitense E., 1958, Z. Astrophysik, 46, 108
- Chen Y., Bressan A., Girardi L., Marigo P., Kong X., Lanza A., 2015, MNRAS, 452, 1068
- Chieffi A., Limongi M., 2013, ApJ, 764, 21
- Crowther P. A., Dessart L., Hillier D. J., Abbott J. B., Fullerton A. W., 2002, A&A, 392, 653
- de Jager C., Nieuwenhuijzen H., van der Hucht K. A., 1988, A&AS, 72, 259
- Ekström S. et al., 2012, A&A, 537, A146
- Eldridge J. J., Stanway E. R., 2009, MNRAS, 400, 1019
- Eldridge J. J., Vink J. S., 2006, A&A, 452, 295
- Farmer R., Renzo M., de Mink S. E., Marchant P., Justham S., 2019, ApJ, 887, 53
- Farrell E., Groh J. H., Hirschi R., Murphy L., Kaiser E., Ekström S., Georgy C., Meynet G., 2021, MNRAS, 502, L40
- Georgy C. et al., 2013, A&A, 558, A103
- Gilkis A., Vink J. S., Eldridge J. J., Tout C. A., 2019, MNRAS, 486, 4451
- Gräfener G., Hamann W. R., 2005, A&A, 432, 633
- Gräfener G., Vink J. S., de Koter A., Langer N., 2011, A&A, 535, A56
- Gräfener G., Owocki S. P., Grassitelli L., Langer N., 2017, A&A, 608, A34
- Grassitelli L., Langer N., Grin N. J., Mackey J., Bestenlehner J. M., Gräfener G., 2018, A&A, 614, A86
- Grevesse N., Sauval A. J., 1998, Space Sci. Rev., 85, 161
- Groh J. H., Oliveira A. S., Steiner J. E., 2008, A&A, 485, 245
- Hainich R. et al., 2014, A&A, 565, A27
- Hainich R., Pasemann D., Todt H., Shenar T., Sander A., Hamann W. R., 2015, A&A, 581, A21
- Hamann W. R., Koesterke L., Wessolowski U., 1995, A&A, 299, 151
- Hamann W. R., Gräfener G., Liermann A., 2006, A&A, 457, 1015
- Hamann W. R. et al., 2019, A&A, 625, A57
- Heger A., Woosley S. E., 2002, ApJ, 567, 532
- Heger A., Langer N., Woosley S. E., 2000, ApJ, 528, 368
- Higgins E. R., Vink J. S., 2020, A&A, 635, A175
- Martins F., Palacios A., 2017, A&A, 598, A56
- Martins F., Hillier D. J., Bouret J. C., Depagne E., Foellmi C., Marchenko S., Moffat A. F., 2009, A&A, 495, 257
- Nugis T., Lamers H. J. G. L. M., 2000, A&A, 360, 227
- Nugis T., Lamers H. J. G. L. M., 2002, A&A, 389, 162
- Nugis T., Crowther P. A., Willis A. J., 1998, A&A, 333, 956
- Paxton B., Bildsten L., Dotter A., Herwig F., Lesaffre P., Timmes F., 2011, ApJS, 192, 3
- Paxton B. et al., 2013, ApJS, 208, 4
- Paxton B. et al., 2015, ApJS, 220, 15
- Podsiadlowski P., Joss P. C., Hsu J. J. L., 1992, ApJ, 391, 246
- Rogers F., Nayfonov A., 2002, ApJ, 576, 1064
- Sander A. A. C., Vink J. S., 2020, MNRAS, 499, 873
- Sander A., Hamann W. R., Todt H., 2012, A&A, 540, A144
- Sander A., Todt H., Hainich R., Hamann W. R., 2014, A&A, 563, A89
- Sander A. A., Hamann W.-R., Todt H., Hainich R., Shenar T., Ramachandran V., Oskinova L. M., 2019, A&A, 621, A92
- Sander A. A. C., Vink J. S., Hamann W. R., 2020, MNRAS, 491, 4406
- Schootemeijer A., Götberg Y., de Mink S. E., Gies D., Zapartas E., 2018, A&A, 615, A30

- Schootemeijer A., Langer N., Grin N. J., Wang C., 2019, A&A, 625, A132
- Shenar T. et al., 2016, A&A, 591, A22 Shenar T. et al., 2019, A&A, 627, A151
- Shenar T., Gilkis A., Vink J. S., Sana H., Sander A. A. C., 2020, A&A, 634, A79
- Tramper F. et al., 2015, A&A, 581, A110
- Tramper F., Sana H., de Koter A., 2016, ApJ, 833, 133
- Vink J. S., 2017, A&A, 607, L8
- Vink J. S., 2019, IAU Symp., 346, 444
- Vink J. S., de Koter A., 2005, A&A, 442, 587
- Vink J. S., de Koter A., Lamers H. J. G. L. M., 2001, A&A, 369, 574
- Vink J. S., Higgins E. R., Sander A. A. C., Sabhahit G. N., 2020, MN-RAS, 504, 146
- Wang L., Gies D. R., Peters G. J., 2018, ApJ, 853, 156
- Woosley S. E., 2017, ApJ, 836, 244
- Woosley S. E., 2019, ApJ, 878, 49
- Woosley S. E., Sukhbold T., Janka H. T., 2020, ApJ, 896, 56
- Yoon S. C., Langer N., 2005, A&A, 443, 643
- Yoon S. C., Langer N., Norman C., 2006, A&A, 460, 199
- Yoon S. C., Gräfener G., Vink J. S., Kozyreva A., Izzard R. G., 2012, A&A, 544, L11
- Yoon S. C., 2017, MNRAS, 470, 3970
- Yusof N. et al., 2013, MNRAS, 433, 1114

APPENDIX A: TABLES OF MODEL GRIDS AND ADDITIONAL FIGURES



Figure A1. Mass–luminosity relation for He ZAMS models with SV20 mass-loss rates (blue triangles), with a comparison of the M-L relation derived from G+11 for pure helium stars (black solid line).





burning stars d solid tracks L00 rates are e models that

Downloaded from https://academic.oup.com/mnras/article/505/4/4874/6289925 by Library user on 21 September 2021

Figure A2. Hertzsprung–Russell diagram of 40–120 M_{\odot} He-burning stars evolving from the He-ZAMS as WRs, calculated at Z_{\odot} . Red solid tracks represent models that employ SV20 mass loss, those with NL00 rates are shown in blue dash–dotted lines, and green dashed lines illustrate models that include H95+ mass loss.

Figure A3. Hertzsprung–Russell diagram of 40–120 M_{\odot} He-burning stars evolving from the He-ZAMS as WRs, calculated at Z_{SMC} . Red solid tracks represent models that employ SV20 mass loss, those with NL00 rates are shown in blue dash–dotted lines, and green dashed lines illustrate models that include H95+ mass loss.

4888 E. R. Higgins et al.

Table A1. Standard Grid of Helium models for Section 4 with zero mass loss during core H-burning, for Z_{\odot} , 0.5 Z_{\odot} , and 0.2 Z_{\odot} . Final masses are provided at core He-exhaustion, as well as the final CO core masses.

Table A2.	Continued.	Standard	Grid of models,	for 0.2 Z _☉ ,	0.1 Z _☉ ,	and 0.	.02
Z_{\odot} .							

Mrecipe	Z/Z_{\odot}	Mini	$M_{ m f}$	Mco	M recipe
			1		H95+
SV20	1	20	12.51	9.65	H95+
SV20	1	40	17.81	14.47	H95+
SV20	1	60	22.46	18.80	H95+
SV20	1	80	26.46	22.90	H95+
SV20	1	100	30.99	26.83	H95+
SV20	1	120	34.97	30.63	H95+
SV20	1	200	50.03	39.84	SV20
NL00	1	20	11.04	8.25	SV20
NL00	1	40	15.04	11.87	SV20
NL00	1	60	18.04	14.62	SV20
NL00	1	80	20.49	16.89	SV20
NL00	1	100	22.56	18.80	SV20
NL00	1	120	24.47	20.51	SV20
NL00	1	200	29.71	25.32	NH 00
1105	1	20	11.00	9.70	NL00
H95+	1	20	11.60	8.79	NL00
H95+	1	40	10.33	13.12	NL00
H95+	1	60	19.23	15.81	NL00
H95+	1	80	21.29	17.80	NL00
H95+	1	100	22.80	19.13	NL00
H95+	1	120	23.99	20.32	NL00
H95+	1	200	26.87	23.07	H95+
SV20	0.5	20	15.45	12.33	H95+
SV20	0.5	40	22.04	18.31	H95+
SV20	0.5	60	27.57	23.55	H95+
SV20	0.5	80	32.80	28.41	H95+
SV20	0.5	100	37.56	32.84	H95+
SV20	0.5	120	41.85	36.90	H95+
SV20	0.5	200	58.87	40.74	SV20
NL00	0.5	20	12.45	9.50	SV20
NL00	0.5	40	17.25	13.85	SV20
NL00	0.5	60	19.90	16.34	SV20
NL00	0.5	80	22.08	18.34	SV20
NL00	0.5	100	24.40	20.50	SV20
NL00	0.5	120	23.54	19.66	SV20
NL00	0.5	200	30.57	26.14	NIL OO
1105	0.5	20	14.25	11.10	NL00
П93+ 1105 -	0.5	20	14.23	11.19	NL00
H95+	0.5	40	22.41	18.05	NL00
П93+ U05 I	0.5	80	20.29	24.09	NL00
H95+	0.5	80	32.98	28.48	NL00
H95+	0.5	100	30.07	32.00	NL00
H95+	0.5	120	39.54	34.57	NL00
H95+	0.5	200	48.37	39.12	H95+
SV20	0.2	20	19.25	16.01	H95+
SV20	0.2	40	28.31	24.07	H95+
SV20	0.2	60	35.60	30.80	H95+
SV20	0.2	80	42.22	37.08	H95+
SV20	0.2	100	48.45	42.86	H95+
SV20	0.2	120	54.34	44.52	H95+
SV20	0.2	200	75.91	64.28	
NL00	0.2	20	12.63	9.66	
NL00	0.2	40	19.91	16.31	
NL00	0.2	60	22.06	18.29	
NL00	0.2	80	24.35	20.38	
NL00	0.2	100	24.88	20.92	
NL00	0.2	120	27.88	23.69	
NL00	0.2	200	31.93	27.33	

$\dot{M}_{\rm recipe}$	Z/Z_{\odot}	$M_{ m ini}$	$M_{ m f}$	$M_{\rm CO}$
H95+	0.2	20	16.02	12.78
H95+	0.2	40	27.09	23.00
H95+	0.2	60	35.92	31.14
H95+	0.2	80	43.28	37.99
H95+	0.2	100	49.70	44.04
H95+	0.2	120	55.34	45.57
H95+	0.2	200	72.38	61.23
SV20	0.1	20	-	-
SV20	0.1	40	32.96	28.33
SV20	0.1	60	41.75	36.51
SV20	0.1	80	49.84	44.01
SV20	0.1	100	57.48	47.18
SV20	0.1	120	65.10	53.88
SV20	0.1	200	92.22	76.16
NL00	0.1	20	16.71	13.45
NL00	0.1	40	26.52	22.41
NL00	0.1	60	30.22	25.82
NL00	0.1	80	32.18	27.61
NL00	0.1	100	32.50	27.93
NL00	0.1	120	27.76	23.58
NL00	0.1	200	32.95	28.31
H95+	0.1	20	18.16	15.41
H95+	0.1	40	33.60	28.95
H95+	0.1	60	47.54	41.61
H95+	0.1	80	60.37	43.87
H95+	0.1	100	72.32	60.02
H95+	0.1	120	83.77	68.80
H95+	0.1	200	122.87	106.88
SV20	0.02	20	_	-
SV20	0.02	40	-	-
SV20	0.02	60	-	-
SV20	0.02	80	74.09	63.27
SV20	0.02	100	85.52	71.00
SV20	0.02	120	96.86	82.96
SV20	0.02	200	140.53	122.81
NL00	0.02	20	13.46	10.47
NL00	0.02	40	35.40	30.79
NL00	0.02	60	47.82	41.83
NL00	0.02	80	56.44	46.35
NL00	0.02	100	62.13	50.28
NL00	0.02	120	66.56	46.86
NL00	0.02	200	71.75	59.62
H95+	0.02	20	19.51	16.03
H95+	0.02	40	38.20	34.28
H95+	0.02	60	56.35	44.43
H95+	0.02	80	74.11	63.41
H95+	0.02	100	91.54	76.72
H95+	0.02	120	108.72	89.51
H95+	0.02	200	174.75	153.65

Table A3. Alternative Grid of models for a range of initial H and He masses (calculated where $X_c < 0.00001$) as a function of initial Z and wind recipe. Final masses are provided at core He-exhaustion, as well as the final CO core masses.

Table A4. Continued. Alternative Grid of models, for 0.2 $Z_{\odot},$ 0.1 $Z_{\odot},$ and 0.02 $Z_{\odot}.$

masses.					M _{recipe}	Z/Z_{\odot}	$M_{ m H}$	M _{He}	$M_{ m f}$	$M_{\rm CO}$	
Mrecipe	Z/Zo	Мн	Мне	Mf	Mco	NL00	0.2	30	16.066	11.584	9.048
	. 0	11	ne	1		NL00	0.2	50	23.554	14.611	11.548
SV20	1	30	10.784	9.301	6.821	NL00	0.2	70	30.854	16.637	13.352
SV20	1	50	12.907	10.153	7.495	NL00	0.2	100	37.007	18.328	14.939
SV20	1	70	15.040	10.887	8.165	NL00	0.2	120	40.164	19.071	15.547
SV20	1	100	17.620	11.742	8.962	NL00	0.2	150	51.357	20.848	17.219
SV20	1	120	18.965	12.162	9.312	NL00	0.2	170	51.551	21.493	17.812
SV20	1	150	21.099	12.806	9.893	H95 +	0.2	30	16.068	12,995	10.301
SV20	1	170	22.101	13.111	10.160	H95+	0.2	50	23.562	17.770	14.421
NL00	1	30	10.771	7.669	5.296	H95+	0.2	70	30.854	21.874	18.205
NL00	1	50	12.889	8.768	6.331	H95+	0.2	100	37.010	25.268	21.288
NL00	1	70	15.038	9.072	6.510	H95+	0.2	120	40.167	26.911	22.841
NL00	1	100	17.619	10.066	7.399	H95 +	0.2	150	51.357	32.243	27.724
NL00	1	120	18.965	10.627	7.912	H95+	0.2	170	51.551	32.407	28.001
NL00	1	150	21.101	11.256	8.508			20			4 4 9 4 9
NL00	1	170	22.103	11.456	8.638	SV20	0.1	30	20.886	20.886	16.919
	-		10 550			SV20	0.1	50	34.844	28.807	24.855
H95+	1	30	10.772	7.837	5.445	SV20	0.1	70	8.436	33.884	29.563
H95+	1	50	12.888	8.912	6.387	SV20	0.1	100	82.126	42.391	37.410
H95+	1	70	15.038	9.780	7.160	SV20	0.1	120	94.747	46.770	41.181
H95+	1	100	17.616	10.764	8.046	SV20	0.1	150	101.778	54.201	-
H95+	1	120	18.961	11.231	8.470	SV20	0.1	170	100.529	55.786	-
H95+	1	150	21.095	11.918	9.114	NL00	0.1	30	20.864	13.702	10.837
H95+	1	170	22.097	12.233	9.372	NL00	0.1	50	34.843	13.017	10.085
SV20	0.5	30	13.572	11.456	10.044	NL00	0.1	70	48.436	16.431	13.224
SV20	0.5	50	18.797	14.773	11.698	NL00	0.1	100	82.055	17.219	13.916
SV20	0.5	70	21.316	15.796	12.655	NL00	0.1	120	94.843	19.228	15.738
SV20	0.5	100	25.699	17.397	14.058	NL00	0.1	150	101.708	25.936	21.926
SV20	0.5	120	29.575	18.724	15.270	NL00	0.1	170	100.536	31.827	27.307
SV20	0.5	150	32.614	19.763	16.225	H05	0.1	30	20.875	17.016	14 400
SV20	0.5	170	34.610	20.382	16.824	H05	0.1	50	20.875	27.260	22 006
NI 00	0.5	30	13 572	0.01/	7 296	H05+	0.1	50 70	18 / 30	35 903	23.000
NL 00	0.5	50	18 788	9.914	8 724	H05	0.1	100	82 213	40 726	13 747
NL00	0.5	50 70	21 315	12 871	0.724	H05+	0.1	120	94 725	49.720 57.530	43.747
NL 00	0.5	100	25 697	14 177	11 123	H05+	0.1	120	101 882	68 791	
NL 00	0.5	120	29.577	15 104	11.125	H95+	0.1	170	100.539	70 527	_
NL 00	0.5	120	32 620	15.848	12 592	11)5+	0.1	170	100.557	10.521	
NL 00	0.5	170	34,610	16.097	12.372	SV20	0.02	50	44.000	44.000	37.443
NLOO	0.5	170	54.010	10.077	12.075	SV20	0.02	100	79.529	65.719	-
H95+	0.5	30	13.572	10.484	7.858	SV20	0.02	120	89.442	71.796	-
H95+	0.5	50	18.790	13.407	10.473	SV20	0.02	150	140.635	91.535	-
H95+	0.5	70	21.315	14.752	11.681	SV20	0.02	170	155.078	99.927	-
H95+	0.5	100	25.697	16.821	13.513	NL00	0.02	30	25.658	10.994	8.299
H95+	0.5	120	29.576	18.484	15.059	NL00	0.02	50	44.272	11.415	8.669
H95+	0.5	150	32.616	19.756	16.254	NL00	0.02	70	62.588	14.701	11.661
H95+	0.5	170	34.610	20.495	16.904	NL00	0.02	100	79.491	18.825	15.358
SV20	0.2	30	16.074	16.070	12.857	NL00	0.02	120	89.442	23.006	19.262
SV20	0.2	50	23 581	20.824	17.317	NL00	0.02	150	140.228	22.461	18.694
SV20	0.2	70	30 854	24.145	20.289	NL 00	0.02	170	155 138	27 149	23 083
SV20	0.2	100	37.014	26.804	22.695		0.02				_0.000
SV20	0.2	120	40.170	28.124	23 983	H95+	0.02	30	25.661	21.938	18.198
SV20	0.2	150	51,356	32,514	27.999	H95+	0.02	50	44.418	32.343	28.744
SV20	0.2	170	51,551	32.613	28,121	H95+	0.02	70	62.588	45.752	40.418
	5.2	170	01.001	52.015	20.121	H95+	0.02	100	79.520	62.506	-
						H95+	0.02	120	89.442	73.388	_

H95+

H95+

0.02

0.02

This paper has been typeset from a $T_{\!E\!}X/I\!\!\!\!\!\!^{A}\!T_{\!E\!}X$ file prepared by the author.

140.426

155.209

150

170

104.398

117.140

_