

Pd-Catalyzed Rearrangement of *N*-Alloc-*N*-allyl Ynamides via Auto-Tandem Catalysis: Evidence for Reversible C–N Activation and Pd(0)-Accelerated Ketenimine Aza-Claisen Rearrangement

Juliana R. Alexander,[‡] Veronika I. Shchepetkina,[‡] Ksenia S. Stankevich,[‡] Rory J. Benedict, Samuel P. Bernhard, Reagan J. Dreiling, and Matthew J. Cook*



Cite This: <https://dx.doi.org/10.1021/acs.orglett.0c04078>



Read Online

ACCESS |




Metrics & More



Article Recommendations



Supporting Information




ABSTRACT: An auto-tandem catalytic double allylic rearrangement of *N*-alloc-*N*-allyl ynamides was developed. This reaction proceeds through two separate and distinct catalytic cycles with both decarboxylative Pd– π -allyl and Pd(0)-promoted aza-Claisen rearrangements occurring. A detailed mechanistic study supported by computations highlights these two separate mechanisms. Previously unreported reversible C–N ionization and a Pd(0)-catalyzed [3,3]-sigmatropic rearrangement were discovered. This study provides new reaction pathways for both π -allyl and sigmatropic rearrangements.

The ability of a single catalytic entity to perform two or more mechanistically distinct reactions is a powerful strategy in synthesis. This approach, termed auto-tandem catalysis (ATC) by Fogg and dos Santos,¹ can provide a very direct and atom-efficient route to complex molecules.² Their definition of an ATC reaction requires a single catalyst, which is present at the reaction outset, to proceed through two different mechanistic cycles without the need for a chemical trigger to alternate between these mechanisms.³ As such, these are mechanistically distinct from cascade reactions that intercept catalyst-bound intermediates and/or use bifunctional/cooperative catalysts.⁴ The use of the ATC approach is especially impactful when sequential C–C bond-forming events occur during the reactions.⁵ Importantly, ATCs allow the *in situ* formation and use of unstable or sensitive organic intermediates that would not be amenable to a linear synthetic approach. Aside from their synthetic utility, these reactions are mechanistically intriguing, with a single catalyst providing orthogonal reactivity between substrates and intermediates in a single reaction sequence. Herein we report the development of a new ATC reaction that performs both π -allyl and sigmatropic allylic rearrangements mediated by a single palladium(0) catalyst. Of particular note are the previously unreported Pd(0)-catalyzed aza-Claisen rearrangement and reversible π -allylic ionization of neutral allyl amines.


We previously disclosed the Pd-catalyzed decarboxylative rearrangement of *N*-alloc ynamides to form ketenimines, which were trapped with nucleophiles (Scheme 1).^{6,7} We envisaged that if *N*-alloc-*N*-allyl ynamides were utilized, two sequential

Scheme 1. Rearrangement of *N*-Alloc-*N*-allyl Ynamides in C–C Bond-Forming ATC Reactions

Previous Work (Reference 6a)



This Work



Received: December 9, 2020

palladium-catalyzed allylic rearrangements could occur through an *N*-allyl ketenimine intermediate. This approach would provide a unique method for the synthesis of quaternary nitriles bearing two independent allyl groups.⁸ More importantly, we hypothesized that both allylic rearrangements could be performed by a single palladium catalyst through two distinct mechanisms, one π -allylic and one sigmatropic. Although the uncatalyzed thermal ketenimine aza-Claisen rearrangement has been previously reported,⁹ we reasoned that a catalyst-accelerated variant could be developed. While electrophilic Pd(II) complexes have been reported to promote a range of sigmatropic rearrangements,¹⁰ these proceed *via* a π -Lewis acidic pathway, requiring either a strongly nucleophilic migrating group or a strongly electrophilic palladium π -Lewis acid.¹¹ Tunge has reported a suite of Pd-catalyzed π -allylic rearrangements to form nitriles; these generate π -allyl intermediates and nitrile α -anions *in situ* (*via* decarboxylative and deacylative routes).¹² While the use of protonated allyl amines as π -allyl donors has been established,¹³ the activation of neutral allyl amines has not been reported and represents a new mode of reactivity. The rearrangement of butyl derivative **1a** was optimized with a Pd(OAc)₂/BINAP catalyst system, with Et₂B(OMe) and *N,O*-BSA proving optimal (Table 1). Both

similar efficiency as **1a**. This reaction provided quaternary nitrile **2o** with high levels of linear regioselectivity.

As uncatalyzed sigmatropic ketenimine aza-Claisen rearrangements were previously reported,⁹ we examined whether both rearrangement reactions were indeed catalyzed by a single palladium catalyst. Two distinct substrate classes were chosen, with butyl (**1a**) and phenyl (**1e**) substrates being the focus of our studies (Scheme 2). Deuterium-labeled analogues (**3** and **4**)

Scheme 2. Substrate Scope




Table 1. Optimization Studies

entry	change from optimized conditions	yield (%) ^a
1	none	72 (56 ^b)
2	no <i>N,O</i> -BSA	24
3	BEt ₃ instead of Et ₂ B(OMe)	42
4	no borane (Et ₂ B(OMe) or BEt ₃)	n.r.
5	(<i>S</i>)-SEGPHOS as the ligand	32
6	Trost ligands (ANDEN or DACH)	n.r.
7	Xantphos as the ligand	63

^aDetermined by ¹H NMR analysis with mesitylene as an internal standard. ^bIsolated yield.


the *N,O*-BSA and Et₂B(OMe) additives were crucial for high efficiency. Et₃B provided products in lower yield, and the absence of borane led to no reaction. Other bidentate ligands, such as SEGPHOS or Xantphos, were effective, whereas both Trost ligand scaffolds led to no reaction.

Following optimization, we embarked on a substrate scope study to examine the generality of the reaction. Quaternary nitriles bearing unsubstituted alkyl groups (**2a**, **2b**), highly substituted alkyl groups (**2c**), and phenethyl (**2d**) were formed in good yields. Aryl alkene substrates (**1e–m**) were particularly efficient for this reaction. Interestingly, there was little difference in efficiency between aryl alkynes substituted by electron-neutral (**1e**), σ -donating (**1f**, **1g**), or σ -withdrawing (**1h–j**) groups. When mesomeric π -withdrawing groups were present (**1k**, **1l**), enhanced reactivity was observed, and excellent yields of the products were obtained. The use of polyaromatic-substituted alkynes was demonstrated, with 2-naphthyl-substituted **1m** and the more hindered 1-naphthyl analogue **1n** reacting with similar reactivity. Substitution at R₂ could also be tolerated, with **1o** bearing a cinnamyl-substituted *N*-alloc group reacting with

^aAverage isolated yield over two reactions. ^bThe reaction was performed on a 1 mmol scale. ^cIsolated as a 10:1 mixture of regioisomers.

were prepared in which CH₂ was independently replaced with CD₂ at the *N*-allyl (**3**) and *N*-alloc (**4**) positions. These were then subjected to the optimized reaction conditions, and the position of the deuterium was tracked (Scheme 3a). When the deuterated *N*-allyl analogues (**3**) were utilized, two reactivity patterns were observed. The butyl substrate (**3a**) provided a single regioisomer **5a** with the deuterium exclusively at the terminal vinylic position. With the phenyl analogue (**3b**), deuterium scrambling was observed in a **5b**:**6b** ratio of 1.2:1. These results clearly demonstrate a divergence in mechanisms between the two substrates (**3a** and **3b**) and indicate that a dissociative allylic pathway occurs for **3b** that is not accessible to **3a**. Of note is the slight excess of the terminal vinylic deuterium isomer (**5b**). The origin of the selectivity in **3b** could be produced by either two competing (concerted and dissociative) mechanisms or a secondary kinetic isotope effect in a rapidly isomerizing (Curtin–Hammett) system. In the latter case, it would suggest the facile formation of a π -allyl intermediate followed by a higher-energy C–C bond-forming step with

Scheme 3. Deuterium-Labeling and Crossover Experiments



^aConditions: 5 mol % Pd(OAc)₂, 6 mol % *rac*-BINAP, Et₃B(OMe) (1 equiv), *N,O*-BSA (1 equiv), toluene, 60 °C, 16 h. ^bRatios were determined by ¹H NMR analyses of unpurified reaction mixtures. ^cPercentages represent the isotopic makeup of individual products and were measured by GCMS/HRMS of unpurified reaction mixtures. Values are corrected for natural isotopic abundance.


rehybridization from sp³ to sp² determining the selectivity (Scheme 3b). When deuteration was installed on the *N*-alloc group (4), isotopic scrambling was observed in both cases. The butyl analogue 4a provided a slight excess of the allylic CD₂ compound 6a, whereas the phenyl analogue 4b provided equimolar quantities of 5b and 6b. These results are consistent with our previous mechanistic experiments and indicate that the *N*-alloc rearrangement proceeds *via* a more classical dissociative π -allylic mechanism.^{6a}

Crossover studies were performed using 0.5 equiv each of 3 and a structurally similar unlabeled compound (1b and 1h) to probe whether any solvent-separated dissociation or π -allyl ligand exchange was occurring (Scheme 3c). The competition reaction of butyl (3a) and hexyl (1b) ynamides provided no evidence of dissociation, with complete retention of the deuterium label in the butyl substrate. Phenyl (3b) and 4-fluorophenyl (1h) ynamides gave trace amounts of deuterium crossover. This suggests that the dissociative pathway for 3b that leads to deuterium scrambling is short-lived and/or that solvent separation is not favored. Thus, we focused our efforts on determining the exact nature of the previously unknown palladium-catalyzed ketenimine aza-Claisen step.

To further elaborate our mechanistic insight, we conducted computational studies on ketenimines 8a and 8e,¹⁴ which are derived from 1a and 1e, respectively.¹⁵ Because of the large ligand, we applied the two-layer quantum-mechanical (QM)/semiempirical (SE) ONIOM model to the palladium complexes.¹⁶ The QM layer was treated with M06L/6-31+G(d) (C, H, P, B, O)/LANL2DZ (ECP Pd)¹⁷ and the SE layer with PM6. Single-point (SP) energies of these optimized structures were calculated utilizing M06L/6-311++G(2d,2p) (C, H, P, B, O)/LANL2DZ (ECP Pd). The solvation by toluene was taken into account using C-PCM.¹⁸ Transition states (TSs) were located using the nudged elastic band (NEB) method implemented in ORCA 4.2.1¹⁹ at the M06L level with the def2-SV(P) basis set followed by the ONIOM/SP process described above. Conformational flexibility was addressed through the mixed torsional/low mode method and relaxed torsion scans (see the Supporting Information (SI) for full details).

Using phenyl ketenimine 8e as an initial model, we explored both catalytic and uncatalyzed pathways for the sigmatropic rearrangement and discovered a complex potential energy surface. The uncatalyzed ketenimine aza-Claisen rearrangement of 8e proceeded through an early transition state analogous to those described by Walters.⁹ This rearrangement is significantly faster with the free ketenimine rather than *via* a Lewis acid–base adduct with the borane additive (not shown in Scheme 3; see the SI for details). In the palladium-catalyzed routes (Scheme 4),²⁰ coordination of ketenimine 8e with the Pd(0) catalyst to form 9 is exergonic by 11.7 kcal/mol. π -Allylic ionization mechanisms were examined with and without the borane additive bound. Direct ionization of 9 to form π -allyl species 10 (*via* TS-ION) was readily accessible with a TS energy of just 11.4 kcal/mol. Once again, the borane adduct route was significantly higher in energy than the direct ionization of 9. As a result, we do not believe that the borane additive participates in the ketenimine rearrangement of 8e and is only necessary to perform the initial π -allyl rearrangement of 1e. The structure of 10 is a heavily distorted η^3 - π -allyl tight ion pair, which can easily isomerize between the η^3 and η^1 haptomers (10 and 12), leading to the deuterium scrambling observed in Scheme 2. Bond rotation is required to form the reactive conformer 13, which can rearrange *via* an inner-sphere C–C bond formatting mechanism (TS-IS). The palladium-bound rearranged nitrile 13 is formed with an overall TS energy of 15.8 kcal/mol from 9.

A concerted Pd(0)-catalyzed pericyclic process was also investigated. This mechanism, which is also consistent with experimental data, proceeds through highly ordered TS-[3,3]-Pd and is 1.9 kcal/mol lower in energy than the TS-IS route. The deuterium scrambling and crossover described in Scheme 3 can be reconciled by rapid and reversible ionization through TS-ION, which is 2.5 kcal/mol lower than TS-[3,3]-Pd. The

Scheme 4. Summary of the DFT Study Examining Catalytic and Uncatalyzed Mechanisms^a

^aGeometries/frequencies were obtained using the QM/SE ONIOM model (M06L/6-31+G(d) (C, H, P, B, O)/LANL2DZ (ECP Pd):PM6). TSs were located using the NEB method in ORCA 4.2.1 at the M06L/def2-SV(P) level and confirmed by intrinsic reaction coordinate calculations. All of the SP energies were calculated using C-PCM (toluene) at the M06L/6-311++G(2d,2p) (C, H, P, B, O)/LANL2DZ (ECP Pd) level. See the SI for full details.

structure of TS-[3,3]-Pd is a hybrid between π -allylic and sigmatropic processes, where the reaction is aided by donation of electron density from the Pd(0) center into the C–N σ^* orbital, leading to a later TS than the uncatalyzed variant TS-[3,3]. When one examines the migrating allylic portion, the three carbon atoms are almost trigonal-planar, and the terminal C–Pd bond lengths (2.36 and 2.53 Å) indicate substantial interactions between the Pd catalyst and both termini. This effect can be seen more clearly when the HOMO and LUMO coefficients are examined (Figure 1). There is significant overlap of the d orbitals of the Pd(0) atom (in the HOMO) with the σ^* orbital of the C–N bond (in the LUMO). Additionally, both the forming C–C and breaking C–N σ bonds are significant contributors to the HOMO, indicative of a sigmatropic process. Furthermore, the HOMO and LUMO contain two and four

nodes, respectively, indicating an aromatic TS. The HOMO/LUMO gap is just -0.87 eV, demonstrating facile transfer of electron density between these two molecular orbitals. As the $\Delta\Delta G^\ddagger$ between the TS-[3,3]-Pd and TS-IS routes is 1.9 kcal/mol, representing an 18:1 difference in relative rates at 60 °C, we believe that the Pd(0)-assisted [3,3]-sigmatropic and inner-sphere mechanisms both occur, with the former being the much faster pathway.

Analogous computational studies performed on butyl ketenimine **8a** indicate that the reaction proceeds through a similar Pd(0)-catalyzed [3,3]-sigmatropic rearrangement, but the ionization pathway is much higher in energy because of the lower stability of the ketenimine anion formed in ionized intermediate **10**. As this dissociative mechanism is no longer in operation, both deuterium isomerization and crossover pathways are no longer accessible (see the SI for details).

In conclusion, we have developed a highly efficient auto-tandem catalytic process to form substituted nitriles from *N*-alloc-*N*-allyl ynamides. This reaction proceeds through two mechanistically distinct allylic rearrangement reactions, with both π -allylic and sigmatropic processes promoted by a single Pd(0) source. Computations and experimental mechanistic data helped elucidate these pathways. This study identified two new reaction mechanisms, including the π -allylic ionization of neutral C–N bonds and a Pd(0)-catalyzed aza-Claisen rearrangement. These Pd(0)-catalyzed sigmatropic rearrangements have potential to be incorporated in many other ATC reactions.




Figure 1. Calculated molecular orbitals for TS-[3,3]-Pd

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.0c04078>.

Experimental procedures, $^1\text{H}/^{13}\text{C}$ NMR spectra, computational methods, and Cartesian coordinates (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Matthew J. Cook – Department of Chemistry and Biochemistry, Montana State University, Bozeman, Montana 59717, United States; orcid.org/0000-0003-4762-8535; Email: matthewcook@montana.edu

Authors

Juliana R. Alexander – Department of Chemistry and Biochemistry, Montana State University, Bozeman, Montana 59717, United States

Veronika I. Shchepetkina – Department of Chemistry and Biochemistry, Montana State University, Bozeman, Montana 59717, United States

Ksenia S. Stankevich – Department of Chemistry and Biochemistry, Montana State University, Bozeman, Montana 59717, United States; orcid.org/0000-0002-6701-7582

Rory J. Benedict – Department of Chemistry and Biochemistry, Montana State University, Bozeman, Montana 59717, United States

Samuel P. Bernhard – Department of Chemistry and Biochemistry, Montana State University, Bozeman, Montana 59717, United States

Reagan J. Dreiling – Department of Chemistry and Biochemistry, Montana State University, Bozeman, Montana 59717, United States

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acs.orglett.0c04078>

Author Contributions

[‡]J.R.A., V.I.S., and K.S.S. contributed equally.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was generously supported by NSF (CHE-1800487) and Montana State University. Calculations were performed on Comet at SDSC through XSEDE (CHE-200093) and the Hyalite High Performance Computing System at MSU. Undergraduate researchers (V.I.S. and R.J.D.) were supported by the NIH-INBRE (P20GM103474) and NSF-REU (CHE-1461218) programs. NSF-MRI (DBI-1532078) and the Murdock Charitable Trust Foundation (2015066:MNL) provided funds for a new 500 MHz NMR spectrometer used in this research. We gratefully acknowledge Professor Sharon R. Neufeldt (MSU) for helpful discussions about DFT studies.

■ REFERENCES

- (1) Fogg, D. E.; dos Santos, E. N. Tandem catalysis: a taxonomy and illustrative review. *Coord. Chem. Rev.* **2004**, *248* (21), 2365–2379.
- (2) For reviews of ATC reactions, see: (a) Shindoh, N.; Takemoto, Y.; Takasu, K. Auto-Tandem Catalysis: A Single Catalyst Activating Mechanistically Distinct Reactions in a Single Reactor. *Chem. - Eur. J.* **2009**, *15* (45), 12168–12179. (b) Patil, N. T.; Shinde, V. S.; Gajula, B. A one-pot catalysis: the strategic classification with some recent examples. *Org. Biomol. Chem.* **2012**, *10* (2), 211–224. (c) Camp, J. E. Auto-Tandem Catalysis: Activation of Multiple, Mechanistically Distinct Process by a Single Catalyst. *Eur. J. Org. Chem.* **2017**, *2017* (3), 425–433.
- (3) For selected examples of ATC reactions, see: (a) Huff, C. A.; Sanford, M. S. Cascade Catalysis for the Homogeneous Hydrogenation of CO_2 to Methanol. *J. Am. Chem. Soc.* **2011**, *133* (45), 18122–18125. (b) Kanbayashi, N.; Takenaka, K.; Okamura, T.-a.; Onitsuka, K. Asymmetric Auto-Tandem Catalysis with a Planar-Chiral Ruthenium Complex: Sequential Allylic Amidation and Atom-Transfer Radical Cyclization. *Angew. Chem., Int. Ed.* **2013**, *52* (18), 4897–4901. (c) Venning, A. R. O.; Kwiatkowski, M. R.; Roque Peña, J. E.; Lainhart, B. C.; Guruparan, A. A.; Alexanian, E. J. Palladium-Catalyzed Carbocyclizations of Unactivated Alkyl Bromides with Alkenes Involving Auto-tandem Catalysis. *J. Am. Chem. Soc.* **2017**, *139* (33), 11595–11600. (d) Zhou, C.; Jiang, J.; Wang, J. Three-Component Synthesis of Isoquinoline Derivatives by a Relay Catalysis with a Single Rhodium(III) Catalyst. *Org. Lett.* **2019**, *21* (13), 4971–4975.
- (4) For reviews of cascade and cooperative catalysis, see: Shibasaki, M.; Kanai, M.; Matsunaga, S.; Kumagai, N. Recent Progress in Asymmetric Bifunctional Catalysis Using Multimetallic Systems. *Acc. Chem. Res.* **2009**, *42* (8), 1117–1127. (b) Siau, W.-Y.; Wang, J. Asymmetric organocatalytic reactions by bifunctional amine-thioureas. *Catal. Sci. Technol.* **2011**, *1* (8), 1298–1310. (c) Allen, A. E.; MacMillan, D. W. C. Synergistic catalysis: A powerful synthetic strategy for new reaction development. *Chemical Science* **2012**, *3* (3), 633–658. (d) Inamdar, S. M.; Shinde, V. S.; Patil, N. T. Enantioselective cooperative catalysis. *Org. Biomol. Chem.* **2015**, *13* (30), 8116–8162. (e) Kim, D.-S.; Park, W.-J.; Jun, C.-H. Metal–Organic Cooperative Catalysis in C–H and C–C Bond Activation. *Chem. Rev.* **2017**, *117* (13), 8977–9015. (f) Chauhan, P.; Mahajan, S.; Kaya, U.; Hack, D.; Enders, D. Bifunctional Amine-Squaramides: Powerful Hydrogen-Bonding Organocatalysts for Asymmetric Domino/Cascade Reactions. *Adv. Synth. Catal.* **2015**, *357* (2–3), 253–281. (g) Wang, J.; Dong, G. Palladium/Norbornene Cooperative Catalysis. *Chem. Rev.* **2019**, *119* (12), 7478–7528.
- (5) For examples of ATC reactions with sequential C–C coupling, see: (a) Kim, B. G.; Snapper, M. L. Preparation of Alkenyl Cyclopropanes through a Ruthenium-Catalyzed Tandem Ene Metathesis–Cyclopropanation Sequence. *J. Am. Chem. Soc.* **2006**, *128* (1), 52–53. (b) Wang, M.-Z.; Zhou, C.-Y.; Che, C.-M. A silver-promoted auto-tandem catalysis for the synthesis of multiply substituted tetrahydrocarbazoles. *Chem. Commun.* **2011**, *47* (4), 1312–1314. (c) Chen, P.; Chen, Z.-C.; Li, Y.; Ouyang, Q.; Du, W.; Chen, Y.-C. Auto-Tandem Cooperative Catalysis Using Phosphine/Palladium: Reaction of Morita–Baylis–Hillman Carbonates and Allylic Alcohols. *Angew. Chem., Int. Ed.* **2019**, *58* (12), 4036–4040. (d) Long, J.; Yu, R.; Gao, J.; Fang, X. Access to 1,3-Dinitriles by Enantioselective Auto-tandem Catalysis: Merging Allylic Cyanation with Asymmetric Hydrocyanation. *Angew. Chem., Int. Ed.* **2020**, *59* (17), 6785–6789.
- (6) (a) Alexander, J. R.; Cook, M. J. Formation of Ketenimines via the Palladium-Catalyzed Decarboxylative π -Allylic Rearrangement of *N*-Alloc Ynamides. *Org. Lett.* **2017**, *19* (21), 5822–5825. Also see: (b) Chen, J.; Cook, M. J. Palladium Catalyzed Decarboxylative Rearrangement of *N*-Alloc Indoles. *Org. Lett.* **2013**, *15* (5), 1088–1091.
- (7) For analogous approaches, see: (a) De Korver, K. A.; Hsung, R. P.; Lohse, A. G.; Zhang, Y. A Divergent Mechanistic Course of Pd(0)-Catalyzed Aza-Claisen Rearrangement and Aza-Rautenstrauch-Type Cyclization of *N*-Allyl Ynamides. *Org. Lett.* **2010**, *12* (8), 1840–1843. (b) DeKorver, K. A.; Wang, X.-N.; Walton, M. C.; Hsung, R. P. Carbocyclization Cascades of Allyl Ketenimines via Aza-Claisen Rearrangements of *N*-Phosphoryl-*N*-allyl-ynamides. *Org. Lett.* **2012**, *14* (7), 1768–1771. (c) Wang, X.-N.; Yeom, H.-S.; Fang, L.-C.; He, S.; Ma, Z.-X.; Kedrowski, B. L.; Hsung, R. P. Ynamides in Ring Forming Transformations. *Acc. Chem. Res.* **2014**, *47* (2), 560–578.
- (8) (a) *The Chemistry of the Cyano Group*; Wiley-Interscience: New York, 1970. (b) North, M. In *Comprehensive Organic Functional Group Transformations II*; Taylor, R. J. K., Ed.; Elsevier: Oxford, U.K., 2005; p

621. (c) Fleming, F. F.; Yao, L.; Ravikumar, P. C.; Funk, L.; Shook, B. C. Nitrile-Containing Pharmaceuticals: Efficacious Roles of the Nitrile Pharmacophore. *J. Med. Chem.* **2010**, *53* (22), 7902–7917. (d) Larock, R. C.; Rozhkov, R. In *Comprehensive Organic Transformations*, 3rd ed.; Larock, R. C., Ed.; John Wiley & Sons, 2018.
- (9) (a) Walters, M. A.; Hoem, A. B.; Arcand, H. R.; Hegeman, A. D.; McDonough, C. S. An extremely mild 3-aza-claisen reaction. 2. New conditions and the rearrangement of α -heteroatom substituted amides. *Tetrahedron Lett.* **1993**, *34* (9), 1453–1456. (b) Walters, M. A.; Hoem, A. B.; McDonough, C. S. Rearrangements of Substituted 3-Aza-1,2,5-hexatrienes. 3. The Scope and Versatility of an Extremely Mild 3-Aza-Cope Reaction. *J. Org. Chem.* **1996**, *61* (1), 55–62.
- (10) For selected examples, see: (a) Lee, E. E.; Batey, R. A. Palladium-Catalyzed Allylic Transposition of (Allyloxy) Iminodiazaphospholindines: A Formal [3,3]-Aza-phospha-oxa-Cope Sigmatropic Rearrangement for the Stereoselective Synthesis of Allylic Amines. *J. Am. Chem. Soc.* **2005**, *127* (42), 14887–14893. (b) Kerrigan, N. J.; Bungard, C. J.; Nelson, S. G. Pd(II)-catalyzed aliphatic Claisen rearrangements of acyclic allyl vinyl ethers. *Tetrahedron* **2008**, *64* (29), 6863–6869. (c) Danowitz, A. M.; Taylor, C. E.; Shrikian, T. M.; Mapp, A. K. Palladium-Catalyzed [3,3]-Rearrangement for the Facile Synthesis of Allenamides. *Org. Lett.* **2010**, *12* (11), 2574–2577. (d) Bao, H.; Tambar, U. K. Catalytic Enantioselective Allylic Amination of Unactivated Terminal Olefins via an Ene Reaction/[2,3]-Rearrangement. *J. Am. Chem. Soc.* **2012**, *134* (45), 18495–18498. (e) Cannon, J. S.; Overman, L. E. Palladium(II)-Catalyzed Enantioselective Reactions Using COP Catalysts. *Acc. Chem. Res.* **2016**, *49* (10), 2220–2231.
- (11) For computational studies, see: (a) Siebert, M. R.; Tantillo, D. J. Transition-State Complexation in Palladium-Promoted [3,3] Sigmatropic Shifts. *J. Am. Chem. Soc.* **2007**, *129* (28), 8686–8687. (b) Watson, M. P.; Overman, L. E.; Bergman, R. G. Kinetic and Computational Analysis of the Palladium(II)-Catalyzed Asymmetric Allylic Trichloroacetimidate Rearrangement: Development of a Model for Enantioselectivity. *J. Am. Chem. Soc.* **2007**, *129* (16), 5031–5044. (c) Gutierrez, O.; Hendrick, C. E.; Kozlowski, M. C. Divergent Reactivity in Pd-Catalyzed [3,3]-Sigmatropic Rearrangement of Allyloxy- and Propargyloxyindoles Revealed by Computation and Experiment. *Org. Lett.* **2018**, *20* (20), 6539–6543.
- (12) (a) Recio, A.; Tunge, J. A. Regiospecific Decarboxylative Allylation of Nitriles. *Org. Lett.* **2009**, *11* (24), 5630–5633. (b) Grenning, A. J.; Tunge, J. A. Deacylative Allylation: Allylic Alkylation via Retro-Claisen Activation. *J. Am. Chem. Soc.* **2011**, *133* (37), 14785–14794. (c) Grenning, A. J.; Tunge, J. A. Deacylative Allylation of Nitroalkanes: Unsymmetric Bisallylation by a Three-Component Coupling. *Angew. Chem., Int. Ed.* **2011**, *50* (7), 1688–1691. (d) Lang, S. B.; Locascio, T. M.; Tunge, J. A. Activation of Alcohols with Carbon Dioxide: Intermolecular Allylation of Weakly Acidic Pronucleophiles. *Org. Lett.* **2014**, *16* (16), 4308–4311. (e) Maji, T.; Tunge, J. A. Catalytic α -Monoallylation of Aryl Acetonitriles. *Org. Lett.* **2014**, *16* (19), 5072–5075. (f) Allegre, K. M.; Brennan, N.; Tunge, J. A. Synthesis of Vinyl Cyclopropanes via Anion Relay Cyclization. *Org. Lett.* **2018**, *20* (14), 4191–4194. (g) Allegre, K.; Tunge, J. Aryl vinyl cyclopropane Cope rearrangements. *Tetrahedron* **2019**, *75* (24), 3319–3329.
- (13) For reviews, see: (a) Ouyang, K.; Hao, W.; Zhang, W.-X.; Xi, Z. Transition-Metal-Catalyzed Cleavage of C–N Single Bonds. *Chem. Rev.* **2015**, *115* (21), 12045–12090. (b) Butt, N. A.; Zhang, W. Transition metal-catalyzed allylic substitution reactions with unactivated allylic substrates. *Chem. Soc. Rev.* **2015**, *44* (22), 7929–7967. For representative examples, see: (c) Murahashi, S.-i.; Makabe, Y. Palladium(0) catalyzed 3-aza-cope rearrangement of N-allylenamines. *Tetrahedron Lett.* **1985**, *26* (45), 5563–5566. (d) Murahashi, S.; Makabe, Y.; Kunita, K. Palladium(0)-catalyzed rearrangement of N-allyl enamines. Synthesis of δ,ϵ -unsaturated imines and γ,δ -unsaturated carbonyl compounds. *J. Org. Chem.* **1988**, *53* (19), 4489–4495. (e) Mukherjee, S.; List, B. Chiral Counteranions in Asymmetric Transition-Metal Catalysis: Highly Enantioselective Pd/Brønsted Acid-Catalyzed Direct α -Allylation of Aldehydes. *J. Am. Chem. Soc.* **2007**, *129* (37), 11336–11337. (f) Li, M.-B.; Wang, Y.; Tian, S.-K. Regioselective and Stereospecific Cross-Coupling of Primary Allylic Amines with Boronic Acids and Boronates through Palladium-Catalyzed C–N Bond Cleavage. *Angew. Chem., Int. Ed.* **2012**, *51* (12), 2968–2971. (g) Xu, Y.-N.; Zhu, M.-Z.; Tian, S.-K. Chiral α -Amino Acid/Palladium-Catalyzed Asymmetric Allylation of α -Branched β -Ketoesters with Allylic Amines: Highly Enantioselective Construction of All-Carbon Quaternary Stereocenters. *J. Org. Chem.* **2019**, *84* (22), 14936–14942.
- (14) *In situ* reaction monitoring (using ^1H NMR) showed no appreciable buildup of **8e**, suggesting that the rearrangement of **8e** is significantly faster than the rearrangement of **1e**.
- (15) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Petersson, G. A.; Nakatsuji, H.; Li, X.; Caricato, M.; Marenich, A.; Bloino, J.; Janesko, B. G.; Gomperts, R.; Mennucci, B.; Hratchian, H. P.; Ortiz, J. V.; Izmaylov, A. F.; Sonnenberg, J. L.; Williams-Young, D.; Ding, F.; Lipparini, F.; Egidi, F.; Goings, J.; Peng, B.; Petrone, A.; Henderson, T.; Ranasinghe, D.; Zakrzewski, V. G.; Gao, J.; Rega, N.; Zheng, G.; Liang, W.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Throssell, K.; Montgomery, J. A., Jr.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E. N.; Kudin, K. N.; Staroverov, V. N.; Keith, T. A.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A. P.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Millam, J. M.; Klene, M.; Adamo, C.; Cammi, R.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Farkas, O.; Foresman, J. B.; Fox, D. J. *Gaussian 16*, rev. A.03; Gaussian, Inc.: Wallingford, CT, 2016.
- (16) (a) Ananikov, V. P.; Musaev, D. G.; Morokuma, K. Real Size of Ligands, Reactants and Catalysts: Studies of Structure, Reactivity and Selectivity by ONIOM and Other Hybrid Computational Approaches. *J. Mol. Catal. A: Chem.* **2010**, *324* (1–2), 104–119. (b) Li, Z.; Liu, L.; Fu, Y.; Guo, Q.-X. Assessing Performance of Diverse ONIOM Methods in Calculation of Structures of Organonickel and Organopalladium Compounds. *J. Mol. Struct.: THEOCHEM* **2005**, *757* (1–3), 69–76. (c) Karami, K.; Abedanzadeh, S.; Farrokhpour, H.; Lipkowsky, J. Synthesis and Characterization of the P,C-Palladacycles with Bridging and Chelating Dinitrogen Ligands and ONIOM Calculations on the Pyrazine-Bridged Organometallic Polymers ($n = 1$ to $n = 10$). *J. Organomet. Chem.* **2016**, *805*, 68–76.
- (17) Zhao, Y.; Truhlar, D. G. The M06 Suite of Density Functionals for Main Group Thermochemistry, Thermochemical Kinetics, Non-covalent Interactions, Excited States, and Transition Elements: Two New Functionals and Systematic Testing of Four M06-Class Functionals and 12 Other Function. *Theor. Chem. Acc.* **2008**, *120* (1–3), 215–241.
- (18) (a) Barone, V.; Cossi, M. Quantum Calculation of Molecular Energies and Energy Gradients in Solution by a Conductor Solvent Model. *J. Phys. Chem. A* **1998**, *102* (11), 1995–2001. (b) Klamt, A.; Schüürmann, G. COSMO: A New Approach to Dielectric Screening in Solvents with Explicit Expressions for the Screening Energy and Its Gradient. *J. Chem. Soc., Perkin Trans. 2* **1993**, No. 5, 799–805.
- (19) (a) Neese, F. The ORCA Program System. *Wiley Interdiscip. Rev.: Comput. Mol. Sci.* **2012**, *2* (1), 73–78. (b) Neese, F. Software Update: The ORCA Program System, Version 4.0. *Wiley Interdiscip. Rev.: Comput. Mol. Sci.* **2018**, *8*, e1327.
- (20) For other detailed DFT studies of π -allyl pathways, see: (a) McPherson, K. E.; Croatt, M. P.; Morehead, A. T.; Sargent, A. L. DFT Mechanistic Investigation of an Enantioselective Tsuji–Trost Allylation Reaction. *Organometallics* **2018**, *37* (21), 3791–3802. (b) Keith, J. A.; Behenna, D. C.; Sherden, N.; Mohr, J. T.; Ma, S.; Marinescu, S. C.; Nielsen, R. J.; Oxgaard, J.; Stoltz, B. M.; Goddard, W. A. The Reaction Mechanism of the Enantioselective Tsuji Allylation: Inner-Sphere and Outer-Sphere Pathways, Internal Rearrangements, and Asymmetric C–C Bond Formation. *J. Am. Chem. Soc.* **2012**, *134* (46), 19050–19060.