

Directed, Nickel-Catalyzed Umpolung 1,2-Carboamination of Alkenyl Carbonyl Compounds

Vincent A. van der Puyl, Joseph Derosa, and Keary M. Engle*

Department of Chemistry, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, California 92037, United States

Supporting Information Placeholder


ABSTRACT: We report a regioselective, nickel-catalyzed *syn*-1,2-carboamination of non-conjugated alkenyl carbonyl compounds with *O*-benzoyl hydroxylamine (N–O) electrophiles and aryl/alkylzinc nucleophiles to afford β - and γ -amino acid derivatives. This method enables preparation of products containing structurally diverse tertiary amine motifs, including heterocycles, and can also be used to form quaternary carbon centers. The reaction takes advantage of a tethered 8-aminoquinoline directing group to control the regiochemical outcome and suppress two-component coupling between the N–O electrophile and organozinc nucleophile. **KEYWORDS:** nickel catalysis, C–N coupling, alkene, directing group, electrophilic nitrogen

Nitrogen-containing small-molecules comprise a significant portion of all known medicines.¹ Thus, novel methods for the formation of carbon–nitrogen (C–N) bonds have been actively pursued.² 1,2-Carboamination represents an appealing strategy for converting readily available alkene starting materials into valuable structurally complex amine products in an expedient manner (Figure 1). This transformation can be carried out using different modes of reactivity, including a classical-polarity approach in which the nitrogen-based reactant functions as a nucleophile (i.e., R_2NH) and an umpolung approach where the nitrogen-based reactant is an electrophile (i.e., R_2NX , X = halide or pseudohalide), as depicted in Figure 1A. Catalytic *intramolecular* (two-component) alkene carboamination involving both polarity types has been extensively studied.³ *Intermolecular* (three-component) variants, on the other hand, remain comparatively unexplored and have typically been limited to conjugated alkenes (e.g., styrenes or acrylates).⁴


In terms of precedents involving non-conjugated, unstrained alkenes,⁵ Liu and coworkers have reported palladium-catalyzed carbonylative 1,2-carboamination using 2-oxazolidone or phthalimide nucleophiles to afford terminal β -amino acids.^{5a} Later, this group reported a similar net transformation involving an azide-containing hypervalent iodine reagent.^{5b} These two reports rely on rapid migratory insertion of CO to outcompete side reactions, such as β -H elimination. Our group has reported a palladium-catalyzed directed 1,2-carboamination of unactivated alkenes via a classical polarity approach (Figure 1D).⁶ In particular, we demonstrated regioselective *anti*-addition of imides, amides, sulfonamides, and various azaheterocycles with aryl iodides across alkenes. These contributions notwithstanding, 1,2-carboamination of non-conjugated alkenes employing aliphatic amines and alkyl carbon coupling partners remain unexplored. The goal of the present study was to address this knowledge gap

through the development of a three-component umpolung carboamination of a non-conjugated alkene using a substrate directivity strategy.

A. general depiction of polarity types for alkene 1,2-carboamination




B. intramolecular umpolung 1,2-carboamination precedents




[Bower, 2015]: cat. Pd, B–[C] ([C] = aryl, acyl, vinyl, alkynyl)
[Zhu, 2017]: cat. Pd, H–[C] ([C] = 1,3,4-oxadiazoles)
[Selander, 2017]: cat. Ni, B–[C] ([C] = aryl)

C. intermolecular umpolung radical-based 1,2-carboamination of styrenes




[Zhang, 2013]: NFSI, TMSCN
[Liu, 2017]: F–NMe(SO₂Ph), B–[C] ([C] = aryl)

D. previous work: directed intermolecular classical 1,2-carboamination



[N]–H = imides, amides, sulfonamides, hydroxamic acid derivatives, and azaheterocycles

E. this work: directed intermolecular umpolung 1,2-carboamination




R^1 – R^4 = alkyl, X = R^4 or Br

Figure 1. Umpolung Carboamination Background.

Electrophilic aminating reagents have a rich history in enabling C–N bond formation.⁷ During the past few years, examples of umpolung carboamination of alkenes and allenes have been described.^{4a,4c} For example, building on seminal reports by Narasaka,⁸ Bower and coworkers described an intramolecular umpolung carboamination of γ,δ -unsaturated oxime esters with arylboronic ester coupling partners.⁹ The Zhu group later described analogous reactivity with 1,3,4-oxadiazole C–H nucleophiles (Figure 1B).¹⁰ Regarding intermolecular examples, in 2013 the Zhang group

described an umpolung radical-based copper-catalyzed aminocycloaddition of styrenes employing N–F reagents (Figure 1C).^{4a} Last year Liu and coworkers published an enantioselective aminoarylation of styrenes also catalyzed by copper using N–F reagents as electrophiles.^{4c}

Table 1. Selected Optimization of Reaction Conditions^a



entry	cat. Ni	Me ₂ Zn (equiv)	2a (equiv)	THF (M)	yield 3a (%)
1	20 mol% Ni(cod) ₂	1.0	1.0	0.10	71
2	20 mol% Ni(cod) ₂	2.0	1.0	0.10	15
3	20 mol% Ni(cod) ₂	3.0	1.0	0.10	n.d.
4	20 mol% Ni(cod) ₂	1.0	1.0	0.25	60
5	20 mol% Ni(cod) ₂	1.0	1.0	0.50	47
6	20 mol% Ni(cod) ₂	1.0	1.0	1.0	10
7	15 mol% Ni(cod) ₂	1.0	1.0	0.10	72
8	10 mol% Ni(cod) ₂	1.0	1.0	0.10	62
10	5 mol% Ni(cod) ₂	1.0	1.0	0.075	68
9	10 mol% Ni(cod) ₂	1.0	1.0	0.075	74
11	15 mol% Ni(cod) ₂	1.0	1.2	0.075	84 (79)
12	15 mol% NiCl ₂	1.0	1.2	0.075	80 (77)
13	15 mol% Ni(acac) ₂	1.0	1.2	0.075	75
14	15 mol% NiBr ₂ ·glyme	1.0	1.2	0.075	60
15	15 mol% NiBr ₂	1.0	1.2	0.075	n.d.

^aReaction conditions: **1a** (0.1 mmol), **2a** (1.0–1.2 equiv), Me₂Zn (1.0 M in heptane), 60 °C, 18–24 h. ¹H NMR yields reported with CH₂Br₂ as internal standard. Isolated yields in parentheses.


In contrast, analogous transformations involving the use of electrophilic aminating reagents in nickel-catalysis have been less extensively studied.^{7g,11} Two-component nickel-catalyzed C–N cross-couplings between organometallic nucleophiles and N–O electrophiles have been described by the Johnson,^{11a} Jarvo,^{11b} and Knochel groups.^{11c} To the best of our knowledge, only a single example of nickel-catalyzed alkene carboamination with organometallic nucleophiles has been reported to date, an intramolecular system developed by Selander and coworkers in 2017 (Figure 1B).^{7g,11c} Realization of an intermolecular nickel-catalyzed carboamination process would present the opportunity to rapidly generate medicinally motifs with dense functionality.

We recently described substrate-directed nickel-catalyzed three-component conjunctive cross-coupling reactions¹² that append differentiated alkyl/aryl fragments to β,γ- and γ,δ-unsaturated carbonyl compounds using aryl/alkyl halides and aryl/alkyl zinc reagents.¹³ The regioselectivity of these reactions is controlled by a tethered 8-aminoquinoline (AQ) directing group that stabilizes 5- or 6-membered metallacycles, thereby suppressing undesired side reactions, such as β-hydride elimination or two-component cross-coupling. Given these results we wondered if it would be possible to employ *O*-benzoylhydroxylamines as electrophiles in lieu of aryl/alkyl halides to synthesize β- and γ-amino acid derivatives under nickel catalysis. We surmised that this approach would complement our previous palladium(II)-catalyzed method (Figure 1D) in several respects.⁶ Namely it would be *syn*-selective, proceed with the opposite sense regioselectivity, enable use of alkyl coupling partners, and potentially be compatible with alkenes distal from the AQ group (Figure 1E).

To test this idea, we elected to use alkene **1a** as the pilot substrate given its unique effectiveness in earlier work^{13–14} and **2a** as the electrophilic nitrogen source based on its success in various

other catalytic methods. These starting materials were combined with commercially available dimethylzinc solution in the presence of catalytic nickel. With 20 mol % Ni(cod)₂ we observed formation of product **3a** (Table 1) in 71% yield (Table 1, Entry 1). DMF, toluene, acetonitrile, and dioxane were also tested under conditions otherwise identical to those in entry

Table 2. Scope of *O*-benzoylhydroxylamines and Organozinc reagents.^a




^aReaction conditions: **1a** (0.2 mmol), **2** (1.2 equiv), Me₂Zn (1.0 equiv, 1.0 M in heptane), THF (0.075 M). ^bEt₂Zn (1.0 M in hexanes). ^cR'ZnBr (2 equiv, 0.5 M in THF). ^d0.1 mmol scale. ^eCy₂Zn (1.0 equiv, 0.4 M in ether). ^fPh₂Zn (1.6 equiv, solid).

1. The reaction proceeded in toluene and dioxane, though yields were attenuated compared to in THF. Considering conditions from our previous work^{13b} we attempted to drive the reaction to completion by using excess Me₂Zn, but in this case we found significantly diminished yields when more than one equivalent was used. We also found that the reaction was higher yielding at lower concentrations, with the optimal concentration being 0.075 M **1a** in THF. Lower catalyst loadings of 10–15 mol % gave comparable yields, though decreasing the catalyst loading further (5 mol %) led to slightly diminished yield. Increasing the amount of **2a** to 1.2 equiv provided **3a** in 84% ¹H NMR yield (79% isolated). We also found that the reaction performs comparably well using several bench-stable Ni(II) salts, enabling a glove-box free protocol. The product **3a** was isolated in 77% yield using NiCl₂ as the precatalyst.

Having optimized the reaction conditions, we proceeded to explore the *O*-benzoylhydroxylamine electrophile scope (Table 2).

We found that several heterocyclic motifs (**3b–3g**) frequently found in bioactive compounds were well-tolerated, including thiomorpholine, *tert*-butoxycarbonyl-protected piperazine, 2-(piperazin-1-yl)pyrimidine, 4,5,6,7-tetrahydrothieno[3,2-*c*]pyridine, piperidine, and pyrrolidine. An array of N–O reagents derived from acyclic amines (**2h–2l**), including *N*-methyl-*N*-benzylamine, diethylamine, dibenzylamine and diallylamine also

Table 3. Scope of Alkene Substrates.^a



^aReaction conditions: **1a** (0.2 mmol), **2** (1.2 equiv), Me_2Zn (1.0 equiv, 1.0 M in heptane), THF (0.075 M). ^b0.1 mmol scale.

reacted under optimized conditions. Sterically hindered or especially reactive N–O reagents could not be used as coupling partners in this reaction; primary-amine-derived electrophiles were also not tolerated. (For a more extensive list of unsuccessful examples, see SI). The product **3l** was obtained in a similar yield using NiCl_2 , highlighting its efficacy as a substitute for Ni(cod)_2 .

The reaction was also compatible with a variety of di-organozinc and organozinc halide nucleophiles, though some reactions were found to proceed in diminished yields. In some cases this could be overcome by slow addition of the organozinc nucleophile, demonstrated in the synthesis of **3m**, which was isolated in 87% yield. Several other primary alkylzinc nucleophiles were compatible, including propyl (**3n**), ethyl propionate (**3o**) and benzyl (**3p**), providing the corresponding products in moderate yields. Secondary alkylzinc nucleophiles such as cyclobutyl (**3q**) and cyclohexyl (**3r**) could also be employed and provided moderate yields. More sterically congested secondary and tertiary carbon nucleophiles were unsuccessful under the optimized reaction conditions. We observed that monoalkylzinc halides were generally lower yielding than dialkylzinc reagents, likely due to their well-known attenuated nucleophilicity. We hypothesize that the need for excess alkylzinc halide (four times more than in the case of dialkylzinc reagents) is due to competitive reduction of the electrophile **2a** before it is able to react in the desired pathway,

leading to decreased yields. In the case of secondary nucleophiles, we believe competing β-hydride elimination pathways generates reducing species in solution that facilitate electrophile decomposition. We have also isolated aminoarylated product **3s** in 27% yield.

We also explored the scope of alkene substrates (Table 3) and found that the reaction was compatible with a variety of substituted alkenes. The relative stereochemistry of **4a** was determined by X-ray crystallography, establishing that the reaction proceeded in a *syn*-selective manner.¹³ The stereochemistry of other products derived from internal 1,2-




Figure 2. Reaction Scale-Up and 8-Aminoquinoline Removal.

disubstituted alkenes were assigned by analogy. Product **4a** was also obtained using NiCl_2 as the precatalyst and was obtained with similar yield. We also found that a phthalimide-protected amine could be tolerated under the reaction conditions to afford carboaminated product **4c** in moderate yield. Given the success of setting quaternary carbon centers in our previously published dialkylation reaction,^{13b} we wondered whether this carboamination reaction could also function in sterically congested environments. We were pleased to find tri- and 1,1-disubstituted alkenes could be used to synthesize compounds **4d** and **4e**, respectively, in good yields, demonstrating the ability of this method to forge quaternary carbon centers at either the β and γ position. The reaction also proceeded in moderate yields with α-methyl substituted alkenyl carbonyl compounds (**4f**), though we found the benzyl-substituted analogue to proceed in significantly reduced yields (<10% isolated). A representative cyclic alkene substrate was unreactive. Beyond β,γ-unsaturated alkenyl carbonyl compounds, we found the reaction could be extended to γ,δ-unsaturated substrates to afford products **4g–4i** and 2-vinylbenzamide-derived product **4j**.

We next performed the reaction on gram scale to demonstrate its synthetic utility. On 5-mmol scale, we were able to isolate 1.30 g of **3a** in 83% yield (Figure 2). We also validated two methods to remove the AQ directing group. Hydrolysis of **3a** afforded β-amino acid **3a'** in 76% yield. Using a method published by Ohshima and coworkers,¹⁵ methanolysis of **3a** afforded ester **3a''** in 79% yield. Furthermore, we found that a stereocenter at the carbon α to the carbonyl did not racemize under the reaction conditions (see SI for details). Moreover, it has previously been demonstrated that α-stereocenters do not epimerize under the aforementioned methanolysis conditions.¹⁵

Regarding the reaction mechanism, we surmised that two plausible redox manifolds could be operative, namely Ni(0)/Ni(II) or Ni(I)/Ni(III) catalysis (depicted in general form as Ni(n)/Ni(n+2)). Moreover, the reaction could proceed via two different orders of events (Figure 3A). In Pathway A, substrate-bound nickel complex **6c** would first undergo oxidative addition with **2** to form intermediate **6a**. Transmetalation followed by insertion to the alkene would form **6b**, which could reductively eliminate to form products **3–4** and regenerate the active catalyst. In the second potential mechanism, Pathway B, intermediate **6c** would first

react via transmetalation, after which migratory insertion would lead to intermediate **6d**. This species could then oxidatively add to **2** to give nickel intermediate **6e**. Reductive elimination would form the key C(sp³)–N bond and regenerate the catalytically active low-valent nickel species. A third mechanistic scenario (see SI) in which C–N bond formation precedes transmetalation and C–C reductive elimination cannot be conclusively ruled out at this stage, though we consider it to be less likely because it would involve formation of larger nickelacycles in preference to smaller nickelacycles with both classes of substrates (6 versus 5 with products **3**, and 7 versus 6 with products **4**).

Though a detailed investigation into the reaction mechanism is outside of the scope of the present study, we nevertheless performed a series of preliminary experiments in an effort to shed light on the nature of the oxidative addition step. We first prepared radical clock electrophile **2m** (Figure 3B). Based on literature precedents, the corresponding aminyl radical that would be formed if SET oxidative addition were operative^{11c,e} (as in a Ni(I)/Ni(III) cycle) is expected to cyclize with a first-order rate constant of approximately 10⁴ s^{−1}.¹⁶ When this electrophile was subjected to standard reaction conditions, only non-cyclized product **3t** was formed in 40% yield. No evidence of cyclization was observed

interpreting this type of experiment due to the ability of the radical inhibitor to modify the catalyst²¹ among other possible side reactions. Nevertheless, if one assumes sufficiently clean reaction profiles and approximates trapping rates for N-centered radical based on available data for C-centered radicals,¹⁸ then these results would corroborate the radical clock experiment in terms of the rates that would be required for an SET pathway to be operative. Though these preliminary mechanistic studies were largely inconclusive, the results and the accompanying analysis will nevertheless serve as a useful foundation for more detailed studies in the future.

In summary, we have developed an intramolecular umpolung carboamination of non-conjugated alkenes that affords a variety of β- and γ-amino acid and ester derivatives. The reaction is enabled by a removable 8-aminoquinoline tethered directing group, which facilitates formation of stabilized 5- or 6-membered nickelacycles, suppresses β-hydride elimination and two-component coupling, and determines the regiochemical outcome. The reaction tolerates a range of alkenes with various substitution patterns and proceeds in the presence of several synthetically important functional groups.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge via the Internet at <http://pubs.acs.org>.

Experimental details, analytical data, and ¹H and ¹³C NMR spectra (PDF)

X-ray data for compound **3a** (CIF)

X-ray data for data (±)-**4a** (CIF)

NMR data (ZIP)




Figure 3. Plausible Pathways and Preliminary Mechanistic Experiments. ^a See Ref. 18 for precedent regarding the background reaction between ZnMe₂ and TEMPO.

by ¹H NMR of the crude reaction mixture. This result is consistent with a two-electron oxidation addition pathway or alternatively with an SET oxidation pathway involving a radical recombination step with a sufficiently large second-order rate constant (>10⁴ M^{−1}s^{−1}).¹⁷

The effect of radical inhibitors was next studied (Figure 3C).¹⁸ The reaction was not inhibited by the addition of BHT (1 equiv). On the other hand, addition of TEMPO (1 equiv) dramatically suppressed product formation; it should be emphasized, however, that interpretation of this latter result is complicated by a fast background reaction between TEMPO and dimethylzinc,¹⁹ as evidenced by formation of TEMPO–Me (¹H NMR and LC-MS). To gain more clarity, we performed an analogous experiment with a representative alkylzinc bromide (cyclobutylzinc bromide). In this case, we again observed almost complete inhibition of the carboamination reaction, this time without concomitant formation of the alkyl–TEMPO adduct (see SI).²⁰ Caution should be taken in

AUTHOR INFORMATION

Corresponding Author

*keary@scripps.edu

ACKNOWLEDGMENT

This work was financially supported by The Scripps Research Institute (TSRI), Pfizer, Inc., Bristol-Myers Squibb (Unrestricted Grant), and the National Science Foundation (CHE-1800280). We further acknowledge the NSF for a Graduate Research Fellowship (NSF/DGE-1346837, J.D.) and a Summer Undergraduate Research Fellowship (NSF/DBI-1359160, V.A.V.). We thank Dr. De-Wei Gao, Mingyu Liu, and Van T. Tran for preparation of alkene substrates and nitrogen electrophiles. Professor Nathan D. Schley (Vanderbilt) is acknowledged for helpful discussion regarding the reaction mechanism. We also thank Professor Arnold L. Rheingold for assistance with X-ray crystallography.

REFERENCES

- (1) (a) Henkel, T.; Brunne, R. M.; Müller, H.; Reichel, F. Statistical Investigation into the Structural Complementarity of Natural Products and Synthetic Compounds. *Angew. Chem. Int. Ed.* **1999**, *38*, 643–647; (b) Vitaku, E.; Smith, D. T.; Njardarson, J. T. Analysis of the Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles Among U.S. FDA Approved Pharmaceuticals. *J. Med. Chem.* **2014**, *57*, 10257–10274.
- (2) (a) Hartwig, J. F. Transition Metal Catalyzed Synthesis of Arylamines and Aryl Ethers from Aryl Halides and Triflates: Scope and Mechanism. *Angew. Chem. Int. Ed.* **1998**, *37*, 2046–2067; (b) Müller, T. E.; Beller, M. Metal-Initiated Amination of Alkenes and Alkynes. *Chem. Rev.* **1998**, *98*, 675–703; (c) Wolfe, J. P.; Wagaw, S.; Marcoux, J.-F.;

- Buchwald, S. L. Rational Development of Practical Catalysts for Aromatic Carbon–Nitrogen Bond Formation. *Acc. Chem. Res.* **1998**, *31*, 805–818;
- (d) Ley, S. V.; Thomas, A. W. Modern Synthetic Methods for Copper-Mediated C(aryl)–O, C(aryl)–N, and C(aryl)–S Bond Formation. *Angew. Chem. Int. Ed.* **2003**, *42*, 5400–5449; (e) Beller, M.; Seayad, J.; Tillack, A.; Jiao, H. Catalytic Markovnikov and anti-Markovnikov Functionalization of Alkenes and Alkynes: Recent Developments and Trends. *Angew. Chem. Int. Ed.* **2004**, *43*, 3368–3398; (f) Wijtmans, R.; Vink, M. K. S.; Schoemaker, H. E.; Delft, F. L. v.; Blaauw, R. H.; Rutje, F. P. J. T. Biological Relevance and Synthesis of C-Substituted Morpholine Derivatives. *Synthesis* **2004**, *5*, 641–662; (g) Hartwig, J. F. Carbon–Heteroatom Bond Formation Catalysed by Organometallic Complexes. *Nature* **2008**, *455*, 314–322; (h) Beriwal, J.; Eycken, E. V. d. C–N Bond Forming Cross-Coupling Reactions: An Overview. *Chem. Soc. Rev.* **2013**, *42*, 9283–9303; (i) Ruiz-Castillo, P.; Buchwald, S. L. Applications of Palladium-Catalyzed C–N Cross-Coupling Reactions. *Chem. Rev.* **2016**, *116*, 12564–12649; (j) Dai, M.; Ye, Z.; Gettys, K. E. Recent Advances in Piperazine Synthesis. *Synthesis* **2017**, *49*, 2589–2604.
- (3) (a) Chemler, S. R. The Enantioselective Intramolecular Aminative Functionalization of Unactivated Alkenes, Dienes, Allenes and Alkynes for the Synthesis of Chiral Nitrogen Heterocycles. *Org. Biomol. Chem.* **2009**, *7*, 3009–3019; (b) McDonald, R. I.; Liu, G.; Stahl, S. S. Palladium(II)-Catalyzed Alkene Functionalization via Nucleopalladation: Stereochemical Pathways and Enantioselective Catalytic Applications. *Chem. Rev.* **2011**, *111*, 2981–3019; (c) Schultz, D. M.; Wolfe, J. P. Recent Developments in Palladium-Catalyzed Alkene Aminoarylation Reactions for the Synthesis of Nitrogen Heterocycles. *Synthesis* **2012**, *44*, 351–361; (d) Garlets, Z. J.; White, D. R.; Wolfe, J. P. Recent Developments in Pd(0)-Catalyzed Alkene–Carboheterofunctionalization Reactions. *Asian J. Org. Chem.* **2016**, *6*, 636–653; (e) Belov, G. M.; Belove, D. S.; Kurkin, A. V.; Yurovskaya, M. A. Palladium-Catalyzed Carboamination in the Synthesis of Nitrogen-Containing Heterocycles. *Chem. Heterocycl. Comp.* **2012**, *48*, 73–81; (f) Shimizu, Y.; Kanai, M. Recent Progress in Copper-Catalyzed Difunctionalization of Unactivated Carbon–Carbon Multiple Bonds. *Tet. Lett.* **2014**, *55*, 3727–3737; (g) Wolfe, J. P. Stereoselective Synthesis of Saturated Heterocycles via Palladium-Catalyzed Alkene Carboetherification and Carboamination Reactions. *Synlett* **2008**, *19*, 2913–2937; (h) Wolfe, J. P. Palladium-Catalyzed Carboetherification and Carboamination Reactions of γ -Hydroxy- and γ -Aminoalkenes for the Synthesis of Tetrahydrofurans and Pyrrolidines. *Eur. J. Org. Chem.* **2007**, 571–582.
- (4) (a) Zhang, H.; Pu, W.; Xiong, T.; Li, Y.; Zhou, X.; Zun, K.; Liu, Q.; Zhang, Q. Copper-Catalyzed Intermolecular Aminocyanation and Diamination of Alkenes. *Angew. Chem. Int. Ed.* **2013**, *52*, 2529–2533; (b) Liu, Y.-Y.; Yang, X.-H.; Song, R.-J.; Luo, S.; Li, J.-H. Oxidative 1,2-Carboamination of Alkenes with Alkyl Nitriles and Amines Toward γ -Amino Alkyl Nitriles. *Nat. Comm.* **2017**, *8*, 14720; (c) Wang, D.; Wu, L.; Wang, F.; Wan, X.; Chen, P.; Lin, Z.; Liu, G. Asymmetric Copper-Catalyzed Intermolecular Aminoarylation of Styrenes: Efficient Access to Optical 2,2-Diarylethylamines. *J. Am. Chem. Soc.* **2017**, *139*, 6811–6814; (d) Wang, D.; Wang, F.; Chen, P.; Lin, Z.; Liu, G. Enantioselective Copper-Catalyzed Intermolecular Amino- and Azidocyanation of Alkenes in a Radical Process. *Angew. Chem. Int. Ed.* **2017**, *56*, 2054–2058; (e) Qian, B.; Chen, S.; Wang, T.; Zhang, X.; Bao, H. Iron-Catalyzed Carboamination of Olefins: Synthesis of Amines and Disubstituted β -Amino Acids. *J. Am. Chem. Soc.* **2017**, *139*, 13076–13082; (f) Gockel, S. N.; Buchanan, T. L.; Hull, K. L. Cu-Catalyzed Three-Component Carboamination of Alkenes. *J. Am. Chem. Soc.* **2018**, *140*, 58–61.
- (5) (a) Cheng, J.; Qi, X.; Li, M.; Chen, P.; Liu, G. Palladium-Catalyzed Intermolecular Aminocarbonylation of Alkenes: Efficient Access of β -Amino Acid Derivatives. *J. Am. Chem. Soc.* **2015**, *137*, 2480–2483; (b) Li, M.; Yu, F.; Chen, P.; Liu, G. Palladium-Catalyzed Intermolecular Azidocarbonylation of Alkenes via a Cooperative Strategy. *J. Org. Chem.* **2017**, *82*, 11682–11690.
- (6) Liu, Z.; Wang, Y.; Wang, Z.; Zeng, T.; Liu, P.; Engle, K. M. Catalytic Intermolecular Carboamination of Unactivated Alkenes via Directed Aminopalladation. *J. Am. Chem. Soc.* **2017**, *139*, 11261–11270.
- (7) (a) Kovacic, P.; Lowery, M. K.; Field, K. W. Chemistry of N-Bromoamines and N-Chloroamines. *Chem. Rev.* **1970**, *70*, 639–665; (b) Erdik, E.; Ay, M. Electrophilic Amination of Carbanions. *Chem. Rev.* **1989**, *89*, 1947–1980; (c) Ciganek, E., Electrophilic Amination of Carbanions, Enolates and Their Surrogates. In *Organic Reactions*, Hoboken, NJ, United States, 2008; Vol. 72, pp 1–366; (d) Barker, T. J.; Jarvo, E. R. Developments in Transition-Metal-Catalyzed Reactions Using Electrophilic Nitrogen Sources. *Synthesis* **2011**, *24*, 3954–3964; (e) Corpet, M.; Gosmini, C. Recent Advances in Electrophilic Amination Reactions. *Synthesis* **2014**, *46*, 2258–2271; (f) Yan, X.; Yang, X.; Xi, C. Recent Progress in Copper-Catalyzed Electrophilic Amination. *Catal. Sci. Technol.* **2014**, *4*, 4169–4177; (g) Dong, X.; Liu, Q.; Dong, Y.; Liu, H. Transition-Metal-Catalyzed Electrophilic Amination: Application of O-Benzoylhydroxylamines in the Construction of the C–N Bond. *Chem. Eur. J.* **2017**, *23*, 2481–2511; (h) Hendrick, C. E.; Wang, Q. Emerging Developments Using Nitrogen–Heteroatom Bonds as Amination Reagents in the Synthesis of Aminoarenes. *J. Org. Chem.* **2017**, *82*, 839–847.
- (8) (a) Tsutsui, H.; Narasaka, K. Synthesis of Pyrrole Derivatives by the Heck-Type Cyclization of γ,δ -Unsaturated ketone O-Pentafluorobenzoyloximes. *Chem. Lett.* **1998**, *28*, 45–46; (b) Kitamura, M.; Zaman, S.; Narasaka, K. Synthesis of Spiro Imines from Oximes by Palladium-Catalyzed Cascade Reaction. *Synlett* **2001**, 974–976; (c) Tsutsui, H.; Kitamura, M.; Narasaka, K. Synthesis of Pyrrole Derivatives by Palladium-Catalyzed Cyclization of γ,δ -Unsaturated Ketone O-Pentafluorobenzoyloximes. *Bull. Chem. Soc. Jpn.* **2002**, *75*, 1451–1460; (d) Zaman, S.; Kitamura, M.; Narasaka, K. Synthesis of Polycyclic Imines by Palladium-Catalyzed Domino Cyclization of Di- and Trienyl Ketone O-Pentafluorobenzoyloximes. *Bull. Chem. Soc. Jpn.* **2003**, *76*, 1055–1062; (e) Kitamura, M.; Narasaka, K. Synthesis of Aza-Heterocycles from Oximes by Amino-Heck Reaction. *Chem. Rec.* **2002**, *2*, 268–277.
- (9) Faulkner, A.; Scott, J. S.; Bower, J. F. An Umpolung Approach to Alkene Carboamination: Palladium Catalyzed 1,2-Amino-Acylation, –Carboxylation, –Arylation, –Vinylolation, and –Alkynylation. *J. Am. Chem. Soc.* **2015**, *137*, 7224–7230.
- (10) Bao, X.; Wang, Q.; Zhu, J. Palladium-Catalyzed Enantioselective Narasaka–Heck Reaction/Direct C–H Alkylation of Arenes: Iminoarylation of Alkenes. *Angew. Chem. Int. Ed.* **2017**, *56*, 9577–9581.
- (11) (a) Berman, A. M.; Johnson, J. S. Nickel-Catalyzed Electrophilic Amination of Organozinc Halides. *Synlett* **2005**, *11*, 1799–1801; (b) Barker, T. J.; Jarvo, E. R. Umpolung Amination: Nickel-Catalyzed Coupling Reactions of *N,N*-Dialkyl-*N*-Chloroamines with Diorganotin Reagents. *J. Am. Chem. Soc.* **2009**, *131*, 15598–15599; (c) Gu, Y.-R.; Duan, X.-H.; Yang, L.; Guo, L.-N. Direct C–H Cyanoalkylation of Heteroaromatic N-Oxides and Quinones via C–C Bond Cleavage of Cyclobutanone Oximes. *Org. Lett.* **2017**, *19*, 5908–5911; (d) Chen, Y.-H.; Graßl, S.; Knochel, P. Cobalt-Catalyzed Electrophilic Amination of Aryl- and Heteroarylzinc Pivalates with *N*-Hydroxylamine Benzoates. *Angew. Chem. Int. Ed.* **2017**, *56*, 1108–1111; (e) Yang, H.-B.; Pathipati, S. R.; Selander, N. Nickel-Catalyzed 1,2-Aminoarylation of Oxime Ester-Tethered Alkenes with Boronic Acids. *ACS Catal.* **2017**, *7*, 8441–8445; (f) Xiao, J.; He, Y.; Ye, F.; Zhu, S. Remote sp^3 C–H Amination of Alkenes with Nitroarenes. *Chem* **2018**, *4*, 1–13. (g) Wang, L.; Wang, C. Ni-Catalyzed 1,2-iminoacylation of alkenes via a reductive strategy. *Org. Chem. Front.* **2018**, *5*, 3476–3482.
- (12) Derosa, J.; Tran, V. T.; van der Puyl, V. A.; Engle, K. M. Carbon–Carbon π bonds as Conjugative Reagents in Cross-Coupling. *Aldrichimica Acta* **2018**, *51*, 21–32.
- (13) (a) Derosa, J.; Tran, V. T.; Boulous, M. N.; Chen, J. S.; Engle, K. M. Nickel-Catalyzed, β -Dicarbofunctionalization of Alkenyl Compounds via Conjugative Cross-Coupling. *J. Am. Chem. Soc.* **2017**, *139*, 10657–10660; (b) Derosa, J.; van der Puyl, V. A.; Tran, V. T.; Liu, M.; Engle, K. M. Directed Nickel-Catalyzed 1,2-Dialkylation of Alkenes. *Chem. Sci.* **2018**, *9*, 5278–5283.
- (14) Lv, H.; Xiao, L.-J.; Zhao, D.; Zhou, Q.-L. Nickel(0)-Catalyzed Linear-Selective Hydroarylation of Unactivated Alkenes and Styrenes with Aryl Boronic Acids. *Chem Sci* **2018**, *9*, 6839–6843.
- (15) Deguchi, T.; Xin, H.-L.; Morimoto, H.; Ohshima, T. Direct Catalytic Alcoholysis of Unactivated 8-Aminoquinoline Amides. *ACS Catal.* **2017**, *7*, 3157–3161.
- (16) Newcomb, M., Radical Kinetics and Clocks. In *Encyclopedia of Radicals in Chemistry, Biology and Materials*, Chatgililoglu, C.; Studer, A., Eds. John Wiley & Sons, Ltd.: 2012.
- (17) Faster radical clocks were also considered. In particular, appropriately substituted aminocyclopropane and aminocyclobutane N-centered radicals would be expected to undergo ring opening with first-order rate constants on the order of 10^8 s $^{-1}$ and 10^{11} s $^{-1}$, respectively (see Ref. 16). However, we found that a model cyclobutylamine-derived electrophile was unreactive under standard conditions (see SI), presumably due to steric hindrance, which precluded this line of inquiry.
- (18) BHT reacts with primary alkyl radicals with a second-order rate constant of roughly 10^3 M $^{-1}$ s $^{-1}$: (a) Franchi, P.; Lucarini, M.; Pedulli, G. F.; Valgimigli, L.; Lunelli, B. Reactivity of Substituted Phenols Toward

Alkyl Radicals. *J. Am. Chem. Soc.* **1999**, *121*, 507–514. In comparison, TEMPO reacts with primary alkyl radicals with a second-order rate constant of roughly $10^9 \text{ M}^{-1}\text{s}^{-1}$: (b) Newcomb, M. Competition Methods and Scales for Alkyl Radical Reaction Kinetics. *Tetrahedron* **1993**, *49*, 1151–1176. To our knowledge trapping corresponding trapping rates for these reagents with nitrogen-centered radicals have not been established.

(19) Budny-Godlewski, K.; Kubicki, D.; Justyniak, I.; Lewiński, J. A New Look at the Reactivity of TEMPO toward Diethylzinc. *Organometallics* **2014**, *33*, 5093–5096.

(20) Alkylzinc bromides have been previously used in radical trapping experiments with nickel and are presumed to have an attenuated background reaction rate with TEMPO compared to dialkylzinc reagents. For leading references, see: (a) Liang, Y.; Fu, G. C. Nickel-Catalyzed Alkyl–Alkyl Cross-Couplings of Fluorinated Secondary Electrophiles: A General Approach to the Synthesis of Compounds having a Perfluoroalkyl Substit-

uent. *Angew. Chem. Int. Ed.* **2015**, *54*, 9047; (b) Soler-Yanes, R.; Arribas-Álvarez, I.; Guisán-Ceinos, M.; Buñuel, E.; Cárdenas, D. J. NiII Catalyzes the Regioselective Cross-Coupling of Alkylzinc Halides and Propargyl Bromides to Allenes. *Chem. Eur. J.* **2017**, *23*, 1584–1590.

(21) For an example where this has been clearly demonstrated, see: Schley, N. D.; Fu, G. C. Nickel-Catalyzed Negishi Arylation of Propargylic Bromides: A Mechanistic Investigation. *J. Am. Chem. Soc.* **2014**, *136*, 16588–16593.

