

# Is Computational Oceanography Coming of Age?

Thomas W. N. Haine\*, Renske Gelderloos, Miguel A. Jimenez-Urias, Ali H. Siddiqui,

*Earth & Planetary Sciences, Johns Hopkins University, Baltimore, MD*

Gerard Lemson, Dimitri Medvedev, Alex Szalay,

*Physics & Astronomy, Johns Hopkins University, Baltimore, MD*

Ryan P. Abernathey,

*Earth & Environmental Sciences, Columbia University, New York, NY*

Mattia Almansì,

*National Oceanography Centre, Southampton, UK*

Christopher N. Hill

*Earth, Atmospheric, & Planetary Sciences, MIT, Cambridge, MA*

<sup>12</sup> \*Corresponding author: Thomas Haine, Thomas.Haine@jhu.edu

## ABSTRACT

13 Computational Oceanography is the study of ocean phenomena by numerical simulation, especially  
14 dynamical and physical phenomena. Progress in information technology has driven exponential  
15 growth in the number of global ocean observations and the fidelity of numerical simulations of the  
16 ocean in the past few decades. The growth has been exponentially faster for ocean simulations,  
17 however. We argue that this faster growth is shifting the importance of field measurements and  
18 numerical simulations for oceanographic research. It is leading to the maturation of Computational  
19 Oceanography as a branch of marine science on par with observational oceanography. One  
20 implication is that ultra-resolved ocean simulations are only loosely constrained by observations.  
21 Another implication is that barriers to analyzing the output of such simulations should be removed.  
22 Although some specific limits and challenges exist, many opportunities are identified for the future  
23 of Computational Oceanography. Most important is the prospect of hybrid computational and  
24 observational approaches to advance understanding of the ocean.

25 *Capsule summary.* Fast growth in the fidelity of ocean general circulation models is driving the  
26 maturation of Computational Oceanography as a branch of marine science on par with observational  
27 oceanography.

## 28 **1. Introduction**

29 Computational Oceanography is the study of ocean phenomena by numerical simulation, es-  
30 pecially dynamical and physical phenomena using ocean general circulation models (OGCMs).  
31 One early pioneer of this field wrote of the 1960s, 1970s, and 1980s as the “birth”, “infancy”,  
32 and “adolescence” of OGCMs, respectively (Bryan 2006, see also Holland and McWilliams 1987;  
33 McWilliams 1996). Similarly, the authors of a comprehensive review of OGCMs wrote at the turn  
34 of the century “this field...has entered an era of healthy adolescence” (Griffies et al. 2000). With  
35 twenty more years of data, this essay explores the continued growth of OGCMs and speculates  
36 on their prospects. We ask: Is Computational Oceanography entering a new era that signifies its  
37 coming of age?

38 For motivation, Fig. 1 compares oceanographic measurements and results from a high-resolution  
39 OGCM. The region of interest is a topographic constriction called the Denmark Strait, between  
40 Greenland and Iceland. The Denmark Strait Overflow (DSO) flows south through this gap and is  
41 an important current for the Atlantic Meridional Overturning Circulation and thus for the ocean’s  
42 role in North Atlantic climate. The two timeseries in Fig. 1a show DSO volume flux (transport).  
43 One timeseries is from *in situ* measurements, the other is from a high resolution regional OGCM  
44 (and they have been processed similarly with similar smoothing). The question is this: Which is  
45 which? Fig. 1b compares *in situ* hydrographic measurements along a section north of Denmark  
46 Strait with a synthetic hydrographic section from the OGCM. And Fig. 1c shows the trajectories  
47 of drifting oceanographic floats approaching Denmark Strait from the north and trajectories of

48 drifting particles in the OGCM released from the same locations. Again, the question is which is  
49 the real data and which is the synthetic data? In each case, the field measurements and the OGCM  
50 results are different, but identifying them is difficult.

51 These are examples of OGCM Turing tests. They are inspired by Alan Turing’s imitation game to  
52 distinguish between, and correctly identify, a person and an intelligent machine. The game involves  
53 asking questions through an interface that obscures whether the responses are from the person or  
54 the machine (Turing 1950). The difficulty of the OGCM Turing tests in Fig. 1 reflects the small  
55 systematic error in the OGCM and therefore its realism. Some OGCM solutions are reaching the  
56 point that they are essentially indistinguishable from observations, so they pass Turing tests like  
57 those in Fig. 1. In the words of Ed Lorenz, numerical experiments will eventually “duplicate the  
58 circulation to any desired degree of accuracy” (Lorenz 1967).<sup>1</sup>

59 With these themes in mind, this essay explores the growth of OGCMs and compares it to the  
60 growth of ocean observations. The focus is on the computer science and information technology  
61 improvements that contribute to the growth. We then speculate on limits, opportunities, and  
62 prospects for OGCMs.

## 63 **2. Unequal Exponential Growth**

64 Two examples illustrate the growth of ocean observations. First, consider temperature observa-  
65 tions in the global deep ocean over the last half century. Fig. 2a shows the cumulative number  
66 of temperature observations deeper than 1000m. They have grown exponentially (notice the y  
67 axis is logarithmic). Averaged over the last century, the exponential growth has a doubling period  
68 of 10.4 years, giving an approximately 60-fold expansion in the deep temperature database since  
69 1960. Technology transitions have maintained this exponential growth, specifically, advances

---

<sup>1</sup>The prescient Lorenz was writing about atmospheric models in the late 1960s, but the message applies to OGCMs today.

70 in micro-electronics and information technology. In the 1990s conductivity-temperature-depth  
71 (CTD) sensors on autonomous profiling floats took over from ship CTD sampling, for example,  
72 leading in the 2000s to the transformative Argo global float network (Argo 2020).

73 Second, consider the history of sea level observations from satellite altimeters. Sea level data  
74 have revolutionized physical oceanography by providing information on the surface circulation,  
75 mesoscale eddies, tides, and sea level change. Fig. 2b shows the sequence of altimeter missions  
76 (colored bars) and the cumulative number of observing days (black line). The number of observing  
77 days reveals the growth in sea level observations (although there is great variety between missions).  
78 The number of sea level observations has grown nearly exponentially since the mid 1980s with a  
79 doubling time of about 8.1 years and a  $\approx 20$ -fold expansion in the sea level database since 1985.  
80 Again, micro-electronic and information technology advances have maintained this growth.

81 Technology advances have also fueled growth in the fidelity of OGCMs. For example, Fig. 2c  
82 shows the history of global OGCM resolution. The black dots show five pioneering (cutting-edge)  
83 models over the last 40 years. The Bryan and Lewis (1979) model had a peak resolution of  $2.4^\circ$   
84 with 12 vertical levels and the Rocha et al. (2016) model had a peak resolution of  $0.02^\circ$  with 90  
85 vertical levels. The growth in OGCM resolution (number of grid points) is exponential with a  
86 doubling time of 2.2 years and a  $10^5$ -fold increase since 1980. We also show the global ocean  
87 models from the Intergovernmental Panel on Climate Change (IPCC) reports. The peak resolution  
88 of the ocean OGCMs in the first IPCC report was  $2.7^\circ$  with 9 vertical levels and the peak resolution  
89 in the latest (sixth) IPCC report is  $0.067^\circ$  with 75 vertical levels.<sup>2</sup> This growth is also exponential  
90 with a doubling time of 2.8 years. For the most highly-resolved models in each assessment, the  
91 doubling time is close to the cutting-edge OGCM doubling time.

---

<sup>2</sup>The AR6 data points on Fig. 2 are from the HighResMIP experiments, which is a sub-project on high-resolution models that does not run the full suite of CMIP6 experiments.

Now compare the horizontal resolution of ocean measurements with OGCM resolution. The Argo profiling float network operates about 4000 floats at any one time. Each float makes a vertical profile from 2000m depth to the surface every ten days. The global average spacing of profiles is therefore 300 km.<sup>3</sup> The spacing between altimeter tracks for the TOPEX/Poseidon and Jason satellite altimeters is also about 315 km (at the equator), with a repeat period of ten days. The present day peak OGCM resolution of  $0.02^\circ \approx 2$  km is therefore 140 times higher.<sup>4</sup>

### 3. Prospects for Future Growth

Looking ahead, the future is bright for the Argo network. The reason is that Argo is part of the Global Ocean and Global Climate Observing System, which implements the Paris Agreement on climate change and United Nations sustainable development goals. New capacities, like deep profiling floats, and new technologies, like biogeochemical sensors, are planned over the next few years (GCOS 2016). It is unclear how the network can double in size in the next decade and maintain long term exponential growth, but it is plausible.

The future is also bright for sea level measurements. The Surface Water and Ocean Topography (SWOT) mission, scheduled for launch in 2022, will start a new era of sea level observation. SWOT will observe sea level over a swath, rather than over a single patch. It will have 15 km resolution, or better, covering most of the global ocean every 21 days (Morrow et al. 2019). It will improve the spatial resolution of sea level data by a factor of about ten. Therefore, the prospects for the altimetry record to continue growing exponentially in the 2020s are good.

---

<sup>3</sup>The vertical resolution of Argo profile data is about 5m, which is about 7 times higher than the best AR6 OGCMs and about 3 times higher than the Poseidon Project run mentioned below.

<sup>4</sup>This comparison avoids the issue of time dependence in the circulation. It simply (and conservatively) imagines the Argo and altimetry data from one ten day period are used to constrain the time-mean OGCM state over that period.

111 For OGCMs, resolution improves as supercomputer technology advances. Historically, that  
112 follows Moore’s “law,” which says that transistor density in microprocessors doubles every two  
113 years (Moore 1975). For instance, machines first achieved petaflop speeds ( $10^{15}$  floating point  
114 operations per second) in 2008 and exaflop speeds ( $10^{18}$ ) in March 2020, a doubling every 1.1 years  
115 (see Fig. 2d). Computers available to the oceanographic, atmospheric, and climate communities  
116 are less powerful. Still, the machines at NCAR and ECMWF<sup>5</sup> also show exponential growth over  
117 recent decades with a doubling every 1.1 years, albeit lagging the cutting-edge machines by about  
118 five years (Fig. 2d). On this basis, the OGCM resolution will probably continue to double every 2.2  
119 years, at least for several more years (assuming funding remains at historic levels). It is reasonable  
120 to expect cutting-edge exascale global OGCMs with horizontal resolution around one kilometer  
121 by the mid 2020s. After that, with widespread anticipation that Moore’s law will end (Waldrop  
122 2016), future growth is uncertain.

#### 123 **4. Maturation of Computational Oceanography**

124 This evidence shows that information technology advances are driving exponential growth in  
125 ocean observations and exponential growth in OGCM resolution. But the OGCM growth rate is  
126 faster. Therefore, OGCM resolution is also growing exponentially faster than the growth in ocean  
127 field data. In 1990, OGCMs were obviously biased compared to measurements, for example, of  
128 deep temperatures or sea level. In 2020, OGCMs are achieving resolutions that are substantially  
129 greater than the gaps between measurements, at least for some regimes, like deep and abyssal  
130 ocean currents. We should expect this trend to continue for the foreseeable future (the next  
131 several years). Therefore, the question arises: When, and in what ways, will OGCMs become as  
132 important as observations for advancing knowledge in physical oceanography? Historically, most

---

<sup>5</sup>Meaning the U.S. National Center for Atmospheric Research and the European Centre for Medium Range Weather Forecasts.

knowledge came from observations of the real ocean.<sup>6</sup> The growth of OGCMs suggests that the field is approaching an era in which numerical circulation models are as important as observations for advancing knowledge. For example, diagnosing and understanding the rectified effects of mesoscale eddy variability on the large-scale, low-frequency circulation will probably rely heavily on high-resolution OGCMs.

What are the criteria to claim that OGCM solutions should be treated, in some cases, as seriously as real measurements? Realizing them would mark the maturation of Computational Oceanography. These criteria are on our checklist:

**1. Confidence in the fidelity of the basic tools and methods.** Consider two types of tool:

First, consider the theoretical definition of the ocean circulation problem. Computational Oceanography relies on software to compute approximations to the incompressible rotating-stratified Navier Stokes equations, with equations for the conservation of dissolved salts and heat (McWilliams 1996; Griffies 2004; Fox-Kemper et al. 2019). There is little doubt that these are the right equations for ocean circulation. The software is mature, stable, and diverse. The issue of unresolved processes, and parametrizing their effects remains an important area of research. For example, it is still unclear how to represent the impacts of unresolved submesoscale processes on the larger scale flow. Although much progress has been made on this problem in the last 30 years (Gent 2011; Le Sommer et al. 2018; Fox-Kemper et al. 2019), resolution improvements have surely played an essential part in refining OGCM accuracy (Griffies et al. 2000). In other words, we believe that the problem of parametrizing unresolved

---

<sup>6</sup>For example, Stewart (2008) writes: “The theory describing a convecting, wind-forced, turbulent fluid in a rotating coordinate system has never been sufficiently well known that important features of the oceanic circulation could be predicted before they were observed. In almost all cases, oceanographers resort to observations to understand oceanic processes.”

scales is not so pathological that it contaminates all of the resolved scales.<sup>7</sup> A corollary is that OGCMs are less complicated than the real ocean, meaning that OGCM variability is a lower bound on the variability in the real system. These are *de facto* working hypotheses of all theoretical and numerical approaches to understanding the ocean circulation.

Second, we need tools to adjust OGCM solutions to agree with observations; that is, to solve the data assimilation and state estimation problem (Bennett 1992; Wunsch 1996, 2006; Kalnay 2002). For example, state estimation is used to produce retrospective reanalyses (hindcasts) of the time-evolving ocean state and data assimilation is used to initialize prospective forecasts of the future. Although many questions remain open, these methods are also now mature, stable, and diverse.

2. **The number of OGCM degrees of freedom exceeds the number of observational constraints.** This criterion concerns the state estimation and data assimilation problems. In essence, it is about whether it is possible (in principle) to adjust an OGCM solution to fit the observations exactly or not. If the OGCM can be adjusted to fit the data exactly, the state estimation problem is under-determined. Otherwise, it is over-determined.<sup>8</sup> The number of OGCM degrees of freedom scales as the number of grid points (for large numbers of grid points). The number of observational constraints scales as the number of distinct measurements. Fig. 2 shows evidence that the number of OGCM degrees of freedom per observational constraint exceeds one because, loosely, the peak OGCM resolution is now 140

---

<sup>7</sup>It is likely that errors in parameterized physics *influence* all resolved scales, not least because of error growth due to deterministic chaos. But the issue is whether the errors in parameterized physics cause systematic errors in the resolved scales, such as biases in statistics of resolved quantities. It is reasonable to suppose that (i) resolution improvements and parameterization improvements reduce these systematic biases towards zero, and (ii) the systematic biases are not so bad as to preclude use of models to understand (and hindcast and predict) the natural system. Of course, these are quantitative (not qualitative) hypotheses that vary from case to case (models, parameterizations, resolved metrics, science questions).

<sup>8</sup>Ignoring the atypical case of the problem being exactly determined.

172 times higher than the Argo and Jason data spacing (see footnote 4). This gap is growing  
173 exponentially because OGCM resolution is growing exponentially faster than data density.  
174 Therefore, the state estimation problem is moving from (in principle) being over-determined  
175 to being under-determined.<sup>9</sup>

176 Crossing this threshold has interesting implications: First, the systematic errors in OGCMs  
177 disappear and they pass Turing or Feigenbaum tests (Turing 1950; Feigenbaum 2003; Harel  
178 2005), like those in Fig. 1. That is, OGCM solutions become indistinguishable from obser-  
179 vations of the real ocean and a subject-matter expert cannot tell them apart. Regional OGCM  
180 simulations of the Denmark Strait Overflow (DSO) at resolutions of 0.5–2 km are approach-  
181 ing this point (Magaldi and Haine 2015; Almansi et al. 2020; Saberi et al. 2020). Similarly,  
182 regional high-resolution state estimates are nearly under-determined (Lea et al. 2006; Dwivedi  
183 et al. 2011). A fair comparison (Turing test) requires that the space-time scales of the ob-  
184 servations and the model results are the same, which means the power spectra should match.  
185 This comparison is a necessary test to realize Lorenz’ vision quoted in the Introduction. It  
186 is not a sufficient test, however, as the OGCM results can resemble the measurements for the  
187 wrong reasons, but we take it as strong evidence of small OGCM bias.

188 Second, the OGCM solutions make accurate, testable predictions about the real ocean. His-  
189 torically, advances from theoretical and numerical research in dynamical oceanography have  
190 lagged advances from observational research (see footnote 6). Once OGCMs become under-  
191 determined by data, it will be common for them to make predictions that can be tested by field

---

<sup>9</sup>It is possible to argue that any inverse problem with real observations is formally under-determined because the observational error can be considered as an unknown parameter to be solved for (Wunsch 1996; Stammer et al. 2002). Regardless, no global ocean circulation state estimate has characterized the null space associated with the indeterminacy (to our knowledge), or presented different solutions that fit the observations equally well. Instead, the practice has been to stop the state estimation once an acceptable fit has been achieved (Stammer et al. 2002; Nguyen et al. 2020).

programs. For example, DSO simulations show exchange of dense water out of the overflow onto the east Greenland continental shelf, and vice versa (Magaldi et al. 2011). They also show entrainment of near-surface waters south of Iceland into the DSO within a few months, at least during hard winters (Saber et al. 2020). It remains to be seen if these predictions occur in the real ocean.

## 5. Limits to Computational Oceanography

Although these opportunities are exciting, there are clear limits to Computational Oceanography. First, direct numerical simulation (DNS) of the global ocean circulation is inconceivable today. DNS in this context means running OGCMs that resolve all scales of motion; from the planetary scale to the dissipation scale (around 1mm), and from centuries to seconds. DNS would avoid the challenge of parametrizing the effects of the unresolved scales, but at vast computational cost. Fig. 3 shows why. It shows the full range of space and time scales relevant to the ocean general circulation, about ten orders of magnitude in both. It also shows the space time volumes accessible to present-day supercomputers, including the best AR6 OGCMs shown in Fig. 2, the Poseidon Project run,<sup>10</sup> and turbulence simulations (DNS and large eddy simulations, LES). To span the entire space time plane, supercomputers would need to resolve about  $10^{25}$  grid points and  $10^{10}$  time steps. That is about 16 orders of magnitude more grid points than is possible today. Extrapolating the doubling time of 2.2 years in Fig. 2c, it would take 120 years to achieve this increase, which is impossible to envision. Clearly, the exponential growth must roll off at some point, and, clearly, OGCM simulations cannot replace observations of the natural ocean.

---

<sup>10</sup>The Poseidon Project intends to run a global OGCM at (nominally) 1 km horizontal resolution ([poseidon.idies.jhu.edu](http://poseidon.idies.jhu.edu)). The Poseidon Project is unrelated to the TOPEX/Poseidon altimeter.

Another potential limit concerns scalability of OGCM codes. Fig. 2 shows that the historic doubling time for the number of OGCM grid points is about twice the doubling time for supercomputer speed. That value is close to the optimal ratio of  $3/2$ , which assumes that machine speedup is spent on increasing horizontal resolution, that model timestep is inversely proportional to the grid spacing (for numerical stability), and that all other factors are equal. In other words, the historic OGCM growth has nearly maintained pace with the supercomputer acceleration. It is unclear how this trend will continue, however, because of the overhead of communication from processor cores to other cores, to memory, and to disk (Le Sommer et al. 2018). Moreover, exascale supercomputers will not resemble petascale supercomputers: they will have different architectures and greater diversity (Giles and Reguly 2014). These changes are driven by physical limits on clock speed and power densities in silicon microprocessors, as well as economic forces. To harness exascale machines OGCM software must radically change (for discussion of this point for atmospheric general circulation models, see Lawrence et al. 2018 and Gropp and Snir 2013). The developers of next generation OGCMs should adopt collaborative, open community habits (Le Sommer et al. 2018). Promising paths are to define and refine modular sub-components, and to develop domain-specific languages, performance tools, and data models that separate different levels in the software stack for optimization by experts (Lawrence et al. 2018). OGCM computational intensity (the fraction of time spent performing floating point calculations versus memory operations) is low: Le Sommer et al. (2018) estimate OGCMs run at 5% peak speed, for example. So there is potential to accelerate OGCMs by reducing this bottleneck (for example by exploiting time parallelism, Schreiber et al. 2017; Hamon et al. 2020). Exploiting new application-specific hardware accelerators and new OGCM solver paradigms, like lower precision (Palmer 2012; Palem 2014), will also be important. These developments will mitigate the saturation of transistor density and the demise of Moore’s Law, and they offer hope to continue the refinement of OGCM meshes.

Finally, there are challenging issues to couple OGCMs to other parts of the Earth system at horizontal resolutions around one kilometer. For example, air/sea interaction, sea ice dynamics, and biogeochemistry are all poorly understood and hard to simulate at these scales.

## 6. Opportunities for Computational Oceanography

The opportunities for Computational Oceanography to advance marine science include:

- Migration from the study of specific instances of phenomena to the study of statistics of these phenomena. The DSO is one of many currents that is affected by rotation, stratification, and bathymetry. It is inconceivable to observe all of them, but they can all be simulated in an exascale OGCM. Empirical characterization of these numerical overflows would be an important step forward.
- Discovery and characterization of intermittent, time-dependent, three-dimensional phenomena, which are hard to observe. Submesoscale currents are in this class, which occur at horizontal scales shorter than several kilometers (Thomas et al. 2008). Diapycnal mixing is another example, which occurs at scales shorter than meters (MacKinnon et al. 2017).
- Comprehensive and illuminating analyses of ocean mass, heat, salt, momentum, energy, and vorticity budgets, in a way that is nearly impossible with direct observations.
- Discovery and characterization of ocean circulation regimes that cannot be observed. Examples include the circulation during the last glacial maximum (paleo-oceanography) or in extra-terrestrial oceans (exo-oceanography). For these ocean circulation problems, the data-sparseness challenge is much worse than for the modern ocean (LeGrand and Wunsch 1995; Amrhein et al. 2018; Way et al. 2017). Criterion 2 was achieved with smaller computational

resources for these fields, and therefore they have already entered the era of Computational Oceanography by the rationale in section 4.

- Robust observing system design using OGCM solutions as synthetic data. These Observing System Simulation Experiments (Errico et al. 2012) should become the best-practice standard for fieldwork design. There are implications for making the OGCM output accessible and easy to work with (see below), but the payoff from engaging observational oceanographers is great.
- Insight from OGCM state estimation to support fieldwork, ideally in real time. The community should recognize the fact that the under-determined state estimates imply an infinite number of OGCM solutions that match the data exactly. This means that techniques are needed to characterize and handle the OGCM null space (indeterminacy). For example, observational oceanographers at sea could make decisions about where, when, and how to observe using OGCM information that captures the range of possible circulation states consistent with data. This practice is common in atmospheric science already.
- More efficient identification of interesting phenomena using automatic methods, like artificial intelligence and data mining (Kutz 2017; Lguensat et al. 2019). In fact, such automatic methods will become essential as the size of OGCM output grows exponentially and overwhelms manual feature identification (see below).
- Increasing transition of dynamical oceanography to an experimental (computational) science. It has long been recognized that idealized models isolate physical mechanisms relevant to the general circulation and thereby build dynamical understanding. We still require idealized models; in particular, we need a hierarchy of models that span the gap between geophysical fluid dynamics problems and realistic simulations of the circulation. This hierarchy will

ensure that the increasing OGCM realism does not outpace understanding of the basic physics (Held 2005; Vallis 2016; Coveney et al. 2016; Emanuel 2020).

## 7. Prospects for Computational Oceanography

How can these priorities be achieved and what are the prospects for Computational Oceanography? We should focus on these issues in the next several years:

- The indeterminacy of OGCM solutions by observations should be recognized—we should “embrace the null space.” Imagine computing an ensemble of high resolution (high degrees of freedom per observation) state estimates that fit the observations (exactly or equally well within instrumental errors). These state estimates would differ, for example, in the characteristics of their eddies, or in their deep circulations, or in their internal wave fields, or in their diapycnal mixing. In such a situation, the different state estimates should all be treated seriously. The ensemble would characterize the null space (indeterminacy) in the inverse problem and therefore quantify the variety of ocean states consistent with observations and ocean circulation physics. This vision for uncertainty quantification echoes the probabilistic practice of ensemble atmospheric model runs to forecast the weather (see also McWilliams 2007; Le Sommer et al. 2018).
- Barriers to dissemination of OGCM simulation output should be lowered—we should “democratize the data.” The output should be freely available, including to non-professional users. Traditionally, effort has focused on the challenges of calculating OGCM solutions with supercomputers. The OGCM output has become increasingly hard to use, because of the massive data volume, and the technical complexities that attend the high-performance computation. Access to high-resolution OGCM output is restricted to a few experts in practice.

The remedy is to build high-performance data science infrastructure to match the high-performance compute infrastructure (Overpeck et al. 2011). These data portals should be open and have low thresholds to getting started. We should be able to sample the simulations the way that we sample the real ocean. For example, it should be easy for an observational oceanographer to plot a synthetic hydrographic section or mooring timeseries. The data portals should include open software and significant compute resources to process and analyze the simulation data. We should avoid the inefficient practice in which users are forced to download voluminous data to their local machines and then write their own code to analyze them. Technologies and infrastructure to achieve these goals are under development, such as the OceanSpy OGCM data analysis package (Almansi et al. 2019), the Pangeo community in geoscience big data ([pangeo.io](http://pangeo.io)), and the SciServer and JASMIN big data science platforms (Medvedev et al. 2016, [www.jasmin.ac.uk](http://www.jasmin.ac.uk)).

- “Benchmark” OGCM reference solutions should be computed using the best available compute resources and served to the public. They are of intrinsic value to all oceanographers, not just ocean modellers, for the reasons stated above. Benchmark solutions for regional ocean circulation problems are valuable for the same reasons, as are idealized simulations of specific ocean dynamical processes. The track record of other fields using this approach is impressive. For instance, the Johns Hopkins Turbulence Database exposes cutting-edge turbulence simulation data to researchers and provides easy-to-use interfaces to retrieve and interact with the data using novel metaphors like immersing virtual sensors into the 4-D data ([turbulence.pha.jhu.edu](http://turbulence.pha.jhu.edu); Perlman et al. 2007; Li et al. 2008).
- OGCMs will migrate to exascale compute resources in the next few years. This migration will involve new paradigms to access the data. For example, with today’s petaflop supercomputers

only about 0.1% of the OGCM solution can be permanently stored for analysis. The problem arises because of the prohibitive time needed to transfer the massive output volume to long-term storage media, and the prohibitive expense of the media. This loss of OGCM data will be much worse on exaflop machines.

To mitigate this problem consider the strategy adopted by the Large Hadron Collider (LHC), the world’s most sophisticated experimental facility. The LHC provides a single source of data on subatomic particle collisions. Several experiments tap the data stream in so-called “beam-lines.” Within each experiment, customized hardware monitors the stream. Only about one event in ten million is retained for storage and detailed analysis. In exascale oceanography the analogous idea (see section 3.3.5 in Asch et al. 2018) is to enable automatic identification of selected circulation events and trigger storage while the OGCM runs. For example, we could target intermittent intense mixing events, plus their antecedents and fates. An implication is that we should build a software interface for community-supplied software plugins to implement the custom triggers. Also, we need to enable posterior re-computation of small space-time chunks of the full solution, with customized diagnostics, and possibly at higher resolutions.

It is instructive to compare Computational Oceanography with computational meteorology, which is the analogous field in atmospheric sciences. Computational meteorology has somewhat different science objectives. Numerical Weather Prediction (NWP) is an important task, for example. The main advances in NWP attributable to growth in computer power are: (i) improved model resolution (now also approaching global 1 km horizontal resolution; Fuhrer et al. 2018), and (ii) improved forecast uncertainty quantification through larger ensembles of forecast runs. Computational meteorology also concerns reanalysis products to hindcast the historical atmospheric state. The

reanalysis state estimation tools tolerate unphysical adjustments (increments), however, which give more accurate fits to observations at lower computational cost. This practice is different to the ocean state estimation tools discussed here, which firmly constrain the model solutions to satisfy the model equations.

Nevertheless, there are several useful lessons from computational meteorology: First, NWP has steadily improved since the 1980s (Bauer et al. 2015). The rate is an improvement in forecast skill of about one day per decade (meaning a 2015 three-day forecast is about as skillful as a 2005 two-day forecast). The improvement derives mainly from better forecast initialization and better atmospheric general circulation models (AGCMs; Magnusson and Källén 2013; see also Simmons and Hollingsworth 2002). In this context, better AGCMs means models that have higher resolution, have more accurate parameterizations and/or complexity, and have larger forecast ensembles that better estimate forecast uncertainty. Computing advances have played an enormous role in these improvements (Bauer et al. 2015). Second, as AGCM resolution increases, new phenomena begin to be resolved. For example, with AGCM grid spacing of a few kilometers convective scales are partly resolved (convective systems) but partly unresolved (convective cells). This partial resolution of convection is called the “gray zone”, akin to eddy-permitting resolution in OGCMs. The best approach to set up convection parameterization schemes in the AGCM gray zone is unclear and forecast skill does not always improve at all lead times as resolution increases (Hong and Dudhia 2012). Moreover, at cloud-resolving resolution, data density is mismatched with AGCM resolution (the number of degrees of freedom exceeds the number of observations) and the model solution is not well constrained (Hong and Dudhia 2012).

## 8. Conclusion

Global OGCMs have a rich history that stretches back to the 1970s and regional OGCMs stretch back to the 1960s (models of the tides stretch back even further; see Cartwright 2012). OGCMs have been valuable to elucidate the ocean circulation since their inception. More broadly, numerical solution of rotating, stratified flow has roots in numerical weather prediction (NWP) from the early twentieth century (Abbe 1901; Bjerknes 1904; Richardson 1922 see also Lynch 2008 and Benjamin et al. 2019 for historical perspectives on NWP and climate models). Since 2000, global OGCMs have continued their exponential improvement in resolution. They are now becoming unconstrained by observations. Benchmark OGCM solutions have increasing value to a growing community and should be permanently archived and freely available. Clear limits, opportunities, and prospects for Computational Oceanography are in sight. For these reasons, our answer to the question posed in the title of this essay is yes: Computational Oceanography is entering a new era and is coming of age.

This field promises powerful new tools to address previously intractable problems. It does not aim to supplant observational oceanography. Indeed, observing the natural ocean must never cease. Instead, the greatest opportunity lies in merging these hitherto disparate branches of marine science. Lasting progress will require that we trust computational insights, verify them with real world observations, and understand them with fundamental theory.

*Acknowledgments.* This material is based upon work supported by the National Science Foundation under Grant Number OAC-1835640 and by the Institute for Data Intensive Engineering and Science at Johns Hopkins University. Comments by Baylor Fox-Kemper and two other reviewers refined the paper.

391 *Data availability statement.* Codes to make the figures are available at  
392 [github.com/hainegroup/Computational-Oceanography-Commentary](https://github.com/hainegroup/Computational-Oceanography-Commentary). For Fig. 2,  
393 the temperature data are from the National Centers for Environmental Information World  
394 Ocean Database, the altimeter mission data are from [www.altimetry.info](http://www.altimetry.info), the IPCC data  
395 are from the IPCC reports and [pcmdi.llnl.gov/CMIP6](http://pcmdi.llnl.gov/CMIP6), and the supercomputer data are from  
396 [en.wikipedia.org/wiki/List\\_of\\_fastest\\_computers](https://en.wikipedia.org/wiki/List_of_fastest_computers). The data for AR6 are from the  
397 HighResMIP project in July 2020, which was incomplete then. The ECMWF and NCAR machine  
398 speed data are from [www.top500.org](http://www.top500.org).

## 399 **References**

- 400 Abbe, C., 1901: The physical basis of long range weather forecasts. *Mon. Weather. Rev.*, **29** (12),  
401 551–561, doi:10.1175/1520-0493(1901)29[551c:tpbolw]2.0.co;2.
- 402 Almansi, M., R. Gelderloos, T. W. N. Haine, A. Saberi, and A. H. Siddiqui, 2019: OceanSpy:  
403 A Python package to facilitate ocean model data analysis and visualization. *Journal of Open*  
404 *Source Software*, **4** (39), 1506, doi:10.21105/joss.01506.
- 405 Almansi, M., T. W. N. Haine, R. Gelderloos, and R. S. Pickart, 2020: Evolution of Denmark  
406 Strait Overflow cyclones and their relationship to overflow surges. *Geophys. Res. Lett.*, doi:  
407 10.1029/2019GL086759.
- 408 Amrhein, D. E., C. Wunsch, O. Marchal, and G. Forget, 2018: A global glacial ocean state  
409 estimate constrained by upper-ocean temperature proxies. *J. Climate*, **31** (19), 8059–8079, doi:  
410 10.1175/jcli-d-17-0769.1.
- 411 Argo, 2020: Argo float data and metadata from global data assembly centre (Argo GDAC).  
412 SEANOE, doi:10.17882/42182.

413 Asch, M., and Coauthors, 2018: Big data and extreme-scale computing. *The International*  
 414 *Journal of High Performance Computing Applications*, **32** (4), 435–479, doi:10.1177/  
 415 1094342018778123.

416 Bauer, P., A. Thorpe, and G. Brunet, 2015: The quiet revolution of numerical weather prediction.  
 417 *Nature*, **525** (7567), 47–55, doi:10.1038/nature14956.

418 Benjamin, S. G., J. M. Brown, G. Brunet, P. Lynch, K. Saito, and T. W. Schlatter, 2019: 100 years  
 419 of progress in forecasting and NWP applications. *Meteorological Monographs*, **59**, 131–1367,  
 420 doi:10.1175/amsmonographs-d-18-0020.1.

421 Bennett, A. F., 1992: *Inverse methods in physical oceanography*. Cambridge University  
 422 Press, Cambridge, United Kingdom and New York, NY, USA, 346 pp., doi:10.1017/  
 423 cbo9780511600807.

424 Bjerknes, V., 1904: Das problem der wettervorhersage, betrachtet vom standpunkte der mechanik  
 425 und der physik (the problem of weather prediction, considered from the viewpoints of mechanics  
 426 and physics). *Meteorologische Zeitschrift*, **21**, 1–7, doi:10.1127/0941-2948/2009/416, translated  
 427 and edited by E. Volken and S. Brönnimann “. – Meteorol. Z. 18 (2009), 663–667.

428 Bryan, K., 2006: Modeling ocean circulation. *Physical Oceanography: Developments since 1950*,  
 429 Springer New York, 29–44, doi:10.1007/0-387-33152-2\_3.

430 Bryan, K., and L. J. Lewis, 1979: A watermass model of the world ocean. *J. Geophys. Res.*, **84**,  
 431 2503–2517, doi:10.1029/jc084ic05p02503.

432 Cartwright, D. E., 2012: *Tides*. Cambridge University Press, Cambridge, United King-  
 433 dom and New York, NY, USA, URL [https://www.ebook.de/de/product/2966034/david\\_edgar\\_](https://www.ebook.de/de/product/2966034/david_edgar_cartwright_tides.html)  
 434 [cartwright\\_tides.html](https://www.ebook.de/de/product/2966034/david_edgar_cartwright_tides.html).

435 Coveney, P. V., E. R. Dougherty, and R. R. Highfield, 2016: Big data need big theory too.  
 436 *Phil. Trans. R. Soc. Lond. A*, **374** (2080), 20160153, doi:10.1098/rsta.2016.0153.

437 Dwivedi, S., T. W. N. Haine, and C. E. Del Castillo, 2011: Upper ocean state estimation in  
 438 the Southern Ocean Gas Exchange Experiment region using the four-dimensional variational  
 439 technique. *J. Geophys. Res.*, **116**, C00F02, doi:10.1029/2009JC005615.

440 Emanuel, K., 2020: The relevance of theory for contemporary research in atmospheres, oceans,  
 441 and climate. *AGU Advances*, **1** (2), doi:10.1029/2019av000129.

442 Errico, R. M., R. Yang, N. C. Privé, K.-S. Tai, R. Todling, M. E. Sienkiewicz, and J. Guo,  
 443 2012: Development and validation of observing-system simulation experiments at NASA's  
 444 global modeling and assimilation office. *Q. J. R. Meteorol. Soc.*, **139** (674), 1162–1178, doi:  
 445 10.1002/qj.2027.

446 Feigenbaum, E. A., 2003: Some challenges and grand challenges for computational intelligence.  
 447 *Journal of the ACM*, **50** (1), 32–40, doi:10.1145/602382.602400.

448 Fox-Kemper, B., and Coauthors, 2019: Challenges and prospects in ocean circulation models.  
 449 *Frontiers in Marine Science*, **6**, doi:10.3389/fmars.2019.00065.

450 Fuhrer, O., and Coauthors, 2018: Near-global climate simulation at 1 km resolution: establishing a  
 451 performance baseline on 4888–GPUs with COSMO 5.0. *Geosci. Mod. Dev.*, **11** (4), 1665–1681,  
 452 doi:10.5194/gmd-11-1665-2018.

453 GCOS, 2016: The global observing system for climate. Tech. Rep. GCOS No. 200, World  
 454 Meteorological Organization. [https://library.wmo.int/index.php?lvl=notice\\_display&id=19838#](https://library.wmo.int/index.php?lvl=notice_display&id=19838#.XxWxPJNKhBw)  
 455 .XxWxPJNKhBw.

- 456 Gent, P. R., 2011: The Gent-McWilliams parameterization: 20/20 hindsight. *Ocean Modelling*,  
457 **39**, 2–9, doi:10.1016/j.ocemod.2010.08.002.
- 458 Giles, M. B., and I. Regulý, 2014: Trends in high-performance computing for engineering calcu-  
459 lations. *Phil. Trans. R. Soc. Lond. A*, **372** (2022), 20130 319, doi:10.1098/rsta.2013.0319.
- 460 Griffies, S. M., 2004: *Fundamentals of Ocean Climate models*. Princeton University Press, Prince-  
461 ton , NJ, USA and Oxford, United Kingdom, 518 pp., doi:10.2307/j.ctv301gzg.
- 462 Griffies, S. M., and Coauthors, 2000: Developments in ocean climate modelling. *Ocean Modelling*,  
463 **2** (3-4), 123–192, doi:10.1016/s1463-5003(00)00014-7.
- 464 Gropp, W., and M. Snir, 2013: Programming for exascale computers. *Computing in Science &*  
465 *Engineering*, **15** (6), 27–35, doi:10.1109/mcse.2013.96.
- 466 Haine, T. W. N., 2010: High-frequency fluctuations in Denmark Strait Overflow transport. *Geo-*  
467 *phys. Res. Lett.*, **37**, L14 601, doi:10.1029/2010GL043272.
- 468 Hamon, F. P., M. Schreiber, and M. L. Minion, 2020: Parallel-in-time multi-level integration  
469 of the shallow-water equations on the rotating sphere. *J. Comput. Phys.*, **407**, 109 210, doi:  
470 10.1016/j.jcp.2019.109210.
- 471 Harel, D., 2005: A Turing-like test for biological modeling. *Nature Biotechnology*, **23** (4), 495–496,  
472 doi:10.1038/nbt0405-495.
- 473 Held, I. M., 2005: The gap between simulation and understanding in climate modeling.  
474 *Bull. Amer. Meteor. Soc.*, **86** (11), 1609–1614, doi:10.1175/bams-86-11-1609.
- 475 Holland, W. R., and J. C. McWilliams, 1987: Computer modeling in physical oceanography from  
476 the global circulation to turbulence. *Physics Today*, **40** (10), 51–57, doi:10.1063/1.881115.

- 477 Hong, S.-Y., and J. Dudhia, 2012: Next-generation numerical weather prediction: Bridging pa-  
478 rameterization, explicit clouds, and large eddies. *Bull. Amer. Meteor. Soc.*, **93** (1), ES6–ES9,  
479 doi:10.1175/2011bams3224.1.
- 480 Kalnay, E., 2002: *Atmospheric Modeling, Data Assimilation and Predictability*. Cambridge  
481 University Press, Cambridge, United Kingdom and New York, NY, USA, doi:10.1017/  
482 cbo9780511802270.
- 483 Klinger, B. A., and T. W. N. Haine, 2019: *Ocean Circulation in Three Dimensions*. 1st ed.,  
484 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, URL  
485 <http://www.cambridge.org/9780521768436>.
- 486 Kutz, J. N., 2017: Deep learning in fluid dynamics. *J. Fluid Mech.*, **814**, 1–4, doi:10.1017/jfm.  
487 2016.803.
- 488 Lawrence, B. N., and Coauthors, 2018: Crossing the chasm: how to develop weather and climate  
489 models for next generation computers? *Geosci. Mod. Dev.*, **11** (5), 1799–1821, doi:10.5194/  
490 gmd-11-1799-2018.
- 491 Le Sommer, J., E. P. Chassignet, and A. J. Wallcraft, 2018: Ocean circulation modeling for  
492 operational oceanography: Current status and future challenges. *New Frontiers in Operational*  
493 *Oceanography*, GODAE OceanView, doi:10.17125/gov2018.ch12.
- 494 Lea, D. J., T. W. N. Haine, and R. F. Gasparovic, 2006: Observability of the Irminger Sea  
495 circulation using variational data assimilation. *Q. J. R. Meteorol. Soc.*, **132** (618), 1545–1576,  
496 doi:10.1256/qj.05.77.

- 497 LeGrand, P., and C. Wunsch, 1995: Constraints from paleotracer data on the North Atlantic  
498 circulation during the Last Glacial Maximum. *Paleoceanography*, **10** (6), 1011–1045, doi:  
499 10.1029/95pa01455.
- 500 Lguensat, R., J. Le Sommer, S. Metref, E. Cosme, and R. Fablet, 2019: Learning generalized  
501 quasi-geostrophic models using deep neural numerical models, 1911.08856v1.
- 502 Li, Y., and Coauthors, 2008: A public turbulence database cluster and applications to study  
503 lagrangian evolution of velocity increments in turbulence. *Journal of Turbulence*, **9**, N31, doi:  
504 10.1080/14685240802376389.
- 505 Lorenz, E. N., 1967: *The Nature and Theory of the General Circulation of the Atmosphere*. WMO:  
506 TP, World Meteorological Organization.
- 507 Lynch, P., 2008: The origins of computer weather prediction and climate modeling. *J. Com-*  
508 *put. Phys.*, **227** (7), 3431–3444, doi:10.1016/j.jcp.2007.02.034.
- 509 MacKinnon, J. A., and Coauthors, 2017: Climate Process Team on internal-wave driven ocean  
510 mixing. *Bull. Amer. Meteor. Soc.*, **98** (11), 2429–2454, doi:10.1175/bams-d-16-0030.1.
- 511 Magaldi, M. G., and T. W. N. Haine, 2015: Hydrostatic and non-hydrostatic simulations of dense  
512 waters cascading off a shelf: the East Greenland case. *Deep Sea Res., Part I*, **96**, 89–104,  
513 doi:10.1016/j.dsr.2014.10.008.
- 514 Magaldi, M. G., T. W. N. Haine, and R. S. Pickart, 2011: On the nature and variability of the  
515 East Greenland Spill Jet: A case study in summer 2003. *J. Phys. Oceanogr.*, **41**, 2307–2327,  
516 doi:10.1175/JPO-D-10-05004.1.
- 517 Magnusson, L., and E. Källén, 2013: Factors influencing skill improvements in the ECMWF  
518 forecasting system. *Mon. Weather. Rev.*, **141** (9), 3142–3153, doi:10.1175/mwr-d-12-00318.1.

- 519 Maltrud, M. E., and J. L. McClean, 2005: An eddy resolving global  $1/10^\circ$  ocean simulation. *Ocean*  
520 *Modelling*, **8 (1-2)**, 31–54, doi:10.1016/j.ocemod.2003.12.001.
- 521 Maltrud, M. E., R. D. Smith, A. J. Semtner, and R. C. Malone, 1998: Global eddy-resolving ocean  
522 simulations driven by 1985-1995 atmospheric winds. *J. Geophys. Res.*, **103 (C13)**, 30 825–  
523 30 853, doi:10.1029/1998jc900013.
- 524 McWilliams, J. C., 1996: Modeling the oceanic general circulation. *Ann. Rev. Fluid Mech.*, **28 (1)**,  
525 215–248, doi:10.1146/annurev.fl.28.010196.001243.
- 526 McWilliams, J. C., 2007: Irreducible imprecision in atmospheric and oceanic simulations.  
527 *Proc. Nat. Acad. Sci.*, **104 (21)**, 8709–8713, doi:10.1073/pnas.0702971104.
- 528 Medvedev, D., G. Lemson, and M. Rippin, 2016: SciServer compute. *Proceedings of the 28th*  
529 *International Conference on Scientific and Statistical Database Management - SSDBM '16*,  
530 ACM Press, doi:10.1145/2949689.2949700.
- 531 Moore, G. E., 1975: Progress in digital integrated electronics IEDM Tech Digest. IEEE.
- 532 Morrow, R., and Coauthors, 2019: Global observations of fine-scale ocean surface topography  
533 with the surface water and ocean topography (SWOT) mission. *Frontiers in Marine Science*, **6**,  
534 doi:10.3389/fmars.2019.00232.
- 535 Nguyen, A. T., H. Pillar, V. Ocaña, A. Bigdeli, T. A. Smith, and P. Heimbach, 2020: The  
536 arctic subpolar gyre sTate estimate (ASTE): Description and assessment of a data-constrained,  
537 dynamically consistent ocean-sea ice estimate for 2002-2017. doi:10.1002/essoar.10504669.1.
- 538 Overpeck, J. T., G. A. Meehl, S. Bony, and D. R. Easterling, 2011: Climate data chal-  
539 lenges in the 21st century. *Science*, **331 (6018)**, 700–702, doi:10.1126/science.1197869, URL

<https://science.sciencemag.org/content/331/6018/700>, <https://science.sciencemag.org/content/331/6018/700.full.pdf>.

Palem, K. V., 2014: Inexactness and a future of computing. *Phil. Trans. R. Soc. Lond. A*, **372** (2018), 20130281, doi:10.1098/rsta.2013.0281.

Palmer, T. N., 2012: Towards the probabilistic earth-system simulator: a vision for the future of climate and weather prediction. *Q. J. R. Meteorol. Soc.*, **138** (665), 841–861, doi:10.1002/qj.1923.

Perlman, E., R. Burns, Y. Li, and C. Meneveau, 2007: Data exploration of turbulence simulations using a database cluster. *Proceedings of the 2007 ACM/IEEE conference on Supercomputing - SC '07*, ACM Press, doi:10.1145/1362622.1362654.

Richardson, L. F., 1922: *Weather Prediction by Numerical Process*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, doi:10.1017/cbo9780511618291, reprinted 2007.

Rocha, C. B., T. K. Chereskin, S. T. Gille, and D. Menemenlis, 2016: Mesoscale to submesoscale wavenumber spectra in Drake Passage. *J. Phys. Oceanogr.*, **46** (2), 601–620, doi:10.1175/jpo-d-15-0087.1.

Saberi, A., T. W. N. Haine, R. Gelderloos, M. F. de Jong, H. Fury, and A. Bower, 2020: Lagrangian perspective on the origins of Denmark Strait Overflow. *J. Phys. Oceanogr.*, **50** (8), 2393–2414, doi:10.1175/JPO-D-19-0210.1.

Schreiber, M., P. S. Peixoto, T. Haut, and B. Wingate, 2017: Beyond spatial scalability limitations with a massively parallel method for linear oscillatory problems. *The International Journal of High Performance Computing Applications*, **32** (6), 913–933, doi:10.1177/1094342016687625.

562 Semtner, A. J., and R. M. Chervin, 1992: Ocean general circulation from a global eddy-resolving  
563 model. *J. Geophys. Res.*, **97 (C4)**, 5493, doi:10.1029/92jc00095.

564 Simmons, A. J., and A. Hollingsworth, 2002: Some aspects of the improvement in skill of  
565 numerical weather prediction. *Q. J. R. Meteorol. Soc.*, **128 (580)**, 647–677, doi:10.1256/  
566 003590002321042135.

567 Stammer, D., and Coauthors, 2002: Global ocean circulation during 1992–1997, estimated from  
568 ocean observations and a general circulation model. *J. Geophys. Res.*, **107 (C9)**, doi:10.1029/  
569 2001jc000888.

570 Stewart, R. H., 2008: *Introduction to physical oceanography*. September 2008 edition ed., pub-  
571 lished online by the author at: [http://oceanworld.tamu.edu/home/course\\_book.htm](http://oceanworld.tamu.edu/home/course_book.htm).

572 Thomas, L. N., A. Tandon, and A. Mahadevan, 2008: Submesoscale processes and dynamics.  
573 *Ocean Modeling in an Eddying Regime*, American Geophysical Union, 17–38, doi:10.1029/  
574 177gm04.

575 Turing, A. M., 1950: I. Computing machinery and intelligence. *Mind*, **LIX (236)**, 433–460,  
576 doi:10.1093/mind/lix.236.433.

577 Vallis, G. K., 2016: Geophysical fluid dynamics: whence, whither and why? *Proc. R. Soc. Lond. A*,  
578 **472 (2192)**, 20160 140, doi:10.1098/rspa.2016.0140.

579 Waldrop, M. M., 2016: The chips are down for Moore’s law. *Nature*, **530 (7589)**, 144–147,  
580 doi:10.1038/530144a.

581 Way, M. J., and Coauthors, 2017: Resolving orbital and climate keys of earth and extraterrestrial  
582 environments with dynamics (ROCKE-3d) 1.0: A general circulation model for simulating

the climates of rocky planets. *The Astrophysical Journal Supplement Series*, **231** (1), 12, doi:  
10.3847/1538-4365/aa7a06.

Wunsch, C., 1996: *The ocean circulation inverse problem*. 1st ed., Cambridge University Press,  
Cambridge, United Kingdom and New York, NY, USA, doi:10.1017/cbo9780511629570, 442  
pp.

Wunsch, C., 2006: *Discrete Inverse and State Estimation Problems*. 1st ed., Cambridge University  
Press, Cambridge, United Kingdom and New York, NY, USA, doi:10.1017/cbo9780511535949,  
371 pp.

## LIST OF FIGURES

- Fig. 1.** OGCM Turing tests. In each of (a)–(c) oceanographic field measurements are compared with OGCM results, but they are unlabeled (and processed similarly). The Turing test is to identify which is which. (a) Denmark Strait Overflow (DSO) volume flux ( $S_v$ ,  $1S_v = 10^6 \text{m}^3 \text{s}^{-1}$ , negative means equatorwards). Adapted from Haine (2010). (b) Salinity (colors) on a section north of Denmark Strait (annual average; the heavy contour is the  $27.80 \sigma_0$  density anomaly). (c) Lagrangian trajectories of RAFOS floats and synthetic RAFOS floats. Adapted from Saberi et al. (2020). . . . . 31
- Fig. 2.** Unequal exponential growth. (a) History of deep (deeper than 1000 m) ocean temperature measurements. The colored dots show different instruments and observing platforms. (b) History of sea level measurements from satellite altimetry expressed by the cumulative number of days of measurement. The satellite missions and their durations are indicated with the colored bars. (c) History of cutting-edge global OGCM and IPCC ocean model resolution expressed by the lengthscale of the horizontal grid and the number of model grid points. Each dot represents one ocean model and the OGCMs are from Bryan and Lewis (1979); Semtner and Chervin (1992); Maltrud et al. (1998); Maltrud and McClean (2005), and Rocha et al. (2016). (d) History of top supercomputers using Rmax speed (FLOPS = floating point operations per second) for fastest machines (open circles) and ECMWF and NCAR machines (closed circles). The lines show best fit exponential growth in each panel ( $\tau_{2\times}$  is the doubling time). . . . . 32
- Fig. 3.** Characteristic space and time scales of the ocean general circulation. Various geophysical and theoretical scales are shown with gray patches and colored lines (for a discussion, see Klinger and Haine 2019). The colored rectangles show cutting-edge circulation models (direct numerical simulation of turbulence, large eddy simulation, the Poseidon Project run, AR6 HighResMIP, and TAR OGCMs). The black dot shows the sampling characteristics of the Argo profiling floats, and the TOPEX/Poseidon-Jason altimeters. The diagram is indicative, not definitive, because it suppresses the anisotropies and inhomogeneities present in the general circulation. . . . . 33




FIG. 1. OGCM Turing tests. In each of (a)–(c) oceanographic field measurements are compared with OGCM results, but they are unlabeled (and processed similarly). The Turing test is to identify which is which. (a) Denmark Strait Overflow (DSO) volume flux (Sv,  $1\text{ Sv} = 10^6\text{ m}^3\text{ s}^{-1}$ , negative means equatorwards). Adapted from Haine (2010). (b) Salinity (colors) on a section north of Denmark Strait (annual average; the heavy contour is the  $27.80\sigma_0$  density anomaly). (c) Lagrangian trajectories of RAFOS floats and synthetic RAFOS floats. Adapted from Saberi et al. (2020).




FIG. 2. Unequal exponential growth. (a) History of deep (deeper than 1000 m) ocean temperature measurements. The colored dots show different instruments and observing platforms. (b) History of sea level measurements from satellite altimetry expressed by the cumulative number of days of measurement. The satellite missions and their durations are indicated with the colored bars. (c) History of cutting-edge global OGCM and IPCC ocean model resolution expressed by the lengthscale of the horizontal grid and the number of model grid points. Each dot represents one ocean model and the OGCMs are from Bryan and Lewis (1979); Semtner and Chervin (1992); Maltrud et al. (1998); Maltrud and McClean (2005), and Rocha et al. (2016). (d) History of top supercomputers using Rmax speed (FLOPS = floating point operations per second) for fastest machines (open circles) and ECMWF and NCAR machines (closed circles). The lines show best fit exponential growth in each panel ( $\tau_{2x}$  is the doubling time).




FIG. 3. Characteristic space and time scales of the ocean general circulation. Various geophysical and theoretical scales are shown with gray patches and colored lines (for a discussion, see Klinger and Haine 2019). The colored rectangles show cutting-edge circulation models (direct numerical simulation of turbulence, large eddy simulation, the Poseidon Project run, AR6 HighResMIP, and TAR OGCMs). The black dot shows the sampling characteristics of the Argo profiling floats, and the TOPEX/Poseidon-Jason altimeters. The diagram is indicative, not definitive, because it suppresses the anisotropies and inhomogeneities present in the general circulation.