

EXACT SEMIDEFINITE PROGRAMMING BOUNDS FOR PACKING PROBLEMS*

MARIA DOSTERT[†], DAVID DE LAAT[‡], AND PHILIPPE MOUSTROU[§]

Abstract. In this paper we give an algorithm to round the floating point output of a semidefinite programming solver to a solution over the rationals or a quadratic extension of the rationals. This algorithm does not require the solution to be strictly feasible and works for large problems. We apply this to get sharp bounds for packing problems, and we use these sharp bounds to prove that certain optimal packing configurations are unique up to rotations. In particular, we show that the configuration coming from the E_8 root lattice is the unique optimal code with minimal angular distance $\pi/3$ on the hemisphere in \mathbb{R}^8 , and we prove that the three-point bound for the $(3, 8, \vartheta)$ -spherical code, where ϑ is such that $\cos \vartheta = (2\sqrt{2} - 1)/7$, is sharp by rounding to $\mathbb{Q}[\sqrt{2}]$. We also use our machinery to compute sharp upper bounds on the number of spheres that can be packed into a larger sphere.

Key words. semidefinite programming, hybrid numeric-symbolic algorithm, spherical codes, packing problems

AMS subject classifications. 52C17, 90C22

DOI. 10.1137/20M1351692

1. Introduction. In this paper we consider the problem of extracting an exact solution from the output of a numerical semidefinite programming solver. A particularly fitting application domain for this is extremal geometry, including packing problems where we ask for the size of a largest independent set in a graph whose set of vertices is a compact space. A typical example is the ϑ -spherical code problem, where the vertex set is the unit sphere, and two distinct vertices x and y are adjacent when the inner product between x and y is at most $\cos \vartheta$. The strongest known upper bounds often use semidefinite programming [2, 14, 16], which is an extension of linear programming where one optimizes over positive semidefinite matrices satisfying linear constraints. Since semidefinite programs are solved in floating point arithmetic, turning the numerical bounds into rigorous upper bounds requires additional work.

To prove that a packing configuration of size N is optimal, we just need any upper bound in the interval $[N, N + 1)$. In this case, extracting a rigorous bound is easy, because it is enough to prove the existence of an exact solution of the semidefinite program whose objective is close to the objective of the floating point solution, which means we simply need to round a strictly feasible solution. However, if we want to prove uniqueness of such a configuration (see section 2.3), or if we aim at solving

*Received by the editors July 9, 2020; accepted for publication (in revised form) February 16, 2021; published electronically May 25, 2021.

<https://doi.org/10.1137/20M1351692>

Funding: This material is based upon work supported by the National Science Foundation under grant DMS-1439786 while the authors were in residence at the Institute for Computational and Experimental Research in Mathematics in Providence, RI, during the “Point Configurations in Geometry, Physics and Computer Science” semester program. The third author received support from the Tromsø Research Foundation grant 17matteCR.

[†]Department of Mathematics, Royal Institute of Technology (KTH), Stockholm, Sweden (maria.dostert@epfl.ch, <https://sites.google.com/view/mariadostert>).

[‡]Department of Applied Mathematics, Delft University of Technology, Delft, The Netherlands (mail@daviddelaat.nl, www.daviddelaat.nl).

[§]Department of Mathematics and Statistics, UiT - The Arctic University of Norway, Tromsø, Norway (philippe.moustrou@uit.no, <https://pmoustro.perso.math.cnrs.fr/>).

more general problems in extremal geometry where the optimal value is not an integer (for instance, energy minimization problems [9, 12]), then we need an exact bound. Extremal geometry thus provides large semidefinite programming problems where we look for an exact optimal solution that may lie on the boundary of the positive semidefinite matrix cone.

For the problem of finding exact sum of squares decompositions of polynomials, there exist several hybrid numeric-symbolic algorithms [37, 8, 23, 24, 31], and some of them [23, 24] can be applied for special instances of polynomials that are not strictly feasible. For general semidefinite programs, as required in our context, there exist symbolic algorithms [20, 33, 21], but they cannot be used for the size of problems we consider. Other approaches rely on facial reduction: While [27] provides an exact method for sum of squares decompositions, more general techniques such as [36] remain purely numerical. We propose a hybrid numeric-symbolic algorithm, which can be related to facial reduction, and is able to tackle large semidefinite problems arising in extremal geometry, where the optimal solutions are not necessarily strictly feasible.

Since we consider problems where we have a candidate optimal configuration for which we have a numerically sharp semidefinite programming bound, we can include the objective value as a linear constraint, and we are only looking for a solution of the corresponding feasibility problem. The semidefinite programming solver returns a near feasible solution of this semidefinite program, given as a list of matrices whose entries are floating point numbers. For this solution the linear constraints are almost satisfied, and the eigenvalues of the matrices are positive or close to zero (but not necessarily nonnegative). If we expect to find a feasible solution with entries in a given field, the challenge consists in turning the floating point values into elements of this field in such a way that the linear constraints are satisfied and the matrices are positive semidefinite.

The solver gives an approximate solution in the relative interior of the feasible set. This means that if the feasible set has the same dimension as the affine space defined by the linear constraints, the near zero eigenvalues will become exactly zero after projecting the approximate solution into the affine space. Since any positive eigenvalue bounded away from zero will remain positive after a small perturbation of the matrix entries, we thus find an exact feasible solution simply by projecting the approximate solution into the affine space. In prior works, the approach has been to include additional linear constraints coming from the complementary slackness conditions of an optimal configuration in the hope that the above condition becomes valid [9, 3]. However, in general, there is no guarantee at all that this works, and we were not able to prove Theorem 4.2 with this approach, which was the original motivation for this project.

In this paper we develop a general procedure to extract an exact optimal solution from the numerical optimal solution of a semidefinite program. Our main idea is the following: We want to understand the eigenvectors corresponding to near-zero eigenvalues, in order to force these eigenvalues to become exactly zero after rounding. This task seems to be challenging, since the kernels can be big, and the solver does not take into account their structure: Even if these linear spaces afford a basis over the algebraic field we want to round to, computing a basis in floating point arithmetic will only provide messy approximations of linear *real* combinations of such vectors. In order to extract a suitable basis from these numerical approximations, we use the Lenstra–Lenstra–Lovász (LLL) algorithm to detect equations that have to be satisfied by the kernel vectors. This provides all additional linear constraints that will ensure the semidefiniteness of the matrices after rounding. In general, even if the constraints

and the objective are rational, an optimal solution of a semidefinite program might require high algebraic degree [34]. However, for every problem we considered, the semidefinite program and the conditions implied by an optimal configuration were defined over the same field (which was either the field of rational numbers or a quadratic field), and we were able to find an exact optimal solution over this field.

A fundamental approach to compute upper bounds on the size of spherical ϑ -codes is the linear programming bound of Delsarte, Goethals, and Seidel [15]. Even if the semidefinite programming *three-point bound* by Bachoc and Vallentin [2] provides stronger bounds, there are very few known cases where the semidefinite programming bound is sharp while the linear programming bound is not. One such example is given in [3], where Bachoc and Vallentin prove optimality and uniqueness of the Petersen code by using adhoc techniques to show that their bound is sharp. Our rounding procedure directly turns a numerical solution of the semidefinite program into an exact rational solution, and we recover the uniqueness of the Petersen code; see section 4.2.

In [40], Schütte and van der Waerden prove optimality of the ϑ -code in S^2 with cardinality 8 and $\cos \vartheta = (2\sqrt{2} - 1)/7$, and Danzer [11] proves uniqueness up to rotations. The proofs of these results are purely geometric and quite technical. Still, in [3] Bachoc and Vallentin mention that their numerical computations suggest that the semidefinite programming bound is tight. However, they do not provide an exact optimal solution, and due to the value of $\cos \vartheta$ it seems that there is no optimal rational solution. With the adaptation of our approach to quadratic fields, we obtain an optimal solution over $\mathbb{Q}[\sqrt{2}]$. Based on our optimal solution, we give a simplified uniqueness proof in section 4.2.

An even more challenging problem is to determine the optimal size of a spherical code in a spherical cap, where the spherical cap with center $e \in S^{n-1}$ and angle ϕ is defined by

$$\text{Cap}^{n-1}(e, \phi) = \{x \in S^{n-1} : e \cdot x \geq \cos(\phi)\}.$$

In this situation, the linear programming bound cannot be applied, but in [4] Bachoc and Vallentin adapt their three-point bound, which becomes a two-point bound in this context, to get semidefinite programming upper bounds. For the hemisphere $\text{Cap}^{n-1}(e, \pi/2)$, they got a numerically sharp bound for $n = 8$, which is closely related to the famous E_8 lattice: The 240 minimal vectors of E_8 give the unique optimal spherical $\pi/3$ -code in dimension 8 [22, 41, 5]. If e is any of these minimal vectors, the intersection of this configuration with $\text{Cap}^7(e, \pi/2)$ is a $\pi/3$ -code of cardinality 183. Bachoc and Vallentin get a numerical bound very close to 183, which proves the optimality of this configuration. Moreover they conjecture that this is the only optimal configuration up to isometry. Here by using our machinery, we provide an exact optimal rational solution and prove this conjecture; see section 4.1.

To find more sharp bounds, we also apply our techniques to the similar problem of packing unit spheres in a larger sphere, which has connections to material science, radio-surgical treatments, and communication theory; see [32]. We use our machinery of rounding to rationals and quadratic fields to find several exact sharp bounds, and we also use this to give families of sharp bounds for all dimensions; see section 4.3.

2. Semidefinite programming bounds for packing problems.

2.1. Characterizations of invariant kernels. In section 2.2 we give a derivation of the semidefinite programming bounds we use in this paper. For this we need to characterize certain invariant positive definite kernels.

We start with the well-known case of $O(n)$ -invariant kernels on the sphere. Let K be a positive definite kernel on S^{n-1} , by which we mean it is a continuous function from $S^{n-1} \times S^{n-1}$ to \mathbb{R} for which the matrices

$$(K(x_i, x_j))_{i,j=1}^m$$

are positive semidefinite for all $m \geq 0$ and $x_1, \dots, x_m \in S^{n-1}$. Such a kernel is said to be $O(n)$ -invariant if $K(\gamma x, \gamma y) = K(x, y)$ for all $\gamma \in O(n)$ and $x, y \in S^{n-1}$. For each $k \geq 0$, let P_k^n be the degree k ultraspherical polynomial for S^{n-1} normalized such that $P_k^n(1) = 1$. These are also known as the Gegenbauer polynomials with parameter $n/2 - 1$. The functions $(x, y) \mapsto P_k^n(x \cdot y)$ are $O(n)$ -invariant positive definite kernels on S^{n-1} , and Schoenberg's characterization says that each positive definite $O(n)$ -invariant kernel on S^{n-1} is of the form

$$K(x, y) = \sum_{k=0}^{\infty} c_k P_k^n(x \cdot y)$$

for nonnegative numbers $c_k \geq 0$, where convergence is uniform and absolute [39].

Bachoc and Vallentin give a characterization for the $O(n-1)$ -invariant kernels on the sphere [2]. As they observe in [4], this also provides a characterization for the $O(n-1)$ -invariant kernels on a spherical cap,

$$\text{Cap}^{n-1}(e, \varphi) = \{x \in S^{n-1} : x \cdot e \geq \cos(\varphi)\},$$

where we view $O(n-1)$ as the stabilizer subgroup of $O(n)$ with respect to some fixed point $e \in S^{n-1}$.

To state the proposition we define the matrix $Y_k^n(u, v, t)$ by

$$Y_k^n(u, v, t)_{i,i'} = u^i v^{i'} ((1-u^2)(1-v^2))^{k/2} P_k^{n-1} \left(\frac{t - uv}{\sqrt{(1-u^2)(1-v^2)}} \right)$$

and its symmetrization

$$\overline{Y}_k^n(u, v, t) = \frac{Y_k^n(u, v, t) + Y_k^n(v, u, t)}{2}.$$

Given a topological space X we use the notation $\mathcal{C}(X \times X)$ for the space of continuous functions $X \times X \rightarrow \mathbb{R}$ and $\mathcal{C}(X \times X)_{\geq 0}$ for the cone of positive definite kernels on X . Furthermore, we denote by $\langle A, B \rangle$ the trace inner product $\text{Trace}(AB^T)$.

The following proposition is by Bachoc and Vallentin [2], where the last part about the uniform limit follows immediately from their work in combination with Theorem A.8 from [12].

PROPOSITION 2.1. *Let $e \in S^{n-1}$. For each integer $d \geq 0$ and positive semidefinite matrices $F_k \in \mathbb{R}^{(d-k+1) \times (d-k+1)}$, $k = 0, \dots, d$, the function*

$$(x, y) \mapsto \sum_{k=0}^d \left\langle F_k, \overline{Y}_k^n(x \cdot e, y \cdot e, x \cdot y) \right\rangle$$

is a $\text{Stab}_{O(n)}(e)$ -invariant positive definite kernel on S^{n-1} , and each $\text{Stab}_{O(n)}(e)$ -invariant positive definite kernel on S^{n-1} is the uniform limit of such kernels.

Define the ball $B^n(R) = \{x \in \mathbb{R}^n : \|x\| \leq R\}$. The following result is an adaptation of the above result with the sphere S^{n-1} replaced by the ball $B^n(R)$. Let

$$(Z_k^n(u, v, t))_{i,i'} = u^i v^{i'} (uv)^k P_k^n\left(\frac{t}{uv}\right),$$

and set

$$\overline{Z}_k^n(u, v, t) = \frac{Z_k^n(u, v, t) + Z_k^n(v, u, t)}{2}.$$

PROPOSITION 2.2. *Let $R > 0$. For each integer $d \geq 0$ and positive semidefinite matrices $F_k \in \mathbb{R}^{(d-k+1) \times (d-k+1)}$, $k = 0, \dots, d$, the function*

$$(x, y) \mapsto \sum_{k=0}^d \left\langle F_k, \overline{Z}_k^n(\|x\|, \|y\|, x \cdot y) \right\rangle$$

is an $O(n)$ -invariant positive definite kernel on $B^n(R)$, and each $O(n)$ -invariant positive definite kernel on $B^n(R)$ is the uniform limit of such kernels.

Proof. By [12, Theorem A.8] we need to show that the \overline{Z}_k^n are the zonal matrices for the space $B^n(R)$ with the action of $O(n)$. For this we define an orthonormal basis $Y_{k,j}^n(x)$, $j \in [d_k]$, for the space of (real-valued) spherical harmonics of degree k , which are the homogeneous polynomials of degree k in n variables that vanish under the Laplacian, and we set

$$e_{k,i,j}(x) = \|x\|^{k+i} Y_{k,j}^n\left(\frac{x}{\|x\|}\right), \quad k \geq 0, i \geq 0, j \in [d_k].$$

The span of the functions $e_{k,i,j}$ is dense in the space of continuous functions on $B^n(R)$. Moreover, these functions are symmetry adapted in the sense that there exists $O(n)$ -equivariant linear maps $T_{i,i'}$ with $T_{k,i,i'} e_{k,i,j} = e_{k,i',j}$. This means these functions form a symmetry adapted system as required by [12, Theorem A.8], except that the functions are orthonormal for an $O(n)$ -invariant inner product as opposed to an $O(n)$ -invariant measure on $B^n(R)$, but the proof in [12] still works with this weaker condition. Alternatively, instead of $\|x\|^{k+i}$, we could use $P_{k+i}(2\|x\|/R - 1)$, where P_{k+i} is the Legendre polynomial of degree $k + i$. Then the polynomials $e_{k,i,j}$ would be orthogonal with respect to the Lebesgue measure on $B^n(R)$. For the sake of simplicity, we use monomials.

The zonal matrices corresponding to the above symmetry adapted system are defined as

$$\overline{Z}_k^n(x, y)_{i,i'} = \sum_{j \in [d_k]} e_{k,i,j}(x) e_{k,i',j}(y),$$

so the proof follows by the addition theorem for spherical harmonics:

$$\sum_{j \in [d_k]} Y_{k,j}^n\left(\frac{x}{\|x\|}\right) Y_{k,j}^n\left(\frac{y}{\|y\|}\right) = P_k^n\left(\frac{x}{\|x\|} \cdot \frac{y}{\|y\|}\right). \quad \square$$

2.2. Derivation of the semidefinite programming bounds. In this paper we give upper bounds for the cardinality of codes in spheres, codes in spherical caps, and codes in balls. The semidefinite programming formulations for spheres and spherical caps are not new, but for completeness we show how these are derived from the general two- and three-point bounds. This puts everything, including our new semidefinite programming bounds for codes in balls, in a common framework. In addition it allows us to discuss in detail and supported by computations how the three-point bounds relate to the two-point bounds.

2.2.1. Two-point bounds. The Lovász ϑ -number is a semidefinite programming upper bound on the independence number of a finite graph, where the *independence number* is the size of a largest subset of the vertices where no two vertices are adjacent. In [1] this is generalized to the *spherical code graph* $G = (V, E)$, which is the graph with vertex set $V = S^{n-1}$ where two distinct vertices x and y are adjacent if $x \cdot y > \cos(\vartheta)$. This immediately generalizes to a compact topological packing graph $G = (V, E)$, which is a graph whose vertex set V is a compact Hausdorff topological space where each finite clique is a subset of an open clique (which in particular forces the independence number to be finite) [14]. This gives the optimization problem

$$(2.1) \quad \begin{aligned} \vartheta(G) = \inf \{ & M \in \mathbb{R} : K \in \mathcal{C}(V \times V)_{\succeq 0}, \\ & K(x, x) \leq M - 1 \text{ for } x \in V, \\ & K(x, y) \leq -1 \text{ for } \{x, y\} \notin E \}. \end{aligned}$$

By integrating over the symmetry group Γ of the graph G , one sees that we can restrict to Γ -invariant kernels without affecting the optimal objective value. In [1] it is shown that for the spherical code graph, Schoenberg's characterization reduces this to the Delsarte linear programming bound [15].

To consider ϑ -codes in a spherical cap one just takes a spherical code graph and restricts the vertex set to $\text{Cap}^{n-1}(e, \varphi)$. By doing this, the symmetry group reduces from $O(n)$ to $O(n-1)$. So instead of Schoenberg's characterization, Bachoc and Vallentin use Proposition 2.1 to derive the following formulation:

$$(2.2) \quad \begin{aligned} \inf \{ & M \in \mathbb{R} : F_0, \dots, F_d \succeq 0, \\ & F(u, u, 1) \leq M - 1 \text{ for } \cos(\varphi) \leq u \leq 1, \\ & F(u, v, t) \leq -1 \text{ for } (u, v, t) \in \Theta \}, \end{aligned}$$

where

$$F(u, v, t) = \sum_{k=0}^d \langle F_k, \overline{Y}_k^n(u, v, t) \rangle$$

and

$$\Theta = \left\{ (u, v, t) : \cos(\varphi) \leq u, v \leq 1, -1 \leq t \leq \cos(\vartheta), \right. \\ \left. 1 + 2uvt - u^2 - v^2 - t^2 \geq 0 \right\}.$$

Problem (2.2) is not yet a semidefinite program because we have infinitely many inequality constraints. However, since these are polynomial inequality constraints we can formulate, for each integer $\delta > 0$, a semidefinite programming upper bound by replacing each constraint of the form

$$p \geq 0 \quad \text{on} \quad S := \{x \in \mathbb{R}^n : g_i(x) \geq 0 \text{ for } i \in [m]\},$$

where p, g_1, \dots, g_m are polynomials, by the condition that there are sum-of-squares polynomials q_0 of degree 2δ and q_i of degree $2\delta - \deg(g_i)$ such that

$$p(x) = q_0(x) + g_1(x)q_1(x) + \dots + g_m(x)q_m(x).$$

This is a semidefinite constraint because if $b_\delta(x)$ is a vector whose entries form a basis for the polynomials in $\mathbb{R}[x_1, \dots, x_n]$ of total degree at most δ , then a polynomial q of degree 2δ is a sum-of-squares polynomial if and only if there is a positive semidefinite

matrix Q such that $q(x) = \langle Q, b_\delta(x)b_\delta(x)^\top \rangle$. It follows from Putinar’s theorem [38] that we get arbitrary good upper approximations by taking δ sufficiently large. In the univariate case, for $\deg(p) = 2d$, it is sufficient to take $\delta = d$, and in practice we set $\delta = d$ also in the multivariate case.

Concretely, we replace the first polynomial inequality constraint in (2.2) by

$$F(u, u, 1) - M + 1 + \langle Q_1, b_\delta(u)b_\delta(u)^\top \rangle + (u - \cos \varphi)(1 - u)\langle Q_2, b_{\delta-1}(u)b_{\delta-1}(u)^\top \rangle = 0,$$

where Q_1 and Q_2 are positive semidefinite matrices.

To replace the second polynomial inequality constraint in (2.2) we first observe that by construction $F(u, v, t) = F(v, u, t)$ for all u, v, t , and we can exploit this symmetry to get a more efficient sums-of-squares characterization.

We have

$$\mathbb{R}[u, v, t] = \mathbb{R}[u + v, uv, t] \oplus (u - v)\mathbb{R}[u + v, uv, t],$$

where $\mathbb{R}[u + v, uv, t]$ is the ring of invariant polynomials. Let $b_\delta = b_\delta(u + v, uv, t)$ be a vector whose entries form a basis for the space of polynomials in $\mathbb{R}[u + v, uv, t]$ of total degree (in the variables u, v, t) at most δ . It follows that any sum-of-squares polynomial $q(u, v, t)$ of degree 2δ that satisfies $q(u, v, t) = q(v, u, t)$ for all u, v, t is of the form

$$(2.3) \quad q(u, v, t) = \langle X_1, b_\delta b_\delta^\top \rangle + (u - v)^2 \langle X_2, b_{\delta-1} b_{\delta-1}^\top \rangle,$$

where X_1 and X_2 are positive semidefinite matrices.

We can now replace the second polynomial constraint in (2.2) by

$$(2.4) \quad \begin{aligned} &F(u, v, t) + 1 + q_1(u, v, t) \\ &+ [(u - \cos(\varphi))(1 - u) + (v - \cos(\varphi))(1 - v)]q_2(u, v, t) \\ &+ (u - \cos(\varphi))(1 - u)(v - \cos(\varphi))(1 - v)q_3(u, v, t) \\ &+ (t + 1)(\cos(\vartheta) - t)q_4(u, v, t) \\ &+ (1 + 2uvt - u^2 - v^2 - t^2)q_5(u, v, t) = 0, \end{aligned}$$

where each q_i is a sum-of-squares polynomial of the form (2.3), with $\delta = d$ for q_1 , $\delta = d - 1$ for q_2 and q_4 , and $\delta = d - 2$ for q_3 and q_5 .

Note that this symmetric formulation is quite a bit better than a naive formulation not exploiting the uv -symmetry. For example, for the $\pi/3$ -code problem in $\text{Cap}^7(e, \pi/2)$ we need to take $d = 9$, and by using the symmetries we reduce the the system from 110376 variables to 37651 variables.

We now give the direct proof that (2.2) gives a valid upper bound, since we will need the details of this proof in section 2.3.

LEMMA 2.3. *The objective value of any feasible solution of (2.2) is an upper bound on the maximal size of a ϑ -code in $\text{Cap}^{n-1}(e, \varphi)$.*

Proof. Suppose C is a ϑ -code in $\text{Cap}^{n-1}(e, \varphi)$ and (M, F_0, \dots, F_d) is a feasible solution to (2.2). On the one hand we have

$$S := \sum_{k=0}^d \left\langle F_k, \sum_{x, y \in C} \overline{Y_k^n}(x \cdot e, y \cdot e, x \cdot y) \right\rangle \geq 0,$$

since for every $k = 0, \dots, d$, the matrices F_k and $\sum_{x, y \in C} \overline{Y_k^n}(x \cdot e, y \cdot e, x \cdot y)$ are positive semidefinite. On the other hand, since the conditions in (2.2) are satisfied, we have

$$\begin{aligned}
 S &= \sum_{x \in C} \sum_{k=0}^d \langle F_k, \overline{Y}_k^n(x \cdot e, x \cdot e, 1) \rangle + \sum_{\substack{x, y \in C \\ x \neq y}} \sum_{k=0}^d \langle F_k, \overline{Y}_k^n(x \cdot e, y \cdot e, x \cdot y) \rangle \\
 &\leq \sum_{x \in C} (M - 1) + \sum_{\substack{x, y \in C \\ x \neq y}} (-1) = |C|(M - 1) - |C|(|C| - 1) = |C|(M - |C|).
 \end{aligned}$$

The inequality $|C| \leq M$ follows immediately. □

Now we consider the problem of packing spheres of radius r into a sphere of radius R . Here we consider the graph with vertex set $B^n(R - r)$, where two distinct vertices x and y are adjacent if $\|x - y\| < 2r$. Using Proposition 2.2, the optimization problem $\vartheta(G)$ reduces to

$$\begin{aligned}
 (2.5) \quad &\inf \{ M \in \mathbb{R} : F_0, \dots, F_d \succeq 0, \\
 &F(u, u, u^2) \leq M - 1 \text{ for } 0 \leq u \leq R - r, \\
 &F(u, v, t) \leq -1 \text{ for } (u, v, t) \in \Omega \},
 \end{aligned}$$

where

$$F(u, v, t) = \sum_{k=0}^d \langle F_k, \overline{Z}_k^n(u, v, t) \rangle$$

and

$$\Omega = \{ (u, v, t) : 0 \leq u, v \leq R - r, -uv \leq t \leq uv, u^2 + v^2 - 2t - 4r^2 \geq 0 \}.$$

The polynomial inequality constraints can now be replaced by sums-of-squares constraints in the same way as above.

2.2.2. Three-point bounds. Fix a point $e \in S^{n-1}$. Since $\text{Cap}^{n-1}(e, \pi) = S^{n-1}$, the linear programming bound (2.1) for spherical codes and the semidefinite programming bound (2.2) for spherical caps give the same bound when $\varphi = \pi$, but of course (2.2) is much more difficult to compute since it uses less symmetry.

The semidefinite programming bound for spherical caps looks rather similar to the three-point bound for spherical codes; both use Proposition 2.1. Before we give the derivation of the three-point bound, we first mention that improved bounds can already be obtained by computing the two-point bound (2.2) for $\text{Cap}^{n-1}(e, \pi - \vartheta)$ and then adding 1 to the resulting value. This gives an upper bound because we can always rotate a spherical code so that the point $-e$ is in the code. Although this usually only gives a small improvement, in dimension 4 it gives the upper bound 24.983 for $d = 10$, which shows that the Lovász ϑ -number is actually already enough to prove the optimality of the 24-cell for the kissing number problem.

Of course, for spherical code problems it is better to use the three-point bound, because it is equally difficult to compute and gives better bounds. To get the full three-point bound we have to also derive constraints coming from functions on $V \times V \times V$, where $V = S^{n-1}$. We follow the derivation from [13], where a general formulation for k -point bounds is given. For $k = 3$ we get

$$\begin{aligned}
 (2.6) \quad &\inf \{ M \in \mathbb{R} : T \in \mathcal{C}(V \times V \times I_1)_{\succeq 0}, \\
 &B_3T(S) \leq M - 1 \text{ for } S \in I_{=1}, \\
 &B_3T(S) \leq -2 \text{ for } S \in I_{=2}, \\
 &B_3T(S) \leq 0 \text{ for } S \in I_{=3} \},
 \end{aligned}$$

where $I_{=k}$ is the set of independent sets of cardinality k in the spherical code graph, and I_k is the union of $I_{=i}$ for $0 \leq i \leq k$. Here $B_3: \mathcal{C}(V \times V \times I_1)_{\text{sym}} \rightarrow \mathcal{C}(I_3 \setminus \{\emptyset\})$ is the operator defined by

$$(2.7) \quad B_3 T(S) = \sum_{\substack{Q \subseteq S \\ |Q| \leq 1}} \sum_{\substack{x, y \in S \\ Q \cup \{x, y\} = S}} T(x, y, Q).$$

In (2.6) we can restrict to $O(n)$ -invariant functions. As shown in [13], using Proposition 2.1, problem (2.6) then reduces to the Bachoc–Vallentin bound: Set

$$T(x, y, \emptyset) = \sum_{k=0}^d a_k P_k^n(x \cdot y)$$

and

$$T(x, y, \{z\}) = \sum_{k=0}^d \left\langle F_k, S_k^n(x \cdot z, y \cdot z, x \cdot y) \right\rangle,$$

where

$$S_k^n = \frac{1}{6} \sum_{\sigma \in S_3} \sigma Y_k^n$$

in which we sum over the permutations on three elements and each element $\sigma \in S_3$ acts on Y_k^n by permuting its arguments. Then, (2.6) reduces to

$$(2.8) \quad \inf \left\{ M \in \mathbb{R} : a_0, \dots, a_d \geq 0, F_0, \dots, F_d \succeq 0, \right. \\ \sum_{k=0}^d a_k + F(1, 1, 1) \leq M - 1, \\ \sum_{k=0}^d a_k P_k^n(u) + 3F(u, u, 1) \leq -1 \text{ for } -1 \leq u \leq \cos(\vartheta), \\ \left. F(u, v, t) \leq 0 \text{ for } (u, v, t) \in \Delta \right\},$$

where

$$F(u, v, t) = \sum_{k=0}^d \left\langle F_k, S_k^n(u, v, t) \right\rangle$$

and

$$\Delta = \{(u, v, t) : -1 \leq u, v, t \leq \cos(\vartheta), 1 + 2uv t - u^2 - v^2 - t^2 \geq 0\}.$$

Analogously to the two-point bound, we can use sums-of-squares relaxations to formulate semidefinite programming upper bounds. For this we write the last condition as

$$F(u, v, t) + q_0(u, v, t) + p(u)q_1(u, v, t) + p(v)q_2(u, v, t) \\ + p(t)q_3(u, v, t) + (1 + 2uv t - u^2 - v^2 - t^2)q_4(u, v, t) = 0,$$

where $p(u) = (\cos(\theta) - u)(1 - u)$ and q_0, \dots, q_4 are sum-of-squares polynomials.

The function $F(u, v, t)$ is symmetric in u, v, t , and as in [30] we can use this symmetry to give a more efficient characterization using [19]. First we reformulate the sums-of-squares factorization as

$$F(u, v, t) + q_0(u, v, t) + \sum_{i=1}^4 s_i q_i(u, v, t) = 0,$$

where

$$\begin{aligned} s_1 &= p(u) + p(v) + p(t), & s_2 &= p(u)p(v) + p(u)p(t) + p(v)p(t), \\ s_3 &= p(u)p(v)p(t), & s_4 &= 1 + 2uvt - u^2 - v^2 - t^2. \end{aligned}$$

Now, without loss of generality, we may assume the sum-of-squares polynomials q_0, \dots, q_4 to be symmetric in u, v, t . Let

$$\theta_1 = u + v + t, \quad \theta_2 = uv + ut + vt, \quad \theta_3 = uvt,$$

and let $b_\delta = b_\delta(\theta_1, \theta_2, \theta_3)$ be a vector whose entries form a basis for the space of polynomials in the invariant ring $\mathbb{R}[\theta_1, \theta_2, \theta_3]$ of total degree (in the variables u, v, t) at most δ . By [19], for each S_3 -invariant sum-of-squares polynomial p of degree 2δ there are positive semidefinite matrices Q_1, Q_2, Q_3 such that

$$p(u, v, t) = \left\langle Q_1, b_\delta b_\delta^T \otimes \Pi_1 \right\rangle + \left\langle Q_2, b_{\delta-3} b_{\delta-3}^T \otimes \Pi_2 \right\rangle + \left\langle Q_3, b_{\delta-2} b_{\delta-2}^T \otimes \Pi_3 \right\rangle,$$

where

$$\begin{aligned} \Pi_1 &= 1, & \Pi_2 &= \theta_1^2 \theta_2^2 - 4\theta_2^3 - 4\theta_1^3 \theta_3 + 18\theta_1 \theta_2 \theta_3 - 27\theta_3^2, \\ \Pi_3 &= \begin{pmatrix} 2\theta_1^2 - 6\theta_2 & -\theta_1 \theta_2 + 9\theta_3 \\ -\theta_1 \theta_2 + 9\theta_3 & 2\theta_2^2 - 6\theta_1 \theta_3 \end{pmatrix}. \end{aligned}$$

LEMMA 2.4. *The optimal value of the semidefinite program given in (2.8) is an upper bound on the maximal size of a ϑ -code in S^{n-1} .*

Proof. Let $(M, a_0, \dots, a_d, F_0, \dots, F_d)$ be a feasible solution of problem (2.8), and let C be a ϑ -code in S^{n-1} .

For each $k \in \{0, \dots, d\}$, the matrices $\sum_{(x,y,z) \in C^3} \overline{Y_k^n}(x \cdot z, y \cdot z, x \cdot y)$ and F_k are positive semidefinite, so

$$S := \sum_{(x,y,z) \in C^3} F(x \cdot z, y \cdot z, x \cdot y) = \sum_{k=0}^d \left\langle F_k, \sum_{(x,y,z) \in C^3} \overline{Y_k^n}(x \cdot z, y \cdot z, x \cdot y) \right\rangle \geq 0.$$

On the other hand, S is equal to

$$|C| F(1, 1, 1) + \sum_{\substack{(x,z) \in C^2 \\ x \neq z}} 3 F(x \cdot z, x \cdot z, 1) + \sum_{\substack{(x,y,z) \in C^3 \\ x \neq z, y \neq z, x \neq y}} F(x \cdot z, y \cdot z, x \cdot y).$$

In the above equation, we use that $F(u, v, t)$ is invariant under the permutations of (u, v, t) . Since F has to satisfy the constraints in (2.8), we get

$$\begin{aligned} S &\leq |C| \left(M - 1 - \sum_{k=0}^d a_k \right) - |C|(|C| - 1) - \sum_{x,y \in C} \left(\sum_{k=0}^d a_k P_k^n(x \cdot y) \right) \\ &\leq |C|(M - 1) - |C|(|C| - 1) - \sum_{k=0}^d a_k \left(|C| + \sum_{\substack{(x,y) \in C^2 \\ x \neq y}} P_k^n(x \cdot y) \right) \end{aligned}$$

$$\begin{aligned} &\leq |C|(M - 1) - |C|(|C| - 1) - \sum_{k=0}^d a_k \sum_{x,y \in C} P_k^n(x \cdot y) \\ &\leq |C|(M - 1) - |C|(|C| - 1). \end{aligned}$$

So we have shown $0 \leq S \leq |C|(M - 1) - |C|(|C| - 1)$, which implies $|C| \leq M$. \square

2.3. Information from exact sharp solutions. Since all our programs give upper bounds on the size of optimal configurations, which has to be an integer, one might wonder why we are interested in exact bounds. Indeed, if we already know a configuration C of M points, any upper bound strictly lower than $M + 1$ ensures the optimality of C .

First, these bounds are a priori only upper bounds on the independence number of the corresponding graph, and it is interesting to point out when these bounds give exactly the independence number. For example, Bachoc and Vallentin proved in [3] that for ϑ -codes in four dimensions, where $\cos \vartheta = 1/6$, the three-point bound gives exactly 10 (even though the two-point bound is not sharp here).

The second interest is geometric: from the proof of Lemma 2.3, we can see that any feasible solution reaching M as objective value provides additional information regarding optimal solutions. For any code C such that $|C| = M$, all the inequalities in the proof of Lemma 2.3 have to be equalities. We get the so-called *complementary slackness* conditions.

COROLLARY 2.5. *Let (M, F_0, \dots, F_d) be a feasible solution to (2.2). If C is a ϑ -code in S^{n-1} with cardinality $|C| = M$, then the following equalities hold:*

(i) for every $k = 0, \dots, d$,

$$\left\langle F_k, \sum_{x,y \in C} \overline{Y_k^n}(x \cdot e, y \cdot e, x \cdot y) \right\rangle = 0;$$

(ii) for every x in C ,

$$F(x \cdot e, x \cdot e, 1) = M - 1;$$

(iii) for every distinct x, y in C ,

$$F(x \cdot e, y \cdot e, x \cdot y) = -1.$$

Conditions (ii) and (iii) are of special interest. If we define the polynomial

$$P(u) = F(u, u, 1) - (M - 1),$$

then for every point x in an optimal code, the inner product $e \cdot x$ is in the set R of roots of P . This gives finitely many possibilities for these inner products. Then, for every $u, v \in R$, we can define

$$P_{u,v}(t) = F(u, v, t) + 1,$$

and for every pair of distinct points (x, y) in an optimal code, the inner product $x \cdot y$ has to be a root of one such polynomial. With this procedure, we get all the possible inner products occurring in the configuration. Once that we get all the possible triples (u, v, t) , we can even use (i) to get the number of occurrences of each triple by solving a linear system.

If we consider the semidefinite programming bound in (2.5) for the number of spheres that can be packed in a given sphere, we will get the possible norms of the centers of the spheres and the possible inner products among these centers.

Analogously, three-point bounds also provide complementary slackness conditions. From Lemma 2.4, we get the following.

COROLLARY 2.6. *Let $(M, a_0, \dots, a_d, F_0, \dots, F_d)$ be a feasible solution of the semidefinite program (2.8). Let C be a ϑ -code in S^{n-1} . If $M = |C|$, then the following properties hold:*

(i)

$$\sum_{k=0}^d a_k + F(1, 1, 1) = M - 1;$$

(ii) for each two distinct $x, y \in C$,

$$\sum_{k=0}^d a_k P_k^n(u) + 3F(x \cdot y, x \cdot y, 1) = -1;$$

(iii) for each three distinct $x, y, z \in C$,

$$F(x \cdot z, y \cdot z, x \cdot y) = 0;$$

(iv) for each $k \in \{0, \dots, d\}$,

$$\left\langle F_k, \sum_{(x,y,z) \in C^3} \overline{Y}_k^n(x \cdot z, y \cdot z, x \cdot y) \right\rangle = 0;$$

(v) for each $k \in \{0, \dots, d\}$,

$$a_k \sum_{x,y \in C} P_k^n(x \cdot y) = 0.$$

In this situation, the procedure described above also gives the possible inner products among points in optimal codes and the distribution of the triples (u, v, t) .

In several cases, this information turns out to be enough to prove the uniqueness of optimal configurations, as we will see in section 4.

On the other hand, if we have a candidate optimal configuration and the bound is sharp, then any optimal solution of the semidefinite program will satisfy the complementary slackness conditions implied by this configuration. Adding these conditions to the semidefinite program usually results in a faster convergence of the interior point method. The complementary slackness conditions can be important to get an exact optimal solution. In the next section, we give an automatic procedure to extract an exact optimal solution from a numerical optimal floating point solution.

3. The rounding procedure. In general, the fact that a semidefinite program is defined over a given algebraic field does not ensure that the optimal solution can be defined over the same field. In fact, even if the constraints and the objective are rational, an optimal solution might require high algebraic degree [34]. However, for each packing problem that we considered where we have a sharp bound, we were able to find an optimal solution over the same field that is required to define the semidefinite program and to formulate the complementary slackness conditions coming from the optimal solution. We first focus on the problem of finding a rational optimal solution and extend this to quadratic fields in section 3.4.

We work with semidefinite programs in block form:

$$\inf \left\{ \sum_{i=1}^l \langle C_i, X_i \rangle : X_1, \dots, X_l \succeq 0, \right. \\ \left. \sum_{i=1}^l \langle A_{i,j}, X_i \rangle = b_j \text{ for } j = 1, \dots, m \right\},$$

where the symmetric matrices C_i and $A_{i,j}$ and the scalars b_j are all rational.

First, since we already know the optimal objective value, we do not solve the semidefinite program as an optimization problem, but we add a linear constraint enforcing the objective value and solve the problem as a feasibility problem. One advantage of this is that by rounding the floating point solution from the semidefinite programming solver to a rational solution, such that all linear equations are satisfied, the objective value will automatically be correct. A second advantage is that the numerical interior point solver we use (SDPA-GMP [18]) gives an approximate solution in the relative interior of the feasible set when solving a feasibility problem, which will be important in section 3.2. When rounding we have to make sure that the obtained rational matrices are positive semidefinite and satisfy all linear equations.

3.1. Rounding in the affine space. Let X_1^*, \dots, X_l^* be the (high precision) floating point output of the semidefinite programming solver. Because of floating point arithmetic they do not satisfy the linear constraints

$$\sum_{i=1}^l \langle A_{i,j}, X_i \rangle = b_j \quad \text{for } j = 1, \dots, m$$

exactly. First we want an exact solution to the above equations that is close to the approximate solution X_1^*, \dots, X_l^* . We rewrite the above linear conditions as the linear system $Ax = b$, so that the concatenation x^* of $\text{vec}(X_i^*)$ for $i = 1, \dots, l$ is an approximate solution of this system. Here vec is the vectorization operator that turns a symmetric matrix into a column vector.

We want to find a vector that satisfies the linear system $Ax = b$ exactly and is close to x^* . One natural way to do it is the following: We transform the system $Ax = b$ into reduced row echelon form, which can be done in rational arithmetic. Then, when solving this system by backsubstitution, for each free variable that we encounter, we use a rational approximation (possibly with some upper bound on the size of the denominator) of the corresponding entry of x^* . We obtain an optimal rational solution of the semidefinite program that satisfies all linear constraints exactly.

In our applications, the linear systems we are dealing with can be large, even after exploiting the symmetries in the sums-of-squares formulation. We therefore write the system as a linear system over the integers and use the Kannan–Bachem algorithm [25] as implemented in [17], which can solve the biggest system we consider in this paper within eight hours on a normal desktop computer.

By continuity of roots, the eigenvalues of the blocks in the rounded solution will be close to the eigenvalues of the blocks in the floating point approximate solution. If the floating point solution did not have near zero eigenvalues, then we would be done. However, in our situation, the matrices X_i^* have many eigenvalues close to zero, so that the rounded solution typically has negative eigenvalues.

Since we already solved the problem as a feasibility problem, the solution we are working with lies in the relative interior of the feasible set of our feasibility problem.

This means there exists no solution that has fewer near zero eigenvalues. Even if we somehow manage to obtain a solver output X_1^*, \dots, X_l^* where all near zero eigenvalues are positive, for example, by adding constraints of the form $X - \varepsilon I \succeq 0$, then the above rounding procedure is likely to turn some of these into negative numbers. Instead of trying to make these eigenvalues positive, our strategy consists in forcing them to be exactly zero by adding some linear constraints.

3.2. Kernel detection. One approach to find these rational linear constraints is to use the complementary slackness condition from Corollary 2.5 or Corollary 2.6 arising from a candidate optimal packing configuration.

In [3], after a slight reformulation of the semidefinite program and some additional work (see section 4.2), these linear constraints are sufficient to get an exact bound. However, in general, especially when the matrices involved are large (see section 4.1), this approach does not provide all the linear constraints, even if we also include constraints coming from the derivatives. We therefore propose an automated procedure to find all necessary linear conditions.

By computing the kernel of X_i^* in high precision floating point arithmetic, we get the eigenvectors corresponding to the near zero eigenvalues. We list these vectors as the columns of the matrix N_i . These vectors are typically not approximations of rational vectors themselves, but approximations of real linear combinations of rational vectors, without any control on the coefficients. Hence there is no point in trying to round them to rational vectors. Instead, we try to extract a rational basis of this kernel by searching for integer equations defining this linear space. To do so, we use the function `lindep` from Nemo [17] that uses the LLL algorithm [28] to find an integer linear combination of the rows of N_i that is close to the zero vector. We remove one row of N_i whose coefficient in this linear combination is nonzero and repeat the procedure to find another integer linear equation, linearly independent from the previous one. We continue until we found the right number of equations. Let M_i be the matrix with these integer relations as its rows. Then we can compute a basis for the nullspace of M_i in rational arithmetic, and these vectors will be the kernel vectors of the rounded version of the matrix X_i^* .

3.3. Rounding and checking. For each kernel basis vector for X_i^* that we find we add the constraints $X_i v = 0$ to the linear system $Ax = b$. If there indeed exist rational bases for the kernel of each matrix in the solution, then by performing the rounding procedure mentioned above on this extended semidefinite program, we find positive semidefinite matrices that satisfy all linear constraints exactly. Due to the new linear equations, we make sure that the rounded matrices have no negative eigenvalues.

We have to verify that this is indeed a solution. First the linear constraints can be verified in rational arithmetic. Then we need to verify that the matrices are positive semidefinite. To do so, we compute their characteristic polynomial and use the property that a real-rooted polynomial $f(t)$ has no negative roots if and only if $(-1)^{\deg f} f(-t)$ has only nonnegative coefficients.

Using this approach, we find an optimal rational solution for all considered packing problems where our semidefinite programming bound is sharp and where the complementary slackness conditions implied by an optimal configuration are rational.

3.4. Extension to quadratic fields. Finally we extend the above approach to the case where we want to round over a quadratic field, which is a natural thing to try

whenever the semidefinite program and/or the complementary slackness conditions arising from an optimal configuration are defined over such a field.

We first describe how the rounding part from section 3.1 can be adapted. In this situation, A and b of the linear system $Ax = b$ are defined over $\mathbb{Q}[\sqrt{\ell}]$. We have an approximate floating point solution x^* , and we are looking for an exact solution $x = x_1 + x_2\sqrt{\ell}$ over $\mathbb{Q}[\sqrt{\ell}]$ that is close to x^* . After writing $A = A_1 + A_2\sqrt{\ell}$ and $b = b_1 + b_2\sqrt{\ell}$ with the entries of A_1, A_2, b_1, b_2 rational, we want to find an exact solution of the following rational linear system:

$$(3.1) \quad \begin{pmatrix} A_1 & \ell A_2 \\ A_2 & A_1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \end{pmatrix}$$

such that $x_1 + x_2\sqrt{\ell} \approx x^*$. To do this, we first need to extract from x^* an approximate solution (x_1^*, x_2^*) of the above linear system such that $x_1^* + x_2^*\sqrt{\ell} \approx x^*$. To find these vectors we write

$$x_1^* = y \quad \text{and} \quad x_2^* = \frac{1}{\sqrt{\ell}}(x^* - y)$$

and solve the linear system

$$\begin{pmatrix} A_1 - \sqrt{\ell}A_2 \\ \sqrt{\ell}A_2 - A_1 \end{pmatrix} y = \begin{pmatrix} b_1 - \sqrt{\ell}A_2x^* \\ \sqrt{\ell}b_2 - A_1x^* \end{pmatrix}$$

for y in floating point arithmetic. We now reduce the rational linear system (3.1) to reduced row echelon form and use backsubstitution to find a rational vector satisfying this system, where for each free variable we use a rational approximation of the corresponding entry in (x_1^*, x_2^*) .

To find the kernel constraints we again first compute an arbitrary basis for the kernel of X_i^* in high precision floating point arithmetic and list the vectors as the columns of the matrix N_i . Then we set

$$M_i = \begin{pmatrix} N_i \\ \sqrt{\ell}N_i \end{pmatrix}$$

and use the LLL algorithm to find an integer linear combination (λ, μ) such that $(\lambda, \mu)^T M_i \approx 0$. Once we find the right number of equations, we can build the matrix H , with rows $(\lambda, \ell\mu)^T$ and $(\mu, \lambda)^T$, for each equation (λ, μ) that we find. By construction, every vector (u, v) in the kernel of H satisfies $X_i^*(u + v\sqrt{\ell}) \approx 0$, and we want $u + v\sqrt{\ell}$ to be in the kernel of the rounded version $X_{i,1} + X_{i,2}\sqrt{\ell}$ of X_i^* . So every vector in a basis of the kernel of H provides two equations

$$X_{i,1}u + \ell X_{i,2}v = 0, \quad X_{i,2}u + X_{i,1}v = 0$$

that we add to the linear system (3.1), taking into account the block structure. Finally, we can apply the rounding procedure in order to get an optimal solution over $\mathbb{Q}[\sqrt{\ell}]$.

There is no theoretical objection to generalizing the above procedure to higher degree algebraic number fields. However, one issue is that the number of variables in the rational system (3.1) will be multiplied by the degree of the extension. It would then be interesting to investigate alternative methods to this step in the procedure, where the linear algebra is done directly over the number field.

4. Applications. In this section we determine exact sharp semidefinite programming bounds for packing problems, and we describe how to prove the uniqueness of some optimal configurations by using the information obtained from complementary slackness.

The code for setting up the semidefinite programs, for running the rounding procedure, and for rigorously checking the rounded solution can be found in the ancillary files from the [arXiv.org](https://arxiv.org) e-print archive. This program runs in Julia 1.1.0 [7] and uses the computer algebra system Nemo [17]. See the included README.txt file for information on how to run the code.

4.1. Codes in spherical caps. In 1979, Odlyzko and Sloane [35] and, independently, Levenshtein [29] proved that the maximal size of a $\pi/3$ -code in S^7 is 240. Such a spherical code C is given by the minimal vectors of the root lattice E_8 . Moreover, in 1981 Bannai and Sloane [6] showed that the maximal $\pi/3$ -code in S^7 is unique up to isometry. Let e be an arbitrary element of C ; then $C \cap \text{Cap}^7(e, \pi/2)$ is a $\pi/3$ -code on the hemisphere $\text{Cap}^7(e, \pi/2)$ with cardinality 183. In 2009, Bachoc and Vallentin [4] proved that the maximal size of such a code is indeed 183. Here we prove that a $\pi/3$ -code in $\text{Cap}^7(e, \pi/2)$ with maximal size is unique up to isometry.

LEMMA 4.1. *Let C be a $\pi/3$ -code in $\text{Cap}^7(e, \pi/2)$ with cardinality 183. Then*

(i) *for every $c \in C$,*

$$e \cdot c \in \{0, 1/2, 1\};$$

(ii) *for any distinct $c, c' \in C$,*

$$c \cdot c' \in \{-1, \pm 1/2, 0\}.$$

Proof. Let (F_0, \dots, F_9) be the exact optimal solution obtained by applying our rounding procedure to the floating point output (available as `183points.jls` in the arXiv version of this paper) that we obtained by solving (2.2) for $d = 9$, and consider

$$F(u, v, t) = \sum_{k=0}^9 \langle F_k, \overline{Y_k^n}(u, v, t) \rangle.$$

Let us first prove (i). Since C is an optimal configuration, following (ii) in Corollary 2.5, every $c \in C$ has to satisfy

$$F(e \cdot c, e \cdot c, 1) = 182.$$

This means that $e \cdot c$ has to be a root of the univariate polynomial

$$g(u) = F(u, u, 1) - 182$$

located in $\Delta_0 = [0, 1]$. By computing its Sturm sequence, we check that the polynomial g has exactly three distinct roots in the interval $(-\varepsilon, 1]$ for a fixed, arbitrary, small enough $\varepsilon > 0$. Since $g(0) = g(1/2) = g(1) = 0$, g cannot have any further roots in $[0, 1]$.

In order to prove (ii), consider any distinct $c, c' \in C$. Following (iii) in Corollary 2.5, we have

$$F(e \cdot c, e \cdot c', c \cdot c') = -1.$$

Due to (i), $e \cdot c$ and $e \cdot c'$ have to be in $\{0, 1/2, 1\}$. As a consequence, $c \cdot c'$ is a root of the univariate polynomial

$$h(t) = F(u_0, v_0, t) + 1$$

for some $u_0, v_0 \in \{0, 1/2, 1\}$. Moreover

$$c \cdot c' \in \{t \in \mathbb{R} : (u_0, v_0, t) \in \Delta\}.$$

Using the same procedure involving the Sturm sequence, we check that all those possible roots lie in $\{-1, \pm 1/2, 0\}$. \square

Due to the properties given in Lemma 4.1, we can prove that E_8 provides the unique optimal $\pi/3$ -code in $\text{Cap}^7(e, \pi/2)$.

THEOREM 4.2. *There is, up to rotations fixing the point e , a unique $\pi/3$ -code in $\text{Cap}^7(e, \pi/2)$ with 183 elements.*

Proof. Let L_0 be the additive subgroup of \mathbb{R}^8 spanned by C . If x and y are two vectors in L_0 , according to (ii) in Lemma 4.1, the inner product $\sqrt{2}x \cdot \sqrt{2}y$ is an integer. So the additive subgroup $L = \sqrt{2}L_0$ is an integral lattice. Moreover, since it is spanned by a set of vectors v such that $v \cdot v = 2$, L has to be a root lattice. Due to the classification of root lattices by Kneser [26], L can be written as a direct sum of the irreducible root lattices A_d, D_d, E_6, E_7, E_8 . Assume that $L = \bigoplus_{i=1}^k L_i$ for some k , where L_i is an irreducible root lattice for every $i = 1, \dots, k$. We denote by r_i the number of roots of L_i and by d_i its rank. The number of roots of the irreducible root lattices is well known (see [10]), and if L_i is not E_8 , then we have $r_i/d_i < 183/8$. Hence if L is not E_8 , its number of roots r satisfies

$$r = \sum_{i=1}^k r_i = \sum_{i=1}^k d_i \frac{r_i}{d_i} < 183.$$

So $L = E_8$.

There are 240 roots in E_8 . Among the corresponding 240 points of L_0 in S^7 , let n_1 and n_2 be the numbers of points lying, respectively, on the equator $\{x \in S^7 : x \cdot e = 0\}$ of S^7 and on the strict upper hemisphere $\{x \in S^7 : x \cdot e > 0\}$. By symmetry we have $n_1 + 2n_2 = 240$. On the other hand, on the same hemisphere we cannot have more than the 183 points coming from C , so $n_1 + n_2 = 183$. Thus the strict upper hemisphere of S^7 contains exactly $n_2 = 57$ elements of C . Suppose that e is not in C . Then, following (i) in Lemma 4.1, 57 elements of the code would lie in $\{x \in S^7 : x \cdot e = 1/2\}$. This would give a ϑ -code in S^6 with 57 elements, where ϑ is such that $\cos \vartheta = 1/3$. Bannai and Sloane [6, Theorem 9] proved that such a code does not exist. So e has to be an element of C , and C is the configuration that we expect. \square

One might wonder whether this approach can be used in dimension 24. However, whereas the Leech lattice Λ_{24} provides a configuration with 144855 points on the hemisphere, the best upper bound that we obtained numerically is only 158611, with $d = 13$.

4.2. Spherical codes. In [3], Bachoc and Vallentin proved the following theorem.

THEOREM 4.3. *The Petersen code is, up to symmetry, the unique ϑ -code in S^3 of cardinality 10, where ϑ is such that $\cos \vartheta = 1/6$.*

Their approach, based on semidefinite optimization, consists of three steps: First, they provide an exact optimal solution for a slightly different formulation of problem (2.8). From this solution, they derive the three-point distance distribution of an optimal code. Finally they prove that the Petersen code is the only code satisfying this distribution. However, the way they obtain their exact optimal solution is not

straightforward. First of all, their new formulation of problem (2.8) gives further information about the solution matrices (F_0, F_1, F_2) , in case they provide an optimal solution. These properties help to determine an exact optimal solution of their semi-definite program, after several steps and various computations. However, the strategy that they use seem to apply only in this specific situation, especially because for this problem the corresponding solution of problem (2.8) is small, since taking $d = 2$ is sufficient.

In our framework, the first step of their approach gets much easier: we can directly solve problem (2.8) and turn the approximate solution from the solver into an exact optimal solution by using our rounding procedure. Then we need to check that our solution also implies the three-point distance distribution of an optimal code.

Proof of Theorem 4.3. Let C be a ϑ -code in S^3 with cardinality 10, where ϑ is such that $\cos \vartheta = 1/6$. The three-point distance distribution of C is defined by

$$\alpha(u, v, t) = \frac{1}{|C|} |\{(c, c', c'') \in C^3 : c \cdot c' = u, c \cdot c'' = v, c' \cdot c'' = t\}|.$$

In order to determine this distribution, we first need to know the possible inner products between the elements of C . Let $(M, a_0, \dots, a_d, F_0, \dots, F_d)$ be the exact optimal solution obtained by applying our rounding procedure to the floating point output (available as `10points.js` in the arXiv version of this paper) that we obtained by solving (2.8) for $d = 6$. Using the complementary slackness condition given in (ii) in Corollary 2.6, the inner product between two distinct elements of C must be a root of the polynomial

$$\sum_{k=0}^d a_k P_k^n(u) + 3 \sum_{k=0}^d \langle F_k, \bar{Y}_k^n(u, u, 1) \rangle + 1$$

located between -1 and $1/6$. By using a Sturm sequence, we check that the only possible roots are $-2/3$ and $1/6$.

Now, due to symmetries, the three-point distance distribution of C is defined by the six values

$$\begin{aligned} &\alpha(1, 1, 1), && \alpha(1/6, 1/6, 1), && \alpha(-2/3, -2/3, 1), \\ &\alpha(1/6, 1/6, 1/6), && \alpha(-2/3, 1/6, 1/6), && \alpha(-2/3, -2/3, 1/6). \end{aligned}$$

We already know $\alpha(1, 1, 1) = 1$. By combining equation $\sum_u \alpha(u, u, 1) = 10$ with the complementary slackness conditions given by (v) in Corollary 2.6, we get $\alpha(1/6, 1/6, 1) = 6$ and $\alpha(-2/3, -2/3, 1) = 3$. Then by solving the linear system involving $\sum_{u,v,t} \alpha(u, v, t) = 100$ together with the equations given by (iv) in Corollary 2.6, we get

$$\alpha(1/6, 1/6, 1/6) = 18, \quad \alpha(-2/3, 1/6, 1/6) = 12, \quad \alpha(-2/3, -2/3, 1/6) = 6.$$

So we recovered the distribution described in [3], and we can apply their last argument to prove that C has to be the Petersen code. \square

The optimal value of the program (2.8) is an upper bound on the cardinality of a ϑ -code in S^2 where ϑ is such that $\cos \vartheta = (2\sqrt{2} - 1)/7$. By using our rounding approach we obtain an exact solution with value 8. Schütte and van der Waerden [40] proved that this is the optimal value. The *square antiprism* provides a spherical code with 8 points in S^2 . Due to Danzer [11] this code is unique up to symmetry. His proof relies on heavy geometric arguments: For example, understanding the angles and

distances which may occur in such a configuration requires a very technical analysis. Here we recover this information thanks to the complementary slackness conditions and provide a new proof of the uniqueness of such a code.

THEOREM 4.4. *The largest ϑ -code in S^2 with $\cos \vartheta = (2\sqrt{2} - 1)/7$ is unique up to symmetry.*

Proof. Let $(M, a_0, \dots, a_d, F_0, \dots, F_d)$ be the exact optimal solution obtained by applying our rounding procedure to the floating point output (available as `8points.jls` in the arXiv version of this paper) that we obtained by solving (2.8) for $d = 7$. Furthermore, let C be an optimal spherical code. Due to the complementary slackness condition given in (ii) in Corollary 2.6, the real roots of the polynomial

$$\sum_{k=0}^d a_k P_k^n(u) + 3 \sum_{k=0}^d \langle F_k, \overline{Y}_k^n(u, u, 1) \rangle + 1$$

located between -1 and $\frac{2\sqrt{2}-1}{7}$ give the possible inner products between two points in C . By using its Sturm sequence, we can check that the only inner products are

$$(2\sqrt{2} - 1)/7, \quad -3(2\sqrt{2} - 1)/7, \quad -(2\sqrt{2} - 1)^2/7.$$

As in the proof of Theorem 4.3, the exact solution, together with the complementary slackness condition (v) in Corollary 2.6 and the properties of the three-point distribution, gives

$$\begin{aligned} \alpha(1, 1, 1) &= 1, & \alpha(v, v, 1) &= 4, \\ \alpha(-3v, -3v, 1) &= 2, & \alpha\left((1 - 2\sqrt{2})v, (1 - 2\sqrt{2})v, 1\right) &= 1, \end{aligned}$$

where $v = (2\sqrt{2} - 1)/7$.

This is already enough information for proving the uniqueness of the configuration. We want to show that C is, up to rotations, the squared antiprism S_8 . We label with c_1, \dots, c_8 the vertices of S_8 , as depicted in Figure 1.

Up to symmetry, we may assume that c_1 is in C . The possible inner products imply that for any $c \in C$, the distance between c and c_1 is among

$$\delta_1 = \sqrt{2 - 2v}, \quad \delta_2 = \sqrt{2 - 2(1 - 2\sqrt{2})v}, \quad \delta_3 = \sqrt{2 + 6v}.$$




FIG. 1. *The squared antiprism.*

Note that $\delta_1 < \delta_2 < \delta_3$. In Figure 1, the points c_2, c_3, c_4 , and c_5 , depicted in red, are at distance δ_1 from c_1 , the blue point c_8 is at distance δ_2 from c_1 , and the green points c_6 and c_7 are at distance δ_3 from c_1 . For every $c \in C$, we define the circle $\mathcal{C}_{\delta_1}(c)$ made of the points of S^2 at distance δ_1 from c_1 . On the one hand, since $\alpha(v, v, 1) = 4$, on average over C , the circle $\mathcal{C}_{\delta_1}(c)$ contains four points of C . On the other hand, it is straightforward to check that one cannot put more than four points on such a circle without violating the distance constraints. This means that for every $c \in C$, there are exactly four elements of the code C in $\mathcal{C}_{\delta_1}(c)$.

In particular, $\mathcal{C}_{\delta_1}(c_1)$ contains four points of C . In fact, there is, up to rotations, only one way to put four compatible points on such a circle. Hence, we may assume that these four points are c_2, c_3, c_4 , and c_5 (see Figure 2).

Three points remain. On the circle $\mathcal{C}_{\delta_1}(c_3)$, we already know three points out of four. This leaves two possibilities for the last codeword on that circle, and the same situation holds in $\mathcal{C}_{\delta_1}(c_4)$. This gives four candidates, but among them, only two are compatible: c_6 and c_7 . Finally, the last remaining point has to lie in the intersection of $\mathcal{C}_{\delta_1}(c_2)$ and $\mathcal{C}_{\delta_1}(c_5)$, and the distance constraints force it to be c_8 . \square

4.3. Sphere in sphere packings. The optimal solution of the semidefinite program (2.5) is an upper bound on the number of spheres of radius r that can be packed into a sphere of radius R . Using our program we determine exact sharp solutions for several values of r and R in various dimensions. In this section we show that we can extract bounds that work for infinitely many different dimensions by extrapolating from the rounded solutions for a few dimensions. In this way we show that the Lovász ϑ -number gives a sharp bound for some families of problems.

THEOREM 4.5. *The Lovász ϑ -number gives a sharp bound on the largest number M of n -dimensional unit spheres that can be packed into a sphere of radius R for*

- (i) $n \geq 2$ with $R = 2$ and $M = 2$,
- (ii) $n \geq 2$ with $R = 2/\sqrt{3} + 1$ and $M = 3$,
- (iii) $n \geq 2$ with $R = \sqrt{2n/(n+1)} + 1$ and $M = n + 1$,
- (iv) $n \geq 2$ with $R = \sqrt{2} + 1$ and $M = 2n$,
- (v) $n = 2$ with $R = 1 + \sqrt{2(1 + 1/\sqrt{5})}$ and $M = 5$,
- (vi) $n = 2$ with $R = 3$ and $M = 7$.

To obtain the proof for (i)–(iv) we first compute an exact sharp solution for several dimensions. From this we notice we can take the same degree d for each dimension n , and we make a guess as to what the general solution of the semidefinite program should be as a function of n . For the sums-of-squares matrices we then compute




FIG. 2. *The only four point configuration in $\mathcal{C}_{\delta_1}(c)$.*

the Cholesky decomposition using symbolic mathematics, which gives us the sums-of-squares decomposition proving that the solution satisfies the right inequalities. Even if we stated the theorem in an unified way with $r = 1$, we present some of the proofs with different scaling in order to simplify the coefficients appearing in the solutions.

Proof. (i) Set $M = 2$. Define the positive semidefinite matrices

$$F_0 = \frac{2}{5} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}, \quad F_1 = (1),$$

and set

$$F(u, v, t) := \sum_{k=0}^1 \langle F_k, \overline{Z}_k^n(u, v, t) \rangle = \frac{2}{5}(uv - u - v + t + 1).$$

Since

$$\begin{aligned} F(u, v, t) = & -1 - \frac{3}{5} \left(1 - \frac{1}{2}(u + v) \right)^2 - \frac{7}{20}(v - u)^2 \\ & - u(1 - u) - v(1 - v) - \frac{1}{2}(u^2 + v^2 - 2t - 4), \end{aligned}$$

we have $F(u, v, t) \leq -1$ for all u, v, t with $0 \leq u, v \leq 1$ and $u^2 + v^2 - 2t - 4 \geq 0$. Furthermore, since

$$F(u, u, u^2) = 1 - \frac{3}{5}(1 - u)^2 - 2u(1 - u),$$

we have $F(u, u, u^2) \leq M - 1$ for all $0 \leq u \leq 1$. The tuple (M, F_0, F_1) thus defines a feasible solution for problem (2.5) with $R = 2$ and $r = 1$, which shows the bound is sharp.

(ii) We construct a feasible solution for problem (2.5) for $n \geq 2$ with $r = 2/\sqrt{3}$ and $R = (2/\sqrt{3} + 1)r = 4/3 + 2/\sqrt{3}$: By considering the positive semidefinite matrices

$$F_0 = \begin{pmatrix} 64/81 & -16/27 \\ -16/27 & 4/9 \end{pmatrix}, \quad F_1 = (9/8),$$

we obtain the function

$$F(u, v, t) = \sum_{k=0}^1 \langle F_k, \overline{Z}_k^n(u, v, t) \rangle = \frac{64}{81} + \frac{4}{9}uv - \frac{16}{27}(u + v) + \frac{9}{8}t.$$

Since the function satisfies the equation

$$\begin{aligned} F(u, v, t) = & -1 - 2(7/9 - 7/24(u + v))^2 - 113/288(v - u)^2 \\ & - 9/8u(4/3 - u) - 9/8v(4/3 - v) - 9/16(u^2 + v^2 - 2t - 16/3), \end{aligned}$$

we have $F(u, v, t) \leq -1$ for all u, v, t with $0 \leq u, v \leq 4/3$ and $u^2 + v^2 - 2t - 16/3 \geq 0$. Furthermore,

$$F(u, u, u^2) = 2 - 2(7/9 - 7/12u)^2 - 9/4u(4/3 - u),$$

so, for all $0 \leq u \leq 4/3$, we have $F(u, u, u^2) \leq 2$. The function $F(u, v, t)$ together with the parameter $M = 3$ gives a feasible solution for problem (2.5) with $r = 2/\sqrt{3}$ and $R = (2/\sqrt{3} + 1)r$.

(iii) We construct a feasible solution for problem (2.5) with $r = \sqrt{2n/(n+1)}$, $R = (\sqrt{2n/(n+1)} + 1)r = 2n/(n+1) + \sqrt{2n/(n+1)}$, and $n \geq 2$: By considering the positive semidefinite matrices

$$F_0 = \begin{pmatrix} 2n/(n+1) & -1 \\ -1 & (n+1)/(2n) \end{pmatrix}, \quad F_1 = \begin{pmatrix} (n+1)^2 \\ 4n \end{pmatrix},$$

we define the function

$$F(u, v, t) = \sum_{k=0}^1 \langle F_k^n, \overline{Z_k^n}(u, v, t) \rangle = \frac{2n}{n+1} - u - v + \frac{n+1}{2n}uv + \frac{(n+1)^2}{4n}t.$$

Since the equation

$$\begin{aligned} -F(u, v, t) - 1 &= \left(\sqrt{\frac{n(n-1)}{n+1}} + \frac{1}{4} \sqrt{\frac{n^2-1}{n}}(u+v) \right)^2 + \frac{n^2+4n+3}{16n}(v-u)^2 \\ &\quad + u \left(\frac{2n}{n+1} - u \right) \frac{(n+1)^2}{4n} + v \left(\frac{2n}{n+1} - v \right) \frac{(n+1)^2}{4n} \\ &\quad + \left(u^2 + v^2 - 2t - 4 \frac{2n}{n+1} \right) \frac{(n+1)^2}{8n} \end{aligned}$$

holds, we have $F(u, v, t) \leq -1$ for all u, v, t with $0 \leq u, v \leq \frac{2n}{n+1}$ and $u^2 + v^2 - 2t - 4 \frac{2n}{n+1} \geq 0$. Moreover,

$$n - F(u, u, u^2) = \left(\sqrt{\frac{n(n-1)}{n+1}} - \sqrt{\frac{n^2-1}{4n}}u \right)^2 + u \left(\frac{2n}{n+1} - u \right) \frac{(n+1)^2}{2n},$$

so $F(u, u, u^2) \leq n$ for all $0 \leq u \leq 2n/(n+1)$.

The function $F(u, v, t)$ together with the parameter $M = n + 1$ gives a feasible solution for problem (2.5) with $r = \sqrt{2n/(n+1)}$ and $R = (\sqrt{2n/(n+1)} + 1)r$.

(iv) The cases $n = 2, 3, 4$ are done separately through our rounding procedure; see `proofs.jl` in the arXiv version of this paper. For $n \geq 5$ we determine a feasible solution for (2.5) with $r = \sqrt{2}$ and $R = (\sqrt{2} + 1)r = 2 + \sqrt{2}$. For this we define the positive semidefinite matrices

$$F_0 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad F_1 = \begin{pmatrix} n & -\frac{1}{4}n \\ -\frac{1}{4}n & \frac{1}{16}n \end{pmatrix}, \quad F_2 = \left(\frac{1}{16}(n-1) \right)$$

and set

$$\begin{aligned} F(u, v, t) &= \sum_{k=0}^2 \langle F_k, \overline{Z_k^n}(u, v, t) \rangle \\ &= nt - \frac{1}{4}nt(u+v) + \frac{1}{16}nuvt + \frac{1}{16}nt^2 - \frac{1}{16}u^2v^2. \end{aligned}$$

To prove that $F(u, v, t) \leq -1$ for all $(u, v, t) \in \Omega$ we want to find sum-of-squares polynomials $q_1(u, v, t), \dots, q_5(u, v, t)$ such that $-1 - F(u, v, t)$ can be written as

$$\begin{aligned} &q_1(u, v, t) + u(2-u)q_2(u, v, t) + v(2-v)q_3(u, v, t) \\ &\quad + (t+uv)(uv-t)q_4(u, v, t) + (u^2+v^2-2t-8)q_5(u, v, t). \end{aligned}$$

Since $0 \leq u, v \leq 2$, the condition $-uv \leq t \leq uv$ in Ω holds if $-4 \leq t \leq 4$. Now we can show that $F(u, v, t) \leq -1$ for all $(u, v, t) \in \Omega$, since $-1 - F(u, v, t)$ is equal to

$$q_1(u, v, t) + u(2 - u)q_2(u, v, t) + v(2 - v)q_3(u, v, t) + (t + 4)(4 - t)q_4(u, v, t) + (u^2 + v^2 - 2t - 8)q_5(u, v, t),$$

where the sum of squares polynomials q_1, \dots, q_4 are displayed in Appendix A.

Next, we show $F(u, u, u^2) \leq 2n - 1$ for all $0 \leq u \leq 2$. For this we define

$$f(u) = F(u, u, u^2) - 2n + 1 = \frac{1}{16} (2n - 1) u^4 - \frac{1}{2} nu^3 + nu^2 - 2n + 1.$$

Note that $f(2) = 0$. It is then sufficient to prove that the polynomial

$$g(u) = f(u)/(u - 2) = \frac{2n - 1}{16} u^3 + \frac{-2n - 1}{8} u^2 + \frac{2n - 1}{4} u + \frac{2n - 1}{2}$$

is positive for every u in $[0, 2]$. Its discriminant

$$\frac{1}{16} (-11n^4 + 21n^3 - 47/4n^2 + 5/2n - 1/4)$$

is negative, since $-11n^4 + 21n^3$ and $-\frac{47}{4}n^2 + \frac{5}{2}n$ are both negative. Hence $g(u)$ has only one real root. This root must be negative, because the leading term of g is positive and $g(0) = n - 1/2 > 0$. Thus g is positive for every $u \geq 0$. In particular $f(u) \leq 0$ for all u in $[0, 2]$, for every $n \geq 5$.

The function $F(u, v, t)$ together with the parameter $M = 2n$ gives a feasible solution for problem (2.5) with $r = \sqrt{2}$ and $R = (\sqrt{2} + 1)r$.

(v), (vi) The exact optimal solution, as well as the verification of the solution, can be obtained by running our program as described in `proofs.jl`. Here the exact solution for (v) is over $\mathbb{Q}[\sqrt{5}]$ and for (vi) over \mathbb{Q} . \square

Appendix A. Sum of squares polynomials for Theorem 4.5.

$$\begin{aligned} q_1(u, v, t) = & \frac{1}{14} \left(\frac{35}{144} t^2 - \frac{151}{432} tu - \frac{125}{432} tv + \frac{9}{4} t \right)^2 \\ & + \frac{25}{238} \left(\frac{25}{144} t^2 + \frac{31}{432} tu + \frac{119}{432} tv \right)^2 + \frac{145}{238} \left(\frac{1}{16} t^2 + \frac{1}{8} tu \right)^2 \\ & + \frac{1}{21} \left(\frac{5}{16} tu + \frac{25}{16} tv + \frac{7}{4} uv - \frac{15}{4} t - \frac{7}{2} u \right)^2 \\ & + \frac{n - 5}{2} \left(\frac{5}{192} (3t^2 - 2tu + 8tv) \right)^2 + \frac{1}{48} (3t^2 - 8tu - 10tv + 48t)^2 \\ & + \frac{21}{64} (t^2 + 2tu)^2 + \frac{1}{4} (tu + 5tv + 8uv - 12t - 16u)^2, \end{aligned}$$

$$\begin{aligned}
q_2(u, v, t) &= \frac{1}{7982} \left(\frac{65}{8}t + \frac{307}{12}u \right)^2 + \frac{2095}{117888}t^2 + \frac{1}{858} \left(\frac{63}{8}u - \frac{65}{16}v \right)^2 \\
&\quad + \frac{1}{22} \left(\frac{5}{24}u - \frac{25}{16}v + \frac{11}{2} \right)^2 + (n-5) \left(\frac{1}{3110} \left(\frac{15}{8}t + \frac{311}{36}u \right)^2 \right. \\
&\quad \left. + \frac{1461}{622} \left(\frac{1}{24}t \right)^2 + \frac{1}{10} \left(\frac{19}{72}u - \frac{5}{16}v \right)^2 + \frac{3}{2} \left(\frac{1}{72}u - \frac{5}{48}v + \frac{1}{2} \right)^2 \right), \\
q_3(u, v, t) &= \frac{7}{51} \left(\frac{15}{16}t + \frac{17}{16}u \right)^2 + \frac{1}{2} \left(\frac{1}{8}t + \frac{5}{16}u - \frac{5}{12}v \right)^2 \\
&\quad + \frac{5}{2} \left(\frac{1}{16}u + \frac{1}{12}v - \frac{1}{2} \right)^2 + \frac{n-5}{430} \left(30 \left(\frac{7}{16}t + \frac{43}{48}u \right)^2 \right. \\
&\quad \left. + 43 \left(\frac{1}{8}t + \frac{5}{16}u - \frac{5}{12}v \right)^2 + \frac{1837}{384}t^2 \right. \\
&\quad \left. + 215 \left(\frac{1}{16}u + \frac{1}{12}v - \frac{1}{2} \right)^2 \right) + \frac{23}{51} \left(\frac{1}{16}t \right)^2, \\
q_4(u, v, t) &= \frac{n-5}{2} \left(\left(\frac{1}{16}t \right)^2 + \left(\frac{13}{96}u + \frac{11}{96}v - \frac{1}{2} \right)^2 + 23 \left(\frac{1}{96}(u-v) \right)^2 \right) \\
&\quad + \frac{5}{512}t^2 + \left(\frac{53}{288}u + \frac{55}{288}v - \frac{3}{4} \right)^2 + 215 \left(\frac{1}{288}(u-v) \right)^2, \\
q_5(u, v, t) &= n \left(\left(\frac{1}{16}t - \frac{1}{24}u - \frac{1}{12}v + \frac{1}{2} \right)^2 + \left(\frac{1}{16}t + \frac{1}{8}u \right)^2 + \frac{1}{72}(u-v)^2 \right).
\end{aligned}$$

Acknowledgments. We are very thankful to Christine Bachoc, Henry Cohn, Mathieu Dutour Sikirić, Frank Vallentin, and Guillermo Fernández Castro for helpful discussions and suggestions. We also thank the referees whose suggestions helped to improve the paper.

REFERENCES

- [1] C. BACHOC, G. NEBE, F. M. DE OLIVEIRA FILHO, AND F. VALLENTIN, *Lower bounds for measurable chromatic numbers*, *Geom. Funct. Anal.*, 19 (2009), pp. 645–661, <https://doi.org/10.1007/s00039-009-0013-7>.
- [2] C. BACHOC AND F. VALLENTIN, *New upper bounds for kissing numbers from semidefinite programming*, *J. Amer. Math. Soc.*, 21 (2008), pp. 909–924, <https://doi.org/10.1090/S0894-0347-07-00589-9>.
- [3] C. BACHOC AND F. VALLENTIN, *Optimality and uniqueness of the (4, 10, 1/6) spherical code*, *J. Combin. Theory Ser. A*, 116 (2009), pp. 195–204, <https://doi.org/10.1016/j.jcta.2008.05.001>.
- [4] C. BACHOC AND F. VALLENTIN, *Semidefinite programming, multivariate orthogonal polynomials, and codes in spherical caps*, *European J. Combin.*, 30 (2009), pp. 625–637, <https://doi.org/10.1016/j.ejc.2008.07.017>.
- [5] E. BANNAI, A. MUNEMASA, AND B. VENKOV, *The nonexistence of certain tight spherical designs*, *Algebra i Analiz*, 16 (2004), pp. 1–23, <https://doi.org/10.1090/S1061-0022-05-00868-X>.
- [6] E. BANNAI AND N. J. A. SLOANE, *Uniqueness of certain spherical codes*, *Canad. J. Math.*, 33 (1981), pp. 437–449, <https://doi.org/10.4153/CJM-1981-038-7>.
- [7] J. BEZANSON, A. EDELMAN, S. KARPINSKI, AND V. B. SHAH, *Julia: A fresh approach to numerical computing*, *SIAM Rev.*, 59 (2017), pp. 65–98, <https://doi.org/10.1137/141000671>.
- [8] D. CIFUENTES, T. KAHLE, AND P. PARRILO, *Sums of squares in Macaulay2*, *J. Softw. Algebra Geom.*, 10 (2020), pp. 17–24, <https://doi.org/10.2140/jsag.2020.10.17>.

- [9] H. COHN AND J. WOO, *Three-point bounds for energy minimization*, J. Amer. Math. Soc., 25 (2012), pp. 929–958, <https://doi.org/10.1090/S0894-0347-2012-00737-1>.
- [10] J. H. CONWAY AND N. J. SLOANE, *Sphere Packings, Lattices and Groups*, 3rd ed., Grundlehren 290 math. wiss., Springer-Verlag, New York, 1999, <https://doi.org/10.1007/978-1-4757-6568-7>.
- [11] L. W. DANZER, *Finite point-sets on S^2 with minimum distance as large as possible*, Discrete Math., 60 (1986), pp. 3–66, [https://doi.org/10.1016/0012-365X\(86\)90002-6](https://doi.org/10.1016/0012-365X(86)90002-6).
- [12] D. DE LAAT, *Moment methods in energy minimization: New bounds for Riesz minimal energy problems*, Trans. Amer. Math. Soc., (2019), <https://doi.org/10.1090/tran/7976>.
- [13] D. DE LAAT, F. CALUZA MACHADO, F. M. DE OLIVEIRA FILHO, AND F. VALLENTIN, *k-Point Semidefinite Programming Bounds for Equiangular Lines*, arXiv e-print, arXiv:1812.06045 [math.OC], 2018, <https://arxiv.org/abs/1812.06045>.
- [14] D. DE LAAT AND F. VALLENTIN, *A semidefinite programming hierarchy for packing problems in discrete geometry*, Math. Program., 151 (2015), pp. 529–553, <https://doi.org/10.1007/s10107-014-0843-4>.
- [15] P. DELSARTE, J.-M. GOETHALS, AND J. J. SEIDEL, *Spherical codes and designs*, Geom. Dedicata, 6 (1977), pp. 363–388.
- [16] M. DOSTERT, C. GUZMÁN, F. M. DE OLIVEIRA FILHO, AND F. VALLENTIN, *New upper bounds for the density of translate packings of three-dimensional convex bodies with tetrahedral symmetry*, Discrete Comput. Geom., 58 (2017), pp. 449–481, <https://doi.org/10.1007/s00454-017-9882-y>.
- [17] C. FIEKER, W. HART, T. HOFMANN, AND F. JOHANSSON, *Nemo/Hecke: Computer algebra and number theory packages for the Julia programming language*, in ISSAC'17—Proceedings of the 2017 ACM International Symposium on Symbolic and Algebraic Computation, ACM, New York, 2017, pp. 157–164.
- [18] K. FUJISAWA, M. FUKUDA, K. KOBAYASHI, M. KOJIMA, K. NAKATA, M. NAKATA, AND M. YAMASHITA, *SDPA (Semidefinite Programming Algorithm) User's Manual – Version 7.0.5*, tech. report, Research Report B-448, Dept. of Mathematical and Computing Sciences, Tokyo Institute of Technology, Tokyo, 2008, <http://sdpa.sourceforge.net>.
- [19] K. GATERMANN AND P. A. PARRILO, *Symmetry groups, semidefinite programs, and sums of squares*, J. Pure Appl. Algebra, 192 (2004), pp. 95–128, <https://doi.org/10.1016/j.jpaa.2003.12.011>.
- [20] D. HENRION, S. NALDI, AND M. SAFEY EL DIN, *Exact algorithms for linear matrix inequalities*, SIAM J. Optim., 26 (2016), pp. 2512–2539, <https://doi.org/10.1137/15M1036543>.
- [21] D. HENRION, S. NALDI, AND M. SAFEY EL DIN, *Exact algorithms for semidefinite programs with degenerate feasible set*, in ISSAC'18—Proceedings of the 2018 ACM International Symposium on Symbolic and Algebraic Computation, ACM, New York, 2018, pp. 191–198, <https://doi.org/10.1145/3208976.3209022>.
- [22] G. A. KABATIANSKY AND V. I. LEVENSHTAIN, *Bounds for packings on the sphere and in space*, Problemy Peredachi Informatsii, 14 (1978), pp. 3–25.
- [23] E. KALTOFEN, B. LI, Z. YANG, AND L. ZHI, *Exact certification of global optimality of approximate factorizations via rationalizing sums-of-squares with floating point scalars*, in Proceedings of ISSAC 2008, ACM, New York, 2008, pp. 155–163, <https://doi.org/10.1145/1390768.1390792>.
- [24] E. L. KALTOFEN, B. LI, Z. YANG, AND L. ZHI, *Exact certification in global polynomial optimization via sums-of-squares of rational functions with rational coefficients*, J. Symbolic Comput., 47 (2012), pp. 1–15, <https://doi.org/10.1016/j.jsc.2011.08.002>.
- [25] R. KANNAN AND A. BACHEM, *Polynomial algorithms for computing the Smith and Hermite normal forms of an integer matrix*, SIAM J. Comput., 8 (1979), pp. 499–507, <https://doi.org/10.1137/0208040>.
- [26] MARTIN KNESER, *Klassenzahlen definitiver quadratischer Formen*, Archiv Math., 8 (1957), pp. 241–250, <https://doi.org/10.1007/BF01898782>.
- [27] S. LAPLAGNE, *Facial reduction for exact polynomial sum of squares decomposition*, Math. Comp., 89 (2020), pp. 859–877, <https://doi.org/10.1090/mcom/3476>.
- [28] A. K. LENSTRA, H. W. LENSTRA, JR., AND L. LOVÁSZ, *Factoring polynomials with rational coefficients*, Math. Ann., 261 (1982), pp. 515–534, <https://doi.org/10.1007/BF01457454>.
- [29] V. I. LEVENSHTAIN, *Boundaries for packings in n-dimensional Euclidean space*, Dokl. Akad. Nauk SSSR, 245 (1979), pp. 1299–1303.
- [30] F. C. MACHADO AND F. M. DE OLIVEIRA FILHO, *Improving the semidefinite programming bound for the kissing number by exploiting polynomial symmetry*, Exp. Math., 27 (2018), pp. 362–369, <https://doi.org/10.1080/10586458.2017.1286273>.

- [31] V. MAGRON AND M. S. E. DIN, *On Exact Reznick, Hilbert-Artin and Putinar's Representations*, preprint: arXiv:1811.10062 [cs.CS], 2018, <https://arxiv.org/abs/1811.10062>.
- [32] R. M'HALLAH, A. ALKANDARI, AND N. MLADENOVIC, *Packing unit spheres into the smallest sphere using VNS and NLP*, *Comput. Oper. Res.*, 40 (2013), pp. 603–615, <https://doi.org/10.1016/j.cor.2012.08.019>.
- [33] S. NALDI, *Solving rank-constrained semidefinite programs in exact arithmetic*, *J. Symbolic Comput.*, 85 (2018), pp. 206–223, <https://doi.org/10.1016/j.jsc.2017.07.009>.
- [34] J. NIE, K. RANESTAD, AND B. STURMFELS, *The algebraic degree of semidefinite programming*, *Math. Program.*, 122 (2010), pp. 379–405, <https://doi.org/10.1007/s10107-008-0253-6>.
- [35] A. M. ODLYZKO AND N. J. A. SLOANE, *New bounds on the number of unit spheres that can touch a unit sphere in n dimensions*, *J. Combin. Theory Ser. A*, 26 (1979), pp. 210–214, [https://doi.org/10.1016/0097-3165\(79\)90074-8](https://doi.org/10.1016/0097-3165(79)90074-8).
- [36] F. PERMENTER AND P. PARRILO, *Partial facial reduction: Simplified, equivalent SDPs via approximations of the PSD cone*, *Math. Program.*, 171 (2018), pp. 1–54, <https://doi.org/10.1007/s10107-017-1169-9>.
- [37] H. PEYRL AND P. A. PARRILO, *Computing sum of squares decompositions with rational coefficients*, *Theoret. Comput. Sci.*, 409 (2008), pp. 269–281, <https://doi.org/10.1016/j.tcs.2008.09.025>.
- [38] M. PUTINAR, *Positive polynomials on compact semi-algebraic sets*, *Indiana Univ. Math. J.*, 42 (1993), pp. 969–984, <https://doi.org/10.1512/iumj.1993.42.42045>.
- [39] I. J. SCHOENBERG, *Positive definite functions on spheres*, *Duke Math. J.*, 9 (1942), pp. 96–108.
- [40] K. SCHÜTTE AND B. L. VAN DER WAERDEN, *Auf welcher Kugel haben 5, 6, 7, 8 oder 9 Punkte mit Mindestabstand Eins Platz?*, *Math. Ann.*, 123 (1951), pp. 96–124, <https://doi.org/10.1007/BF02054944>.
- [41] N. J. A. SLOANE, *Tables of sphere packings and spherical codes*, *IEEE Trans. Inform. Theory*, 27 (1981), pp. 327–338, <https://doi.org/10.1109/TIT.1981.1056351>.