



Four-loop collinear anomalous dimensions in QCD and $\mathcal{N} = 4$ super Yang-Mills

Bakul Agarwal^{a,*}, Andreas von Manteuffel^{a,*}, Erik Panzer^{b,*}, Robert M. Schabinger^{a,*}

^a Department of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA

^b Mathematical Institute, University of Oxford, OX2 6GG, Oxford, UK

ARTICLE INFO

Article history:

Received 20 April 2021

Received in revised form 24 June 2021

Accepted 2 July 2021

Available online 12 July 2021

Editor: A. Volovich

ABSTRACT

We calculate the collinear anomalous dimensions in massless four-loop QCD and $\mathcal{N} = 4$ supersymmetric Yang-Mills theory from the infrared poles of vertex form factors. We give very precise numerical approximations and a conjecture for the complete analytic results in both models we consider.

© 2021 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

Over the last several years, significant attention has been given to the calculation of cusp and collinear anomalous dimensions in massless perturbation theory. The light-like *cusp anomalous dimensions* [1] enter the leading infrared poles of massless scattering amplitudes and have recently been calculated to four-loop order both in Quantum Chromodynamics (QCD) and $\mathcal{N} = 4$ supersymmetric Yang-Mills theory ($\mathcal{N} = 4$ SYM) [2–17]. The *collinear anomalous dimensions* enter the subleading infrared poles and can be extracted from the $1/\epsilon$ poles of vertex *form factors* [18–28]. Partial results are available at four-loop order both in QCD [3,6,7,11,8,29,30,17] and $\mathcal{N} = 4$ supersymmetric Yang-Mills theory ($\mathcal{N} = 4$ SYM) [31–33]. In this article, we consider the analytically unknown four-loop contributions to the collinear anomalous dimensions.

In QCD, the basic quark and gluon form factors are the normalized amplitudes for, respectively, a virtual photon decaying into a pair of massless quarks, $\gamma^*(q) \rightarrow q(p_1)\bar{q}(p_2)$, and a Higgs boson decaying into two gluons in the limit of infinite top quark mass, $h(q) \rightarrow g(p_1)g(p_2)$, whereas in $\mathcal{N} = 4$ SYM, the Sudakov form factor is the normalized amplitude

$$\bar{\mathcal{F}}^{\mathcal{N}=4} = \frac{1}{N} \int d^4x e^{-iq \cdot x} \langle \phi_{12}^a(p_1) \phi_{12}^b(p_2) | [\phi_{34}^c \phi_{34}^c](x) | 0 \rangle, \quad (1)$$

where field superscripts denote adjoint $SU(N_c)$ color indices, field subscripts denote $SU(4)_R$ indices, and the constant N is chosen such that $\bar{\mathcal{F}}^{\mathcal{N}=4}$ is one at leading order.

In QCD, the perturbative expansion of the bare form factors is

$$\bar{\mathcal{F}}_{\text{bare}}^r = 1 + \sum_{L=1}^{\infty} \left(\frac{\alpha_s^{\text{bare}}}{4\pi} \right)^L \left(\frac{4\pi\mu_\epsilon^2}{-q^2 e^{\gamma_E}} \right)^{L\epsilon} \bar{\mathcal{F}}_L^r(\epsilon), \quad (2)$$

where $r = q$ or g for quarks or gluons, α_s^{bare} is the bare coupling, $q^2 = (p_1 + p_2)^2$ is the virtuality, μ_ϵ is the 't Hooft scale, $\epsilon = (4 - d)/2$ is the parameter of dimensional regularization, and γ_E is Euler's constant. In $\mathcal{N} = 4$ SYM, the expansion is

$$\bar{\mathcal{F}}^{\mathcal{N}=4} = 1 + \sum_{L=1}^{\infty} \lambda^L \left(\frac{\mu_\epsilon^2}{-q^2} \right)^{L\epsilon} \bar{\mathcal{F}}_L^{\mathcal{N}=4}(\epsilon), \quad (3)$$

in terms of the modified bare 't Hooft coupling

$$\lambda = \frac{N_c g_{\mathcal{N}=4}^2}{16\pi^2} (4\pi e^{-\gamma_E})^\epsilon, \quad (4)$$

and $g_{\mathcal{N}=4}$ is the bare coupling of the $\mathcal{N} = 4$ SYM model.

The collinear anomalous dimensions of QCD receive the four-loop contributions

$$\gamma_4^r = G_4^r[0] - \beta_0 G_3^r[1] - \beta_1 G_2^r[1] - \beta_2 G_1^r[1] + \beta_0^2 G_2^r[2] + 2\beta_0 \beta_1 G_1^r[2] - \beta_0^3 G_1^r[3] + 8\beta_3 \delta_{gr}, \quad (5)$$

where we follow [25], see also [34]. In Eq. (5), $G_L^r[k]$ denotes the ϵ^k coefficient of the *resummation function* $G_L^r(\epsilon)$ as defined in Eqs.

* Corresponding authors.

E-mail addresses: agarwalb@msu.edu (B. Agarwal), vmante@msu.edu (A. von Manteuffel), erik.panzer@maths.ox.ac.uk (E. Panzer), schabing@msu.edu (R.M. Schabinger).

(2.14)–(2.17) of [25], and β_{L-1} denotes the massless QCD beta function coefficient of order L , see e.g. [35,36] for explicit results. In $\mathcal{N} = 4$ SYM, the absence of a running coupling implies

$$\gamma_4^{\mathcal{N}=4} = \gamma_4^{\mathcal{N}=4}[0]. \quad (6)$$

The $1/\epsilon$ poles of the form factors allow for a determination of the resummation functions and thus the collinear anomalous dimensions.

The remainder of this article is organized as follows. In Section 2, we describe our computational methods based on integration by parts reductions and direct integration of Feynman parametric representations; we also give results for some master integrals. In Section 3, we present analytic results for the $1/\epsilon$ poles of the four-loop form factors $\bar{\mathcal{F}}_4^q(\epsilon)$, $\bar{\mathcal{F}}_4^g(\epsilon)$, and $\bar{\mathcal{F}}_4^{\mathcal{N}=4}(\epsilon)$, and the corresponding collinear anomalous dimensions γ_4^q , γ_4^g , and $\gamma_4^{\mathcal{N}=4}$. Our analytic results are expressed in terms of zeta values and a single leading-order-in- ϵ coefficient of a finite master integral, which could not be straightforwardly handled by the HyperInt program [37]. In Section 4, we employ a very precise numerical approximation of this integral coefficient to provide complete numerical results for all form factors and collinear anomalous dimensions. In Section 5, we perform a PSLQ analysis [38] to lift our precise numerical data to conjectured analytic results and describe a number of plausibility arguments which support our conjecture. Finally, in Section 6, we conclude.

2. Computational methods

Our analytic calculation of the form factors follows that of [17], employing a primary integration by parts reduction [39–46,547–49] and a subsequent rotation [50–52] to a judiciously-chosen basis of finite master integrals [53–56], computed with a private implementation, Finred. We make heavy use of HyperInt for the analytic evaluation of the master integrals. However, for the two topologies shown in Fig. 1 the generic Feynman parametric representation that we use is not *linearly reducible* [57,58], that is, it cannot be directly integrated with the algorithm [37]. We have been able to find linearly reducible integrands only for the *leading-order* ϵ -expansion coefficients of *specific* integrals in these topologies as will be explained in the following. However, we do not know whether generic integrals in these topologies can be rendered linearly reducible to all orders by changing variables.

For the topology on the left-hand side of Fig. 1, we found a basis of two finite integrals in $d = 6 - 2\epsilon$ dimensions, which appear for the first time at the level of the $1/\epsilon$ poles and which each have 15 as the sum of their propagator exponents. Due to this choice, the exponent of the Symanzik polynomial $\mathcal{U}$ vanishes at zeroth order in the ϵ expansion of these integrals – the *only* order which we need to obtain results for γ_4^q , γ_4^g , and $\gamma_4^{\mathcal{N}=4}$. The remaining polynomial $\mathcal{F}$, it turns out, is by itself linearly reducible for this topology, therefore allowing for a straightforward application of HyperInt.

For the topology on the right-hand side of Fig. 1, the situation is more complicated, despite the fact that we find a change of variables which renders the $\mathcal{F}$ polynomial linearly reducible in this case as well. First, we count four master integrals for the topology in d dimensions, but we were able to choose a basis such that only three of them contribute to the $1/\epsilon$ poles of the form factors. Unfortunately, we did not find a suitable basis of integrals for the $1/\epsilon$ pole such that all of them are independent of the $\mathcal{U}$ polynomial, e.g. by choosing 15 for the sum of the propagator exponents for all integrals in $d = 6 - 2\epsilon$ dimensions.

Instead, the best-case scenario seems to allow for a straightforward treatment of two out of three finite integrals only; the remaining finite integral which contributes to the $1/\epsilon$ poles can be




Fig. 1. Top-level topologies whose Feynman parametric representation is not linearly reducible and therefore not directly accessible to HyperInt for generic integrals.

chosen to have 13, the minimal number, as the sum of its propagator exponents. Unfortunately, an additional complication arises due to the fact that making such a choice of finite integrals requires the computation of some finite integrals in subtopologies to higher orders in the ϵ expansion than would have been necessary in the finite integral basis of [17], i.e. one constructed to be compatible with a basis of uniform weight as suggested in [55].

Our auxiliary results include, for example, the $\mathcal{O}(\epsilon^2)$ term of

$$\begin{aligned} & \text{Diagram (Left)} \\ &= -\frac{53}{6}\zeta_3^2 - \frac{1579}{630}\zeta_2^3 + \frac{535}{36}\zeta_5 + \frac{5}{6}\zeta_3\zeta_2 \\ & - \frac{17}{12}\zeta_2^2 + \frac{121}{18}\zeta_3 + \frac{8}{9}\zeta_2 + \epsilon \left(-\frac{1033}{6}\zeta_7 - \frac{53}{3}\zeta_5\zeta_2 + \frac{21}{5}\zeta_3\zeta_2^2 \right. \\ & + \frac{139}{18}\zeta_3^2 + \frac{1609}{126}\zeta_2^3 + \frac{6235}{108}\zeta_5 - \frac{145}{18}\zeta_3\zeta_2 - \frac{49}{12}\zeta_2^2 + \frac{1478}{27}\zeta_3 \\ & + \frac{565}{54}\zeta_2 \left. \right) + \epsilon^2 \left(\frac{10403}{30}\zeta_{5,3} + \frac{2887}{6}\zeta_5\zeta_3 + \frac{307}{2}\zeta_3^2\zeta_2 \right. \\ & - \frac{247041}{1000}\zeta_2^4 + \frac{78607}{144}\zeta_7 + \frac{110}{9}\zeta_5\zeta_2 + \frac{2461}{180}\zeta_3\zeta_2^2 - \frac{18467}{216}\zeta_3^2 \\ & + \frac{30287}{2268}\zeta_2^3 + \frac{79117}{324}\zeta_5 - \frac{30455}{108}\zeta_3\zeta_2 - \frac{1495}{216}\zeta_2^2 + \frac{112325}{324}\zeta_3 \\ & + \frac{6854}{81}\zeta_2 \left. \right) + \mathcal{O}(\epsilon^3), \end{aligned} \quad (7)$$

where a dot indicates a squared propagator, the integral normalization follows the conventions of [54], and


$$\zeta_{5,3} = \sum_{m=1}^{\infty} \frac{1}{m^5} \sum_{n=1}^{m-1} \frac{1}{n^3} \approx 0.0377076729848\dots \quad (8)$$

To obtain this and other results, we ran HyperInt in a highly parallelized setup, accumulating several CPU years in total. For the most complicated topology, we determined two integrals with a propagator exponent sum of 15 analytically,

$$\begin{aligned} & \text{Diagram (Right)} \\ &= -\frac{4221}{16}\zeta_7 + \frac{159}{2}\zeta_5\zeta_2 + \frac{25}{2}\zeta_3\zeta_2^2 \\ & - 14\zeta_3^2 - \frac{631}{70}\zeta_2^3 + \frac{535}{2}\zeta_5 - 69\zeta_3\zeta_2 + \mathcal{O}(\epsilon), \end{aligned} \quad (9)$$

$$\begin{aligned} & \text{Diagram (Right)} \\ &= \frac{252}{5}\zeta_{5,3} + 195\zeta_5\zeta_3 - 18\zeta_3^2\zeta_2 \\ & - \frac{202807}{10500}\zeta_2^4 - \frac{959}{8}\zeta_7 + 50\zeta_5\zeta_2 - \frac{51}{5}\zeta_3\zeta_2^2 + \mathcal{O}(\epsilon), \end{aligned} \quad (10)$$

leaving only a single, finite integral, that we choose as



$$\equiv \mathcal{H} + \mathcal{O}(\epsilon). \quad (11)$$

We were able to evaluate $\mathcal{H}$ to high precision numerically by running the `pySecDec` program [59] for several months in a distributed manner on high-performance GPUs,

$$\mathcal{H} \approx -0.7015802723647 \pm 6.98 \cdot 10^{-11}, \quad (12)$$

where, with some foresight, we have appropriately rounded our numerical result and the provided estimate of its statistical uncertainty. While we expect our numerical results to suffice for phenomenological applications, we will see later that the precision of our results even allows us to put forth a plausible conjecture for the analytical form of $\mathcal{H}$ and, therefore, of the four-loop collinear anomalous dimensions.

In contrast to the case of QCD, the reduced integrand for the Sudakov form factor of the $\mathcal{N} = 4$ SYM model is known as a simple linear combination of (conjecturally) uniform weight Feynman integrals defined in [32] and evaluated through to weight six in [16]:

$$\begin{aligned} \bar{\mathcal{F}}_4^{\mathcal{N}=4}(\epsilon) = & 16I_{p,1}^{(1)} + 4I_{p,2}^{(2)} - 4I_{p,3}^{(3)} + 4I_{p,4}^{(4)} + I_{p,5}^{(5)} + 4I_{p,6}^{(6)} \\ & + 8I_{p,7}^{(7)} + 4I_{p,8}^{(9)} - 4I_{p,9}^{(10)} + 2I_{p,10}^{(12)} + 2I_{p,11}^{(12)} + 4I_{p,12}^{(13)} \\ & + 4I_{p,13}^{(14)} - 4I_{p,14}^{(17)} + 4I_{p,15}^{(17)} - 4I_{p,16}^{(19)} + 2I_{p,17}^{(19)} + 2I_{p,18}^{(21)} \\ & + I_{p,19}^{(25)} + 4I_{p,20}^{(30)} + 4I_{p,21}^{(13)} + 8I_{p,22}^{(14)} - 4I_{p,23}^{(14)} - 2I_{p,24}^{(14)} \\ & + 8I_{p,25}^{(17)} - 2I_{p,26}^{(17)} - 4I_{p,27}^{(17)} - 4I_{p,28}^{(19)} - 2I_{p,29}^{(19)} - 2I_{p,30}^{(19)} \\ & + 2I_{p,31}^{(19)} - I_{p,32}^{(30)} \\ & + \frac{12}{N_c^2} \left[2I_1^{(21)} + 2I_2^{(22)} + 2I_3^{(23)} - 4I_4^{(24)} + I_5^{(25)} - I_6^{(26)} \right. \\ & - I_7^{(26)} + 8I_8^{(27)} + 4I_9^{(28)} + 16I_{10}^{(29)} + 4I_{11}^{(30)} + 4I_{12}^{(27)} \\ & - 2I_{13}^{(28)} + 4I_{14}^{(29)} + 4I_{15}^{(29)} + 4I_{16}^{(30)} + 4I_{17}^{(30)} + 4I_{18}^{(30)} \\ & \left. + 4I_{19}^{(22)} + 4I_{20}^{(22)} - 4I_{21}^{(24)} + I_{22}^{(24)} + 2I_{23}^{(28)} \right]. \quad (13) \end{aligned}$$

The integral $\mathcal{H}$ enters both $I_6^{(26)}$ and $I_7^{(26)}$ in the non-planar-color part of Eq. (13), allowing for an over-determination of the lower-weight power products of zeta values which enter $\mathcal{H}$. Let us emphasize that a rotation to a basis of finite integrals is still of paramount importance for our $\mathcal{N} = 4$ SYM calculation, but, due to the uniform weight property, it is convenient to organize the calculation of the $\mathcal{N} = 4$ Sudakov form factor in terms of the master integrals entering Eq. (13).

Note that, while the most complicated, non-linearly reducible integral topologies appeared in all four-loop form factors we calculated through to weight seven, the calculation of the QCD master integrals to sufficiently high orders in ϵ was harder overall, because it involved a significant number of computationally challenging, non-planar integral topologies that do not appear in the four-loop $\mathcal{N} = 4$ SYM Sudakov form factor.

3. Analytical results

In this section, we present results for the $1/\epsilon$ poles of the four-loop form factors we consider, $\bar{\mathcal{F}}_4^q(\epsilon)$, $\bar{\mathcal{F}}_4^g(\epsilon)$, and $\bar{\mathcal{F}}_4^{\mathcal{N}=4}(\epsilon)$, and the corresponding four-loop collinear anomalous dimensions, γ_4^q , γ_4^g , and $\gamma_4^{\mathcal{N}=4}$, as a function of $\mathcal{H}$. In fact, we write $\delta \equiv \mathcal{H} - \mathcal{H}_{\text{conj}}$ for the difference between $\mathcal{H}$ and our conjectured expression discussed in section 5, such that $\delta = 0$ if our conjecture is true. Further, we employ the $SU(N_c)$ color factors

$$\begin{aligned} C_F &= \frac{N_c^2 - 1}{2N_c}, \quad C_A = N_c, \\ N_A &= N_c^2 - 1, \quad N_F = N_c, \\ \frac{d_A^{abcd} d_F^{abcd}}{N_F} &= \frac{(N_c^2 - 1)(N_c^2 + 6)}{48}, \\ \frac{d_A^{abcd} d_A^{abcd}}{N_A} &= \frac{N_c^2(N_c^2 + 36)}{24} \end{aligned} \quad (14)$$

and set $T_F = 1/2$. The number of light quark flavors is denoted by N_f and their charge-weighted sum, normalized to the charge of the external quark q , by $N_{q\gamma} \equiv \sum_{q'} e_{q'}/e_q$.

For the $1/\epsilon$ poles of the four-loop form factors, we find

$$\begin{aligned} \bar{\mathcal{F}}_4^q(\epsilon) \Big|_{1/\epsilon} = & \frac{d_A^{abcd} d_F^{abcd}}{N_F} \left[-\frac{871}{2} \zeta_7 - 128 \zeta_5 \zeta_2 + \frac{92}{5} \zeta_3 \zeta_2^2 \right. \\ & + \frac{418}{3} \zeta_3^2 - \frac{3476}{315} \zeta_2^3 + \frac{230}{9} \zeta_5 + 224 \zeta_3 \zeta_2 - \frac{28}{15} \zeta_2^2 + \frac{976}{9} \zeta_3 \\ & + \frac{272}{3} \zeta_2 - 24 + 80\delta \Big] + C_A^3 C_F \left[\frac{45511}{48} \zeta_7 + \frac{1033}{30} \zeta_3 \zeta_2^2 \right. \\ & - \frac{206}{3} \zeta_5 \zeta_2 - \frac{52547}{36} \zeta_3^2 - \frac{44204}{135} \zeta_2^3 + \frac{77285}{108} \zeta_5 + \frac{1034}{9} \zeta_3 \zeta_2 \\ & + \frac{504629}{270} \zeta_2^2 + \frac{1195223}{54} \zeta_3 - \frac{6051515}{648} \zeta_2 - \frac{651546401}{23328} \\ & \left. - \frac{70}{3} \delta \right] + C_A^2 C_F^2 \left[-\frac{38323}{18} \zeta_7 + \frac{7921}{9} \zeta_5 \zeta_2 + \frac{35767}{135} \zeta_3 \zeta_2^2 \right. \\ & + \frac{703139}{81} \zeta_3^2 - \frac{10901}{63} \zeta_2^3 - \frac{4775068}{405} \zeta_5 - \frac{99463}{27} \zeta_3 \zeta_2 \\ & - \frac{24831149}{4860} \zeta_2^2 - \frac{376873993}{5832} \zeta_3 + \frac{347179283}{11664} \zeta_2 \\ & + \frac{29277646423}{419904} + 60\delta \Big] + C_A C_F^3 \left[6172 \zeta_7 - \frac{11548}{5} \zeta_5 \zeta_2 \right. \\ & - \frac{10508}{15} \zeta_3 \zeta_2^2 - \frac{302497}{27} \zeta_2^3 + \frac{672709}{945} \zeta_2^2 + \frac{1151047}{54} \zeta_5 \\ & + \frac{203729}{27} \zeta_3 \zeta_2 + \frac{5489227}{1620} \zeta_2^2 + \frac{100425413}{1944} \zeta_3 - \frac{38043757}{1296} \zeta_2 \\ & - \frac{6800926313}{139968} - 40\delta \Big] + C_F^4 \left[-\frac{14162}{21} \zeta_7 + \frac{5792}{5} \zeta_5 \zeta_2 \right. \\ & + \frac{6208}{9} \zeta_3 \zeta_2^2 + \frac{14060}{9} \zeta_2^3 - \frac{16786}{15} \zeta_2^2 - \frac{235816}{15} \zeta_5 \\ & - \frac{32966}{9} \zeta_3 \zeta_2 + \frac{479}{15} \zeta_2^2 - \frac{87481}{18} \zeta_3 + \frac{26425}{3} \zeta_2 + \frac{94257}{8} \Big] \\ & + \text{fermionic terms}, \quad (15) \end{aligned}$$

$$\begin{aligned} \bar{\mathcal{F}}_4^g(\epsilon) \Big|_{1/\epsilon} = & \frac{d_A^{abcd} d_A^{abcd}}{N_A} \left[-\frac{871}{2} \zeta_7 - 128 \zeta_5 \zeta_2 + \frac{92}{5} \zeta_3 \zeta_2^2 \right. \\ & + \frac{418}{3} \zeta_3^2 - \frac{4972}{315} \zeta_2^3 - \frac{340}{9} \zeta_5 + 292 \zeta_3 \zeta_2 + \frac{226}{15} \zeta_2^2 + \frac{3676}{9} \zeta_3 \\ & - 8 \zeta_2 - \frac{32}{3} + 80\delta \Big] + C_A^4 \left[\frac{4351243}{1008} \zeta_7 - \frac{15289}{45} \zeta_5 \zeta_2 \right. \\ & + \frac{77927}{270} \zeta_3 \zeta_2^2 - \frac{1109647}{324} \zeta_2^3 - \frac{1090111}{945} \zeta_2^2 - \frac{617401}{405} \zeta_5 \\ & + \frac{95669}{162} \zeta_3 \zeta_2 + \frac{1870843}{4860} \zeta_2^2 + \frac{7649891}{1458} \zeta_3 + \frac{196729}{72} \zeta_2 \\ & \left. - \frac{749534537}{104976} - \frac{10}{3} \delta \right] + \text{fermionic terms}, \quad (16) \end{aligned}$$

$$\begin{aligned} \bar{\mathcal{F}}_4^{\mathcal{N}=4}(\epsilon) \Big|_{1/\epsilon} = & \frac{541619}{126} \zeta_7 - \frac{15529}{45} \zeta_5 \zeta_2 + \frac{39067}{135} \zeta_3 \zeta_2^2 \\ & + \frac{1}{N_c^2} \left[-\frac{2613}{4} \zeta_7 - 192 \zeta_5 \zeta_2 + \frac{138}{5} \zeta_3 \zeta_2^2 + 120\delta \right]. \quad (17) \end{aligned}$$

For the quark and gluon form factors, we omitted non-singlet and singlet pole terms involving one, two or three closed fermion loops, they can be obtained from Ref. [17]. To the best of our

knowledge, we provide exact and unconditional results for the $1/\epsilon$ pole of the C_F^4 color structure, and therefore massless Quantum Electrodynamics, for the first time.

For the collinear anomalous dimensions of QCD and $\mathcal{N} = 4$ SYM we obtain

$$\begin{aligned} \gamma_4^q = & \frac{d_A^{abcd} d_F^{abcd}}{N_F} \left[3484\zeta_7 + 1024\zeta_5\zeta_2 - \frac{736}{5}\zeta_3\zeta_2^2 - \frac{3344}{3}\zeta_3^2 \right. \\ & + \frac{27808}{315}\zeta_2^3 - \frac{1840}{9}\zeta_5 - 1792\zeta_3\zeta_2 + \frac{224}{15}\zeta_2^2 - \frac{7808}{9}\zeta_3 \\ & - \frac{2176}{3}\zeta_2 + 192 - 640\delta \left. \right] + C_A^3 C_F \left[\frac{1648}{3}\zeta_5\zeta_2 - \frac{45511}{6}\zeta_7 \right. \\ & - \frac{4132}{15}\zeta_3\zeta_2^2 + \frac{5126}{9}\zeta_3^2 - \frac{77152}{315}\zeta_2^3 + \frac{175166}{27}\zeta_5 \\ & + \frac{15400}{9}\zeta_3\zeta_2 + \frac{186742}{135}\zeta_2^2 - \frac{1751224}{243}\zeta_3 + \frac{1062149}{729}\zeta_2 \\ & + \frac{7179083}{26244} + \frac{560}{3}\delta \left. \right] + C_A^2 C_F^2 \left[17220\zeta_7 - 4208\zeta_5\zeta_2 \right. \\ & - \frac{128}{5}\zeta_3\zeta_2^2 - \frac{14204}{3}\zeta_3^2 - \frac{43976}{35}\zeta_2^3 + \frac{10708}{9}\zeta_5 + \frac{4192}{9}\zeta_3\zeta_2 \\ & - \frac{48680}{27}\zeta_2^2 + \frac{259324}{27}\zeta_3 - \frac{93542}{27}\zeta_2 + \frac{29639}{18} - 480\delta \left. \right] \\ & + C_A C_F^3 \left[-21840\zeta_7 + 4128\zeta_5\zeta_2 + \frac{512}{5}\zeta_3\zeta_2^2 + 6440\zeta_3^2 \right. \\ & + \frac{634376}{315}\zeta_2^3 - 1952\zeta_5 - \frac{3976}{3}\zeta_3\zeta_2 + \frac{8668}{5}\zeta_2^2 - 6520\zeta_3 \\ & + 2334\zeta_2 - \frac{2085}{2} + 320\delta \left. \right] + C_F^4 \left[11760\zeta_7 - 768\zeta_5\zeta_2 \right. \\ & + \frac{256}{5}\zeta_3\zeta_2^2 - 2304\zeta_3^2 - \frac{33776}{35}\zeta_2^3 - 5040\zeta_5 - 240\zeta_3\zeta_2 \\ & - \frac{1368}{5}\zeta_2^2 + 4008\zeta_3 - 900\zeta_2 + \frac{4873}{12} \left. \right] + N_f\text{-terms}, \end{aligned} \quad (18)$$

$$\begin{aligned} \gamma_4^g = & \frac{d_A^{abcd} d_A^{abcd}}{N_A} \left[3484\zeta_7 + 1024\zeta_5\zeta_2 - \frac{736}{5}\zeta_3\zeta_2^2 - \frac{3344}{3}\zeta_3^2 \right. \\ & + \frac{39776}{315}\zeta_2^3 + \frac{2720}{9}\zeta_5 - 2336\zeta_3\zeta_2 - \frac{1808}{15}\zeta_2^2 - \frac{12512}{9}\zeta_3 \\ & + 64\zeta_2 + \frac{128}{9} - 640\delta \left. \right] + C_A^4 \left[-\frac{2671}{6}\zeta_7 - \frac{2212}{15}\zeta_3\zeta_2^2 \right. \\ & - \frac{896}{3}\zeta_5\zeta_2 - \frac{286}{9}\zeta_3^2 - \frac{674696}{945}\zeta_2^3 + \frac{19232}{27}\zeta_5 + \frac{1588}{3}\zeta_3\zeta_2 \\ & + \frac{249448}{135}\zeta_2^2 + \frac{36380}{243}\zeta_3 - \frac{1051411}{729}\zeta_2 + \frac{10672040}{6561} + \frac{80}{3}\delta \left. \right] \\ & + N_f\text{-terms}, \end{aligned} \quad (19)$$

$$\begin{aligned} \gamma_4^{\mathcal{N}=4} = & -300\zeta_7 - 256\zeta_5\zeta_2 - 384\zeta_4\zeta_3 \\ & + \frac{1}{N_c^2} \left[5226\zeta_7 + 1536\zeta_5\zeta_2 - 552\zeta_4\zeta_3 - 960\delta \right]. \end{aligned} \quad (20)$$

We used the Supplementary Material of [17] to extract the resummation functions in Eq. (5) from our results for the poles of the form factors. The omitted fermionic terms in γ_4^q and γ_4^g are due to non-singlet contributions involving one, two, or three closed fermion loops; they are available from Ref. [17]. To the best of our knowledge, our planar-color $\mathcal{N} = 4$ SYM calculation is the first to independently confirm the analytic analysis of [33].

Let us now compare the maximal transcendental weight (seven) part of the collinear anomalous dimensions in QCD to the $\mathcal{N} = 4$ SYM result. Here, the QCD contributions involving a closed fermion loop do not contribute since they are of weight six at most. Selecting the weight seven terms in the expression (19) for γ_4^g , replacing C_A and $d_A^{abcd} d_A^{abcd}/N_A$ by their $SU(N_c)$ values (14), and dividing by N_c^4 due to the different normalization used for the $\mathcal{N} = 4$ SYM reproduces the expression (20) for $\gamma_4^{\mathcal{N}=4}$. In a similar way, the

maximal weight terms in (18) for γ_4^q coincide with the result (20) for $\gamma_4^{\mathcal{N}=4}$, if one first changes from the fundamental to the adjoint color representation by substituting $C_F \rightarrow C_A$ and $d_F^{abcd} d_A^{abcd}/N_F \rightarrow d_A^{abcd} d_A^{abcd}/N_A$. We note that these relations are fulfilled without putting any constraint on the maximal weight part of δ , assuming its weight is not greater than seven. In summary, we observe that the principle of maximal transcendentality [60,61] is fulfilled for the four-loop collinear anomalous dimensions, at leading and also at subleading color.

4. Numerical results

In this section, we compile the explicit numerical results for the $1/\epsilon$ poles of the form factors and collinear anomalous dimensions, which we obtain using the numerical approximation for $\mathcal{H}$ (12). We provide twelve digits for the color structures where exact analytic results are available (i.e. from [17] for the N_f -dependent terms). For the form factors we have

$$\begin{aligned} \bar{\mathcal{F}}_4^q(\epsilon) \Big|_{1/\epsilon} \approx & \frac{d_A^{abcd} d_F^{abcd}}{N_F} \left[274.4588169341 \pm 5.6 \cdot 10^{-9} \right] \\ & + C_A^3 C_F \left[-13274.5995371593 \pm 1.6 \cdot 10^{-9} \right] \\ & + C_A^2 C_F^2 \left[19661.7351772000 \pm 4.2 \cdot 10^{-9} \right] \\ & + C_A C_F^3 \left[-1602.0556057677 \pm 2.8 \cdot 10^{-9} \right] \\ & + C_F^4 \left[-2212.79784915... \right] \\ & + N_f \frac{d_F^{abcd} d_F^{abcd}}{N_F} \left[53.1274437988... \right] \\ & + N_f C_A^2 C_F \left[10203.6391859... \right] \\ & + N_f C_A C_F^2 \left[-12551.5075480... \right] \\ & + N_f C_F^3 \left[2095.46596925... \right] \\ & + N_{q\gamma} C_A \frac{d_F^{abc} d_F^{abc}}{N_F} \left[-235.381562912... \right] \\ & + N_{q\gamma} C_F \frac{d_F^{abc} d_F^{abc}}{N_F} \left[-243.738662819... \right] \\ & + N_f^2 C_A C_F \left[-2304.68219272... \right] \\ & + N_f^2 C_F^2 \left[1604.85115658... \right] \\ & + N_{q\gamma} N_f \frac{d_F^{abc} d_F^{abc}}{N_F} \left[42.7966478022... \right] \\ & + N_f^3 C_F \left[158.065537245... \right], \end{aligned} \quad (21)$$

$$\begin{aligned} \bar{\mathcal{F}}_4^g(\epsilon) \Big|_{1/\epsilon} \approx & \frac{d_A^{abcd} d_A^{abcd}}{N_A} \left[579.5738867755 \pm 5.6 \cdot 10^{-9} \right] \\ & + C_A^4 \left[-1081.02280574667 \pm 2.3 \cdot 10^{-10} \right] \\ & + N_f \frac{d_A^{abcd} d_A^{abcd}}{N_A} \left[-604.701004352... \right] \\ & + N_f C_A^3 \left[-532.481107793... \right] \\ & + N_f C_A^2 C_F \left[-650.895247054... \right] \\ & + N_f C_A C_F^2 \left[14.9978706950... \right] \\ & + N_f C_F^3 \left[17.25 \right] \\ & + N_f^2 \frac{d_F^{abcd} d_F^{abcd}}{N_A} \left[95.1966169377... \right] \\ & + N_f^2 C_A^2 \left[1574.06171919... \right] \end{aligned}$$

$$\begin{aligned}
& + N_f^2 C_A C_F [282.052204632...] \\
& + N_f^2 C_F^2 [23.4647858335...] \\
& + N_f^3 C_A [-261.955705460...] \\
& + N_f^3 C_F [-12.7425646335...], \quad (22)
\end{aligned}$$

$$\begin{aligned}
\bar{\mathcal{F}}_4^{\mathcal{N}=4}(\epsilon) \Big|_{1/\epsilon} & \approx 4687.07846404... \\
& + \frac{1}{N_c^2} [-896.4243270825 \pm 8.4 \cdot 10^{-9}]. \quad (23)
\end{aligned}$$

Comparing Eq. (21) and Eq. (22) to, respectively, Eq. (3.27) of [29] and Eq. (11) of [30], we find that our results agree completely to within their given error estimates. We note that our Eqs. (21) and (22) significantly improve upon the QCD results of [29,30], as the numerical approximations provided therein are, depending on the color structure, accurate to, at best, six significant digits (e.g. $C_A^2 C_F^2$) and, at worst, one significant digit (e.g. $d_A^{abcd} d_A^{abcd} / N_A$). For the collinear anomalous dimensions we have

$$\begin{aligned}
\gamma_4^q & \approx \frac{d_A^{abcd} d_F^{abcd}}{N_F} [-2195.670535473 \pm 4.5 \cdot 10^{-8}] \\
& + C_A^3 C_F [-13.809312037 \pm 1.3 \cdot 10^{-8}] \\
& + C_A^2 C_F^2 [2438.569338812 \pm 3.3 \cdot 10^{-8}] \\
& + C_A C_F^3 [-1373.764650948 \pm 2.2 \cdot 10^{-8}] \\
& + C_F^4 [392.899478384...] \\
& + N_f \frac{d_A^{abcd} d_F^{abcd}}{N_F} [-425.019550390...] \\
& + N_f C_A^2 C_F [-274.147360589...] \\
& + N_f C_A C_F^2 [-912.844845636...] \\
& + N_f C_F^3 [151.933788877...] \\
& + N_f^2 C_A C_F [109.081415293...] \\
& + N_f^2 C_F^2 [-12.5342425083...] \\
& + N_f^3 C_F [4.88682798281...], \quad (24)
\end{aligned}$$

$$\begin{aligned}
\gamma_4^g & \approx \frac{d_A^{abcd} d_A^{abcd}}{N_A} [-2451.040712450 \pm 4.5 \cdot 10^{-8}] \\
& + C_A^4 [1557.4287417889 \pm 1.9 \cdot 10^{-9}] \\
& + N_f \frac{d_A^{abcd} d_F^{abcd}}{N_A} [-41.2080190194...] \\
& + N_f C_A^3 [-1033.98729659...] \\
& + N_f C_A^2 C_F [-57.9377499658...] \\
& + N_f C_A C_F^2 [-100.315097910...] \\
& + N_f C_F^3 [46] \\
& + N_f^2 \frac{d_A^{abcd} d_F^{abcd}}{N_A} [253.857645167...] \\
& + N_f^2 C_A^2 [70.7744401902...] \\
& + N_f^2 C_A C_F [73.9372035966...] \\
& + N_f^2 C_F^2 [-21.9767440643...] \\
& + N_f^3 C_A [0.405507202650...] \\
& + N_f^3 C_F [1.26748971193...], \quad (25)
\end{aligned}$$

$$\begin{aligned}
\gamma_4^{\mathcal{N}=4} & \approx -1238.74771725... \\
& + \frac{1}{N_c^2} [7171.394616660 \pm 6.7 \cdot 10^{-8}]. \quad (26)
\end{aligned}$$

The errors provided in the above equations are all statistical errors from the `pySecDec` evaluation of $\mathcal{H}$ given in Eq. (12). For the non-planar-color part of the $\mathcal{N}=4$ SYM collinear anomalous dimension, Eq. (26) is consistent with the result of [32] within their provided error estimate, but it is a vast improvement over Eq. (5.8) of [32] as the latter gives essentially only an order of magnitude estimate for this quantity. Besides our comparisons to the existing literature, we also carried out direct numerical cross-checks on all of the contributing four-loop form factor master integral expansion coefficients using either `pySecDec` or `FIESTA 4` [62].

5. Lifting numerical data to analytic expressions

In this section, we analyze the unknown analytic form of $\mathcal{H}$. First, assuming that the ϵ -expansion coefficients of $I_6^{(26)}$ and $I_7^{(26)}$ are multiple zeta values of uniform weight implies that $\mathcal{H}$ has the form

$$\begin{aligned}
\mathcal{H}_{\text{conj}} & = a\zeta_7 + b\zeta_5\zeta_2 + c\zeta_3\zeta_2^2 + 10\zeta_3^2 + \frac{223}{210}\zeta_2^3 \\
& - 25\zeta_5 - 6\zeta_3\zeta_2 + \frac{3}{10}\zeta_2^2 + 9\zeta_3 \quad (27)
\end{aligned}$$

for some rational numbers a , b and c . This constraint will be central to the following discussion.

Let us consider the unknown part of Eq. (27),

$$a\zeta_7 + b\zeta_5\zeta_2 + c\zeta_3\zeta_2^2 \approx 6.279370861144 \pm 7.0 \cdot 10^{-11}, \quad (28)$$

and fit the constants a , b , and c using the PSLQ algorithm. Our first task will be to assess the uncertainty of our high-precision run of `pySecDec` by comparing approximation (12) to a preliminary run of the program at a somewhat lower precision,

$$\mathcal{H} \approx -0.7015802399 \pm 2.75 \cdot 10^{-8}. \quad (29)$$

Subtracting the central value of (12) from the central value of (29) and dividing by the uncertainty of (29), we find a ratio of 1.2. This indicates that the initial estimate produced by `pySecDec` was a bit too large but that the uncertainty estimate produced by the program seems trustworthy. We will proceed under the assumption that the given statistical error faithfully represents the actual uncertainty of the approximation, excluding in particular the logical possibility of a substantial but hidden systematic shift. In particular, we consider 11 digits in (28) to be significant and 10 digits to be safe.

In order to obtain a rough estimate for the required number of digits for a successful fit, we considered the complexity of rational numbers appearing in a sample set of analytically known integrals from other topologies. Here, we selected all 318 integrals whose leading term in the ϵ expansion involves weight 7 zeta values, and which might therefore be similar to the unknown integral. We find that, on average, 10 digits were required to successfully reconstruct the rational coefficients of weight 7 zeta values. It therefore seems possible that the right-hand side of (28) could suffice to fit the rational constants on the left-hand side.

While retaining ten digits of approximation (28) is not quite good enough, retaining eleven digits results in the very promising fit:

$$a = \frac{161}{16}, \quad b = \frac{5}{2}, \quad c = -\frac{5}{2}. \quad (30)$$

We now attempt to quantify whether these putative values for a , b , and c are reasonable or not using statistical arguments and numerical extrapolation. This analysis is of considerable importance,

as our fit stretches approximation (28) to its limit. As a start, note that subtracting the presumptive exact value $\mathcal{H}_{\text{conj}}$ obtained from fit (30) from the central value of (12) leaves the difference

$$\delta = \mathcal{H} - \mathcal{H}_{\text{conj}} \approx 7.75 \cdot 10^{-11} \pm 6.98 \cdot 10^{-11}. \quad (31)$$

The ratio of δ divided by its uncertainty is 1.1, which seems reasonable and compatible with $\delta = 0$ in light of the discussion after Eq. (29).

Our PSLQ fit (30) seems plausible in the sense, that the rational numbers involve relatively small integers and are rather similar in structure to what we observe for the rational prefactors of $\{\zeta_7, \zeta_5\zeta_2, \zeta_3\zeta_2^2\}$ in our sample of 318 superficially-similar leading-order integral expansion coefficients.¹ Indeed, we observe that for all 318 elements of our sample, the denominators of the rational prefactors in front of $\{\zeta_7, \zeta_5\zeta_2, \zeta_3\zeta_2^2\}$ are *never* larger than $\{75, 12, 240\}$ and the prime factors which appear in the prime factorizations of the denominators are *never* larger than $\{5, 3, 5\}$. Moreover, the number 161/16 we found for a is actually a quite typical one with respect to our 318 samples of ζ_7 coefficients; its numerator appears in front of ζ_7 10 times in our sample and its denominator is actually the most common one in our sample, appearing 111 times as the denominator of the rational prefactor of ζ_7 .

In order to explore the robustness of the fit (30), we systematically extrapolated the ten significant digits of approximation (28). To be precise, we continue its decimal expansion in all possible ways out to a maximum of fifteen significant digits.² After each digit is added by our trial code, a PSLQ fit is attempted and then judged according to the simplicity and similarity criteria of the previous paragraph. We find it remarkable that, all the way out to fifteen decimal digits, no other possible PSLQ fit looks nearly as natural as (30) above. For this extrapolation analysis, we only considered decimal expansions lying within plus or minus two times the reported statistical uncertainty on $\mathcal{H}$. In an abundance of caution, we repeated our analysis with only nine digits of approximation (28) and found nothing different for extrapolations out to fourteen decimal digits lying within plus or minus five times the reported statistical uncertainty on $\mathcal{H}$.

In terms of $\mathcal{H} = \delta + \mathcal{H}_{\text{conj}}$, our result (20) for the collinear anomalous dimension in $\mathcal{N} = 4$ SYM reads

$$\gamma_4^{\mathcal{N}=4} = -300\zeta_7 - 256\zeta_5\zeta_2 - \frac{768}{5}\zeta_3\zeta_2^2 + \frac{1}{N_c^2} \left[14886\zeta_7 + 3936\zeta_5\zeta_2 - \frac{13104}{5}\zeta_3\zeta_2^2 + 9600\zeta_3^2 + \frac{7136}{7}\zeta_3^3 - 24000\zeta_5 - 5760\zeta_3\zeta_2 + 288\zeta_2^2 + 8640\zeta_3 - 960\mathcal{H} \right].$$

We see that our conjecture, corresponding to $\delta = 0$ in Eq. (20), achieves two simplifications: The expression becomes homogeneous in the transcendental weight of the zeta values, and their coefficients are integers if one chooses the basis element $\zeta_4\zeta_3$ instead of $\zeta_3\zeta_2^2$.

6. Conclusion

In this work, we obtained precise numerical approximations for the four-loop collinear anomalous dimensions of QCD and $\mathcal{N} = 4$ SYM as well as conjectures for the full analytic results in terms

of zeta values. We observe that the principle of maximal transcendentality holds for both the leading and the subleading color contributions. Our experiments with pySecDec suggest that it should be possible to numerically evaluate the finite parts of the QCD form factors to sufficiently high precision for phenomenological purposes.

Note added

We are grateful to Roman N. Lee, Alexander V. Smirnov, Vladimir A. Smirnov, and Matthias Steinhauser for sharing insights regarding the analytic evaluation of $\mathcal{H}$ after we submitted the initial preprint of this article. Indeed, the result of [63] confirms the analytic expression conjectured for $\mathcal{H}$ in this work.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We would like to thank the anonymous referee for helpful suggestions regarding checks of our results and the principle of maximal transcendentality. BA and AvM are supported in part by the National Science Foundation through Grant 2013859. EP is funded as a Royal Society University Research Fellow through grant URF\R1\201473, and previously by All Souls College Oxford. We acknowledge the High Performance Computing Center at Michigan State University for computing resources and thank the team for their help and support. We also thank the Mathematical Institute at the University of Oxford for computing resources.

Appendix A. Supplementary material

Supplementary material related to this article can be found online at <https://doi.org/10.1016/j.physletb.2021.136503>.

References

- [1] G.P. Korchemsky, A.V. Radyushkin, Renormalization of the Wilson loops beyond the leading order, Nucl. Phys. B 283 (1987) 342–364.
- [2] A. Grozin, J.M. Henn, G.P. Korchemsky, P. Marquard, The three-loop cusp anomalous dimension in QCD and its supersymmetric extensions, J. High Energy Phys. 01 (2016) 140, arXiv:1510.07803.
- [3] J.M. Henn, A.V. Smirnov, V.A. Smirnov, M. Steinhauser, A planar four-loop form factor and cusp anomalous dimension in QCD, J. High Energy Phys. 05 (2016) 066, arXiv:1604.03126.
- [4] B. Ruijl, T. Ueda, J.A.M. Vermaseren, J. Davies, A. Vogt, First Forcer results on deep-inelastic scattering and related quantities, PoS LL2016 (2016) 071, arXiv:1605.08408.
- [5] A. von Manteuffel, R.M. Schabinger, Quark and gluon form factors to four-loop order in QCD: the N_f^3 contributions, Phys. Rev. D 95 (2017) 034030, arXiv:1611.00795.
- [6] J.M. Henn, A.V. Smirnov, V.A. Smirnov, M. Steinhauser, R.N. Lee, Four-loop photon quark form factor and cusp anomalous dimension in the large- N_c limit of QCD, J. High Energy Phys. 03 (2017) 139, arXiv:1612.04389.
- [7] R.N. Lee, A.V. Smirnov, V.A. Smirnov, M. Steinhauser, The N_f^2 contributions to fermionic four-loop form factors, Phys. Rev. D 96 (2017) 014008, arXiv:1705.06862.
- [8] S. Moch, B. Ruijl, T. Ueda, J.A.M. Vermaseren, A. Vogt, Four-loop non-singlet splitting functions in the planar limit and beyond, J. High Energy Phys. 10 (2017) 041, arXiv:1707.08315.
- [9] S. Moch, B. Ruijl, T. Ueda, J.A.M. Vermaseren, A. Vogt, On quartic colour factors in splitting functions and the gluon cusp anomalous dimension, Phys. Lett. B 782 (2018) 627–632, arXiv:1805.09638.
- [10] A. Grozin, Four-loop cusp anomalous dimension in QED, J. High Energy Phys. 06 (2018) 073, arXiv:1805.05050, Addendum: J. High Energy Phys. 01 (2019) 134.

¹ By experimenting with our sample expressions, we found that fewer significant digits are generally required for a fit in terms of $\{\zeta_7, \zeta_5\zeta_2, \zeta_4\zeta_3\}$, which is therefore the default ansatz for our actual PSLQ runs.

² It turns out that *all* weight seven terms of our 318 sample expansion coefficients can be reconstructed by running the PSLQ algorithm with fifteen significant digit input precision.

- [11] R.N. Lee, A.V. Smirnov, V.A. Smirnov, M. Steinhauser, Four-loop quark form factor with quartic fundamental colour factor, *J. High Energy Phys.* 02 (2019) 172, arXiv:1901.02898.
- [12] J.M. Henn, T. Peraro, M. Stahlhofen, P. Wasser, Matter dependence of the four-loop cusp anomalous dimension, *Phys. Rev. Lett.* 122 (2019) 201602, arXiv:1901.03693.
- [13] R. Brüser, A. Grozin, J.M. Henn, M. Stahlhofen, Matter dependence of the four-loop QCD cusp anomalous dimension: from small angles to all angles, *J. High Energy Phys.* 05 (2019) 186, arXiv:1902.05076.
- [14] A. von Manteuffel, R.M. Schabinger, Quark and gluon form factors in four loop QCD: the N_f^2 and $N_{q\gamma} N_f$ contributions, *Phys. Rev. D* 99 (2019) 094014, arXiv:1902.08208.
- [15] J.M. Henn, G.P. Korchemsky, B. Mistlberger, The full four-loop cusp anomalous dimension in $\mathcal{N} = 4$ super Yang-Mills and QCD, *J. High Energy Phys.* 04 (2020) 018, arXiv:1911.10174.
- [16] T. Huber, A. von Manteuffel, E. Panzer, R.M. Schabinger, G. Yang, The four-loop cusp anomalous dimension from the $\mathcal{N} = 4$ Sudakov form factor, *Phys. Lett. B* 807 (2020) 135543, arXiv:1912.13459.
- [17] A. von Manteuffel, E. Panzer, R.M. Schabinger, Cusp and collinear anomalous dimensions in four-loop QCD from form factors, *Phys. Rev. Lett.* 124 (2020) 162001, arXiv:2002.04617.
- [18] A.H. Mueller, On the asymptotic behavior of the Sudakov form-factor, *Phys. Rev. D* 20 (1979) 2037.
- [19] J.C. Collins, Algorithm to compute corrections to the Sudakov form-factor, *Phys. Rev. D* 22 (1980) 1478.
- [20] A. Sen, Asymptotic behavior of the Sudakov form-factor in QCD, *Phys. Rev. D* 24 (1981) 3281.
- [21] L. Magnea, G.F. Sterman, Analytic continuation of the Sudakov form-factor in QCD, *Phys. Rev. D* 42 (1990) 4222–4227.
- [22] G.F. Sterman, M.E. Tejeda-Yeomans, Multi-loop amplitudes and resummation, *Phys. Lett. B* 552 (2003) 48–56, arXiv:hep-ph/0210130.
- [23] V. Ravindran, J. Smith, W.L. van Neerven, Two-loop corrections to Higgs boson production, *Nucl. Phys. B* 704 (2005) 332–348, arXiv:hep-ph/0408315.
- [24] Z. Bern, L.J. Dixon, V.A. Smirnov, Iteration of planar amplitudes in maximally supersymmetric Yang-Mills theory at three loops and beyond, *Phys. Rev. D* 72 (2005) 085001, arXiv:hep-th/0505205.
- [25] S.-O. Moch, J.A.M. Vermaseren, A. Vogt, The quark form-factor at higher orders, *J. High Energy Phys.* 08 (2005) 049, arXiv:hep-ph/0507039.
- [26] S. Moch, J.A.M. Vermaseren, A. Vogt, Three-loop results for quark and gluon form-factors, *Phys. Lett. B* 625 (2005) 245–252, arXiv:hep-ph/0508055.
- [27] L.J. Dixon, L. Magnea, G.F. Sterman, Universal structure of sub-leading infrared poles in gauge theory amplitudes, *J. High Energy Phys.* 08 (2008) 022, arXiv:0805.3515.
- [28] G. Falcioni, E. Gardi, C. Milloy, Relating amplitude and PDF factorisation through Wilson-line geometries, *J. High Energy Phys.* 11 (2019) 100, arXiv:1909.00697.
- [29] G. Das, S.-O. Moch, A. Vogt, Soft corrections to inclusive deep-inelastic scattering at four loops and beyond, *J. High Energy Phys.* 03 (2020) 116, arXiv:1912.12920.
- [30] G. Das, S.-O. Moch, A. Vogt, Approximate four-loop QCD corrections to the Higgs-boson production cross section, *Phys. Lett. B* 807 (2020) 135546, arXiv:2004.00563.
- [31] F. Cachazo, M. Spradlin, A. Volovich, Four-loop collinear anomalous dimension in $\mathcal{N} = 4$ Yang-Mills theory, *Phys. Rev. D* 76 (2007) 106004, arXiv:0707.1903.
- [32] R.H. Boels, T. Huber, G. Yang, The Sudakov form factor at four loops in maximal super Yang-Mills theory, *J. High Energy Phys.* 01 (2018) 153, arXiv:1711.08449.
- [33] L.J. Dixon, The principle of maximal transcendentality and the four-loop collinear anomalous dimension, *J. High Energy Phys.* 01 (2018) 075, arXiv:1712.07274.
- [34] V. Ravindran, Higher-order threshold effects to inclusive processes in QCD, *Nucl. Phys. B* 752 (2006) 173–196, arXiv:hep-ph/0603041.
- [35] T. van Ritbergen, J.A.M. Vermaseren, S.A. Larin, The four loop beta function in quantum chromodynamics, *Phys. Lett. B* 400 (1997) 379–384, arXiv:hep-ph/9701390.
- [36] M. Czakon, The four-loop QCD beta-function and anomalous dimensions, *Nucl. Phys. B* 710 (2005) 485–498, arXiv:hep-ph/0411261.
- [37] E. Panzer, Algorithms for the symbolic integration of hyperlogarithms with applications to Feynman integrals, *Comput. Phys. Commun.* 188 (2015) 148–166, arXiv:1403.3385.
- [38] S. Arno, D.H. Bailey, H.R.P. Ferguson, Analysis of PSLQ, an integer relation finding algorithm, *Math. Comput.* 68 (225) (1999) 351.
- [39] F. Tkachov, A theorem on analytical calculability of four loop renormalization group functions, *Phys. Lett. B* 100 (1981) 65–68.
- [40] K. Chetyrkin, F. Tkachov, Integration by parts: the algorithm to calculate beta functions in 4 loops, *Nucl. Phys. B* 192 (1981) 159–204.
- [41] T. Gehrmann, E. Remiddi, Differential equations for two loop four point functions, *Nucl. Phys. B* 580 (2000) 485–518, arXiv:hep-ph/9912329.
- [42] S. Laporta, High precision calculation of multi-loop Feynman integrals by difference equations, *Int. J. Mod. Phys. A* 15 (2000) 5087–5159, arXiv:hep-ph/0102033.
- [43] J. Gluza, K. Kajda, D.A. Kosower, Towards a basis for planar two-loop integrals, *Phys. Rev. D* 83 (2011) 045012, arXiv:1009.0472.
- [44] D. Cabarcas, J. Ding, Linear algebra to compute syzygies and Gröbner bases, in: *Proceedings of the 36th International Symposium on Symbolic and Algebraic Computation (ISSAC '11)*, ACM Press, 2011, pp. 67–74.
- [45] R.M. Schabinger, A new algorithm for the generation of unitarity-compatible integration by parts relations, *J. High Energy Phys.* 01 (2012) 077, arXiv:1111.4220.
- [46] A. von Manteuffel, R.M. Schabinger, A novel approach to integration by parts reduction, *Phys. Lett. B* 744 (2015) 101–104, arXiv:1406.4513.
- [47] H. Ita, Two-loop integrand decomposition into master integrals and surface terms, *Phys. Rev. D* 94 (2016) 116015, arXiv:1510.05626.
- [48] K.J. Larsen, Y. Zhang, Integration-by-parts reductions from unitarity cuts and algebraic geometry, *Phys. Rev. D* 93 (2016) 041701(R), arXiv:1511.01071.
- [49] J. Böhm, A. Georgoudis, K.J. Larsen, M. Schulze, Y. Zhang, Complete sets of log-arithmic vector fields for integration-by-parts identities of Feynman integrals, *Phys. Rev. D* 98 (2018) 025023, arXiv:1712.09737.
- [50] R.N. Lee, A.A. Pomeransky, Critical points and number of master integrals, *J. High Energy Phys.* 11 (2013) 165, arXiv:1308.6676.
- [51] R.N. Lee, LiteRed 1.4: a powerful tool for reduction of multi-loop integrals, *J. Phys. Conf. Ser.* 523 (2014) 012059, arXiv:1310.1145.
- [52] T. Bitoun, C. Bogner, R.P. Klausen, E. Panzer, Feynman integral relations from parametric annihilators, *Lett. Math. Phys.* 109 (2019) 497–564, arXiv:1712.09215.
- [53] A. von Manteuffel, E. Panzer, R.M. Schabinger, A quasi-finite basis for multi-loop Feynman integrals, *J. High Energy Phys.* 02 (2015) 120, arXiv:1411.7392.
- [54] A. von Manteuffel, E. Panzer, R.M. Schabinger, On the computation of form factors in massless QCD with finite master integrals, *Phys. Rev. D* 93 (2016) 125014, arXiv:1510.06758.
- [55] R.M. Schabinger, Constructing multi-loop scattering amplitudes with manifest singularity structure, *Phys. Rev. D* 99 (2019) 105010, arXiv:1806.05682.
- [56] A. von Manteuffel, R.M. Schabinger, Planar master integrals for four-loop form factors, *J. High Energy Phys.* 05 (2019) 073, arXiv:1903.06171.
- [57] F. Brown, The massless higher-loop two-point function, *Commun. Math. Phys.* 287 (2009) 925–958, arXiv:0804.1660.
- [58] F. Brown, On the periods of some Feynman integrals, arXiv:0910.0114.
- [59] S. Borowka, G. Heinrich, S. Jahn, S.P. Jones, M. Kerner, J. Schlenk, T. Zirke, pySecDec: a toolbox for the numerical evaluation of multi-scale integrals, *Comput. Phys. Commun.* 222 (2018) 313–326, arXiv:1703.09692.
- [60] A.V. Kotikov, L.N. Lipatov, DGLAP and BFKL evolution equations in the $\mathcal{N} = 4$ supersymmetric gauge theory, arXiv:hep-ph/0112346.
- [61] A.V. Kotikov, L.N. Lipatov, DGLAP and BFKL equations in the $\mathcal{N} = 4$ supersymmetric gauge theory, *Nucl. Phys. B* 661 (2003) 19–61, arXiv:hep-ph/0208220, Erratum: *Nucl. Phys. B* 685 (2004) 405–407.
- [62] A.V. Smirnov, FIESTA 4: optimized Feynman integral calculations with GPU support, *Comput. Phys. Commun.* 204 (2016) 189–199, arXiv:1511.03614.
- [63] R.N. Lee, A. von Manteuffel, R.M. Schabinger, A.V. Smirnov, V.A. Smirnov, M. Steinhauser, Fermionic corrections to quark and gluon form factors in four-loop QCD, arXiv:2105.11504, 2021.