

1 Plant Derived Tissue and Soil Nutrient Concentration for Plantations of Four Conifer  
2 Species Growing Under Different Site and Vegetation Management Conditions

3

4 Callan Cannon <sup>1,2</sup>, Carlos Gonzalez-Benecke <sup>2,\*</sup> and Maxwell Wightman <sup>2</sup>

5 <sup>1</sup> U.S.D.A. Forest Service; callanfcannon@gmail.com

6 <sup>2</sup> Department of Forest Engineering, Resources and Management, Oregon State

7 University, Corvallis, OR 97331, USA; carlos.gonzalez@oregonstate.edu (C.A.G-B);

8 maxwell.wightman@oregonstate.edu (M.G.W.)

9 \* Corresponding author (carlos.gonzalez@oregonstate.edu); Tel.: +1-541-737-2103

10 Received: date; Accepted: date; Published: date

11

12 **Keywords:** Douglas-fir; western hemlock; western redcedar; grand fir; intensive  
13 silviculture; stand productivity; macronutrients; micronutrients

14

## 15 **Abstract**

16 This study investigates the long-term effects of vegetation management on nutrient  
17 concentration of various tissues and ecosystem components of 16 to 18 year-old Douglas-  
18 fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) stands  
19 growing in Oregon's central Coast Range (CR) and DF and WRC growing in Oregon's  
20 Cascade mountain foothills (CF) under two contrasting vegetation management (VM)  
21 treatments. The treatments consist of: Control, which received no herbicide application post  
22 planting, and VM, which received five years of spring release herbicide application. Both  
23 treatments include a fall site preparation herbicide application. The ecosystem was broken  
24 down into crop trees (separated into foliage, live branches, bark, and stemwood), midstory

25 species (separated into foliage and stem), understory, forest floor, fine roots, and mineral  
26 soil (with depth increments 0.0-0.2 m, 0.2-0.4 m, 0.4-0.6 m, and 0.6-1.0 m). All samples  
27 were analyzed for concentration of total carbon, nitrogen, phosphorus, potassium, calcium,  
28 magnesium, sulfur, boron, copper, iron, manganese, sodium, and zinc. This study design  
29 resulted in 1,740 unique nutrient concentration results being reported. The effect of VM  
30 (treatment) on tissue concentration varied by nutrient, overstory crop species (species),  
31 ecosystem component, and site. Forest floor and crop tree bark, followed by fine roots, were  
32 the ecosystem component nutrient concentrations that showed the greatest number of  
33 treatment effects across all species. Soil concentrations showed large variation across sites  
34 but were generally unaffected by treatment and species. At the CR site, magnesium and  
35 calcium soil concentrations were higher in VM plots across species, while zinc  
36 concentrations were lower. There were no other effects of treatment on soil nutrient  
37 concentrations, but there were some significant treatment x crop species interactions. Most  
38 notably, at the CF site, the concentration of C and N were higher in VM plots than control  
39 plots of DF, while the opposite was true for WRC. While total soil concentrations were  
40 generally unaffected by treatment and are unlikely to be adversely affected in the long term,  
41 it is possible that VM can reduce soil nitrogen for slow growing species like WRC.

## 42 **Introduction**

43 Tissue and soil nutrient concentrations are useful measures in order to determine the  
44 nutrient status of a stand as well as potential for nutrient deficiencies or soil nutrient depletion  
45 ( Turner et al., 1977; Stone, 1990; Slesak et al., 2016; DeBruler et al., 2019). They are the  
46 basis for various nutrient management guidelines such as Diagnosis and Integrated  
47 Recommendation system (DRIS) and the Kinsey regime which allow development of site-  
48 specific fertilization prescriptions (Beaufils, 1973; Mainwaring et al., 2014). Nutrient  
49 concentrations are useful in this respect because they indicate how much of a resource is

50 available in the exploitable soil as well as whether plant foliage is optimally equipped to meet  
51 a plant's physiological needs. If a plant is lacking a particular nutrient or set of nutrients such  
52 that its physiological processes are limited, it will have a suboptimal concentration of  
53 nutrients in its foliage. The lowest foliar concentration where nutrients do not significantly  
54 limit growth is known as the critical concentration (Ulrich, 1952).

55 Plants distribute nutrients throughout their tissues in order to satisfy their physiological  
56 needs. These nutrients are often divided into two categories, macronutrients and  
57 micronutrients, based on the relative requirements of plants. The following are considered  
58 macronutrients and are required in larger amounts: carbon (C), nitrogen (N), phosphorous  
59 (P), potassium (K), calcium (Ca), magnesium (Mg), and sulfur (S). The following are  
60 considered micronutrients and are required in much smaller amounts: boron (B), copper (Cu),  
61 iron (Fe), manganese (Mn), sodium (Na), and zinc (Zn).

62 Foliage is generally the tissue type that contains the greatest concentration of nutrients  
63 with the exception of Ca which may be higher in the branches, trunk and phloem (Cole and  
64 Gessel, 1992; Augusto et al., 2008; Marschner and Marschner, 2012). While foliage  
65 comprises approximately 4% of aboveground biomass in a 40-year-old Douglas-fir stand, it  
66 contains roughly 70% of the total aboveground nitrogen (Turner and Long, 1975; Turner,  
67 1981; Cole and Gessel, 1992).

68 Silvicultural treatments, such as vegetation management (VM), during the establishment  
69 phase set the trajectory for stand development. These treatments may affect plants by altering  
70 the concentration of nutrients in a tissue or in soil (Burger and Pritchett, 1988; Powers and  
71 Reynolds, 1999; Powers et al., 2005). Looking at the content of a tissue may not reveal  
72 physiologically important changes and may only show trends in biomass if concentrations  
73 remain the same. A decrease in tissue nutrient concentration may mean that an organism is  
74 having difficulty meeting its physiological needs for that nutrient, whereas a decrease in

75 content can be the result of a number of factors such as reduced biomass or changes in  
76 allocation.

77 The effects of VM on plant nutrient concentrations has been studied, although generally  
78 in younger tree seedlings. VM allows trees greater access to site resources and commonly  
79 affects nutrient content as the treatments often produce significantly more biomass in most  
80 tissues (Petersen et al., 2008; Devine et al., 2011), whereas its effects on nutrient  
81 concentration vary by study and tree age. Five-year-old Douglas-fir seedlings have shown  
82 increased foliar N content and concentration with vegetation control ( Slesak et al., 2010;  
83 Devine et al., 2011). These trends varied between sites and concentration effects were only  
84 significant at the study level and not at the site level (Devine et al., 2011). A study in the  
85 Oregon Coast Range showed N was higher in VM treated Douglas-fir seedlings after the first  
86 year of growth but not the second (Rose and Ketchum, 2002). In contrast, B showed a  
87 significant decrease in VM treated plots but only after the second year of growth (Rose and  
88 Ketchum, 2002). Differences in concentrations are not always observed, as Petersen et al.  
89 found that there were no differences in foliar N, P, K, S, Ca and Mg in five-year-old Douglas-  
90 fir seedlings (Petersen et al., 2008). A recent study at the Long Term Soil Productivity (LTSP)  
91 sites in the Pacific Northwest found that effects on plant nutrition in 15- and 20-year-old  
92 Douglas-fir stands varied by site and soil properties (Littke et al., 2020a). One site with  
93 historically low base cations showed reduced foliar Ca with sustained VM. Another site with  
94 historically higher cations and lower N displayed increased foliar Al and Mg and lower foliar  
95 N at a second site with sustained VM, and no detectable differences in foliar nutrients at a  
96 third site (Littke et al., 2020a).

97 The effects of VM on foliar nutrients change over time. Across a gradient of site  
98 conditions, foliar N and P concentrations were greater for treated plots early in stand  
99 development. These differences disappeared at ages 7 and 9 for all sites, except for N  
100 concentrations at the site that had lowest N levels and untreated trees displayed signs of N

101 deficiency (Powers and Reynolds, 1999). One study of loblolly pine conducted at mid-  
102 rotation found that eradication of herbaceous vegetation during stand establishment resulted  
103 in a decrease in foliar N and K (Miller et al., 2006). They found that all available soil nutrients  
104 declined over time but this decline was greater for C, N and Ca.

105 The effect of silvicultural management on soil concentration has also been studied, with  
106 most studies focusing on different forms of N or P. The LTSP study has investigated the  
107 effects of different intensive management practices across the US, including sites in the PNW  
108 (Powers et al., 2005). Sites in Oregon show that after planting, soil nutrients (exchangeable  
109 Ca, Mg, K, and total N) tend to increase after 10 years in the top 0.3 m of soil, although the  
110 increase is greater when there is no vegetation control after planting (Slesak et al., 2016).  
111 Total soil P is more variable, tending to decrease 10 years after planting in the top 0.3 m. At  
112 one site the decrease was less when harvest residues were left on site and there was no  
113 vegetation control after planting, while at the other site the decrease was less with annual  
114 vegetation control after planting (Slesak et al., 2016). A follow up study looked at total P and  
115 different pools of labile to less labile P 10 years after planting which all showed roughly the  
116 same result: at one site, when there was a detectable difference in P concentrations of any  
117 pool, concentrations were higher with no annual vegetation control while the other site  
118 showed the opposite trend (DeBruler et al., 2019). A similar study from the Fall River LTSP  
119 site in Washington showed that total soil N concentrations in the top 0.15 m of soil decreased  
120 10 years after planting (Knight et al., 2014). A recent study at the same sites showed a general  
121 decrease in soil base cations and reduced simulated nitrate uptake at 15 or 20 years with  
122 annual VM, with forest floor samples showing similar trends (Littke et al., 2020a, 2020b).

123 Most studies look at only a few nutrients and tend to focus on younger trees and only  
124 one or two crop species (typically Douglas-fir and ponderosa pine in Oregon). In this study  
125 we investigated how vegetation management affected various nutrients (7 macro, including  
126 C, and 6 micro) on multiple conifer species (Douglas-fir, western hemlock, western redcedar,

127 and grand fir) in two important timber producing ecoregions in Oregon (the Oregon Cascade  
128 foothills and the Oregon Coast Range). The specific objectives of this study were: to quantify  
129 nutrient concentrations of all ecosystem components, explore how these varied by overstory  
130 species, site, and VM treatment, and explore whether crop tree foliar concentrations were  
131 correlated with soil concentrations.

## 132 Materials and Methods

### 133 Description of Sites

134 Two contrasting study sites were selected for this study. The Coastal Range (CR) site is  
135 located at 44.616°N, 123.574°W near Summit, OR, approximately 40 km from the coast. The  
136 site was planted in the year 2000 and experiences a mean annual temperature of 11.1°C and  
137 average annual rainfall of 1,707 mm. The CR site was planted with coast Douglas-fir (DF,  
138 *Pseudotsuga menziesii* var. *menziesii* (Mirbel) Franco) and western hemlock (WH, *Tsuga*  
139 *heterophylla* (Raf.) Sarg.) (four replicates each, eight plots per species), and western redcedar  
140 (WRC, *Thuja plicata* Donn ex D. Don) and grand fir (GF, *Abies grandis* (Dougl. ex D. Don)  
141 Lindl.). (three replicates each, six plots per species). Soils at the CR site are part of the  
142 Preacher-Bohannon complex which is derived from siltstone and sandstone, and has a fine  
143 and loamy texture (Flamenco et al. 2019, USDA 2009). This soil complex is classified as an  
144 Andic Dystrudept, meaning that while it is not an Andosol, it has high aluminum and iron  
145 activity (Soil Survey Staff 2015). This site sits near the western edge of the Tyee formation,  
146 a sedimentary rock formation that is composed largely of marine micaceous sandstone and  
147 siltstone.

148 The Cascade Foothills (CF) site is located at 44.476°N, 122.726°W near Sweet Home,  
149 OR, and was planted in the year 2001 only with DF and WRC (four replicates each). The site  
150 has a mean annual temperature of 12.4°C and an average annual rainfall of 1,179 mm. Soils  
151 at the CF site are from the Bellpine series which is derived from sedimentary rock, and have

152 a fine and loamy texture (Flamenco, et al. 2019, Soil Survey Staff 2015, Ulrich, 1952). Soils  
153 of this series are classified as Xeric Haplohumults, indicating an Ultisol with high organic  
154 matter content that experiences seasonal drought. These soils are well drained and  
155 characterized by a more xeric moisture regime than the CR site. The bedrock is a mixture of  
156 basalt, sedimentary rocks, and tuff. Similar to the CR site, these soils are derived from  
157 sedimentary bedrock, however tuff and mafic intrusions will lend different chemical  
158 characteristics to these soils. Mafic rocks tend to be higher in iron and magnesium than  
159 sandstone. This site was formerly agricultural land that was not sufficiently productive.

160 *Study Design*

161 A randomized complete block design with eight VM regimes (treatments) was  
162 implemented at each of the two sites. The eight different VM treatments consisted of spring  
163 release applications that differed in the number and timing of herbicide treatments applied  
164 during the first 5 years after planting, see Chen 2004 for more details (Chen, 2004). Similar  
165 to Flamenco et al. (2019), for this study we used only the control (Control; only pre-planting  
166 vegetation control) and the 5 consecutive years of spring release vegetation management  
167 treatments (VM). Each treatment plot was 24.4 m x 24.4 m (0.06 ha) in size and was planted  
168 with 64 seedlings (8 rows of 8 trees) with 3 m x 3 m spacing, resulting in a planting density  
169 of 1,111 trees ha<sup>-1</sup>. Measurement plots consisted of the internal six rows of six trees allowing  
170 for a one tree buffer on all sides. All plots were planted with a single tree species, and the  
171 experimental unit was the plot. All DF plots received pre-commercial thinning at age 12 years  
172 to reduce stocking by 25% and thinning residues were left on site. A summary of stand  
173 attributes at age 18-years is provided in Table A1.

174 The ecosystem was divided into soil layers and plant derived tissues. The plant derived  
175 components were broken down into overstory (planted crop trees), midstory (hardwoods and  
176 natural conifer regeneration), understory (shrubs, grasses, forbs, ferns and moss) and forest  
177 floor (including coarse woody debris). The overstory was divided into foliage, live branches,

178 stemwood, bark, and fine roots. The midstory was broken down into foliage and bole  
179 (stemwood and bark). The soil was divided into four layers (0-0.2 m, 0.2-0.4 m, 0.4-0.6 m,  
180 and 0.6-1 m).

181 Tissue samples were collected from both overstory crop trees and midstory hardwood  
182 species. The crop tree canopy was above that of the midstory species and tree sizes are  
183 reported in Flamenco et al. (2019). Overstory tissue for nutrient analysis were obtained from  
184 samples collected by Flamenco et al. (2019), who destructively sampled 4 trees for each crop  
185 species and treatment at each site (48 trees total). Sampled trees were chosen to represent the  
186 range of stem diameters present at both sites. Stemwood samples were collected by removing  
187 a stem section (or cookie) at DBH. Stem bark samples were obtained by removing the bark  
188 from the cookie taken at DBH. Branch and foliage samples were collected from the middle  
189 of the living crown (see Flamenco et al., 2019) for further details on crop tree sampling).

190 As dominant midstory species are the same across sites, samples for nutritional analysis  
191 were taken only at the CR site without respect to treatment (only few midstory individuals  
192 were found in the VM plots). Midstory tissue samples for nutrient analysis (foliage and  
193 stemwood) were collected from midstory trees during July 2019. Only the four most  
194 prevalent species were sampled: red alder (*Alnus rubra* Bong.), bigleaf maple (*Acer*  
195 *macrophyllum* Pursh), Oregon cherry (*Prunus emarginata* (Douglas ex Hook.) D. Dietr.),  
196 and cascara buckthorn (*Frangula purshiana* DC.). These four species account for 98% of the  
197 midstory biomass (Flamenco et al., 2019). Stemwood samples were collected at DBH using  
198 a 12-mm increment borer from four different individuals from each species. Foliage samples  
199 were also taken from four different individuals from each species.

200 Understory, forest floor and fine roots were collected from 6 subplots (0.6 m x 0.6 m)  
201 per plot. All vegetation in or hanging over these plots was collected. The forest floor was  
202 manually removed down to the organic horizon and included woody debris, duff, and litter.  
203 Researchers then collected a core of the top 0.2 m of mineral soil and used a 2 mm sieve to

204 collect fine roots (Flamenco et al., 2019). Within a plot, all six subsamples were combined  
205 for nutrient analysis. One sample from each of the lower soil layers (from 0.2 m to 1.0 m  
206 depth) was collected in the spring 2019 from each plot using 50 mm x 50 mm soil cores  
207 (AMS, bulk density soil sampling kit). Fine roots were collected from these soil samples  
208 using a 2 mm sieve.

209 *Nutrient Analysis*

210 All plant samples were oven-dried at 65°C until reaching constant weight and ground to  
211 pass a 0.425 mm sieve. These tissues were then prepared for nutrient extraction by overnight  
212 combustion in quartz tubes at 580°C. Samples were extracted in 20% v/v HCl for 15 minutes  
213 and then diluted 1:1 with distilled water. These extracts were filtered and stored at 4°C until  
214 analysis. Total soil nutrients were extracted by microwave digestion. Samples were heated to  
215 175°C in an Anton-Paar MicrowaveGO and held at that temperature for 4.5 minutes in a  
216 solution of 70% HNO3. Digested samples were diluted 1:1 with distilled water, filtered, and  
217 stored at 4°C until analysis. Concentrations of C, N and S were determined by dry combustion  
218 using an Elementar vario MACRO cube. All other nutrients (P, K, Mg, Ca, B, Cu, Fe, Mn,  
219 Na, and Zn), were determined by analyzing extracts with an Agilent ICP-OES 5110. All  
220 analyses were carried out at the Central Analytical Laboratory at Oregon State University.

221 *Statistical Analysis*

222 The Statistical Analysis Software version 9.4 (SAS Institute Inc. Cary, NC) was used for  
223 all statistical analysis. Analysis of variance, including Tukey multiple comparisons tests, was  
224 used to test the effects of site, species and treatments on all soil and plant derived  
225 concentrations (PROC MIXED, SAS Institute Inc. Cary, NC), where block was included as  
226 a random effect. Significance was determined using  $\alpha=0.05$ . As not all species were planted  
227 at each site- site, site x species, site x treatment, and site x species x treatment effects were  
228 calculated using a reduced dataset including only Douglas-fir and western redcedar. Pearson

229 correlation coefficients between plant nutrient concentrations and soil nutrient concentrations  
230 were determined across treatments, species and sites (PROC CORR, SAS Institute. Cary,  
231 NC). SigmaPlot version 14 (Systat Software, Inc. San Jose, CA) was used to create all figures.

232 **Results**

233 The results from this study are extensive, with results of 13 different nutrient  
234 concentrations for 11 ecosystem components of conifer plantations of four different crop  
235 species growing under two contrasting VM treatments at two sites. The second site only  
236 contained two of the four species, two nutrients (K and Na) were below detectable levels in  
237 stemwood, and S was not measured for soil components- resulting in 1,644 unique nutrient  
238 concentration results. Nutrient concentrations for the foliage and stem of four midstory  
239 species growing at the CR site are also reported for an additional 96 unique results. We  
240 focused on treatment, site, and species effects of N, P, K, Mg, B, Mn, Zn, and Cu as well as  
241 correlations between soil nutrients and plant derived nutrients. Results for each of the 1,740  
242 unique ecosystem component nutrient concentrations can be found in the appendix. Tables  
243 A2-A14 provide values for each of the 13 nutrients for all ecosystem components, crop  
244 species, VM treatments, and sites. Tables A15 and A16 provide values for the midstory and  
245 understory.

246 *Crop Species and Vegetation Management Effects*

247 A summary of ANOVA results for the effects of crop species, treatment, and crop species  
248 x treatment interaction on nutrient concentrations are provided in Table 1 for the CR site and  
249 Table 2 for the CF site. We considered P-values less than 0.05 to be significant but have also  
250 included values between 0.05 and 0.1 for reader's consideration. In general, crop species had  
251 a larger effect on nutrient concentrations than treatment or crop species x treatment  
252 interaction. At the CR site, 30%, 12%, and 7% of nutrient concentrations (n=137) were  
253 affected by crop species, treatment, and crop species x treatment interaction, respectively,

254 while 51% were unaffected by these factors (Table 1). For the CF site, 23%, 5%, and 8% of  
255 nutrient concentrations were affected by crop species, treatment, and crop species x treatment  
256 interaction, respectively, while 64% were unaffected by these factors (Table 2).

257

**Table 1.** P values for the effect of crop species (SPP), treatment (TRT), and their interaction (SPP\*TRT) on the concentration of C, N, P, K, Mg, Ca, S, B, Cu, Fe, Mn, Na, and Zn in plant derived and soil ecosystem components of 16-18 year-old Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) stands growing on a site located in the central Coast Range (CR) of western Oregon. P values only included when  $P < 0.1$  and values below 0.05 are presented in bold.

**Table 2.** P values for the effect of crop species (SPP), treatment (TRT), and their interaction (SPP\*TRT) on the concentration of C, N, P, K, Mg, Ca, S, Cu, Fe, Mn, Na, and Zn in plant derived and soil ecosystem components of 16-18 year-old Douglas-fir (DF) and western redcedar (WRC) stands growing on a site located in Cascade foothills (CF) of western Oregon. P values only included when  $P < 0.1$  and values below 0.05 are presented in bold.


268

269 At both sites, plant derived nutrient concentrations (crop trees, understory, and forest  
270 floor, n = 89) were more affected by crop species and treatment than the soil components  
271 (n=48). At CR, 61% of plant derived nutrient concentrations and 27% of soil nutrient  
272 concentrations were affected by crop species, treatment, or their interaction. At CF, 46% of  
273 plant derived nutrient concentrations and 19% of soil nutrient concentrations were affected.  
274 The understory was largely unaffected by crop species and treatment, only showing  
275 significant effects for C and Fe at the CR site and S and Zn at the CF site. Mg, Mn, and C  
276 were the nutrients most effected by treatments at CR and N, Ca, and Mn were the most  
277 affected nutrients at CF.

278 Within crop tree tissues, fine roots showed the lowest C concentration, ranging between  
279 27.4 to 33.9% (indicating that the fine root sample likely included dead roots), while all other  
280 crop tree tissues ranged from 46% to 50% (standard for living plant tissue (Ågren, 2008) ).  
281 The concentration of C in crop tree branches, bark, stemwood, and roots varied by species at  
282 the CR site and was generally higher for DF than the other species, except for roots (Table  
283 1). The only effect of treatment on crop tree C concentration at CR was VM plots having  
284 higher branch C than control plots (P=0.044). VM treatment did not affect crop tree C at CF,  
285 but DF had higher bark and foliage C than WRC (Table 2, P<0.003).


286

**Figure 1.** Concentrations of N, P, K, and Mg for foliage, branches, bark, stemwood, fine roots and understory of 16-18 year-old stands of Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) at sites in the central Coast Range (CR, shown above dotted line) and Cascade foothills (CF, shown below dotted line) of Western Oregon. Concentrations of control plots are shown with a white symbol and treatment plots are shown with a filled symbol.



288 For all species, the largest N, P and K concentrations were observed in foliage, ranging  
289 between 0.978 to 1.252% N, 0.116 to 0.255% P, and 0.381 to 0.607% K (Figure 1). At the  
290 CF site, DF has higher foliar concentrations of N, P, and K than WRC ( $P<0.012$ ). Foliar N  
291 was not affected by crop species or treatment at the CR site, however foliar P was  
292 significantly higher in WH than all other species ( $P<0.031$ ). There was a significant crop  
293 species x treatment interaction for foliar K at the CR site ( $P=0.005$ ) such that WH growing  
294 in the control had higher foliar K than WRC growing under either treatment ( $P<0.047$ ). Foliar  
295 Mg was not affected by crop species at CR but was significantly higher in control plots  
296 ( $P=0.014$ ). At CF, foliar Mg of DF was higher in VM plots than control plots ( $P=0.027$ ),  
297 while WRC was unaffected by treatment (Figure 1). The concentration of N, P, K and Mg  
298 were lower in branches, bark, stemwood, and roots than foliage and often varied by species,  
299 and to a lesser extent treatment, except for K at the CF site (Tables 1 and 2). Bark was to  
300 most sensitive to crop species and treatment followed by branches and stemwood.

**Figure 2.** Concentrations of N, P, K, and Mg for forest floor and the 0-20 cm, 20-40 cm, 40-60 cm and 60-100 cm soil layers of 16-18 year-old stands of Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) at sites in the central Coast Range (CR, shown above dotted line) and Cascade foothills (CF, shown below dotted line) of Western Oregon. Concentrations of control plots are shown with a white symbol and treatment




302

303 The concentration of B, Cu, Mn, and Zn in crop tree tissues were not affected by treatment  
304 at either site except for fine root Cu at both sites and bark Cu and Zn at CR (Tables 1 and 2).  
305 In each of these cases, nutrient concentrations were higher in control plots than VM plots  
306 (Figure 3). The effect of crop species was more pronounced than that of treatment. At the CR  
307 site, foliar, branch, bark, and stemwood B, Mn, and Zn all varied by species except for  
308 stemwood Mn. For example, WH foliar B was higher than DF and WRC while GF foliar Zn  
309 was higher than DF and WH. Crop tree Cu concentrations were generally unaffected by  
310 species with the exception of DF having higher stemwood Cu than WH and GF. When the  
311 effect of crop species was significant for crop tree tissue B, Cu, Mn, and Zn at CF (Table 2),  
312 concentrations tended to be higher in DF than WRC except for bark B and fine root Cu.

313


**Figure 3.** Concentrations of B, Mn, Zn, and Cu for foliage, branches, bark, stemwood, fine roots and understory of 16-18 year-old stands of Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) at sites in the central Coast Range (CR, shown above dotted line) and Cascade foothills (CF, shown below dotted line) of Western Oregon. Concentrations of control plots are shown with a white symbol and treatment plots are shown with a filled symbol.



317       Soil nutrient concentrations were mostly unaffected by crop species or treatment (Tables  
318   1 and 2). At both sites, six of the thirteen nutrients did not show any crop species, treatment,  
319   or crop species x treatment interaction for any soil depth. Additionally, three nutrients only  
320   showed an effect for one of the four soil layers at CR while this was true for five nutrients at  
321   CF. Soil Mg was the most impacted nutrient at CR with WH having lower soil Mg than WRC  
322   in all three of the upper soil layers and all other species in the 0.2-0.4 m layer ( $P<0.050$ ). Soil  
323   Mg in the 0.6-1.0 m layer was not affected by species but was higher in VM plots than control  
324   plots at CR ( $P=0.046$ ). It should be noted that there were no detectable species differences in  
325   the deepest layer (0.6-1.0 m) for any nutrient at CR. There was a treatment x crop species  
326   interaction for soil C and N at CF such that the concentration of these elements in the 0.4-0.6  
327   m and 0.6-1.0 m layers was higher in VM plots than control plots for DF while the opposite  
328   was true for WRC (Figure 2).

329       There were a few significant site x crop species x treatment interactions. Notably, there  
330   were two depths (0.2-0.4 m and 0.4-0.6 m) for WRC at the CF site where there was  
331   significantly lower soil N in treated plots than Control plots ( $P<0.05$ ) and one layer (0.6-1.0  
332   m) for which this trend was marginally significant ( $P=0.07$ ). For DF at the CF site soil N  
333   concentrations were higher in treated plots than Control plots for the 0.6-1.0 m depth  
334   ( $P<0.05$ ). Soil C concentrations were higher in the 0.4-0.6 m depth for VM plots of DF at the  
335   CR site ( $P<0.05$ ).

**Figure 4.** Concentrations of B, Mn, Zn, and Cu for forest floor and the 0-20 cm, 20-40 cm, 40-60 cm and 60-100 cm soil layers of 16-18 year-old stands of Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) at sites in the central Coast Range (CR, shown above dotted line) and Cascade foothills (CF, shown below dotted line) of Western Oregon. Concentrations of control plots are shown with a white symbol



339

340 A more general review of the results showed some interesting trends. Nutrient  
341 concentration in mineral soil decreased with depth for C, N, P and Ca, but no clear trend was  
342 observed for K and Mg (Figure 2). Micronutrient concentrations of soils decreased with  
343 increasing depth for Mn and Zn while other micronutrients showed no pattern. For Na, the  
344 top layer of soil contained the lowest concentration across all species (Figure 4).  
345 Concentrations of Mg, Ca and S in forest floor were relatively high, ranging between 0.116  
346 to 0.146% Mg, 0.754 to 1.600% Ca and 0.090 to 0.111% S. Both, Cu and Fe, had the highest  
347 concentrations in fine roots ranging between 4.8 and 6.4 ppm Cu and 1209 and 1554 ppm Fe.  
348 The forest floor also contained a notably high concentration of Fe ranging from 914 to 1281  
349 ppm. The concentrations of Mn were highest in the forest floor for all species except WRC,  
350 with concentrations ranging from 449 to 833 ppm. The concentration of B was highest in  
351 foliage for all species except for WRC, with concentrations averaging between 22.3 and 12.4  
352 ppm. Each species had highest Zn concentrations in a different tissue. The concentration of  
353 Na was highest in fine roots and forest floor, averaging between 118 and 162 ppm. In WRC,  
354 concentrations of Zn, B, and Mn were highest in fine roots.

355 *Site Effects*

356 The effect of site on nutrient concentrations (averaged across VM treatments) is provided  
357 in Table 3. The effect of site was more pronounced in DF plots than WRC plots. 39% of plant  
358 derived nutrient concentrations (crop trees, understory, and forest floor, n=89) in DF plots  
359 were significantly affected by site compared to 29% in WRC plots. 73% and 46% of soil  
360 nutrient concentrations (n=48) were affected by site in DF and WRC plots, respectively. Soil  
361 nutrient concentrations were highly site dependent for all depths, with the exception of C, N  
362 and Zn. 57% of the 119 significant site effects indicated that the nutrient concentration was  
363 higher at the CF site than the CR site.

364

365 **Table 3.** P values of site effect for concentration of C, N, P, K, Mg, Ca, S, B, Cu, Fe, Mn,  
 366 Na, and Zn for each nutrient tissue type and soil layer for 16-18 year-old Douglas-fir (DF)  
 367 and western redcedar (WRC) stands growing on sites located in the central Coast Range (CR)  
 368 and the Cascade foothills (CF) of western Oregon (data averaged between Control and VM  
 369 treatments). Green cells indicate that the concentration was higher at the CR site and white  
 370 cells indicate the concentration was higher at the CF site. Blank cells indicate no significant  
 371 differences across sites.

Spp	Tissue	C	N	P	K	Mg	Ca	S	B	Cu	Fe	Mn	Na	Zn
DF	Foliage*				0.005				0.009		0.039		0.002	
	Branch*		0.032					0.028					0.020	
	Bark*					0.004					0.003		0.042	
	Wood*								0.040	0.001	0.002		0.040	
	Root*	0.014				<0.001	0.001	<0.001		0.001	0.002	0.001	0.022	
	Understory*	0.011	0.013	0.033				0.034		0.015				
	Forest floor*	0.033			0.029		0.006		0.004	0.006		<0.001	0.022	0.018
	Soil 0.0-0.2 m*			<0.001	0.001	<0.001		<0.001		<0.001	0.001	<0.001	<0.001	
	Soil 0.2-0.4 m**			<0.001	0.002	<0.001		0.002		<0.001	0.004	<0.001	0.004	
	Soil 0.4-0.6 m**			0.001	0.001	<0.001		0.016		<0.001	0.008	<0.001	0.038	<0.001
	Soil 0.6-1.0 m**			0.001	<0.001	<0.001		0.047		<0.001	0.003	<0.001	0.013	0.001
WRC	Foliage*					0.024								
	Branch*										<0.001			
	Bark*	0.010			0.002		0.001						<0.001	
	Wood*	<0.001												
	Root*	0.033			0.009	0.029	0.017			0.012	0.019	<0.001		
	Understory*						0.009		0.022			0.036		
	Forest floor*					0.036		0.049					<0.001	
	Soil 0.0-0.2 m*			0.002	<0.001		0.008		0.002	0.002	<0.001	0.006		
	Soil 0.2-0.4 m**			0.001	<0.001		0.014		0.001	<0.001	<0.001	0.019		
	Soil 0.4-0.6 m**					<0.001	<0.001		0.007	<0.001	<0.001	0.038	<0.001	
	Soil 0.6-1.0 m**					0.001	<0.001	0.021		0.016	0.018	0.001	0.020	<0.001

372 \*: sampled at age 16 years; \*\*: sampled at age 18 years

373

374 Concentrations of N were lower at the CR site for the understory in DF plots and the  
 375 bark and stemwood of WRC, but the concentration was higher for the branches of DF (Table  
 376 3). Concentrations of B were lower at the CR site in the forest floor, foliage, roots, and  
 377 understory of DF. Concentrations of C at the CR site were lower in the forest floor and  
 378 understory of DF, but higher for roots of both DF and WRC. For Ca, concentrations were  
 379 lower at the CR site for roots and understory of both DF and WRC, but higher in the bark of  
 380 WRC. Concentrations of Fe were lower at the CR site for the bark, foliage, and stemwood of  
 381 DF and lower in the fine roots of both DF and WRC. For K, concentrations were lower at the  
 382 CR site for the forest floor and foliage of DF, but higher for the bark of WRC and the fine


383 roots of both DF and WRC. Concentrations of Mg were higher at the CR site for the bark and  
384 roots of DF and the forest floor, foliage and roots of WRC. Concentrations of Mn were lower  
385 at the CR site for the forest floor, roots, and stemwood of DF and for the roots and understory  
386 of WRC. Concentrations of Na were higher at the CR site for the bark, branches, forest floor,  
387 foliage, and fine roots of DF and for the bark and forest floor of WRC.


388 Generally, soil nutrient concentrations of Ca, B, Cu, Fe, and Mn were higher at the CF  
389 site and soil nutrient concentrations of K, Mg and Na were higher at the CR site for both  
390 species (Table 3). Soil nutrient concentrations tended to show similar patterns for all depths,  
391 with all layers being significantly higher at one site or displaying no significant difference.  
392 Na and K were the only nutrient concentrations that were significantly different between sites  
393 in some layers but not others. Phosphorous was the only nutrient that had differences in soil  
394 nutrient across sites for one species but not the other, being significantly higher in all layers  
395 at the CF site for DF but not for WRC.

396 *Correlations between Soil Nutrient Concentration and Crop Tree Foliar Nutrient  
397 Concentration*

398 Soil nutrient concentrations (weighted averaged across depths) were correlated with foliar  
399 nutrient concentrations for several nutrients and species. DF was the species that showed the  
400 greatest number of significant correlations, with foliar concentrations of P, B, Na, and Fe  
401 increasing with increasing soil concentrations (Figures 5 and 6). Significant positive  
402 correlations were also observed between soil and foliar concentrations of Mg for WRC and  
403 between soil and foliar concentrations of Zn for GF. It is likely that the correlations observed  
404 for DF and WRC are driven by differences between sites. When the sites are analyzed  
405 separately, the only one of the above correlations for DF and WRC that remains marginally  
406 significant is the relationship between soil and foliar concentrations of Fe for DF at the CR  
407 site ( $P=0.093$ , data not shown).

Figure 5. Plots of foliar vs soil concentrations of C, N, P, K, Ca, and Mg of 16-18 year-old stands of Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) at sites in the central Coast Range (CR) and Cascade foothills (CF) of Western Oregon. Correlations are shown when the model is significant across both sites ( $P < 0.1$ ). Correlations for DF are shown with a solid line and the correlations for WRC are shown with a dotted line. Soil nutrient concentrations represent a weighted average for the top 1 m of soil.





438 *Midstory Species Nutrient concentrations*

439 Nutrient concentrations for the stem and leaves of midstory species sampled at the CR  
440 site are presented in Tables A14 and A15. As with crop trees, foliage had higher  
441 concentrations of all nutrients than the stem with the exception of C, which was  
442 approximately 50% for both the foliage and stem. When compared to crop trees, several foliar  
443 nutrients tended to be higher in hardwood foliage. Midstory foliar nutrients generally had  
444 higher concentrations of N, K, Mg, and Cu than all crop species. P was generally higher in  
445 midstory foliage with ACMA, PREM, and FRPU average concentrations ranging from  
446 0.331-0.412 ppm, ALRU was the exception to this trend with an average foliar concentration  
447 of 0.147 ppm which falls within the range of crop tree foliage.

448 **Discussion**

449 Treatment effects on nutrient concentration varied by site, tissue, and nutrient. Bark and  
450 forest floor were the two tissue types most affected by vegetation control treatment, followed  
451 by fine roots. Crop tree foliage, branches, and stemwood all showed no treatment differences  
452 for all species at both sites, except for foliar Mg at CR, branch N at CF, and stemwood Ca at  
453 CF. The forest floor was the tissue type most affected by treatment. This makes sense as the  
454 litter from the VM plots was almost entirely composed of conifer litter, with some inclusion  
455 of understory litter, whereas the forest floor of the C plots contained litter from midstory  
456 species, whose foliar nutrition differs significantly from the conifers. Concentrations of Cu  
457 and Mg were higher in the forest floor for control plots, although this trend was less  
458 pronounced for DF and WRC at the CF site, since untreated plots had less robust midstory  
459 development (Flamenco et al. 2019). Concentrations of K in forest floor were also higher in  
460 Control plots, but this trend was more pronounced for WRC. These trends of higher base  
461 cations in the forest floor without annual vegetation control agree with similar findings in 15  
462 to 20-year-old Douglas-fir (Littke et al., 2020a, 2020b). Concentrations of Mn were higher

463 in forest floor of VM plots, which makes sense because conifers are accumulators of this  
464 nutrient. As observed elsewhere, cascara buckthorn also accumulated high concentrations of  
465 Mn in its foliage, but other midstory species did not (Zasoski et al., 1990).

466 Bark was the tissue type second most often affected by treatment, with effects seen for  
467 P, K, Mg, and Ca. Generally, with the exception of DF K concentrations, bark nutrient  
468 concentrations were higher in Control plots at the CF site. Based on comparisons with a  
469 dataset that separated bark, phloem, and stemwood, it is likely that the bark samples in this  
470 study contained the phloem, which contains a significant portion of stem nutrients (Augusto  
471 et al., 2008). While the current foliage of trees tends to represent the current nutritional status,  
472 the bark is accumulated over the lifespan of the tree. P and K are highly mobile in tree tissues  
473 and are easily translocated, and Mg concentrations show similar patterns in bark tissue  
474 implying that it is also somewhat mobile (Helmisaari and Siltala, 1989). The fact that these  
475 concentrations are higher in Control plots may indicate that they had higher nutrient  
476 concentrations in the inner bark at the time of sampling or may suggest a larger portion of  
477 live inner bark. Generally, if this were the case it would be expected that foliage  
478 concentrations would show a similar pattern which they do not. While difficult to study in  
479 depth due to the small annual increment in bark tissues, it has been shown that certain  
480 nutrients (N, P, K, Ca, Mg and possibly Zn) are retranslocated from the bark, although this  
481 is likely a small overall source of nutrients (Helmisaari and Siltala, 1989; Hendrickson, 1987;  
482 Laclau et al., 2003). Thus, higher bark concentrations may indicate that these nutrients were  
483 poorly retranslocated from the outer bark before the tissue became dormant. This would  
484 suggest that the trees in the Control plots were less stressed for these nutrients over their  
485 lifetime resulting in a lower retranslocation efficiency.

486 The vegetation management treatments produced some differences in soil nutrient  
487 concentrations, although not many. Unlike other studies, results presented here are total  
488 concentrations of soil nutrients as opposed to exchangeable concentrations (with the

exception of C, which is often presented as total). Total soil nutrients concentrations are larger than exchangeable concentrations and as some of the nutrients quantified are not accessible to plants or mobile enough to leach, total soil nutrients are less likely to change due to biotic or abiotic factors. Soil N concentration was affected by treatment differently for different species and soil depths. For all species, generally, N decreased with soil depth. N is a common limiting element in these forests and this indicates that for this slow growing species, sustained vegetation control may reduce the ability of the ecosystem to retain N, as was shown by Miller et al. (2006). Concentrations of C in the soil was, generally, not affected by VM treatment. Only one species and showed higher soil C in one layer in Control plots. Across all species, soil concentrations of Ca were higher in the 0.0-0.2 m layer of the VM treated plots, although a similar study that measured exchangeable Ca in soil of 15-20 year old Douglas-fir showed the opposite trend (Harrington et al., 2020; Little et al., 2020b). The Matlock site of the LTSP displayed less C and N at both years 10 and 15, which agrees with our results, although they interpret this as due to Scotch broom infestation in the control plots (Harrington et al., 2020; Slesak et al., 2016). These studies also noted greater increases/concentrations in soil cations in plots without control of competing vegetation (although it should be noted they were measuring exchangeable cation pools and not total soil cations). Our study did not note any treatment differences in P and K concentrations, both of which were noted in Slesak et al. (2016). Another study of similar design conducted in western Washington noted no treatment differences in total soil N for all depths, but did note more C in the 0.6-1.0 m layer in herbicide treated plots (Knight et al., 2014). Additionally, this study did not note any difference in total soil P concentrations between vegetation management treatments.

The foliar nutrient concentrations measured here generally agree with published values. Moore et al. (2004) measured foliar concentrations of unfertilized GF and DF in the Intermountain West, calculating percentiles for each nutrient. DF foliar nutrient

515 concentrations in this study generally fell within the ranges published for N, P, Mn, Fe, and  
516 Cu. Measured concentrations for K, Mg, and B ranged from 40<sup>th</sup> percentile to below levels  
517 measured in the study, whereas S concentrations ranged from 80<sup>th</sup> percentile to greater than  
518 observed concentrations. Measured concentrations for two of the elements were entirely  
519 outside of these published ranges- Ca concentrations being higher than the highest reported  
520 value, and Zn concentrations being lower. These differences may be due to different nutrient  
521 availabilities in different soil types- as the measured Ca concentrations in DF foliage agree  
522 better with data from sites in Oregon (Mainwaring et al., 2014). A study of old growth DF  
523 showed similar trends for N, P, Mg, and K. However, our reported Ca values were lower,  
524 although by less than a factor of 2 (Cross and Perakis, 2011). According to a nutrient  
525 diagnosis guide for Douglas-fir in western British Columbia, there are possible deficiencies  
526 of K, Mg (at the CR site), S, B (at the CF site), Cu, Fe, and Zn (Ballard and Carter, 1986).  
527 Additionally, this reference suggested that DF at both sites were severely to slightly-  
528 moderately deficient in N, though this guide was developed for current year foliage as  
529 opposed to composite samples (Ballard and Carter, 1986).

530 Nutrient concentrations of GF were less in line with concentrations in the Intermountain  
531 West as reported by Moore et al. (2004), although GF was the most variable of the species  
532 measured. Only N, P, S, Mg, and Zn fall entirely in the reported ranges. All other nutrients  
533 fell outside the published range, with Ca, Mn, Fe, and Cu being greater and K and B being  
534 lower (Moore et al., 2004).

535 Foliar nutrients of WRC also generally agree with published literature values. Radwan  
536 and Harrington (2011) measured foliar concentrations of WRC trees sampled from a range  
537 of different sites in Washington and British Columbia, with a couple of sites in Oregon. The  
538 concentrations measured here are generally within the published range for N, P, K, Mg, and,  
539 S- although the lowest concentrations measured by this study were lower than those of  
540 Radwan and Harrington (1986). However, the Ca concentrations measured in this study were

541 almost two-fold higher than their published data. When compared to foliar concentrations  
542 from a different study in British Columbia- measured N, P, K, S, and Mg concentrations were  
543 lower than published values, whereas Ca concentrations are higher (Kranabetter et al., 2003).

544 As with the other species, most published foliar values of WH report concentrations in  
545 current year foliage. Foliar N was lower than values from old growth specimens in the coast  
546 range and stands in western Washington (Cross and Perakis, 2011; Radwan and DeBell,  
547 1980). Concentrations of P, however, were higher than those reported for old growth  
548 specimens, slightly higher than coastal stands reported by Radwan and DeBell (1980), but  
549 fitting with stands in the Cascades. Ca values, as with other species, were higher than other  
550 published values (Cross and Perakis, 2011; Kranabetter et al., 2003; Radwan and DeBell,  
551 1980).

552 Soil concentrations of C, N, and P are in line with other studies in the Oregon Coast  
553 Range (Cromack et al., 1999; Cross and Perakis, 2011). Concentrations of C and N from both  
554 sites are similar to the STR and CTC sites in Mainwaring et al. (2014), which are  
555 geographically very close to the CR and CF sites respectively. Soil concentrations of Cu, Mn  
556 and Zn are in or near the ranges predicted by the USGS, with Cu and Zn concentrations  
557 slightly lower than the predicted ranges. Concentrations of Ca, K and Mg are lower than  
558 USGS predictions by approximately an order of magnitude. Measurements of Ca in soil  
559 residue (<2 mm) in the Oregon Coast Range averaged 0.25% on sedimentary bedrock to  
560 0.77% on basaltic bedrock (Hynicka et al., 2016). These values are only two-fold higher than  
561 the 0.13% average at the CR site (located on sedimentary bedrock) and 0.35% at the CF site  
562 (located on basaltic bedrock). It should be noted that the Basaltic bedrock sites in Hynicka et  
563 al. (2016) were from basaltic sites in the Oregon Coast range and not in the Cascade foothills.

564

565 Differences in nutrient concentration between site varied by nutrient and tissue type.  
566 Similar trends were noticed for both species, although DF displayed more site dependent

567 nutrient differences. Most of the differences in tissue nutrient concentration were associated  
568 with differences in total soil nutrient concentration. Generally, soils at the CR site had higher  
569 concentrations of K, Mg, and Na while soils at the CF site had higher concentrations of Ca,  
570 B, Cu, Fe, and Mn. When there were differences in tissue concentrations, they generally  
571 followed similar trends, with the exception of branch Cu and bark Ca in WRC as well as  
572 forest floor and foliage K in DF. This suggests that, while the soil nutrients measured were  
573 total concentrations as opposed to accessible concentrations, they may be indicative of trends  
574 in available concentrations between sites.

575 Differences in parent material are able to explain some of the soil concentration  
576 differences between the two sites. Basaltic rocks tend to have higher concentrations of Fe,  
577 Mg, and Ca than sedimentary rock, although this can change depending upon the nature of  
578 the sedimentary material. This study found that there were higher soil concentrations of Fe  
579 at the CF site which is more volcanic, but less Mg. It is possible that this is due to the nature  
580 of sedimentary rock at the CR site or land use history at the CF site. The CF site was  
581 previously agricultural land that was relatively low yielding. It may be that farming  
582 procedures decreased soil Mg. It has been shown that application of lime in the form of Ca  
583 carbonate depletes the exchangeable Mg, although this may only be a small portion of the  
584 total Mg at a site. Additionally, studies of soils formed on the Tyee formation (which the CR  
585 site is located on) show that these sites contain a large amount of montmorillonite, a clay  
586 which commonly has Mg isomorphous substitutions in the A1 layer (McBride, 1994;  
587 McWilliams, 1973; Metson, 1974).

588 P is almost entirely sourced from bedrock, with soil reserves declining with age. The  
589 bedrock from the Tyee formation formed in the middle Eocene, somewhere between 54 and  
590 36 Ma. The bedrock that the CF site is located on is estimated to be between 32 and 11 Ma  
591 in various parts of the range. Additionally, the Oregon Coast Range (CR site) generally  
592 experiences greater rainfall and higher biomass production than the West Cascades (CF site)

593 (Hudiburg et al., 2009). Both plant activity and moisture are important soil forming factors.  
594 Given this information it is reasonable to suspect that soils at the CR site are more developed  
595 which may have resulted in less soil P than the CF site.

596 Soil K levels in the PNW are low compared to the rest of the country due to a lack of K  
597 feldspar in the parent material. According to the USGS, concentrations near the study sites  
598 should range from 0.8 to 1.2% in the top 0.05 m and A horizon, although soil at 1 m depth  
599 by the CR site may have lower concentrations (Smith et al., 2019). Cu concentrations are  
600 high in the areas near both sites, ranging from 30 to 300 ppm or more in the top meter of soil  
601 (Smith et al., 2019). Soil Mn is high, ranging from 880-1210 ppm through A horizon, with  
602 samples at 1 m depth have higher concentrations near the CR site (Smith et al., 2019). Zn  
603 concentrations are also high, ranging from 80-100 ppm at both sites with possible higher  
604 concentration in the A horizon of the CF site. Soil Fe concentrations are also high in Oregon,  
605 ranging from 3 ppm to 14 ppm (Smith et al., 2019). Concentrations of Mg near the CF site  
606 range from 1 to 13% in the top 0.05 m and A horizon, whereas they range from 0.7 to 1.2 %  
607 near the CR site (Smith et al., 2019).

608 Species differences in concentrations were more common than treatment differences and  
609 showed notably different, but expected, patterns when compared to site differences. Species  
610 differences in soil concentration were most common in the top 0.2 m, which is to be expected  
611 as this is where the greatest quantity of fine roots are found. The species effect was significant  
612 across all species for 5 nutrients (Table 4). However, when comparing one species to another,  
613 these trends were often not significant (Figure 3). Lower soil C for DF may reflect a lower  
614 rate of fine root turnover or a higher rate of microbial respiration. Mg generally had the lowest  
615 concentrations under WH. This may indicate that there is greater uptake or leaching of this  
616 nutrient under this species. Even although root samples are a composite of fine roots from all  
617 vegetation within each plot, the higher concentrations of Zn, B, and Mn in fine roots of WRC  
618 may suggest that WRC invests more micronutrients to fine roots than the other species.

619 It is difficult to draw general trends for species differences in aboveground tissue  
620 concentrations. Elements such as B and Zn did not have strong trends that indicate the  
621 tendency of one species to accumulate more of a nutrient across all tissue types. Similarly,  
622 no tissue type tended to have higher concentrations of all or most nutrients in any given  
623 species. Mn had significantly higher tissue concentrations in the stemwood, bark, branches,  
624 and foliage of WH, which indicates that this species may accumulate more Mn than other  
625 species. WH, as a species, is capable of growing at lower soil pH than other conifers and soil  
626 Mn becomes more available at lower pH. The trend observed here may indicate that WH has  
627 adapted to survive with higher tissue concentrations of Mn due to its preference for acidic  
628 soils. Concentrations of P were highest for stemwood, bark and foliage of WH. This differs  
629 from old growth species in the Oregon Coast Range which showed DF species as having not  
630 significantly higher foliar concentrations than WH (Cross and Perakis, 2011). A study of  
631 WRC and WH in coastal British Columbia showed no differences across species on a number  
632 of different site types (Kranabetter et al., 2003)

633 **Conclusions**

634 Effects of VM on nutrient concentrations of plant derived tissue at ages 16-18 varied by  
635 site, species, nutrient, and tissue. Bark and forest floor were the two tissue types that were  
636 most sensitive to VM treatment. Differences in forest floor nutrient concentrations are likely  
637 driven by the changes in plant species composition between VM and Control plots, with  
638 midstory and understory species contributing chemically distinct litter in many Control plots.  
639 Differences in bark concentrations may indicate differences in nutrient retranslocation over  
640 the lives of the different stands. Since the treatment had little effect on foliar nutrient  
641 concentrations, we expect the physiology, including photosynthetic efficiency of the foliage,  
642 to also be similar between competing vegetation control treatments. This means that crop tree

643 growth differences between Control and VM treatments cannot be explained by the foliar  
644 nutrient status at ages 16-18.

645 Few treatment effects on soil were discovered and varied by species, site, and depth.  
646 When differences were detectable, soil concentrations of N and Mg were higher in VM plots.  
647 The one exception was that soil N and Ca concentrations for WRC at the CR site were  
648 significantly lower for 0.2-0.4 m and 0.4-0.6 m depth increments in VM plots. Additionally,  
649 deep soil C (0.4-0.6 m) showed a significant decrease under VM for DF at the CR site.  
650 Generally, tissue concentrations were most affected by species and soil concentrations were  
651 most affected by site. This study does not indicate the potential for total soil nutrient reserves  
652 to be depleted by even sustained vegetation management treatment. WRC at the CR site was  
653 a notable exception, where VM plots showed significantly lower N concentrations. This may  
654 indicate the potential for reduced N retention on a slow growing species, such as WRC,  
655 receive five years of post-planting herbicide application. This study did not attempt to  
656 quantify fluxes between various available and unavailable soil nutrient pools, and as such  
657 there may be treatment differences in nutrient availability that cannot be observed from this  
658 data.

659

660 **Funding:** This research was supported by the Starker Forests Inc, Cascade Timber  
661 Consulting Inc., the Oregon State University Forest Engineering, Resources and  
662 Management Department and the Vegetation Management Research Cooperative at Oregon  
663 State University. Callan Cannon's graduate education was funded in part by the Konnie  
664 Family Forest Engineering Fellowship and the Lee Harris Memorial Fellowship.

665 **Acknowledgments:** We want to acknowledge all people who helped to install the study,  
666 especially Dr. Robin Rose, Mr. Mark Gourley and Mr. Bill Marshall. Special thanks to Gloria  
667 Ambrowiak, Adam Fund, and Marcus Kleber in the Central Analytical Laboratory at Oregon  
668 State for advice on analytical techniques, and help with chemical analysis. The authors would

669 also like to thank Jeff Hatten and Adrian Gallo with help designing soil sampling regimes  
670 and lending equipment.

671

672 **Conflicts of Interest:** The authors declare no conflict of interest

673 **References**

674 Ågren, G.I., 2008. Stoichiometry and nutrition of plant growth in natural communities. *Annu.*  
675 *Rev. Ecol. Evol. Syst.* 39, 153–170.  
676 <https://doi.org/10.1146/annurev.ecolsys.39.110707.173515>

677 Augusto, L., Meredieu, C., Bert, D., Trichet, P., Porté, A., Bosc, A., Lagane, F., Loustau, D.,  
678 Pellerin, S., Danjon, F., Ranger, J., Gelpe, J., 2008. Improving models of forest nutrient  
679 export with equations that predict the nutrient concentration of tree compartments. *Ann.*  
680 *For. Sci.* 65. <https://doi.org/10.1051/forest:2008059>

681 Ballard, T.M., Carter, R.E., 1986. Evaluating forest stand nutrient status, *Land Manage. Rep.*  
682 20. Victoria, BC.

683 Beaufils, E.R., 1973. Diagnosis and recommendation integrated system (DRIS). *Soil Sci.*  
684 *Bull.* 1, 1–132.

685 Burger, J.A., Pritchett, W.L.L., 1988. Site preparation effects on soil moisture and available  
686 nutrients in a pine plantation in the Florida flatwoods. *For. Sci.* 34, 77–87.

687 Chen, F.-H., 2004. Effects of Weed Control on Vegetation Dynamics in Pacific Northwest  
688 Conifer Plantations.

689 Cole, D.W., Gessel, S.P., 1992. Fundamentals of Tree Nutrition, in: Chappell, H.N.,  
690 Weetman, G.F., Miller, R.E. (Eds.), *Forest Fertilization: Sustaining and Improving*

691      Nutrition and Growth of Western Forests. College of Forest Resorces, University of  
692      Washington, Seattle, pp. 7–16.

693      Cromack, K., Miller, R.E., Anderson, H.W., Helgerson, O.T., Smith, R.B., 1999. Soil Carbon  
694      and Nutrients in a Coastal Oregon Douglas-Fir Plantation with Red Alder. *Soil Sci. Soc.*  
695      *Am. J.* 63, 232–239. <https://doi.org/10.2136/sssaj1999.03615995006300010034x>

696      Cross, A., Perakis, S.S., 2011. Tree species and soil nutrient profiles in old-growth forests of  
697      the Oregon Coast Range. *Can. J. For. Res.* 41, 195–210. <https://doi.org/10.1139/x10->  
698      199

699      DeBruler, D.G., Schoenholtz, S.H., Slesak, R.A., Strahm, B.D., Harrington, T.B., 2019. Soil  
700      phosphorus fractions vary with harvest intensity and vegetation control at two  
701      contrasting Douglas-fir sites in the Pacific northwest. *Geoderma* 350, 73–83.  
702      <https://doi.org/10.1016/j.geoderma.2019.04.038>

703      Devine, W.D., Harrington, T.B., Terry, T.A., Harrison, R.B., Slesak, R.A., Peter, D.H.,  
704      Harrington, C.A., Shilling, C.J., Schoenholtz, S.H., 2011. Five-year vegetation control  
705      effects on aboveground biomass and nitrogen content and allocation in Douglas-fir  
706      plantations on three contrasting sites. *For. Ecol. Manage.* 262, 2187–2198.  
707      <https://doi.org/10.1016/j.foreco.2011.08.010>

708      Flamenco, H.N., Gonzalez-Benecke, C.A., Wightman, M.G., 2019. Long-term effects of  
709      vegetation management on biomass stock of four coniferous species in the Pacific  
710      Northwest United States. *For. Ecol. Manage.* 432, 276–285.  
711      <https://doi.org/10.1016/j.foreco.2018.09.033>

712      Harrington, T.B., Slesak, R.A., Dollins, J.P., Schoenholtz, S.H., Peter, D.H., 2020. Logging-

713 debris and vegetation-control treatments influence competitive relationships to limit 15-  
714 year productivity of coast Douglas-fir in western Washington and Oregon. *For. Ecol.*  
715 *Manage.* 473. <https://doi.org/10.1016/j.foreco.2020.118288>

716 Helmisaari, H.S., Siltala, T., 1989. Variation in nutrient concentrations of *pinus sylvestris*  
717 stems. *Scand. J. For. Res.* 4, 443–451. <https://doi.org/10.1080/02827588909382580>

718 Hendrickson, O., 1987. Winter Branch Nutrients in Northern Conifers and Hardwoods. *For.*  
719 *Sci.* 33, 1068–1074.

720 Hudiburg, T., Law, B., Turner, D.P., Campbell, J., Donato, D., Duane, M., 2009. Carbon  
721 dynamics of Oregon and Northern California forests and potential land-based carbon  
722 storage. *Ecol. Appl.* 19, 163–180. <https://doi.org/10.1890/07-2006.1>

723 Hynicka, J.D., Pett-Ridge, J.C., Perakis, S.S., 2016. Nitrogen enrichment regulates calcium  
724 sources in forests. *Glob. Chang. Biol.* 22, 4067–4079.  
725 <https://doi.org/10.1111/gcb.13335>

726 Knight, E., Footen, P., Harrison, R., Terry, T., Holub, S., 2014. Competing Vegetation  
727 Effects on Soil Carbon and Nitrogen in a Douglas-fir Plantation. *Soil Sci. Soc. Am. J.*  
728 78, S146–S151. <https://doi.org/10.2136/sssaj2013.07.0320nafsc>

729 Kranabetter, J.M., Banner, A., Shaw, J., 2003. Growth and nutrition of three conifer species  
730 across site gradients of north coastal British Columbia. *Can. J. For. Res.* 33, 313–324.  
731 <https://doi.org/10.1139/x02-188>

732 Laclau, J.P., Deleporte, P., Ranger, J., Bouillet, J.P., Kazotti, G., 2003. Nutrient dynamics  
733 throughout the rotation of Eucalyptus clonal stands in Congo. *Ann. Bot.* 91, 879–892.  
734 <https://doi.org/10.1093/aob/mcg093>

735 Littke, K.M., Harrington, T.B., Holub, S.M., Littke, W.R., Harrison, R.B., Turnblom, E.C.,

736 2020a. Douglas-fir biomass allocation and net nutrient pools 15-20 years after organic

737 matter removal and vegetation control. *Forests* 11, 1–21.

738 <https://doi.org/10.3390/F11091022>

739 Littke, K.M., Harrington, T.B., Slesak, R.A., Holub, S.M., Hatten, J.A., Gallo, A.C., Littke,

740 W.R., Harrison, R.B., Turnblom, E.C., 2020b. Impacts of organic matter removal and

741 vegetation control on nutrition and growth of Douglas-fir at three Pacific Northwestern

742 Long-Term Soil Productivity sites. *For. Ecol. Manage.* 468, 118176.

743 <https://doi.org/10.1016/j.foreco.2020.118176>

744 Mainwaring, D.B., Maguire, D.A., Perakis, S.S., 2014. Three-year growth response of young

745 Douglas-fir to nitrogen, calcium, phosphorus, and blended fertilizers in Oregon and

746 Washington. *For. Ecol. Manage.* 327, 178–188.

747 <https://doi.org/10.1016/j.foreco.2014.05.005>

748 McBride, M.B., 1994. Environmental chemistry of soils. Oxford University Press, New York

749 (USA).

750 McWilliams, R.G., 1973. Stratigraphic and biostratigraphic relationships of the Tyee and

751 Yamhill Formations in central-western Oregon, in: The Ore Bin. regon Department of

752 Geology and Mineral Industries, Cambridge, pp. 169–186.

753 <https://doi.org/10.1017/CBO9781107415324.004>

754 Metson, A.J., 1974. Some factors governing the availability of soil magnesium: A review.

755 *New Zeal. J. Exp. Agric.* 2, 277–319. <https://doi.org/10.1080/03015521.1974.10427689>

756 Miller, J.H., Allen, H.L., Zutter, B.R., Zedaker, S.M., Newbold, R.A., 2006. Soil and pine

757 foliage nutrient responses 15 years after competing-vegetation control and their  
758 correlation with growth for 13 loblolly pine plantations in the southern United States.  
759 Can. J. For. Res. 36, 2412–2425. <https://doi.org/10.1139/x06-164>

760 Moore, J.A., Mika, P.G., Shaw, T.M., Garrison-Johnston, M.I., 2004. Foliar nutrient  
761 characteristics of four conifer species in the interior Northwest United States. West. J.  
762 Appl. For. 19, 13–24. <https://doi.org/10.1093/wjaf/19.1.13>

763 Powers, R.F., Reynolds, P.E., 1999. Ten-year responses of ponderosa pine plantations to  
764 repeated vegetation and nutrient control along an environmental gradient. Can. J. For.  
765 Res. 29, 1027–1038. <https://doi.org/10.1139/x99-104>

766 Powers, R.F., Scott, D.A., Sanchez, F.G., Voldseth, R.A., Page-dumroese, D., Elioff, J.D.,  
767 Stone, D.M., 2005. The North American long-term soil productivity experiment :  
768 Findings from the first decade of research 220, 31–50.  
769 <https://doi.org/10.1016/j.foreco.2005.08.003>

770 Radwan, M.A., DeBell, D.S., 1980. Site index, growth, and foliar chemical composition  
771 relationships in western hemlock. For. Sci. 26, 283–290.

772 Radwan, M.A., Harrington, C.A., 1986. Foliar chemical concentrations, growth, and site  
773 productivity relations in western red cedar. Can. J. For. Res. 16, 1069–1075.  
774 <https://doi.org/10.1139/x86-185>

775 Rose, R., Ketchum, J.S., 2002. Interaction of vegetation control and fertilization on conifer  
776 species across the Pacific Northwest. Can. J. For. Res. 32, 136–152.  
777 <https://doi.org/10.1139/x01-180>

778 Slesak, R.A., Harrington, T.B., Peter, D.H., DeBruler, D.G., Schoenholtz, S.H., Strahm,

779 B.D., 2016. Effects of intensive management practices on 10-year Douglas-fir growth,  
780 soil nutrient pools, and vegetation communities in the Pacific Northwest, USA. For.  
781 Ecol. Manage. 365, 22–33. <https://doi.org/10.1016/j.foreco.2016.01.019>

782 Slesak, R.A., Harrington, T.B., Schoenholtz, S.H., 2010. Soil and Douglas-fir (*Pseudotsuga*  
783 *menziesii*) foliar nitrogen responses to variable logging-debris retention and competing  
784 vegetation control in the Pacific Northwest. Can. J. For. Res. 40, 254–264.  
785 <https://doi.org/10.1139/X09-188>

786 Smith, D.B., Solano, F., Woodruff, L.G., Cannon, W.F., Ellefsen, K.J., 2019. Geochemical  
787 and mineralogical maps, with interpretation, for soils of the conterminous United States.  
788 <https://doi.org/https://doi.org/10.3133/sir20175118>.

789 Stone, E.L., 1990. Boron deficiency and excess in forest trees: a review. For. Ecol. Manage.  
790 1, 49–75.

791 Turner, J., 1981. Nutrient cycling in an age sequence of western Washington douglas-fir  
792 stands. Ann. Bot. 48, 159–170. <https://doi.org/10.1093/oxfordjournals.aob.a086109>

793 Turner, J., Lambert, M.J., Gessel, S.P., 1977. Use of foliage sulphate concentration to predict  
794 response to urea application by Douglas-fir. Can. J. For. Res. 7, 476–480.  
795 <https://doi.org/https://doi.org/10.1139/x77-061>

796 Turner, J., Long, J.N., 1975. Accumulation of Organic Matter in a Series of Douglas-fir  
797 Stands. Can. J. For. Res. 5, 681–690.

798 Ulrich, A., 1952. Physiological Bases for Assessing the Nutritional Requirements of Plants.  
799 Annu. Rev. Plant Physiol. 3, 207–228.  
800 <https://doi.org/10.1146/annurev.pp.03.060152.001231>

801 Zasoski, R.J., Porada, H.J., Ryan, P.J., Greenleaf-Jenkins, J., Gessel, S.P., 1990.  
802 Observations of copper, zinc, iron and manganese status in western Washington forests.  
803 For. Ecol. Manage. 37, 7–25. [https://doi.org/10.1016/0378-1127\(90\)90043-B](https://doi.org/10.1016/0378-1127(90)90043-B)  
804  
805

806 **Appendix A**

807 Tables A1-A14: Control: no post-planting vegetation control, VM: sustained vegetation  
808 control for first 5 years post planting. Trt: Effect of vegetation management treatment; Site:  
809 Effect of site; Site x Trt: Interactive effect of treatment and site. The P-value shown is in  
810 bold if the difference in concentration was significant at  $\alpha=0.05$ .

811

812      **Table A1.** Average trees per ha (TPHA,  $\text{ha}^{-1}$ ), mean height (height, m), quadratic  
 813      mean diameter (QMD, cm), crop tree basal area ( $\text{BA}_{\text{CT}}$ ,  $\text{m}^2 \text{ ha}^{-1}$ ) and midstory basal  
 814      area ( $\text{BA}_{\text{M}}$ ,  $\text{m}^2 \text{ ha}^{-1}$ ), for 18 year-old Douglas-fir (DF), western hemlock (WH),  
 815      western redcedar (WRC), and grand fir (GF) planted stands growing under  
 816      contrasting treatments of vegetation management on sites located in the central  
 817      Coast Range (CR) and the Cascade foothills (CF) of western Oregon.

Site	Species	Treatment	TPHA ( $\text{ha}^{-1}$ )	Height (m)	QMD (cm)	$\text{BA}_{\text{CT}}$ ( $\text{m}^2 \text{ ha}^{-1}$ )	$\text{BA}_{\text{M}}$ ( $\text{m}^2 \text{ ha}^{-1}$ )
CR	DF	Control	681	17.1	8.5	25.1	0.0
		VM	725	18.1	9.2	31.0	0.0
	WH	Control	868	13.5	6.7	19.4	16.1
		VM	1032	17.2	9.0	42.6	0.0
	WRC	Control	748	6.2	4.1	7.0	29.3
		VM	967	10.7	7.0	24.0	0.7
	GF	Control	907	11.8	5.9	16.5	17.7
		VM	987	15.6	9.2	42.5	0.0
CF	DF	Control	696	14.8	7.2	18.4	4.5
		VM	718	17.1	8.9	28.5	0.0
	WRC	Control	352	8.7	6.4	7.0	2.7
		VM	935	9.6	6.3	19.1	0.0

818

819

820  
821  
822  
823  
824

**Table A2.** Concentration (ppm) of Boron (B) of tree and ecosystem components for 16-18 year-old Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) stands growing under contrasting treatments of vegetation management on sites located in the central Coast Range (CR) and Cascade Foothills (CF) of western Oregon. SE is the standard error.

Species	Tissue	CR Control		CR VM		CF Control		CF VM		P-value		
		ppm	SE	ppm	SE	ppm	SE	ppm	SE	Trt	Site	Site x Trt
<b>DF</b>	Foliage	10.91	1.93	11.62	1.18	23.10	2.61	22.50	2.85	0.973	<b>0.001</b>	0.797
	Branches	9.90	1.82	9.69	0.54	10.69	1.19	10.85	1.93	0.905	0.548	0.818
	Bark	7.50	0.05	7.48	0.45	8.18	0.69	7.35	0.31	0.313	0.484	0.266
	Wood	1.72	0.14	1.66	0.12	2.52	0.69	1.96	0.19	0.421	0.162	0.511
	Understory	19.12	3.23	15.20	2.46	41.66	7.80	38.09	5.53	0.225	<b>0.006</b>	0.952
	Forest Floor	11.99	0.44	13.10	0.41	18.41	1.44	16.37	1.22	0.622	<b>0.001</b>	0.118
	Fine Roots	11.36	0.76	12.49	1.35	17.23	0.57	15.49	1.35	0.764	<b>0.002</b>	0.180
	Soil 0.0-0.2 m	31.00	1.54	33.55	2.03	68.33	5.28	61.98	3.67	0.575	< <b>0.001</b>	0.225
	Soil 0.2-0.4 m	36.97	1.13	39.07	2.62	77.37	7.35	80.97	1.70	0.486	< <b>0.001</b>	0.852
	Soil 0.4-0.6 m	39.87	1.57	38.51	4.63	80.37	6.93	78.39	2.85	0.716	< <b>0.001</b>	0.946
	Soil 0.6-1.0 m	34.42	3.03	34.87	2.70	89.67	19.77	91.31	7.43	0.924	< <b>0.001</b>	0.957
<b>WH</b>	Foliage	22.99	4.66	21.52	1.82	-	-	-	-	0.772	-	-
	Branches	10.33	0.59	9.93	1.14	-	-	-	-	0.692	-	-
	Bark	11.57	2.01	8.72	0.31	-	-	-	-	0.211	-	-
	Wood	2.75	0.22	2.01	0.02	-	-	-	-	<b>0.015</b>	-	-
	Understory	22.64	4.80	16.54	1.89	-	-	-	-	0.282	-	-
	Forest Floor	11.91	0.27	14.68	1.30	-	-	-	-	0.082	-	-
	Fine Roots	12.14	1.26	8.74	0.49	-	-	-	-	<b>0.046</b>	-	-
	Soil 0.0-0.2 m	33.44	1.55	34.01	1.41	-	-	-	-	0.477	-	-
	Soil 0.2-0.4 m	45.46	4.24	41.49	1.44	-	-	-	-	0.268	-	-
	Soil 0.4-0.6 m	45.09	0.92	45.17	1.82	-	-	-	-	0.970	-	-
	Soil 0.6-1.0 m	44.22	2.65	40.31	2.49	-	-	-	-	0.356	-	-
<b>WRC</b>	Foliage	11.89	2.35	11.50	1.03	12.81	0.27	13.36	1.30	0.954	0.355	0.751
	Branches	8.40	1.47	8.63	0.63	7.59	1.73	8.41	1.77	0.726	0.731	0.844
	Bark	15.46	2.15	14.34	2.86	12.98	1.41	15.25	1.33	0.783	0.705	0.420
	Wood	2.71	0.14	2.86	0.41	3.27	0.15	3.03	0.56	0.448	0.184	0.315
	Understory	15.22	3.04	20.71	4.39	40.55	9.78	50.86	15.73	0.496	<b>0.033</b>	0.833
	Forest Floor	17.85	0.59	13.23	3.28	14.52	2.77	14.78	3.67	0.492	0.777	0.444
	Fine Roots	14.13	1.84	12.47	0.93	20.45	2.55	17.28	1.77	0.152	0.071	0.622
	Soil 0.0-0.2 m	31.93	1.86	33.28	0.99	85.15	5.95	88.32	8.50	0.276	< <b>0.001</b>	0.645
	Soil 0.2-0.4 m	36.82	1.11	39.02	5.06	88.39	6.71	78.68	5.89	0.413	< <b>0.001</b>	0.215
	Soil 0.4-0.6 m	44.92	0.00	43.44	1.70	76.37	7.20	80.35	6.83	0.672	< <b>0.001</b>	0.369
	Soil 0.6-1.0 m	47.37	4.16	40.43	3.88	77.81	2.86	76.73	13.98	0.643	<b>0.014</b>	0.733
<b>GF</b>	Foliage	15.36	1.70	15.95	3.79	-	-	-	-	0.925	-	-
	Branches	11.81	0.95	17.37	4.21	-	-	-	-	0.245	-	-
	Bark	13.15	2.70	10.61	1.38	-	-	-	-	0.282	-	-
	Wood	2.73	0.73	2.94	0.49	-	-	-	-	0.822	-	-
	Understory	26.54	6.63	17.11	3.28	-	-	-	-	0.271	-	-
	Forest Floor	14.31	1.22	12.49	0.83	-	-	-	-	0.285	-	-
	Fine Roots	13.50	2.28	10.29	0.92	-	-	-	-	0.263	-	-
	Soil 0.0-0.2 m	23.79	5.27	29.40	1.21	-	-	-	-	0.358	-	-
	Soil 0.2-0.4 m	28.73	3.51	39.27	3.23	-	-	-	-	0.092	-	-
	Soil 0.4-0.6 m	40.80	10.37	41.39	1.74	-	-	-	-	0.953	-	-
	Soil 0.6-1.0 m	33.69	9.36	40.34	4.58	-	-	-	-	0.442	-	-

825  
826  
827

**Table A3.** Concentration (%) of carbon (C) of tree and ecosystem components for 16-18 year-old Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) stands growing under contrasting treatments of vegetation management on sites located in the central Coast Range (CR) and Cascade Foothills (CF) of western Oregon. SE is the standard error.

Species	Tissue	CR Control		CR VM		CF Control		CF VM		P-value		
		%	SE	%	SE	%	SE	%	SE	Trt	Site	Site x Trt
DF	Foliage	49.20	0.23	49.26	0.57	50.00	0.04	49.38	0.25	0.439	0.267	0.310
	Branches	47.32	0.24	47.45	0.11	47.07	0.19	47.10	0.25	0.700	0.172	0.800
	Bark	48.32	0.74	49.21	1.11	49.74	0.49	49.63	0.40	0.702	0.182	0.595
	Wood	47.77	0.08	47.57	0.18	47.76	0.26	48.03	0.08	0.827	0.234	0.190
	Understory	37.22	5.54	31.50	1.87	42.31	1.72	44.34	0.76	0.559	<b>0.013</b>	0.231
	Forest Floor	31.22	4.08	35.34	2.26	41.64	1.97	41.37	3.06	0.527	<b>0.017</b>	0.472
	Fine Roots	31.74	2.06	30.57	2.01	23.86	2.82	26.85	1.32	0.676	<b>0.027</b>	0.350
	Soil 0.0-0.2 m	3.80	0.12	4.20	1.04	4.26	0.46	4.53	0.36	0.584	0.521	0.925
	Soil 0.2-0.4 m	2.87	0.58	2.43	0.49	2.80	0.53	3.14	0.49	0.849	<b>0.038</b>	0.168
	Soil 0.4-0.6 m	2.37	0.24	1.09	0.26	0.82	0.14	1.36	0.11	0.102	<b>0.012</b>	<b>0.002</b>
	Soil 0.6-1.0 m	0.65	0.23	0.82	0.02	0.54	0.12	0.81	0.13	0.166	0.945	0.711
WH	Foliage	49.22	0.53	49.41	0.19	-	-	-	-	0.719	-	-
	Branches	46.68	0.10	46.33	0.18	-	-	-	-	0.060	-	-
	Bark	47.08	1.06	45.18	0.45	-	-	-	-	0.152	-	-
	Wood	47.63	0.41	47.85	0.13	-	-	-	-	0.544	-	-
	Understory	42.05	1.25	42.19	0.76	-	-	-	-	0.928	-	-
	Forest Floor	40.93	1.70	41.01	2.27	-	-	-	-	0.976	-	-
	Fine Roots	32.73	3.51	35.08	0.93	-	-	-	-	0.542	-	-
	Soil 0.0-0.2 m	4.64	0.42	5.28	0.44	-	-	-	-	0.324	-	-
	Soil 0.2-0.4 m	2.75	0.89	2.64	0.22	-	-	-	-	0.906	-	-
	Soil 0.4-0.6 m	1.11	0.22	1.92	0.54	-	-	-	-	0.234	-	-
	Soil 0.6-1.0 m	0.71	0.16	0.61	0.14	-	-	-	-	0.659	-	-
WRC	Foliage	48.92	0.47	48.07	0.56	48.89	0.36	48.81	0.23	0.288	0.415	0.378
	Branches	46.97	0.11	46.69	0.12	46.06	0.24	42.87	3.87	0.353	0.288	0.429
	Bark	48.35	0.19	48.95	0.63	47.04	0.60	47.15	0.37	0.302	<b>0.015</b>	0.338
	Wood	48.62	0.04	48.52	0.10	46.97	0.95	46.88	0.80	0.793	<b>0.001</b>	0.592
	Understory	43.59	0.20	36.95	5.72	41.62	1.83	44.05	0.55	0.437	0.439	0.124
	Forest Floor	39.32	0.95	38.15	4.02	42.55	4.35	39.15	4.44	0.368	0.434	0.650
	Fine Roots	34.52	4.10	29.84	1.72	24.28	2.88	23.19	0.73	0.290	<b>0.026</b>	0.490
	Soil 0.0-0.2 m	5.29	0.88	4.62	0.17	4.75	0.35	4.85	0.69	0.646	0.799	0.535
	Soil 0.2-0.4 m	3.91	1.13	2.71	0.43	3.16	0.47	2.66	0.57	0.087	0.250	0.419
	Soil 0.4-0.6 m	2.23	0.44	1.33	0.25	1.68	0.39	0.78	0.12	<b>0.018</b>	0.114	0.998
	Soil 0.6-1.0 m	0.62	0.17	0.50	0.08	0.74	0.06	0.56	0.08	0.156	0.374	0.745
GF	Foliage	48.73	0.13	48.61	0.23	-	-	-	-	0.650	-	-
	Branches	46.70	0.31	45.99	0.30	-	-	-	-	0.153	-	-
	Bark	46.81	0.55	46.61	0.59	-	-	-	-	<b>0.044</b>	-	-
	Wood	47.28	0.12	47.44	0.16	-	-	-	-	0.374	-	-
	Understory	44.33	0.13	40.32	1.38	-	-	-	-	0.099	-	-
	Forest Floor	34.17	3.39	36.81	1.88	-	-	-	-	0.534	-	-
	Fine Roots	27.79	2.74	26.95	2.71	-	-	-	-	0.839	-	-
	Soil 0.0-0.2 m	6.71	1.42	5.53	0.34	-	-	-	-	0.462	-	-
	Soil 0.2-0.4 m	2.47	0.07	2.61	0.49	-	-	-	-	0.787	-	-
	Soil 0.4-0.6 m	1.55	0.21	1.70	0.49	-	-	-	-	0.790	-	-
	Soil 0.6-1.0 m	0.78	0.26	0.51	0.13	-	-	-	-	0.415	-	-

**Table A4.** Concentration (%) of calcium (Ca) of tree and ecosystem components for 16-18 year-old Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) stands growing under contrasting treatments of vegetation management on sites located in the central Coast Range (CR) and Cascade Foothills (CF) of western Oregon. SE is the standard error.

Species	Tissue	CR Control		CR VM		CF Control		CF VM		P-value		
		%	SE	%	SE	%	SE	%	SE	Trt	Site	Site x Trt
DF	Foliage	0.561	0.108	0.573	0.090	0.627	0.046	0.641	0.069	0.878	0.426	0.994
	Branches	0.352	0.076	0.344	0.073	0.293	0.031	0.285	0.067	0.638	0.532	0.752
	Bark	0.324	0.040	0.228	0.055	0.355	0.071	0.294	0.042	0.121	0.228	0.717
	Wood	0.086	0.047	0.036	0.001	0.045	0.006	0.036	0.000	0.232	0.402	0.409
	Understory	0.649	0.114	0.606	0.055	1.216	0.312	1.037	0.121	0.545	<b>0.016</b>	0.712
	Forest Floor	0.707	0.065	0.749	0.038	0.987	0.054	0.996	0.061	0.660	<b>0.003</b>	0.774
	Fine Roots	0.387	0.045	0.320	0.037	0.679	0.059	0.521	0.028	<b>0.024</b>	<b>&lt;0.001</b>	0.322
	Soil 0.0-0.2 m	0.151	0.034	0.133	0.022	0.371	0.049	0.378	0.034	0.878	<b>&lt;0.001</b>	0.748
	Soil 0.2-0.4 m	0.108	0.030	0.113	0.048	0.339	0.057	0.376	0.019	0.512	<b>&lt;0.001</b>	0.606
	Soil 0.4-0.6 m	0.103	0.037	0.062	0.016	0.231	0.047	0.221	0.034	0.369	<b>0.038</b>	0.587
	Soil 0.6-1.0 m	0.048	0.013	0.041	0.008	0.183	0.054	0.141	0.049	0.336	0.118	0.479
WH	Foliage	0.577	0.120	0.761	0.092	-	-	-	-	0.295	-	-
	Branches	0.252	0.042	0.265	0.022	-	-	-	-	0.790	-	-
	Bark	0.347	0.027	0.439	0.022	-	-	-	-	<b>0.040</b>	-	-
	Wood	0.085	0.014	0.066	0.004	-	-	-	-	0.232	-	-
	Understory	0.803	0.068	0.747	0.056	-	-	-	-	0.565	-	-
	Forest Floor	0.759	0.059	0.749	0.061	-	-	-	-	0.905	-	-
	Fine Roots	0.451	0.023	0.367	0.048	-	-	-	-	0.166	-	-
	Soil 0.0-0.2 m	0.106	0.018	0.116	0.022	-	-	-	-	0.736	-	-
	Soil 0.2-0.4 m	0.080	0.034	0.076	0.011	-	-	-	-	0.927	-	-
	Soil 0.4-0.6 m	0.083	0.032	0.057	0.006	-	-	-	-	0.466	-	-
	Soil 0.6-1.0 m	0.034	0.002	0.040	0.003	-	-	-	-	0.184	-	-
WRC	Foliage	1.138	0.158	1.265	0.103	1.446	0.152	1.350	0.142	0.950	0.339	0.609
	Branches	0.563	0.095	0.666	0.074	0.774	0.121	0.493	0.031	0.337	0.846	0.052
	Bark	1.230	0.077	1.078	0.047	0.936	0.079	0.790	0.053	<b>0.042</b>	<b>0.001</b>	0.959
	Wood	0.125	0.003	0.210	0.082	0.124	0.003	0.107	0.006	0.369	0.194	0.251
	Understory	0.896	0.086	0.785	0.018	1.313	0.139	1.106	0.128	0.206	<b>0.011</b>	0.690
	Forest Floor	1.084	0.107	1.080	0.130	0.897	0.165	1.411	0.257	0.117	0.796	0.114
	Fine Roots	0.495	0.093	0.412	0.013	0.683	0.089	0.655	0.086	0.519	<b>0.027</b>	0.753
	Soil 0.0-0.2 m	0.169	0.063	0.147	0.037	0.418	0.052	0.317	0.023	0.212	<b>0.004</b>	0.398
	Soil 0.2-0.4 m	0.137	0.009	0.082	0.022	0.342	0.044	0.260	0.044	<b>0.007</b>	<b>0.002</b>	0.414
	Soil 0.4-0.6 m	0.110	0.022	0.045	0.002	0.270	0.010	0.238	0.023	<b>0.019</b>	<b>&lt;0.001</b>	0.371
	Soil 0.6-1.0 m	0.055	0.005	0.042	0.006	0.181	0.040	0.250	0.105	0.683	<b>0.031</b>	0.550
GF	Foliage	1.254	0.192	1.156	0.075	-	-	-	-	0.836	-	-
	Branches	0.438	0.074	0.468	0.050	-	-	-	-	0.751	-	-
	Bark	0.869	0.128	0.562	0.021	-	-	-	-	0.055	-	-
	Wood	0.092	0.011	0.082	0.008	-	-	-	-	0.466	-	-
	Understory	0.743	0.175	0.915	0.015	-	-	-	-	0.399	-	-
	Forest Floor	1.408	0.322	1.792	0.120	-	-	-	-	0.350	-	-
	Fine Roots	0.531	0.039	0.404	0.068	-	-	-	-	0.190	-	-
	Soil 0.0-0.2 m	0.187	0.023	0.186	0.017	-	-	-	-	0.961	-	-
	Soil 0.2-0.4 m	0.108	0.017	0.110	0.037	-	-	-	-	0.950	-	-
	Soil 0.4-0.6 m	0.107	0.028	0.063	0.015	-	-	-	-	0.160	-	-
	Soil 0.6-1.0 m	0.082	0.031	0.060	0.015	-	-	-	-	0.302	-	-

841      **Table A5.** Concentration (%) of calcium (Cu) of tree and ecosystem components  
 842      for 16-18 year-old Douglas-fir (DF), western hemlock (WH), western redcedar  
 843      (WRC), and grand fir (GF) stands growing under contrasting treatments of  
 844      vegetation management on sites located in the central Coast Range (CR) and  
 845      Cascade Foothills (CF) of western Oregon. SE is the standard error.

Species	Tissue	CR Control		CR VM		CF Control		CF VM		P-value		
		%	SE	%	SE	%	SE	%	SE	Trt	Site	Site x Trt
DF	Foliage	2.821	0.296	2.760	0.171	3.125	0.199	3.135	0.382	0.977	0.214	0.990
	Branches	4.097	0.348	3.679	0.157	3.321	0.242	3.380	0.312	0.497	0.119	0.383
	Bark	3.163	0.219	3.558	0.226	3.728	0.339	3.982	0.049	0.188	0.055	0.767
	Wood	0.980	0.193	0.668	0.124	1.623	0.451	1.974	0.359	0.965	0.016	0.289
	Understory	7.299	2.644	5.558	1.081	5.588	0.282	5.629	0.629	0.574	0.587	0.556
	Forest Floor	2.591	0.368	2.717	0.307	5.497	0.597	3.991	0.268	0.050	0.001	0.027
	Fine Roots	5.472	0.462	4.780	0.504	6.205	0.272	5.060	0.211	0.015	0.204	0.462
	Soil 0.0-0.2 m	21.488	1.366	23.100	2.299	34.138	2.300	36.947	1.454	0.239	<0.001	0.739
	Soil 0.2-0.4 m	23.435	1.899	22.904	1.560	39.889	3.648	44.601	4.543	0.110	<0.001	0.055
	Soil 0.4-0.6 m	25.346	2.176	25.383	1.879	40.184	4.020	48.353	5.205	0.034	<0.001	0.036
	Soil 0.6-1.0 m	26.786	1.575	25.605	1.904	32.147	3.515	52.624	2.422	0.001	<0.001	0.001
WH	Foliage	3.219	0.415	3.482	1.004	-	-	-	-	0.817	-	-
	Branches	5.036	0.225	4.516	0.343	-	-	-	-	0.098	-	-
	Bark	4.270	0.464	3.117	0.318	-	-	-	-	0.032	-	-
	Wood	1.374	0.228	1.672	0.102	-	-	-	-	0.181	-	-
	Understory	6.412	0.734	3.327	0.262	-	-	-	-	0.022	-	-
	Forest Floor	4.497	0.644	3.441	0.448	-	-	-	-	0.227	-	-
	Fine Roots	4.689	0.332	4.168	0.265	-	-	-	-	0.164	-	-
	Soil 0.0-0.2 m	23.383	0.642	22.839	2.147	-	-	-	-	0.764	-	-
	Soil 0.2-0.4 m	25.657	1.331	25.124	2.029	-	-	-	-	0.793	-	-
	Soil 0.4-0.6 m	25.295	0.936	28.228	1.313	-	-	-	-	0.143	-	-
	Soil 0.6-1.0 m	25.344	1.860	26.789	1.701	-	-	-	-	0.052	-	-
WRC	Foliage	3.766	0.401	3.107	0.207	4.580	0.893	3.672	0.650	0.013	0.036	0.679
	Branches	3.131	0.400	3.173	0.133	1.984	0.198	1.851	0.119	0.854	<0.001	0.723
	Bark	3.104	0.111	3.008	0.235	3.487	0.349	3.465	0.585	0.941	0.363	0.873
	Wood	1.309	0.224	1.121	0.117	1.515	0.303	1.137	0.131	0.207	0.531	0.447
	Understory	6.137	1.438	4.660	1.062	6.453	0.761	7.421	0.804	0.673	0.179	0.077
	Forest Floor	5.269	1.067	2.943	0.333	3.767	0.256	3.512	0.991	0.107	0.597	0.181
	Fine Roots	5.202	0.571	4.553	0.795	8.444	1.194	6.605	0.483	0.181	0.012	0.507
	Soil 0.0-0.2 m	25.250	0.545	25.630	0.749	37.249	1.868	37.493	1.642	0.754	<0.001	0.946
	Soil 0.2-0.4 m	28.743	1.705	27.367	0.770	43.444	2.803	42.652	2.547	0.658	<0.001	0.905
	Soil 0.4-0.6 m	29.174	1.398	29.736	1.692	47.158	3.378	46.073	2.921	0.927	<0.001	0.774
	Soil 0.6-1.0 m	28.440	0.316	30.504	2.314	47.089	4.222	39.943	7.768	0.643	0.025	0.407
GF	Foliage	3.604	0.251	3.561	0.386	-	-	-	-	0.929	-	-
	Branches	4.350	0.246	8.449	3.691	-	-	-	-	0.224	-	-
	Bark	4.917	0.619	3.371	0.556	-	-	-	-	0.113	-	-
	Wood	1.492	0.196	1.717	0.143	-	-	-	-	0.384	-	-
	Understory	5.797	1.227	5.459	0.006	-	-	-	-	0.797	-	-
	Forest Floor	3.588	0.204	3.166	0.345	-	-	-	-	0.101	-	-
	Fine Roots	5.211	0.727	4.452	0.550	-	-	-	-	0.445	-	-
	Soil 0.0-0.2 m	19.904	2.373	22.552	0.117	-	-	-	-	0.362	-	-
	Soil 0.2-0.4 m	22.744	1.796	25.057	0.925	-	-	-	-	0.361	-	-
	Soil 0.4-0.6 m	25.506	2.670	27.250	0.732	-	-	-	-	0.489	-	-
	Soil 0.6-1.0 m	25.516	3.262	28.778	0.993	-	-	-	-	0.316	-	-

846

847

848

849

**Table A6.** Concentration (%) of Iron (Fe) of tree and ecosystem components for 16-18 year-old Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) stands growing under contrasting treatments of vegetation management on sites located in the central Coast Range (CR) and Cascade Foothills (CF) of western Oregon. SE is the standard error.

Species	Tissue	CR Control		CR VM		CF Control		CF VM		P-value		
		ppm	SE	ppm	SE	ppm	SE	ppm	SE	Trt	Site	Site x Trt
DF	Foliage	40.374	3.729	49.468	4.865	52.809	2.079	68.790	11.367	0.079	<b>0.032</b>	0.608
	Branches	32.965	6.804	31.799	7.397	19.074	3.130	41.479	21.153	0.386	0.862	0.338
	Bark	32.919	2.797	33.206	4.116	54.570	10.560	60.521	8.608	0.675	<b>0.006</b>	0.703
	Wood	15.673	2.541	12.119	0.381	24.705	4.188	25.887	2.668	0.679	<b>0.002</b>	0.414
	Understory	746.5	321.0	1199.1	241.2	1016.8	787.2	439.7	99.0	0.874	0.865	0.212
	Forest Floor	1356	157	1244	131	1204	84	1321	206	0.986	0.809	0.465
	Fine Roots	1332	87	1313	68	1854	214	1697	66	0.493	<b>0.003</b>	0.589
	Soil 0.0-0.2 m	18052	933	18671	534	25711	546	25234	492	0.915	<0.001	0.417
	Soil 0.2-0.4 m	19436	243	19776	770	26252	432	26066	85	0.870	<0.001	0.578
	Soil 0.4-0.6 m	20096	451	19478	1125	26735	478	26931	123	0.745	<0.001	0.536
	Soil 0.6-1.0 m	19962	957	19936	769	26681	1329	27979	664	0.522	<0.001	0.505
WH	Foliage	42.439	6.768	67.957	29.388	-	-	-	-	0.430	-	-
	Branches	25.985	3.878	33.484	3.966	-	-	-	-	0.225	-	-
	Bark	38.260	12.915	54.005	11.879	-	-	-	-	0.378	-	-
	Wood	26.125	8.471	13.569	0.725	-	-	-	-	0.217	-	-
	Understory	691.6	195.1	1070.4	146.1	-	-	-	-	0.171	-	-
	Forest Floor	837	178	991	282	-	-	-	-	0.587	-	-
	Fine Roots	1365	251	1230	90	-	-	-	-	0.629	-	-
	Soil 0.0-0.2 m	19842	228	19509	764	-	-	-	-	0.636	-	-
	Soil 0.2-0.4 m	21842	466	20587	447	-	-	-	-	<b>0.046</b>	-	-
	Soil 0.4-0.6 m	21574	433	21462	233	-	-	-	-	0.829	-	-
	Soil 0.6-1.0 m	22408	344	21730	267	-	-	-	-	0.170	-	-
WRC	Foliage	55.945	2.035	49.319	3.265	108.640	35.892	95.675	33.281	0.483	0.515	0.929
	Branches	35.320	6.059	49.661	16.892	64.994	32.138	17.001	0.599	0.408	0.893	0.127
	Bark	41.827	3.792	59.508	21.698	67.886	12.104	56.334	7.116	0.796	0.515	0.300
	Wood	17.690	4.188	46.565	28.724	18.277	3.498	22.328	1.780	0.283	0.435	0.413
	Understory	710.9	161.8	1269.6	262.2	1488.4	799.3	436.7	99.0	0.628	0.956	0.134
	Forest Floor	1009	156	1276	151	1076	448	1179	255	0.534	0.958	0.779
	Fine Roots	1045	273	1207	233	1827	162	1923	73	0.459	<b>0.009</b>	0.843
	Soil 0.0-0.2 m	18754	527	19125	216	25953	587	26381	596	0.073	<0.001	0.878
	Soil 0.2-0.4 m	20710	454	19802	1025	26331	415	26317	336	0.344	<0.001	0.358
	Soil 0.4-0.6 m	20918	241	21072	308	27077	489	26862	307	0.895	<0.001	0.435
	Soil 0.6-1.0 m	22655	1529	20397	1158	27408	407	26421	1494	0.212	<b>0.001</b>	0.613
GF	Foliage	75.395	13.104	69.281	10.755	-	-	-	-	0.796	-	-
	Branches	42.971	9.619	33.885	6.139	-	-	-	-	0.456	-	-
	Bark	100.507	48.424	96.906	48.725	-	-	-	-	0.224	-	-
	Wood	12.196	1.599	26.540	4.913	-	-	-	-	<b>0.032</b>	-	-
	Understory	524.4	347.0	1514.9	66.2	-	-	-	-	0.075	-	-
	Forest Floor	1357	199	1061	109	-	-	-	-	0.263	-	-
	Fine Roots	1365	157	1404	83	-	-	-	-	0.670	-	-
	Soil 0.0-0.2 m	16258	2412	19209	300	-	-	-	-	0.292	-	-
	Soil 0.2-0.4 m	16941	1953	20432	386	-	-	-	-	0.154	-	-
	Soil 0.4-0.6 m	18975	2910	20353	364	-	-	-	-	0.643	-	-
	Soil 0.6-1.0 m	18841	3190	21689	327	-	-	-	-	0.451	-	-

856  
857  
858  
859  
860

**Table A7.** Concentration (%) of potassium (K) of tree and ecosystem components for 16-18 year-old Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) stands growing under contrasting treatments of vegetation management on sites located in the central Coast Range (CR) and Cascade Foothills (CF) of western Oregon. SE is the standard error.

Species	Tissue	CR Control		CR VM		CF Control		CF VM		P-value		
		%	SE	%	SE	%	SE	%	SE	Trt	Site	Site x Trt
DF	Foliage	0.580	0.042	0.446	0.026	0.750	0.056	0.654	0.069	<b>0.043</b>	<b>0.003</b>	0.720
	Branches	0.235	0.061	0.229	0.016	0.266	0.033	0.215	0.052	0.523	0.850	0.611
	Bark	0.230	0.013	0.283	0.034	0.200	0.012	0.175	0.027	0.576	0.052	0.065
	Wood	-	-	-	-	-	-	-	-	-	-	-
	Understory	1.465	0.652	0.770	0.368	1.702	0.233	1.528	0.412	0.345	0.310	0.565
	Forest Floor	0.155	0.025	0.117	0.014	0.251	0.024	0.179	0.016	<b>0.014</b>	<b>0.004</b>	0.326
	Fine Roots	0.208	0.026	0.199	0.044	0.091	0.010	0.075	0.012	0.645	<b>0.001</b>	0.906
	Soil 0.0-0.2 m	0.166	0.000	0.169	0.021	0.095	0.004	0.087	0.006	0.841	< <b>0.001</b>	0.647
	Soil 0.2-0.4 m	0.160	0.020	0.145	0.019	0.081	0.009	0.069	0.005	0.345	< <b>0.001</b>	0.900
	Soil 0.4-0.6 m	0.164	0.019	0.159	0.017	0.059	0.003	0.068	0.006	0.896	< <b>0.001</b>	0.581
	Soil 0.6-1.0 m	0.173	0.009	0.162	0.019	0.057	0.002	0.064	0.007	0.838	< <b>0.001</b>	0.383
WH	Foliage	0.620	0.062	0.478	0.036	-	-	-	-	0.094	-	-
	Branches	0.157	0.007	0.169	0.043	-	-	-	-	0.791	-	-
	Bark	0.333	0.045	0.232	0.032	-	-	-	-	0.113	-	-
	Wood	-	-	-	-	-	-	-	-	-	-	-
	Understory	1.175	0.401	0.840	0.182	-	-	-	-	0.476	-	-
	Forest Floor	0.210	0.048	0.148	0.024	-	-	-	-	0.295	-	-
	Fine Roots	0.174	0.027	0.144	0.030	-	-	-	-	<b>0.037</b>	-	-
	Soil 0.0-0.2 m	0.153	0.015	0.135	0.006	-	-	-	-	0.289	-	-
	Soil 0.2-0.4 m	0.146	0.012	0.138	0.007	-	-	-	-	0.585	-	-
	Soil 0.4-0.6 m	0.135	0.012	0.144	0.009	-	-	-	-	0.579	-	-
	Soil 0.6-1.0 m	0.124	0.012	0.146	0.014	-	-	-	-	0.286	-	-
WRC	Foliage	0.339	0.071	0.346	0.031	0.358	0.042	0.482	0.045	0.068	0.243	0.423
	Branches	0.133	0.027	0.116	0.036	0.191	0.026	0.145	0.023	0.212	0.608	0.488
	Bark	0.162	0.041	0.140	0.017	0.146	0.020	0.170	0.030	0.995	0.468	0.933
	Wood	-	-	-	-	-	-	-	-	-	-	-
	Understory	1.024	0.488	0.790	0.370	1.147	0.206	1.540	0.171	0.641	0.762	0.112
	Forest Floor	0.431	0.148	0.160	0.025	0.185	0.070	0.149	0.087	0.122	0.186	0.226
	Fine Roots	0.177	0.016	0.129	0.010	0.108	0.011	0.078	0.010	<b>0.009</b>	<b>0.001</b>	0.469
	Soil 0.0-0.2 m	0.176	0.011	0.180	0.022	0.127	0.019	0.094	0.006	0.364	<b>0.002</b>	0.265
	Soil 0.2-0.4 m	0.171	0.005	0.166	0.013	0.082	0.011	0.086	0.011	0.967	< <b>0.001</b>	0.380
	Soil 0.4-0.6 m	0.164	0.005	0.172	0.020	0.074	0.013	0.078	0.008	0.602	< <b>0.001</b>	0.872
	Soil 0.6-1.0 m	0.182	0.036	0.201	0.036	0.079	0.018	0.061	0.010	0.990	<b>0.001</b>	0.371
GF	Foliage	0.433	0.034	0.586	0.072	-	-	-	-	0.064	-	-
	Branches	0.277	0.048	0.469	0.063	-	-	-	-	0.074	-	-
	Bark	0.371	0.071	0.240	0.032	-	-	-	-	0.140	-	-
	Wood	-	-	-	-	-	-	-	-	-	-	-
	Understory	1.263	0.254	0.627	0.063	-	-	-	-	0.082	-	-
	Forest Floor	0.153	0.021	0.111	0.017	-	-	-	-	0.068	-	-
	Fine Roots	0.130	0.015	0.142	0.018	-	-	-	-	0.438	-	-
	Soil 0.0-0.2 m	0.178	0.023	0.174	0.014	-	-	-	-	0.892	-	-
	Soil 0.2-0.4 m	0.141	0.009	0.159	0.010	-	-	-	-	0.241	-	-
	Soil 0.4-0.6 m	0.141	0.004	0.166	0.022	-	-	-	-	0.337	-	-
	Soil 0.6-1.0 m	0.157	0.017	0.196	0.037	-	-	-	-	0.396	-	-

**Table A8.** Concentration (%) of magnesium (Mg) of tree and ecosystem components for 16-18 year-old Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) stands growing under contrasting treatments of vegetation management on sites located in the central Coast Range (CR) and Cascade Foothills (CF) of western Oregon. SE is the standard error.

Species	Tissue	CR Control		CR VM		CF Control		CF VM		P-value		
		%	SE	%	SE	%	SE	%	SE	Trt	Site	Site x Trt
DF	Foliage	0.108	0.016	0.100	0.008	0.079	0.003	0.103	0.007	0.311	0.184	0.161
	Branches	0.052	0.015	0.039	0.004	0.032	0.003	0.035	0.003	0.305	0.393	0.168
	Bark	0.046	0.004	0.038	0.004	0.029	0.003	0.032	0.002	0.502	<b>0.004</b>	0.143
	Wood	0.011	0.002	0.010	0.001	0.009	0.001	0.007	0.000	0.196	0.144	0.594
	Understory	0.244	0.046	0.188	0.037	0.289	0.026	0.321	0.031	0.697	0.166	0.180
	Forest Floor	0.121	0.004	0.102	0.002	0.146	0.005	0.096	0.010	<0.001	0.051	<b>0.016</b>
	Fine Roots	0.083	0.002	0.070	0.004	0.064	0.003	0.053	0.001	<b>0.001</b>	<0.001	0.819
	Soil 0.0-0.2 m	0.287	0.009	0.287	0.013	0.146	0.009	0.157	0.015	0.650	<0.001	0.641
	Soil 0.2-0.4 m	0.296	0.009	0.297	0.007	0.153	0.010	0.172	0.010	0.290	<0.001	0.351
	Soil 0.4-0.6 m	0.304	0.012	0.303	0.008	0.143	0.011	0.155	0.019	0.505	<0.001	0.454
	Soil 0.6-1.0 m	0.315	0.013	0.306	0.015	0.142	0.011	0.145	0.026	0.866	<0.001	0.748
WH	Foliage	0.130	0.013	0.097	0.005	-	-	-	-	<b>0.042</b>	-	-
	Branches	0.035	0.003	0.031	0.005	-	-	-	-	0.258	-	-
	Bark	0.050	0.006	0.035	0.000	-	-	-	-	<b>0.048</b>	-	-
	Wood	0.017	0.002	0.013	0.000	-	-	-	-	0.125	-	-
	Understory	0.343	0.076	0.228	0.037	-	-	-	-	0.226	-	-
	Forest Floor	0.168	0.019	0.123	0.018	-	-	-	-	0.129	-	-
	Fine Roots	0.100	0.006	0.088	0.008	-	-	-	-	0.297	-	-
	Soil 0.0-0.2 m	0.257	0.006	0.258	0.006	-	-	-	-	0.816	-	-
	Soil 0.2-0.4 m	0.267	0.007	0.274	0.010	-	-	-	-	0.567	-	-
	Soil 0.4-0.6 m	0.272	0.011	0.278	0.015	-	-	-	-	0.743	-	-
	Soil 0.6-1.0 m	0.235	0.030	0.286	0.012	-	-	-	-	0.161	-	-
WRC	Foliage	0.138	0.016	0.108	0.015	0.092	0.009	0.083	0.015	0.149	0.063	0.417
	Branches	0.037	0.005	0.040	0.004	0.039	0.005	0.026	0.004	0.316	0.230	0.101
	Bark	0.064	0.006	0.049	0.005	0.049	0.004	0.046	0.004	0.079	0.065	0.223
	Wood	0.017	0.001	0.021	0.004	0.016	0.001	0.015	0.001	0.532	0.182	0.261
	Understory	0.242	0.065	0.264	0.092	0.274	0.074	0.329	0.021	0.486	0.614	0.764
	Forest Floor	0.232	0.033	0.111	0.014	0.098	0.018	0.089	0.030	<b>0.030</b>	<b>0.019</b>	0.054
	Fine Roots	0.091	0.017	0.095	0.010	0.071	0.007	0.067	0.008	0.997	<b>0.045</b>	0.741
	Soil 0.0-0.2 m	0.289	0.012	0.306	0.020	0.156	0.011	0.149	0.006	0.666	<0.001	0.344
	Soil 0.2-0.4 m	0.298	0.003	0.314	0.015	0.160	0.010	0.161	0.012	0.364	<0.001	0.456
	Soil 0.4-0.6 m	0.295	0.009	0.324	0.016	0.160	0.019	0.168	0.012	0.271	<0.001	0.526
	Soil 0.6-1.0 m	0.276	0.014	0.329	0.030	0.147	0.027	0.171	0.024	0.162	<b>0.000</b>	0.585
GF	Foliage	0.137	0.019	0.124	0.015	-	-	-	-	0.306	-	-
	Branches	0.042	0.003	0.065	0.019	-	-	-	-	0.273	-	-
	Bark	0.062	0.007	0.048	0.004	-	-	-	-	0.151	-	-
	Wood	0.016	0.002	0.016	0.001	-	-	-	-	0.850	-	-
	Understory	0.326	0.062	0.203	0.027	-	-	-	-	0.143	-	-
	Forest Floor	0.129	0.005	0.116	0.004	-	-	-	-	0.112	-	-
	Fine Roots	0.109	0.005	0.101	0.010	-	-	-	-	0.476	-	-
	Soil 0.0-0.2 m	0.262	0.016	0.306	0.010	-	-	-	-	0.134	-	-
	Soil 0.2-0.4 m	0.290	0.004	0.309	0.003	-	-	-	-	0.057	-	-
	Soil 0.4-0.6 m	0.289	0.018	0.318	0.009	-	-	-	-	0.211	-	-
	Soil 0.6-1.0 m	0.295	0.025	0.321	0.009	-	-	-	-	0.378	-	-

**Table A9.** Concentration (ppm) of manganese (Mn) of tree and ecosystem components for 16-18 year-old Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) stands growing under contrasting treatments of vegetation management on sites located in the central Coast Range (CR) and Cascade Foothills (CF) of western Oregon. SE is the standard error.

Species	Tissue	CR Control		CR VM		CF Control		CF VM		P-value		
		ppm	SE	ppm	SE	ppm	SE	ppm	SE	Trt	Site	Site x Trt
DF	Foliage	271.57	61.57	345.00	67.39	432.70	38.33	426.92	27.68	0.539	<b>0.039</b>	0.460
	Branches	129.63	71.13	106.28	28.22	121.68	7.84	120.08	8.28	0.394	0.483	0.422
	Bark	98.11	18.56	90.84	22.47	191.67	47.24	138.88	14.17	0.259	<b>0.022</b>	0.337
	Wood	11.54	1.83	14.20	2.19	33.35	5.44	23.11	4.39	0.335	<b>0.002</b>	0.113
	Understory	758.15	473.96	472.71	140.72	654.28	75.68	767.36	153.95	0.702	0.979	0.386
	Forest Floor	392.41	38.08	496.48	47.81	771.35	56.15	892.64	77.42	0.070	<0.001	0.882
	Fine Roots	178.05	29.82	203.47	19.50	632.16	106.62	516.00	40.72	0.463	<0.001	0.259
	Soil 0.0-0.2 m	1080.93	150.39	1060.01	196.26	3232.82	184.29	3249.19	473.06	0.994	<0.001	0.948
	Soil 0.2-0.4 m	870.24	175.56	804.17	165.54	2366.88	384.96	3112.45	699.82	0.426	<b>0.002</b>	0.346
	Soil 0.4-0.6 m	723.95	155.04	654.69	114.49	811.16	192.02	1887.14	343.45	<b>0.033</b>	<b>0.046</b>	<b>0.020</b>
	Soil 0.6-1.0 m	225.84	28.98	395.10	97.23	966.85	461.01	1064.67	246.35	0.625	<b>0.021</b>	0.896
WH	Foliage	799.00	168.96	1073.50	242.65	-	-	-	-	0.389	-	-
	Branches	189.01	25.44	247.77	28.19	-	-	-	-	0.152	-	-
	Bark	249.82	28.86	286.52	24.11	-	-	-	-	0.191	-	-
	Wood	51.26	27.15	93.46	7.96	-	-	-	-	0.176	-	-
	Understory	501.81	80.44	463.83	65.70	-	-	-	-	0.727	-	-
	Forest Floor	551.48	69.67	1115.21	217.07	-	-	-	-	<b>0.045</b>	-	-
	Fine Roots	235.67	20.23	209.59	9.72	-	-	-	-	0.230	-	-
	Soil 0.0-0.2 m	1062.13	167.14	1039.16	56.49	-	-	-	-	0.888	-	-
	Soil 0.2-0.4 m	923.23	201.51	940.08	92.12	-	-	-	-	0.942	-	-
	Soil 0.4-0.6 m	828.76	166.23	737.71	147.34	-	-	-	-	0.696	-	-
	Soil 0.6-1.0 m	524.78	135.60	301.76	38.43	-	-	-	-	0.165	-	-
WRC	Foliage	160.70	16.86	164.77	22.53	202.88	45.27	180.77	28.59	0.771	0.355	0.673
	Branches	37.94	5.15	45.61	12.22	54.78	8.18	32.66	1.73	0.425	0.623	0.113
	Bark	64.10	10.46	56.80	13.90	64.91	21.09	58.20	7.81	0.520	0.772	0.966
	Wood	11.51	2.55	5.12	1.29	9.01	1.79	7.44	0.70	0.128	0.810	0.297
	Understory	232.24	27.57	292.00	41.97	685.59	121.57	761.36	168.16	0.369	<b>0.013</b>	0.912
	Forest Floor	260.36	64.68	258.04	87.90	638.73	300.78	543.66	113.71	0.781	0.201	0.792
	Fine Roots	196.30	36.94	287.89	54.12	706.62	71.83	782.38	18.06	0.137	<0.001	0.881
	Soil 0.0-0.2 m	1393.03	206.25	1625.14	256.60	4086.64	682.47	4759.76	421.65	0.175	<b>0.000</b>	0.479
	Soil 0.2-0.4 m	1616.05	3.18	1409.30	245.83	3546.61	625.99	3806.35	505.71	0.924	<b>0.003</b>	0.416
	Soil 0.4-0.6 m	1298.11	204.81	1076.09	289.72	2603.45	397.55	1762.51	529.58	0.264	0.085	0.493
	Soil 0.6-1.0 m	525.45	147.26	554.84	230.62	1166.43	211.36	931.87	181.01	0.620	<b>0.029</b>	0.525
GF	Foliage	554.18	72.75	544.60	117.14	-	-	-	-	0.947	-	-
	Branches	121.21	15.72	151.75	56.15	-	-	-	-	0.619	-	-
	Bark	250.71	81.15	237.25	44.46	-	-	-	-	0.526	-	-
	Wood	38.85	8.02	31.82	8.36	-	-	-	-	0.478	-	-
	Understory	589.42	295.82	380.89	2.29	-	-	-	-	0.520	-	-
	Forest Floor	579.36	28.53	793.76	88.66	-	-	-	-	0.088	-	-
	Fine Roots	258.02	30.46	268.60	21.84	-	-	-	-	0.792	-	-
	Soil 0.0-0.2 m	1085.16	125.94	1802.15	131.35	-	-	-	-	<b>0.017</b>	-	-
	Soil 0.2-0.4 m	1047.57	145.00	1274.89	286.61	-	-	-	-	0.518	-	-
	Soil 0.4-0.6 m	757.21	230.04	1117.80	266.10	-	-	-	-	0.363	-	-
	Soil 0.6-1.0 m	437.78	201.84	449.65	143.25	-	-	-	-	0.964	-	-

876  
877  
878  
879  
880

**Table A10.** Concentration (%) of nitrogen (N) of tree and ecosystem components for 16-18 year-old Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) stands growing under contrasting treatments of vegetation management on sites located in the central Coast Range (CR) and Cascade Foothills (CF) of western Oregon. SE is the standard error.

Species	Tissue	CR Control		CR VM		CF Control		CF VM		P-value		
		%	SE	%	SE	%	SE	%	SE	Trt	Site	Site x Trt
DF	Foliage	1.174	0.044	1.348	0.267	1.232	0.061	1.255	0.028	0.495	0.901	0.598
	Branches	0.336	0.041	0.313	0.018	0.222	0.019	0.223	0.032	0.708	<b>0.004</b>	0.696
	Bark	0.338	0.044	0.313	0.042	0.256	0.017	0.270	0.036	0.874	0.109	0.594
	Wood	0.043	0.005	0.042	0.007	0.134	0.024	0.091	0.025	0.273	<b>0.007</b>	0.273
	Understory	1.344	0.378	1.034	0.034	1.378	0.134	1.638	0.091	0.902	0.181	0.181
	Forest Floor	0.938	0.201	1.080	0.074	0.968	0.025	0.958	0.078	0.571	0.719	0.511
	Fine Roots	0.647	0.027	0.611	0.066	0.580	0.054	0.525	0.017	0.291	0.114	0.824
	Soil 0.0-0.2 m	0.239	0.007	0.249	0.050	0.263	0.026	0.249	0.022	0.943	0.703	0.709
	Soil 0.2-0.4 m	0.162	0.023	0.113	0.004	0.179	0.027	0.206	0.026	0.547	<b>0.016</b>	0.061
	Soil 0.4-0.6 m	0.137	0.022	0.069	0.016	0.084	0.010	0.115	0.008	0.161	0.602	<b>0.004</b>
	Soil 0.6-1.0 m	0.046	0.003	0.061	0.004	0.048	0.008	0.066	0.009	<b>0.011</b>	0.144	0.791
WH	Foliage	1.065	0.024	1.002	0.059	-	-	-	-	0.237	-	-
	Branches	0.270	0.026	0.252	0.025	-	-	-	-	0.622	-	-
	Bark	0.283	0.049	0.320	0.020	-	-	-	-	0.506	-	-
	Wood	0.076	0.007	0.069	0.005	-	-	-	-	0.412	-	-
	Understory	1.625	0.219	1.319	0.074	-	-	-	-	0.275	-	-
	Forest Floor	1.125	0.027	0.680	0.028	-	-	-	-	<0.001	-	-
	Fine Roots	0.538	0.053	0.506	0.014	-	-	-	-	0.592	-	-
	Soil 0.0-0.2 m	0.197	0.015	0.215	0.019	-	-	-	-	0.378	-	-
	Soil 0.2-0.4 m	0.148	0.036	0.156	0.013	-	-	-	-	0.817	-	-
	Soil 0.4-0.6 m	0.084	0.010	0.127	0.030	-	-	-	-	0.224	-	-
	Soil 0.6-1.0 m	0.059	0.006	0.053	0.006	-	-	-	-	0.502	-	-
WRC	Foliage	0.982	0.059	1.137	0.218	1.037	0.213	0.757	0.070	0.700	0.327	0.196
	Branches	0.262	0.046	0.264	0.033	0.223	0.023	0.117	0.015	0.120	<b>0.012</b>	0.111
	Bark	0.234	0.019	0.221	0.004	0.304	0.021	0.291	0.041	0.618	<b>0.017</b>	0.995
	Wood	0.114	0.013	0.097	0.020	0.390	0.050	0.350	0.096	0.345	<b>0.006</b>	0.570
	Understory	1.600	0.367	1.243	0.199	1.181	0.261	1.502	0.081	0.926	0.712	0.115
	Forest Floor	1.227	0.217	0.690	0.065	0.638	0.077	0.635	0.109	0.054	<b>0.026</b>	0.056
	Fine Roots	0.589	0.078	0.555	0.065	0.545	0.026	0.530	0.011	0.597	0.466	0.833
	Soil 0.0-0.2 m	0.299	0.029	0.223	0.019	0.240	0.022	0.244	0.010	0.107	0.359	0.077
	Soil 0.2-0.4 m	0.196	0.038	0.155	0.014	0.198	0.033	0.162	0.027	<b>0.041</b>	0.362	0.851
	Soil 0.4-0.6 m	0.140	0.024	0.089	0.016	0.128	0.027	0.078	0.006	<b>0.034</b>	0.580	0.998
	Soil 0.6-1.0 m	0.063	0.012	0.051	0.001	0.078	0.010	0.053	0.009	0.070	0.284	0.436
GF	Foliage	1.150	0.085	1.072	0.047	-	-	-	-	0.525	-	-
	Branches	0.281	0.045	0.476	0.170	-	-	-	-	0.312	-	-
	Bark	0.408	0.045	0.288	0.047	-	-	-	-	0.116	-	-
	Wood	0.096	0.010	0.086	0.020	-	-	-	-	0.708	-	-
	Understory	1.307	0.285	1.323	0.107	-	-	-	-	0.959	-	-
	Forest Floor	1.033	0.154	0.977	0.201	-	-	-	-	0.834	-	-
	Fine Roots	0.637	0.119	0.557	0.040	-	-	-	-	0.560	-	-
	Soil 0.0-0.2 m	0.322	0.064	0.262	0.011	-	-	-	-	0.411	-	-
	Soil 0.2-0.4 m	0.156	0.004	0.167	0.032	-	-	-	-	0.753	-	-
	Soil 0.4-0.6 m	0.105	0.010	0.107	0.026	-	-	-	-	0.965	-	-
	Soil 0.6-1.0 m	0.067	0.014	0.050	0.006	-	-	-	-	0.315	-	-

881

882

883  
884  
885  
886  
887

**Table A11.** Concentration (ppm) of sodium (Na) of tree and ecosystem components for 16-18 year-old Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) stands growing under contrasting treatments of vegetation management on sites located in the central Coast Range (CR) and Cascade Foothills (CF) of western Oregon. SE is the standard error.

Species	Tissue	CR Control		CR VM		CF Control		CF VM		P-value		
		ppm	SE	ppm	SE	ppm	SE	ppm	SE	Trt	Site	Site x Trt
DF	Foliage	148.58	25.34	229.92	41.50	94.07	12.77	70.10	3.90	0.277	<b>0.001</b>	0.059
	Branches	47.097	20.326	33.214	4.105	2.848	3.269	6.462	7.574	0.619	<b>0.018</b>	0.417
	Bark	252.50	122.42	102.55	3.745	28.500	4.201	41.981	8.322	0.288	0.073	0.212
	Wood	-	-	-	-	-	-	-	-	-	-	-
	Understory	205.89	48.98	171.86	25.79	177.92	30.71	148.38	34.07	0.309	0.460	0.941
	Forest Floor	177.49	13.68	160.98	8.77	133.90	1.00	128.11	12.63	0.193	<b>0.005</b>	0.511
	Fine Roots	157.85	28.11	139.01	8.50	92.85	13.78	80.67	7.66	0.232	<b>0.005</b>	0.786
	Soil 0.0-0.2 m	173.16	14.92	186.22	5.97	122.03	2.45	124.80	3.52	0.360	< <b>0.001</b>	0.548
	Soil 0.2-0.4 m	171.23	8.21	173.98	4.29	174.77	17.92	185.24	21.00	0.544	0.200	0.720
	Soil 0.4-0.6 m	217.99	12.33	255.96	15.92	137.91	3.83	135.33	9.93	0.055	< <b>0.001</b>	<b>0.034</b>
	Soil 0.6-1.0 m	161.13	6.46	174.59	5.52	123.29	7.92	116.91	6.87	0.469	< <b>0.001</b>	0.066
WH	Foliage	117.29	10.84	142.83	36.38	-	-	-	-	0.526	-	-
	Branches	1.778	4.232	6.563	2.151	-	-	-	-	0.165	-	-
	Bark	89.351	21.525	76.812	8.700	-	-	-	-	0.610	-	-
	Wood	-	-	-	-	-	-	-	-	-	-	-
	Understory	246.05	42.27	191.50	31.67	-	-	-	-	0.341	-	-
	Forest Floor	163.11	11.97	160.60	3.07	-	-	-	-	0.845	-	-
	Fine Roots	119.53	8.76	130.86	6.55	-	-	-	-	0.211	-	-
	Soil 0.0-0.2 m	153.37	3.94	158.00	7.24	-	-	-	-	0.595	-	-
	Soil 0.2-0.4 m	187.92	9.38	204.82	15.36	-	-	-	-	0.384	-	-
	Soil 0.4-0.6 m	188.64	9.19	189.55	15.14	-	-	-	-	0.954	-	-
	Soil 0.6-1.0 m	149.90	1.42	149.69	9.37	-	-	-	-	0.983	-	-
WRC	Foliage	130.18	36.39	96.33	5.04	66.37	29.55	66.15	9.66	0.409	0.122	0.416
	Branches	14.398	14.133	11.174	2.879	2.072	6.913	-7.412	2.992	0.450	0.082	0.707
	Bark	63.242	6.253	55.987	5.957	22.849	4.542	20.074	5.938	0.397	< <b>0.001</b>	0.702
	Wood	-	-	-	-	-	-	-	-	-	-	-
	Understory	164.66	35.07	237.94	68.48	171.04	46.74	179.00	55.39	0.247	0.111	0.338
	Forest Floor	203.49	19.58	164.50	3.94	94.85	8.27	101.09	8.60	0.164	< <b>0.001</b>	0.074
	Fine Roots	180.45	58.79	132.43	4.16	131.37	31.66	78.64	7.98	0.126	0.193	0.937
	Soil 0.0-0.2 m	156.19	19.24	158.06	7.61	132.26	9.62	136.84	9.04	0.786	0.116	0.909
	Soil 0.2-0.4 m	229.57	16.06	212.81	22.44	189.91	12.14	235.59	11.64	0.365	0.592	0.068
	Soil 0.4-0.6 m	190.92	12.75	169.88	6.10	133.48	5.52	135.68	9.11	0.301	< <b>0.001</b>	0.209
	Soil 0.6-1.0 m	180.45	9.67	188.02	7.09	116.35	8.89	119.86	6.11	0.510	< <b>0.001</b>	0.808
GF	Foliage	67.127	15.166	72.742	10.920	-	-	-	-	0.774	-	-
	Branches	30.592	17.391	36.721	9.497	-	-	-	-	0.665	-	-
	Bark	62.789	6.010	50.629	7.701	-	-	-	-	0.083	-	-
	Wood	-	-	-	-	-	-	-	-	-	-	-
	Understory	248.70	21.53	222.58	1.61	-	-	-	-	0.337	-	-
	Forest Floor	135.31	4.48	140.73	8.22	-	-	-	-	0.302	-	-
	Fine Roots	122.17	9.84	158.62	27.13	-	-	-	-	0.268	-	-
	Soil 0.0-0.2 m	166.06	10.84	196.64	1.72	-	-	-	-	<b>0.050</b>	-	-
	Soil 0.2-0.4 m	206.18	4.78	212.21	22.39	-	-	-	-	0.812	-	-
	Soil 0.4-0.6 m	185.03	1.59	188.50	8.36	-	-	-	-	0.675	-	-
	Soil 0.6-1.0 m	178.57	16.87	159.85	21.69	-	-	-	-	0.353	-	-

888

889

890  
891  
892  
893  
894

**Table A12.** Concentration (%) of phosphorus (P) of tree and ecosystem components for 16-18 year-old Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) stands growing under contrasting treatments of vegetation management on sites located in the central Coast Range (CR) and Cascade Foothills (CF) of western Oregon. SE is the standard error.

Species	Tissue	CR Control		CR VM		CF Control		CF VM		P-value		
		%	SE	%	SE	%	SE	%	SE	Trt	Site	Site x Trt
DF	Foliage	0.163	0.018	0.169	0.019	0.213	0.005	0.193	0.023	0.664	0.084	0.462
	Branches	0.060	0.014	0.056	0.007	0.065	0.009	0.061	0.010	0.610	0.484	0.942
	Bark	0.059	0.005	0.059	0.006	0.055	0.002	0.053	0.003	0.780	0.417	0.927
	Wood	0.004	0.000	0.003	0.000	0.007	0.002	0.004	0.001	0.155	0.092	0.189
	Understory	0.146	0.046	0.105	0.012	0.207	0.027	0.254	0.034	0.895	<b>0.008</b>	0.088
	Forest Floor	0.087	0.009	0.089	0.006	0.093	0.008	0.096	0.006	0.730	0.371	0.991
	Fine Roots	0.078	0.004	0.080	0.012	0.068	0.002	0.081	0.009	0.242	0.334	0.378
	Soil 0.0-0.2 m	0.526	0.028	0.555	0.041	1.018	0.078	0.954	0.054	0.752	<b>&lt;0.001</b>	0.404
	Soil 0.2-0.4 m	0.561	0.120	0.380	0.024	0.879	0.086	0.877	0.080	0.276	<b>0.001</b>	0.285
	Soil 0.4-0.6 m	0.504	0.077	0.349	0.026	0.662	0.058	0.687	0.054	0.273	<b>0.001</b>	0.138
	Soil 0.6-1.0 m	0.273	0.028	0.334	0.028	0.736	0.193	0.647	0.048	0.892	<b>0.002</b>	0.475
WH	Foliage	0.280	0.044	0.230	0.031	-	-	-	-	0.397	-	-
	Branches	0.043	0.004	0.044	0.005	-	-	-	-	0.875	-	-
	Bark	0.095	0.009	0.077	0.005	-	-	-	-	0.127	-	-
	Wood	0.010	0.003	0.010	0.002	-	-	-	-	0.973	-	-
	Understory	0.197	0.044	0.157	0.005	-	-	-	-	0.405	-	-
	Forest Floor	0.105	0.004	0.109	0.005	-	-	-	-	0.413	-	-
	Fine Roots	0.086	0.006	0.083	0.012	-	-	-	-	0.695	-	-
	Soil 0.0-0.2 m	0.708	0.154	0.674	0.074	-	-	-	-	0.814	-	-
	Soil 0.2-0.4 m	0.637	0.174	0.617	0.103	-	-	-	-	0.921	-	-
	Soil 0.4-0.6 m	0.541	0.083	0.505	0.061	-	-	-	-	0.710	-	-
	Soil 0.6-1.0 m	0.362	0.071	0.332	0.039	-	-	-	-	0.408	-	-
WRC	Foliage	0.127	0.009	0.113	0.007	0.113	0.012	0.109	0.015	0.440	0.441	0.650
	Branches	0.031	0.006	0.039	0.004	0.044	0.005	0.042	0.007	0.700	0.361	0.404
	Bark	0.053	0.007	0.039	0.002	0.052	0.002	0.050	0.006	0.140	0.319	0.228
	Wood	0.005	0.000	0.006	0.001	0.007	0.001	0.006	0.001	0.694	0.133	0.250
	Understory	0.164	0.041	0.168	0.045	0.176	0.040	0.246	0.030	0.357	0.344	0.403
	Forest Floor	0.100	0.018	0.090	0.021	0.064	0.012	0.064	0.021	0.713	0.060	0.726
	Fine Roots	0.082	0.003	0.078	0.013	0.064	0.003	0.081	0.005	0.371	0.310	0.162
	Soil 0.0-0.2 m	0.763	0.084	0.826	0.076	1.083	0.161	1.150	0.180	0.438	<b>0.005</b>	0.981
	Soil 0.2-0.4 m	0.642	0.079	0.602	0.056	0.805	0.102	0.869	0.151	0.838	<b>0.009</b>	0.404
	Soil 0.4-0.6 m	0.560	0.008	0.484	0.030	0.683	0.111	0.575	0.103	0.112	<b>0.018</b>	0.749
	Soil 0.6-1.0 m	0.400	0.018	0.364	0.032	0.548	0.071	0.536	0.156	0.819	0.202	0.909
GF	Foliage	0.136	0.008	0.155	0.016	-	-	-	-	0.348	-	-
	Branches	0.064	0.009	0.112	0.025	-	-	-	-	0.122	-	-
	Bark	0.078	0.010	0.052	0.006	-	-	-	-	0.073	-	-
	Wood	0.006	0.000	0.007	0.002	-	-	-	-	0.547	-	-
	Understory	0.175	0.053	0.144	0.005	-	-	-	-	0.594	-	-
	Forest Floor	0.098	0.012	0.111	0.011	-	-	-	-	0.468	-	-
	Fine Roots	0.098	0.008	0.111	0.020	-	-	-	-	0.586	-	-
	Soil 0.0-0.2 m	0.732	0.174	0.923	0.166	-	-	-	-	0.471	-	-
	Soil 0.2-0.4 m	0.464	0.057	0.649	0.157	-	-	-	-	0.331	-	-
	Soil 0.4-0.6 m	0.481	0.148	0.498	0.091	-	-	-	-	0.925	-	-
	Soil 0.6-1.0 m	0.382	0.161	0.365	0.006	-	-	-	-	0.923	-	-

895

896

897  
898  
899  
900  
901

**Table A13.** Concentration (%) of Sulfur (S) of tree and ecosystem components for 16-18 year-old Douglas-fir (DF), western hemlock (WH), western redcedar (WRC), and grand fir (GF) stands growing under contrasting treatments of vegetation management on sites located in the central Coast Range (CR) and Cascade Foothills (CF) of western Oregon. SE is the standard error.

Species	Tissue	CR Control		CR VM		CF Control		CF VM		P-value		
		%	SE	%	SE	%	SE	%	SE	Trt	Site	Site x Trt
DF	Foliage	0.121	0.006	0.154	0.039	0.115	0.004	0.111	0.002	0.463	0.229	0.367
	Branches	0.065	0.005	0.061	0.004	0.049	0.002	0.052	0.003	0.244	<b>0.006</b>	<b>0.024</b>
	Bark	0.065	0.011	0.051	0.002	0.049	0.003	0.049	0.002	0.291	0.188	0.308
	Wood	0.032	0.001	0.034	0.003	0.030	0.000	0.033	0.001	0.244	0.509	0.790
	Understory	0.157	0.029	0.113	0.004	0.147	0.012	0.168	0.017	0.554	0.279	0.099
	Forest Floor	0.106	0.012	0.112	0.007	0.114	0.006	0.113	0.002	0.592	0.994	0.494
	Fine Roots	0.079	0.005	0.080	0.001	0.071	0.005	0.075	0.002	0.524	0.103	0.661
WH	Foliage	0.110	0.018	0.092	0.019	-	-	-	-	0.481	-	-
	Branches	0.052	0.003	0.050	0.003	-	-	-	-	0.370	-	-
	Bark	0.055	0.004	0.057	0.006	-	-	-	-	0.529	-	-
	Wood	0.032	0.000	0.032	0.001	-	-	-	-	0.878	-	-
	Understory	0.160	0.015	0.172	0.006	-	-	-	-	0.452	-	-
	Forest Floor	0.104	0.006	0.107	0.006	-	-	-	-	0.784	-	-
	Fine Roots	0.075	0.004	0.068	0.002	-	-	-	-	0.234	-	-
WRC	Foliage	0.077	0.011	0.058	0.001	0.084	0.003	0.078	0.003	0.071	0.071	0.399
	Branches	0.048	0.004	0.050	0.001	0.045	0.002	0.036	0.003	0.301	<b>0.011</b>	0.081
	Bark	0.063	0.010	0.055	0.004	0.048	0.000	0.047	0.001	0.429	<b>0.012</b>	0.449
	Wood	0.035	0.001	0.036	0.002	0.036	0.001	0.035	0.002	0.663	0.680	0.119
	Understory	0.148	0.014	0.150	0.016	0.121	0.012	0.122	0.003	0.886	<b>0.026</b>	0.957
	Forest Floor	0.116	0.017	0.100	0.015	0.080	0.005	0.074	0.005	0.302	<b>0.027</b>	0.648
	Fine Roots	0.079	0.005	0.080	0.005	0.074	0.003	0.076	0.002	0.524	0.103	0.661
GF	Foliage	0.102	0.015	0.090	0.011	-	-	-	-	0.900	-	-
	Branches	0.053	0.003	0.070	0.008	-	-	-	-	0.113	-	-
	Bark	0.062	0.002	0.053	0.002	-	-	-	-	<b>0.019</b>	-	-
	Wood	0.033	0.002	0.034	0.003	-	-	-	-	0.873	-	-
	Understory	0.187	0.019	0.150	0.005	-	-	-	-	0.134	-	-
	Forest Floor	0.101	0.007	0.099	0.011	-	-	-	-	0.885	-	-
	Fine Roots	0.084	0.004	0.074	0.004	-	-	-	-	11.520	-	-

902

903      **Table A14.** Concentration (ppm) of zinc (Zn) of tree and ecosystem components for  
 904      16-18 year-old Douglas-fir (DF), western hemlock (WH), western redcedar (WRC),  
 905      and grand fir (GF) stands growing under contrasting treatments of vegetation  
 906      management on sites located in the central Coast Range (CR) and Cascade Foothills  
 907      (CF) of western Oregon. SE is the standard error.

Species	Tissue	CR Control		CR VM		CF Control		CF VM		P-value		
		ppm	SE	ppm	SE	ppm	SE	ppm	SE	Trt	Site	Site x Trt
DF	Foliage	11.031	2.015	10.416	0.856	12.386	1.079	11.922	2.705	0.897	0.841	0.881
	Branches	19.187	2.456	18.013	1.075	14.970	2.112	13.522	2.780	0.422	0.092	0.965
	Bark	16.825	0.555	18.194	1.248	17.993	3.263	16.904	1.087	0.927	0.997	0.515
	Wood	3.674	0.681	3.001	0.380	4.635	0.741	4.906	0.793	0.769	0.053	0.494
	Understory	26.617	11.655	16.493	6.024	21.821	2.580	33.762	5.748	0.903	0.408	0.155
	Forest Floor	9.793	1.488	10.135	1.217	15.676	1.857	14.442	1.495	0.777	<b>0.020</b>	0.620
	Fine Roots	10.979	0.876	12.399	1.642	12.767	1.252	10.983	0.367	0.836	0.799	0.102
	Soil 0.0-0.2 m	58.228	3.024	65.472	7.464	67.605	3.764	69.057	11.592	0.527	0.534	0.671
	Soil 0.2-0.4 m	61.790	2.428	56.924	6.136	62.827	3.904	69.478	11.360	0.892	0.471	0.396
	Soil 0.4-0.6 m	62.669	2.541	57.902	7.540	50.524	3.791	57.838	9.597	0.846	0.405	0.375
	Soil 0.6-1.0 m	50.815	4.120	54.975	5.893	49.460	12.250	51.591	8.244	0.679	0.658	0.893
WH	Foliage	11.686	0.896	10.341	0.983	-	-	-	-	0.351	-	-
	Branches	9.741	2.491	7.155	1.007	-	-	-	-	0.373	-	-
	Bark	7.075	1.283	5.218	1.207	-	-	-	-	0.333	-	-
	Wood	3.645	0.942	3.168	0.487	-	-	-	-	0.683	-	-
	Understory	27.572	7.117	13.608	0.456	-	-	-	-	0.131	-	-
	Forest Floor	14.366	0.997	12.490	1.538	-	-	-	-	0.346	-	-
	Fine Roots	10.428	0.565	10.376	1.370	-	-	-	-	0.973	-	-
	Soil 0.0-0.2 m	67.528	3.716	66.495	2.980	-	-	-	-	0.836	-	-
	Soil 0.2-0.4 m	70.611	4.061	65.751	4.357	-	-	-	-	0.446	-	-
	Soil 0.4-0.6 m	67.979	5.162	64.724	2.614	-	-	-	-	0.594	-	-
	Soil 0.6-1.0 m	53.941	5.279	53.569	1.731	-	-	-	-	0.935	-	-
WRC	Foliage	16.022	1.474	12.557	0.827	15.489	1.906	13.358	1.630	0.089	0.931	0.667
	Branches	10.508	1.384	9.513	0.823	9.059	2.663	5.889	0.947	0.225	0.145	0.516
	Bark	18.825	5.188	12.305	2.538	11.587	2.558	9.807	0.453	0.215	0.150	0.469
	Wood	2.012	0.044	2.494	0.350	2.889	0.592	3.038	0.564	0.492	0.136	0.716
	Understory	19.021	6.003	14.034	3.098	20.366	5.086	31.775	5.296	0.549	0.095	0.145
	Forest Floor	13.283	1.445	14.650	7.188	15.039	2.856	10.771	2.520	0.691	0.708	0.448
	Fine Roots	13.898	1.484	10.095	0.611	35.193	17.267	19.389	4.405	0.287	0.365	0.499
	Soil 0.0-0.2 m	67.375	7.071	71.251	4.990	80.082	9.555	86.671	4.025	0.292	<b>0.036</b>	0.772
	Soil 0.2-0.4 m	77.650	6.002	68.045	5.955	71.207	6.946	78.539	6.081	0.749	0.535	0.051
	Soil 0.4-0.6 m	75.434	9.748	66.819	4.836	62.587	6.037	68.148	5.730	0.823	0.604	0.334
	Soil 0.6-1.0 m	46.951	8.450	57.207	4.117	50.765	4.232	61.533	12.002	0.238	0.570	0.975
GF	Foliage	24.702	3.402	24.965	5.025	-	-	-	-	0.300	-	-
	Branches	14.174	3.192	16.743	2.596	-	-	-	-	0.562	-	-
	Bark	17.857	2.810	8.741	0.635	-	-	-	-	<b>0.019</b>	-	-
	Wood	4.353	0.877	3.680	0.325	-	-	-	-	0.499	-	-
	Understory	19.750	7.024	17.388	5.177	-	-	-	-	0.800	-	-
	Forest Floor	17.965	3.157	20.271	2.256	-	-	-	-	0.584	-	-
	Fine Roots	14.379	0.757	16.935	5.214	-	-	-	-	0.624	-	-
	Soil 0.0-0.2 m	65.707	1.521	83.900	6.691	-	-	-	-	0.078	-	-
	Soil 0.2-0.4 m	75.079	3.157	75.342	10.140	-	-	-	-	0.976	-	-
	Soil 0.4-0.6 m	66.434	2.161	75.266	8.629	-	-	-	-	0.377	-	-
	Soil 0.6-1.0 m	56.841	5.840	57.043	4.607	-	-	-	-	0.980	-	-

908

909

910

911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932

**Table A15.** Average concentrations in percent and standard errors (SE) of the macronutrients carbon, nitrogen, phosphorous, potassium, magnesium, calcium, and sulfur for the understory and the foliage and wood of the midstory species: bigleaf maple (ACMA), red alder (ALRU), Oregon bitter cherry (PREM), and cascara buckthorn (FRPU). Understory average was taken across sites, species, and treatments. BLD stands for below detectable levels.

		Carbon		Nitrogen		Phosphorous		Potassium		Magnesium		Calcium		Sulfur	
		%	SE	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
<b>ACMA</b>	Foliage	44.917	0.380	3.010	0.116	0.371	0.043	1.496	0.034	0.344	0.014	1.163	0.134	0.204	0.010
	Stem	45.472	0.127	0.117	0.002	0.017	0.005	BLD	-	0.036	0.001	0.082	0.003	0.043	0.000
<b>ALRU</b>	Foliage	47.240	0.330	2.938	0.154	0.147	0.007	1.083	0.142	0.253	0.003	0.590	0.052	0.087	0.010
	Stem	53.112	0.448	0.552	0.004	0.009	0.001	BLD	-	0.026	0.002	0.087	0.015	0.048	0.003
<b>PREM</b>	Foliage	44.108	0.576	3.248	0.100	0.412	0.032	1.506	0.049	0.397	0.018	1.147	0.126	0.160	0.002
	Stem	44.916	0.088	0.110	0.003	0.021	0.010	BLD	-	0.027	0.005	0.124	0.017	0.049	0.003
<b>FRPU</b>	Foliage	42.813	0.101	2.678	0.049	0.331	0.009	1.709	0.056	0.351	0.012	1.213	0.082	0.117	0.002
	Stem	45.212	0.095	0.143	0.005	0.007	0.002	BLD	-	0.042	0.001	0.125	0.008	0.045	0.001
<b>Understory</b>	Total	40.834	0.845	1.375	0.059	0.180	0.011	1.177	0.104	0.272	0.015	0.907	0.049	0.000	0.000

**Table A16.** Average concentrations in  $\text{mg kg}^{-1}$  (ppm) and standard errors (SE) of the micronutrients boron, copper, iron, sodium, and zinc for the understory and the foliage and wood of the midstory species: bigleaf maple (ACMA), red alder (ALRU), Oregon bitter cherry (PREM), and cascara buckthorn (FRPU). Understory average was taken across sites, species, and treatments. BLD stands for below detectable levels.

		Boron		Copper		Iron		Manganese		Sodium		Zinc	
		ppm	SE	ppm	SE	ppm	SE	ppm	SE	ppm	SE	ppm	SE
<b>ACMA</b>	Foliage	14.66	2.91	6.589	0.381	100.00	17.40	180.98	15.05	192.61	31.74	52.05	4.65
	Stem	2.70	0.10	3.332	0.598	100.46	10.53	24.51	4.02	BLD	-	5.66	0.81
<b>ALRU</b>	Foliage	15.50	0.42	8.179	0.458	62.62	4.17	229.03	16.38	166.26	23.17	26.30	2.61
	Stem	2.36	0.08	2.239	0.097	135.46	29.79	17.74	0.69	BLD	-	5.09	0.32
<b>PREM</b>	Foliage	18.53	1.29	5.681	0.478	156.25	1.79	58.01	3.16	93.55	19.47	11.73	1.66
	Stem	3.62	0.44	2.121	0.246	89.29	5.48	4.74	0.11	BLD	-	2.49	0.19
<b>FRPU</b>	Foliage	32.07	0.34	6.845	0.684	138.71	9.10	833.53	12.75	157.24	11.48	23.73	1.44
	Stem	2.72	0.09	2.377	0.412	109.86	38.47	31.32	1.13	BLD	-	2.37	0.39
<b>Understory</b>	Total	27.67	2.59	5.839	0.330	918.54	115.52	562.36	55.27	195.19	11.32	22.24	1.84