

Many-Body Signatures of Collective Decay in Atomic Chains

Stuart J. Masson,^{1,*} Igor Ferrier-Barbut^{1,2} Luis A. Orozco,³ Antoine Browaeys,² and Ana Asenjo-Garcia^{1,†}

¹Department of Physics, Columbia University, New York, New York 10027, USA

²Université Paris-Saclay, Institut d'Optique Graduate School, CNRS, Laboratoire Charles Fabry, 91127 Paris, France

³Joint Quantum Institute, Department of Physics and NIST, University of Maryland, College Park, Maryland 20742, USA



(Received 26 August 2020; accepted 16 November 2020; published 23 December 2020)

Fully inverted atoms placed at exactly the same location synchronize as they deexcite, and light is emitted in a burst (known as “Dicke’s superradiance”). We investigate the role of finite interatomic separation on correlated decay in mesoscopic chains and provide an understanding in terms of collective jump operators. We show that the superradiant burst survives at small distances, despite Hamiltonian dipole-dipole interactions. However, for larger separations, competition between different jump operators leads to dephasing, suppressing superradiance. Collective effects are still significant for arrays with lattice constants of the order of a wavelength, and lead to a photon emission rate that decays nonexponentially in time. We calculate the two-photon correlation function and demonstrate that emission is correlated and directional, as well as sensitive to small changes in the interatomic distance. These features can be measured in current experimental setups, and are robust to realistic imperfections.

DOI: 10.1103/PhysRevLett.125.263601

Collective effects in the interaction between light and matter have attracted interest since the seminal work of Dicke in the 1950s, who studied the problem of photon emission by many atoms at identical locations [1]. In this purely dissipative scenario, the atomic dipoles become phase locked during their decay and emit collectively. This leads to an initial increase in the photon emission rate—the famous “superradiant burst” or “superfluorescence”—rather than the typical exponential decay for independent atoms.

Dicke’s scenario ignores coherent dipole-dipole interactions between atoms, which are relevant for finite interatomic distances and have been predicted to wash out superradiant decay [2,3]. However, signatures of collective behavior persist even in systems of size much larger than the resonance wavelength. For example, theoretical studies of ordered arrays of emitters have shown the existence of extremely subradiant (i.e., dark) few-excitation states [4–12], as well as directional collective emission [4,13–21].

Recent experimental realizations of ordered atomic arrays, both in optical lattices [22–26] and tweezer arrays [27–33], open the door for investigation of these predictions. These platforms have already allowed for the demonstration of a two-dimensional atomic mirror [26] and the measurement of collective frequency shifts in a one-dimensional (1D) atomic array [33]. Current experimental capabilities offer the possibility of measuring statistics of the emitted photons. This raises the question of whether collective decay imprints correlations on the emitted photons. This would allow for the potentially tunable generation of nonclassical states of light (and maybe of a superradiant laser [34,35]), critical for quantum

technologies. Conversely, connecting the correlations in the light back to the atomic quantum state would offer a unique light-based probe to characterize these dissipative many-body systems.

Here, we make an important step in this direction by investigating collective decay and superradiance in mesoscopic 1D ordered arrays. We find that a superradiant burst survives for short interatomic distances ($d \lesssim \lambda_0/4$). We use a quantum jump approach to connect the sequence of collective jumps to the statistics of the light emitted. We show that strong spatial correlations between emitted photons are imparted by the collective decay and that they persist even for $d \sim \lambda_0$.

We consider N atoms of resonance frequency ω_0 , arranged in an ordered chain along the z -axis with lattice constant d , as shown in Fig. 1(a). Interactions between atoms are obtained by tracing out the electromagnetic field under a Born-Markov approximation [36,37]. The atomic density matrix, ρ , evolves under the master equation

$$\dot{\rho} = -\frac{i}{\hbar} [\mathcal{H}, \rho] + \sum_{i,j=1}^N \frac{\Gamma^{ij}}{2} (2\hat{\sigma}_{ge}^j \rho \hat{\sigma}_{eg}^i - \rho \hat{\sigma}_{eg}^i \hat{\sigma}_{ge}^j - \hat{\sigma}_{eg}^i \hat{\sigma}_{ge}^j \rho), \quad (1)$$

where the Hamiltonian is

$$\mathcal{H} = \hbar \sum_{i=1}^N \omega_0 \hat{\sigma}_{ee}^i + \hbar \sum_{i,j=1}^N J^{ij} \hat{\sigma}_{eg}^i \hat{\sigma}_{ge}^j. \quad (2)$$

Here, $\hat{\sigma}_{ge}^i = |g_i\rangle \langle e_i|$ is the atomic coherence operator, with $|e_i\rangle$ and $|g_i\rangle$ the excited and ground states of the cycling




FIG. 1. A chain of excited atoms decays collectively, emitting correlated photons. (a) Schematic of the considered setup. Atoms have resonance frequency ω_0 , and are separated by a constant distance d . The relevant transition is selected via a small magnetic field. (b) Decay rates of the jump operators $\{\hat{O}_\nu\}$ for $N = 10$ atoms. Each operator approximates a spin wave with wave vector k_ν , with the darkest (lightest) lines corresponding to spin waves with minimum (maximum) wave vector. Inset: Angular emission pattern (in arbitrary units) following action of the most subradiant operator on a fully inverted array, measured by detectors of width $\Delta\theta = 0.01\pi$. Dashed lines are analytically obtained angles of peak emission, $\theta_{\max} = \arccos(\pm k_\nu/k_0)$ with $k_\nu = 0(\pi/d)$ for $d = 0.9(1.1)\lambda_0$.

transition of the i th atom at position $\mathbf{r}_i = \{x_i, y_i, z_i\}$. The coherent and dissipative interactions between atoms i and j are [38,39]

$$J^{ij} - i \frac{\Gamma^{ij}}{2} = -\frac{\mu_0 \omega_0^2}{\hbar} \varphi^* \cdot \mathbf{G}_0(\mathbf{r}_i, \mathbf{r}_j, \omega_0) \cdot \varphi, \quad (3)$$

where $\varphi = (|\varphi|/\sqrt{2})(\hat{x} + i\hat{y})$ is the dipole matrix element of the circularly polarized transition σ^+ , and $\mathbf{G}_0(\mathbf{r}_i, \mathbf{r}_j, \omega_0)$ is the propagator of the electromagnetic field between positions \mathbf{r}_i and \mathbf{r}_j [9]. The scattered field along the axis of the chain is σ^+ polarized and the atoms behave as two-level systems even in the presence of complex hyperfine structure [40–42].

An ensemble of N atoms decays collectively, via a set of N jump operators, $\{\hat{O}_\nu\}$, with associated decay rates $\{\Gamma_\nu\}$. These operators are eigenstates of the dissipative interaction matrix Γ with elements Γ^{ij} [13–15]. The master equation can be written in terms of these operators as

$$\dot{\rho} = -\frac{i}{\hbar} [\mathcal{H}, \rho] + \sum_{\nu=1}^N \frac{\Gamma_\nu}{2} (2\hat{O}_\nu \rho \hat{O}_\nu^\dagger - \rho \hat{O}_\nu^\dagger \hat{O}_\nu - \hat{O}_\nu^\dagger \hat{O}_\nu \rho). \quad (4)$$

The operators' decay rates can be superradiant, i.e., $\Gamma_\nu > \Gamma_0$, or subradiant, i.e., $\Gamma_\nu < \Gamma_0$, with $\Gamma_0 \equiv \Gamma^{ii}$ the single-atom spontaneous emission rate.

Jumps happen stochastically, and operators act on all atoms with a set of amplitudes and phases sensitive to d and the atomic quantization axis. The states $\{\hat{O}_\nu^\dagger |g\rangle^{\otimes N}\}$ ($\{\hat{O}_\nu^\dagger |e\rangle^{\otimes N}\}$) form an orthonormal basis for the single-excitation (“single-hole”) system. The jump operators can be classified according to the symmetries of the Lindblad operator. For an infinite 1D array, these are discrete translations along \hat{z} . Thus, the operators correspond to Bloch waves with a wave vector along the chain direction k_ν , i.e., $\hat{O}_\nu = (1/\sqrt{N}) \sum_{i=1}^N e^{-ik_\nu z_i} \hat{\sigma}_{ge}^i$. In 1D geometries, the collective decay rates Γ_ν change with d featuring sharp oscillations at $d \simeq n\lambda_0/2$ with $n \in \mathbb{N}$ [7,43,44], as shown in Fig. 1(b). For $d = n\lambda_0/2 + \epsilon$ ($d = n\lambda_0/2 - \epsilon$), with $\epsilon \rightarrow 0^+$, there are a small number of superradiant (subradiant) operators, with the majority of rates weakly subradiant (superradiant). These oscillations arise from 1D lattice sums and can be understood by analogy with the decay of a dipole in a cavity [7].

When a jump occurs, a photon is emitted. We calculate the emission angle of the radiated photons by means of directed-detection operators, following Carmichael and coworkers [13–15]. Photon detection at a point $\mathbf{R} = (r, \theta, \phi)$ corresponds to action of the operator

$$\hat{D}(\theta, \phi) = \sqrt{\frac{3\Gamma_0}{8\pi} \left(1 - \frac{\sin^2 \theta}{2}\right)} d\Omega \sum_{j=1}^N e^{-ik_0 z_j \cos \theta} \hat{\sigma}_{ge}^j, \quad (5)$$

where $d\Omega$ is a solid-angle differential. The detectors are assumed to be in far field, such that $|\mathbf{R}| \gg \lambda_0, Nd$. The probability of a photon detection in direction (θ, ϕ) is $P(\theta, \phi) d\Omega = \langle \hat{D}^\dagger(\theta, \phi) \hat{D}(\theta, \phi) \rangle$. A (square) photon detector of finite solid angle $\Delta\Omega$ and angular width $\Delta\theta$ sees intensity

$$\mathfrak{I}(\theta) = \frac{\Delta\Omega}{\Delta\theta} \int_{\theta-\Delta\theta/2}^{\theta+\Delta\theta/2} P(\theta') d\theta'. \quad (6)$$

Photon emission caused by action of a jump operator is directional. Figure 1(b) shows the angular distribution of a photon emitted during the action of the most subradiant operator on the fully inverted array. The maximal emission angles, calculated by considering correlations between jump and directed-detection operators, are [see Supplemental Material (SM) [45]]

$$\theta_{\max} = \arccos \left(\frac{n\lambda_0}{d} \pm \frac{k_\nu}{k_0} \right), \quad n \in \mathbb{Z}. \quad (7)$$




FIG. 2. Decay of a fully inverted chain and stochastic action of collective jump operators. (a) Photon emission rate normalized by atom number $[-d\langle n_{\text{exc}}(t) \rangle / d(N\Gamma_0 t)]$ of an array of $N = 8$ atoms for different interatomic distances. As d is increased, the emission rate at early times shows a transition from an increase (superradiant burst) to a decrease. Inset: Boundary between regions. We estimate a burst occurs if the emission rate is larger at $N\Gamma_0 t = 10^{-4}$ than at $t = 0$. (b) Illustrations of jump operator action during the decay, for different lattice constants. The star represents the initial state $|e\rangle^{\otimes N}$. Circles represent action of one of the N different jump operators \hat{O}_ν , colored and displayed in order from most superradiant (white) on the left to most subradiant (black) on the right. Line thickness represents the likelihood of a particular path, based on a set of 1000 trajectories. For $d = 0.1\lambda_0$, some trajectories are extremely subradiant (23 trajectories not fully deexcited by $\Gamma_0 t = 500$ are omitted).

Here, the \pm accounts for the mirror reflection symmetry of a finite chain (whose jump operators carry $\pm k_\nu$ wave-vector components). Jump operators cannot be expressed in terms of directed-detection operators [14]. For multiple holes and excitations, the intensity pattern may contain additional lobes due to atomic correlations.

A fully inverted array develops correlations as it decays. The rate of change of the atomic population, $\langle \hat{n}_{\text{exc}} \rangle = \sum_{i=1}^N \langle \hat{\sigma}_{ee}^i \rangle$, dictates the photon emission rate

$$R(t) = -\frac{d\langle \hat{n}_{\text{exc}}(t) \rangle}{dt} = \sum_{\nu=1}^N \Gamma_\nu \langle \hat{O}_\nu^\dagger \hat{O}_\nu \rangle. \quad (8)$$

At $t = 0$, all atoms are excited, and there are not any correlations between them ($\langle \hat{O}_\nu^\dagger \hat{O}_\nu \rangle = 1 \forall \nu$). By definition, $\sum_\nu \Gamma_\nu = \text{Tr } \mathbf{\Gamma} = N\Gamma_0$, and $R(t = 0) = N\Gamma_0$. Since the atoms are uncorrelated, the initial decay is the sum of N independently decaying atoms.

Dicke superradiance ($d = 0$) is a unique situation, as there is only one jump operator with nonzero decay rate, \hat{O}_D . That operator has rate $N\Gamma_0$ and acts identically on all atoms, i.e., $\hat{O}_D = (1/\sqrt{N}) \sum_{i=1}^N \hat{\sigma}_{ge}^i$. The decay rate $R_D(t) = N\Gamma_0 \langle \hat{O}_D^\dagger \hat{O}_D \rangle$ is maximized with half the atoms excited. This gives rise to the superradiant burst seen in Fig. 2(a), where the peak rate of photon emission scales as N^2 and occurs at some finite time [2]. Since there is a single jump operator, decay is never subradiant. Dicke superradiance seems not to be recovered as $d \rightarrow 0$ [see Fig. 2(a) and the SM [45]]: the emission rate saturates to a different asymptotic curve, suggesting that Dicke superradiance is not analytically connected to this regime. Rings show similar behavior [45]. We attribute this saturation to a complex interplay of stimulation and competition between

the set of jump operators that deexcites the array, as discussed below.

In extended arrays, jump operators enhance their own action [1,13,14] but compete with each other. This occurs due to correlations induced by photon emission, irrespective of whether the photon is detected or not. For the fully inverted array, the normalized probability of two different successive jumps (\hat{O}_ν and \hat{O}_μ) can be calculated analytically. In the large N limit, it yields [45]

$$\tilde{g}^{(2)}(\tau = 0)|_{\nu,\mu} = \frac{\langle \hat{O}_\nu^\dagger \hat{O}_\mu^\dagger \hat{O}_\mu \hat{O}_\nu \rangle}{\langle \hat{O}_\nu^\dagger \hat{O}_\nu \rangle \langle \hat{O}_\mu^\dagger \hat{O}_\mu \rangle} \simeq 1 + \delta_{\nu\mu} - \frac{2}{N}. \quad (9)$$

While this is a process of spontaneous emission, each jump operator enhances itself (but not others), and thus, an effective stimulated emission of radiation occurs at certain angles, as jump operators are directional. The last term in the equation is a fermionic correction that illustrates that a single atom can only host a single excitation (there is a $1/N$ probability for two excitations to overlap in a chain of N atoms, and the factor of 2 arises because there are two identical ways to assign two excitations). For $N \geq 4$, $\tilde{g}^{(2)}(\tau = 0)|_{\nu,\nu} > 1 \forall \nu$, i.e., even in very short chains, all operators enhance their own action.

The competition between different jump operators causes dephasing of the atomic state, reducing and eventually blocking superradiance. Following this argument, atoms in other geometries must also dephase, since for $d \neq 0$, there are always multiple operators and, thus, competition between decay paths. The set of likely deexcitation paths diversifies as d increases, as shown in Fig. 2(b), leading to faster dephasing. This reduces the intensity of the superradiant burst and brings it forward in time—no longer happening when half of the atoms are




FIG. 3. Directional photon emission from a fully inverted chain of $N = 10$ atoms. (top) Intensity, normalized by value at $t = 0$, at far-field detectors of angular width $\Delta\theta = 0.01\pi$. Curves represent evenly spaced snapshots of the intensity profile for $\Gamma_0 t \in [0, 20]$. (bottom) Directional two-photon correlation function, $g^{(2)}(\tau = 0)|_{\theta, \theta}$ as defined in Eq. (10).

excited, but earlier—until the burst disappears. Each path has a likelihood dictated by the operator decay rates and the correlations induced as the ensemble deexcites. The only forbidden paths are those where the cumulative effect of the jump operators breaks the mirror symmetry about the center of the array.

We find that the superradiant burst survives at small enough interatomic distances despite being suppressed, as shown in Fig. 2(a). While the peak intensity is fainter and does not scale as N^2 [45], the photon emission rate initially increases. In the inset, we show the distance for which the superradiant burst disappears. We estimate that a superradiant burst will occur if the derivative of the photon emission rate at $t = 0$ is positive. We perform calculations for large atom numbers by truncating the Hilbert space to subspaces with up to two atoms in the ground state (i.e., maximum of two photons emitted). This captures well the early dynamics. As shown in the inset of Fig. 2(a), for long chains, superradiant features are retained for $d \lesssim 0.25\lambda_0$. At this upper limit, competition between jump operators becomes so strong that the burst becomes marginal and is eventually suppressed.

The action of different jump operators throughout the evolution leads to changes in the directionality of photon emission at different times, as shown in Fig. 3. The fully excited ensemble emits quite broadly in space. Without correlations, the atoms emit as independently radiating dipoles [13,14]. However, at late times, subradiant operators become dominant and emission is strongly peaked in a direction dictated by d . Angular emission is narrow for

$d = 0.9\lambda_0$, as there is one dominant subradiant mode, but broad for $d = 1.1\lambda_0$ where multiple subradiant modes are important [see Fig. 1(b)]. Radiation in different directions is correlated [20]: emission at angle θ_1 enhances emission in directions that satisfy $\cos\theta_2 = \cos\theta_1 - n\lambda_0/d$, $n \in \mathbb{Z}$, as jump operator emission patterns are multilobed [45].

Self-enhancing (or stimulated) emission is confirmed by calculating the direction-dependent second order correlation function

$$g^{(2)}(\tau = 0)|_{\theta, \theta} = \frac{\langle \hat{D}^\dagger(\theta)\hat{D}^\dagger(\theta)\hat{D}(\theta)\hat{D}(\theta) \rangle}{\langle \hat{D}^\dagger(\theta)\hat{D}(\theta) \rangle^2}. \quad (10)$$

Figure 3 shows large, direction-dependent bunching in the field radiated by the array under evolution according to Eq. (4). At $t = 0$, $g^{(2)}(\tau = 0)|_{\theta, \theta}$ can be calculated analytically, yielding a spatially uniform value of $2 - 2/N$ [45], reproducing Dicke's result [1,15]. At late times, there are large peaks at intensity minima. While single photon emission is very unlikely, conditioned on one photon, a second is significantly more likely, such that pairs are relatively enhanced [17,19]. At late times, the signal can be sub-Poissonian in the direction of peak intensity [see plot for $d = 0.9\lambda_0$ in Fig. 3], as subradiance is predominantly a single-excitation effect and photon pairs are suppressed. Evidence of such directional statistics has been observed for two emitters [51].

Signatures of collective decay can be observed without fully inverting the array, but preparing spin coherent states, instead [52],




FIG. 4. Role of initial number of excitations on photon emission rate from spin coherent states. Photon emission (solid lines) from $|\varphi = 0.3, \mathbf{k} = k_0 \hat{z}\rangle$ with average excitation number $\langle \hat{n}_{\text{exc}} \rangle = \varphi N$ [see Eq. (11)]. Dashed lines show exponential fits, from which one extracts decay rate coefficients γ_{early} (dashed) and γ_{late} (dot-dashed). Inset: Early and late fitted decay rates as a function of the initial number of excitations. For all plots, $N = 10$. For $d = 0.9\lambda_0$, γ_{early} is fitted for $\Gamma_0 t \in [0, 4]$ and γ_{late} for $\Gamma_0 t \in [10, 20]$. For $d = 1.1\lambda_0$, γ_{early} is fitted for $\Gamma_0 t \in [0, 2]$ and γ_{late} for $\Gamma_0 t \in [5, 10]$.

$$|\varphi, \mathbf{k}\rangle = \prod_{i=1}^N \sqrt{1-\varphi} |g_i\rangle + e^{i\mathbf{k} \cdot \mathbf{r}_i} \sqrt{\varphi} |e_i\rangle, \quad (11)$$

where $0 \leq \varphi \leq 1$. These states have a binomially distributed excitation number defined by probability φ , and an excitation expectation value $\langle \hat{n}_{\text{exc}} \rangle = \varphi N$. Coherent spin states can be prepared experimentally by exciting the array with an intense pulse of duration $\tau \ll (N\Gamma_0)^{-1}$, $(J^{12})^{-1}$, to prevent collective effects.

Coherent spin states exhibit nonexponential temporal decay due to the interplay of multiple jump operators. The subradiant tail survives at distances accessible with current experimental capabilities, as shown in Fig. 4. This can be characterized by separately fitting the early and late dynamics, as demonstrated in experiments with atomic clouds in free space [53,54] and near a nanofiber [55]. The fitted decay rates of early dynamics depend on the initial atomic state; the decay is a many-body problem dependent on the density of excitations. Fits at late times do not depend so markedly on the number of excitations, since late dynamics are predominantly a single-excitation phenomenon independent of initial conditions. For large numbers of excitations, γ_{early} converges for all geometries as the fully excited state is uncorrelated. The contrast between early and late fits is significantly larger for $d = n\lambda_0/2 - \epsilon$, due to the differences in decay rates across each resonance. As shown in the SM [45], the drive can be used to imprint correlations on the array, some of which survive at long times and impact the radiation pattern.

Experimental realizations in the regime $d \sim \lambda_0$ should be feasible with current technologies. However, reaching the superradiant regime requires shorter interparticle distances. The limit of $d \lesssim 0.25\lambda_0$ established here could be satisfied with long-wavelength transitions (such that $\lambda_0 > 800$ nm) of lanthanide atoms trapped in optical lattices with wavelength near their strong transitions ~ 400 nm [56,57]. Even shorter interatomic distances can be reached in disordered ensembles [58,59], which constitute an interesting prospect for future work. Arrays of solid state emitters, such as localized excitonic quantum dots or strain-generated defects in 2D materials [60,61], are an additional playground for collective decay.

In conclusion, we have studied the collective decay of ordered chains of atoms in free space. We have found that superradiance survives significant interatomic separations, though Dicke's perfectly symmetric decay is lost at any finite distance in all ordered geometries. For separations comparable to the resonance wavelength, strong signatures of collective decay remain, and photon emission has directional features and a subradiant tail. These phenomena are robust to realistic experimental imperfections, namely, finite filling fraction and classical position noise [45]. Geometry can be used to control the many-body optical response of arrays. With conditional feedback control [62] (assisted by directional detection), it may pave the way toward the preparation of target entangled states, such as metrologically useful subradiant states.

Research on superradiance and many-body physics was supported by Programmable Quantum Materials, an Energy Frontier Research Center funded by the U.S. Department of Energy (DOE), Office of Science, Basic Energy Sciences (BES), under Award No. DE-SC0019443. Research on late-time dynamics and subradiance was supported by the National Science Foundation QII-TAQS (Grant No. 1936359). I. F. B. and A. B. acknowledge financial support by the Région Île-de-France in the framework of Domaine d'Intérêt Majeur SIRTEQ (Project DSHAPE).

* s.j.masson@columbia.edu

† ana.asenjo@columbia.edu

- [1] R. H. Dicke, Coherence in spontaneous radiation processes, *Phys. Rev.* **93**, 99 (1954).
- [2] M. Gross and S. Haroche, Superradiance: An essay on the theory of collective spontaneous emission, *Phys. Rep.* **93**, 301 (1982).
- [3] M. G. Benedict, A. M. Ermolaev, V. A. Malyshev, I. V. Sokolov, and E. D. Trifonov, *Super-Radiance: Multiaatomic Coherent Emission* (CRC Press, Boca Raton, 1996).
- [4] H. Zoubi and H. Ritsch, Metastability and directional emission characteristics of excitons in 1D optical lattices, *Europ. Lett.* **90**, 23001 (2010).

[5] R. J. Bettles, S. A. Gardiner, and C. S. Adams, Enhanced Optical Cross Section Via Collective Coupling of Atomic Dipoles in a 2D Array, *Phys. Rev. Lett.* **116**, 103602 (2016).

[6] R. T. Sutherland and F. Robicheaux, Collective dipole-dipole interactions in an atomic array, *Phys. Rev. A* **94**, 013847 (2016).

[7] R. J. Bettles, S. A. Gardiner, and C. S. Adams, Cooperative eigenmodes and scattering in one-dimensional atomic arrays, *Phys. Rev. A* **94**, 043844 (2016).

[8] G. Facchinetto, S. D. Jenkins, and J. Ruostekoski, Storing Light with Subradiant Correlations in Arrays of Atoms, *Phys. Rev. Lett.* **117**, 243601 (2016).

[9] A. Asenjo-Garcia, M. Moreno-Cardoner, A. Albrecht, H. J. Kimble, and D. E. Chang, Exponential Improvement in Photon Storage Fidelities Using Subradiance and Selective Radiance in Atomic Arrays, *Phys. Rev. X* **7**, 031024 (2017).

[10] L. Henriet, J. S. Douglas, D. E. Chang, and A. Albrecht, Critical open-system dynamics in a one-dimensional optical-lattice clock, *Phys. Rev. A* **99**, 023802 (2019).

[11] S. J. Masson and A. Asenjo-Garcia, Atomic-waveguide quantum electrodynamics, *Phys. Rev. Research* **2**, 043213 (2020).

[12] L. A. Williamson, M. O. Borgh, and J. Ruostekoski, Superatom Picture of Collective Nonclassical Light Emission and Dipole Blockade in Atom Arrays, *Phys. Rev. Lett.* **125**, 073602 (2020).

[13] H. J. Carmichael and K. Kim, A quantum trajectory unraveling of the superradiance master equation, *Opt. Commun.* **179**, 417 (2000).

[14] J. P. Clemens, L. Horvath, B. C. Sanders, and H. J. Carmichael, Collective spontaneous emission from a line of atoms, *Phys. Rev. A* **68**, 023809 (2003).

[15] J. P. Clemens, L. Horvath, B. C. Sanders, and H. J. Carmichael, Shot-to-shot fluctuations in the directed superradiant emission from extended atomic samples, *J. Opt. B* **6**, S736 (2004).

[16] M. O. Scully, E. S. Fry, C. H. Raymond Ooi, and K. Wódkiewicz, Directed Spontaneous Emission from an Extended Ensemble of n Atoms: Timing is Everything, *Phys. Rev. Lett.* **96**, 010501 (2006).

[17] D. Bhatti, J. von Zanthier, and G. S. Agarwal, Superbunching and nonclassicality as new hallmarks of superradiance, *Sci. Rep.* **5**, 17335 (2015).

[18] E. Shahmoon, D. S. Wild, M. D. Lukin, and S. F. Yelin, Cooperative Resonances in Light Scattering from Two-Dimensional Atomic Arrays, *Phys. Rev. Lett.* **118**, 113601 (2017).

[19] Qurrat-ul-Ain Gulfam and Z. Ficek, Highly directional photon superbunching from a few-atom chain of emitters, *Phys. Rev. A* **98**, 063824 (2018).

[20] I. Liberal, I. Ederra, and R. W. Ziolkowski, Grating lobes in higher-order correlation functions of arrays of quantum emitters: Directional photon bunching versus correlated directions, *Photonics* **6**, 14 (2019).

[21] K. E. Ballantine and J. Ruostekoski, Subradiance-protected excitation spreading in the generation of collimated photon emission from an atomic array, *Phys. Rev. Research* **2**, 023086 (2020).

[22] W. S. Bakr, A. Peng, M. E. Tai, R. Ma, J. Simon, J. I. Gillen, S. Fölling, L. Pollet, and M. Greiner, Probing the superfluid-to-Mott insulator transition at the single-atom level, *Science* **329**, 547 (2010).

[23] J. F. Sherson, C. Weitenberg, M. Endres, M. Cheneau, I. Bloch, and S. Kuhr, Single-atom-resolved fluorescence imaging of an atomic Mott insulator, *Nature (London)* **467**, 68 (2010).

[24] D. Greif, M. F. Parsons, A. Mazurenko, C. S. Chiu, S. Blatt, F. Huber, G. Ji, and M. Greiner, Site-resolved imaging of a fermionic Mott insulator, *Science* **351**, 953 (2016).

[25] A. Kumar, T.-Y. Wu, F. Giraldo, and D. S. Weiss, Sorting ultracold atoms in a three-dimensional optical lattice in a realization of Maxwell's demon, *Nature (London)* **561**, 83 (2018).

[26] J. Rui, D. Wei, A. Rubio-Abadal, S. Hollerith, J. Zeiher, D. M. Stamper-Kurn, C. Gross, and I. Bloch, A subradiant optical mirror formed by a single structured atomic layer, *Nature (London)* **583**, 369 (2020).

[27] B. J. Lester, N. Luick, A. M. Kaufman, C. M. Reynolds, and C. A. Regal, Rapid Production of Uniformly Filled Arrays of Neutral Atoms, *Phys. Rev. Lett.* **115**, 073003 (2015).

[28] H. Kim, W. Lee, H.-G. Lee, H. Jo, Y. Song, and J. Ahn, *In situ* single-atom array synthesis using dynamic holographic optical tweezers, *Nat. Commun.* **7**, 13317 (2016).

[29] M. Endres, H. Bernien, A. Keesling, H. Levine, E. R. Anschuetz, A. Krajenbrink, C. Senko, V. Vuletic, M. Greiner, and M. D. Lukin, Atom-by-atom assembly of defect-free one-dimensional cold atom arrays, *Science* **354**, 1024 (2016).

[30] D. Barredo, S. de Léséleuc, V. Lienhard, T. Lahaye, and A. Browaeys, An atom-by-atom assembler of defect-free arbitrary two-dimensional atomic arrays, *Science* **354**, 1021 (2016).

[31] D. Barredo, V. Lienhard, S. de Léséleuc, T. Lahaye, and A. Browaeys, Synthetic three-dimensional atomic structures assembled atom by atom, *Nature (London)* **561**, 79 (2018).

[32] A. Browaeys and T. Lahaye, Many-body physics with individually controlled Rydberg atoms, *Nat. Phys.* **16**, 132 (2020).

[33] A. Glicenstein, G. Ferioli, N. Šibalić, L. Brossard, I. Ferrier-Barbut, and A. Browaeys, Collective Shift in Resonant Light Scattering by a One-Dimensional Atomic Chain, *Phys. Rev. Lett.* **124**, 253602 (2020).

[34] D. Meiser, J. Ye, D. R. Carlson, and M. J. Holland, Prospects for a Millihertz-Linewidth Laser, *Phys. Rev. Lett.* **102**, 163601 (2009).

[35] J. G. Bohnet, Z. Chen, J. M. Weiner, D. Meiser, M. J. Holland, and J. K. Thompson, A steady-state superradiant laser with less than one intracavity photon, *Nature (London)* **484**, 78 (2012).

[36] T. Gruner and D.-G. Welsch, Green-function approach to the radiation-field quantization for homogeneous and inhomogeneous Kramers-Kronig dielectrics, *Phys. Rev. A* **53**, 1818 (1996).

[37] H. T. Dung, L. Knöll, and D.-G. Welsch, Resonant dipole-dipole interaction in the presence of dispersing and absorbing surroundings, *Phys. Rev. A* **66**, 063810 (2002).

[38] M. J. Stephen, First-order dispersion forces, *J. Chem. Phys.* **40**, 669 (1964).

[39] R. H. Lehmberg, Radiation from an N -atom system. I. General formalism, *Phys. Rev. A* **2**, 883 (1970).

[40] M. Hebenstreit, B. Kraus, L. Ostermann, and H. Ritsch, Subradiance Via Entanglement in Atoms with Several Independent Decay Channels, *Phys. Rev. Lett.* **118**, 143602 (2017).

[41] A. Piñeiro Orioli and A. M. Rey, Dark States of Multilevel Fermionic Atoms in Doubly Filled Optical Lattices, *Phys. Rev. Lett.* **123**, 223601 (2019).

[42] A. Asenjo-Garcia, H. J. Kimble, and D. E. Chang, Optical waveguiding by atomic entanglement in multilevel atom arrays, *Proc. Natl. Acad. Sci. U.S.A.* **116**, 25503 (2019).

[43] G. Nienhuis and F. Schuller, Spontaneous emission and light scattering by atomic lattice models, *J. Phys. B* **20**, 23 (1987).

[44] R. G. DeVoe and R. G. Brewer, Observation of Superradiant and Subradiant Spontaneous Emission of Two Trapped Ions, *Phys. Rev. Lett.* **76**, 2049 (1996).

[45] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.125.263601> for further information on correlation functions between operators, the limit $d \rightarrow 0$, the impact of geometry on the decay, and a more detailed discussion of experimental considerations, which includes Refs. [46–50].

[46] L. Ostermann, D. Plankensteiner, H. Ritsch, and C. Genes, Protected subspace Ramsey spectroscopy, *Phys. Rev. A* **90**, 053823 (2014).

[47] N. Schlosser, G. Reymond, and P. Grangier, Collisional Blockade in Microscopic Optical Dipole Traps, *Phys. Rev. Lett.* **89**, 023005 (2002).

[48] T. Grünzweig, A. Hilliard, M. McGovern, and M. F. Andersen, Near-deterministic preparation of a single atom in an optical microtrap, *Nat. Phys.* **6**, 951 (2010).

[49] Y. H. Fung and M. F. Andersen, Efficient collisional blockade loading of a single atom into a tight microtrap, *New J. Phys.* **17**, 073011 (2015).

[50] M. O. Brown, T. Thiele, C. Kiehl, T.-W. Hsu, and C. A. Regal, Gray-Molasses Optical-Tweezer Loading: Controlling Collisions for Scaling Atom-Array Assembly, *Phys. Rev. X* **9**, 011057 (2019).

[51] S. Wolf, S. Richter, J. von Zanthier, and F. Schmidt-Kaler, Light of Two Atoms in Free Space: Bunching or Anti-bunching?, *Phys. Rev. Lett.* **124**, 063603 (2020).

[52] F. T. Arecchi, E. Courtens, R. Gilmore, and H. Thomas, Atomic coherent states in quantum optics, *Phys. Rev. A* **6**, 2211 (1972).

[53] W. Guerin, M. O. Araújo, and R. Kaiser, Subradiance in a Large Cloud of Cold Atoms, *Phys. Rev. Lett.* **116**, 083601 (2016).

[54] A. Cipris, N. A. Moreira, T. S. do Espírito Santo, P. Weiss, C. J. Villas-Boas, R. Kaiser, W. Guerin, and R. Bachelard, Subradiance with saturated atoms: Population enhancement of the long-lived states, [arXiv:2009.05172](https://arxiv.org/abs/2009.05172).

[55] P. Solano, P. Barberis-Blostein, F. K. Fatemi, L. A. Orozco, and S. L. Rolston, Super-radiance reveals infinite-range dipole interactions through a nanofiber, *Nat. Commun.* **8**, 1857 (2017).

[56] M. Lepers, J.-F. Wyart, and O. Dulieu, Anisotropic optical trapping of ultracold erbium atoms, *Phys. Rev. A* **89**, 022505 (2014).

[57] H. Li, J.-F. Wyart, O. Dulieu, S. Nascimbène, and M. Lepers, Optical trapping of ultracold dysprosium atoms: Transition probabilities, dynamic dipole polarizabilities and van der Waals c_6 coefficients, *J. Phys. B* **50**, 014005 (2016).

[58] J. Pellegrino, R. Bourgain, S. Jennewein, Y. R. P. Sortais, A. Browaeys, S. D. Jenkins, and J. Ruostekoski, Observation of Suppression of Light Scattering Induced by Dipole-Dipole Interactions in a Cold-Atom Ensemble, *Phys. Rev. Lett.* **113**, 133602 (2014).

[59] L. Corman, J. L. Ville, R. Saint-Jalm, M. Aidelsburger, T. Bienaimé, S. Nascimbène, J. Dalibard, and J. Beugnon, Transmission of near-resonant light through a dense slab of cold atoms, *Phys. Rev. A* **96**, 053629 (2017).

[60] C. Palacios-Berraquero, D. M. Kara, A. R.-P. Montblanch, M. Barbone, P. Latawiec, D. Yoon, A. K. Ott, M. Loncar, A. C. Ferrari, and M. Atatüre, Large-scale quantum-emitter arrays in atomically thin semiconductors, *Nat. Commun.* **8**, 053629 (2017).

[61] N. V. Proscia, Z. Shotan, H. Jayakumar, P. Reddy, C. Cohen, M. Dollar, A. Alkauskas, M. Doherty, C. A. Meriles, and V. M. Menon, Near-deterministic activation of room-temperature quantum emitters in hexagonal boron nitride, *Optica* **5**, 1128 (2018).

[62] W. P. Smith, J. E. Reiner, L. A. Orozco, S. Kuhr, and H. M. Wiseman, Capture and Release of a Conditional State of a Cavity QED System by Quantum Feedback, *Phys. Rev. Lett.* **89**, 133601 (2002).