



TeV Emission of Galactic Plane Sources with HAWC and H.E.S.S.

H. Abdalla¹, F. Aharonian^{2,3,4}, F. Ait Benkhali³, E. O. Angüner⁵, C. Arcaro⁶, C. Armand⁷, T. Armstrong⁸, H. Ashkar⁹, M. Backes^{1,6}, V. Baghmanyan^{10,*}, V. Barbosa Martins¹¹, A. Barnacka¹², M. Barnard⁶, Y. Becherini¹³, D. Berge¹¹, K. Bernlöhr³, B. Bi¹⁴, M. Böttcher⁶, C. Boisson¹⁵, J. Bolmont¹⁶, M. de Bonyde Lavergne⁷, M. Breuhaus³, R. Brose², F. Brun^{9,†}, P. Brun⁹, M. Bryan¹⁷, M. Büchele¹⁸, T. Bulik¹⁹, T. Bylund¹³, S. Caroff⁷, A. Carosi⁷, T. Chand⁶, S. Chandra⁶, A. Chen²⁰, G. Cotter⁸, M. Curylo¹⁹, J. Damascene Mbarubucyeye¹¹, I. D. Davids¹, J. Davies⁸, C. Deil³, J. Devin²¹, L. Dirson²², A. Djannati-Atai²¹, A. Dmytriiev¹⁵, A. Donath³, V. Doroshenko¹⁴, L. Dreyer⁶, C. Duffy²³, J. Dyks²⁴, K. Egberts²⁵, F. Eichhorn¹⁸, S. Einecke²⁶, G. Emery¹⁶, J.-P. Ernenwein⁵, K. Feijen²⁶, S. Fegan²⁷, A. Fiasson⁷, G. Fichet de Clairfontaine¹⁵, G. Fontaine²⁷, S. Funk¹⁸, M. Fülling¹¹, S. Gabici²¹, Y. A. Gallant²⁸, G. Giavitto¹¹, L. Giunti^{9,21}, D. Glawion¹⁸, J. F. Glicenstein⁹, D. Gottschall¹⁴, M.-H. Grondin²⁹, J. Hahn³, M. Haupt¹¹, G. Hermann³, J. A. Hinton^{3,*}, W. Hofmann³, C. Hoischen²⁵, T. L. Holch¹¹, M. Holler³⁰, M. Hörbe⁸, D. Horns²², D. Huber³⁰, M. Jamroz¹², D. Jankowsky¹⁸, F. Jankowsky³¹, I. Jung-Richardt¹⁸, E. Kasai¹, M. A. Kastendieck²², K. Katarzyński³², U. Katz¹⁸, D. Khangulyan³³, B. Khélifi²¹, S. Klepser¹¹, W. Kluźniak²⁴, Nu. Komin²⁰, R. Konno¹¹, K. Kosack⁹, D. Kostunin¹¹, M. Kreter⁶, G. Lamanna⁷, A. Lemièrre²¹, M. Lemoine-Goumard²⁹, J.-P. Lenain¹⁶, F. Leuschner¹⁴, C. Levy¹⁶, T. Lohse³⁴, I. Lypova¹¹, J. Mackey², J. Majumdar¹¹, D. Malyshev¹⁴, D. Malyshev¹⁸, V. Marandon^{3,*}, P. Marchegiani²⁰, A. Marcowith²⁸, A. Mares²⁹, G. Martí-Devesa³⁰, R. Marx^{3,31}, G. Maurin⁷, P. J. Meintjes³⁵, M. Meyer¹⁸, A. M. W. Mitchell³, R. Moderski²⁴, L. Mohrmann¹⁸, A. Montanari⁹, C. Moore²³, P. Morris⁸, E. Moulin⁹, J. Muller²⁷, T. Murach¹¹, K. Nakashima¹⁸, A. Nayerhoda¹⁰, M. de Naurois²⁷, H. Ndiyavala⁶, J. Niemiec¹⁰, L. Oakes³⁴, P. O'Brien²³, H. Odaka³⁶, S. Ohm¹¹, L. Olivera-Nieto^{3,*}, E. de Ona Wilhelmi¹¹, M. Ostrowski¹², M. Panter³, S. Panny³⁰, R. D. Parsons³⁴, G. Peron³, B. Peyaud⁹, Q. Piel⁷, S. Pita²¹, V. Poireau⁷, A. Priyana Noel¹², D. A. Prokhorov¹⁷, H. Prokoph¹¹, G. Pühlhofer¹⁴, M. Punch^{13,21}, A. Quirrenbach³¹, S. Raab¹⁸, R. Rauth³⁰, P. Reichherzer⁹, A. Reimer³⁰, O. Reimer³⁰, Q. Remy³, M. Renaud²⁸, F. Rieger³, L. Rinchiuso⁹, C. Romoli³, G. Rowell²⁶, B. Rudak²⁴, V. Sahakian³⁷, S. Sailer³, H. Salzmann¹⁴, D. A. Sanchez⁷, A. Santangelo¹⁴, M. Sasaki¹⁸, J. Schäfer¹⁸, F. Schüssler⁹, H. M. Schutte⁶, U. Schwanke³⁴, M. Seglar-Arroyo⁹, M. Senniappan¹³, A. S. Seyffert⁶, N. Shafi²⁰, J. N. S. Shapopi¹, K. Shiningayamwe¹, R. Simoni¹⁷, A. Sinha²¹, H. Sol¹⁵, A. Specovius¹⁸, S. Spencer⁸, M. Spir-Jacob²¹, Ł. Stawarz¹², L. Sun¹⁷, R. Steenkamp¹, C. Stegmann^{11,25}, S. Steinmassl³, C. Steppa²⁵, T. Takahashi³⁸, T. Tavernier⁹, A. M. Taylor¹¹, R. Terrier²¹, J. H. E. Thiersen⁶, D. Tiziani¹⁸, M. Tluczykont²², L. Tomankova¹⁸, C. Trichard²⁷, M. Tsiros³, R. Tuffs³, Y. Uchiyama³³, D. J. van der Walt⁶, C. van Eldik¹⁸, C. van Rensburg¹, B. van Soelen³⁵, G. Vasileiadis²⁸, J. Veh¹⁸, C. Venter⁶, P. Vincent¹⁶, J. Vink¹⁷, H. J. Völk³, Z. Wadiasinh⁶, S. J. Wagner³¹, J. Watson⁸, F. Werner^{3,*}, R. White³, A. Wiercholska^{10,31}, Yu Wun Wong¹⁸, A. Yusafzai¹⁸, M. Zacharias^{6,15}, R. Zanin³, D. Zargaryan^{2,4}, A. A. Zdziarski²⁴, A. Zech¹⁵, S. J. Zhu¹¹, A. Zmija¹⁸, J. Zorn³, S. Zouari²¹, N. Żywucka⁶

(H.E.S.S. Collaboration),

and

A. Albert³⁹, R. Alfaro⁴⁰, C. Alvarez⁴¹, J. C. Arteaga-Velázquez⁴², K. P. Arunbabu⁴³, D. Avila Rojas⁴⁰, E. Belmont-Moreno⁴⁰, S. Y. BenZvi⁴⁴, C. Brisbois⁴⁵, K. S. Caballero-Mora⁴¹, T. Capistrán⁴⁶, A. Carramiñana⁴⁷, S. Casanova^{48,*}, U. Cotti⁴², J. Cotzomi⁴⁹, S. Coutiño de León⁴⁷, E. De la Fuente^{50,51}, C. de León⁴², R. Diaz Hernandez⁴⁷, J. C. Díaz-Vélez⁵⁰, B. L. Dingus³⁹, M. A. DuVernois⁵², M. Durocher³⁹, R. W. Ellsworth⁴⁵, K. Engel⁴⁵, C. Espinoza⁴⁰, K. L. Fan⁴⁵, M. Fernández Alonso⁵³, N. Fraija⁴⁶, A. Galván-Gámez⁴⁶, D. Garcia⁴⁰, J. A. García-González⁵⁴, F. Garfias⁴⁶, G. Giacinti^{3,*}, M. M. González⁴⁶, J. A. Goodman⁴⁵, J. P. Harding³⁹, S. Hernandez⁴⁰, B. Hona⁵⁵, D. Huang⁵⁶, F. Hueyotl-Zahuantitla⁴¹, P. Hüntemeyer⁵⁶, A. Iriarte⁴⁶, A. Jardin-Blicq^{3,57,58,*}, V. Joshi^{59,*}, D. Kieda⁵⁵, W. H. Lee⁴⁶, H. León Vargas⁴⁰, J. T. Linnemann⁶⁰, A. L. Longinotti⁴⁷, G. Luis-Raya⁶¹, R. López-Coto⁶², K. Malone³⁹, O. Martinez⁶³, I. Martinez-Castellanos⁴⁵, J. Martínez-Castro⁶⁴, J. A. Matthews⁶⁵, P. Miranda-Romagnoli⁶⁶, J. A. Morales-Soto⁴², E. Moreno⁶³, M. Mostafa⁵³, A. Nayerhoda⁴⁸, L. Nellen⁶⁷, M. Newbold⁵⁵, M. U. Nisa⁶⁰, R. Noriega-Papaqui⁶⁶, N. Omodei⁶⁸, A. Peisker⁶⁰, Y. Pérez Araujo⁴⁶, E. G. Pérez-Pérez⁶¹, C. D. Rho⁶⁹, D. Rosa-González⁴⁷, E. Ruiz-Velasco^{3,*}, F. Salesa Greus^{48,69}, A. Sandoval⁴⁰, M. Schneider⁴⁵, H. Schoorlemmer³, J. Serna-Franco⁴⁰, A. J. Smith⁴⁵, R. W. Springer⁵⁵, P. Surajbali³, K. Tollefson⁶⁰, I. Torres⁴⁷, R. Torres-Escobedo⁵⁰, R. Turner⁷⁰, F. Ureña-Mena⁴⁷, L. Villaseñor⁶³, T. Weisgarber⁷¹, E. Willox⁴⁵, and H. Zhou⁷²

(HAWC Collaboration)

¹ University of Namibia, Department of Physics, Private Bag 13301, Windhoek 10005, Namibia² Dublin Institute for Advanced Studies, 31 Fitzwilliam Place, Dublin 2, Ireland³ Max-Planck Institute for Nuclear Physics, D-69117 Heidelberg, Germany; armelle.jardin-blicq@mpi-hd.mpg.de, contact.hess@hess-experiment.eu⁴ High Energy Astrophysics Laboratory, RAU, 123 Hovsep Emin Street, Yerevan 0051, Armenia⁵ Aix Marseille Université, CNRS/IN2P3, CPPM, Marseille, France⁶ Centre for Space Research, North-West University, Potchefstroom 2520, South Africa⁷ Laboratoire d'Annecy de Physique des Particules, Université Grenoble Alpes, Université Savoie Mont Blanc, CNRS, LAPP, F-74000 Annecy, France

- ⁸ University of Oxford, Department of Physics, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, UK
⁹ IRFU, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France; contact.hess@hess-experiment.eu
¹⁰ Instytut Fizyki Jądrowej PAN, ul. Radzikowskiego 152, 31-342 Kraków, Poland
¹¹ DESY, D-15738 Zeuthen, Germany
¹² Obserwatorium Astronomiczne, Uniwersytet Jagielloński, ul. Orla 171, 30-244 Kraków, Poland
¹³ Department of Physics and Electrical Engineering, Linnaeus University, 351 95 Växjö, Sweden
¹⁴ Institut für Astronomie und Astrophysik, Universität Tübingen, Sand 1, D-72076 Tübingen, Germany
¹⁵ Laboratoire Univers et Théories, Observatoire de Paris, Université PSL, CNRS, Université de Paris, F-92190 Meudon, France
¹⁶ Sorbonne Université, Université Paris Diderot, Sorbonne Paris Cité, CNRS/IN2P3, Laboratoire de Physique Nucléaire et de Hautes Energies, LPNHE, 4 Place Jussieu, F-75252 Paris, France
¹⁷ GRAPPA, Anton Pannekoek Institute for Astronomy, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands
¹⁸ Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen Centre for Astroparticle Physics, Erwin-Rommel-Straße 1, D-91058 Erlangen, Germany
¹⁹ Astronomical Observatory, The University of Warsaw, Al. Ujazdowskie 4, 00-478 Warsaw, Poland
²⁰ School of Physics, University of the Witwatersrand, 1 Jan Smuts Avenue, Braamfontein, Johannesburg, 2050 South Africa
²¹ Université de Paris, CNRS, Astroparticule et Cosmologie, F-75013 Paris, France
²² Universität Hamburg, Institut für Experimentalphysik, Luruper Chaussee 149, D-22761 Hamburg, Germany
²³ Department of Physics and Astronomy, The University of Leicester, University Road, Leicester, LE1 7RH, UK
²⁴ Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences, ul. Bartycka 18, 00-716 Warsaw, Poland
²⁵ Institut für Physik und Astronomie, Universität Potsdam, Karl-Liebknecht-Strasse 24/25, D-14476 Potsdam, Germany
²⁶ School of Physical Sciences, University of Adelaide, Adelaide 5005, Australia
²⁷ Laboratoire Leprince-Ringuet, École Polytechnique, CNRS, Institut Polytechnique de Paris, F-91128 Palaiseau, France
²⁸ Laboratoire Univers et Particules de Montpellier, Université Montpellier, CNRS/IN2P3, CC 72, Place Eugène Bataillon, F-34095 Montpellier Cedex 5, France
²⁹ Université Bordeaux, CNRS/IN2P3, Centre d'Études Nucléaires de Bordeaux Gradignan, F-33175 Gradignan, France
³⁰ Institut für Astro- und Teilchenphysik, Leopold-Franzens-Universität Innsbruck, A-6020 Innsbruck, Austria
³¹ Landessternwarte, Universität Heidelberg, Königstuhl, D-69117 Heidelberg, Germany
³² Institute of Astronomy, Faculty of Physics, Astronomy and Informatics, Nicolaus Copernicus University, Grudziadzka 5, 87-100 Torun, Poland
³³ Department of Physics, Rikkyo University, 3-34-1 Nishi-Ikebukuro, Toshima-ku, Tokyo 171-8501, Japan
³⁴ Institut für Physik, Humboldt-Universität zu Berlin, Newtonstr. 15, D-12489 Berlin, Germany
³⁵ Department of Physics, University of the Free State, PO Box 339, Bloemfontein 9300, South Africa
³⁶ Department of Physics, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan
³⁷ Yerevan Physics Institute, 2 Alikhanian Brothers Street, 375036 Yerevan, Armenia
³⁸ Kavli Institute for the Physics and Mathematics of the Universe (WPI), The University of Tokyo Institutes for Advanced Study (UTIAS), The University of Tokyo, 5-1-5 Kashiwa-no-Ha, Kashiwa, Chiba, 277-8583, Japan
³⁹ Physics Division, Los Alamos National Laboratory, Los Alamos, NM, USA
⁴⁰ Instituto de Física, Universidad Nacional Autónoma de México, Ciudad de México, Mexico
⁴¹ Universidad Autónoma de Chiapas, Tuxtla Gutiérrez, Chiapas, Mexico
⁴² Universidad Michoacana de San Nicolás de Hidalgo, Morelia, Mexico
⁴³ Instituto de Geofísica, Universidad Nacional Autónoma de México, Ciudad de México, Mexico
⁴⁴ Department of Physics & Astronomy, University of Rochester, Rochester, NY, USA
⁴⁵ Department of Physics, University of Maryland, College Park, MD, USA
⁴⁶ Instituto de Astronomía, Universidad Nacional Autónoma de México, Ciudad de México, Mexico
⁴⁷ Instituto Nacional de Astrofísica, Óptica y Electrónica, Puebla, Mexico
⁴⁸ Institute of Nuclear Physics Polish Academy of Sciences, PL-31342 IFJ-PAN, Krakow, Poland
⁴⁹ Facultad de Ciencias Físico Matemáticas, Benemérita Universidad Autónoma de Puebla, Puebla, Mexico
⁵⁰ Departamento de Física, Centro Universitario de Ciencias Exactas e Ingenierías, Universidad de Guadalajara, Guadalajara, Mexico
⁵¹ Institute for Cosmic Ray Research, University of Tokyo, 277-8582 Chiba, Kashiwa, Kashiwanoha, 5 Chome-1-5, Japan
⁵² Department of Physics, University of Wisconsin-Madison, Madison, WI, USA
⁵³ Department of Physics, Pennsylvania State University, University Park, PA, USA
⁵⁴ Tecnológico de Monterrey, Escuela de Ingeniería y Ciencias, Ave. Eugenio Garza Sada 2501, Monterrey, N.L., 64849, Mexico
⁵⁵ Department of Physics and Astronomy, University of Utah, Salt Lake City, UT, USA
⁵⁶ Department of Physics, Michigan Technological University, Houghton, MI, USA
⁵⁷ Department of Physics, Faculty of Science, Chulalongkorn University, 254 Phayathai Road, Pathumwan, Bangkok 10330, Thailand
⁵⁸ National Astronomical Research Institute of Thailand (Public Organization), Don Kao, Mae Rim, Chiang Mai 50180, Thailand
⁵⁹ Erlangen Centre for Astroparticle Physics, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, Germany
⁶⁰ Department of Physics and Astronomy, Michigan State University, East Lansing, MI, USA
⁶¹ Universidad Politécnica de Pachuca, Pachuca, Hgo, Mexico
⁶² INFN and Università di Padova, via Marzolo 8, I-35131, Padova, Italy
⁶³ Facultad de Ciencias Físico Matemáticas, Benemérita Universidad Autónoma de Puebla, Puebla, Mexico
⁶⁴ Centro de Investigación en Computación, Instituto Politécnico Nacional, México City, Mexico
⁶⁵ Department of Physics and Astronomy, University of New Mexico, Albuquerque, NM, USA
⁶⁶ Universidad Autónoma del Estado de Hidalgo, Pachuca, Mexico
⁶⁷ Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Ciudad de México, Mexico
⁶⁸ Department of Physics, Stanford University, Stanford, CA 94305060, USA
⁶⁹ Instituto de Física Corpuscular, CSIC, Universitat de València, E-46980, Paterna, Valencia, Spain
⁷⁰ Department of Physics, Michigan Technological University, Houghton, MI, USA
⁷¹ Department of Physics, University of Wisconsin-Madison, Madison, WI, USA
⁷² Tsung-Dao Lee Institute & School of Physics and Astronomy, Shanghai Jiao Tong University, Shanghai, People's Republic of China

Received 2021 March 8; revised 2021 April 7; accepted 2021 April 7; published 2021 August 9

* Author in both collaborations.

† H.E.S.S. corresponding author. For questions concerning H.E.S.S. results contact contact.hess@hess-experiment.eu.

‡ HAWC corresponding author. For questions concerning HAWC results contact armelle.jardin-blicq@mpi-hd.mpg.de.

Abstract

The High Altitude Water Cherenkov (HAWC) observatory and the High Energy Stereoscopic System (H.E.S.S.) are two leading instruments in the ground-based very-high-energy γ -ray domain. HAWC employs the water Cherenkov detection (WCD) technique, while H.E.S.S. is an array of Imaging Atmospheric Cherenkov Telescopes (IACTs). The two facilities therefore differ in multiple aspects, including their observation strategy, the size of their field of view, and their angular resolution, leading to different analysis approaches. Until now, it has been unclear if the results of observations by both types of instruments are consistent: several of the recently discovered HAWC sources have been followed up by IACTs, resulting in a confirmed detection only in a minority of cases. With this paper, we go further and try to resolve the tensions between previous results by performing a new analysis of the H.E.S.S. Galactic plane survey data, applying an analysis technique comparable between H.E.S.S. and HAWC. Events above 1 TeV are selected for both data sets, the point-spread function of H.E.S.S. is broadened to approach that of HAWC, and a similar background estimation method is used. This is the first detailed comparison of the Galactic plane observed by both instruments. H.E.S.S. can confirm the γ -ray emission of four HAWC sources among seven previously undetected by IACTs, while the three others have measured fluxes below the sensitivity of the H.E.S.S. data set. Remaining differences in the overall γ -ray flux can be explained by the systematic uncertainties. Therefore, we confirm a consistent view of the γ -ray sky between WCD and IACT techniques.

Unified Astronomy Thesaurus concepts: [High energy astrophysics \(739\)](#); [Gamma-ray observatories \(632\)](#); [Gamma-ray astronomy \(628\)](#); [Surveys \(1671\)](#)

1. Introduction

The High Energy Stereoscopic System (H.E.S.S.) telescope array and the High Altitude Water Cherenkov (HAWC) observatory are examples of instruments using the two main techniques developed in very high energy (VHE) γ -ray astronomy. H.E.S.S. is an array of five Imaging Atmospheric Cherenkov Telescopes (IACTs) that image the Cherenkov light produced by the atmospheric air showers. H.E.S.S. is located in Namibia at a latitude of approximately 23° south, and at an altitude of 1800 m. Its collection area is of the order of 10^5 m^2 above 1 TeV. H.E.S.S. has a field of view of 5° in diameter, with a γ -ray sensitivity roughly uniform for the innermost 2° that gradually drops toward the edges. Its point-spread function (PSF) is better than 0.1° above 1 TeV for zenith angles less than 30° (Aharonian et al. 2006). H.E.S.S. is a pointing instrument observing only at night. Including CT5, the biggest telescope at the center of the array, it is sensitive in the energy range 30 GeV–100 TeV, and its energy resolution is better than 15% above 1 TeV. H.E.S.S. surveyed the Galactic plane from 2004 to 2013, producing the H.E.S.S. Galactic Plane Survey (HGPS; H.E.S.S. Collaboration et al. 2018a) based on 2700 hours of selected data, with 78 detected γ -ray sources. The survey covers the inner Galactic plane from 250° to 65° in longitude and from -3° to $+3^\circ$ in latitude, extending up to -5° and $+5^\circ$ in some regions. This observation program was carried out by the four middle-size telescopes, known as H.E.S.S. phase I. The energy threshold for the hard cuts used in making emission maps is, for most ranges of Galactic longitude, between 400 and 700 GeV (see Figure 2 of H.E.S.S. Collaboration et al. (2018a)). The achieved sensitivity is better than 2% of the Crab flux, assuming a point source with a spectral index of -2.3 , and improves to 0.5% in some regions with high observation time, like in the vicinity of the Galactic Center.

The second observatory, HAWC, is an array of 300 water Cherenkov detectors (WCDs) located at a latitude of about 19° north in Mexico, at an altitude of 4100 m. The collection area of HAWC is of the order of the physical area of the detector, $\sim 10^4 \text{ m}^2$ above 500 GeV for a source at the Crab decl. HAWC detects the Cherenkov light produced in the water by the individual charged particles of air showers. Since HAWC is a survey instrument, it continuously monitors the sky above it. It has

an instantaneous field of view of $\sim 2 \text{ sr}$ and can observe two-thirds of the sky from $\sim -20^\circ$ to $\sim +60^\circ$ in decl., every day. The achieved sensitivity is $\sim 2\%$ of the Crab flux for sources with decl. $0^\circ < \delta < 40^\circ$ assuming a spectral index of -2.5 , and degrades to 8% of the Crab flux for sources at the edge of HAWC observable sky ($\delta = -20^\circ$ and $+60^\circ$). HAWC is sensitive in the energy range between $\sim 300 \text{ GeV}$ to above 100 TeV, but is optimized to γ rays with a primary energy above $\sim 1 \text{ TeV}$. The detected events are allocated to one of the nine analysis bins, depending on the fraction of the array that triggered, from analysis bin 1 gathering events triggering 7%–10% of the array, to analysis bin 9 for events hitting 84%–100% of the array. This binning is linked to the energy of the incoming particle, since events triggering only a small fraction of the array are likely to have lower energy than events triggering most of the array. However, there is some degeneracy and each analysis bin spans roughly one order of magnitude of energy, making HAWC's energy resolution quite poor as shown by Abeysekara et al. (2017a), Figure 3. The PSF of HAWC is $\sim 1^\circ$ for the first analysis bin and improves to $\sim 0.2^\circ$ for the ninth analysis bin, as depicted in Figure 9 of Abeysekara et al. (2017a).

The HAWC Collaboration published its 2HWC catalog based on 507 days of data (Abeysekara et al. 2017b) as the result of the first source search performed with the complete HAWC detector. More recently, using three times more data, the HAWC Collaboration published the 3HWC catalog (Albert et al. 2020). In this search, 65 sources have been reported, among them 20 detected for the first time at TeV energies. The newly detected unassociated sources were subsequently studied by IACT collaborations who looked for counterparts. The VERITAS Collaboration (Very Energetic Radiation Imaging Telescope Array System), for example, looked at 14 2HWC sources and confirmed only one (Abeysekara et al. 2018). Archival data taken with the Major Atmospheric Gamma Imaging Cherenkov (MAGIC) telescopes were analyzed again to look for three unassociated 2HWC sources, without detecting any significant emission (Ahnen et al. 2019). One explanation would be that these sources might be relatively extended, with a low surface brightness and possibly a hard spectrum. IACTs, with their limited field of view and their very good angular resolution, may not be sensitive to angularly extended objects with low surface brightness, using their standard analysis. We show in this paper

that a dedicated analysis of IACT data is required to detect the extended and faint HAWC sources.

IACTs and WCDs have already demonstrated that they complement each other, with their overlapping observable sky and similar energy range yet different field of view and angular resolution. One example is the source MGRO J1908+06, first detected by the Milagro observatory (Abdo et al. 2007), the predecessor of HAWC, and confirmed by H.E.S.S. (Aharonian et al. 2009). In this paper we go further in exploiting their complementarity. Using a new analysis for the H.E.S.S. data, we present a comparison of the part of the Galactic plane common to H.E.S.S. and HAWC. After describing the data sets in Section 2, Section 3 presents the new analysis method that aims at producing a H.E.S.S. Galactic plane map in the most similar way to HAWC. The results are presented in Section 4 and the conclusion in Section 5.

2. Data Set

2.1. HAWC Data

The analysis presented here uses 1523 days of data, taken between 2014 November 26 and 2019 June 3 and makes use of events falling in the analysis bins 4–9. As defined in Abeysekara et al. (2017a), these are events that triggered more than $\sim 25\%$ of the array. The main motivation for this selection is to have a reasonably good reconstruction, sufficient statistics, and a reasonable PSF. These cuts result in an energy threshold of approximately 1 TeV for a source at the Crab decl., increasing toward the central Galaxy. The PSF of the instrument is 0.4° for bin 4 and decreases to less than 0.2° for the highest bin.

2.2. H.E.S.S. Data

The maps used to perform the H.E.S.S. analysis presented here have been made using the HGPS data set (H.E.S.S. Collaboration et al. 2018a) for Galactic longitude $10^\circ < \ell < 60^\circ$, reconstructed with the Image Pixel-wise fit for Atmospheric Cherenkov Telescopes (ImPACT) algorithm (Parsons & Hinton 2014). ImPACT uses a maximum likelihood approach to determine the best-fit shower parameters instead of using the traditional parameterization of the camera images (Cawley et al. 1985), which was used in the published HGPS (H.E.S.S. Collaboration et al. 2018a). Using ImPACT reconstruction, the PSF of H.E.S.S. is about 0.06° at 1 TeV for zenith angles smaller than 30° and its energy resolution is $\sim 10\%$ above 1 TeV. In this analysis, events above 1 TeV are selected to match the HAWC energy range.

The results presented here have been verified with a different calibration, reconstruction, and gamma/hadron separation method using a semianalytical description of the air shower (de Naurois & Rolland 2009).

3. Analysis

For both data sets described in the previous section, a test statistic (TS) map is created with Gammapy (Deil et al. 2017) using a data map, a background map, and an exposure map. The TS is obtained by computing the likelihood ratio between the hypothesis of a source model, and a null hypothesis assuming no source but background only, for the test source centered on each pixel of the sky map. The source image through the instrument is assumed to have the shape of a Gaussian with a fixed angular extent σ . The flux normalization is the only free parameter. According to Wilks’s theorem (Wilks 1938), the distribution of

the TS under the null hypothesis follows asymptotically a χ^2 distribution with one degree of freedom (see Appendix D). The statistical significance is then equal to $\pm\sqrt{\text{TS}}$. Note that the significance mentioned thereafter is not corrected for trials. The flux is obtained by dividing the excess that maximizes the likelihood by the γ -ray exposure. The first row of Figure 1 shows the significance maps of the Galactic plane detected by H.E.S.S., above ~ 1 TeV, made in a similar way to the published HGPS: using a Gaussian of size $\sigma = 0.1^\circ$ and the ring background method (described in H.E.S.S. Collaboration et al. (2018a)). The background estimate is normalized in a ring region surrounding the sources, where detected sources or suspected γ -ray emission have been excluded. A spectral index of -2.5 is used, while it is -2.3 in the HGPS.

For the HAWC map, due to the large variation of the PSF with analysis bin, the TS map is produced by a maximum likelihood method over all analysis bins 4–9. A simple power law with a spectral index of -2.5 is used as the spectral assumption. More details can be found in Abeysekara et al. (2017b). A new catalog search has been performed on the HAWC data selected for this analysis, following the procedure described in Albert et al. (2020) and assuming sources that are point-like on the scale of HAWC angular resolution. The HAWC sources presented here, using the prefix HAWC at the beginning of their name, result from this new source search. They are listed in Table 2, in Appendix A, with their fitted position and their counterpart in the 3HWC catalog.

In the spirit of making the Galactic plane maps from both instruments in the most similar way, a new H.E.S.S. TS map is computed using a Gaussian of $\sigma = 0.4^\circ$, comparable to the size of the largest HAWC PSF for our data set. It corresponds to the PSF of the analysis bin 4 where the most events are expected. This has as direct consequence to decrease the flux sensitivity as more background is included, which is visible in the Figure 2. The corresponding significance map is displayed in the second row of Figure 1. Finally, another background estimation method called the field-of-view background method (Berge et al. 2007) is used, as it is the closest one to the HAWC direct integration background method (Abeysekara et al. 2017a). The field-of-view method uses as background the H.E.S.S. exposure map, i.e., the expected events from cosmic-ray background, taking advantage of the whole field of view of H.E.S.S. and not only a small area contained in a ring around the source position. The shape of the background map is estimated from regions of the sky at similar zenith angle, excluding known VHE γ -ray sources. The map is normalized to the number of events outside the exclusion regions. However, note that using an average exposure assumes that its shape does not change with time and is not an instantaneous measurement as it does not fully account for variations in the instrument response over time, contrary to the HAWC direct integration method. Moreover, the required normalization step uses the number of events outside exclusion regions in the field of view, which has a limited size. The third map of Figure 1 shows the H.E.S.S. significance map produced using a Gaussian of 0.4° , similarly to the map in the second row, but using the field-of-view background method instead of the ring background method. Furthermore, an additional exclusion region defined by a 2° wide strip centered on the Galactic plane ($-1^\circ < b < 1^\circ$) is used, to avoid any contamination from the Galactic plane large scale γ -ray emission to the background estimation. This is the H.E.S.S. map produced with the most similar analysis method to the HAWC one, visible on the fourth row.




Figure 1. HAWC and H.E.S.S. Galactic plane maps. The green circles are the 68% containment of the HGPS sources and the black dots are the location of the HAWC sources. The regions in the white boxes are highlighted in Figure 3. The green dotted lines point to H.E.S.S. sources undetected by HAWC. The orange dotted lines point to HAWC sources previously undetected by H.E.S.S. (see Section 4.2). From top to bottom: (1) H.E.S.S. Galactic plane map for $E > 1$ TeV, using ImPACT reconstruction and a Gaussian kernel of size 0.1° ; the ring background method is applied on each observation run separately, with an adaptive radius. The standard exclusion regions around sources are used. (2) Same H.E.S.S. data, using a Gaussian kernel of 0.4° . (3) Same H.E.S.S. data, using the field-of-view background method and a Gaussian kernel of 0.4° . For the background normalization, in addition to the standard exclusion regions around sources, a 2° wide exclusion band covering the Galactic plane is used. (4) HAWC Galactic plane map with 1523 days of data, using events in the analysis bins 4–9.

4. Results and Discussion

4.1. HAWC and H.E.S.S. Original Maps

The H.E.S.S. significance map in the first row of Figure 1, produced in a traditional way with a Gaussian size $\sigma = 0.1^\circ$ and the ring background method, shows some important differences compared to the HAWC significance map, in the last row of the same figure. The corresponding point-like sensitivity curves at $b = 0^\circ$ are displayed in the left panel of Figure 2 in dark blue and dashed orange, respectively, for the part of the Galactic plane studied here. The right panel of Figure 2 shows the detection horizon corresponding to the sensitivity curves plotted in the left panel of the same figure, for a source luminosity of 10^{34} erg s $^{-1}$. It illustrates the difference in exposure between both instruments: H.E.S.S. has an exposure time that varies significantly from one

region to another, while the HAWC exposure changes very smoothly across the sky. It also illustrates the part of the sky covered by HAWC and by H.E.S.S. for the HGPS.

Among the 29 sources detected by H.E.S.S. in the selected region of the HGPS, HAWC detected significant emission at the location of 27 of them, even though only 15 sources are actually resolved. The two HGPS sources that do not show significant emission in the HAWC map are HESS J1943+213, which is likely to be an extragalactic source seen in the Galactic plane (H.E.S.S. Collaboration et al. 2018a) and HESS J1911+090 associated with the supernova remnant (SNR) W49B. This is expected because the flux of these sources lies below the HAWC sensitivity curve, as shown in Figure 2. On the other hand, HAWC detected 22 sources and only 15 have a counterpart in the HGPS catalog. Hence, seven sources are detected only by HAWC.




Figure 2. Left: sensitivity curves at $b = 0^\circ$ corresponding to the first (H.E.S.S. point-like source analysis, dark blue curve), third (new H.E.S.S. extended source analysis, light blue curve), and fourth maps (HAWC, dashed orange) of Figure 1. The orange and blue points represent the flux in Crab units of the seven HAWC sources that were previously undetected by H.E.S.S., obtained using the analysis presented here at the location of the HAWC sources. The black borders highlight the four HAWC sources newly detected by H.E.S.S., and they lie above the sensitivity curve corresponding to this analysis. The flux values can be found in Table 2 of Appendix A. The green points correspond to the flux of the two H.E.S.S. sources not detected by HAWC and they lie below the HAWC sensitivity curve. Note that 1 crab unit is equal to $2.26 \times 10^{-11} \text{ cm}^{-2} \text{ s}^{-1}$ for H.E.S.S. (from H.E.S.S. Collaboration et al. (2018a)) and $2.24 \times 10^{-11} \text{ cm}^{-2} \text{ s}^{-1}$ for HAWC (from Albert et al. (2020)). Right: face-on view of the galaxy. The H.E.S.S. horizon at $b = 0^\circ$ for a 5σ detection of a point-like source is depicted in dark blue. The H.E.S.S. horizon corresponding to this analysis is depicted in light blue. The HAWC horizon is shown in dashed orange. The horizons are produced using the sensitivity curves from the left plot (plotted here for the longitude range corresponding to the comparison) for a source luminosity of $10^{34} \text{ erg s}^{-1}$. The spiral arms of the galaxy (Vallée 2014) are schematically drawn in gray. The green and orange triangles represent the sources detected by H.E.S.S. and HAWC, respectively, for which a distance is known. The sources located further than the horizon must be brighter than $10^{34} \text{ erg s}^{-1}$. Adapted from the H.E.S.S. Collaboration et al. (2018a).

There are several possible explanations for the different number of sources detected by HAWC and H.E.S.S.:

1. The resolution of H.E.S.S. at 1 TeV is several times better than the resolution of HAWC for analysis bin 4. Since its PSF is wider, HAWC cannot resolve all the H.E.S.S. sources and several H.E.S.S. sources are seen as only one by HAWC.
2. With its 0.1° kernel and the ring background subtraction, the HGPS analysis is optimized for sources that are point-like or have a modest extension. Extended sources with low surface brightness are challenging for this kind of analysis. The variation of sensitivity as a function of the source extension is illustrated by H.E.S.S. Collaboration et al. (2018a), Figure 13. The seven HAWC sources undetected by H.E.S.S. lie above the H.E.S.S. sensitivity curve (dark blue curve in Figure 2), which means that if their image was a Gaussian of the order of 0.1° in size, H.E.S.S. would have detected them. Therefore, they likely have a larger extension.
3. The H.E.S.S. sensitivity peaks at lower energy than HAWC, as illustrated by Abeysekara et al. (2017a), Figure 15: above ~ 10 TeV, gamma/hadron separation in HAWC becomes very efficient and detection is not limited by the background anymore but by the number of arriving photons. Hence, continuous observing instruments like HAWC take advantage over short exposure instruments like H.E.S.S.

4.2. HAWC Sources Newly Detected by H.E.S.S.

Using the new analysis presented in Section 3, a new significance map is produced, shown in the third row of Figure 1. Its corresponding sensitivity curve is plotted in light blue in the left panel of Figure 2. Statistically significant emission $> 5\sigma$ is now visible next to four of the previously undetected HAWC sources: HAWC J1928+178, HAWC J1914+118, HAWC J1907+085, and HAWC J1852+013. These sources are visible in the four panels of Figure 3, with their possible counterparts. White contours indicate the significance level. For each new H.E.S.S. source, a maximum of emission is identified in the flux map produced together with the TS map, to give a preliminary estimate of the location of the H.E.S.S. source. In the case of the H.E.S.S. source corresponding to HAWC J1852+013, the location of maximum flux is identified by increasing the energy threshold to 2 and 5 TeV in order to lower the contamination from the extended emission from the neighboring sources and possibly from large scale diffuse γ -ray emission. The four new H.E.S.S. sources are located 0.3° – 0.5° away from the HAWC source.

HESS J1928+181 is the H.E.S.S. detection of the source of unknown origin HAWC J1928+178. It may be associated with the middle-aged pulsar (82 kyr) PSR J1928+1746 (Lopez-Coto et al. 2017; Jardin-Blicq 2019), located 4.3 kpc away. Detected in radio wavelength (Cordes et al. 2006), it has a spin-down power of $\dot{E} = 1.6 \times 10^{36} \text{ erg s}^{-1}$. In that case, the γ -ray




Figure 4. Longitude profiles of the integrated flux above 1 TeV at $b = 0^\circ$ for the new H.E.S.S. map, using the 0.4° Gaussian and the field-of-view background method in cyan, and the HAWC map in dashed orange. The bottom panel shows the difference between the H.E.S.S. and HAWC flux and the corresponding histogram (mean and standard deviation given on top of the histogram). The systematic uncertainties are derived for each parameter used to produce the maps (listed in Appendix B) and added quadratically, are represented by the dotted envelop. More details on the derivation of the systematic uncertainties can be found Appendix B.




Figure 5. Profiles of the integrated flux above 1 TeV for H.E.S.S. and HAWC, along the latitude axis for different ranges in longitude ($10^\circ < \ell < 60^\circ$, $10^\circ < \ell < 30^\circ$, and $30^\circ < \ell < 60^\circ$). The shaded bands represent the 1σ statistic uncertainties. The systematic uncertainties are not shown here but from the study presented in Appendix B they can reach $\sim 50\%$, depending on the longitude. Note that the sources have not been subtracted. The latitude profile extracted from the ^{12}CO survey of Dame et al. (2001) and the profile of the H I emission from the HI4PI survey (HI4PI Collaboration et al. 2016) are also shown for the corresponding range in longitude, with an arbitrary normalization for display purposes.

them and the corresponding histogram is plotted on the bottom panel. The integrated flux measured by H.E.S.S. is on average slightly higher than that of HAWC by $3.7 \times 10^{-13} \text{ cm}^{-2} \text{ s}^{-1}$. Nevertheless, the flux values are compatible within the systematic uncertainties, described in Appendix B, over most of the Galactic plane.

Figure 5 shows the latitude profiles of the integrated flux above 1 TeV for H.E.S.S. and HAWC. In this case, a sliding box of 0.4° width is defined with different lengths in longitude, as indicated in the different plots of the figure, that scans the Galactic plane in latitude between -3.5° and $+3.5^\circ$. In each of these boxes the surface brightness is derived by normalizing the

average flux by the solid angle. The ^{12}CO profile using data from Dame et al. (2001) and the H I profile using the HI4PI survey from HI4PI Collaboration et al. (2016) are also superimposed with an arbitrary normalization for display. The H.E.S.S. profile peaks at slightly negative latitude with a small asymmetry and seems to follow the ^{12}CO profile, similar to what has been reported by H.E.S.S. Collaboration et al. (2018a), Figure 11, when comparing with the distribution of sources, or with Abramowski et al. (2014), Figure 2. The HAWC profile is more symmetric and centered around $b = 0^\circ$, like the H I profile. The origin of this difference is not clear. Since the H I emission is very broad around the Galactic plane, the diffuse γ -ray emission coming from these high-latitude regions is likely removed in the H.E.S.S. analysis due to the background estimation method, but is conserved in the HAWC analysis because its very large field of view allows for a large exclusion region of $\pm 3^\circ$ around the Galactic plane. Hence, it could be the indication of an underlying diffuse emission component removed in the H.E.S.S. map. This may also explain that the H.E.S.S. profile tends to be slightly negative for $|b| > 2^\circ$. A more detailed study of the large scale diffuse emission is beyond the scope of this paper and will be the topic of a future paper. Another possible explanation for the shift in the H.E.S.S. profile with respect to $b = 0^\circ$ could be related to the HAWC bias in the source location reconstruction as a function of decl., reported by Albert et al. (2020), Figure 11. This would affect the overall shape toward positive latitude as it gets closer to the Galactic Center.

Moreover, the latitude profile for $10^\circ < \ell < 60^\circ$ (visible in the left plot of Figure 5) shows that H.E.S.S. detects more flux than HAWC, while the profiles over smaller longitude bands (visible in the right plots of Figure 5 for $10^\circ < \ell < 60^\circ$ and $30^\circ < \ell < 60^\circ$ Figure 6, in Appendix C, for 10° wide bands) show that HAWC detects more flux than H.E.S.S. for $\ell < 40^\circ$. This can be explained by the fact that the exposure time is taken into account: more weight is given to regions with more exposure time, and regions with $\ell < 40^\circ$ have a low exposure time for HAWC and a high exposure time for H.E.S.S.

5. Conclusion

HAWC and H.E.S.S. are two complementary instruments using different analysis techniques. This study is the first attempt to reconcile their data and this analysis shows consistent results: constructing the maps by using comparable procedures, a common energy range and comparable angular resolution leads to two remarkably similar maps. Overall, HAWC and H.E.S.S. measure similar fluxes. Nonetheless, there are some remaining differences that are only partially understood. The difference in sensitivity as a function of energy may play a role since HAWC is more sensitive than H.E.S.S. at the highest energies, it will be more sensitive to sources with a hard spectrum. In the part of the Galactic plane common to H.E.S.S. and HAWC, four HAWC sources previously undetected by H.E.S.S. show significant emission above the detection level of 5σ , using the analysis presented here. The main ingredients leading to this result are the use of a large Gaussian smearing factor, followed by the field-of-view background estimation method together with a large exclusion band to avoid contamination from the Galactic plane. It is also an excellent demonstration of the capability for an IACT to be sensitive to extended sources with low-surface-brightness emission. Each of these sources will require a dedicated analysis in order to study them in more detail and

attempt to identify the possible origin of the observed γ -ray emission. Future work would benefit from combining the data from both instruments in a joint analysis, which will be supported by ongoing efforts toward multi-instrument open source tools like Gammapy (Deil et al. 2017) or the Multi-Mission Maximum Likelihood (3ML) framework (Vianello et al. 2015), together with a more realistic model of the exposure map as presented in Mohrmann et al. (2019) for the background estimation. The number of new extended γ -ray emission regions is expected to increase further by applying this new analysis to the southern part of the HGPS, which is out of HAWC's reach. The future observatories SWGO (Southern Wide-field Gamma-ray Observatory (Schoorlemmer 2019)) and CTA (Cherenkov Telescope Array; Cherenkov Telescope Array Consortium et al. 2019) can take advantage of the complementarity of the two detection techniques illustrated here. SWGO will observe the southern hemisphere with an improved sensitivity and angular resolution in comparison with HAWC. Combination of SWGO data with CTA data and their better control over systematic errors will allow broadband spectrum analysis and provide a deeper understanding of the sources.

We acknowledge the support from the US National Science Foundation (NSF); the US Department of Energy Office of High-Energy Physics; the Laboratory Directed Research and Development (LDRD) program of Los Alamos National Laboratory; Consejo Nacional de Ciencia y Tecnología (CONACyT), México, grants 271051, 232656, 260378, 179588, 254964, 258865, 243290, 132197, A1-S-46288, and A1-S-22784, cátedras 873, 1563, 341, and 323, Red HAWC, México; DGAPA-UNAM grants IG101320, IN111315, IN111716-3, IN111419, IA102019, and IN110621; VIEP-BUAP; PIFI 2012, 2013, PROFOCIE 2014, 2015; the University of Wisconsin Alumni Research Foundation; the Institute of Geophysics, Planetary Physics, and Signatures at Los Alamos National Laboratory; Polish Science Centre grant, DEC-2017/27/B/ST9/02272; Coordinación de la Investigación Científica de la Universidad Michoacana; Royal Society—Newton Advanced Fellowship 180385; Generalitat Valenciana, grant CIDEGENT/2018/034; Chulalongkorn University's CUNiverse (CUAASC) grant; Coordinación General Académica e Innovación (CGAI-UdeG) and PRODEP-SEP through UDG-CA-499; Institute of Cosmic Ray Research (ICRR), University of Tokyo. Thanks to Scott Delay, Luciano Díaz and Eduardo Murrieta for technical support.

The support of the Namibian authorities and of the University of Namibia in facilitating the construction and operation of H.E.S.S. is gratefully acknowledged, as is the support by the German Ministry for Education and Research (BMBF), the Max Planck Society, the German Research Foundation (DFG), the Helmholtz Association, the Alexander von Humboldt Foundation, the French Ministry of Higher Education, Research and Innovation, the Centre National de la Recherche Scientifique (CNRS/IN2P3 and CNRS/INSU), the Commissariat à l'énergie atomique et aux énergies alternatives (CEA), the U.K. Science and Technology Facilities Council (STFC), the Knut and Alice Wallenberg Foundation, the National Science Centre, Poland grant No. 2016/22/M/ST9/00382, the South African Department of Science and Technology and National Research Foundation, the University of Namibia, the National Commission on Research, Science & Technology of Namibia (NCRST), the Austrian Federal Ministry of Education, Science and Research

and the Austrian Science Fund (FWF), the Australian Research Council (ARC), the Japan Society for the Promotion of Science, and the University of Amsterdam. We appreciate the excellent work of the technical support staff in Berlin, Zeuthen, Heidelberg, Palaiseau, Paris, Saclay, Tübingen, and Namibia in the construction and operation of the equipment. This work benefited from services provided by the H.E.S.S. Virtual Organisation, supported by the national resource providers of the EGI Federation.

Facility: HAWC (<https://www.hawc-observatory.org/>) H.E.S.S. (<https://www.mpi-hd.mpg.de/hfm/HESS/>).

Software: Astropy (Astropy Collaboration et al. 2013; Price-Whelan et al. 2018), Gammapy-0.13 and Gammapy-0.17 (Deil et al. 2017; Nigro et al. 2019).

Appendix A HAWC Sources

The HAWC sources in this analysis are the result of the HAWC source search performed on the latest HAWC data using

1523 days and events falling in the analysis bins 4–9 only. According to the definition given in Abeysekara et al. (2017a), they correspond to events triggering more than $\sim 25\%$ of the array that would have an energy greater than approximately 1 TeV for a source at the Crab decl. The procedure is the same as the one used for the 3HWC catalog (Albert et al. 2020). Table 1 shows the resulting list of sources with their position and the association with the nearby TeVCat⁷⁶ (Wakely & Horan 2008) sources within 1° of the HAWC source. Table 2 gives the integrated flux above 1 TeV from the HAWC flux map and the H.E.S.S. flux map using the new analysis at the location of the HAWC source reported in Table 1. The sources highlighted in bold are found both in the HAWC search and in the HGPS catalog. The sources highlighted in italic are the HAWC sources discussed in this analysis having no H.E.S.S. counterpart in the HGPS. The sources with a dagger ($\dagger$) are called *secondary sources*. They correspond to local maxima that have reached the detection threshold but not the separation criteria based on the TS gap between two neighbor sources, as defined in the 3HWC catalog (Albert et al. 2020).

Table 1
HAWC Source List with Their Position and TeVCat Association

Name	TS	R.A. (deg)	Decl. (deg)	ℓ (deg)	b (deg)	1σ Stat. Unc. (deg)	TeVCat Source
HAWC J1809-190	222.8	272.46	−19.04	11.33	0.18	0.11	3HWC J1809-190, HESS J1809-193
HAWC J1813-125	42.5	273.30	−12.56	17.40	2.59	0.15	3HWC J1813-125, HESS J1813-126
HAWC J1813-174	354.1	273.43	−17.47	13.15	0.13	0.06	3HWC J1813-174, HESS J1813-178
HAWC J1819-151	69.4	274.79	−15.17	15.79	0.07	0.14	3HWC J1819-150 $\dagger$, HESS J1818-154
HAWC J1825-134	1856.6	276.46	−13.40	18.12	−0.53	0.06	3HWC J1825-134, HESS J1825-137
HAWC J1831-096	176.4	277.91	−9.63	22.11	−0.03	0.11	3HWC J1831-095, HESS J1831-098
HAWC J1837-066	1111.9	279.40	−6.62	25.47	0.04	0.06	3HWC J1837-066, HESS J1837-069
HAWC J1843-034	705.0	280.99	−3.47	28.99	0.08	0.06	3HWC J1843-034, HESS J1843-033
HAWC J1847-017	226.8	281.91	−1.79	30.90	0.03	0.09	3HWC J1847-017, HESS J1848-018
HAWC J1849+001	372.4	282.31	0.11	32.78	0.54	0.06	3HWC J1849+001, HESS J1849-000 , IGR J18490-0000
<i>HAWC J1852+013</i>	93.7	283.01	1.38	34.23	0.50	0.10	3HWC J1852+013 $\dagger$
HAWC J1857+027	507.4	284.33	2.80	36.09	−0.03	0.06	3HWC J1857+027, HESS J1857+026
<i>HAWC J1859+057$\dagger$</i>	33.0	284.85	5.72	38.93	0.84	0.13	3HWC J1857+051 $\dagger$
<i>HAWC J1907+085</i>	48.2	286.79	8.57	42.35	0.44	0.10	3HWC J1907+085
HAWC J1908+063	1068.5	287.05	6.39	40.53	−0.80	0.06	3HWC J1908+063, HESS J1908+063 , MGRO J1908+06
HAWC J1912+103	130.8	288.02	10.39	44.52	0.20	0.07	3HWC J1912+103, HESS J1912+101
<i>HAWC J1914+118</i>	64.2	288.68	11.87	46.13	0.32	0.09	3HWC J1914+118
<i>HAWC J1920+147</i>	34.4	290.17	14.79	49.39	0.39	0.10	3HWC J1920+147 $\dagger$, W 51
HAWC J1922+140	138.1	290.70	14.09	49.01	−0.38	0.06	3HWC J1922+140, HESS J1923+140 , W 51
<i>HAWC J1923+169$\dagger$</i>	28.2	290.79	16.96	51.58	0.89	0.11	3HWC J1923+169 $\dagger$
<i>HAWC J1928+178</i>	172.5	292.10	17.82	52.93	0.20	0.09	3HWC J1928+178
HAWC J1930+188	86.1	292.54	18.84	54.03	0.32	0.09	3HWC J1930+188, HESS J1930+188 , SNR G054.1+00.3

Note. The position uncertainty reported here is statistical only. The systematic uncertainty varies with decl. and is discussed in more detail in Appendix B. For each source, the nearby known TeV sources within 1° of the HAWC source, listed in TeVCat (Wakely & Horan 2008), are also given. See the text for more details.

⁷⁶ <http://tevcat.uchicago.edu/>

Table 2
HAWC Source List with the HAWC and H.E.S.S. Integrated Flux above 1 TeV

Name	$F_{\text{HAWC}} > 1 \text{ TeV}$ ($10^{-12} \text{ cm}^{-2} \text{ s}^{-1}$)	$F_{\text{HESS}} > 1 \text{ TeV}$ ($10^{-12} \text{ cm}^{-2} \text{ s}^{-1}$)	HGPS Counterpart (ℓ, b) Integrated Flux above 1 TeV	$F_{\text{HESS}} > 1 \text{ TeV}$ at HGPS Location ($10^{-12} \text{ cm}^{-2} \text{ s}^{-1}$)
HAWC J1809-190	5.98 ± 0.47	4.15 ± 0.17	HESS J1809-193 (11.11, -0.02)	5.37 ± 0.45
HAWC J1813-125	1.29 ± 0.22	1.11 ± 0.25	HESS J1813-126 (17.31, 2.49)	1.04 ± 0.21
HAWC J1813-174	6.49 ± 0.41	5.74 ± 0.33	HESS J1813-178 (12.82, -0.025)	2.12 ± 0.40
HAWC J1819-151	2.13 ± 0.28	1.16 ± 0.17	HESS J1818-154 (15.41, 0.16)	0.23 ± 0.05
HAWC J1825-134	11.4 ± 0.34	9.78 ± 0.20	HESS J1825-137 (17.52, -0.62)	19.15 ± 1.85
HAWC J1831-096	2.29 ± 0.20	2.28 ± 0.18	...	...
HAWC J1837-066	5.65 ± 0.21	7.32 ± 0.25	HESS J1837-069 (25.15, -0.087)	11.55 ± 0.49
HAWC J1843-034	3.64 ± 0.16	4.27 ± 0.24	HESS J1843-033 (28.90, 0.075)	3.04 ± 0.20
HAWC J1847-017	1.82 ± 0.14	2.66 ± 0.19	HESS J1848-018 (30.92, -0.20)	1.11 ± 0.15
HAWC J1849+001	2.11 ± 0.13	2.03 ± 0.17	HESS J1849-000 (32.61, 0.53)	0.57 ± 0.07
HAWC J1852+013	0.97 ± 0.11	1.22 ± 0.16	...	...
HAWC J1857+027	2.54 ± 0.13	4.75 ± 0.26	HESS J1857+026 (36.06, -0.06)	4.0 ± 0.29
HAWC J1859+057 [†]	0.48 ± 0.091	0.39 ± 0.31	...	...
HAWC J1907+085	0.56 ± 0.089	1.03 ± 0.16	...	...
HAWC J1908+063	3.52 ± 0.13	3.71 ± 0.22	HESS J1908+063 (40.55, -0.84)	8.35 ± 0.57
HAWC J1912+103	0.98 ± 0.096	2.01 ± 0.13	HESS J1912+101 (44.46-0.13)	2.49 ± 0.34
HAWC J1914+118	0.61 ± 0.085	1.09 ± 0.20	...	...
HAWC J1920+147	0.43 ± 0.080	0.54 ± 0.22	...	...
HAWC J1922+140	0.92 ± 0.088	1.38 ± 0.22	HESS J1923+141 (49.08, -0.40)	0.69 ± 0.11
HAWC J1923+169 [†]	0.39 ± 0.079	0.51 ± 0.31	...	...
HAWC J1928+178	1.01 ± 0.09	1.49 ± 0.24	...	...
HAWC J1930+188	0.74 ± 0.087	1.77 ± 0.23	HESS J1930+188 (54.06, 0.27)	0.32 ± 0.068

Note. The flux values are reported at the location of the HAWC sources from Table 1. The H.E.S.S. flux from this analysis is also reported at the location of the HGPS source when there is an association. The positions of the HGPS sources are given in galactic coordinates in parentheses. The flux uncertainty reported here is statistical only. The systematic uncertainties on the flux are estimated to be 30% for H.E.S.S. (H.E.S.S. Collaboration et al. 2018a) and for HAWC (Abeysekara et al. 2019). The HAWC flux reported here is slightly different from the output from the HAWC source search from Albert et al. (2020) on the data set considered here because it assumes a simple power-law spectrum with a constant spectral index of -2.5 for all sources, while in the catalog search it is a free parameter from the fit. Moreover, the flux reported for H.E.S.S. is expected to differ from the one reported in the HGPS due to the hypotheses of 0.4° on the Gaussian size and the use of an spectral index of -2.5 instead of -2.3 , and the different background estimation method.

Appendix B Systematic Uncertainties

The difference between HAWC and H.E.S.S. flux profiles may be explained, at least partially, by evaluating the systematic uncertainties. First, each instrument has its own absolute systematic uncertainties on the flux: for H.E.S.S., they are mainly due to atmospheric effects (weather condition and seasonal effect), but also to the modeling of the instrument, and have been evaluated to $\sim 30\%$ (H.E.S.S. Collaboration et al. 2018a). HAWC is dominated by the uncertainties on the modeling of the instrument, which results in $\sim 30\%$ uncertainty on the flux (Abeysekara et al. 2019). Second, the variation in flux resulting from a variation of each of the parameters listed below, used to produce the maps, is evaluated. For most of them, the median flux variation over the part of the Galactic plane used for the comparison ($10^\circ < \ell < 60^\circ$) and local effects may have to be taken into account for specific studies. When a parameter highly depends on the longitude, the flux variation is derived for all positions along the Galactic plane. All systematic uncertainties are then added quadratically and represented as the dotted envelope in Figure 4.

1. The spectral index: An index of -2.5 is used both for HAWC and H.E.S.S. for the entire part of the Galactic plane considered for the comparison. However, H.E.S.S. used -2.3 when producing the HGPS, as being the mean index of TeV Galactic sources. Another H.E.S.S. map was produced (not presented here) using the new analysis

presented in this paper and an index of -2.3 . Using an index of -2.3 instead of -2.5 leads to a median flux variation of $\sim 1.7\%$ over the part of the Galactic plane considered here.

2. The analysis energy threshold: While it can be set quite precisely to be 1 TeV for H.E.S.S. data, with an uncertainty of $\sim 10\%$ (Parsons & Hinton 2014), the uncertainty on the HAWC energy threshold is much larger due to the use of the analysis bins. This can cause HAWC to be dominated by events of a few TeV instead of 1 TeV, depending on the zenith angle. A H.E.S.S. map was produced using the new analysis presented in this paper and an energy threshold of 2 TeV instead of 1 TeV. Using 2 TeV would lead to a median variation of $\sim 14\%$.
3. The background method: A H.E.S.S. map was produced (not presented here) using the new analysis presented in this paper but using the ring background method. Using the field-of-view background method instead of the ring background method leads to a systematic increase of the flux by a median value of $1.9 \times 10^{-13} \text{ cm}^{-2} \text{ s}^{-1}$.
4. The imperfect knowledge of the H.E.S.S. background, due to the local nature of the background estimation and the sensitivity to instrumental conditions: Assuming that the background is known with a 5% precision, the corresponding error on the flux at each point in longitude is evaluated to range between $1.5 \times 10^{-13} \text{ cm}^{-2} \text{ s}^{-1}$ at $\ell = 10^\circ$ and $5 \times 10^{-13} \text{ cm}^{-2} \text{ s}^{-1}$ at $\ell = 60^\circ$.

5. H.E.S.S. analysis chain: A H.E.S.S. map was produced (not presented here) using the new analysis presented in this paper with the other H.E.S.S. analysis chain to cross check the results. The median variation in flux seen by the two different H.E.S.S. analysis chains is $\sim 25\%$.
6. Using the HAWC flux map: In the study presented in this paper, the flux are obtained by reading the flux map, making the assumption that the spectral index is -2.5 for all sources. The flux values can be compared with the one obtained from the HAWC catalog search, where both the index and the flux normalization are fitted. The median difference in flux between both is $\sim 48\%$ of the flux.
7. The source size hypotheses: A constant size of 0.4° was used in this analysis to produce the H.E.S.S. TS map, whereas the PSF of HAWC varies with zenith angle and as a function of the analysis bin. The average HAWC PSF over the analysis bins 4–9 is actually closer to 0.3° . Another H.E.S.S. map using the new analysis presented in this paper was produced (not shown here) using a Gaussian of 0.3° instead of 0.4° . Decreasing the size of the Gaussian to 0.3° leads to a flux 39% lower.
8. The systematic bias on the source position reconstruction (visible in Figure 11 of Albert et al. (2020)) has also been taken into account in the systematic uncertainties: Similarly, a relation has been derived to describe the difference in the reconstructed location of the sources in

common to the HAWC data set used this analysis and the HGPS (Table 2). Multiple profiles have then been derived and averaged according to this relation and its errors, given a mean systematic uncertainty that depends on the longitude. The systematic error is then estimated to be $9.3 \times 10^{-14} \text{ cm}^{-2} \text{ s}^{-1}$ (median), which is negligible compare to the other systematic uncertainties.

An in-depth study with joint analysis of common sources would be needed to quantify these effects more precisely; this is beyond the scope of the first approach presented here.

Appendix C Latitude Profiles in 10° Longitude Bands

The latitude profiles of the integrated flux above 1 TeV for H.E.S.S. and HAWC, as described in Section 4.3, are plotted in Figure 6 in bands of 10° in longitude as indicated on the plots. HAWC detects more flux than H.E.S.S. in all bands with $\ell < 40^\circ$. In the band with $40^\circ < \ell < 50^\circ$ HAWC and H.E.S.S. detect a similar level of flux, and for $50^\circ < \ell < 60^\circ$, H.E.S.S. detects slightly more flux. However, when taking into account the exposure time of each instrument over each latitude range, in the average over $10^\circ < \ell < 60^\circ$, H.E.S.S. detects more flux as shown in the left plot of Figure 5. Note that the systematic uncertainties are not shown here but can reach 50% of the flux.




Figure 6. Profiles of the integrated flux above 1 TeV for H.E.S.S. and HAWC along the latitude axis in bands of 10° in longitude. The shaded bands represent the 1σ uncertainties. The systematic uncertainties are not shown here.

Appendix D

Applicability of Wilks's Theorem on HAWC and H.E.S.S. Data

To validate the use of Wilks's theorem and check the significance distribution, synthetic maps have been created for HAWC and H.E.S.S. by generating random Poisson realizations assuming background only. The significance maps are

then produced in the same way as the maps presented in the paper, as described in Section 3. Figure 7 shows an example for one synthetic map for H.E.S.S. with the new analysis and one for HAWC using the standard analysis. The distribution of significance in the resulting null hypothesis maps is found to be consistent with a Gaussian centered on 0 with a standard deviation of 1, as expected.




Figure 7. H.E.S.S. (top) and HAWC (bottom) synthetic significance map with their distribution.

ORCID iDs

F. Aharonian <https://orcid.org/0000-0003-1157-3915>
T. Armstrong <https://orcid.org/0000-0001-5067-2620>
H. Ashkar <https://orcid.org/0000-0002-2153-1818>
M. Backes <https://orcid.org/0000-0002-9326-6400>
V. Baghmanyar <https://orcid.org/0000-0003-0477-1614>
V. Barbosa Martins <https://orcid.org/0000-0002-5085-8828>
A. Barnacka <https://orcid.org/0000-0001-5655-4158>
M. Barnard <https://orcid.org/0000-0003-1720-7959>
D. Berge <https://orcid.org/0000-0002-2918-1824>
K. Bernlöhr <https://orcid.org/0000-0001-8065-3252>
M. Böttcher <https://orcid.org/0000-0002-8434-5692>
C. Boisson <https://orcid.org/0000-0001-5893-1797>
M. Breuhaus <https://orcid.org/0000-0003-0268-5122>
F. Brun <https://orcid.org/0000-0003-0770-9007>
P. Brun <https://orcid.org/0000-0002-0207-958X>
T. Bulik <https://orcid.org/0000-0003-2045-4803>
T. Bylund <https://orcid.org/0000-0003-2946-1313>
S. Caroff <https://orcid.org/0000-0002-1103-130X>
T. Chand <https://orcid.org/0000-0002-1833-3749>
S. Chandra <https://orcid.org/0000-0002-8776-1835>
A. Chen <https://orcid.org/0000-0001-6425-5692>
G. Cotter <https://orcid.org/0000-0002-9975-1829>
J. Damascene Mbarubucyeye <https://orcid.org/0000-0002-4991-6576>
I. D. Davids <https://orcid.org/0000-0002-6476-964X>
J. Davies <https://orcid.org/0000-0002-2394-4720>
J. Devin <https://orcid.org/0000-0003-1018-7246>
A. Donath <https://orcid.org/0000-0003-4568-7005>
L. Dreyer <https://orcid.org/0000-0002-4971-3672>
S. Einecke <https://orcid.org/0000-0001-9687-8237>
G. Fichet de Clairfontaine <https://orcid.org/0000-0003-1143-3883>
G. Fontaine <https://orcid.org/0000-0002-6443-5025>
S. Funk <https://orcid.org/0000-0002-2012-0080>
G. Giavitto <https://orcid.org/0000-0002-7629-6499>
D. Glawion <https://orcid.org/0000-0003-4865-7696>
J. F. Glicenstein <https://orcid.org/0000-0003-2581-1742>

M.-H. Grondin <https://orcid.org/0000-0002-8383-251X>
J. A. Hinton <https://orcid.org/0000-0002-1031-7760>
C. Hoischen <https://orcid.org/0000-0001-6661-2278>
T. L. Holch <https://orcid.org/0000-0001-5161-1168>
D. Horns <https://orcid.org/0000-0003-1945-0119>
M. Jamroz <https://orcid.org/0000-0002-0870-7778>
U. Katz <https://orcid.org/0000-0002-7063-4418>
D. Khangulyan <https://orcid.org/0000-0002-7576-7869>
B. Khélifi <https://orcid.org/0000-0001-6876-5577>
S. Klepser <https://orcid.org/0000-0002-8949-4275>
Nu. Komin <https://orcid.org/0000-0003-3280-0582>
R. Konno <https://orcid.org/0000-0003-1892-2356>
K. Kosack <https://orcid.org/0000-0001-8424-3621>
D. Kostunin <https://orcid.org/0000-0002-0487-0076>
M. Kreter <https://orcid.org/0000-0002-3092-3506>
M. Lemoine-Goumard <https://orcid.org/0000-0002-4462-3686>
J.-P. Lenain <https://orcid.org/0000-0001-7284-9220>
F. Leuschner <https://orcid.org/0000-0001-9037-0272>
T. Lohse <https://orcid.org/0000-0002-9751-7633>
J. Mackey <https://orcid.org/0000-0002-5449-6131>
D. Malyshev <https://orcid.org/0000-0001-9689-2194>
D. Malyshev <https://orcid.org/0000-0002-9102-4854>
V. Marandon <https://orcid.org/0000-0001-9077-4058>
P. Marchegiani <https://orcid.org/0000-0001-7487-8287>
A. Marcowith <https://orcid.org/0000-0002-3971-0910>
A. Mares <https://orcid.org/0000-0003-3345-7800>
G. Martí-Devesa <https://orcid.org/0000-0003-0766-6473>
R. Marx <https://orcid.org/0000-0002-6557-4924>
M. Meyer <https://orcid.org/0000-0002-0738-7581>
A. M. W. Mitchell <https://orcid.org/0000-0003-3631-5648>
L. Mohrmann <https://orcid.org/0000-0002-9667-8654>
A. Montanari <https://orcid.org/0000-0002-3620-0173>
P. Morris <https://orcid.org/0000-0002-8533-8232>
E. Moulin <https://orcid.org/0000-0003-4007-0145>
J. Muller <https://orcid.org/0000-0003-0004-4110>
T. Murach <https://orcid.org/0000-0003-1128-5008>
M. de Naurois <https://orcid.org/0000-0002-7245-201X>

- H. Ndiyavala  <https://orcid.org/0000-0001-9279-1775>
 J. Niemiec  <https://orcid.org/0000-0001-6036-8569>
 S. Ohm  <https://orcid.org/0000-0002-3474-2243>
 L. Olivera-Nieto  <https://orcid.org/0000-0002-9105-0518>
 M. Ostrowski  <https://orcid.org/0000-0002-9199-7031>
 S. Panny  <https://orcid.org/0000-0001-5770-3805>
 R. D. Parsons  <https://orcid.org/0000-0003-3457-9308>
 G. Peron  <https://orcid.org/0000-0003-3255-0077>
 V. Poireau  <https://orcid.org/0000-0002-4768-0256>
 M. Punch  <https://orcid.org/0000-0002-4710-2165>
 P. Reichherzer  <https://orcid.org/0000-0003-4513-8241>
 A. Reimer  <https://orcid.org/0000-0001-8604-7077>
 O. Reimer  <https://orcid.org/0000-0001-6953-1385>
 F. Rieger  <https://orcid.org/0000-0003-1334-2993>
 L. Rinchiuso  <https://orcid.org/0000-0003-0540-9967>
 C. Romoli  <https://orcid.org/0000-0003-2541-4499>
 G. Rowell  <https://orcid.org/0000-0002-9516-1581>
 B. Rudak  <https://orcid.org/0000-0003-0452-3805>
 S. Sailer  <https://orcid.org/0000-0001-8273-8495>
 A. Santangelo  <https://orcid.org/0000-0003-4187-9560>
 M. Sasaki  <https://orcid.org/0000-0001-5302-1866>
 F. Schüssler  <https://orcid.org/0000-0003-1500-6571>
 H. M. Schutte  <https://orcid.org/0000-0002-1769-5617>
 U. Schwanke  <https://orcid.org/0000-0002-1229-278X>
 M. Seglar-Arroyo  <https://orcid.org/0000-0001-8654-409X>
 M. Senniappan  <https://orcid.org/0000-0001-6734-7699>
 J. N. S. Shapopi  <https://orcid.org/0000-0002-7130-9270>
 A. Sinha  <https://orcid.org/0000-0002-9238-7163>
 S. Spencer  <https://orcid.org/0000-0001-5516-1205>
 S. Steinmassl  <https://orcid.org/0000-0002-2865-8563>
 C. Steppa  <https://orcid.org/0000-0003-0116-8836>
 A. M. Taylor  <https://orcid.org/0000-0001-9473-4758>
 R. Terrier  <https://orcid.org/0000-0002-8219-4667>
 C. van Eldik  <https://orcid.org/0000-0001-9669-645X>
 B. van Soelen  <https://orcid.org/0000-0003-1873-7855>
 C. Venter  <https://orcid.org/0000-0002-2666-4812>
 J. Vink  <https://orcid.org/0000-0002-4708-4219>
 H. J. Völk  <https://orcid.org/0000-0003-2386-8067>
 Z. Wadiasingh  <https://orcid.org/0000-0002-9249-0515>
 S. J. Wagner  <https://orcid.org/0000-0002-7474-6062>
 J. Watson  <https://orcid.org/0000-0003-4282-7463>
 F. Werner  <https://orcid.org/0000-0002-6941-1073>
 A. Wiercholska  <https://orcid.org/0000-0003-4472-7204>
 M. Zacharias  <https://orcid.org/0000-0001-5801-3945>
 R. Zanin  <https://orcid.org/0000-0001-6320-1801>
 D. Zargaryan  <https://orcid.org/0000-0002-2876-6433>
 A. A. Zdziarski  <https://orcid.org/0000-0002-0333-2452>
 A. Zech  <https://orcid.org/0000-0002-4388-5625>
 S. J. Zhu  <https://orcid.org/0000-0002-6468-8292>
 J. Zorn  <https://orcid.org/0000-0001-9309-0700>
 S. Zouari  <https://orcid.org/0000-0002-5333-2004>
 N. Żywucka  <https://orcid.org/0000-0003-2644-6441>
 A. Albert  <https://orcid.org/0000-0003-0197-5646>
 R. Alfaro  <https://orcid.org/0000-0001-8749-1647>
 K. P. Arunbabu  <https://orcid.org/0000-0002-3032-663X>
 D. Avila Rojas  <https://orcid.org/0000-0002-4020-4142>
 E. Belmont-Moreno  <https://orcid.org/0000-0003-3207-105X>
 S. Y. BenZvi  <https://orcid.org/0000-0001-5537-4710>
 C. Brisbois  <https://orcid.org/0000-0002-5493-6344>
 K. S. Caballero-Mora  <https://orcid.org/0000-0002-4042-3855>
 T. Capistrán  <https://orcid.org/0000-0003-2158-2292>
 A. Carramiñana  <https://orcid.org/0000-0002-8553-3302>
 S. Casanova  <https://orcid.org/0000-0002-6144-9122>
 U. Cotti  <https://orcid.org/0000-0002-7607-9582>
 J. Cotzomi  <https://orcid.org/0000-0002-1132-871X>
 S. Coutiño de León  <https://orcid.org/0000-0002-7747-754X>
 E. De la Fuente  <https://orcid.org/0000-0001-9643-4134>
 C. de León  <https://orcid.org/0000-0002-8528-9573>
 R. Diaz Hernandez  <https://orcid.org/0000-0001-8487-0836>
 J. C. Díaz-Vélez  <https://orcid.org/0000-0002-0087-0693>
 B. L. Dingus  <https://orcid.org/0000-0001-8451-7450>
 M. A. DuVernois  <https://orcid.org/0000-0002-2987-9691>
 M. Durocher  <https://orcid.org/0000-0003-2169-0306>
 R. W. Ellsworth  <https://orcid.org/0000-0003-2338-0344>
 K. Engel  <https://orcid.org/0000-0001-5737-1820>
 C. Espinoza  <https://orcid.org/0000-0001-7074-1726>
 N. Fraija  <https://orcid.org/0000-0002-0173-6453>
 A. Galván-Gómez  <https://orcid.org/0000-0001-5193-3693>
 J. A. García-González  <https://orcid.org/0000-0002-4188-5584>
 F. Garfías  <https://orcid.org/0000-0003-1122-4168>
 G. Giacinti  <https://orcid.org/0000-0001-9745-5738>
 M. M. González  <https://orcid.org/0000-0002-5209-5641>
 J. A. Goodman  <https://orcid.org/0000-0002-9790-1299>
 J. P. Harding  <https://orcid.org/0000-0001-9844-2648>
 S. Hernandez  <https://orcid.org/0000-0002-2565-8365>
 B. Hona  <https://orcid.org/0000-0002-7609-343X>
 D. Huang  <https://orcid.org/0000-0002-5447-1786>
 F. Hueyotl-Zahuanitla  <https://orcid.org/0000-0002-5527-7141>
 A. Iriarte  <https://orcid.org/0000-0001-5811-5167>
 A. Jardin-Blicq  <https://orcid.org/0000-0002-6738-9351>
 V. Joshi  <https://orcid.org/0000-0003-4467-3621>
 D. Kieda  <https://orcid.org/0000-0003-4785-0101>
 W. H. Lee  <https://orcid.org/0000-0002-2467-5673>
 H. León Vargas  <https://orcid.org/0000-0001-5516-4975>
 J. T. Linnemann  <https://orcid.org/0000-0003-2696-947X>
 A. L. Longinotti  <https://orcid.org/0000-0001-8825-3624>
 G. Luis-Raya  <https://orcid.org/0000-0003-2810-4867>
 R. López-Coto  <https://orcid.org/0000-0002-3882-9477>
 K. Malone  <https://orcid.org/0000-0001-8088-400X>
 O. Martinez  <https://orcid.org/0000-0001-9052-856X>
 I. Martinez-Castellanos  <https://orcid.org/0000-0001-9035-1290>
 J. Martínez-Castro  <https://orcid.org/0000-0002-2824-3544>
 J. A. Matthews  <https://orcid.org/0000-0002-2610-863X>
 P. Miranda-Romagnoli  <https://orcid.org/0000-0002-8390-9011>
 J. A. Morales-Soto  <https://orcid.org/0000-0001-9361-0147>
 E. Moreno  <https://orcid.org/0000-0002-1114-2640>
 M. Mostafá  <https://orcid.org/0000-0002-7675-4656>
 A. Nayerhoda  <https://orcid.org/0000-0003-0587-4324>
 L. Nellen  <https://orcid.org/0000-0003-1059-8731>
 M. Newbold  <https://orcid.org/0000-0001-9428-7572>
 M. U. Nisa  <https://orcid.org/0000-0002-6859-3944>
 R. Noriega-Papaqui  <https://orcid.org/0000-0001-7099-108X>
 N. Omodei  <https://orcid.org/0000-0002-5448-7577>
 Y. Pérez Araujo  <https://orcid.org/0000-0002-8774-8147>
 E. G. Pérez-Pérez  <https://orcid.org/0000-0001-5998-4938>
 C. D. Rho  <https://orcid.org/0000-0002-6524-9769>
 D. Rosa-González  <https://orcid.org/0000-0003-1327-0838>

E. Ruiz-Velasco  <https://orcid.org/0000-0001-6939-7825>
 F. Salesa Greus  <https://orcid.org/0000-0002-8610-8703>
 A. Sandoval  <https://orcid.org/0000-0001-6079-2722>
 M. Schneider  <https://orcid.org/0000-0001-8644-4734>
 H. Schoorlemmer  <https://orcid.org/0000-0002-8999-9249>
 A. J. Smith  <https://orcid.org/0000-0002-1012-0431>
 R. W. Springer  <https://orcid.org/0000-0002-1492-0380>
 P. Surajbali  <https://orcid.org/0000-0002-8516-6469>
 K. Tollefson  <https://orcid.org/0000-0001-9725-1479>
 I. Torres  <https://orcid.org/0000-0002-1689-3945>
 R. Turner  <https://orcid.org/0000-0003-1068-6707>
 F. Ureña-Mena  <https://orcid.org/0000-0002-2748-2527>
 L. Villaseñor  <https://orcid.org/0000-0001-6876-2800>
 E. Willox  <https://orcid.org/0000-0002-6623-0277>
 H. Zhou  <https://orcid.org/0000-0003-0513-3841>

References

- Abdo, A. A., Allen, B., Berley, D., et al. 2007, *ApJL*, **664**, L91
 Abeysekara, A. U., Albert, A., Alfaro, R., et al. 2017a, *ApJ*, **843**, 39
 Abeysekara, A. U., Albert, A., Alfaro, R., et al. 2017b, *ApJ*, **843**, 40
 Abeysekara, A. U., Albert, A., Alfaro, R., et al. 2019, *ApJ*, **881**, 134
 Abeysekara, A. U., Archer, A., Benbow, W., et al. 2018, *ApJ*, **866**, 24
 Abramowski, A., Aharonian, F., Ait Benkhali, F., et al. 2014, *PhRvD*, **90**, 122007
 Aharonian, F., Akhperjanian, A. G., Anton, G., et al. 2009, *A&A*, **499**, 723
 Aharonian, F., Akhperjanian, A. G., Bazer-Bachi, A. R., et al. 2006, *A&A*, **457**, 899
 Ahnen, M. L., Ansoldi, S., Antonelli, L. A., et al. 2019, *MNRAS*, **485**, 356
 Albert, A., Alfaro, R., Alvarez, C., et al. 2020, *ApJ*, **905**, 76
 Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, **558**, A33
 Berge, D., Funk, S., & Hinton, J. 2007, *A&A*, **466**, 1219
 Cawley, M. F., Fegan, D. J., Gibbs, K., et al. 1985, Proc. ICRC (San Diego), **19**, 453
 Cherenkov Telescope Array Consortium, Acharya, B. S., Agudo, I., et al. 2019, Science with the Cherenkov Telescope Array (Singapore: World Scientific)
 Cordes, J. M., Freire, P. C. C., Lorimer, D. R., et al. 2006, *AJ*, **637**, 446
 Dame, T. M., Hartmann, D., & Thaddeus, P. 2001, *ApJ*, **547**, 792
 de Naurois, M., & Rolland, L. 2009, *APh*, **32**, 231
 Deil, C., Zanin, R., Lefaucheur, J., et al. 2017, Proc. ICRC (Busan), **35**, 766
 Fuerst, E., Reich, W., Reich, P., Handa, T., & Sofue, Y. 1987, *A&AS*, **69**, 403
 H.E.S.S. Collaboration, Abdalla, H., Abramowski, A., et al. 2018a, *A&A*, **612**, A1
 H.E.S.S. Collaboration, Abdalla, H., Abramowski, A., et al. 2018b, *A&A*, **612**, A3
 H14PI Collaboration, Ben Bekhti, N., Flöer, L., et al. 2016, *A&A*, **594**, A116
 Hnatyk, B., Hnatyk, R., Zhdanov, V., & Voitsekhovskiy, V. 2020, arXiv:2009.06081
 Jardin-Blicq, A. 2019, PhD thesis, Ruprecht-Karls-Universität, <http://hdl.handle.net/21.11116/0000-0004-F359-6>
 Kargaltsev, O., Durant, M., Pavlov, G. G., & Garmire, G. 2012, *ApJS*, **201**, 37
 Lazarus, P., Brazier, A., Hessels, J. W. T., et al. 2015, *ApJ*, **812**, 81
 Lopez-Coto, R., Marandon, V., & Brun, F. 2017, Proc. ICRC (Busan), **35**, 732
 Mazets, E. P., Golenetskij, S. V., & Guryan, Y. A. 1979, *SvAL*, **5**, 343
 Mohrmann, L., Specovius, A., Tiziani, D., et al. 2019, *A&A*, **632**, A72
 Mori, K., An, H., Feng, Q., et al. 2020, *ApJ*, **897**, 129
 Nigro, C., Deil, C., Zanin, R., et al. 2019, *A&A*, **625**, A10
 Parsons, R. D., & Hinton, J. A. 2014, *APh*, **56**, 26
 Price-Whelan, A. M., Sipőcz, B. M., Günther, H. M., et al. 2018, *AJ*, **156**, 123
 Schoorlemmer, H. 2019, Proc. ICRC (Madison), **36**, 785
 Vallée, J. P. 2014, *ApJS*, **215**, 1
 Vianello, G., Lauer, R. J., Younk, P., et al. 2015, arXiv:1507.08343
 Wakely, S. P., & Horan, D. 2008, Proc. ICRC (Mérida), **30**, 1341
 Wilks, S. S. 1938, *Ann. Math. Stat.*, **9**, 60
 Yao, J. M., Manchester, R. N., & Wang, N. 2017, *ApJ*, **835**, 29