

GALOIS-EQUIVARIANT MCKAY BIJECTIONS FOR PRIMES DIVIDING $q - 1$

BY

A. A. SCHAEFFER FRY*

*Department of Mathematics and Statistics, Metropolitan State University of Denver
Denver, CO 80217, USA
e-mail: aschae6@msudenver.edu*

ABSTRACT

We prove that for most groups of Lie type, the bijections used by Malle and Späth in the proof of Isaacs–Malle–Navarro’s inductive McKay conditions for the prime 2 and odd primes dividing $q - 1$ are also equivariant with respect to certain Galois automorphisms. In particular, this shows that these bijections are candidates for proving Navarro–Späth–Vallejo’s recently-posed inductive Galois–McKay conditions. On the way, we show that several simple groups of Lie type satisfy the McKay–Navarro conjecture for the prime 2.

* The author was supported in part by grants from the Simons Foundation (Award No. 351233) and the National Science Foundation (Award No. DMS-1801156). Part of the work was completed while the author was in residence at the Mathematical Sciences Research Institute in Berkeley, California during the Spring 2018 semester program “Group Representation Theory and Applications”, supported by the National Science Foundation under Award No. DMS-1440140. Received August 6, 2020 and in revised form February 12, 2021

1. Introduction

One of the main sets of questions of interest in the representation theory of finite groups falls under the umbrella of the so-called local-global conjectures, which relate the character theory of a finite group G to that of certain local subgroups. Another rich topic in the area is the problem of determining how certain other groups (for example, the group of automorphisms $\text{Aut}(G)$ of G , or the Galois group $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$) act on the set of irreducible characters of G and the related question of determining the fields of values of characters of G . This paper concerns the McKay–Navarro conjecture, which incorporates both of these main problems.

For a finite group G , the field $\mathbb{Q}(e^{2\pi i/|G|})$, obtained by adjoining the $|G|$ th roots of unity in some algebraic closure $\overline{\mathbb{Q}}$ of $\mathbb{Q}$, is a splitting field for G . The group

$$\mathcal{G} := \text{Gal}(\mathbb{Q}(e^{2\pi i/|G|})/\mathbb{Q})$$

acts naturally on the set of irreducible ordinary characters, $\text{Irr}(G)$, of G via $\chi^\sigma(g) := \chi(g)^\sigma$ for $\chi \in \text{Irr}(G)$, $g \in G$, $\sigma \in \mathcal{G}$.

In [Nav04], G. Navarro conjectured a refinement to the well-known McKay conjecture that incorporates this action of $\mathcal{G}$. Let ℓ be a prime and write $\mathcal{H}_\ell \leq \mathcal{G}$ for the subgroup consisting of $\sigma \in \mathcal{G}$ satisfying that there is some nonnegative integer r such that $\zeta^\sigma = \zeta^{\ell^r}$ for every ℓ' -root of unity ζ . Specifically, the McKay–Navarro conjecture (also sometimes known as the “Galois–McKay” conjecture) posits that if $P \in \text{Syl}_\ell(G)$ is a Sylow ℓ -subgroup of G , then for every $\sigma \in \mathcal{H}_\ell$, the number of characters in $\text{Irr}_{\ell'}(G)$ that are fixed by σ is the same as the number of characters in $\text{Irr}_{\ell'}(\text{N}_G(P))$ fixed by σ . Here for a finite group X , we write

$$\text{Irr}_{\ell'}(X) = \{\chi \in \text{Irr}(X) \mid \ell \nmid \chi(1)\}.$$

A stronger version of the McKay–Navarro conjecture says that, further, there should be an $\mathcal{H}_\ell$ -equivariant bijection between the sets $\text{Irr}_{\ell'}(G)$ and $\text{Irr}_{\ell'}(\text{N}_G(P))$. In particular, this would imply that the corresponding fields of values are preserved over the field of ℓ -adic numbers (see also [Tur08]). This version of the conjecture was recently reduced by Navarro–Späth–Vallejo in [NSV20] to proving certain inductive conditions for finite simple groups. These “inductive Galois–McKay conditions” can roughly be described as an “equivariant” condition and an “extension” condition. In this article, we are concerned with the first part of [NSV20, Definition 3.1] (the equivariant bijection part of the inductive Galois–McKay conditions):

CONDITION ‡: *Let G be a finite quasisimple group and let $Q \in \text{Syl}_\ell(G)$. Then there is a proper $\text{Aut}(G)_Q$ -stable subgroup M of G with $N_G(Q) \leq M$ and an $\text{Aut}(G)_Q \times \mathcal{H}_\ell$ -equivariant bijection*

$$\text{Irr}_{\ell'}(G) \rightarrow \text{Irr}_{\ell'}(M)$$

such that corresponding characters lie over the same character of $Z(G)$.

In [SF16, SFT18, SF19], the author proves a consequence of the McKay–Navarro conjecture in the case that $\ell = 2$. On the way to proving this consequence, the author described in [SF19] the action of Galois automorphisms on characters of groups of Lie type, in terms of the Howlett–Lehrer parametrization of Harish-Chandra series. The main obstruction to the action of $\mathcal{G}$ on these parameters being well-behaved is the presence of three characters of the so-called relative Weyl group that occur in the description. In [SFT20, Sections 3–4], one of these characters is studied in some detail on the way to describing the fields of values for characters of symplectic and special orthogonal groups. Here, we study the other two, especially for the principal series, which allows us to obtain our main results.

In [MS16], G. Malle and B. Späth complete the proof of the ordinary McKay conjecture for the prime $\ell = 2$. This groundbreaking result builds on previous work of Malle, Späth, and others, showing that simple groups satisfy the so-called “inductive McKay” conditions provided by Isaacs–Malle–Navarro in [IMN07, Section 10]. In particular, much of this work built upon [Mal07, Mal08], in which Malle provided a set of bijections in the case of groups of Lie type that served as candidates for the inductive McKay conditions. In the current article, we adapt the methods and results of [SF19] to show that in most cases for the prime $\ell = 2$ or odd primes ℓ dividing $q - 1$, these bijections used in [Mal07, Mal08, MS16] to prove the inductive McKay conditions for groups of Lie type defined over $\mathbb{F}_q$ are also $\mathcal{H}_\ell$ -equivariant, and therefore satisfy Condition ‡. In particular, these bijections are therefore candidates for eventually proving the inductive Galois–McKay conditions. The main result is the following:

THEOREM A: *Let q be a power of a prime and assume G is a group of Lie type of simply connected type defined over $\mathbb{F}_q$ and such that G is not of type A_n nor of Suzuki or Ree type. Let ℓ be a prime not dividing q and write $d_\ell(q)$ for the multiplicative order of q modulo ℓ if ℓ is odd, respectively modulo 4 if $\ell = 2$. Assume that one of the following holds:*

- (1) $d_\ell(q) = 1$ and either $\ell \neq 3$ or G is not $G_2(q)$ with $q \equiv 4, 7 \pmod{9}$;
 (2) $\ell = 2$, $d_\ell(q) = 2$, and G is not $D_n(q)$ or ${}^2D_n(q)$ with $n \geq 4$.

Then Condition $\nmid$ holds for G and the prime ℓ .

We remark that the omitted cases require additional techniques, which we plan to address in future work. Further, the “extension” condition in [NSV20, Definition 3.1] (i.e., parts (iii) and (iv) of the relation on $\mathcal{H}_\ell$ -triples [NSV20, Definition 1.5]) presents a separate set of challenges that we do not address here. However, in the cases covered by Theorem A in which $\text{Out}(G)$ is trivial (and hence this condition is trivial) and the simple group $G/Z(G)$ does not have an exceptional Schur multiplier, we obtain as an immediate corollary that $G/Z(G)$ satisfies the inductive Galois–McKay conditions. In particular, Theorem A yields that for $q = p$ a prime, the groups $G_2(p)$ with $p > 3$ and $F_4(p)$ and $E_8(p)$ with p odd satisfy the inductive Galois–McKay conditions for all primes ℓ that divide $p - 1$, with the possible exception of $\ell = 3$ for $G_2(p)$.

Setting aside for the moment the inductive conditions, even the statement of the McKay–Navarro conjecture has only been proved for a very limited number of simple groups, most notably the case of groups of Lie type in defining characteristic [Ruh17]. On the way to proving Theorem A, we show that if G is of type B or is exceptional but not of type E_6 , 2E_6 , nor Suzuki or Ree type, then all odd-degree characters of G are rational-valued. As a consequence, we obtain the statement of the McKay–Navarro conjecture for these groups.

THEOREM B: *Let S be as in one of the following cases:*

- q is any power of an odd prime and S is a simple group $G_2(q)$, ${}^3D_4(q)$, $F_4(q)$, $E_7(q)$, $E_8(q)$, or $B_n(q)$ with $n \geq 3$;
- q is a power of an odd prime such that $q \equiv \pm 1 \pmod{8}$ and S is a simple group $C_n(q)$ with $n \geq 2$;
- S is one of the exceptional covering groups $3.B_3(3)$ or $3.G_2(3)$.

Then the McKay–Navarro conjecture holds for the finite group S for the prime $\ell = 2$.

The structure of the paper is as follows. In Section 2, we set some notation and hypotheses to be used throughout the paper, and recall some previous results from [MS16] about odd-degree characters and from [SF19] regarding the action of $\mathcal{G}$ on characters of groups of Lie type. In Sections 3–5, we discuss in some detail the three characters of the relative Weyl group that arise in the

description of this action as mentioned above. In Section 6, we prove Theorem A(1) with the exception of $\ell = 2$ and $G = \mathrm{Sp}_{2n}(q)$ for certain q . In Section 7, we complete the proof of Theorem A when $\ell = 2$ and $G = \mathrm{Sp}_{2n}(q)$. Finally in Section 8, we prove Theorem A(2) for the remaining cases, as well as Theorem B.

2. Additional notation and preliminaries

In this section, we set some notation to be used throughout the paper, before recalling the main result of [SF19, Section 3] and some key ingredients from [MS16].

2.1. BASIC NOTATION. Let $X \leq Y$ be a subgroup of a finite group Y . For an irreducible character $\chi \in \mathrm{Irr}(Y)$, we denote by $\chi \downarrow X$ the restriction of χ to a character of X . For $\varphi \in \mathrm{Irr}(X)$, we denote by $\mathrm{Ind}_X^Y(\varphi)$ the induced character of Y . The notation $\mathrm{Irr}(X|\chi)$ for $\chi \in \mathrm{Irr}(Y)$ and $\mathrm{Irr}(Y|\varphi)$ for $\varphi \in \mathrm{Irr}(X)$ will denote the set of irreducible constituents of $\chi \downarrow X$, respectively of $\mathrm{Ind}_X^Y(\varphi)$. For a prime ℓ , the notation $\mathrm{Irr}_{\ell'}(\bullet)$ will always mean the subset of $\mathrm{Irr}(\bullet)$ consisting of characters with degree relatively prime to ℓ .

If $X \triangleleft Y$ is normal and $\varphi \in \mathrm{Irr}(X)$, we will denote by Y_φ the stabilizer (also known as the inertia subgroup) of φ in Y . As in [Spä09], if $\mathcal{X}$ is a subset of $\mathrm{Irr}(X)$ such that every $\varphi \in \mathcal{X}$ extends to its inertia group Y_φ in Y , we define an **extension map for $\mathcal{X}$ with respect to $X \triangleleft Y$** to be any map

$$\mathcal{X} \rightarrow \bigcup_{\varphi \in \mathcal{X}} \mathrm{Irr}(Y_\varphi)$$

that sends each $\varphi \in \mathcal{X}$ to one of its extensions to Y_φ .

If Y is a cyclic group of even order, we will denote by -1_Y the unique character of Y of order 2.

2.2. GROUPS OF LIE TYPE AND RELEVANT SUBGROUPS. Let $q = p^a$ be a power of a prime p . Throughout, unless otherwise specified, we let $\mathbf{G}$ be a simple, simply connected algebraic group defined over $\overline{\mathbb{F}}_q$.

Let $\mathbf{T} \leq \mathbf{B}$ be a maximal torus and Borel subgroup of $\mathbf{G}$, respectively. Let Φ be the root system of $\mathbf{G}$ with respect to $\mathbf{T} \leq \mathbf{B}$ and Δ a set of simple roots chosen with respect to $\mathbf{T}$. We will at times do computations in $\mathbf{G}$ using the Chevalley generators and relations, as in [GLS98, Theorem 1.12.1]. In particular, $x_\alpha(t)$, $n_\alpha(t)$, and $h_\alpha(t)$ for $t \in \overline{\mathbb{F}}_q$ and $\alpha \in \Phi$ are as defined there, and $\mathbf{T}$ may be written as the direct product of groups $\prod_{\alpha \in \Delta} \{h_\alpha(t) \mid t \in \overline{\mathbb{F}}_q^\times\}$ since $\mathbf{G}$ is simply connected.

Now let $G = \mathbf{G}^F$ be the group of fixed points of $\mathbf{G}$ under a Frobenius endomorphism F defined over $\mathbb{F}_q$. That is, $F: \mathbf{G} \rightarrow \mathbf{G}$ is of the form $F_q \circ \tau$, where τ is a graph automorphism defined by $\tau: x_\alpha(t) \mapsto x_{\tau'(\alpha)}(t)$ for $\alpha \in \Delta, t \in \overline{\mathbb{F}}_q$, and τ' a length-preserving symmetry of the Dynkin diagram associated to Δ ; and $F_q = F_p^a$ where $F_p: \mathbf{G} \rightarrow \mathbf{G}$ is defined by $F_p: x_\alpha(t) \mapsto x_\alpha(t^p)$ for $t \in \overline{\mathbb{F}}_q$ and $\alpha \in \Phi$. (In particular, we omit the cases of Suzuki and Ree groups.) Then $\mathbf{T}$ and $\mathbf{B}$ are F -stable.

As in [Spä09, Setting 2.1] or [MS16, Section 3.A], we let

$$V := \langle n_\alpha(\pm 1) \mid \alpha \in \Phi \rangle \leq N_{\mathbf{G}}(\mathbf{T})$$

be the **extended Weyl group** and let H be the group

$$H := V \cap \mathbf{T} = \langle h_\alpha(-1) \mid \alpha \in \Phi \rangle,$$

which is elementary abelian of size $(2, q-1)^{|\Delta|}$. Then we have $N_{\mathbf{G}}(\mathbf{T}) = \langle \mathbf{T}, V \rangle$. Let $\mathbf{W} := N_{\mathbf{G}}(\mathbf{T})/\mathbf{T}$ be the Weyl group of $\mathbf{G}$. (Note that $\mathbf{W} = V$ and $H = 1$ if $p = 2$.) Throughout, we further write $T := \mathbf{T}^F$ and $N := N_{\mathbf{G}}(\mathbf{T})^F$.

Now, we let $v \in \mathbf{G}$ be the canonical representative in V of the longest element of the Weyl group $\mathbf{W}$ of $\mathbf{G}$, as in [MS16, Section 3.A] or [Spä10, Definition 3.2]. Note that a suitable conjugation in $\mathbf{G}$ induces an automorphism of $\mathbf{G}$ mapping G to $\mathbf{G}^{vF}$. Write $T_1 = \mathbf{T}^{vF}$, $V_1 := V^{vF}$, and $H_1 := H^{vF}$. Then by [MS16, Lemma 3.2] (see also [Spä09, Proposition 6.4 and Table 2] for the case ${}^3\mathbf{D}_4(q)$ and [Spä09, Proposition 5.1 and Lemma 6.1] in the cases $\mathbf{G}_2(q)$, $\mathbf{F}_4(q)$, and $\mathbf{E}_8(q)$), we have $\mathbf{T} = C_{\mathbf{G}}(\mathbf{S})$ for some Sylow 2-torus $\mathbf{S}$ of $(\mathbf{G}, vF)$, and we have $N = TV$ and $N_1 = T_1V_1$, where $N_1 := N_{\mathbf{G}}(\mathbf{S})^{vF}$. Further, [Mal07, Proposition 5.11] yields that N controls G -fusion in T and N_1 controls $\mathbf{G}^{vF}$ -fusion in T_1 , and the results of [Mal07] imply that in most cases being considered in Theorem A, the groups N and N_1 are the natural choices to play the role of M in Condition $\S$.

We note the following, which follows from the fact that the longest element of $\mathbf{W}$ is central in the stated cases (see the proofs of [Spä09, Lemma 6.1] and [MS16, Lemma 3.2]).

LEMMA 2.1: *Assume $\mathbf{G}$ is of type $\mathbf{B}_n, \mathbf{C}_n, \mathbf{D}_{2n}, \mathbf{G}_2, \mathbf{F}_4, \mathbf{E}_7, \mathbf{E}_8$ or that $G = {}^2\mathbf{D}_n(q)$ or ${}^3\mathbf{D}_4(q)$. Then $V_1 = V$ and $H_1 = H$.*

If ℓ is a prime different from p , we will write $d_\ell(q)$ for the order of q modulo ℓ if ℓ is odd, and the order of q modulo 4 if $\ell = 2$. The role of T_1 and N_1 will be important in the case $d_2(q) = 2$, i.e. $q \equiv 3 \pmod{4}$.

2.3. GALOIS AUTOMORPHISMS AND HARISH-CHANDRA SERIES. Let $\mathbf{L} \leq \mathbf{P}$ be an F -stable Levi subgroup and parabolic subgroup of $\mathbf{G}$, respectively, satisfying $\mathbf{T} \leq \mathbf{L}$ and $\mathbf{B} \leq \mathbf{P}$, so that $L := \mathbf{L}^F$ is a split standard Levi subgroup in the parabolic subgroup $P := \mathbf{P}^F$ of $G = \mathbf{G}^F$. Let $\lambda \in \text{Irr}(L)$ be a cuspidal character and write $W(\lambda) := N(L)_\lambda/L$, where $N(L) := (N_G(L) \cap N)L$. Recall that such a pair (L, λ) is called a cuspidal pair for G , and the Harish-Chandra induced character $R_L^G(\lambda)$ is defined as $\text{Ind}_P^G(\text{Inf}_L^P(\lambda))$, where $\text{Inf}_L^P(\lambda)$ denotes the inflation of λ to a character of P , viewing L as a quotient of P by its unipotent radical. We will write $\mathcal{E}(G, L, \lambda) \subseteq \text{Irr}(G)$ for the set of constituents of $R_L^G(\lambda)$, known as a Harish-Chandra series.

Using the fact that every cuspidal $\psi \in \text{Irr}(L)$ extends to $N(L)_\psi$ due to [Gec93] and [Lus84, Theorem 8.6], we will apply the concept of extension maps to the case of cuspidal characters of L and $L \triangleleft N(L)$. Thanks to this and the work of Howlett and Lehrer [HL80, HL83], for a cuspidal pair (L, λ) of G , we have a bijection between $\mathcal{E}(G, L, \lambda)$ and $\text{Irr}(W(\lambda))$ induced by a bijection

$$f: \text{Irr}(\text{End}_{\overline{\mathbb{Q}}_G}(R_T^G(\lambda))) \rightarrow \text{Irr}(W(\lambda)),$$

where by $\text{End}_{\overline{\mathbb{Q}}_G}(R_T^G(\lambda))$ we mean the endomorphism algebra of a fixed module affording $R_T^G(\lambda)$. We write $R_L^G(\lambda)_\eta$ for the character of G in $\mathcal{E}(G, L, \lambda)$ corresponding to $\eta \in \text{Irr}(W(\lambda))$. In particular, every character of G can be written in such a way.

Now, there is a subsystem $\Phi_\lambda \subseteq \Phi$ with simple roots $\Delta_\lambda \subseteq \Phi_\lambda \cap \Phi^+$ such that $W(\lambda)$ can be decomposed as a semidirect product $R(\lambda) \rtimes C(\lambda)$, where $R(\lambda) = \langle s_\alpha \mid \alpha \in \Phi_\lambda \rangle$ is a Weyl group with root system Φ_λ and $C(\lambda)$ is the stabilizer of Δ_λ in $W(\lambda)$. (See [Car93, Section 10.6] or [HL80, Section 2] for more details.) Here for $\alpha \in \Phi$, s_α is the corresponding reflection in $W := \mathbf{W}^F = N/T$ induced by the element $n_\alpha(-1)$ of N . The root system Φ_λ is obtained as follows. Set

$$\Phi^b = \{\alpha \in \Phi \setminus \Phi_L \mid w(\Delta_L \cup \{\alpha\}) \subseteq \Delta \text{ for some } w \in W \text{ and } (w_0^L w_0^\alpha)^2 = 1\}.$$

Here w_0^L , and w_0^α are the longest elements in $W(L) := N(L)/L$ and $\langle W(L), s_\alpha \rangle$, respectively, and $\Phi_L \subseteq \Phi$ is the root system of $W(L)$ with simple system $\Delta_L \subseteq \Delta$. Then for $\alpha \in \Phi^b$, letting L_α denote the standard Levi subgroup of G with simple system $\Delta_L \cup \{\alpha\}$, L is a standard Levi subgroup of L_α and $p_{\alpha, \lambda} \geq 1$ is defined to be the ratio between the degrees of the two constituents of $R_{L_\alpha}^{L_\alpha}(\lambda)$. Then

$$\Phi_\lambda = \{\alpha \in \Phi^b \mid s_\alpha \in W(\lambda) \text{ and } p_{\alpha, \lambda} \neq 1\}.$$

Theorem 3.8 of [SF19] describes the action of $\mathcal{G}$ on $\text{Irr}(G)$ with respect to the parameters (L, λ, η) . We restate it here for the convenience of the reader, although we first must establish the necessary notation.

NOTATION FOR THEOREM 2.2. Let (L, λ) be a cuspidal pair for G . Fix an extension map Λ for cuspidal characters with respect to $L \triangleleft N(L)$, so that for each cuspidal $\psi \in \text{Irr}(L)$, we have $\Lambda(\psi)$ is an extension of ψ to $N(L)_\psi$. For $\sigma \in \mathcal{G}$, note that $N(L)_\lambda = N(L)_{\lambda^\sigma}$, and define $\delta_{\lambda, \sigma}$ to be the linear character of $W(\lambda)$ such that $\Lambda(\lambda)^\sigma = \delta_{\lambda, \sigma} \Lambda(\lambda^\sigma)$, guaranteed by Gallagher's theorem [Isa06, Corollary 6.17]. Further, let $\delta'_{\lambda, \sigma} \in \text{Irr}(W(\lambda))$ be the character such that $\delta'_{\lambda, \sigma}(w) = \delta_{\lambda, \sigma}(w)$ for $w \in C(\lambda)$ and $\delta'_{\lambda, \sigma}(w) = 1$ for $w \in R(\lambda)$. Let $\gamma_{\lambda, \sigma}$ be the function on $W(\lambda)$ such that

$$\gamma_{\lambda, \sigma}(w) = \frac{\sqrt{\text{ind}(w_c)}^\sigma}{\sqrt{\text{ind}(w_c)}} \quad \text{where } w = w_r w_c \text{ for } w_c \in C(\lambda) \text{ and } w_r \in R(\lambda).$$

Here

$$\text{ind}(w_c) := |U_0 \cap U_0^{w_0 w_c}|,$$

where U_0 is the unipotent radical of B and w_0 is the longest element of W . (See Lemma 3.1 below for a description of $\gamma_{\lambda, \sigma}$ more conducive to our purposes.)

For $\eta \in \text{Irr}(W(\lambda))$, we denote by $\eta^{(\sigma)}$ the character $\mathfrak{f}(\tilde{\eta}^\sigma)$, where

$$\tilde{\eta} \in \text{Irr}(\text{End}_{\overline{\mathbb{Q}}_G}(\mathbf{R}_T^G(\lambda)))$$

is such that $\mathfrak{f}(\tilde{\eta}) = \eta$. (See [SF19, Section 3.5].) Note that this is not necessarily the same as η^σ , although we show below in Section 5 that this is often the case.

THEOREM 2.2 ([SF19, Theorem 3.8]): *Let $\sigma \in \mathcal{G}$, let (L, λ) be a cuspidal pair for G , and keep the notation above. Let $\eta \in \text{Irr}(W(\lambda))$. Then*

$$(\mathbf{R}_L^G(\lambda)_\eta)^\sigma = \mathbf{R}_L^G(\lambda^\sigma)_{\eta'},$$

where $\eta' \in \text{Irr}(W(\lambda)) = \text{Irr}(W(\lambda^\sigma))$ is defined by

$$\eta'(w) = \gamma_{\lambda, \sigma}(w) \delta'_{\lambda, \sigma}(w^{-1}) \eta^{(\sigma)}(w)$$

for each $w \in W(\lambda)$.

Theorem 2.2 shows that the characters $\gamma_{\lambda, \sigma}$, $\delta'_{\lambda, \sigma}$, and $\eta^{(\sigma)}$ are the obstructions to $\mathcal{G}$ acting on the parameters (L, λ, η) in an equivariant way. We discuss these characters in some detail below in Sections 3, 4, and 5, respectively.

2.4. LUSZTIG SERIES AND ODD-DEGREE CHARACTERS. We will write G^* for the dual $G^* = \mathbf{G}^{*F^*}$ of G , where $(\mathbf{G}^*, F^*)$ is dual to $(\mathbf{G}, F)$ as in, for example, [Car93, Section 4.2]. Let $\mathbf{T}^*$ be an F^* -stable maximal torus of $\mathbf{G}^*$ dual to $\mathbf{T}$. Write $T = \mathbf{T}^F$ and $T^* = (\mathbf{T}^*)^{F^*}$. We write $W = \mathbf{W}^F$, where $\mathbf{W} = \mathbf{N}_{\mathbf{G}}(\mathbf{T})/\mathbf{T}$, and similarly for W^* . This duality induces an isomorphism $\text{Irr}(T) \rightarrow T^*$ (see, for example, [Car93, Proposition 4.4.1]). For $\lambda \in \text{Irr}(T)$, the following lemma allows us to view $W(\lambda)$ and $R(\lambda)$ as the groups $W(s)$ and $W^\circ(s)$ for a certain semisimple element $s \in G^*$, where we write $W(s)$ and $W^\circ(s)$ for the fixed points of the Weyl groups of $\mathbf{C}_{\mathbf{G}^*}(s)$ and $\mathbf{C}_{\mathbf{G}^*}^\circ(s)$, respectively, under F^* , and implies that $C(\lambda)$ is isomorphic to a subgroup of $(\mathbf{Z}(\mathbf{G})/\mathbf{Z}(\mathbf{G})^\circ)^F$. (See also [SFT20, Lemma 3.8].)

LEMMA 2.3 ([SF19, Lemma 4.5]): *Let $\lambda \in \text{Irr}(T)$ and let $s \in T^*$ correspond to λ in the sense of [Car93, Proposition 4.4.1]. Then $W(\lambda)$ is isomorphic to $W(s)$. Further, if $\mathbf{G}$ is simple of simply connected type, not of type A_n , then there is an isomorphism $\kappa: W(\lambda) \rightarrow W(s)$ such that $\kappa(R(\lambda)) = W^\circ(s)$. In particular, in this case $W(\lambda)/R(\lambda)$ is isomorphic to $(\mathbf{C}_{\mathbf{G}^*}(s)/\mathbf{C}_{\mathbf{G}^*}^\circ(s))^{F^*}$.*

Recall that $\text{Irr}(G)$ may be partitioned into so-called **rational Lusztig series** $\mathcal{E}(G, s)$ ranging over the G^* -classes of semisimple elements s in G^* . Each $\mathcal{E}(G, s)$ is further a disjoint union of Harish-Chandra series $\mathcal{E}(G, L, \lambda)$ satisfying $\lambda \in \mathcal{E}(L, s)$ (see [Bon06, 11.10]). The following results from [MS16] describe the Lusztig and Harish-Chandra series containing odd-degree characters.

THEOREM 2.4 ([MS16, Theorem 7.7]): *Let q be odd and let $\mathbf{G}$ be simple, of simply connected type, not of type A_n . Let $\chi \in \text{Irr}_{2'}(G)$. Then either χ lies in the principal series of G , or $q \equiv 3 \pmod{4}$, $G = \text{Sp}_{2n}(q)$ with $n \geq 1$ odd, $\chi \in \mathcal{E}(G, s)$ with $\mathbf{C}_{G^*}(s) = \mathbf{B}_{2k}(q) \cdot {}^2\mathbf{D}_{n-2k}(q).2$, where $0 \leq k \leq (n-3)/2$ and χ lies in the Harish-Chandra series of a cuspidal character of degree $(q-1)/2$ of a Levi subgroup $\text{Sp}_2(q) \times (q-1)^{n-1}$.*

LEMMA 2.5 ([MS16, extension of Lemma 7.9]): *Let q be odd and let $\mathbf{G}$ be simple, of simply connected type, not of type A_n . Let $\chi \in \text{Irr}_{2'}(G)$. Then $\chi = \mathbf{R}_T^G(\lambda)_\eta$, where T is a maximally split torus of G , $\lambda \in \text{Irr}(T)$ is such that $2 \nmid [W : W(\lambda)]$, and $\eta \in \text{Irr}_{2'}(W(\lambda))$, except possibly in the case $\mathbf{G}$ is type C_n with n odd and $q \equiv 3 \pmod{4}$. In the latter case, χ may also be of the form $\chi = \mathbf{R}_L^G(\lambda)_\eta$ with (L, λ) as in Theorem 2.4, $2 \nmid [W(L) : W(\lambda)]$ and $\eta \in \text{Irr}_{2'}(W(\lambda))$.*

The following is [MS16, Lemma 7.5], extended to some additional exceptional groups.

LEMMA 2.6: *Let $G = \mathbf{G}^F$ where $\mathbf{G}$ is simple of simply connected type, not of type A_n , C_n , E_6 , or D_{2n+1} , and such that F is a Frobenius endomorphism defining G over $\mathbb{F}_q$ with q odd. Then if $\chi \in \text{Irr}_{2'}(G)$ lies in the rational series $\mathcal{E}(G, s)$, then $s^2 = 1$.*

Proof. Let $s \in G^*$ be such that $\chi \in \text{Irr}_{2'}(G)$ lies in the series $\mathcal{E}(G, s)$. Recall that $\chi \in \text{Irr}_{2'}(G)$ implies that s centralizes a Sylow 2-subgroup of G^* . Then for types B_n, D_{2n}, E_7 , this is [MS16, Lemma 7.5]. For the remaining cases, the statement can be obtained by analyzing the possible centralizer structures of G^* , which are listed, for example, at [Lüb07], or in [DM87] for type 3D_4 . In particular, if s is non-central, then we have $2 \mid [W : W(s)]$ unless $C_{G^*}(s)$ is of type $A_1(q) + A_1(q)$ if $G = G_2(q)$; $B_4(q)$ if $G = F_4(q)$; $D_8(q)$ if $G = E_8(q)$; and $A_1(q^3) + A_1(q)$ if $G = {}^3D_4(q)$, in which cases $s^2 = 1$. ■

3. Square roots and the character $\gamma_{\lambda, \sigma}$

Let p be a prime and q a power of p . Keep the notation of Theorem 2.2. The following, found as [SFT20, Lemma 3.11], shows that $\gamma_{\lambda, \sigma}$ is indeed a character of $W(\lambda)$ (and hence of $C(\lambda)$) and is closely related to the action of $\sigma \in \mathcal{G}$ on $\sqrt{p}$. We remark that here we do not require the assumption that $\mathbf{G}$ is simply connected.

LEMMA 3.1: *Assume (L, λ) is a cuspidal pair for a finite reductive group $G = \mathbf{G}^F$ defined over $\mathbb{F}_q$ with q a power of a prime p , and let $\sigma \in \mathcal{G}$. Then in the notation of Theorem 2.2, $\gamma_{\lambda, \sigma}$ is a character of $W(\lambda)$, and is moreover trivial if and only if at least one of the following holds:*

- q is a square;
- the length $l(w_c)$, with respect to the Weyl group of $\mathbf{G}$, is even for each $w_c \in C(\lambda)$; or
- σ fixes $\sqrt{p}$.

Otherwise,

$$\gamma_{\lambda, \sigma}(w) = (-1)^{l(w_c)},$$

where $w = w_r w_c$ with $w_r \in R(\lambda)$ and $w_c \in C(\lambda)$.

Let $\ell \neq p$ be another prime. In [SFT20, Section 4], we discuss the action of $\mathcal{H}_\ell$ on $\sqrt{p}$ (and on $\gamma_{\lambda,\sigma}$) when p is odd. We record the following observations, which extend the discussion to the case $p = 2$.

LEMMA 3.2: *Let ℓ be an odd prime and let $\sigma \in \mathcal{H}_\ell$ with corresponding integer r such that $\sigma(\zeta) = \zeta^{\ell^r}$ for every ℓ' -root of unity ζ . Let $p \neq \ell$ be a prime. Then $\sqrt{p}^\sigma = \sqrt{p}$ if and only if r is even or p is a square modulo ℓ .*

Proof. For p odd, this is [SFT20, Lemma 4.4]. Note that $\sqrt{2}$ may be written

$$\sqrt{2} = \zeta_8 + \zeta_8^{-1}$$

for some fixed primitive 8th root of unity ζ_8 . Then $\sqrt{2}^\sigma = \sqrt{2}$ if and only if ℓ^r is $\pm 1 \pmod{8}$. This proves the claim, since the condition $\ell \equiv \pm 1 \pmod{8}$ is equivalent to 2 being square modulo ℓ . ■

COROLLARY 3.3: *Keep the hypothesis of Lemma 3.1, and assume that $\ell \neq p$ is an odd prime dividing $q - 1$. Then $\mathbb{Q}(\sqrt{q})$ is contained in the fixed field of $\mathcal{H}_\ell$ and the character $\gamma_{\lambda,\sigma}$ is trivial.*

Proof. This follows from Lemmas 3.1 and 3.2, since the condition $q \equiv 1 \pmod{\ell}$ forces p to be a square modulo ℓ if q is not a square. ■

In the case $\ell = 2$, we have the following, returning to our default assumption that $\mathbf{G}$ is simply connected.

PROPOSITION 3.4: *Let $G = \mathbf{G}^F$ where $\mathbf{G}$ is simple of simply connected type, not of type A_n or C_n , and such that F is a Frobenius endomorphism defining G over $\mathbb{F}_q$ with q odd. Let $\chi = R_T^G(\lambda)_\eta \in \text{Irr}_{2'}(G)$, as in Lemma 2.5. Then*

$$\gamma_{\lambda,\sigma} = 1$$

for every $\sigma \in \mathcal{G}$.

Proof. If $|\mathbf{Z}(\mathbf{G})^F|$ is odd, this follows from Lemma 2.3 and the fact that $\gamma_{\lambda,\sigma}$ has order dividing 2. If $\mathbf{G}$ is type B_n ($n \geq 3$) or D_n ($n \geq 4$) and $G \neq {}^3D_4(q)$, then every element of $C(\lambda)$ has even length in the Weyl group of $\mathbf{G}$, by [SF19, Proposition 4.11], and hence the statement holds using Lemma 3.1. This leaves the case $\mathbf{G}$ is of type E_7 . As in Lemma 2.6, we may analyze the possible centralizer structures and see that either $s = 1$ or $C_{\mathbf{G}^*}(s)$ is connected, of type $D_6 + A_1$. In either case, this again yields $C(\lambda) = 1$, and the result follows. ■

4. The extension map Λ

Keep the notation of Section 2. In particular, recall that $\mathbf{G}$ is simple of simply connected type and $G = \mathbf{G}^F$ is defined over $\mathbb{F}_q$ with q a power of the prime p , not of Suzuki or Ree type. We have

$$V := \langle n_\alpha(\pm 1) \mid \alpha \in \Phi \rangle \leq N_{\mathbf{G}}(\mathbf{T}),$$

$H := V \cap \mathbf{T}$ is an elementary abelian 2-group (and trivial when $p = 2$), $N = TV$, and $N_1 = T_1V_1$. For $\lambda \in \text{Irr}(T)$, note that $N_\lambda = TV_\lambda$.

Let $\iota: \mathbf{G} \hookrightarrow \tilde{\mathbf{G}}$ be a regular embedding such that F extends to a Frobenius endomorphism of $\tilde{\mathbf{G}}$, which we will also denote by F . (See, e.g., [GM20, Section 1.7] for more details on regular embeddings.) We will write $\tilde{G} := \tilde{\mathbf{G}}^F$. Let D denote the subgroup of outer automorphisms generated by the field automorphism F_p and graph automorphisms commuting with F , so that $\text{Aut}(G)$ is induced by $\tilde{G} \rtimes D$ (see, e.g., [GLS98, Theorem 2.5.1]).

Our goal in this section is to obtain an extension map Λ with respect to $T \triangleleft N$ that is “close to” $\mathcal{G}$ -equivariant. In particular, we make use of the existing maps from [Spä09, MS16]. We begin with the case of some exceptional groups.

LEMMA 4.1: *Let G be as above, and assume either $p = 2$ or $\mathbf{G}$ is of type G_2, F_4, E_6 , or E_8 . Then there is an $ND \times \mathcal{G}$ -equivariant extension map Λ with respect to $T \triangleleft N$.*

Proof. The construction of the map Λ is exactly as in [MS16, Corollary 3.13] in the case $d = 1$ and v is trivial, using the construction from [Spä09, Section 5] for a VD -equivariant extension map with respect to $H \triangleleft V$. We provide the construction for the convenience of the reader, and to show that it is $\mathcal{G}$ -equivariant.

(1) A $\mathcal{G}$ -EQUIVARIANT EXTENSION MAP WITH RESPECT TO $H \triangleleft V$. First, note that when $p = 2$, we have $H = 1$. Hence we may assume p is odd. According to [Spä09, Lemma 5.3], for each $\delta \in \text{Irr}(H)$, there is a subset $R(\delta)$ of Φ such that $V_\delta = \langle H, n_\alpha(-1) \mid \alpha \in R(\delta) \rangle$. By [Spä09, Proof of Lemma 5.4] there is an extension map Λ'_0 with respect to $H \triangleleft V$ such that $\Lambda'_0(\delta)$ is trivial on $n_\alpha(-1)$ for each $\alpha \in R(\delta)$. Note that H is an elementary abelian 2-group, so $\delta^\sigma = \delta$ for all $\delta \in \text{Irr}(H)$ and all $\sigma \in \mathcal{G}$. Hence we see that

$$\Lambda'_0(\delta)^\sigma = \Lambda'_0(\delta) = \Lambda'_0(\delta^\sigma)$$

for each $\delta \in \text{Irr}(H)$ and $\sigma \in \mathcal{G}$.

(2) A $VD \times \mathcal{G}$ -EQUIVARIANT EXTENSION MAP WITH RESPECT TO $H \triangleleft V$. Let $\delta_1, \dots, \delta_r$ be representatives of the V -orbits on $\text{Irr}(H)$. For each $x \in V$ and $i \in \{1, \dots, r\}$, there is some linear $\mu_{x,i} \in \text{Irr}(V_{\delta_i}/H)$ such that

$$\Lambda'_0(\delta_i)^x = \mu_{x,i} \Lambda'_0(\delta_i^x).$$

Note that for each x and i , $\mu_{x,i}$ is fixed by each $\sigma \in \mathcal{G}$, since V_{δ_i}/H is generated by involutions and $\mu_{x,i}$ is a linear character. Then we define another extension map with respect to $H \triangleleft V$ via $\Lambda_0(\delta_i) := \Lambda'_0(\delta_i)$ for each i and

$$\Lambda_0(\delta_i^x) := \Lambda'_0(\delta_i)^x = \mu_{x,i} \Lambda'_0(\delta_i^x)$$

for each $x \in V$. This map is well-defined, since if $\delta_i^x = \delta_i^y$, then $xy^{-1} \in V_{\delta_i}$ fixes δ_i and $\Lambda'_0(\delta_i)$. By construction, Λ_0 is V -equivariant. Further, we have for each $x \in V$, $i \in \{1, \dots, r\}$, and $\sigma \in \mathcal{G}$,

$$\Lambda_0(\delta_i^x)^\sigma = \mu_{x,i}^\sigma \Lambda'_0(\delta_i^x)^\sigma = \mu_{x,i} \Lambda'_0(\delta_i^x) = \Lambda_0(\delta_i^x).$$

Hence Λ_0 is a V -equivariant extension map with respect to $H \triangleleft V$ which still satisfies $\Lambda_0(\delta)^\sigma = \Lambda_0(\delta) = \Lambda_0(\delta^\sigma)$ for each $\delta \in \text{Irr}(H)$ and $\sigma \in \mathcal{G}$. Further, since F_0 acts trivially on V , the extension map Λ_0 is $VD \times \mathcal{G}$ -equivariant by the proofs of [MS16, Proposition 3.9] and [Spä09, Lemma 8.2].

(3) THE $ND \times \mathcal{G}$ -EQUIVARIANT EXTENSION MAP WITH RESPECT TO $T \triangleleft N$. Finally, we use the strategy of [Spä09, 4.2] to obtain the desired extension map with respect to $T \triangleleft N$. For $\lambda \in \text{Irr}(T)$, we define an extension $\Lambda(\lambda) = \Lambda_\lambda$ of λ to N_λ via $\Lambda_\lambda(vt) = \Lambda_0(\lambda \downarrow_H)(v)\lambda(t)$ for each $v \in V_\lambda$ and $t \in T$, since $N = TV$. Then Λ is ND -equivariant by construction, and

$$\Lambda_\lambda^\sigma(vt) = \Lambda_0(\lambda \downarrow_H)^\sigma(v)\lambda^\sigma(t) = \Lambda_0(\lambda \downarrow_H^\sigma)(v)\lambda^\sigma(t) = \Lambda_{\lambda^\sigma}(vt)$$

for each $\sigma \in \mathcal{G}$, $v \in V_\lambda$, and $t \in T$. ■

Remark 4.2: When $G = {}^3\mathbf{D}_4(q)$, we similarly have an ND -equivariant extension map Λ with respect to $T \triangleleft N$, by [CS13, Theorem 3.6].

For the remainder of the article, we let Λ denote the ND -equivariant extension map with respect to $T \triangleleft N$ as in Lemma 4.1, Remark 4.2, or [MS16, Corollary 3.13]. We remark that for the overlapping case that $\mathbf{G}$ is type $\mathbf{E}_6$, these maps agree.

Recall that for $\lambda \in \text{Irr}(T)$ and $\sigma \in \mathcal{G}$, we let $\delta_{\lambda,\sigma} \in \text{Irr}(W(\lambda))$ denote the linear character of $W(\lambda) = W(\lambda^\sigma)$ such that $\Lambda(\lambda)^\sigma = \delta_{\lambda,\sigma} \Lambda(\lambda^\sigma)$. It will also sometimes be useful to denote $\Lambda_\lambda := \Lambda(\lambda)$.

Let R_λ be the subgroup of N_λ given by

$$R_\lambda := \langle T, n_\alpha(-1) \mid \alpha \in \Phi_\lambda \rangle,$$

so that $R_\lambda/T \cong R(\lambda)$. Note that $\Phi_\lambda = \Phi_{\lambda^\sigma}$ (see [SF19, Lemma 3.2]). The following extends [SFT20, Lemma 3.13] and implies that the character $\delta_{\lambda,\sigma}$ of $W(\lambda)$ can be viewed as a character of $C(\lambda)$, and hence is equal to $\delta'_{\lambda,\sigma}$.

LEMMA 4.3: *Keep the notation from the beginning of the section, and assume $\mathbf{G}$ is not of type A_n . Let $\lambda \in \text{Irr}(T)$ and $\sigma \in \mathcal{G}$. Then*

$$\Lambda_\lambda^\sigma \downarrow_{R_\lambda} = \Lambda_{\lambda^\sigma} \downarrow_{R_\lambda}.$$

In particular,

$$R(\lambda) \leq \ker \delta_{\lambda,\sigma}.$$

Proof. Let $\lambda_0 := \lambda \downarrow_H$ and note that $\lambda_0^\sigma = \lambda_0$, since H is an elementary abelian 2-group. By the construction in the proofs of [MS16, Corollary 3.13] and [Spä09, Lemma 4.2], we see that

$$\Lambda_\lambda(x) = \Lambda_{\lambda^\sigma}(x) \quad \text{for } x \in V_\lambda,$$

and hence for $x \in V(\lambda) := \langle n_\alpha(-1) \mid \alpha \in \Phi_\lambda \rangle$. Then for $t \in T$ and $x \in V(\lambda)$, we have

$$\Lambda_\lambda^\sigma(tx) = \Lambda_\lambda^\sigma(t)\Lambda_\lambda^\sigma(x) = \lambda^\sigma(t)\Lambda_\lambda(x) = \lambda^\sigma(t)\Lambda_{\lambda^\sigma}(x) = \Lambda_{\lambda^\sigma}(tx),$$

where the second equality is since $\Lambda_\lambda(x) \in \{\pm 1\}$ for $x \in V(\lambda)$, thanks to [SFT20, Lemma 3.13]. ■

COROLLARY 4.4: *Let G be $G_2(q)$, $F_4(q)$, ${}^3D_4(q)$, $E_6(q)$, ${}^2E_6(q)$, or $E_8(q)$. Then Λ is $ND \times \mathcal{G}$ -equivariant. Further, there is an $N_1D \times \mathcal{G}$ -equivariant extension map Λ_1 with respect to $T_1 \triangleleft N_1$.*

Proof. The statement about Λ is just a restatement of Lemma 4.1, except for the case ${}^3D_4(q)$. In the latter case, since $C(\lambda) = 1$ by Lemma 2.3, Lemma 4.3 immediately yields that the map Λ in Remark 4.2 is also $\mathcal{G}$ -equivariant. Hence we assume $q \equiv 3 \pmod{4}$ and prove the second statement.

If $\mathbf{G}$ is not of type E_6 , this follows analogously to Lemma 4.1 (and appealing to Lemma 4.3 in the case ${}^3D_4(q)$ since $C(\lambda) = 1$), since $V_1 = V$, $H_1 = H$, and $N_1 = T_1V$ in these cases by Lemma 2.1. If $\mathbf{G}$ is type E_6 , then the proof of [Spä09, Lemma 6.1] shows that either again $V_1 = V$ and $H_1 = H$ or $V_1 \cong V'$ and $H_1 \cong H'$, where $H' \triangleleft V'$ are the corresponding groups for a root system of type F_4 . In either case, the proof of Lemma 4.1 again yields the result. ■

COROLLARY 4.5: *In the situation of Lemma 4.3, we have $\delta_{\lambda,\sigma}^2 = 1$.*

Proof. If $\mathbf{G}$ is of type $\mathbf{E}_6$ or $(Z(\mathbf{G})/Z^\circ(\mathbf{G}))^F$ is trivial or an elementary abelian 2-group, then Lemmas 2.3, 4.1, and 4.3 imply the result, since $\delta_{\lambda,\sigma}$ is trivial for $\mathbf{E}_6$ and can be thought of as a character of $C(\lambda)$ in each case.

This leaves the case $\mathbf{G}$ is of type $\mathbf{D}_n$ with n odd. In this case, along the lines of the illustration in Lemma 4.1(3), the extension map Λ is constructed using an extension map Λ_0 with respect to $H \triangleleft V$, which is constructed in [MS16, Proposition 3.10]. Using an embedding of $\mathbf{G}$ into a group $\overline{\mathbf{G}}$ of type $\mathbf{B}_n$, the map Λ_0 in [MS16, Proposition 3.10] is moreover defined in terms of an extension map Λ_B with respect to $\overline{H} \triangleleft \overline{V}$ in the case of type $\mathbf{B}$, where $\overline{V}$ is the corresponding extended Weyl group for $\overline{\mathbf{G}}$ and $\overline{H} = H$ is the toral subgroup. Specifically, we have $\Lambda_0(\lambda) = (\Lambda_B(\lambda) \downarrow_{V_\lambda})^t$ for a specific $t \in \mathbf{T}$. Here to alleviate notation, we let

$$\Lambda_0(\lambda) := \Lambda_0(\lambda \downarrow_H) \quad \text{and} \quad \Lambda_B(\lambda) := \Lambda_B(\lambda \downarrow_H).$$

Let $\sigma \in \mathcal{G}$. Letting $\delta_B \in \text{Irr}(\overline{V}_{\lambda \downarrow_H}/H)$ be such that $\Lambda_B(\lambda)^\sigma = \delta_B \Lambda_B(\lambda^\sigma)$, we then have

$$\begin{aligned} \Lambda_0(\lambda^\sigma) &= (\Lambda_B(\lambda^\sigma) \downarrow_{V_\lambda})^t = ((\delta_B^{-1} \Lambda_B(\lambda)^\sigma) \downarrow_{V_\lambda})^t \\ &= (\delta_B^{-1} \downarrow_{V_\lambda})^t (\Lambda_B(\lambda)^\sigma \downarrow_{V_\lambda})^t \\ &= (\delta_B^{-1} \downarrow_{V_\lambda})^t \Lambda_0(\lambda)^\sigma. \end{aligned}$$

Then since $\Lambda_\lambda(tv) = \lambda(t)\Lambda_0(\lambda)(v)$ for $t \in T$ and $v \in V_\lambda$, we see that

$$\delta_{\lambda,\sigma} = (\delta_B) \downarrow_{V_\lambda}^t.$$

But since δ_B has order dividing 2, this completes the proof. $\blacksquare$

We next show that Λ is further $\mathcal{G}$ -equivariant in the case of $\mathbf{E}_7$ or $\mathbf{B}_n$, when restricted to the members of $\text{Irr}(T)$ such that $\mathcal{E}(G, T, \lambda)$ contains odd-degree characters.

PROPOSITION 4.6: *Let $G = \mathbf{G}^F$ such that $\mathbf{G}$ is simple or simply connected type $\mathbf{B}_n$ with $n \geq 3$ or $\mathbf{E}_7$. Assume q is odd and that $\mathcal{E}(G, T, \lambda) \cap \text{Irr}_{2'}(G)$ is nonempty. Then $\delta_{\lambda,\sigma} = 1$ for any $\sigma \in \mathcal{G}$.*

Proof. Note that if $\mathbf{G}$ is type $\mathbf{E}_7$, then the proof of Proposition 3.4 yields

$$R(\lambda) = W(\lambda)$$

for any $\lambda \in \text{Irr}(T)$ such that $\mathcal{E}(G, T, \lambda) \cap \text{Irr}_{2'}(G) \neq \emptyset$, and hence $\delta_{\lambda,\sigma} = 1$ for any $\sigma \in \mathcal{G}$ in this case.

So let $\mathbf{G}$ be of type B_n , and assume that $R(\lambda) \neq W(\lambda)$, as otherwise we have $\delta_{\lambda,\sigma} = 1$ trivially by appealing to Lemma 4.3. Then $\mathbf{G}^*$ is type C_n and $W(\lambda) \cong W(s)$ is type $(C_{n/2} \times C_{n/2}) \cdot 2$ with $n = 2^a \geq 4$, from the proof of [SF19, Lemma 4.11]. (See also [MS16, Lemma 7.6].)

Let $\Delta = \{\alpha_1, \dots, \alpha_n\}$ be a system of simple roots for $\mathbf{G}$, with labeling as in [GLS98, Remark 1.8.8], so that $\alpha_i = e_i - e_{i+1}$ for $1 \leq i < n$ and $\alpha_n = e_n$, where $\{e_1, \dots, e_n\}$ is an orthonormal basis for $\mathbb{R}^n$. The following can be seen using the relevant Chevalley relations, found for example in [GLS98, Theorem 1.12.1].

Let ζ be a generator for $\mathbb{F}_q^\times$. Then λ can be described by $\lambda(h_{\alpha_{n/2}}(\zeta)) = -1$ and $\lambda(h_{\alpha_i}(\zeta)) = 1$ for $i \neq n/2$. Hence $R(\lambda) \cong C_{n/2}^2$ is generated by the image in $\mathbf{W}$ of the elements $n_{e_i}(-1)$ for $1 \leq i \leq n$, together with the elements $n_{\alpha_i}(-1)$ for $1 \leq i < n$ with $i \neq n/2$ and $C(\lambda)$ is generated by the image in $\mathbf{W}$ of the element

$$c := n_{e_1 - e_{n/2+1}}(-1) n_{e_2 - e_{n/2+2}}(-1) \cdots n_{e_n - e_n}(-1).$$

Now, we have $h_{e_i - e_{n/2+i}}(-1) = h_{\alpha_i}(-1) \cdots h_{\alpha_{n/2+i-1}}(-1)$ for each $1 \leq i \leq n/2$, so that

$$\lambda(h_{e_i - e_{n/2+i}}(-1)) = \lambda(h_{\alpha_{n/2}}(-1)) = \lambda(h_{\alpha_{n/2}}(\zeta)^{(q-1)/2}) = (-1)^{(q-1)/2}.$$

Writing $\Lambda_\lambda := \Lambda(\lambda)$, this yields that

$$\Lambda_\lambda(c)^2 = \lambda(c^2) = \prod_{i=1}^{n/2} \lambda(h_{e_i - e_{n/2+i}}(-1)) = (-1)^{n(q-1)/4} = 1.$$

In particular, $\Lambda_\lambda(c) \in \{\pm 1\}$ is necessarily fixed by any $\sigma \in \mathcal{G}$, so $\delta_{\lambda,\sigma} = 1$ for all $\sigma \in \mathcal{G}$ by applying Lemma 4.3. ■

5. The generic algebra and the character $\eta^{(\sigma)}$

Here we recall some relevant details regarding the construction of generic algebras and their specializations. For more, we refer the reader to [HL83], though we remark that here we work over $\overline{\mathbb{Q}}$, rather than $\mathbb{C}$.

Keep the notation of Section 2.3. Let (L, λ) , with $\lambda \in \text{Irr}(L)$, be a cuspidal pair for G . Let $\mathbf{u} := \{u_\alpha \mid \alpha \in \Delta_\lambda\}$ be a set of indeterminates such that $u_\alpha = u_\beta$ if and only if α and β are $W(\lambda)$ -conjugate. For $m \in \mathbb{Q}$, we will write $\mathbf{u}^m$ for the set $\{u_\alpha^m \mid \alpha \in \Delta_\lambda\}$ and set $\sqrt{\mathbf{u}} := \mathbf{u}^{1/2}$. Let $\mathbb{A}_0 := \overline{\mathbb{Q}}[\mathbf{u}, \mathbf{u}^{-1}]$, let $\mathbb{K}$ be an algebraic closure of the quotient field of $\mathbb{A}_0$ containing $\mathbb{Q}(\sqrt{\mathbf{u}})$, and let $\mathbb{A}$ be the integral closure of $\mathbb{A}_0$ in $\mathbb{K}$.

We construct the so-called **generic algebra** $\mathcal{H}$ as the free $\mathbb{A}$ -module with basis

$$\{a_w \mid w \in W(\lambda)\},$$

together with the unique $\mathbb{A}$ -bilinear associative multiplication satisfying [HL83, Theorem 4.1]. Let $\mathbb{K}\mathcal{H}$ denote the $\mathbb{K}$ -algebra $\mathbb{K} \otimes_{\mathbb{A}} \mathcal{H}$, which is semisimple by [HL83, Corollary 4.6]. For a ring homomorphism $h: \mathbb{A} \rightarrow \overline{\mathbb{Q}}$, we obtain the so-called **specialized algebra**

$$\mathcal{H}^h := \overline{\mathbb{Q}} \otimes_{\mathbb{A}} \mathcal{H}$$

with basis $\{1 \otimes a_w \mid w \in W(\lambda)\}$ and structure constants obtained by applying h to the structure constants of $\mathcal{H}$. If h is the extension to $\mathbb{A}$ of a homomorphism $h_0: \mathbb{A}_0 \rightarrow \overline{\mathbb{Q}}$ and $\mathcal{H}^h$ is semisimple, then [HL83, Proposition 4.7] yields that there is a bijection $\varphi \mapsto \varphi^h$ between $\mathbb{K}$ -characters corresponding to simple $\mathbb{K}\mathcal{H}$ -modules and $\overline{\mathbb{Q}}$ -characters of simple $\mathcal{H}^h$ -modules, where φ^h is defined by

$$\varphi^h(1 \otimes a_w) = h(\varphi(a_w))$$

for each $w \in W(\lambda)$.

In particular, the homomorphisms $f_0, g_0: \mathbb{A}_0 \rightarrow \overline{\mathbb{Q}}$ defined by

$$f_0(u_\alpha) = p_{\alpha, \lambda} \quad \text{and} \quad g_0(u_\alpha) = 1,$$

respectively, yield specializations $\mathcal{H}^f$ and $\mathcal{H}^g$ satisfying

$$\mathcal{H}^f \cong \text{End}_{\overline{\mathbb{Q}}G}(\mathbf{R}_L^G(\lambda)) \quad \text{and} \quad \mathcal{H}^g \cong \overline{\mathbb{Q}}W(\lambda),$$

using [HL83, Lemma 4.2]. (We remark that the 2-cocycle in the original bijection may be taken to be trivial by [Gec93].)

For the remainder of this section, let $L = T$ be a maximally split torus of G , so that $\mathcal{E}(G, L, \lambda) = \mathcal{E}(G, T, \lambda)$ is a principal series. In this situation, we may follow the treatment of [HK80]. In particular, let $\mathcal{H}_0$ denote the subalgebra of $\mathcal{H}$ generated by $\{a_w \mid w \in R(\lambda)\}$, so that $\mathcal{H}_0$ is the generic Iwahori–Hecke algebra corresponding to the Coxeter group $R(\lambda)$.

We remark that since $R(\lambda)$ is a Weyl group, all of its characters are rational-valued, using for example [GP00, Theorems 5.3.8, 5.4.5, 5.5.6, and Corollary 5.6.4]. We will use this fact throughout the remainder of the paper, and our next observations aim to give an analogue to this fact for the characters of $\mathcal{H}_0$.

LEMMA 5.1: *With the notation above, let $\mathbb{K}_0$ be $\mathbb{Q}(\mathbf{u})$ in the case that $R(\lambda)$ contains no component of type G_2, E_7 , or E_8 , and let $\mathbb{K}_0 := \mathbb{Q}(\sqrt{\mathbf{u}})$ otherwise. Then $\mathbb{K}_0$ is a splitting field for $\mathcal{H}_0$.*

Proof. This follows directly from [GP00, Example 9.3.4 and Theorems 6.3.8 and 9.3.5]. ■

Now, using the discussion preceding [HK80, Theorem 3.7], we may identify the group algebra $\mathbb{K}C(\lambda)$ with the subalgebra of $\mathbb{K}\mathcal{H}$ generated by $\{a_d \mid d \in C(\lambda)\}$. In this situation, $C(\lambda)$ acts on $\mathbb{K}\mathcal{H}_0$ via

$$d(a_w) = a_{dwd^{-1}}$$

for $w \in R(\lambda)$ and $d \in C(\lambda)$, and on the irreducible $\mathbb{K}$ -characters ψ_0 of $\mathbb{K}\mathcal{H}_0$ via

$$\psi_0^d(a_w) = \psi_0(a_{dwd^{-1}}).$$

For an irreducible $\mathbb{K}$ -character ψ_0 of $\mathbb{K}\mathcal{H}_0$, let $\mathbb{K}C(\lambda)_{\psi_0}\mathcal{H}_0$ denote the subalgebra generated by $\{a_{dw} \mid d \in C(\lambda)_{\psi_0}; w \in R(\lambda)\}$. Then [HK80, Theorem 3.7] implies that ψ_0 extends to an irreducible character of $\mathbb{K}C(\lambda)_{\psi_0}\mathcal{H}_0$.

Further, [HK80, Lemma 3.12] and the discussion preceding it provide an analogue of Clifford theory and Gallagher's theorem for the characters of $\mathbb{K}\mathcal{H}$. Namely, if ψ is an extension of ψ_0 as above, then any extension of ψ_0 to $\mathbb{K}C(\lambda)_{\psi_0}\mathcal{H}_0$ is of the form $\beta\psi$ for $\beta \in \text{Irr}(C(\lambda)_{\psi_0})$, where

$$(\beta\psi)(a_{dw}) := \beta(d)\psi(a_{dw})$$

for $d \in C(\lambda)_{\psi_0}$ and $w \in R(\lambda)$. Further, any irreducible character of $\mathbb{K}\mathcal{H}$ is induced from such a character where, for ψ a character of $\mathbb{K}C(\lambda)_{\psi_0}\mathcal{H}_0$, the induced character $\text{Ind}_{C(\lambda)_{\psi_0}\mathcal{H}_0}^{\mathcal{H}}\psi$ is defined by

$$(1) \quad \text{Ind}_{C(\lambda)_{\psi_0}\mathcal{H}_0}^{\mathcal{H}}\psi(a_w) = \frac{1}{|C(\lambda)_{\psi_0}|} \sum \psi(a_{dwd^{-1}})$$

for $w \in W(\lambda)$, where the sum is taken over those $d \in C(\lambda)$ such that

$$a_{dwd^{-1}} \in C(\lambda)_{\psi_0}\mathcal{H}_0.$$

LEMMA 5.2: *Let ρ_0 be an irreducible $\mathbb{K}_0$ -representation of $\mathbb{K}_0\mathcal{H}_0$ affording the character ψ_0 , with notation as in Lemma 5.1. Then there is an extension ρ of ρ_0 to an irreducible representation of $\mathbb{K}C(\lambda)_{\psi_0}\mathcal{H}_0$ that can be afforded over $\mathbb{K}_0$.*

Proof. Let $\rho_0^{\mathbb{K}}$ denote ρ_0 viewed as a $\mathbb{K}$ -representation. In [HK80, Theorems 3.7 and 3.8], the $\mathbb{K}$ -character ψ_0 is shown to extend to a $\mathbb{K}$ -character ψ of $\mathbb{K}C(\lambda)_{\psi_0}\mathcal{H}_0$ by constructing a representation ρ extending $\rho_0^{\mathbb{K}}$. In particular, $\rho(d)$ for $d \in C(\lambda)_{\psi_0}$ is constructed using the equivalence properties of $\rho_0^{\mathbb{K}}$. Since $\psi_0 = \psi_0^d$ for every $d \in C(\lambda)_{\psi_0}$, we have $\rho_0^{\mathbb{K}}$ is equivalent over $\mathbb{K}$ to $(\rho_0^{\mathbb{K}})^d$, and hence ρ_0 and ρ_0^d are also equivalent over $\mathbb{K}_0$, using for example [CR62, Theorem 29.7]. But in the proof of [HK80, Theorem 3.8], this means that the construction for ρ can be done over $\mathbb{K}_0$, rather than $\mathbb{K}$, yielding the statement. ■

COROLLARY 5.3: *Keep the notation as above. Let ψ_0^f be an irreducible character of $\mathcal{H}_0^f$. Then there is an irreducible character of $\mathcal{H}^f$ lying over ψ_0^f whose values on the basis elements $1 \otimes a_w$ lie in $\mathbb{Q}(\sqrt{q})$. If $R(\lambda)$ contains no component of type G_2, E_7 , or E_8 , then there is further such a character with values in $\mathbb{Q}$.*

Similarly, if ψ_0^g is an irreducible character of $\mathcal{H}_0^g$, then there is an irreducible character of $\mathcal{H}^g$ lying over ψ_0^g with values in $\mathbb{Q}$.

Proof. Let ψ_0 be the corresponding character of $\mathbb{K}\mathcal{H}_0$ and let ψ be the $\mathbb{K}_0$ -character of $\mathbb{K}C(\lambda)_{\psi_0}\mathcal{H}_0$ extending ψ_0 guaranteed by Lemma 5.2. Then the statement follows immediately if $C(\lambda)_{\psi_0} = C(\lambda)$, since $g(u_\alpha) = 1$ and $f(u_\alpha) = p_{\alpha,\lambda}$ is a power of q for each $\alpha \in \Delta_\lambda$, using [HK80, Lemma 2.6]. Otherwise, we see by (1) that the character τ of $\mathcal{H}$ induced from ψ will also have values in $\mathbb{K}_0$ on the basis elements a_w , and hence τ^f and τ^g will also take its values in the stated fields. ■

Recall that we denote by $\mathfrak{f}$ the bijection $\mathfrak{f}: \text{Irr}(\text{End}_{\overline{\mathbb{Q}}_G}(\mathcal{R}_T^G(\lambda))) \rightarrow \text{Irr}(W(\lambda))$ induced by the specializations. In other words, $\mathfrak{f}: \text{Irr}(\mathcal{H}^f) \rightarrow \text{Irr}(\mathcal{H}^g)$ is given by $\mathfrak{f}(\tau^f) = \tau^g$ for τ an irreducible character of the generic algebra $\mathcal{H}$. Recall that for $\eta \in \text{Irr}(W(\lambda))$ and $\sigma \in \mathcal{G}$, if $\phi \in \text{Irr}(\text{End}_{\overline{\mathbb{Q}}_G}(\mathcal{R}_T^G(\lambda)))$ is such that $\mathfrak{f}(\phi) = \eta$, then we write $\eta^{(\sigma)}$ for the character of $W(\lambda)$ such that $\mathfrak{f}(\phi^\sigma) = \eta^{(\sigma)}$. Although one might hope that $\eta^{(\sigma)} = \eta^\sigma$ in general, this is known not to be the case. (For example there are situations in which every character of the Weyl group has values in $\mathbb{Q}$, but characters in the corresponding Iwahori–Hecke algebra take values in $\mathbb{Q}(\sqrt{q})$.) However, our main result of this section implies that this does hold in certain situations.

PROPOSITION 5.4: *Let $\lambda \in \text{Irr}(T)$ and $\eta \in \text{Irr}(W(\lambda))$. Let $\sigma \in \mathcal{G}$, and further assume that σ fixes $\sqrt{q}$ in the case that $R(\lambda)$ contains a component of type G_2, E_7 , or E_8 . Then $\eta^\sigma = \eta^{(\sigma)}$.*

Proof. Let $\eta \in \text{Irr}(W(\lambda))$ lie over $\eta_0 \in \text{Irr}(R(\lambda))$. Note that both η^σ and $\eta^{(\sigma)}$ also lie over η_0 , since the characters of the Weyl group $R(\lambda)$ are all rational-valued.

Write $\eta_0 = \psi_0^g$ for ψ_0 a character of $\mathcal{H}_0$, with the notation as above, and let ψ be the character of $C(\lambda)_{\psi_0} \mathcal{H}_0$ with values on basis elements in $\mathbb{K}_0$ ensured by Lemma 5.2. Let τ be the character induced from ψ to $\mathcal{H}$, so that τ also has values in $\mathbb{K}_0$ on the basis elements a_w , and hence τ^g and τ^f are fixed by σ . Note that, using Gallagher's theorem (and its analogue discussed before (1)) and [Isa06, Problem (5.3)], together with the fact that $C(\lambda)$ is either cyclic or Klein-four, we have $\eta = (\beta\tau)^g = \beta\tau^g$ for some $\beta \in \text{Irr}(C(\lambda))$, since g does not affect the values of β . Then $\eta^\sigma = \beta^\sigma \tau^g = (\beta^\sigma \tau)^g$. But also, $(\beta\tau)^f = \beta\tau^f$, and hence $\mathfrak{f}(\beta\tau^f) = \eta$, and we similarly have $(\beta\tau^f)^\sigma = \beta^\sigma \tau^f = (\beta^\sigma \tau)^f$. Hence

$$\mathfrak{f}((\beta\tau^f)^\sigma) = \eta^\sigma,$$

so $\eta^{(\sigma)} = \eta^\sigma$ by definition. $\blacksquare$

5.1. THE CASE $\ell = 2$. We next consider the situation of $\ell = 2$. In the following observation, for $R \triangleleft H$ we denote by $\text{Irr}_{2',H}(R) \subseteq \text{Irr}_{2'}(R)$ the set of $\theta \in \text{Irr}(R)$ such that $\theta \in \text{Irr}(R|\eta)$ for some $\eta \in \text{Irr}_{2'}(H)$.

LEMMA 5.5: *Let H be a finite group such that $H = R \rtimes C$ with C an elementary abelian 2-group.*

(a) *Let $\sigma \in \mathcal{G}$ and let $\eta \in \text{Irr}_{2'}(H|\theta)$, where $\theta \in \text{Irr}(R)$ is fixed by σ . Then*

$$\eta^\sigma = \eta.$$

(b) *If $C = C_2$ is cyclic, then there is a $\mathcal{G}$ -equivariant map*

$$\Xi: \text{Irr}_{2',H}(R) \rightarrow \text{Irr}_{2'}(H)$$

such that $\Xi(\theta)$ is an extension of θ to H for each $\theta \in \text{Irr}_{2',H}(R)$.

Proof. First note that for $\eta \in \text{Irr}_{2'}(H)$, it follows from Clifford theory that η restricts irreducibly to some $\theta \in \text{Irr}_{2'}(R)$ since H/R is a 2-group.

Now consider part (a). Note that the statement is true for linear characters, since if η is linear, we have

$$\eta^\sigma(rc) = \eta(r)^\sigma \eta(c)^\sigma = \theta(r)^\sigma \eta(c)^\sigma = \theta(r) \eta(c) = \eta(rc)$$

for any $r \in R$ and $c \in C$, since $c^2 = 1$ and $\theta^\sigma = \theta$ by assumption.

Now, the character $\det \theta$ (obtained by composing the determinant with a representation affording θ) is a linear character fixed by σ . Hence any extension μ of $\det \theta$ to H is fixed by σ by the previous paragraph. But [Isa06, Lemma 6.24] implies that there is a unique extension $\eta' \in \text{Irr}(H)$ of θ such that $\det \eta' = \mu$, and hence η' is fixed by σ by uniqueness. Then since $\eta = \eta' \beta$ for some $\beta \in \text{Irr}(C)$ by Gallagher's theorem, we see $\eta^\sigma = \eta$ as well, since β is a character of an elementary abelian 2-group. This completes part (a).

We now consider part (b). Note that again the statement holds for linear characters, since the extension is determined by its value on a generator c of C . In particular, we may define $\Xi_0: \text{Irr}_{2',H}(R/R') \rightarrow \text{Irr}_{2'}(H/R')$ to be an extension map such that $\Xi_0(\theta)$ is an extension to H satisfying $\Xi_0(\theta)(c) = 1$ for each linear $\theta \in \text{Irr}_{2',H}(R)$. Then

$$\Xi_0(\theta)^\sigma = \Xi_0(\theta^\sigma)$$

for each $\sigma \in \mathcal{G}$ and each linear $\theta \in \text{Irr}_{2',H}(R)$.

Now, for general $\theta \in \text{Irr}_{2',H}(R)$, we may define $\Xi(\theta)$ to be the unique extension to H such that $\det(\Xi(\theta)) = \Xi_0(\det \theta)$, guaranteed by [Isa06, Lemma 6.24]. Then since

$$\det(\Xi(\theta^\sigma)) = \Xi_0(\det \theta^\sigma) = \Xi_0(\det \theta)^\sigma = \det(\Xi(\theta))^\sigma = \det(\Xi(\theta)^\sigma),$$

this forces $\Xi(\theta)^\sigma = \Xi(\theta^\sigma)$ for each $\sigma \in \mathcal{G}$ and $\theta \in \text{Irr}_{2',H}(R)$. ■

This yields the following extensions of [SF19, Proposition 4.13 and Corollary 4.16].

COROLLARY 5.6: *Let $G = \mathbf{G}^F$ for a connected reductive algebraic group $\mathbf{G}$ and F a Frobenius morphism defining G over $\mathbb{F}_q$ with q odd. Let $\lambda \in \text{Irr}(T)$ such that $C(\lambda)$ is a (possibly trivial) elementary abelian 2-group. Let $\sigma \in \mathcal{G}$ and $\eta \in \text{Irr}_{2'}(W(\lambda))$. Then*

$$\eta^{(\sigma)} = \eta = \eta^\sigma.$$

Proof. Since $R(\lambda)$ is a Weyl group, every $\theta \in \text{Irr}(R(\lambda))$ is rational-valued, so the statement $\eta = \eta^\sigma$ follows directly from Lemma 5.5, since $W(\lambda) = R(\lambda) \rtimes C(\lambda)$. If $R(\lambda)$ contains no component $\mathbf{G}_2$, $\mathbf{E}_7$, or $\mathbf{E}_8$, Proposition 5.4 yields $\eta^\sigma = \eta^{(\sigma)}$. However, note that the exceptions to $\mathbb{K}_0 = \mathbb{Q}(\mathbf{u})$ in Lemma 5.1 in case $\mathbf{G}_2$, $\mathbf{E}_7$, and $\mathbf{E}_8$ have even degree, using [GP00, Theorem 8.3.1, Examples 9.3.2 and 9.3.4], and hence we may remove the assumption that σ fixes $\sqrt{q}$ in this case in Proposition 5.4. ■

6. The bijection for the principal series

In this section, we prove part (1) of Theorem A, with the exception of the case of $\mathrm{Sp}_{2n}(q)$ when $\ell = 2$ and $q \equiv 5 \pmod{8}$. This exception will be completed in Section 7.

As before, let $\mathbf{G}$ be simple of simply connected type and let $F: \mathbf{G} \rightarrow \mathbf{G}$ be a Frobenius morphism defining $G = \mathbf{G}^F$ over $\mathbb{F}_q$, and $G \triangleleft \tilde{G}$ as a result of a regular embedding.

We keep the notation from Section 2. In particular, we let $T = \mathbf{T}^F$ be a maximally split torus of G and let $N = N_{\mathbf{G}}(\mathbf{T})^F = N_G(\mathbf{T})$ and

$$\tilde{N} := N_{\tilde{G}}(\mathbf{T}).$$

Further, $D \leq \mathrm{Aut}(G)$ denotes the group of automorphisms generated by graph and field automorphisms. For $\lambda \in \mathrm{Irr}(T)$, recall that $\mathcal{E}(G, T, \lambda)$ denotes the set of irreducible constituents of $R_T^G(\lambda)$ and that Λ is the ND -equivariant extension map with respect to $T \triangleleft N$ discussed in Section 4.

According to [MS16, Theorem 5.2], the map

$$\Omega: \bigcup_{\lambda \in \mathrm{Irr}(T)} \mathcal{E}(G, T, \lambda) \rightarrow \mathrm{Irr}(N)$$

given by

$$(2) \quad R_T^G(\lambda)_\eta \mapsto \mathrm{Ind}_{N_\lambda}^N(\Lambda(\lambda)_\eta)$$

defines an $\tilde{N}D$ -equivariant bijection (see also [CS13, Theorem 4.5] and its proof when $Z(G) = 1$).

Using the results of the previous sections, we may show that Ω is equivariant with respect to the Galois automorphisms in $\mathcal{H}_\ell$ when ℓ is an odd prime dividing $q - 1$.

PROPOSITION 6.1: *Let Ω be as in (2) and assume $G = \mathbf{G}^F$ is as above, not of type A. Let ℓ be an odd prime dividing $q - 1$. Then*

$$\Omega(\chi^\sigma) = \Omega(\chi)^\sigma$$

for any $\chi \in \bigcup_{\lambda \in \mathrm{Irr}(T)} \mathcal{E}(G, T, \lambda)$ and $\sigma \in \mathcal{H}_\ell$.

Proof. Let $\chi = R_T^G(\lambda)_\eta$. Then by Theorem 2.2 and the definition (2) of Ω , we have

$$\Omega(\chi^\sigma) = \Omega(R_T^G(\lambda^\sigma)_{\gamma_{\lambda, \sigma}(\delta'_{\lambda, \sigma})^{-1}\eta(\sigma)}) = \mathrm{Ind}_{N_\lambda}^N(\Lambda(\lambda^\sigma)\gamma_{\lambda, \sigma}(\delta'_{\lambda, \sigma})^{-1}\eta(\sigma)).$$

But $\eta^{(\sigma)} = \eta^\sigma$ and $\gamma_{\lambda,\sigma} = 1$, by Proposition 5.4 and Corollary 3.3. Further,

$$\delta_{\lambda,\sigma} = \delta'_{\lambda,\sigma}$$

by Lemma 4.3. Hence, we have

$$\Omega(\chi^\sigma) = \text{Ind}_{N_\lambda}^N (\Lambda(\lambda^\sigma) \delta_{\lambda,\sigma}^{-1} \eta^\sigma).$$

On the other hand,

$$\Omega(\chi)^\sigma = (\text{Ind}_{N_\lambda}^N (\Lambda(\lambda) \eta))^\sigma = \text{Ind}_{N_\lambda}^N (\Lambda(\lambda)^\sigma \eta^\sigma) = \text{Ind}_{N_\lambda}^N (\Lambda(\lambda^\sigma) \delta_{\lambda,\sigma} \eta^\sigma).$$

The result now follows, since $\delta_{\lambda,\sigma} = \delta_{\lambda,\sigma}^{-1}$ by Corollary 4.5. $\blacksquare$

By [Mal07, Theorems 5.14 and 5.19], when $d_\ell(q) = 1$, we have N contains $N_G(Q)$ for some Sylow ℓ -subgroup Q of G (and hence N is also $\text{Aut}(G)_Q$ -stable by [CS13, Proposition 2.5]), with the following exceptions: $\ell = 3$ and $G = \text{SL}_3(q)$ with $q \equiv 4, 7 \pmod{9}$, $\ell = 3$ and $G = G_2(q)$ with $q \equiv 4, 7 \pmod{9}$, or $\ell = 2$ and $G = \text{Sp}_{2n}(q)$ with $q \equiv 5 \pmod{8}$. We now complete the proof of Theorem A(1), with these exceptions.

THEOREM 6.2: *Let q be a power of a prime p and let $\mathbf{G}$ be simple of simply connected type, not of type A. Let $F: \mathbf{G} \rightarrow \mathbf{G}$ be a Frobenius morphism such that $G := \mathbf{G}^F$ is defined over $\mathbb{F}_q$ and let ℓ be a prime such that $d_\ell(q) = 1$. Further assume that $q \equiv 1 \pmod{9}$ if $\ell = 3$ and $\mathbf{G}$ is type $\mathbf{G}_2$, and $q \equiv 1 \pmod{8}$ if $\ell = 2$ and $\mathbf{G}$ is of type C. Then the map defined by Equation (2) induces an $\tilde{N}D \times \mathcal{H}_\ell$ -equivariant bijection*

$$\Omega_\ell: \text{Irr}_{\ell'}(G) \rightarrow \text{Irr}_{\ell'}(N).$$

In particular, Condition $\ddagger$ holds for G , taking $M = N$.

Proof. From [Mal07, Proposition 7.3], every $\chi \in \text{Irr}_{\ell'}(G)$ lies in a Harish-Chandra series $\mathcal{E}(G, T, \lambda)$ for T a maximally split torus of G and by [Mal07, Proposition 7.8], the bijection defined by (2) indeed yields a bijection

$$\Omega_\ell: \text{Irr}_{\ell'}(G) \rightarrow \text{Irr}_{\ell'}(N)$$

such that corresponding characters lie over the same character of $Z(G)$. By [Mal07, Theorems 5.14 and 5.19] and [CS13, Proposition 2.5], N is $\text{Aut}(G)_Q$ -stable and contains $N_G(Q)$ for some Sylow ℓ -subgroup Q . If ℓ is odd, the result now follows from Proposition 6.1.

Hence we assume $\ell = 2$, $q \equiv 1 \pmod{4}$, and further $q \equiv 1 \pmod{8}$ if G is $\mathrm{Sp}_{2n}(q)$. Let $\sigma \in \mathcal{H}_2$. We must show that $\Omega(\chi)^\sigma = \Omega(\chi^\sigma)$ for each $\chi \in \mathrm{Irr}_{2'}(G)$. Note that Lemma 4.3 and Corollary 4.5 imply that $\delta'_{\lambda,\sigma} = \delta_{\lambda,\sigma} = \delta_{\lambda,\sigma}^{-1}$.

By Lemma 2.5, we have $\chi = R_T^G(\lambda)_\eta$ for some $\lambda \in \mathrm{Irr}(T)$ and $\eta \in \mathrm{Irr}_{2'}(W(\lambda))$. Then we have $\gamma_{\lambda,\sigma} = 1$, using [SFT20, Lemma 4.11] for $\mathrm{Sp}_{2n}(q)$ in the case being considered, and Proposition 3.4 otherwise. Further, we have $\eta^\sigma = \eta^{(\sigma)}$ by Proposition 5.4 or Corollary 5.6. Then with this, the statement follows by the same calculation as the proof of Proposition 6.1. ■

This proves Theorem A(1) with the exception of the case $G = \mathrm{Sp}_{2n}(q)$, $\ell = 2$, and $q \equiv 5 \pmod{8}$, which we consider in the next section.

7. $\mathrm{Sp}_{2n}(q)$ with $\ell = 2$

In this section, we prove Theorem A in the case $\ell = 2$ and $G = \mathrm{Sp}_{2n}(q)$. Throughout this section, let $\mathcal{H} := \mathcal{H}_2$ and let $G = \mathrm{Sp}_{2n}(q)$ with q odd. Since the case $q \equiv 1 \pmod{8}$ is completed in Theorem 6.2 above, we assume throughout that q is an odd power of an odd prime. In [Mal08, Section 4.4], Malle shows that G satisfies the inductive McKay conditions for $\ell = 2$ in this case. In particular, he constructs a proper $\mathrm{Aut}(G)_Q$ -stable subgroup $M \leq G$ containing $N_G(Q)$, where Q is a Sylow 2-subgroup of G , such that there is an $\mathrm{Aut}(G)_Q$ -equivariant bijection $\mathrm{Irr}_{2'}(G) \leftrightarrow \mathrm{Irr}_{2'}(M)$ satisfying that corresponding characters lie over the same character of $Z(G)$, in addition to the stronger properties required in the inductive McKay conditions. We will show that this bijection can be chosen to further be equivariant with respect to $\mathcal{H}$.

The following, found in [SFT20, Corollary 14.3], describes the character $\gamma_{\lambda,\sigma}\delta_{\lambda,\sigma}$ in the case of $\mathrm{Sp}_{2n}(q)$, and will be needed to show the $\mathcal{H}$ -equivariance of the bijection.

LEMMA 7.1 (Corollary 14.3 of [SFT20]): *Let $G = \mathrm{Sp}_{2n}(q)$ with q odd, and let $\lambda \in \mathrm{Irr}(T)$ be a nontrivial character of a maximally split torus T of G such that $\lambda^2 = 1$. Let $\ell = 2$ and $\sigma \in \mathcal{H}_2$. Then $\gamma_{\lambda,\sigma}\delta_{\lambda,\sigma} \in \mathrm{Irr}(C(\lambda)) \cong \{\pm 1\}$ satisfies:*

- *If $q \equiv \pm 1 \pmod{8}$, then $\gamma_{\lambda,\sigma}\delta_{\lambda,\sigma} = 1$.*
- *If $q \equiv \pm 3 \pmod{8}$, then $\gamma_{\lambda,\sigma}\delta_{\lambda,\sigma} = (-1)^r$, where r is the integer such that σ sends each odd root of unity to the 2^r power.*

In particular, $\gamma_{\lambda,\sigma}\delta_{\lambda,\sigma}$ is nontrivial if and only if $\sqrt{\omega q}$ is moved by σ , where $\omega = (-1)^{(q-1)/2}$.

We begin by recalling the group M and bijection from [Mal08, Section 4.4], which depends on whether or not n is a power of 2. Write

$$n = \sum_{j \in J} 2^j$$

for the 2-adic decomposition of n .

7.1. THE CASE $n = 2^j \geq 2$. Here suppose that $|J| = 1$, so that n is a power of 2. In this case, the group M from [Mal08, Section 4.4] is a wreath product $\mathrm{Sp}_n(q) \wr 2$. Write X for the base group $\mathrm{Sp}_n(q) \times \mathrm{Sp}_n(q)$, embedded naturally as 2-by-2 block matrices. The members of $\mathrm{Irr}_{2'}(M)$ are the extensions to M of the characters $\mu \otimes \mu$ of X , where $\mu \in \mathrm{Irr}_{2'}(\mathrm{Sp}_n(q))$. For $\mu \in \mathrm{Irr}_{2'}(\mathrm{Sp}_n(q))$, we let $\Xi(\mu)$ denote the corresponding extension of $\mu \otimes \mu$ to M such that $\Xi(\mu^\sigma) = \Xi(\mu)^\sigma$ for all $\sigma \in \mathcal{G}$, guaranteed by Lemma 5.5(b). Note that by Gallagher's theorem, the other extension of $\mu \otimes \mu$ is $\Xi(\mu)\beta$, where $\beta \in \mathrm{Irr}(M/X)$ has order 2, and

$$\Xi(\mu^\sigma)\beta = \Xi(\mu)^\sigma\beta = (\Xi(\mu)\beta)^\sigma$$

since β is necessarily fixed by $\mathcal{G}$.

LEMMA 7.2: *Let $G = \mathrm{Sp}_{2n}(q)$ for q an odd power of an odd prime and $n \geq 2$ a power of 2, and let $\ell = 2$. Then Condition $\nmid$ holds for G , taking $M = \mathrm{Sp}_n(q) \wr 2$ as above.*

Proof. Keeping the notation as above, the bijection $\mathrm{Irr}_{2'}(G) \rightarrow \mathrm{Irr}_{2'}(M)$ is constructed in [Mal08, Theorem 4.10] as any bijection sending odd-degree unipotent characters of G to the characters of the form $\Xi(\mu)$ and $\Xi(\mu)\beta$ for μ an odd-degree unipotent character of $\mathrm{Sp}_n(q)$, and sending non-unipotent characters to those of the form $\Xi(\mu)$ and $\Xi(\mu)\beta$ for μ non-unipotent. From [Lus02, Corollary 1.12], we know the unipotent characters of G and X are rational-valued, and hence fixed by every $\sigma \in \mathcal{G}$. Then the same is true for the characters of $\mathrm{Irr}_{2'}(M)$ lying above unipotent characters of X , by Lemma 5.5. In particular, the bijection is $\mathcal{H}$ -equivariant on unipotent characters.

Using [Mal08, Lemma 4.1 and Proposition 4.6] and Theorem 2.4, we see that for any positive integer k , the non-unipotent characters of $\mathrm{Sp}_{2k}(q)$ of odd degree lie in a single Harish-Chandra series, namely a principal series $\mathcal{E}(\mathrm{Sp}_{2k}(q), T, \lambda)$ with $\lambda^2 = 1$. By Proposition 5.4, $\eta = \eta^{(\sigma)} = \eta^\sigma$ for all $\eta \in \mathrm{Irr}(W(\lambda))$ here. Hence the action of $\mathcal{H}$ on $\mathcal{E}(\mathrm{Sp}_{2k}(q), T, \lambda)$ is determined by the character $\gamma_{\lambda, \sigma} \delta_{\lambda, \sigma}$, by Theorem 2.2.

From Lemma 7.1, we note that $\gamma_{\lambda,\sigma}\delta_{\lambda,\sigma}$ is independent of k , so that $\mathcal{H}$ permutes pairs of non-unipotent members of $\text{Irr}_{2'}(G)$ exactly when it permutes pairs of non-unipotent members of $\text{Irr}_{2'}(X)$ (namely, when $\sqrt{\omega q}^\sigma \neq \sqrt{\omega q}$), and in an analogous way. Then the bijection $\text{Irr}_{2'}(G) \rightarrow \text{Irr}_{2'}(M)$ may be chosen to be $\mathcal{H}$ -equivariant by ensuring that each pair $\{\mathbf{R}_T^G(\lambda)_\eta, \mathbf{R}_T^G(\lambda)_{(-1_{C(\lambda)})\eta}\}$ is mapped to pairs of the form

$$\{\Xi(\mathbf{R}_{T'}^{\text{Sp}_n(q)}(\lambda')_{\eta'}), \Xi(\mathbf{R}_{T'}^{\text{Sp}_n(q)}(\lambda')_{(-1_{C(\lambda_1)})\eta'})\}$$

or of the form

$$\{\Xi(\mathbf{R}_{T'}^{\text{Sp}_n(q)}(\lambda')_{\eta'})\beta, \Xi(\mathbf{R}_{T'}^{\text{Sp}_n(q)}(\lambda')_{(-1_{C(\lambda_1)})\eta'})\beta\}$$

for $\eta \in \text{Irr}_{2'}(W(\lambda))$ and $\eta' \in \text{Irr}_{2'}(W(\lambda'))$, where $\mathcal{E}(G, T, \lambda)$ and $\mathcal{E}(\text{Sp}_n(q), T', \lambda')$ are the unique Harish-Chandra series of G and $\text{Sp}_n(q)$, respectively, containing non-unipotent odd-degree characters. This proves the statement. ■

7.2. THE CASE $n \neq 2^j$. When $|J| \geq 2$, let $m = \max_{j \in J} \{2^j\}$. In this case, Malle defines the subgroup M in [Mal08, Theorem 4.11] as

$$\text{Sp}_{2(n-m)}(q) \times \text{Sp}_{2m}(q),$$

naturally embedded as block matrices. Writing

$$M_1 := \text{Sp}_{2(n-m)}(q) \quad \text{and} \quad M_2 := \text{Sp}_{2m}(q),$$

we see using [Mal08, Propositions 4.2, 4.5] that the members of $\text{Irr}_{2'}(X)$ for any $X \in \{G, M_1, M_2\}$ are found in Lusztig series $\mathcal{E}(X, t)$ indexed by semisimple classes corresponding to the identity $t = 1$ and classes of 2-central involutions t of the dual group X^* , and that these semisimple classes are then in bijection with subsets of J , $J \setminus \{m\}$, and $\{m\}$, respectively. Here the empty subset corresponds to the series of unipotent characters. Write $\mathcal{E}_{2'}(X, t)$ for the set $\mathcal{E}(X, t) \cap \text{Irr}_{2'}(X)$.

PROPOSITION 7.3: *Let $G = \text{Sp}_{2n}(q)$ for q an odd power of an odd prime and $n \geq 2$, and let $\ell = 2$. Then Condition $\ddagger$ holds for G , taking M to be the subgroup of G described above.*

Proof. By Lemma 7.2, we may assume n is not a power of 2 and keep the notation above. Let $s_1 \in M_1^*$ and $s_2 \in M_2^*$ correspond to subsets I_1 and I_2 of $J \setminus \{m\}$ and $\{m\}$, respectively. Then the bijection from [Mal08, Theorem 4.11] sends $\mathcal{E}_{2'}(M_1, s_1) \otimes \mathcal{E}_{2'}(M_2, s_2) \subset \text{Irr}_{2'}(M)$ to the set $\mathcal{E}_{2'}(G, s) \subset \text{Irr}_{2'}(G)$,

where s corresponds to the subset $I_1 \cup I_2$ of J . This bijection naturally sends products of unipotent characters to a unipotent character, and hence is $\mathcal{H}$ -equivariant on unipotent characters since they are again rational-valued.

Let $\chi \in \text{Irr}_{2'}(X)$ be a non-unipotent character in a series parametrized by a semisimple element corresponding to the subset I , where

$$X := \text{Sp}_{2k}(q) \in \{G, M_1, M_2\}.$$

By Theorem 2.4, either $\chi \in \mathcal{E}(X, T, \lambda)$ where T is a maximally split torus of X and $\lambda \in \text{Irr}(T)$ satisfies $\lambda^2 = 1$, or $q \equiv 3 \pmod{4}$, $0 \in I$, and $\chi \in \mathcal{E}(X, L, \lambda)$, where $L \cong \text{Sp}_2(q) \times T_0$ with T_0 a maximally split torus of $\text{Sp}_{2(k-1)}(q)$. In the latter case, we further have

$$\lambda = \psi \otimes \lambda_0,$$

where $\lambda_0 \in \text{Irr}(T_0)$ satisfies $\lambda_0^2 = 1$ and ψ is one of the two characters ψ_1 or ψ_2 of $\text{Sp}_2(q)$ of degree $\frac{q-1}{2}$.

Let $\omega \in \{\pm 1\}$ be such that $q \equiv \omega \pmod{4}$. Note that $\sigma \in \mathcal{H}$ interchanges ψ_1 and ψ_2 if and only if $\sqrt{\omega q}^\sigma \neq \sqrt{\omega q}$. Further, [SFT20, Theorem B] and its proof yields that more generally, the non-unipotent members of $\text{Irr}_{2'}(X)$ are fixed if $\sqrt{\omega q}^\sigma = \sqrt{\omega q}$ and are permuted by σ analogously in pairs via

$$(\text{R}_T^G(\lambda)_\eta)^\sigma = \text{R}_T^G(\lambda)_{(-1_{C(\lambda)})\eta} \quad \text{and} \quad (\text{R}_L^G(\psi_1 \otimes \lambda_0)_\eta)^\sigma = \text{R}_L^G(\psi_2 \otimes \lambda_0)_{(-1_{C(\lambda_0)})\eta},$$

if $\sqrt{\omega q}^\sigma \neq \sqrt{\omega q}$.

Recall that non-unipotent characters in $\text{Irr}_{2'}(G)$ are mapped to

$$\chi_1 \otimes \chi_2 \in \text{Irr}_{2'}(M_1 \times M_2)$$

such that at least one of χ_1 or χ_2 is also non-unipotent. Then this yields that the bijection $\text{Irr}_{2'}(G) \rightarrow \text{Irr}_2(M)$ may be fixed to be $\mathcal{H}$ -equivariant by sending pairs $\{\text{R}_T^G(\lambda)_\eta, \text{R}_T^G(\lambda)_{(-1_{C(\lambda)})\eta}\}$ or $\{\text{R}_L^G(\psi_1 \otimes \lambda_0)_\eta, \text{R}_L^G(\psi_2 \otimes \lambda_0)_{(-1_{C(\lambda_0)})\eta}\}$ to pairs in $\text{Irr}_{2'}(M)$ whose non-unipotent components are analogous pairs in $\text{Irr}_{2'}(M_1)$ and $\text{Irr}_{2'}(M_2)$ and whose unipotent components (if applicable) are equal. Then the characters in these pairs are necessarily interchanged by $\sigma \in \mathcal{H}$ if and only if the same is true for the pairs in $\text{Irr}_{2'}(G)$ mapped to them. ■

This completes the proof of Theorem A(1), as well as the case of $G = \text{Sp}_{2n}(q)$ in Theorem A(2).

8. Remaining cases for $\ell = 2$ and $d_2(q) = 2$

In this section, we complete the proof of Theorem A(2). In particular, we will consider the cases $G = \mathbf{G}_2(q)$, ${}^3\mathbf{D}_4(q)$, $\mathbf{F}_4(q)$, $\mathbf{E}_6^\epsilon(q)_{sc}$, $\mathbf{E}_7(q)_{sc}$, $\mathbf{E}_8(q)$, or $\mathbf{B}_n(q)_{sc}$ with $n \geq 3$, especially when $q \equiv 3 \pmod{4}$. Here we write $\mathbf{E}_6^\epsilon$ for $\epsilon \in \{\pm 1\}$ to denote $\mathbf{E}_6$ in the case $\epsilon = +1$ and ${}^2\mathbf{E}_6$ in the case $\epsilon = -1$.

Recall that we have $\mathbf{T} = \mathbf{C}_{\mathbf{G}}(\mathbf{S})$ for some Sylow 2-torus $\mathbf{S}$ of $(\mathbf{G}, vF)$ and $N_1 := \mathbf{N}_{\mathbf{G}}(\mathbf{S})^{vF} = \mathbf{N}^{vF}$. By [Mal07, Theorem 7.8], if $q \equiv 3 \pmod{4}$, then N_1 contains $\mathbf{N}_G(Q)$ for a Sylow 2-subgroup Q of G (and hence N_1 is also $\text{Aut}(G)_Q$ -stable by [CS13, Proposition 2.5]), and there is a bijection

$$\Omega_1: \text{Irr}_{2'}(G) \rightarrow \text{Irr}_{2'}(N_1),$$

which is moreover $\text{Aut}(G)_Q$ -equivariant by [CS13, Proposition 4.5] and [MS16, Theorem 6.3] combined with [Spä12, Theorem 2.12], and where corresponding characters lie over the same character of $Z(G)$. Hence it suffices to show that this bijection can be chosen to further be $\mathcal{H}_2$ -equivariant. In fact, for the listed groups aside from $\mathbf{E}_6^\epsilon(q)$, we will show that every member of $\text{Irr}_{2'}(G)$ and of $\text{Irr}_{2'}(N_1)$ is rational-valued.

PROPOSITION 8.1: *Let q be odd and let $G = \mathbf{G}_2(q)$, ${}^3\mathbf{D}_4(q)$, $\mathbf{F}_4(q)$, $\mathbf{E}_7(q)_{sc}$, $\mathbf{E}_8(q)$, or $\mathbf{B}_n(q)_{sc}$ with $n \geq 3$. Then every member of $\text{Irr}_{2'}(G)$ is rational-valued. Further, keeping the notation above, we have every member of $\text{Irr}_{2'}(N_1)$ is rational-valued when $q \equiv 3 \pmod{4}$.*

Before we prove this statement, we introduce a little more notation for the situation $q \equiv 3 \pmod{4}$. Let $(\mathbf{G}^*, (vF)^*)$ be dual to $(\mathbf{G}, vF)$. By [Mal07, Lemma 3.3], the torus $\mathbf{T}^*$ dual to $\mathbf{T}$ can be identified with $\mathbf{C}_{\mathbf{G}^*}(\mathbf{S}^*)$ and we write $\mathbf{N}^* := \mathbf{N}_{\mathbf{G}^*}(\mathbf{S}^*)$, $N_1^* := (\mathbf{N}^*)^{(vF)^*}$, and $T_1^* := (\mathbf{T}^*)^{(vF)^*}$.

By the proof of [Mal07, Theorem 7.8], we see that both $\text{Irr}_{2'}(\mathbf{G}^{vF})$ (and hence $\text{Irr}_{2'}(G)$ since $G \cong \mathbf{G}^{vF}$) and $\text{Irr}_{2'}(N_1)$ are in bijection with pairs (λ_1, η_1) or (s, η_1) , where $\lambda_1 \in \text{Irr}(T_1)$ satisfies $[N_1 : (N_1)_{\lambda_1}]$ is odd and $\lambda_1 \in \mathcal{E}(T_1, s)$ with $s \in T_1^*$ a semisimple element centralizing a Sylow 2-subgroup of N_1^* (and hence of $\mathbf{G}^{*(vF)^*}$), up to N_1^* -conjugation; and $\eta_1 \in \text{Irr}_{2'}(W_1(\lambda_1))$. Here we define

$$W_1(\lambda_1) := (N_1)_{\lambda_1}/T_1,$$

which by [Mal07, Proposition 7.7] is isomorphic to

$$W_{N_1^*}(s) := C_{N_1^*}(s)/T_1^*.$$

Now, the member of $\text{Irr}_{2'}(N_1)$ corresponding to (λ_1, η_1) is of the form

$$\text{Ind}_{(N_1)_{\lambda_1}}^{N_1}(\Lambda_1(\lambda_1)\eta_1),$$

where Λ_1 is an extension map with respect to $T_1 \triangleleft N_1$. Note that $W_{N_1^*}(s)$ can also be thought of as the fixed points under $(vF)^*$ of the Weyl group of the possibly disconnected group $C_{\mathbf{G}^*}(s)$, which in turn is the set of $w \in (\mathbf{W}^*)^{(vF)^*}$ such that $s^w = s$. Write $W_1(s)$ for this group, so that $W_1(s) \cong W_1(\lambda_1)$. Further, write $W_1^\circ(s)$ for the fixed points under $(vF)^*$ of the Weyl group of the connected component $C_{\mathbf{G}^*}(s)^\circ$. The group $W_1^\circ(s)$ is then a true Weyl group (in the sense of (B, N) pairs and Coxeter groups) and is the subgroup of $W_1(s)$ generated by the simple reflections corresponding to $\alpha^* \in \Phi^*$ such that $\alpha^*(s) = 1$, where we write Φ^* for the root system of $\mathbf{G}^*$. (See, e.g. [DM91, Remark 2.4].) Then by the isomorphism $\text{Irr}(T_1) \cong T_1^*$ given by [Car93, Propositions 4.2.3, 4.4.1], it follows that

$$(3) \quad W_1^\circ(s) \cong R_1(\lambda_1) \quad \text{and} \quad W_1(s)/W_1^\circ(s) \cong W_1(\lambda_1)/R_1(\lambda_1),$$

where we define $R_1(\lambda_1)$ to be the reflection group generated by the simple reflections s_α for $\alpha \in \Phi$ such that $s_\alpha \in W_1(\lambda_1)$ and $T_1 \cap \langle X_\alpha, X_{-\alpha} \rangle$ is in the kernel of λ_1 . Here X_α denotes the root subgroup of G associated to α .

Proof of Proposition 8.1. By [MS16] (see Theorem 2.4 and Lemma 2.5), we know every member of $\text{Irr}_{2'}(G)$ is of the form

$$\chi = R_T^G(\lambda)_\eta \in \mathcal{E}(G, T, \lambda),$$

where $[W : W(\lambda)]$ and $\eta(1)$ are both odd. Note that using Lemma 2.6, we see $s^2 = 1$, and it follows that $\lambda^2 = 1$ and hence $\lambda^\sigma = \lambda$. Now, Proposition 3.4 yields that $\gamma_{\lambda, \sigma}$ is trivial for any $\sigma \in \mathcal{G}$. Further, $\eta = \eta^{(\sigma)} = \eta^\sigma$ for any $\eta \in \text{Irr}_{2'}(W(\lambda))$ and $\sigma \in \mathcal{G}$, using Corollary 5.6. Further, Corollary 4.4 and Proposition 4.6 yields that $\delta_{\lambda, \sigma} = 1$ for any $\sigma \in \mathcal{G}$. From this, we now see that $\chi^\sigma = \chi$ for each $\chi \in \text{Irr}_{2'}(G)$ and $\sigma \in \mathcal{G}$, using Theorem 2.2, and we turn our attention to $\text{Irr}_{2'}(N_1)$.

Let $q \equiv 3 \pmod{4}$. As in the preceding discussion, $\chi \in \text{Irr}_{2'}(G)$ and $\Omega_1(\chi)$ are also parametrized by a pair (λ_1, η_1) with the properties described above. Let $\sigma \in \mathcal{G}$ and write $\text{Ind}_{(N_1)_{\lambda_1}}^{N_1}(\Lambda_1(\lambda_1)\eta_1)$ for $\Omega_1(\chi)$, so that

$$\Omega_1(\chi)^\sigma = \text{Ind}_{(N_1)_{\lambda_1}}^{N_1}(\Lambda_1(\lambda_1)^\sigma \eta_1^\sigma).$$

Note that Lemma 5.5 gives $\eta_1^\sigma = \eta_1$, since $R_1(\lambda_1)$ is a Weyl group by (3) and hence has rational-valued characters. So, noting that $\lambda_1^\sigma = \lambda_1$ since $s^2 = 1$, it now suffices to show $\delta = 1$, where δ is the character of $W_1(\lambda_1)$ such that

$$\Lambda_1(\lambda_1)^\sigma = \Lambda_1(\lambda_1^\sigma)\delta$$

guaranteed by Gallagher's theorem. In the case $G = \mathbf{G}_2(q)$, ${}^3\mathbf{D}_4(q)$, $\mathbf{F}_4(q)$, or $\mathbf{E}_8(q)$, this is accomplished by Corollary 4.4, completing the proof in these cases.

We may therefore assume G is type $\mathbf{E}_7$ or $\mathbf{B}_n$. Now, δ is trivial on $R_1(\lambda_1)$ by the same proof as [SFT20, Lemma 3.13]. Then taking into consideration (3), we may argue analogously to Proposition 4.6, replacing ζ with a generator of the cyclic group of size $q + 1$ in $\mathbb{F}_{q^2}^\times$ in the case $\mathbf{B}_n$, to obtain $\delta = 1$. ■

PROPOSITION 8.2: *Let $q \equiv 3 \pmod{4}$ and let $G = \mathbf{E}_6^\epsilon(q)_{sc}$ and keep the notation from before. Then the $\text{Aut}(G)_Q$ -equivariant bijection*

$$\Omega_1 : \text{Irr}_{2'}(G) \rightarrow \text{Irr}_{2'}(N_1)$$

is also $\mathcal{H}_2$ -equivariant.

Proof. Throughout, we identify G with $\mathbf{G}^{vF}$. By analyzing the possible centralizer structures $C_{G^*}(s)$ of semisimple elements s of G^* (see also [NT16a, Lemma 4.13]), we see that the only possibilities for s yielding $\chi \in \mathcal{E}(G, s)$ with odd degree are $s = 1$ and those with $C_{G^*}(s)$ of type $\mathbf{D}_5^\epsilon(q) \times (q - \epsilon)$. By [DM91, Remark 13.24], odd-degree characters in $\mathcal{E}(G, s)$ are in bijection with odd-degree unipotent characters of $C_{G^*}(s)$ in such a way that if χ corresponds to the unipotent character ψ , we have $\chi(1) = [G^* : C_{G^*}(s)]_{p'} \psi(1)$.

The unipotent characters of odd degree of G are all fixed by $\mathcal{G}$, using [SF19, Proposition 4.4]. In the non-unipotent cases, $C_{G^*}(s)$ is connected, and the eight odd-degree unipotent characters of $C_{G^*}(s)$ have distinct degrees. Hence, the image of $\chi \in \mathcal{E}(G, s) \cap \text{Irr}_{2'}(G)$ under $\sigma \in \mathcal{H}_2$ is completely determined by the image $\mathcal{E}(G, s)^\sigma$ of $\mathcal{E}(G, s)$ under σ . Further, [SFT18, Lemma 3.4] implies that $\mathcal{E}(G, s)^\sigma = \mathcal{E}(G, s^\sigma)$, where if $\sigma \in \mathcal{H}_2$ sends odd roots of unity to the 2^r power and 2-power roots of unity to the power b , we define $s^\sigma := s_2^{2^r} s_2^b$, where $s = s_{2'} s_2 = s_2 s_{2'}$ with $|s_{2'}|$ odd and $|s_2|$ a power of 2.

Then if $\chi \in \text{Irr}_{2'}(G)$ corresponds to the pair (s, η_1) or (λ_1, η_1) , we see that for $\sigma \in \mathcal{H}_2$, χ^σ corresponds to the pair $(s^\sigma, \eta_1) = (s^\sigma, \eta_1^\sigma)$. (Note that since $C_{G^*}(s)$ is connected, $W_1(\lambda_1) \cong W_1(s)$ is a Weyl group in this case, and hence η_1 is rational-valued.)

On the other hand, let $\Omega_1(\chi) = \text{Ind}_{(N_1)_{\lambda_1}}^{N_1}(\Lambda_1(\lambda_1)\eta_1)$ correspond to (λ_1, η_1) or (s, η_1) and let $\sigma \in \mathcal{H}_2$. As before, we have $\eta_1^\sigma = \eta_1$. Further, by Corollary 4.4, Λ_1 is $\mathcal{G}$ -equivariant, and certainly $(N_1)_{\lambda_1} = (N_1)_{\lambda_1^\sigma}$, hence

$$\Omega(\chi)^\sigma = \text{Ind}_{(N_1)_{\lambda_1}}^{N_1}(\Lambda_1(\lambda_1)^\sigma \eta_1^\sigma) = \text{Ind}_{(N_1)_{\lambda_1}}^{N_1}(\Lambda_1(\lambda_1^\sigma) \eta_1).$$

Further, note that $\lambda_1^\sigma \in \mathcal{E}(T_1, s)^\sigma = \mathcal{E}(T_1, s^\sigma)$ is also completely determined by s^σ . Then again here we have $\Omega_1(\chi)^\sigma$ corresponds to the pair (s^σ, η_1) , yielding that $\Omega_1(\chi)^\sigma = \Omega_1(\chi^\sigma)$, as desired. ■

Together, Propositions 8.1 and 8.2 yield the desired result:

COROLLARY 8.3: *Theorem A(2) holds for $G = \mathbf{G}_2(q)$, ${}^3\mathbf{D}_4(q)$, $\mathbf{F}_4(q)$, $\mathbf{E}_6^\epsilon(q)_{sc}$, $\mathbf{E}_7(q)_{sc}$, $\mathbf{E}_8(q)$, or $\mathbf{B}_n(q)_{sc}$ with $n \geq 3$, taking $M = N_1$.*

We end by discussing briefly the Sylow 2-subgroups of the groups under consideration here to arrive at Theorem B. By the Corollary of the main theorems of [Kon05], the Sylow 2-subgroups of the simple groups listed in Theorem B are self-normalizing. Further, by [NT16b], the abelianization $P/[P, P]$ of the Sylow 2-subgroups in these cases are elementary abelian.

PROPOSITION 8.4 ([NT16b]): *Let q be a power of an odd prime and let S be a simple group $\mathbf{G}_2(q)$, ${}^3\mathbf{D}_4(q)$, $\mathbf{F}_4(q)$, $\mathbf{E}_7(q)$, $\mathbf{E}_8(q)$, $\mathbf{C}_n(q)$ with $n \geq 2$, or $\mathbf{B}_n(q)$ with $n \geq 3$. Let P be a Sylow 2-subgroup of S . Then $P/[P, P]$ is elementary abelian:*

Proof. This is directly from Propositions 3.5, 3.7, and 4.1 of [NT16b]. ■

Proof of Theorem B. Recall that the McKay conjecture holds for $\ell = 2$ by [MS16], so it suffices to show that there is a $\mathcal{H}_2$ -equivariant bijection between $\text{Irr}_{2'}(S)$ and $\text{Irr}_{2'}(N_S(P))$ where P is a Sylow 2-subgroup of S . By combining Proposition 8.4 with Proposition 8.1 and the fact that $P = N_S(P)$ in the stated cases for which S is simple, we immediately obtain the result, except possibly for the exceptional covering groups and $\mathbf{C}_n(q) = \text{PSp}_{2n}(q)$. However, in the latter case, [SFT20, Lemma 4.10 and Theorem B], together with Lemma 2.6, implies that every member of $\text{Irr}_{2'}(S)$ is again fixed by $\mathcal{H}_2$. For the groups $3.\mathbf{B}_3(3)$ and $3.\mathbf{G}_2(3)$, we have $N_S(P) = P \times C_3$ and hence $\text{Irr}_{2'}(N_S(P))$ can be viewed as the irreducible characters of $P/[P, P] \times C_3$. Using GAP, we see the members of $\text{Irr}_{2'}(S)$ without $Z(S)$ in the center have field of values $\mathbb{Q}(e^{2\pi i/3})$ and that a $\mathcal{H}_2$ -equivariant bijection may again be chosen in this case. ■

ACKNOWLEDGEMENTS. The author began this work during her stay at the Mathematical Sciences Research Institute in Berkeley, California, as part of the Spring 2018 MSRI semester program “Group Representation Theory and Applications”. She thanks the MSRI and the organizers of the program for making her stay possible and providing a collaborative and productive work environment. She would also like to thank G. Navarro, B. Späth, and C. Vallejo for helpful discussions and an early preprint of their reduction theorem. She further thanks C. Vallejo and G. Malle for their comments on an earlier draft of this work. Finally, she thanks the anonymous referee for their helpful remarks.

References

- [Bon06] C. Bonnafé, *Sur les caractères des groupes réductifs finis à centre non connexe: applications aux groupes spéciaux linéaires et unitaires*, Astérisque **306** (2006).
- [Car93] R. W. Carter, *Finite Groups of Lie Type*. Wiley Classics Library, John Wiley & Sons, Chichester, 1993.
- [CR62] C. W. Curtis and I. Reiner, *Representation Theory of Finite Groups and Associative Algebras*, Pure and Applied Mathematics, Vol. 11, Interscience, New York–London, 1962.
- [CS13] M. Cabanes and B. Späth, *Equivariance and extendibility in finite reductive groups with connected center*, Mathematische Zeitschrift **275** (2013), 689–713.
- [DM87] D. I. Deriziotis and G. O. Michler, *Character table and blocks of finite simple triality groups ${}^3D_4(q)$* , Transactions of the American Mathematical Society **303** (1987), 39–70.
- [DM91] F. Digne and J. Michel, *Representations of Finite Groups of Lie Type*, London Mathematical Society Student Texts, Vol. 21, Cambridge University Press, Cambridge, 1991.
- [Gec93] M. Geck, *A note on Harish-Chandra induction*, Manuscripta Mathematica **80** (1993), 393–401.
- [GLS98] D. Gorenstein, R. Lyons and R. Solomon, *The Classification of the Finite Simple Groups. Number 3. Part I. Chapter A*, Mathematical Surveys and Monographs, Vol. 40, American Mathematical Society, Providence, RI, 1998.
- [GM20] M. Geck and G. Malle, *The Character Theory of Finite Groups of Lie Type*, Cambridge Studies in Advanced Mathematics, Vol. 187, Cambridge University Press, Cambridge, 2020.
- [GP00] M. Geck and G. Pfeiffer, *Characters of Finite Coxeter Groups and Iwahori–Hecke Algebras*, London Mathematical Society Monographs, Vol. 21, The Clarendon Press, Oxford University Press, New York, 2000.
- [HK80] R. B. Howlett and R. W. Kilmoyer, *Principal series representations of finite groups with split BN pairs*, Communications in Algebra **8** (1980), 543–583.
- [HL80] R. B. Howlett and G. I. Lehrer, *Induced cuspidal representations and generalised Hecke rings*, Inventiones Mathematicae **58** (1980), 37–64.

- [HL83] R. B. Howlett and G. I. Lehrer, *Representations of generic algebras and finite groups of Lie type*, Transactions of the American Mathematical Society **280** (1983), 753–779.
- [IMN07] I. M. Isaacs, G. Malle and G. Navarro, *A reduction theorem for the McKay conjecture*, Inventiones Mathematicae **170** (2007), 33–101.
- [Isa06] I. M. Isaacs, *Character Theory of Finite Groups*, AMS Chelsea Publishing, Providence, RI, 2006.
- [Kon05] A. S. Kondrat'ev, *Normalizers of Sylow 2-subgroups in finite simple groups*, Matematicheskie Zametki **78** (2005), 368–376.
- [Lüb07] F. Lübeck, *Data for finite groups of lie type and related algebraic groups*, <http://www.math.rwth-aachen.de/~Frank.Luebeck/chev/index.html>,
- [Lus84] G. Lusztig, *Characters of Reductive Groups over a Finite Field*, Annals of Mathematics Studies, Vol. 107, Princeton University Press, Princeton, NJ, 1984.
- [Lus02] G. Lusztig, *Rationality properties of unipotent representations*, Journal of Algebra **258** (2002), 1–22.
- [Mal07] G. Malle, *Height 0 characters of finite groups of Lie type*, Representation Theory **11** (2007), 192–220.
- [Mal08] G. Malle, *Extensions of unipotent characters and the inductive McKay condition*, Journal of Algebra **320** (2008), 2963–2980.
- [MS16] G. Malle and B. Späth, *Characters of odd degree*, Annals of Mathematics **184** (2016), 869–908.
- [Nav04] G. Navarro, *The McKay conjecture and Galois automorphisms*, Annals of Mathematics **160** (2004), 1129–1140.
- [NSV20] G. Navarro, B. Späth and C. Vallejo, *A reduction theorem for the Galois–McKay conjecture*, Transactions of the American Mathematical Society **373** (2020), 6157–6183.
- [NT16a] G. Navarro and P. H. Tiep, *Irreducible representations of odd degree*, Mathematische Annalen **365** (2016), 1155–1185.
- [NT16b] G. Navarro and P. H. Tiep, *Real groups and sylow 2-subgroups*, Advances in Mathematics **299** (2016), 331–360.
- [Ruh17] L. Ruhstorfer, *The Navarro refinement of the McKay conjecture for finite groups of Lie type in defining characteristic*, Journal of Algebra **582** (2021), 177–205.
- [SF16] A. A. Schaeffer Fry, *Odd-degree characters and self-normalizing Sylow subgroups: A reduction to simple groups*, Communications in Algebra **44** (2016), 1882–1904.
- [SF19] A. A. Schaeffer Fry, *Galois automorphisms on Harish-Chandra series and Navarro's self-normalizing Sylow 2-subgroup conjecture*, Transactions of the American Mathematical Society **372** (2019), 457–483.
- [SFT18] A. A. Schaeffer Fry and J. Taylor, *On self-normalising Sylow 2-subgroups in type A*, Journal of Lie Theory **28** (2018), 139–168.
- [SFT20] A. A. Schaeffer Fry and J. Taylor, *Galois automorphisms and classical groups*, <https://arxiv.org/abs/2005.14088>.
- [Spä09] B. Späth, *The McKay conjecture for exceptional groups and odd primes*, Mathematische Zeitschrift **261** (2009), 571–595.

- [Spä10] B. Späth, *Sylow d -tori of classical groups and the McKay conjecture, I*, Journal of Algebra **323** (2010), 2469–2493.
- [Spä12] B. Späth, *Inductive McKay condition in defining characteristic*, Bulletin of the London Mathematical Society **44** (2012), 426–438.
- [Tur08] A. Turull, *Strengthening the McKay conjecture to include local fields and local Schur indices*, Journal of Algebra **319** (2008), 4853–4868.