

Clockwork mechanism to remove superradiance limits

Anubhav Mathur[✉], Surjeet Rajendran, and Erwin H. Tanin[✉]

*Department of Physics and Astronomy, Johns Hopkins University,
3400 N. Charles Street, Baltimore, Maryland 21218, USA*



(Received 12 May 2020; accepted 31 August 2020; published 22 September 2020)

The superradiant instability of black hole space-times has been used to place limits on ultra-light bosonic particles. While the initial growth of the mode is gravitational and thus model independent, the ability to place a limit on new particles requires the mode to grow unhindered to a large number density. Nonlinear interactions between the particle and other light degrees of freedom that are mediated through higher-dimension operators can damp this growth, eliminating the limit. However, these nonlinearities may also destroy a cosmic abundance of these light particles, an attractive avenue for their discovery in several experiments. We study the specific example of the QCD axion and show that it is easy to construct models where these nonlinearities eliminate limits from superradiance while preserving their cosmic abundance.

DOI: [10.1103/PhysRevD.102.055015](https://doi.org/10.1103/PhysRevD.102.055015)

I. INTRODUCTION

Ultra-light bosonic particles such as axions, axion-like particles, relaxions and hidden photons have attracted significant attention. These particles are naturally light, emerging as messengers of the deep ultraviolet. They can solve fundamental problems in particle physics such as the strong CP [1–3], hierarchy [4] and cosmological constant problems [5,6]. These particles can also be the dark matter of the Universe, since they can be copiously produced either in the high-temperature Universe or during a period of inflation [7–12]. This exceptionally strong physics case has motivated a plethora of experiments [13–19] to directly search for these particles, along with attempts to impose phenomenological constraints on the viable parameter space of these particles. Many of these constraints arise from astrophysical measurements [20]. This is not a surprise: these particles have suppressed interactions with the Standard Model and these suppressions are most easily overcome in extreme astrophysical environments. But, these constraints are not robust. While gross properties of astrophysical environments are reasonably understood, we do not have direct measurements of the microphysics of these environments.¹ It is thus plausible

that while these constraints may apply to simple models, small modifications can completely remove them [22].

One interesting class of constraints on light particles arises from superradiance. In this process, if a particle can be absorbed by a body, when that body rotates, it can become kinematically favorable for that body to emit that particle and spin down. This process can be efficient for bosonic particles around extreme astrophysical bodies such as black holes [23,24] and neutron stars [25]. In these cases, superradiance can populate the gravitationally bound states of the boson around the star leading to stimulated emission into that state, causing the system to rapidly drain the rotational kinetic energy of the star. Observations of rotating bodies such as black holes and neutron stars can then be used to place limits on the existence of such particles.

In the case of superradiance from black holes, the needed absorptive process is provided by gravitation. Due to the universal nature of gravity, it is often claimed that these bounds are “model independent.” This is of course not the case: while it is true that the initial gravitational growth of the superradiant state is model independent, for these bounds to apply, nonlinearities in the bound state must be sufficiently small. When nonlinear interactions become large, the superradiant amplification can stop, preventing this process from efficiently draining the rotational energy of the system. While some nonlinearities may eliminate the superradiance limit on the existence of light particles, they may also significantly change the cosmology of such particles, potentially eliminating them as dark matter candidates. Given the number of experiments searching for ultra-light dark matter over a wide range of parameters, it is thus important to know if nonlinearities that would damp superradiance would also preserve the dark matter abundance of such particles.

¹This is especially true of bounds placed using the superradiance phenomenon in black holes. The black hole information problem poses a serious challenge to confident claims about the physics of black hole horizons. It is plausible that there are classical singularities (or firewalls) at the horizon [21], significantly affecting the physics.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

In this paper, we show that this is possible. We will specifically study the example of the QCD axion, although our model building approach should be applicable to other scenarios as well. We point out that if the QCD axion has an additional technically natural axion-like interaction with a dark sector that consists of a light (massless) dark photon and a fermion, these additional interactions can completely eliminate superradiance limits on the axion from rotating black holes, while preserving its cosmic abundance. To achieve this, the axion is assumed to couple more strongly with this dark photon than to QCD. This enhances nonlinearities when the superradiant mode begins to grow and thus damps it. The fermions in the dark sector are assumed to have a cosmic abundance; this abundance yields a plasma mass for the hidden photon in the early Universe, preventing these nonlinearities from affecting the evolution of the dark matter axion. This does not happen around black holes today since these fermions do not have a significant number density around them.² The rest of this paper is organized as follows. In Sec. II we present our model and parameters. In Sec. III we show how this parameter space evades superradiance constraints and in Sec. IV we show that it preserves axion dark matter. We then conclude in Sec. V.

II. MODEL

We consider the following Lagrangian for the QCD axion ϕ :

$$\mathcal{L} \supset \frac{\phi}{f_a} G\tilde{G} + \frac{\phi}{f_{\gamma'}} F'\tilde{F}' + qA'_\mu \bar{\psi}\gamma^\mu\psi + m_\psi \bar{\psi}\psi + \frac{1}{2}m_a^2\phi^2. \quad (1)$$

Here f_a is the coupling of the QCD axion to gluons (G), $f_{\gamma'}$ is the coupling to a new massless U(1) (F') and ψ are fermions of charge q under F' with mass m_ψ . We will be interested in QCD axions in the mass range

$$6 \times 10^{-13} \text{ eV} \lesssim m_a \lesssim 2 \times 10^{-11} \text{ eV} \quad (2)$$

that are claimed to have been excluded by measurements of the spins of black holes [24], corresponding to $f_a \approx 10^{18}$ GeV. Our intent is to simply identify a part of parameter space where this model is self-consistent and achieves our desired goal. A broader analysis may reveal additional parameters where this goal could be achieved, but this is beyond the scope of our work. To that end, we take $f_{\gamma'} \approx 10^{11}$ GeV,³ with q and m_ψ in the range shown in

²The damping of superradiance due to nonlinearities was studied in Refs. [26–28]. But, they were not concerned with preserving axion dark matter.

³Axions can naturally possess couplings of varying strengths with different gauge groups. This can be accomplished in the so-called aligned axion or clockwork models [29–31].




FIG. 1. The range of parameter space (m_ψ, q) for $f_a = 10^{18}$ GeV and $f_{\gamma'} = 10^{11}$ GeV satisfying all of the constraints and assumptions in Sec. IV.

Fig. 1. In the following sections, we show that we can accomplish our goals with these parameters.

III. AXION MASS BOUNDS FROM BLACK HOLE SUPERRADIANCE

Superradiance is possible whenever the rotation rate (ω) of the black hole is larger than the mass (m_a) of the emitted particle. But, this process is efficient only when the Bohr radius of the superradiant mode has significant overlap with the ergoregion ($\sim$ the Schwarzschild radius r_s) where superradiant amplification is possible. Together, these conditions imply that superradiance is efficient only when $m_a \approx \omega$ and $\omega r_s \approx 1$ i.e., the bound requires a nearly extremal black hole and it applies to particles whose masses are close to the rotation rate of the black hole. When these conditions are satisfied, superradiance could extract a significant fraction of the rotational energy $\sim M(\omega r_s)^2$ of the black hole (whose mass is M). This energy is carried by the bosonic field ϕ . Since efficient superradiance requires the Bohr radius of this field to be $\sim r_s$, we must then have $m_a^2 \phi^2 r_s^3 \sim M(\omega r_s)^2$. But, since $m_a \approx \omega$ and $\omega r_s \approx 1$, $\phi \sim M_{\text{pl}}$. Given this large field value (and implied number density), limits from superradiance are subject to unknowable nonlinear instabilities that could arise due to new physics at very high energies. We now discuss one such instability that arises from the model in Eq. (1).

Consider the coupling

$$\mathcal{L}_{a\gamma'} = \frac{\phi}{f_{\gamma'}} F'_{\mu\nu} \tilde{F}'^{\mu\nu} \quad (3)$$

between the axion and the dark photon that we introduced in Eq. (1). In the presence of such a coupling, efficient dark

photon production may occur through processes such as tachyonic instability or parametric resonance (narrow or broad). Regardless of the details of the dark photon production, when enough dark photons are produced, their backreaction to the axion will inevitably thwart the exponential axion growth in order to conserve energy. As we will see, this reduces the maximal Bose enhancement of the axion growth rate, making the whole black hole spindown process effectively slower and weakening the superradiance bounds from black hole spin-mass observations.

At scales much smaller than the spatial extent $\sim m_a^{-1}$ of the axion bound states,⁴ the axion field is roughly uniform and can be approximated as

$$\phi(t) = \phi_0 \cos(m_a t). \quad (4)$$

The equation of motion of the circularly polarized dark photon modes $A_k^\pm$ then takes the form of the Mathieu equation [27,32,33]

$$\frac{d^2 A_k^\pm}{dz^2} + [p_k - 2q_k \cos(2z)] A_k^\pm = 0 \quad (5)$$

with

$$p_k = \frac{4k^2}{m_a^2}, \quad q_k = \pm \frac{2k\phi_0}{f_{\gamma'} m_a}, \quad z = \frac{m_a t}{2}. \quad (6)$$

Exponential growth of the dark photon field may proceed through narrow parametric resonance, broad parametric resonance, and tachyonic instability. In order for the produced dark photons to accumulate, their growth rate must be significantly faster than their escape rate out of the axion cloud, i.e., the inverse of the time it takes for a photon to traverse the size of the axion cloud $\sim m_a^{-1}$ [34]

$$\Gamma_{\text{esc}} \sim m_a. \quad (7)$$

It is immediately clear that narrow and broad parametric resonance will not be efficient in draining the axion cloud. While the narrow resonance condition $|q_k| \lesssim 1$ for the fastest growing modes (those with $p_k \simeq 1$) translates to the condition $\phi_0 \lesssim f_{\gamma'}$, it takes $\phi_0 \gg f_{\gamma'}$ for the narrow resonance growth rate $\sim m_a \phi_0 / f_{\gamma'}$ to beat Γ_{esc} . Meanwhile, broad parametric resonance occurs at the rate of $\sim m_a$, which is at best comparable to Γ_{esc} .

Tachyonic instabilities on the other hand are not similarly inhibited. This instability happens when the squared frequency $\omega_\pm^2 = k^2 \mp k \frac{\phi}{f_{\gamma'}}$ of one of the circularly polarized dark photon modes becomes negative. Throughout most of an axion oscillation, during which $|\dot{\phi}| \sim m_a \phi_0$, the fastest

tachyonic growth occurs for the modes $k \simeq |\dot{\phi}| / (2f_{\gamma'})$ at the rate

$$\Gamma_{\text{tach}} \sim \frac{m_a \phi_0}{f_{\gamma'}}. \quad (8)$$

As we can see, $\Gamma_{\text{tach}} \gg \Gamma_{\text{esc}}$ is satisfied⁵ when the axion field amplitude ϕ_0 exceeds $f_{\gamma'}$. This sets an upper bound on the axion amplitude

$$\phi_0 \lesssim f_{\gamma'} \quad (9)$$

since if ϕ_0 were to exceed $f_{\gamma'}$, rapid dark photon production would take place, draining energy from the axion field and preventing ϕ_0 from growing. This suppresses the ability of superradiance to drain rotational energy from the black hole. Suppose the field needed to grow to ϕ_{max} in order to noticeably spin down the black hole.⁶ In the presence of the coupling with the dark photon, this rate is now limited to $\Gamma_{\text{SR}} \left(\frac{f_{\gamma'}}{\phi_{\text{max}}}\right)^2$, where Γ_{SR} is the unsuppressed superradiance rate in the absence of the axion interaction with the dark photon. Let us now see how this relaxes bounds on axions from superradiance.

A. Relaxation of bounds from observed black holes

When superradiance occurs at a rate Γ_{SR} , it needs a time τ_{BH} where

$$\Gamma_{\text{SR}} \tau_{\text{BH}} \gtrsim 100 \quad (10)$$

in order for the superradiant mode to sufficiently grow and damp the rotation of the black hole. Reference [24] used this condition to derive the axion mass bound (2) from the reported spins, masses, and ages of five stellar mass black holes: M33 X-7, LMC X-1, GRO J1655–40, Cygnus X-1, and GRS 1915 + 105. They obtained the superradiance rate Γ_{SR} using the semianalytical method detailed in Ref. [35] and adopted

$$\tau_{\text{BH}} = \min(\tau_{\text{age}}, \tau_{\text{Edd}}) \quad (11)$$

where τ_{age} is the age of the black hole and $\tau_{\text{Edd}} = 4 \times 10^8$ years is the Eddington time. This is tantamount to assuming that the black hole accretes at most at the Eddington rate since its birth.

⁵In order for the dark photon field to grow, the tachyonic growth rate also needs to be much faster than the oscillation rate of the axion, m_a , or else the most tachyonic dark photon mode would see a temporally averaged-out axion field. Nevertheless this condition is parametrically similar to $\Gamma_{\text{tach}} \gg \Gamma_{\text{esc}}$.

⁶As per our earlier arguments, qualitatively, $\phi_{\text{max}} \sim M_{\text{pl}}$. But there are $\mathcal{O}(1)$ factors in the estimate; we will thus use the actual field value obtained from the superradiance analysis in our quantitative arguments.

⁴Here and henceforth we take the “gravitational fine-structure constant” to be $\alpha \equiv m_a M / M_{\text{pl}}^2 \sim 1$ because only then is the superradiance growth rate considerably fast.

In order to place limits, one needs measurements of black hole spin. For the purposes of this section, we take the measurements of black hole spin quoted in Ref. [24] at face value (we comment on the quality of these measurements and the robustness of inferred bounds in Sec. V). In the absence of a coupling to dark photons, the axion masses (2) to which a given black hole is sensitive are simply those values of m_a for which $\Gamma_{\text{SR}}(m_a, M)$ satisfies Eq. (10). In the absence of an analytic solution for the spectrum of superradiant levels, we follow Ref. [36] in using the “semianalytical” Γ_{SR} shown in Fig. 5 of Ref. [35], which agrees with analytical formulas under the $r_s m_a \ll \ell$ and WKB approximations, as well as with numerical results.

When the dark photon coupling is added, the maximum rate of rotational energy extracted from the black hole is suppressed by a factor $(\frac{f_{\gamma'}}{\phi_{\text{max}}})^2$. Using this suppressed rate, we find the largest value of $f_{\gamma'}$ for which $\Gamma_{\text{SR}} \tau_{\text{BH}} \lesssim \max[100, (\frac{\phi_{\text{max}}}{f_{\gamma'}})^2]$ for all black holes considered in Ref. [24]. It suffices to consider the highest superradiant mode $n = 1$, $\ell = m = 1$. We find that a coupling of $f_{\gamma'} \sim 10^{11}$ GeV results in the total removal of the superradiance bounds.

The axion to dark photon coupling scale $f_{\gamma'}$ is highly model dependent and can in principle go to much lower values, especially if one invokes the clockwork scenario [29–31]. Further, since $f_{\gamma'}$ couples two dark sectors (namely, axions and dark photons), there are currently no model-independent constraints on its value.

IV. AXION DARK MATTER

The arguments of Sec. III show that there are no model-independent limits from superradiance on the existence of axions. But, many experimental methods to discover the axion are aimed at searching for a cosmic abundance of these particles. This has been considered an appealing path since generic initial conditions of the Universe end up producing such a cosmic abundance. It is thus interesting to ask if the nonlinear interactions introduced in Eq. (1) would also similarly deplete the cosmic abundance of the axion. Indeed, such depletion is possible and it can be used to eliminate the mild fine tuning of the initial axion angle necessary in QCD axion models with $f_a \gtrsim 10^{12}$ GeV. Since our goal is to simply eliminate superradiance limits while preserving axion dark matter, we present a simple scenario where the cosmic abundance is preserved.

To that end, we add a fermion charged under the dark photon. This fermion is assumed to have a cosmic abundance, giving the dark photon a plasma mass in the early Universe. This plasma mass suppresses the cosmological tachyonic instability, preserving axion dark matter. Since it does not accumulate in significant numbers around black holes, the damping of superradiance around black

holes is unaltered. We now show a parameter space that accomplishes this goal.

As shown in Eq. (1), consider the interactions of the dark photon A'_μ with a dark fermion ψ :

$$\mathcal{L} \supset q A'_\mu \bar{\psi} \gamma^\mu \psi + m_\psi \bar{\psi} \psi. \quad (12)$$

In the presence of a plasma of these fermions with temperature T_ψ and number density n_ψ , the dark photons acquire a plasma mass [37]

$$\omega_p \sim \begin{cases} q T_\psi, & T_\psi \gtrsim m_\psi, \\ q \sqrt{n_\psi / m_\psi}, & T_\psi \lesssim m_\psi. \end{cases} \quad (13)$$

Tachyonic instabilities are suppressed as long as

$$\omega_p \gg \frac{\dot{\phi}}{f_{\gamma'}}. \quad (14)$$

Before demonstrating a viable parameter space, we make the following simplifying assumption. We will assume that the charge q of the fermion is

$$q \lesssim \frac{m_\psi^2}{m_a f_{\gamma'}} \sim 10^{-9} \left(\frac{m_\psi}{1 \text{ eV}} \right)^2 \left(\frac{f_a}{10^{18} \text{ GeV}} \right) \left(\frac{10^{11} \text{ GeV}}{f_{\gamma'}} \right). \quad (15)$$

We do this in order to ignore the complication that the superradiant process around a black hole that produces the dark photon may also simultaneously produce these charged fermions. One may worry that if such charged fermions are produced around the black hole, they could contribute to a residual plasma mass that may suppress the tachyonic instability around the black hole as well. While this is a possibility, further analysis is required to see if this is actually an issue: in the absence of confining forces, it is possible that the produced fermions will simply leave the black hole without creating a plasma that would block the effect. We simply choose to avoid this problem. The condition (15) is equivalent to requiring that there is no Schwinger pair production of the dark fermions from the dark electric field building around the black hole during superradiance i.e., Eq. (15) implies that the dark electric field E' , which is at most comparable to the square root of the maximum energy density of the axion field, $m_a f_{\gamma'}$, does not exceed the Schwinger limit $E'_{\text{crit}} \sim m_\psi^2 / q$.

The dark photon and fermion are uncoupled to the Standard Model except through the highly suppressed axion portal. Observational limits on them are thus weak: they are limited by bounds on dark radiation and dark matter. Given this freedom, we make further simplifying assumptions:

- (1) The dark fermions ψ are in thermal equilibrium among themselves with a temperature

$$T_\psi = \kappa T \quad (16)$$

where T is the temperature of the Standard Model sector and κ is $O(1)$ but small enough to avoid violating the effective relativistic degrees of freedom N_{eff} bound, $\Delta N_{\text{eff}} \lesssim 0.3$ [38].

- (2) $m_\psi \ll 1$ eV.
- (3) The dark fermions are thermally decoupled from the dark photons. This forces⁷

$$q \lesssim 10^{-6} \left(\frac{m_\psi}{1 \text{ eV}} \right)^{1/8}. \quad (17)$$

Assumption 1 ensures that the energy density of ψ is negligible compared to that of the Standard Model radiation when ψ is relativistic, assumption 2 ensures that the energy density of ψ is negligible compared to that of the dark matter when ψ is nonrelativistic, and assumption 3 helps ensure that tachyonic instability is blocked throughout cosmic history by preventing Boltzmann suppression of the ψ abundance.

At the beginning of axion oscillation, the requirement that tachyonic instability be blocked amounts to $\dot{\phi}/f_{\gamma'} < \omega_p$, which imposes the constraint

$$q \gtrsim \frac{m_a f_a \theta_i}{T_{\text{osc}} f_{\gamma'}} \sim 10^{-16} \left(\frac{f_a}{10^{18} \text{ GeV}} \right)^{1/8} \left(\frac{10^{11} \text{ GeV}}{f_{\gamma'}} \right) \quad (18)$$

where $T_{\text{osc}} \sim 0.3 \sqrt{m_a M_{\text{pl}}}$ is the temperature at which the axion begins to oscillate [39] and θ_i is the initial axion angle. As long as ψ is relativistic, the plasma mass $\omega_p \propto T$ decays slower than $\dot{\phi}/f_{\gamma'} \propto T^{-3/2}$, and tachyonic instability remains blocked. Furthermore, thanks to the nonzero chemical potential of ψ that comes with assumption 3, the ψ abundance always follows a Fermi-Dirac distribution with total number density $n_\psi \sim T^3$, even when $T \ll m_\psi$. Consequently, the two quantities of interest, $\dot{\phi}/f_{\gamma'} \propto T^{-3/2}$ and $\omega_p \propto \sqrt{n_\psi} \propto T^{-3/2}$, get Hubble diluted at the same rate when the ψ fermions become nonrelativistic and the tachyonic instability continues to be blocked.

Accounting for all of the above constraints and assumptions, the remaining parameter space is shown in Fig. 1. That said, there is certainly more viable parameter space beyond what we have identified within the boundaries of our simplifying assumptions.

⁷The annihilation rate of the dark fermions ψ to dark photons γ' is $\Gamma_{\text{ann}} \sim 10^{-3} q^4 T \min(1, \frac{T^2}{m_\psi^2})$. When ψ is relativistic, the Hubble rate $H \propto T^2$ scales faster than Γ_{ann} . Conversely, when it is nonrelativistic, $H \propto T^{3/2}$ scales slower than Γ_{ann} . Correspondingly, the ratio Γ_{ann}/H peaks at $T \sim m_\psi$ and requiring Γ_{ann} to be greater than the matter-domination expression of H at $T \sim m_\psi$ imposes an upper bound on the dark fermion charge, $q \lesssim 10^{-6} (m_\psi/1 \text{ eV})^{1/8}$.

Given the above discussion, the reader may wonder if a dark sector was necessary to accomplish our goals, or if they could have been accomplished with the photon. Much of our discussion applies to the photon as well. The extremely dense plasma in the early Universe would inhibit the conversion of axions to photons, preserving axion dark matter. While the photon is better constrained than the dark photon, current limits [40] on the coupling would still allow us to eliminate much of the superradiance bound on axions. However, there is a key uncertainty: we do not know the plasma mass (or its spatial distribution) around the black hole. This plasma mass could range between 10^{-11} – 10^{-2} eV, depending on the details of the plasma environment around the black hole [41,42]. In regions where the plasma mass is small enough, our mechanism will kick in and suppress superradiance. But, absent a better understanding of the spatial distribution of the plasma mass around the black hole, we cannot place robust exclusions.

V. CONCLUSIONS

We have shown that superradiance limits on light particles from measurements of black hole spins are model dependent. While the initial growth of the superradiant modes is due to gravity, the bound requires a large number density (with field values $\sim M_{\text{pl}}$) of the new particle around the black hole. This density could trigger instabilities that dampen superradiance. We have also shown that these instabilities can exist only around black holes without affecting the cosmic abundance of axion dark matter.

The parameter choices we made in this paper were obtained by taking the black hole spin measurements quoted in Ref. [24] at face value. However, the spin measurements used in Ref. [24] are not conservative. There is inherent astrophysical uncertainty in these measurements. Existing measurements of black hole spins are based on the inference of the innermost stable circular orbit of the accretion disk surrounding the black hole which is expected to be a monotonically decreasing function of the black hole spin. Unfortunately, this method is highly dependent on the astrophysical model used. At this time, there is not even community agreement on the measured spin. This is a critical issue for bounds on superradiance from black hole spin measurements: this process is efficient only for nearly extremal black holes. If the spin was 20–30% smaller, the bounds would be significantly different and in some cases entirely absent. At present, it is not clear if the astrophysical model can attain the level of precision needed for these bounds.

Interestingly, if the actual spin was somewhat lower than the nearly extremal values used in Ref. [24], it might be possible to eliminate superradiance limits with just a single dark photon instead of also additionally requiring dark fermions. At lower spin rates, the superradiance rate is significantly slower, enabling $f_{\gamma'}$ to be higher. This larger value of $f_{\gamma'}$ would deplete the cosmic abundance of the

axion, but it is possible that this depletion is at the right level required to avoid over production of axion dark matter, solving the problems of the so-called “anthropic” axion window without fine tuning. In this paper, we focused on demonstrating the model dependence of super-radiance limits on axions. Similar limits have also been placed on other scalars and dark photons. It would be interesting to see if similar models can be constructed in these scenarios where nonlinearities suppress superradiant

growth while preserving a cosmic abundance of these particles.

ACKNOWLEDGMENTS

We would like to thank Emanuele Berti, Michael Fedderke, Peter Graham, David E. Kaplan and Julian Krolik for discussions. S. R. was supported in part by the NSF under Grants No. PHY-1818899 and No. PHY-1638509.

-
- [1] R. D. Peccei and H. R. Quinn, *CP Conservation in the Presence of Instantons*, *Phys. Rev. Lett.* **38**, 1440 (1977).
 - [2] S. Weinberg, *A New Light Boson?*, *Phys. Rev. Lett.* **40**, 223 (1978).
 - [3] F. Wilczek, *Problem of Strong P and T Invariance in the Presence of Instantons*, *Phys. Rev. Lett.* **40**, 279 (1978).
 - [4] P. W. Graham, D. E. Kaplan, and S. Rajendran, *Cosmological Relaxation of the Electroweak Scale*, *Phys. Rev. Lett.* **115**, 221801 (2015).
 - [5] P. W. Graham, D. E. Kaplan, and S. Rajendran, *Born again universe*, *Phys. Rev. D* **97**, 044003 (2018).
 - [6] P. W. Graham, D. E. Kaplan, and S. Rajendran, *Relaxation of the cosmological constant*, *Phys. Rev. D* **100**, 015048 (2019).
 - [7] J. Preskill, M. B. Wise, and F. Wilczek, *Cosmology of the invisible axion*, *Phys. Lett. B* **120**, 127 (1983).
 - [8] M. Dine and W. Fischler, *The not so harmless axion*, *Phys. Lett. B* **120**, 137 (1983).
 - [9] P. W. Graham, J. Mardon, and S. Rajendran, *Vector dark matter from inflationary fluctuations*, *Phys. Rev. D* **93**, 103520 (2016).
 - [10] Y. Nomura, S. Rajendran, and F. Sanches, *Axion Isocurvature and Magnetic Monopoles*, *Phys. Rev. Lett.* **116**, 141803 (2016).
 - [11] P. W. Graham and A. Scherlis, *Stochastic axion scenario*, *Phys. Rev. D* **98**, 035017 (2018).
 - [12] F. Takahashi, W. Yin, and A. H. Guth, *QCD axion window and low-scale inflation*, *Phys. Rev. D* **98**, 015042 (2018).
 - [13] D. Budker, P. W. Graham, M. Ledbetter, S. Rajendran, and A. Sushkov, *Proposal for a Cosmic Axion Spin Precession Experiment (CASPEr)*, *Phys. Rev. X* **4**, 021030 (2014).
 - [14] S. Chaudhuri, P. W. Graham, K. Irwin, J. Mardon, S. Rajendran, and Y. Zhao, *Radio for hidden-photon dark matter detection*, *Phys. Rev. D* **92**, 075012 (2015).
 - [15] P. W. Graham and S. Rajendran, *New observables for direct detection of axion dark matter*, *Phys. Rev. D* **88**, 035023 (2013).
 - [16] S. Beurthey *et al.*, *MADMAX status report*, [arXiv:2003.10894](https://arxiv.org/abs/2003.10894).
 - [17] P. Sikivie, *Invisible axion search methods*, [arXiv:2003.02206](https://arxiv.org/abs/2003.02206).
 - [18] P. W. Graham, D. E. Kaplan, J. Mardon, S. Rajendran, and W. A. Terrano, *Dark matter direct detection with accelerometers*, *Phys. Rev. D* **93**, 075029 (2016).
 - [19] P. W. Graham, D. E. Kaplan, J. Mardon, S. Rajendran, W. A. Terrano, L. Trahms, and T. Wilkason, *Spin precession experiments for light axionic dark matter*, *Phys. Rev. D* **97**, 055006 (2018).
 - [20] G. G. Raffelt, *Astrophysical axion bounds*, *Lect. Notes Phys.* **741**, 51 (2008).
 - [21] D. E. Kaplan and S. Rajendran, *Firewalls in general relativity*, *Phys. Rev. D* **99**, 044033 (2019).
 - [22] M. Ahlers, H. Gies, J. Jaeckel, J. Redondo, and A. Ringwald, *Light from the hidden sector*, *Phys. Rev. D* **76**, 115005 (2007).
 - [23] A. Arvanitaki, S. Dimopoulos, S. Dubovsky, N. Kaloper, and J. March-Russell, *String axiverse*, *Phys. Rev. D* **81**, 123530 (2010).
 - [24] A. Arvanitaki, M. Baryakhtar, and X. Huang, *Discovering the QCD axion with black holes and gravitational waves*, *Phys. Rev. D* **91**, 084011 (2015).
 - [25] D. E. Kaplan, S. Rajendran, and P. Riggins, *Particle probes with superradiant pulsars*, [arXiv:1908.10440](https://arxiv.org/abs/1908.10440).
 - [26] H. Fukuda and K. Nakayama, *Aspects of nonlinear effect on black hole superradiance*, *J. High Energy Phys.* **01** (2020) 128.
 - [27] T. Ikeda, R. Brito, and V. Cardoso, *Blasts of Light from Axions*, *Phys. Rev. Lett.* **122**, 081101 (2019).
 - [28] R. Brito, V. Cardoso, and P. Pani, *Superradiance: Energy Extraction, Black-Hole Bombs and Implications for Astrophysics and Particle Physics* (Springer, New York, 2015), Vol. 906.
 - [29] K. Choi, H. Kim, and S. Yun, *Natural inflation with multiple sub-Planckian axions*, *Phys. Rev. D* **90**, 023545 (2014).
 - [30] K. Choi and S. Hui Im, *Realizing the relaxion from multiple axions and its UV completion with high scale supersymmetry*, *J. High Energy Phys.* **01** (2016) 149.
 - [31] D. E. Kaplan and R. Rattazzi, *Large field excursions and approximate discrete symmetries from a clockwork axion*, *Phys. Rev. D* **93**, 085007 (2016).
 - [32] L. Kofman, A. D. Linde, and A. A. Starobinsky, *Towards the theory of reheating after inflation*, *Phys. Rev. D* **56**, 3258 (1997).
 - [33] P. Agrawal, N. Kitajima, M. Reece, T. Sekiguchi, and F. Takahashi, *Relic abundance of dark photon dark matter*, *Phys. Lett. B* **801**, 135136 (2020).

- [34] M. P. Hertzberg and E. D. Schiappacasse, Dark matter axion clump resonance of photons, *J. Cosmol. Astropart. Phys.* **11** (2018) 004.
- [35] A. Arvanitaki and S. Dubovsky, Exploring the string axiverse with precision black hole physics, *Phys. Rev. D* **83**, 044026 (2011).
- [36] A. Arvanitaki, M. Baryakhtar, and X. Huang, Discovering the qcd axion with black holes and gravitational waves, *Phys. Rev. D* **91**, 084011 (2015).
- [37] E. Braaten and D. Segel, Neutrino energy loss from the plasma process at all temperatures and densities, *Phys. Rev. D* **48**, 1478 (1993).
- [38] R. H. Cyburt, B. D. Fields, K. A. Olive, and T.-H. Yeh, Big bang nucleosynthesis: 2015, *Rev. Mod. Phys.* **88**, 015004 (2016).
- [39] P. Agrawal, G. Marques-Tavares, and W. Xue, Opening up the QCD axion window, *J. High Energy Phys.* **03** (2018) 049.
- [40] L. Chen and J. P. Conlon, Constraints on massive axion-like particles from x-ray observations of NGC 1275, *Mon. Not. R. Astron. Soc.* **479**, 2243 (2018).
- [41] S. Sen, Plasma effects on lasing of a uniform ultralight axion condensate, *Phys. Rev. D* **98**, 103012 (2018).
- [42] M. Boskovic, R. Brito, V. Cardoso, T. Ikeda, and H. Witek, Axionic instabilities and new black hole solutions, *Phys. Rev. D* **99**, 035006 (2019).