

Progerin-expressing endothelial cells are unable to adapt to shear stress

Brooke E. Danielsson,¹ Hannah C. Peters,¹ Kranthi Bathula,¹ Lindsay M. Spear,¹ Natalie A. Noll,¹ Kris N. Dahl,^{2,3} and Daniel E. Conway^{1,*}

¹Department of Biomedical Engineering, Virginia Commonwealth University, Richmond, Virginia; ²Department of Chemical Engineering, Carnegie Mellon University, Pittsburgh, Pennsylvania; and ³Department, Thornton Tomasetti, New York City, New York

ABSTRACT Hutchinson-Gilford progeria syndrome (HGPS) is a rare premature aging disease caused by a single-point mutation in the lamin A gene, resulting in a truncated and farnesylated form of lamin A. This mutant lamin A protein, known as progerin, accumulates at the periphery of the nuclear lamina, resulting in both an abnormal nuclear morphology and nuclear stiffening. Patients with HGPS experience rapid onset of atherosclerosis, with death from heart attack or stroke as teenagers. Progerin expression has been shown to cause dysfunction in both vascular smooth muscle cells and endothelial cells (ECs). In this study, we examined how progerin-expressing endothelial cells adapt to fluid shear stress, the principal mechanical force from blood flow. We compared the response to shear stress for progerin-expressing, wild-type lamin A overexpressing, and control endothelial cells to physiological levels of fluid shear stress. Additionally, we also knocked down ZMPSTE24 in endothelial cells, which results in increased farnesylation of lamin A and similar phenotypes to HGPS. Our results showed that endothelial cells either overexpressing progerin or with ZMPSTE24 knockdown were unable to adapt to shear stress, experiencing significant cell loss at a longer duration of exposure to shear stress (3 days). Endothelial cells overexpressing wild-type lamin A also exhibited similar impairments in adaptation to shear stress, including similar levels of cell loss. Quantification of nuclear morphology showed that progerin-expressing endothelial cells had similar nuclear abnormalities in both static and shear conditions. Treatment of progerin-expressing cells and ZMPSTE24 KD cells with lonafarnib and methystat, drugs previously shown to improve HGPS nuclear morphology, resulted in improvements in adaptation to shear stress. Additionally, the prealignment of cells to shear stress before progerin-expression prevented cell loss. Our results demonstrate that changes in nuclear lamins can affect the ability of endothelial cells to properly adapt to shear stress.

SIGNIFICANCE Endothelial cells are an especially mechanoresponsive cell type, adapting to the physiological forces of blood flow, including cyclic stretch and fluid shear stress. In this study we show that pathological changes in lamin A, associated with the disease Hutchinson-Gilford progeria syndrome (HGPS), impair the ability of these cells to adapt to fluid shear stress, leading to cell loss. In addition to providing a better understanding of the pathology of HGPS, our results illustrate the importance of the nuclear lamina in mechanoadaptation of endothelial cells.

INTRODUCTION

The nuclear lamina, a fibrous lamin-protein network, located between the inner membrane of the nuclear envelope and chromatin, provides structural support for the nucleus and plays a major role in nuclear shape, gene regulation, as well as the assembly and disassembly of the nucleus during cell division (1–4). Numerous studies have

shown the mechanoadaptive and mechanoresponsive nature of the nuclear lamina under force (5–9).

HGPS is a rare, premature aging disease in children caused by an autosomal dominant mutation in the *LMNA* gene, resulting in an aberrant form of lamin A termed progerin (10). The predominant mutation in HGPS involves a de novo point mutation (1824C > T) that activates a cryptic donor splice site resulting in an internal deletion of 50 amino acids. These amino acids include a cleavage site for ZMPSTE24, a protease that removes the farnesyl group from mature lamin A—the loss of this cleavage site leads to the permeant farnesylation of progerin. Permeant farnesylation of progerin is thought to be the critical feature of

Submitted August 20, 2021, and accepted for publication January 5, 2022.

*Correspondence: dconway@vcu.edu

Editor: Tom Misteli.

<https://doi.org/10.1016/j.bpj.2022.01.004>

© 2022 Biophysical Society.

HGPS, which is supported by the observation that ZMPSTE24 knockout mice phenocopy HGPS (11,12). Similar to lamin A, progerin accumulates at the nuclear periphery. However, progerin expression has been shown to alter the nuclear lamina structure and leads to several downstream nuclear defects, including abnormal nuclear morphology, increased nuclear stiffness, redistribution of heterochromatin, modified nuclear pore structure, alterations in gene expression, and nuclear structural instability (13–17).

Patients with HGPS have accelerated atherosclerosis, leading to premature death as a result of heart attack and stroke (18). Cardiovascular alterations in patients with HGPS are similar to atherosclerosis of aging individuals such as exhibiting hypertension, vascular stiffening and calcification, and plaque formation (19–21). Yet these patients do not experience traditional risk factors for atherosclerosis, such as hypercholesterolemia or increased serum levels of C-reactive protein (22). Thus, an important question is why and how progerin expression affects the vasculature to lead to rapid onset of atherosclerosis. One of the hallmarks of the disease is the loss of vascular smooth muscle cells (vSMCs) in large arterial vessels (23,24). Several studies, using both induced pluripotent stem cells and HGPS mouse models have shown that expression of progerin in vSMCs impairs cell proliferation (25), impairs cell response to physiological levels of strain (stretch) (26,27), decreases vasoreactivity (28), and accelerates atherosclerosis by inducing endoplasmic reticulum stress (29,30).

In addition to vSMCs, progerin also alters endothelial cell function. Using an HGPS mouse model with progerin expression only in endothelial cells, it was shown that this resulted in increased inflammation, impaired vascularization, and a shortened life span (31). In a similar HGPS mouse endothelial model, it was observed that progerin-expressing endothelial cells caused cardiac pathologies, and that these endothelial cells had impaired mechanoresponsivity (32). We hypothesized that an important aspect of progerin expression in endothelial cells would be impaired mechanoadaptation to shear stress, which was suggested in a previous study (32). To investigate this hypothesis, we developed an HGPS model using human umbilical vein endothelial cells (HUVECs) expressing progerin, as well as HUVECs with ZMPSTE24 short hairpin RNA (shRNA) knockdown. Progerin-expressing cells, as well as ZMPSTE24 knockdown cells, failed to adapt to physiological levels of fluid shear stress, exhibiting cell loss at longer timepoints of shear stress exposure. Cell loss was rescued by treatment with the farnesyltransferase inhibitor lonafarnib, DNA demethylase inhibitor methylstat, as well as preadaptation of cells to fluid shear stress before progerin expression. Collectively our results show that endothelial cells expressing progerin cannot adapt to the mechanical forces of fluid shear stress, which may be

an important aspect of the rapid onset of atherosclerosis in patients with HGPS.

MATERIALS AND METHODS

Cell culture and transfection

Commercially available primary HUVECs (pooled, passages 3–5, Lonza, Basel, Switzerland) were grown in EGM-2 medium (Lonza, Basel, Switzerland). To express progerin in HUVECs, a previously described HA-tagged progerin (33) was expressed using adenovirus (34). The lowest level of adenovirus that infected nearly 100% of cells was used. To overexpress wild-type lamin A in HUVECs, lamin A adenovirus (based on RefSeq BC014507) was purchased from Vector Biolabs and used at an identical titer level as progerin, as previously described (34).

shRNA knockdown

To knockdown ZMPSTE24, the pLKO-1 vector was used (Sigma Aldrich, St Louis, MO; clone ID TRCN0000294124, target sequence TGGTAAGGCCAATGTTATTTA). Lentivirus was prepared using HEK 293 cells, with second-generation packaging plasmids, pSPAX2 (Addgene 12260) and pMD2.G (Addgene 12259). HUVECs were transduced with lentivirus shRNA and selected with puromycin (1 μ g/mL). A nontargeting shRNA (Sigma SHC216) was used as a control.

Lonafarnib and methylstat treatment

HUVECs were treated with either 1.0 μ M methylstat (Sigma-Aldrich, SML0343), a histone demethyltransferase inhibitor, or 0.5 μ M lonafarnib (Tocris, Bristol, UK; 6265), a farnesyltransferase inhibitor. HUVECs were treated with a daily dose of 1.0 μ M methylstat (Sigma-Aldrich, SML0343) for 48 h. HUVECs were treated with a daily dose of 0.5 μ M lonafarnib (Tocris, 6265) for 72 h.

Fluid shear stress

HUVECs were seeded onto ibidi chamber slides (ibidi-treated μ -slides I^{0.6}, cat #80186 or ibidi-treated μ -slides V^{0.4}, cat#80606, Gräfelting, Germany), coated with 60 ng/mL fibronectin (Sigma-Aldrich, F1141). At 80% confluency, HUVECs were exposed to laminar or oscillatory shear stress using the ibidi pump system (ibidi, cat #10902, Germany), at 12 dynes/cm² for 1, 3, or 6 days with or without modifications perfused in the media.

Cell fixation and labeling

After fluid shear stress experiments were finished, HUVECs were washed two times with PBS and fixed for 10 min at room temperature with 4% paraformaldehyde in PBS. After three washes with PBS, the cells were permeabilized for 10 min at room temperature with 0.2% Triton X-100 in PBS and blocked with 5% BSA for 1 h at room temperature. Cells were then incubated overnight at 4°C room temperature with the primary Ab diluted in blocking solution, using either anti-lamin A antibody (cat # sc-7292, Santa Cruz Biotechnology, Santa Cruz, CA) for control cells, or anti-HA antibody (cat # 901,501, Biolegend) for progerin-expressing cells and anti-lamin B1 antibody (cat # ab16048, Abcam, Cambridge, UK). Three more washes with PBS were then followed by incubation with the secondary Ab (Alexa Fluor 647-conjugated donkey anti-mouse IgG; Thermo Fisher Scientific, Waltham, MA) and stained with rhodamine phalloidin (cat # PHDR1, Cytoskeleton) for 45 min followed by three additional PBS washes. Samples were stained with Hoechst 33342 (Thermo Fisher Scientific) and mounted with ibidi mounting medium (ibidi, cat #50001, Germany). For cell death

assays, an apoptosis/necrosis Detection kit (Abcam, ab176749) was used following the manufacturer's instructions. The same antibodies, as well as ZMPSTE24 (Novus Biologicals, Abingdon, UK; NB100-2387ss) and tubulin (Proteintech, Rosemont, IL; mouse monoclonal), were used for Western blotting quantification of overexpression and knockdown.

Quantification and imaging analysis

Samples were imaged on Zeiss LSM 710 confocal microscope at 20 $\times$ and 63 $\times$. Image analysis was completed using Fuji Image J (Minato City, Japan). We analyzed the progerin-expressing endothelial cell response to the force of shear stress by quantifying abnormally structured nuclear lamina (through lamin A, HA-progerin, and lamin B staining), nuclear lamina outward blebs (through lamin A and HA-progerin staining), and the presence or absence of micronuclei (through Hoechst staining). Cell loss was quantified using a cell counting macros on Image J.

Statistical analysis

Statistical significance was measured using an unpaired, two-tailed Student's *t*-test for data containing two groups. For data involving more than two groups, the analysis of variance (ANOVA) test was performed in order to obtain the statistical analysis for the data sets concerned. A further comparison of the groups was performed using the Tukey test to determine significant differences between groups. All statistical tests were conducted at a 5% significance level ($p < 0.05$). Prism GraphPad

was used for statistical analyses. Each experiment was done in triplicate. Every experiment was completed 3 times, and between 5 and 10 images from each group were quantified.

RESULTS

Endothelial cells expressing progerin or overexpressing wild-type lamin A are unable to adapt to shear stress

Previous work has shown that HGPS cells exhibit both a combination of a stiffened nucleoskeleton and a softened nuclear interior (14), which in turn can cause mechanical irregularities and impaired mechanoadaptation (14,34). We therefore sought to understand how progerin expression would affect endothelial cell responses to fluid shear stress, the frictional drag force created by blood flow. We expressed progerin in HUVECs using an adenovirus, and as a control we also examined the effects of overexpression (OE) of wild-type lamin A. HUVECs expressing either progerin or overexpressing wild-type lamin A had a significant decrease in cell count after 72 h of arterial levels of shear stress (12 dynes/cm²) (Fig. 1 A and B). Western blotting was used to




FIGURE 1 Progerin expression causes cell loss in HUVECs under laminar shear stress. Confocal images taken at 20 $\times$. Scale bar, 50 μ m. Arrow represents direction of fluid shear stress at 12 dynes/cm². LSS, laminar shear stress; OSS, oscillatory shear stress. HUVECs stained with HA-progerin (Progerin), laminA/C (Lamin A-OE, Control, ZMPSTE24, nonsilencing shRNA). Ordinary one-way ANOVA. $N = 3$. (A) Cell loss in progerin-expressing HUVECs under laminar shear stress for 0, 24, and 72 h. (B) Cell count for HUVECs under laminar shear stress for 0, 24, and 72 h. (C) Western blot for protein expression of HA-progerin, lamin a/c and tubulin. (D) Progerin-expressing cells undergo apoptosis under 24 h of laminar shear stress. (E) Cell loss in ZMPSTE24 KD HUVECs under laminar shear stress for 0 and 24 h. (F) Cell count for ZMPSTE24 KD HUVECs under laminar shear stress for 0 and 24 h. (G) Western blot for ZMPSTE24 and tubulin showing the knockdown of ZMPSTE24 in HUVECs. (H and I) No difference was observed between 24 h of oscillatory and laminar shear forces in progerin-expressing HUVECs. Images taken with confocal microscope at 10 $\times$. Scale bar, 50 μ m. A cell death assay showed progerin-expressing HUVECs under 24 h of laminar shear stress experience apoptosis, indicated by the presence of the phosphatidylserine, an apoptotic marker.

confirm both lamin A and progerin OE (Fig. 1 C), which was similar to prior published findings (34). Interestingly there was minimal cell loss for both groups at 24 h of shear stress, indicating that cell loss is not a rapid event. Nontransduced control cells had no cell loss and were able to characteristically align in the direction of shear stress. Additionally, no cell loss was observed for progerin-expressing or lamin A OE HUVECs when grown in static culture (0-h images), indicating that fluid shear stress is necessary for cell loss. Additionally, we examined surface expression of phosphatidylserine as a marker of apoptosis. Progerin-expressing HUVECs had a substantial number of cells with surface expression of phosphatidylserine when exposed to 24 h of shear stress (Fig. 1 D), suggesting that apoptosis is occurring before cell loss.

As an alternate approach to model HGPS in endothelial cells, we developed HUVECs in which ZMPSTE24 was knocked down by shRNA. Densitometry analysis of Western blot (Fig. 1 G) showed that the cells with ZMPSTE24 KD had 40% less ZMPSTE24 than the shRNA control. ZMPSTE24 is a protease involved in the post-translational cleavage of farnesylated prelamin A. The loss of ZMPSTE24 leads to the accumulation of farnesylated lamin A and phenocopies features of HGPS (11,12). HUVECs with knockdown of ZMPSTE24 showed increased nuclear wrinkles and also experienced similar cell loss when exposed to shear stress, (Fig. 1 E and F). To see if oscillatory flow would cause more cell death events in the progerin-expressing cells we exposed the HUVECs to 12 dynes/cm² of laminar and oscillatory shear stress. No significant difference in cell loss was

observed between the different flow types (Fig. 1 H and I). Taken together, these results show OE of lamin A or the expression of progerin, can affect the ability of endothelial cells to adapt to the forces of fluid flow.

Progerin-expressing endothelial cells have increased nuclear abnormalities in both static and shear conditions

Progerin-expressing HUVECs under both flow (Fig. 1 A) and static conditions displayed exaggerated nuclear morphologies. Previous work has shown that the lamina in HGPS cells has a significantly reduced ability to rearrange under mechanical stress (13). We quantified nuclear abnormalities (including dysmorphic nuclear lamina, outward blebs, and micronuclei) in progerin-expressing HUVECs under shear stress and static conditions, and compared results with controls and the lamin OE groups. Consistent with prior observations in nonendothelial cell types, nuclear abnormalities were present in progerin-expressing endothelial cells. Fig. 2 A shows the examples of dysmorphic nuclear lamina in both progerin-expressing and lamin OE cells. In the progerin-expressing cells, we also observed disruptions in lamin B structure. We also examined the incidence of outward blebs (Fig. 2 B) and micronuclei (Fig. 2 C). The quantification of dysmorphic nuclear lamina, outward blebs, and micronuclei showed similar levels of occurrence both under shear stress (Fig. 2 D) and static culture conditions (Fig. 2 E) for progerin-expressing cells.




FIGURE 2 Laminar shear stress causes nuclear envelope disruption in Progerin-expressing HUVECs. Confocal images taken at 20 $\times$, 40 $\times$, and 63 $\times$. Scale bar, 10 μ m. Arrow represents the direction of fluid shear stress at 12 dynes/cm². LSS, laminar shear stress. HUVECs HA-progerin (HA, lamin B, lamin A/C, and Hoechst stained), control, and lamin A-OE (lamin A/C, lamin B, and Hoechst stained). Ordinary one-way ANOVA. N = 3. (A) Dysmorphic nuclear lamina characterized by abnormal structure of lamin a/c, HA-progerin and lamin B staining in progerin-expressing HUVECs under 24 h of laminar shear stress. (B and C) Lamin A/C blebs and micronuclei was observed in progerin-expressing cells after 24 h of shear stress. (D) Quantification of nuclear abnormalities in 24-h laminar shear stress condition. (E) Quantification of nuclear abnormalities in static (0 HR shear stress) condition.

Lonafarnib rescues cell loss and nuclear envelope dysmorphia in progerin-expressing HUVECs under shear stress

In progerin-expressing cells, the attachment of farnesyl groups causes the nuclear envelope to have lobes instead of a round shape. This lobulation of the nuclear envelope is due to the accumulation of progerin within the nucleus and dramatically changes the nuclear architecture as well as its stability (21). Previously it has been shown that the farnesyltransferase inhibitor lonafarnib can be used to prevent progerin accumulation and improve nuclear shape (35) and is currently used in clinical trials as a treatment for HGPS. To inhibit farnesylation of lamin A, ZMPSTE24 KD and progerin-expressing HUVECs were treated with lonafarnib for 72 h.

These treated cells were then subjected to 24 h of laminar shear stress. We hypothesized that inhibiting progerin farnesylation would improve the ability of these cells to adapt to shear stress. Our results showed that lonafarnib significantly prevented cell loss in the ZMPSTE24 KD cells (Fig. 3 A and B), but did not lead to a significant improvement in the progerin-expressing cells. Furthermore, the results showed that lonafarnib treatment prevented nuclear envelope disruptions in the HA-progerin expressing cells exposed to shear stress (Fig. 3 C and D). Taken together, these improvements show the effects of lonafarnib on cell loss after exposure to fluid shear stress, suggesting that lonafarnib enhances the ability of cells to respond and adapt to changes in mechanical forces.

Methylstat rescues cell loss in progerin-expressing and ZMPSTE24 KD HUVECs under shear stress

It has been previously shown that progerin-expressing cells have alterations in histone modifications, including a loss of peripheral heterochromatin, decreased levels of H3K9me3, and increased levels of trimethylation of H4K20, an epigenetic mark for constitutive heterochromatin on H4 (17). Notably, pharmacological-induced increases in heterochromatin have been shown to rescue nuclear morphology in an HGPS patient cells (36). To examine if increases in heterochromatin would improve the ability of progerin-expressing endothelial cells to adapt to shear stress, we used the drug methylstat, an inhibitor of histone trimethyl demethylases. ZMPSTE24 KD and progerin-expressing HUVECs were treated with methylstat for 72 h and exposed to 24 h of laminar shear stress. Cell loss (Fig. 4 A and B) and nuclear envelope disruptions were prevented (Fig. 4 C and D) in cells treated with methylstat. Taken together, these results show that increases in DNA methylation in progerin-expressing endothelial cells rescue nuclear morphology and ability to adapt to shear stress.

Prealignment of progerin-expressing HUVECs prevents cell loss under shear stress

We hypothesized that progerin-expressing endothelial cells which did not have to undergo cellular and nuclear shape changes would be less affected by exposure to shear stress.




FIGURE 3 Lonafarnib rescues cell loss and nuclear envelope disruptions in progerin-expressing HUVECs under shear stress. Confocal images taken at 20× and 40×. Scale bar, 50 μm. Arrow represents direction of fluid shear stress at 12 dynes/cm². LSS, laminar shear stress. HUVECs progerin and ZMPSTE24 KD cells stained with HA-progerin and Hoechst N = 3. HUVECs were treated with a daily dose of 0.5 μM lonafarnib (Tocris, 6265) for 72 h and then exposed to laminar shear stress for 24 h. (A and B) Lonafarnib rescues cell loss in ZMPSTE24 KD cells under 24 h of shear stress compared to static group (0 h). Ordinary one-way ANOVA. N = 3. (C and D) Lonafarnib rescues nuclear envelope disruptions in progerin-expressing HUVECs under shear stress. Unpaired *t* test. N = 3.




FIGURE 4 Methylnstat rescues cell loss in Progerin-expressing HUVECs and ZMPSTE24 KD cells under shear stress. Confocal images taken at 20 $\times$ and 40 $\times$. Scale bar, 50 μ m. Arrow represents direction of fluid shear stress at 12 dynes/cm². LSS, laminar shear stress. HUVECs progerin and ZMPSTE24 KD cells stained with HA-progerin and Hoechst N = 3. HUVECs were treated with a daily dose of 1.0 μ M methylnstat (Sigma-Aldrich, SML0343) for 48 h and then exposed to fluid flow. (A and B) Methylnstat rescues cell loss in progerin-expressing HUVECs, as well as in ZMPSTE24 KD HUVECs, under shear stress compared to the static group (0 h). Ordinary one-way ANOVA. N = 3. (C and D) Methylnstat rescues nuclear envelope disruptions in progerin-expressing HUVECs under shear stress. Unpaired *t* test. N = 3.

To investigate how progerin expression would affect aligned endothelial cells, we first exposed nontransduced HUVECs to shear stress for 72 h to induce alignment. Afterward, these cells were then transduced with progerin or a control adenovirus (GFP), and exposed to an additional 72 h of shear stress. Our results showed that progerin expression in prealigned cells (Fig. 5 A) resulted in reduced cell loss (Fig. 5 C) and nuclear envelope disruptions (Fig. 5 D) when compared with progerin-expressing cells not adapted to shear stress (Fig. 5 B).

DISCUSSION

In this work, we developed an *in vitro* model of HGPS endothelium by (1) expressing progerin and (2) knockdown of ZMPSTE24 in HUVEC. In our model, the expression of progerin could be higher than what is observed in patients with HGPS. Even if the expression of progerin were above physiological levels, it may still be relevant given evidence that progerin expression increases over time, even in healthy cells (37). Additionally, we also assume that knockdown of ZMPSTE24 serves as an additional approach that may result in a more modest, and potentially physiological level of progerin. With either progerin OE or ZMPSTE24 knockdown HUVEC exhibited characteristic progerin-induced changes in nuclear morphology, indicating that the endothelial cell is sensitive to progerin expression. Strikingly, the application of fluid shear stress resulted in dramatic cell loss, occur-

ring between 1 and 3 days of fluid shear stress (Fig. 1). Interestingly, prealignment of endothelial cells to fluid shear stress before expression of progerin (Fig. 5) prevented the cell loss, suggesting that progerin-expression prevents the ability of endothelial cells to adapt to changes in mechanical forces, but has less of an effect in cells already adapted and aligned to shear stress.

Although a number of prior studies have focused on the effects of progerin in fibroblasts and vSMCs (20,24,29,30,38–40), our results add to growing evidence that the endothelium is also sensitive to progerin expression (31,32,41), including a recent study showing that progeria endothelial cells exhibit an altered response to shear stress (42). Although two prior mouse models expressing progerin in the endothelium did not report a dramatic loss of endothelial cells (31,32), our finding that prealigned cells are less sensitive to progerin expression suggests that progerin expression may be most significant when endothelial cells are required to adapt to changes in mechanical environments. Supporting this hypothesis Sun et al. (31) showed that there was defective neovascularization of progerin-expressing endothelial cells in response to ischemia. Additionally, Osmanagic-Myers et al. (32) showed impaired endothelial alignment in response to short-term (3 h) fluid shear stress. The loss of endothelial cells is especially significant in the context of HGPS early onset atherosclerosis, as endothelial dysfunction and damage is considered an initial step in the onset of atherosclerosis (43). Recently, Pitrez




FIGURE 5 Prealignment of Progerin-Expressing HUVECs prevents cell loss under shear stress. Confocal images taken at 20 $\times$. Scale bar, 50 μ m. Arrow represents direction of fluid shear stress at 12 dynes/cm². LSS, laminar shear stress. HUVECs stained with HA-progerin and control stained with lamin A/C. Both groups stained with Phalloidin. Ordinary one-way ANOVA. N = 3. (A) HUVECs were exposed to shear stress for 3 days and once actin alignment (actin fibers oriented parallel to the applied direction of force) and nuclear remodeling was observed (nuclear shape appeared to have changed from more circular to more elongated) progerin expression was induced and the cells were exposed to shear stress of an additional 3 days. (B) Progerin-expressing HUVECs (cells treated with progerin adenovirus 24 h before shear) and the control groups were exposed to shear stress for 3 days total. When progerin was induced after actin and nuclear remodeling had occurred, cell loss (C) and nuclear envelope disruptions were prevented (D).

et al. (44) showed HGPS patient-derived vSMCs cultured under arterial flow conditions detached from a microchip after a few days of culture and that this process is mediated by the upregulation of the enzyme metalloprotease 13.

We showed that two pharmacological treatments previously shown to improve HGPS nuclear morphology in epithelial and fibroblast cell lines, lonafarnib (farnesyltransferase inhibitor) (35) and methylstat (inhibitor of histone trimethyl demethylases) (36) also restored normal nuclear morphology to progerin-expressing endothelial cells (Figs. 3 and 4). Interestingly, increasing chromatin methylation with methylstat significantly improved the ability of progerin-expressing endothelial cells to adapt to fluid shear stress (Fig. 4). However, lonafarnib did not rescue cell loss for progerin-expressing cells (but did rescue ZMPSTE24 knockdown cells). This result may be in part due to differences in the level of progerin expression, which may be lower in the ZMPSTE24 knockdown cells. Thus, improvements in nuclear morphology did not uniformly result in improved mechanoadaptation to shear stress.

An unanswered question in our work is the specific biochemical or physical mechanisms that account for the loss of progerin-expressing cells under fluid shear stress. Cell loss could be a direct result of increased apoptosis under shear stress (Fig. 1 D). Related, prior work by Bidault et al. (41) showed progerin-expressing in endothelial cells have increased markers of DNA damage as well as upregulation of p53 and p21 which induce cellular senescence. The reasons for increased apoptosis are not yet known. One possibility is that apoptosis may result from the inability of progerin expressing cells to remodel their nucleus to adapt to shear stress. Nuclear circularity, elongation, and area rapidly change upon exposure to shear stress (45), suggesting that the nuclear lamina must undergo remodeling. Interestingly impaired lamin A phosphorylation has been observed in HGPS iPS-mesenchymal stem cells (46). Since lamin A phosphorylation is associated with the ability of the nuclear lamina to remodel, it is possible that the impaired lamin A phosphorylation prevents the progerin lamina from remodeling. Our results showing minimal effects of

progerin-expression in prealigned cells (Fig. 5) supports this remodeling hypothesis.

We define cell loss as the decrease in cell count from the start of shear stress (0 h group). In addition to apoptosis, cell loss could be the result of slower proliferation or mechanical instability. Prior work has shown that cells expressing progerin proliferate slower (47); however, given that cells were seeded at confluence, proliferation is not expected to be a major contributing factor to cell count. Cell detachment, perhaps as a result of weakened focal adhesions or cell-cell adhesions, could be a cause for the decreased cell count. Our group recently published work showing that the LINC complex is a necessary structure in endothelial cell mechanoadaptation, in which we observed that disruption of the LINC complex resulted in rapid loss of cells under fluid shear stress (<24 h), due in part to impaired cell-substrate attachment (48). We do not believe the progerin-expressing cells have weaker or impaired attachment to the substrate, as we observed no cell loss or detachment in static culture, as well as minimal cell loss after one day of shear stress.

Interestingly, we observed that similar levels of cells were lost under shear stress when wild-type lamin A was overexpressed as compared with progerin expression (Fig. 1 A). However, lamin A overexpressing cells had significantly less dysmorphic nuclear lamina and blebbing when compared with progerin-expressing cells (Fig. 2 D). Thus, changes in nuclear morphology do not completely account for the cell loss or an inability to adapt to shear stress. OE of lamin A has been shown to increase nuclear stiffness (49), similar to progerin expression (14). It is, therefore, tempting to speculate that stiffer nuclei are less able to adapt to changing mechanical forces. Prior work has shown that there are substantial changes in nuclear shape and stiffness when endothelial cells adapt to shear stress (50), indicating that the nucleus undergoes significant remodeling in response to shear stress.

This work highlights the nuclear lamina as a critical feature for endothelial adaptability to fluid shear stress. We hypothesize that other factors that control nuclear stiffness beyond nuclear lamins, such as chromatin stiffness, may also impact how readily endothelial cells adapt to shear stress. We also note that aging can induce similar nuclear phenotypes to HGPS, including expression of progerin, increased nuclear stiffness, and altered nuclear morphology (51). An important question will be to determine if aging-associated changes in nuclear stiffness similarly impair endothelial cell adaptation to mechanical forces.

AUTHOR CONTRIBUTIONS

B.D., K.D., and D.C. designed the study. N.N., L.S., K.B., H.P., and B.D. performed experiments. H.P. and B.D. analyzed collected data. B.D. and D.C. wrote the manuscript.

ACKNOWLEDGMENTS

We acknowledge Andrew Stephens for helpful discussions. This work was supported by an NSF Graduate Research Fellowship (to B.D.), NIH grant R35 GM119617 (to D.C.), and NSF CAREER award CMMI 1653299 (to D.C.) and American Heart Association grant 16SDG27370007 (to D.C.). Services in support of the research project were generated by the VCU Microscopy Shared Resource, in part, with funding from NIH-NCI Cancer Center support grant P30 CA016059.

REFERENCES

1. Gruenbaum, Y., and O. Medalia. 2015. Lamins: the structure and protein complexes. *Curr. Opin. Cell Biol.* 32:7–12.
2. Gruenbaum, Y., and R. Foisner. 2015. Lamins: nuclear intermediate filament proteins with fundamental functions in nuclear mechanics and genome regulation. *Annu. Rev. Biochem.* 84:131–164.
3. Burke, B., and C. L. Stewart. 2014. Functional architecture of the cell's nucleus in development, aging, and disease. *In* Current Topics in Developmental Biology. Academic Press Inc, pp. 1–52.
4. Eriksson, M., W. T. Brown, ..., F. S. Collins. 2003. Recurrent de novo point mutations in lamin A cause Hutchinson–Gilford progeria syndrome. *Nature*. 423:293–298.
5. Guilluy, C., L. D. Osborne, ..., K. Burridge. 2014. Isolated nuclei adapt to force and reveal a mechanotransduction pathway in the nucleus. *Nat. Cell Biol.* 16:376–381.
6. Guilluy, C., and K. Burridge. 2015. Nuclear mechanotransduction: forcing the nucleus to respond. *Nucleus*. 6:19–22.
7. Cho, S., J. Irianto, and D. E. Discher. 2017. Mechanosensing by the nucleus: from pathways to scaling relationships. *J. Cell Biol.* 216:305–315.
8. Szczesny, S. E., and R. L. Mauck. 2017. The nuclear option: evidence implicating the cell nucleus in mechanotransduction. *J. Biomech. Eng.* 139:0210061.
9. Kirby, T. J., and J. Lammerding. 2018. Emerging views of the nucleus as a cellular mechanosensor. *Nat. Cell Biol.* 20:373–381.
10. De Sandre-Giovannoli, A., R. Bernard, ..., N. Lévy. 2003. Lamin A truncation in Hutchinson–Gilford progeria. *Science*. 300:2055.
11. Bergo, M. O., B. Gavino, ..., S. G. Young. 2002. Zmpste24 deficiency in mice causes spontaneous bone fractures, muscle weakness, and a prelamin A processing defect. *Proc. Natl. Acad. Sci. U S A*. 99:13049–13054.
12. Fong, L. G., J. K. Ng, ..., S. G. Young. 2004. Heterozygosity for Lmna deficiency eliminates the progeria-like phenotypes in Zmpste24-deficient mice. *Proc. Natl. Acad. Sci. U S A*. 101:18111–18116.
13. Dahl, K. N., P. Scaffidi, ..., T. Misteli. 2006. Distinct structural and mechanical properties of the nuclear lamina in Hutchinson–Gilford progeria syndrome. *Proc. Natl. Acad. Sci. U S A*. 103:10271–10276.
14. Booth, E. A., S. T. Spagnol, ..., K. N. Dahl. 2015. Nuclear stiffening and chromatin softening with progerin expression leads to an attenuated nuclear response to force. *Soft Matter*. 11:6412–6418.
15. Goldman, R. D., D. K. Shumaker, ..., F. S. Collins. 2004. Accumulation of mutant lamin A progressive changes in nuclear architecture in Hutchinson–Gilford progeria syndrome. *Proc. Natl. Acad. Sci. U S A*. 101:8963–8968.
16. Verstraeten, V. L. R. M., J. Y. Ji, ..., J. Lammerding. 2008. Increased mechanosensitivity and nuclear stiffness in Hutchinson–Gilford progeria cells: effects of farnesyltransferase inhibitors. *Aging Cell*. 7:383–393.
17. Shumaker, D. K., T. Dechat, ..., R. D. Goldman. 2006. Mutant nuclear lamin A leads to progressive alterations of epigenetic control in premature aging. *Proc. Natl. Acad. Sci. U S A*. 103:8703–8708.
18. Baker, P. B., N. Baba, and C. P. Boesel. 1981. Cardiovascular abnormalities in progeria. Case report and review of the literature. *Arch. Pathol. Lab. Med.* 105:384–386.

19. Olive, M., I. Harten, ..., L. B. Gordon. 2010. Cardiovascular pathology in Hutchinson-Gilford progeria: correlation with the vascular pathology of aging. *Arterioscler. Thromb. Vasc. Biol.* 30:2301–2309.
20. Aliper, A. M., A. B. Csoka, ..., A. Zhavoronkov. 2015. Signaling pathway activation drift during aging: Hutchinson-Gilford progeria syndrome fibroblasts are comparable to normal middle-age and old-age cells. *Aging (Albany NY)*. 7:26–37.
21. Gonzalo, S., R. Kreienkamp, and P. Askjaer. 2017. Hutchinson-Gilford progeria syndrome: a premature aging disease caused by LMNA gene mutations. *Ageing Res. Rev.* 33:18–29.
22. Gordon, L. B., I. A. Harten, ..., A. H. Lichtenstein. 2005. Reduced adiponectin and HDL cholesterol without elevated C-reactive protein: clues to the biology of premature atherosclerosis in Hutchinson-Gilford progeria syndrome. *J. Pediatr.* 146:336–341.
23. Gerhard-herman, M., L. B. Smoot, ..., L. B. Gordon. 2012. Mechanisms of premature vascular aging in child. *Hypertension*. 59:92–97.
24. Stehbens, W. E., B. Delahunt, ..., E. Gilbert-Barness. 2001. Smooth muscle cell depletion and collagen types in progeric arteries. *Cardiovasc. Pathol.* 10:133–136.
25. Zhang, H., Z. M. Xiong, and K. Cao. 2014. Mechanisms controlling the smooth muscle cell death in progeria via down-regulation of poly(ADP-ribose) polymerase 1. *Proc. Natl. Acad. Sci. U S A.* 111:E2261–E2270.
26. Ribas, J., Y. S. Zhang, ..., A. Khademhosseini. 2017. Biomechanical strain exacerbates inflammation on a progeria-on-a-chip model. *Small*. 13:1–13.
27. Kim, P. H., J. Luu, ..., L. G. Fong. 2018. Disrupting the LINC complex in smooth muscle cells reduces aortic disease in a mouse model of Hutchinson-Gilford progeria syndrome. *Sci. Transl. Med.* 10:1–13.
28. Atchison, L., H. Zhang, ..., G. A. Truskey. 2017. A tissue engineered blood vessel model of Hutchinson-Gilford progeria syndrome using human iPSC-derived smooth muscle cells. *Sci. Rep.* 7:1–12.
29. Hamczyk, M. R., R. Villa-Bellosta, ..., V. Andrés. 2018. Vascular smooth muscle-specific progerin expression accelerates atherosclerosis and death in a mouse model of Hutchinson-Gilford progeria syndrome. *Circulation*. 138:266–282.
30. Hamczyk, M. R., R. Villa-Bellosta, ..., V. Andrés. 2019. Progerin accelerates atherosclerosis by inducing endoplasmic reticulum stress in vascular smooth muscle cells. *EMBO Mol. Med.* 11:e9736.
31. Sun, S., W. Qin, ..., B. Liu. 2020. Vascular endothelium-targeted Sirt7 gene therapy rejuvenates blood vessels and extends life span in a Hutchinson-Gilford progeria model. *Sci. Adv.* 6:eaay5556.
32. Osmanagic-Myers, S., A. Kiss, ..., R. Foisner. 2018. Endothelial progerin expression causes cardiovascular pathology through an impaired mechanoresponse. *J. Clin. Invest.* 129:531.
33. Datta, S., C. J. Snow, and B. M. Paschal. 2014. A pathway linking oxidative stress and the Ran GTPase system in progeria. *Mol. Biol. Cell.* 25:1202–1215.
34. Danielsson, B. E., K. V. Tieu, K. Bathula, T. J. Armiger, P. S. Vellala, R. E. Taylor, K. N. Dahl, and D. E. Conway. 2020. Lamin microaggregates lead to altered mechanotransmission in progerin-expressing cells. *Nucleus*. 11:194–204.
35. Capell, B. C., M. R. Erdos, ..., F. S. Collins. 2005. Inhibiting farnesylation of progerin prevents the characteristic nuclear blebbing of Hutchinson-Gilford progeria syndrome. *Proc. Natl. Acad. Sci. U S A.* 102:12879–12884.
36. Stephens, A. D., P. Z. Liu, ..., J. F. Marko. 2018. Chromatin histone modifications and rigidity affect nuclear morphology independent of lamins. *Mol. Biol. Cell.* 29:220–233.
37. Cao, K., C. D. Blair, ..., F. S. Collins. 2011. Progerin and telomere dysfunction collaborate to trigger cellular senescence in normal human fibroblasts. *J. Clin. Invest.* 121:2833–2844.
38. Gabriel, D., D. D. Shafry, ..., K. Djabali. 2017. Intermittent treatment with farnesyltransferase inhibitor and sulforaphane improves cellular homeostasis in Hutchinson-Gilford progeria fibroblasts. *Oncotarget*. 8:64809–64826.
39. Bridger, J. M., and I. R. Kill. 2004. Aging of Hutchinson-Gilford progeria syndrome fibroblasts is characterised by hyperproliferation and increased apoptosis. *Exp. Gerontol.* 39:717–724.
40. Yang, S. H., M. O. Bergo, ..., L. G. Fong. 2005. Blocking protein farnesyltransferase improves nuclear blebbing in mouse fibroblasts with a targeted Hutchinson-Gilford progeria syndrome mutation. *Proc. Natl. Acad. Sci. U S A.* 102:10291–10296.
41. Bidault, G., M. Garcia, ..., V. Bérézat. 2020. Progerin expression induces inflammation, oxidative stress and senescence in human coronary endothelial cells. *Cells*. 9:1201.
42. Atchison, L., N. O. Abutaleb, ..., G. A. Truskey. 2020. iPSC-derived endothelial cells affect vascular function in a tissue-engineered blood vessel model of Hutchinson-Gilford progeria syndrome. *Stem Cell Rep.* 14:325–337.
43. Rafieian-Kopaei, M., M. Setorki, ..., H. Nasri. 2014. Atherosclerosis: process, indicators, risk factors and new hopes. *Int. J. Prev. Med.* 5:927.
44. Pitrez, P. R., L. Estronca, ..., L. Ferreira. 2020. Vulnerability of progeroid smooth muscle cells to biomechanical forces is mediated by MMP13. *Nat. Commun.* 11:1–16.
45. Jiang, Y., and J. Y. Ji. 2018. Expression of nuclear lamin proteins in endothelial cells is sensitive to cell passage and fluid shear stress. *Cell. Mol. Bioeng.* 11:53–64.
46. Cho, S., A. Abbas, ..., D. E. Discher. 2018. Progerin phosphorylation in interphase is lower and less mechanosensitive than lamin-a,c in ipsc-derived mesenchymal stem cells. *Nucleus*. 9:235–250.
47. Dechat, T., T. Shimi, ..., R. D. Goldman. 2007. Alterations in mitosis and cell cycle progression caused by a mutant lamin A known to accelerate human aging. *Proc. Natl. Acad. Sci. U S A.* 104:4955–4960.
48. Denis, K. B., J. I. Cabe, ..., D. E. Conway. 2021. The LINC complex is required for endothelial cell adhesion and adaptation to shear stress and cyclic stretch. *Mol Biol Cell.* 32:1654–1663.
49. Harada, T., J. Swift, ..., D. E. Discher. 2014. Nuclear lamin stiffness is a barrier to 3D migration, but softness can limit survival. *J. Cell Biol.* 204:669–682.
50. Deguchi, S., K. Maeda, ..., M. Sato. 2005. Flow-induced hardening of endothelial nucleus as an intracellular stress-bearing organelle. *J. Biomech.* 38:1751–1759.
51. Ashapkin, V. V., L. I. Kutueva, ..., I. I. Kireev. 2019. Are there common mechanisms between the Hutchinson-Gilford progeria syndrome and natural aging? *Front. Genet.* 10:455.