



# HHS Public Access

## Author manuscript

*J Alzheimers Dis.* Author manuscript; available in PMC 2021 July 16.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Published in final edited form as:

*J Alzheimers Dis.* 2021; 82(1): 47–57. doi:10.3233/JAD-201129.

## Classifying Non-Dementia and Alzheimer's disease/ Vascular Dementia Patients Using Kinematic, Time-Based, and Visuospatial Parameters: The Digital Clock Drawing Test

Anis Davoudi<sup>1</sup>, Catherine Dion<sup>2</sup>, Shawna Amini<sup>2</sup>, Patrick J. Tighe<sup>3</sup>, Catherine C. Price<sup>2,3</sup>, David J. Libon<sup>#4</sup>, Parisa Rashidi<sup>#1</sup>

<sup>1</sup>Department of Biomedical Engineering, University of Florida, Gainesville, Florida, USA

<sup>2</sup>Department of Clinical and Health Psychology, University of Florida, Gainesville, FL, USA

<sup>3</sup>Department of Anesthesiology, University of Florida, Gainesville, Florida, USA

<sup>4</sup>Departments of Geriatrics and Gerontology and Psychology, New Jersey Institute for Successful Aging, School of Osteopathic Medicine, Rowan University, New Jersey, USA

# These authors contributed equally to this work.

### Abstract

**Background:** Advantages of digital clock drawing metrics for dementia subtype classification needs examination.

**Objective:** To assess how well kinematic, time-based, and visuospatial features extracted from the digital Clock Drawing Test (dCDT) can classify a combined group of Alzheimer's disease/ Vascular Dementia patients versus healthy controls (HC), and classify dementia patients with Alzheimer's disease (AD) versus vascular dementia (VaD).

**Methods:** Healthy, community-dwelling control participants (n= 175), patients diagnosed clinically with Alzheimer's disease (n= 29), and vascular dementia (n= 27) completed the dCDT to command and copy with hands to *10 after 11*”. Thirty-seven dCDT command and 37 copy dCDT features were extracted and subjected to Random Forest machine learning analysis.

**Results:** When HC participants were compared to participants with dementia, optimal area under the curve was achieved using models that combined both command and copy dCDT features (AUC= 91.52%). Similarly, when AD versus VaD participants were compared, optimal area under the curve was, again, achieved with models that combined both command and copy features (AUC= 76.94%). Subsequent follow-up analyzes of a corpus of 10 variables of interest complied using a Gini Index found that groups could be dissociated based on kinematic, time-based, and visuospatial features.

**Conclusions:** The dCDT is able to operationally define graphomotor output that cannot be measured using traditional paper and pencil test administration in older health controls and

---

**Correspondence:** Catherine Price, Ph.D., Associate Professor, Clinical and Health Psychology, University of Florida, Gainesville, Florida, 32610; 352-273-5929; cep23@phhp.ufl.edu.

**Conflict of interest:** The authors report no conflict of interest.

participants with dementia. These data suggest that kinematic, time-based, and visuospatial behavior obtained using the dCDT may provide much needed neurocognitive biomarkers that may be able to identify and tract dementia syndromes.

## Keywords

Alzheimer's disease; vascular dementia; digital Clock Drawing Test; executive control; Alzheimer's disease/ Vascular Spectrum Dementia; kinematics

## Introduction

Alzheimer's disease (AD) and vascular dementia (VaD) are two common types of dementia. Current diagnostic studies for the diagnosis and classification of AD and VaD include cerebrospinal fluid assays, neuroimaging studies of degenerative alterations, and comprehensive neuropsychological testing. Yet, these methods can be expensive, invasive, and time-consuming, which limits their potential utility of widespread screening for neurodegenerative disease [1, 2]. A short, noninvasive test that can identify and screen patients at risk for neuropsychological impairment associated with these dementia disorders could help with diagnostic decision-making; and may be useful for treatment planning as disease-modifying medication becomes available.

Previous research has investigated how the Clock Drawing Test (CDT) can detect underlying neurocognitive impairment in dementia [3–6]. The traditional CDT is comprised of two conditions. In the command condition, the patient is asked, *“to draw the clock face, put in all of the numbers and set the hands to read 10 after 11.”* This is followed by the copy condition where patients are asked to copy a model clock [7]. Regarding dementia identification, the standard paper and pencil administration of the clock drawing test has high specificity, but low sensitivity [6, 8]. Recently, Spenciere and colleagues (2017) examined a multitude of paper and pencil clock scoring systems and concluded that many scoring systems are able to distinguish between clinical groups. However, many paper and pencil scoring systems are difficult to operationalize, and there is no consensus regarding optimal clock drawing criteria. This latter observation is illustrated by comparing the scoring systems used by Rouleau et al [9], Royal et al [10], and Cosentino et al. [11]. These clock drawing systems differ with respect to administration procedures, scoring criteria, and the clinical groups that were assessed. Nonetheless, all three studies show meaningful between-group differences that are able to define neurocognitive constructs that underlie dementia syndromes[12].

Over the past decade, a digital version of the clock drawing test (dCDT) has been introduced by the Clock Sketch Consortium [13]. Digital clock drawing protocols use inexpensive digital pen technology. This obviates many of the problems associated with traditional pencil and paper clock drawing scoring systems. Moreover, this technology is able to capture a multitude of behaviors that cannot otherwise be extracted using traditional pen and paper tests. For example, all pen strokes are time stamped and latencies between pen strokes are recorded. This results in precise measurement of all constructional elements such as size, length, pen pressure, and drawing velocity throughout the entire test. The richness of this

information allows for a very detailed analysis of process and errors [14, 15] that can inform researchers and clinicians regarding the underlying neurocognitive impairment.

While previous studies have evaluated the use of the traditional clock drawing test for the detection and classification of dementia subtypes [9–11], the advantages of the dCDT have yet to be fully explored for this purpose. A previous research study conducted by Muller and colleagues used several digitally-obtained features measuring kinematic behavior such as pen pressure and drawing velocity extracted from digital pen data. Logistic regression analyses were able to differentiate patients with amnestic mild cognitive impairment (aMCI) versus mild Alzheimer's dementia (mAD) [16]. Subsequent research [17] using machine-learning models were able to differentiate patients with memory impairment, vascular cognitive disorders, and healthy individuals. In a third study, researchers extracted 350 features from digital clock drawings from 163 patients. Neural networks and information theory-based feature selection methods were able to classify 91.42% of AD versus non-MCI patients [18].

In the present study, we evaluated how well machine learning analysis using Random Forest [19] models can differentiate between dementia patients diagnosed with Alzheimer's Disease (AD), Vascular Dementia (VaD) associated with MRI evidence of subcortical white matter alterations, and non-demented elder controls.

## Methods

### Participants.

Data were acquired from two prospective research investigations approved by the University of Florida's Institutional Review Board; and, from an investigation conducted at Rowan University. Written informed consent was obtained from all participants with all investigations conducted in accordance with the Declaration of Helsinki.

### Healthy Control (HC) Participants.

Inclusion criteria were: age 55 or older, English as primary language, intact instrumental activities of daily living (IADLs), and baseline neuropsychological testing negative for cognitive impairment per Diagnostic and Statistical Manual of Mental Disorders – Fifth Edition [20]. Exclusion criteria were: the presence of neurodegenerative disorders; major medical illness including head trauma or heart disease that could induce encephalopathy; major psychiatric disorders; documented learning disabilities; a seizure disorder or other significant neurological illness; less than a sixth-grade education, and history of substance abuse. The Telephone Interview for Cognitive Status [21] was used to screen for dementia. An in-person appointment was completed where neuropsychological and clock drawing protocols were administered along with an assessment of medical comorbidities (Charlson Comorbidity Index [22]). The presence of anxiety, depression, and ADL/IADL abilities were also assessed. Data were reviewed by a licensed clinical neuropsychologist, double scored, and double data entered for accuracy.

### Dementia Participants.

Individuals with dementia were evaluated at the New Jersey Institute for Successful Aging (NJISA), Memory Assessment Program, School of Osteopathic Medicine, Rowan University. Individuals were seen by a neuropsychologist, psychiatrist, and social worker. Inclusion criteria included age 55 and up, with exclusion criteria the same as that for HC participants except neither the TICS nor the Charlson Comorbidity Index were administered. In addition to neuropsychological assessment, all dementia participants were evaluated with the Mini Mental State Exam [23], serum studies, and an MRI scan of the brain. Additional exclusion criteria included vitamin B12, folate, or thyroid deficiency. Individuals with AD and VaD associated with this study have been described in prior research studies [24]. As described in prior reports [25, 26], these individuals were diagnosed with either AD (n= 29) or VaD (n= 27) using standard diagnostic criteria, respectively.

### Digital Clock Drawing Test (dCDT) Parameters.

The dCDT yields a corpus of over 2000 variables. In the current research 37 dCDT features were used for classification. This corpus of dCDT variables was chosen based on prior research [18] demonstrating the capacity of these variables to dissociate between clinical groups (Table 1).

Clock face area was [27] calculated with the formula  $(\pi * (\text{average of two radii})^2)$ . The presence of anchor digits [28] was determined whether participants initially drew the digits 12, 3, 6, and 9 inside the clock face before drawing the remaining digits. Kinematic parameters including mean pen pressure, and the ratio of mean pen pressure divided by mean drawing velocity. All dCDT features described in Table 1 were extracted from the digitally acquired drawings from both the command and the copy test conditions. When a feature could not be calculated (e.g., the ratio of the hour hand length and minute hand length when a hand is missing) the feature was classified as missing. We handled missing data using Multivariate Imputation by Chained Equations (MICE) [29].

### Machine Learning Analysis.

As stated above, our primary goal was to use Random Forest as a classification model to determine how well dCDT features from the three data sets (1) command, (2) copy, and (3) combined command/copy test conditions could classify individuals into their respective diagnostic categories based upon two binary classification problems, e.g., (1) all dementia participants versus HC classification, and (b) AD versus VaD (Figure 1). Random Forest is a machine learning algorithm that has shown strong performance in public health studies [30, 31]. Random Forest models generally do not make any assumption about the attribute of distributions and also consider the interactions between the attributes. Unlike logistic regression models, Random Forest models are not limited to learning linear decision boundaries. Additionally, their ensemble and boosted approach can accommodate overfitting.

We used nested cross-validation for the development of Random Forest classifiers and five-fold cross-validation for both inner and outer folds in classifying dementia versus HC. Due to the modest sample size, we used three-fold cross-validation for both inner and outer folds

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

to classify AD versus VaD. In addition to accuracy, we reported several other performance metrics including area under the ROC curve (AUC), sensitivity, specificity, and the F1 score, since accuracy alone is not sufficient to reflect the performance of the model for imbalanced datasets. We repeated the experiments 50 times and reported confidence intervals for all performance metrics. We also reported the ranking of variables' importance in the final models in terms of mean decrease in Gini index. The Gini index was used to express the probability that a random element in a data set is randomly and incorrectly labeled according to the distribution of the labels. Thus, the importance of variables are ranked based on how much they each decrease the Gini index in splitting trees in a random forest model.

### Descriptive Statistics.

Chi-Square analyses or *t*-tests were used to compare groups on demographic and the top-ten clock variables selected using variable importance from Random Forest models. Significance was set at  $p < 0.05$ . Bonferroni correction was applied to account for multiple comparisons. All analyses were performed using R version 3.5.2.

## Results

### Demographic and Clinical Data.

Our final dataset comprised 231 individuals (56 dementia, 175 HC; Table 2). Dementia individuals were older (dementia age = 80.04, HC age = 68.37;  $p < 0.001$ ); had fewer years of education (dementia = 12.75, HC = 16.39,  $p < 0.001$ ); and scored lower on the MMSE (dementia total = 22.35, HC total = 28.81,  $p < 0.001$ ). The AD and VaD dementia groups did not differ for age. However, the VaD group was comprised of more female individuals (female/male in VaD: 85.18%, in AD: 51.72%;  $p < 0.016$ ) with fewer years of education (VaD = 11.85 vs. AD = 13.61;  $p < 0.033$ ; Table 3).

### Machine Learning Analysis: HC versus dementia.

None of the participants produced perseverations when drawing digits; or drew digits past the number 12, or a single line connecting the numbers 11 and 2 (i.e., the ten-eleven stroke). These errors, along with missing hands and more-than-two hands variables were excluded because they were absent or occurred very infrequently (command = 2.16% and 0.43%; copy = 0% and 1.30%, respectively). Random Forest classifiers were developed using clock features extracted from command clocks, copy clocks, and combining command and copy clocks. All three models performed similarly in detecting dementia versus HC participants. However, models developed using data from both the command and copy test condition performed best; e.g., area under the curve medians differentiating between HC and dementia were command (89.76%), copy (87.54%), and command/copy (91.52%; Table 4).

### Machine Learning Analysis: AD versus VaD.

The models performed similarly in classifying AD and VaD patients. Models developed using both the command and copy test conditions performed slightly better than models developed using data from command or copy clocks (AUC medians - command = 74.82%, copy = 73.70%, and command & copy = 76.94%; Table 5). Figures 2 and 3 show the median and 95% confidence interval of the top five variables for models using a single clock

condition; and the top 10 variables for models using both clock conditions according to their Gini index (ranked based on the median value).

### **Clock Features of Interest: HC versus Dementia.**

Figure 2 shows the median and 95% confidence interval of the 10 top variables of interest for models using both command and copy features according to their Gini index (ranked based on the median value). Table 6 displays between-group analyses for these 10 variables. Features that statistically differentiated between HC versus all dementia participants were approximately equally represented between the two clock drawing test conditions (command= 4; copy= 3). As displayed in Table 6, groups were differentiated based on a combination of features measuring kinematic operations, decision-making latency, and the location of features within the clock face. Pen pressure (command & copy) was greater for HC compared to dementia. Post-clock face latency was slower for dementia as compared to HC. Finally, precision regarding the location of the hour and minute hands in relation to the clock face, center dot was more accurate for HC compared to dementia.

### **Clock Features of Interest: AD versus VaD.**

Figure 3 shows the median and 95% confidence interval of the 10 top variables of interest for models using both clock conditions together according to their Gini index (ranked based on the median value). In this analysis, there were perhaps, more features from the copy as compared to the command test condition that differentiated between groups (copy= 4; command= 2). As displayed in Table 7 groups were differentiated based on a combination of features measuring kinematic operations and drawing speed. Pen pressure for both test conditions was greater for VaD compared to AD patients. The ratio of pen pressure and drawing velocity was greater for VaD compared to AD consistent with reduced pen pressure along with slower drawing speed. Finally, total drawing time was slower for VaD compared to AD in the copy test condition.

## **Discussion**

The purpose of the current research was to examine how well features from the dCDT analyzed with machine learning using Random Forrest could correctly classify HC versus AD/ VaD dementia patients; and, patients diagnosed clinically with AD versus VaD. It is widely believed as symptoms and behavior suggesting either MCI or dementia emerge decades before sufficient functional and neuropsychological impairment occurs that would permit a clinical diagnosis. At such time that disease-modifying medication is available, the putative target population will most certainly be individuals early in the course of their illness. An important question revolves around the neuropsychological methods or techniques that are able to identify emergent AD/ VaD syndromes.

The MMSE and MoCA are commonly administered neuropsychological tests used to screen for neurocognitive impairment. The problems associated with the MMSE and MoCA include the lack of granularity of the behavior that can be assessed and scored. The dCDT obviates this problem. The time necessary to administer the test is very reasonable; and, as shown above, a large corpus of highly nuanced behavior can be measured. As compared to

prior digital clock drawing research, the data described above achieved comparable or better rates of classifying HC versus dementia individuals [15, 25]. Our classification rates are also comparable to prior research comparing HC versus individuals diagnosed with MCI recruited from a memory clinic [18].

### **Differences between Healthy Control and Dementia Participants -**

From the entire corpus of approximately 2000 dCDT features, a circumscribed corpus of 37 command/ copy dCDT features were selected and assessed with machine learning analyses. Ten variables of interest emerged. The decision to use the corpus of the 37 features described above was guided by the fact that these variables measure very common and necessary behavior for successful test performance.

In the analysis classifying HC versus all dementia individuals, features that were able to dissociate between groups included kinematic parameters that measured command and copy pen pressure. In both clock drawing test conditions, pen pressure was lower when dementia patients were compared to HC participants. These data are consistent with prior clock drawing research [15]. These data are also similar to prior research associating reduced pen pressure among patients with Parkinson's disease (PD) versus HC [32] when PD patients were asked to draw an Archimedean spiral. Accurate output using a pen requires the production of comparatively gross strokes such as the clock face, and the production of discrete strokes such as digits and the clock hands [33]. The current research was able to only analyze mean pen pressure for the entire drawing. However, it is possible that greater between-group differences would have emerged to the extent pen pressure data could have been obtained from each portion of the drawing (i.e., clock face, number placement, time setting).

### **Time-Based dCDT Parameters -**

In the copy condition only, participants with dementia produced lower Post-Clock Face Latency (PCFL). PCFL is one of a number of decision-making latencies obtained from the dCDT [34] and measures the time necessary to transition from one portion of the test to the next portion of the test. Most often PCFL measures the time between the completion of the clock face and drawing either the center dot or the number 12. Slower PCFL for dementia versus HC participants in the copy test condition might suggest the need for greater scanning back and forth from the clock model to the patient's drawing. If this is true, then slower copy PCFL suggests that copying a model of a clock might not be as automatized as expected. Concomitant measures of visual scanning behavior while patients are drawing would help disambiguate this issue. Also, in the copy condition only dementia, as compared to HC individuals were less accurate in drawing both hands in relation to the center dot.

### **AD versus VaD Patients -**

Many of the variables of interest described above also dissociated AD versus VaD patients. Patients with AD obtained lower scores for command/ copy pen pressure compared to VaD. However, participants with VaD, relative to AD, appeared to exert greater pen pressure in the context of slower velocity. VaD patients also produced slow total time to completion in the copy test condition compared to AD patients. Both of these findings are consistent with

previous research showing that VaD patients produce less output as a function of time [35]. Prior research has argued that the neurocognitive constructs underlying clock drawing to command and clock are complimentary but not identical. These observations are consistent with the data described above showing that optimal classification was achieved using a combination of command and copy features. Finally, a question that needs to be addressed revolves around the practicality of extracting the behavior described above. Does the analysis of nuanced graphomotor output truly offer a clinical advantage? The data described above appears to answer this question in the affirmative. However, we acknowledge that additional research examining other clinical groups needs to be undertaken.

The current research is not without weakness. First, we recognize the need for replication in larger samples and correction for age/education differences between dementia and the HC participants. There is a small but statistically significant group difference in education. It is possible that lower education may negatively impact clock drawing performance; education is considered neuroprotective against the emergence of dementia [36, 37]. There is also an age difference between dementia and HC, with the HC group younger. A post-hoc analysis showed us, however, a similar model (lower AUC, 87.89 for command and copy; 87.24 command; 86.16 copy) for participants age 70 and up (67 HC, and 52 dementia). Although dementia groups did not differ for age, this is another factor that can potentially influence clock drawing behavior between and within dementia participants. Second, while we opted to use Random Forest classification models because of their overall strong performance in other health domain studies, the performance of other machine learning models needs to be evaluated. While Random Forest model performance is not affected significantly by collinearity among variables, variable importance will, nonetheless, be impacted; the reported variable importance values need to be interpreted with caution. Because of the modest and uneven sample size, weighting the outcome classes and under-sampling/oversampling techniques may also improve the performance of the models. Third, marginally different recruitment procedures were used to recruit healthy control and dementia participants (e.g., the HC group did not complete laboratory measurements examining vitamin B12, folate, or thyroid deficiency, that may have resulted in exclusion). Finally, most of the participants in the current research were white and right-handed, suggesting the need to replicate the findings reported above with participants from other ethnic/demographic communities.

Despite these limitations, the current research has several strengths including data from two common dementia subtypes and the analysis of features that measure kinematic behavior. These data suggest that machine learning analyses of digitally acquired output using the dCDT could provide much needed neuropsychological biomarkers that might be used to identify and monitor changes in cognition over time for dementia syndromes. Future studies need to examine dCDT variables for validity and cross comparison across different test modalities (pen capture versus tablet) and dCDT variable value for differentiating HC from neurodegenerative diseases including those with and without dementia (e.g., PD; [36]).

## Acknowledgements:

We sincerely thank the participants who provided time and effort towards this investigation and allowed us to improve our understanding of clock drawing behaviors. We also sincerely thank Dana Penney, Ph.D., Director of Neuropsychology, Lahey Hospital, Boston, Massachusetts, and Randall Davis, Ph.D., Professor, Massachusetts Institute of Technology, Cambridge, Massachusetts - without whom this study would not have been possible due to their development of the dCDT. Drs. Penney and Davis are the originators of the digital clock drawing software used within the current investigation. We also sincerely thank the research coordinators and administrative staff members who assisted with data collection, Institutional Research Board processing, and administrative funding paperwork.

### Funding:

NIH grants from the National Institutes of Health (R01 AG055337, R01NS082386; R01 NR014810) and National Science Foundation (NSF 13-543). P.R. was supported by CAREER award, NSF-IIS 1750192, from the National Science Foundation (NSF), Division of Information and Intelligent Systems (IIS), NIH NIBIB R21EB027344-01, and by NIH R01AG055337.

## Bibliography

- [1]. Bernstein A, Rogers KM, Possin KL, Steele NZ, Ritchie CS, Kramer JH, Geschwind M, Higgins JJ, Wohlgemuth J, Pesano R (2019) Dementia assessment and management in primary care settings: a survey of current provider practices in the United States. *BMC health services research* 19, 919. [PubMed: 31783848]
- [2]. Zverova M (2018) Alzheimer's disease and blood-based biomarkers - potential contexts of use. *Neuropsychiatr Dis Treat* 14, 1877–1882. [PubMed: 30050302]
- [3]. Kirby M, Denihan A, Bruce I, Coakley D, Lawlor BA (2001) The clock drawing test in primary care: sensitivity in dementia detection and specificity against normal and depressed elderly. *International Journal of Geriatric Psychiatry* 16, 935–940. [PubMed: 11607936]
- [4]. Reiner K, Eichler T, Hertel J, Hoffmann W, Thyrian JR (2018) The Clock Drawing Test: A Reasonable Instrument to Assess Probable Dementia in Primary Care? *Current Alzheimer Research* 15, 38–43. [PubMed: 28891446]
- [5]. Ahn T-B, Na B, Lee D (2018) Visuospatial dysfunction in cube copying test and clock drawing test in non-demented Parkinson's disease: a volumetric analysis. *Parkinsonism & Related Disorders* 46, e50.
- [6]. Palsetia D, Rao GP, Tiwari SC, Lodha P, De Sousa A (2018) The clock drawing test versus mini-mental status examination as a screening tool for dementia: A clinical comparison. *Indian journal of psychological medicine* 40, 1. [PubMed: 29403122]
- [7]. Mainland BJ, Shulman KI (2017) Clock drawing test In *Cognitive Screening Instruments* Springer, pp. 67–108.
- [8]. Roalff DR, Moberg PJ, Xie SX, Wolk DA, Moelter ST, Arnold SE (2013) Comparative accuracies of two common screening instruments for classification of Alzheimer's disease, mild cognitive impairment, and healthy aging. *Alzheimer's & dementia : the journal of the Alzheimer's Association* 9, 529–537.
- [9]. Royall DR, Cordes JA, Polk M (1998) CLOX: an executive clock drawing task. *Journal of Neurology, Neurosurgery & Psychiatry* 64, 588.
- [10]. Rouleau I, Salmon DP, Butters N, Kennedy C, McGuire K (1992) Quantitative and qualitative analyses of clock drawings in Alzheimer's and Huntington's disease. *Brain and Cognition* 18, 70–87. [PubMed: 1543577]
- [11]. Cosentino S, Jefferson A, Chute DL, Kaplan E, Libon DJ (2004) Clock Drawing Errors in Dementia: Neuropsychological and Neuroanatomical Considerations. *Cognitive and Behavioral Neurology* 17.
- [12]. Spenciere B, Alves H, Charchat-Fichman H (2017) Scoring systems for the Clock Drawing Test: A historical review. *Dementia & neuropsychologia* 11, 6–14. [PubMed: 29213488]

[13]. Libon DJ, Penney DL, Davis R, Tabby DS, Eppig J, Nieves C, Bloch A, Donohue J, Brennan L, Rife K (2014) Deficits in processing speed and decision making in relapsing-remitting multiple-sclerosis: the digital clock drawing test (dCDT). *J Mult Scler* 1, 2.

[14]. Kaplan E (1988) The process approach to neuropsychological assessment. *Aphasiology* 2, 309–311.

[15]. Kaplan E (1990) The process approach to neuropsychological assessment of psychiatric patients. *The Journal of Neuropsychiatry and Clinical Neurosciences*.

[16]. Muller S, Herde L, Preische O, Zeller A, Heymann P, Robens S, Elbing U, Laske C (2019) Diagnostic value of digital clock drawing test in comparison with CERAD neuropsychological battery total score for discrimination of patients in the early course of Alzheimer's disease from healthy individuals. *Sci Rep* 9, 3543. [PubMed: 30837580]

[17]. Souillard-Mandar W, Davis R, Rudin C, Au R, Libon DJ, Swenson R, Price CC, Lamar M, Penney DL (2016) Learning Classification Models of Cognitive Conditions from Subtle Behaviors in the Digital Clock Drawing Test. *Mach Learn* 102, 393–441. [PubMed: 27057085]

[18]. Binaco R, Calzaretto N, Epifano J, McGuire S, Umer M, Emrani S, Wasserman V, Libon DJ, Polikar R Machine Learning Analysis of Digital Clock Drawing Test Performance for Differential Classification of Mild Cognitive Impairment Subtypes Versus Alzheimer's Disease. *Journal of the International Neuropsychological Society*, 1–11.

[19]. Breiman L (2001) Random Forests. *Machine Learning* 45, 5–32.

[20]. American Psychiatric Association., American Psychiatric Association. DSM-5 Task Force. (2013) Diagnostic and statistical manual of mental disorders : DSM-5, American Psychiatric Association, Washington, D.C.

[21]. Cook SE, Marsiske M, McCoy KJ (2009) The use of the Modified Telephone Interview for Cognitive Status (TICS-M) in the detection of amnestic mild cognitive impairment. *Journal of geriatric psychiatry and neurology* 22, 103–109. [PubMed: 19417219]

[22]. Charlson ME, Pompei P, Ales KL, MacKenzie CR (1987) A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 40, 373–383. [PubMed: 3558716]

[23]. Folstein MF, Folstein SE, McHugh PR (1975) "Mini-mental state"; A practical method for grading the cognitive state of patients for the clinician. *Journal of psychiatric research* 12, 189–198. [PubMed: 1202204]

[24]. Emrani S, Lamar M, Price CC, Wasserman V, Matusz E, Au R, Swenson R, Nagele R, Heilman KM, Libon DJ (2020) Alzheimer's/Vascular Spectrum Dementia: Classification in Addition to Diagnosis. *Journal of Alzheimer's Disease*, 1–9.

[25]. Price CC, Jefferson AL, Merino JG, Heilman KM, Libon DJ (2005) Subcortical vascular dementia: integrating neuropsychological and neuroradiologic data. *Neurology* 65, 376–382. [PubMed: 16087901]

[26]. Price CC, Garrett KD, Jefferson AL, Cosentino S, Tanner JJ, Penney DL, Swenson R, Giovannetti T, Bettcher BM, Libon DJ (2009) Leukoaraiosis severity and list-learning in dementia. *The Clinical neuropsychologist* 23, 944–961. [PubMed: 19370451]

[27]. Hizel LP, Warner ED, Wiggins ME, Tanner JJ, Parvataneni H, Davis R, Penney DL, Libon DJ, Tighe P, Garvan CW, Price CC (2019) Clock Drawing Performance Slows for Older Adults After Total Knee Replacement Surgery. *Anesth Analg* 129, 212–219. [PubMed: 30273231]

[28]. Lamar M, Ajilore O, Leow A, Charlton R, Cohen J, GadElkarim J, Yang S, Zhang A, Davis R, Penney D, Libon DJ, Kumar A (2016) Cognitive and connectome properties detectable through individual differences in graphomotor organization. *Neuropsychologia* 85, 301–309. [PubMed: 27037044]

[29]. Buuren SV, Groothuis-Oudshoorn K (2010) mice: Multivariate imputation by chained equations in R. *Journal of statistical software*, 1–68.

[30]. Tu W, Chen PA, Koenig N, Gomez D, Fujiwara E, Gill MJ, Kong L, Power C (2020) Machine learning models reveal neurocognitive impairment type and prevalence are associated with distinct variables in HIV/AIDS. *Journal of NeuroVirology* 26, 41–51. [PubMed: 31520320]

[31]. Kanerva N, Kontto J, Erkkola M, Nevalainen J, Mannisto S (2018) Suitability of random forest analysis for epidemiological research: Exploring sociodemographic and lifestyle-related risk

factors of overweight in a cross-sectional design. *Scand J Public Health* 46, 557–564. [PubMed: 29082809]


[32]. Zham P, Kumar DK, Dabnichki P, Poosapadi Arjunan S, Raghav S (2017) Distinguishing Different Stages of Parkinson's Disease Using Composite Index of Speed and Pen-Pressure of Sketching a Spiral. *Front Neurol* 8, 435. [PubMed: 28932206]

[33]. HoRrE S, Shibata K (2018) Quantitative evaluation of handwriting: factors that affect pen operating skils. *The Journal of Physical Therapy Science* 30, 971–975. [PubMed: 30154583]

[34]. Piers RJ, Devlin KN, Ning B, Liu Y, Wasserman B, Massaro JM, Lamar M, Price CC, Swenson R, Davis R (2017) Age and graphomotor decision making assessed with the digital clock drawing test: the Framingham Heart Study. *Journal of Alzheimer's Disease* 60, 1611–1620.


[35]. Lamar M, Price CC, Davis KL, Kaplan E, Libon DJ (2002) Capacity to maintain mental set in dementia. *Neuropsychologia* 40, 435–445. [PubMed: 11684176]

[36]. Dion C, Frank BE, Crowley SJ, Hizel LP, Rodriguez K, Tanner J, Libon DJ, Price CC (2021) Parkinson's Disease Cognitive Phenotypes Show Unique Clock Drawing Features when Measured with Digital Technology. *J Parkinsons Dis* 11(2), 1–13.. [PubMed: 33492249]




**Figure 1.**

Analysis flow. The analysis in the study included preprocessing the raw pen movement data, extracting features, and developing and testing classifiers.

**Figure 2.**

Ranking of the top ten variables ranked based on their median in 50 runs for Random Forest models for classification of dementia vs HC using A) command data, B) copy data, and C) command and copy data together.

**Figure 3.**

Ranking of the top ten variables ranked based on their median in 50 runs for Random Forest models for classification of dementia subtypes using A) command data, B) copy data, and C) command and copy data together.

**Table 1.**

Digital Clock Drawing Test Feature description.

| Variable Name  | Variable Description   |
|--|--|
| <b>Total number of strokes</b>                           | Total number of strokes used in drawing all components of this clock   |
| <b>Total completion time (sec)</b>                       | Total time to draw all clock components  |
| <b>Clockface number of strokes</b>                       | Total number of strokes used to draw the clockface   |
| <b>Clockface total time (sec)</b>                        | Time to draw the clockface   |
| <b>Clockface overshoot distance (mm)</b>                 | The minimum distance between start of the longest clockface stroke and the end, divided by the total length of the stroke. Gives a normalized measure of how far apart or overlapped the closest points near the end of the stroke are |
| <b>Clockface overshoot angle (degrees)</b>               | Number of degrees between the start and end of the longest stroke in the first clockface, measured from the center of the ellipse. Positive: end of stroke goes past the start, negative: end does not go past the start               |
| <b>Hour Hand length (mm)</b>                             | Distance from innermost to outermost point on the hand   |
| <b>Hour Hand distance from center (mm)</b>               | Distance from inner end of hour hand to center of ellipse  |
| <b>Hour Hand difference from ideal angle (degrees)</b>   | Difference between hour hand angle and the perfect angle   |
| <b>Minute Hand length (mm)</b>                           | Distance from innermost to outermost point on the hand   |
| <b>Minute Hand distance from center (mm)</b>             | Distance from inner end of minute hand to center of ellipse  |
| <b>Minute Hand difference from ideal angle (degrees)</b> | Difference between minute hand angle and the perfect angle   |
| <b>Hour Hand length/Minute Hand length</b>               | Represents the ratio of hour hand length the minute hand length  |
| <b>Post-Clock Face Latency (PCFL) (sec)</b>              | Delay between finishing the last stroke in the clockface and whatever is drawn next  |
| <b>Pre-First Hand Latency (PFHL) (sec)</b>               | Delay between whatever was drawn before the first clock hand and starting to draw that clock hand  |
| <b>Pre-Second Hand Latency (PSHL) (sec)</b>              | Delay between whatever was drawn before the second clock hand and starting to draw that clock hand   |
| <b>Total Drawing time (sec)</b>                          | Total time with the pen drawing on the paper   |
| <b>Drawing length (mm)</b>                               | Summation of total ink length of the drawn components  |
| <b>Average digit height (mm)</b>                         | Average height of the bounding boxes for all numbers on the clock  |
| <b>Average digit width (mm)</b>                          | Average width of the bounding boxes for all numbers on the clock   |
| <b>Average digit distance from circumference (mm)</b>    | Average distance from each digit to the nearest point on the clock face  |
| <b>Average digit misplacement (degrees)</b>              | A summation of each digit's distance from ideal placement, divided by the number of digits   |
| <b>Clockface area (mm<sup>2</sup>)</b>                   | Area of a circle fitted to the drawn clockface   |
| <b>Symmetry</b>  | Measure of how circular or oblong the clockface is, using horizontal and vertical dimensions   |
| <b>Velocity (mm/sec)</b>                                 | Drawing length/Drawing time  |
| <b>Average pen pressure</b>                              | Average pressure of the pen on the paper during the entire test  |
| <b>Standard deviation of pressure</b>                    | Standard deviation of pressure of the pen on the paper during the entire test  |
| <b>Average pressure/velocity</b>                         | Average pressure/Velocity  |
| <b>Any digit missing</b>                                 | Any digits not appearing on the clock  |
| <b>Any digit outside</b>                                 | Any digits placed outside the clockface  |
| <b>Any digit perseveration</b>                           | Any digits repeated  |
| <b>Anchoring</b>   | Anchoring  |

| Variable Name                      | Variable Description                                 |
|------------------------------------|--|
| <b>Any hand missing</b>            | Any of the hour hand or minute hand missing          |
| <b>More than one for each hand</b> | More than one hour-hand or more than one minute-hand |
| <b>Hand crossed Out</b>            | Any of the hands crossed out                         |
| <b>Any digit over 12</b>           | Any digits over 12 drawn                             |
| <b>Ten-eleven stroke</b>           | Single line drawn connecting the numbers 11 and 2    |

**Table 2.**

Participants Demographic/ Clinical Characteristics.

| Variable             | All (n=231)  | Dementia (n=56) | HC (n=175)   | p-value |
|----------------------|--------------|-----------------|--------------|---------|
| Age, mean (SD)       | 71.20 (7.72) | 80.04 (6.28)    | 68.37 (5.76) | <0.0001 |
| Female, N (%)        | 118 (51.08%) | 38 (67.86%)     | 80 (45.71%)  | 0.0063  |
| Race: white N (%)    | 223 (96.54%) | 55 (98.21%)     | 168 (96.00%) | 0.7122  |
| Handedness           | 218 (94.78%) | 56 (100.00%)    | 162 (93.10%) | 0.0943  |
| Education, mean (SD) | 15.52 (3.09) | 12.75 (3.07)    | 16.39 (2.54) | <0.0001 |
| MMSE, mean (SD)      | 27.33 (3.16) | 22.35 (2.71)    | 28.81 (1.07) | <0.0001 |

**Table 3.**

Demographic and Clinical Characteristics of Dementia Patients.

| Variable             | All (n=56)   | AD (n=29)    | VaD (n=27)   | p-value |
|----------------------|--------------|--------------|--------------|---------|
| Age, mean (SD)       | 80.04 (6.28) | 80.34 (6.03) | 79.72 (6.65) | 0.7163  |
| Female, N (%)        | 38 (67.86%)  | 15 (51.72%)  | 23 (85.18%)  | 0.0167  |
| Race: white N (%)    | 55 (98.21%)  | 28 (96.55%)  | 27 (100.00%) | 1       |
| Handedness           | 56 (100.00%) | 29 (100.00%) | 27 (100.00%) | -       |
| Education, mean (SD) | 12.75 (3.07) | 13.61 (2.81) | 11.85 (3.12) | 0.0330  |
| MMSE, mean (SD)      | 22.35 (2.71) | 22.24 (3.35) | 22.44 (2.02) | 0.7931  |

**Table 4.**

Model performance reported for the dementia vs. HC classification – AUC: Area under the ROC curve.

| Model: Dementia vs. Healthy | Accuracy             | AUC                  | Specificity          | Sensitivity          | F1 score             |
|-----------------------------|----------------------|----------------------|----------------------|----------------------|----------------------|
| <b>Command</b>              | 90.48 (85.91, 93.74) | 89.76 (84.22, 93.92) | 97.71 (94.86, 98.86) | 69.64 (52.19, 82.14) | 77.95 (64.66, 86.32) |
| <b>Copy</b>                 | 89.18 (84.18, 92.54) | 87.54 (81.54, 92.83) | 96.57 (93.84, 98.29) | 66.07 (51.79, 78.17) | 73.84 (61.35, 83.13) |
| <b>Command &amp; Copy</b>   | 91.34 (87.64, 94.71) | 91.52 (86.74, 95.17) | 97.71 (96.0, 99.43)  | 71.43 (57.95, 83.53) | 80.21 (69.61, 88.37) |

**Table 5.**

Model performance reported for the VaD vs. AD subtype classification – AUC: Area under the ROC curve.

| Model: Dementia subtypes  | Accuracy             | AUC                  | Specificity          | Sensitivity          | F1 score             |
|---------------------------|----------------------|----------------------|----------------------|----------------------|----------------------|
| <b>Command</b>            | 74.11 (53.57, 85.31) | 74.82 (53.39, 85.70) | 72.41 (55.17, 86.21) | 74.07 (48.15, 95.46) | 74.58 (51.07, 85.19) |
| <b>Copy</b>               | 73.21 (58.93, 85.31) | 73.70 (59.08, 85.31) | 79.31 (62.07, 89.66) | 66.67 (48.15, 58.19) | 72.00 (53.86, 84.70) |
| <b>Command &amp; Copy</b> | 76.78 (60.71, 87.10) | 76.94 (60.75, 87.48) | 79.31 (55.17, 88.87) | 74.07 (56.39, 92.59) | 76.00 (58.54, 87.38) |

**Table 6.**

Ranking of the top ten command and copy clocks variables - All dementia patients versus HC participants

| Variable   | All (n=231)   | Dementia (n=56) | HC (n=175)    | t-value  | df     | p-value           |
|--|---------------|-----------------|---------------|----------|--------|-------------------|
| <b>1. Mean pressure (command)</b>                        | 95.16 (32.23) | 52.14 (40.95)   | 108.93 (7.54) | 10.322   | 56.197 | <0.0001 *<br>D<HC |
| <b>2. Standard deviation of pressure (command)</b>       | 23.27 (5.87)  | 19.38 (9.00)    | 24.51 (3.68)  | 4.1556   | 60.976 | 0.0001 *<br>D<HC  |
| <b>3. Mean pressure (copy)</b>                           | 97.61 (29.99) | 57.31 (38.02)   | 110.51 (6.46) | 10.425   | 56.021 | <0.0001 *<br>D<HC |
| <b>4. Average pressure/velocity (command)</b>            | 3.03 (1.82)   | 2.90 (3.02)     | 3.07 (1.21)   | 0.42607  | 60.738 | 0.6716<br>ns      |
| <b>5. Average pressure/velocity (copy)</b>               | 3.33 (2.03)   | 3.30 (3.51)     | 3.35 (1.24)   | 0.10497  | 59.47  | 0.9168<br>ns      |
| <b>6. Post clockface latency (PCFL) (copy)</b>           | 1.50 (1.48)   | 2.23 (2.57)     | 1.27 (0.76)   | -2.7643  | 58.095 | 0.0076<br>D>HC    |
| <b>7. Minute hand difference from ideal angle (copy)</b> | -2.19 (12.30) | -0.34 (20.96)   | -2.79 (7.73)  | -8.85617 | 59.861 | 0.3953<br>ns      |
| <b>8. Hour hand distance from center (copy)</b>          | 3.26 (2.51)   | 3.98 (2.58)     | 3.04 (2.45)   | -2.4081  | 88.833 | 0.0181<br>D>HC    |
| <b>9. Total completion time (copy)</b>                   | 30.86 (14.43) | 42.50 (21.68)   | 27.14 (8.32)  | -5.1806  | 60.265 | <0.0001 *<br>D>HC |
| <b>10. Minute hand distance from center (command)</b>    | 4.51 (4.56)   | 6.12 (4.97)     | 4.03 (4.33)   | -2.7389  | 75.445 | 0.0077<br>D>HC    |

means and standard deviations;

\* signifies variables that remain significant after Bonferroni correction

**Table 7.**

Ranking of the top ten command and copy clocks variables comparing AD and VaD participants

| Variable   | AD (n=29)     | VaD (n=27)    | t-value | Df     | p-value            |
|--|---------------|---------------|---------|--------|--------------------|
| <b>1. Average pressure (command)</b>               | 21.57 (31.56) | 84.98 (17.66) | 9.3618  | 44.575 | <0.0001*<br>AD<VaD |
| <b>2. Average pressure (copy)</b>                  | 30.23 (29.89) | 86.39 (19.99) | 8.3175  | 49.15  | <0.0001*<br>AD<VaD |
| <b>3. Average pressure/velocity (copy)</b>         | 1.26 (1.31)   | 5.48 (3.83)   | 5.438   | 31.606 | <0.0001*<br>AD<VaD |
| <b>4. Average pressure/velocity (command)</b>      | 0.92 (1.28)   | 5.01 (2.93)   | 6.6897  | 34.974 | <0.0001*<br>AD<VaD |
| <b>5. Average digit height (command)</b>           | 5.67 (1.47)   | 5.15 (1.33)   | -1.385  | 53.959 | 0.1718<br>ns       |
| <b>6. Total drawing time (copy)</b>                | 14.36 (6.68)  | 20.73 (9.32)  | 2.922   | 46.872 | 0.0053<br>AD<VaD   |
| <b>7. Standard deviation of pressure (copy)</b>    | 18.36 (7.18)  | 23.63 (4.27)  | 3.3576  | 46.153 | 0.0016*<br>AD<VaD  |
| <b>8. Clockface overshoot angle (command)</b>      | 2.29 (1.50)   | 3.17 (3.05)   | 1.4703  | 40.065 | 0.1828<br>Ns       |
| <b>9. Hour hand distance from center (command)</b> | 5.85 (4.79)   | 7.25 (6.00)   | 0.9462  | 47.838 | 0.3488<br>Ns       |
| <b>10. Hour hand distance from center (copy)</b>   | 3.37 (2.41)   | 4.63 (2.64)   | 1.861   | 52.598 | 0.0683<br>ns       |

mean and standard deviation;

\* signifies variables that remain significant after Bonferroni correction