

A new assemblage of plant mesofossils (late Turonian–middle Santonian; Upper Cretaceous) from the Tamagawa Formation, Kuji Group, in northeastern Japan

MASAMICHI TAKAHASHI¹, PATRICK S. HERENDSEEN², FABIANY HERRERA², REN HIRAYAMA³, HISAO ANDO⁴, KAZUHISA SASAKI⁵ AND PETER R. CRANE^{6,7}

¹Department of Environmental Sciences, Niigata University, 8050, 2-cho, Ikarashi, Nishi-ku, Niigata 950-2181, Japan
(e-mail: masamichi@env.sc.niigata-u.ac.jp)

²Chicago Botanic Garden, 1000 Lake Cook Road, Glencoe, Illinois 60022, USA

³School of International Liberal Studies, Waseda University, 1-7-14, Nishiwaseda, Shinjuku-ku, Tokyo 169-0051, Japan

⁴Department of Earth Sciences, Faculty of Science, Ibaraki University, 2-1-1, Bunkyo, Mito 310-8512, Japan

⁵Kuji Amber Museum, 19-156-133, Kokujicho, Kuji, Iwate 028-0071, Japan

⁶Yale School of the Environment, Yale University, 195, Prospect Street, New Haven, Connecticut 06511, USA

⁷Oak Spring Garden Foundation, 1776, Loughborough Lane, Upperville, Virginia 20184, USA

Received November 14, 2019; Revised manuscript accepted March 4, 2020

Abstract. A preliminary description is provided of a new assemblage of small, three-dimensional and charcoalified mesofossils from the Tamagawa Formation (late Turonian–middle Santonian; Upper Cretaceous) of the Kuji Group in northeastern Japan. The new mesofossils yield excellent structural details and include well-preserved circinate shoots of ferns together with conifer leafy-shoots, seeds and probable pollen cones, and variety of angiosperm fruits and seeds, including fruits of Cornales and seeds of Nymphaeales. The new mesofossil assemblage is complementary to the previously published macrofossil flora from the Kuji Group.

Key words: charcoalified, Cretaceous, Kuji Group, late Turonian–middle Santonian, mesofossils, Tamagawa Formation

Introduction

Since the early classic papers by Friis and Skarby (1981, 1982), numerous paleobotanical studies of small three-dimensionally preserved flowers, fruits and seeds (mesofossils) recovered by bulk sieving techniques from Cretaceous sediments, have greatly enhanced our understanding of the pattern of angiosperm diversification through the first 70 million years of their evolutionary history. The bulk of the angiosperm mesofossil record has been recovered from Cretaceous sediments collected in eastern North America, central Portugal, and southern Sweden (Atkinson *et al.*, 2018; Friis and Pedersen, 2012; Friis *et al.*, 2013a, b, 2015, 2017a, b, 2018a, b, c, d, e, 2019; Herendsean *et al.*, 2017; Martinez *et al.*, 2016; Mendes and Friis, 2018; Mendes *et al.*, 2014; Schönenberger *et al.*, 2012). In contrast, the mesofossil record from Asia is much less extensive, and so far angiosperm mesofossils have been recovered only from Kazakhstan and Japan (Frumin and Friis, 1996, 1999; Frumin *et al.*,

2004; Takahashi *et al.*, 1999a, b, 2001a, b, 2002, 2007, 2008a, b, 2014, 2017).

In Japan, current knowledge of Cretaceous angiosperms is derived from a variety of permineralized angiosperm flowers and fruiting structures recovered from the Upper Cretaceous (Coniacian–Santonian) Yezo Group of Hokkaido (Nishida, 1985, 1991, 1994; Nishida and Nishida, 1988; Nishida *et al.*, 1996). In addition, a mesofossil assemblage from the Kamikitaba locality (Ashizawa Formation, early Coniacian, *ca.* 89 Ma) of the Futaba Group of northeastern Honshu in Japan preserves a mesofossil flora that includes three-dimensional angiosperm flowers, as well as several kinds of angiosperm fruits and seeds. The fossils recovered include flowers or fruits of *Futabanthus* (Annonaceae, Takahashi *et al.*, 2008b), *Esgueiria* (Combretaceae, Takahashi *et al.*, 1999b), *Hironoia* (Cornaceae, Takahashi *et al.*, 2002), *Archaeofagacea* (Fagaceae, Takahashi *et al.*, 2008a), *Lauranthus*, *Microlaurus* (Lauraceae, Takahashi *et al.*, 2001a, 2014), and *Archaeostella* (Trochodendraceae, Takahashi *et al.*, 2017).

Fossil seeds assigned to *Symphaenale* (Nymphaeaceae) have also been recovered from another mesofossil assemblage, Gokurakuzawa locality, (Tamayama Formation, Santonian, ca. 85 Ma) in the Futaba Group (Takahashi *et al.*, 2007).

In this study, a preliminary description is provided of a new assemblage of plant mesofossils from sediments of the Kuji Group exposed about 350 km north of the Futaba Group along the Pacific coast of northeastern Japan that includes small, three-dimensional and charcoaled mesofossils.

The Kuji Group has yielded abundant fossil leaves (Tanai, 1979) from the Tamagawa and Sawayama formations, and has yielded a palynoflora similar to that from the Futaba Group, but with the important addition of *Aquilapollenites* pollen in the Kuji assemblages (Miki, 1972, 1977; Umetsu and Kurita, 2007). Takahashi *et al.* (2001b) have described fossil megaspores from the Tamagawa Formation. The new mesofossils described here include well-preserved circinate coiled fern shoots (fiddleheads), conifer leafy-shoots, seeds and pollen cones, and a variety of angiosperm fruits and seeds. This preliminary report highlights the potential for future discoveries of informative fossil material, including well-preserved angiosperm flowers, from the Tamagawa Formation of the Kuji Group exposed in a part of Japan where sediments potentially rich in mesofossils have not been fully explored.

Materials and methods

The Kuji Group includes the Tamagawa, Kunitan and Sawayama formations in ascending order (Ando, 1997). Arimoto *et al.* (2018) suggested that the age of the Kuji Group ranges from the middle Turonian to the middle Campanian based on the $\delta^{13}\text{C}$ wood chemostratigraphy and U-Pb radiometric analysis of intercalating tuff layers. The Tamagawa Formation, the lowermost of the three formations in the Kuji Group, consists of a non-marine and shallow marine sedimentary succession, characterized by alluvial fan conglomerates with lacustrine mudstones, fluvial sandstone and mudstone, and estuary to shoreface sandstones (Ando, 1997). Outcrops of the Tamagawa Formation comprise a sequence of sandy to subordinatedly silty clastic sediments deposited in fluvial to shallow-marine environments (Ando, 1997; Umetsu and Kurita, 2007; Umetsu *et al.*, 2013). The deposits are regarded as of middle Turonian–middle Campanian age (from 92.6 ± 0.6 Ma to 79.1 ± 1.2 Ma) based on the U-Pb radiometric dating of intercalating tuff layers and carbon isotope chemostratigraphy (Arimoto *et al.*, 2018).

The plant mesofossil assemblage was isolated from a black carbonaceous, poorly-sorted sandy mudstone from

the upper part of the Tamagawa Formation that is exposed along the sea cliff at Tamagawa, Noda village, Iwate Prefecture, in northeastern Honshu (lat. $40^\circ 5' 12.1776''$ N, long. $141^\circ 49' 42.8082''$ E). The samples were collected by MT in 2016–2018 from a horizon corresponding to the carbonaceous mudstone recorded at 230 m at the Tamagawa section (column 1 of fig. 3 of Arimoto *et al.*, 2018). This layer is broadly correlated to a vertebrate bone bed in the Osawada section, that includes well-preserved terrestrial fossils of turtles (*Adocus*), crocodylomorphs, and dinosaurs (Hirayama *et al.*, 2010; Umetsu *et al.*, 2013; column 2 on fig. 3 of Arimoto *et al.*, 2018). Arimoto *et al.* (2018) estimated the depositional age of the dinosaur-bearing bone bed of the Tamagawa Formation in the Kuji Group to be the early - middle Santonian (85.9 ± 0.7 Ma). On the other hand, Hirayama *et al.* (2021) estimated the age of the same bone bed of the Tamagawa Formation to be the late Turonian (90.51 ± 0.54 Ma) based on U-Pb dating for zircon grains of an intercalating tuff bed.

Plant mesofossils were isolated by sieving from unconsolidated sediment samples. The samples were disaggregated in water, then cleaned and rinsed using standard techniques (e.g., Friis *et al.*, 2011; Takahashi *et al.*, 1999a). The mesofossil specimens were examined with a Hitachi S-4300 field emission SEM. Some specimens were photographed (LM) using a Canon Rebel camera with 100 mm macrolens attached to a stack-shot system, and digital images were merged using Helicon Focus software.

The specimens illustrated in this article are deposited in the paleobotanical collections of the National Science Museum (NSM-PP), Tsukuba, 305-0005 Japan.

Results

Ferns

Ferns are represented in the collection by about seven specimens of charcoaled, tightly curled, circinate shoots (fiddleheads), approximately 2–4 mm in diameter. The shoots have trichomes (Figure 1A). The systematic affinities of the fiddleheads are uncertain, although the specimens are similar in size and trichomes to the fiddleheads of the filmy fern, *Hymenophyllum iwatsukii* (Hymenophyllaceae) from the Khukhteg Formation (Aptian–Albian; Lower Cretaceous) at Tevshii Govi in Mongolia (Herrera *et al.*, 2017). Tanai (1979) reported megafossils of fronds of Osmundaceae, Schizaeaceae, and Gleicheniaceae from the Tamagawa Formation of the Kuji Group. Other pteridophytes recovered from the Tamagawa Formation include abundant megaspores of *Molaspora* (Marsileales), *Erlansonisporites* (Selaginellales), as well as forms assigned to *Verrutrilletes*, *Bacutri-*




Figure 1. Mesofossils from the Tamagawa Formation (Kuji Group). A, Charcoalified circinnately curled fern fronds (fiddleheads) showing pinnules; Inset, trichomes, NSM-PP 12140; B, Conifer leafy shoot with terminal buds, NSM-PP 12141; C, Seed cone type 1, NSM-PP 12142; D, Seed cone type 2, NSM-PP 12143; E, cf. *Hironoia fusiformis* Takahashi *et al.* (Cornales), NSM-PP 12144; F, Fruit (Endocarp), lateral view of specimen with leathery outer layer (exocarp) missing, showing reticulate surface of the inner sclerotic layer, NHM-PP 12145; G–I, cf. *Symphaenale futabensis* Takahashi *et al.* (Nymphaeales), NHM-PP 12146; G, Seed in lateral view showing micropyle hilum and raphe; H, Detail of micropyle (M) and raphe (R); I, Seed surface showing the digitate outlines of the columnar sclerenchyma that comprise the exotesta sclerenchyma.

letes and *Trileites* (Takahashi *et al.*, 2001b).

Conifers

Conifers are abundant in the assemblage and are represented by about 30 leafy shoots with terminal vegetative buds, as well as pollen cones, seed cones and seeds (Figure 1B–D). The leafy shoots have spirally arranged acicular scale-like leaves with acute apices (Figure 1B), and terminal rounded shoot apices (Figure 1B). The buds range from 2.0–2.2 mm in width, and 1.3–1.8 mm in length. The leafy buds appear similar to those present in Cupressaceae (Schulz *et al.*, 2005).

Seed cone type 1 is represented in the collection by a single specimen (Figure 1C). The cone is ovoid-oblong, 2.0–2.3 mm long, 1.5 mm wide, with numerous helically arranged bract-scale complexes. The complexes are linear-lanceolate, 0.5–0.6 mm long and 0.2–0.3 mm wide. At the base the scales are subtruncate (Figure 1C). Apices of the bract-scale complexes are abraded and poorly preserved. The seeds are unknown.

Seed cone type 2 is represented in the collection by a single specimen (Figure 1D). The seed cone type 2 is elongate-cylindric, 9.5 mm long, 6.0 mm wide, with 12–18 helically arranged tightly, persistent and imbricated bract-scale complexes. The apices of the ovuliferous scales are rhomboidal, 2.2–3.8 mm high and 3.8–5.0 mm wide, with an truncate apex (ovuliferous scale umbo). The specimen shows a peduncle that bears only small, thick, rudimentary bracts. The systematic affinity of the female cone is likely to the Cupressaceae (Schulz *et al.*, 2005). A permineralized conifer seed cone, *Archicupressus*, has been recovered from the Yezo Group (Coniacian – Santonian) in Hokkaido (Ohsawa *et al.*, 1992). *Archicupressus* is ellipsoid, 3.8 cm long and 1.7 cm in diameter, consisting of more than 12 bract-scale complexes. The permineralized seed cone from the Yezo Group is much larger than the seed cone type 2 from the Kuji Group.

Angiosperms

A preliminary survey of the angiosperms represented in the mesofossil assemblage has recovered five kinds of fruits and a single kind of dispersed seed.

Fruit type 1 is represented in the collection by 20 specimens (Figure 1E). The fruits are globose to elliptical, 4.0–4.2 mm long and 2.2–2.5 mm wide. The endocarps are trimerous, consisting of three fused carpels. The surface of the endocarp varies from smooth to corrugated. Each locule opens by a single dorsal valve that often becomes detached first at the apex of the endocarp (Figure 1E). The endocarp tapers gradually toward the apex and narrows at the base into a slender stalk. Fruit type 1 is similar to *Hironoia fusiformis* Takahashi *et al.* (Cornales) from the Kamikitaba assemblage (Takahashi *et al.*, 2002).

Fruit type 2 (endocarp) is represented in the collection by two specimens (Figure 1F). Both are flattened woody structures, *ca.* 1.7–2.0 mm in diameter. The outer layer is missing and exposes an inner sclerotic layer with a distinctive reticulate surface (Figure 1F). This type of endocarp is recovered frequently from the Kamikitaba assemblage (Takahashi *et al.*, 1999a, fig. 9A, B). The systematic affinity of these fruits is unknown. However, they are similar in general form to specimens described as *Sabia* (Sabiaceae) (Knobloch and Mai, 1986; Bodor *et al.*, 2012) or *Porosia* (Rutaceae) (Manchester and Kodrul, 2014).

Seed type 1 is represented in the collection by a single specimen (Figure 1G–I). The seed is anatropous, ovoid, and 1.2–1.5 mm long. The raphe enters the seed obliquely at a hilum that is adjacent to, but separated from the micropyle. The micropyle has a distinct plug with a papillate surface. The exotesta is composed of columnar palisade sclerenchyma cells with anticlinal walls that are strongly undulate. This seed is similar to *Symphaenale futabensis* Takahashi *et al.* (Nymphaeales) from the Tamayama Formation of the Futaba Group (Takahashi *et al.*, 2007).

Discussion

The fossil flora of the Kuji Group

The Kuji Group has yielded abundant plant macrofossils preserved as impressions, and a comprehensive account of the leaf flora documented pteridophytes (*Gleichenites*, *Asplenium*, *Sachalinia*, Osmundaceae, Schizaeaceae, Salviniaceae and *Cladophlebis*), gymnosperms (*Araucarites*, *Taxodium* and *Glyptostrobus*, *Metasequoia*, Cycadopsida, Nilssoniaceae, Cupressaceae), and angiosperms (Magnoliaceae, Lauraceae, Menispermaceae, Trochodendraceae, Platanaceae, Fagaceae, Dilleniaceae, Salicaceae, Sapindaceae, Vitaceae, Rhamnaceae, and Trapaceae [Tanai, 1979]). Only a subset of these plants occurs in the Tamagawa flora, from which 11 pteridophyte, six conifer and ten angiosperm species have been described (Tanai, 1979).

Miki (1977) showed that palynomorphs from the Tamagawa and Sawayama formations consist of 56% pteridophyte spores dominated by Schizaeaceous spores, 20% gymnosperm pollen dominated by Pinaceous winged and Taxodiaceous inaperturate pollen, and 26% angiosperm pollen dominated by tricolpate type and *Aquillapollenites* group. Umetsu and Kurita (2007) showed that gymnosperm pollen dominates the palynoflora of the upper part of the Tamagawa Formation (54–68%) with frequent *Taxodiaceapollenites* and coniferous winged pollen. Pollen of the *Aquillapollenites* group also occurs frequently from the upper part of the Tamagawa Formation (Umetsu and Kurita, 2007).

Takahashi *et al.* (2001b) described an assemblage of dispersed megaspores of Marsileales and Selaginellales from the Tamagawa Formation. The well-preserved mesofossil assemblage recovered in the present study demonstrates that the flora of the Tamagawa Formation is dominated by conifers, but also includes a diversity of pteridophytes and angiosperms. The mesofossil assemblage is complementary to the macrofossil and microfossil floras. Results from the present preliminary study suggest that further investigation of mesofossil floras may help to elucidate the early evolution of flowering plants in northeastern Asia. A possibility of special interest would be the recovery of flowers or fruits that could be attributed to the *Aquilapollenites* group, an important and enigmatic group of angiosperms of Late Cretaceous age that experienced significant extinction at the Cretaceous-Paleogene boundary. Finding fossil flowers bearing *Aquilapollenites* pollen would be particularly significant because no mesofossils have yet been documented with this type of pollen.

Comparison between the fossil floras of the Kuji Group and the Futaba Group

The Kuji and Futaba groups are regarded as the western marginal facies of Late Cretaceous forearc basin sediments that are now exposed along the Pacific coast of northeastern Japan (Ando, 2003; Ando and Takahashi, 2017). Their geological sedimentary environments are generally similar, consisting of fluvial, estuarine to shallow-marine sandstone-dominated sedimentary successions, associated with alluvial fan conglomerates and fluvial-marsh carbonaceous mudstone bearing mesofossils and amber (Uchida *et al.*, 2018).

The mesofossil horizons from the lower part of the Ashizawa Formation of the Futaba Group (Kamikitaba locality: Takahashi *et al.*, 1999a) and the upper part of the Tamagawa Formation of the Kuji Group are estimated to be of a similar late Turonian to middle Santonian age. This is also reflected in components of the mesoflora. The cornalean fruits, *Hironoia*, occur in both the Futaba Group and the Kuji Group. In both cases the fossils recovered from the upper part of the Tamagawa Formation (85.9 ± 0.7 Ma: Arimoto *et al.*, 2018; 90.51 ± 0.54 Ma: Hirayama *et al.*, 2021) in the Kuji Group are very similar to those recovered from the Kamikitaba locality (about 89 Ma) in the Futaba Group (Takahashi *et al.*, 2002, 2007; Ando and Takahashi, 2017). The nymphaealean seeds, *Symphaenale*, also occur both in the lower part of the Tamayama Formation of the Futaba Group (Santonian, Gokurakuzawa locality: Takahashi *et al.*, 2007), and the Tamagawa Formation of the Kuji Group.

Based on fossil pollen, Miki (1977) identified 57 species (48%) of pteridophytes, 26 species (22%) of gymnosperms and 37 species (30%) of angiosperms from

the Futaba Group, and suggested that these microfloras are very similar to each other in their floristic composition, except for the *Aquilapollenites* group from the Kuji Group.

Some difference between the two is also shown in the gymnosperm components and their inferred vegetation. Fragments of leafy shoots showing acicular leaves with acute apices are very common in the Kamikitaba assemblage (conifer leaf type 1: Takahashi *et al.*, 1999a), do not occur frequently from the Kuji Group. Additional studies of mesofossil assemblages from the Kuji Group have the potential to provide critical new data toward resolving the history of angiosperm and gymnosperm diversification in eastern Asia during the Late Cretaceous and clarifying biogeographic patterns.

Acknowledgments

We thank Steven R. Manchester for his suggestions and Tatsuo Ushiki and Manabu Hayatsu for permission to use FE-SEM, S-4300.

References

- Ando, H., 1997: Apparent stacking patterns of depositional sequences in the Upper Cretaceous shallow-marine to fluvial successions, Northeast Japan. *Memoir of the Geological Society of Japan*, vol. 48, p. 43–59.
- Ando, H., 2003: Stratigraphic correlation of Upper Cretaceous to Paleocene forearc basin sediments in Northeast Japan: cyclic sedimentation and basin evolution. *Journal of Asian Earth Sciences*, vol. 21, p. 921–935.
- Ando, H. and Takahashi, M., 2017: Reconstruction of the Cretaceous Paleo-Japan continental arc-trench system reconsidered from Cretaceous geologic records in the Japanese Islands. *Fossils (Palaeontological Society of Japan)*, vol. 102, p. 43–62. (in Japanese with English abstract)
- Arimoto, J., Takashima, R., Nishi, H., Yamanaka, T., Orihashi, Y., Jo, S., Yamamoto, K. and Umetsu, K., 2018: Constraining the depositional age of an Upper Cretaceous non-marine and shallow marine siliciclastic succession, Kuji Group, northeastern Japan, based on carbon isotope stratigraphy and U–Pb radiometric dating. *Cretaceous Research*, vol. 92, p. 264–278.
- Atkinson, B. A., Martinez, C. and Crepet, W. L., 2018: Cretaceous asterid evolution: fruits of *Eydeia jerseyensis* sp. nov. (Cornales) from the upper Turonian of eastern North America. *Annals of Botany*, vol. 20, p. 1–10.
- Bodor, E. R., Baranyi, V. and Heřmanova, Z., 2012: The earliest Sabiaceae fruit remains of Hungary. *Hatkeniana*, vol. 7, p. 11–18.
- Friis, E. M., Crane, P. R. and Pedersen, K. R., 2011: *Early Flowers and Angiosperm Evolution*, 600 p. Cambridge University Press, Cambridge.
- Friis, E. M., Crane, P. R. and Pedersen, K. R., 2017a: *Saportanthus*, an extinct genus of Laurales from the Early Cretaceous of Portugal. *International Journal of Plant Sciences*, vol. 178, p. 650–672.
- Friis, E. M., Crane, P. R. and Pedersen, K. R., 2018b: *Tanispermum*, a new genus of distinctive hemi-orthotropous to hemi-anatropous angiosperm seeds from the Early Cretaceous of eastern North America. *American Journal of Botany*, vol. 105, p. 1369–1388.

- Friis, E. M., Crane, P. R. and Pedersen, K. R., 2018c: Fossil seeds with affinities to Austrobaileyales and Nymphaeales from the Early Cretaceous (early-middle Albian) of Virginia and Maryland, U.S.A.: new evidence for extensive extinction near the base of the angiosperm tree. In: Krings, M., Harper, C. J., Cuneo, N. R., Rothwell, G. W. eds., *Transformative Paleobotany: Papers to Commemorate the Life and Legacy of Thomas N. Taylor*, p. 417–435. Elsevier, Amsterdam.
- Friis, E. M., Crane, P. R. and Pedersen, K. R., 2018d: Extinct taxa of exotestal seeds close to Austrobaileyales and Nymphaeales from the Early Cretaceous of Portugal. *Fossil Imprint*, vol. 74, p. 135–158.
- Friis, E. M., Crane, P. R. and Pedersen, K. R., 2018e: *Rightcania* and *Kvacekispermum*: Early Cretaceous seeds from eastern North America and Portugal provide further evidence of the early chloranthoid diversification. *Fossil Imprint*, vol. 74, p. 65–76.
- Friis, E. M., Crane, P. R. and Pedersen, K. R., 2019: Extinct diversity among Early Cretaceous angiosperms: mesofossil evidence of early Magnoliales from Portugal. *International Journal of Plant Sciences*, vol. 180, p. 93–127.
- Friis, E. M., Crane, P. R., Pedersen, K. R., Stampanoni, M. and Marone, F., 2015: Exceptional preservation of tiny embryos documents seed dormancy in early angiosperms. *Nature*, vol. 528, p. 551–554.
- Friis, E. M., Mendes, M. M. and Pedersen, K. R., 2018a: *Paisia*, an Early Cretaceous eudicot angiosperm flower with pantoporate pollen from Portugal. *Grana*, vol. 57, p. 1–15.
- Friis, E. M. and Pedersen, K. R., 2012: *Bertilanthus scanicus*, a new asterid flower from the Late Cretaceous (late Santonian-early Campanian) of Scania, Sweden. *International Journal of Plant Sciences*, vol. 173, p. 318–330.
- Friis, E. M., Pedersen, K. R. and Crane, P. R., 2013a: New diversity among chlamydospermous seeds from the Early Cretaceous of Portugal and North America. *International Journal of Plant Sciences*, vol. 174, p. 530–558.
- Friis, E. M., Pedersen, K. R. and Crane, P. R., 2017b: *Kenilanthus*, a new eudicot flower with tricolpate pollen from the Early Cretaceous (early-middle Albian) of eastern North America. *Grana*, vol. 56, p. 161–173.
- Friis, E. M., Pedersen, K. R. and Endress, P. K., 2013b: Floral structure of extant *Quintinia* (Paracryphiales, campanulids) compared with the Late Cretaceous *Silvianthemum* and *Bertilanthus*. *International Journal of Plant Sciences*, vol. 174, p. 647–664.
- Friis, E. M. and Skarby, A., 1981: Structurally preserved angiosperm flowers from the Upper Cretaceous of southern Sweden. *Nature*, vol. 291, p. 485–486.
- Friis, E. M. and Skarby, A., 1982: *Scandianthus* gen. nov. angiosperm flowers of saxifragalean affinity from the Upper Cretaceous of southern Sweden. *Annals of Botany*, vol. 50, p. 569–583.
- Frumin, S., Eklund, H. and Friis, E. M., 2004: *Mauldinia hirsuta* sp. nov., a new member of the extinct genus *Mauldinia* (Lauraceae) from the Late Cretaceous (Cenomanian-Turonian) of Kazakhstan. *International Journal of Plant Science*, vol. 165, p. 883–895.
- Frumin, S. and Friis, E. M., 1996: Liriodendroid seeds from the Late Cretaceous of Kazakhstan and North Carolina, USA. *Review of Palaeobotany and Palynology*, vol. 94, p. 39–55.
- Frumin, S. and Friis, E. M., 1999: Magnoliid reproductive organs from the Cenomanian-Turonian of north-western Kazakhstan: Magnoliaceae and Illiciaceae. *Plant Systematics and Evolution*, vol. 216, p. 265–288.
- Herendeen, P. S., Friis, E. M., Pedersen, K. R. and Crane, P. R., 2017: Palaeobotanical redux: revisiting the age of the angiosperms. *Nature Plants*, vol. 3, p. 1–8.
- Herrera, F., Moran, R. C., Shi, G., Ichinnorov, N., Takahashi, M., Crane, P. R. and Herendeen, P. S., 2017: An exquisitely preserved filmy fern (Hymenophyllaceae) from the Early Cretaceous of Mongolia. *American Journal of Botany*, vol. 104, p. 1370–1381.
- Hirayama, R., Kobayashi, Y., Sonoda, T. and Sasaki, K., 2010: Preliminary report of terrestrial vertebrates from the Late Cretaceous Tamagawa Formation of the Kuji Group of Kuji City, Iwate Prefecture, northeastern Japan. *Journal of Fossil Research*, vol. 42, p. 74–82. (in Japanese with English abstract)
- Hirayama, R., Sonoda, T., Uno, H., Tsutsumi, Y., Sasaki, K. and Takisawa, T., 2021: *Adocus kohaku*, a new species of aquatic turtle (Testudines: Cryptodira: Adocidae) from the Late Cretaceous of Kuji, Iwate Prefecture, northeast Japan. *International Journal of Paleobiology & Paleontology*, vol. 4, p. 1–12.
- Knobloch, E. and Mai, D. H., 1986: Monographie der Früchte und Samen in der Kreide von Mitteleuropa. *Rozprawy Źródłowe geologicznego, vol. 47, p. 1–219.*
- Manchester, S. R. and Kodrul, T. M., 2014: Morphology, affinities and phytogeographic history of *Porosia Hickey* in the Cretaceous and Paleocene of North America and Asia. *Acta Palaeobotanica*, vol. 54, p. 77–99.
- Martinez, C., Choo, T. Y. S., Allevato, D., Nixon, K. C., Crepet, W. L., Harbert, R. S. and Daghljan, C. P., 2016: *Rariglanda jerseyensis*, a new ericalean fossil flower from the Late Cretaceous of New Jersey. *Botany*, vol. 94, p. 747–758.
- Mendes, M. M. and Friis, E. M., 2018: The Nossa Senhora da Luz flora from the Early Cretaceous (early Aptian-late Albian) of Juncal in the western Portuguese Basin. *Acta Palaeobotanica*, vol. 58, p. 159–174.
- Mendes, M. M., Grimm, G. W., Pais, J. and Friis, E. M., 2014: Fossil *Kajanthus lusitanicus* gen. et sp. nov. from Portugal: Floral evidence for Early Cretaceous Lardizabalaceae (Ranunculales, basal eudicot). *Grana*, vol. 53, p. 283–301.
- Miki, A., 1972: Palynological study of the Kuji Group in northeastern Honshu, Japan. *Journal of the Faculty of Science, Hokkaido University, Series 4, Geology and Mineralogy*, vol. 15, p. 513–604.
- Miki, A., 1977: Late Cretaceous pollen and spore floras of northern Japan: composition and interpretation. *Journal of the Faculty of Science, Hokkaido University, Series 4, Geology and Mineralogy*, vol. 17, p. 399–436.
- Nishida, H., 1985: A structurally preserved magnolialean fructification from the mid-Cretaceous of Japan. *Nature*, vol. 318, p. 58–59.
- Nishida, H., 1991: Diversity and significance of Late Cretaceous permineralized plant remains from Hokkaido, Japan. *Botanical Magazine, Tokyo*, vol. 104, p. 253–273.
- Nishida, H., 1994: *Elsemaria*, a Late Cretaceous angiosperm fructification from Hokkaido, Japan. *Plant Systematics and Evolution, Supplement*, vol. 8, p. 123–135.
- Nishida, H. and Nishida, M., 1988: *Protomonimia kasai-nakajhongii* gen. et sp. nov.: A permineralized magnolialean fructification from the mid-Cretaceous of Japan. *Botanical Magazine, Tokyo*, vol. 101, p. 397–437.
- Nishida, M., Ohsawa, T., Nishida, H., Yoshida, A. and Kanie, Y., 1996: A permineralized magnolialean fructification from the Upper Cretaceous of Hokkaido, Japan. *Research Institute of Evolutionary Biology, Scientific Report*, vol. 8, p. 19–30.
- Ohsawa, T., Nishida, H. and Nishida, M., 1992: Structure and affinities of the petrified plants from the Cretaceous of northern Japan and Saghalien, XI, A cupressoid seed cone from the Upper Cretaceous of Hokkaido. *Botanical Magazine, Tokyo*, vol. 105, p. 125–133.
- Schöenberger, J., von Balthazar, M., Takahashi, M., Xiao, X., Crane, P. R. and Herendeen, P. S., 2012: *Glandulocalyx upatoiensis*, a fossil flower of Ericales (Actinidiaceae/ Clethraceae) from the Late Cretaceous (Santonian) of Georgia, USA. *Annals of Botany*, vol.

- 109, p. 921–936.
- Schulz, C., Knopf, P. and Stützel, T., 2005: Identification key to the Cypress family (Cupressaceae). *Feddes Repertorium*, vol. 116, p. 96–146.
- Takahashi, M., Crane, P. R. and Ando, H., 1999a: Fossil flowers and associated plant fossils from the Kamikitaba locality (Ashizawa Formation, Futaba Group, lower Coniacian, Upper Cretaceous) of Northeast Japan. *Journal of Plant Research*, vol. 112, p. 187–206.
- Takahashi, M., Crane, P. R. and Ando, H., 1999b: *Esgueiria futabaensis* sp. nov., a new angiosperm flower from the Upper Cretaceous (lower Coniacian) of northeastern Honshu, Japan. *Paleontological Research*, vol. 3, p. 81–87.
- Takahashi, M., Crane, P. R. and Ando, H., 2001b: Fossil megaspores of Marsileales and Selaginellales from the Santonian (Upper Cretaceous) of the Tamagawa Formation (Kuji Group) in Northeast Japan. *International Journal of Plant Sciences*, vol. 162, p. 431–439.
- Takahashi, M., Crane, P. R. and Manchester, S. R., 2002: *Hironoia fusiformis* gen. et sp. nov.; a cornalean fruit from the Kamikitaba locality (Upper Cretaceous, Lower Coniacian) in northeastern Japan. *Journal of Plant Research*, vol. 115, p. 463–473.
- Takahashi, M., Friis, E. M. and Crane, P. R., 2007: Fossil seeds of Nymphaeales from the Tamayama Formation (Futaba Group), Upper Cretaceous (early Santonian) of northeastern Honshu, Japan. *International Journal of Plant Sciences*, vol. 168, p. 341–350.
- Takahashi, M., Friis, E. M., Herendeen, P. S. and Crane, P. R., 2008a: Fossil flowers of Fagales from the Kamikitaba locality (early Coniacian; Late Cretaceous) of northeastern Japan. *International Journal of Plant Sciences*, vol. 169, p. 899–907.
- Takahashi, M., Friis, E. M., Uesugi K., Suzuki, Y. and Crane, P. R., 2008b: Floral evidence of Annonaceae from the Late Cretaceous of Japan. *International Journal of Plant Sciences*, vol. 169, p. 908–917.
- Takahashi, M., Herendeen, P. S. and Crane, P. R., 2001a: Lauraceous fossil flower from the Kamikitaba locality (lower Coniacian; Upper Cretaceous) in northeastern Japan. *Journal of Plant Research*, vol. 114, p. 429–434.
- Takahashi, M., Herendeen, P. S. and Xiao, X., 2017: Two early eudicot fossil flowers from the Kamikitaba assemblage (Coniacian, Late Cretaceous) in northeastern Japan. *Journal of Plant Research*, vol. 130, p. 809–826.
- Takahashi, M., Herendeen, P. S., Xiao, X. and Crane, P. R., 2014: Lauraceous fossil flowers from the Kamikitaba assemblage (Coniacian, Late Cretaceous) of northeastern Japan. *Systematic Botany*, vol. 39, p. 715–724.
- Tanai, T., 1979: Late Cretaceous floras from the Kuji district, northeastern Honshu, Japan. *Journal of the Faculty of Science, Hokkaido University, Series 4, Geology and Mineralogy*, vol. 19, p. 75–136.
- Uchida, T., Koshino, H., Takahashi, S., Shimizu, E., Takahashi, H., Yoshida, J., Shinden, H., Tsujimura, M., Kofujita, H., Uesugi, S. and Kimura, K., 2018: Ca^{2+} -signal transduction inhibitors, kujiol A and kujigamberol B, isolated from Kuji amber using a mutant yeast. *Journal of Natural Products*, vol. 81, p. 1070–1074.
- Umetsu, K., Hirayama, R., Sonoda, T. and Takashima, R., 2013: Marine and nonmarine deposits of the Cretaceous Miyako and Kuji groups and Late Cretaceous terrestrial vertebrates in Iwate Prefecture, northeast Japan. *Journal of the Geological Society of Japan, Supplement*, vol. 119, p. 82–95. (in Japanese with English abstract)
- Umetsu, K. and Kurita, Y., 2007: Palynostratigraphy and age of the Upper Cretaceous Kuji Group, northeast Iwate Prefecture, Northeast Japan. *Journal of the Japanese Association for Petroleum Technology*, vol. 72, p. 215–223. (in Japanese with English abstract)

Author contributions

M. T. initiated the study and carried out all process of the study, such as collecting samples, cleaning, bulk sieving process, sorting, and SEM work. P. H. and F. H. have contributed to the taxonomic aspects. R. H and H. A are responsible for the geological aspects. K. S. has contributed to the field work and collecting samples. All authors contributed to the writing of the paper.