



Genome Sequence of *Synechococcus* sp. Strain LA31, Isolated from a Temperate Estuary

 Joshua D. Kling,^a  Eric A. Webb,^a  David A. Hutchins^a

^aDepartment of Biological Sciences, University of Southern California, Los Angeles, California, USA

ABSTRACT Cluster 5 *Synechococcus* species are widely acknowledged for their broad distribution and biogeochemical importance. In particular, subcluster 5.2 strains inhabit freshwater, estuarine, and marine environments but are understudied, compared to other subclusters. Here, we present the genome for *Synechococcus* sp. strain LA31, a strain that was recently isolated from Narragansett Bay, Rhode Island, USA.

Unicellular picocyanobacteria from the genus *Synechococcus* are found throughout aquatic environments (1–3) and play a key role in global carbon cycling (4). However, the biodiversity and biogeochemical contributions of nonmarine *Synechococcus* strains are only beginning to be understood (3). Here, we present the genome of *Synechococcus* sp. strain LA31, which was recently isolated from a temperate estuary.

LA31 was isolated from the Narragansett Bay Time Series in Rhode Island, USA (latitude, 41.47; longitude, −71.40). To enrich for phytoplankton, surface water (22°C; salinity, 28.84 psu) was spiked with nutrients comparable to F/40 medium for 10 days (5) and given 150 $\mu\text{mol photons/m}^{-2} \cdot \text{s}^{-1}$ of light (12-h light/12-h dark cycle). Single cells were sorted into 96-well plates using an Influx flow cytometer (BD, San Jose, CA, USA) and were maintained under the same conditions. Wells showing growth after 2 weeks were streaked onto F/2 agar plates for isolation, and one colony was transferred to F/2 medium (5). DNA was extracted using a series of freeze-thaw cycles in N_2 , proteinase K incubations, and final extraction with the Qiagen DNeasy PowerBiofilm kit (Hilden, Germany), as described previously (6). MR DNA (Shallowater, TX, USA) performed library preparation (SMRTbell Express template preparation kit v2.0; Pacific Biosciences), sequencing (Sequel system), read quality control, and assembly (Hierarchical Genome Assembly Process [HGAP] in single-molecule real-time [SMRT] Analysis v9.0). Assembly completeness and contamination were measured with CheckM v1.1.3 (7). Gene calling and annotation were done with Prodigal v2.6.3 (8) and KofamScan v1.3.0 (9). antiSMASH v6.0 (10) was used for detection of secondary metabolite genes, tRNAscan-SE v2.0 for detection of tRNA sequences (11), and Barrnap v0.9 for detection of rRNA sequences (12). A phylogenetic tree of all unicellular picocyanobacterial assemblies available in the NCBI RefSeq database (13) was constructed using GToTree v1.5.51 with the included cyanobacterial marker gene set (14–18). Default parameters were used for all software unless otherwise noted.

A total of 1,209,177 reads (average length, 3,532.17 bp) were assembled into a single contig 2,752,051 bp in length (GC content, 63%; completeness, 99.46%; contamination, 0.54%). A total of 2,965 coding regions were detected (1,501 annotated), with 47 tRNAs and three 5S, 16S, and 23S rRNA genes. A phylogenetic tree of 251 conserved proteins from 219 genomes placed LA31 in *Synechococcus* subcluster 5.2 (Fig. 1), branching with both brackish (CB0101 [19]) and freshwater (*Vulcanococcus limneticus* LL [20]) isolates.

Nitrogen transporters for nitrate/nitrite (*NRT* [$n = 1$]), ammonium (*AMT* [$n = 2$]), urea (*urtABCDE* and *Dur3*), and 15 amino acids (21) were detected in LA31. A purine/

Editor Frank J. Stewart, Montana State University

Copyright © 2022 Kling et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Joshua D. Kling, joshuakl@usc.edu, or David A. Hutchins, dahutch@usc.edu.

The authors declare no conflict of interest.

Received 30 July 2021

Accepted 9 November 2021




FIG 1 Maximum likelihood tree created with 251 concatenated amino acid sequences found in all currently available cluster 5 *Synechococcus*, *Cyanobium*, *Vulcanococcus*, and *Prochlorococcus* genomes. The distribution of the three subclusters within cluster 5 is indicated with brackets on the right side. Strain LA31 (presented here) is shown in bold. *Gloeobacter violaceus* PCC 7421 is included as an outgroup. Node numbers represent local support values derived from 1,000 resamplings as part of FastTree. Genbank accession numbers are listed for each genome.

hypoxanthine transporter was also detected; such transporters are upregulated in eukaryotes under N deprivation but have not been reported in cyanobacteria (22). High-affinity phosphate uptake systems (*pstABC* and *pstS* [$n = 3$]) and phosphonate transporters (*phnD* and *phnE*) were detected for P, and Fe transporters were detected for both ferrous (*feoAB*) and ferric (*fbpAB* or *idiAB*) (23). Four toxin-antitoxin gene pairs were detected, similar to closely related CB0101 (24). Thirteen secondary metabolite coding regions were identified for ribosomally synthesized and posttranslationally modified peptides (RiPPs) ($n = 10$), terpene synthesis ($n = 2$), and hieridrin B ($n = 1$), a potential antimalarial compound (25).

Data availability. This assembly is available under GenBank accession number [CP075523](#) and RefSeq assembly accession number [GCF_018502385.1](#). The version described is the first version. Reads are available under SRA accession number [SRR14511408](#).

ACKNOWLEDGMENTS

Funding for this work was provided by National Science Foundation grant OCE1638804 to D.A.H. and grant OCE1851222 to D.A.H. and E.A.W.

REFERENCES

- Lee MD, Ahlgren NA, Kling JD, Walworth NG, Rocap G, Saito MA, Hutchins DA, Webb EA. 2019. Marine *Synechococcus* isolates representing globally abundant genomic lineages demonstrate a unique evolutionary path of genome reduction without a decrease in GC content. *Environ Microbiol* 21:1677–1686. <https://doi.org/10.1111/1462-2920.14552>.
- Cabello-Yeves PJ, Haro-Moreno JM, Martin-Cuadrado A-B, Ghai R, Picazo A, Camacho A, Rodríguez-Valera F. 2017. Novel *Synechococcus* genomes reconstructed from freshwater reservoirs. *Front Microbiol* 8:1151. <https://doi.org/10.3389/fmicb.2017.01151>.
- Chen F, Wang K, Kan J, Bachoon DS, Lu J, Lau S, Campbell L. 2004. Phylogenetic diversity of *Synechococcus* in the Chesapeake Bay revealed by ribulose-1,5-bisphosphate carboxylase-oxygenase (RuBisCO) large subunit gene (*rbcL*) sequences. *Aquat Microb Ecol* 36:153–164. <https://doi.org/10.3354/ame036153>.
- Flombaum P, Gallegos JL, Gordillo RA, Rincón J, Zabala LL, Jiao N, Karl DM, Li WKW, Lomas MW, Veneziano D, Vera CS, Vrugt JA, Martiny AC. 2013. Present and future global distributions of the marine cyanobacteria *Prochlorococcus* and *Synechococcus*. *Proc Natl Acad Sci U S A* 110: 9824–9829. <https://doi.org/10.1073/pnas.1307701110>.
- Guillard RRL. 1975. Culture of phytoplankton for feeding marine invertebrates, p 29–60. In Smith WL, Chanley MH (ed), *Culture of marine invertebrate animals*. Plenum Press, New York, NY.
- Florea KM, Thrash JC, Webb EA. 2020. Metagenome-assembled genome sequence of *Dolichospermum circinale* strain Clear-D4, assembled from a harmful cyanobacterial bloom enrichment culture. *Microbiol Resour Announc* 9:e01123-20. <https://doi.org/10.1128/MRA.01123-20>.
- Parks DH, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
- Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. <https://doi.org/10.1186/1471-2105-11-119>.
- Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, Ogata H. 2020. KofamKOALA: KEGG ortholog assignment based on profile HMM and adaptive score threshold. *Bioinformatics* 36:2251–2252. <https://doi.org/10.1093/bioinformatics/btz859>.
- Blin K, Shaw S, Kloosterman AM, Charlop-Powers Z, van Wezel GP, Medema MH, Weber T. 2021. antiSMASH 6.0: improving cluster detection and comparison capabilities. *Nucleic Acids Res* 49:W29–W35. <https://doi.org/10.1093/nar/gkab335>.
- Lowe TM, Chan PP. 2016. tRNAscan-SE On-line: integrating search and context for analysis of transfer RNA genes. *Nucleic Acids Res* 44:W54–W57. <https://doi.org/10.1093/nar/gkw413>.
- Seemann T. 2013. barnap 0.9: rapid ribosomal RNA prediction. <https://github.com/tseemann/barnap>.
- Pruitt KD, Maglott DR. 2001. RefSeq and LocusLink: NCBI gene-centered resources. *Nucleic Acids Res* 29:137–140. <https://doi.org/10.1093/nar/29.1.137>.
- Lee MD. 2019. GToTree: a user-friendly workflow for phylogenomics. *Bioinformatics* 35:4162–4164. <https://doi.org/10.1093/bioinformatics/btz188>.
- Eddy SR. 2011. Accelerated profile HMM searches. *PLoS Comput Biol* 7: e1002195. <https://doi.org/10.1371/journal.pcbi.1002195>.
- Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25:1972–1973. <https://doi.org/10.1093/bioinformatics/btp348>.
- Price MN, Dehal PS, Arkin AP. 2009. FastTree: computing large minimum evolution trees with profiles instead of a distance matrix. *Mol Biol Evol* 26: 1641–1650. <https://doi.org/10.1093/molbev/msp077>.
- Letunic I, Bork P. 2007. Interactive Tree Of Life (iTOL): an online tool for phylogenetic tree display and annotation. *Bioinformatics* 23:127–128. <https://doi.org/10.1093/bioinformatics/btl529>.
- Marsan D, Wommack KE, Ravel J, Chen F. 2014. Draft genome sequence of *Synechococcus* sp. strain CB0101, isolated from the Chesapeake Bay estuary. *Genome Announc* 2:e01111-13. <https://doi.org/10.1128/genomeA.01111-13>.
- Di Cesare A, Cabello-Yeves PJ, Chrismas NAM, Sánchez-Baracaldo P, Salcher MM, Callieri C. 2018. Genome analysis of the freshwater planktonic *Vulcanococcus limneticus* sp. nov. reveals horizontal transfer of nitrogenase operon and alternative pathways of nitrogen utilization. *BMC Genomics* 19:259. <https://doi.org/10.1186/s12864-018-4648-3>.
- Zubkov MV, Fuchs BM, Tarran GA, Burkill PH, Amann R. 2003. High rate of uptake of organic nitrogen compounds by *Prochlorococcus* cyanobacteria as a key to their dominance in oligotrophic oceanic waters. *Appl Environ Microbiol* 69:1299–1304. <https://doi.org/10.1128/AEM.69.2.1299-1304.2003>.
- Terrado R, Monier A, Edgar R, Lovejoy C. 2015. Diversity of nitrogen assimilation pathways among microbial photosynthetic eukaryotes. *J Phycol* 51:490–506. <https://doi.org/10.1111/jpy.12292>.
- Ahlgren NA, Belisle BS, Lee MD. 2020. Genomic mosaicism underlies the adaptation of marine *Synechococcus* ecotypes to distinct oceanic iron niches. *Environ Microbiol* 22:1801–1815. <https://doi.org/10.1111/1462-2920.14893>.
- Marsan D, Place A, Fucich D, Chen F. 2017. Toxin-antitoxin systems in estuarine *Synechococcus* strain CB0101 and their transcriptomic responses to environmental stressors. *Front Microbiol* 8:1213. <https://doi.org/10.3389/fmicb.2017.01213>.
- Costa M, Sampaio-Dias IE, Castelo-Branco R, Scharfenstein H, Rezende de Castro R, Silva A, Schneider MPC, Araújo MJ, Martins R, Domingues VF, Nogueira F, Camões V, Vasconcelos VM, Leão PN. 2019. Structure of hieridins C, synthesis of hieridins B and C, and evidence for prevalent alkylresorcinol biosynthesis in picocyanobacteria. *J Nat Prod* 82:393–402. <https://doi.org/10.1021/acs.jnatprod.8b01038>.