

THE F_p -SELBERG INTEGRAL OF TYPE A_n

RICHARD RIM' ANYI' ¹ AND ALEXANDER VARCHENKO²

^{1,2} *Department of Mathematics, University of North Carolina at Chapel Hill Chapel Hill,
NC 27599-3250, USA*

² *Faculty of Mathematics and Mechanics, Lomonosov Moscow State University
Leninskiye Gory 1, 119991 Moscow GSP-1, Russia*

² *Moscow Center of Fundamental and Applied Mathematics Leninskiye
Gory 1, 119991 Moscow GSP-1, Russia*

Key words: KZ and dynamical equations, reduction modulo p , Selberg integrals, F_p -integrals

2010 Mathematics Subject Classification: 13A35 (33C60, 32G20)

Abstract. We present an F_p -Selberg integral formula of type A_n , in which the F_p -Selberg integral is an element of the finite field F_p , where p is an odd prime. The formula is motivated by analogy between multidimensional hypergeometric solutions of the KZ equations and polynomial solutions of the same equations reduced modulo p . The A_1 -type formula was proved in a previous paper by the authors. The A_2 -type formula is proved in this paper. We also sketch the proof of the A_n -type formula for $n > 2$.

Contents

1.	Introduction	2
2.	Preliminary remarks	3
2.1.	Cancellation of factorials	4
2.2.	Dyson's formula	4
2.3.	F_p -Integrals	4
2.4.	F_p -Beta integral	5
3.	F_p -Selberg integral of type A_n	5
3.1.	Admissible parameters	5
3.2.	Main result	6
3.3.	Relation between the F_p -Selberg integrals of types A_{n-1} - and A_n	7
4.	The A_2 -type Selberg integral over $\mathbb{C}$	9
4.1.	The A_2 -formula over $\mathbb{C}$	9
4.2.	Weight functions	9

¹ *E-mail:* rimanyi@email.unc.edu, supported in part by Simons Foundation grant 523882

² *E-mail:* anv@email.unc.edu, supported in part by NSF grant DMS-1954266

4.3. Representations of $\mathfrak{sl}_3$	10
5. The A_2 -type Selberg integrals over F_p	12
5.1. Relations between F_p -integrals	12
5.2. Proof of Theorem 3.4	14
5.3. Evaluation of $I_{0,0,0}(a,b_1,b_2,c)$	17
5.4. Sketch of the proof of formula (3.11) for $n > 2$	18
References	19

1. Introduction

In 1944 Atle Selberg proved the following integral formula: (1.1)

$$\int_0^1 \cdots \int_0^1 \prod_{1 \leq i < j \leq k} |x_i - x_j|^{2\gamma} \prod_{i=1}^k x_i^{\alpha-1} (1-x_i)^{\beta-1} dx_1 \cdots dx_k$$

Selberg proved the following integral formula: (1.1)

$$= \prod_{j=1}^k \frac{\Gamma(1+j\gamma)}{\Gamma(1+\gamma)} \frac{\Gamma(\alpha+(j-1)\gamma) \Gamma(\beta+(j-1)\gamma)}{\Gamma(\alpha+\beta+(k+j-2)\gamma)}$$

integral formula: (1.1)

where α, β, γ are complex numbers such that $\operatorname{Re} \alpha > 0$, $\operatorname{Re} \beta > 0$, and $\operatorname{Re} \gamma > -\min[(\operatorname{Re} \alpha)/(n-1), (\operatorname{Re} \beta)/(n-1)]$. See [Se, AAR]. Hundreds of papers are devoted to the generalizations of the Selberg integral formula and its applications, see for example [AAR, FW] and references therein. There are q -analysis versions of the formula, the generalizations associated with Lie algebras, elliptic versions, finite field versions, see some references in [AAR, FW, As, Ha, Ka, Op, Ch, TV1, TV2, TV4, Wa1, Wa2, Sp, R, FSV, An, Ev]. In the finite field versions, one considers additive and multiplicative characters of a finite field, which map the field to the field of complex numbers, and forms an analog of equation (1.1), in which both sides are complex numbers. The simplest of such formulas is the classical relation between Jacobi and Gauss sums, see [AAR, An, Ev].

In [RV] we suggested another version of the Selberg integral formula, in which the F_p -Selberg integral is an element of the finite field F_p with an odd prime number p of elements.

Our motivation in [RV] came from the theory of Knizhnik-Zamolodchikov (KZ) equations, see [KZ, EFK]. These are the systems of linear differential equations, satisfied by conformal blocks on the sphere in the WZW model of conformal field theory. The KZ equations were solved in multidimensional hypergeometric integrals in [SV1], see also [V1, V2]. The following general principle was formulated in [MuV]: if an example of the KZ-type equations has a one-dimensional space of solutions, then the corresponding multidimensional hypergeometric integral can be evaluated explicitly. As an illustration of that principle in

[MuV], an example of the $\mathfrak{sl}_2$ differential KZ equations with a one-dimensional space of solutions was considered, the corresponding multidimensional hypergeometric integral was reduced to the Selberg integral and then evaluated by formula (1.1). See other illustrations in [FV, FSV, TV1, TV2, TV4, V3, RTVZ].

Recently in [SV2] the KZ equations were considered modulo a prime number p and polynomial solutions of the reduced equations were constructed, see also [SIV, V3, V4, V5, V6, V7]. The construction is analogous to the construction of the multidimensional hypergeometric solutions, and the constructed polynomial solutions were called the F_p -hypergeometric solutions.

In [RV] we considered the reduction modulo p of the same example of the $\mathfrak{sl}_2$ differential KZ equations, that led in [MuV] to the Selberg integral. We evaluated the corresponding F_p -hypergeometric solution by analogy with the evaluation of the Selberg integral and obtained the F_p -Selberg integral formula in [RV, Theorem 4.1].

In [TV4, Theorem 3.3] the Selberg integral formula of type A_2 was proposed and proved,

$$\begin{aligned}
& \int_{C^{k_1, k_2}[0, 1]} \prod_{i=1}^{k_1} t_i^{\alpha-1} (1-t_i)^{\beta_1-1} \prod_{j=1}^{k_2} (1-s_j)^{\beta_2-1} \prod_{i=1}^{k_1} \prod_{j=1}^{k_2} |s_j - t_i|^{-\gamma} \\
& \quad \times \prod_{1 \leq i < i' \leq k_1} |t_i - t_{i'}|^{2\gamma} \prod_{1 \leq j < j' \leq k_2} |s_j - s_{j'}|^{2\gamma} dt_1 \dots dt_{k_1} ds_1 \dots ds_{k_2} \\
& = \prod_{i=1}^{k_1-k_2} \frac{\Gamma(\beta_1 + (i-1)\gamma)}{\Gamma(\alpha + \beta_1 + (i+k_1-2)\gamma)} \\
& \quad \times \prod_{i=1}^{k_2} \frac{\Gamma(\beta_2 + (i-1)\gamma)}{\Gamma(1 + \beta_2 + (i+k_2-k_1-2)\gamma)} \frac{\Gamma(\beta_1 + \beta_2 + (i-2)\gamma)}{\Gamma(\alpha + \beta_1 + \beta_2 + (i+k_2-3)\gamma)} \\
& \quad \times \prod_{i=1}^{k_2} \frac{\Gamma(1 + (i-k_1-1)\gamma) \Gamma(i\gamma)}{\Gamma(\gamma)} \prod_{i=1}^{k_1} \frac{\Gamma(\alpha + (i-1)\gamma) \Gamma(i\gamma)}{\Gamma(\gamma)}.
\end{aligned}$$

Here $\operatorname{Re} \alpha > 0$, $\operatorname{Re} \beta_1 > 0$, $\operatorname{Re} \beta_2 > 0$, $\operatorname{Re} \gamma < 0$ and $|\operatorname{Re} \gamma|$ is sufficiently small. The integration cycle $C^{k_1, k_2}[0, 1]$ is defined in [TV4, Section 3], also see its definition in [Wa1, Wa2, FW].

The starting point of this formula was an example of the joint system of the $\mathfrak{sl}_3$ trigonometric differential KZ equations and associated dynamical difference equations, an example in which the space of solutions is one-dimensional. The A_n -type Selberg integral formula for arbitrary n was obtained in [Wa1, Wa2], see also [FW].

In this paper we consider the reduction modulo p of the same example of the joint system of the $\mathfrak{sl}_{n+1}$ trigonometric differential KZ equations and associated dynamical difference equations, which led in [TV4, Wa1] to the A_n -type Selberg integral formula. Using the reduction modulo p of these differential and difference equations we obtain our A_n -type F_p -Selberg integral formula for $n > 1$, see (3.11). For $n = 1$ the formula is proved in [RV,

Theorem 4.1]. For $n = 2$ the formula is proved in Theorem 3.4 below. We sketch the proof of the formula for $n > 2$ in Section 5.4. The details of that sketch will appear elsewhere.

The paper is organized as follows. In Section 2 we collect useful facts. In Section 3 we introduce the notion of F_p -integral and discuss the integral formula for the F_p -beta integral. In Section 3 we define the A_n -type F_p -Selberg integral and present its evaluation formula. Theorem 3.4 states that the formula holds for $n = 2$. In Section 3 we also prove Theorem 3.7, which is used in the transition from the A_{n-1} -type formula to the A_n -type formula, in particular, in the transition from the known A_1 -type formula to the new A_2 -type formula. In Section 4 we sketch the proof of formula (1.2) following [TV4]. In Section 5 we adapt this proof to prove Theorem 3.4.

The authors thank I.Cherednik, P.Etingof, E.Rains for useful discussions.

2. Preliminary remarks

In this paper p is an odd prime number.

2.1. Cancellation of factorials.

Lemma 2.1. *If a, b are non-negative integers and $a + b = p - 1$, then in F_p we have*

$$(2.1) \quad a!b! = (-1)^{a+1}.$$

Proof. We have $a! = (-1)^a(p-1) \dots (p-a)$ and $p-a = b+1$. Hence $a!b! = (-1)^a(p-1)! = (-1)^{a+1}$ by Wilson's Theorem.

2.2. Dyson's formula. We shall use Dyson's formula

$$\text{C.T.} \prod_{1 \leq i < j \leq k} (1 - x_i/x_j)^c (1 - x_j/x_i)^c = \frac{(kc)!}{(c!)^k}$$

where C.T. denotes the constant term. See the formula in [AAR, Section 8.8].

2.3. F_p -Integrals. Let M be an F_p -module. Let $P(x_1, \dots, x_k)$ be a polynomial with coefficients in M ,

$$(2.3) \quad P(x_1, \dots, x_k) = \sum_d c_d x_1^{d_1} \dots x_k^{d_k}.$$

Let $l = (l_1, \dots, l_k) \in \mathbb{Z}_{>0}^k$. The coefficient $c_{l_1 p - 1, \dots, l_k p - 1}$ is called the F_p -integral over the p -cycle $[l_1, \dots, l_k]_p$ and is denoted by $R_{[l_1, \dots, l_k]_p} P(x_1, \dots, x_k) dx_1 \dots dx_k$.

Lemma 2.2. For $i = 1, \dots, k-1$ we have

$$(2.4) \quad \int_{[l_1, \dots, l_{i+1}, l_i, \dots, l_k]_p} P(x_1, \dots, x_{i+1}, x_i, \dots, x_k) dx_1 \dots dx_k \\ = \int_{[l_1, \dots, l_k]_p} P(x_1, \dots, x_k) dx_1 \dots dx_k.$$

Lemma 2.3. For any $i = 1, \dots, k$, we have

$$\int_{[l_1, \dots, l_k]_p} \frac{\partial P}{\partial x_i}(x_1, \dots, x_k) dx_1 \dots dx_k = 0 \\ = (k_1, \dots, k_n) \in \mathbb{Z}_{>0}^n$$

$$(2.5) \quad [k]_p := [(1)_{k_1}; (k_1)_{k_2}; \dots; (k_{n-1})_{k_n}]_p,$$

where x_y denotes the y -tuple $(x_1, \dots, x_y)$. For example for $n = 2$, $k = (3, 2)$, we have $[k]_p = [1, 1, 1; 3, 3]_p$.

2.4. $\mathbb{F}_p$ -Beta integral. For non-negative integers the classical beta integral formula says

$$\int_0^1 x^a (1-x)^b dx = \frac{a! b!}{(a+b+1)!}$$

Theorem 2.4 ([V7]). Let $a < p$, $b < p$, $p-1 \nmid a+b$. Then in $\mathbb{F}_p$ we have

$$\int_{[1]_p} x^a (1-x)^b dx = - \frac{a! b!}{(a+b+1-p)!}$$

If $a+b < p-1$, then

$$\int_{[1]_p} x^a (1-x)^b dx = 0$$

3. $\mathbb{F}_p$ -Selberg integral of type A_n

3.1. Admissible parameters. Let $k = (k_1, \dots, k_n) \in \mathbb{Z}_{>0}^n$ and $k_i > k_{i+1}$, $i = 1, \dots, n-1$.

Set $k_0 = k_{n+1} = 0$.

Let $a, b_1, \dots, b_n, c \in \mathbb{Z}_{>0}$. Denote $b = (b_1, \dots, b_n)$ and

(3.1)

$$\begin{aligned}
R_{\mathbf{k}}(a, b, c) &= \\
&= \prod_{1 \leq s \leq r \leq n} \prod_{i=1}^{k_r - k_{r+1}} \frac{(r - s + b_s + \cdots + b_r + (i + s - r - 1)c)!}{(r - s + 1 + a_s + b_s + \cdots + b_r + (i + s - r + k_s - k_{s-1} - 2)c - \delta_{s,1}p)!} \\
&\times (-1)^{\sum_{i=1}^n k_i} \left(\prod_{i=1}^{k_1} (a_1 + (i - 1)c)! \right) \left(\prod_{r=1}^n \prod_{i=1}^{k_r} \frac{(ic)!}{c!} \right) \left(\prod_{r=2}^n \prod_{i=1}^{k_r} (p + (i - k_{r-1} - 1)c)! \right),
\end{aligned}$$

where $a_1 = a, a_2 = \cdots = a_n = 0$; $\delta_{s,1}$ is 1 if $s = 1$ and is zero otherwise.

We say that $a, b_1, \dots, b_n, c \in \mathbb{Z}_{>0}$ are *admissible* if $a + (k_1 - 1)c < p - 1$ and for any factorial $x!$ on the right-hand side of (3.1) we have $0 \leq x < p$. The set of all admissible (a, b, c) is denoted by A_k .

$$\mathbb{Z}_{>0}^{n+2}$$

(3.2)

$$\begin{aligned}
0 \leq r - s + b_s + \cdots + b_r + (s - r)c, \\
r - s + b_s + \cdots + b_r + (k_r - k_{r+1} + s - r - 1)c \leq p - 1,
\end{aligned}$$

for $1 \leq s \leq r \leq n$;

(3.3)

$$\begin{aligned}
0 \leq r - s + 1 + b_s + \cdots + b_r + (s - r + k_s - k_{s-1} - 1)c, \\
r - s + 1 + b_s + \cdots + b_r + (s - r + k_r - k_{r+1} + k_s - k_{s-1} - 2)c \leq p - 1,
\end{aligned}$$

for $2 \leq s \leq r \leq n$;

(3.4)

$$\begin{aligned}
p \leq r + a + b_1 + \cdots + b_r + (k_1 - r)c, \\
r + a + b_1 + \cdots + b_r + (k_r - k_{r+1} + k_1 - r - 1)c < 2p,
\end{aligned}$$

for $1 \leq r \leq n$;

(3.5)

$$a + (k_1 - 1)c < p - 1, \quad b_1 > p - 1 - (a + (k_1 - 1)c), \quad 0 < k_1 c < p.$$

Lemma 3.2. Assume that $(a, b, c) \in A_k$. Then

(3.6)

$$b_1 > p - 1 - (a + (k_1 - 1)c), \quad b_s > (k_{s-1} - k_s + 1)c - 1, \quad s = 2, \dots, n.$$

Proof. The inequality $b_s > (k_{s-1} - k_s + 1)c - 1$ for $s = 2, \dots, n$ follows from the first inequality in (3.3) for $r = s$. The inequality $b_1 > p - 1 - (a + (k_1 - 1)c)$ follows from the first inequality in (3.4) for $r = 1$.

Example. Let $n = 1, k = (k_1)$. Then

$$R_{(k_1)}(a, b_1, c) = \prod_{i=1}^{k_1} \frac{(ic)!}{c!} \frac{(a + (i-1)c)! (b_1 + (i-1)c)!}{(1 + a + b_1 + (i + k_1 - 2)c - p)!}$$

and $A_{(k_1)}$ consists of $a, b, c \in \mathbb{Z}_{>0}$ such that

$$(3.8) \quad a + (k_1 - 1)c < p - 1, \quad b_1 + (k_1 - 1)c \leq p - 1, \quad k_1 c \leq p - 1, \quad p - 1 \leq a + b_1 + (k_1 - 1)c, \quad a + b_1 + (2k_1 - 2)c < 2p - 1.$$

$$= (k_1, \dots, k_n) \in \mathbb{Z}_{>0}^n$$

$$t^{(i)} = (t_1^{(i)}, \dots, t_{k_i}^{(i)}), \quad i = 1, \dots, n.$$

Define the *master polynomial*

$$\Phi_k(t; a, b, c) = \prod_{i=1}^n \left(\prod_{j=1}^{k_i} (t_j^{(i)})^{a_i} (1 - t_j^{(i)})^{b_i} \prod_{1 \leq j < j' \leq k_i} (t_j^{(i)} - t_{j'}^{(i)})^{2c} \right) \prod_{i=1}^{n-1} \prod_{j=1}^{k_{i+1}} \prod_{j'=1}^{k_i} (t_j^{(i+1)} - t_{j'}^{(i)})^{p-c}$$

Denote

$$(3.10) \quad S_k(a, b, c) = \int_{[k]_p} \Phi_k dt.$$

The F_p -integral $S_k(a, b, c)$ is called the *F_p -Selberg integral of type A_n* .

Conjecture 3.3. *Let n be a positive integer. Let $k = (k_1, \dots, k_n) \in \mathbb{Z}_{>0}^n$, $k_i > k_{i+1}$, $i = 1, \dots, n-1$. Then for any $(a, b, c) \in A_k$ we have the equality in F_p :*

$$(3.11) \quad S_k(a, b, c) = R_k(a, b, c).$$

For $n = 1$ formula (3.11) is proved in [RV, Theorem 4.1]. For $n = 2$ formula (3.11) is proved in the next theorem.

Theorem 3.4. *Let $k = (k_1, k_2) \in \mathbb{Z}_{>0}^2$, $k_1 > k_2$. Then for any $(a, b, c) \in A_k$ we have the equality in F_p :*

$$(3.12) \quad S_k(a, b, c) = R_k(a, b, c).$$

Formula (3.11) for $n = 2$ is deduced from formula (3.11) for $n = 1$ in Section 5.

More generally, for any k formula (3.11) for $n = k$ can be deduced from formula (3.11) for $n = k - 1$ similarly, see the sketch of that in Section 5.4. Details of that deduction will appear elsewhere. Because of that formula (3.11) for any n is formulated as a conjecture and not as a theorem.

Remark. Theorem 3.4 can be extended to the case of k such that $k_1 > k_2$, but the structure of inequalities in Lemma 3.1 will depend on the appearance of the equality $k_1 = k_2$ in k , and the proof of Theorem 3.4 will split into different sub-cases. To shorten the exposition we restrict ourselves to k such that $k_1 > k_2$.

Example. Here is the simplest A_2 -type F_p -Selberg integral formula with $k_1 = k_2 = 1$.

Theorem 3.5. Assume that a, b_1, b_2, c are integers such that

$$\begin{aligned} 0 \leq a < p, \quad 0 \leq c \leq p, \quad 0 \leq b_2 - c + 1 < p, \\ 0 \leq b_1 + b_2 - c + 1 < p, \quad p - 1 \leq a + b_1 + b_2 - c + 1 < 2p - 1. \end{aligned}$$

Then in F_p we have

$$(3.13) \quad \int_{[1;1]_p} t^a (1-t)^{b_1} (s-t)^{p-c} (1-s)^{b_2} dt ds = \frac{a! (b_1 + b_2 - c + 1)!}{(a + b_1 + b_2 - c + 2 - p)!} \frac{(p-c)! (b_2)!}{(b_2 - c + 1)!}$$

Proof. Change variables $s = t + (1-t)v$, then the F_p -integral becomes equal to

$$\int_{[1,1]_p} t^a (1-t)^{b_1+b_2-c+1} v^{p-c} (1-v)^{b_2} dt dv.$$

Applying the F_p -beta integral formula we obtain the theorem.

The simplest A_3 -type F_p -Selberg integral formula $k_1 = k_2 = k_3 = 1$ is given by the next theorem.

Theorem 3.6. Let a, b_1, b_2, b_3, c be integers such that all factorials on the right-hand side of formula (3.14) are factorials of non-negative integers less than p . Then in F_p we have

$$(3.14) \quad \int_{[1;1;1]_p} t^a (1-t)^{b_1} (s-t)^{p-c} (1-s)^{b_2} (u-s)^{p-c} (1-u)^{b_3} dt ds du \\ = - \frac{a! (b_1 + b_2 + b_3 - 2c + 2)!}{(a + b_1 + b_2 + b_3 - 2c + 3 - p)!} \frac{(p-c)! (b_2 + b_3 - c + 1)!}{(b_2 + b_3 - 2c + 2)!} \frac{(p-c)! (b_3)!}{(b_3 - c + 1)!}$$

Proof. The proof is the same as the proof of the previous theorem.

The versions of identities (3.13), (3.14) over complex numbers see in [MuV, Theorem 1].

3.3. Relation between the F_p -Selberg integrals of types A_{n-1} - and A_n .

Theorem 3.7. Let $n > 1$ and $k = (k_1, \dots, k_n)$, $k^0 = (k_1, \dots, k_{n-1})$, $b = (b_1, \dots, b_n)$, $b^0 = (b_1, \dots, b_{n-1})$. Assume that formula (3.11) holds for the F_p -Selberg integral $S_{[k^0]_p}(a, b^0, c)$ of type A_{n-1} . Also assume that $b_n = (k_{n-1} - k_n + 1)c - 1$. Then formula (3.11) holds for the F_p -Selberg integral $S_{[k]_p}(a, b, c)$ of type A_n .

$$t_j^{(n)}$$

$$\prod_{j=1}^{k_n} (t_j^{(n)})^{k_{n-1}p-1} \\ t_j^{(i)}$$

$\Phi_{[k^0]_p}(t^{(1)}, \dots, t^{(n-1)}; a, b^0, c)$ of $\Phi_{[k]_p}(t; a, b, c)$ and are used to calculate the coefficient of

$$\prod_{j=1}^{k_1} (t_j^{(1)})^{p-1} \prod_{i=2}^{n-1} \prod_{j=1}^{k_i} (t_j^{(i)})^{k_{i-1}p-1}$$

More precisely, under the assumptions of the theorem we have

$$S_{\mathbf{k}}(a, b, c) = (-1)^{b_n k_n + c k_n (k_n - 1)/2} \frac{(k_n c)!}{(c!)^{k_n}} S_{\mathbf{k}'}(a', b', c)$$

$$(-1)^{b_n k_n + c k_n (k_n - 1)/2} \frac{(k_n c)!}{(c!)^{k_n}} \prod_{j=1}^{k_n} (t_j^{(n)})^{k_{n-1} p - 1}$$

$$k_n k_{n-1}$$

$$\begin{matrix} Y Y & (n) & (n-1) p - c & Y & (n) & (n) 2c \\ (t_j & - t_{j0} &) & (t_j & - t_{j0} &) \end{matrix}, j=1, j^0=1$$

$$16j < j^0 \leq k_n$$

see Dyson's formula. We have $(-1)^{b_n k_n + c k_n (k_n - 1)/2} = (-1)^{(k_{n-1} k_n - k_n (k_n + 1)/2) c - k_n}$. Hence

$$S_{\mathbf{k}}(a, b, c) = (-1)^{(k_{n-1} k_n - k_n (k_n + 1)/2) c - k_n} \frac{(k_n c)!}{(c!)^{k_n}} S_{\mathbf{k}'}(a, b', c)$$

$$= (-1)^{(k_{n-1} k_n - k_n (k_n + 1)/2) c - k_n} \frac{(k_n c)!}{(c!)^{k_n}} R_{\mathbf{k}'}(a, b', c)$$

where $S_{k_0}(a, b^0, c) = R_{k_0}(a, b^0, c)$ holds by assumptions. To prove the theorem we need to show that

$$R_{[\mathbf{k}]_p}(a, (b', b_n), c) = (-1)^{(k_{n-1} k_n - k_n (k_n + 1)/2) c - k_n} \frac{(k_n c)!}{(c!)^{k_n}} R_{\mathbf{k}'}(a, b', c)$$

Indeed we have

$$R_{\mathbf{k}}(a, (b', b_n), c) = R_{\mathbf{k}'}(a, b', c) \prod_{i=1}^{k_n} \frac{(ic)!}{c!} \prod_{i=1}^{k_n} (p + (i - k_{n-1} - 1)c)!$$

$$\times \prod_{1 \leq s \leq n} \prod_{i=1}^{k_n} \frac{(n - s + b_s + \cdots + b_n + (i + s - n - 1)c)!}{(n - s + 1 + a_s + b_s + \cdots + b_n + (i + s - n + k_s - k_{s-1} - 2)c - \delta_{s,1} p)!}$$

$$\times \prod_{1 \leq s \leq n-1} \prod_{i=1}^{k_n} \frac{(n - s + a_s + b_s + \cdots + b_{n-1} + (i + s - n + k_s - k_{s-1} - 3)c - \delta_{s,1} p)!}{(n - 1 - s + b_s + \cdots + b_{n-1} + (i + s - r - 1)c)!}$$

$$= R_{\mathbf{k}'}(a, b', c) \prod_{i=1}^{k_n} \frac{(ic)!}{c!} \prod_{i=1}^{k_n} \frac{(b_n + (i - 1)c)! (p + (i - k_{n-1} - 1)c)!}{(1 + b_n + (i + k_n - k_{n-1} - 2)c)!}$$

$$= R_{\mathbf{k}'}(a, b', c) \prod_{i=1}^{k_n} \frac{(ic)!}{c!} \prod_{i=1}^{k_n} \frac{((i + k_{n-1} - k_n)c - 1)! (p + (i - k_{n-1} - 1)c)!}{((i - 1)c)!}$$

$$= R_{\mathbf{k}'}(a, b', c) (-1)^{(k_{n-1} k_n - k_n (k_n + 1)/2) c - k_n} \frac{(k_n c)!}{(c!)^{k_n}}$$

where in the last step we use the cancellation Lemma 2.1. The theorem is proved.

Corollary 3.8. Let $n > 1$, $k = (k_1, \dots, k_n)$, and $(a, b_1, c) \in A_{(k_1)}$. Let $b = (b_1, \dots, b_n)$, where $b_i = (k_{i-1} - k_i + 1)c - 1$ for $i = 2, \dots, n$. Then formula (3.11) holds for the F_p -Selberg integral $S_{[\mathbf{k}]_p}(a, b, c)$ of type A_n .

Proof. Formula (3.11) for the F_p -Selberg integrals of type A_1 is proved in [RV]. Hence the corollary follows from Theorem 3.7 by induction on n .

4. The A_2 -type Selberg integral over $\mathbb{C}$

In this section we formulate the A_2 -type Selberg integral formula over $\mathbb{C}$, formulated and proved in [TV4], and sketch the proof of the formula, following [TV4]. In Section 5 we adapt this proof to prove the A_2 -type F_p -Selberg integral formula, that is, formula (3.11) for $n = 2$.

4.1. The A_2 -formula over $\mathbb{C}$. For $k_1 > k_2 > 0$ let $t = (t_1, \dots, t_{k_1})$, $s = (s_1, \dots, s_{k_2})$. Define the *master function*

$$(4.1) \quad \begin{aligned} \Phi(t; s) = & \prod_{i=1}^{k_1} t_i^{\alpha_i - 1} (1 - t_i)^{\beta_1 - 1} \prod_{j=1}^{k_2} (1 - s_j)^{\beta_2 - 1} \prod_{i=1}^{k_1} \prod_{j=1}^{k_2} |s_j - t_i|^{-\gamma} \\ & \times \prod_{i=1}^{k_1} |t_i - t_{i_0}|_{2\gamma} \prod_{j=1}^{k_2} |s_j - s_{j_0}|_{2\gamma} \prod_{1 \leq i < i_0 \leq k_1} \prod_{1 \leq j < j_0 \leq k_2} \end{aligned}$$

and the integral

$$(4.2) \quad \tilde{S}(\alpha, \beta_1, \beta_2, \gamma) = \int_{C^{k_1, k_2}[0, 1]} \Phi(t; s) dt ds,$$

where the integration cycle $C^{k_1, k_2}[0, 1]$ is defined in [TV4, Section 3]. The explicit description of this cycle is of no importance in this paper.

Theorem 4.1 ([TV4, Theorem 3.3]). *Let $\alpha, \beta_1, \beta_2, \gamma$ be complex numbers such that $\operatorname{Re} \alpha > 0$, $\operatorname{Re} \beta_1 > 0$, $\operatorname{Re} \beta_2 > 0$, $\operatorname{Re} \gamma < 0$ and $|\operatorname{Re} \gamma|$ sufficiently small. Then*

$$\begin{aligned} \tilde{S}(\alpha, \beta_1, \beta_2, \gamma) = & \prod_{i=1}^{k_1 - k_2} \frac{\Gamma(\beta_1 + (i - 1)\gamma)}{\Gamma(\alpha + \beta_1 + (i + k_1 - 2)\gamma)} \\ & \times \prod_{i=1}^{k_2} \frac{\Gamma(\beta_2 + (i - 1)\gamma)}{\Gamma(1 + \beta_2 + (i + k_2 - k_1 - 2)\gamma)} \frac{\Gamma(\beta_1 + \beta_2 + (i - 2)\gamma)}{\Gamma(\alpha + \beta_1 + \beta_2 + (i + k_2 - 3)\gamma)} \\ & \times \prod_{i=1}^{k_2} \frac{\Gamma(1 + (i - k_1 - 1)\gamma) \Gamma(i\gamma)}{\Gamma(\gamma)} \prod_{i=1}^{k_1} \frac{\Gamma(\alpha + (i - 1)\gamma) \Gamma(i\gamma)}{\Gamma(\gamma)}. \end{aligned}$$

In the next Sections 4.2, 4.3 we sketch the proof of formula (4.3) following [TV4].

4.2. Weight functions. To evaluate $\tilde{S}(\alpha, \beta_1, \beta_2, \gamma)$ we introduce a collection of new integrals $J_{l_1, l_2, m}(\alpha, \beta_1, \beta_2, \gamma)$, which also can be evaluated explicitly, see [TV4]. For a function $f(t_1, \dots, t_k)$ set

$$_{t_1, \dots, t_k} f(t_1, \dots, t_k) = \frac{1}{k!} \sum_{\sigma \in S_k} f(t_{\sigma_1}, \dots, t_{\sigma_k})$$

Given $k_1 > k_2 > 0$, we say that a triple of non-negative integers (l_1, l_2, m) is *allowable* if $l_1 \leq k_1 - k_2 + l_2$, $l_2 \leq k_2$ and $m \leq \min(l_1, l_2)$. For any allowable triple (l_1, l_2, m) define the weight function

$$\begin{aligned} W_{l_1, l_2, m}(t_1, \dots, t_{k_1}; s_1, \dots, s_{k_2}) = \\ \prod_{a=1}^{l_1} t_a \prod_{b=1}^{l_2} (1 - t_b) \prod_{c=1}^m \frac{1 - s_c}{s_c - t_c} \prod_{d=1}^{k_2 - m} \frac{1 - s_d}{s_d - t_{b+k_1-k_2}} \\ = \text{Sym}_{t_1, \dots, t_{k_1}} \text{Sym}_{s_1, \dots, s_{k_2}} \end{aligned}$$

and the integral

$$J_{l_1, l_2, m}(\alpha, \beta_1, \beta_2, \gamma) = \int_{C^{k_1, k_2}[0, 1]} \Phi(t; s) W_{l_1, l_2, m}(t; s) dt ds.$$

In particular,

$$(4.4) \quad J_{0, k_2, 0}(\alpha, \beta_1, \beta_2, \gamma) = \tilde{S}(\alpha, \beta_1 + 1, \beta_2, \gamma).$$

4.3. Representations of $\mathfrak{sl}_3$. Consider the complex Lie algebra $\mathfrak{sl}_3$ with standard generators $f_1, f_2, e_1, e_2, h_1, h_2$, simple roots σ_1, σ_2 , fundamental weights ω_1, ω_2 . Let $V_{\lambda_1}, V_{\lambda_2}$ be the irreducible $\mathfrak{sl}_3$ -modules with highest weights

$$\lambda_1 = -\frac{\alpha}{\gamma} \omega_1, \quad \lambda_2 = -\frac{\beta_1}{\gamma} \omega_1 - \frac{\beta_2}{\gamma} \omega_2$$

and highest weight vectors v_1, v_2 . For $k_1 > k_2 > 0$ consider the weight subspace

$V_{\lambda_1} \otimes V_{\lambda_2}[\lambda_1 + \lambda_2 - k_1 \sigma_1 - k_2 \sigma_2]$ of the tensor product $V_{\lambda_1} \otimes V_{\lambda_2}$ and the singular weight subspace $\text{Sing} V_{\lambda_1} \otimes V_{\lambda_2}[\lambda_1 + \lambda_2 - k_1 \sigma_1 - k_2 \sigma_2]$ consisting of the vectors $w \in V_{\lambda_1} \otimes V_{\lambda_2}[\lambda_1 + \lambda_2 - k_1 \sigma_1 - k_2 \sigma_2]$ such that $e_1 w = 0, e_2 w = 0$. A basis of $V_{\lambda_1} \otimes V_{\lambda_2}[\lambda_1 + \lambda_2 - k_1 \sigma_1 - k_2 \sigma_2]$ is formed by the vectors

$$v_{l_1, l_2, m} = \frac{f_1^{k_1 - k_2 - l_1 + l_2} [f_1, f_2]^{k_2 - l_2} v_1 \otimes f_1^{l_1 - m} [f_1, f_2]^m f_2^{l_2 - m} v_2}{(k_1 - k_2 - l_1 + l_2)! (k_2 - l_2)! (l_1 - m)! m! (l_2 - m)!}$$

labeled by allowable triples (l_1, l_2, m) . It is known from the theory of KZ equations that the vector

$$J = \sum_{l_1, l_2, m} (-1)^{l_1} J_{l_1, l_2, m}(\alpha, \beta_1, \beta_2, \gamma) v_{l_1, l_2, m}$$

is a singular vector, see [M, Theorem 2.4], [MaV, Corollary 10.3], cf. [RSV].

The singular vector equations $e_1 J = 0$, $e_2 J = 0$ are calculated with the help of the formulas:

$$\begin{aligned} h_1 v_1 &= -\frac{\alpha}{\gamma} v_1, & h_2 v_1 &= 0, & h_1 v_2 &= -\frac{\beta_1}{\gamma} v_2, & h_2 v_2 &= -\frac{\beta_2}{\gamma} v_2, \\ [h_1, f_1] &= -2f_1, & [h_1, f_2] &= f_2, & [h_2, f_1] &= f_1, & [h_2, f_2] &= -2f_2 \\ [e_1, f_1] &= h_1, & [e_1, f_2] &= [e_2, f_1] = 0, & [e_2, f_2] &= h_2, \end{aligned}$$

$$[h_1, [f_1, f_2]] = -[f_1, f_2], \quad [h_2, [f_1, f_2]] = -[f_1, f_2], \quad [e_1, [f_1, f_2]] = f_2, \quad [e_2, [f_1, f_2]] = -f_1.$$

Here are some of the singular vector relations.

Theorem 4.2 (cf. [TV4, Theorem 5.2]). *We have*

$$J_{0, l_2, 0} = (-1)^{l_2} J_{0, 0, 0} \prod_{i=0}^{l_2-1} \frac{(k_1 - k_2 + i + 1)\gamma}{\beta_2 + i\gamma}$$

Proof. We have

$$\begin{aligned} e_2 \frac{f_1^{k_1-k_2+i} [f_1, f_2]^{k_2-i} v_1 \otimes f_2^i v_2}{(k_1 - k_2 + i)! (k_2 - i)! i!} &= - (k_1 - k_2 + i + 1) \frac{f_1^{k_1-k_2+i+1} [f_1, f_2]^{k_2-i-1} v_1 \otimes f_2^i v_2}{(k_1 - k_2 + i + 1)! (k_2 - i - 1)! i!} \\ &\quad + \left(-\frac{\beta_2}{\gamma} - i + 1 \right) \frac{f_1^{k_1-k_2+i} [f_1, f_2]^{k_2-i} v_1 \otimes f_2^{i-1} v_2}{(k_1 - k_2 + i)! (k_2 - i)! (i - 1)!}. \end{aligned}$$

Calculating the coefficient of $\frac{f_1^{k_1-k_2+i+1} [f_1, f_2]^{k_2-i-1} v_1 \otimes f_2^i v_2}{(k_1 - k_2 + i + 1)! (k_2 - i - 1)! i!}$ in $e_2 J = 0$ we obtain

$$(4.6) \quad (k_1 - k_2 + i + 1)\gamma J_{0, i, 0} + (\beta_2 + i\gamma) J_{0, i+1, 0} = 0.$$

This implies the theorem.

Hence

$$J_{0, k_2, 0}(\alpha, \beta_1, \beta_2, \gamma) = (-1)^{k_2} J_{0, 0, 0}(\alpha, \beta_1, \beta_2, \gamma) \prod_{i=0}^{k_2-1} \frac{(k_1 - k_2 + i + 1)\gamma}{\beta_2 + i\gamma}$$

Combining (4.4) and (4.7) we observe that formula (4.3) is equivalent to the formula

$$\begin{aligned}
J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma) &= \prod_{i=1}^{k_1-k_2} \frac{\Gamma(1 + \beta_1 + (i-1)\gamma)}{\Gamma(1 + \alpha + \beta_1 + (i+k_1-2)\gamma)} \\
&\times \prod_{i=1}^{k_2} \frac{\Gamma(1 + \beta_2 + (i-1)\gamma)}{\Gamma(1 + \beta_2 + (i+k_2-k_1-2)\gamma)} \frac{\Gamma(1 + \beta_1 + \beta_2 + (i-2)\gamma)}{\Gamma(1 + \alpha + \beta_1 + \beta_2 + (i+k_2-3)\gamma)} \\
&\times \prod_{i=1}^{k_2} \frac{\Gamma((i-k_1-1)\gamma) \Gamma(i\gamma)}{\Gamma(\gamma)} \prod_{i=1}^{k_1} \frac{\Gamma(\alpha + (i-1)\gamma) \Gamma(i\gamma)}{\Gamma(\gamma)}.
\end{aligned}$$

Denote by $R_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma)$ the right-hand side of (4.8).

To prove (4.8) we use the following observation. The weight subspace $V_{\lambda_1}[\lambda_1 - k_1\sigma_1 - k_2\sigma_2] \subset V_{\lambda_1}$ is one-dimensional with a basis vector

$$v_{0,0,0} = \frac{f_1^{k_1-k_2} [f_1, f_2]^{k_2} v_1 \otimes v_2}{(k_1 - k_2)! (k_2)!}$$

By [MaV, Theorem 5.1] the vector-valued function $J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma) v_{0,0,0}$ satisfies the dynamical difference equations introduced in [TV3],

$$(4.9) \quad J_{0,0,0}(\alpha, \beta_1 - 1, \beta_2, \gamma) v_{0,0,0} = J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma) B_1 v_{0,0,0},$$

$$(4.10) \quad J_{0,0,0}(\alpha, \beta_1, \beta_2 - 1, \gamma) v_{0,0,0} = J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma) B_2 v_{0,0,0}.$$

Here B_1, B_2 are certain linear operators acting on V_{λ_1} and preserving the weight decomposition of V_{λ_1} , see formulas for these operators in the example in [MaV, Section 7.1] and in [MaV, Section 3.1], also see [TV3, Formula (8)].

Written explicitly equations (4.9), (4.10) give us the difference equations for the scalar function $J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma)$ with respect to the shift of the variables $\beta_1 \rightarrow \beta_1 - 1$ and $\beta_2 \rightarrow \beta_2 - 1$,

$$(4.11)$$

$$\begin{aligned}
&\times \prod_{i=1} \frac{1}{\beta_1 + \beta_2 + (i-2)\gamma}, \\
J_{0,0,0}(\alpha, \beta_1, \beta_2 - 1, \gamma) &= J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma) \\
&\times \prod_{i=1}^{k_2} \frac{\beta_2 + (i + k_2 - k_1 - 2)\gamma}{\beta_2 + (i-1)\gamma} \frac{\alpha + \beta_1 + \beta_2 + (i + k_2 - 3)\gamma}{\beta_1 + \beta_2 + (i-2)\gamma} \\
J_{0,0,0}(\alpha, \beta_1 - 1, \beta_2, \gamma) &= J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma) \prod_{i=1}^{k_1-k_2} \frac{\alpha + \beta_1 + (i + k_1 - 2)\gamma}{\beta_1 + (i-1)\gamma} \\
&\quad \times \frac{\alpha + \beta_1 + \beta_2 + (i + k_1 - 3)\gamma}{\beta_1 + \beta_2 + (i-2)\gamma}
\end{aligned}$$

$$(4.12)$$

The difference equations for $J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma)$ are the same as the difference equations for the function $R_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma)$ with respect to the shift of the variables $\beta_1 \rightarrow \beta_1 - 1$ and $\beta_2 \rightarrow \beta_2 - 1$. Therefore, the functions $J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma)$ and $R_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma)$ are proportional up to a periodic function of β_1, β_2 . The periodic function can be fixed by comparing asymptotics as $\text{Re}\beta_1 \rightarrow \infty$, $\text{Re}\beta_2 \rightarrow \infty$. This finishes the proof in [TV4] of formulas (4.8) and (4.3).

5. The A_2 -type Selberg integrals over F_p

5.1. Relations between F_p -integrals. For $k = (k_1, k_2)$, $k_1 > k_2 > 0$ and integers $0 < a, b_1, b_2, c < p$ define the master polynomial

$$(5.1) \quad \Phi_k(t; s; a, b_1, b_2, c) = \prod_{i=1}^{k_1} t a_i (1 - t_i)^{b_1} \prod_{j=1}^{k_2} Y(1 - s_j)^{b_2} \prod_{i=1}^{k_1} \prod_{j=1}^{k_2} Y(s_j - t_i)^{p-c} \\ \times \prod_{i=1}^{k_1} Y(t_i - t_{i_0})^{2c} \prod_{j=1}^{k_2} Y(s_j - s_{j_0})^{2c}$$

and the F_p -integral

$$(5.2) \quad S_k(a, b_1, b_2, c) = \int_{[k]_p} \Phi_k(t; s; a, b_1, b_2, c) dt ds,$$

where the p -cycle $[k]_p$ is defined in (2.5). This is the A_2 -type F_p -Selberg integral, see (3.10).

For an allowable triple (l_1, l_2, m) define the F_p -integral

$$(5.3) \quad I_{l_1, l_2, m}(a, b_1, b_2, c) = \int_{[k]_p} \Phi_k(t; s; a, b_1, b_2, c) \frac{W_{l_1, l_2, m}(t; s)}{\prod_{i=1}^{k_1} t_i (1 - t_i) \prod_{j=1}^{k_2} (1 - s_j)} dt ds,$$

where $W_{l_1, l_2, m}(t; s)$ is the weight function defined in Section 4.2. Clearly we have

$$(5.4) \quad I_{0, k_2, 0}(a, b_1, b_2, c) = S_k(a - 1, b_1, b_2 - 1, c).$$

Denote

$$\begin{aligned}
(5.5) \quad B_0(a, b_1, b_2, c) &= (-1)^{k_2} \prod_{i=0}^{k_2-1} \frac{(k_1 - k_2 + i + 1)c}{b_2 + ic}, \\
B_1(a, b_1, b_2, c) &= \prod_{i=1}^{k_1-k_2} \frac{a + b_1 + (i + k_1 - 2)c}{b_1 + (i - 1)c} \prod_{i=1}^{k_2} \frac{a + b_1 + b_2 + (i + k_2 - 3)c}{b_1 + b_2 + (i - 2)c} \\
B_2(a, b_1, b_2, c) &= \prod_{i=1}^{k_2} \frac{b_2 + (i + k_2 - k_1 - 2)c}{b_2 + (i - 1)c} \frac{a + b_1 + b_2 + (i + k_2 - 3)c}{b_1 + b_2 + (i - 2)c}.
\end{aligned}$$

Theorem 5.1. Assume that $k_1 < p$.

- (i) Assume that every factor in B_0 in the numerator or denominator is a nonzero element of $\mathbb{F}_p$. Then

$$(5.6) \quad I_{0,k_2,0}(a, b_1, b_2, c) = B_0(a, b_1, b_2, c) I_{0,0,0}(a, b_1, b_2, c).$$

- (ii) Assume that every factor in B_1 in the numerator or denominator is a nonzero element of $\mathbb{F}_p$ and $b_1 > 1$, then

$$(5.7) \quad I_{0,0,0}(a, b_1 - 1, b_2, c) = B_1(a, b_1, b_2, c) I_{0,0,0}(a, b_1, b_2, c).$$

- (iii) Assume that every factor in B_2 in the numerator or denominator is a nonzero element of $\mathbb{F}_p$ and $b_2 > 1$, then

$$(5.8) \quad I_{0,0,0}(a, b_1, b_2 - 1, c) = B_2(a, b_1, b_2, c) I_{0,0,0}(a, b_1, b_2, c).$$

Proof. Equation (5.6) is an $\mathbb{F}_p$ -analog of equation (4.7) and its proof is analogous to the proof of equation (4.7).

More precisely, consider the complex Lie algebra $\mathfrak{sl}_3$ with standard generators $f_1, f_2, e_1, e_2, h_1, h_2$, simple roots σ_1, σ_2 , fundamental weights ω_1, ω_2 . Let $V_{\lambda_1}, V_{\lambda_2}$ be the irreducible $\mathfrak{sl}_3$ modules with highest weights

$$\begin{aligned}
\lambda_1 &= -\frac{a}{c+p} \omega_1, & \lambda_2 &= -\frac{b_1}{c+p} \omega_1 - \frac{b_2}{c+p} \omega_2 \\
& & & f_1^{r_1} [f_1, f_2]^{r_2} v_1 \\
& & & f_1^{r_1} [f_1, f_2]^{r_2} f_2^{r_3} v_2
\end{aligned}$$

$$-\frac{a}{c+p} - \frac{b_1}{c+p}, -\frac{b_2}{c+p}$$

$V_{\lambda_1}^{\mathbb{F}_p}$

$$f_1^{r_1} [f_1, f_2]^{r_2} v_1$$

$$V_{\lambda_2}^{\mathbb{F}_p}$$

$$V_{\lambda_1}^{\mathbb{F}_p} \otimes V_{\lambda_2}^{\mathbb{F}_p} [\lambda_1 + \lambda_2 -$$

$$v_{l_1, l_2, m} = \frac{V_{\lambda_1}^{\mathbb{F}_p} \otimes V_{\lambda_2}^{\mathbb{F}_p} f_1^{k_1 - k_2 - l_1 + l_2} [f_1, f_2]^{k_2 - l_2} v_1 \otimes f_1^{l_1 - m} [f_1, f_2]^m f_2^{l_2 - m} v_2}{(k_1 - k_2 - l_1 + l_2)! (k_2 - l_2)! (l_1 - m)! m! (l_2 - m)!}$$

labeled by allowable triples (l_1, l_2, m) .

Lemma 5.2. *The vector*

$$I = \sum_{l_1, l_2, m} (-1)^{l_1} I_{l_1, l_2, l_m}(a, b_1, b_2, c) v_{l_1, l_2, m}$$

$$V_{\lambda_1}^{\mathbb{F}_p} \otimes V_{\lambda_2}^{\mathbb{F}_p}$$

Proof. Equations $e_1 I = 0$, $e_2 I = 0$ are $\mathbb{F}_p$ -analogs of equations $e_1 J = 0$, $e_2 J = 0$ over $\mathbb{C}$.

For $i = 1, 2$, the vector $e_i J$ is the integral of a certain differential $k_1 + k_2$ -form μ_i . It is shown in [SV2, Theorems 6.16.2], [M, Theorem 2.4] that $\mu_i = dv_i$, where v_i is some explicitly written differential $k_1 + k_2 - 1$ -form. This implies $e_i J = 0$ by Stokes' theorem.

The vector $e_i I$ is the $\mathbb{F}_p$ -integral of the same μ_i reduced modulo p . It is explained in [SV2, Section 4] that the differential form v_i also can be reduced modulo p and this implies that the $\mathbb{F}_p$ -integral $e_i I$ is zero by Lemma 2.3. Cf. the proof of [SV2, Theorem 2.4].

Lemma 5.2 implies the equations

$$(5.9) \quad (k_1 - k_2 + i + 1) c I_{0, i, 0} + (b_2 + ic) I_{0, i+1, 0} = 0$$

for $i = 0, \dots, k_2 - 1$, similarly to the proof of equations (4.6). The iterated application of equation (5.9) implies equation (5.6).

The proof of equations (5.7), (5.8) is parallel to the proof of equations (4.11), (4.12). We prove (5.7). The proof of (5.8) is similar.

Equation (4.11) follows from equation (4.9):

$$V_{\lambda_1}^{\mathbb{F}_p}$$

The proof of (5.10) in [MaV] goes as follows. The left-hand side of (5.10) is a vector-valued integral of a suitable differential $k_1 + k_2$ -form μ . It is shown in [M, Theorem 5.1] that $\mu = dv$, where v is some explicitly written differential $k_1 + k_2 - 1$ -form. This implies (5.10) by Stokes' theorem.

The p -analog of the left-hand side of (5.10) is the element

$$I_{0,0,0}(a, b_1 - 1, b_2, c)v_{0,0,0} - I_{0,0,0}(a, b_1, b_2, c)\mathbb{B}_1 v_{0,0,0} \in V_{\lambda_1}^{\mathbb{F}_p}$$

This element is the F_p -integral of the same μ reduced modulo p . It is explained in [SV2, Section 4] that the differential form v also can be reduced modulo p and this implies that the F_p -integral in the left-hand side of (5.11) equals zero by Lemma 2.3. Hence equation (5.7) is proved and Theorem 5.1 is proved.

5.2. Proof of Theorem 3.4. Recall the set of admissible parameters A_k introduced in Section 3.1 for $k = (k_1, k_2)$, $k_1 > k_2 > 0$.

Lemma 5.3. Assume that $(a, b_1 - 1, b_2, c), (a, b_1, b_2, c) \in A_k$. Then

$$(5.12) \quad S_k(a, b_1 - 1, b_2, c) = S_k(a, b_1, b_2, c) \prod_{i=1}^{k_1-k_2} \frac{1 + a + b_1 + (i + k_1 - 2)c - p}{b_1 + (i - 1)c} \\ \times \prod_{i=1}^{k_2} \frac{2 + a + b_1 + b_2 + (i + k_1 - 3)c - p}{1 + b_1 + b_2 + (i - 2)c}.$$

Assume that $(a, b_1, b_2 - 1, c), (a, b_1, b_2, c) \in A_k$. Then

$$S_k(a, b_1, b_2 - 1, c) = S_k(a, b_1, b_2, c) \prod_{i=1}^{k_2} \frac{1 + b_2 + (i + k_2 - k_1 - 2)c}{b_2 + (i - 1)c} \\ \times \prod_{i=1}^{k_2} \frac{2 + a + b_1 + b_2 + (i + k_1 - 3)c - p}{1 + b_1 + b_2 + (i - 2)c}.$$

Proof. The lemma follows from formulas (5.4) and (5.6) and Theorem 5.1. For $n = 2$

formula (3.1) takes the form:

$$R_k(a, b_1, b_2, c) = (-1)^{k_1+k_2} \prod_{i=1}^{k_1-k_2} \frac{(b_1 + (i - 1)c)!}{(1 + a + b_1 + (i + k_1 - 2)c - p)!} \\ \times \prod_{i=1}^{k_2} \frac{(b_2 + (i - 1)c)!}{(1 + b_2 + (i + k_2 - k_1 - 2)c)!} \frac{(1 + b_1 + b_2 + (i - 2)c)!}{(2 + a + b_1 + b_2 + (i + k_1 - 3)c - p)!} \\ \times \prod_{i=1}^{k_1} (a + (i - 1)c)! \prod_{i=1}^{k_2} (p + (i - k_1 - 1)c)! \prod_{r=1}^2 \prod_{i=1}^{k_r} \frac{(ic)!}{c!}.$$

Lemma 5.4. Assume that $(a, b_1 - 1, b_2, c), (a, b_1, b_2, c) \in A_k$. Then

$$(5.15) \quad R_{\mathbf{k}}(a, b_1 - 1, b_2, c) = R_{\mathbf{k}}(a, b_1, b_2, c) \prod_{i=1}^{k_1-k_2} \frac{1 + a + b_1 + (i + k_1 - 2)c - p}{b_1 + (i - 1)c} \\ \times \prod_{i=1}^{k_2} \frac{2 + a + b_1 + b_2 + (i + k_1 - 3)c - p}{1 + b_1 + b_2 + (i - 2)c}.$$

$$R_{\mathbf{k}}(a, b_1, b_2 - 1, c) = R_{\mathbf{k}}(a, b_1, b_2, c) \prod_{i=1}^{k_2} \frac{1 + b_2 + (i + k_2 - k_1 - 2)c}{b_2 + (i - 1)c} \\ \times \prod_{i=1}^{k_2} \frac{2 + a + b_1 + b_2 + (i + k_1 - 3)c - p}{1 + b_1 + b_2 + (i - 2)c}.$$

By Lemmas 5.3 and 5.4 the functions $S_{\mathbf{k}}(a, b_1, b_2, c)$ and $R_{\mathbf{k}}(a, b_1, b_2, c)$ defined on $A_{\mathbf{k}}$ satisfy the same difference equations with respects to the shifts of variables $b_1 \rightarrow b_1 - 1$ and $b_2 \rightarrow b_2 - 1$.

Lemma 5.5. Assume that a, c are positive integers such that $0 < k_1 c \leq p - 1$, $a + (k_1 - 1)c < p - 1$. Then the point

$$(5.17) \quad (a, b_1, b_2, c) = (a, p - 1 - (a + (k_1 - 1)c), (k_1 - k_2 + 1)c - 1, c)$$

lies in $A_{\mathbf{k}}$.

Proof. If (a, b_1, b_2, c) is given by (5.17), then

$$R_{\mathbf{k}} = (-1)^{k_1+k_2} \prod_{i=1}^{k_1-k_2} \frac{(p - 1 - (a + (k_1 - i)c))!}{((i - 1)c)!} \\ \times \prod_{i=1}^{k_2} \frac{((k_1 - k_2 + i)c - 1)!}{((i - 1)c)!} \frac{(p - 1 - (a + (k_2 - i)c))!}{((k_1 - k_2 + i - 1)c)!} \\ \times \prod_{i=1}^{k_1} (a + (i - 1)c)! \prod_{i=1}^{k_2} (p + (i - k_1 - 1)c)! \prod_{r=1}^2 \prod_{i=1}^{k_r} \frac{(ic)!}{c!}$$

This proves the lemma.

Lemma 5.6. Assume that $\tilde{a}, \tilde{c}$ are non-negative integers such that $0 < k_1 \tilde{c} \leq p - 1$, $\tilde{a} + (k_1 - 1)\tilde{c} \leq p - 1$. Denote by $A_{\mathbf{k}}(\tilde{a}, \tilde{c})$ the set of all $(a, b_1, b_2, c) \in A_{\mathbf{k}}$ such that $a = \tilde{a}$, $c = \tilde{c}$.

Then $A_{\mathbf{k}}(\tilde{a}, \tilde{c})$ consists of the pairs (b_1, b_2) of non-negative integers satisfying the inequalities

$$(5.18) \quad p - 1 - (\tilde{a} + (k_1 - 1)\tilde{c}) \leq b_1, \quad (k_1 - k_2 + 1)\tilde{c} - 1 \leq b_2$$

and some other inequalities of the form

$$(5.19) \quad b_1 \leq A_1, \quad b_2 \leq A_2, \quad b_1 + b_2 \leq A_{12},$$

where A_1, A_2, A_{12} are some integers such that

$$A_1 > p - 1 - (a + (k_1 - 1)c), \quad A_2 > (k_1 - k_2 + 1)c - 1, \quad A_{12} > p - 1 - (a + (k_1 - 1)c) + (k_1 - k_2 + 1)c - 1.$$

Proof. The lemma follows from Lemmas 3.1 and 3.2.

Corollary 5.7. Any point $(a, b_1, b_2, c) \in A_k(a, c)$ can be connected with the point $(a, p - 1 - (a + (k_1 - 1)c), (k_1 - k_2 + 1)c - 1, c) \in A_k(a, c)$ by a piece-wise linear path in $A_k(a, c)$ consisting of the vectors $(0, -1, 0, 0)$ or $(0, 0, -1, 0)$.

Proof of Theorem 3.4. For $n = 1$, $k = (k_1)$ and the point $(a, p - 1 - (a + (k_1 - 1)c), c)$ formula (3.11) holds by [RV, Theorem 4.1].

For $n = 2$, $k = (k_1, k_2)$ and the point $(a, p - 1 - (a + (k_1 - 1)c), (k_1 - k_2 + 1)c - 1, c)$ formula (3.11) holds by Lemma 5.5 and Theorem 3.7.

For $n = 2$, $k = (k_1, k_2)$ and arbitrary $(a, b_1, b_2, c) \in A_k(a, c)$ formula (3.11) holds by Lemmas 5.3, 5.4 and Corollary 5.7. Theorem 3.4 for $n = 2$ is proved. **Corollary 5.8.** Let $n > 2$, $k = (k_1, \dots, k_n)$, and $(a, (b_1, b_2), c) \in A_{(k_1, k_2)}$. Let $b = (b_1, \dots, b_n)$, where $b_i = (k_{i-1} - k_i + 1)c - 1$ for $i = 3, \dots, n$. Then formula (3.11) holds for the F_p -Selberg integral $S_{[k]_p}(a, b, c)$ of type A_n .

Proof. Formula (3.11) for the F_p -Selberg integrals of type A_2 is proved in Theorem 3.4. Hence the corollary follows from Theorem 3.7 by induction on n .

5.3. Evaluation of $I_{0,0,0}(a, b_1, b_2, c)$. In this section we evaluate $I_{0,0,0}(a, b_1, b_2, c)$ without using the evaluation of $S_k(a, b_1, b_2, c)$.

Theorem 5.9. Let $k = (k_1, k_2)$, $k_1 > k_2 > 0$ and $a, b_1, b_2, c \in \mathbb{Z}_{>0}$. Assume that $a + (k_1 - 1)c < p$ and all factorials on the right-hand side of the next formula are factorials of the non-negative integers less than p . Then

$$\begin{aligned} I_{0,0,0}(a, b_1, b_2, c) &= (-1)^{k_1+k_2} \prod_{i=1}^{k_1-k_2} \frac{(b_1 + (i-1)c)!}{(a + b_1 + (i + k_1 - 2)c - p)!} \\ &\times \prod_{i=1}^{k_2} \frac{(b_2 + (i-1)c)!}{(b_2 + (i + k_2 - k_1 - 2)c)!} \frac{(b_1 + b_2 + (i-2)c)!}{(a + b_1 + b_2 + (i + k_1 - 3)c - p)!} \\ &\times \prod_{i=1}^{k_1} (a + (i-1)c - 1)! \prod_{i=1}^{k_2} (p + (i - k_1 - 1)c - 1)! \prod_{r=1}^2 \prod_{i=1}^{k_i} \frac{(ic)!}{c!}. \end{aligned}$$

Proof. The proof is parallel to the proof of Theorem 3.4 for $n = 2$.

Denote by A_k^I the set of all $a, b_1, b_2, c \in \mathbb{Z}_{>0}$ satisfying the assumptions of Theorem 5.9. Notice that if $(a, b_1, b_2, c) \in A_k^I$, then

$$(5.21) \quad b_1 > p - (a + (k_1 - 1)c), \quad b_2 > (k_1 - k_2 + 1)c. \\ (a, b_1, (k_1 - k_2 + 1)c, c) \in \mathcal{A}_k^I$$

$$\prod_{j=1}^{k_n} s_j^{k_1 p - 1}$$

$$k_1$$

$$\Phi_{(k_1)}(t_1, \dots, t_{k_1}, a-1, b_1, c) = \prod_{16i < i^0 6k_1} Y(t_i - t_{i0})^{2c} \prod_{i=1} Y t_{ia-1} (1 - t_i)^{b_1}$$

$$\prod_{j=1}^{k_1} t_i^{p-1}$$

More precisely, under the assumptions of the theorem we have

$$I_{0,0,0}(a, b_1, (k_1 - k_2 + 1)c, c) = (-1)^{b_2 k_2 + c k_2 (k_2 - 1)/2} \frac{(k_2 c)!}{(c!)^{k_2}} S_{(k_1)}(a - 1, b_1, c)$$

cf. the proof of Theorem 3.7. We have $(-1)^{b_2 k_2 + c k_2 (k_2 - 1)/2} = (-1)^{(k_1 k_2 - k_2 (k_2 + 1)/2)c - k_2}$. By [RV,

Theorem 4.1] we have $S_{(k_1)}(a - 1, b_1, c) = R_{(k_1)}(a - 1, b_1, c)$. Hence

$$(5.22) \quad I_{0,0,0}(a, b_1, (k_1 - k_2 + 1)c, c) = (-1)^{(k_1 k_2 - k_2 (k_2 + 1)/2)c} (-1)^{k_1 + k_2} \\ \times \frac{(k_2 c)!}{(c!)^{k_2}} \prod_{j=1}^{k_1} \frac{(j c)!}{c!} \frac{(a + (j - 1)c - 1)! (b + (j - 1)c)!}{(a + b + (k_1 + j - 2)c - p)!}$$

$$R_{\mathbf{k}}^I(a, b_1, b_2, c$$

$$(5.23)$$

$$R_{\mathbf{k}}^I(a, b_1, (k_1 - k_2 + 1)c, c) = (-1)^{k_1 + k_2} \prod_{i=1}^{k_1 - k_2} \frac{(b_1 + (i - 1)c)!}{(a + b_1 + (i + k_1 - 2)c - p)!} \\ \times \prod_{i=1}^{k_2} \frac{((k_1 - k_2 + 1)c + (i - 1)c)!}{((k_1 - k_2 + 1)c + (i + k_2 - k_1 - 2)c)!} \\ \times \prod_{i=1}^{k_2} \frac{(b_1 + (k_1 - k_2 + 1)c + (i - 2)c)!}{(a + b_1 + (k_1 - k_2 + 1)c + (i + k_1 - 3)c - p)!} \\ \times \prod_{i=1}^{k_1} (a + (i - 1)c - 1)! \prod_{i=1}^{k_2} (p + (i - k_1 - 1)c - 1)! \prod_{r=1}^2 \prod_{i=1}^{k_i} \frac{(i c)!}{c!}.$$

$$= (-1)^{k_1+k_2} (-1)^{(k_1 k_2 - k_2(k_2+1)/2)c} \frac{(k_2 c)!}{(c!)^{k_2}} \prod_{j=1}^{k_1} \frac{(ic)!}{c!} \frac{(a + (j-1)c - 1)! (b + (j-1)c)!}{(a + b + (k_1 + j - 2)c - p)!}$$

where we used the cancellation Lemma 2.1 in the last step. Hence

$$I_{0,0,0}(a, b_1, (k_1 - k_2 + 1)c, c) = R_{\mathbf{k}}^I(a, b_1, (k_1 - k_2 + 1)c, c$$

$$R_{\mathbf{k}}^I(a, b_1, b_2, c$$

$$R_{\mathbf{k}}^I(a, b_1, b_2, c) \text{ on } \mathcal{A}_{\mathbf{k}}^I$$

Corollary 5.7.

5.4. Sketch of the proof of formula (3.11) for $n > 2$. The proof is parallel to the proof of Theorem 3.4.

$$\begin{aligned} V_{\lambda_1}^{\mathbb{F}_p} & \quad V_{\lambda_2}^{\mathbb{F}_p} \\ \lambda_1 &= -\frac{a}{c+p} \omega_1, \quad \lambda_2 = -\frac{b_1}{c+p} \omega_1 - \cdots - \frac{b_n}{c+p} \omega_n \end{aligned}$$

and highest weight vectors v_1, v_2 . Consider the PBW basis $B = (u)$ of the weight subspace

$$V_{\lambda_1}^{\mathbb{F}_p} \otimes V_{\lambda_2}^{\mathbb{F}_p} [\lambda_1 + \lambda_2 - \sum_{i=1}^n k_i \sigma_i$$

of that basis:

$$\begin{aligned} u_1 &= \frac{f_1^{k_1-k_2} [f_1, f_2]^{k_2-k_3} \cdots [f_1, [f_2, \dots, [f_{n-1}, f_n] \dots]]^{k_n} v_1 \otimes v_2}{(k_1 - k_2)! (k_2 - k_3)! \cdots (k_n)!} \\ u_2 &= \frac{f_1^{k_1} v_1 \otimes f_2^{k_2} \cdots f_n^{k_n} v_2}{(k_1)! (k_2)! \cdots (k_n)!}. \end{aligned}$$

For $n = 2$ these vectors are the vectors $v_{0,0,0}$ and $v_{0,k_2,0}$ in the proof of Theorem 5.1.

To any basis vector $u \in B$ we assign the weight function $W_u(t)$ defined in [RSV, Section 6.1], here t is the collection of variables defined in (3.9). Then we consider the $\mathbb{F}_p$ -integrals Z

$$I_u(a, b, c) = \int_{[k]_p} \Phi(t, a, b, c) W_u(t) dt.$$

It follows from the formulas for the weight functions that

$$I_{u_2}(a, b, c) = S_k(a - 1, b_1, b_2 - 1, \dots, b_n - 1, c),$$

cf. (5.4). It is known from the theory of KZ equations that the vector

$$I(a, b, c) = \sum_{u \in B} X_{I_u}(a, b, c) u$$

$$V_{\lambda_1}^{\mathbb{F}_p} \otimes V_{\lambda_2}^{\mathbb{F}_p} [\lambda_1 + \lambda_2 - \sum_{i=1}^n k_i \sigma_i$$

(5.24) $I_{u_2}(a, b, c) = B_0(a, b, c)I_{u_1}(a, b, c)$, where $B_0(a, b, c)$ is an explicit expression like in (5.6).

$$\sum_{i=1}^n k_i \sigma_i \quad V_{\lambda_1}^{\mathbb{F}_p} [\lambda_1 + \lambda_2 -$$

(5.25) $I_{u_1}(a, b_1, \dots, b_i - 1, \dots, b_n, c) = B_i(a, b, c)I_{u_1}(a, b, c)$, $i = 1, \dots, n$, where $B_i(a, b, c)$ are explicit products like in (5.7) and (5.8).

Equation (5.24) and difference equations (5.25) imply that the two functions $S_k(a, b, c)$ and $R_k(a, b, c)$, defined on the set A_k , satisfy the same difference equations with respect to the shift of variables $b_i \rightarrow b_i - 1$ for $i = 1, \dots, n$. By Corollary 3.8 we also know that the two functions are equal at the distinguished point

$$(a, p - 1 - (a + (k_1 - 1)c), (k_1 - k_2 + 1)c - 1, \dots, (k_{n-1} - k_n + 1)c - 1, c) \in A_k.$$

This implies that the two functions are equal (cf. Corollary 5.7) and formula (3.11) holds for any n . The details of this sketch will be published elsewhere.

References

- [AAR] G.E.Andrews, R.Askey, R.Roy, *Special functions*, Cambridge University Press, 1999
- [An] G.W. Anderson, *The evaluation of Selberg sums*, C. R. Acad. Sci. Paris S'ér. I Math. **311** (1990), no. 8, 469–472
- [Ao] K.Aomoto, *Jacobi polynomials associated with Selberg integral*, Siam J. Math, **18** (1987), No. 2, 545–549
- [As] R.Askey, *Some basic hypergeometric extensions of integrals of Selberg and Andrews*, SIAM J. Math. Anal. **11** (1980), 938–951
- [Ch] I.Cherednik, *From double Hecke algebra to analysis*, Doc.Math.J.DMV, Extra Volume ICM 1998,II, 527–531
- [EFK] P.Etingof, I.Frenkel, A.Kirillov, *Lectures on representation theory and Knizhnik-Zamolodchikov equations*, Mathematical Surveys and Monographs, 58, AMS, Providence, RI, 1998. xiv+198 pp. ISBN: 0-8218-0496-0
- [Ev] R.J.Evans, *The evaluation of Selberg character sums*, L'Enseign. Math. **37** (1991), 235–248
- [FSV] G.Felder, L.Stevens, and A.Varchenko, *Elliptic Selberg integrals and conformal blocks*, Math. Res. Lett. **10** (2003), no. 5–6, 671–684
- [FV] G.Felder and A.Varchenko, *Integral representation of solutions of the elliptic Knizhnik-Zamolodchikov-Bernard equations*, Int. Math. Res. Notices, 1995, N. **5**, 221–233

- [FW] P.J.Forrester and S.O.Warnaar, *The importance of the Selberg integral*, Bull. Amer. Math. Soc. (N.S.) **45** (2008), 489–534
- [Ha] L.Habsieger, *Une q -int'egrale de Selberg et Askey*, SIAM J. Math. Anal. **19** (1988), 1475–1489
- [Ka] J.Kaneko, *q -Selberg integrals and Macdonald polynomials*, Ann. Sci. Ecole Norm. Sup. **29** (1996), 583–637
- [KZ] V.Knizhnik and A.Zamolodchikov, *Current algebra and the Wess-Zumino model in two dimensions*, Nucl. Phys. **B247** (1984), 83–103
- [M] A.Matsuo, *An application of Aomoto-Gelfand Hypergeometric Functions to the $SU(n)$ Kniznik-Zamolodchikov Equation*, Comm. Math. Phys. **134** (1990), 65–77 (1990).
- [MaV] Y.Markov and A.Varchenko, *Hypergeometric Solutions of Trigonometric KZ Equations satisfy Dynamical Difference Equations*, Adv. Math. **166** (2002), no. 1, 100–147
- [MuV] E.Mukhin and A.Varchenko, *Remarks on critical points of phase functions and norms of Bethe vectors*, Adv. Stud. Pure Math. **27** (2000), 239–246
- [Op] E.M.Opdam, *Some applications of hypergeometric shift operators*, Invent. Math. **98** (1989), 1–18
- [R] E.Rains, *Multivariate Quadratic Transformations and the Interpolation Kernel*, Contribution to the Special Issue on Elliptic Hypergeometric Functions and Their Applications, SIGMA **14** (2018), 019, 69 pages, <https://doi.org/10.3842/SIGMA.2018.019>
- [RSV] R.Rimányi, L.Stevens, A. Varchenko, *Combinatorics of rational functions and Poincare-Birkhoff-Witt expansions of the canonical $U(n)$ -valued differential form*, Ann. Comb. **9** (2005), no. 1, 57–74
- [RTVZ] R.Rimányi, V.Tarasov, A.Varchenko, P.Zinn-Justin, *Extended Joseph polynomials, quantized conformal blocks, and a q -Selberg type integral*, Journal of Geometry and Physics, **62** (2012), pp. 2188–2207
- [RV] R.Rimányi, A. Varchenko, *The F_p -Selberg integral*, arXiv:2011.14248, 1–19
- [Se] A.Selberg, *Bemerkninger om et multipelt integral*, Norsk Mat. Tidsskr., **26** (1944), 71–78
- [SIV] A.Slinkin, A.Varchenko *Hypergeometric Integrals Modulo p and Hasse–Witt Matrices*, arXiv:2001.06869, 1–36
- [Sp] V.Spiridonov, *On the elliptic beta function*, (Russian) Uspekhi Mat. Nauk **56** (2001), no. 1, 181–182; translation in Russian Math. Surveys **56** (2001), no. 1, 185–186
- [SV1] V.Schechtman, A.Varchenko, *Arrangements of Hyperplanes and Lie Algebra Homology*, Invent. Math. **106** (1991), 139–194

- [SV2] V.Schechtman, A.Varchenko, *Solutions of KZ differential equations modulo p* , The Ramanujan Journal, **48** (3), 2019, 655–683, <https://doi.org/10.1007/s11139-018-0068-x>
- [TV1] V.Tarasov and A.Varchenko, *Geometry of q -hypergeometric functions as a bridge between Yangians and quantum affine algebras*, Invent. Math. **128** (1997), 501–588
- [TV2] V.Tarasov and A.Varchenko, *Geometry of q -hypergeometric functions, quantum affine algebras and elliptic quantum groups*, Asterisque, **246** (1997), 1–135
- [TV3] V.Tarasov and A.Varchenko, *Difference Equations Compatible with Trigonometric KZ Differential Equations*, IMRN, **15** (2000), 801–829
- [TV4] V.Tarasov and A.Varchenko, *Selberg-type integrals associated with $\mathfrak{sl}_3$* , Lett. Math. Phys. **65** (2003), 173–185
- [V1] A.Varchenko, *The Euler beta-function, the Vandermonde determinant, the Legendre equation, and critical values of linear functions on a configuration of hyperplanes, I.* (Russian) Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. **6**, 1206–1235, 1337; translation in Math. USSR-Izv. **35** (1990), no. 3, 543–571
- [V2] A.Varchenko, *Special functions, KZ type equations, and Representation theory*, CBMS, Regional Conference Series in Math., n. 98, AMS (2003)
- [V3] A.Varchenko, *Remarks on the Gaudin model modulo p* , arXiv:1708.06264, Journal of Singularities, **18** (2018), 486–499
- [V4] A.Varchenko, *Solutions modulo p of Gauss-Manin differential equations for multidimensional hypergeometric integrals and associated Bethe ansatz*, arXiv:1709.06189, Mathematics 2017, 5(4), 52; doi:10.3390/math5040052, 1–18
- [V5] A.Varchenko, *Hyperelliptic integrals modulo p and Cartier-Manin matrices*, arXiv:1806.03289, 1–16
- [V6] A.Varchenko, *An invariant subbundle of the KZ connection mod p and reducibility of $\mathfrak{sl}_{C^2}$ Verma modules mod p* , arXiv:2002.05834, 1–14
- [V7] A.Varchenko, *Determinant of F_p -hypergeometric solutions under ample reduction*, arXiv:2010.11275, 1–22
- [Wa1] S.Ole Warnaar, *A Selberg integral for the Lie algebra A_n* , Acta Math. **203** (2009), no. 2, 269–304
- [Wa2] S.Ole Warnaar, *The $\mathfrak{sl}_3$ Selberg integral*, Adv. Math. **224** (2010), no. 2, 499–524