

Determinacy from strong compactness of ω_1

Nam Trang*
Trevor M. Wilson†

January 14, 2021

Abstract

In the absence of the Axiom of Choice, the “small” cardinal ω_1 can exhibit properties more usually associated with large cardinals, such as strong compactness and supercompactness. For a local version of strong compactness, we say that ω_1 is X -strongly compact (where X is any set) if there is a fine, countably complete measure on $\wp_{\omega_1}(X)$. Working in $\text{ZF} + \text{DC}$, we prove that the $\wp(\omega_1)$ -strong compactness and $\wp(\mathbb{R})$ -strong compactness of ω_1 are equiconsistent with AD and $\text{AD}_{\mathbb{R}} + \text{DC}$ respectively, where AD denotes the Axiom of Determinacy and $\text{AD}_{\mathbb{R}}$ denotes the Axiom of Real Determinacy. The $\wp(\mathbb{R})$ -supercompactness of ω_1 is shown to be slightly stronger than $\text{AD}_{\mathbb{R}} + \text{DC}$, but its consistency strength is not computed precisely. An equiconsistency result at the level of $\text{AD}_{\mathbb{R}}$ without DC is also obtained.

Key words: strong compactness, supercompactness, core model induction, square sequences, coherent sequences, HOD, large cardinals, determinacy

2010 MSC: 03E15, 03E45, 03E55, 03E60

*University of North Texas, Denton, TX, USA. Email: ntrang@math.unt.edu

†Miami University, Oxford, Ohio, USA. Email: twilson@miamioh.edu

Contents

1	Introduction	3
2	Framework for the core model induction	8
2.1	Ω -premice, strategy premice, and g -organized Ω -premice	8
2.2	A very brief tale of hod mice	11
2.3	Core model induction operators	12
3	From Ω to $\mathcal{M}_1^{\sharp, \Omega}$	14
4	The maximal model of $\text{AD}^+ + \Theta = \theta_\Sigma$	21
5	A model of $\text{AD}^+ + \Theta > \theta_\Sigma$	27
6	Ω^* is constructibly closed	34
7	Further results, questions, and open problems	39

1 Introduction

We assume $\text{ZF} + \text{DC}$ as our background theory unless otherwise stated. (However, we will sometimes weaken our choice principle to a fragment of DC .) In this setting, it is possible for ω_1 to exhibit “large cardinal” properties such as strong compactness. The appropriate definition of strong compactness is made in terms of measures (ultrafilters) on sets of the form $\wp_{\omega_1}(X)$.

Definition 1.1. *Let X be an uncountable set. A measure μ on $\wp_{\omega_1}(X)$ is **countably complete** if it is closed under countable intersections and **fine** if it contains the set $\{\sigma \in \wp_{\omega_1}(X) : x \in \sigma\}$ for all $x \in X$. We say that ω_1 is X -strongly compact if there is a countably complete fine measure on $\wp_{\omega_1}(X)$.*

For uncountable sets X and Y , we will often use the elementary fact that if ω_1 is X -strongly compact and there is a surjection from X to Y , then ω_1 is Y -strongly compact as witnessed by a push-forward measure.

In the absence of AC , it may become necessary to consider degrees X of strong compactness that are not wellordered. The first and most important example is $X = \mathbb{R}$. The theory $\text{ZFC} + \text{“there is a measurable cardinal”}$ is equiconsistent with the theory $\text{ZF} + \text{DC} + \text{“}\omega_1\text{ is } \mathbb{R}\text{-strongly compact.”}$ (For a proof of the forward direction, see Trang [19]. The reverse direction is proved by noting that ω_1 is ω_1 -strongly compact, hence measurable, and considering an inner model $L(\mu)$ where μ is a measure on ω_1 .)

Another way to obtain \mathbb{R} -strong compactness of ω_1 that is more relevant to this paper is by the Axiom of Determinacy. If AD holds then by Martin’s cone theorem, for every set $A \in \wp_{\omega_1}(\mathbb{R})$ the property $\{x \in \mathbb{R} : x \leq_T d\} \in A$ either holds for a cone of Turing degrees d or fails for a cone of Turing degrees d , giving a countably complete fine measure on $\wp_{\omega_1}(\mathbb{R})$.

Besides \mathbb{R} , another relevant degree of strong compactness is the cardinal Θ , which is defined as the least ordinal that is not a surjective image of \mathbb{R} . In other words, Θ is the successor of the continuum in the sense of surjections. If the continuum can be wellordered then this is the same as the successor in the sense of injections (that is, \mathfrak{c}^+ .) However in general it can be much larger. For example, if AD holds then Θ is strongly inaccessible by Moschovakis’s coding lemma, but on the other hand there is no injection from ω_1 into \mathbb{R} .

If ω_1 is \mathbb{R} -strongly compact, then pushing forward a measure witnessing this by surjections, we see that ω_1 is λ -strongly compact for every uncountable cardinal $\lambda < \Theta$. In general all we can say is $\Theta \geq \omega_2$ and so this does not give anything beyond measurability of ω_1 . However, it does suggest two marginal strengthenings of the hypothesis “ ω_1 is \mathbb{R} -strongly compact” with the potential to increase the consistency strength beyond measurability. Namely, we may add the hypothesis “ ω_1 is ω_2 -strongly compact” or the hypothesis “ ω_1 is Θ -strongly

compact.” We will consider both strengthenings and obtain equiconsistency results in both cases.

In order to state and obtain sharper results, we first recall some combinatorial consequences of strong compactness. Let λ be an infinite cardinal and let $\vec{C} = (C_\alpha : \alpha \in \lim(\lambda))$ be a sequence such that each set C_α is a club subset of α . The sequence \vec{C} is *coherent* if for all $\beta \in \lim(\lambda)$ and all $\alpha \in \lim(C_\beta)$ we have $C_\alpha = C_\beta \cap \alpha$. A *thread* for a coherent sequence \vec{C} is a club subset $D \subset \lambda$ such that for all $\alpha \in \lim(D)$ we have $C_\alpha = D \cap \alpha$. An infinite cardinal λ is called *threadable* if every coherent sequence of length λ has a thread. Threadability of λ is also known as $\neg\Box(\lambda)$.

The following result is a well-known consequence of the “discontinuous ultrapower” characterization of strong compactness. However, without **AC** Łoś’s theorem may fail for ultrapowers of V , so we must verify that the argument can be done using ultrapowers of appropriate inner models instead.

Lemma 1.2. *Assume $\text{ZF} + \text{DC} + \text{“}\omega_1 \text{ is } \lambda\text{-strongly compact”}$ where λ is a cardinal of uncountable cofinality. Then λ is threadable.*

Proof. Let $\vec{C} = (C_\alpha : \alpha \in \lim(\lambda))$ be a coherent sequence such that each set C_α is a club in α . Consider the **ZFC** model $L[\{(\alpha, \beta) : \alpha \in C_\beta\}]$, which we abbreviate as $L[\vec{C}]$. Let μ be a countably complete fine measure on $\wp_{\omega_1}(\lambda)$ and let $j : L[\vec{C}] \rightarrow \text{Ult}(L[\vec{C}], \mu)$ be the corresponding ultrapower map, where the ultrapower is defined using all functions $\wp_{\omega_1}(\lambda) \rightarrow L[\vec{C}]$ in V . The ultrapower is wellfounded by countable completeness and **DC**, so it has the form $L[j(\vec{C})]$. Note that j is discontinuous at λ : for any ordinal $\alpha < \lambda$, we have $j(\alpha) \leq [\sigma \mapsto \sup \sigma]_\mu < j(\lambda)$ where the first inequality holds because μ is fine and the second inequality holds because λ has uncountable cofinality.

Now the argument continues as usual. We define the ordinal $\gamma = \sup j[\lambda]$ and note that $\gamma < j(\lambda)$ and that $j[\lambda]$ is an ω -club in γ . Therefore the set $j[\lambda] \cap \lim(j(\vec{C})_\gamma)$ is unbounded in γ , so its preimage $S = j^{-1}[\lim(j(\vec{C})_\gamma)]$ is unbounded in λ . Note that the club C_α is an initial segment of C_β whenever $\alpha, \beta \in S$ and $\alpha < \beta$; this is easy to check using the elementarity of j and the coherence of $j(\vec{C})$. Therefore the union of clubs $\bigcup_{\alpha \in S} C_\alpha$ threads the sequence \vec{C} . \square

If $\lambda < \Theta$ then **DC_R** suffices in place of **DC**:

Lemma 1.3. *Assume $\text{ZF} + \text{DC}_\mathbb{R} + \text{“}\omega_1 \text{ is } \mathbb{R}\text{-strongly compact.”}$ Let $\lambda < \Theta$ be a cardinal of uncountable cofinality. Then λ is threadable.*

Proof. Let \vec{C} be a coherent sequence of length λ . First, note that we may pass to an inner model containing \vec{C} where **DC** holds in addition to our other hypotheses. Namely,

let $f : \mathbb{R} \rightarrow \lambda$ be a surjection, let μ be a fine, countably complete measure on $\wp_{\omega_1}(\mathbb{R})$, let $C = \{(\alpha, \beta) : \alpha \in C_\beta\}$, and consider the model $M = L(\mathbb{R})[f, \mu, C]$, where the square brackets indicate that we are constructing from f , μ and C as predicates. (In the case of μ , this distinction is important: we are not putting all elements of μ into the model.)

It can be easily verified that all of our hypotheses are downward absolute to the model M , and that our desired conclusion that \vec{C} has a thread is upward absolute from M to V . In the model M every set is a surjective image of $\mathbb{R} \times \alpha$ for some ordinal α , so DC follows from $\text{DC}_{\mathbb{R}}$ by a standard argument. Moreover, ω_1 is λ -strongly compact in M by pushing forward the measure μ (restricted to M) by the surjection f , so the desired result follows from Lemma 1.2. \square

A further combinatorial consequence of strong compactness of ω_1 is the failure of Jensen's square principle \square_{ω_1} . In fact $\neg\square_{\omega_1}$ follows from the assumption that ω_2 is threadable or singular (note that successor cardinals may be singular in the absence of AC.)

Lemma 1.4. *Assume ZF. If ω_2 is singular or threadable, then $\neg\square_{\omega_1}$.*

Proof. Suppose toward a contradiction that ω_2 is singular or threadable and we have a \square_{ω_1} -sequence $(C_\alpha : \alpha \in \lim(\omega_2))$. If ω_2 is singular, we do not need coherence of the sequence to reach a contradiction. Take any cofinal set C_{ω_2} in ω_2 of order type $\leq\omega_1$ and recursively define a sequence of functions $(f_\alpha : \alpha \in [\omega_1, \omega_2])$ such that each function f_α is a surjection from ω_1 onto α , using our small cofinal sets C_α at limit stages. Then the function f_{ω_2} is a surjection from ω_1 onto ω_2 , a contradiction. On the other hand, if ω_2 is regular and threadable, take a thread C_{ω_2} through the square sequence. Then by the usual argument the order type of C_{ω_2} is at most $\omega_1 + \omega$, contradicting the regularity of ω_2 . \square

Now we can state our equiconsistency results and prove their easier directions.

Theorem 1.5. *The following theories are equiconsistent:*

1. ZF + DC + AD.
2. ZF + DC + “ ω_1 is $\wp(\omega_1)$ -strongly compact.”
3. ZF + DC + “ ω_1 is \mathbb{R} -strongly compact and ω_2 -strongly compact.”
4. ZF + DC + “ ω_1 is \mathbb{R} -strongly compact and $\neg\square_{\omega_1}$.”

Proof. (1) \implies (2): Under AD, Martin's cone theorem implies that ω_1 is \mathbb{R} -strongly compact. There is a surjection from \mathbb{R} onto $\wp(\omega_1)$ by Moschovakis's coding lemma, so ω_1 is $\wp(\omega_1)$ -strongly compact as well.

(2) \implies (3): This follows from the existence of surjections from $\wp(\omega_1)$ onto \mathbb{R} and ω_2 .

(3) \implies (4): This follows from Lemmas 1.2 and 1.4.

Con (4) \implies Con (1): In Sections 3 and 4, we will show that statement (4) implies $\text{AD}^{L(\mathbb{R})}$. \square

Moving up the consistency strength hierarchy, the next natural target for an equiconsistency result is the theory $\text{ZF} + \text{AD}_{\mathbb{R}}$. Here $\text{AD}_{\mathbb{R}}$ denotes the Axiom of Determinacy for real games, which has higher consistency strength than AD and cannot hold in $L(\mathbb{R})$. To get a model of $\text{AD}_{\mathbb{R}}$ we will need to augment our strong compactness hypothesis somehow, for example with a hypothesis on Θ or $\wp(\mathbb{R})$. For any set X , we write DC_X for the fragment of DC that allows us to choose ω -sequences of subsets of X .

Theorem 1.6. *The following theories are equiconsistent:*

1. $\text{ZF} + \text{AD}_{\mathbb{R}}$.

2. $\text{ZF} + \text{DC}_{\wp(\omega_1)} + \text{``}\omega_1 \text{ is } \mathbb{R}\text{-strongly compact and } \Theta \text{ is singular.``}$

Proof. Con (1) \implies Con (2): By Solovay [11], if $\text{ZF} + \text{AD}_{\mathbb{R}}$ is consistent then so is $\text{ZF} + \text{AD}_{\mathbb{R}} + \text{``}\Theta \text{ is singular.``}$ (In particular Solovay showed that the cofinality of Θ can be countable, which implies the failure of DC .) Under $\text{AD}_{\mathbb{R}}$ we have that ω_1 is \mathbb{R} -strongly compact by Martin's measure (this just follows from AD) and we have $\text{DC}_{\mathbb{R}}$ (this follows from uniformization for total relations on \mathbb{R} .) Moreover there is a surjection from \mathbb{R} to $\wp(\omega_1)$ by the coding lemma, so $\text{DC}_{\mathbb{R}}$ can be strengthened to $\text{DC}_{\wp(\omega_1)}$.

Con (2) \implies Con (1): In Sections 5 and 6, we will show that if statement (2) holds, then statement (1) holds in an inner model of the form $L(\Omega^*, \mathbb{R})$ where $\Omega^* \subset \wp(\mathbb{R})$. Note that statement (2) implies that ω_2 is either singular (if $\omega_2 = \Theta$) or threadable (if $\omega_2 < \Theta$, by Lemma 1.3) so in either case we have $\neg\Box_{\omega_1}$ by Lemma 1.4. Therefore we can make some use of the argument for Con (4) \implies Con (1) of Theorem 1.5 here, once we check that $\text{DC}_{\wp(\omega_1)}$ suffices in place of DC for this argument. \square

Finally, we will obtain an equiconsistency result at the level of $\text{ZF} + \text{DC} + \text{AD}_{\mathbb{R}}$. Note that this theory has strictly higher consistency strength than $\text{ZF} + \text{AD}_{\mathbb{R}}$. (By contrast, $\text{ZF} + \text{DC} + \text{AD}$ and $\text{ZF} + \text{AD}$ are equiconsistent by a theorem of Kechris.)

Theorem 1.7. *The following theories are equiconsistent:*

1. $\text{ZF} + \text{DC} + \text{AD}_{\mathbb{R}}$.

2. $\text{ZF} + \text{DC} + \text{``}\omega_1 \text{ is } \wp(\mathbb{R})\text{-strongly compact.``}$

3. $\text{ZF} + \text{DC} + \text{``}\omega_1 \text{ is } \mathbb{R}\text{-strongly compact and } \Theta\text{-strongly compact.``}$

4. $\text{ZF} + \text{DC} + \text{``}\omega_1 \text{ is } \mathbb{R}\text{-strongly compact and } \Theta \text{ is singular.``}$

Proof. $\text{Con}(1) \implies \text{Con}(2)$: By Solovay [11], under $\text{ZF} + \text{AD}_{\mathbb{R}}$ we have DC if and only if Θ has uncountable cofinality, and in a minimal model of $\text{ZF} + \text{DC} + \text{AD}_{\mathbb{R}}$ we have that Θ is singular of cofinality ω_1 . Assume that we are in such a minimal model of $\text{ZF} + \text{DC} + \text{AD}_{\mathbb{R}}$ and take a cofinal increasing function $\pi : \omega_1 \rightarrow \Theta$.

We can express $\wp(\mathbb{R})$ as an increasing union $\bigcup_{\alpha < \omega_1} \Gamma_\alpha$ where the pointclass Γ_α consists of all sets of reals of Wadge rank at most $\pi(\alpha)$. For each $\alpha < \omega_1$ there is a surjection from \mathbb{R} onto Γ_α , so ω_1 is Γ_α -strongly compact. Moreover, $\text{AD}_{\mathbb{R}}$ implies that there is a uniform way to choose, for each $\alpha < \omega_1$, a countably complete fine measure μ_α on $\wp_{\omega_1}(\Gamma_\alpha)$ witnessing this fact (namely the unique normal fine measure; see Woodin [23, Theorem 4].)

Using a countably complete nonprincipal measure ν on ω_1 (which exists because ω_1 is ω_1 -strongly compact) we can assemble these measures μ_α into a countably complete fine measure μ^* on $\wp_{\omega_1}(\wp(\mathbb{R}))$ as follows: for $A \subseteq \wp_{\omega_1}(\wp(\mathbb{R}))$, we say

$$A \in \mu^* \iff \forall^*_\nu \alpha A \cap \wp_{\omega_1}(\Gamma_\alpha) \in \mu_\alpha.$$

It's easy to verify that μ^* is countably complete because ν and the μ_α 's are countably complete. Likewise, it's easy to verify that μ^* is fine because ν is uniform and the μ_α 's are fine. Therefore the measure μ^* witnesses that ω_1 is $\wp(\mathbb{R})$ -strongly compact, so statement (2) holds (in our minimal model of $\text{ZF} + \text{DC} + \text{AD}_{\mathbb{R}}$.)

(2) \implies (3): This follows from the existence of surjections from $\wp(\mathbb{R})$ onto \mathbb{R} and Θ .

$\text{Con}(1) \implies \text{Con}(4)$: This follows by the aforementioned result of Solovay that in a minimal model of $\text{ZF} + \text{DC} + \text{AD}_{\mathbb{R}}$ the cardinal Θ is singular of cofinality ω_1 (and of course ω_1 is \mathbb{R} -strongly compact by Martin's measure.)

$\text{Con}(3) \vee \text{Con}(4) \implies \text{Con}(1)$: We will show in Sections 5 and 6 that if either statement (3) or statement (4) holds, then statement (1) holds in an inner model of the form $L(\Omega^*, \mathbb{R})$ where $\Omega^* \subset \wp(\mathbb{R})$. The proof of $\text{Con}(4) \implies \text{Con}(1)$ is similar to the proof of $\text{Con}(2) \implies \text{Con}(1)$ in Theorem 1.7, although one should note that the inner model $L(\Omega^*, \mathbb{R})$ does not simply absorb DC from V ; a bit more argument is required. \square

The authors would like to thank the referee for a careful reading of this article and for pointing out several minor errors. The first author would like to thank the NSF for its generous support through grant DMS-1849295.

2 Framework for the core model induction

This section is an adaptation of the framework for the core model induction developed in [10] and [9], which in turn build on earlier formulations in [7]. For more detailed discussions on the notions defined below as well as results concerning them, see [10] and [9]. The first subsection imports some terminology from the theory of hybrid mice developed in [10] and [9]. The terminology in this subsection will be used in Subsection 2.3 to define core model induction operators and will be needed in many other places in the paper. The reader may skip them on the first read and come back when needed. Subsection 2.2 summarizes the theory of hod mice developed in [3]. Subsection 2.3 defines core model induction operators which are the operators we will construct in this paper.

2.1 Ω -premice, strategy premice, and g -organized Ω -premice

For a complete theory of \mathcal{F} -premice for operators \mathcal{F} , the reader is advised to read [9]; for a detailed treatment of strategy mice, the reader is advised to read [10, Sections 2,3]. We will use the terminology from these sources from now on.¹

The definition below is essentially [10, Definition 3.8]. For explanations about the notations, see [10, Sections 2,3]. In the following definition, the objects $\Omega, \varphi, X, A, \kappa$ are defined as in [10, Section 3]. Roughly, Ω is either a κ -strategy or a mouse operator with nice condensation properties defined on a cone of H_κ above $A \in HC$, φ is a formula in the language of strategy premice, and X codes the pair (Ω, φ) .

Definition 2.1. *Let $t = (\Omega, \varphi, X, A, \kappa)$ be suitable and $\mathfrak{M} = \mathcal{M}_1^{X, \#}(A)$. We say that \mathfrak{M} generically interprets Ω ² iff there are formulas Φ, Ψ in \mathcal{L}^+ and some $\gamma > \delta^{\mathfrak{M}}$ such that $\mathfrak{M}|\gamma \vDash \Phi$ and for any non-dropping $\Lambda_{\mathfrak{M}}^{X, \kappa}$ -iterate \mathcal{N} of \mathfrak{M} via a countable tree \mathcal{T} based on $\mathfrak{M}|\delta^{\mathfrak{M}}$,³ any \mathcal{N} -cardinal δ , any $\gamma \in Ord$ such that $\mathcal{N}|\gamma \vDash \Phi \ \& \ \text{"}\delta \text{ is Woodin"\}$, and any g which is set-generic over $\mathcal{N}|\gamma$ (with $g \in V$), we have that $\mathcal{R} =_{\text{def}} (\mathcal{N}|\gamma)[g]$ is closed under Ω , and $\Omega \upharpoonright \mathcal{R}$ is defined over \mathcal{R} by Ψ . We say such a pair (Φ, Ψ) generically determines t (or just Ω).*

Let $A \in HC$ and let Ω be either an operator or an iteration strategy. We say that (Ω, A) (or just Ω) is nice iff (Ω, A) is suitable and $(t_{\Omega, A})_2$ generically interprets Ω .⁴ We say that

¹The theory of strategy mice can be developed as a special case of the general theory of operator mice in [9] but the authors of the papers decided to define strategy mice as \mathcal{J} -structures as this approach seems more convenient and gave the right notation for proving strong condensation properties of strategy mice like [10, Lemma 4.1].

²In [10, Definition 3.8], the terminology is: t determines itself on generic extensions. We will later define a notion of generic determination which is slightly different.

³ $\delta^{\mathfrak{M}}$ is the Woodin cardinal of \mathfrak{M} and $\Lambda_{\mathfrak{M}}^{X, \kappa}$ denotes the unique X -(0, κ)-iteration strategy for \mathfrak{M} .

⁴ $t_{\Omega, A}$ is a 5-tuple defined [10, page 27] and $(t_{\Omega, A})_2$ is the third component of $t_{\Omega, A}$.

(Φ, Ψ) generically determines (Ω, A) iff (Φ, Ψ) generically determines $t_{\Omega, A}$.

We fix a nice (Ω, A) (or just nice Ω ; we will at times ignore A), $X = (t_{\Omega, A})_2$, \mathfrak{M} , $\Lambda_{\mathfrak{M}} = \Lambda$, and (Φ, Ψ) for the rest of the section. We define $\mathcal{M}_1^X(A)$ from \mathfrak{M} in the standard way.

See [10, Section 3] for a proof that if $\Omega = \Sigma$ is a strategy (of a hod mouse or suitable mouse) with branch condensation and is fullness preserving with respect to mice in some sufficiently closed, determined pointclass Γ or if Σ is the unique strategy of a sound Y -mouse for some operator Y , $\mathcal{M}_1^{Y,\#}$ generically interprets Y , and Y condenses finely (see [9, Definition 3.18]) then \mathfrak{M} generically interprets Ω .

Definition 2.2 (Sargsyan, [3]). *Let M be a transitive structure. Let \dot{G} be the name for the generic $G \subseteq \text{Col}(\omega, M)$ and let $\dot{x}_{\dot{G}}$ be the canonical name for the real coding $\{(n, m) \mid G(n) \in G(m)\}$, where we identify G with $\bigcup G$. The tree \mathcal{T}_M for making M generically generic is the iteration tree \mathcal{T} on \mathfrak{M} of maximal length such that:*

1. \mathcal{T} is via Λ and is everywhere non-dropping.
2. $\mathcal{T} \upharpoonright o(M) + 1$ is the tree given by linearly iterating the first total measure of \mathfrak{M} and its images.
3. Suppose $\text{lh}(\mathcal{T}) \geq o(M) + 2$ and let $\alpha + 1 \in (o(M), \text{lh}(\mathcal{T}))$. Let $\delta = \delta(\mathcal{M}_\alpha^\mathcal{T})$ and let $\mathbb{B} = \mathbb{B}(M_\alpha^\mathcal{T})$ be the extender algebra of $M_\alpha^\mathcal{T}$ at δ . Then $E_\alpha^\mathcal{T}$ is the extender E with least index in $M_\alpha^\mathcal{T}$ such that for some condition $p \in \text{Col}(\omega, M)$, $p \Vdash \text{"There is a } \mathbb{B}\text{-axiom induced by } E\text{ which fails for } \dot{x}_{\dot{G}}\text{"}$.

Assuming that \mathfrak{M} is sufficiently iterable, then \mathcal{T}_M exists and has successor length.

The operator ${}^g\Omega$, defined in [10, Definition 3.42], and used in building g -organized Ω -premice, feeds in branches for such \mathcal{T}_M 's for various $\mathcal{M} \triangleleft \mathcal{N}$, where \mathcal{N} is a g -organized Ω -premouse. We will also ensure that being such a structure is first-order — other than wellfoundedness and the correctness of the branches — by allowing sufficient spacing between these branches (see [10, Remark 3.37]).

[10] also defines the notion Θ - g -organized Ω -premouse. The difference between the two hierarchies is very minor (see the remark 2.5). The main difference is that in the latter hierarchy, say \mathcal{N} is a Θ - g -organized Ω -premouse, and $\mathcal{M} \triangleleft \mathcal{N}$ is an “activation level”, i.e. branch information of \mathcal{T}_M is to be fed into the branch predicate of \mathcal{N} , if there is a level $\mathcal{M} \trianglelefteq \mathcal{R} \trianglelefteq \mathcal{N}$ such that $\mathcal{R} = (\mathcal{M}|o(\mathcal{M}) + \gamma, \mathbb{E}^{\mathcal{M}|o(\mathcal{M}) + \gamma}, [0, \gamma)_{\mathcal{T}_M})$, where $\gamma < \text{lh}(\mathcal{T}_M)$ and $\mathcal{R} \models \text{"}\Theta\text{ doesn't exist"\}$, then we stop feeding in further branch information of \mathcal{T}_M beyond the least such \mathcal{R} . The reader can again see [10, Section 3] for a more extensive treatment of these notions.

If x is a transitive set, then $o(x)$ is defined to be $x \cap \text{Ord}$. If \mathcal{M} is a (hybrid) premouse over a transitive set x , then $\rho_k(\mathcal{M})$ is the least ordinal ρ such that there is some set $A \subseteq [x \times \rho]^{<\omega}$ such that A is $r\Sigma_k(\mathcal{M})$ but $A \notin \mathcal{M}$.

Suppose (Ω, A) is nice (Ω can be a mouse operator or an iteration strategy).⁵ Suppose Γ is an inductive-like pointclass that is determined. Let $\mathfrak{M} = \mathcal{M}_1^{X,\sharp}(A)$ where $X = (t_{(\Omega,A)})_2$; later on in the paper, we occasionally write $\mathcal{M}_1^{\Omega,\sharp}(A)$ for \mathfrak{M} . $\text{Lp}^{\text{g}\Omega}(x)$ is defined as the stack of $\text{g}\Omega$ -premice \mathcal{M} over x such that \mathcal{M} is x -sound, there is some n such that $\rho_{n+1}(\mathcal{M}) \leq o(x) < \rho_n(\mathcal{M})$ and every countable, transitive \mathcal{M}^* embeddable into \mathcal{M} has an $\text{g}\Omega$ - (n, ω_1+1) -iteration strategy⁶ Δ for a transitive x . We define $\text{Lp}^{\text{g}\Omega, \Gamma}(x)$ similarly but demand additionally that $\Delta \in \Gamma$. For \mathcal{N} a $\text{g}\Omega$ -premouse, let $\text{Lp}_+^{\text{g}\Omega}(\mathcal{N})$ denotes the stack of all g -organized Ω -premice \mathcal{M} such that either $\mathcal{M} = \mathcal{N}$, or $\mathcal{N} \triangleleft \mathcal{M}$, \mathcal{N} is a strong cutpoint of \mathcal{M} , \mathcal{M} is $o(\mathcal{N})$ -sound, and there is $n < \omega$ such that $\rho_{n+1}(\mathcal{M}) \leq o(\mathcal{N}) < \rho_n(\mathcal{M})$ and \mathcal{M} is countably $\text{g}\Omega$ - (n, ω_1+1) -iterable above $o(\mathcal{N})$. We define $\text{Lp}_+^{\text{g}\Omega, \Gamma}(\mathcal{N})$ similarly. These notions can be generalized to $\text{g}\Omega$ or any other operator in an obvious way (cf. [10, Definition 2.43]). We define $\text{Lp}^{\text{g}\Omega}(x)$ etc similarly. Θ - g -organized Ω -mice over \mathbb{R} are important in the scales analysis generalizing Steel's work in $\text{Lp}(\mathbb{R})$ (see the remark below).

Definition 2.3. Let $Y \subseteq \mathbb{R}$. We say that Y is self-scaled iff there are scales on Y and $\mathbb{R} \setminus Y$ which are projective (i.e., Σ_n^1 for some $n < \omega$) in Y .

Definition 2.4. Suppose Ω is nice and $Y \subseteq \mathbb{R}$ is self-scaled. We define $\text{Lp}^{\text{g}\Omega}(\mathbb{R}, Y)$ as the stack of all g -organized Ω -mice \mathcal{N} over (H_{ω_1}, Y) (with parameter \mathfrak{M}). We similarly define $\text{Lp}^{\text{g}\Omega}(\mathbb{R}, Y)$ as the stack of all Θ - g -organized Ω -mice \mathcal{N} over (H_{ω_1}, Y) (with parameter \mathfrak{M}). We also say (Θ - g -organized) Ω -premouse over (\mathbb{R}, Y) to in fact mean over (H_{ω_1}, Y) .

Remark 2.5. Switching from the g -organized hierarchy to the Θ - g -organized hierarchy was for a purely technical purpose, so that various proofs concerning the scales analyses work out (it is not known to work for the g -organized hierarchy). The two hierarchies are very closely related. In fact, for Ω and Y as in Definition 2.4, $\wp(\mathbb{R}) \cap \text{Lp}^{\text{g}\Omega}(\mathbb{R}, Y) = \wp(\mathbb{R}) \cap \text{Lp}^{\text{g}\Omega}(\mathbb{R}, Y)$. Suppose \mathcal{M} is an initial segment of the first hierarchy and \mathcal{M} is E -active. Note that $\mathcal{M} \models \text{"}\Theta\text{ exists"}$ and $\mathcal{M} \upharpoonright \Theta$ is Ω -closed. By induction below $\Theta^{\mathcal{M}}$, $\mathcal{M} \upharpoonright \Theta^{\mathcal{M}}$ can be rearranged into an initial segment \mathcal{N}' of the second hierarchy. Above $\Theta^{\mathcal{M}}$, we simply copy the E -sequence and B -sequence⁷ from \mathcal{M} over to obtain an $\mathcal{N} \triangleleft \text{Lp}^{\text{g}\Omega}(\mathbb{R}, X)$ extending \mathcal{N}' . The converse is

⁵From now on, we typically say “let Ω be a nice operator” in place of this. So Ω is either a mouse operator in the sense of [9] or an iteration strategy as in [10].

⁶This implies, among other things, that iterates of \mathcal{M}^* according to the strategy are g -organized Ω -premice.

⁷The E -sequence is the extender sequence of \mathcal{M} and the B -sequence codes fragments of the strategy of \mathfrak{M} .

similar. Similarly, if Ω is such that $\text{Lp}^\Omega(\mathbb{R}, Y)$ is well-defined and Ω relativizes well, then $\wp(\mathbb{R}) \cap \text{Lp}^{\mathbb{g}\Omega}(\mathbb{R}, Y) = \wp(\mathbb{R}) \cap \text{Lp}^\Omega(\mathbb{R}, Y)$. See [10, Remark 4.11].

Let $\Omega = \Sigma$, where Σ is a nice iteration strategy that appears in core model induction applications, $A \in \text{HC}$ transitive such that $\mathcal{P} \in \mathcal{J}_1(A)$, X be defined from (Ω, A) as above, and suppose $\mathfrak{M} = \mathcal{M}_1^{X,\#}(A)$ exists. We have that \mathfrak{M} generically interprets (Ω, A) . Also, the core model induction will give us that the code of Ω , $\text{Code}(\Omega)$ (under a natural coding of subsets of HC by subsets of \mathbb{R}) is self-scaled. Thus, we can define $\text{Lp}^{\mathbb{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))$ as above (assuming sufficient iterability of \mathfrak{M}). A core model induction is then used to prove that there is a maximal constructibly closed initial segment \mathcal{M} of $\text{Lp}^{\mathbb{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))$ that satisfies AD^+ . What's needed to prove this is the scales analysis of $\text{Lp}^{\mathbb{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))$ from the optimal hypothesis (similar to those used by Steel; see [15] and [14]). This is carried out in [10]; we will not go into details here.

2.2 A very brief tale of hod mice

In this paper, a hod premouse \mathcal{P} is one defined as in [3]. The reader is advised to consult [3] for basic results and notations concerning hod premice and mice. Let us mention some basic first-order properties of a hod premouse \mathcal{P} . There is an ordinal $\lambda^\mathcal{P}$ and sequences $\langle(\mathcal{P}(\alpha), \Sigma_\alpha^\mathcal{P}) \mid \alpha < \lambda^\mathcal{P}\rangle$ and $\langle\delta_\alpha^\mathcal{P} \mid \alpha \leq \lambda^\mathcal{P}\rangle$ such that

1. $\langle\delta_\alpha^\mathcal{P} \mid \alpha \leq \lambda^\mathcal{P}\rangle$ is increasing and continuous and if $\alpha = 0$ or is a successor ordinal then $\mathcal{P} \models \delta_\alpha^\mathcal{P}$ is Woodin; no other \mathcal{P} -cardinals are Woodin cardinals of \mathcal{P} ;
2. $\mathcal{P}(0) = \text{Lp}_\omega(\mathcal{P}|\delta_0)^\mathcal{P}$; for $\alpha < \lambda^\mathcal{P}$, $\mathcal{P}(\alpha + 1) = (\text{Lp}_\omega^{\mathbb{g}\Sigma_\alpha^\mathcal{P}}(\mathcal{P}|\delta_\alpha))^\mathcal{P}$; for limit $\alpha \leq \lambda^\mathcal{P}$, $\mathcal{P}(\alpha) = (\text{Lp}_\omega^{\mathbb{g}\oplus_{\beta < \alpha}\Sigma_\beta^\mathcal{P}}(\mathcal{P}|\delta_\alpha))^\mathcal{P}$;
3. $\mathcal{P} \models \Sigma_\alpha^\mathcal{P}$ is a $(\omega, o(\mathcal{P}), o(\mathcal{P}))$ ⁸-strategy for $\mathcal{P}(\alpha)$ with hull condensation;
4. if $\alpha < \beta < \lambda^\mathcal{P}$ then $\Sigma_\beta^\mathcal{P}$ extends $\Sigma_\alpha^\mathcal{P}$.

Hod mice in this paper are g-organized; this is so that S -constructions work out smoothly as in the pure $L[\mathbb{E}]$ -case. We will write $\delta^\mathcal{P}$ for $\delta_{\lambda^\mathcal{P}}^\mathcal{P}$ and $\Sigma^\mathcal{P} = \bigoplus_{\beta < \lambda^\mathcal{P}} \Sigma_\beta^\mathcal{P}$. Note that $\mathcal{P}(0)$ is a pure extender model. Suppose \mathcal{P} and \mathcal{Q} are two hod premice. Then $\mathcal{P} \trianglelefteq_{\text{hod}} \mathcal{Q}$ if there is $\alpha \leq \lambda^\mathcal{Q}$ such that $\mathcal{P} = \mathcal{Q}(\alpha)$. We say then that \mathcal{P} is a *hod initial segment* of \mathcal{Q} . (\mathcal{P}, Σ) is a *hod pair* if \mathcal{P} is a hod premouse and Σ is a strategy for \mathcal{P} (acting on countable stacks of countable normal trees) such that $\Sigma^\mathcal{P} \subseteq \Sigma$ and this fact is preserved under Σ -iterations. Typically, we will construct hod pairs (\mathcal{P}, Σ) such that Σ has hull condensation, branch condensation, and is Γ -fullness preserving for some pointclass Γ .

⁸This just means $\Sigma_\alpha^\mathcal{P}$ acts on all stacks of ω -maximal, normal trees in \mathcal{P} .

The reader should consult [3] for the definition of $B(\mathcal{Q}, \Sigma)$, and $I(\mathcal{Q}, \Sigma)$. Roughly speaking, $B(\mathcal{Q}, \Sigma)$ is the collection of all hod pairs which are strict hod initial segments of a Σ -iterate of \mathcal{Q} and $I(\mathcal{Q}, \Sigma)$ is the collection of all Σ -iterates of Σ . In the case $\lambda^{\mathcal{Q}}$ is limit, $\Gamma(\mathcal{Q}, \Sigma)$ is the collection of $A \subseteq \mathbb{R}$ such that A is Wadge reducible to some Ψ for which there is some \mathcal{R} such that $(\mathcal{R}, \Psi) \in B(\mathcal{Q}, \Sigma)$. See [3] for the definition of $\Gamma(\mathcal{Q}, \Sigma)$ in the case $\lambda^{\mathcal{Q}}$ is a successor ordinal.

[3] constructs under AD^+ and the hypothesis that there are no models of “ $\text{AD}_{\mathbb{R}} + \Theta$ is regular” hod pairs that are fullness preserving, positional, commuting, and have branch condensation. Such hod pairs are particularly important for our computation as they are points in the direct limit system giving rise to HOD of AD^+ models. Under AD^+ , for hod pairs $(\mathcal{M}_{\Sigma}, \Sigma)$, if Σ is a strategy with branch condensation and $\vec{\mathcal{T}}$ is a stack on \mathcal{M}_{Σ} with last model \mathcal{N} , $\Sigma_{\mathcal{N}, \vec{\mathcal{T}}}$ is independent of $\vec{\mathcal{T}}$. Therefore, later on we will omit the subscript $\vec{\mathcal{T}}$ from $\Sigma_{\mathcal{N}, \vec{\mathcal{T}}}$ whenever Σ is a strategy with branch condensation and \mathcal{M}_{Σ} is a hod mouse. In a core model induction, we don’t quite have, at the moment $(\mathcal{M}_{\Sigma}, \Sigma)$ is constructed, an AD^+ -model M such that $(\mathcal{M}_{\Sigma}, \Sigma) \in M$ but we do know that every $(\mathcal{R}, \Lambda) \in B(\mathcal{M}_{\Sigma}, \Sigma)$ belongs to such a model. We then can show (using our hypothesis) that $(\mathcal{M}_{\Sigma}, \Sigma)$ belongs to an AD^+ -model.

2.3 Core model induction operators

Let

$$\Omega^* = \{A \subseteq \mathbb{R} \mid L(A, \mathbb{R}) \models \text{AD}^+\}.$$

We assume

(†): There is no model M containing all reals and ordinals such that $M \models \text{AD}_{\mathbb{R}} + \text{“}\Theta \text{ is regular”}$.

Under this smallness assumption, by work of G. Sargsyan in [3], Ω^* is a Wadge hierarchy and furthermore, if M is a model of AD^+ then M is a model of Strong Mouse Capturing (SMC). Operators that we construct in the core model induction will also have the following additional properties (besides being nice).

In the following, a transitive structure N is *closed* under an operator Ω if whenever $x \in \text{dom}(\Omega) \cap N$, then $\Omega(x) \in N$.

Definition 2.6 (relativizes well). *Let Ω be an a Y -mouse operator for some operator Y .⁹ We say that Ω relativizes well if there is a formula $\phi(x, y, z)$ such that for any $a, b \in \text{dom}(\Omega)$ such that $a \in L_1(b)$, whenever N is a transitive model of ZFC^- such that N is closed under Y and $a, b, \Omega(b) \in N$, then $\Omega(a) \in N$ and is the unique $x \in N$ such that $N \models \phi[x, a, \Omega(b)]$.*

⁹ Y may be the rud operator, in which case Ω is just a mouse operator in the usual sense.

Definition 2.7 (determines itself on generic extensions). *Suppose Ω is an operator. We say that Ω determines itself on generic extensions if there is a formula $\phi(x, y, z)$ and a parameter $c \in HC$ such that for any countable transitive structure N of ZFC^- such that N contains c and is closed under Ω , for any generic extension $N[g]$ of N in V , $\Omega \cap N[g] \in N[g]$ and is definable over $N[g]$ via (ϕ, c) , i.e. for any $e \in N[g] \cap \text{dom}(\Omega)$, $\Omega(e) = d$ if and only if d is the unique $d' \in N[g]$ such that $N[g] \models \phi[c, d', e]$.*

We are now in a position to introduce the core model induction operators that we will need in this paper. These are particular kinds of mouse operators (in the sense of [9, 3.43]) that are constructed during the course of the core model induction. These operators can be shown to satisfy the sort of condensation described in [9, Section 3] (e.g. condense finely), relativize well, and determine themselves on generic extensions.

Definition 2.8. *Let Γ be an inductive-like pointclass. For $x \in \mathbb{R}$, $C_\Gamma(x)$ denotes the set of all $y \in \mathbb{R}$ such that for some ordinal $\gamma < \omega_1$, y (as a subset of ω) is $\Delta_\Gamma(\{\gamma, x\})$.*

Let $x \in HC$ be transitive and let $f : \omega \rightarrow x$ be a surjection. Then $c_f \in \mathbb{R}$ denotes the code for (x, \in) determined by f . And $C_\Gamma(x)$ denotes the set of all $y \in HC \cap \wp(x)$ such that for all surjections $f : \omega \rightarrow x$ we have $f^{-1}(y) \in C_\Gamma(c_f)$.

Definition 2.9. *Let (Ω, A) be as above, $t \in HC$ with $\mathfrak{M} \in \mathcal{J}_1(t)$. Let $1 \leq k < \omega$. A premouse \mathcal{N} over t is Ω - Γ - k -suitable (or just k -suitable if Γ and Ω are clear from the context) iff there is a strictly increasing sequence $\langle \delta_i \rangle_{i < k}$ such that*

1. $\forall \delta \in \mathcal{N}, \mathcal{N} \models \text{“}\delta \text{ is Woodin}\text{”}$ if and only if $\exists i < k (\delta = \delta_i)$.
2. $o(\mathcal{N}) = \sup_{i < \omega} (\delta_{k-1}^{+i})^\mathcal{N}$.
3. If $\mathcal{N}|\eta$ is a strong cutpoint of \mathcal{N} then $\mathcal{N}|(\eta^+)^\mathcal{N} = \text{Lp}_+^{\text{g}\Omega, \Gamma}(\mathcal{N}|\eta)$.
4. Let $\xi < o(\mathcal{N})$, where $\mathcal{N} \models \text{“}\xi \text{ is not Woodin}\text{”}$. Then $C_\Gamma(\mathcal{N}|\xi) \models \text{“}\xi \text{ is not Woodin}\text{”}$.

We write $\delta_i^\mathcal{N} = \delta_i$; also let $\delta_{-1}^\mathcal{N} = 0$ and $\delta_k^\mathcal{N} = o(\mathcal{N})$.¹⁰

Let \mathcal{N} be 1-suitable and let $\xi \in o(\mathcal{N})$ be a limit ordinal, such that $\mathcal{N} \models \text{“}\xi \text{ isn't Woodin}\text{”}$. Let $Q \triangleleft \mathcal{N}$ be the Q-structure for ξ . If ξ is a strong cutpoint of \mathcal{N} then $Q \triangleleft \text{Lp}_+^{\text{g}\Omega, \Gamma}(\mathcal{N}|\xi)$ by 3. Assume now that \mathcal{N} is reasonably iterable. If ξ is a strong cutpoint of Q , our mouse capturing hypothesis combined with 4 gives that $Q \triangleleft \text{Lp}_+^{\text{g}\Omega, \Gamma}(\mathcal{N}|\xi)$. If ξ is an \mathcal{N} -cardinal then indeed ξ is a strong cutpoint of Q , since \mathcal{N} has only finitely many Woodins. If ξ is not a strong cutpoint of Q , then by definition, we do not have $Q \triangleleft \text{Lp}_+^{\text{g}\Omega, \Gamma}(\mathcal{N}|\xi)$. However, using

¹⁰We could also define a suitable premouse \mathcal{N} as a Θ -g-organized \mathcal{F} -premouse and all the results that follow in this paper will be unaffected.

$*$ -translation (see [13]), one can find a level of $\text{Lp}_+^{g\Omega, \Gamma}(\mathcal{N}|\xi)$ which corresponds to Q (and this level is in $C_\Gamma(\mathcal{N}|\xi)$).

If Ω is a nice operator and Σ is an iteration strategy for a Ω - Γ -1-suitable premouse \mathcal{P} such that Σ has branch condensation and is Γ -fullness preserving (for some pointclass Γ), then we say that (\mathcal{P}, Σ) is a Ω - Γ -suitable pair or just Γ -suitable pair or just suitable pair if the pointclass and/or the operator Ω is clear from the context (this notion of suitability is not related to the one mentioned in Definition 2.1).

Definition 2.10 (Core model induction operators). *Suppose (\mathcal{P}, Σ) is a \mathcal{G} - Ω^* -suitable pair for some nice operator \mathcal{G} or a hod pair such that Σ has branch condensation and is Ω^* -fullness preserving. Let $\Omega = \Sigma$ (note that Ω is suitable). Assume $\text{Code}(\Omega)$ is self-scaled. We say J is a Σ -core model induction operator or just a Σ -cmi operator if one of the following holds:*

1. *J is a nice Ω -mouse operator (or g -organized Ω -mouse operator) defined on a cone of H_{ω_1} above some $a \in H_{\omega_1}$. Furthermore, J condenses finely, relativizes well and determines itself on generic extensions.*
2. *For some $\alpha \in \text{OR}$ such that α ends either a weak or a strong gap in the sense of [15] and [10], letting $M = \text{Lp}^{g\Omega}(\mathbb{R}, \text{Code}(\Omega))||\alpha$ and $\Gamma = (\Sigma_1)^M$, $M \models \text{AD}^+ + \text{MC}(\Sigma)$.¹¹ For some transitive $b \in H_{\omega_1}$ and some 1-suitable (or more fully Ω - Γ -1-suitable) Ω -premouse \mathcal{Q} over b , $J = \Lambda$, where Λ is an (ω_1, ω_1) -iteration strategy for \mathcal{Q} which is Γ -fullness preserving, has branch condensation and is guided by some self-justifying-system (sjs) $\vec{A} = (A_i : i < \omega)$ such that for some real x , for each i , $A_i \in \text{OD}_{b, \Sigma, x}^M$ and \vec{A} seals the gap that ends at α .¹²*

3 From Ω to $\mathcal{M}_1^{\sharp, \Omega}$

Suppose (\mathcal{P}, Σ) is a \mathcal{G} - Ω^* -suitable pair for some nice operator \mathcal{G} such that Σ has branch condensation and is Ω^* -fullness preserving. (Recall that Ω^* is the pointclass of all sets of reals A such that $L(A, \mathbb{R}) \models \text{AD}^+$.) As a special case we also allow $(\mathcal{P}, \Sigma) = (\emptyset, \emptyset)$; the analysis of this special case is enough to prove Theorem 1.5. In this section we assume the strong hypothesis

$$\text{ZF} + \text{DC}_{\wp(\omega_1)} + \text{``}\omega_1 \text{ is } \mathbb{R}\text{-strongly compact and } \neg \square_{\omega_1}\text{.''}$$

¹¹ $\text{MC}(\Sigma)$ stands for Mouse Capturing relative to Σ which says that for $x, y \in \mathbb{R}$, x is $\text{OD}(\Sigma, y)$ (or equivalently x is $\text{OD}(\Omega, y)$) iff x is in some g -organized Ω -mouse over y . SMC is the statement that for every hod pair (\mathcal{P}, Σ) such that Σ is fullness preserving and has branch condensation, $\text{MC}(\Sigma)$ holds.

¹²This implies that \vec{A} is Wedge cofinal in $\text{Env}(\Gamma)$, where $\Gamma = \Sigma_1^M$. Note that $\text{Env}(\Gamma) = \wp(\mathbb{R})^M$ if α ends a weak gap and $\text{Env}(\Gamma) = \wp(\mathbb{R})^{\text{Lp}^\Sigma(\mathbb{R})|(\alpha+1)}$ if α ends a strong gap.

Note that this follows from any of the hypotheses of Theorems 1.5, 1.6, and 1.7.

Let Ω be a Σ -CMI operator. (If $(\mathcal{P}, \Sigma) = (\emptyset, \emptyset)$ then Ω is an ordinary CMI operator of the kind typically used in proving $\text{AD}^{L(\mathbb{R})}$.) We will use our strong hypothesis to obtain the $\mathcal{M}_1^{\Omega, \sharp}$ operator, which is the relativization of the \mathcal{M}_1^\sharp operator to a fine-structural hierarchy where the levels are obtained by repeated applications of the Ω operator (rather than the rud operator, as in ordinary mice. Basically, for each x in $\text{dom}(\Omega)$, if Ω is a strategy, $\mathcal{M}_1^{\Omega, \sharp}(x)$ is $\mathcal{M}_1^{X, \sharp}(x)$, where $X = (\Omega, \varphi_{\min})$ and φ_{\min} is defined as in [10, Definition 3.2] and otherwise $\mathcal{M}_1^{\Omega, \sharp}(x)$ is defined as in [9].)

The argument is similar to that used to obtain the ordinary \mathcal{M}_1^\sharp operator from the failure of square at a measurable cardinal in ZFC. The relativization of the standard arguments from \mathcal{M}_1^\sharp to $\mathcal{M}_1^{\Omega, \sharp}$ presents no special problems, but working without the Axiom of Choice requires a bit of care because ultrapowers of V may fail to satisfy Łoś's theorem. However, Łoś's theorem does hold for ultrapowers of wellordered inner models of V , and more generally for ultraproducts of families of inner models that are uniformly wellordered in the sense that there is a function associating to each model a wellordering of that model.

The relevance of Jensen's square principle \square_κ here is that it holds for all infinite cardinals κ in all Mitchell–Steel extender models (mice) by Schimmerling and Zeman [6, Theorem 2]. The proof of this result is sufficiently abstract that it relativizes from mice to Ω -mice in a straightforward manner. Therefore if \square_κ fails in V , we get a failure of covering: the successor of κ cannot be computed correctly by any Ω -mouse.

Because we are not assuming the Axiom of Choice, we will not construct the core model in V but rather in an inner model H of V satisfying ZFC. This model H will be obtained as a kind of HOD. A method used by Schimmerling and Steel [5] to prove covering results for the core model of V can be adapted to the core model of H , provided that we can show that H is close enough to V in the relevant sense. We show this closeness by using Vopěnka's theorem, similar to Schindler [8].

The following lemma is the main result of this section. It will form the “successor step” in the proofs of the main theorems.

Lemma 3.1. *Assume $\text{ZF} + \text{DC}_{\wp(\omega_1)} + \text{“}\omega_1 \text{ is } \mathbb{R}\text{-strongly compact and } \neg \square_{\omega_1}\text{.”}$ Let (\mathcal{P}, Σ) be a \mathcal{G} - Ω^* -suitable pair for some nice operator \mathcal{G} , a hod pair such that Σ has branch condensation and is Ω^* -fullness preserving, or (\emptyset, \emptyset) . Let Ω be a Σ -CMI operator defined on a cone in $H(\omega_1)$ over some element $a \in H(\omega_1)$. Then for every element x of this cone, $\mathcal{M}_1^{\Omega, \sharp}(x)$ exists.*

Proof. First, note that we may assume without loss of generality that full DC holds, by passing to the inner model $L(\wp(\omega_1), \Sigma, \Omega)[\mu]$ where we are constructing relative to a predicate μ for a fine countable complete measure on $\wp_{\omega_1}(\mathbb{R})$. The hypothesis and conclusion are

absolute to this inner model. In particular the model satisfies $\neg\Box_{\omega_1}$ because it computes ω_2 correctly, and it satisfies $DC_{\wp(\omega_1)}$ because it contains all countable sequences from $\wp(\omega_1)$. In the inner model, this fragment of DC implies full DC by a standard argument using the fact that every set is the surjective image of $\wp(\omega_1) \times \alpha$ for some ordinal α . Therefore we may use DC in the argument that follows.

Note that because ω_1 is measurable, the operators Ω^\sharp and $\Omega^{\sharp\sharp}$ are also defined on the cone in $H(\omega_1)$ over a . Let $x \in H(\omega_1)$ be in the cone over a . Take a countably complete fine measure μ on $\wp_{\omega_1}(\mathbb{R})$. Because μ -almost every set σ contains a real coding x , for such σ we can define the inner model

$$H_\sigma = \text{HOD}_{\{\Omega, x\}}^{L^{\Omega^\sharp}(\sigma)}.$$

A few remarks on notation: The model $L^{\Omega^\sharp}(\sigma)$ is the proper class model that is obtained by iterating the top measure of $\Omega^{\sharp\sharp}(\sigma)$ out of the universe. It is closed under its version of Ω even above the point ω_1^V up to which Ω was originally defined; however, we will only ever use the Ω operator of the model $L^{\Omega^\sharp}(\sigma)$ up to the least indiscernible of that model, which is the critical point of the top measure of $\Omega^{\sharp\sharp}(\sigma)$ and is countable in V . By the parameter Ω in the definition of H_σ , we really mean the restriction of Ω to the model $L^{\Omega^\sharp}(\sigma)$, which is amenable to that model because Ω relativizes well. There will not be any incompatibility between the various restrictions and extensions of Ω that we use, so we denote them all by “ Ω ”.

Let ξ_σ denote the least indiscernible of $L^{\Omega^\sharp}(\sigma)$. Note that in the model H_σ we can do core model theory below ξ_σ : it is well-known that the existence of an external measure can substitute for measurability of ξ_σ in this regard. The operator Ω is amenable to H_σ (again because it relativizes well) and we can attempt the $K^{c,\Omega}(x)$ construction in H_σ up to the cardinal ξ_σ . This is like the ordinary K^c construction, except relativized to Ω and built over the set x (see [9, Definition 3.28] and [10, Definition 2.46]). By the K^Ω existence dichotomy (see Schindler and Steel [7]) applied in the various models H_σ , one of the following two cases holds:

1. For μ -almost every set $\sigma \in \wp_{\omega_1}(\mathbb{R})$, the model H_σ satisfies the statement that $\mathcal{M}_1^{\Omega,\sharp}(x)$ exists and is ξ_σ -iterable by the (unique) Ω^\sharp -guided strategy.
2. For μ -almost every set $\sigma \in \wp_{\omega_1}(\mathbb{R})$, the model K_σ , defined as the core model $(K^\Omega(x))^{H_\sigma}$ built up to ξ_σ , exists and has no Woodin cardinals.

Claim 3.2. *If case (1) of the K^Ω existence dichotomy holds, then $\mathcal{M}_1^{\Omega,\sharp}(x)$ exists in V .*

Proof. For μ -almost every set $\sigma \in \wp_{\omega_1}(\mathbb{R})$, the premouse $(\mathcal{M}_1^{\Omega,\sharp}(x))^{H_\sigma}$ exists by the case hypothesis. It is sound and projects to x , so it codes itself as a subset of x , which is

countable. Therefore by the countable completeness of μ we can fix a single Ω -premouse \mathcal{M} over x such that $\mathcal{M} = (\mathcal{M}_1^{\Omega,\sharp}(x))^{H_\sigma}$ for μ -almost every set σ . We will show that \mathcal{M} is ω_1 -iterable in V by the (unique) Ω^\sharp -guided iteration strategy. Then $(\omega_1 + 1)$ -iterability will follow by the measurability of ω_1 .

Let \mathcal{T} be a countable Ω^\sharp -guided putative iteration tree on \mathcal{M} in V , where by “putative” we mean that its last model, if it has one, may fail to be an Ω -premouse. (Note that an Ω -premouse is required in particular to be wellfounded, and this is the only requirement if $\Omega = \text{rud}$.) We want to show that if \mathcal{T} has successor length, then its last model is an Ω -premouse, and if it has limit length, then it has a cofinal branch b such that $\mathcal{M}_b^{\mathcal{T}}$ is an Ω -premouse and $\mathcal{Q}(b, \mathcal{T}) \trianglelefteq \Omega^\sharp(\mathcal{M}(\mathcal{T}))$.

Take a real t that codes \mathcal{T} . Then for μ -almost every set σ we have $t \in \sigma$ by the fineness of μ . Fix a set σ such that H_σ satisfies the statement “ $\mathcal{M}_1^{\Omega,\sharp}(x)$ exists and is ξ_σ -iterable,” $(\mathcal{M}_1^{\Omega,\sharp}(x))^{H_\sigma} = \mathcal{M}$, and $t \in \sigma$. By Vopěnka’s theorem applied in the model $L^{\Omega^\sharp}(\sigma)$, the real t is contained in a generic extension $H_\sigma[g]$ of H_σ . In fact because ξ_σ is inaccessible in $L^{\Omega^\sharp}(\sigma)$ the poset from the proof of Vopěnka’s theorem (see, for example, Jech [2, Theorem 15.46]) is in $(V_{\xi_\sigma})^{H_\sigma}$.

In H_σ the Ω -premouse \mathcal{M} is ξ_σ -iterable by the Ω^\sharp -guided strategy, by our assumptions. Because the Ω^\sharp operator condenses finely (cf. [9, Section 3])¹³ and determines itself on generic extensions,¹⁴ a standard argument (see Schindler and Steel [7, Lemma 2.7.2]) shows that \mathcal{M} is still ξ_σ -iterable in $H_\sigma[g]$ by the Ω^\sharp -guided iteration strategy there. We note here that since ξ_σ is countable, we really apply generic interpretability of Ω^\sharp to a countable submodel of H_σ , namely $V_{\xi_\sigma}^{H_\sigma}$.

The model $H_\sigma[g]$ sees that the tree \mathcal{T} is Ω^\sharp -guided. Therefore in $H_\sigma[g]$, if \mathcal{T} has successor length, then the last model of \mathcal{T} is a wellfounded Ω -premouse, and if \mathcal{T} has limit length, then it has a cofinal branch b such that $\mathcal{M}_b^{\mathcal{T}}$ is an Ω -premouse and $\mathcal{Q}(b, \mathcal{T}) \trianglelefteq \Omega^\sharp(\mathcal{M}(\mathcal{T}))$. In either case this fact about \mathcal{T} is absolute to V , giving the desired iterability. \square

Claim 3.3. *Case (2) of the K^Ω existence dichotomy cannot hold.*

Proof. This case is where the hypothesis $\neg \square_{\omega_1}$ is used. Because H_σ is defined as the $\text{HOD}_{\{\Omega, x\}}$ of $L^{\Omega^\sharp}(\sigma)$, we can define the Vopěnka poset $\mathbb{P}_\sigma \in H_\sigma$ to make every countable set of countable ordinals in $L^{\Omega^\sharp}(\sigma)$ generic over H_σ . For a countable set of countable ordinals a of $L^{\Omega^\sharp}(\sigma)$, let $g_{\sigma,a}$ denote the H_σ -generic filter over \mathbb{P}_σ induced by a , which has the property that

¹³This is a more detailed version of “condenses well” in the literature.

¹⁴In the “gap in scales” case, the proof that the Ω^\sharp operator determines itself on generic extensions is given by Schindler and Steel [7, Section 5.6, proof of Claim 1 in case $n = 0$]. The proof in the other cases is a straightforward induction.

$a \in H_\sigma[g_{\sigma,a}]$.¹⁵ Note that $\mathbb{P}_\sigma \in (V_{\xi_\sigma})^{H_\sigma}$ because ξ_σ is inaccessible in $L^{\Omega^\sharp}(\sigma)$.

Define the ultraproducts

$$\begin{aligned} H &= [\sigma \mapsto H_\sigma]_\mu & \Xi &= [\sigma \mapsto \xi_\sigma]_\mu \\ K &= [\sigma \mapsto K_\sigma]_\mu & \mathbb{P} &= [\sigma \mapsto \mathbb{P}_\sigma]_\mu. \end{aligned}$$

Every countable set of countable ordinals a in V is seen as a countable set of countable ordinals in $L^{\Omega^\sharp}(\sigma)$ for μ -almost every σ (by fineness applied to a real coding a) so we can define the ultraproduct

$$g_a = [\sigma \mapsto g_{\sigma,a}]_\mu.$$

Then applying Loś's theorem to uniformly wellordered families of structures is enough to establish the following facts.¹⁶

- H is an inner model of ZFC with a cardinal $\Xi > \omega_1^V$ that is large enough to do core model theory below it.
- K is the core model of H built up to Ξ , and it has no Woodin cardinals.
- $\mathbb{P} \in (V_\Xi)^H$ is a poset.
- To each countable set of countable ordinals a in V we have assigned an H -generic filter $g_a \subset \mathbb{P}$ such that $a \in H[g_a]$.

Now let $\kappa = \omega_1^V$ and define the μ -ultrapower map

$$j : V \rightarrow \text{Ult}(V, \mu), \quad \text{crit}(j) = \kappa.$$

Recall that j itself is not elementary, but its restrictions to wellordered inner models are elementary. (We remark that one could use any ultrapower map with critical point κ here; the measurability of ω_1^V suffices for the following argument in place of \mathbb{R} -strong compactness of ω_1^V , although it is not clear that it would suffice for the previous argument.)

Note that to every set $A \subset \kappa$ in V we can assign a $j(H)$ -generic filter $g_A \subset j(\mathbb{P})$ such that $A \in j(H)[g_A]$. To see this, consider the sequence of generic filters $\vec{g}_A = (g_{A \cap \alpha} : \alpha < \kappa)$, use the elementarity of the map $j \upharpoonright L[H, A, \vec{g}]$, and define $g_A = j(\vec{g}_A)_\kappa$.

¹⁵Unlike in case (1), it is important here that the Vopěnka generic filter $g_{\sigma,a}$ is induced by a itself and does not depend on the choice of a real coding a .

¹⁶If the measure μ were normal, then Loś's theorem could be applied to the models $L^{\Omega^\sharp}(\sigma)$ themselves to yield a model $L^{\Omega^\sharp}(\mathbb{R})$ in which H , K , Ξ , and \mathbb{P} could then be defined. But this is not possible in general, for example under $\text{AD} + V = L(\mathbb{R})$, where the hypothesis of the lemma holds for $\Omega = \text{rud}$ but \mathbb{R}^\sharp does not exist.

Because \square_κ fails in V , we have

$$(\kappa^+)^{j(K)} < (\kappa^+)^V$$

by a result of Schimmerling and Zeman [6, Theorem 2] relativized to the operator $j(\Omega)$ and applied to the model $j(K)$, which is the core model of $j(H)$.

Take a set $A \subset \kappa$ in V coding a wellordering of κ of order type $(\kappa^+)^{j(K)}$ and define $g = g_A$. Because $A \in j(H)[g]$ we get

$$j(H)[g] \vDash (\kappa^+)^{j(K)} < \kappa^+.$$

Because g was added by a small forcing below the large cardinal $j(\Xi)$ where $j(K)$ was constructed, we have that $j(K)$ is still the core model of $j(H)[g]$.¹⁷ Therefore (and this is the crucial point) the model $j(H)[g]$ sees the failure of covering for its own core model at κ , so we can apply the map j once more to get a contradiction by a standard argument, outlined below.

Consider the restriction

$$j \upharpoonright j(H)[g] : j(H)[g] \rightarrow j(j(H))[j(g)],$$

which is an elementary embedding. Because the domain $j(H)[g]$ satisfies $(\kappa^+)^{j(K)} < \kappa^+$, the further restriction $j \upharpoonright \wp(\kappa)^{j(K)}$ is in the codomain $j(j(H))[j(g)]$ by a standard argument due to Kunen. Therefore we have

$$F \in j(j(H))[j(g)]$$

where F is the $(\kappa, j(\kappa))$ -extender over $j(K)$ derived from the map $j \upharpoonright \wp(\kappa)^{j(K)}$. Note that $K|\kappa = j(K)|\kappa$, and κ is an inaccessible cardinal in both ZFC models K and $j(K)$ because it is a measurable cardinal in V . Therefore $j(K)|j(\kappa) = j(j(K))|j(\kappa)$, and $j(\kappa)$ is an inaccessible cardinal in both models $j(K)$ and $j(j(K))$, so we have

$$\begin{aligned} (\kappa^+)^{j(K)} &= (\kappa^+)^{j(j(K))} < j(\kappa) \quad \text{and} \\ \wp(\kappa)^{j(K)} &= \wp(\kappa)^{j(j(K))}. \end{aligned}$$

Therefore the extender F can also be considered as an extender over $j(j(K))$, and it coheres with $j(j(K))$. Note that $j(j(K))$ is the core model of $j(j(H))[j(g)]$

This extender F has superstrong type, and we can apply the maximality property of the

¹⁷To make sense of the core model of $j(H)[g]$ we are using the fact that $j(H)$'s version of the operator Ω determines itself on generic extensions. Any failure of this gets reflected to a countable substructure N . By fine condensation, the version of Ω in N is in fact $\Omega \cap N$. Now we apply the fact that Ω determines itself on generic extensions of N to get a contradiction.

core model [5, Theorem 2.3] in the model $j(j(H))[j(g)]$ to show that every proper initial segment $F \upharpoonright \nu$ of F , where $\nu < j(\kappa)$, is on the sequence of the core model $j(j(K))$. Then in the core model $j(j(K))$, these initial segments will witness that κ is a Shelah cardinal. This will contradict our case hypothesis, which says that there are no Woodin cardinals in K .

Let $\mathcal{M} = j(j(K))$ and let $F \upharpoonright \nu$, where $\nu < j(\kappa)$, be a proper initial segment of F . We want to see that the extender $F \upharpoonright \nu$ is on the \mathcal{M} -sequence. Without loss of generality we may assume that ν is at least the common κ^+ of the models $j(K)$ and \mathcal{M} . It suffices to show that the pair $(\mathcal{M}, F \upharpoonright \nu)$ is weakly countably certified [5, Definition 2.2]. Working in the model $j(H)[g]$, take a transitive, power admissible set N such that $N^\omega \subset N$, $V_\kappa \cup j(K) \cap ((\kappa^+)^{j(K)} + 1) \subset N$, and $|N| = \kappa$. Stepping out to V for a moment and applying Kunen's argument again, we have

$$G \in j(j(H))[j(g)]$$

where G is the $(\kappa, j(\kappa))$ -extender over N derived from $j \upharpoonright \wp(\kappa)^N$. Now in the model $j(j(H))[j(g)]$ it is easy to verify that the pair (N, G) is a weak \mathcal{A} -certificate [5, Definition 2.1] for $(\mathcal{M}, F \upharpoonright \nu)$ whenever \mathcal{A} is a countable subset of $\bigcup_{n < \omega} \wp([\kappa]^n) \cap \mathcal{M} \upharpoonright \nu$,¹⁸ noting that $\mathcal{M} \upharpoonright \nu$, \mathcal{M} , and $j(K)$ all have the same subsets of $[\kappa]^n$ (because ν is greater than or equal to the common κ^+ of $j(K)$ and \mathcal{M}). \square

We have shown that if case (1) of the K^Ω existence dichotomy holds, then the conclusion of the lemma holds, and we have shown that case (2) contradicts the hypothesis of the lemma, so the proof of the lemma is complete. \square

We remark that because Ω is a Σ -CMI operator, the operator $\mathcal{M}_1^{\Omega, \sharp}$ given by the lemma is also a Σ -CMI operator.

Corollary 3.4. *Assume $\text{ZF} + \text{DC}_{\wp(\omega_1)} + \text{``}\omega_1 \text{ is } \mathbb{R}\text{-strongly compact and } \neg \square_{\omega_1}\text{''}$. Then PD holds.*

Proof. We show by induction on $n < \omega$ that the \mathcal{M}_n^\sharp operator is total on $H(\omega_1)$. The base case is the \mathcal{M}_0^\sharp operator, meaning the ordinary sharp operator, which is total on $H(\omega_1)$ because ω_1 is measurable. For the induction step we apply Lemma 3.1 to go from the operator $\Omega = \mathcal{M}_n^\sharp$ to the operator $\mathcal{M}_1^{\Omega, \sharp}$, which is stronger than \mathcal{M}_{n+1}^\sharp . It follows from the existence of $\mathcal{M}_n^\sharp(x)$ for every $n < \omega$ and $x \in \mathbb{R}$ that Projective Determinacy holds. \square

In the next section we will strengthen this conclusion to $\text{AD}^{L(\mathbb{R})}$ and thereby obtain an equiconsistency result (Theorem 1.5.)

¹⁸Or indeed if \mathcal{A} is equal to $\bigcup_{n < \omega} \wp([\kappa]^n) \cap \mathcal{M} \upharpoonright \nu$ itself; we don't need countability, and we don't need to choose the certificate (N, G) differently depending on \mathcal{A} (or on ν , for that matter.)

4 The maximal model of $\text{AD}^+ + \Theta = \theta_\Sigma$

Throughout this section, we assume the hypothesis of Lemma 3.1, namely we assume

$$\text{ZF} + \text{DC}_{\wp(\omega_1)} + \text{``}\omega_1 \text{ is } \mathbb{R}\text{-strongly compact and } \neg \square_{\omega_1}\text{.''}$$

Suppose (\mathcal{P}, Σ) is a \mathcal{G} - Ω^* -suitable pair for some nice operator \mathcal{G} such that Σ has branch condensation and is Ω^* -fullness preserving. As a special case we also allow $(\mathcal{P}, \Sigma) = (\emptyset, \emptyset)$; the analysis of this special case is enough to prove Theorem 1.5. We first define the “maximal pointclass of $\text{AD}^+ + \Theta = \theta_\Sigma$ ”.

Definition 4.1. *Let (\mathcal{P}, Σ) be as above. Let*

$$\Omega_\Sigma = \bigcup \{\wp(\mathbb{R}) \cap L(A, \mathbb{R}) \mid A \subseteq \mathbb{R} \text{ and } L(A, \mathbb{R}) \models \text{AD}^+ + \Theta = \theta_\Sigma + \text{MC}(\Sigma)\}.$$

We note that by (†), Ω_Σ is a Wadge hierarchy. In the case $(\mathcal{P}, \Sigma) = (\emptyset, \emptyset)$, substitute θ_0 for θ_Σ and ordinary mouse capturing MC for $\text{MC}(\Sigma)$. In this section, we will prove that

$$L(\Omega_\Sigma, \mathbb{R}) \cap \wp(\mathbb{R}) = \Omega_\Sigma. \quad (4.1)$$

This has the consequence that $L(\Omega_\Sigma, \mathbb{R}) \models \text{AD}^+ + \Theta = \theta_\Sigma$. The model $L(\Omega_\Sigma, \mathbb{R})$ is called the “maximal model of $\text{AD}^+ + \Theta = \theta_\Sigma$ ”.

Let $\Omega = \Sigma$. The proof of (4.1) depends on understanding models of $\text{ZF} + \text{AD}^+ + V = L(\wp(\mathbb{R})) + \Theta = \theta_\Sigma + \text{MC}(\Sigma)$ as hybrid mice over \mathbb{R} , Θ -g-organized as in Section 2.1. (In the case $(\mathcal{P}, \Sigma) = (\emptyset, \emptyset)$, we consider ordinary mice over \mathbb{R} , namely levels of $L(\mathbb{R})$, and we do not need Θ -g-organization by Remark 2.5. To keep the notations uniform in this section, we will use the notation $Lp^{\text{c}\Omega}(\mathbb{R}, \text{Code}(\Omega))$ to denote $Lp(\mathbb{R})$ in the case $(\mathcal{P}, \Sigma) = (\emptyset, \emptyset)$.)

Ω is suitable and $\mathcal{M}_1^{\Omega, \sharp}$ generically interprets Ω .¹⁹ Let Λ be the unique $(\omega_1 + 1)$ - Ω -iteration strategy for $\mathcal{M}_1^{\Omega, \sharp}$. It can be shown to follow from the hypotheses of Theorems 1.6 and 1.7 (in particular using the fact that every uncountable regular cardinal $\leq \Theta$ is threadable) that the iteration strategy Λ can be extended to a unique $(\Theta + 1)$ -iteration strategy with branch condensation, which we will also call Λ . (This “strategy extension” step is not necessary for the case $(\mathcal{P}, \Sigma) = (\emptyset, \emptyset)$, so we postpone its proof until Section 5.)

As in [10], we use Λ to define $Lp^{\text{c}\Omega}(\mathbb{R}, \text{Code}(\Omega))$. The only thing to check is that $(\Theta + 1)$ -iterability is sufficient to run the definition of $Lp^{\text{c}\Omega}(\mathbb{R}, \text{Code}(\Omega))$ in [10]. Suppose by induction, we have defined a level $\mathcal{M} \triangleleft Lp^{\text{c}\Omega}(\mathbb{R}, \text{Code}(\Omega))$ (in general, the following argument

¹⁹By results of [10], $\mathcal{M}_1^{\Omega, \sharp}$ generically interprets Ω for (\mathcal{P}, Σ) being a \mathcal{G} - Ω -suitable pair or a hod pair where Σ has branch condensation and is Ω -fullness preserving.

works for any transitive structure M containing \mathbb{R} such that there is a surjection from \mathbb{R} onto M) and without loss of generality, we assume \mathcal{M} is a tree activation level $\mathcal{N}_{\alpha+1}$ and we are trying to define the level $\mathcal{M}_{\alpha+1}$ (in the notation of [10, Definition 3.38]); this just means that $\mathcal{M}_{\alpha+1}$ is the first level above \mathcal{M} by which we have fed in all necessary branch information about $\mathcal{T}_{\mathcal{M}}$. It comes down to defining $\mathcal{T}_{\mathcal{M}}$ as in Definition 2.2. Working in the model $N = L(\mathcal{M}, \mathbb{R}, f)[\Sigma]$,²⁰ where f is a surjection from \mathbb{R} onto \mathcal{M} , we need to see that the genericity iteration that defines $\mathcal{T}_{\mathcal{M}}$ terminates in less than Θ many steps. Suppose not, letting $\mathcal{T} \in N$ be the corresponding tree of length $\Theta + 1$. In N , letting γ be a large regular cardinal $> \Theta$, we can construct some $X \prec L_{\gamma}(\mathcal{M}, \mathbb{R}, f)[\Sigma]$ that contains all relevant objects (in particular, $\mathbb{R} \cup \mathcal{M} \cup \{\mathcal{M}\} \subset X$) and such that there is a surjection from \mathbb{R} onto X . Let $\pi : M_X \rightarrow X$ be the uncollapse map and let $\xi = \text{crit}(\pi)$; then $\xi < \Theta$ and $\pi(\xi) \leq \Theta$. We note that π can be canonically extended to a map $\pi^+ : M_X[G] \rightarrow L_{\gamma}(\mathcal{M}, \mathbb{R}, f)[\Sigma][G]$, where $G \subseteq \text{Col}(\omega, \mathbb{R})$ is $L(\mathcal{M}, \mathbb{R}, f)[\Sigma]$ -generic. We also note that since $\mathcal{M} \cup \{\mathcal{M}\} \subset X$, $\xi > o(\mathcal{M})$. We can then use standard arguments (cf. [16, Theorem 3.11]), where $X[G]$ plays the role of the countable hull X there, to conclude that $\text{lh}(\mathcal{T}) < \Theta$. Contradiction. So $\mathcal{T}_{\mathcal{M}}$ is defined and has length $< \Theta$.

To prove (4.1), we need the following definition.

Definition 4.2. *We define $\text{sLp}^{\mathcal{G}\Omega}(\mathbb{R}, \text{Code}(\Omega))$ to be the union of those $\mathcal{M} \triangleleft \text{Lp}^{\mathcal{G}\Omega}(\mathbb{R}, \text{Code}(\Omega))$ such that whenever $\pi : \mathcal{M}^* \rightarrow \mathcal{M}$ is elementary, $\mathcal{P} \in \pi^{-1}(\text{HC})$, and \mathcal{M}^* is countable and transitive, then \mathcal{M}^* is $\mathcal{G}\Omega$ - $(\omega_1 + 1)$ -iterable with unique strategy Λ such that $\Lambda \upharpoonright \text{HC} \in \mathcal{M}$.*

We note that $\text{sLp}^{\mathcal{G}\Omega}(\mathbb{R}, \text{Code}(\Omega))$ is an initial segment of $\text{Lp}^{\mathcal{G}\Omega}(\mathbb{R}, \text{Code}(\Omega))$ ²¹ and it is trivially constructibly closed. Also, $\text{sLp}^{\mathcal{G}\Omega}(\mathbb{R}, \text{Code}(\Omega)) \models \Theta = \theta_{\Sigma}$ and the extender sequence of $\text{sLp}^{\mathcal{G}\Omega}(\mathbb{R}, \text{Code}(\Omega))$ is definable over $\text{sLp}^{\mathcal{G}\Omega}(\mathbb{R}, \text{Code}(\Omega))$ from Ω , which in turn is definable from Σ . In this section, we outline the core model induction up to the “last gap” of $\text{sLp}^{\mathcal{G}\Omega}(\mathbb{R}, \text{Code}(\Omega))$. This will show that

$$\text{sLp}^{\mathcal{G}\Omega}(\mathbb{R}, \text{Code}(\Omega)) \models \text{AD}^+ + \text{MC}(\Sigma). \quad (4.2)$$

From [13, Theorem 17.1] and [4], we know that if $M \models V = L(\wp(\mathbb{R}))+\text{AD}^++\text{MC}(\Sigma)+\Theta = \theta_{\Sigma}$, then $M \models V = L(\text{sLp}^{\mathcal{G}\Omega}(\mathbb{R}, \text{Code}(\Omega)))$. This and equation 4.2 imply equation 4.1. It then suffices to prove equation 4.2.²³ The rest of the section is devoted to this task.

²⁰By “ Σ ”, we mean the set $\{(\mathcal{T}, \beta) : \beta \in \Sigma(\mathcal{T})\}$.

²¹The initial segment may be strict.

²²Ordinal definability from Σ in the definition of $\text{MC}(\Sigma)$ is in the language of set theory, not in the language of $\text{sLp}^{\mathcal{G}\Omega}(\mathbb{R}, \text{Code}(\Omega))$, but by the paragraph above 4.2, this will not make a difference.

²³Note that the statement “ $\mathcal{N} \triangleleft \text{sLp}^{\mathcal{G}\Omega}(\mathbb{R}, \text{Code}(\Omega))$ ” is absolute between models containing \mathbb{R}, \mathcal{N} and closed under Ω .

The following definitions are obvious generalizations of those defined in [7].

Definition 4.3. *We say that the coarse mouse witness condition $W_{\gamma}^{*,\text{g}\Omega}$ holds if, whenever $U \subseteq \mathbb{R}$ and both U and its complement have scales in $\text{Lp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))|\gamma$, then for all $k < \omega$ and $x \in \mathbb{R}$ there is a coarse (k, U) -Woodin-mouse M containing x , closed under the strategy Λ of $\mathcal{M}_1^{\Omega,\sharp}$ with an $(\omega_1 + 1)$ -iteration strategy whose restriction to H_{ω_1} is in $\text{Lp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))|\gamma$.²⁴*

Remark 4.4. *By the proof of [7, Lemma 3.3.5], $W_{\gamma}^{*,\text{g}\Omega}$ implies $\text{Lp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))|\gamma \models \text{AD}^+$.*

Definition 4.5. *An ordinal γ is a critical ordinal in $\text{Lp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))$ if there is some $U \subseteq \mathbb{R}$ such that U and $\mathbb{R} \setminus U$ have scales in $\text{Lp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))|(\gamma + 1)$ but not in $\text{Lp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))|\gamma$. In other words, γ is critical in $\text{Lp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))$ just in case $W_{\gamma+1}^{*,\text{g}\Omega}$ does not follow trivially from $W_{\gamma}^{*,\text{g}\Omega}$.*

To any Σ_1 formula $\theta(v)$ in the language of $\text{Lp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))$ we associate formulae $\theta_k(v)$ for $k \in \omega$, such that θ_k is Σ_k , and for any γ and any real x ,

$$\text{Lp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))|(\gamma + 1) \models \theta[x] \iff \exists k < \omega \text{ Lp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))|\gamma \models \theta_k[x].$$

Definition 4.6. *Suppose $\theta(v)$ is a Σ_1 formula (in the language of set theory expanded by a name for \mathbb{R} and a predicate for $\text{g}\Omega$), and z is a real; then a $\langle \theta, z \rangle$ -prewitness is an ω -sound g -organized Ω -premouse N over z in which there are $\delta_0 < \dots < \delta_9$, S , and T such that N satisfies the formulae expressing*

- (a) ZFC ,
- (b) $\delta_0, \dots, \delta_9$ are Woodin,
- (c) S and T are trees on some $\omega \times \eta$ which are absolutely complementing in $V^{\text{Col}(\omega, \delta_9)}$, and
- (d) For some $k < \omega$, $p[T]$ is the Σ_{k+3} -theory (in the language with names for each real and predicate for $\text{g}\Omega$) of $\text{Lp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))|\gamma$, where γ is least such that $\text{Lp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))|\gamma \models \theta_k[z]$.

If N is also $(\omega, \omega_1, \omega_1 + 1)$ -iterable (as a g -organized Ω -mouse), then we call it a $\langle \theta, z \rangle$ -witness.

Definition 4.7. *We say that the fine mouse witness condition $W_{\gamma}^{\text{g}\Omega}$ holds if whenever $\theta(v)$ is a Σ_1 formula (in the language \mathcal{L}^+ of g -organized Ω -premice (cf. [10])), z is a real, and $\text{Lp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))|\gamma \models \theta[z]$, then there is a $\langle \theta, z \rangle$ -witness \mathcal{N} whose ${}^g\Omega$ -iteration strategy, when restricted to countable trees on \mathcal{N} , is in $\text{Lp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega))|\gamma$.*

²⁴We demand the strategy has the property that iterates of M according to the strategy are closed under Λ .

Lemma 4.8. $W_\gamma^{*,g\Omega} \rightarrow W_\gamma^{g\Omega}$ for limit γ .

The proof of the above lemma is a straightforward adaptation of that of [7, Lemma 3.5.4]. One main point is the use of the g -organization: g -organized Ω -mice behave well with respect to generic extensions in the sense that if \mathcal{P} is a g -organized Ω -mouse and h is set generic over \mathcal{P} then $\mathcal{P}[h]$ can be rearranged to a g -organized Ω -mouse over h .

The induction is guided by the pattern of scales in $\text{Lp}^{g\Omega}(\mathbb{R}, \text{Code}(\Omega))$ as analyzed in [10]. To show $\text{AD}^+ + \text{MC}(\Sigma)$ holds in $\text{sLp}^{g\Omega}(\mathbb{R}, \text{Code}(\Omega))$, we show $\text{sLp}^{g\Omega}(\mathbb{R}, \text{Code}(\Omega)) \vDash \forall \alpha (\alpha \text{ is critical} \rightarrow W_\alpha^{*,g\Omega})$. Our plan is to show $W_{\alpha+1}^{*,g\Omega}$ assuming $W_\alpha^{*,g\Omega}$ for α critical. Lemma 3.1 and the subsequent corollary provide the base case for our induction. For $\alpha > 0$, we have three cases:

1. α is a successor of a critical ordinal, or α is a limit of critical ordinals and $\text{cf}(\alpha) = \omega$.
2. α is an inadmissible limit of critical ordinals and $\text{cf}(\alpha) > \omega$.
3. α ends a weak gap or is the successor of an ordinal that ends a strong gap. Say the gap is $[\gamma, \alpha^*]$, where $\alpha^* = \alpha$ if the gap is weak and $\alpha^* + 1 = \alpha$ if the gap is strong. Furthermore, $\text{sLp}^{g\Omega}(\mathbb{R}, \text{Code}(\Omega)) \upharpoonright \alpha \vDash \text{MC}(\Sigma) + \text{AD}^+ + \Theta = \theta_\Sigma$.

We deal with the easy case (1) first. In this case, let $\Gamma = \Sigma_1^{\text{sLp}^{g\Omega}(\mathbb{R}, \text{Code}(\Omega)) \upharpoonright \alpha}$. Then $C_\Gamma = \bigcup_{n < \omega} C_{\Gamma_n}$ for some increasing sequence of scaled pointclasses $\langle \Gamma_n \mid n < \omega \rangle$. By $W_\alpha^{*,g\Omega}$, for each n , we have Σ -cmi operators $\langle K_m \mid m < \omega \rangle$ that collectively witness $\text{Det}(\bigcup_n \Gamma_n)$. Say each K_m is defined on a cone above some fixed $a \in \text{HC}$. The desired mouse operator K_0 is defined as follows: For each transitive and self-wellordered $A \in \text{HC}$ coding a , $J_0(A)$ is the shortest initial segment $\mathcal{M} \triangleleft \text{Lp}^{g\Omega}(A)$ such that $\mathcal{M} \vDash \text{ZFC}^-$ and \mathcal{M} is closed under K_m for all m . J_0 is total and trivially relativizes well and determines itself on generic extensions because the K_m 's have these properties. We then use Lemma 3.1 to get that $J_1 = \mathcal{M}_1^{\sharp, J_0}$ is defined on the cone above a by arguments in the previous section. Inductively, we get that $J_{n+1} = \mathcal{M}_1^{\sharp, J_n}$ is defined on the cone above a for all n and one easily gets that these operators are Σ -cmi operators. By Lemma 4.1.3 of [7], this implies $W_{\alpha+1}^{*,g\Omega}$.

Now we're on to the case where α is inadmissible and $\text{cf}(\alpha) > \omega$. Let $\phi(v_0, v_1)$ be a Σ_1 formula and $x \in \mathbb{R}$ be such that

$$\forall y \in \mathbb{R} \exists \beta < \alpha \text{sLp}^{g\Omega}(\mathbb{R}, \text{Code}(\Omega)) \upharpoonright \beta \vDash \phi[x, y],$$

and letting $\beta(x, y)$ be the least such β ,

$$\alpha = \sup \{ \beta(x, y) \mid y \in \mathbb{R} \}.$$

We first define J_0 on transitive and self-wellordered $A \in \text{HC}$ coding x . For $n < \omega$, let

$$\phi_n^* \equiv \exists \gamma (\text{Lp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega)) \mid \gamma \models \forall i \in \omega (i > 0 \Rightarrow \phi((v)_0, (v)_i) \wedge (\gamma + \omega n) \text{ exists})).$$

For such an A as above, let \mathcal{M} be an A -premouse and let G be a $\text{Col}(\omega, A)$ -generic filter over \mathcal{M} . Then $\mathcal{M}[G]$ can be regarded as a g -organized Ω -mouse over $z(G, A)$ where $z(G, A)$ is a real coding G, A and is obtained from G, A in some simple fashion.²⁵ Also, let σ_A be a term defined uniformly (in \mathcal{M}) from A, x such that

$$(\sigma_A^G)_0 = x$$

and

$$\{(\sigma_A^G)_i \mid i > 0\} = \{\rho^G \mid \rho \in L_1(A) \wedge \rho^G \in \mathbb{R}\}.$$

Let φ be a sentence in the language of A -premice such that for any A -premouse \mathcal{M} , $\mathcal{M} \models \varphi$ iff whenever G is \mathcal{M} -generic for $\text{Col}(\omega, A)$, then for any n there is a $\gamma < o(\mathcal{M})$ such that

$$\mathcal{M}[z(G, A)] \mid \gamma \text{ is a } \langle \phi_n^*, \sigma_A^G \rangle\text{-prewitness.}$$

Then $J_0(A)$ is the shortest initial segment of $\text{Lp}^{\text{g}\Omega}(A)$ which satisfies φ , if it exists, and is undefined otherwise. Using the fact that $W_\alpha^{\text{g}\Omega}$ holds, we get that $J_0(A)$ exists for all $A \in \text{HC}$ coding x because α has uncountable cofinality and there are only countably many $\langle \phi_n^*, \rho_A^G \rangle$. Also we can then define J_n as before. It's easy to show again that the J_n 's relativize well and determine themselves on generic extensions, so they are Σ -cmi operators. This implies $W_{\alpha+1}^{*, \text{g}\Omega}$.

Lastly, we consider the gap case. Using the notations as in case 3 above, let $\Gamma = \Sigma_1^{\text{sLp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega)) \mid \gamma}$. If $[\gamma, \alpha^*]$ is a weak gap, then by the scales analysis at the end of a weak gap from [14] and [10], we can construct a self-justifying system (sjs) \mathcal{A} Wadge-cofinal in $\wp(\mathbb{R}) \cap \text{sLp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega)) \mid \alpha^*$.²⁶ If $[\gamma, \alpha^*]$ is a strong gap, then by the Kechris–Woodin theorem, AD^+ holds in $\text{sLp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega)) \mid \alpha$, and again by results of [14], [10], and [21], we also get a self-justifying system \mathcal{A} Wadge-cofinal in $\text{sLp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega)) \mid \alpha \cap \wp(\mathbb{R})$. From \mathcal{A} and arguments in [7, Section 5], there is a pair (\mathcal{Q}, Λ) such that \mathcal{Q} is Γ -suitable and Λ is the (ω_1, ω_1) -strategy for \mathcal{Q} guided by \mathcal{A} (see the next section for more details on self-justifying systems). Let $J_0 = \Lambda$. We assume that \mathcal{A} contains the universal Γ -set and hence the universal $\check{\Gamma}$ -set.

Claim 4.9. J_0 determines itself on generic extensions.

²⁵This is one of the main reasons that we consider $\text{g}\Omega$ -mice; this is so that generic extensions of $\text{g}\Omega$ -mice can be rearranged to $\text{g}\Omega$ -mice.

²⁶This means \mathcal{A} is a countable collection containing a universal $\Sigma_1^{\text{sLp}^{\text{g}\Omega}(\mathbb{R}, \text{Code}(\Omega)) \mid \gamma}$ set, closed under complements and whenever $A \in \mathcal{A}$, then there is a scale whose individual norms are coded by sets in \mathcal{A} .

Proof. Let N be a countable transitive structure of ZFC^- such that N is closed under J_0 (and hence under Λ). We simply describe a procedure that determines Λ on generic extensions of N ; the reader may gladly verify that this is enough to prove the claim. Let $g \in V$ be generic over N ; we assume without loss of generality that $g \subseteq \text{Col}(\omega, \kappa)$ for some N -cardinal κ and N has a $\text{Col}(\omega, \kappa)$ -name $\dot{\mathcal{A}}$ for \mathcal{A} . The existence of $\dot{\mathcal{A}}$ follows from [7, Lemma 3.7.5] and the fact that $C_\Gamma(H_{\kappa^+}^N) \subset N$. The latter follows from a simple genericity iteration argument using the fact that N is closed under Λ . Let $i : \mathcal{Q} \rightarrow \mathcal{R}$ be a $H_{\kappa^+}^N$ -genericity iteration in N according to Λ . Since R is Γ -suitable, $C_\Gamma(H_{\kappa^+}^N) \subset \mathcal{R}[H_{\kappa^+}^N]$. Since $\mathcal{R}[H_{\kappa^+}^N] \subset N$, $C_\Gamma(H_{\kappa^+}^N) \subset N$ as claimed.

Let \mathcal{T} be a tree according to Λ of limit length in $N[g]$ (the argument for stacks is similar). If \mathcal{T} is short, we can find the \mathcal{Q} -structure $\mathcal{Q}(\mathcal{T})$ for \mathcal{T} and this in turn determines the branch $b = \Lambda(\mathcal{T}) \in N[g]$. The \mathcal{Q} -structure $\mathcal{Q}(\mathcal{M}(\mathcal{T}))$ belongs to $C_\Gamma(\mathcal{M}(\mathcal{T}))$ and can be computed using $\dot{\mathcal{A}}_g$; the point is the universal $\check{\Gamma}$ -set belongs to \mathcal{A} , so $N[g]$ can use $\dot{\mathcal{A}}_g$ to compute the C_Γ -operator correctly.

Suppose \mathcal{T} is maximal. By boolean comparison (cf. [7, Section 5.4]), we can find a tree $\mathcal{U} \in N$ according to Λ such that

- (i) \mathcal{U} is non-dropping with last model $\mathcal{M}^\mathcal{U}$ and branch embedding $\pi^\mathcal{U}$;
- (ii) $\Lambda(\mathcal{T}) = b$ is the unique branch in $N[g]$ with last model $\mathcal{M}^\mathcal{T}$ and branch embedding $\pi^\mathcal{T}$ such that there is an elementary embedding $\tau : \mathcal{M}^\mathcal{T} \rightarrow \mathcal{M}^\mathcal{U}$ with $\pi^\mathcal{U} = \tau \circ \pi^\mathcal{T}$.²⁷

□

Furthermore, J_0 is suitable (we can construct $\mathcal{M}_1^{J_0, \sharp}$ by arguments in the previous section) and $\mathcal{M}_1^{J_0, \sharp}$ generically interprets J_0 by [10, Lemma 4.8]. Note that J_0 and \mathcal{A} are projectively equivalent in any reasonable coding. We can use Lemma 3.1 to show $W_{\alpha+1}^{*, g_\Omega}$ by constructing a sequence of operators $(J_n : n < \omega)$, where $J_{n+1} = \mathcal{M}_1^{J_n, \sharp}$ for all n .²⁸ This concludes the outline of the proof of 4.2 and 4.1.

It now follows easily that we can strengthen the conclusion of Projective Determinacy in Corollary 3.4 to obtain the following result.

Corollary 4.10. *Assume $ZF + DC_{\wp(\omega_1)} + \text{``}\omega_1 \text{ is } \mathbb{R}\text{-strongly compact and } \neg \square_{\omega_1}\text{.''}$ Then AD holds in $L(\mathbb{R})$.*

This corollary completes the proof of Theorem 1.5. It also forms a significant first step in the proofs of Theorems 1.6 and 1.7.

²⁷The map τ is a fine-structural embedding. Typically, it is a k -embedding in the sense of [16] where k is the degree of the tree \mathcal{T} .

²⁸These operators, again, can be shown to be Σ -cmi operators. Here and elsewhere, we suppress the formula φ_{\min} defined in [10, Definition 3.2] from the definition of $J_1 = \mathcal{M}_1^{J_0, \sharp}$; to be entirely correct, according to [10], J_1 should be $\mathcal{M}_1^{(J_0, \varphi_{\min}), \sharp}$.

5 A model of $\text{AD}^+ + \Theta > \theta_\Sigma$

Suppose (\mathcal{P}, Σ) is a \mathcal{G} - Ω^* -suitable pair for some nice operator \mathcal{G} such that Σ has branch condensation and is Ω^* -fullness preserving. As a special case we also allow $(\mathcal{P}, \Sigma) = (\emptyset, \emptyset)$. In the previous section we showed (under our strong hypotheses plus a smallness assumption) that there is a maximal model of $\text{AD}^+ + V = L(\wp(\mathbb{R})) + \Theta = \theta_\Sigma$ containing all reals and ordinals. This model has the form $L(\Omega_\Sigma, \mathbb{R})$ where $L(\Omega_\Sigma, \mathbb{R}) \cap \wp(\mathbb{R}) = \Omega_\Sigma$. In this section, we will go just beyond this model to obtain a model of $\text{AD}^+ + \Theta > \theta_\Sigma$ containing all reals and ordinals.

Define the pointclass

$$\Gamma = (\Sigma_1^2(\text{Code}(\Sigma)))^{\Omega^*}.$$

Note that we have $\Gamma = (\Sigma_1^2(\text{Code}(\Sigma)))^{\Omega_\Sigma}$; this is because if a set of reals $A \in \Omega^*$ witnesses a $\Sigma_1^2(\text{Code}(\Sigma))$ fact about a real x , then there is a set of reals in $\Delta_1^2(\text{Code}(\Sigma), x)^{L(A, \mathbb{R})}$ witnessing the same fact about x by Woodin's Δ_1^2 basis theorem relativized to x and $\text{Code}(\Sigma)$ and applied in the model $L(A, \mathbb{R})$, and such a set of reals can be shown to be in Ω_Σ .

Recall from Section 4 that (under our smallness assumption) the maximal model $L(\Omega_\Sigma, \mathbb{R})$ of $\text{AD}^+ + \Theta = \theta_\Sigma$ is, up to its Θ , a hybrid mouse over \mathbb{R} of the form $\text{sLp}^{\mathcal{G}_\Omega}(\mathbb{R}, \text{Code}(\Omega))$ where we have defined the operator $\Omega = \Sigma$. We remind the reader that $\text{Code}(\Omega)$ is self-scaled.

In particular we have

$$\Omega_\Sigma = \wp(\mathbb{R}) \cap \text{sLp}^{\mathcal{G}_\Omega}(\mathbb{R}, \text{Code}(\Omega)),$$

so we can reformulate our pointclass as

$$\Gamma = (\Sigma_1^2)^{\text{sLp}^{\mathcal{G}_\Omega}(\mathbb{R}, \text{Code}(\Omega))} = (\Sigma_1^2)^{\text{sLp}^{\mathcal{G}_\Omega}(\mathbb{R}, \text{Code}(\Omega))|_\alpha}$$

where $\alpha = (\delta_1^2)^{\text{sLp}^{\mathcal{G}_\Omega}(\mathbb{R}, \text{Code}(\Omega))}$ is the ordinal beginning the last gap of $\text{sLp}^{\mathcal{G}_\Omega}(\mathbb{R}, \text{Code}(\Omega))$. (Recall that by Σ_1^2 we mean to include Ω , or equivalently Σ , as a parameter. By self-iterability it makes no difference whether we also include the extender sequence as a parameter.)

Like the pointclass considered in the “gap in scales” case of the core model induction in Section 4, the pointclass Γ is an inductive-like pointclass with the scale property. Our next task is to find the next scaled pointclass, or (what is roughly equivalent) to build a scale on a complete $\check{\Gamma}$ set. Unlike in Section 4, this next scaled pointclass cannot be found within $\text{sLp}^{\mathcal{G}_\Omega}(\mathbb{R}, \text{Code}(\Omega))$. The reason is that the complete $\check{\Gamma}$ set $\{(x, y) \in \mathbb{R} \times \mathbb{R} : y \notin \text{OD}_x^{\text{sLp}^{\mathcal{G}_\Omega}(\mathbb{R}, \text{Code}(\Omega))}\}$ cannot have any uniformization in $\text{sLp}^{\mathcal{G}_\Omega}(\mathbb{R}, \text{Code}(\Omega))$, and therefore cannot have any scale in $\text{sLp}^{\mathcal{G}_\Omega}(\mathbb{R}, \text{Code}(\Omega))$, by a standard argument.

We will use our strong hypotheses (as in Theorems 1.6 and 1.7) to build a scale on a complete $\check{\Gamma}$ set. Each prewellordering of this scale will be in $L(\Omega_\Sigma, \mathbb{R})$, or equivalently in $sLp^{\mathbb{C}_\Omega}(\mathbb{R}, \text{Code}(\Omega))$, although the sequence of prewellorderings cannot be, as we just saw.

More directly, what we will show is that the prewellorderings are in a pointclass $\text{Env}(\Gamma)$, the envelope of Γ . The notion of envelope was used by Martin to identify the next scaled pointclass after an inductive-like scaled pointclass in the **AD** context; see Jackson [1]. We will need its adaptation to the partial determinacy context as defined in the second author's thesis [20] (see also the subsequent article [21].)

It turns out that $\text{Env}(\Gamma) \subset L(\Omega_\Sigma, \mathbb{R})$, and in fact $\text{Env}(\Gamma)$ consists exactly of the sets of reals that are ordinal-definable from Σ in the model $L(\Omega_\Sigma, \mathbb{R})$, but we will not be able to see this until later. For now we must use the following "local" definition of the envelope in terms of the ambiguous pointclass $\Delta_\Gamma = \Gamma \cap \check{\Gamma}$ and in terms of the notion of " Δ_Γ in an ordinal parameter." This notion can be defined in general, but here we can take the following characterization as a definition: a set of reals is Δ_Γ in an ordinal parameter if and only if it is Δ_1 -definable over $sLp^{\mathbb{C}_\Omega}(\mathbb{R}, \text{Code}(\Omega)) \mid \alpha$ from ordinals (and Ω , or equivalently Σ .)

Definition 5.1. *The envelope of Γ , denoted by $\text{Env}(\Gamma)$, is the pointclass consisting of all pointsets A such that, for every countable $\sigma \subset \mathbb{R}$, there is a pointset A' that is Δ_Γ in an ordinal parameter and satisfies $A \cap \sigma = A' \cap \sigma$.*

The boldface pointclass $\text{Env}(\Gamma)$ is defined similarly but allowing a real parameter. That is, $A \in \text{Env}(\Gamma)$ if there is a real x such that for every countable $\sigma \subset \mathbb{R}$ there is a pointset A' that is $\Delta_\Gamma(x)$ in an ordinal parameter and satisfies $A \cap \sigma = A' \cap \sigma$.

The following fact about envelopes is crucial for our argument. It is essentially proved in the thesis [20] (which deals with generic large cardinal properties of ω_1 in **ZFC** rather than with large cardinal properties of ω_1 in **ZF** + **DC**, but the argument carries over to the present context.) An easier version with "scale" replaced by "semiscale" is proved in the article [21], and a special case of the scale construction appears in another article [22].

Lemma 5.2 (Wilson). *Assume **ZF** + **DC**. Let Γ be an inductive-like pointclass with the scale property. Suppose that ω_1 is $\text{Env}(\Gamma)$ -strongly compact. Then there is a scale on a universal $\check{\Gamma}$ set, each of whose prewellorderings is in $\text{Env}(\Gamma)$.*

We will also need the fact that if **ZF** + **DC** $_{\mathbb{R}}$ holds and the boldface ambiguous part Δ_Γ of the pointclass Γ is determined, as it is here, then $\text{Env}(\Gamma)$ is determined and projectively closed (Wilson [20, 21]; based on work of Kechris, Woodin, and Martin.) Therefore Wadge's lemma applies to it, as one can easily verify that the relevant games are determined. Moreover, the Wadge preordering of $\text{Env}(\Gamma)$ ²⁹ is a prewellordering: otherwise by **DC** $_{\mathbb{R}}$ we could

²⁹Really a preordering of pairs $\{B, \neg B\}$ where $B \in \text{Env}(\Gamma)$.

choose a sequence of pointsets in $\mathbf{Env}(\Gamma)$ that was strictly decreasing in the Wadge ordering, but then by the proof of the Martin–Monk theorem we get a contradiction. (Again one can easily verify that the relevant games are determined.)

Note that the prewellorderings of a scale as in Lemma 5.2 must be Wadge-cofinal in $\mathbf{Env}(\Gamma)$; otherwise the sequence of prewellorderings itself would be coded by a set of reals in $\mathbf{Env}(\Gamma)$, which is impossible as mentioned above. From such a scale, it then follows by a general argument (see Jackson [1] and the straightforward adaptation [20, Section 4.3] to the partial determinacy context) that we can obtain a self-justifying system contained in $\mathbf{Env}(\Gamma)$:³⁰

Lemma 5.3. *Assume $\mathbf{ZF} + \mathbf{DC}$. Let Γ be an inductive-like pointclass with the scale property such that Δ_Γ is determined. Suppose that ω_1 is $\mathbf{Env}(\Gamma)$ -strongly compact. Then there is a self-justifying system $\mathcal{A} \subset \mathbf{Env}(\Gamma)$ containing a universal Γ set.*

We will use this lemma together with the hypotheses of Theorems 1.6 or 1.7 to obtain a self-justifying system $\mathcal{A} \subset \mathbf{Env}(\Gamma)$ containing a universal Γ set. We begin with the observation that the length of the Wadge prewellordering of $\mathbf{Env}(\Gamma)$ is at most Θ by the usual argument: the initial segment corresponding to a set $B \in \mathbf{Env}(\Gamma)$ is the image of \mathbb{R} under the function $y \mapsto g_y^{-1}[B]$, where g_y denotes the continuous function coded by the real y . Moreover, the lightface envelope $\mathbf{Env}(\Gamma)$ admits a wellordering (essentially an ultrapower of the canonical wellordering of the Δ_Γ -in-an-ordinal sets by Martin’s cone measure, which measures the relevant sets by $\mathbf{Env}(\Gamma)$ -determinacy.)

Lemma 5.4. *Let Γ be an inductive-like pointclass with the scale property such that Δ_Γ is determined. Assume $\mathbf{ZF} + \mathbf{DC} + \omega_1$ is Θ -strongly compact.” Then there is a self-justifying system $\mathcal{A} \subset \mathbf{Env}(\Gamma)$ containing a universal Γ set.*

Proof. Consider the restriction of the Wadge prewellordering of $\mathbf{Env}(\Gamma)$ to the lightface envelope $\mathbf{Env}(\Gamma)$. We can refine this prewellordering to a wellordering by taking its lexicographical product with a wellordering of $\mathbf{Env}(\Gamma)$, which exists, as mentioned above. This refinement has the property that its length is at most Θ , because its initial segment below any set $A \in \mathbf{Env}(\Gamma)$ is contained in the Wadge-initial segment $\{B \in \mathbf{Env}(\Gamma) : B \leq_w A\}$. (It’s not clear whether the original wellordering of $\mathbf{Env}(\Gamma)$ described above has this property.) Therefore our hypothesis implies that ω_1 is $\mathbf{Env}(\Gamma)$ -strongly compact, and the desired conclusion follows by Lemma 5.3. \square

³⁰We don’t know if it is possible to obtain a self-justifying system contained in the lightface envelope, but this will not matter for our application.

Lemma 5.5. *Let Γ be an inductive-like pointclass with the scale property such that Δ_Γ is determined. Assume $\text{ZF} + \text{DC}_\mathbb{R} + \text{``}\omega_1 \text{ is } \mathbb{R}\text{-strongly compact and } \Theta \text{ is singular.}''$ Then there is a self-justifying system $\mathcal{A} \subset \mathbf{Env}(\Gamma)$ containing a universal Γ set.*

Proof. Let $<$ be a wellordering of $\text{Env}(\Gamma)$ that refines the Wadge prewellordering (as in the previous proof) and therefore has length at most Θ . Using the hypothesis that Θ is singular to deal with the apparent possibility that $<$ has length equal to Θ , we can obtain a function $f : \mathbb{R} \rightarrow \text{Env}(\Gamma)$ that is cofinal with respect to $<$ and therefore also cofinal with respect to the Wadge prewellordering of $\text{Env}(\Gamma)$. Then we can define a partial surjection from $\mathbb{R} \times \mathbb{R}$ onto $\text{Env}(\Gamma)$ by mapping $(x, y) \in \mathbb{R} \times \mathbb{R}$ to the preimage of the set $f(x)$ under the continuous function coded by the real y , whenever this preimage happens to be in $\text{Env}(\Gamma)$.

Therefore there is a surjection from \mathbb{R} onto $\text{Env}(\Gamma)$, and by our hypothesis that ω_1 is \mathbb{R} -strongly compact, it follows that ω_1 is $\text{Env}(\Gamma)$ -strongly compact. We could now apply Lemma 5.3 to obtain the desired conclusion, except for the problem that we only have $\text{DC}_\mathbb{R}$ in place of DC . This problem can be solved by passing to an inner model.

Take a fine, countably complete measure μ on $\wp_{\omega_1}(\text{Env}(\Gamma))$ and consider the model $L(X)[\mu]$ where $X = \text{Env}(\Gamma)^\omega \cup \mathbb{R}$. In V we have $\text{DC}_\mathbb{R}$ and we have a surjection from \mathbb{R} to X , so we have DC_X . Because an ω -sequence of elements of X can be coded by a single element of X , we have DC_X in $L(X)[\mu]$ as well. In $L(X)[\mu]$ every set is a surjective image of $X \times \xi$ for some ordinal ξ , so DC follows from DC_X by a standard argument. Then we can apply Lemma 5.3 in $L(X)[\mu]$ and note that the conclusion is upward absolute to V . \square

Now that we have obtained a self-justifying system $\mathcal{A} = (A_i : i < \omega)$ sealing the envelope of Γ , we may proceed as in the “gap in scales” case of Section 4 to get a pair (\mathcal{Q}, Λ) such that \mathcal{Q} is a Γ -suitable g-organized Ω -premouse and Λ is the (ω_1, ω_1) -iteration strategy for \mathcal{Q} guided by \mathcal{A} . A slight difference from Section 4 is caused by the fact that, at this stage in the argument, we do not know how to rule out the possibility that the pointclass $\mathbf{Env}(\Gamma)$ is strictly larger than the pointclass $\Omega_\Sigma = \wp(\mathbb{R}) \cap \text{sLp}^{\text{c}\Omega}(\mathbb{R}, \text{Code}(\Omega))$.

However, this difference does not create any problem because the important thing is that every set $A \in \mathbf{Env}(\Gamma)$ (and in particular every set A_i in our self-justifying system \mathcal{A}) has the property that, for a cone of $b \in \text{HC}$, the hybrid lower part mouse $\text{Lp}^{\text{g}\Omega, \Gamma}(b)$ has a $\text{Col}(\omega, b)$ -term for a set of reals that locally captures A . (If A is in the lightface envelope then the base of the cone is \emptyset and this holds for all $b \in \text{HC}$.) For a proof, see Wilson [20, Section 4.2]. This local term-capturing property is sufficient to make sense of the notion of A -iterability, to prove the existence of A -iterable g-organized and Θ -g-organized Ω -premice, and to get an iteration strategy Λ guided by the self-justifying system \mathcal{A} . The adaptation of existing proofs to this context is straightforward.

Defining the Σ -CMI operator $\mathcal{F} = \Lambda$, we can then use Lemma 3.1 to construct a sequence of Σ -CMI operators $(J_n : n < \omega)$, where $J_0 = \mathcal{F}$ and $J_{n+1} = \mathcal{M}_1^{\sharp, J_n}$ for all $n > 0$. Because \mathcal{A} and \mathcal{F} are projectively equivalent (in any reasonable coding) this shows the existence of a determined projective-like hierarchy just beyond $\mathbf{Env}(\Gamma)$, and therefore beyond the maximal model of $\mathbf{AD}^+ + \Theta = \theta_\Sigma$.

To continue further and get a model of $\mathbf{ZF} + \mathbf{AD}^+ + \Theta > \theta_\Sigma$, we proceed along the lines of Section 4. The difference is that now the operator \mathcal{F} is here to stay: we must consider \mathcal{F} -hybrid mice from this point on, and never return to considering Ω -hybrid mice because they cannot give us anything new.

Our model of $\mathbf{AD}^+ + \Theta > \theta_\Sigma$ will be obtained as the maximal model of $\mathbf{AD}^+ + \Theta = \theta_\Lambda$ and θ_Σ will be the penultimate member of its Solovay sequence. The existence of this maximal model is established by the results of Section 4 with the suitable pair (\mathcal{Q}, Λ) and its associated operator \mathcal{F} in place of the hod pair (or suitable pair, or empty pair) (\mathcal{P}, Σ) and its associated operator Ω . (For this reason it is important that we allowed suitable pairs as well as hod pairs and empty pairs in Sections 3 and 4.)

To obtain the maximal model of $\mathbf{AD}^+ + \Theta = \theta_\Lambda$, it remains only to show that Λ can be extended to a $(\Theta + 1)$ -iteration strategy with branch condensation. (In fact, we will show that it can be extended to a Θ^+ -iteration strategy with branch condensation.) As remarked in Section 4, this strategy extension is necessary to define the model $\text{sLp}^{\mathcal{C}\mathcal{F}}(\mathbb{R}, \text{Code}(\mathcal{F}))$ via g-organization, which in turn is necessary to analyze the pattern of scales in this model.

Note that because the iteration strategy Λ is guided by a self-justifying system, it has branch condensation and hull condensation and the set of reals coding it is Suslin. Accordingly, we can use the following lemma to extend Λ . Our argument is based on Schindler and Steel [7, Lemmas 2.1.11 and 2.1.12], but some adaptations are necessary in the absence of AC. A similar argument is also found in Steel [12].

Before proving the lemma (which will take the remainder of this section) let us note that the hypothesis that every uncountable regular cardinal $\leq \Theta$ is threadable follows from the hypotheses of Theorems 1.6 and 1.7. In particular, it follows from the hypothesis $\mathbf{ZF} + \mathbf{DC} + \text{``}\omega_1 \text{ is } \Theta\text{-strongly compact''}$ and also from the hypothesis $\mathbf{ZF} + \mathbf{DC}_\mathbb{R} + \text{``}\omega_1 \text{ is } \mathbb{R}\text{-strongly compact and } \Theta \text{ is singular.}''$ Note also that the conclusion that the extension of Λ has hull condensation, together with the fact that the original ω_1 -iteration strategy Λ has branch condensation, implies that the extended strategy also has branch condensation by an easy Skolem hull argument. (We can take the Skolem hull in an inner model of \mathbf{ZFC} , so that no choice is required.)

Lemma 5.6. *Assume that \mathbf{ZF} holds and let Λ be an ω_1 -iteration strategy with hull con-*

densation for a premouse³¹ \mathcal{Q} . Assume that $\text{Code}(\Lambda)$ is Suslin. Let η be an uncountable cardinal and assume that every uncountable regular cardinal $\leq \eta$ is threadable. Then Λ has a (necessarily unique) extension to an η^+ -iteration strategy with hull condensation.

Proof. Let \mathcal{T} be a putative iteration tree on \mathcal{Q} of length less than η^+ and such that every countable hull of \mathcal{T} is by Λ . (A putative iteration tree is like an iteration tree except that its last model, if it has one, is allowed to be illfounded.) What we want to show is that if \mathcal{T} has a last model, then this last model is wellfounded, and if \mathcal{T} has limit length, then it has a unique cofinal wellfounded branch b such that every countable hull $\bar{\mathcal{T}}^{\bar{b}}$ of $\mathcal{T}^{\bar{b}}$ is also by Λ (in which case our extension of Λ can and must choose this branch.)

In the case that \mathcal{T} has a last model, it is easy to see that the last model must be wellfounded; otherwise by taking a Skolem hull (of $L_{\eta^+}[\mathcal{Q}, \mathcal{T}]$, say, so that no choice is required) we may obtain a countable hull of \mathcal{T} whose last model is illfounded, but the last model of the hull must be wellfounded because the hull is by the iteration strategy Λ .

Now suppose that \mathcal{T} has limit length. This case will require a bit more work. First we note that it suffices to find some cofinal branch b of \mathcal{T} such that every countable hull $\bar{\mathcal{T}}^{\bar{b}}$ of $\mathcal{T}^{\bar{b}}$ is by Λ ; then a Skolem hull argument shows that there can be at most one such branch and that any such branch is wellfounded. Let q be a real coding the premouse \mathcal{Q} . We consider two subcases.

1. $\text{lh}(\mathcal{T})$ has uncountable cofinality.

In this subcase, we use the general fact about iteration trees that the sequence of branches $[0, \alpha)_{\mathcal{T}}$ for limit ordinals $\alpha < \text{lh}(\mathcal{T})$ is a coherent sequence of clubs. Here $\text{lh}(\mathcal{T})$ is threadable (equivalently, has threadable cofinality,) so the tree \mathcal{T} has a unique cofinal branch b obtained by threading this coherent sequence. Let $\bar{\mathcal{T}}^{\bar{b}}$ be a countable hull of $\mathcal{T}^{\bar{b}}$. We want to show that $\bar{\mathcal{T}}^{\bar{b}}$ is by Λ .

Let x be a real coding $\bar{\mathcal{T}}^{\bar{b}}$. The model $N = L[q, \mathcal{T}, b, \Lambda, x]$ ³² satisfies **AC** and therefore \square_{ω} , whereas V satisfies “ ω_1 is threadable” and therefore $\neg\square_{\omega}$, so $\omega_1^N < \omega_1$. Note that the model N sees that $\bar{\mathcal{T}}^{\bar{b}}$ is a hull of $\mathcal{T}^{\bar{b}}$ by the absoluteness of wellfoundedness for the tree of attempts to build a map $\text{lh}(\bar{\mathcal{T}}) \rightarrow \text{lh}(\mathcal{T})$ witnessing this (or we could just put such a map into the model.) The model N also sees, of course, that $\text{lh}(\mathcal{T})$ has uncountable cofinality.

Working in N , by a Skolem hull argument we can take a hull $\mathcal{T}^{* \cap b^*}$ of $\mathcal{T}^{\bar{b}}$ such that $\text{lh}(\mathcal{T}^*)$ has cardinality and cofinality ω_1 and $\bar{\mathcal{T}}^{\bar{b}}$ is a hull of $\mathcal{T}^{* \cap b^*}$. Because the tree \mathcal{T}^*

³¹By a premouse here we mean an \mathcal{F} -premouse where \mathcal{F} is an operator that condenses finely (such as the core model induction operators that we consider in this paper.) Alternatively we could use coarse mice here, because we will only need the extended strategy for genericity iterations.

³²We are abusing notation here. For example, instead of Λ itself as a predicate we mean $\{(\mathcal{U}, \xi) : \xi \in \Lambda(\mathcal{U})\}$.

is countable in V the branch $\Lambda(\mathcal{T}^*)$ is defined, and the model N can see it. In N the tree \mathcal{T}^* can have at most one cofinal branch because its length has uncountable cofinality, so $\Lambda(\mathcal{T}^*) = b^*$. Therefore the hull $\mathcal{T}^* \cap b^*$ is by Λ , and by hull condensation *its* hull $\bar{\mathcal{T}} \cap \bar{b}$ is also by Λ , as desired.

2. $\text{lh}(\mathcal{T})$ has countable cofinality.

In this subcase, we define an elementary substructure $X \prec L_{\eta^+}[\mathcal{Q}, \mathcal{T}]$ in V to be *appropriate* if $\mathcal{Q} \cup \{\mathcal{Q}, \mathcal{T}\} \subset X$, X is countable, and $X \cap \text{lh}(\mathcal{T})$ is cofinal in $\text{lh}(\mathcal{T})$. For an appropriate elementary substructure $X \prec L_{\eta^+}[\mathcal{Q}, \mathcal{T}]$, let $\sigma_X : M_X \rightarrow X$ denote the uncollapse map of X , define the tree $\mathcal{T}_X = \sigma_X^{-1}(\mathcal{T})$ on \mathcal{Q} , and note that \mathcal{T}_X is a hull of \mathcal{T} as witnessed by the map $\sigma_X \upharpoonright \text{lh}(\mathcal{T}_X)$.

Furthermore, for any two appropriate elementary substructures $X, Y \prec L_{\eta^+}[\mathcal{Q}, \mathcal{T}]$ such that $X \subset Y$, let $\sigma_{XY} : M_X \rightarrow M_Y$ denote the factor map $\sigma_Y^{-1} \circ \sigma_X$ and note that \mathcal{T}_X is a hull of \mathcal{T}_Y as witnessed by the map $\sigma_{XY} \upharpoonright \text{lh}(\mathcal{T}_X)$.

We say that an elementary substructure $X \prec L_{\eta^+}[\mathcal{Q}, \mathcal{T}]$ is *stable* if it is appropriate and for every appropriate elementary substructure $Y \prec L_{\eta^+}[\mathcal{Q}, \mathcal{T}]$ such that $X \subset Y$ we have

$$\Lambda(\mathcal{T}_X) = \sigma_{XY}^{-1}[\Lambda(\mathcal{T}_Y)].$$

Note that an equivalent condition would be $\sigma_X[\Lambda(\mathcal{T}_X)] \subset \sigma_Y[\Lambda(\mathcal{T}_Y)]$ because distinct cofinal branches are eventually disjoint.

Assume for the moment that there is a stable elementary substructure $X \prec L_{\eta^+}[\mathcal{Q}, \mathcal{T}]$. Then we can define the branch b of \mathcal{T} to be the downward closure of the set $\sigma_X[\Lambda(\mathcal{T}_X)]$ in the \mathcal{T} -ordering. For every appropriate elementary substructure $Y \prec L_{\eta^+}[\mathcal{Q}, \mathcal{T}]$ such that $X \subset Y$, we have $\sigma_Y^{-1}[b] = \Lambda(\mathcal{T}_Y)$. Moreover, the tree $\mathcal{T}_Y \cap \sigma_Y^{-1}[b]$ is a hull of $\mathcal{T} \cap b$.³³ Therefore club many countable hulls of $\mathcal{T} \cap b$ are by Λ and we can argue as in subcase (1) that every countable hull of $\mathcal{T} \cap b$ is by Λ .

So assume toward a contradiction that there is no stable X . Let S be a tree on $\omega \times \text{Ord}$ that projects to $\text{Code}(\Lambda)$, let $f : \omega \rightarrow \text{lh}(\mathcal{T})$ be a cofinal map, and define the model $N' = L[q, \mathcal{T}, S, f]$. (Recall that q is a real coding the premouse \mathcal{Q} .) Note that the model N' satisfies the statement ‘‘there is no stable X ’’ as well as V does: for any appropriate elementary substructure $X \prec L_{\eta^+}[\mathcal{Q}, \mathcal{T}]$ in N' , we may use the absoluteness of wellfoundedness of the tree of attempts to find an appropriate elementary substructure $Y \prec L_{\eta^+}[\mathcal{Q}, \mathcal{T}]$ such that $X \subset Y$ but $\Lambda(\mathcal{T}_X) \neq \sigma_{XY}^{-1}[\Lambda(\mathcal{T}_Y)]$. (We may use the tree S to witness values of Λ .)

³³In general if $\bar{\mathcal{U}}$ is a hull of an iteration tree \mathcal{U} as witnessed by a map $\sigma : \text{lh}(\bar{\mathcal{U}}) \rightarrow \text{lh}(\mathcal{U})$, c is a cofinal branch of \mathcal{U} , and $c \cap \text{range}(\sigma)$ is cofinal in $\text{lh}(\mathcal{U})$, then $\bar{\mathcal{U}} \cap \sigma^{-1}[c]$ is a hull of $\mathcal{U} \cap c$.

Define $\gamma = \omega_1^{N'}$ and note that $\gamma < \omega_1$, just as for the model N in the uncountable cofinality case. In the model N' we can build a continuous, \subset -increasing sequence $(X_\alpha : \alpha \leq \gamma)$ of appropriate elementary substructures of $L_{\eta^+}[\mathcal{Q}, \mathcal{T}]$ such that

$$\Lambda(\mathcal{T}_\alpha) \neq \sigma_{\alpha, \alpha+1}^{-1}[\Lambda(\mathcal{T}_{\alpha+1})]$$

for all $\alpha < \gamma$, where we define $\mathcal{T}_\alpha = \mathcal{T}_{X_\alpha}$, $\sigma_\alpha = \sigma_{X_\alpha}$, etc.

Define the cofinal branch $b = \Lambda(\mathcal{T}_\gamma)$ of \mathcal{T}_γ and note that this branch is in the model N' because it can be computed using the tree $S \in N'$. For all sufficiently large $\alpha < \gamma$ the intersection $b \cap \sigma_{\alpha, \gamma}[\text{lh}(\mathcal{T}_\alpha)]$ is cofinal in $\text{lh}(\mathcal{T}_\gamma)$, which implies that the tree $\mathcal{T}_\alpha^\frown \sigma_{\alpha, \gamma}^{-1}[b]$ is a hull of $\mathcal{T}_\gamma^\frown b$. So by hull condensation we have $\sigma_{\alpha, \gamma}^{-1}[b] = \Lambda(\mathcal{T}_\alpha)$ for all such α , and by considering such an α and its successor we get $\Lambda(\mathcal{T}_\alpha) = \sigma_{\alpha, \alpha+1}^{-1}[\Lambda(\mathcal{T}_{\alpha+1})]$, a contradiction. \square

Remark 5.7. *The proof above is given in the case \mathcal{Q} is a coarse premouse. In the case \mathcal{Q} is a (fine-structural) \mathcal{F} -premouse for some \mathcal{F} , one only needs slight modifications. In particular, one needs to require that the last model of the tree \mathcal{T} in the proof is a well-founded \mathcal{F} -premouse.*

6 Ω^* is constructibly closed

The main theorem of this section is the following.

Theorem 6.1 (ZF + DC $_{\mathbb{R}}$). *Assume there is no transitive AD $^+$ model M containing $\mathbb{R} \cup \text{OR}$ such that there is a pointclass $\Gamma \subsetneq \wp(\mathbb{R})^M$ with $L(\Gamma) \cap \wp(\mathbb{R}) = \Gamma$ and $L(\Gamma) \models \text{AD}_{\mathbb{R}} + \text{DC}$. Then $L(\Omega^*) \cap \wp(\mathbb{R}) = \Omega^*$.*

Remark 6.2. *We note that the smallness assumption in Theorem 6.1 is stronger than (\dagger) . It allows for the existence of a minimal model of “AD $_{\mathbb{R}}$ + DC” but not much more. The Solovay sequence of the minimal model of “AD $_{\mathbb{R}}$ + DC” has length ω_1 . We will use (\dagger^+) to denote this hypothesis.*

We assume (\dagger^+) throughout this section. Suppose the Solovay sequence of Ω^* is of successor length.³⁴ Then by Section 4, $\Omega^* = \wp(\mathbb{R}) \cap M$, where for some operator \mathcal{F} ,

³⁴The Solovay sequence $(\theta_\alpha : \alpha < \gamma)$ of a pointclass Ω^* with the property that if $A \in \Omega^*$, then $L(A, \mathbb{R}) \models \text{AD}^+$ and $\wp(\mathbb{R}) \cap L(A, \mathbb{R}) \subseteq \Omega^*$ is defined as follows. θ_0 is the supremum of α such that there is some $A \in \Omega^*$ and some OD $^{L(A, \mathbb{R})}$ surjection $\pi : \mathbb{R} \rightarrow \alpha$. If $\lambda < \gamma$ is limit, then $\theta_\gamma = \sup_{\alpha < \lambda} \theta_\alpha$. If θ_α has been defined and $\alpha + 1 < \gamma$, then letting $A \in \Omega^*$ be of Wadge rank θ_α , $\theta_{\alpha+1}$ is the supremum of β such that there is some $B \in \Omega^*$ and some OD $(A)^{L(B, \mathbb{R})}$ surjection $\pi : \mathbb{R} \rightarrow \beta$.

$$M = \bigcup \{\mathcal{M} \triangleleft \text{Lp}^{\mathcal{F}}(\mathbb{R}, \text{Code}(\mathcal{F})) \mid \mathcal{M} \models \text{AD}^+ \wedge \mathcal{M} \text{ is self-iterable}\},^{35}$$

and furthermore, Section 4 also shows that

$$\wp(\mathbb{R}) \cap M = \wp(\mathbb{R}) \cap L(M).$$

Clearly, this then shows that $\Omega^* = \wp(\mathbb{R}) \cap L(\Omega^*)$.

Suppose now the Solovay sequence of Ω^* is of limit length. Let \mathcal{H} be the direct limit of all hod pairs $(\mathcal{Q}, \Lambda) \in \Omega^*$ such that Λ has branch condensation and is Ω^* -fullness preserving. \mathcal{H} is a union of hod premice and by (†) and [3], \mathcal{H} has ordinal height Θ^{Ω^*} .³⁶ Let λ be the length of the Solovay sequence of Ω^* , so λ is a limit ordinal. By the smallness assumption of the theorem, $\lambda \leq \omega_1$. From now on, we write Θ^* for Θ^{Ω^*} and θ_α^* for each $\theta_\alpha^{\Omega^*}$ on the Solovay sequence of Ω^* .

The following is the main lemma.

Lemma 6.3 (ZF + DC $_{\mathbb{R}}$). *There is no $\mathcal{M} \trianglelefteq L[\mathcal{H}]$ such that $\mathcal{H} \in \mathcal{M}$ and $\rho_\omega(\mathcal{M}) < \Theta^*$.*

Proof. Suppose not. Let $\mathcal{N} \trianglelefteq L[\mathcal{H}]$ be least such that $\rho_\omega(\mathcal{N}) < \Theta^*$. Let $B \in \Omega^*$ be of Wadge rank θ_{n+1}^* where $n < \lambda$ is such that $\rho_\omega(\mathcal{N}) \leq \theta_n^*$ and $\theta_n^* \geq v$, where v is the \mathcal{N} -cofinality of λ . Suppose k is the least such that $\rho_{k+1}(\mathcal{N}) < \Theta^*$; we may assume $\rho_{k+1}(\mathcal{N}) \leq \theta_n^*$. Let $M = L_\gamma(\mathbb{R}, B, \mathcal{N})$, where γ is some sufficiently large cardinal so that $L_\gamma(\mathbb{R}, B, \mathcal{N}) \models \text{ZF}^- + \text{DC}$.

For countable $\sigma \prec M$ containing all relevant objects, let $\pi_\sigma : M_\sigma \rightarrow M$ be the transitive uncollapse map whose range is σ . Such a σ exists by DC in $L(\mathbb{R}, B, \mathcal{N})$. For each such σ , let $\pi_\sigma(\mathcal{H}_\sigma, \Theta_\sigma, \lambda_\sigma, \mathcal{N}_\sigma, B_\sigma, v_\sigma) = (\mathcal{H}, \Theta^*, \lambda, \mathcal{N}, B, v)$. Let $\Sigma_\sigma^- = \bigoplus_{\alpha < \lambda_\sigma} \Sigma_{\mathcal{H}_\sigma(\alpha)}$. Note that for each $\alpha < \lambda_\sigma$, $\Sigma_{\mathcal{H}_\sigma(\alpha)}$ acts on all countable stacks as it is the pullback of some hod pair (\mathcal{R}, Λ) with the property that $\mathcal{M}_\infty(\mathcal{R}, \Lambda) = \mathcal{H}(\pi_\sigma(\alpha))$.

Let $\sigma \prec M$ be such that $\omega_1^{M_\sigma} > n$; this is possible since $n < \lambda \leq \omega_1$. $\Sigma_{\mathcal{H}_\sigma(n+1)}$ is Ω^* -fullness preserving and has branch condensation. This follows from the choice of B , which gives that $(\mathcal{H}_\sigma(n+1), \Sigma_{\mathcal{H}_\sigma(n+1)})$ is a tail of some hod pair $(\mathcal{Q}, \Lambda) \in M_\sigma$ such that \mathcal{Q} has $n+1$ Woodin cardinals and Λ has branch condensation and is Ω^* -fullness preserving. We let Σ_σ^n be the fragment of Σ_σ^- for stacks on \mathcal{N}_σ above $\delta_n^{\mathcal{N}_\sigma}$. Note that Σ_σ^n is an iteration strategy of \mathcal{N}_σ above $\delta_n^{\mathcal{N}_\sigma}$ since Σ_σ^n -iterations are above v_σ , which may be measurable in \mathcal{N}_σ , and hence does not create new Woodin cardinals. Σ_σ^n has branch condensation. We then have that $\Sigma_\sigma^n \in \Omega^*$; otherwise, by results in the previous sections, we can show $L(\Sigma_\sigma^n, \mathbb{R}) \models \text{AD}^+$ and

³⁵This means whenever \mathcal{M}^* is countable and transitive and there is an elementary embedding from \mathcal{M}^* into \mathcal{M} , then \mathcal{M}^* is $(\omega, \omega_1 + 1)$ - \mathcal{F} -iterable in \mathcal{M} .

³⁶In fact, the universe of \mathcal{H} is precisely the set of all bounded subsets A of Θ^{Ω^*} such that A is OD in $L(B, \mathbb{R})$ for some $B \in \Omega^*$.

this contradicts the definition of Ω^* .³⁷ Also, by [3, Theorem 3.26], Σ_σ^n is Γ -fullness preserving where $\Gamma =_{\text{def}} \Gamma(\mathcal{N}_\sigma, \Sigma_\sigma^n)$.

We then consider the directed system \mathcal{F} of tuples (\mathcal{Q}, Λ) where \mathcal{Q} agrees with \mathcal{N}_σ up to $\delta_n^{\mathcal{N}_\sigma}$, and (\mathcal{Q}, Λ) is Dodd–Jensen equivalent to $(\mathcal{H}_\sigma, \Sigma_\sigma^n)$, that is (\mathcal{Q}, Λ) and $(\mathcal{H}_\sigma, \Sigma_\sigma^n)$ coiterate (above $\delta_n^{\mathcal{N}_\sigma}$) to a hod pair (\mathcal{R}, Ψ) . \mathcal{F} can be characterized as the directed system of hod pairs (\mathcal{Q}, Λ) extending $(\mathcal{N}_\sigma(n), \Sigma_{\mathcal{N}_\sigma(n)})$ such that $\Gamma(\mathcal{Q}, \Lambda) = \Gamma$, Λ has branch condensation and is Γ -fullness preserving. We note that \mathcal{F} is $\text{OD}_{\Sigma_{\mathcal{H}_\sigma(n)}}$ in $L(C, \mathbb{R})$ for some $C \in \Omega^*$. We fix such a C ; so $L(C, \mathbb{R}) \models \text{AD}^+ + \text{SMC}$. Let $A \subseteq \delta_n^{\mathcal{N}_\sigma}$ witness $\rho_{k+1}(\mathcal{N}_\sigma) \leq \delta_n^{\mathcal{N}_\sigma}$. Then A is $\text{OD}_{\Sigma_{\mathcal{H}_\sigma(n)}}$ in $L(C, \mathbb{R})$. By SMC in $L(C, \mathbb{R})$ and the fact that $\mathcal{N}_\sigma(n+1)$ is Ω^* -full, $A \in \text{Lp}^{\Sigma_{\mathcal{H}_\sigma(n)}}(\mathcal{N}_\sigma | \delta_n^{\mathcal{N}_\sigma}) \in \mathcal{N}_\sigma$. This contradicts the definition of A . \square

For $\alpha < \lambda$, let us write $\wp_{\theta_\alpha}(\mathbb{R})$ for $(\wp_{\theta_\alpha}(\mathbb{R}))^{\Omega^*}$ and Σ_α for $\Sigma_\alpha^\mathcal{H}$. We also need the following notation: let $(\mathcal{P}, \Sigma) \in \Omega^*$ be a hod pair, let $\mathcal{M}_{\mathcal{P}, \Sigma}^\sharp = \mathcal{M}_\omega^{\Sigma, \sharp}$ be the minimal \mathcal{P} -sound, active Σ -mouse with ω many Woodin cardinals $\delta_0^{\mathcal{M}_{\mathcal{P}, \Sigma}} < \delta_1^{\mathcal{M}_{\mathcal{P}, \Sigma}} < \dots$, and let $\delta_\omega^{\mathcal{M}_{\mathcal{P}, \Sigma}} = \sup_i \delta_i^{\mathcal{M}_{\mathcal{P}, \Sigma}}$.³⁸ Finally, we let $\mathcal{M}_{\mathcal{P}, \Sigma} = \mathcal{M}_\omega^\Sigma$ be the corresponding proper class mouse obtained from $\mathcal{M}_{\mathcal{P}, \Sigma}^\sharp$ by iterating the top extender OR many times. We remind the reader that at this point, we assume that λ is a limit ordinal.

Lemma 6.4 (ZF + DC $_{\mathbb{R}}$). *Fix $s \in (\Theta^*)^{<\omega}$ and let $\alpha < \lambda$ be such that $s \in (\theta_\alpha^*)^{<\omega}$. Then for any formula ψ and any hod pair $(\mathcal{Q}, \Lambda) \in \Omega^*$ such that Λ is Ω^* -fullness preserving, has branch condensation, and $\Gamma(\mathcal{Q}, \Lambda) = \wp_{\theta_\alpha}(\mathbb{R})$,*

$$L(\Lambda, \mathbb{R}) \models \psi[s] \iff \mathcal{M}_{\alpha, \infty} \models \text{"the derived model satisfies } \psi[i_{\mathcal{H}(\alpha), \infty}^{\Sigma_\alpha}(s)]\text{"} \quad (*)$$

where $\mathcal{M}_{\alpha, \infty}$ is the direct limit of all iterates of $\mathcal{M}_{\mathcal{Q}, \Lambda}$ below $\delta_0^{\mathcal{M}_{\mathcal{Q}, \Lambda}}$ via its canonical strategy and the derived model is computed at $\delta_\omega^{\mathcal{M}_{\alpha, \infty}}$.

Proof. Fix s, ψ, α , and (\mathcal{Q}, Λ) as in the statement of the lemma. First we note that Σ_α is a tail of Λ . Let $\mathcal{P} = \mathcal{M}_{\mathcal{Q}, \Lambda}$ and let Σ be the canonical strategy of \mathcal{P} extending Λ . Note that for any Σ -iterate \mathcal{P}^* of \mathcal{P} , we can iterate \mathcal{P}^* using Σ to some \mathcal{P}' such that $L(\Lambda, \mathbb{R})$ is the derived model of \mathcal{P}' at $\delta_\omega^{\mathcal{P}'}$.³⁹ We may assume also that s is in the range of the direct limit map from \mathcal{P} into $\mathcal{M}_{\alpha, \infty}$.

Suppose the left hand side of the equivalence fails, that is

$$L(\Lambda, \mathbb{R}) \models \neg\psi[s].$$

³⁷We also have that Σ_σ^n is the join of countably many sets of reals, each of which is in Ω^* and hence is Suslin co-Suslin. This implies that Σ_σ^n is self-scaled.

³⁸Sections 3 and 4 show that $\mathcal{M}_{\mathcal{P}, \Sigma}^\sharp$ exists and its canonical strategy is in Ω^* .

³⁹This is analogous to the fact that $L(\mathbb{R})$ is the derived model of an iterate of \mathcal{M}_ω .

Work in $V^{\text{Col}(\omega, \mathbb{R})}$ and let $\{(\mathcal{P}_n, \Sigma_n) \mid n < \omega \wedge (\mathcal{P}_n, \Sigma_n) \in I(\mathcal{P}, \Sigma)\}$ be cofinal in the directed system of Σ -iterates below $\delta_0^\mathcal{P}$; here we take $(\mathcal{P}_0, \Sigma_0) = (\mathcal{P}, \Sigma)$.⁴⁰ For $m \leq n < \omega$, let $i_n : \mathcal{P}_n \rightarrow \mathcal{P}_{n+1}$ be the iteration map and let $i_{m,n} : \mathcal{P}_m \rightarrow \mathcal{P}_n$ and $i_{m,\infty} : \mathcal{P}_m \rightarrow \mathcal{M}_{\alpha,\infty}$ be the natural maps. Set $s_0 = i_{0,\infty}^{-1}(s)$ and let $s_n = i_{0,n}(s_0)$. Let $(\mathcal{P}_k^\omega : k < \omega)$ and $(\pi_{k,l} : \mathcal{P}_k^\omega \rightarrow \mathcal{P}_l^\omega : k \leq l < \omega)$ come from the simultaneous \mathbb{R} -genericity iteration construction described in [17, Lemma 6.50]. We also let $j_i : \mathcal{P}_i \rightarrow \mathcal{P}_i^\omega$ be the iteration map; here the iterations are above the s_i 's, i.e.

$$j_i(s_i) = s_i.$$

By properties of the construction, for $k \leq l < \omega$

$$j_l \circ i_{k,l} = \pi_{k,l} \circ j_k.$$

Let $\mathcal{P}_\omega^\omega$ be the direct limit of \mathcal{P}_k^ω under the embeddings $\pi_{k,l}$ and let $\pi_{i,\omega} : \mathcal{P}_i^\omega \rightarrow \mathcal{P}_\omega^\omega$ and $j_\omega : \mathcal{M}_{\alpha,\infty} \rightarrow \mathcal{P}_\omega^\omega$ be the natural maps. Note that $j_\omega(s) = s$.

By our assumptions, for each i ,

$$\mathcal{P}_i^\omega \models 1 \Vdash \text{“the derived model satisfies } \neg\psi[s] \text{ and } s = i_{\mathcal{P}_i^\omega, \infty}(s_i) \text{”}.$$

Let k be such that for all $l \geq k$, $\pi_{l,l+1}(s) = s$ (k exists because $\mathcal{P}_\omega^\omega$ is well-founded), and let $s^* = \pi_{k,\omega}(s)$. By elementarity,

$$\mathcal{P}_\omega^\omega \models 1 \Vdash \text{“the derived model satisfies } \neg\psi[s^*] \text{ and } s^* = i_{\mathcal{P}_\omega^\omega, \infty}(s) \text{”}.$$

By elementarity of j_ω and the fact that $j_\omega(s) = s$, we get

$$\mathcal{M}_{\alpha,\infty} \models 1 \Vdash \text{“the derived model satisfies } \neg\psi[i_{\mathcal{H}(\alpha), \infty}^{\Sigma_\alpha}(s)] \text{”}.$$

Contradiction. The other direction is proved similarly. \square

Remark 6.5. *The right hand side of $(*)$ can be defined in \mathcal{H} from Σ_α uniformly in Σ_α . This is because the right hand side of $(*)$ is equivalent to the statement: in the derived model of $L[\mathcal{H}]$ at the supremum of its Woodin cardinals $\psi[i_{\mathcal{H}(\alpha), \infty}^{\Sigma_\alpha}(s)]$ holds, where \mathbb{R}^* is the $\text{Col}(\omega, < \Theta)$ -symmetric reals over $L[\mathcal{H}]$ induced by some $g \subseteq \text{Col}(\omega, < \Theta)$. This is because $\mathcal{M}_{\alpha,\infty} \in \mathcal{H}$ and in $L[\mathcal{H}][g]$, we can naturally embed the direct limit of Σ_α -iterates of $\mathcal{H}(\alpha)$ in the derived model of $\mathcal{M}_{\alpha,\infty}$ into the direct limit of Σ_α -iterates of $\mathcal{H}(\alpha)$ in the derived model of $L[\mathcal{H}]$.*

⁴⁰There is an awkward point here. We don't know that (\mathcal{P}, Σ) is iterable in $V^{\text{Col}(\omega, \mathbb{R})}$, but we can run the argument below inside an $L[T, x]$ where T is a tree projecting to some universal Γ set A and Γ is an inductive-like, scaled pointclass beyond $\wp_{\theta_\alpha}(\mathbb{R})$ and $x \in \mathbb{R}$ codes \mathcal{P} as well as the reduction of A to $\text{Code}(\Sigma)$. We may also assume $(*)$ is absolute between Ω and the model $L[T, x]$'s version of Ω . Since $\mathbb{R} \cap L[T, x]$ is countable, we can proceed with the argument below pretending that V is $L[T, x]$, with appropriate definitions of objects $\mathcal{M}_{\alpha,\infty}, \Lambda$ etc.

Recall from [18] the following version of the Vopěnka algebra. For each $\alpha < \lambda$, let \mathbb{P}_α^* be the boolean algebra $(\{A \subseteq \wp(\xi)^n \mid n < \omega \wedge \xi < \theta_\alpha \wedge \exists B \in \Omega^* A \in \text{OD}^{L(B, \mathbb{R})}\}, \subseteq)$ and let $\mathbb{P}_\alpha \in \mathcal{H} \cap \wp(\theta_\alpha)$ be the isomorphic copy of \mathbb{P}_α^* . It's clear that for each α , \mathbb{P}_α^* and \mathbb{P}_α are OD in $L(\wp_{\theta_\beta}(\mathbb{R}))$ for any $\beta > \alpha$ and the definition is uniform in α . Furthermore, for $\alpha < \beta$, there is a natural embedding of \mathbb{P}_α^* into \mathbb{P}_β^* (and hence from \mathbb{P}_α into \mathbb{P}_β) and these embeddings are also OD in $L(\wp_{\theta_\gamma}(\mathbb{R}))$ for any $\gamma > \beta$ and again, the definition is uniform in α and β . Let \mathbb{P} be the direct limit of the \mathbb{P}_α 's under the natural embeddings. The following corollary of Lemma 6.4 shows that $\mathbb{P} \in L[\mathcal{H}]$. We note that in the corollary below, the language of the structure $L[\mathcal{H}]$ has the predicate for the sequence of strategies $\{\Sigma_\alpha \mid \alpha < \lambda\}$.

Corollary 6.6. *For each $\alpha < \lambda$, \mathbb{P}_α is definable in $L[\mathcal{H}]$ from $\{\theta_{\alpha+1}, \Sigma_{\alpha+1}\}$, uniformly in α . Similarly, for $\alpha < \beta$, the natural embedding from \mathbb{P}_α into \mathbb{P}_β is definable in $L[\mathcal{H}]$ from $\{\theta_{\alpha+1}, \theta_{\beta+1}, \Sigma_{\alpha+1}, \Sigma_{\beta+1}\}$, uniformly in α and β . Consequently, $\mathbb{P} \in L[\mathcal{H}]$.*

Proof. We just prove the first clause; the proof of the second clause is similar. Fix any $\beta > \alpha$ and let (\mathcal{Q}, Λ) , (\mathcal{P}, Σ) , and $\mathcal{M}_{\beta, \infty}$ be defined as in the proof of Lemma 6.4 but for Σ_β . Note that $\mathbb{P}_\alpha \in \mathcal{H}(\beta)$.⁴¹ By Lemma 6.4 and the subsequent remark,

$L[\mathcal{H}] \models 1 \Vdash$ in the derived model, $L(\Sigma_\beta, \mathbb{R}^*)$ satisfies “ $i_{\mathcal{H}(\beta), \infty}(\mathbb{P}_\alpha)$ is the Vopěnka algebra at $i_{\mathcal{H}(\beta), \infty}(\theta_\alpha)$ ”.

The above gives a uniform definition of \mathbb{P}_α from $\{\theta_\beta, \Sigma_\beta\}$ inside $L[\mathcal{H}]$ for any $\beta > \alpha$.

Clearly, the third clause follows from the first two clauses. \square

Using Corollary 6.6 and [18, Theorem 4.3.19], we can conclude that

- $L[\mathcal{H}](\Omega^*)$ is a symmetric extension of $L[\mathcal{H}]$ via \mathbb{P} .
- $\wp(\mathbb{R}) \cap L[\mathcal{H}](\Omega^*) = \Omega^*$.

These, in particular, imply $L(\Omega^*) \cap \wp(\mathbb{R}) = \Omega^*$. This completes the proof of Theorem 6.1.

Lemma 6.3 shows that $V_{\Theta^\Omega} \cap L[\mathcal{H}] = |\mathcal{H}|$. In the case $L[\mathcal{H}] \models$ “the set of Woodin cardinals has limit order type”, let M be the derived model of $L[\mathcal{H}]$ (at the supremum of $L[\mathcal{H}]$'s Woodin cardinals). Then $M \models \text{AD}_\mathbb{R}$ (cf. [3, Section 3.3]). This, combined with the result of the previous section, proves Theorem 1.6; Theorem 6.1 proves something stronger, namely, Ω^* is constructibly closed.

Lemma 6.7. *If DC holds and the order type of the Woodin cardinals of \mathcal{H} is a limit ordinal, then $\text{cf}(\Theta^*) > \omega$ and $L[\mathcal{H}](\Omega^*) \models \text{AD}_\mathbb{R} + \text{DC}$.*

⁴¹ \mathbb{P}_α need not be in $\mathcal{H}(\alpha)$.

Proof. Suppose $\text{cf}(\Theta^*) = \omega$. Let M be a transitive structure containing $\mathcal{H}^+ \cup \Omega^*$ for $\mathcal{H}^+ = L[\mathcal{H}]|\gamma$, where $\gamma > \Theta^*$ is a regular cardinal in $L[\mathcal{H}]$. Let $\sigma \prec M$ be countable such that σ is cofinal in Θ^* ; the existence of such a σ follows from DC. Now the π_σ -realizable strategy Σ_σ defined in the proof of Lemma 6.3 acts on $\pi_\sigma^{-1}(\mathcal{H}^+)$. Σ_σ on stacks below Θ_σ is simply Σ_σ^- in this case; by replacing $(\mathcal{N}_\sigma, \Sigma_\sigma)$ by an iterate, we may assume Σ_σ has branch condensation. We can show then that $\Sigma_\sigma \in \Omega^*$ as before. Furthermore, letting i be the direct limit map from $\pi_\sigma^{-1}(\mathcal{H}^+)$ into the direct limit \mathcal{M}_∞ of all of its Σ_σ -iterates in Ω^* , then by elementarity

$$\pi_\sigma \upharpoonright \Theta_\sigma = i \upharpoonright \Theta_\sigma.$$

So i is cofinal in Θ^* and \mathcal{H} is the direct limit of hod initial segments of $\pi_\sigma^{-1}(\mathcal{H})$ via Σ_σ . Let $\Omega = \Sigma_\sigma$. By a core model induction through $\text{sLp}^{\mathcal{G}_\Omega}(\mathbb{R}, \text{Code}(\Omega))$ like in the previous sections, we get a hod pair $(\mathcal{Q}, \Lambda) \in \Omega^*$ such that letting \mathcal{Q}_∞ be the direct limit of all Λ -iterates of \mathcal{Q} , $\mathcal{H} \triangleleft \mathcal{Q}_\infty$. This contradicts the definition of \mathcal{H} .

The second clause follows immediately from the first clause and [11]. \square

We have completed the proof of the following theorems.

Theorem 6.8 (ZF + DC $_{\mathbb{R}}$). *Suppose $\Omega^* = \{A \subseteq \mathbb{R} \mid L(A, \mathbb{R}) \models \text{AD}^+\}$ and (\dagger^+) holds. Suppose $\Omega^* \neq \emptyset$ and for every suitable pair (\mathcal{P}, Σ) or hod pair (\mathcal{P}, Σ) such that Σ has branch condensation and is Ω^* -fullness preserving, $\Sigma \in \Omega^*$. If the Solovay sequence of Ω^* has limit length, then $\Omega^* = L(\Omega^*, \mathbb{R}) \cap \wp(\mathbb{R})$ and $L(\Omega^*, \mathbb{R}) \models \text{AD}_{\mathbb{R}}$.*

Theorem 6.9 (ZF + DC). *Suppose $\Omega^* = \{A \subseteq \mathbb{R} \mid L(A, \mathbb{R}) \models \text{AD}^+\}$ and (\dagger^+) holds. Suppose $\Omega^* \neq \emptyset$ and for every suitable pair (\mathcal{P}, Σ) or hod pair (\mathcal{P}, Σ) such that Σ has branch condensation and is Ω^* -fullness preserving, $\Sigma \in \Omega^*$. If the Solovay sequence of Ω^* has limit length, then $L(\Omega^*, \mathbb{R}) \cap \wp(\mathbb{R}) = \Omega^*$ and $L(\Omega^*, \mathbb{R}) \models \text{AD}_{\mathbb{R}} + \text{DC}$.*

Together with the results of the previous section, the above theorems complete the proof of Theorems 1.6 and 1.7.

7 Further results, questions, and open problems

We first mention a few natural questions regarding possible weakenings of the hypotheses of Theorems 1.5 and 1.7. (In some cases one could also formulate versions with fragments of DC along the lines of 1.6.)

Question 7.1. *What are the consistency strengths of the following theories:*

1. ZF + DC + “ ω_1 is ω_2 -strongly compact”?

2. $\text{ZF} + \text{DC} + \text{``}\omega_1 \text{ is } \Theta\text{-strongly compact''}$?

Are they equiconsistent with $\text{ZF} + \text{DC} + \text{AD}$ and $\text{ZF} + \text{DC} + \text{AD}_{\mathbb{R}}$ respectively?

One could try to weaken the compactness hypotheses further:

Question 7.2. *What are the consistency strengths of the following theories:*

1. $\text{ZF} + \text{DC} + \text{``}\omega_1 \text{ is threadable and } \neg \square_{\omega_1}\text{''}$?
2. $\text{ZF} + \text{DC} + \text{``every uncountable regular cardinal } \leq \Theta \text{ is threadable''}$?

Are they equiconsistent with $\text{ZF} + \text{DC} + \text{AD}$ and $\text{ZF} + \text{DC} + \text{AD}_{\mathbb{R}}$ respectively?

However, it may be overly ambitious at present to seek a positive answer especially in case 2; one could try to answer the following question first:

Question 7.3. *What is the consistency strength of the theory $\text{ZF} + \text{DC} + \text{``}\omega_1 \text{ is } \mathbb{R}\text{-strongly compact and } \Theta \text{ is threadable''}$? Is it equiconsistent with $\text{ZF} + \text{DC} + \text{AD}_{\mathbb{R}}$?*

We mention a corollary of the proof of Theorem 1.5.

Theorem 7.4. *The following theories are equiconsistent:*

1. $\text{ZF} + \text{DC} + \text{AD}$
2. $\text{ZF} + \text{DC} + \text{``}\omega_1 \text{ is } \mathbb{R}\text{-strongly compact and } \Theta > \omega_2\text{''}$

Proof. (1) \implies (2): As mentioned in Section 1, the statement “ ω_1 is \mathbb{R} -strongly compact” is a consequence of the existence of the Turing cone measure, which follows from AD, and the statement $\Theta > \omega_2$ follows from the Moschovakis coding lemma.

(2) \implies (1): Using a push-forward measure, it’s easy to see that statement (2) above implies statement (3) of Theorem 1.5. \square

If we strengthen statement (2) above to “ ω_1 is \mathbb{R} -supercompact and $\Theta > \omega_2$ ”,⁴² then we obtain an equiconsistency with “there are ω^2 many Woodin cardinals”, which is strictly stronger than AD. This is a result of Woodin (see [19]). Similarly, if we strengthen statement (2) of Theorem 1.7 to $\text{ZF} + \text{DC} + \text{``}\omega_1 \text{ is } \wp(\mathbb{R})\text{-supercompact''}$ then we obtain the sharp for a model of $\text{AD}_{\mathbb{R}} + \text{DC}$. To see this, note that from the result of Theorem 1.7, we get a model $L(\Omega^*, \mathbb{R}) \models \text{AD}_{\mathbb{R}} + \text{DC}$, where $\Omega^* \subseteq \wp(\mathbb{R})$. Fix a countably complete, fine, normal measure μ on $\wp_{\omega_1}(\wp(\mathbb{R}))$. Then note that by normality,

⁴²We say that ω_1 is X -supercompact if there is a countably complete, fine, normal measure μ on $\wp_{\omega_1}(X)$. μ is normal on $\wp_{\omega_1}(X)$ if whenever $F : \wp_{\omega_1}(X) \rightarrow \wp_{\omega_1}(X)$ is such that $\{\sigma \mid F(\sigma) \subseteq \sigma \wedge F(\sigma) \neq \emptyset\} \in \mu$ then there is some $x \in X$ such that the set $\{\sigma \mid x \in F(\sigma)\} \in \mu$.

$$\forall_\mu^* \sigma L(\Omega_\sigma^*, \mathbb{R}_\sigma) \models \text{AD}_\mathbb{R} + \text{DC},$$

where we have that $\Omega^* = [\sigma \mapsto \Omega_\sigma^*]_\mu$ and $\mathbb{R} = [\sigma \mapsto \mathbb{R}_\sigma]_\mu$. Now, $\forall_\mu^* \sigma (\Omega_\sigma^*, \mathbb{R}_\sigma)^\sharp$ exists; by normality again, the sharp for $L(\Omega^*, \mathbb{R})$ exists. This demonstrates that the theory $\text{ZF} + \text{DC} + \text{``}\omega_1 \text{ is } \wp(\mathbb{R})\text{-supercompact''}$ is strictly stronger than $\text{ZF} + \text{DC} + \text{``}\omega_1 \text{ is } \wp(\mathbb{R})\text{-strongly compact''}$. However, we don't know its exact consistency strength.

Question 7.5. *What is the exact consistency strength of $\text{ZF} + \text{DC} + \text{``}\omega_1 \text{ is } \wp(\mathbb{R})\text{-supercompact''}$?*

We end with the following set of questions.

Question 7.6. *What are the consistency strengths of the following theories:*

1. “ $\text{ZF} + \text{DC} + \text{``}\omega_1 \text{ is } \wp(\wp(\mathbb{R}))\text{-strongly compact''}$ ”?
2. “ $\text{ZF} + \text{DC} + \text{``}\omega_1 \text{ is } \wp(\wp(\mathbb{R}))\text{-supercompact''}$ ”?
3. “ $\text{ZF} + \text{DC} + \omega_1 \text{ is strongly compact}$ ”?
4. “ $\text{ZF} + \text{DC} + \omega_1 \text{ is supercompact}$ ”?

In particular, are the theories (3) and (4) equiconsistent?

It's worth noting that Woodin (unpublished) has shown the theory “ $\text{ZF} + \text{DC} + \omega_1$ is supercompact” is consistent relative to a proper class of Woodin limits of Woodin cardinals. We hope the techniques in this paper when combined with the theory of hod mice would allow us to make significant progress in answering these questions.

References

- [1] Steve Jackson. Structural consequences of AD. In Matthew Foreman and Akihiro Kanamori, editors, *Handbook of Set Theory*, volume 3, pages 1753–1876. Springer, 2010.
- [2] Thomas Jech. *Set theory*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003. The third millennium edition, revised and expanded.
- [3] G. Sargsyan. *Hod Mice and the Mouse Set Conjecture*, volume 236 of *Memoirs of the American Mathematical Society*. American Mathematical Society, 2015.
- [4] Grigor Sargsyan and John Steel. The Mouse Set Conjecture for sets of reals. *The Journal of Symbolic Logic*, pages 671–683, 2015.

- [5] E. Schimmerling and J. R. Steel. The maximality of the core model. *Trans. Amer. Math. Soc.*, 351(8):3119–3141, 1999.
- [6] Ernest Schimmerling and Martin Zeman. Square in core models. *Bulletin of Symbolic Logic*, 7(03):305–314, 2001.
- [7] Ralf Schindler and John Steel. The core model induction. Available at https://ivv5hpp.uni-muenster.de/u/rds/core_model_induction.pdf, 2014.
- [8] Ralf-Dieter Schindler. Successive weakly compact or singular cardinals. *The Journal of Symbolic Logic*, 64(01):139–146, 1999.
- [9] F. Schlutzenberg and N. Trang. The fine structure of operator mice. *arXiv:1604.00083v1*, 2016.
- [10] F. Schlutzenberg and N. Trang. Scales in hybrid mice over \mathbb{R} . *arXiv:1210.7258v4*, 2016.
- [11] Robert M. Solovay. The independence of DC from AD. In *Cabal Seminar 76–77*, pages 171–183. Springer, 1978.
- [12] John R. Steel. PFA implies $\text{AD}^{\text{L}(\mathbb{R})}$. *J. Symbolic Logic*, 70(4):1255–1296, 2005.
- [13] John R. Steel. Derived models associated to mice. In *Computational prospects of infinity. Part I. Tutorials*, volume 14 of *Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap.*, pages 105–193. World Sci. Publ., Hackensack, NJ, 2008.
- [14] John R. Steel. Scales in $K(\mathbb{R})$ at the end of a weak gap. *J. Symbolic Logic*, 73(2):369–390, 2008.
- [15] John R. Steel. Scales in $\mathbf{K}(\mathbb{R})$. In *Games, scales, and Suslin cardinals. The Cabal Seminar. Vol. I*, volume 31 of *Lect. Notes Log.*, pages 176–208. Assoc. Symbol. Logic, Chicago, IL, 2008.
- [16] John R. Steel. An outline of inner model theory. In Matthew Foreman and Akihiro Kanamori, editors, *Handbook of Set Theory*, volume 3, pages 1595–1684. Springer, 2010.
- [17] John R. Steel and W. Hugh Woodin. HOD as a core model. In *Ordinal definability and recursion theory: the Cabal Seminar*, volume 3, pages 257–346, 2016.
- [18] N. Trang. *Generalized Solovay measures, the HOD analysis, and the core model induction*. PhD thesis, Berkeley, 2013.

- [19] N. Trang. Structure theory of $L(\mathbb{R}, \mu)$ and its applications. *The Journal of Symbolic Logic*, 80(01):29–55, 2015.
- [20] Trevor M. Wilson. *Contributions to descriptive inner model theory*. PhD thesis, University of California, Berkeley, 2012.
- [21] Trevor M. Wilson. The envelope of a pointclass under a local determinacy hypothesis. *Annals of Pure and Applied Logic*, 166(10):991–1018, 2015.
- [22] Trevor M. Wilson. Scales on Π_1^2 sets. *Mathematical Research Letters*, 22(1):301–316, 2015.
- [23] W. H. Woodin. AD and the uniqueness of the supercompact measures on $P_{\omega_1}(\lambda)$. In *Cabal Seminar 79–81*, pages 67–71. Springer, 1983.