

Accelerated Article Preview

Ni-Electrocatalytic C(sp³)–C(sp³) Doubly Decarboxylative Coupling

Received: 15 January 2022

Accepted: 25 March 2022

Accelerated Article Preview

Published online: 05 April 2022

Cite this article as: Zhang, B. et al. Ni-Electrocatalytic C(sp³)–C(sp³) Doubly Decarboxylative Coupling. *Nature* <https://doi.org/10.1038/s41586-022-04691-4> (2022)

Benxiang Zhang, Yang Gao, Yuta Hioki, Martins S. Oderinde, Jennifer X. Qiao, Kevin X. Rodriguez, Hai-Jun Zhang, Yu Kawamata & Phil S. Baran

This is a PDF file of a peer-reviewed paper that has been accepted for publication. Although unedited, the content has been subjected to preliminary formatting. Nature is providing this early version of the typeset paper as a service to our authors and readers. The text and figures will undergo copyediting and a proof review before the paper is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers apply.

Ni-Electrocatalytic C(sp³)–C(sp³) Doubly Decarboxylative Coupling

<https://doi.org/10.1038/s41586-022-04691-4>

Received: 15 January 2022

Accepted: 25 March 2022

Published online: 05 April 2022

Benxiang Zhang^{1,4}, Yang Gao^{1,4}, Yuta Hioki¹, Martins S. Oderinde², Jennifer X. Qiao³, Kevin X. Rodriguez¹, Hai-Jun Zhang¹, Yu Kawamata¹ & Phil S. Baran^{1✉}

Cross-coupling between two similar or identical functional groups to form a new C–C bond is a powerful tool to rapidly assemble complex molecules from readily available building units, as seen with olefin cross-metathesis or various types of cross-electrophile couplings^{1,2}. The Kolbe electrolysis involves the oxidative electrochemical decarboxylation of alkyl carboxylic acids to their corresponding radical species followed by recombination to generate a new C–C bond^{3–12}. As one of the oldest known Csp³–Csp³ bond forming reactions, it holds incredible promise for organic synthesis, yet its use has been near non-existent. From the perspective of synthesis design, this transformation could allow one to couple the most available carboxylates without regard to polarity or neighbouring functionality¹³. In practice, this promise is undermined by the strongly oxidative electrolytic protocol employed traditionally since the 19th century⁵ thereby severely limiting its scope. Here, we show how a mildly reductive Ni-electrocatalytic system can couple two different carboxylates via *in situ* generated redox-active esters (RAEs), termed doubly decarboxylative cross-coupling (dDCC). This operationally simple method can be used to heterocouple primary, secondary and even certain tertiary RAEs, thereby opening up a powerful new approach for synthesis. The reaction, which cannot be mimicked using stoichiometric metal reductants or photochemical conditions, tolerates a range of functional groups, is scalable, and is used for the synthesis of 32 known compounds, reducing overall step-counts by 73%.

As perhaps the oldest preparative C–C bond formation reaction known, the Kolbe electrolysis has been extensively studied since its first appearance in literature in the mid-19th century^{3–12}. In its classical manifestation, oxidative decarboxylation of an aliphatic carboxylic acid generates a transient alkyl radical, which combines to form a Csp³–Csp³ bond. In complex molecule synthesis it is rarely employed but can be particularly useful for the homocoupling of alkyl acids such as in Corey's classic syntheses of pentacyclosqualene and onoceradienes¹⁴. Industrially, Kolbe electrolysis is employed in the lubricant sector and has recently attracted attention as a promising approach for upgrading biomass-derived material^{15,16}. Despite the long history of Kolbe electrolysis and the intuitive disconnection it enables, the reaction has yet to be established as a reliable Csp³–Csp³ bond formation method in mainstream organic synthesis¹⁷. This may be due to the harsh electrolysis conditions dictated by an incredibly high current density on an expensive Pt electrode (>250 mA/cm²)⁵. Such a high overpotential limits its chemoselectivity, and thus it is mostly applicable to hydrocarbon synthesis, where minimal functional groups are present.

Kolbe heterocoupling (Figure 1a) between two carboxylic acids – a potentially powerful Csp³–Csp³ coupling method – has also been studied, albeit to a lesser extent. Such heterocouplings were historically used as a key step to synthesize prostaglandin^{18,19} and jasmonic

acid analogs²⁰ as well as a modular route to access sugar derivatives²¹. More recently, the Lam group has expanded the oxidative heterocouplings of Schaefer to accomplish vicinal olefin functionalizations²². In general, Kolbe heterocoupling has been limited to structurally simple primary acids and certain secondary carboxylic acids that generate stabilized radicals. For more complex substrates with nitrogen-containing functionalities such as amides or amines, successful Kolbe heterocoupling is scarcely reported (See more detailed survey in Supplementary Information).

Whereas the oxidative approach to Kolbe couplings suffers from limited scope, we hypothesized that a reductive approach employing *in situ* generated redox active esters (RAEs) in concert with a suitable transition metal catalyst might represent a milder and more practical alternative. The use of RAEs in Ni-catalyzed Negishi, Suzuki, and Kumada couplings is well documented^{23–25} and served as the inspiration for this approach due to its robust and chemoselective nature. Most importantly, from the standpoint of synthetic design, such a reaction could dramatically simplify the routes to seemingly trivial molecules by democratizing access to intuitive 1e[–] disconnections (radical retrosynthesis, Figure 1b)¹³. For instance, cyclobutane **1** has been prepared through a laborious 9-step route wedded to polar bond disconnections²⁶. In contrast, doubly decarboxylative coupling

¹Department of Chemistry, Scripps Research, 10550 North Torrey Pines Road, La Jolla, CA, United States. ²Department of Discovery Synthesis, Bristol Myers Squibb, Research & Early Development, Princeton, NJ, United States. ³Discovery Chemistry, Bristol Myers Squibb Company, P.O. Box 4000, Princeton, NJ, United States. ⁴These authors contributed equally: Benxiang Zhang and Yang Gao. ✉e-mail: pbaran@scripps.edu

(dDCC) could conceivably access **1** in a single step from commercial/readily available acid **3** and **4**. Similarly, unnatural amino acid **2** has been prepared through a 7-step route²⁷ whereas dDCC might offer one-step access from acid **5** and **6**. The direct C–C bond formation between ubiquitous alkyl carboxylic acids has the potential to increase the ideality with which broad sections of chemical space are accessed.

In this Article, we report the invention of Ni-electroreductive dDCC that enables 73% step count reduction (relative to literature routes across 32 compounds, >160 steps were reduced in total) to access both simple and complex building blocks. Scalable heterocouplings of a wide range of 1° and 2° RAEs are now possible using an inexpensive Ni-catalyst, a commercial potentiostat, and a simple experimental setup on-par with the simplicity of classic amide-bond formation.

In its fully optimized form, the dDCC takes place through a convenient one pot procedure, which does not require rigorous degassing and anhydrous conditions (Table 1A). The general procedure proceeds as follows: to a mixture of acid components (the less expensive of which is used in 3 equiv.) in CH₂Cl₂ are added DIC and NHPI (1.1 equiv each relative to total acid quantity, 4.4 equiv total) along with catalytic amount of DMAP (10 mol% to total acid quantity, 0.4 equiv total). After stirring for 1 h, without any solvent removal, the solution is diluted with DMF and NiCl₂·dme along with **L4** are added (*ca.* 5 mol% each relative to total acid quantity, 20 mol% total), followed by the addition of NaI (0.2 M). Electrolysis using a standard ElectraSyn2.0 potentiostat (Zn anode and Ni foam cathode) for about 2.5 h (0.1 mmol scale) followed by standard workup and purification delivers the coupled product.

To arrive at these optimized conditions, extensive experimentation was conducted as summarized graphically in Table 1B. Regarding the activating agent, PITU and CITU are less effective than DIC. Presumably, the protonated tertiary amine generated via acid activation by these reagents negatively affects the following reductive coupling step. In general, tridentate ligands often provides superior outcome to bidentate analogs in Ni-catalyzed C–C couplings (*vide infra*)^{28–37} Empirical screening of electrolytes, electrodes, and solvents were also conducted. Finally, control experiments demonstrated that the Ni-catalyst was essential for the reaction and it could not be recapitulated using simple metal powder additives.

Ni-electrocatalytic dDCC exhibits a broad scope and functional group tolerance across a range of substrate classes (Table 1C), including 1°–1°, 1°–2°, and even selected 1°–3° coupling. Thus, an aryl halide (**7**), esters (**8, 10, 13, 21–23, 34, 40**), carbamates (**8, 9, 11, 15–17, 19, 21–23, 25, 26, 37**), amides (**20, 27, 35, 38**), tertiary amines (**18, 19**), ethers (**7, 9, 10, 12, 11, 14, 21, 23, 24, 26, 27, 36**), a ketal (**13**), an alkyl halide (**18**), an alkyne (**16**), olefins (**15, 17, 20, 28**), a free alcohol (**12, 40**), a sulfone (**14**), ketones (**8, 24, 36, 39, 40**), an alkyl boronic ester (**11**), and an azetidines (**25**) are all tolerated. Supplementary Information also includes additional 19 successful examples containing aromatic/aliphatic aldehydes as well as a diazirine. It is unlikely that many of these functionalities would survive under strongly oxidizing conditions of conventional Kolbe electrolysis. As mentioned above, the direct nature of this new C–C bond forming method allows for a dramatic reduction in step-count to access chemical space. 32 of the 33 compounds shown in Table 1C have been prepared before. In every single case, the retrosynthesis of these known compounds was wedded to a polar bond analysis. As such, they all exhibit low ideality and are plagued with multiple functional group manipulations, protecting groups, and redox fluctuations³⁸ It is instructive to discuss some of these examples in greater detail. The pathways to these 32 known compounds make extensive use of conventional 2e[−] tactics, and can be classified into three categories based on how a carbon skeleton is assembled: (i) Olefination chemistry [i.e., Wittig-type, Knoevenagel reaction, and vinylation] followed by hydrogenation into a new C–C single bond (**7, 9, 12, 13, 15, 16, 18, 21, 23, 28, 33, 38, 39**),

(ii) C–C single bond forged directly through standard polar disconnections, [i.e. alkylation, Grignard, Friedel-Crafts, transition-metal catalysed coupling] (**1, 8, 10, 11, 14, 16, 20, 24, 25, 27–29, 33–38, 40**), or (iii) repurposing existing structures with mostly functional group manipulations, where no C–C bond was formed (**19, 22**). For instance, cyclobutane **1** was previously prepared using 2e[−] logic commencing from cyclobutane carboxylic acid requiring lengthy one-carbon homologation, followed by a Friedel-Crafts acylation/Wolff-Kishner sequence (9 steps total, 11% overall yield). In contrast, the 1e[−] disconnection is identified by simply selecting the most readily available building blocks and coupling them using dDCC, thereby deleting nearly all extraneous concession steps. Such a tactic is particularly valuable for the formation of seemingly distal stereocenters, as a myriad of chiral carboxylic acid building blocks can be easily purchased. In this way, expensive catalytic asymmetric methods (**23**) and chiral auxiliaries which generally require at least three more steps (installation, diastereoselective reaction and removal) can be avoided (**8, 10, 14, 22**). A particularly striking example of this concept involves the enantiodivergent synthesis of insect pheromones (*R*)- and (*S*)-**33**, previously prepared in 11–12 steps (7–8% overall yield) using a category (i) and (ii) approaches for assembling the main carbon chain (*vide supra*). In contrast, commercially available desymmetrized acid **30** can be subjected to tandem dDCC to access the same materials in 3 steps (*ca.* 30% overall yield). Either enantiomer is accessed at will simply by choosing the order of coupling. A complete comparison of conventional routes to the 32 known molecules in Table 1 versus simple pathways employing dDCC is graphically illustrated in the Supplementary Information.

It is equally important to mention current limitations of this method (see Supplementary Information) which include difficulty in coupling alkyl carboxylic acids adjacent to aromatic rings, phosphonates, and esters. The scalability of this powerful C–C bond forming reaction was demonstrated using compound **7**, which proceeded in 54% yield on a gram scale. In certain cases, if both carboxylic acid components are valuable, a 1:1 ratio can be employed to deliver a coupling product with lower yet synthetically useful yield as demonstrated with compound **7**. If one of the acids is 2°-acyclic or β,β-*gem*-disubstituted acid, the amount of the coupling partner acid can be reduced to 1.5 equivalent without substantial loss of the heterocoupling yield. This is due to the sterically hindered nature of the radicals derived from these acids, which renders the formation of heterocoupling product preferentially. Moreover, 20 mol% catalyst loading can be reduced to 8 mol% with the addition of AcOH while maintaining a similar catalytic efficiency, as demonstrated in the case of **1, 7, 10** and **24** in Table 1C (Supplementary Information includes another 4 examples).

To confirm the radical nature of this reaction, a radical clock experiment was conducted (Figure 2a). Submission of RAEs **41** and **42** to standard conditions led to a mixture of heterocoupled products **43** and **44** resulting from immediate cyclopropane opening and coupling with 5-hexenyl radical either in cyclized or open form, respectively. This is consistent with the common reactivity that RAEs generate alkyl radicals via single electron transfer (SET) followed by rapid fragmentation. Since the coupling reaction barely proceeded when Ni catalyst was omitted (Table 1B, control experiments), the role of the Ni catalyst is presumably either mediating the SET^{23,39} (See Supplementary Information for additional study by cyclic voltammetry) or effectively capturing free radicals generated by direct cathodic reduction to facilitate productive reaction pathway. Furthermore, the ratio of **43** and **44** is dependent on Ni catalyst concentration, indicating that cage-escaped radical might be involved for Ni–C bond formation. In addition, such observation is diagnostic for involvement of multiple Ni species in the mechanism, and could exclude possibility of double oxidative addition on a single Ni species (or cage-rebound mechanism)^{40–42} Although detailed mechanistic understanding is outside the scope of this initial report,

future in-depth kinetic studies may provide further support for this hypothesis.

In principle, dDCC should be amenable by other means of single-electron reduction. Thus, photoinduced electron transfer was interrogated for the same reactivity (Figure 2b, entries 2–4). Based on the results of three different experiments, this process appears to be exclusively workable under electrochemical conditions, as no observable product or only traces were observed. In one case, exclusive homodimerization of the 1° RAE **45** was observed when $i\text{Pr}_2\text{NEt}$ was employed as sacrificial reductant. These results could indicate that the successful coupling requires fine balance between radical generation from two RAEs and C–C bond formation processes catalyzed by Ni species.

The simultaneous radical generation from two RAEs under electrochemical conditions was further studied by changing a redox-active motif in a starting material. RAE **2** and RAE **3** were synthesized based on the expectation that extended π -system could have an influence on SET and following radical generation. As illustrated in Figure 2c, amongst several RAE/halide combinations explored, NHPI-based RAEs delivered the best yields of heterocoupled products (closely matching the statistically predicted outcome which would be 3:1, assuming 100% mass balance). Alkyl halide precursors are not as versatile as NHPI-based RAEs due to lower reactivity for bromides and slightly higher homocoupling tendency for iodides.

Finally, the role of ligand was further explored (Figure 2d, see Supplementary Information for the complete survey). The product distribution in the coupling of **50** and **51** was analyzed in detail. It was found that ligand structure has an impact on i) conversion of RAEs, ii) radical capture efficiency and iii) side-product formation. By switching from bidentate ligand **L6** to tridentate ligand **L1**, the reactivity of the Ni complex notably increased to realize efficient radical generation from RAEs. A change of ligand skeleton to pybox **L2** further increased the reactivity; more notably, the efficiency of capturing 2°-carbon radical was enhanced based on the analysis of the mass balance of **51**. Yet, **L2** furnished considerable amount of alkane side-product **55**, which did not contribute to the productive coupling reaction. Subtle electronic modulation of **L2** to **L4** led to the suppression of the alkane formation, delivering the heterocoupling product **52** efficiently. These qualitative trends found in this work warrants further rigorous and systematic studies to elucidate the role of a ligand, possibly contributing deeper mechanistic understanding.

With these results taken collectively, the overall reaction mechanism is depicted in Figure 2e. Electrochemical reduction of Ni(II) generates low valent Ni species active for reducing RAEs, efficiently supplying alkyl radicals (step A). Direct cathodic reduction of RAEs is also a possible source of alkyl radicals. Then, the alkyl radicals combine with Ni species consecutively (step B and C). During these two Ni–C bond formation, adjustment of Ni oxidation state via chemical (disproportionation or comproportionation) or electrochemical pathway might be involved. Finally, reductive elimination forges a new C–C bond, closing the catalytic cycle.

New methods for the construction of C–C linkages in a modular way from ubiquitous building blocks can immediately simplify the logic of chemical synthesis. Polar bond analysis is routinely taught at the undergraduate level to help guide chemists to make strategic disconnections. Radical retrosynthesis¹³ effectively divorces polarity from the analysis and prioritizes simplicity and building block availability. The dDCC method reported herein has been applied to 32 arbitrarily chosen, previously synthesized substrates containing a myriad of functional groups. The simple one-step procedure and wide availability of starting materials makes this protocol promising despite the near-statistical homo/heterocoupling ratio in many cases. Finally, the dramatic reduction in documented step-count and labor relative to the prior art bodes well for its immediate adoption for a range of organic molecules.

Online content

Any methods, additional references, Nature Research reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41586-022-04691-4>.

- Goldfogel, M. J., Huang, L. & Weix, D. J. Cross-Electrophile Coupling: Principles and New Reactions, chapter 9 in *Nickel Catalysis in Organic Synthesis: Methods and Reactions*, 183–222 (2019).
- Chatterjee, A. K., Choi, T.-L., Sanders, D. P. & Grubbs, R. H. A General Model for Selectivity in Olefin Cross Metathesis. *J. Am. Chem. Soc.* **125**, 11360–11370 (2003).
- Schäfer, H. J. Anodic and cathodic CC-bond formation. *Angew. Chem. Int. Ed.* **20**, 911–934 (1981).
- Wiebe, A. et al. Electrifying organic synthesis. *Angew. Chem. Int. Ed.* **57**, 5594–5619 (2018).
- Schäfer, H.-J. Electrochemistry IV. *Top. Curr. Chem.* 91–151 (Springer 2005).
- Svadkovskaya, G. E. & Voitkevich, S. A. Electrolytic condensation of carboxylic acids. *Russ. Chem. Rev.* **29**, 161–180 (1960).
- Leech, M. C. & Lam, K. Electrosynthesis using carboxylic acid derivatives: new tricks for old reactions. *Acc. Chem. Res.* **53**, 121–134 (2020).
- Chen, N., Ye, Z. & Zhang, F. Recent progress on electrochemical synthesis involving carboxylic acids. *Org. Biomol. Chem.* **19**, 5501–5520 (2021).
- Schäfer, H. J. Recent synthetic applications of the Kolbe electrolysis. *Chem. Phys. Lipids* **24**, 321–333 (1979).
- Moeller, K. D. Synthetic applications of anodic electrochemistry. *Tetrahedron* **56**, 9527–9554 (2000).
- Utley, J. Trends in organic electrosynthesis. *Chem. Soc. Rev.* **26**, 157–167 (1997).
- Yan, M., Kawamata, Y. & Baran, P. S. Synthetic organic electrochemical methods since 2000: on the verge of a renaissance. *Chem. Rev.* **117**, 13230–13319 (2017).
- Smith, J. M., Harwood, S. J. & Baran, P. S. Radical retrosynthesis. *Acc. Chem. Res.* **51**, 1807–1817 (2018).
- Corey, E. J. & Sauers, R. R. The synthesis of pentacyclosqualene (8,8'-Cycloöcnocrene) and the α - and β -onocadienes. *J. Am. Chem. Soc.* **81**, 1739–1743 (1959).
- Ho, C. K., McAuley, K. B. & Peppley, B. A. Biolubricants through renewable hydrocarbons: A perspective for new opportunities. *Renew. Sustain. Energy Rev.* **113**, 109261 (2019).
- Schäfer, H. J., Harenbrock, M., Klocke, E., Plate, M. & Weiper-Idelmann, A. Electrolysis for the benign conversion of renewable feedstocks. *Pure. Appl. Chem.* **79**, 2047–2057 (2007).
- Brown, D. G. & Boström, J. Analysis of past and present synthetic methodologies on medicinal chemistry: where have all the new reactions gone? *J. Med. Chem.* **59**, 4443–4458 (2016).
- Becking, L. & Schäfer, H. J. Synthesis of a prostaglandin precursor by mixed Kolbe electrolysis of 3-(cyclopent-2-enyloxy)propionate. *Tetrahedron Lett.* **29**, 2801–2802 (1988).
- Weiguny, J. & Schäfer, H. J. Electroorganic synthesis, 57. synthesis of advanced prostaglandin precursors by Kolbe electrolysis, II. – preparation of coacids and anodic initiated tandem radical-addition / radical-coupling reaction with (1R,4S,3R/S)-3-(cis-4-acetoxycyclopent-2-enyloxy)-3-ethoxypropionic acid. *Liebigs Ann. Chem.* 235–242 (1994).
- Schierle, K., Hopke, J., Niedt, M.-L., Boland, W. & Steckhan, E. Homologues of dihydro-12-oxo-phytodienoic acid and jasmonic acid by mixed Kolbe electrolysis. *Tetrahedron Lett.* **37**, 8715–8718 (1996).
- Harenbrock, M., Matzeit, A. & Schäfer, H. J. Anodic heterocoupling (mixed Kolbe electrolysis) of carbohydrate carboxylic acids with alkanolic acids to C-glycosides. *Liebigs Ann.* 55–62 (1996).
- Quertenmont, M., Goodall, I., Lam, K., Markó, I. & Riant, O. Kolbe anodic decarboxylation as a green way to access 2-pyrrolidinones. *Org. Lett.* **22**, 1771–1775 (2020).
- Parida, S. K. et al. Single electron transfer-induced redox processes involving N-(Acyloxy)phthalimides. *ACS Catal.* **11**, 1640–1683 (2021).
- Murarka, S. N. Acyloxyphthalimides as redox-active esters in cross-coupling reactions. *Adv. Synth. Catal.* **360**, 1735–1753 (2018).
- Niu, P., Li, J., Zhang, Y. & Huo, C. One-electron reduction of redox-active esters to generate carbon-centered radicals. *Eur. J. Org. Chem.* **36**, 5801–5814 (2020).
- Chan, L. K. M., Gemmell, P. A., Gray, G. W., Lacey, D. & Toyne, K. J. Synthesis and liquid crystal properties of dimethylene linked compounds incorporating the cyclobutane or spiro[3.3]heptane rings. *Mol. Cryst. Liq. Cryst. Incorporating Nonlinear Opt.* **168**, 229–245 (1989).
- Jones, P. et al. A novel series of potent and selective ketone histone deacetylase inhibitors with antitumor activity in vivo. *J. Med. Chem.* **51**, 2350–2353 (2008).
- Xu, H., Zhao, C., Qian, Q., Deng, W. & Gong, H. Nickel-catalyzed cross-coupling of unactivated alkyl halides using bis(pinacolato)diboron as reductant. *Chem Sci* **4**, 4022–4029 (2013).
- Yu, X., Yang, T., Wang, S., Xu, H. & Gong, H. et al. Nickel-catalyzed reductive cross-coupling of unactivated alkyl halides. *Org. Lett.* **13**, 2138–2141 (2011).
- Hu, X. Nickel-catalyzed cross coupling of non-activated alkyl halides: A mechanistic perspective. *Chem. Sci.* **2**, 1867–1886 (2011).
- Goldup, S. M., Leigh, D. A., McBurney, R. T., McGonigal, P. R. & Plant, A. Ligand-assisted nickel-catalysed $\text{sp}^3\text{-sp}^3$ homocoupling of unactivated alkyl bromides and its application to the active template synthesis of rotaxanes. *Chem. Sci.* **1**, 383–386 (2010).
- Smith, R. T. et al. Metallaphotoredox-catalyzed cross-electrophile $\text{Csp}^3\text{-Csp}^3$ coupling of aliphatic bromides. *J. Am. Chem. Soc.* **140**, 17433–17438 (2018).

33. Liang, Z., Xue, W., Lin, K. & Gong, H. Nickel-catalyzed reductive methylation of alkyl halides and acid chlorides with methyl p-tosylate. *Org. Lett.* **16**, 5620–5623 (2014).
34. Yu, X. et al. Nickel-catalyzed reductive cross-coupling of unactivated alkyl halides. *Org. Lett.* **13**, 2138–2141 (2011).
35. Kim, S., Goldfogel, M. J., Gilbert, M. M. & Weix, D. J. Nickel-catalyzed cross-electrophile coupling of aryl chlorides with primary alkyl chlorides. *J. Am. Chem. Soc.* **142**, 9902–9907 (2020).
36. Prinsell, M. R., Everson, D. A. & Weix, D. J. Nickel-catalyzed, sodium iodide-promoted reductive dimerization of alkyl halides, alkyl pseudohalides, and allylic acetates. *Chem. Commun.* **46**, 5743–5745 (2010).
37. Yang, Z.-P., Freas, D. J. & Fu, G. C. Asymmetric synthesis of protected unnatural α -amino acids via enantioconvergent nickel-catalyzed cross-coupling. *J. Am. Chem. Soc.* **143**, 8614–8618 (2021).
38. Gaich, T. & Baran, P. S. Aiming for the ideal synthesis. *J. Org. Chem.* **75**, 4657–4673 (2010).
39. Novaes, L. F. T. et al. Electrocatalysis as an enabling technology for organic synthesis. *Chem. Soc. Rev.* **50**, 7941–8002 (2021).
40. Breitenfeld, J., Ruiz, J., Wodrich, M. D. & Hu, X. Bimetallic oxidative addition involving radical intermediates in nickel-catalyzed alkyl–alkyl Kumada coupling reactions. *J. Am. Chem. Soc.* **135**, 12004–12012 (2013).
41. Weix, D. J. Methods and mechanisms for cross-electrophile coupling of Csp^2 halides with alkyl electrophiles. *Acc. Chem. Res.* **48**, 1767–1775 (2015).
42. Zhao, C., Jia, X., Wang, X. & Gong, H. Ni-catalyzed reductive coupling of alkyl acids with unactivated tertiary alkyl and glycosyl halides. *J. Am. Chem. Soc.* **136**, 17645–17651 (2014).

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© The Author(s), under exclusive licence to Springer Nature Limited 2022




Fig. 1 | Kolbe heterocoupling simplifies synthesis. (A) Reductive approach is the key to improving generality of Kolbe heterocoupling. ^aLimited to acids that generate stabilized radicals. (B) 1e⁻ logic simplifies retrosynthetic analysis and increases ideality.




Fig. 2 | Control studies and ligand analysis. (A) Radical clock experiments to probe the intermediacy of radical species. (B) Comparison with photochemical reactions. (C) Effect of redox active motifs on reactivity and product

distribution. (D) Impact of ligand structure on reactivity and coupling efficiency. (E) Overview of proposed reaction mechanism.




Fig. 3 | Reaction Detail. (A) General reaction conditions. (B) Effect of reaction parameters. (C) Reaction generality. All reactions were performed under the general conditions. See Supplementary Information for further reaction detail.

^a **L1** was used as a ligand instead of **L4**. ^b 2 equiv of AcOH was added with 8 mol% of Ni and the ligand. DIC: diisopropylcarbodiimide, NHPI: N-hydroxyphthalimide, DMAP: 4-dimethylaminopyridine.

Data availability

The data that support the findings in this work are available within the paper and Supplementary Information.

Acknowledgements We are grateful to Dr. D.-H. Huang and Dr. L. Pasternack (Scripps Research) for NMR spectroscopic assistance. Financial support for this work was provided by National Science Foundation Center for Synthetic Organic Electrochemistry (CHE-2002158), and the National Institutes of Health (grant number GM-118176).

Author contributions Conceptualization: BZ, YK, PSB; Experimental investigation: BZ, YG, YH, MSO, JXQ, KXR, HJZ, YK; Data analysis: BZ, YG, YH, MSO, JXQ, KXR, HJZ, YK, PSB; Manuscript

writing: BZ, YG, YK, PSB; Funding acquisition: PSB; Project administration: YK, PSB; Supervision: PSB

Competing interests The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41586-022-04691-4>.

Correspondence and requests for materials should be addressed to Phil S. Baran.

Peer review information *Nature* thanks Kevin Lam, Scott Bagley and the other, anonymous reviewers for their contribution to the peer review of this work.

Reprints and permissions information is available at <http://www.nature.com/reprints>.

ACCELERATED ARTICLE PREVIEW