

THEMES

Mitochondrial Biology in Health, Aging, and Disease

Mitochondrial transplantation in cardiomyocytes: foundation, methods, and outcomes

Paria Ali Pour,^{1,2} Sina Hosseinian,^{1,3} and Arash Kheradvar^{1,2,3}

¹Edwards Lifesciences Center for Advanced Cardiovascular Technology, Irvine, California; ²Department of Biomedical Engineering, University of California, Irvine, California; and ³School of Medicine, University of California, Irvine, California

Abstract

Mitochondrial transplantation is emerging as a novel cellular biotherapy to alleviate mitochondrial damage and dysfunction. Mitochondria play a crucial role in establishing cellular homeostasis and providing cell with the energy necessary to accomplish its function. Owing to its endosymbiotic origin, mitochondria share many features with their bacterial ancestors. Unlike the nuclear DNA, which is packaged into nucleosomes and protected from adverse environmental effects, mitochondrial DNA are more prone to harsh environmental effects, in particular that of the reactive oxygen species. Mitochondrial damage and dysfunction are implicated in many diseases ranging from metabolic diseases to cardiovascular and neurodegenerative diseases, among others. While it was once thought that transplantation of mitochondria would not be possible due to their semiautonomous nature and reliance on the nucleus, recent advances have shown that it is possible to transplant viable functional intact mitochondria from autologous, allogenic, and xenogeneic sources into different cell types. Moreover, current research suggests that the transplantation could positively modulate bioenergetics and improve disease outcome. Mitochondrial transplantation techniques and consequences of transplantation in cardiomyocytes are the theme of this review. We outline the different mitochondrial isolation and transfer techniques. Finally, we detail the consequences of mitochondrial transplantation in the cardiovascular system, more specifically in the context of cardiomyopathies and ischemia.

ischemia reperfusion injury; mitochondrial diseases; mitochondrial transplantation; mitochondrial transfer; mitochondrial cardiomyopathy

INTRODUCTION

Providing adequate contractile force to propel blood throughout the body is the myocardium's primary function, which requires energy, primarily in the form of ATP supplemented by the mitochondria. Chemiosmotic hypothesis has long established that electron transport system (ETS) creates a proton gradient across the mitochondrion's inner membrane, which is used to drive ATP synthesis (1, 2). Before this, most were convinced of the role of ETS in ATP synthesis with a provision that it is achieved based on a direct relationship through substrate-level phosphorylation (3–5). Reactive oxygen species (ROS) at complexes I and III are the by-products of ETS, which can contribute to mitochondrial damage when in excess, and are part of the signaling sequence leading to apoptosis (6). In cardiomyocytes, mitochondria are also responsible for regulation of calcium homeostasis to avoid irreversible ischemic heart damage due to cellular calcium overload (7). Given the mitochondria's various functions in cardiomyocytes, together with cardiomyocytes' heavy energy demand, mitochondrial damage or dysfunction is associated

with a variety of heart diseases. Mitochondrial dysfunction is commonly observed in cardiomyopathies, which can be due to ischemic damage or genetic mutations (8).

Proper cardiac function depends on collective cellular bioenergetics and thus mitochondrial quality. Myocardium is the highest energy-demanding tissue in the body (9), with mitochondria accounting for up to one third of the total cell volume (10). Cardiomyocytes require high energy to sustain their cellular function, and their demands increase during exercise or stress. The major source of energy for cardiomyocytes are carbohydrates in the fed state and fatty acids in the fasted state (11). Fatty acid beta oxidation is less energy-efficient than glucose; previous studies have shown that supra-physiological free fatty acid (FFA) levels observed in heart failure or myocardial infarction are due to high levels of nor-epinephrine (11).

A cellular byproduct of oxidative phosphorylation is reactive oxygen species (ROS). Normal levels of ROS are crucial for cellular communication, mediating different signaling pathways that are ultimately responsible for maintaining homeostasis and adaptation to physiological stress (12). Hence,

it is the amount of ROS that determines whether a beneficial or harmful role is played. ROS play a role in oxidative modification of receptors, kinases, phosphatases, caspases, ion channels, and transcription factors. In addition, they are implicated in signaling cascades that are important in cell fate, such as the protein kinase B (Akt), AMP-activate protein kinase (AMPK), and mitogen-activated protein kinase (MAPK) pathways (13–15). Other recent studies have demonstrated the role of ROS signaling in a variety of other signaling pathways, activating p53, c-jun NH₂-terminal kinase 1, and NF- κ B (16–18). Under physiological hypoxia, mitochondria release ROS, which subsequently activate hypoxia inducible factor 1, a crucial mediator for metabolic adaptation under low oxygen circumstances (19). ROS are also important in regulating synaptic plasticity-related signaling molecules, receptors, and channels and critical for long-term potentiation, which is beyond the scope of the current review. Large amounts of reactive oxygen, nitrogen, and lipid species can lead to mitochondrial (mt)DNA damage and irreversible impairment of mitochondria, reduced ATP generation, and eventually cell death. ROS-mediated activation of NHE-1 and CaMKII in cardiac hypertrophy and heart failure, respectively, and ROS burst in ischemia-reperfusion injury are some examples of ROS involvement in the context of cardiovascular pathologies (13).

Mitochondria also play important roles in calcium homeostasis. Calcium influx into the mitochondria is primarily regulated by mitochondria calcium uniporter. Ryanodine receptor 1 and leucine zipper-EF-hand-containing transmembrane protein 1 (LETM1) are other mechanisms that have been proposed for calcium entry into mitochondria. A remaining question is whether LETM1 is in fact a Ca²⁺/H⁺ exchanger or a K⁺/H⁺ exchanger (20). Calcium exits mitochondria through Na⁺/Ca²⁺/Li⁺ exchanger (20). The intracellular sodium level is maintained below the cytosolic sodium by sodium-proton exchangers to create the electrochemical gradient necessary for the influx of sodium that is coupled to the efflux of calcium (20).

Calcium plays a regulatory role in mitochondria and modulates the activity of mitochondrial dehydrogenases such as pyruvate dehydrogenase, isocitrate dehydrogenase, and α -ketoglutarate dehydrogenase. Moreover, calcium leads to activation of complex V of the ETS, which converts the proton motive force to ATP. As such, an increase in work, leads to increased cytosolic and subsequently matrix calcium that in turn leads to increased ATP production (20). While under increased work demand, increased mitochondrial calcium leads to increased ATP production; it is also calcium overload that triggers cellular necrosis under ischemia-reperfusion injury (21). Furthermore, it has been shown that in heart failure an increase in cytosolic sodium is accompanied by increased matrix calcium, which leads to calcium efflux from mitochondria and reduced ATP production (22).

Given that the dysfunctional mitochondria have a devastating role in the pathophysiology of variety of diseases, mitochondrial transplantation may offer exciting therapeutic applications. Although research on mitochondrial transplantation is currently dedicated predominantly to treatment of ischemic diseases in the context of cardiology and neurology, mitochondria play an important role in function of the kidney, gastrointestinal system, reproductive system, respiratory

system, and hematological system, which is beyond the scope of this review. Here we summarize the current research surrounding mitochondrial transplantation in the context of therapies for cardiovascular diseases.

Mitochondrial transplantation includes the process of isolation (Fig. 1), transfer, and internalization of exogenous intact mitochondria into targets cell or tissue (Fig. 2). Intracellular transplantation of these functional mitochondria has been proposed to repair damaged tissue in the brain, respiratory system, cardiovascular system, and renal system (23–27) to name a few. It is hypothesized that owing to their endosymbiotic origin, the recipient cells can uptake and integrate exogenous mitochondria and in doing so rescue cellular function and restore cellular homeostasis. Ali Pour et al. (28) and others have also shown the possibility of non-autologous/allogenic and even interspecies/xenogeneic mitochondrial transplantation (28–31).

MITOCHONDRIAL ISOLATION

Several techniques for isolation of mitochondria, based on differential centrifugation (DC) (32), density gradient high-speed ultracentrifugation (UC) (33), and magnetic cell separation (MACS) (34) have been practiced so far (Fig. 1).

Original method of isolation relies on centrifugation at differential speeds to fractionate the harvesting cell's nuclear and cytoplasmic components and purify isolated mitochondria. This is achieved by initially centrifuging the cell suspension or tissue homogenate to pellet the nuclear fraction and cellular debris, followed by centrifugation at higher speeds to extract and purify mitochondria from the cytosol fraction. Another centrifugation-based technique uses a Ficoll sucrose density gradient and exceedingly high-speed centrifugation and buoyancy to isolate mitochondria (34). However, with centrifugation techniques, it is difficult to achieve both high yield and purity. Since yield is high in DC, purity is lower compared with UC with roughly ~20% lysosomal and peroxisomal contaminants (35), and similarly, since purity is higher in UC, yield is lower.

A more recent method known as fractionated mitochondrial magnetic separation (FMMS) (34) seems to lead to a high-yield and high-purity solution. Referred to as FMMS here, MACS, which is a method typically used for separation of cell populations based on their surface antigens, has also been used in the context of mitochondrial isolation (34). In FMMS, mitochondrion is first labeled with anti-TOM22, which is a magnetically tagged antibody against the translocase of the outer mitochondrial membrane 22. After this incubation, the labeled lysate is passed through a column within a magnetic field. Subsequently, the magnetically tagged mitochondria will be trapped in the column and eluted, which leads to a higher purity isolate compared with the centrifugation techniques (34).

Another method proposed by McCully et al. (27) relies on collection of autologous tissue samples massing <0.1 g from a subject through biopsy and homogenization of extracted tissues via an automated homogenizer, followed by brief digestion and filtration to break down cellular membranes and free organelles. This method may better meet the needs of clinical settings where an interventional time below 60 min is desired. Lastly, centrifugation is used to concentrate




Figure 1. Mitochondrial isolation approaches in a glance. Current mitochondrial isolation techniques include **A**: differential centrifugation (DC). **B**: density gradient high-speed ultracentrifugation (UC). **C**: fractionated mitochondrial magnetic separation (FMMS). **D**: isolation for clinical use. Figure created with BioRender.

mitochondria, yielding $\sim 1 \times 10^9$ mitochondria from a 0.1-g mass (27). After the isolation, regardless of the used techniques, mitochondria can be immediately used for transfer and internalization (Fig. 1).

MITOCHONDRIAL TRANSFER AND INTERNALIZATION

Mitochondrial transfer or internalization as a natural process has been observed in living cells, where mitochondria from one cell can be passed on to another via tunneling nanotubes, usually under conditions of cellular distress (31, 36–39). However, in biomedical literature, mitochondrial transfer has been used interchangeably with transplantation, for artificial internalization of exogenous mitochondria. Here, we refer to exogenous delivery of mitochondria as

transplantation (38, 40–42). Mitochondrial transplantation has been attempted through direct injection of isolated mitochondria to the recipient tissue (31, 43–45), coincubation (28, 29, 31, 46–50), centrifugation (51), magnetomitotransfer (use of electromagnetic forces) (40), and photothermal nanoblade (Fig. 2) (41).

Direct injection has been used in the *in vitro* settings, animal models, and even in clinical settings where the isolated mitochondria are directly injected into the recipient cells (through microinjection) or tissue (via needles) (43–45).

Coincubation, as the name indicates, involves the incubation of the isolated mitochondria and recipient cells in the vicinity of one another to allow the recipient cells uptake mitochondria, perhaps similar to the way in which ancestral mitochondrion, once a bacterium, was taken up by the ancestral eukaryotic cell (Fig. 3 and Fig. 4).




A Direct Injection**B Co-incubation****C Centrifugation****D Electromagnetic Forces (Magnetomitotransfer)****E Photothermal (Nanoblade)**

Figure 2. Mitochondrial transplantation approaches in a glance. Mitochondrial transplantation techniques include direct injection (A), co-incubation (B), centrifugation (C), electromagnetic forces (D; known as magnetomitotransfer), and photothermal (E; nanoblade). Figure created with BioRender.

Centrifugation at 1,500 g for 5 min has been shown to enhance the delivery of exogenous mitochondria when compared with the simple co-incubation method. In this technique, once the isolated mitochondria are mixed with the recipient cells, the co-incubated sample is briefly centrifuged to enhance and accelerate the mitochondrial uptake (Fig. 3) (51).

Magnetomitotransfer is accomplished first by labeling mitochondria with anti-TOM22 and then exposing these magnetically labeled mitochondria to a magnetic plate. Magnetomitotransfer has a reported efficiency of 78–92% compared with co-incubation's efficiency of 17% (40). Additionally, this technique can increase mitochondrial internalization evidenced by a higher ratio of cells and higher density of transferred mitochondria within the first day of the recipient cell culture (40).

Finally, a photothermal nanoblade procedure has been reported to be used to transfer isolated mitochondria into somatic mammalian cells (41). In their proof-of-principle study, Wu et al. (41) showed that their nanoblade apparatus

was successful in transferring isolated mitochondria, leading to restoration of metabolic gene expression, global energetic, and metabolite profiles. Although the application of such a mechanism may be limited, their protocol using a precise apparatus has the potential to protect against the transfer of other unwanted cytosolic biomolecules such as microRNAs and abhorrent signaling molecules that can impact cellular biologic functions (41).

Nevertheless, mitochondrial transplantation studies are still in their infancy and more needs to be done to assess its efficiency and efficacy. Evaluating the efficiency of mitochondrial transplantation is challenging particularly considering discrimination and proper quantification of native and transplanted mitochondrial population. Although these organelles seem to be independent, they are in fact interconnected through a dynamic reticulated network, allowing to share their contents (31, 52). Mitochondria can assume different sizes and shapes through fission and fusion processes, and lack of fluorescently stabilized cell lines that allow continuous monitoring of native and transplanted mitochondria




Figure 3. Mitochondrial transplantation. Nonautologous (allogenic) mitochondria from skeletal muscle (rat L6; ATCC, Manassas, VA) cells were isolated using a commercially available isolation kit (ThermoFisher Scientific, Waltham, MA), labeled with MitoTracker Green FM (ThermoFisher Scientific, Waltham, MA), and transplanted to cardiomyocytes (rat H9c2; ATCC, Manassas, VA) via combination of centrifugation and coincubation. Success of mitochondrial transplantation was visualized using holotomography microscopy (Nanolive SA, Ecublens, Switzerland) 24 h posttransplantation. The data are from Kheradvar Laboratory.

make the comparison of native and transplanted population (and hence determining transplantation efficiency) challenging. Given the recent availability of human-induced pluripotent stem cells with stabilized fluorescently tagged mitochondria (mECFP and mTagRFP), in our opinion, further quantitative studies with more consistent results can be performed in the near future.

Maeda et al. (53) attempted to increase mitochondrial transplantation efficiency using TransActivator of Transcription dextran (TAT-dextran) complexes by enhancing cellular uptake of exogenous mitochondria. They suggest that TAT-dextran complexes enhance cellular uptake of exogenous mitochondria and can improve the protective effect of mitochondrial replenishment in rat cardiomyocytes against oxidative stress. When compared with coincubation, TAT-dextran transferred mitochondria showed a lower expression of apoptosis-promoting gene *Bax* to apoptosis-suppressing gene *Bcl-2*, indicating a more effective anti-apoptotic shift. TAT-dextran modification also increased internalization of exogenous mitochondria as confirmed by flow cytometric analysis ($183 \pm 18.4\%$, $P < 0.01$ vs. coincubation group) (53).

Actin-dependent endocytosis and macropinocytosis have been speculated as possible mechanisms of internalization of the transplanted mitochondria into a recipient cell. More studies are necessary to establish the mechanism or mechanisms of mitochondrial internalization. In our opinion, perhaps more than one mitochondrial internalization mechanism could be involved in internalization of exogenous mitochondria by recipient cells with potential dependency on the recipient cell/tissue type and mode of mitochondrial delivery. Pacak et al. (26) investigated blocking caveolae-mediated, clathrin-mediated

endocytosis, actin-dependent endocytosis, macropinocytosis, and tunneling nanotubes using pharmacological blockers of endocytosis pathways, i.e., methyl- β -cyclodextrin (MbCD), cytochalasin D (CytoD), 5-(*N*-ethyl-*N*-isopropyl) amiloride (EIPA), and nocodazole (Noco), respectively. Before mitochondrial transplantation, the cells were incubated with one of the above-mentioned pharmacological blockers. Mitochondrial internalization was only significantly decreased in the presence of CytoD, which suggests the involvement of actin-dependent endocytosis pathway (26). A more recent work by the same group, Cowan et al. (54) used three-dimensional superresolution microscopy and transmission electron microscopy to visualize exogenous human mitochondria posttransplantation and looked more closely at the dynamic of intracellular trafficking of mitochondria. With respect to mitochondrial internalization through endocytosis, mitochondria are taken by the cells through endocytic vesicles formed by the invagination of the cellular membrane during the cellular uptake process. These vesicles are then fused with early endosomes as a major sorting compartment. Once the endosome reaches the cells, it should ideally release its contents (i.e., mitochondria). However, endosomal escape can be a limiting step if the mitochondria get trapped inside the endosomes. If mitochondria are destined to be degraded, they will be transported to late endosomes, which




Figure 4. Native vs. transplanted mitochondrial population. Mitochondrial transplantation into cardiomyocytes visualized by structured illumination fluorescence microscopy (Keyence, Itasca, IL). Transplanted mitochondrial population labeled with pHrodo Red SE (ThermoFisher Scientific, Waltham, MA), native mitochondria labeled with MitoTracker Green FM (ThermoFisher Scientific, Waltham, MA), and cell nucleus labeled with NucBlue Live (ThermoFisher Scientific, Waltham, MA). Figure adapted from an image previously published by Ali Pour et al. (28).

together with Golgi's lysosomal compartments will form the endolysosomes that can provide the proper environmental condition for degradation and recycling of material. Yet, some cargo might be able to escape the late endosomes. In their studies, Cowan et al. (54) labeled early and late endosomes with early endosome antigen 1 and late endocytic marker Rab7, respectively. They assumed some of the transplanted mitochondria were degraded based on their fragmented appearance in some lysosomes. However, the majority of internalized green fluorescent protein-labeled mitochondria successively progressed through the endolysosomal system, successfully escaped the endosomal and lysosomal compartments, and effectively fused with the endogenous mitochondria based on their association with the cardiomyocyte mitochondrial network (54).

Although studies by Pacak et al. (26) and Cowan et al. (54) suggest actin-dependent endocytosis as the mechanism for mitochondrial internalization in cardiomyocytes, previous studies by Kitani et al. (42) and Kesner et al. (55) suggest macro-pinocytosis involvement in the context of internalization of mitochondria in human uterine endometrial cancer cells. In studies by Kesner et al. (55) using transmission electron microscopy, they observed mitochondria were engulfed by the recipient cells through cellular extensions and cell surface ruffling. This led them to hypothesize macropinocytosis involvement, which was confirmed when mitochondrial uptake was blocked upon use of micropinocytosis inhibitors except the clathrin-mediated endocytosis inhibitors.

With respect to this disagreement on the underlying mechanism of mitochondrial internalization, Patel et al. (56) argue that with the exception of the work by Kesner et al. (55), current knowledge may not suggest a conclusive and reliable decision since the studies only rely on use of endocytotic inhibitors, which can inhibit multiple endocytic pathways.

CONSEQUENCES OF MITOCHONDRIAL TRANSPLANTATION IN THE CARDIOVASCULAR SYSTEM

Mitochondrial Transplantation Improves Outcomes in Ischemia-Reperfusion Injury

Ischemia-reperfusion injury in the context of ischemic heart disease is defined as the paradoxical exacerbation of cellular dysfunction and death following restoration of blood flow to previously hypoxic tissues (57). Ischemic damage to mitochondria leads to their malfunction. Since mitochondria's DNA encodes most of their components, prolonged period of ischemia can permanently damage the ability of cardiomyocytes' mitochondria to generate ATP (58, 59).

It is currently believed that irreversible ischemic damage may be due to 1) loss of ATP (58, 59); 2) supraphysiological free fatty acids (FFA) levels (11); 3) formation of excess reactive oxygen species (ROS) (60–62); 4) cellular calcium overload (63–66); and finally, 5) sodium pump inhibition (Fig. 5) (67, 68).

In unstressed healthy individuals, although ROS are continually generated during cellular metabolism, they are effectively removed by cellular antioxidant systems. An example of this antioxidant system is superoxide dismutase

that removes superoxide radicals produced during cellular metabolism (69). Under normal circumstances, these antioxidant systems are efficient at maintaining ROS at a healthy level. However, it is during ischemia when ROS production exceeds the “buffering” capacity of the antioxidant system, leading to disruption of calcium homeostasis, mitochondrial and cellular DNA damage, and mitochondrial lipid peroxidation (70, 71). Ultimately, this leads to cell death by apoptosis through the intrinsic apoptosis pathway (60–62).

In the classical manifestation of ischemia, occlusions in the arterial supply leads to the hypoxic effects described earlier. Restoration of blood flow, or reperfusion, is frequently found to exacerbate tissue injury. In fact, reperfusion can lead to an additional 40% cardiac cell death (72). This may be due to the altered metabolism of ischemia-damaged mitochondria, leading to excessive ROS production and disruption of the mitochondria membrane (73). Reperfusion also elicits rapid alterations in ion flux, and this rapid change of pH may lead to enhanced cytotoxicity (67, 68). To return to normal pH levels, the cell utilizes sodium-dependent pH regulators, such as the Na^+/H^+ exchanger. Once activated, hydrogen is pumped out of the cell, leading to intracellular sodium accumulation. High sodium levels activate $\text{Na}^+/\text{Ca}^{2+}$ exchangers, leading to Ca^{2+} overload, which in turn causes ATP depletion and damage to mitochondria (63–66).

Metabolic and ionic changes secondary to the lack of oxygen are major contributors to the mitochondrial damage in the myocardium during ischemia, which significantly affect cardiac function and cell survival during ischemia reperfusion injuries. Masuzawa et al. (43) hypothesized and tested that in ischemia, cell survival will improve by transplantation of autologously derived mitochondria immediately before reperfusion. In their in vivo and in vitro studies in New Zealand White rabbits and Lewis rats, autologously derived mitochondria were injected into regional zones of ischemia, following 29 min of reperfusion injury, and the hearts were allowed to recover for 4 wk. Their results showed that the transplanted mitochondria are internalized by the tissue in 2 to 8 h. Moreover, cardiac enzymes, creatine kinase MB and cardiac troponin-I, apoptosis, and infarct size significantly decreased posttransplantation in the reperfusion injury zone. Additionally, within 10 min after reperfusion, normal contraction was reestablished in the transplant group, while in the control group, hypokinesia in the reperfusion injury zone persisted until 4-wk recovery timepoint (Fig. 6) (43).

Cowan et al. (46) expanded on their previous work (43) and demonstrated the effectiveness of transplantation in an intracoronary delivery model. They effectively delivered mitochondria through coronary vasculature in Langendorff-perfused rabbit hearts that were subjected to 30 min of ischemia and then reperfused for 10 min. They concluded that mitochondrial transplantation through coronary arteries was practicable and occurred within minutes. Mitochondria delivered via coronary arteries spread throughout the cardiac tissue, whereas those locally injected were predominantly localized to the interstitium of the injection site. Vascular delivery of mitochondria reduced infarct size similar to direct injection of mitochondria, which indicates a clinically relevant method, since direct injection may not be feasible in all interventional settings (46).




Figure 5. Mitochondrial dysfunction in ischemia. It is currently believed that irreversible ischemic damage is due to 1) loss of ATP, 2) supraphysiological free fatty acids levels, 3) formation of excess reactive oxygen species (ROS), 4) cellular calcium overload, and finally, 5) sodium pump inhibition. Figure created with BioRender.

Previous work has demonstrated that mitochondria from diabetic rats are dysfunctional, having reduced ATP production and respiratory capacity (74). Doulamis et al. (29) investigated whether the source of allogenic mitochondria played a part in improved clinical outcomes, by comparing mitochondria isolated from either nondiabetic or diabetic rats. Their results demonstrate that mitochondrial transplantation enhances postischemic myocardial function and reduces cardiomyocyte injury in diabetic rat hearts regardless of the source of mitochondria (Fig. 7). Both groups of diabetic rats receiving mitochondria exhibited an increase in left ventricular (LV) pressure after transplantation. In diabetic rats that received allogenic mitochondria, mean LV pressure was 92.8 ± 5.2 mmHg, 110.7 ± 2.4 mmHg, and 44.3 ± 5.9 mmHg in rats that received mitochondria from diabetic, nondiabetic, and healthy control donors, respectively. Similar improvements were observed for infarct size, with sizes of 11.8 ± 0.7 , $9.9 \pm 0.5\%$, and $52.0 \pm 1.4\%$, in rats that received mitochondria from diabetic, nondiabetic, and healthy control donors, respectively.

There was no significant difference in cardiac function or myocellular viability between mitochondria-receiving groups, suggesting that mitochondrial transplantation has great potential to reduce morbidity and mortality in diabetic patients subjected to myocardial ischemia-reperfusion injury (29).

McCully's group (29) continued their mitochondrial transplantation studies in swine models; Kaza et al. (44) performed mini-thoracotomies on Yorkshire pigs with pectoralis major dissected and removed to be used for isolation of autologous mitochondria. After 24 min of ischemia-reperfusion injury, the hearts received multiple injections of either a sham control or vehicle containing mitochondria into the area at risk. Levels of creatine kinase-MB and troponin I were significantly decreased in the treatment group compared with the sham group, which persisted for at least 4 wk posttransplantation. Immune, inflammatory, and cytokine markers showed no difference among the groups. While there was no significant difference in the area at risk between the sham control and vehicle containing mitochondria ($P = 0.48$), the infarct size was significantly reduced ($P = 0.004$) in the group receiving mitochondria. Multiple doses of mitochondria were tested, with no significant difference in decreasing the infarct size despite the number of mitochondria injected (44).

The swine model was also used to test the prophylactic effect of mitochondrial transplantation for the hearts exposed to ischemia reperfusion injury (Fig. 7). Guariento et al. (49) cannulated 26 Yorkshire pigs with sham control buffer or mitochondria, either serially, having 10 injections over 60 min, or with a single bolus. The hearts were




Figure 6. Mitochondrial dysfunction in mitochondrial diseases. Imbalance in mitochondrial dynamics (decrease in fusion, increase in fission, and decrease in mitophagy) leads to accumulation of damage/mutant mitochondria (*top*) and Clonal replication and expansion of damaged/mutant mtDNA beyond tissue threshold level for phenotype expression (*bottom*). ROS, reactive oxygen species; nDNA, nuclear DNA; mtDNA, mitochondrial DNA. Figure created with BioRender.

subjected to regional ischemia, 15 min after injection, for 30 min and were reperfused for 120 min. They found that both groups of pigs receiving mitochondria had significantly increased coronary blood flow, ejection fraction, and improved LV pressure. Like the previous findings, there was no difference in the size of the area at risk, but pigs receiving mitochondria did have a significantly smaller infarct size, suggesting mitochondrial transplantation provides prophylactic cardioprotection from ischemic reperfusion injury. Most notably, there was no significant difference between those who received a single bolus of mitochondria versus those who received serial injections, suggesting no advantages for the latter (49).

Although previous studies focus on immediate injection or cannulation of mitochondria, in a typical manifestation of ischemia reperfusion injury, patients will typically not seek treatment in the minutes or hours following an ischemic event. Blitzer et al. (48) investigated the efficacy of delayed transplantation of mitochondria for cardioprotection in a swine model by subjecting Yorkshire pigs to 30 min of ischemia, allowing for 120 min of reperfusion, and then introducing either sham control or autologous mitochondria to the coronary ostium. The

authors suggest that delayed transplantation still provides cardioprotection, as pigs that received mitochondria after the delay had a significantly enhanced ejection fraction ($P = 0.02$) and significantly lower infarct size ($P < 0.001$), which further underscores the clinical application of mitochondrial transplantation (48).

Mitochondrial transplantation may also play a role in improving the efficacy of organ transplantation. Moskowitzova et al. (50) hypothesized that mitochondrial transplantation may increase cold ischemia time in heart transplants. Cold ischemia time is defined as the time between the chilling of a tissue after its blood supply has been cut off and the time it is warmed by having blood supply restored. Moskowitzova et al. (50) suggest that this type of ischemia-reperfusion injury also benefits from mitochondrial transplantation. Pre- and posttransplantation injections of either sham control (heparin) or vehicle containing mitochondria to transplanted rat hearts with a cold ischemic time of 29 h, followed for the next 24 h, suggest that beating score, ejection fraction, and shortening fraction were significantly enhanced ($P < 0.05$). Neutrophil infiltration and necrosis were significantly decreased ($P < 0.05$), ultimately suggesting that their




Figure 7. Mitochondrial transplantation in animal models. Allogenic mitochondrial transplantation has been shown to increase cardioprotection in diabetic rat hearts (*left*), autologous mitochondria improve cardioprotection and decrease infarct size in rabbits undergone regional ischemia (*middle*), and autologous mitochondrial transplantation before ischemic damage in pigs leads to improved cardiac function and decreased infarct size, demonstrating mitochondrial transplantation's prophylactic effects (*right*). The diagrams summarize the work by Masuzawa et al. (44), Doulamis et al. (29), and Guariento et al. (50). Figure created with BioRender.

protocol can enhance graft function and reduce graft failure in transplant patients (50).

In many countries, donation after circulatory death as a measure for organ transplantation involves regulations related to the retrieval of organs for the purpose of transplantation that follows death. Warm ischemia is a serious concern to this procedure as it affects the entire organ (75). Guariento et al. (76) recently investigated whether mitochondrial transplantation can help address this concern by increasing cardiac efficiency. They had 29 Yorkshire pigs undergo circulatory death by cessation of mechanical ventilation. After 20 min of warm ischemia and 15 min of reperfusion via an ex situ blood perfusion system, the hearts received either a sham control or vehicle containing autologous mitochondria. A third group received a second injection of mitochondria after 2 h of ex situ heart perfusion. The hearts which received autologous mitochondria showed a significantly increased left ventricular pressure, fractional shortening, and myocardial oxygen consumption. Infarct size was also significantly reduced in hearts receiving mitochondria when

compared with the control group with sham treatment (76).

The question of efficacy and safety of mitochondrial transplantation remains to be addressed. Accordingly, Shin et al. (47) developed a novel method for delivering mitochondria through an intracoronary catheter. To investigate mitochondrial biodistribution, mitochondria were tagged using ^{72}F -rhodamine-6G and were visualized using positron emission tomography scan. Visualization revealed global distribution of the mitochondria. These studies also found that intracoronary injection of mitochondria at any tested concentration did not affect heart rate, mean arterial pressure, or cardiac rhythm but enhanced regional and global LV function. Intracoronary injection increased coronary blood flow in a concentration-dependent scheme without causing any detectable lesions or blockages. The authors used antidiuretic hormone (ADH) to induce myocardial stress and increase myocardial demand. Injection of ADH resulted in coronary vasoconstriction and decrease in coronary blood flow, which was rescued by intracoronary injection of mitochondria. Ultimately, this study reveals the safety and efficacy of

mitochondrial transplantation via coronary catheters. This procedure of bolus delivery makes mitochondrial transplantation more applicable for use in clinical settings (47).

The first clinical study of mitochondrial transplantation recruited pediatric patients with ischemia-reperfusion injury who were on extracorporeal membrane oxygenation (ECMO) support (77). Autologous mitochondria from nonischemic skeletal muscle were isolated, suspended in 1 mL respiration buffer, and administered to the affected myocardium through direct epicardial injection with a tuberculin syringe, and monitored by epicardial echocardiography to assess hypokinesis. Following autologous mitochondrial transplantation, four in five patients were successfully removed from ECMO support, although ultimately only three survived. Additionally, the autopsy of patients who could not be removed from ECMO indicated no difference in pre- and postinjection markers of systemic inflammatory syndrome suggesting no inflammation or rejection due to mitochondrial treatment (77).

Although this study was limited in its sample size to draw any statistically significant conclusion, it paved the way for their next pilot study examining autologous mitochondrial transplantation in pediatric patients (45). In this retrospective study of patients requiring ECMO, 14 patients underwent revascularization alone and 10 underwent revascularization with subsequent mitochondrial transplantation. While 80% of the patients who received mitochondrial transplantation were removed from ECMO within 1 wk, only 29% of the patients in the revascularization alone group became ECMO free ($P = 0.02$) (45). Ventricular strain was significantly lower in the mitochondrial recipient group. Median functional recovery time after revascularization was also significantly shorter with fewer adverse cardiovascular events (45). Ultimately, this study suggested that mitochondrial transplantation is associated with successful separation from ECMO and better cardiac function overall (45).

Mitochondrial Transplantation Improves Cellular Bioenergetics in Healthy Cardiomyocytes

One of the many aims of mitochondrial transplantation is to modulate cellular bioenergetics, and in turn function, in the recipient cells. Recent studies by Ali Pour et al. (28) demonstrate short and long term (up to 28 days) posttransplantation bioenergetics in healthy cardiomyocytes. They show that cellular bioenergetics *in vitro* is acutely improved posttransplantation; however, this enhancement is short lived and the improved bioenergetic indexes (basal respiration and ATP production) return to normal biological levels in long-term. This observation suggests that mitochondrial transplantation may boost healthy cellular energetics, which can be clinically useful in mitigating acute myocardial stress (28). However, more research is needed to investigate potential and efficacy of mitochondrial transplantation in treatment of more chronic conditions.

Mitochondrial Diseases and Dysfunction

Mitochondrial cardiomyopathy is characterized by abnormal myocardium secondary to genetic defects that leads to impairment or loss of the factors in the mitochondrial

respiratory chain, in the absence of other cardiac diseases (Fig. 6) (8). Cardiomyopathies are some of the most frequent manifestations of mitochondrial diseases, occurring in 20–40% of children with mitochondrial disease (78, 79). Clinical manifestations of mitochondrial cardiomyopathies can vary in severity from asymptomatic to severe symptoms, most notably heart failure and arrhythmias. Mortality in children with mitochondrial diseases is higher in those with cardiomyopathy than in those without (79).

It is speculated that the etiology of mitochondrial cardiomyopathy lies in the impairment or loss of the electron transport chain complex subunits and CoQ₁₀ synthesis (8). Cardiomyopathy has been reported in patients with mutations in mitochondrial and nuclear genes encoding complex I subunits, such as MTND1 and MTND5 (80). Mutations in genes encoding for complex II subunits, such as SDHA and SDHD, have also been reported in patients with cardiomyopathies (81).

The prevalence of mitochondrial diseases is much higher than previously thought. It is currently estimated that mitochondrial diseases occur in 1 in 4,000 individuals, which could still be an underestimate due to misdiagnosis (82). Presently, no clinically proven therapy is available for mitochondrial diseases, which highlights the unmet clinical needs.

Potential therapies envisioned are broadly divided into gene-based and cellular therapies. Cell-based therapies are established on the transplantation or delivery of intact new organelles from healthy cells to diseased cells, which may potentially improve outcome for patients who suffer from the consequences of acquired or congenital mitochondrial defects merely due to mitochondrial mutations. This can be achieved through intercellular transfer of exogenous mitochondria, potentially from allogenic sources. Several studies have also shown the efficacy of intercellular transfer from stem cells to diseased cells (36, 83–86), and transfer of exogenous mitochondria via transplantation is currently being investigated (31, 87).

Mitochondrial genome, similar to the nuclear genome, is constantly exposed to external and internal agents that can result in damage and mutation (88). Mitochondrial diseases develop as a result of the clonal expansion of the mutated/damaged mtDNA (Fig. 6). Different tissues have different thresholds for mtDNA mutation load that can lead to cellular mitochondrial dysfunction, with the threshold being lower for cells that depend heavily on the energy supplied by mitochondria, such as cardiac cells (89). Since mitochondria play a critical role in the cell, the maintenance of mtDNA integrity is vital to the well-being of the cell. Thus mitochondria constantly undergo fission and fusion processes to maintain their fitness (90).

mtDNA is packaged into DNA-protein assemblies called nucleoids (91). Fusion as the name implies is the process where two mitochondria join and exchange their nucleoids (92, 93). Fission is the reverse process, in which there is an increase in number of mitochondria (92, 93). Fusion is believed to have a protective role in pathologies caused by mtDNA mutations (92). It has been shown that fusion allows for complementation effect where two nucleoids, each with different mutations in the same mitochondria, display the wild-type phenotype (94, 95). However, when the mutation

is severe, it would be best to isolate and remove the mutation through fission, in which the mutant nucleoids are segregated into a mitochondrion (or few mitochondria) that can be marked for degradation by the mitochondrial autophagy, commonly referred to as mitophagy (93).

With the accumulation of deleterious mutation, nucleoid replication also increases by the retrograde signaling to produce more nucleoids so that the mitochondria would be able to meet the demand of the cell, which is another mechanism mitochondrion employs in maintaining its proper function and providing the cell with energy (96).

Although mitochondria lack a true DNA repair mechanism, they use fission and fusion processes to maintain a healthy network of mitochondria and this characteristic is the reason why we believe mitochondrial transplantation may have the potential to serve as a cellular biotherapy in the context of mitochondrial diseases due to mtDNA mutations. If the transplanted mitochondria are adopted by the recipient cells for long term, transplantation of healthy mitochondria can dilute the mutation load. Moreover, with an effective treatment plan (appropriate mitochondrial dose and frequency of treatment), which can be established using mathematical modeling of mitochondrial dynamics in conjunction with experimental verification, the mutation load can be modulated to reduce it to lower levels than those which lead to clinical manifestations. Diseases that involve mitochondrial defects or mt-nucleus signaling defects, but not the ones with nucleus involvement, will benefit from such therapies.

CHALLENGES OF MITOCHONDRIAL TRANSPLANTATION

Although the studies summarized above show promising results in animal models, there are still many unknowns and challenges around mitochondrial transplantation, which need to be addressed. This could potentially also explain the reason behind the sparse number of clinical studies involving mitochondrial transplantation.

One such challenge lies in the source of the mitochondria being used. Given that most studies utilize autologous mitochondria, this greatly limits the clinical use of mitochondrial transplantation. In our opinion, fibroblasts may be the best source in terms of ease of access for mitochondrial transplantation for various purposes. However, the number of mitochondria in fibroblasts is not comparable to myocytes or cardiomyocytes. Platelets could potentially serve as another source; while platelets have a limited number of mitochondria, they are widely available. However, for more specialized cells such as those in the muscles, retina or cochlea, the mitochondria source may be restricted to the same cell type. The clinical usage of mitochondrial transplantation is also limited due to necessity and challenges of mitochondrial harvest. In clinical studies thus far, autologous mitochondria from either the abdominal rectus muscle or the pectoralis major muscles have been used. It has been reported that the activity of isolated mitochondria greatly decreases when stored on ice for more than 1 h (27). This short timeframe demands a rapid procedure, which may not be ideal for complex surgeries where timing is a crucial deterministic factor.

However, if mitochondria could instead be used from a stored source instead of one that is needed to be rapidly extracted and prepared, clinical applications would be greatly broadened.

Another challenge for mitochondrial transplantation revolves around mtDNA haplotypes. mtDNA haplotypes are specific regions of mitochondrial DNA that cluster with other mitochondrial sequences and show origins of maternal lineages. mtDNA haplotypes are often associated with a variety of phenotypes and disease. Bevers et al. (97) outline that mitochondrial haplotypes have no direct impact on mitochondrial respiration in *drosophila* but may be associated with metabolism-related phenotypes. Through swapping of mitochondrial genomes, they demonstrated that a mitochondrial haplotype associated with high-food intake can rescue a low food intake phenotype, having drastic phenotypic implications (97).

mtDNA haplotypes have been shown to influence mtDNA replication. Liu et al. (98) report that in swine models, different mitochondrial haplotypes are associated with different numbers of mtDNA copies. They also show that haplotypes with higher mtDNA copy numbers have an increased transcript abundance of mitochondrial-encoded subunits ND1, DN6 and ATP6. Ultimately, their data suggest that variations in mitochondrial haplotypes may affect mitochondrial content (98). This is corroborated by findings from Lee et al. (99). They demonstrate that mtDNA haplotypes influence changes in gene expression and affinity for mtDNA replication factors to modulate mtDNA copy number. They also show that different haplotypes induce distinct DNA methylation patterns, which skew genetic expression patterns (99).

It is evident that mitochondrial haplotypes may have drastic effects on genomic expression, and ultimately, phenotype. To our knowledge, we are not aware of any studies that examines the effects of different mitochondrial haplotypes in the context of mitochondrial transplantation. Such studies are needed in the field, as transplanted mitochondria will conserve their DNA haplotype and may affect respiratory capacity and metabolism.

Although sometimes practical, allogenic mitochondrial transplantation poses challenges of its own, particularly that it may mediate an autoimmune response in the host. The studies by McCully's group (43, 44, 47) examined the role of immune system in response to mitochondrial transplantation, and several of their studies in animal models suggest that transplantation of autologous or allogenic mitochondria does not result in a significant alloreactivity or an increased damage-associated molecular pattern (DAMPs) response. Their results did not suggest an increase in cytokine or chemokine levels, which are crucial mediators of immune responses (43, 44, 47).

Conversely, other studies have shown that mitochondria may induce an inflammatory response. In fact, some DAMPs are associated with the mitochondria, likely secondary to its bacterial origin (100–103), mtDNA, cytochrome c, and the mitochondrial enzyme carbamoyl phosphate synthase have all been found to induce inflammation through DAMPs (100–103). Lin et al. (104) demonstrated the ability of extracellular mitochondria to activate endothelial cells, resulting in an upregulation of endothelial cell adhesion molecules and the production of cytokines and chemokines. They also

showed that the administration of allogenic mitochondria increased rejection of a xenograft heart transplant in a mouse model (104). Pollara et al. (105) reported that mitochondria-associated DAMPs were abundant in the systemic circulation of post mortem organ transplant donors. It was also concomitantly reported that proinflammatory cytokines and chemokines were elevated (105). In the case of necrosis through direct tissue trauma, Zhang et al. (106, 107) reported that mitochondrial particles released during cellular necrosis can activate neutrophils, having immunostimulatory effects. These reports could pose a challenge to the therapeutic potential of mitochondria transplantation. More studies in the field are needed to examine the alloreactivity of allogenic mitochondria. Of particular use would be additional studies examining the mechanisms responsible for mitochondrial-mediated immune responses.

CONCLUSIONS

The studies summarized here collectively demonstrate the success and potential of mitochondrial transplantation in ameliorating mitochondrial function and ultimately in improving cardiac function. Despite advances in the field, challenges remain; the conceptual challenge is to better understand how mitochondrial transplantation affects intracellular processes, bioenergetics, and signaling. Additional studies are needed to investigate the role of mtDNA haplotypes and the potential for auto-immune reactions. Establishing the best route of access and method of administering mitochondria to achieve the desired effect is of equal significance to institute mitochondrial transplantation as a viable clinical solution. Although direct injection of mitochondria is practical in surgical approaches, vascular access may serve as a more clinically appropriate mechanism of mitochondria distribution. Looking to the future, mitochondrial transplantation holds the potential to serve as a subcellular biotherapy to rejuvenate damaged or dysfunctional mitochondria and in doing so restoring homeostasis to the cells, protecting the heart tissue from damage, and, ultimately, improving cardiac function.

ACKNOWLEDGMENTS

We thank Mathew Bell from Nanolive Microscope Team for assistance with imaging studies that contributed to the image presented in Fig. 3.

GRANTS

A. Kheradvar was funded by National Science Foundation Grant 2109959.

DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the authors.

AUTHOR CONTRIBUTIONS

P.A. and A.K. conceived and designed research; P.A. performed experiments; P.A. analyzed data; P.A. interpreted results of experiments; P.A. and S.H. prepared figures; P.A., S.H. and A.K.

drafted manuscript; P.A., S.H., and A.K. edited and revised manuscript; P.A., S.H. and A.K. approved final version of manuscript.

REFERENCES

- Mitchell P. Chemiosmotic coupling in oxidative and photosynthetic phosphorylation. *Biol Rev Camb Philos Soc* 41: 445–502, 1966. doi:10.1111/j.1469-185X.1966.tb01501.x.
- Mitchell P. Coupling of phosphorylation to electron and hydrogen transfer by a chemi-osmotic type of mechanism. *Nature* 191: 144–148, 1961. doi:10.1038/191144a0.
- King TE. Reconstitution of the respiratory chain. *Adv Enzymol Relat Areas Mol Biol* 28: 155–236, 1966. doi:10.1002/9780470122730.ch3.
- Keilin D. On cytochrome, a respiratory pigment, common to animals, yeast, and higher plants. *The Roy Soc London Biolo Sci* 98: 312–339, 1925. doi:10.1098/rspb.1925.0039.
- Mitchell P. Keilin's respiratory chain concept and its chemiosmotic consequences. *Science* 206: 1148–1159, 1979. doi:10.1126/science.388618.
- Vilahur G, Juan-Babot O, Pena E, Onate B, Casani L, Badimon L. Molecular and cellular mechanisms involved in cardiac remodeling after acute myocardial infarction. *J Mol Cell Cardiol* 50: 522–533, 2011. doi:10.1016/j.yjmcc.2010.12.021.
- Lubbe WF, Podzuweit T, Opie LH. Potential arrhythmogenic role of cyclic adenosine monophosphate (AMP) and cytosolic calcium overload: implications for prophylactic effects of beta-blockers in myocardial infarction and proarrhythmic effects of phosphodiesterase inhibitors. *J Am Coll Cardiol* 19: 1622–1633, 1992. doi:10.1016/0735-1097(92)90629-2.
- Meyers DE, Basha HI, Koenig MK. Mitochondrial cardiomyopathy: pathophysiology, diagnosis, and management. *Tex Heart Inst J* 40: 385–394, 2013.
- Goffart S, von Kleist-Retzow JC, Wiesner RJ. Regulation of mitochondrial proliferation in the heart: power-plant failure contributes to cardiac failure in hypertrophy. *Cardiovasc Res* 64: 198–207, 2004. doi:10.1016/j.cardiores.2004.06.030.
- David H, Bozner A, Meyer R, Wassilew G. Pre- and postnatal development and ageing of the heart. Ultrastructural results and quantitative data. *Exp Pathol Suppl (1981)* 7: 1–176, 1981.
- Cohn JN, Levine TB, Olivari MT, Garberg V, Lura D, Francis GS, Simon AB, Rector T. Plasma norepinephrine as a guide to prognosis in patients with chronic congestive heart failure. *N Engl J Med* 311: 819–823, 1984. doi:10.1056/NEJM198409273111303.
- Collins Y, Chouchani ET, James AM, Menger KE, Cocheme HM, Murphy MP. Mitochondrial redox signalling at a glance. *J Cell Sci* 125: 801–806, 2012 [Erratum in *J Cell Sci* 2012 125: 1837, 2012]. doi:10.1242/jcs.098475.
- Zhao RZ, Jiang S, Zhang L, Yu ZB. Mitochondrial electron transport chain, ROS generation and uncoupling (Review). *Int J Mol Med* 44: 3–15, 2019. doi:10.3892/ijmm.2019.4188.
- Son Y, Cheong YK, Kim NH, Chung HT, Kang DG, Pae HO. Mitogen-activated protein kinases and reactive oxygen species: how can ROS activate MAPK pathways? *J Signal Transduct* 2011: 792639, 2011. doi:10.1155/2011/792639.
- Zhang C, Jia X, Bao J, Chen S, Wang K, Zhang Y, Li P, Wan JB, Su H, Wang Y, Mei Z, He C. Polyphyllin VII induces apoptosis in HepG2 cells through ROS-mediated mitochondrial dysfunction and MAPK pathways. *BMC Complement Altern Med* 16: 58, 2016. doi:10.1186/s12906-016-1036-x.
- Chandel NS, Vander Heiden MG, Thompson CB, Schumacker PT. Redox regulation of p53 during hypoxia. *Oncogene* 19: 3840–3848, 2000. doi:10.1038/sj.onc.1203727.
- Chandel NS, Trzyna WC, McClintock DS, Schumacker PT. Role of oxidants in NF-kappa B activation and TNF-alpha gene transcription induced by hypoxia and endotoxin. *J Immunol* 165: 1013–1021, 2000. doi:10.4049/jimmunol.165.2.1013.
- Nemoto S, Takeda K, Yu ZX, Ferrans VJ, Finkel T. Role for mitochondrial oxidants as regulators of cellular metabolism. *Mol Cell Biol* 20: 7311–7318, 2000. doi:10.1128/MCB.20.19.7311-7318.2000.
- Chandel NS, Maltepe E, Goldwasser E, Mathieu CE, Simon MC, Schumacker PT. Mitochondrial reactive oxygen species trigger hypoxia-induced transcription. *Proc Natl Acad Sci U S A* 95: 11715–11720, 1998. doi:10.1073/pnas.95.20.11715.

20. Finkel T, Menazza S, Holmstrom KM, Parks RJ, Liu J, Sun J, Liu J, Pan X, Murphy E. The ins and outs of mitochondrial calcium. *Circ Res* 116: 1810–1819, 2015. doi:10.1161/CIRCRESAHA.116.305484.
21. Murphy E, Steenbergen C. Preconditioning: the mitochondrial connection. *Annu Rev Physiol* 69: 51–67, 2007. doi:10.1146/annurev.physiol.69.031905.163645.
22. Liu T, Takimoto E, Dimaano VL, DeMazumder D, Kettlewell S, Smith G, Sidor A, Abraham TP, O'Rourke B. Inhibiting mitochondrial Na⁺/Ca²⁺ exchange prevents sudden death in a Guinea pig model of heart failure. *Circ Res* 115: 44–54, 2014. doi:10.1161/CIRCRESAHA.115.303062.
23. Jabbari H, Roushandeh AM, Rostami MK, Razavi-Toosi MT, Shokrgozar MA, Jahanian-Najafabadi A, Kuwahara Y, Roudkenar MH. Mitochondrial transplantation ameliorates ischemia/reperfusion-induced kidney injury in rat. *Biochim Biophys Acta Mol Basis Dis* 1866: 165809, 2020. doi:10.1016/j.bbadis.2020.165809.
24. Moskowitsova K, Orfany A, Liu K, Ramirez-Barbieri G, Thedsanamoorthy JK, Yao R, Guariento A, Doulamis IP, Blitzer D, Shin B, Snay ER, Inkster JA, Iken K, Packard AB, Cowan DB, Visner GA, Del Nido PJ, McCully JD. Mitochondrial transplantation enhances murine lung viability and recovery after ischemia-reperfusion injury. *Am J Physiol Lung Cell Mol Physiol* 318: L78–L88, 2020. doi:10.1152/ajplung.00221.2019.
25. Zhang Z, Ma Z, Yan C, Pu K, Wu M, Bai J, Li Y, Wang Q. Muscle-derived autologous mitochondrial transplantation: A novel strategy for treating cerebral ischemic injury. *Behav Brain Res* 356: 322–331, 2019. doi:10.1016/j.bbr.2018.09.005.
26. Pacak CA, Preble JM, Kondo H, Seibel P, Levitsky S, Del Nido PJ, Cowan DB, McCully JD. Actin-dependent mitochondrial internalization in cardiomyocytes: evidence for rescue of mitochondrial function. *Biol Open* 4: 622–626, 2015. doi:10.1242/bio.20151478.
27. McCully JD, Cowan DB, Emani SM, Del Nido PJ. Mitochondrial transplantation: from animal models to clinical use in humans. *Mitochondrion* 34: 127–134, 2017. doi:10.1016/j.mito.2017.03.004.
28. Ali Pour P, Kenney MC, Kheradvar A. Bioenergetics consequences of mitochondrial transplantation in cardiomyocytes. *J Am Heart Assoc* 9: e014501, 2020. doi:10.1161/JAHA.119.014501.
29. Doulamis IP, Guariento A, Duignan T, Orfany A, Kido T, Zurakowski D, Del Nido PJ, McCully JD. Mitochondrial transplantation for myocardial protection in diabetic hearts. *Eur J Cardiothorac Surg* 57: 836–845, 2020. doi:10.1093/ejcts/ezz326.
30. Jiang XP, Elliott RL, Head JF. Exogenous normal mammary epithelial mitochondria suppress glycolytic metabolism and glucose uptake of human breast cancer cells. *Breast Cancer Res Treat* 153: 519–529, 2015. doi:10.1007/s10549-015-3583-0.
31. Popov LD. One step forward: extracellular mitochondria transplantation. *Cell Tissue Res* 384: 607–612, 2021. doi:10.1007/s00441-021-03428-5.
32. Bensley RR, Hoerr NL. Studies on cell structure by the freezing-drying method. VI. The preparation and properties of mitochondria. *Anat Record* 60: 449–455, 1934. doi:10.1002/ar.1090600408.
33. Sims NR. Rapid isolation of metabolically active mitochondria from rat brain and subregions using Percoll density gradient centrifugation. *J Neurochem* 55: 698–707, 1990. doi:10.1111/j.1471-4159.1990.tb04189.x.
34. Hubbard WB, Harwood CL, Prajapati P, Springer JE, Saatman KE, Sullivan PG. Fractionated mitochondrial magnetic separation for isolation of synaptic mitochondria from brain tissue. *Sci Rep* 9: 9656, 2019. doi:10.1038/s41598-019-45568-3.
35. de Duve C. Tissue fraction-past and present. *J Cell Biol* 50: 20, 1971.
36. Spees JL, Olson SD, Whitney MJ, Prockop DJ. Mitochondrial transfer between cells can rescue aerobic respiration. *Proc Natl Acad Sci U S A* 103: 1283–1288, 2006. doi:10.1073/pnas.051051103.
37. Lou E, Fujisawa S, Morozov A, Barlas A, Romin Y, Dogan Y, Gholami S, Moreira AL, Manova-Todorova K, Moore MA. Tunneling nanotubes provide a unique conduit for intercellular transfer of cellular contents in human malignant pleural mesothelioma. *PLoS One* 7: e33093, 2012. doi:10.1371/journal.pone.0033093.
38. Islam MN, Das SR, Emin MT, Wei M, Sun L, Westphalen K, Rowlands DJ, Quadri SK, Bhattacharya S, Bhattacharya J. Mitochondrial transfer from bone-marrow-derived stromal cells to pulmonary alveoli protects against acute lung injury. *Nat Med* 18: 759–765, 2012. doi:10.1038/nm.2736.
39. Huang X, Sun L, Ji S, Zhao T, Zhang W, Xu J, Zhang J, Wang Y, Wang X, Franzini-Armstrong C, Zheng M, Cheng H. Kissing and nanotunneling mediate intermitochondrial communication in the heart. *Proc Natl Acad Sci U S A* 110: 2846–2851, 2013. doi:10.1073/pnas.1300741110.
40. Macheiner T, Fengler VH, Agreiter M, Eisenberg T, Madeo F, Kolb D, Huppertz B, Ackbar R, Sargsyan K. Magnetomitotransfer: an efficient way for direct mitochondria transfer into cultured human cells. *Sci Rep* 6: 35571, 2016. doi:10.1038/srep35571.
41. Wu TH, Sagullo E, Case D, Zheng X, Li Y, Hong JS, TeSlaa T, Patananan AN, McCaffery JM, Niazi K, Braas D, Koehler CM, Graeber TG, Chiou PY, Teitell MA. Mitochondrial transfer by photo-thermal nanoblade restores metabolite profile in mammalian cells. *Cell Metab* 23: 921–929, 2016. doi:10.1016/j.cmet.2016.04.007.
42. Kitani T, Kami D, Matoba S, Gojo S. Internalization of isolated functional mitochondria: involvement of macropinocytosis. *J Cell Mol Med* 18: 1694–1703, 2014. doi:10.1111/jcmm.12316.
43. Masuzawa A, Black KM, Pacak CA, Ericsson M, Barnett RJ, Drumm C, Seth P, Bloch DB, Levitsky S, Cowan DB, McCully JD. Transplantation of autologously derived mitochondria protects the heart from ischemia-reperfusion injury. *Am J Physiol Heart Circ Physiol* 304: H966–H982, 2013. doi:10.1152/ajpheart.00883.2012.
44. Kaza AK, Wamala I, Friehs I, Kuebler JD, Rathod RH, Berra I, Ericsson M, Yao R, Thedsanamoorthy JK, Zurakowski D, Levitsky S, Del Nido PJ, Cowan DB, McCully JD. Myocardial rescue with autologous mitochondrial transplantation in a porcine model of ischemia/reperfusion. *J Thorac Cardiovasc Surg* 153: 934–943, 2017. doi:10.1016/j.jtcvs.2016.10.077.
45. Guariento A, Piekarski BL, Doulamis IP, Blitzer D, Ferraro AM, Harild DM, Zurakowski D, Del Nido PJ, McCully JD, Emani SM. Autologous mitochondrial transplantation for cardiogenic shock in pediatric patients following ischemia-reperfusion injury. *J Thorac Cardiovasc Surg*. In press. doi:10.1016/j.jtcvs.2020.10.151.
46. Cowan DB, Yao R, Akurathi V, Snay ER, Thedsanamoorthy JK, Zurakowski D, Ericsson M, Friehs I, Wu Y, Levitsky S, Del Nido PJ, Packard AB, McCully JD. Intracoronary delivery of mitochondria to the ischemic heart for cardioprotection. *PLoS One* 11: e0160889, 2016. doi:10.1371/journal.pone.0160889.
47. Shin B, Saeed MY, Esch JJ, Guariento A, Blitzer D, Moskowitsova K, Ramirez-Barbieri G, Orfany A, Thedsanamoorthy JK, Cowan DB, Inkster JA, Snay ER, Staffa SJ, Packard AB, Zurakowski D, Del Nido PJ, McCully JD. A novel biological strategy for myocardial protection by intracoronary delivery of mitochondria: safety and efficacy. *JACC Basic Transl Sci* 4: 871–888, 2019. doi:10.1016/j.jacbs.2019.08.007.
48. Blitzer D, Guariento A, Doulamis IP, Shin B, Moskowitsova K, Barbieri GR, Orfany A, Del Nido PJ, McCully JD. Delayed transplantation of autologous mitochondria for cardioprotection in a porcine model. *Ann Thorac Surg* 109: 711–719, 2020. doi:10.1016/j.athoracsurg.2019.06.075.
49. Guariento A, Blitzer D, Doulamis I, Shin B, Moskowitsova K, Orfany A, Ramirez-Barbieri G, Staffa SJ, Zurakowski D, Del Nido PJ, McCully JD. Preischemic autologous mitochondrial transplantation by intracoronary injection for myocardial protection. *J Thorac Cardiovasc Surg* 160: e15–e29, 2020. doi:10.1016/j.jtcvs.2019.06.111.
50. Moskowitsova K, Shin B, Liu K, Ramirez-Barbieri G, Guariento A, Blitzer D, Thedsanamoorthy JK, Yao R, Snay ER, Inkster JA, Orfany A, Zurakowski D, Cowan DB, Packard AB, Visner GA, Del Nido PJ, McCully JD. Mitochondrial transplantation prolongs cold ischemia time in murine heart transplantation. *J Heart Lung Transplant* 38: 92–99, 2019. doi:10.1016/j.healun.2018.09.025.
51. Kim MJ, Hwang JW, Yun CK, Lee Y, Choi YS. Delivery of exogenous mitochondria via centrifugation enhances cellular metabolic function. *Sci Rep* 8: 3330, 2018. doi:10.1038/s41598-018-21539-y.
52. Bereiter-Hahn J, Voth M. Dynamics of mitochondria in living cells: shape changes, dislocations, fusion, and fission of mitochondria. *Microsc Res Tech* 27: 198–219, 1994. doi:10.1002/jemt.1070270303.
53. Maeda H, Kami D, Maeda R, Murata Y, Jo JI, Kitani T, Tabata Y, Matoba S, Gojo S. TAT-dextran-mediated mitochondrial transfer enhances recovery from models of reperfusion injury in cultured cardiomyocytes. *J Cell Mol Med* 24: 5007–5020, 2020. doi:10.1111/jcmm.15120.
54. Cowan DB, Yao R, Thedsanamoorthy JK, Zurakowski D, Del Nido PJ, McCully JD. Transit and integration of extracellular mitochondria

- in human heart cells. *Sci Rep* 7: 17450, 2017. doi:10.1038/s41598-017-17813-0.
55. Kesner EE, Saada-Reich A, Lorberboum-Galski H. Characteristics of mitochondrial transformation into human cells. *Sci Rep* 6: 26057, 2016. doi:10.1038/srep26057.
56. Patel D, Rorbach J, Downes K, Szukszo MJ, Pekalski ML, Minczuk M. Macropinocytic entry of isolated mitochondria in epidermal growth factor-activated human osteosarcoma cells. *Sci Rep* 7: 12886, 2017. doi:10.1038/s41598-017-13227-0.
57. Fitridge R, Thompson M. Mechanisms of vascular disease: a reference book for vascular specialists. In: *Mechanisms of Vascular Disease: a Reference Book for Vascular Specialists*, edited by Fitridge R, Thompson M. Adelaide, Australia: University of Adelaide, 2011.
58. Pagliaro P, Femminò S, Popara J, Penna C. Mitochondria in Cardiac Postconditioning. *Front Physiol* 9: 287, 2018. doi:10.3389/fphys.2018.00287.
59. Piper HM, Noll T, Siegmund B. Mitochondrial function in the oxygen depleted and reoxygenated myocardial cell. *Cardiovasc Res* 28: 1–15, 1994. doi:10.1093/cvr/28.1.1.
60. Murphy MP. How mitochondria produce reactive oxygen species. *Biochem J* 417: 1–13, 2009. doi:10.1042/BJ20081386.
61. Minutoli L, Puzzolo D, Rinaldi M, Irrera N, Marini H, Arcoraci V, Bitto A, Crea G, Pisani A, Squadrito F, Trichilo V, Bruschetta D, Micali A, Altavilla D. ROS-mediated NLRP3 inflammasome activation in brain, heart, kidney, and testis ischemia/reperfusion injury. *Oxid Med Cell Longev* 2016: 2183026, 2016. doi:10.1155/2016/2183026.
62. Wu CC, Bratton SB. Regulation of the intrinsic apoptosis pathway by reactive oxygen species. *Antioxid Redox Signal* 19: 546–558, 2013. doi:10.1089/ars.2012.4905.
63. Turer AT, Hill JA. Pathogenesis of myocardial ischemia-reperfusion injury and rationale for therapy. *Am J Cardiol* 106: 360–368, 2010. doi:10.1016/j.amjcard.2010.03.032.
64. Tani M, Neely JR. Role of intracellular Na⁺ in Ca²⁺ overload and depressed recovery of ventricular function of reperfused ischemic rat hearts. Possible involvement of H⁺-Na⁺ and Na⁺-Ca²⁺ exchange. *Circ Res* 65: 1045–1056, 1989. doi:10.1161/01.res.65.4.1045.
65. Nayler WG. The role of calcium in the ischemic myocardium. *Am J Pathol* 102: 262–270, 1981.
66. Kusuoka H, Porterfield JK, Weisman HF, Weisfeldt ML, Marban E. Pathophysiology and pathogenesis of stunned myocardium. Depressed Ca²⁺ activation of contraction as a consequence of reperfusion-induced cellular calcium overload in ferret hearts. *J Clin Invest* 79: 950–961, 1987. doi:10.1172/JCI112906.
67. Lemasters JJ, Bond JM, Chacon E, Harper IS, Kaplan SH, Ohata H, Trollinger DR, Herman B, We C. The pH paradox in ischemia-reperfusion injury to cardiac myocytes. *EXS* 76: 99–114, 1996. doi:10.1007/978-3-0348-8988-9_7.
68. Bond JM, Herman B, Lemasters JJ. Protection by acidotic pH against anoxia/reoxygenation injury to rat neonatal cardiac myocytes. *Biochem Biophys Res Commun* 179: 798–803, 1991. doi:10.1016/0006-291x(91)91887-i.
69. Winterbourn CC. The biological chemistry of hydrogen peroxide. *Methods Enzymol* 528: 3–25, 2013. doi:10.1016/B978-0-12-405881-1.00001-X.
70. Murphy E, Steenbergen C. Mechanisms underlying acute protection from cardiac ischemia-reperfusion injury. *Physiol Rev* 88: 581–609, 2008. doi:10.1152/physrev.00024.2007.
71. Mattson MP, Culmsee C, Yu ZF. Apoptotic and antiapoptotic mechanisms in stroke. *Cell Tissue Res* 301: 173–187, 2000. doi:10.1007/s004419900154.
72. Zhou T, Prather ER, Garrison DE, Zuo L. Interplay between ROS and antioxidants during ischemia-reperfusion injuries in cardiac and skeletal muscle. *Int J Mol Sci* 19: 417, 2018. doi:10.3390/ijms19020417.
73. Lesnefsky EJ, Chen Q, Tandler B, Hoppel CL. Mitochondrial dysfunction and myocardial ischemia-reperfusion: implications for novel therapies. *Annu Rev Pharmacol Toxicol* 57: 535–565, 2017. doi:10.1146/annurev-pharmtox-010715-103335.
74. Raza H, John A, Howarth FC. Increased oxidative stress and mitochondrial dysfunction in Zucker diabetic rat liver and brain. *Cell Physiol Biochem* 35: 1241–1251, 2015. doi:10.1159/000373947.
75. Manara AR, Murphy PG, O'Callaghan G. Donation after circulatory death. *Br J Anaesth* 108, Suppl 1: i108–121, 2012. doi:10.1093/bja/aer357.
76. Guariento A, Doulamis IP, Duignan T, Kido T, Regan WL, Saeed MY, Hoganson DM, Emani SM, Fynn-Thompson F, Matte GS, Del Nido PJ, McCully JD. Mitochondrial transplantation for myocardial protection in ex-situ perfused hearts donated after circulatory death. *J Heart Lung Transplant*. In press. doi:10.1016/j.healun.2020.06.023.
77. Emani SM, Piekarski BL, Harild D, Del Nido PJ, McCully JD. Autologous mitochondrial transplantation for dysfunction after ischemia-reperfusion injury. *J Thorac Cardiovasc Surg* 154: 286–289, 2017. doi:10.1016/j.jtcvs.2017.02.018.
78. Scaglia F, Towbin JA, Craigen WJ, Belmont JW, Smith EO, Neish SR, Ware SM, Hunter JV, Fernbach SD, Vladutiu GD, Wong LJ, Vogel H. Clinical spectrum, morbidity, and mortality in 113 pediatric patients with mitochondrial disease. *Pediatrics* 114: 925–931, 2004. doi:10.1542/peds.2004-0718.
79. Holmgren D, Wahlander H, Eriksson BO, Oldfors A, Holme E, Tulinius M. Cardiomyopathy in children with mitochondrial disease: clinical course and cardiological findings. *Cardiovasc Pathol* 24: 107371–107288, 2021. doi:10.1016/S0195-668X(02)00387-1.
80. Fassone E, Rahman S. Complex I deficiency: clinical features, biochemistry and molecular genetics. *J Med Genet* 49: 578–590, 2012 [Erratum in *J Med Genet* 49: 668, 2012]. doi:10.1136/jmedgenet-2012-101159.
81. Alston CL, Ceccatelli Berti C, Blakely EL, Olahova M, He L, McMahon CJ, Olpin SE, Hargreaves IP, Nolli C, McFarland R, Goffrini P, O'Sullivan MJ, Taylor RW. A recessive homozygous p. Asp92Gly SDHD mutation causes prenatal cardiomyopathy and a severe mitochondrial complex II deficiency. *Hum Genet* 134: 869–879, 2015. doi:10.1007/s00439-015-1568-z.
82. Schaefer AM, Taylor RW, Turnbull DM, Chinnery PF. The epidemiology of mitochondrial disorders—past, present and future. *Biochim Biophys Acta* 1659: 115–120, 2004. doi:10.1016/j.bbabi.2004.09.005.
83. Yang H, Borg TK, Ma Z, Xu M, Wetzel G, Saraf LV, Markwald R, Runyan RB, Gao BZ. Biochip-based study of unidirectional mitochondrial transfer from stem cells to myocytes via tunneling nanotubes. *Biofabrication* 8: 015012, 2016. doi:10.1088/1758-5090/8/1/015012.
84. Wang X, Gerdes HH. Transfer of mitochondria via tunneling nanotubes rescues apoptotic PC12 cells. *Cell Death Differ* 22: 1181–1191, 2015. doi:10.1038/cdd.2014.211.
85. Li CJ, Chen PK, Sun LY, Pang CY. Enhancement of mitochondrial transfer by antioxidants in human mesenchymal stem cells. *Oxid Med Cell Longev* 2017: 8510805, 2017. doi:10.1155/2017/8510805.
86. Feng Y, Zhu R, Shen J, Wu J, Lu W, Zhang J, Zhang J, Liu K. Human bone marrow mesenchymal stem cells rescue endothelial cells experiencing chemotherapy stress by mitochondrial transfer via tunneling nanotubes. *Stem Cells Dev* 28: 674–682, 2019. doi:10.1089/scd.2018.0248.
87. Ikeda G, Santoso MR, Tada Y, Li AM, Vaskova E, Jung JH, O'Brien C, Egan E, Ye J, Yang PC. Mitochondria-rich extracellular vesicles from autologous stem cell-derived cardiomyocytes restore energetics of ischemic myocardium. *J Am Coll Cardiol* 77: 1073–1088, 2021. doi:10.1016/j.jacc.2020.12.060.
88. Alexeyev M, Shokolenko I, Wilson G, LeDoux S. The maintenance of mitochondrial DNA integrity—critical analysis and update. *Cold Spring Harb Perspect Biol* 5: a012641, 2013. doi:10.1101/cshperspect.a012641.
89. El-Hattab AW, Scaglia F. Mitochondrial cardiomyopathies. *Front Cardiovasc Med* 3: 25, 2016. doi:10.3389/fcvm.2016.00025.
90. Seo AY, Joseph AM, Dutta D, Hwang JC, Aris JP, Leeuwenburgh C. New insights into the role of mitochondria in aging: mitochondrial dynamics and more. *J Cell Sci* 123: 2533–2542, 2010. doi:10.1242/jcs.070490.
91. Jacobs HT, Lehtinen SK, Spelbrink JN. No sex please, we're mitochondria: a hypothesis on the somatic unit of inheritance of mammalian mtDNA. *Bioessays* 22: 564–572, 2000. doi:10.1002/(SICI)1521-1878(200006)22:6<564::AID-BIES9>3.0.CO;2-4.
92. DiMauro S. Mitochondrial diseases. *Biochim Biophys Acta* 1658: 80–88, 2004. doi:10.1016/j.bbabi.2004.03.014.

93. **Detmer SA, Chan DC.** Functions and dysfunctions of mitochondrial dynamics. *Nat Rev Mol Cell Biol* 8: 870–879, 2007. doi:10.1038/nrm2275.
94. **Detmer SA, Chan DC.** Complementation between mouse Mfn1 and Mfn2 protects mitochondrial fusion defects caused by CMT2A disease mutations. *J Cell Biol* 176: 405–414, 2007. doi:10.1083/jcb.200611080.
95. **Ono T, Isobe K, Nakada K, Hayashi JI.** Human cells are protected from mitochondrial dysfunction by complementation of DNA products in fused mitochondria. *Nat Genet* 28: 272–275, 2001. doi:10.1038/90116.
96. **Herbst A, Pak JW, McKenzie D, Bua E, Bassiouni M, Aiken JM.** Accumulation of mitochondrial DNA deletion mutations in aged muscle fibers: evidence for a causal role in muscle fiber loss. *J Gerontol A Biol Sci Med Sci* 62: 235–245, 2007. doi:10.1093/gerona/62.3.235.
97. **Bevers RP, Litovchenko M, Kapopoulou A, Braman VS, Robinson MR, Auwerx J, Hollis B, Deplancke B.** Mitochondrial haplotypes affect metabolic phenotypes in the Drosophila Genetic Reference Panel. *Nat Metab* 1: 1226–1242, 2019 [Erratum in *Nat Metab* 2: 381, 2020]. doi:10.1038/s42255-019-0147-3.
98. **Liu X, Trakooljul N, Hadlich F, Murani E, Wimmers K, Ponsuksili S.** Mitochondrial-nuclear crosstalk, haplotype and copy number variation distinct in muscle fiber type, mitochondrial respiratory and metabolic enzyme activities. *Sci Rep* 7: 14024, 2017. doi:10.1038/s41598-017-14491-w.
99. **Lee WT, Sun X, Tsai TS, Johnson JL, Gould JA, Garama DJ, Gough DJ, McKenzie M, Trounce IA, St John JC.** Mitochondrial DNA haplotypes induce differential patterns of DNA methylation that result in differential chromosomal gene expression patterns. *Cell Death Discov* 3: 17062, 2017. doi:10.1038/cddiscovery.2017.62.
100. **Struck J, Uhlein M, Morgenthaler NG, Furst W, Hoflich C, Bahrami S, Bergmann A, Volk HD, Redl H.** Release of the mitochondrial enzyme carbamoyl phosphate synthase under septic conditions. *Shock* 23: 533–538, 2005.
101. **Iyer SS, Pulsikens WP, Sadler JJ, Butter LM, Teske GJ, Ulland TK, Eisenbarth SC, Florquin S, Flavell RA, Leemans JC, Sutterwala FS.** Necrotic cells trigger a sterile inflammatory response through the Nlrp3 inflammasome. *Proc Natl Acad Sci U S A* 106: 20388–20393, 2009. doi:10.1073/pnas.0908698106.
102. **Collins LV, Hajizadeh S, Holme E, Jonsson IM, Tarkowski A.** Endogenously oxidized mitochondrial DNA induces in vivo and in vitro inflammatory responses. *J Leukoc Biol* 75: 995–1000, 2004. doi:10.1189/jlb.0703328.
103. **Gollihue JL, Rabchevsky AG.** Prospects for therapeutic mitochondrial transplantation. *Mitochondrion* 35: 70–79, 2017. doi:10.1016/j.mito.2017.05.007.
104. **Lin L, Xu H, Bishawi M, Feng F, Samy K, Truskey G, Barbas AS, Kirk AD, Brennan TV.** Circulating mitochondria in organ donors promote allograft rejection. *Am J Transplant* 19: 1917–1929, 2019. doi:10.1111/ajt.15309.
105. **Pollara J, Edwards RW, Lin L, Bendersky VA, Brennan TV.** Circulating mitochondria in deceased organ donors are associated with immune activation and early allograft dysfunction. *JCI Insight* 3: e121622, 2018. doi:10.1172/jci.insight.121622.
106. **Zhang Q, Raoof M, Chen Y, Sumi Y, Sursal T, Junger W, Brohi K, Itagaki K, Hauser CJ.** Circulating mitochondrial DAMPs cause inflammatory responses to injury. *Nature* 464: 104–107, 2010. doi:10.1038/nature08780.
107. **Zhang Q, Itagaki K, Hauser CJ.** Mitochondrial DNA is released by shock and activates neutrophils via p38 map kinase. *Shock* 34: 55–59, 2010. doi:10.1097/SHK.0b013e3181cd8c08.