

# Electrophilic Sulfur Reagent Design Enables Directed *syn*-Carbosulfonylation of Unactivated Alkenes

Zi-Qi Li<sup>†</sup>, Yilin Cao<sup>†</sup>, Taeho Kang<sup>†</sup>, and Keary M. Engle<sup>†\*</sup>

<sup>†</sup> Department of Chemistry, Scripps Research, 10550 North Torrey Pines Road, La Jolla, California 92037, United States

**ABSTRACT:** A multi-component approach to structurally complex organosulfur products is described via the nickel-catalyzed 1,2-carbosulfonylation of unactivated alkenes with organoboron nucleophiles and tailored organosulfur electrophiles. Key to the development of this transformation is the identification of a modular *N*-alkyl-*N*-(arylsulfonyl)arenesulfonamide family of sulfur electrophiles. Tuning the electronic and steric properties of the leaving group in these reagents controls pathway selectivity, favoring three-component coupling and suppressing side reactions, as examined via computational studies. The unique *syn*-stereoselectivity differs from traditional electrophilic sulfonyl transfer processes involving a thiiranium ion intermediate and arises from the directed arylnickel(I) migratory insertion mechanism, as elucidated through reaction kinetics and control experiments. Reactivity and regioselectivity are facilitated by a collection of monodentate, weakly coordinating native directing groups, including sulfonamides, alcohols, amines, amides, and azaheterocycles.

## INTRODUCTION

Organosulfur compounds have diverse functions and find applications as pharmaceuticals<sup>1</sup> and functional materials.<sup>2</sup> Organosulfides, in particular, can also be readily converted into other functional groups, making them versatile building blocks in synthesis.<sup>3</sup> While reliable approaches for catalytic two-component C–S bond formation<sup>4</sup> via cross-coupling reactions<sup>5</sup> and C–H functionalization<sup>6</sup> have emerged during the past two decades, multicomponent C–S bond-forming reactions remain less explored.<sup>7</sup> In this context, reactions that merge an alkene, a sulfur-based reaction partner, and a carbogenic group combine C–C skeletal formation and C–S installation into a single operation, representing an attractive means of generating complex, stereochemically dense organosulfur products from simple chemical inputs. Pioneered by Trost<sup>8</sup>, Denmark<sup>9</sup> and others<sup>10</sup>, the most well-established method for 1,2-carbosulfonylation of alkenes involves formation of a thiiranium ion intermediate<sup>11</sup> through electrophilic sulfonyl transfer and subsequent nucleophilic ring opening, with *anti*-stereoselectivity arising from the S<sub>N</sub>2 nature of the ring-opening step and regioselectivity dictated by alkene substitution pattern (scheme 1A). While enabling in its own right, existing methodology for three-component *anti*-1,2-carbosulfonylation of alkenes is limited to a small collection of carbon-nucleophiles, namely cyanide<sup>8a</sup>, acetylides<sup>8b</sup>, and organozinc reagents<sup>10d</sup>, and in the latter case, the transformation is only compatible with styrene substrates. Complementing this existing methodology with a *syn*-stereoselective counterpart that proceeds via a distinct mechanistic pathway would be highly enabling (scheme 1B). By bypassing thiiranium ion formation, we envisioned that it would not only be possible to achieve the opposite stereochemical outcome but also to expand the alkene scope to include nucleophilic functional groups that are prone to intramolecular cyclization in established thiiranium chemistry.<sup>9g, 12</sup>


Recently, nickel-catalyzed redox-neutral 1,2-functionalization of unactivated alkenes<sup>13</sup> has emerged as a powerful method for joining together a nucleophile, an electrophile, and an alkene in

a selective fashion.<sup>14</sup> Depending on the identity of the coupling partners, two closely related yet distinct mechanisms can operate (scheme 1C). A Ni(0)/Ni(II) cycle occurs by an oxidative-addition-first pathway, where the electrophile is incorporated distal to the directing group.<sup>15</sup> Meanwhile, a Ni(I)/Ni(III) cycle involving a transmetalation-first mechanism typically manifests in reversed regioselectivity.<sup>16</sup> In both cases, *syn*-selectivity is dictated by inner-sphere migratory insertion. With these considerations in mind and informed by our previous experience in developing nickel-catalyzed 1,2-difunctionalization of alkenes directed by native functional groups<sup>15c</sup>, we reasoned that it would be possible to develop a *syn*-selective 1,2-carbosulfonylation by employing sulfur-based electrophiles within this mode of catalysis. While attractive in principle, the envisioned three-component coupling would require surmounting several challenges, including premature sulfonyl transfer to the alkene or carbon nucleophile, competitive β-hydride elimination from the alkynickel intermediate, and the potential inhibition from the sulfur-containing products or leaving group generated upon sulfonyl transfer.


Herein we report a nickel-catalyzed *syn*-selective 1,2-carbosulfonylation reaction<sup>17</sup> of simple unactivated alkenes<sup>18</sup>. Critical to the success of this reaction is the identification of *N*-sulfonyl-*N*-alkyl sulfonamides as N–S electrophiles (scheme 1D). Taking inspiration from various electrophiles in the literature that contain nitrogen-based leaving groups for fluorination<sup>19</sup>, trifluoromethylthiolation<sup>20</sup>, cyanation<sup>21</sup>, and acylation<sup>22</sup>, this design takes advantages of the sterically and electronically tunable nature of the sulfonamide leaving group (N<sub>LC</sub>), which can be readily modified along each of its two vectors to control pathway selectivity and thereby maximize the yield of desired three-component coupling.

## Scheme 1. Background and Synopsis of Current Work


A. Thiiiranum-ion-mediated anti-selective 1,2-carbosulfonylation.




B. This work: *syn*-selective 1,2-carbosulfonylation of unactivated alkenes.



C. Nickel-catalyzed redox-neutral alkene 1,2-difunctionalization.



D. Carbon- and heteroatom electrophiles containing nitrogen leaving groups.



## RESULTS AND DISCUSSION

**Reaction discovery.** To begin the investigation, we focused on the model reaction of alkanyl sulfonamide **1<sup>23</sup>** with *p*-tolylboronic acid neopentyl glycol ester (**2a**) and various N-S electrophiles (scheme 2A). In effort to find a suitable sulfonyl electrophile with the proper reactivity profile—appropriately tempered to not transfer the sulfonyl group to the alkene or arylnickel(I) intermediate yet sufficiently reactive to engage the alkynickel(I) intermediate with a rate faster than that of  $\beta$ -hydride elimination—we first attempted established sulfonyl electrophiles from the literature (**S1–S7**). Modest success was found with *N*-sulfonyl lactams (17–51% yield, **S5–S7**)<sup>7a</sup>; however, desired three-component coupling was accompanied by formation of nearly equimolar quantities of oxidative Heck byproducts, indicating similar rates of intramolecular  $\beta$ -hydride elimination and intermolecular N-S oxidative addition. Extensive screening of the reaction conditions did not improve yields or product selectivity with leaving groups **NLG5–NLG7**. In contrast, with an *N*-Me benzamide as leaving group, better yields (53–64%) and product selectivities (7–11% combined byproducts) were observed (**S8–S10**). We then examined **S11–S23** with *N*-substituted arylsulfonamides as leaving groups, taking advantage of the easily tunable nature of this scaffold, which allows numerous modifications to be quickly assayed. Although electron-withdrawing groups (as in **S11–S13**)<sup>24</sup> proved deleterious for the reaction, improved yields (69–94%) were observed with **S14–S17**. When a *para*-methoxy group was used (**S17**), the desired product was obtained in 94% yield and with <5% oxidative Heck byproducts. Steric modification to either the arylsulfonyl or *N*-alkyl vectors revealed no further improvement (**S18–S22**).

Interestingly, when the *N*-alkyl group was changed to a *N*-phenyl group, no desired product was observed (**S23**).<sup>25</sup>


We performed computational parameterization of the tested reagents to understand the origins of the high product yield with **S17** (scheme 2B). As N-S reagents were expected to be involved mainly during the oxidative addition step, LUMO energy and bond dissociation energy (BDE)<sup>26</sup> were computed and plotted against percentage yields. The former value relates to kinetics and thermodynamics of organonickel coordination to the N-S reagent, while the latter reflects the subsequent oxidative addition process. By visualizing the data using a 3D-plot, we were pleased to find the maximum yield is obtained when both parameters are within certain threshold (50 to 52 kcal/mol for BDE and -0.6 to -0.1 eV for LUMO energy). Within this data set, analysis of several poor-performing electrophiles is informative. Reagents **S11–S13** possess BDEs near the optimal range but have low LUMO energy (mean of -0.9 eV), reflecting the presence of the electron-withdrawing groups. On the other hand, **S23**, which contains the same leaving group as a previously reported radical fluorinating agent,<sup>25</sup> has a LUMO energy within the typical range of high-yielding variants but features a BDE of 39.6 kcal/mol—as low as the well accepted SOMO-phile **S1** (39.8 kcal/mol)<sup>24</sup>. In this case, single-electron transfer (SET) may take place to generate off-cycle species (i.e., a stable nitrogen centered radical and a sulfur radical). This also agrees with our hypothesis that **S17** functions as a 2e- covalent electrophile. The moderate yields obtained with **S5–S10** could reflect the slightly higher than optimal BDE of 52.7 kcal/mol, with the sulfur possessing diminished electrophilicity due to the weaker electron-withdrawing effect of the *N*-alkylamide versus *N*-alkylsulfonamide leaving group. Interestingly, **S4**, which has been applied in Lewis-base-catalyzed sulfonyl transfer reactions involving thiiranum intermediates<sup>9, 12</sup>, was too stable to react (BDE = 79.2 kcal/mol).

X-ray crystallographic analysis (scheme 2C) of representative N-S electrophiles revealed a non-pyramidalized N-atom<sup>27</sup> stemming from delocalization of the lone pair electron from nitrogen to the S=O  $\pi^*$  orbitals (**S17**, **S18**). Meanwhile, in the case of **S23**, a more pyramidalized nitrogen and longer N-S bond indicates diminished delocalization and thus a weaker N-S bond. **S6** on the other hand, features an approximately trigonal planar structure which is responsible for its higher BDE (55.3 kcal/mol) and lower reactivity.

**Substrate scope.** Having identified an optimal class of N-S electrophiles, we examined the scope and limitations of the method (Table 1). Regarding the electrophile scope, arylsulfonyl moieties with electron-neutral or -withdrawing substituents (**3aa–3ag**) could be efficiently introduced with the optimal leaving group (**NLG17**). On the other hand, the highly electron-donating *p*-methoxy case (**3ab**) benefited from using a less electron-rich leaving group (**NLG15**), indicating that it is possible to further fine-tune the leaving group to match the electronic properties of the arylsulfonyl moiety as needed (see SI for comparative data). Sterically bulky arylsulfonyl groups did not significantly inhibit the reaction (**3ah–3ak**), and halo-substituted arylsulfonyl coupling partners were tolerated (**3ac–3ae**, **3ai–3aj**), offering the opportunity for further derivatization by cross-coupling. In preliminary experiments, we found that an alkylsulfonyl group could also be introduced onto two representative alkanyl sulfonamide substrates, though yields were low in these cases (**3al–3am**).

**Scheme 2. Discovery of nickel-catalyzed 1,2-carbosulfonylation of unactivated alkenes**

A. Optimization of reaction conditions<sup>a</sup>




<sup>a</sup>Optimization of reaction conditions.  $\text{Ni}(\text{cod})_2/1/2\text{a}/[\text{N-S}]/\text{NaOH} = 0.01/0.1/0.2/0.2/0.2$  (mmol). Reaction was run in THF (0.05 M) at 60 °C. All percentages represent <sup>1</sup>H NMR yields with benzyl 4-fluorobenzoate as internal standard. <sup>b</sup>Surface of Yield/LUMO energy/BDE plot generated by 3D mathematical fitting. For detailed 2D Yield/LUMO energy and Yield/BDE plots with computed data points, see supporting information. Gas phase LUMO energy calculated at M05-2X/6-311+G(d,p) level of theory. Solution phase BDE calculated at SMD(acetonitrile)/M05-2X/6-311+G(d,p), SMD(acetonitrile)/M05-2X/6-311+G(d) level of theory. <sup>c</sup>Hydrogen atoms omitted for clarity. Introduction of an electron-withdrawing group on the arylsulfonyl group (*p*-CF<sub>3</sub>) with Nu17 led to a more pyramidalized nitrogen ( $\chi_N = 32.17856$ ) and longer N-S bond (1.69376 Å); See supporting information for crystal structure of **S28** (CCDC 2110120).

Next, we surveyed different organoboron nucleophiles. Arylboron coupling partners substituted at the *para*- or *meta*-positions with electron-donating or -withdrawing groups were generally well tolerated, with the former slightly higher-yielding in general (**3ba**–**3bo**). Potentially inhibitory or reactive groups, such as  $-\text{NHBOC}$ ,  $-\text{CHO}$ , and  $-\text{CN}$ , were all compatible (**3bg**–**3bi**, **3bm**–**3bn**). Furthermore, with an *ortho*-methyl substituent, **3bp** was obtained in 56% yield. Bicyclic coupling partners also proved suitable, giving **3bq**–**3bt** in moderate to good yield. Six-membered heteroaryl nucleophiles, including quinoline, pyridine, and purine moiety, could be installed though a substituent at the 2-position was required to attenuate the coordinating ability of the basic N(*sp*<sup>2</sup>) atom (**3bu**–**3bw**). On the other hand, unmasked pyridine and purine as well as electron-rich five-membered heterocycles were incompatible in this reaction. Alkenyl-B(nep) reactants constitute another limitation, as only 20% of the corresponding product was observed in the case of a representative styrenyl coupling partner.

A series of substituted alkenyl sulfonamides were evaluated. Sulfonamides derived from both homoallyl and bishomoallyl

amines were tolerated (**4a**–**4h**), with the former giving slightly lower yield. Gratifyingly, benzenesulfonyl groups with different substituents at the *para*-position (**4a**–**4g**) and a methanesulfonyl group (**4h**) all furnished the corresponding products in moderate to good yield. 4-Cyano benzenesulfonamide (Cs) can serve the dual role of directing group and amine activating group (see below) and gave **4f** and **4g** in 53% and 85% yield, respectively.<sup>28</sup> Moderate yield and diastereoselectivity were obtained with substrates containing  $\alpha$ - or  $\beta$ -branching (**4i**–**4k**). With internal alkenes (**4l**–**4r**), *Z*-configured alkenes provided higher yield, as well as greater and opposite diastereoselectivity, compared with the corresponding *E*-isomers. With a skipped diene substrate, the reaction took place selectively at the  $\gamma,\delta$ -position, due to the formation of a relatively stable 5-membered metallacycle (**4q**). With a 4-trifluoromethylbenzenesulfonyl directing group,<sup>23</sup> **4r** was obtained in 78% yield and >20:1 d.r. through a 6-membered metallacycle. A sulfonamide derived from *ortho*-allylaniline furnished **4s** in 59% yield with no chain-walking byproducts detected.

**Table 1. Electrophile, Nucleophile, and Alkenyl Sulfonamide Scope<sup>a</sup>**




<sup>a</sup>Reactions performed on 0.1 mmol scale. Percentages represent isolated yields. <sup>b</sup>Yield in brackets was obtained from an experiment performed on 1.0 mmol scale. <sup>c</sup>Yield in parentheses was obtained with **S34** (featuring **N<sub>Li</sub>11** as leaving group, see SI for details). <sup>d</sup>Percentage yields represent <sup>1</sup>H NMR yields with benzyl 4-fluorobenzoate as internal standard. N.D.=not detected. <sup>e</sup>Yield obtained with **S35** (see SI for details).

We next investigated alkenes containing other synthetically useful native functional groups (Table 2). This seemingly straightforward extension is complicated by the fact that minor changes to the directing group structure have a significant impact on coordination strength and metallacycle stability. To our delight, after brief optimization of base and solvent (see SI for details), a diverse collection of native directing groups was found to facilitate this transformation. Starting from homoallyl alcohol, **5a** and **5b** were obtained in 64% and 72% yield, respectively.<sup>16d</sup> A large-scale experiment on 4 mmol scale furnished **5b** in 59% yield. Substituents at the  $\alpha$ -position with respect to the hydroxyl group furnished **5c** and **5d** in 49% and 61% yields, respectively, with negligible diastereoselectivity. Late-stage functionalization of allylestrenol, a progestin medication, furnished the corresponding product in 34% yield

with 1.5:1 diastereoselectivity (**5e**). Free alkenyl amines bearing a basic  $N(sp^3)$  atom are traditionally challenging substrates owing to their ability to sequester the catalyst off-cycle. Gratifyingly, **5f**–**5j** were obtained in moderate to good yield. With a chiral benzyl amine directing auxiliary, **5h** was obtained in 60% yield and 1.1:1 diastereoselectivity. In addition, an alkene bearing an amide group proved compatible, giving **5k** in 35% yield. Motivated by our previous success in developing a 1,2-allylmethylation reaction of alkenes directed by diverse azaheterocycles<sup>29</sup>, we attempted these useful substrate families and were pleased to find that pyridine (**5l**), pyrazoles (**5m**–**5o**), triazoles (**5p**–**5q**), tetrazole (**5r**), indazoles (**5s**–**5t**), and benzotriazole (**5u**) were all compatible directing groups.

**Table 2. Alkene Scope<sup>a</sup>**



<sup>a</sup>Reactions performed on 0.1 mmol scale. Percentages represent isolated yields. Diastereomeric ratios were determined by <sup>1</sup>H NMR analysis of the isolated products.

<sup>b</sup>Results in brackets was obtained on 4.0 mmol scale.

**Synthetic versatility.** The N-S reagents used in this study can readily be prepared on decagram scale using modular and operationally convenient chemistry. If desired, the sulfonamide leaving group can be recovered in nearly quantitative yield and recycled (see SI for detail). The reaction can be performed outside of the glovebox using the air-stable precatalyst Ni(cod)(DQ) under Schlenk technique, as demonstrated using the model reaction in scheme 3A (80% yield, entry 1a).<sup>30</sup> A series of stress tests showed that this protocol is fairly robust. Simply purging with nitrogen over the solvent surface was sufficient (entry 1b), and even running the reaction under air gave 44% yield (entry 1c). When 1 mol% catalyst loading was used with extended reaction time of 40 h, 70% yield was obtained (entry 2). Reducing the electrophile and nucleophile loading led to lower yet synthetically useful yields (entry 3). A series of product derivatizations were performed to showcase the synthetic utility of the method (scheme 3B). From a  $-NH(Cs)$  product, *N*-alkylation and *-arylation* followed by mild deprotecting furnished **6a** and **6b**.<sup>28</sup>

This sequence allows straightforward incorporation of groups that are problematic in the nickel-catalyzed 1,2-carbosulfonylation (e.g., nitroarenes). Hofmann–Löffler–Freytag (HLF) cyclization furnished 1,2-disubstituted pyrrolidine **6c** in 62% overall yield over 2 steps.<sup>31</sup> Controlled oxidation of the thioether to the sulfoxide or sulfone and successive oxidation to the sulfonyl carboxylic acid using different oxidants furnished **6d**–**6f** in excellent yields<sup>9</sup>, with **6e** representing a key intermediate in the synthesis of matrix metalloproteinases (MMP) inhibitor.<sup>32</sup> Finally, a thianthrene-based radical precursor **6g** was prepared over 3 steps in 72% overall yield.

**Mechanistic studies.** A control experiment with *N*-methyl alkenyl sulfonamide (**7a**) under standard conditions did not lead to product formation, suggesting that the sulfonamide group coordinates through nitrogen (scheme 3C). Similarly, substrates without a suitable directing group such as 1-phenylbutene and 1-dodecene were unreactive (see SI for detail). No desired product was formed from styrenyl

sulfonamide (**7b**), which rules out a stepwise oxidative Heck/hydrosulfenylation pathway. The relative stereochemistry of **4o'** (major) was determined by single-crystal X-ray diffraction of an oxidized derivative, indicating a *syn*-addition mechanism. Kinetic studies were performed using the conversion of **1** to **3bd** as the model reaction applying the method of initial rates (scheme 3D, see SI for detail). The reaction was positive order in  $[\text{Ni}(\text{cod})_2]$  and **[S17]**. Saturation kinetics were observed in **[1]**, as it showed positive order at low **[1]** and zero-order behavior at high **[1]**. The other two


components,  $[\text{ArB}(\text{nep})]$  and  $[\text{NaOH}]$ , were zero-order. Collectively, the data are consistent with the  $\text{Ni}^{\text{II}}/\text{Ni}^{\text{III}}$  cycle shown in scheme 3E, involving a sequence of transmetalation, migratory insertion, N–S oxidative addition, and C–S reductive elimination.<sup>16, 33</sup> The kinetic data indicate that the turnover-limiting step at standard or elevated alkene concentration is N–S oxidative addition, whereas at low alkene concentration, an earlier step becomes turnover-limiting, namely alkene coordination or irreversible migratory insertion following reversible alkene binding.

### Scheme 3. Synthetic Versatility and Mechanistic Studies


#### A. Stress tests

entry	deviation from standard <sup>a</sup>	yield (%) <sup>b</sup>	Note
1	$\text{Ni}(\text{cod})_2$ in place of $\text{Ni}(\text{cod})_2$		
1a	Schlenk Technique	80	
1b	purge $\text{N}_2$ for 5 sec	79	
1c	Under Air	44	
2	with 1 mol% $\text{Ni}(\text{cod})_2$	70	40 h
3	with low <b>S17</b> and <i>p</i> -TolB(nep) loading	82/78	1.5/1.2 equiv


#### B. Product derivatization



#### C. Mechanistic experiments



#### D. Kinetic experiments



<sup>a</sup>Standard conditions:  $\text{Ni}(\text{cod})_2/1/\text{S17}/2\text{a}/\text{NaOH} = 0.01/0.1/0.2/0.2/0.2$  (mmol). Reaction was run in THF (0.05 M) at 60 °C. <sup>b</sup>All percentages represent  $^1\text{H}$  NMR yields with benzyl 4-fluorobenzoate as internal standard. <sup>c</sup>TT = thianthrene. <sup>d</sup>Standard conditions:  $\text{Ni}(\text{cod})_2/1/\text{S17}/4\text{-FC}_6\text{H}_4\text{B}(\text{nep})/\text{NaOH} = 0.015/0.15/0.27/0.3/0.3$  (mmol). Reaction was run in THF (0.05 M) at 35 °C. Normalized initial rate was calculated by defining initial rate of the standard conditions as 1.0 within each set of experiments. For details see supporting information.

## CONCLUSION

In conclusion, a novel family of N–S reagents with *N*-alkylsulfonamide leaving groups was developed to enable 1,2-carbosulfenylation of unactivated alkenes under nickel catalysis. The synthetic versatility of the method stems from the broad scope of compatible alkenes bearing different native directing groups. The identification of N–S reagents based upon modular components allows for fine tuning of the electronic and steric properties, which ultimately affect stability, reactivity, and catalytic performance. DFT calculations indicate that optimal yield is achieved when parameters such as BDE and LUMO energy fall within a narrow range of values. This finding is expected to guide the development of other heteroatom-based electrophiles for use in multicomponent catalytic couplings.

## ASSOCIATED CONTENT

Supporting Information. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Experimental procedures, spectral and crystallographic data (PDF)

NMR data (MNova format) (ZIP)

**Accession Codes** CCDC 2110122 (**S6**), 2114518 (**S17**), 2110121 (**S18**), 2117151 (**S23**), 2110120 (**S28**), 2110123 ( $\pm$ -**6c**), 2117152 ( $\pm$ -**6h**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), or by emailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk), or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

## AUTHOR INFORMATION

### Corresponding Author

Keary M. Engle – Department of Chemistry, The Scripps Research Institute, La Jolla, California 92037, United States; orcid.org/0000-0003-2767-6556; Email: keary@scripps.edu

## Authors

Zi-Qi Li – Department of Chemistry, The Scripps Research Institute, La Jolla, California 92037, United States; orcid.org/0000-0002-1353-3004.

Yilin Cao – Department of Chemistry, The Scripps Research Institute, La Jolla, California 92037, United States; orcid.org/0000-0002-4964-0565.

Taeho Kang – Department of Chemistry, The Scripps Research Institute, La Jolla, California 92037, United States; orcid.org/0000-0003-4549-262X.

## Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENT

This work was financially supported by the National Science Foundation (CHE-2102550), the Alfred P. Sloan Fellowship program, and the Camille Dreyfus Teacher-Scholar program. We thank Bristol Myers Squibb for a Graduate Fellowship (Z.-Q.L.), the National Science Foundation (DBI-1759544) for a Summer Undergraduate Research Fellowship (Y.C.), and Kwanjeong Educational Foundation for a Graduate Fellowship (T.K.). Dr. Milan Gembicky, Dr. Jake B. Bailey, and Dr. Erika Samolova are acknowledged for X-ray crystallographic analysis. Dr. Jason Chen is acknowledged for HRMS analysis. Prof. Peng Liu (Pittsburg), Dr. Luo-Yan Liu (Scripps Research), Dr. Zhen Liu (Caltech), and Dr. Michael A. Schmidt (Bristol Myers Squibb) are acknowledged for helpful discussion.

## REFERENCES

1. (a) Islardi, E. A.; Vitaku, E.; Njardarson, J. T. Data-mining for sulfur and fluorine: an evaluation of pharmaceuticals to reveal opportunities for drug design and discovery. *J. Med. Chem.* **2014**, *57*, 2832–2842; (b) Zhao, C.; Rakesh, K. P.; Ravidar, L.; Fang, W. Y.; Qin, H. L. Pharmaceutical and medicinal significance of sulfur (S(VI))-Containing motifs for drug discovery: A critical review. *Eur. J. Med. Chem.* **2019**, *162*, 679–734.
2. Takimiya, K.; Osaka, I.; Mori, T.; Nakano, M. Organic semiconductors based on [1]benzothieno[3,2-b][1]benzothiophene substructure. *Acc. Chem. Res.* **2014**, *47*, 1493–1502.
3. Modha, S. G.; Mehta, V. P.; van der Eycken, E. V. Transition metal-catalyzed C–C bond formation via C–S bond cleavage: an overview. *Chem. Soc. Rev.* **2013**, *42*, 5042–5055.
4. (a) Kondo, T.; Mitsudo T. Metal-catalyzed carbon-sulfur bond formation. *Chem. Rev.* **2000**, *100*, 3205–3220. (b) Beletskaya, I. P.; Ananikov, V. P. Transition-metal-catalyzed C–S, C–Se, and C–Te bond formation via cross-coupling and atom-economic addition reactions. *Chem. Rev.* **2011**, *111*, 1596–1636.
5. Mann, G.; Baranano, D.; Hartwig, J. F.; Rheingold, A. L.; Guzei, I. A. Carbon–sulfur bond-forming reductive elimination involving  $sp$ -,  $sp^2$ -, and  $sp^3$ -hybridized carbon. mechanism, steric effects, and electronic effects on sulfide formation. *J. Am. Chem. Soc.* **1998**, *120*, 9205–9219.
6. Shen, C.; Zhang, P.; Sun, Q.; Bai, S.; Hor, T. S. A.; Liu, X. Recent advances in C–S bond formation via C–H bond functionalization and decarboxylation. *Chem. Soc. Rev.* **2015**, *44*, 291–314.
7. (a) Li, R. H.; Zhou, Y.; Yoon, K. Y.; Dong, Z.; Dong, G. B. Sulfenamide-enabled *ortho* thiolation of aryl iodides via palladium/norbornene cooperative catalysis. *Nat. Commun.* **2019**, *10*, 8; (b) Cai, W. Q.; Gu, Z. H. Selective *ortho* thiolation enabled by tuning the ancillary ligand in palladium/norbornene catalysis. *Org. Lett.* **2019**, *21*, 3204–3209.
8. (a) Trost, B. M.; Shibata, T.; Martin, S. J. Nucleophilic attack on olefins initiated by dimethylmethyliosulfonium fluoroborate (DMTSF). Cyanosulfonylation and oxy- and oxosulfonylation. *J. Am. Chem. Soc.* **1982**, *104*, 3228–3230; (b) Trost, B. M.; Martin, S. J. Alkynyl sulfonylation. A direct approach for nucleophilic addition and substitution of olefins by carbanions. *J. Am. Chem. Soc.* **1984**, *106*, 4263–4265.
9. (a) Denmark, S. E.; Jaunet, A., Catalytic, Enantioselective, Intramolecular Carbosulfonylation of Olefins. *J. Am. Chem. Soc.* **2013**, *135*, 6419–6422; (b) Denmark, S. E.; Jaunet, A., Catalytic, Enantioselective, Intramolecular Carbosulfonylation of Olefins. Preparative and Stereochemical Aspects. *J. Org. Chem.* **2014**, *79*, 140–171; (c) Denmark, S. E.; Hartmann, E.; Kornfilt, D. J.; Wang, H. Mechanistic, crystallographic, and computational studies on the catalytic, enantioselective sulfonylfunctionalization of alkenes. *Nat. Chem.* **2014**, *6*, 1056–1064; (d) Denmark, S. E.; Chi, H. M., Catalytic, Enantioselective, Intramolecular Carbosulfonylation of Olefins. Mechanistic Aspects: A Remarkable Case of Negative Catalysis. *J. Am. Chem. Soc.* **2014**, *136*, 3655–3663; (e) Tao, Z.; Robb, K. A.; Panger, J. L.; Denmark, S. E., Enantioselective, Lewis Base-Catalyzed Carbosulfonylation of Alkenylboronates by 1,2-Boronate Migration. *J. Am. Chem. Soc.* **2018**, *140*, 15621–15625; (f) Tao, Z.; Robb, K. A.; Zhao, K.; Denmark, S. E., Enantioselective, Lewis Base-Catalyzed Sulfenocyclization of Polyenes. *J. Am. Chem. Soc.* **2018**, *140*, 3569–3573; (g) Matviitsuk, A.; Panger, J. L.; Denmark, S. E., Catalytic, Enantioselective Sulfenofunctionalization of Alkenes: Development and Recent Advances. *Angew. Chem. Int. Ed.* **2020**, *59*, 19796–19819.
10. (a) Luo, J.; Cao, Q.; Cao, X.; Zhao, X., Selenide-catalyzed Enantioselective Synthesis of Trifluoromethylthiolated Tetrahydronaphthalenes by Merging Desymmetrization and Trifluoromethylthiolation. *Nat. Commun.* **2018**, *9*, 527; (b) Song, X. F.; Ye, A. H.; Xie, Y. Y.; Dong, J. W.; Chen, C.; Zhang, Y.; Chen, Z. M., Lewis-Acid-Mediated Thiocyanato Semipinacol Rearrangement of Allylic Alcohols for Construction of alpha-Quaternary Center beta-Thiocyanato Carbonyls. *Org. Lett.* **2019**, *21*, 9550–9554; (c) Xie, Y. Y.; Chen, Z. M.; Luo, H. Y.; Shao, H.; Tu, Y. Q.; Bao, X.; Cao, R. F.; Zhang, S. Y.; Tian, J. M., Lewis Base/Bronsted Acid Co-catalyzed Enantioselective Sulfonylation/Semipinacol Rearrangement of Di- and Trisubstituted Allylic Alcohols. *Angew. Chem. Int. Ed.* **2019**, *58*, 12491–12496; (d) Tang, M.; Han, S.; Huang, S.; Huang, S.; Xie, L. G. Carbosulfonylation of alkenes with organozinc reagents and dimethyl(methylthio)sulfonium trifluoromethanesulfonate. *Org. Lett.* **2020**, *22*, 9729–9734.
11. Lin, S.; Jacobsen, E. N. Thiourea-catalysed ring opening of episulfonium ions with indole derivatives by means of stabilizing non-covalent interactions. *Nat. Chem.* **2012**, *4*, 817–824.
12. (a) Denmark, S. E.; Chi, H. M. Lewis base catalyzed, enantioselective, intramolecular sulfenoamination of olefins. *J. Am. Chem. Soc.* **2014**, *136*, 8915–8918; (b) Denmark, S. E.; Rossi, S.; Webster, M. P.; Wang, H., Catalytic, Enantioselective Sulfonylation of Ketone-derived Enoxysilanes. *J. Am. Chem. Soc.* **2014**, *136*, 13016–13028; (c) Matviitsuk, A.; Denmark, S. E., Enantio- and Diastereoselective, Lewis Base Catalyzed,

Cascade Sulfenoacetalization of Alkenyl Aldehydes. *Angew. Chem. Int. Ed.* **2019**, *58*, 12486–12490; (d) Roth, A.; Denmark, S. E., Enantioselective, Lewis Base-Catalyzed, Intermolecular Sulfenoamination of Alkenes. *J. Am. Chem. Soc.* **2019**, *141*, 13767–13771.

13. (a) Derosa, J.; Apolinar, O.; Kang, T.; Tran, V. T.; Engle, K. M. Recent developments in nickel-catalyzed intermolecular dicarbofunctionalization of alkenes. *Chem. Sci.* **2020**, *11*, 4287–4296; (b) Diccianni, J.; Lin, Q.; Diao, T. Mechanisms of nickel-catalyzed coupling reactions and applications in alkene functionalization. *Acc. Chem. Res.* **2020**, *53*, 906–919. (c) Qi, X.; Diao, T., Nickel-Catalyzed Dicarbofunctionalization of Alkenes. *ACS Catal.* **2020**, *10*, 8542–8556.

14. (a) Giri, R.; Kc, S., Strategies toward Dicarbofunctionalization of Unactivated Olefins by Combined Heck Carbometalation and Cross-Coupling. *J. Org. Chem.* **2018**, *83*, 3013–3022; (b) Zhu, S.; Zhao, X.; Li, H.; Chu, L. Catalytic three-component dicarbofunctionalization reactions involving radical capture by nickel. *Chem. Soc. Rev.* **2021**, *50*, 10836–10856; (c) Wickham, L. M.; Giri, R., Transition Metal (Ni, Cu, Pd)-Catalyzed Alkene Dicarbofunctionalization Reactions. *Acc. Chem. Res.* **2021**, *54*, 3415–3437.

15. (a) Derosa, J.; Tran, V. T.; Boulous, M. N.; Chen, J. S.; Engle, K. M. Nickel-catalyzed  $\beta,\gamma$ -dicarbofunctionalization of alkenyl carbonyl compounds via conjunctive cross-coupling. *J. Am. Chem. Soc.* **2017**, *139*, 10657–10660; (b) Li, W.; Boon, J. K.; Zhao, Y. Nickel-catalyzed difunctionalization of allyl moieties using organoboronic acids and halides with divergent regioselectivities. *Chem. Sci.* **2018**, *9*, 600–607; (c) Derosa, J.; Kleinmans, R.; Tran, V. T.; Karunandana, M. K.; Wisniewski, S. R.; Eastgate, M. D.; Engle, K. M. Nickel-catalyzed 1,2-diarylation of simple alkenyl amides. *J. Am. Chem. Soc.* **2018**, *140*, 17878–17883; (d) Basnet, P.; Kc, S.; Dhungana, R. K.; Shrestha, B.; Boyle, T. J.; Giri, R. Synergistic bimetallic Ni/Ag and Ni/Cu catalysis for regioselective  $\gamma,\delta$ -diarylation of alkenyl ketimines: addressing  $\beta$ -H elimination by *in situ* generation of cationic Ni(II) catalysts. *J. Am. Chem. Soc.* **2018**, *140*, 15586–15590.

16. (a) Logan, K. M.; Sardini, S. R.; White, S. D.; Brown, M. K. Nickel-catalyzed stereoselective arylation of unactivated alkenes. *J. Am. Chem. Soc.* **2018**, *140*, 159–162; (b) Derosa, J.; van der Puyl, V. A.; Tran, V. T.; Liu, M. Y.; Engle, K. M. Directed nickel-catalyzed 1,2-dialkylation of alkenyl carbonyl compounds. *Chem. Sci.* **2018**, *9*, 5278–5283; (c) van der Puyl, V. A.; Derosa, J.; Engle, K. M. Directed, Nickel-catalyzed umpolung 1,2-carboamination of alkenyl carbonyl compounds. *ACS Catal.* **2019**, *9*, 224–229; (d) Kang, T.; Kim, N.; Cheng, P. T.; Zhang, H.; Foo, K.; Engle, K. M. Nickel-catalyzed 1,2-carboamination of alkenyl alcohols. *J. Am. Chem. Soc.* **2021**, *143*, 13962–13970; (e) Dhungana, R. K.; Aryal, V.; Niroula, D.; Sapkota, R. R.; Lakomy, M. G.; Giri, R. Nickel-catalyzed regioselective alkenylarylation of  $\gamma,\delta$ -alkenyl ketones via carbonyl coordination. *Angew. Chem. Int. Ed.* **2021**, *60*, 19092–19096. (f) Kang, T.; González, J. M.; Li, Z.-Q.; Foo, K.; Cheng, P. T. W.; Engle, K. M. Alkene Difunctionalization Directed by Free Amines: Diamine Synthesis via Nickel-Catalyzed 1,2-Carboamination. *ACS Catal.* **2022**, DOI: 10.1021/acscatal.2c00373.

17. (a) Sugoh, K.; Kuniyasu, H.; Sugae, T.; Ohtaka, A.; Takai, Y.; Tanaka, A.; Machino, C.; Kambe, N.; Kurosawa, H. A prototype of transition-metal-catalyzed carbothiolation of alkynes. *J. Am. Chem. Soc.* **2001**, *123*, 5108–5109; (b) Hooper, J. F.; Chaplin, A. B.; Gonzalez-Rodriguez, C.; Thompson, A. L.; Weller, A. S.; Willis, M. C. Aryl methyl sulfides as substrates for rhodium-catalyzed alkyne carbothiolation: arene functionalization with activating group recycling. *J. Am. Chem. Soc.* **2012**, *134*, 2906–2909; (c) Iwasaki, M.; Topolovcan, N.; Hu, H.; Nishimura, Y.; Gagnot, G.; Nakorn, R. N.; Yuvacharaskul, R.; Nakajima, K.; Nishihara, Y. Palladium-catalyzed regio- and stereoselective carbothiolation of terminal alkynes with azolyl sulfides. *Org. Lett.* **2016**, *18*, 1642–1645.

18. Hosoya, Y.; Kobayashi, I.; Mizoguchi, K.; Nakada, M. Palladium-catalyzed carbothiolation via trapping of the  $\sigma$ -alkyl palladium intermediate with RSTIPS. *Org. Lett.* **2019**, *21*, 8280–8284.

19. (a) Furuya, T.; Kamlet, A. S.; Ritter, T. Catalysis for fluorination and trifluoromethylation. *Nature* **2011**, *473*, 470–477; (b) Umemoto, T.; Yang, Y.; Hammond, G. B. Development of N-F fluorinating agents and their fluorinations: Historical perspective. *Beilstein J. Org. Chem.* **2021**, *17*, 1752–1813.

20. Shao, X.; Xu, C.; Lu, L.; Shen, Q. Shelf-stable electrophilic reagents for trifluoromethylthiolation. *Acc. Chem. Res.* **2015**, *48*, 1227–1236.

21. Anbarasan, P.; Neumann, H.; Beller, M. A general rhodium-catalyzed cyanation of aryl and alkenyl boronic acids. *Angew. Chem. Int. Ed.* **2011**, *50*, 519–522.

22. Hie, L.; Fine Nathel, N. F.; Shah, T. K.; Baker, E. L.; Hong, X.; Yang, Y. F.; Liu, P.; Houk, K. N.; Garg, N. K. Conversion of amides to esters by the nickel-catalysed activation of amide C–N bonds. *Nature* **2015**, *524*, 79–83.

23. Apolinar, O.; Tran, V. T.; Kim, N.; Schmidt, M. A.; Derosa, J.; Engle, K. M., Sulfonamide directivity enables Ni-catalyzed 1,2-diarylation of diverse alkenyl amines. *ACS Catal.* **2020**, *10*, 14234–14239.

24. Mao, R.; Bera, S.; Turla, A. C.; Hu, X. Copper-catalyzed intermolecular functionalization of unactivated C(sp<sup>3</sup>)–H bonds and aliphatic carboxylic acids. *J. Am. Chem. Soc.* **2021**, *143*, 14667–14675.

25. Meyer, D.; Jangra, H.; Walther, F.; Zippe, H.; Renaud, P. A third generation of radical fluorinating agents based on N-fluoro-N-arylsulfonamides. *Nat. Commun.* **2018**, *9*, 4888.

26. Yang, J. D.; Wang, Y.; Xue, X. S.; Cheng, J. P. A Systematic evaluation of the N-F bond strength of electrophilic N-F reagents: hints for atomic fluorine donating ability. *J. Org. Chem.* **2017**, *82*, 4129–4135.

27. Szostak, M.; Aube, J. Chemistry of bridged lactams and related heterocycles. *Chem. Rev.* **2013**, *113*, 5701–5765.

28. Schmidt, M. A.; Stokes, R. W.; Davies, M. L.; Roberts, F. 4-Cyanobenzenesulfonamides: amine synthesis and protecting strategy to compliment the nosyl group. *J. Org. Chem.* **2017**, *82*, 4550–4560.

29. Tran, V. T.; Li, Z.-Q.; Gallagher, T. J.; Derosa, J.; Liu, P.; Engle, K. M. Integrating allyl electrophiles into nickel-catalyzed conjunctive cross-coupling. *Angew. Chem. Int. Ed.* **2020**, *59*, 7029–7034.

30. (a) Tran, V. T.; Li, Z.-Q.; Apolinar, O.; Derosa, J.; Joannou, M. V.; Wisniewski, S. R.; Eastgate, M. D.; Engle, K. M. Ni(COD)(DQ): An air-stable 18-electron nickel(0)-olefin precatalyst. *Angew. Chem. Int. Ed.* **2020**, *59*, 7409–7413. (b) Tran, V. T.; Kim, N.; Rubel, C. Z.; Wu, X.; Kang, T.; Jankins, T. C.; Li, Z.-Q.; Joannou, M. V.; Ayers, S.; Gembicky, M.; Bailey, J.; Sturgell, E. J.; Sanchez, B. B.; Chen, J. S.; Lin, S.; Eastgate, M. D.; Wisniewski, S. R.; Engle, K. M. Structurally Diverse Bench-Stable Nickel(0) Pre-Catalysts: A Practical Toolkit for In Situ Ligation Protocols. Submission Date: 2022-02-11. *ChemRxiv* **2022**, DOI: 10.26434/chemrxiv-2022-7zvh. (accessed 2022-03-08).


31. O'Broin, C. Q.; Fernández, P.; Martínez, C.; Muñiz, K. N-Iodosuccinimide-promoted Hofmann–Löffler reactions of sulfonimides under visible light. *Org. Lett.* **2016**, *18*, 436–439.

32. Freskos, J. N.; Mischke, B. V.; DeCrescenzo, G. A.; Heintz, R.; Getman, D. P.; Howard, S. C.; Kishore, N. N.; McDonald, J. J.; Munie, G. E.; Rangwala, S.; Swearingen, C. A.

Voliva, C.; Welsch, D. J. Discovery of a novel series of selective MMP inhibitors: Identification of the  $\gamma$ -sulfone-thiols. *Bioorg. Med. Chem. Lett.* **1999**, *9*, 943–948.

33. (a) Kc, S.; Dhungana, R. K.; Shrestha, B.; Thapa, S.; Khanal, N.; Basnet, P.; Lebrun, R. W.; Giri, R., Ni-Catalyzed Regioselective Alkylation of Vinylarenes via  $C(sp^3)-C(sp^3)/C(sp^3)-C(sp^2)$  Bond Formation and Mechanistic Studies. *J. Am. Chem. Soc.* **2018**, *140*, 9801–9805; (b) Lin, Q.;

Diao, T., Mechanism of Ni-Catalyzed Reductive 1,2-Dicarbofunctionalization of Alkenes. *J. Am. Chem. Soc.* **2019**, *141*, 17937–17948; (c) Dhungana, R. K.; Sapkota, R. R.; Wickham, L. M.; Niroula, D.; Shrestha, B.; Giri, R., Ni-Catalyzed Arylbenzylolation of Alkenylarenes: Kinetic Studies Reveal Autocatalysis by  $ZnX_2$ . *Angew. Chem. Int. Ed.* **2021**, *60*, 22977–22982.



Insert Table of Contents artwork here

---