

Isolation of cuspidal spectrum, with application to the Gan–Gross–Prasad conjecture

By RAPHAËL BEUZART-PLESSIS, YIFENG LIU, WEI ZHANG, and XINWEN ZHU

Abstract

We introduce a new technique for isolating components on the spectral side of the trace formula. By applying it to the Jacquet–Rallis relative trace formula, we complete the proof of the global Gan–Gross–Prasad conjecture and its refinement Ichino–Ikeda conjecture for $U(n) \times U(n+1)$ in the stable case.

Contents

1. Introduction	520
1.1. Isolation of cuspidal spectrum	520
1.2. Application to the Gan–Gross–Prasad conjecture	522
1.3. Strategy of proofs	525
1.4. Notation and conventions	528
2. Multipliers of Schwartz convolution algebra	530
2.1. Preliminaries on holomorphic functions	530
2.2. Multiplier functions	531
2.3. Multipliers of Schwartz algebra	533
2.4. Proof of Theorem 2.13	536
3. Isolation of spectrum	547
3.1. Statement of results	547
3.2. Cuspidal data and cuspidal components	549
3.3. Annihilation of all but finitely many components	551
3.4. Proof of results	556
4. Application to the Gan–Gross–Prasad conjecture	560
4.1. Jacquet–Rallis relative trace formulae, d’après Zydor	560
4.2. Smooth transfer of Schwartz test functions	562

Keywords: trace formula, multipliers, isolation of spectrum, cuspidal automorphic representations, Gan–Gross–Prasad conjecture

AMS Classification: Primary: 11F67, 11F70, 11F72.

© 2021 Department of Mathematics, Princeton University.

4.3. Weak automorphic base change	568
4.4. Gan–Gross–Prasad and Ichino–Ikeda conjectures	573
Appendix A. Extending a result of Delorme to reductive groups	577
References	580

1. Introduction

1.1. *Isolation of cuspidal spectrum.* Let F be a number field and G a connected reductive group over F . In this subsection, we describe a general method for annihilating the non-cuspidal spectrum, while keeping certain prescribed representations of $G(\mathbb{A}_F)$. We fix an open compact subgroup K of $G(\mathbb{A}_F^\infty)$. Denote by $\mathcal{S}(G(\mathbb{A}_F))_K$ the space of bi- K -invariant Schwartz test functions on $G(\mathbb{A}_F)$; that is, we allow the archimedean component to be a Schwartz function rather than just a compactly supported smooth function. Recall that a Schwartz function on $G(F_\infty)$ is a smooth function f such that Df is bounded for every algebraic differential operator D on $G(F_\infty)$.¹ Then $\mathcal{S}(G(\mathbb{A}_F))_K$ is an algebra under convolution and acts continuously on the L^2 -spectrum $L^2(G(F)\backslash G(\mathbb{A}_F)/K)$ via the right regular representation R .

THEOREM 1.1 (see Theorem 3.6 for a more general version). *Let $\pi = \otimes_v \pi_v$ be an irreducible admissible representation of $G(\mathbb{A}_F)$ with unitary automorphic central character ω . Suppose that there does not exist a pair (P, σ) , where P is a proper parabolic subgroup of G (defined over F) and σ is a cuspidal automorphic representation of $M(\mathbb{A}_F)$ with M the Levi quotient of P , such that π_v is a constituent of $\text{Ind}_P^G(\sigma_v)$ for all but finitely many places v of F . Then there is a multiplier $\mu \star$ of the algebra $\mathcal{S}(G(\mathbb{A}_F))_K$ such that for every $f \in \mathcal{S}(G(\mathbb{A}_F))_K$,*

- (1) $R(\mu \star f)$ maps $L^2(G(F)\backslash G(\mathbb{A}_F)/K, \omega)$ into $L_{\text{cusp}}^2(G(F)\backslash G(\mathbb{A}_F)/K, \omega)_\pi$ — the π -nearly isotypic subspace of $L_{\text{cusp}}^2(G(F)\backslash G(\mathbb{A}_F)/K, \omega)$, that is, the direct sum of K -invariants of irreducible subrepresentations $\tilde{\pi}$ such that $\tilde{\pi}_v \simeq \pi_v$ for all but finitely many places v of F ;
- (2) $\pi(\mu \star f) = \pi(f)$.

Recall that a multiplier of a complex linear algebra $\mathcal{S}$ is a complex linear operator $\mu \star: \mathcal{S} \rightarrow \mathcal{S}$ that commutes with both left and right multiplications. Finding multipliers is most interesting when $\mathcal{S}$, such as $\mathcal{S}(G(\mathbb{A}_F))_K$, is non-commutative and non-unital.

Remark 1.2. We have

¹Readers should not confuse it with a Harish-Chandra Schwartz function, which is defined by a much weaker condition.

- (1) [Theorem 1.1](#), together with its application to the stable case of the Gan–Gross–Prasad conjecture discussed in the next subsection, breaks the long-standing impression that to understand every (non-CAP) cuspidal automorphic representation, one has to understand the full L^2 -spectrum on the spectral side of the (relative) trace formula.
- (2) When G is anisotropic modulo center, $L^2(G(F)\backslash G(\mathbb{A}_F)/K, \omega)$ coincides with $L^2_{\text{cusp}}(G(F)\backslash G(\mathbb{A}_F)/K, \omega)$, which is a Hilbert direct sum of subspaces of the form $L^2_{\text{cusp}}(G(F)\backslash G(\mathbb{A}_F)/K, \omega)_{\pi'}$ for irreducible (cuspidal) automorphic representations π' of $G(\mathbb{A}_F)$ with central character ω and non-zero K -invariants up to near equivalence. In this case, [Theorem 1.1](#) implies that one can use multipliers to modify $f \in \mathcal{S}(G(\mathbb{A}_F))_K$ such that the effect is the same as composing the projection map to an arbitrarily given factor of the form $L^2_{\text{cusp}}(G(F)\backslash G(\mathbb{A}_F)/K, \omega)_{\pi}$.
- (3) It is crucial that we work with the algebra $\mathcal{S}(G(\mathbb{A}_F))_K$ of Schwartz test functions in [Theorem 1.1](#). The method does not work if one works with compactly supported test functions.
- (4) In [Theorem 1.1](#), we do not even require π to be cuspidal automorphic. For example, it is possible that $\pi^K \neq \{0\}$ but $L^2_{\text{cusp}}(G(F)\backslash G(\mathbb{A}_F)/K, \omega)_{\pi} = 0$. Then the theorem provides a uniform way to modify $f \in \mathcal{S}(G(\mathbb{A}_F))_K$, without changing its action on π^K , but annihilating the entire L^2 -spectrum.

In fact, we also obtain a result for isolating general cuspidal components of the L^2 -spectrum. For simplicity, here we only state the theorem for $G = \text{Res}_{F'/F} \text{GL}_n$ for a finite extension F'/F . For such G , we have the result on the classification of automorphic representations [JS81, Th. 4.4].

THEOREM 1.3 (special case of [Theorem 3.19](#)). *In the situation of [Theorem 1.1](#), suppose that $G = \text{Res}_{F'/F} \text{GL}_n$ and that π is an irreducible constituent of $\text{Ind}_P^G(\sigma)$ for a parabolic subgroup P of G and a cuspidal automorphic representation σ of the Levi quotient of P . Then there is a multiplier $\mu \star$ of the algebra $\mathcal{S}(G(\mathbb{A}_F))_K$ such that for every $f \in \mathcal{S}(G(\mathbb{A}_F))_K$,*

- (1) $R(\mu \star f)$ maps $L^2(G(F)\backslash G(\mathbb{A}_F)/K, \omega)$ into $L^2_{(M, \sigma)}(G(F)\backslash G(\mathbb{A}_F)/K, \omega)$, where the latter is the cuspidal component of $L^2(G(F)\backslash G(\mathbb{A}_F)/K, \omega)$ associated to (M, σ) ;
- (2) $\pi(\mu \star f) = \pi(f)$.

In the process of proving [Theorems 1.1](#) and [1.3](#), we need a sufficient supply for multipliers of the archimedean component of $\mathcal{S}(G(\mathbb{A}_F))_K$. Now we temporarily switch the notation so that $G = \underline{G}(\mathbb{R})$ for a connected reductive algebraic group $\underline{G}$ over $\mathbb{R}$. Let $\mathfrak{h}^*$ be the real vector space spanned by the weight lattice of the abstract Cartan group of $\underline{G}_{\mathbb{C}}$, and let W be the Weyl group of $\underline{G}_{\mathbb{C}}$, which acts on $\mathfrak{h}^*$. By the Harish-Chandra isomorphism, the infinitesimal character χ_{π} of an irreducible admissible representation π of G gives

rise to a W -orbit in $\mathfrak{h}_{\mathbb{C}}^*$. In [Definition 2.8](#), we will define a space $\mathcal{M}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)$ of holomorphic functions on $\mathfrak{h}_{\mathbb{C}}^*$ satisfying certain growth conditions, stable under the action of W . Let $\mathcal{S}(G)$ be the convolution algebra of Schwartz functions on G . The following is our theorem on multipliers of $\mathcal{S}(G)$.

THEOREM 1.4 (Theorem [2.13](#)). *For every function $\mu \in \mathcal{M}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$, there is a unique linear operator $\mu \star: \mathcal{S}(G) \rightarrow \mathcal{S}(G)$, such that*

$$\pi(\mu \star f) = \mu(\chi_{\pi}) \cdot \pi(f)$$

holds for every $f \in \mathcal{S}(G)$ and every irreducible admissible representation π of G . In particular, $\mu \star$ is a multiplier of $\mathcal{S}(G)$.

Now it is a good time to explain the crucial difference between $\mathcal{S}(G)$ and $C_c^\infty(G)$. As pointed out in the final remark of the article [\[Del84\]](#), if G has no compact factors, then the only W -invariant holomorphic functions on $\mathfrak{h}_{\mathbb{C}}^*$ that give rise to (continuous) multipliers of $C_c^\infty(G)$ are polynomials, that is, elements in the center of the universal enveloping algebra. Even if one considers only the subalgebra $C_c^\infty(G)_{(K)}$ of bi- K -finite compactly supported smooth functions for a fixed maximal compact subgroup K of G , the W -invariant holomorphic functions on $\mathfrak{h}_{\mathbb{C}}^*$ that give rise to (continuous) multipliers of $C_c^\infty(G)_{(K)}$ have to be of exponential type, a property not required for $\mathcal{S}(G)$. The removal of the restriction of being of exponential type will vastly increase the collection of multipliers, making it possible to obtain results like [Theorem 1.1](#), as long as the Casimir eigenvalues of cuspidal automorphic representations are distributed in a certain discrete way, while the latter is indeed fulfilled by a result of Donnelly [\[Don82\]](#).

1.2. Application to the Gan–Gross–Prasad conjecture. In this subsection, we describe the results obtained by applying [Theorem 1.1](#) and its variants to the Jacquet–Rallis relative trace formula [\[JR11\]](#), [\[Zyd20\]](#). Let E/F be a quadratic extension of number fields, with c the Galois involution. Let $n \geq 1$ be an integer.

Definition 1.5. We let Π be an isobaric automorphic representation of $\mathrm{GL}_n(\mathbb{A}_E)$.

- (1) We say that Π is *conjugate self-dual* if its contragredient $\Pi^\vee$ is isomorphic to $\Pi \circ c$.
- (2) We say that Π is *hermitian* if Π is an isobaric sum of mutually non-isomorphic cuspidal automorphic representations $\Pi_1, \dots, \Pi_d$ in which each factor Π_i is conjugate self-dual and satisfies that $L(s, \Pi_i, \mathrm{As}^{(-1)^n})$ is regular at $s = 1$.

Let V be a (non-degenerate) hermitian space over E of rank n (with respect to the involution c). Put $G := \mathrm{U}(V)$, which is a reductive group over F .

Definition 1.6. Let π be an irreducible admissible representation of $G(\mathbb{A}_F)$. We say that an isobaric automorphic representation Π of $\mathrm{GL}_n(\mathbb{A}_E)$ is a *weak automorphic base change* of π if for all but finitely many places v of F split in E , the (split) local base change of π_v is isomorphic to Π_v . If weak automorphic base change of π exists, then it is unique up to isomorphism by [Ram18, Th. A]; we denote it by $\mathrm{BC}(\pi)$.

Remark 1.7. In most of the literature, weak automorphic base change of π requires the local-global compatibility for all but finitely many places, so in some sense our notion of weak automorphic base change should really be “very weak automorphic base change,” though by the endoscopic classification for unitary groups [Art13], [Mok15], [KMSW14], we now know that these two definitions are equivalent. The reason we use this weaker notion is that we want to make our argument independent of the endoscopic theory for unitary groups. In fact, under our weaker notion of automorphic base change, we can prove, as a byproduct of the Jacquet–Rallis relative trace formulae, the following result:

- The weak automorphic base change (in the sense of Definition 1.6) of π exists as long as there exist infinitely many places v of F split in E such that π_v is generic.

Our proof is completely free of using endoscopic trace formulae. The method can be used to show the local-global compatibility at all places where π_v is unramified as well, but the argument implicitly relies on [Mok15], [KMSW14]. See Remark 4.15 for more details.

The following theorem confirms the global Gan–Gross–Prasad conjecture [GGP12] for $\mathrm{U}(n) \times \mathrm{U}(n+1)$ in the stable case completely, which improves previous results in [Zha14b], [Xue19], [BP20a].

THEOREM 1.8. *Let Π_n and Π_{n+1} be hermitian cuspidal automorphic representations of $\mathrm{GL}_n(\mathbb{A}_E)$ and $\mathrm{GL}_{n+1}(\mathbb{A}_E)$, respectively. Then the following two statements are equivalent:*

- (1) *We have $L(\frac{1}{2}, \Pi_n \times \Pi_{n+1}) \neq 0$.*
- (2) *There exist*
 - *a hermitian space V_n over E of rank n , which gives another hermitian space $V_{n+1} := V_n \oplus E \cdot e$ over E of rank $n+1$ in which e has norm 1;*
 - *for $m = n, n+1$, an irreducible subrepresentation $\pi_m \subseteq \mathcal{A}_{\mathrm{cusp}}(G_m)$ of $G_m(\mathbb{A}_F)$, where $G_m := \mathrm{U}(V_m)$, satisfying $\mathrm{BC}(\pi_m) \simeq \Pi_m$, and a cusp form $\varphi_m \in \pi_m$,*

such that

$$\mathcal{P}(\varphi_n, \varphi_{n+1}) := \int_{G_n(F) \backslash G_n(\mathbb{A}_F)} \varphi_n(h) \varphi_{n+1}(h) \, dh \neq 0,$$

where dh is the Tamagawa measure on $G_n(\mathbb{A}_F)$.

Before our current work, the above theorem was known under the restriction that both Π_n and Π_{n+1} are supercuspidal at some prime of E that is split over F . However, removing this (last) local restriction is crucial for arithmetic application to certain motives like symmetric powers of elliptic curves; see [LTX⁺19].

Using the above theorem, we can obtain the following non-vanishing result on central L -values, which improves [Zha14b, Th. 1.2].

THEOREM 1.9. *Let Π_{n+1} be a hermitian cuspidal automorphic representation of $\mathrm{GL}_{n+1}(\mathbb{A}_E)$. Then there exists a hermitian cuspidal automorphic representation Π_n of $\mathrm{GL}_n(\mathbb{A}_E)$ such that*

$$L(\tfrac{1}{2}, \Pi_n \times \Pi_{n+1}) \neq 0.$$

The last application is the Ichino–Ikeda conjecture [II10], which is a refinement of the Gan–Gross–Prasad conjecture by giving an explicit formula for $|\mathcal{P}(\varphi_n, \varphi_{n+1})|^2$. In the case of $\mathrm{U}(n) \times \mathrm{U}(n+1)$, it is formulated in [Har14, Conj. 1.2] (see also [Zha14a, Conj. 1.1]). The following theorem confirms the Ichino–Ikeda conjecture for $\mathrm{U}(n) \times \mathrm{U}(n+1)$ in the stable case completely, which improves previous results in [Zha14a], [BP20a], [BP21b].

THEOREM 1.10. *Let the situation be as in (2) of Theorem 1.8. If, moreover, π_n and π_{n+1} are both everywhere tempered, then the Ichino–Ikeda conjecture holds for π_n and π_{n+1} .*

The proofs of Theorems 1.8, 1.9, and 1.10 will be given in Section 4.4.

Remark 1.11. The results on isolating components of the L^2 -spectrum obtained in this article can vastly simplify the computation on the spectral side toward the endoscopic case of the Gan–Gross–Prasad and the Ichino–Ikeda conjectures for $\mathrm{U}(n) \times \mathrm{U}(n+1)$ as well. Indeed, on the unitary side, by Theorems 3.6 and 4.14, it suffices to understand the π -nearly isotypic subspace of the L^2 -spectrum, which is contained in the *cuspidal* spectrum; on the general linear side, by Theorem 3.19, it suffices to understand the *single* cuspidal component of the L^2 -spectrum that corresponds to $\mathrm{BC}(\pi)$.²

Remark 1.12. Using similar ideas, one can improve the results on the Gan–Gross–Prasad conjecture and the Ichino–Ikeda conjecture for $\mathrm{U}(n) \times \mathrm{U}(n)$ previously obtained by Hang Xue [Xue14], [Xue16], based on the relative trace formulae developed in [Liu14], after establishing analogous results in [BP21b], [CZ21], [Zyd20].

²In fact, during the referee process of the current article, the endoscopic case has already been worked out in [BPCZ20].

1.3. *Strategy of proofs.* The main part of the article has three sections, responsible for three clusters of results: existence of multipliers of Schwartz convolution algebras, isolation of cuspidal components in the L^2 -spectrum, and confirmation of the Gan–Gross–Prasad conjecture in the stable case and related results, respectively. In this subsection, we will briefly explain the strategy of proving these results.

In [Section 2](#), we prove [Theorem 1.4](#). First, we give a slightly more precise description of the space $\mathcal{M}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)$. Let $(X^*, \Phi, X_*, \Phi^\vee)$ be the root datum of $\underline{G}_{\mathbb{C}}$ so that $\mathfrak{h}^* = X^* \otimes_{\mathbb{Z}} \mathbb{R}$. For every automorphism ϑ of $(X^*, \Phi, X_*, \Phi^\vee)$ of order at most two, we can formulate two types of growth conditions — *moderate growth* and *rapid decay* with respect to ϑ — for holomorphic functions on $\mathfrak{h}_{\mathbb{C}}^*$. (See [Definition 2.8](#) for the accurate definition.) In particular, when $\vartheta = 1$, they coincide with the usual notions of moderate growth and rapid decay on vertical strips, respectively. The (inner form class) of $\underline{G}$ gives rise to a set θ of automorphisms of $(X^*, \Phi, X_*, \Phi^\vee)$ of order at most two, stable under W -conjugation. Then we define $\mathcal{M}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)$ (resp. $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)$) to be the space of holomorphic functions on $\mathfrak{h}_{\mathbb{C}}^*$ that have moderate growth (resp. rapid decay) with respect to all elements in $\theta \cup \{1\}$. It will be clear from the definition (and suggested by the terminology) that $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*) \subseteq \mathcal{M}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)$. We will first prove [Theorem 1.4](#) for the smaller space $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$ ([Proposition 2.24](#)), and then use a limit process to pass to $\mathcal{M}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$.

For $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$, note that $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)$ is contained in $\mathcal{N}_{\{1\}}(\mathfrak{h}_{\mathbb{C}}^*)$, the space of holomorphic functions on $\mathfrak{h}_{\mathbb{C}}^*$ that have rapid decay on vertical strips. Essentially by a result of Delorme ([Proposition A.1](#)), elements in $\mathcal{N}_{\{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$ will give multipliers for the subalgebra $\mathcal{S}(G)_{(K)}$ of $\mathcal{S}(G)$ of bi- K -finite Schwartz functions. Therefore, to construct $\mu \star f$ for $\mu \in \mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$ and $f \in \mathcal{S}(G)$, we may choose a sequence $\{f_n\} \subseteq C_c^\infty(G)_{(K)}$ approaching f in $\mathcal{S}(G)$ and show that $\{\mu \star f_n\}$ converges. It is a crucial observation that the Fréchet topology of $\mathcal{S}(G)$ is also induced from the semi-norms $f \mapsto \|Df\|_{L^2}$ for all algebraic differential operators D on G . Thus, it suffices for us to show that $\|D(\mu \star -)\|_{L^2}$ is continuous on the subspace $C_c^\infty(G)_{(K)}$ with respect to the subspace topology for all D . When D is (left and right) invariant, we have $D(\mu \star f) = \mu \star (Df)$, hence the continuity is clear. It remains to consider the case where D is a polynomial on G , or equivalently, a matrix coefficient of a finite dimensional algebraic representation of G . In this case, we can show that there are finitely many pairs (S_i, L_i) , depending on the matrix coefficient presentation of D only, in which S_i is a linear operator of $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$ and L_i is an algebraic differential operator of $\mathcal{S}(G)$ preserving $C_c^\infty(G)_{(K)}$, such that $D(\mu \star f) = \sum_i S_i(\mu) \star L_i(f)$. Thus, we obtain the desired continuity for $\|D(\mu \star -)\|_{L^2}$, hence [Theorem 1.4](#) for the smaller space $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$ is confirmed. However, in order to pass to

$\mathcal{M}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$, we will need more precise bounds on the linear operators S_i with respect to a certain natural family of semi-norms on $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)$, as stated in [Lemma 2.20](#).

In [Section 3](#), we prove all results related to the isolation of the L^2 -spectrum. To make the discussion here more explicit, we restrict ourselves to the number field $F = \mathbb{Q}$ and the group $G = \mathrm{PGL}_{n,\mathbb{Q}}$ with $n \geq 2$, and we only consider [Theorem 1.1](#) with π cuspidal automorphic. In particular, $Z = 1$, and we will omit the necessarily trivial character ω in the following notation. For every tuple $\underline{n} = (n_1, n_2, \dots)$ of integers $0 < n_1 \leq n_2 \leq \dots$ summing up to n , we let $M_{\underline{n}}$ be the standard diagonal Levi subgroup of G of block sizes $\underline{n}$ and $W_{\underline{n}}$ the subgroup of the Weyl group stabilizing $M_{\underline{n}}$. Then we have the coarse Langlands decomposition

$$L^2(G(\mathbb{Q}) \backslash G(\mathbb{A})/K) = \widehat{\bigoplus_{(M,\sigma)} L^2_{(M,\sigma)}(G(\mathbb{Q}) \backslash G(\mathbb{A})/K)},$$

where $M = M_{\underline{n}}$ for some $\underline{n}$ and σ is a cuspidal automorphic representation of $M(\mathbb{A})$ up to twist and $W_{\underline{n}}$ -conjugation. Our goal, in view of the strong multiplicity one property, is to find μ such that for every $f \in \mathcal{S}(G(\mathbb{A}))_K$, $\mu \star f$ annihilates all components but $L^2_{(G,\pi)}(G(\mathbb{Q}) \backslash G(\mathbb{A})/K)$ and maintains the action of f on π .

For the first step, it is not hard to construct a function $\mu_{\infty}^0 \in \mathcal{M}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$ (notation with respect to the real reductive group $G \otimes_{\mathbb{Q}} \mathbb{R}$) satisfying $\mu_{\infty}^0(\chi_{\pi_{\infty}}) = 1$ and the following condition: there exists a finite set $\mathfrak{T}$ of K_{∞} -types, where K_{∞} is the standard maximal compact subgroup of $G(\mathbb{R}) = \mathrm{PGL}_n(\mathbb{R})$, such that for every $f \in \mathcal{S}(G(\mathbb{R}))$, $\mu_{\infty}^0 \star f$ annihilates the component $L^2_{(M,\sigma)}(G(\mathbb{Q}) \backslash G(\mathbb{A})/K)$ if it has no K_{∞} -types in $\mathfrak{T}$. This will exclude all but finitely many components $L^2_{(M,\sigma)}(G(\mathbb{Q}) \backslash G(\mathbb{A})/K)$ when $M \in \{M_{(1,\dots,1)}, M_{(n)}\}$. However, for other M , there are still infinitely many components remaining.

The second step is to annihilate all but finitely many components

$$L^2_{(M,\sigma)}(G(\mathbb{Q}) \backslash G(\mathbb{A})/K)$$

with a K_{∞} -type in $\mathfrak{T}$ for every M . To explain the idea, we consider the simplest non-trivial case where $M = M_{(1,\dots,1,2)}$ (hence $n \geq 3$). It is easy to see that there exist a finite set $\mathfrak{T}_M$ of $(K_{\infty} \cap M(\mathbb{R}))$ -types and an open compact subgroup $K_M \subseteq M(\mathbb{A}^{\infty})$ such that if $L^2_{(M,\sigma)}(G(\mathbb{Q}) \backslash G(\mathbb{A})/K)$ has a K_{∞} -type in $\mathfrak{T}$, then σ must have a $(K_{\infty} \cap M(\mathbb{R}))$ -type in $\mathfrak{T}_M$ and non-trivial K_M -invariants. Note that there are infinitely many such σ up to twist and $W_{(1,\dots,1,2)}$ -conjugation! However, we observe that the Casimir operator for the derived subgroup of M , which is simply $\mathrm{SL}_{2,\mathbb{Q}}$, gives a polynomial function λ on $\mathfrak{h}_{\mathbb{C}}^*$. It is a well-known result of Harish-Chandra that for any given $\lambda_0 \in \mathbb{C}$, there are only finitely many cuspidal automorphic representations σ of $M(\mathbb{A})$ up to twist that have a $(K_{\infty} \cap M(\mathbb{R}))$ -type in $\mathfrak{T}_M$ and non-trivial K_M -invariants, such that $\lambda(\chi_{\sigma_{\infty}}) = \lambda_0$. By another well-known result on the distribution of Casimir eigenvalues of SL_2

(or [Don82] for general semisimple groups), one can find a holomorphic function ν_M on $\mathfrak{h}_{\mathbb{C}}^*$ that has moderate growth on vertical strips with zeros exactly $\lambda^{-1}(\Lambda \setminus \lambda(\chi_{\pi_{\infty}}))$, where $\Lambda \subseteq \mathbb{C}$ is the set of Casimir eigenvalues of cuspidal automorphic representations σ of $M(\mathbb{A})$ that have a $(K_{\infty} \cap M(\mathbb{R}))$ -type in $\mathfrak{T}_M$ and non-trivial K_M -invariants.³ Using ν_M , it is not hard to construct an element $\mu_{\infty}^M \in \mathcal{M}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$ such that for every $f \in \mathcal{S}(G(\mathbb{R}))$, $\mu_{\infty}^M \star f$ annihilates all but finitely components $L_{(M,\sigma)}^2(G(\mathbb{Q}) \backslash G(\mathbb{A})/K)$ with given $M = M_{(1,\dots,1,2)}$ and maintains the action of f on π . In fact, we can achieve this for every M .

The last step is to annihilate, without changing π , every single component $L_{(M,\sigma)}^2(G(\mathbb{Q}) \backslash G(\mathbb{A})/K)$ that is not isomorphic to π . This is easy if $M = M_{(n)} = G$, since we can use spherical Hecke operators at unramified (non-archimedean) places. When $M \neq G$, $L_{(M,\sigma)}^2(G(\mathbb{Q}) \backslash G(\mathbb{A})/K)$ is a “continuous” space of induced representations. However, there is a secret correlation between (non-archimedean) Hecke eigenvalues and (archimedean) infinitesimal characters for all representations that contribute to $L_{(M,\sigma)}^2(G(\mathbb{Q}) \backslash G(\mathbb{A})/K)$. This motivates us to construct, for every component $L_{(M,\sigma)}^2(G(\mathbb{Q}) \backslash G(\mathbb{A})/K)$ that is not isomorphic to π , a multiplier $\mu_{(M,\sigma)}$ that is “mixed” from $\mathcal{M}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$ and spherical Hecke operators, such that for every $f \in \mathcal{S}(G(\mathbb{A}))_K$, $\mu_{(M,\sigma)} \star f$ annihilates $L_{(M,\sigma)}^2(G(\mathbb{Q}) \backslash G(\mathbb{A})/K)$ and maintains the action of f on π . We remark that this step is inspired by the work [LV07].

To conclude the proof of Theorem 1.1, we only need to take the product of μ_{∞}^0 , $\{\mu_{\infty}^M\}_M$, and (finitely many) $\mu_{(M,\sigma)}$.

In Section 4, we prove the Gan–Gross–Prasad conjecture in the stable case and other related results. Here, we explain how to apply Theorem 1.1 to the Jacquet–Rallis relative trace formulae to attack the Gan–Gross–Prasad conjecture. The Jacquet–Rallis relative trace formulae have two sides: the group $G' := \text{Res}_{E/F} \text{GL}_{n,E} \times \text{Res}_{E/F} \text{GL}_{n+1,E}$, and the group $G^V := \text{U}(V_n) \times \text{U}(V_{n+1})$ where $V = (V_n, V_{n+1})$ is a pair of hermitian spaces over E as in Theorem 1.8. The two sides share the same space of orbits, which is an affine variety B over F ; in other words, we have surjective morphisms $G' \rightarrow B \leftarrow G^V$. By Zydor’s extension of the Jacquet–Rallis relative trace formulae, we have a relative trace formula on each side: For every $f' \in \mathcal{S}(G'(\mathbb{A}_F))$ that annihilates the entire non-cuspidal part of $L^2(G'(F) \backslash G'(\mathbb{A}_F), 1)$, we have the identity

$$\sum_{\Pi \text{ cuspidal}} I_{\Pi}(f') = \sum_{\gamma \in B(F)} I_{\gamma}(f'),$$

where I_{Π} and I_{γ} are certain invariant functionals on $\mathcal{S}(G'(\mathbb{A}_F))$ associated to a cuspidal automorphic representation Π and an element $\gamma \in B(F)$ defined

³However, we cannot require ν_M to be of exponential type at the same time.

via relative characters and (regularized) relative orbital integrals, respectively. Similarly, for every $f^V \in \mathcal{S}(G^V(\mathbb{A}_F))$ that annihilates the entire non-cuspidal part of $L^2(G^V(F) \backslash G^V(\mathbb{A}_F), 1)$, we have the identity

$$\sum_{\pi^V \text{ cuspidal}} J_{\pi^V}(f^V) = \sum_{\delta^V \in B(F)} J_{\delta^V}(f^V).$$

In practice, we have to consider all pairs V up to isomorphism. The starting point for the comparison of trace formulae is to find a pair of test functions $(f', \{f^V\}_V)$ that have *matching orbital integrals*, so that $I_\gamma(f') = J_{\delta^V}(f^V)$ when $\gamma = \delta^V$. The common strategy of finding $(f', \{f^V\}_V)$ that annihilate non-cuspidal spectra is to take f' such that f'_v annihilates all non-supercuspidal representations at some non-archimedean place v of F split in E , and similarly for f^V . However, by doing this, we necessarily annihilate all cuspidal automorphic representations that are nowhere supercuspidal. The new invention, which is enabled by [Theorem 1.1](#), is to modify an arbitrary pair $(f', \{f^V\}_V)$ of test functions that have matching orbital integrals by multipliers, so that the resulting test functions annihilate non-cuspidal spectra and maintain their actions on any prescribed representations $(\Pi, \{\pi^V\})$ in which Π is cuspidal automorphic and is isomorphic to $\mathrm{BC}(\pi^V)$. Thus, it is a natural question to find multipliers $(\mu', \{\mu^V\}_V)$ from [Theorem 1.1](#), such that $(\mu' \star f', \{\mu^V \star f^V\}_V)$ still have matching orbital integrals. The answer turns out to be quite elegant: there is a natural “base change” map from those multipliers for G^V (obtained in the way of [Theorem 1.1](#)) to those for G' ; and we show in [Proposition 4.8](#) that if μ' is the base change of μ^V for all V , then $(\mu' \star f', \{\mu^V \star f^V\}_V)$ have matching orbital integrals as long as $(f', \{f^V\}_V)$ do. Such multipliers $(\mu', \{\mu^V\}_V)$ are not hard to find. Therefore, we can compare the above two relative trace formulae without sacrificing any prescribed representations $(\Pi, \{\pi^V\})$ as above. The rest of the argument is a standard business in trace formulae approach.

The article has an appendix ([Appendix A](#)) in which we extend a result of Delorme to reductive groups, which is only used in the proof of [Proposition 2.24](#).

1.4. Notation and conventions.

- For a set S , we denote by $\mathbf{1}_S$ the characteristic function of S .
- In the main text, if we do not specify the base ring of a tensor product $\otimes$, then the base ring is $\mathbb{C}$.
- For a real vector space U , we put $U_{\mathbb{C}} := U \otimes_{\mathbb{R}} \mathbb{C}$, and $iU := U \otimes_{\mathbb{R}} i\mathbb{R}$, which is a subspace of the underlying real vector space of $U_{\mathbb{C}}$. We have the $\mathbb{R}$ -linear map $\mathrm{Re}: U_{\mathbb{C}} \rightarrow U$ by taking the real part.
- For a finite dimensional complex vector space U , we denote by $\mathcal{O}(U)$ the ring of holomorphic functions on U , $\mathcal{O}_{\mathrm{exp}}(U) \subseteq \mathcal{O}(U)$ the subring of holomorphic functions of exponential type, and $\mathbb{C}[U] \subseteq \mathcal{O}_{\mathrm{exp}}(U)$ the subring of

polynomial functions. A lattice (resp. full lattice) of U is a subgroup L of U such that the induced map $L \otimes_{\mathbb{Z}} \mathbb{C} \rightarrow U$ is injective (resp. bijective).

- For a complex linear algebra $\mathcal{S}$, we denote by $\text{Mul}(\mathcal{S})$ the $\mathbb{C}$ -algebra of multipliers of $\mathcal{S}$, that is, complex linear operators $\mu \star: \mathcal{S} \rightarrow \mathcal{S}$ satisfying $\mu \star (f * g) = (\mu \star f) * g = f * (\mu \star g)$ for every $f, g \in \mathcal{S}$, where $*$ denotes the multiplication in $\mathcal{S}$.
- By a prime of a number field, we mean a non-archimedean place. In [Sections 3 and 4](#), we will encounter various sets of places of a number field F . To summarize,
 - $\mathbf{S}$ will always be a finite set consisting of primes;
 - $\mathbf{T}$ will always be a (possibly infinite) set consisting of primes;
 - $\square$ will always be a finite set containing all archimedean places.
- For an algebraic group G over a number field F , we put $G_{\infty} := G(F \otimes_{\mathbb{Q}} \mathbb{R})$ for short.
- A subgroup of an algebraic group defined over a field is by default defined over the same field.
- Let P be a parabolic subgroup of a reductive group G .
 - We denote by $N_P \subseteq P$ the unipotent radical.
 - When σ is an admissible representation of $P(R)$ for an appropriate ring R for which admissibility makes sense, we denote by $\text{I}_P^G(\sigma)$ the normalized parabolic induction as an admissible representation of $G(R)$.
 - When σ is an admissible representation of $M(R)$ for a Levi subgroup M of P , we also write $\text{I}_P^G(\sigma)$ by regarding σ as a representation of $P(R)$ through inflation.

Acknowledgements. We would like to thank the American Institute of Mathematics for supporting three of us (Y. L., W. Z., and X. Z.) via the AIM SQuaREs project *Geometry of Shimura varieties and arithmetic application to L-functions* from 2017 to 2019, where an initial discussion toward this article happened in our last SQuaREs workshop in the spring of 2019. We thank Yichao Tian and Liang Xiao, the other two members of our SQuaREs project, since the project itself is a crucial motivation for the current article. We thank Jean-Loup Waldspurger for reading an early draft of the article and providing several useful comments. We thank the anonymous referee for careful reading and many helpful suggestions.

The project leading to this publication of R. B.-P. has received funding from Excellence Initiative of Aix-Marseille University-A*MIDEX, a French “Investissements d’Avenir” programme. The research of Y. L. is partially supported by the NSF grant DMS-1702019 and a Sloan Research Fellowship. The research of W. Z. is partially supported by the NSF grant DMS-1838118 and DMS-1901642. The research of X. Z. is partially supported by the NSF grant DMS-1902239 and a Simons Fellowship.

2. Multipliers of Schwartz convolution algebra

In this section, we construct sufficiently many multipliers of the convolution algebra of Schwartz functions for a real connected reductive algebraic group.

In [Section 2.1](#), we record some lemmas on constructing holomorphic functions on complex vector spaces with special properties. In [Section 2.2](#), we define various spaces of functions related to the multipliers of Schwartz convolution algebras. In [Section 2.3](#), we state our result ([Theorem 2.13](#)) on the existence of sufficiently many multipliers of Schwartz convolution algebras. In [Section 2.4](#), we prove [Theorem 2.13](#), following the strategy described in [Section 1.3](#).

2.1. Preliminaries on holomorphic functions. We first review some facts about entire functions on the complex plane. Recall that the order of an entire function $\Psi: \mathbb{C} \rightarrow \mathbb{C}$ is defined as

$$\inf\{e \in [0, +\infty) \mid \exists C_e > 0, \text{ such that } |\Psi(z)| < C_e \exp(|z|^e) \text{ for all } z \in \mathbb{C}\}.$$

Here, $|z| = \sqrt{z\bar{z}}$. If the above set is empty, then we say that Ψ has infinite order.

Next, we review the construction of entire functions with prescribed zeroes. Let Λ be a subset of $\mathbb{C}$, and let $p \geq 0$ be an integer. We define the (formal) Weierstrass product

$$\Psi_{\Lambda,p}(z) := z^\delta \prod_{\substack{\lambda \in \Lambda \\ \lambda \neq 0}} E_p(z/\lambda),$$

where $E_p(z) := (1-z)\exp(z + \cdots + z^p/p)$ is the elementary function, and $\delta = 1$ (resp. $\delta = 0$) if 0 belongs (resp. does not belong) to Λ .

Definition 2.1. Let $p \geq 0$ be an integer. We say that Λ has *rank* p if

- (1) Λ is countable;
- (2) p is the least non-negative integer such that $\sum_{\lambda \in \Lambda, \lambda \neq 0} |\lambda|^{-(p+1)}$ converges.

LEMMA 2.2. *Let $\Lambda \subseteq \mathbb{C}$ be a subset of rank p for some integer $p \geq 0$. Then $\Psi_{\Lambda,p}$ is a well-defined entire function of (finite) order at most $p+1$, with the set of zeroes exactly Λ .*

Proof. It is well known that there exist constants $C, C' > 0$ such that $|\Psi_{\Lambda,p}(z)| < C \exp(C'|z|^{p+1})$ for $z \in \mathbb{C}$. Thus, $\Psi_{\Lambda,p}$ is of finite order at most $p+1$ by definition. The set of zeroes is clear from the construction. $\square$

Now we consider a finite dimensional real vector space U .

Definition 2.3. Let ν be a holomorphic function on $U_{\mathbb{C}}$. We say that ν has *moderate vertical growth* if for every $M > 0$, there exists $r_M \in \mathbb{R}$ such that

$$\sup_{\|\operatorname{Re} z\| < M} |\nu(z)| \cdot (1 + \|z\|)^{r_M} < \infty$$

holds for some, hence every, Euclidean norm $\|\cdot\|$ on $U_{\mathbb{C}}$.

PROPOSITION 2.4. *Let $\lambda: U_{\mathbb{C}} \rightarrow \mathbb{C}$ be a polynomial function. For every entire function Ψ of finite order, there exists a holomorphic function ν on $U_{\mathbb{C}}$ that has moderate vertical growth, such that the set of zeroes of ν is exactly the inverse image of the set of zeroes of Ψ along λ .*

Proof. We choose an isomorphism $U \simeq \mathbb{R}^r$ and write elements in $U_{\mathbb{C}}$ as $z = (z_1, \dots, z_r)$. Suppose that Ψ is of order e and λ has degree d . Take an odd integer $q > de$. Define ν by the formula

$$\nu(z) := \exp(z_1^{2q} + \dots + z_r^{2q}) \cdot \Psi(\lambda(z))$$

for $z \in U_{\mathbb{C}}$, which is holomorphic. It is straightforward to check that $\nu(z)$ has moderate vertical growth. Moreover, we have $\nu(z) = 0$ if and only if $\lambda(z)$ is a zero of Ψ . The proposition follows. $\square$

COROLLARY 2.5. *Let $\lambda: U_{\mathbb{C}} \rightarrow \mathbb{C}$ be a polynomial function. For every subset $\Lambda \subseteq \mathbb{C}$ of finite rank, there exists a holomorphic function ν on $U_{\mathbb{C}}$ that has moderate vertical growth and whose set of zeroes is exactly $\lambda^{-1}\Lambda$.*

Proof. The proof follows from Lemma 2.2 and Proposition 2.4. $\square$

COROLLARY 2.6. *Let L be a lattice of $U_{\mathbb{C}}$ and $A \subseteq U_{\mathbb{C}}$ a finite subset. Then there exists a holomorphic function ν on $U_{\mathbb{C}}$ that has moderate vertical growth, vanishes on $L \setminus A$, and is nowhere vanishing on A .*

Proof. Let r be the dimension of U . We may choose linearly independent complex linear maps $\lambda_1, \dots, \lambda_r: U_{\mathbb{C}} \rightarrow \mathbb{C}$ such that L is contained in $\bigcap_{i=1}^r \lambda_i^{-1}\mathbb{Z}$. For every $1 \leq i \leq r$, the subset $\mathbb{Z} \setminus \lambda_i(A) \subseteq \mathbb{C}$ is of finite rank. By Corollary 2.5, we may find a holomorphic function ν_i on $U_{\mathbb{C}}$ that has moderate vertical growth, such that the set of zeroes of ν_i is exactly $\lambda_i^{-1}(\mathbb{Z} \setminus \lambda_i(A))$. Put $\nu := \prod_{i=1}^r \nu_i$. Then ν has moderate vertical growth, vanishes on all but finitely many elements in L , and is nowhere vanishing on A . Let $z_1, \dots, z_s$ be the finitely many elements in $L \setminus A$ at which ν is non-vanishing. For each $1 \leq j \leq s$, we may choose an affine function $l_j: U_{\mathbb{C}} \rightarrow \mathbb{C}$ such that $l_j(z_j) = 0$ and that l_j is nowhere vanishing on A . Then the holomorphic function $\nu \cdot \prod_{j=1}^s l_j$ satisfies the requirement in the corollary. $\square$

Remark 2.7. In fact, from the proof of Proposition 2.4, we may even require ν in Corollaries 2.5 and 2.6 to have exponential decay on vertical strips. However, we do not need this in what follows.

2.2. *Multiplier functions.* In this subsection, we introduce the spaces of multiplier functions that will give multipliers of the convolution algebra of Schwartz functions. We fix a root datum $(X^*, \Phi, X_*, \Phi^\vee)$ and

- let W be the Weyl group of $(X^*, \Phi, X_*, \Phi^\vee)$;

- let $\text{Aut}(\mathbf{X}^*, \Phi, \mathbf{X}_*, \Phi^\vee)^\heartsuit$ be the set of automorphisms of $(\mathbf{X}^*, \Phi, \mathbf{X}_*, \Phi^\vee)$ of order at most 2, which is finite and stable under the conjugation action of W ;
- put $\mathfrak{h}^* := \mathbf{X}^* \otimes_{\mathbb{Z}} \mathbb{R}$;
- for every $\vartheta \in \text{Aut}(\mathbf{X}^*, \Phi, \mathbf{X}_*, \Phi^\vee)^\heartsuit$, let $\mathfrak{h}_\vartheta^*$ (resp. $\mathfrak{h}_\vartheta^{*-}$) be the $+1$ -eigenspace (resp. -1 -eigenspace) of the action of ϑ on $\mathfrak{h}^*$; and let Y_ϑ^* be the projection of $\rho + \mathbf{X}^* \subseteq \mathfrak{h}^*$ onto $\mathfrak{h}_\vartheta^{*-}$, where ρ is the half sum of positive roots in Φ with respect to an arbitrary base of Φ .⁴

It is clear that Y_ϑ^* is a translation of a discrete subgroup of $\mathfrak{h}_\vartheta^{*-}$. Though Y_ϑ^* is not necessarily a subgroup, in what follows, we will sometimes write $H \oplus Y_\vartheta^*$ for a subgroup H of $\mathfrak{h}_{\vartheta, \mathbb{C}}^*$ as the subset of $\mathfrak{h}_\mathbb{C}^*$ consisting of elements of the form $\alpha + \varpi$ for $\alpha \in H$ and $\varpi \in Y_\vartheta^*$.

We take a subset $\theta \subseteq \text{Aut}(\mathbf{X}^*, \Phi, \mathbf{X}_*, \Phi^\vee)^\heartsuit$ that is stable under the conjugation action of W . In the following definition, we introduce several important spaces of holomorphic functions on $\mathfrak{h}_\mathbb{C}^*$ that will be related to multipliers of Schwartz convolution algebras.

Definition 2.8. We define several spaces of holomorphic functions on $\mathfrak{h}_\mathbb{C}^*$.

- (1) Define $\mathcal{M}_\theta(\mathfrak{h}_\mathbb{C}^*)$ to be the space of holomorphic functions μ on $\mathfrak{h}_\mathbb{C}^*$ such that for every $\vartheta \in \theta$ and every $M > 0$, there exists $r_{\vartheta, M} \in \mathbb{R}$ such that

$$(2.1) \quad \sup_{\substack{\alpha \in \mathfrak{h}_{\vartheta, \mathbb{C}}^*, \|\text{Re } \alpha\| < M \\ \varpi \in Y_\vartheta^*}} |\mu(\alpha + \varpi)| \cdot (1 + \|\alpha + \varpi\|)^{r_{\vartheta, M}} < \infty$$

holds for some, hence every, Euclidean norm $\|\cdot\|$ on $\mathfrak{h}_\mathbb{C}^*$.

- (2) Define $\mathcal{N}_\theta(\mathfrak{h}_\mathbb{C}^*)$ to be the space of holomorphic functions μ on $\mathfrak{h}_\mathbb{C}^*$ such that for every $\vartheta \in \theta$, every $M > 0$, and every $r \in \mathbb{R}$,

$$\sup_{\substack{\alpha \in \mathfrak{h}_{\vartheta, \mathbb{C}}^*, \|\text{Re } \alpha\| < M \\ \varpi \in Y_\vartheta^*}} |\mu(\alpha + \varpi)| \cdot (1 + \|\alpha + \varpi\|)^r < \infty$$

holds for some, hence every, Euclidean norm $\|\cdot\|$ on $\mathfrak{h}_\mathbb{C}^*$.

- (3) Define $\mathcal{M}_\theta^\sharp(\mathfrak{h}_\mathbb{C}^*)$ to be the subspace of $\mathcal{M}_{\theta \cup \{1\}}(\mathfrak{h}_\mathbb{C}^*)$ consisting of μ satisfying that for every $\vartheta \in \theta$, $\mu|_{\mathfrak{h}_{\vartheta, \mathbb{C}}^* + \varpi} = 0$ for all but finitely many elements $\varpi \in Y_\vartheta^*$.

When $\theta = \{1\}$, we suppress it in the subscripts in the above notation.

Remark 2.9. We have the following concerning [Definition 2.8](#):

- (1) In (1) (resp. (2)), for every $\vartheta \in \theta$ and every $\varpi \in Y_\vartheta^*$, the function $\alpha \mapsto \mu(\alpha + \varpi)$ has moderate vertical growth ([Definition 2.3](#)) (resp. has rapid decay on vertical strips) on $\mathfrak{h}_{\vartheta, \mathbb{C}}^*$. On the other hand, for every $\vartheta \in \theta$, if we

⁴It is easy to see that Y_ϑ^* does not depend on the choice of the base.

restrict the growth condition (2.1) to $i\mathfrak{h}_\vartheta^* \oplus \mathbf{Y}_\vartheta^*$, then it means that μ has polynomial growth.

- (2) The spaces $\mathcal{M}_\theta(\mathfrak{h}_\mathbb{C}^*)$, $\mathcal{N}_\theta(\mathfrak{h}_\mathbb{C}^*)$, and $\mathcal{M}_\theta^\sharp(\mathfrak{h}_\mathbb{C}^*)$ are all closed under multiplication and the action of $\mathbf{W}$.
- (3) If $\mu \in \mathcal{O}(\mathfrak{h}_\mathbb{C}^*)$ has moderate vertical growth, then $\mu \cdot \mathcal{M}_\theta^\sharp(\mathfrak{h}_\mathbb{C}^*) \subseteq \mathcal{M}_\theta^\sharp(\mathfrak{h}_\mathbb{C}^*)$. In particular, we have $\mathcal{M}(\mathfrak{h}_\mathbb{C}^*) \cdot \mathcal{M}_\theta^\sharp(\mathfrak{h}_\mathbb{C}^*) \subseteq \mathcal{M}_\theta^\sharp(\mathfrak{h}_\mathbb{C}^*)$. Note that $\mathcal{M}(\mathfrak{h}_\mathbb{C}^*) = \mathcal{M}_{\{1\}}(\mathfrak{h}_\mathbb{C}^*)$ according to Definition 2.8.

The following lemma demonstrates the existence of elements in $\mathcal{M}_\theta^\sharp(\mathfrak{h}_\mathbb{C}^*)^\mathbf{W}$.

LEMMA 2.10. *For every element $\alpha_0 \in \mathfrak{h}_\mathbb{C}^*$, there exists an element $\mu \in \mathcal{M}_\theta^\sharp(\mathfrak{h}_\mathbb{C}^*)^\mathbf{W}$ such that $\mu(\alpha_0) \neq 0$.*

Proof. For each $\vartheta \in \theta$, we can find, by Corollary 2.6, a holomorphic function ν_ϑ on $\mathfrak{h}_\mathbb{C}^*$ that has moderate vertical growth and vanishes on $\mathfrak{h}_{\vartheta, \mathbb{C}}^* + \varpi$ for all but finitely many elements $\varpi \in \mathbf{Y}_\vartheta^*$, such that $\nu(w\alpha_0) \neq 0$ for every $w \in \mathbf{W}$. Now since θ is a finite set, we can take the product $\nu := \prod_{\vartheta \in \theta} \nu_\vartheta \in \mathcal{M}_\theta^\sharp(\mathfrak{h}_\mathbb{C}^*)$, satisfying $\nu(w\alpha_0) \neq 0$ for every $w \in \mathbf{W}$. Put $\mu := \prod_{w \in \mathbf{W}} \nu \circ w$. Then μ belongs to $\mathcal{M}_\theta^\sharp(\mathfrak{h}_\mathbb{C}^*)^\mathbf{W}$ and satisfies $\mu(\alpha_0) \neq 0$. The lemma follows. $\square$

2.3. *Multipliers of Schwartz algebra.* We now consider a connected reductive algebraic group $\underline{G}$ over $\mathbb{R}$. Let $(\mathbf{X}^*, \Phi, \mathbf{X}_*, \Phi^\vee)$ be the root datum associated to $\underline{G}_\mathbb{C}$, namely, $\mathbf{X}^*$ and $\mathbf{X}_*$ are the weight and coweight lattices of the abstract Cartan group of $\underline{G}_\mathbb{C}$, with Φ and $\Phi^\vee$ the subsets of roots and coroots of $\underline{G}_\mathbb{C}$, respectively. We keep the notation in the previous subsection.

We let θ be the subset of $\text{Aut}(\mathbf{X}^*, \Phi, \mathbf{X}_*, \Phi^\vee)^\heartsuit$ consisting of elements of the form $w\vartheta$ in which $w \in \mathbf{W}$ and $\vartheta \in \text{Aut}(\mathbf{X}^*, \Phi, \mathbf{X}_*, \Phi^\vee)^\heartsuit$ is an element that induces the real form $\underline{G}$ of $\underline{G}_\mathbb{C}$. Then θ is stable under the conjugation action of $\mathbf{W}$.

Denote by $\mathfrak{g}$ the complex Lie algebra of $\underline{G}_\mathbb{C}$ and by $\mathcal{U}(\mathfrak{g})$ the universal enveloping algebra of $\mathfrak{g}$ with the center $\mathcal{Z}(\mathfrak{g})$. By the Harish-Chandra isomorphism $\mathcal{Z}(\mathfrak{g}) \simeq \mathbb{C}[\mathfrak{h}_\mathbb{C}^*]^\mathbf{W}$ [HC51], we obtain a character χ_α of $\mathcal{Z}(\mathfrak{g})$ for every element α of $\mathfrak{h}_\mathbb{C}^*$. Conversely, every character χ of $\mathcal{Z}(\mathfrak{g})$ gives rise to a $\mathbf{W}$ -orbit in $\mathfrak{h}_\mathbb{C}^*$, hence $\mu(\chi)$ is well defined for an element $\mu \in \mathcal{O}(\mathfrak{h}_\mathbb{C}^*)^\mathbf{W}$.

Put $G := \underline{G}(\mathbb{R})$. We fix a maximal compact subgroup K of G and a Haar measure dg on G . Denote by $\mathbb{C}[G]$ and $\mathcal{D}[G]$ the complex algebras of algebraic functions and algebraic differential operators on G , respectively.

By an *admissible representation* of G , we mean a smooth admissible Fréchet representation of moderate growth of G in the sense of Casselman–Wallach ([Cas89], [Wal92, §11]). The category of admissible representations of G is equivalent to the category of Harish-Chandra $(\mathfrak{g}, K)$ -modules by the functor of taking K -finite vectors. For an irreducible admissible representation π of G ,

we denote by $\chi_\pi: \mathcal{Z}(\mathfrak{g}) \rightarrow \mathbb{C}$ its infinitesimal character, which is identified with a $\mathbf{W}$ -orbit in $\mathfrak{h}_{\mathbb{C}}^*$.

We recall some definitions and facts from [Wal83, §§2.5, 2.6]. A *Schwartz function* on G is a smooth function f such that Df is bounded for every $D \in \mathcal{D}(G)$. Let $\mathcal{S}(G)$ be the convolution algebra of Schwartz functions on G , equipped with its natural Fréchet topology, under which the convolution product $*$, defined by the formula

$$(f_1 * f_2)(g) := \int_G f_1(gh^{-1})f_2(h) \, dh,$$

is continuous. For every admissible representation (or more generally Fréchet representation of moderate growth) (π, V_π) of G , the expression

$$\pi(f)v := \int_G f(g)\pi(g)v \, dg$$

is absolutely convergent for every $f \in \mathcal{S}(G)$ and $v \in V_\pi$, hence defines a continuous operator $\pi(f) \in \text{End}(V_\pi)$.

Remark 2.11. We have $\mathcal{S}(G) \subseteq \bigcap_{p>0} \mathcal{C}^p(G)$, where $\mathcal{C}^p(G)$ denotes the convolution algebra of Harish-Chandra L^p -Schwartz functions on G ; the inclusion is an equality if and only if the center of G is compact.

Remark 2.12. Let $L^2(G)$ be the L^2 -space of G . Using the Sobolev lemma, it is easy to see that $\mathcal{S}(G)$ is also the space of $f \in L^2(G)$ such that $Df \in L^2(G)$ for every $D \in \mathcal{D}(G)$, where Df is understood in the sense of distributions. Moreover, the Fréchet topology of $\mathcal{S}(G)$ is also induced from the semi-norms $f \mapsto \|Df\|_{L^2}$ for all $D \in \mathcal{D}(G)$.

The following theorem provides many multipliers of the algebra $\mathcal{S}(G)$.

THEOREM 2.13. *For every element $\mu \in \mathcal{M}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^{\mathbf{W}}$, there is a unique linear operator*

$$\mu \star: \mathcal{S}(G) \rightarrow \mathcal{S}(G),$$

such that

$$\pi(\mu \star f) = \mu(\chi_\pi) \cdot \pi(f)$$

holds for every $f \in \mathcal{S}(G)$ and every irreducible admissible representation π of G . In particular, $\mu \star \in \text{Mul}(\mathcal{S}(G))$ is a multiplier of $\mathcal{S}(G)$.

The proof of this theorem will be given in the next subsection.

Remark 2.14. The subset θ defined above only determines the inner form class of $\underline{G}$. We may define a refined invariant $\theta_{\underline{G}}$ associated to $\underline{G}$ to be the subset of $\text{Aut}(\mathbf{X}^*, \Phi, \mathbf{X}_*, \Phi^\vee)^\heartsuit$ of elements induced from all maximal tori $\underline{T}$ of $\underline{G}$ in the way described in Lemma 2.19 below, which is contained in θ and stable under the conjugation action of $\mathbf{W}$. Note that

- $\theta_{\underline{G}} = \theta$ if and only if $\underline{G}$ is quasi-split;
- $\theta_{\underline{G}}$ contains 1 if and only if $\underline{G}$ is split;

- $\theta_{\underline{G}}$ contains -1 if and only if $\underline{G}$ admits an anisotropic maximal torus;
- $\theta_{\underline{G}} = \{-1\}$ if and only if $\underline{G}$ is anisotropic.

Example 2.15. Suppose that $\underline{G}_{\mathbb{C}} = \mathrm{SL}_{2,\mathbb{C}}$. In this case, we may identify X^* with $\mathbb{Z}$ under which $\Phi = \{2, -2\}$, hence $W = \mathrm{Aut}(X^*, \Phi, X_*, \Phi^\vee) = \{\pm 1\}$. There are two cases of $\underline{G}$ up to isomorphism.

- (1) When $\underline{G}$ is split, we have $\theta = \theta_{\underline{G}} = \{\pm 1\}$. In this case, $\mathcal{M}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W = \mathcal{M}_{\theta_{\underline{G}}}(\mathfrak{h}_{\mathbb{C}}^*)^W = \mathcal{M}_{\{\pm 1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$, which consists of even holomorphic functions on $\mathbb{C}$ that have moderate vertical growth, and have polynomial growth on the subset $\mathbb{Z} \subseteq \mathbb{C}$.
- (2) When $\underline{G}$ is anisotropic, we have $\theta_{\underline{G}} = \{-1\} \subseteq \theta = \{\pm 1\}$. In this case, $\mathcal{M}_{\theta_{\underline{G}}}(\mathfrak{h}_{\mathbb{C}}^*)^W = \mathcal{M}_{\{-1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$, which consists of even holomorphic functions on $\mathbb{C}$ that have polynomial growth on the subset $\mathbb{Z} \subseteq \mathbb{C}$.

Remark 2.16. It is natural to ask when a holomorphic function μ on $\mathfrak{h}_{\mathbb{C}}^*$ gives a multiplier as in [Theorem 2.13](#). We conjecture that μ does this if and only if it belongs to $\mathcal{M}_{\theta_{\underline{G}}}(\mathfrak{h}_{\mathbb{C}}^*)^W$; in particular, the conjecture gives rise to a homomorphism $\mathcal{M}_{\theta_{\underline{G}}}(\mathfrak{h}_{\mathbb{C}}^*)^W \rightarrow \mathrm{Mul}(\mathcal{S}(G))$ of complex algebras. By a result of Harish-Chandra, the subset of $\mathfrak{h}_{\mathbb{C}}^*$ of all infinitesimal characters of irreducible admissible representations of G is the union $\bigcup_{\vartheta \in \theta_{\underline{G}}} \mathfrak{h}_{\vartheta, \mathbb{C}}^* \oplus Y_{\vartheta}^*$. Thus, the kernel of the previous homomorphism consists exactly of those μ that vanish on $\bigcup_{\vartheta \in \theta_{\underline{G}}} \mathfrak{h}_{\vartheta, \mathbb{C}}^* \oplus Y_{\vartheta}^*$.

This conjecture can be easily checked when $\underline{G}$ is anisotropic, that is, when G is compact. On the other hand, when $\underline{G}$ is split, [Theorem 2.13](#) implies the existence of the homomorphism $\mathcal{M}_{\theta_{\underline{G}}}(\mathfrak{h}_{\mathbb{C}}^*)^W \rightarrow \mathrm{Mul}(\mathcal{S}(G))$, as in this case $\theta \cup \{1\} = \theta = \theta_{\underline{G}}$ ([Remark 2.14](#)). We also remark that when $\underline{G} = \mathrm{SL}_{2,\mathbb{R}}$, this conjecture was known as a consequence of the work [[Bar88](#)].

Remark 2.17. [Theorem 2.13](#) together with [Definition 2.8\(3\)](#) provide us with a homomorphism

$$\mathcal{M}_{\theta}^{\sharp}(\mathfrak{h}_{\mathbb{C}}^*)^W \rightarrow \mathrm{Mul}(\mathcal{S}(G))$$

of complex algebras.

To end this subsection, we record a property for elements in $\mathcal{M}_{\theta}^{\sharp}(\mathfrak{h}_{\mathbb{C}}^*)^W$ that will be used later. As usual, by a K -type, we mean an isomorphism class of irreducible smooth representations of K .

LEMMA 2.18. *Let μ be an element in $\mathcal{M}_{\theta}^{\sharp}(\mathfrak{h}_{\mathbb{C}}^*)^W$. Then there is a finite set $\mathfrak{T}(\mu)$ of K -types such that for every irreducible admissible representation π of G satisfying $\mu(\chi_{\pi}) \neq 0$, we have that $\pi|_K$ contains some member from $\mathfrak{T}(\mu)$.*

Proof. By [Definition 2.8\(3\)](#) together with Harish-Chandra's description of the infinitesimal characters of discrete series of Levi subgroups of G [[HC66](#)],

there are only finitely many pairs (M, σ) where M is a Levi subgroup of G and σ is an irreducible discrete series representation of M , up to conjugation and unramified twists, such that μ is non-vanishing on the infinitesimal character of the parabolic induction of (M, σ) . By the Langlands classification and the fact that for a standard module, the Langlands quotient inherits its minimal K -types [Kna01, Th. 15.10], we can take $\mathfrak{T}(\mu)$ to be the minimal K -types of the parabolic induction of those finitely many pairs (M, σ) . $\square$

2.4. *Proof of Theorem 2.13.* In this subsection, we prove Theorem 2.13. We fix a W -invariant Euclidean norm $\|\cdot\|$ on $\mathfrak{h}_{\mathbb{C}}^*$. For $\vartheta \in \text{Aut}(\mathbf{X}^*, \Phi, \mathbf{X}_*, \Phi^\vee)^\heartsuit$, $M > 0$, and $r \in \mathbb{R}$, we put

$$p_{\vartheta, M, r}(\mu) := \sup_{\substack{\alpha \in \mathfrak{h}_{\vartheta, \mathbb{C}}^*, \|\text{Re } \alpha\| < M \\ \varpi \in \mathbf{Y}_{\vartheta}^*}} |\mu(\alpha + \varpi)| \cdot (1 + \|\alpha + \varpi\|)^r$$

for $\mu \in \mathcal{O}(\mathfrak{h}_{\mathbb{C}}^*)$. Clearly, we have $p_{\vartheta, M, r}(\mu) \leq p_{\vartheta, M', r}(\mu)$ if $M \leq M'$ and $p_{\vartheta, M, r}(\mu) \leq p_{\vartheta, M, r'}(\mu)$ if $r \leq r'$.

We start with some discussion on maximal tori of $\underline{G}$. Let H be the abstract Cartan group of $\underline{G}_{\mathbb{C}}$, whose weight lattice is $\mathbf{X}^*$. Recall that the abstract Cartan group H of $\underline{G}_{\mathbb{C}}$ represents the presheaf on the opposite category of complex tori whose value on (a complex torus) S is the set of collections $\{q_B: B \rightarrow S\}$ of homomorphisms for every Borel subgroup B of $\underline{G}_{\mathbb{C}}$ satisfying that $q_B = q_{B'} \circ \alpha_{B, B'}$ for every inner automorphism $\alpha_{B, B'}$ of $\underline{G}_{\mathbb{C}}$ mapping B into B' . In particular, for every maximal torus T of $\underline{G}$, there is a canonical W -conjugacy class of isomorphisms $\underline{T}_{\mathbb{C}} \xrightarrow{\sim} H$ from the universal object, which we refer as universal isomorphisms.

Let $\underline{T}$ be a maximal torus of $\underline{G}$. Put $T := \underline{T}(\mathbb{R})$. Then T admits a decomposition $T = A_T T_c$ in which A_T and T_c are the maximal split and the anisotropic (analytic) sub-tori, respectively; it induces a decomposition $\mathfrak{t} = \mathfrak{a}_T \oplus \mathfrak{t}_c$ for the corresponding Lie algebras.

LEMMA 2.19. *For every universal isomorphism $\underline{T}_{\mathbb{C}} \xrightarrow{\sim} H$, which induces a decomposition $\mathfrak{h}^* = \mathfrak{a}_T^* \oplus \mathfrak{it}_c^*$, there is a unique element $\vartheta \in \theta$ such that $\mathfrak{a}_T^* = \mathfrak{h}_{\vartheta}^*$ and $\mathfrak{it}_c^* = \mathfrak{h}_{\vartheta}^{*-}$.*

Proof. We define an automorphism $\vartheta_{\underline{T}}$ of $\underline{T}$ as follows. Let d be the rank of $\underline{T}$, and fix an isomorphism $\iota_{\underline{T}}: \underline{T}_{\mathbb{C}} \xrightarrow{\sim} \mathbf{G}_{m, \mathbb{C}}^d$. Then we set $\vartheta_{\underline{T}} := \overline{\iota_{\underline{T}}}^{-1} \circ \iota_{\underline{T}}$, where $\overline{\iota_{\underline{T}}}$ denotes the complex conjugate of $\iota_{\underline{T}}$.

It is easy to see that $\vartheta_{\underline{T}}$ does not depend on the choice of $\iota_{\underline{T}}$ and, moreover, that on the Lie algebra $\mathfrak{t}$ of $\underline{T}$, $\vartheta_{\underline{T}}$ acts by $+1$ on $\mathfrak{a}_T$ and by -1 on $\mathfrak{t}_c$. Take a universal isomorphism $\underline{T}_{\mathbb{C}} \xrightarrow{\sim} H$, which gives rise to a decomposition $\mathfrak{h}^* = \mathfrak{a}_T^* \oplus \mathfrak{it}_c^*$. Then $\vartheta_{\underline{T}}$ induces an element $\vartheta \in \text{Aut}(\mathbf{X}^*, \Phi, \mathbf{X}_*, \Phi^\vee)$ that belongs to θ , which satisfies $\mathfrak{a}_T^* = \mathfrak{h}_{\vartheta}^*$ and $\mathfrak{it}_c^* = \mathfrak{h}_{\vartheta}^{*-}$. The lemma is proved as the uniqueness is clear. $\square$

As we have explained in [Section 1.3](#), the key step is to bound $\|D(\mu \star f)\|_{L^2}$ when $D \in \mathcal{D}(G)$ is a polynomial, which boils down to the study of certain linear operators on $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^{\mathbb{W}}$. Those operators turn out to be from the algebra $\mathcal{S}$ below, which we study now.

Denote by $\widetilde{\mathbb{W}} := X^* \rtimes \mathbb{W}$ the extended affine Weyl group and by $\widetilde{\mathbb{W}}^{\heartsuit} \subseteq \widetilde{\mathbb{W}}$ the subset of reflections. Recall that an element $s \in \widetilde{\mathbb{W}}$ is a *reflection* if the locus H_s of fixed points of s on $\mathfrak{h}_{\mathbb{C}}^*$ is an affine hyperplane. For every $s \in \widetilde{\mathbb{W}}^{\heartsuit}$, we fix an affine function ℓ_s on $\mathfrak{h}_{\mathbb{C}}^*$ with zero locus H_s .

For every $w \in \widetilde{\mathbb{W}}$, we denote by t_w the operator on $\mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]$ given by $(t_w P)(\alpha) = P(w^{-1}\alpha)$. Let $\mathcal{R}$ be the algebra of endomorphisms of $\mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]$ generated by t_w for all $w \in \widetilde{\mathbb{W}}$ and multiplications by elements of $\mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]$. We consider $\mathcal{R}$ as a left $\mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]$ -module in the obvious way. Put

$$\begin{aligned}\mathcal{S} &:= \mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*][\ell_s^{-1} \mid s \in \widetilde{\mathbb{W}}^{\heartsuit}] \otimes_{\mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]} \mathcal{R}, \\ \mathcal{T} &:= \mathbb{C}(\mathfrak{h}_{\mathbb{C}}^*) \otimes_{\mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]} \mathcal{R},\end{aligned}$$

where $\mathbb{C}(\mathfrak{h}_{\mathbb{C}}^*)$ denotes the function field of $\mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]$. We have natural maps $\mathcal{R} \rightarrow \mathcal{S} \rightarrow \mathcal{T}$ of algebras. The algebra $\mathcal{S}$ is independent of the choices of $\{\ell_s \mid s \in \widetilde{\mathbb{W}}^{\heartsuit}\}$. The algebra $\mathcal{T}$ can be considered as the algebra of endomorphisms of $\mathbb{C}(\mathfrak{h}_{\mathbb{C}}^*)$ generated by t_w for all $w \in \widetilde{\mathbb{W}}$ and multiplications by elements of $\mathbb{C}(\mathfrak{h}_{\mathbb{C}}^*)$.

The following lemma tells us when an element $S \in \mathcal{S}$ extends to a linear operator on $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)$ and how it behaves with respect to the family of seminorms $\{p_{\vartheta, M, r}\}$.

LEMMA 2.20. *Let $S \in \mathcal{S}$ be an element satisfying $S\mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*] \subset \mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]$. Then S extends uniquely to a continuous endomorphism of $\mathcal{O}(\mathfrak{h}_{\mathbb{C}}^*)$ with respect to the topology of uniform convergence on compact subsets, which we still denote by S . Moreover, there exist $M_S > 0$ and $r_S > 0$ such that for every $M > 0$ and every $r \in \mathbb{R}$, there exists $C_{S, M, r} > 0$ such that*

$$\begin{aligned}p_{1, M, r}(S\mu) &\leq C_{S, M, r} \cdot p_{1, M+M_S, r+r_S}(\mu), \\ \max_{\vartheta \in \theta} p_{\vartheta, M, r}(S\mu) &\leq C_{S, M, r} \cdot \max_{\vartheta \in \theta} p_{\vartheta, M+M_S, r+r_S}(\mu)\end{aligned}$$

hold for every $\mu \in \mathcal{O}(\mathfrak{h}_{\mathbb{C}}^*)$. In particular,

- (1) S preserves the subspace $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)$;
- (2) if $S\mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*] \subset \mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]^{\mathbb{W}}$, then S preserves the subspace $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^{\mathbb{W}}$.

Proof. Note that for the topology of uniform convergence on compact subsets, the subspace $\mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]$ is dense in $\mathcal{O}(\mathfrak{h}_{\mathbb{C}}^*)$, hence the uniqueness of the extension is clear.

Now we show the existence of the extension, with the estimate on $p_{\vartheta, M, r}(S\mu)$ together. By definition, S is of the form $\Theta^{-1}S'$, where $S' \in \mathcal{R}$ and Θ is a finite product (possibly with multiplicities) of ℓ_s for $s \in \widetilde{\mathbb{W}}^{\heartsuit}$. It is immediate that

the action of S' on $\mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]$ extends by continuity to an action on $\mathcal{O}(\mathfrak{h}_{\mathbb{C}}^*)$ with the following property: there exist $M_{S'} > 0$ and $r_{S'} > 0$ such that for every $M > 0$ and every $r \in \mathbb{R}$, there exists $C_{S',M,r} > 0$ such that

$$\begin{aligned} p_{1,M,r}(S'\mu) &\leq C_{S',M,r} \cdot p_{1,M+M_{S'},r+r_{S'}}(\mu), \\ \max_{\vartheta \in \theta} p_{\vartheta,M,r}(S'\mu) &\leq C_{S',M,r} \cdot \max_{\vartheta \in \theta} p_{\vartheta,M+M_{S'},r+r_{S'}}(\mu) \end{aligned}$$

hold for every $\mu \in \mathcal{O}(\mathfrak{h}_{\mathbb{C}}^*)$. In particular, S' preserves the subspace $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)$. Now the similar properties for S follow from [Lemma 2.21](#) below.

For the last two claims, (1) is an immediate consequence of the estimate on $p_{\vartheta,M,r}(S\mu)$, and (2) follows from the uniqueness of the extension. $\square$

LEMMA 2.21. *For every $s \in \widetilde{W}^\vee$, there exists $M_s > 0$ such that for every $M > 0$ and every $r \in \mathbb{R}$, there exists $C_{s,M,r} > 0$ such that*

$$\begin{aligned} p_{1,M,r}(\ell_s^{-1}\mu) &\leq C_{s,M,r} \cdot p_{1,M+M_s,r}(\mu), \\ \max_{\vartheta \in \theta} p_{\vartheta,M,r}(\ell_s^{-1}\mu) &\leq C_{s,M,r} \cdot \max_{\vartheta \in \theta} p_{\vartheta,M+M_s,r}(\mu) \end{aligned}$$

hold for every $\mu \in \mathcal{O}(\mathfrak{h}_{\mathbb{C}}^*)$ that vanishes on H_s . In particular, for a (holomorphic) function $\mu \in \mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)$ that vanishes on H_s , the function $\ell_s^{-1}\mu$ belongs to $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)$ as well.

Proof. For every fixed $\vartheta \in \theta \cup \{1\}$, we claim that there exists $M_s > 0$ such that for every $M > 0$ and $r \in \mathbb{R}$, there exists $C_{s,M,r} > 0$ such that

$$p_{\vartheta,M,r}(\ell_s^{-1}\mu) \leq \begin{cases} C_{s,M,r} \cdot p_{\vartheta,M+M_s,r}(\mu) & \text{if } \vartheta = 1, \\ C_{s,M,r} \cdot \max_{\vartheta' \in \theta} p_{\vartheta',M+M_s,r}(\mu) & \text{if } \vartheta \neq 1 \end{cases}$$

holds for every $\mu \in \mathcal{O}(\mathfrak{h}_{\mathbb{C}}^*)$ that vanishes on H_s . The lemma then follows from this claim.

To prove the claim, let ℓ_s° be the linear part of ℓ_s . Take a function $\mu \in \mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)$ that vanishes on H_s .

First suppose that $\ell_s^\circ|_{\mathfrak{h}_{\vartheta,\mathbb{C}}^*} \neq 0$. Then the claim is an easy consequence of Cauchy's integral formula. Indeed, take an element $\alpha_s \in \mathfrak{h}_{\vartheta,\mathbb{C}}^*$ such that $\ell_s^\circ(\alpha_s) = 1$. When $|\ell_s(\alpha + \varpi)| > 1/2$, we have

$$\left| \frac{\mu(\alpha + \varpi)}{\ell_s(\alpha + \varpi)} \right| \leq 2|\mu(\alpha + \varpi)|,$$

whereas when $|\ell_s(\alpha + \varpi)| \leq 1/2$, we have

$$\left| \frac{\mu(\alpha + \varpi)}{\ell_s(\alpha + \varpi)} \right| = \left| \int_{u \in \mathbb{C}^\times, |u|=1} \frac{\mu(\alpha + u\alpha_s + \varpi)}{\ell_s(\alpha + u\alpha_s + \varpi)} du \right| \leq 2 \sup_{u \in \mathbb{C}^\times, |u|=1} |\mu(\alpha + u\alpha_s + \varpi)|.$$

Together, we obtain

$$(2.2) \quad \left| \frac{\mu(\alpha + \varpi)}{\ell_s(\alpha + \varpi)} \right| \leq 2 \sup_{|u| \leq 1} \|\mu(\alpha + u\alpha_s + \varpi)\|.$$

Choose $C_{s,M,r} > 0$ such that

$$2(1 + \|\alpha + \varpi\|)^r \leq C_{s,M,r} \cdot \min_{u \in \mathbb{C}^\times, |u|=1} (1 + \|\alpha + u\alpha_s + \varpi\|)^r$$

holds for every $\alpha \in \mathfrak{h}_{\vartheta, \mathbb{C}}^*$ with $\|\operatorname{Re} \alpha\| < M$ and every $\varpi \in \mathbf{Y}_{\vartheta}^*$. Then we have

$$p_{\vartheta, M, r}(\ell_s^{-1} \mu) \leq C_{s, M, r} \cdot p_{\vartheta, M+M_s, r}(\mu)$$

with $M_s := \|\alpha_s\|$ depending only on s . Thus, the claim holds.

Now suppose that $\ell_s^\circ|_{\mathfrak{h}_{\vartheta, \mathbb{C}}^*} = 0$, hence $\vartheta \neq 1$. Since the image of $\mathfrak{h}_{\vartheta, \mathbb{C}}^* \oplus \mathbf{Y}_{\vartheta}^*$ under ℓ_s is a discrete subset of $\mathbb{C}$, there exists $C_s > 0$ such that $|\ell_s| > C_s$ on $(\mathfrak{h}_{\vartheta, \mathbb{C}}^* \oplus \mathbf{Y}_{\vartheta}^*) \setminus H_s$. It follows that

$$(2.3) \quad \sup_{\substack{\alpha \in \mathfrak{h}_{\vartheta, \mathbb{C}}^*, \|\operatorname{Re} \alpha\| < M \\ \varpi \in \mathbf{Y}_{\vartheta}^*, \alpha + \varpi \notin H_s}} \left| \frac{\mu(\alpha + \varpi)}{\ell_s(\alpha + \varpi)} \right| \cdot (1 + \|\alpha + \varpi\|)^r \leq C_s^{-1} \cdot p_{\vartheta, M, r}(\mu).$$

It remains to bound $\ell_s^{-1} \mu$ on vertical strips of $\mathfrak{h}_{\vartheta, \mathbb{C}}^* \oplus (\mathbf{Y}_{\vartheta}^* \cap H_s) = (\mathfrak{h}_{\vartheta, \mathbb{C}}^* \oplus \mathbf{Y}_{\vartheta}^*) \cap H_s$. Let $w_s \in \widetilde{W}^\vee \cap W$ be the reflection that is the linear part of s . As $\ell_s^\circ|_{\mathfrak{h}_{\vartheta, \mathbb{C}}^*} = 0$, the actions of w_s and ϑ on $\mathfrak{h}_{\mathbb{C}}^*$ commute. Thus, the element $\vartheta' := w_s \vartheta$ satisfies $\vartheta'^2 = 1$, hence belongs to θ . However for ϑ' , we have $\mathfrak{h}_{\vartheta', \mathbb{C}}^{*-} = \mathfrak{h}_{\vartheta, \mathbb{C}}^{*-} \cap \ker \ell_s^\circ$ and $\mathfrak{h}_{\vartheta, \mathbb{C}}^* \subseteq \mathfrak{h}_{\vartheta', \mathbb{C}}^*$.

Since the map ℓ_s° induces an isomorphism $\mathfrak{h}_{\vartheta', \mathbb{C}}^* \cap \mathfrak{h}_{\vartheta, \mathbb{C}}^{*-} \xrightarrow{\sim} \mathbb{C}$, there exists an element $\alpha'_s \in \mathfrak{h}_{\vartheta', \mathbb{C}}^*$ such that $\varpi' := \varpi - \alpha'_s \in \mathfrak{h}_{\vartheta', \mathbb{C}}^{*-}$ for every $\varpi \in \mathbf{Y}_{\vartheta}^* \cap H_s$. Since $\ell_s^\circ|_{\mathfrak{h}_{\vartheta', \mathbb{C}}^*} \neq 0$, we obtain from (2.2) that there exists $\alpha_s \in \mathfrak{h}_{\vartheta', \mathbb{C}}^*$ such that

$$\left| \frac{\mu(\alpha + \varpi)}{\ell_s(\alpha + \varpi)} \right| \leq 2 \sup_{|u| \leq 1} |\mu(\alpha + u\alpha_s + \alpha'_s + \varpi')|$$

holds for $\alpha + \varpi \in (\mathfrak{h}_{\vartheta, \mathbb{C}}^* \oplus \mathbf{Y}_{\vartheta}^*) \cap \ker \ell_s$. Note that the projection of $\mathbf{Y}_{\vartheta}^*$ onto $\mathfrak{h}_{\vartheta', \mathbb{C}}^{*-}$ coincides with $\mathbf{Y}_{\vartheta'}^*$. Thus, as in the previous case, we may find a constant $C_{s, M, r} \geq C_s^{-1}$ such that

$$\sup_{\substack{\alpha \in \mathfrak{h}_{\vartheta, \mathbb{C}}^*, \|\operatorname{Re} \alpha\| < M \\ \varpi \in \mathbf{Y}_{\vartheta}^*, \alpha + \varpi \in H_s}} \left| \frac{\mu(\alpha + \varpi)}{\ell_s(\alpha + \varpi)} \right| \cdot (1 + \|\alpha + \varpi\|)^r \leq C_{s, M, r} \cdot p_{\vartheta', M+M_s, r}(\mu)$$

with $M_s := \|\alpha_s\| + \|\alpha'_s\|$. Thus, the claim holds after combining with (2.3). $\square$

Now we relate the algebra $\mathcal{S}$ to finite dimensional algebraic representations of G . Let (τ, W_τ) be a finite dimensional algebraic representation of G , and let

$$\delta: \mathcal{U}(\mathfrak{g}) \rightarrow \mathcal{U}(\mathfrak{g}) \otimes \operatorname{End}(W_\tau)$$

be the homomorphism that sends $X \in \mathfrak{g}$ to $X \otimes 1 + 1 \otimes \tau(X)$. Let $\mathcal{R}_\tau$ be the centralizer of the image of δ . By [Kos75, Th. 4.8], $\mathcal{R}_\tau$ is a finitely generated free $\mathcal{Z}(\mathfrak{g})$ -module, where $\mathcal{Z}(\mathfrak{g})$ acts on $\mathcal{U}(\mathfrak{g}) \otimes \text{End}(W_\tau)$ by $z \cdot (u \otimes A) = (zu) \otimes A$. Recall that we have identified $\mathcal{Z}(\mathfrak{g})$ with $\mathbb{C}[\mathfrak{h}_\mathbb{C}^*]^W$ via the Harish-Chandra isomorphism.

LEMMA 2.22. *For every basis $v_1, \dots, v_r$ of the finite free $\mathcal{Z}(\mathfrak{g})$ -module $\mathcal{R}_\tau$, there exist elements $S_1, \dots, S_r \in \mathcal{S}$ sending $\mathbb{C}[\mathfrak{h}_\mathbb{C}^*]$ to $\mathbb{C}[\mathfrak{h}_\mathbb{C}^*]^W$ such that*

$$\delta(z) = \sum_{i=1}^r S_i(z) v_i$$

for every $z \in \mathcal{Z}(\mathfrak{g})$.

Proof. The basis $v_1, \dots, v_r$ induces an isomorphism $\mathcal{R}_\tau \simeq \mathcal{Z}(\mathfrak{g})^{\oplus r}$, through which the right multiplication by $\delta(z)$ for $z \in \mathcal{Z}(\mathfrak{g})$ is represented by a matrix $S(z) = (S_{i,j}(z))_{1 \leq i,j \leq r} \in \text{Mat}_r(\mathcal{Z}(\mathfrak{g}))$. By Lemma 2.23 below, it suffices to show that for every $1 \leq i, j \leq r$, the map $z \in \mathcal{Z}(\mathfrak{g}) \mapsto S_{i,j}(z)$ is induced from an element of $\mathcal{S}$ sending $\mathbb{C}[\mathfrak{h}_\mathbb{C}^*]$ to $\mathbb{C}[\mathfrak{h}_\mathbb{C}^*]^W$.

Let $\Lambda(\tau) \subset \mathbf{X}^*$ be the set of weights of τ . For every $z \in \mathcal{Z}(\mathfrak{g})$, put

$$P_z(X) := \prod_{\lambda \in \Lambda(\tau)} (X - t_\lambda z),$$

which belongs to $\mathcal{Z}(\mathfrak{g})[X] = \mathbb{C}[\mathfrak{h}_\mathbb{C}^*]^W[X]$, as $\Lambda(\tau)$ is W -invariant. According to [Kos75, Th. 4.9], we have $P_z(\delta(z)) = 0$, or equivalently $P_z(S(z)) = 0$ for every $z \in \mathcal{Z}(\mathfrak{g})$. As the characters $z \in \mathcal{Z}(\mathfrak{g}) \mapsto t_\lambda z \in \mathbb{C}[\mathfrak{h}_\mathbb{C}^*]$ for $\lambda \in \Lambda(\tau)$ are all distinct, there exist an element $P \in \text{GL}_r(\mathbb{C}[\mathfrak{h}_\mathbb{C}^*])$ and a family of weights $\lambda_1, \dots, \lambda_r \in \Lambda(\tau)$ such that

$$S(z) = P \begin{pmatrix} t_{\lambda_1} z & & \\ & \ddots & \\ & & t_{\lambda_r} z \end{pmatrix} P^{-1}$$

holds in $\text{Mat}_r(\mathbb{C}[\mathfrak{h}_\mathbb{C}^*])$ for every $z \in \mathcal{Z}(\mathfrak{g})$. This shows that for every $1 \leq i, j \leq r$, the map $S_{i,j}$ is induced from an element $S'_{i,j} \in \mathcal{S}$ sending $\mathbb{C}[\mathfrak{h}_\mathbb{C}^*]^W$ to $\mathbb{C}[\mathfrak{h}_\mathbb{C}^*]^W$. However, up to replacing $S'_{i,j}$ by $|W|^{-1} \sum_{w \in W} S'_{i,j} t_w$, we see that $S'_{i,j}$ also sends $\mathbb{C}[\mathfrak{h}_\mathbb{C}^*]$ to $\mathbb{C}[\mathfrak{h}_\mathbb{C}^*]^W$.

The lemma is proved. $\square$

LEMMA 2.23. *If an element $T \in \mathcal{S}$ satisfies $T\mathbb{C}[\mathfrak{h}_\mathbb{C}^*] \subset \mathbb{C}[\mathfrak{h}_\mathbb{C}^*]$, then T is contained in the image of $\mathcal{S} \rightarrow \mathcal{S}$.*

This lemma is essentially [KK86, Lemma 4.9]. For the readers' convenience, we give a detailed proof.

Proof. The lemma amounts to the following: For an irreducible polynomial $P \in \mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]$, and a family $(P_w)_{w \in \widetilde{W}}$ of elements of $\mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]$ with $P_w = 0$ for all but finitely many w , if

$$\left(\sum_{w \in \widetilde{W}} P_w t_w \right) \mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*] \subset P \mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*],$$

then either P divides P_w for every $w \in \widetilde{W}$, or $P \in \ell_s \mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]$ for some $s \in \widetilde{W}^\vee$.

Suppose that $P \notin \ell_s \mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]$ for every $s \in \widetilde{W}^\vee$. Then, the zero set of P is not contained in the union $\bigcup_{w \in \widetilde{W} \setminus \{1\}} \ker(w - 1)$. This implies that the stabilizer of a generic element of the zero set $Z(P) := \{\alpha \in \mathfrak{h}_{\mathbb{C}}^* \mid P(\alpha) = 0\}$ in $\widetilde{W}$ is trivial. Let $\alpha \in Z(P)$ be such an element. Note that for every $Q \in \mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]$, we have

$$\sum_{w \in \widetilde{W}} P_w(\alpha) Q(w^{-1}\alpha) = 0.$$

As the linear forms $Q \mapsto Q(w^{-1}\alpha)$ on $\mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]$ are linearly independent, it follows that $P_w(\alpha) = 0$ for every generic element $\alpha \in Z(P)$. Thus, P divides P_w for every $w \in \widetilde{W}$. The lemma is proved. $\square$

As we have explained in [Section 1.3](#), we will first prove [Theorem 2.13](#) for the smaller space $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$, together with bounds on $\|D(\mu \star f)\|_{L^2}$, which is the content of the next proposition.

PROPOSITION 2.24. *[Theorem 2.13](#) holds for $\mu \in \mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$. Moreover, for every $D \in \mathcal{D}(G)$, there exists a real number $M_D > 0$ such that for every $f \in \mathcal{S}(G)$ and every $r \in \mathbb{R}$, there exists $C_{D,f,r} > 0$ such that*

$$\|D(\mu \star f)\|_{L^2} \leq C_{D,f,r} \cdot \max_{\vartheta \in \theta} p_{\vartheta, M_D, r}(\mu)$$

holds for every $\mu \in \mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^)^W$.*

Proof. Let $C_c^\infty(G)_{(K)}$ be the subalgebra of $C_c^\infty(G)$ of bi- K -finite functions. As $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W \subseteq \mathcal{N}(\mathfrak{h}_{\mathbb{C}}^*)^W$, the existence of the function $\mu \star f$ for $f \in C_c^\infty(G)_{(K)}$ was essentially proved by Delorme (see [Proposition A.1](#)). As $C_c^\infty(G)_{(K)}$ is dense in $\mathcal{S}(G)$, it suggests we show that the map $C_c^\infty(G)_{(K)} \rightarrow \mathcal{S}(G)$, $f \mapsto \mu \star f$, extends continuously to an endomorphism of $\mathcal{S}(G)$.

We claim that for every $D \in \mathcal{D}(G)$, there exists $M_D > 0$ such that for every $r \in \mathbb{R}$, there exists a continuous semi-norm $\nu_{D,r}$ on $\mathcal{S}(G)$ such that

$$(2.4) \quad \|D(\mu \star f)\|_{L^2} \leq \nu_{D,r}(f) \cdot \max_{\vartheta \in \theta} p_{\vartheta, M_D, r}(\mu)$$

holds for every $\mu \in \mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$ and $f \in C_c^\infty(G)_{(K)}$.

We deduce the proposition assuming the claim. Fix $\mu \in \mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$, and for every $f \in \mathcal{S}(G)$, choose a sequence $\{f_n\}$ in $C_c^\infty(G)_{(K)}$ that converges to f . Then by [Remark 2.12](#), (2.4) implies that $\mu \star f_n$ converges to an element in $\mathcal{S}(G)$,

which we denote by $\mu \star f$. It is easy to see that the map $\mu \star: \mathcal{S}(G) \rightarrow \mathcal{S}(G)$ satisfies the requirement in [Theorem 2.13](#). The second part of the proposition follows from (2.4) by taking $C_{D,f,r} = \nu_{D,r}(f)$.

Now we show the claim. Note that we have a canonical isomorphism $\mathbb{C}[G] \otimes \mathcal{U}(\mathfrak{g}) \simeq \mathcal{D}(G)$ given by $P \otimes X \mapsto P \cdot L(X)$, where $L(X)$ stands for the left invariant differential operator on G associated to X . For $X \in \mathcal{U}(\mathfrak{g})$, we have $L(X)(\mu \star f) = \mu \star L(X)f$. Since the action of $L(X)$ preserves the set of continuous semi-norms on $\mathcal{S}(G)$, it suffices to show (2.4) for $D = P \in \mathbb{C}[G]$.

First, we consider the case $P = 1$. By the Plancherel formula of Harish-Chandra [\[HC76\]](#), there exists a Borel measure $d\pi$ on the tempered dual $\text{Temp}(G)$ of G such that

$$\|f\|_{L^2}^2 = \int_{\text{Temp}(G)} \|\pi(f)\|_{\text{HS}}^2 d\pi$$

for every $f \in \mathcal{S}(G)$, where $\|\cdot\|_{\text{HS}}$ stands for the Hilbert-Schmidt norm. Thus, we have

$$\|\mu \star f\|_{L^2}^2 = \int_{\text{Temp}(G)} |\mu(\chi_\pi)|^2 \|\pi(f)\|_{\text{HS}}^2 d\pi$$

for every $f \in C_c^\infty(G)_{(K)}$. Let $\theta_{\underline{G}}$ be the subset of θ obtained from maximal tori of $\underline{G}$ in the way of [Lemma 2.19](#) (see also [Remark 2.14](#)). By Harish-Chandra's description of the infinitesimal characters of tempered representations [\[HC75\]](#), the union of χ_π for $\pi \in \text{Temp}(G)$ is exactly the W -stable subset

$$\bigcup_{\vartheta \in \theta_{\underline{G}}} i\mathfrak{h}_{\vartheta}^* \oplus \Upsilon_{\vartheta}^*$$

of $\mathfrak{h}_{\mathbb{C}}^*$. In particular, we have

$$|\mu(\chi_\pi)| \leq \left(\max_{\vartheta \in \theta_{\underline{G}}} p_{\vartheta,1,r}(\mu) \right) \cdot (1 + \|\chi_\pi\|)^{-r}$$

for every $\pi \in \text{Temp}(G)$. Choose elements $z_1, \dots, z_N \in \mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]^W = \mathcal{Z}(\mathfrak{g})$ (depending on r) such that

$$(1 + \|\chi_\pi\|)^{-2r} \leq |z_1(\chi_\pi)|^2 + \dots + |z_N(\chi_\pi)|^2$$

holds for every $\pi \in \text{Temp}(G)$. It follows that

$$\begin{aligned} \|\mu \star f\|_{L^2}^2 &\leq \left(\max_{\vartheta \in \theta_{\underline{G}}} p_{\vartheta,1,r}(\mu) \right)^2 \sum_{i=1}^N \int_{\text{Temp}(G)} |z_i(\chi_\pi)|^2 \cdot \|\pi(f)\|_{\text{HS}}^2 d\pi \\ &= \left(\max_{\vartheta \in \theta_{\underline{G}}} p_{\vartheta,1,r}(\mu) \right)^2 \sum_{i=1}^N \|z_i \star f\|_{L^2}^2 \end{aligned}$$

for every $f \in C_c^\infty(G)_{(K)}$. Thus, we obtain (2.4) with the semi-norm $\nu_{1,r}$ given by

$$\nu_{1,r}(f) := \left(\sum_{i=1}^N \|z_i \star f\|_{L^2}^2 \right)^{\frac{1}{2}}.$$

Now we treat the case for general $P \in \mathbb{C}[G]$. As P is a finite sum of matrix coefficients of finite dimensional algebraic representations of G , by linearity, we may as well assume that there exists a finite dimensional algebraic representation (τ, W_τ) of G , $w \in W_\tau$ and $w^* \in W_\tau^*$ such that

$$P(g) = \langle \tau(g)w, w^* \rangle, \quad g \in G.$$

Let $v_1, \dots, v_r$ be a basis of the finite free $\mathcal{Z}(\mathfrak{g})$ -module $\mathcal{R}_\tau$, and let $S_1, \dots, S_r$ be elements of $\mathcal{S}$ as in Lemma 2.22. By Lemma 2.20, $S_1, \dots, S_r$ extend continuously to endomorphisms of $\mathcal{O}(\mathfrak{h}_\mathbb{C}^*)^W$ for the topology of uniform convergence on compact subsets, which preserve $\mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_\mathbb{C}^*)^W$, for which we use the same notation. Since the map $\mathcal{U}(\mathfrak{g}) \otimes \text{End}(W_\tau) \rightarrow \mathcal{U}(\mathfrak{g}) \otimes \text{End}(W_\tau)$ given by $u \otimes A \mapsto (1 \otimes A)\delta(u)$ is an isomorphism, each element v_i in the basis can be written as a finite sum

$$(2.5) \quad v_i = \sum_j (1 \otimes A_{ij})\delta(u_{ij})$$

for some $A_{ij} \in \text{End}(W_\tau)$ and $u_{ij} \in \mathcal{U}(\mathfrak{g})$. Let $P_{ij} \in \mathbb{C}[G]$ be the element defined by $P_{ij}(g) := \langle A_{ij}\tau(g)w, w^* \rangle$ for $g \in G$. Note that S_i , u_{ij} , and P_{ij} depend on the data (τ, W_τ, w, w^*) only. We make the following claim:

(*) For every $f \in C_c^\infty(G)_{(K)}$,

$$P(\mu \star f) = \sum_{i,j} S_i(\mu) \star P_{ij}L(u_{ij})f$$

holds.

Assuming (*), by the $P = 1$ case, we have

$$\begin{aligned} \|P(\mu \star f)\|_{L^2} &\leq \sum_{i,j} \|S_i(\mu) \star P_{ij}L(u_{ij})f\|_{L^2} \\ &\leq \sum_{i,j} \nu_{1,r'}(P_{ij}L(u_{ij})f) \cdot \max_{\vartheta \in \theta} p_{\vartheta,1,r'}(S_i(\mu)) \end{aligned}$$

for every $f \in C_c^\infty(G)_{(K)}$ and every $r' \in \mathbb{R}$. By Lemma 2.20, there exist $M_P > 1$, $r_P > 0$, and $C_P > 0$ such that

$$\max_{\vartheta \in \theta} p_{\vartheta,1,r-r_P}(S_i(\mu)) \leq C_P \cdot \max_{\vartheta \in \theta} p_{\vartheta,M_P,r}(\mu)$$

holds for every i and every $\mu \in \mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^W$. Thus, we obtain (2.4) with the semi-norm $\nu_{P,r}$ given by

$$\nu_{P,r}(f) := C_P \sum_{i,j} \nu_{1,r-r_P}(P_{ij}L(u_{ij})f).$$

It remains to confirm (*). By the injectivity of the operator valued Fourier transform, it suffices to check that

$$(2.6) \quad \pi(P(\mu \star f)) = \sum_{i,j} \pi(S_i(\mu) \star P_{ij}L(u_{ij})f) = \sum_{i,j} S_i(\mu)(\chi_\pi) \cdot \pi(P_{ij}L(u_{ij})f)$$

for every irreducible admissible representation π of G . By the subquotient theorem, it is enough to check (2.6) when π is a principal series whose underlying space we denote by V_π . We have the equality

$$(2.7) \quad \pi(P(\mu \star f)) = c_{w^*} \circ (\pi \otimes \tau)(\mu \star f) \circ b_w$$

in $\text{End}(V_\pi)$, where $b_w: V_\pi \rightarrow V_\pi \otimes W_\tau$ is the map defined by $b_w(v) = v \otimes w$, and $c_{w^*}: V_\pi \otimes W_\tau \rightarrow V_\pi$ is the map defined by $c_{w^*}(v \otimes w') = \langle w', w^* \rangle v$. By a density argument, it is even sufficient to establish (2.6) for a principal series in general position, for which, by Lemma 2.25 below, we can assume that $\pi \otimes \tau$ is semisimple.

Assume now that $\pi \otimes \tau$ is semisimple. Let $(z_n)_n$ be a sequence of elements of $\mathcal{Z}(\mathfrak{g})$ converging to μ for the topology of uniform convergence on compact subsets. As $\pi \otimes \tau$ decomposes as a direct sum of irreducible admissible representations of G , $(\pi \otimes \tau)(z_n \star f)$ converges to $(\pi \otimes \tau)(\mu \star f)$ for the topology of pointwise convergence. The action of G induces an action of $\mathcal{U}(\mathfrak{g})$ on (the smooth vectors in) V_π , which further induces an action of $\mathcal{R}_\tau \subseteq \mathcal{U}(\mathfrak{g}) \otimes \text{End}(W_\tau)$ on $V_\pi \otimes W_\tau$. Note that under this action, $\delta(u)$ acts by $(\pi \otimes \tau)(u)$ for every $u \in \mathcal{U}(\mathfrak{g})$, and $z \otimes 1$ acts by $\pi(z) \otimes 1_{W_\tau}$ for every $z \in \mathcal{Z}(\mathfrak{g})$. Thus, by (2.5) and Lemma 2.22, we have

$$\begin{aligned} (\pi \otimes \tau)(z_n) &= \delta(z_n) = \sum_{i=1}^r S_i(z_n)v_i = \sum_{i,j} S_i(z_n)(1 \otimes A_{ij})\delta(u_{ij}) \\ &= \sum_{i,j} (\pi(S_i(z_n)) \otimes 1_{W_\tau}) \circ (1_{V_\pi} \otimes A_{ij}) \circ (\pi \otimes \tau)(u_{ij}) \\ &= \sum_{i,j} S_i(z_n)(\chi_\pi) \cdot (1_{V_\pi} \otimes A_{ij}) \circ (\pi \otimes \tau)(u_{ij}) \end{aligned}$$

in $\text{End}(V_\pi \otimes W_\tau)$. Pre-composing $(\pi \otimes \tau)(f)$ and passing to the limit, we get

$$(\pi \otimes \tau)(\mu \star f) = \sum_{i,j} S_i(\mu)(\chi_\pi) \cdot (1_{V_\pi} \otimes A_{ij}) \circ (\pi \otimes \tau)(L(u_{ij})f),$$

which, together with (2.7), implies (2.6). The claim (*) is proved.

The proof of the proposition is now complete. $\square$

LEMMA 2.25. *Let $P_0 = M_0 N_0$ be a minimal parabolic subgroup of G , σ an irreducible (finite dimensional) representation of M_0 , and put $\sigma_\xi := \sigma \otimes \xi$ for ξ an unramified character of M_0 . Then for ξ in general position, the admissible representation $I_{P_0}^G(\sigma_\xi) \otimes \tau$ is semisimple, that is, a direct sum of irreducible admissible representations.*

Proof. By the Frobenius reciprocity, $I_{P_0}^G(\sigma_\xi) \otimes \tau$ is isomorphic to $I_{P_0}^G(\sigma_\xi \otimes \tau|_{P_0})$, where $\tau|_{P_0}$ denotes the restriction of τ to P_0 (so that the inducing representation is now non-trivial on N_0). Let A_0 be the split center of M_0 and $\mathfrak{a}_0$ its Lie algebra. The representation $\sigma \otimes \tau|_{P_0}$ of P_0 admits a filtration indexed by the characters of $\mathfrak{a}_0$ for the partial order defined by the cone of positive roots with respect to P_0 , such that in the associated graded vector space $\bigoplus_{\lambda \in \mathfrak{a}_{0,\mathbb{C}}^*} V_\lambda$, $\mathfrak{a}_0$ acts on V_λ by the character λ . This implies that N_0 acts trivially on each of the graded pieces. Thus, $\{V_\lambda \mid \lambda \in \mathfrak{a}_{0,\mathbb{C}}^*\}$ are semisimple representations of M_0 with distinct central characters. Therefore, $I_{P_0}^G(\sigma_\xi) \otimes \tau$ admits a filtration with associated graded representations $\bigoplus_{\lambda \in \mathfrak{a}_{0,\mathbb{C}}^*} I_{P_0}^G(V_\lambda \otimes \xi)$, and for generic ξ , the representations $\{I_{P_0}^G(V_\lambda \otimes \xi) \mid \lambda \in \mathfrak{a}_{0,\mathbb{C}}^*\}$ are all semisimple with distinct infinitesimal characters, hence $I_{P_0}^G(\sigma_\xi) \otimes \tau$ is itself semisimple. The lemma follows. $\square$

Now we deduce Theorem 2.13 from Proposition 2.24 by a limit process.

Proof of Theorem 2.13. First, we claim that there exists $P \in \mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]^W$ that is homogeneous and takes positive real values on $(\bigcup_{\vartheta \in \theta \cup \{1\}} i\mathfrak{h}_\vartheta^* \oplus \mathfrak{h}_\vartheta^{*-}) \setminus \{0\}$. Indeed, such a polynomial can be obtained as follows. Let W_θ be the group of linear automorphisms of $\mathfrak{h}^*$ generated by W and θ . Let $P_1, \dots, P_N$ be homogeneous generators of the kernel of $\mathbb{R}[\mathfrak{h}^*]^{W_\theta} \rightarrow \mathbb{R}$ sending P to $P(0)$ (as an ideal of $\mathbb{R}[\mathfrak{h}^*]^{W_\theta}$) with $d_j := \deg P_j > 0$ for $1 \leq j \leq N$. Put

$$P := P_1^{4d'_1} + \dots + P_N^{4d'_N},$$

where $d'_j := d_1 \cdots \widehat{d_j} \cdots d_N$. Since P is homogeneous of degree $4d_1 \cdots d_N$, we have $P(i\alpha) = P(\alpha)$ for every $\alpha \in \mathfrak{h}_{\mathbb{C}}^*$. As the only common zero of $P_1, \dots, P_N$ is zero, it suffices to show that P_j takes real values on $\mathfrak{h}_\vartheta^* \oplus i\mathfrak{h}_\vartheta^{*-}$ for every $1 \leq j \leq N$ and every $\vartheta \in \theta \cup \{1\}$. Since $\vartheta(\bar{\alpha}) = \alpha$ for $\alpha \in \mathfrak{h}_\vartheta^* \oplus i\mathfrak{h}_\vartheta^{*-}$ and since P_j is ϑ -invariant, we have $P_j(\alpha) = P_j(\vartheta(\bar{\alpha})) = P_j(\bar{\alpha}) = \overline{P_j(\alpha)}$, hence $P_j(\alpha) \in \mathbb{R}$ for $\alpha \in \mathfrak{h}_\vartheta^* \oplus i\mathfrak{h}_\vartheta^{*-}$.

We now choose such a polynomial P as above, of degree $d > 0$. It has the following property: there exist $\delta > 0$ and $\epsilon > 0$ such that for every $\vartheta \in \theta \cup \{1\}$, every $\alpha \in \mathfrak{h}_{\vartheta,\mathbb{C}}^*$, and every $\varpi \in \mathfrak{h}_\vartheta^{*-}$ satisfying $\|\alpha + \varpi\| = 1$,

$$\operatorname{Re} P(\alpha + \varpi) \geq \epsilon$$

holds as long as $\|\operatorname{Re} \alpha\| \leq \delta$. In particular, for every $\vartheta \in \theta \cup \{1\}$ and $M > 0$,

$$\operatorname{Re} P(\alpha + \varpi) = \|\alpha + \varpi\|^d \cdot \operatorname{Re} P\left(\frac{\alpha + \varpi}{\|\alpha + \varpi\|}\right) \geq \epsilon \|\alpha + \varpi\|^d$$

holds if $\|\operatorname{Re} \alpha\| < M$ and $\|\alpha + \varpi\| \geq \delta^{-1}M$. This implies the following statements:

- (1) For every $M > 0$ and every $r < 0$, there exists $C_{M,r} > 0$ such that

$$|\exp(-P(\alpha + \varpi))| < C_{M,r}(1 + \|\alpha + \varpi\|)^r$$

holds for every $\vartheta \in \theta \cup \{1\}$, every $\alpha \in \mathfrak{h}_{\vartheta, \mathbb{C}}^*$ with $\|\operatorname{Re} \alpha\| < M$, and every $\varpi \in \mathbf{Y}_{\vartheta}^*$.

- (2) For every $\epsilon > 0$ that is sufficiently small and every $M > 0$, there exists $C_{\epsilon, M} > 0$ such that

$$|\exp(-P(\alpha + \varpi)) - 1| \leq C_{\epsilon, M} \|\alpha + \varpi\|^\epsilon$$

holds for every $\vartheta \in \theta \cup \{1\}$, every $\alpha \in \mathfrak{h}_{\vartheta, \mathbb{C}}^*$ with $\|\operatorname{Re} \alpha\| < M$, and every $\varpi \in \mathbf{Y}_{\vartheta}^*$.

Now we take an element $\mu \in \mathcal{M}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^{\mathbf{W}}$ and define functions μ_n for every integer $n \geq 1$ by the formula

$$\mu_n(\alpha) = \exp\left(-P\left(\frac{\alpha}{n}\right)\right) \cdot \mu(\alpha), \quad \alpha \in \mathfrak{h}_{\mathbb{C}}^*.$$

By [Definition 2.8\(2\)](#), for every $M > 0$, we may choose $r_M \in \mathbb{R}$ such that $p_{\vartheta, M, r_M}(\mu) < \infty$ holds for every $\vartheta \in \theta \cup \{1\}$.

From (1), we obtain $p_{\vartheta, M, r}(\mu_n) < \infty$ for every $\vartheta \in \theta \cup \{1\}$, every $M > 0$, every $r > r_M$, and every $n \geq 1$. Thus, $\mu_n \in \mathcal{N}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)^{\mathbf{W}}$ for every $n \geq 1$. From (2), we obtain

$$\lim_{n \rightarrow \infty} p_{\vartheta, M, r}(\mu_n - \mu) = 0$$

for every $\vartheta \in \theta \cup \{1\}$, every $M > 0$, and every $r < r_M$. Then, by [Proposition 2.24](#) and [Remark 2.12](#), for every $f \in \mathcal{S}(G)$, $\{\mu_n \star f\}_{n \geq 1}$ is a Cauchy sequence in $\mathcal{S}(G)$; we denote its limit (in $\mathcal{S}(G)$) by $\mu \star f$. It follows immediately that $\pi(\mu \star f) = \mu(\chi_\pi) \cdot \pi(f)$ holds for every irreducible admissible representation π of G . The theorem is proved. $\square$

Remark 2.26. In fact, from the proof of [Theorem 2.13](#), we see that for every $\mu \in \mathcal{M}_{\theta \cup \{1\}}(\mathfrak{h}_{\mathbb{C}}^*)$, $\mu \star$ is continuous and preserves the subalgebra $\mathcal{S}(G)_{(K)}$ of bi- K -finite functions in $\mathcal{S}(G)$. Moreover, our argument of deducing [Theorem 2.13](#) from [Proposition 2.24](#) can be applied to deduce from [Proposition A.1](#) that every element of $\mathcal{M}(\mathfrak{h}_{\mathbb{C}}^*)$ gives a multiplier of $\mathcal{S}(G)_{(K)}$.

3. Isolation of spectrum

In this section, we state and prove various results on the isolation of cuspidal components of the L^2 -spectrum. In [Section 3.1](#), we introduce several important definitions and state the results. In [Section 3.2](#), we recall the coarse Langlands decomposition, cuspidal data, and cuspidal components. In [Section 3.3](#), we show how to annihilate all but finitely many cuspidal components. We finish the proof in [Section 3.4](#).

Let F be a number field.

3.1. Statement of results. We consider a connected reductive algebraic group G over F . Let $\mathfrak{g}$ be the complex Lie algebra of $G \otimes_{\mathbb{Q}} \mathbb{C}$, and let Z be the maximal F -split torus in the center of G . Let $\mathbf{S}_G$ be the set of primes of F such that G_v is ramified. We fix a maximal compact subgroup K_0 of $G(\mathbb{A}_F)$, and a Haar measure $dg = \prod_v dg_v$ on $G(\mathbb{A}_F)$, such that $K_{0,v}$ is hyperspecial maximal with volume 1 under dg_v for every prime v not in $\mathbf{S}_G$.

We first recall the definition of the L^2 -spectrum for G . Take a unitary automorphic character

$$\omega: Z(\mathbb{A}_F) \rightarrow \mathbb{C}^\times.$$

We define $L^2(G(F) \backslash G(\mathbb{A}_F), \omega)$ to be the L^2 completion of the subspace of smooth functions φ on $G(\mathbb{A}_F)$ satisfying

- $\varphi(z\gamma g) = \omega(z)\varphi(g)$ for every $z \in Z(\mathbb{A}_F)$, $\gamma \in G(F)$, and $g \in G(\mathbb{A}_F)$;
- $|\varphi|^2$, regarded as a function on $Z(\mathbb{A}_F)G(F) \backslash G(\mathbb{A}_F)$, is integrable.

The group $G(\mathbb{A}_F)$ acts on $L^2(G(F) \backslash G(\mathbb{A}), \omega)$ via the right regular representation R , which preserves the subspace $L^2_{\text{cusp}}(G(F) \backslash G(\mathbb{A}), \omega)$ of cuspidal functions.

Definition 3.1. We define the space of *Schwartz test functions* to be

$$\mathcal{S}(G(\mathbb{A}_F)) := \mathcal{S}(G_\infty) \otimes \mathcal{S}(G(\mathbb{A}_F^\infty)),$$

which is endowed with the convolution product with respect to the fixed Haar measure dg . Here, $\mathcal{S}(G(\mathbb{A}_F^\infty))$ denotes the restricted tensor product of $\mathcal{S}(G(F_v))$ for $v \nmid \infty$, where $\mathcal{S}(G(F_v))$ is nothing but $C_c^\infty(G(F_v))$.

The algebra $\mathcal{S}(G(\mathbb{A}_F))$ acts on $L^2(G(F) \backslash G(\mathbb{A}_F), \omega)$ continuously via the right regular representation R , with respect to the Haar measure dg .

Definition 3.2. We say a Schwartz test function $f \in \mathcal{S}(G(\mathbb{A}_F))$ is ω -*quasi-cuspidal* (or simply *quasi-cuspidal* when $\omega = 1$ is the trivial character) if the image of the endomorphism $R(f)$ on $L^2(G(F) \backslash G(\mathbb{A}_F), \omega)$ is contained in $L^2_{\text{cusp}}(G(F) \backslash G(\mathbb{A}_F), \omega)$.

For every prime $v \notin \mathbf{S}_G$, we let

$$\mathcal{H}_{G,v} := \mathbb{C}[K_{0,v} \backslash G(F_v) / K_{0,v}]$$

be the spherical Hecke algebra of G_v (with respect to $K_{0,v}$) with the unit $\mathbf{1}_{K_{0,v}}$. For a (possibly infinite) set $\mathbf{T}$ of primes of F containing $\mathbf{S}_G$, we let $\mathcal{H}_G^{\mathbf{T}}$ be the restricted tensor product of $\mathcal{H}_{G,v}$ for primes $v \notin \mathbf{T}$.

Definition 3.3. A $\mathbf{T}$ -character (for G) is a pair $\chi = (\chi_\infty, \chi^{\infty, \mathbf{T}})$ in which χ_∞ is a character of $\mathcal{Z}(\mathfrak{g})$ and $\chi^{\infty, \mathbf{T}}$ is a character of $\mathcal{H}_G^{\mathbf{T}}$.⁵

For a $\mathbf{T}$ -character χ and a prime $v \notin \mathbf{T}$, its v -component χ_v , which is a character of $\mathcal{H}_{G,v}$, gives rise to a $(K_{0,v})$ -spherical (irreducible admissible) representation of $G(F_v)$, unique up to isomorphism. For an irreducible admissible representation $\pi = \otimes_v \pi_v$ of $G(\mathbb{A}_F)$ such that $\pi_v^{K_{0,v}} \neq \{0\}$ for every prime $v \notin \mathbf{T}$, we have the induced $\mathbf{T}$ -character $\chi_{\pi^{\mathbf{T}}} = (\chi_{\pi_\infty}, \chi_{\pi^{\infty, \mathbf{T}}})$.

The following definition mimics the original notion of CAP representations by Piatetski-Shapiro [PS82]. In this section, we will only use it for $G' = G$, while the more general case will be used in [Section 4](#).

Definition 3.4. Let G' be an inner form of G (over F). We say that a $\mathbf{T}$ -character χ is $(G', \mathbf{T})$ -CAP if there exist a proper parabolic subgroup P' of G' and a cuspidal automorphic representation σ of $M_{P'}(\mathbb{A}_F)$, where $M_{P'} := P'/N_{P'}$, such that for all but finitely many primes v of F not in $\mathbf{T}$ for which $G_v \simeq G'_v$, the spherical representation corresponding to χ_v is a constituent of $\mathrm{I}_{P'}^{G'}(\sigma_v)$. For an irreducible admissible representation π of $G(\mathbb{A}_F)$, we say that π is $(G', \mathbf{T})$ -CAP if $\chi_{\pi^{\mathbf{T}}}$ is.

Now we fix

- a subset $\mathbf{T}$ of primes of F containing $\mathbf{S}_G$ and a $\mathbf{T}$ -character $\chi = (\chi_\infty, \chi^{\infty, \mathbf{T}})$;
- a finite set $\mathbf{S}$ of primes of F satisfying $\mathbf{S}_G \subseteq \mathbf{S} \subseteq \mathbf{T}$; and
- a subgroup $K \subseteq K_0^\infty$ of finite index of the form $K = K_{\mathbf{S}} \times \prod_{v \notin \mathbf{S}} K_{0,v}$.

We denote by $\mathcal{S}(G(\mathbb{A}_F^\infty))_K$ the subalgebra of $\mathcal{S}(G(\mathbb{A}_F^\infty))$ of bi- K -invariant functions, and we put

$$\mathcal{S}(G(\mathbb{A}_F))_K := \mathcal{S}(G_\infty) \otimes \mathcal{S}(G(\mathbb{A}_F^\infty))_K.$$

For $? = \emptyset, \text{cusp}$, let $L^2_?(G(F) \backslash G(\mathbb{A}_F) / K, \omega) \subseteq L^2_?(G(F) \backslash G(\mathbb{A}_F), \omega)$ be the subspace consisting of functions that are invariant under $R(K)$, on which $\mathcal{S}(G(\mathbb{A}_F))_K$ acts continuously via the right regular representation R . We denote by

$$L^2_{\text{cusp}}(G(F) \backslash G(\mathbb{A}_F) / K, \omega)[\chi] \subseteq L^2_{\text{cusp}}(G(F) \backslash G(\mathbb{A}_F) / K, \omega)$$

⁵Warning: $\mathbf{T}$ -characters are actually characters away from $\mathbf{T}$; same for $\mathbf{T}$ -multipliers later.

the maximal closed subspace of which $\mathcal{Z}(\mathfrak{g}) \otimes \mathcal{H}_G^T$ acts on its smooth vectors by the character $\chi_\infty \otimes \chi^{\infty, T}$.

The connected reductive group $G \otimes_{\mathbb{Q}} \mathbb{R}$ over $\mathbb{R}$ determines a root datum $(X^*, \Phi, X_*, \Phi^\vee)$ and a subset $\theta \subseteq \text{Aut}(X^*, \Phi, X_*, \Phi^\vee)^\vee$ as at the beginning of [Section 2.3](#). We adopt the notation from [Section 2.2](#). For an element $\mu \in \mathcal{O}(\mathfrak{h}_{\mathbb{C}}^*)^W \otimes \mathcal{H}_G^T$, we may evaluate μ on χ to obtain a complex number $\mu(\chi)$. [Theorem 2.13](#) provides us with a linear map $\mathcal{M}_\theta^\sharp(\mathfrak{h}_{\mathbb{C}}^*)^W \rightarrow \text{Mul}(\mathcal{S}(G_\infty))$ (see [Remark 2.17](#)).

Definition 3.5. A T -multiplier of $\mathcal{S}(G(\mathbb{A}_F))_K$ is an element in $\mathcal{M}_\theta^\sharp(\mathfrak{h}_{\mathbb{C}}^*)^W \otimes \mathcal{H}_G^T$.

As $\mathcal{H}_G^T$ is contained in the center of $\mathcal{S}(G(\mathbb{A}_F^\infty))_K$, a T -multiplier μ induces a multiplier

$$\mu \star \in \text{Mul}(\mathcal{S}(G(\mathbb{A}_F))_K)$$

of $\mathcal{S}(G(\mathbb{A}_F))_K$. Now we can state our main results on the isolation of the spectrum.

THEOREM 3.6. *Suppose that χ is not (G, T) -CAP ([Definition 3.4](#)). Then there exists a T -multiplier μ of $\mathcal{S}(G(\mathbb{A}_F))_K$ such that for every $f \in \mathcal{S}(G(\mathbb{A}_F))_K$,*

- (1) $R(\mu \star f)$ maps $L^2(G(F) \backslash G(\mathbb{A}_F)/K, \omega)$ into $L_{\text{cusp}}^2(G(F) \backslash G(\mathbb{A}_F)/K, \omega)[\chi]$;
- (2) $\mu(\chi) = 1$.

In particular, $\mu \star f$ is an ω -quasi-cuspidal Schwartz test function ([Definition 3.2](#)).

In particular, [Theorem 1.1](#) follows by taking $T = S$ and $\chi = \chi_{\pi^T}$.

For general χ , we have the following theorem. In fact, we will prove a stronger result in [Theorem 3.19](#) below.

THEOREM 3.7. *There exists a T -multiplier μ of $\mathcal{S}(G(\mathbb{A}_F))_K$ such that for every $f \in \mathcal{S}(G(\mathbb{A}_F))_K$,*

- (1) $R(\mu \star f)$ maps $L_{\text{cusp}}^2(G(F) \backslash G(\mathbb{A}_F)/K, \omega)$ into $L_{\text{cusp}}^2(G(F) \backslash G(\mathbb{A}_F)/K, \omega)[\chi]$;
- (2) $\mu(\chi) = 1$.

Remark 3.8. In the above two theorems, we do not require that $\chi = \chi_{\pi^T}$ for an irreducible admissible representation π of $G(\mathbb{A}_F)$.

The remaining part of this section will be devoted to the proof of the above two theorems.

3.2. Cuspidal data and cuspidal components. We recall the notion of cuspidal data and cuspidal components for the group G . We fix

- a minimal Levi subgroup M_0 of (a parabolic subgroup of) G ;
- a minimal parabolic subgroup P_0 of G containing M_0 ; and
- a maximal torus T of $G \otimes_{\mathbb{Q}} \mathbb{R}$ over $\mathbb{R}$ that is contained in $M_0 \otimes_{\mathbb{Q}} \mathbb{R}$.

Without loss of generality, we assume that the fixed maximal compact subgroup K_0 of $G(\mathbb{A}_F)$ is admissible relative to $M_0(\mathbb{A}_F)$ (in the sense of [Art81, §1]). We identify X^* with the weight lattice of $T_{\mathbb{C}}$. Let $W(G, M_0)$ be the Weyl group of the pair (G, M_0) . We say that a subgroup M of G is a *standard Levi subgroup* if there exists a parabolic subgroup of G containing P_0 , of which M is the unique Levi subgroup containing M_0 .

For a standard Levi subgroup M of G , we denote by

- Z_M the center of M (in particular, Z is the maximal F -split torus in Z_G);
- P_M the unique parabolic subgroup of G containing P_0 , of which M is a Levi subgroup;
- $\Omega_M(\omega)$ the set of unitary automorphic characters $\omega_M: Z_M(\mathbb{A}_F) \rightarrow \mathbb{C}^\times$ satisfying $\omega_M|_{Z(\mathbb{A}_F)} = \omega$;
- $M(\mathbb{A}_F)^1$ the intersection of the kernels of all automorphic characters $M(\mathbb{A}_F) \rightarrow \mathbb{R}_+^\times$, and
- $\mathfrak{a}_M$ the real vector space $Z(\mathbb{A}_F)M(\mathbb{A}_F)^1 \backslash M(\mathbb{A}_F)$.⁶

For $s \in \mathfrak{a}_{M, \mathbb{C}}^*$, we denote by $\xi_s: Z(\mathbb{A}_F)M(\mathbb{A}_F)^1 \backslash M(\mathbb{A}_F) \rightarrow \mathbb{C}^\times$ the corresponding (automorphic) character obtained by composing s with the exponential map $\exp: \mathbb{C} \rightarrow \mathbb{C}^\times$, which is unitary if and only if $s \in i\mathfrak{a}_M^*$. For an admissible representation σ of $M(\mathbb{A}_F)$, we put $\sigma_s := \sigma \otimes \xi_s$ for $s \in \mathfrak{a}_{M, \mathbb{C}}^*$.

Definition 3.9. For a cuspidal automorphic representation σ of $M(\mathbb{A}_F)$ with central character $\omega_M \in \Omega_M(\omega)$, we denote by

$$L^2(M, \sigma) \subseteq L_{\text{cusp}}^2(M(F) \backslash M(\mathbb{A}_F), \omega_M)$$

the maximal closed σ -isotypic subspace.

We define $\mathfrak{C}(M, \omega)$ to be the set of isomorphism classes of cuspidal automorphic representations of $M(\mathbb{A}_F)$ whose central character belongs to $\Omega_M(\omega)$, and an equivalence relation $\sim$ on $\mathfrak{C}(M, \omega)$ by the following rule: $\sigma \sim \sigma'$ if there exists $s \in i\mathfrak{a}_M^*$ such that $\sigma' = \sigma_s$. Let $\mathfrak{C}(M, \omega)^\heartsuit$ be the quotient of $\mathfrak{C}(M, \omega)$ by $\sim$.

We define $\mathfrak{D}(G, \omega)$ to be the set of pairs (M, σ) , where M is a standard Levi subgroup of G and $\sigma \in \mathfrak{C}(M, \omega)$, and an equivalence relation $\approx$ on $\mathfrak{D}(G, \omega)$ by the following rule: $(M, \sigma) \approx (M', \sigma')$ if there exists $w \in W(G, M_0)$ such that $M' = M^w$ and $\sigma' \sim \sigma^w$ in the sense above. We denote by $\mathfrak{D}(G, \omega)^\heartsuit$ the quotient of $\mathfrak{D}(G, \omega)$ by $\approx$.

By [Lan76, Lemma 4.6(i)] or [MW95, II.2.4], we have the decomposition

$$(3.1) \quad L^2(G(F) \backslash G(\mathbb{A}_F), \omega) = \widehat{\bigoplus_{(M, \sigma) \in \mathfrak{D}(G, \omega)^\heartsuit} L_{(M, \sigma)}^2(G(F) \backslash G(\mathbb{A}_F), \omega)}$$

⁶In fact, $\mathfrak{a}_M$ is usually denoted by $\mathfrak{a}_M^G$ (for example in [Art05, Art13]); it is the real Lie algebra of the maximal $\mathbb{Q}$ -split torus contained in $\text{Res}_{F/\mathbb{Q}} Z \backslash M$.

of Hilbert spaces, known as the coarse Langlands decomposition. We call an element (M, σ) in $\mathfrak{D}(G, \omega)^\vee$ a *cuspidal datum*, and we call $L^2_{(M, \sigma)}(G(F) \backslash G(\mathbb{A}_F), \omega)$ the (*automorphic*) *cuspidal component* associated to (M, σ) .

For the readers' convenience, we recall the construction of the cuspidal component $L^2_{(M, \sigma)}(G(F) \backslash G(\mathbb{A}_F), \omega)$. For $(M, \sigma) \in \mathfrak{D}(G, \omega)$, we denote by $\mathcal{A}^G_{(M, \sigma)}$ the space of $R(K_0)$ -finite smooth functions ϕ on $N_{P_M}(\mathbb{A}_F) \backslash G(\mathbb{A}_F)$ satisfying that for every $x \in K_0$, the function $\phi_x: m \mapsto \delta_{P_M}^{-1/2}(m)\phi(mx)$ is an automorphic form on $M(\mathbb{A}_F)$ contained in $L^2(M, \sigma)$, where δ_{P_M} is the modulus character of P_M . For every Paley–Wiener function

$$\Phi: \mathfrak{a}^*_{M, \mathbb{C}} \rightarrow \mathcal{A}^G_{(M, \sigma)}$$

valued in a finite dimensional subspace of $\mathcal{A}^G_{(M, \sigma)}$, we put

$$\tilde{\Phi}(g) := \int_{i\mathfrak{a}^*_M} \Phi_s(g) \cdot \xi_s(m(g)) \, ds$$

for $g \in P(F) \backslash G(\mathbb{A}_F)$, where $m(g) \in Z(\mathbb{A}_F)M(\mathbb{A}_F)^1 \backslash M(\mathbb{A}_F)$ denotes the image of the $M(\mathbb{A}_F)$ -component of g under the Iwasawa decomposition $G(\mathbb{A}_F) = N_{P_M}(\mathbb{A}_F)M(\mathbb{A}_F)K_0$. We have the *pseudo-Eisenstein series*

$$E(g, \Phi) := \sum_{\gamma \in P(F) \backslash G(F)} \tilde{\Phi}(\gamma g),$$

which belongs to $L^2(G(F) \backslash G(\mathbb{A}_F), \omega)$. Then $L^2_{(M, \sigma)}(G(F) \backslash G(\mathbb{A}_F), \omega)$ is defined to be the closure of the subspace spanned by $E(-, \Phi)$ for all Paley–Wiener functions Φ as above. We have that

$$L^2_{(M, \sigma)}(G(F) \backslash G(\mathbb{A}_F), \omega) = L^2_{(M', \sigma')}(G(F) \backslash G(\mathbb{A}_F), \omega)$$

if and only if $(M, \sigma) \approx (M', \sigma')$.

Let $L^2_{(M, \sigma)}(G(F) \backslash G(\mathbb{A}_F)/K, \omega) \subseteq L^2_{(M, \sigma)}(G(F) \backslash G(\mathbb{A}_F), \omega)$ be the subspace consisting of functions that are invariant under $R(K)$. Taking $R(K)$ -invariants, (3.1) induces the following decomposition of Hilbert spaces:

$$(3.2) \quad L^2(G(F) \backslash G(\mathbb{A}_F)/K, \omega) = \widehat{\bigoplus_{(M, \sigma) \in \mathfrak{D}(G, \omega)^\vee} L^2_{(M, \sigma)}(G(F) \backslash G(\mathbb{A}_F)/K, \omega)}.$$

3.3. Annihilation of all but finitely many components. We start to prove the results in Section 3.1. In particular, we fix the T-character $\chi = (\chi_\infty, \chi^{\infty, T})$, the subsets $\mathbf{S}_G \subseteq \mathbf{S} \subseteq \mathbf{T}$ of primes of F , and the subgroup $K \subseteq K_0^\infty$. In this subsection, we construct an element $\mu_\infty^\chi \in \mathcal{M}_\theta^\#(\mathfrak{h}_\mathbb{C}^*)^W$ satisfying $\mu_\infty^\chi(\chi_\infty) \neq 0$ and such that $R(\mu_\infty^\chi \star f)$ annihilates all but finitely many cuspidal components in (3.2).

Take a standard Levi subgroup M of G . Let M_{der} be the derived subgroup of M . Let $\mathbf{X}_M^*$ be the weight lattice of $(T \cap M_{\text{der}} \otimes_{\mathbb{Q}} \mathbb{R})_{\mathbb{C}}$, and put $\mathfrak{b}_M^* := \mathbf{X}_M^* \otimes_{\mathbb{Z}} \mathbb{R}$.

Then we have a canonical map $X^* \rightarrow X_M^*$, which induces a surjective map $\mathfrak{h}^* \rightarrow \mathfrak{b}_M^*$. Denote by $\mathfrak{c}_M^*$ the kernel of $\mathfrak{h}^* \rightarrow \mathfrak{b}_M^*$, which canonically contains $\mathfrak{a}_M^*$.

Remark 3.10. In fact, $\mathfrak{c}_M^*/\mathfrak{a}_M^*$ is canonically the real cotangent space of $Z_M(\mathbb{A}_F)^1 Z(\mathbb{A}_F)$ at the identity. Thus, $\mathfrak{c}_M^* = \mathfrak{a}_M^*$ holds if and only if $Z = 1$ and $Z_M(\mathbb{A}_F)^1$ is discrete, and $\mathfrak{c}_M^* = \mathfrak{a}_M^*$ holds for every standard Levi subgroup M if and only if G is semisimple and split over $\mathbb{Q}$.

Denote by $\gamma_M: \mathfrak{h}^* \rightarrow \mathfrak{h}^*/\mathfrak{a}_M^*$ the quotient map. We have a canonical decomposition $\mathfrak{h}^* = \mathfrak{c}_M^* \oplus \mathfrak{b}_M^*$, which gives rise to two linear maps

$$\gamma_M^+: \mathfrak{h}^* \rightarrow \mathfrak{c}_M^*/\mathfrak{a}_M^*, \quad \gamma_M^-: \mathfrak{h}^* \rightarrow \mathfrak{b}_M^*,$$

so that $\gamma_M = \gamma_M^+ \oplus \gamma_M^-$. Let W_M be the Weyl group of the pair $(M \otimes_{\mathbb{Q}} \mathbb{C}, T_{\mathbb{C}})$, which is canonically a subgroup of W and acts on $\mathfrak{c}_M^*$ trivially. For every element $\sigma \in \mathfrak{C}(M, \omega)$ and every $s \in \mathfrak{a}_{M, \mathbb{C}}^*$, the infinitesimal character $\chi_{\sigma s, \infty}$ is a W_M -orbit in $\mathfrak{h}_{\mathbb{C}}^*$, satisfying that $\gamma_M(\chi_{\sigma s, \infty}) \subseteq \mathfrak{h}_{\mathbb{C}}^*/\mathfrak{a}_{M, \mathbb{C}}^*$ does not depend on s , hence only on the class of σ in $\mathfrak{C}(M, \omega)^{\heartsuit}$. We also denote by $\chi_{\sigma s, \infty}^G$ the infinitesimal character of $\text{Ind}_{P_M}^G(\chi_{\sigma s, \infty})$, which is simply the W -orbit of $\chi_{\sigma s, \infty}$ in $\mathfrak{h}_{\mathbb{C}}^*$.

The Casimir operator for $M_{\text{der}} \otimes_{\mathbb{Q}} \mathbb{R}$ defines a map

$$\lambda_M: \mathfrak{b}_{M, \mathbb{C}}^* \rightarrow \mathbb{C},$$

which is a W_M -invariant polynomial function.

Definition 3.11. We say that an element $\sigma \in \mathfrak{C}(M, \omega)^{\heartsuit}$ is χ_{∞} -typical if both

- $\gamma_M^+(\chi_{\sigma \infty}) \in \gamma_M^+(\chi_{\infty})$ and
- $\lambda_M(\gamma_M^-(\chi_{\sigma \infty})) \in \lambda_M(\gamma_M^-(\chi_{\infty}))$

hold. Denote by $\mathfrak{C}(M, \omega)_{\chi_{\infty}}^{\heartsuit}$ the subset of $\mathfrak{C}(M, \omega)^{\heartsuit}$ consisting of χ_{∞} -typical elements.

Informally speaking, χ_{∞} -typical elements are those whose infinitesimal characters cannot be distinguished from χ_{∞} via the two maps γ_M^+ and $\lambda_M \circ \gamma_M^-$, hence their associated cuspidal components will not be annihilated by the method described below in this subsection.

Now we start the construction of μ_{∞} , following the strategy (the first two steps) described in [Section 1.3](#). We first choose an element $\mu_{\infty}^0 \in \mathcal{M}_{\theta}^{\sharp}(\mathfrak{h}_{\mathbb{C}}^*)^W$ such that $\mu_{\infty}^0(\chi_{\infty}) \neq 0$, which is possible by [Lemma 2.10](#). We denote by $\mathfrak{T} := \mathfrak{T}(\mu_{\infty}^0)$ the finite set of $K_{0, \infty}$ -types from [Lemma 2.18](#). For every standard Levi subgroup M of G , we

- put $K_{0, \infty}^M := M_{\infty} \cap K_{0, \infty}$, which is a maximal compact subgroup of M_{∞} ;
- fix a finite set $\mathfrak{T}_M$ of $K_{0, \infty}^M$ -types satisfying the following property: if σ is an irreducible admissible representation of M_{∞} such that $I_{P_M}^G(\sigma)|_{K_{\infty}}$ contains

a member from $\mathfrak{T}$, then $\sigma|_{K_{0,\infty}^M}$ contains a member from $\mathfrak{T}_M$ (when $M = G$, we take $\mathfrak{T}_M = \mathfrak{T}$);

- fix an open compact subgroup $K_{M,\mathbf{S}}$ of $M(F_{\mathbf{S}})$ satisfying the following property: if σ is an irreducible admissible representation of $M(F_{\mathbf{S}})$ such that $I_{P_M}^G(\sigma)^{K_{\mathbf{S}}} \neq \{0\}$, then $\sigma^{K_{M,\mathbf{S}}} \neq \{0\}$ (when $M = G$, we take $K_{M,\mathbf{S}} = K_{\mathbf{S}}$);
- put $K_{M,v} := M(F_v) \cap K_v$ for every prime v of F not in $\mathbf{S}$, which is a hyperspecial maximal subgroup of $M(F_v)$;
- put $K_M := K_{M,\mathbf{S}} \times \prod_{v \notin \mathbf{S}} K_{M,v}$.

Let $\mathfrak{C}(M, \omega; K_M, \mathfrak{T}_M)$ be the subset of $\mathfrak{C}(M, \omega)$ consisting of σ satisfying $\sigma^{K_M} \neq \{0\}$ and that $\sigma|_{K_{0,\infty}^M}$ contains a member from $\mathfrak{T}_M$. It is clear that $\mathfrak{C}(M, \omega; K_M, \mathfrak{T}_M)$ is closed under the equivalence relation $\sim$ in [Definition 3.9](#). Let $\mathfrak{C}(M, \omega; K_M, \mathfrak{T}_M)^\heartsuit$ be the quotient of $\mathfrak{C}(M, \omega; K_M, \mathfrak{T}_M)$ by $\sim$.

Put $K_{M_{\text{der}}} := K_M \cap M_{\text{der}}(\mathbb{A}_F^\infty)$ and $K_{0,\infty}^{M_{\text{der}}} := K_{0,\infty}^M \cap M_{\text{der},\infty}$. Let $\mathfrak{T}_{M_{\text{der}}}$ be the (finite) set of $K_{0,\infty}^{M_{\text{der}}}$ -types that appear in the restriction of members in $\mathfrak{T}_M$ to $K_{0,\infty}^{M_{\text{der}}}$. Denote by $\mathfrak{C}(M_{\text{der}}; K_{M_{\text{der}}}, \mathfrak{T}_{M_{\text{der}}})$ the set of isomorphism classes of cuspidal automorphic representations ς of $M_{\text{der}}(\mathbb{A}_F)$ satisfying $\varsigma^{K_{M_{\text{der}}}} \neq \{0\}$ and that $\varsigma|_{K_{0,\infty}^{M_{\text{der}}}}$ contains a member from $\mathfrak{T}_{M_{\text{der}}}$.

The following lemma is crucial for us to construct functions in $\mathcal{M}(\mathfrak{h}_{\mathbb{C}}^*)^W$ that vanish on χ_{σ_∞} for σ not χ_∞ -typical.

LEMMA 3.12. *The subset*

$$\Lambda(M, \omega; K_M, \mathfrak{T}_M) := \{\lambda_M(\gamma_M^-(\chi_{\sigma_\infty})) \mid \sigma \in \mathfrak{C}(M, \omega; K_M, \mathfrak{T}_M)\} \subseteq \mathbb{C}$$

is of finite rank ([Definition 2.1](#)).

Proof. It suffices to show that the subset of Casimir eigenvalues of elements in $\mathfrak{C}(M_{\text{der}}; K_{M_{\text{der}}}, \mathfrak{T}_{M_{\text{der}}})$ is of finite rank. As $\mathfrak{T}_{M_{\text{der}}}$ is a finite set, this is a direct consequence of [[Don82](#), Th. 9.1]. $\square$

The following lemma shows that there are only finitely many χ_∞ -typical elements in $\mathfrak{C}(M, \omega; K_M, \mathfrak{T}_M)^\heartsuit$.

LEMMA 3.13. *The set $\mathfrak{C}(M, \omega; K_M, \mathfrak{T}_M)^\heartsuit \cap \mathfrak{C}(M, \omega)_{\chi_\infty}^\heartsuit$ is finite.*

Proof. There are only finitely many elements in $\mathfrak{C}(M_{\text{der}}; K_{M_{\text{der}}}, \mathfrak{T}_{M_{\text{der}}})$ whose Casimir eigenvalue belongs to the set $\lambda_M(\gamma_M^-(\chi_\infty))$ by [[Don82](#), Th. 9.1]. It follows that there are only finitely many elements in $\mathfrak{C}(M, \omega; K_M, \mathfrak{T}_M)^\heartsuit$ that are χ_∞ -typical. In other words, the set $\mathfrak{C}(M, \omega; K_M, \mathfrak{T}_M)^\heartsuit \cap \mathfrak{C}(M, \omega)_{\chi_\infty}^\heartsuit$ is finite. $\square$

The following lemma achieves the goal in the second step of the strategy described in [Section 1.3](#).

LEMMA 3.14. *For every standard Levi subgroup M of G , there exists an element $\mu_\infty^M \in \mathcal{M}_\theta^\sharp(\mathfrak{h}_\mathbb{C}^*)^W$ satisfying $\mu_\infty^M(\chi_\infty) \neq 0$, and such that for every $\sigma \in \mathfrak{C}(M, \omega; K_M, \mathfrak{T}_M)^\heartsuit \setminus \mathfrak{C}(M, \omega)_{\chi_\infty}^\heartsuit$, μ_∞^M vanishes on $\chi_{\sigma s, \infty}^G$, the infinitesimal character of $\text{Ind}_{P_M}^G(\sigma s, \infty)$, for every $s \in \mathfrak{a}_{M, \mathbb{C}}^*$.*

Proof. We first consider the special case where $\mathfrak{c}_M = \mathfrak{a}_M$. In this case, we have $\gamma_M = \gamma_M^-: \mathfrak{h}^* \rightarrow \mathfrak{b}_M^*$. By Lemma 3.12, we may apply Corollary 2.5 to the subset $\Lambda = \Lambda(M, \omega; K_M, \mathfrak{T}_M) \setminus \lambda_M(\gamma_M(\chi_\infty))$, hence obtain a holomorphic function ν on $\mathfrak{b}_{M, \mathbb{C}}^*$ that has moderate vertical growth, and vanishes exactly on $\lambda_M^{-1}(\Lambda(M, \omega; K_M, \mathfrak{T}_M) \setminus \lambda_M(\gamma_M(\chi_\infty)))$. We regard ν as a function on $\mathfrak{h}_\mathbb{C}^*$ via the quotient map $\gamma_M: \mathfrak{h}_\mathbb{C}^* \rightarrow \mathfrak{b}_{M, \mathbb{C}}^*$. By construction, ν is nowhere vanishing on χ_∞ , and vanishes on $\chi_{\sigma s, \infty}$ for every $\sigma \in \mathfrak{C}(M, \omega; K_M, \mathfrak{T}_M)^\heartsuit \setminus \mathfrak{C}(M, \omega)_{\chi_\infty}^\heartsuit$ and $s \in \mathfrak{a}_{M, \mathbb{C}}^*$. Finally, put

$$\mu_\infty^M := \mu_\infty^0 \cdot \prod_{w \in W} \nu \circ w,$$

which belongs to $\mathcal{M}_\theta^\sharp(\mathfrak{h}_\mathbb{C}^*)^W$ by Remark 2.9(3); it satisfies the requirements in the lemma.

Now we treat the general case where $\mathfrak{c}_M/\mathfrak{a}_M$ might be non-trivial. The extra work is to treat the factor γ_M^+ . For every element $\sigma \in \mathfrak{C}(M, \omega; K_M, \mathfrak{T}_M)^\heartsuit$, the restriction of the central character of σ_s to $Z_M(\mathbb{A}_F)^1 Z(\mathbb{A}_F)$, say ω_σ , is independent of s . Moreover, the infinitesimal character of $\omega_{\sigma s, \infty}$, which is canonically an element in $\mathfrak{c}_{M, \mathbb{C}}^*/\mathfrak{a}_{M, \mathbb{C}}^*$ by Remark 3.10, is simply $\gamma_M^+(\chi_{\sigma s, \infty})$. Since $\omega_\sigma|_{Z(\mathbb{A}_F)} = \omega$, ω_σ is invariant under $Z_M(F) \cdot (K_M \cap Z_M(\mathbb{A}_F^\infty))$, and $Z_M(F) \backslash Z_M(\mathbb{A}_F)^1 / K_M \cap Z_M(\mathbb{A}_F^\infty)$ is a compact abelian real Lie group, there exists a lattice L_M of $\mathfrak{c}_{M, \mathbb{C}}^*/\mathfrak{a}_{M, \mathbb{C}}^*$ such that $\gamma_M^+(\chi_{\sigma s, \infty})$ is contained in L_M for every $\sigma \in \mathfrak{C}(M, \omega; K_M, \mathfrak{T}_M)^\heartsuit$. Applying Corollary 2.6 to $U = \mathfrak{c}_M^*/\mathfrak{a}_M^*$, $L = L_M$, and $A = \gamma_M^+(\chi_\infty)$, we obtain a holomorphic function ν^+ on $\mathfrak{c}_{M, \mathbb{C}}^*/\mathfrak{a}_{M, \mathbb{C}}^*$ that has moderate vertical growth, vanishes on $L_M \setminus \gamma_M^+(\chi_\infty)$, and is nowhere vanishing on $\gamma_M^+(\chi_\infty)$. On the other hand, similar to the special case, we have a holomorphic function ν^- on $\mathfrak{b}_{M, \mathbb{C}}^*$ that has moderate vertical growth, and vanishes exactly on $\lambda_M^{-1}(\Lambda(M, \omega; K_M, \mathfrak{T}_M) \setminus \lambda_M(\gamma_M^-(\chi_\infty)))$. Put

$$\nu := \nu^+ \times \nu^-: \mathfrak{c}_{M, \mathbb{C}}^*/\mathfrak{a}_{M, \mathbb{C}}^* \oplus \mathfrak{b}_{M, \mathbb{C}}^* \rightarrow \mathbb{C},$$

which we regard as a function on $\mathfrak{h}_\mathbb{C}^*$ via the quotient map $\mathfrak{h}_\mathbb{C}^* \rightarrow \mathfrak{h}_\mathbb{C}^*/\mathfrak{a}_{M, \mathbb{C}}^*$. As in the special case, we put

$$\mu_\infty^M := \mu_\infty^0 \cdot \prod_{w \in W} \nu \circ w,$$

which belongs to $\mathcal{M}_\theta^\sharp(\mathfrak{h}_\mathbb{C}^*)^W$ and satisfies the requirements in the lemma. $\square$

By definition, we have an identification $\coprod_M \mathfrak{C}(M, \omega) = \mathfrak{D}(G, \omega)$, which induces a surjective map

$$(3.3) \quad \coprod_M \mathfrak{C}(M, \omega)^\heartsuit \rightarrow \mathfrak{D}(G, \omega)^\heartsuit$$

by passing to the equivalence relations. We denote by $\mathfrak{D}(G, \omega; K, \mathfrak{T})^\heartsuit$ the image of

$$\coprod_M \mathfrak{C}(M, \omega; K_M, \mathfrak{T}_M)^\heartsuit$$

under the map (3.3), and by $\mathfrak{D}(G, \omega; K, \mathfrak{T})_{\chi_\infty}^\heartsuit$ we denote the image of

$$\coprod_M \mathfrak{C}(M, \omega; K_M, \mathfrak{T}_M)^\heartsuit \cap \mathfrak{C}(M, \omega)_{\chi_\infty}^\heartsuit$$

under the map (3.3). Now we can construct the desired function μ_∞ in the following proposition.

PROPOSITION 3.15. *The function*

$$\mu_\infty := \mu_\infty^0 \cdot \prod_M \mu_\infty^M,$$

in which the product is taken over standard Levi subgroups of G and μ_∞^M is a function obtained from Lemma 3.14,⁷ belongs to $\mathcal{M}_\theta^\#(\mathfrak{h}_\mathbb{C}^*)^\mathbb{W}$ and satisfies $\mu_\infty(\chi_\infty) \neq 0$. Moreover, for every $f \in \mathcal{S}(G(\mathbb{A}_F))_K$, the endomorphism $R(\mu_\infty \star f)$ of $L^2(G(F) \backslash G(\mathbb{A}_F)/K, \omega)$ annihilates the subspace $L^2_{(M, \sigma)}(G(F) \backslash G(\mathbb{A}_F)/K, \omega)$ if (M, σ) does not belong to $\mathfrak{D}(G, \omega; K, \mathfrak{T})_{\chi_\infty}^\heartsuit$.

Note that by Lemma 3.13, $\mathfrak{D}(G, \omega; K, \mathfrak{T})_{\chi_\infty}^\heartsuit$ is a finite set.

Proof. The first statement is clear from the construction.

By the description of $L^2_{(M, \sigma)}(G(F) \backslash G(\mathbb{A}_F)/K, \omega)$ recalled in the last subsection, it suffices to show that $R(\mu_\infty \star f)E(-, \Phi) = 0$ for every Paley–Wiener function Φ . By Remark 2.26, it suffices to consider the case where f , hence $\mu_\infty \star f$, are bi- $K_{0, \infty}$ -finite. Then since $R(\mu_\infty \star f)E(-, \Phi) = E(-, R(\mu_\infty \star f)\Phi)$, it suffices to show that $\text{Ind}_{P_M}^G(\sigma_s)(\mu_\infty \star f) = 0$ for every $s \in \mathfrak{a}_{M, \mathbb{C}}^*$. Note that we have

$$\text{Ind}_{P_M}^G(\sigma_s)(\mu_\infty \star f) = \mu_\infty(\chi_{\sigma_{s, \infty}}^G) \cdot \text{Ind}_{P_M}^G(\sigma_s)(f).$$

Take an element $(M, \sigma) \in \mathfrak{D}(G, \omega)$ whose image in $\mathfrak{D}(G, \omega)^\heartsuit$ does not belong to $\mathfrak{D}(G, \omega; K, \mathfrak{T})_{\chi_\infty}^\heartsuit$. If $\sigma^{K_M} = \{0\}$, then we have $\text{Ind}_{P_M}^G(\sigma_s)(f) = 0$ by our choice of K_M . If $\sigma|_{K_{0, \infty}^M}$ does not contain any member from $\mathfrak{T}_M$, then we have $\mu_\infty(\chi_{\sigma_{s, \infty}}^G) = \mu_\infty^0(\chi_{\sigma_{s, \infty}}^G) = 0$ for every $s \in \mathfrak{a}_{M, \mathbb{C}}^*$ by our choice of $\mathfrak{T}_M$.

⁷In fact, it suffices to take the product over a set of representatives of standard Levi subgroups of G with respect to the conjugation action of $W(G, M_0)$.

Otherwise, we must have $\mu_\infty(\chi_{\sigma_{s,\infty}}^G) = 0$ for every $s \in \mathfrak{a}_{M,\mathbb{C}}^*$ by the property of μ_∞^M from Lemma 3.14. The proposition follows. $\square$

3.4. *Proof of results.* To annihilate components $L_{(M,\sigma)}^2(G(F)\backslash G(\mathbb{A}_F)/K, \omega)$ in which σ is possibly χ_∞ -typical, mixed multipliers from both archimedean and non-archimedean places are needed.

Definition 3.16. For an element $(M, \sigma) \in \mathfrak{D}(G, \omega)$, we say the $\mathbf{T}$ -character χ occurs in (M, σ) if there exists $s \in \mathfrak{a}_{M,\mathbb{C}}^*$ such that

- $\chi_{\sigma_{s,\infty}}^G = \chi_\infty$;
- for every prime v of F not in $\mathbf{T}$, the spherical representation corresponding to χ_v is a constituent of $\text{Ind}_{P_M}^G(\sigma_{s,v})$.

It is clear that such property depends only on the equivalence class of (M, σ) in $\mathfrak{D}(G, \omega)^\heartsuit$. We denote by $\mathfrak{D}(G, \omega)_\chi^\heartsuit$ the subset of $\mathfrak{D}(G, \omega)^\heartsuit$ in which χ occurs.

We need to annihilate every element $(M, \sigma) \in \mathfrak{D}(G, \omega)$ in which χ does not occur, which is possible by the following proposition.

PROPOSITION 3.17. *Let (M, σ) be an element in $\mathfrak{D}(G, \omega)$. Assume that χ does not occur in (M, σ) . Then there exists a $\mathbf{T}$ -multiplier $\mu_{(M,\sigma)}$ of $\mathcal{S}(G(\mathbb{A}_F))_K$ satisfying*

- (1) *the endomorphism $R(\mu_{(M,\sigma)} \star f)$ of $L^2(G(F)\backslash G(\mathbb{A}_F)/K, \omega)$, for every $f \in \mathcal{S}(G(\mathbb{A}_F))_K$, annihilates $L_{(M,\sigma)}^2(G(F)\backslash G(\mathbb{A}_F)/K, \omega)$;*
- (2) $\mu_{(M,\sigma)}(\chi) \neq 0$.

Proof. We may assume that $\sigma^{K_M} \neq \{0\}$; otherwise

$$L_{(M,\sigma)}^2(G(F)\backslash G(\mathbb{A}_F)/K, \omega) = \{0\},$$

hence the proposition is trivial. Then for every $s \in \mathfrak{a}_{M,\mathbb{C}}^*$ and every prime v of F not in $\mathbf{T}$, we have $\text{I}_{P_M}^G(\sigma_{s,v})^{K_v} \neq \{0\}$, which gives rise to a character $\chi_{\sigma_{s,v}}^G : \mathcal{H}_{G,v} \rightarrow \mathbb{C}$. Then we have a character

$$\chi_{\sigma_\infty, \mathbf{T}}^G : \mathcal{H}_G^\mathbf{T} \rightarrow \mathbb{C}.$$

We also recall the infinitesimal character $\chi_{\sigma_{s,\infty}}^G$ of $\text{I}_{P_M}^G(\sigma_{s,\infty})$, which is simply the W -orbit of $\chi_{\sigma_{s,\infty}}$ in $\mathfrak{h}_{\mathbb{C}}^*$. Thus, we obtain a $\mathbf{T}$ -character $\chi_{\sigma_s^\mathbf{T}}^G = (\chi_{\sigma_{s,\infty}}^G, \chi_{\sigma_\infty, \mathbf{T}}^G)$ for G . We suppress the superscript G when $M = G$.

We first consider the easy case where $M = G$. Since χ does not occur in (G, σ) , we have either $\chi_{\sigma_\infty} \neq \chi_\infty$ or $\chi_{\sigma_\infty, \mathbf{T}} \neq \chi_\infty^\mathbf{T}$. In the first case, by Lemma 2.10, we can find an element $\mu_{(M,\sigma)} \in \mathcal{M}_\theta^\sharp(\mathfrak{h}_{\mathbb{C}}^*)^W$ satisfying $\mu_{(M,\sigma)}(\chi_\infty) \neq 0$. After multiplying by a suitable element in $\mathbb{C}[\mathfrak{h}_{\mathbb{C}}^*]^W$, we may further require that $\mu_{(M,\sigma)}(\chi_{\sigma_\infty}) = 0$. In the second case, we can certainly find

an element $\mu_{(M,\sigma)} \in \mathcal{H}_G^T$ satisfying $\mu_{(M,\sigma)}(\chi_{\sigma^\infty, \tau}) = 0$ but $\mu_{(M,\sigma)}(\chi^{\infty, \tau}) \neq 0$. The proposition follows.

We now consider the hard case where M is a proper standard Levi subgroup. Recall that $\mathfrak{a}_M^*$ is canonically a subspace of $\mathfrak{h}^*$. We fix

- an element α_σ in $\chi_{\sigma^\infty} \subseteq \mathfrak{h}_\mathbb{C}^*$,
- an element α_χ in $\chi_\infty \subseteq \mathfrak{h}_\mathbb{C}^*$,
- a linear splitting map $\ell: \mathfrak{h}^* \rightarrow \mathfrak{a}_M^*$ of the subspace $\mathfrak{a}_M^* \subseteq \mathfrak{h}^*$.

For $w \in W$, put $s_w := \ell(w\alpha_\chi) - \ell(\alpha_\sigma) \in \mathfrak{a}_{M,\mathbb{C}}^*$. Since χ does not occur in (M, σ) , for every $w \in W$, we can take $v[w]$ to be either ∞ or a prime of F not in T such that $\chi_{\sigma_{s_w, v[w]}^G} \neq \chi_{v[w]}$. It allows us to choose an element $\nu_w \in \mathcal{H}_{G, v[w]}$ (we regard $\mathcal{H}_{G, \infty}$ as $\mathbb{C}[\mathfrak{h}_\mathbb{C}^*]^W$) such that

$$(3.4) \quad \nu_w(\chi_{v[w]}) \neq \nu_w(\chi_{\sigma_{s_w, v[w]}^G}).$$

Note that for every $w' \in W$, the function $\alpha \mapsto \nu_w(\chi_{\sigma_{\ell(w'\alpha) - \ell(\alpha_\sigma), v[w]}^G})$ is of exponential type, hence belongs to $\mathcal{O}_{\exp}(\mathfrak{h}_\mathbb{C}^*)$. We put

$$\nu_{w, w'}^\dagger := \nu_w - \nu_w(\chi_{\sigma_{\ell \circ w'(-) - \ell(\alpha_\sigma), v[w]}^G}),$$

regarded as an element in $\mathcal{O}_{\exp}(\mathfrak{h}_\mathbb{C}^*) \otimes \mathcal{H}_G^T$. Then (3.4) simply says $\nu_{w, w}^\dagger(\alpha_\chi, \chi^{\infty, \tau}) \neq 0$. From this, it is elementary to see that there exist complex constants $\{C_w\}_{w \in W}$ such that

$$\sum_{w \in W} C_w \nu_{w, w'}^\dagger(\alpha_\chi, \chi^{\infty, \tau}) \neq 0$$

for every $w' \in W$. Put

$$\nu_{w'}^\dagger := \sum_{w \in W} C_w \nu_{w, w'}^\dagger.$$

Then put

$$\nu^\dagger := \prod_{w' \in W} \nu_{w'}^\dagger,$$

which is an element in $\mathcal{O}_{\exp}(\mathfrak{h}_\mathbb{C}^*)^W \otimes \mathcal{H}_G^T$, satisfying $\nu^\dagger(\chi) \neq 0$.

Now we claim that $\nu^\dagger(\chi_{\sigma_s^T}^G) = 0$ for every $s \in \mathfrak{a}_{M,\mathbb{C}}^*$. Note that the element $\alpha_\sigma + s \in \mathfrak{h}_\mathbb{C}^*$ belongs to $\chi_{\sigma_{s, \infty}^G}$. Since we have

$$\begin{aligned} \nu_1^\dagger(\alpha_\sigma + s, \chi_{\sigma_{s, \infty}^G}^T) &= \sum_{w \in W} C_w \nu_{w, 1}^\dagger(\alpha_\sigma + s, \chi_{\sigma_{s, \infty}^G}^T) \\ &= \sum_{w \in W} C_w \left(\nu_w(\chi_{\sigma_{s, v[w]}^G}^T) - \nu_w(\chi_{\sigma_{\ell(\alpha_\sigma + s) - \ell(\alpha_\sigma), v[w]}^G}^T) \right) = 0, \end{aligned}$$

the claim follows. It is easy to find an element $\mu^\dagger \in \mathcal{M}(\mathfrak{h}_\mathbb{C}^*)^W$ satisfying $\mu^\dagger(\chi_\infty) \neq 0$, and such that $\mu^\dagger \cdot \nu^\dagger$ belongs to $\mathcal{M}(\mathfrak{h}_\mathbb{C}^*)^W \otimes \mathcal{H}_G^T$. Then by Remark 2.9(3),

$$\mu_{(M,\sigma)} := \mu_\infty^0 \cdot \mu^\dagger \cdot \nu^\dagger$$

is a T -multiplier of $\mathcal{S}(G(\mathbb{A}_F))_K$, satisfying $\mu_{(M,\sigma)}(\chi) \neq 0$.

It remains to show that for every $f \in \mathcal{S}(G(\mathbb{A}_F))_K$, the endomorphism $R(\mu_{(M,\sigma)} \star f)$ annihilates $L^2_{(M,\sigma)}(G(F) \backslash G(\mathbb{A}_F)/K, \omega)$. However, this follows from the same argument as in the proof of [Proposition 3.15](#) since for every $s \in \mathfrak{a}_{M,\mathbb{C}}^*$, we have $\mu_{(M,\sigma)}(\chi_{\sigma_s^\tau}^G) = 0$. Thus, the proposition follows. $\square$

Remark 3.18. The idea of constructing the element $\nu^\dagger$ in the proof of [Proposition 3.17](#) is inspired by [LV07]. The similar construction also appeared in [YZ17] in the case where $G = \mathrm{PGL}_2$ and F is a function field.

The following theorem is the most general form on isolating cuspidal components, from which we will deduce [Theorems 3.6](#) and [3.7](#).

THEOREM 3.19. *Let χ be a $\mathbf{T}$ -character. There exists a $\mathbf{T}$ -multiplier μ of $\mathcal{S}(G(\mathbb{A}_F))_K$ such that*

(1) *for every $f \in \mathcal{S}(G(\mathbb{A}_F))_K$, the image of the endomorphism*

$$R(\mu \star f): L^2(G(F) \backslash G(\mathbb{A}_F)/K, \omega) \rightarrow L^2(G(F) \backslash G(\mathbb{A}_F)/K, \omega)$$

is contained in

$$\bigoplus_{(M,\sigma) \in \mathfrak{D}(G,\omega)_\chi^\heartsuit} L^2_{(M,\sigma)}(G(F) \backslash G(\mathbb{A}_F)/K, \omega)$$

under the decomposition (3.9), where $\mathfrak{D}(G,\omega)_\chi^\heartsuit$ is as introduced in [Definition 3.16](#);

(2) $\mu(\chi) = 1$.

In particular, [Theorem 1.3](#) follows by taking $\mathbf{T} = \mathbf{S}$ and $\chi = \chi_{\pi^\tau}$, and the strong multiplicity one property for $G = \mathrm{Res}_{F'/F} \mathrm{GL}_n$ [PS79].

Proof. By [Lemma 3.13](#), we know that $\mathfrak{D}(G, \omega; K, \mathfrak{T})_{\chi_\infty}^\heartsuit$ is a finite set. Thus, we may choose a finite subset $\mathfrak{D} \subseteq \mathfrak{D}(G, \omega)$ that maps surjectively to the (finite) set $\mathfrak{D}(G, \omega; K, \mathfrak{T})_{\chi_\infty}^\heartsuit \setminus \mathfrak{D}(G, \omega)_\chi^\heartsuit$. Now we put

$$\mu := \mu_\infty \cdot \prod_{(M,\sigma) \in \mathfrak{D}} \mu_{(M,\sigma)}$$

from [Propositions 3.15](#) and [3.17](#), which is a $\mathbf{T}$ -multiplier of $\mathcal{S}(G(\mathbb{A}_F))_K$. Then (1) is satisfied. For (2), we only need to replace μ by $\mu(\chi)^{-1}\mu$ as $\mu(\chi) \neq 0$. The theorem follows. $\square$

Proof of [Theorems 3.6](#) and [3.7](#). Note that the intersection

$$L^2_{\mathrm{cusp}}(G(F) \backslash G(\mathbb{A}_F)/K, \omega) \bigcap \widehat{\bigoplus_{(M,\sigma) \in \mathfrak{D}(G,\omega)_\chi^\heartsuit} L^2_{(M,\sigma)}(G(F) \backslash G(\mathbb{A}_F)/K, \omega)}$$

is exactly $L^2_{\mathrm{cusp}}(G(F) \backslash G(\mathbb{A}_F)/K, \omega)[\chi]$. Thus, [Theorem 3.7](#) immediately follows from [Theorem 3.19](#). Moreover, [Theorem 3.6](#) follows from [Theorem 3.19](#)

and the relation

$$\widehat{\bigoplus_{(M,\sigma) \in \mathfrak{D}(G,\omega)_\chi^\vee}} L^2_{(M,\sigma)}(G(F) \backslash G(\mathbb{A}_F)/K, \omega) \subseteq L^2_{\text{cusp}}(G(F) \backslash G(\mathbb{A}_F)/K, \omega)$$

since χ is not G -CAP. $\square$

Remark 3.20. In fact, we may use the Langlands decomposition of the L^2 -spectrum [Lan76] (as formulated in [Art05, Th. 7.2]) instead of the decomposition (3.1), and [Mül89, Th. 0.1] instead of [Don82, Th. 9.1] in the proof of Lemma 3.12, to obtain a version of Theorem 3.19 in terms of the Langlands components, which is slightly stronger than the current version of Theorem 3.19.

More precisely, let $\mathfrak{D}'(G, \omega)$ be the set of pairs (M, σ) in which σ is now a discrete automorphic representation of $M(\mathbb{A}_F)$ whose central character belongs to $\Omega_M(\omega)$, with a similarly defined equivalence relation $\approx$. We have the following Langlands decomposition,

$$L^2(G(F) \backslash G(\mathbb{A}_F), \omega) = \widehat{\bigoplus_{(M,\sigma) \in \mathfrak{D}'(G,\omega)^\vee} L^2_{(M,\sigma)}(G(F) \backslash G(\mathbb{A}_F), \omega)'},$$

which refines (3.1), in which $L^2_{(M,\sigma)}(G(F) \backslash G(\mathbb{A}_F), \omega)'$ denotes the Langlands component associated to (M, σ) . The conclusion of Theorem 3.19 can be strengthened to that there exists a T-multiplier μ of $\mathcal{S}(G(\mathbb{A}_F))_K$ such that for every $f \in \mathcal{S}(G(\mathbb{A}_F))_K$,

(1) $R(\mu \star f)$ maps $L^2(G(F) \backslash G(\mathbb{A}_F)/K, \omega)$ into

$$\widehat{\bigoplus_{(M,\sigma) \in \mathfrak{D}'(G,\omega)_\chi^\vee} L^2_{(M,\sigma)}(G(F) \backslash G(\mathbb{A}_F)/K, \omega)'}$$

in which $\mathfrak{D}'(G, \omega)_\chi^\vee$ is similarly defined as $\mathfrak{D}(G, \omega)_\chi^\vee$ in Definition 3.16;

(2) $\mu(\chi) = 1$.

Note that we have the inclusion

$$\begin{aligned} & \widehat{\bigoplus_{(M,\sigma) \in \mathfrak{D}'(G,\omega)_\chi^\vee} L^2_{(M,\sigma)}(G(F) \backslash G(\mathbb{A}_F)/K, \omega)'} \\ & \subseteq \widehat{\bigoplus_{(M,\sigma) \in \mathfrak{D}(G,\omega)_\chi^\vee} L^2_{(M,\sigma)}(G(F) \backslash G(\mathbb{A}_F)/K, \omega)}. \end{aligned}$$

However, it could be strict. For example, when $G = \text{PGL}_2$, M is the standard diagonal Levi subgroup, and σ is the trivial character, we have

$$\begin{aligned} L^2_{(M,\sigma)}(G(F) \backslash G(\mathbb{A}_F)/K, 1) &= L^2_{(M,\sigma)}(G(F) \backslash G(\mathbb{A}_F)/K, 1)' \\ &\oplus L^2_{(G,\bar{\sigma})}(G(F) \backslash G(\mathbb{A}_F)/K, 1)', \end{aligned}$$

where $\tilde{\sigma}$ is the trivial character of $G(\mathbb{A}_F)$. It is possible to find χ such that $(M, \sigma) \in \mathfrak{D}'(G, \omega)_\chi^\vee$, hence $(M, \sigma) \in \mathfrak{D}(G, \omega)_\chi^\vee$, but $(G, \tilde{\sigma}) \notin \mathfrak{D}'(G, \omega)_\chi^\vee$. However, one cannot find χ such that $(G, \tilde{\sigma}) \in \mathfrak{D}'(G, \omega)_\chi^\vee$ but $(M, \sigma) \notin \mathfrak{D}'(G, \omega)_\chi^\vee$.

4. Application to the Gan–Gross–Prasad conjecture

In this section, we discuss the application of the results in previous sections to the global Gan–Gross–Prasad conjecture and the Ichino–Ikeda conjecture for $U(n) \times U(n+1)$ in the stable case. In [Section 4.1](#), we recall the Jacquet–Rallis relative trace formulae and their extension by Zydor. In [Section 4.2](#), we introduce the notion of smooth transfer and study its relation with multipliers. In [Section 4.3](#), we deduce some results concerning weak base change using Jacquet–Rallis relative trace formulae, which are necessary for the proof of the Gan–Gross–Prasad conjecture. In [Section 4.4](#), we prove the Gan–Gross–Prasad conjecture in the stable case and other related results.

Let E/F be a quadratic extension of number fields and $\eta_{E/F}: \mathbb{A}_F^\times \rightarrow \mathbb{C}^\times$ the associated quadratic automorphic character. Let $n \geq 1$ be an integer.

4.1. *Jacquet–Rallis relative trace formulae, d’après Zydor.* In this subsection, we collect some results from Zydor’s extension of the Jacquet–Rallis relative trace formulae. We start with the general linear groups. Put

$$G'_n := \text{Res}_{E/F} \text{GL}_{n,E}, \quad G'_{n+1} := \text{Res}_{E/F} \text{GL}_{n+1,E}, \quad G' := G'_n \times G'_{n+1}.$$

We have two reductive subgroups H'_1 and H'_2 of G' as at the beginning of [\[BP20a, §3\]](#), in which H'_1 is the graph of the natural embedding $G'_n \rightarrow G'_{n+1}$ via the first n coordinates, and H'_2 is the subgroup $\text{GL}_{n,F} \times \text{GL}_{n+1,F}$. Let $G'_{\text{rs}} \subseteq G'$ be the Zariski open subset of regular semisimple elements. Recall that an element γ of G' is regular semisimple if $H'_1 \gamma H'_2$ is Zariski closed in G' and the natural map $H'_1 \times H'_2 \rightarrow H'_1 \gamma H'_2$ is an isomorphism. Put $B' := H'_1 \backslash G' / H'_2$, which is an affine variety over F , and let $B'_{\text{rs}} \subseteq B'$ be the image of G'_{rs} . Let Z' be the maximal F -split center of G' , which is also the center of H'_2 . We denote by

$$\eta: H'_2(\mathbb{A}_F) = \text{GL}_n(\mathbb{A}_F) \times \text{GL}_{n+1}(\mathbb{A}_F) \rightarrow \mathbb{C}^\times$$

the character $(\eta_{E/F}^{n-1} \circ \det) \boxtimes (\eta_{E/F}^n \circ \det)$, which is trivial on $Z'(\mathbb{A}_F)H'_2(F)$.

For every element $\Pi \in \mathfrak{C}(G', 1)$ ([Definition 3.9](#)), there is a distribution I_Π on $\mathcal{S}(G'(\mathbb{A}_F))$ [[BP20a, §3.1](#)] such that for $f' \in \mathcal{S}(G'(\mathbb{A}_F))$,

$$(4.1) \quad I_\Pi(f') = \sum_{\phi \in \mathcal{B}(\Pi)} \left(\int_{H'_1(F) \backslash H'_1(\mathbb{A}_F)} (\Pi(f')\phi)(h_1) dh_1 \right) \cdot \overline{\left(\int_{Z'(\mathbb{A}_F)H'_2(F) \backslash H'_2(\mathbb{A}_F)} \phi(h_2)\eta(h_2) dh_2 \right)}$$

for an arbitrary orthonormal basis $\mathcal{B}(\Pi)$ of Π with respect to the Peterson inner product over G'/Z' . Here, dh_1 and dh_2 are the Tamagawa measures on $H'_1(\mathbb{A}_F)$ and $Z'(\mathbb{A}_F)\backslash H'_2(\mathbb{A}_F)$, respectively.

For every element $\gamma \in B'(F)$, there is a distribution I_γ on $\mathcal{S}(G'(\mathbb{A}_F))$ [Zyd20, Th. 5.9].⁸ When $\gamma \in B'_{\text{rs}}(F)$, I_γ is defined by relative orbital integrals; more precisely, for $f' \in \mathcal{S}(G'(\mathbb{A}_F))$ we have

$$(4.2) \quad I_\gamma(f') = O_\eta(\tilde{\gamma}, f') := \int_{H'_1(\mathbb{A}_F) \times H'_2(\mathbb{A}_F)} f'(h_1^{-1} \tilde{\gamma} h_2) \eta(h_2) dh_1 dh_2,$$

where $\tilde{\gamma} \in G'_{\text{rs}}(F)$ is an arbitrary lift of γ . In general, the distribution I_γ is a certain regularization of relative orbital integrals, whose precise definition is complicated and will not be used in our later discussion, hence we will not discuss it here. The following proposition is Zydor's extension of the Jacquet–Rallis trace formula for G' .

PROPOSITION 4.1. *Let $f' \in \mathcal{S}(G'(\mathbb{A}_F))$ be a quasi-cuspidal Schwartz test function (Definition 3.2). Then we have the following identity of absolutely convergent sums:*

$$\sum_{\Pi \in \mathfrak{C}(G', 1)} I_\Pi(f') = \sum_{\gamma \in B'(F)} I_\gamma(f').$$

Proof. This follows from the same proof of [Zyd20, Th. 5.10]. $\square$

Now we consider unitary groups. We denote by $\mathfrak{V}$ the set of isomorphism classes of pairs $V = (V_n, V_{n+1})$ of (non-degenerate) hermitian spaces over E , where V_n has rank n and $V_{n+1} = V_n \oplus E.e$ in which e has norm 1. For every place v of F , we have a local analogue $\mathfrak{V}_v$ and a localization map $\mathfrak{V} \rightarrow \mathfrak{V}_v$ sending V to V_v . For every $V \in \mathfrak{V}$, put

$$G_n^V := \text{U}(V_n), \quad G_{n+1}^V := \text{U}(V_{n+1}), \quad G^V := G_n^V \times G_{n+1}^V.$$

We have a reductive subgroup H^V of G^V as at the beginning of [BP20a, §3], which is the graph of the natural embedding $G_n^V \rightarrow G_{n+1}^V$. Let $G_{\text{rs}}^V \subseteq G^V$ be the Zariski open subset of regular semisimple elements. Recall that an element δ^V of G^V is regular semisimple if $H^V \delta^V H^V$ is Zariski closed in G^V and the natural map $H^V \times H^V \rightarrow H^V \delta^V H^V$ is an isomorphism. Put $B^V := H^V \backslash G^V / H^V$, which is an affine variety over F , and let $B_{\text{rs}}^V \subseteq B^V$ be the image of G_{rs}^V .

⁸Although [Zyd20, Th. 5.9] only states for test functions in $C_c^\infty(G'(\mathbb{A}_F))$, its proof works for $\mathcal{S}(G'(\mathbb{A}_F))$ as well. We have a similar situation below concerning [Zyd20, Th. 6.6].

For every element $\pi^V \in \mathfrak{C}(G^V, 1)$ (Definition 3.9), there is a distribution J_{π^V} on $\mathcal{S}(G^V(\mathbb{A}_F))$ [BP20a, §3.1] such that for $f^V \in \mathcal{S}(G^V(\mathbb{A}_F))$,

$$(4.3) \quad J_{\pi^V}(f^V) = \sum_{\varphi \in \mathcal{B}(\pi^V)} \left(\int_{H^V(F) \backslash H^V(\mathbb{A}_F)} (\pi^V(f^V)\varphi)(h) dh \right) \cdot \overline{\left(\int_{H^V(F) \backslash H^V(\mathbb{A}_F)} \varphi(h) dh \right)}$$

for an arbitrary orthonormal basis $\mathcal{B}(\pi^V)$ of π^V with respect to the Peterson inner product over G^V . Here, dh is the Tamagawa measure on $H^V(\mathbb{A}_F)$.

For every element $\delta^V \in B^V(F)$, there is a distribution J_{δ^V} on $\mathcal{S}(G^V(\mathbb{A}_F))$ [Zyd20, Th. 6.6]. When $\delta^V \in B_{\text{rs}}^V(F)$, J_{δ^V} is defined by relative orbital integrals; more precisely, we have for $f^V \in \mathcal{S}(G^V(\mathbb{A}_F))$,

$$(4.4) \quad J_{\delta^V}(f^V) = O(\tilde{\delta}^V, f^V) := \int_{H^V(\mathbb{A}_F) \times H^V(\mathbb{A}_F)} f^V(h_1^{-1} \tilde{\delta}^V h_2) dh_1 dh_2,$$

where $\tilde{\delta}^V \in G_{\text{rs}}^V(F)$ is an arbitrary lift of δ^V .⁹ In general, the distribution I_γ is a certain regularization of relative orbital integrals, whose precise definition is complicated and will not be used in our later discussion; hence we will not discuss it here. The following proposition is Zydor's extension of the Jacquet–Rallis trace formula for G^V .

PROPOSITION 4.2. *Let $f^V \in \mathcal{S}(G^V(\mathbb{A}_F))$ be a quasi-cuspidal Schwartz test function (Definition 3.2). Then we have the following identity of absolutely convergent sums:*

$$\sum_{\pi^V \in \mathfrak{C}(G^V, 1)} J_{\pi^V}(f^V) = \sum_{\delta^V \in B^V(F)} J_{\delta^V}(f^V).$$

Proof. This follows from the same proof of [Zyd20, Th. 6.7]. $\square$

The affine varieties B' and B^V are canonically isomorphic (see [Zha14b, §3.1]). For an F -algebra R , we say that $\gamma \in G'_{\text{rs}}(R)$ and $\delta^V \in G_{\text{rs}}^V(R)$ are a *matching pair* if their images in $B'(R) = B^V(R)$ coincide.

4.2. Smooth transfer of Schwartz test functions. We now discuss the smooth transfer of Schwartz test functions. We keep the notation from the previous subsection.

Definition 4.3. Let v be a non-archimedean place of F that is unramified in E . We say that

⁹In this case, the map $G_{\text{rs}}^V(F) \rightarrow B_{\text{rs}}^V(F)$ is not surjective. If δ^V does not lift, then we simply set $J_{\delta^V}(f^V) = 0$.

- K'_v is a *relative hyperspecial maximal subgroup* of $G'(F_v)$ if it is $\mathrm{GL}_n(O_{E_v}) \times \mathrm{GL}_{n+1}(O_{E_v})$;
- K_v^V is a *relative hyperspecial maximal subgroup* of $G^V(F_v)$ if it is of the form $\mathrm{U}(L_n) \times \mathrm{U}(L_{n+1})$ in which L_n is a self-dual lattice of $V_{n,v}$ and $L_{n+1} = L_n \oplus O_{E_v} \cdot e$.

For $G \in \{G', G^V\}$ and every finite set $\square$ of places of F containing all archimedean ones, we define the algebra

$$\mathcal{S}(G(F_\square)) := \mathcal{S}(G_\infty) \otimes \bigotimes_{v \in \square, v \nmid \infty} \mathcal{S}(G(F_v)).$$

The following definition slightly extends the notion of smooth transfer for Schwartz test functions that are not necessarily pure tensors.

Definition 4.4 (Smooth transfer). Let $f' \in \mathcal{S}(G'(\mathbb{A}_F))$ be a Schwartz test function that is not necessarily a pure tensor.

- (1) Let $\square$ be a finite set of places of F containing all archimedean ones, and take $V \in \mathfrak{V}$. We say that $f'_\square \in \mathcal{S}(G'(F_\square))$ and $f_\square^V \in \mathcal{S}(G^V(F_\square))$ have *matching orbital integrals* if for every matching pair $\gamma \in G'_{\mathrm{rs}}(F_\square)$ and $\delta^V \in G^V_{\mathrm{rs}}(F_\square)$, we have

$$O(\delta^V, f_\square^V) = \Omega_\square(\gamma) \cdot O_\eta(\gamma, f'_\square).$$

Here, $O(\delta^V, f_\square^V)$ and $O_\eta(\gamma, f'_\square)$ are relative orbital integrals defined in the same way as (4.4) and (4.2) after replacing $\mathbb{A}_F$ by $F_\square$, respectively; and $\Omega_\square(\gamma) := \prod_{v \in \square} \Omega_v(\gamma_v) \in \{\pm 1\}$ is a certain transfer factor (see [BP20a, §§3.3, 3.4] for more details).

- (2) For $V \in \mathfrak{V}$ and a Schwartz test function $f^V \in \mathcal{S}(G^V(\mathbb{A}_F))$, we say that f' and f^V are *smooth transfer* if there exists a sufficiently large finite set $\square$ of places of F containing all archimedean ones and those ramified in E such that
 - (a) $f' = f'_\square \otimes \bigotimes_{v \notin \square} \mathbf{1}_{K'_v}$ with $f'_\square \in \mathcal{S}(G'(F_\square))$ and K'_v a relative hyperspecial maximal subgroup of $G'(F_v)$;
 - (b) $f^V = f_\square^V \otimes \bigotimes_{v \notin \square} \mathbf{1}_{K_v^V}$ with $f_\square^V \in \mathcal{S}(G^V(F_\square))$ and K_v^V a relative hyperspecial maximal subgroup of $G^V(F_v)$;
 - (c) $f'_\square \in \mathcal{S}(G'(F_\square))$ and $f_\square^V \in \mathcal{S}(G^V(F_\square))$ have matching orbital integrals in the sense of (1).
- (3) Given a collection $(f^V)_{V \in \mathfrak{V}}$ with $f^V \in \mathcal{S}(G^V(\mathbb{A}_F))$ among which all but finitely many are zero, we say that f' and $(f^V)_{V \in \mathfrak{V}}$ are *complete smooth transfer* if f' and f^V are smooth transfer for every $V \in \mathfrak{V}$.

Remark 4.5. By the relative fundamental lemma [Yun11, BP21a], Definition 4.4(2) is independent of the choice of $\square$.

The existence of smooth transfer is ensured by the following proposition.

PROPOSITION 4.6. *Let v be a place of F .*

- (1) *There is a dense subspace $\mathcal{S}_{\text{tr}}(G'(F_v))$ of $\mathcal{S}(G'(F_v))$, equal to $\mathcal{S}(G'(F_v))$ if v is non-archimedean or split in E , such that for every $f'_v \in \mathcal{S}_{\text{tr}}(G'(F_v))$, one can find $(f_v^{V_v})_{V_v \in \mathfrak{V}_v}$ with $f_v^{V_v} \in \mathcal{S}(G^{V_v}(F_v))$ such that f'_v and $f_v^{V_v}$ have matching orbital integrals for every $V_v \in \mathfrak{V}_v$.*
- (2) *For every $V_v \in \mathfrak{V}_v$, there is a dense subspace $\mathcal{S}_{\text{tr}}(G^{V_v}(F_v))$ of $\mathcal{S}(G^{V_v}(F_v))$, equal to $\mathcal{S}(G^{V_v}(F_v))$ if v is non-archimedean or split in E , such that for every $f_v^{V_v} \in \mathcal{S}_{\text{tr}}(G^{V_v}(F_v))$, one can find $f'_v \in \mathcal{S}(G'(F_v))$ such that f'_v and $f_v^{V_v}$ have matching orbital integrals for every $W_v \in \mathfrak{V}_v$ where $f_v^{W_v} = 0$ for $W_v \neq V_v$.*

Proof. This follows from the combination of [Zha14b, Th. 2.6] (for v non-archimedean) and [Xue19, Th. 2.7] (for v archimedean). $\square$

The following proposition is a deep theorem of Chaudouard and Zydor.

PROPOSITION 4.7. *Suppose that f' and $(f^V)_{V \in \mathfrak{V}}$ are complete smooth transfer in the sense of Definition 4.4(3). Then we have*

$$\sum_{\gamma \in B'(F)} I_\gamma(f') = \sum_{V \in \mathfrak{V}} \sum_{\delta^V \in B^V(F)} J_{\delta^V}(f^V),$$

in which the summation over V is in fact finite.

Proof. This is [CZ21, Th. 16.2.4.1]. Note that the authors in [CZ21] only prove this for pure tensor Schwartz test functions, but their proof works for general Schwartz test functions as well, using the fact that $\bigotimes_{v|\infty} \mathcal{S}(G'(F_v))$ and $\bigotimes_{v|\infty} \mathcal{S}(G^V(F_v))$ are dense in $\mathcal{S}(G'_\infty)$ and $\mathcal{S}(G_\infty^V)$, respectively, by [AG10, Cor. 2.6.3]. $\square$

Now we study smooth transfer of Schwartz test functions under the action of $\mathbf{T}$ -multipliers. Let $\mathbf{T}_0$ be the set of primes of F that are *non-split* in E . We take

- a finite set $\mathbf{S}$ of primes of F containing those ramified in E ;
- an open compact subgroup $K' \subseteq G'(\mathbb{A}_F^\infty)$ of the form $K' = K'_\mathbf{S} \times \prod_{v \notin \mathbf{S}} K'_v$ in which K'_v is a relative hyperspecial maximal subgroup of $G'(F_v)$ for every prime $v \notin \mathbf{S}$;
- an element $V \in \mathfrak{V}$; and
- an open compact subgroup $K^V \subseteq G^V(\mathbb{A}_F^\infty)$ of the form

$$K^V = K_\mathbf{S}^V \times \prod_{v \notin \mathbf{S}} K_v^V$$

in which K_v^V is a relative hyperspecial maximal subgroup of $G^V(F_v)$ for every prime $v \notin \mathbf{S}$.

Put $\mathbf{T} := \mathbf{S} \cup \mathbf{T}_0$.

The connected real reductive group $G' \otimes_{\mathbb{Q}} \mathbb{R}$ determines a root datum $(X'^*, \Phi', X'_*, \Phi'^{\vee})$ and a subset $\theta' \subseteq \text{Aut}(X'^*, \Phi', X'_*, \Phi'^{\vee})^{\vee}$ as at the beginning of [Section 2.3](#), together with $\mathfrak{h}'^*$ and W' from [Section 2.2](#). Similarly, for $G^V \otimes_{\mathbb{Q}} \mathbb{R}$, we have corresponding objects $(X^{V*}, \Phi^V, X_*^V, \Phi^{V\vee})$, θ^V , $\mathfrak{h}^{V*}$, and W^V .¹⁰ By the base change homomorphism on the dual groups, we have a canonical map $\mathfrak{h}^{V*} \rightarrow \mathfrak{h}'^*$ that is injective, by which we will identify $\mathfrak{h}^{V*}$ as a subspace of $\mathfrak{h}'^*$. Moreover, taking restriction induces a ring homomorphism

$$(4.5) \quad \text{bc}_{\infty}^V: \mathcal{O}(\mathfrak{h}'^*)^{W'} \rightarrow \mathcal{O}(\mathfrak{h}_C^{V*})^{W^V}.$$

For primes away from $\mathbf{T}$, we have a similar homomorphism

$$(4.6) \quad \text{bc}_V^{\mathbf{T}}: \mathcal{H}_{G'}^{\mathbf{T}} \rightarrow \mathcal{H}_{G^V}^{\mathbf{T}}$$

given by unramified base change (which is simply a convolution product since primes away from $\mathbf{T}$ are all split in E). Taking tensor product, we obtain a homomorphism

$$(4.7) \quad \text{bc}^V := \text{bc}_{\infty}^V \otimes \text{bc}_V^{\mathbf{T}}: \mathcal{O}(\mathfrak{h}'^*)^{W'} \otimes \mathcal{H}_{G'}^{\mathbf{T}} \rightarrow \mathcal{O}(\mathfrak{h}_C^{V*})^{W^V} \otimes \mathcal{H}_{G^V}^{\mathbf{T}}.$$

PROPOSITION 4.8. *Let $f' = \bigotimes_v f'_v \in \mathcal{S}(G'(\mathbb{A}_F))_{K'}$ and $f^V = \bigotimes_v f_v^V \in \mathcal{S}(G^V(\mathbb{A}_F))_{K^V}$ be two pure tensor Schwartz test functions such that f'_v and f_v^V have matching orbital integrals for every place v of F . Let μ' and μ^V be $\mathbf{T}$ -multipliers of $\mathcal{S}(G'(\mathbb{A}_F))_{K'}$ and $\mathcal{S}(G^V(\mathbb{A}_F))_{K^V}$ ([Definition 3.5](#)), respectively, such that $\mu^V = \text{bc}^V(\mu')$. Then $\mu' \star f'$ and $\mu^V \star f^V$ are smooth transfer in the sense of [Definition 4.4\(2\)](#).*

Informally speaking, the proposition asserts that multipliers that are compatible under base change preserve smooth transfer.

Proof. Let $\square$ be a sufficiently large finite set of places of F containing all archimedean ones and disjoint from $\mathbf{T}$, so that μ' and μ^V are of the form $\mu'_{\square} \otimes (\bigotimes_{v \notin \mathbf{T} \cup \square} \mathbf{1}_{K'_v})$ and $\mu^V_{\square} \otimes (\bigotimes_{v \notin \mathbf{T} \cup \square} \mathbf{1}_{K^V_v})$ respectively. Below we only care about $\square$ -components, hence will write μ' for $\mu'_{\square}$ and μ^V for $\mu^V_{\square}$, respectively.

Let $\mathcal{S}(G'(F_{\square}))_{K'}$ be the subalgebra of $\mathcal{S}(G'(F_{\square}))$ consisting of functions that are bi-invariant under $\prod_{v \in \square, v \nmid \infty} K'_v$, and similarly for $\mathcal{S}(G^V(F_{\square}))_{K^V}$. To prove the proposition, it is enough to show that if $f' \in \mathcal{S}(G'(F_{\square}))_{K'}$ and $f^V \in \mathcal{S}(G^V(F_{\square}))_{K^V}$ have matching orbital integrals (in the sense of [Definition 4.4\(1\)](#)), then $\mu' \star f'$ and $\mu^V \star f^V$ have matching orbital integrals as well. In fact, by [Lemma 4.9](#) below, f' and f^V have matching relative characters. Since $\mu^V = \text{bc}^V(\mu')$, by [Theorem 2.13](#), we know that $\mu' \star f'$ and $\mu^V \star f^V$ also have

¹⁰In fact, the objects $(X^{V*}, \Phi^V, X_*^V, \Phi^{V\vee})$, θ^V , $\mathfrak{h}^{V*}$, and W^V do not depend on V . However, we still keep V in superscripts in order to make notation more consistent.

matching relative characters. Thus again by [Lemma 4.9](#) below, we know that $\mu' \star f'$ and $\mu^V \star f^V$ have matching orbital integrals. The proposition follows. $\square$

LEMMA 4.9. *Let $\square$ be a finite set of places of F containing all archimedean ones, in which all primes are split in E . Take $f' \in \mathcal{S}(G'(F_\square))$ and $f^V \in \mathcal{S}(G^V(F_\square))$. Then f' and f^V have matching orbital integrals (in the sense of [Definition 4.4\(1\)](#)) if and only if they have matching relative characters, that is,*

$$\kappa_{V_\square} J_\pi(f^V) = I_{\text{BC}(\pi)}(f')$$

for every $\pi \in \text{Temp}_{H_\square^V}(G_\square^V)$, where we follow the notation in [\[BP21b, Th. 5.4.1\]](#).

The term $\kappa_{V_\square} := \prod_{v \in \square} \kappa_{V_v}$ is a constant depending only on $V_\square$, and the distributions $J_\pi(f^V)$ and $I_{\text{BC}(\pi)}(f')$ are local analogues of [\(4.3\)](#) and [\(4.1\)](#), which depend linearly on $\pi(f^V)$ and $\text{BC}(\pi)(f')$, respectively.

Proof. As $\bigotimes_{v|\infty} \mathcal{S}(G'(F_v))$ is dense in $\mathcal{S}(G'_\infty)$ by [\[AG10, Cor. 2.6.3\]](#), applying [\[BP21b, \(5.5.10\)\]](#) to local fields F_v for $v \in \square$ and by continuity, we obtain a similar identity

$$(4.8) \quad \int_{B'_{\text{rs}}(F_\square)} O_\eta(\gamma, f'_1) \overline{O_\eta(\gamma, f'_2)} d\gamma \\ = |\tau|_{E_\square}^{-n(n-1)/2} \int_{\text{Temp}(G_\square^{\text{qs}})/\text{stab}} I_{\text{BC}(\pi)}(f'_1) \overline{I_{\text{BC}(\pi)}(f'_2)} \frac{|\gamma^*(0, \pi, \text{Ad}, \psi')|}{|S_\pi|} d\pi$$

for every $f'_1, f'_2 \in \mathcal{S}(G'(F_\square))$. Similarly, as $\bigotimes_{v|\infty} \mathcal{S}(G^V(F_v))$ is dense in $\mathcal{S}(G^V_\infty)$ by [\[AG10, Cor. 2.6.3\]](#), applying [\[BP21b, \(5.5.3\)\]](#) to local fields F_v for $v \in \square$ and by continuity, we obtain a similar identity

$$(4.9) \quad \int_{B'_{\text{rs}}(F_\square)} O(\delta^V, f_1^V) \overline{O(\delta^V, f_2^V)} d\delta^V = \int_{\text{Temp}_{H_\square^V}(G_\square^V)} J_\pi(f_1^V) \overline{J_\pi(f_2^V)} \mu_{G_\square^V}^*(\pi) d\pi$$

for every $f_1^V, f_2^V \in \mathcal{S}(G^V(F_\square))$.¹¹

We have the following observations:

- (1) On the geometric side, we may identify $B'_{\text{rs}}(F_\square)$ as a subspace, say $B'_{\text{rs}}(F_\square)_V$, of $B'_{\text{rs}}(F_\square)$ under which the measures are compatible.
- (2) On the spectral side, by the local Gan–Gross–Prasad at places in $\square$ [\[BP20b\]](#), the local Jacquet–Langlands transfer map

$$\text{Temp}_{H_\square^V}(G_\square^V) \rightarrow \text{Temp}(G_\square^{\text{qs}})/\text{stab}$$

¹¹We remark that in [\(4.8\)](#), the product $O_\eta(\gamma, f'_1) \overline{O_\eta(\gamma, f'_2)}$ descends to a function on $B'_{\text{rs}}(F_\square)$, while in [\(4.9\)](#), both $O(\delta^V, f_1^V)$ and $\overline{O(\delta^V, f_2^V)}$ descend to functions on $B'_{\text{rs}}(F_\square)$.

is injective, under which the restriction of the measure $\frac{|\gamma^*(0, \pi, \text{Ad}, \psi')|}{|\mathcal{S}_\pi|} d\pi$ coincides with $\mu_{G_\square^V}^*(\pi) d\pi$ by [BP21b, Th. 5.4.3]. We also note that $\kappa_{V_\square} \overline{\kappa_{V_\square}} = |\tau|_{E_\square}^{n(n-1)/2}$.

Now we show the lemma.

First, suppose that f' and f^V have matching orbital integrals. Take $g^V \in \bigotimes_{v \in \square} \mathcal{S}_{\text{tr}}(G^V(F_v))$, and let $g' \in \bigotimes_{v \in \square} \mathcal{S}(G'(F_v))$ be an element that has matching orbital integrals with $(g^V, 0, \dots)$ from Proposition 4.6(2). By [BP21b, Th. 5.4.1], g' and $(g^V, 0, \dots)$ have matching relative characters as well. In particular, $O_\eta(\gamma, g') = 0$ unless $\gamma \in B'_{\text{rs}}(F_\square)_V$, and $I_{\text{BC}(\pi)}(g') = 0$ unless $\pi \in \text{Temp}_{H_\square^V}(G_\square^V)$. Then combining (4.8) and (4.9) and using the condition that f' and f^V have matching orbital integrals, we obtain

$$\begin{aligned} & \int_{\text{Temp}_{H_\square^V}(G_\square^V)} \kappa_{V_\square} J_\pi(f^V) \overline{J_\pi(g^V)} \mu_{G_\square^V}^*(\pi) d\pi \\ &= \int_{\text{Temp}_{H_\square^V}(G_\square^V)} I_{\text{BC}(\pi)}(f') \overline{J_\pi(g^V)} \mu_{G_\square^V}^*(\pi) d\pi \end{aligned}$$

for every $g^V \in \bigotimes_{v \in \square} \mathcal{S}_{\text{tr}}(G^V(F_v))$. Now we can use the technique in [BP21b, §5.7.3] to separate π , so that we obtain $\kappa_{V_\square} J_\pi(f^V) = I_{\text{BC}(\pi)}(f')$ for every $\pi \in \text{Temp}_{H_\square^V}(G_\square^V)$. In other words, f' and f^V have matching relative characters.

Second, suppose that f' and f^V have matching relative characters. The proof is similar, and we arrive at the identity

$$\begin{aligned} & \int_{B_{\text{rs}}^V(F_\square)} O(\delta^V, f^V) \overline{O(\delta^V, g^V)} d\delta^V \\ &= \int_{B_{\text{rs}}^V(F_\square)} \Omega_\square(\gamma(\delta^V)) O(\gamma(\delta^V), f') \overline{O(\delta^V, g^V)} d\delta^V \end{aligned}$$

for every $g^V \in \bigotimes_{v \in \square} \mathcal{S}_{\text{tr}}(G^V(F_v))$. Here, $\gamma(\delta^V) \in G'_{\text{rs}}(F_\square)$ is an arbitrary element whose image in $B'(F_\square)$ coincides with δ^V . Moreover, it is clear that the term $\Omega_\square(\gamma(\delta^V)) O(\gamma(\delta^V), f')$ is independent of the choice of such $\gamma(\delta^V)$. It is easy to see that locally at every given $\delta^V \in B_{\text{rs}}^V(F_\square)$, the functions $O(-, g^V)$ for $g^V \in \bigotimes_{v \in \square} \mathcal{S}_{\text{tr}}(G^V(F_v))$ span a dense subset in the L^2 -space of $B_{\text{rs}}^V(F_\square)$. Thus, we have $O(\delta^V, f^V) = \Omega_\square(\gamma(\delta^V)) O(\gamma(\delta^V), f')$ for every $\delta^V \in B_{\text{rs}}^V(F_\square)$, as both sides are continuous functions in δ^V . In other words, f' and f^V have matching orbital integrals.

The lemma is proved. $\square$

Remark 4.10. Lemma 4.9 holds by the same proof without assuming that primes in $\square$ are split in E . However, in this case, the argument (more precisely, [BP21b, Th. 5.4.1]) relies on results from [Mok15], [KMSW14].

The following proposition will not be used in this article, but might be useful for other purposes. In particular, it shows the existence of Gaussian test functions (in the space of Schwartz functions) in the sense of [RSZ20, Def. 7.9]. We record it here as it is essentially a corollary of Proposition 4.8.

PROPOSITION 4.11. *In the situation of Proposition 4.6(2), when v is an archimedean place of F that is non-split in E and V_v is positive definite, the subspace $\mathcal{S}_{\text{tr}}(G^{V_v}(F_v))$ contains $f_v^{V_v}$ as long as $J_{\pi_v}(f_v^{V_v}) = 0$ for all but finitely many (tempered) irreducible admissible representations π_v of $G^{V_v}(F_v)$.*

Proof. Since the question is local, we may just assume that E is imaginary quadratic, hence $v = \infty$ is the unique archimedean place of $F = \mathbb{Q}$. To ease notation, we suppress v in the proof. By linearity, it suffices to consider f^V such that $J_{\pi}(f^V) \neq 0$ for exactly one irreducible admissible representation π_0 of $G^V(F)$ and that $J_{\pi_0}(f^V) = 1$.

By Proposition 4.6(2), we may find an element $f_1^V \in \mathcal{S}_{\text{tr}}(G^V(F))$ satisfying $J_{\pi_0}(f_1^V) = 1$, so that we may take an element $f'_1 \in \mathcal{S}(G'(F))$ such that f'_1 and f_1^W have matching orbital integrals for every $W \in \mathfrak{V}$ where $f_1^W = 0$ for $W \neq V$. Now since $G^V(F)$ is compact, the set of infinitesimal characters of irreducible representations of $G^V(F)$ is a lattice. It follows easily from Corollary 2.6 that we can find an element $\mu' \in \mathcal{M}_{\theta}^{\sharp}(\mathfrak{h}_{\mathbb{C}}^*)^W$ such that if we put $\mu^V := \text{bc}^V(\mu')$, then $\mu^V(\chi_{\pi}) \neq 0$ only when $\pi = \pi_0$ and $\mu^V(\chi_{\pi_0}) = 1$. By Theorem 2.13, we have new elements $f_2^V := \mu^V \star f_1^V \in \mathcal{S}(G^V(F))$ and $f'_2 := \mu' \star f'_1 \in \mathcal{S}(G'(F))$. By Lemma 4.9 (for $\square = \{\infty\}$), f'_1 and f_1^W have matching relative characters, so for f'_2 and f_2^W where $f_2^W = 0$ for $W \neq V$, hence f'_2 and f_2^W have matching orbital integrals. Now since $J_{\pi}(f^V) = J_{\pi}(f_2^V)$ for every irreducible admissible representation π of $G^V(F)$, using Lemma 4.9 twice, we know that f'_2 and f^W have matching orbital integrals for every $W \in \mathfrak{V}$ where $f^W = 0$ for $W \neq V$. In other words, f^V belongs to $\mathcal{S}_{\text{tr}}(G^V(F))$.

The proposition is proved. $\square$

4.3. *Weak automorphic base change.* We keep the setup from the previous subsection.

We denote by $\mathfrak{V}_{(\mathbf{S})}$ the subset of $\mathfrak{V}$ consisting of V such that G_v^V is unramified for every prime v of F not in $\mathbf{S}$. Then $\mathfrak{V}_{(\mathbf{S})}$ is a finite set, and for $V, W \in \mathfrak{V}_{(\mathbf{S})}$, we have $V_v = W_v \in \mathfrak{V}_v$ for every prime v of F not in $\mathbf{S}$. We consider

- a non-empty subset $\mathfrak{V}_{(\mathbf{S})}^{\circ}$ of $\mathfrak{V}_{(\mathbf{S})}$;
- a T-character χ' for G' (Definition 3.3); and
- for each $V \in \mathfrak{V}_{(\mathbf{S})}^{\circ}$, a T-character χ^V for G^V (Definition 3.3) such that the base change of χ^V coincides with χ' , and also we fix an open compact subgroup $K^V \subseteq G^V(\mathbb{A}_F^{\infty})$ as in the previous subsection.

The following lemma tells us how to find multipliers that are compatible under base change.

LEMMA 4.12. *Given a $\mathbf{T}$ -multiplier μ' of $\mathcal{S}(G'(\mathbb{A}_F))_{K'}$ satisfying $\mu'(\chi') = 1$ and a $\mathbf{T}$ -multiplier μ^V of $\mathcal{S}(G^V(\mathbb{A}_F))_{K^V}$ satisfying $\mu^V(\chi^V) = 1$ for each $V \in \mathfrak{V}_{(\mathbf{s})}^\circ$, we can find new $\mathbf{T}$ -multipliers $\tilde{\mu}'$ of $\mathcal{S}(G'(\mathbb{A}_F))_{K'}$ and $\tilde{\mu}^V$ of $\mathcal{S}(G^V(\mathbb{A}_F))_{K^V}$ for each $V \in \mathfrak{V}_{(\mathbf{s})}^\circ$ satisfying the following:*

- (1) $\tilde{\mu}'(\chi') = 1$, and $\tilde{\mu}'$ is a multiple of μ' by a $\mathbf{T}$ -multiplier of $\mathcal{S}(G'(\mathbb{A}_F))_{K'}$;
- (2) for every $V \in \mathfrak{V}_{(\mathbf{s})}^\circ$, $\tilde{\mu}^V(\chi^V) = 1$, $\tilde{\mu}^V$ is a multiple of μ^V by a $\mathbf{T}$ -multiplier of $\mathcal{S}(G^V(\mathbb{A}_F))_{K^V}$, and $\tilde{\mu}^V = \mathbf{bc}^V(\tilde{\mu}')$.

Proof. For every $V \in \mathfrak{V}_{(\mathbf{s})}^\circ$, recall the maps $\mathbf{bc}_\infty^V$ (4.5), $\mathbf{bc}_V^{\mathbf{T}}$ (4.6), and $\mathbf{bc}^V$ (4.7) from the previous subsection, which are all surjective. We choose a section

$$\mathbf{cb}_V^{\mathbf{T}}: \mathcal{H}_{G^V}^{\mathbf{T}} \rightarrow \mathcal{H}_{G'}^{\mathbf{T}}$$

of the linear map $\mathbf{bc}_V^{\mathbf{T}}$ of vector spaces. We may choose a linear splitting map $\ell_V: \mathfrak{h}^{V*} \rightarrow \mathfrak{h}^{V*}$ of the subspace $\mathfrak{h}^{V*} \subseteq \mathfrak{h}^*$ that sends every W' -orbit into a W^V -orbit, which induces a section

$$\mathbf{cb}_\infty^V: \mathcal{O}(\mathfrak{h}_{\mathbb{C}}^{V*})^{W^V} \rightarrow \mathcal{O}(\mathfrak{h}_{\mathbb{C}}^{*})^{W'}$$

of the linear map $\mathbf{bc}_\infty^V$ of vector spaces. Taking tensor product, we obtain a linear map

$$\mathbf{cb}^V := \mathbf{cb}_\infty^V \otimes \mathbf{cb}_V^{\mathbf{T}}: \mathcal{O}(\mathfrak{h}_{\mathbb{C}}^{V*})^{W^V} \otimes \mathcal{H}_{G^V}^{\mathbf{T}} \rightarrow \mathcal{O}(\mathfrak{h}_{\mathbb{C}}^{*})^{W'} \otimes \mathcal{H}_{G'}^{\mathbf{T}},$$

which is a section of $\mathbf{bc}^V$. To construct the desired $\mathbf{T}$ -multipliers, we define

$$\tilde{\mu}' := (\mu')^2 \cdot \prod_{V \in \mathfrak{V}_{(\mathbf{s})}^\circ} \mathbf{cb}^V((\mu^V)^2), \quad \tilde{\mu}^V := \mathbf{bc}^V(\tilde{\mu}').$$

Note that $\mathbf{bc}_\infty^V$ sends $\mathcal{M}(\mathfrak{h}_{\mathbb{C}}^{*})^{W'}$ into $\mathcal{M}(\mathfrak{h}_{\mathbb{C}}^{V*})^{W^V}$, and $\mathbf{cb}_V^{\mathbf{T}}$ sends $\mathcal{M}(\mathfrak{h}_{\mathbb{C}}^{V*})^{W^V}$ into $\mathcal{M}(\mathfrak{h}_{\mathbb{C}}^{*})^{W'}$. Then it follows easily from Remark 2.9(3) that $\tilde{\mu}'$ and $\tilde{\mu}^V$ for $V \in \mathfrak{V}_{(\mathbf{s})}^\circ$ are $\mathbf{T}$ -multipliers satisfying (1) and (2). The lemma is proved. $\square$

The next proposition reveals a relation between the Gan–Gross–Prasad period integral and weak automorphic base change.

PROPOSITION 4.13. *Consider an element $V \in \mathfrak{V}_{(\mathbf{s})}$ and an element $\pi^V \in \mathfrak{C}(G^V, 1)$. Suppose that we can find a cuspidal automorphic form $\varphi \in L^2(G^V, \pi^V)$ satisfying*

$$\mathscr{P}(\varphi) := \int_{H^V(F) \backslash H^V(\mathbb{A}_F)} \varphi(h) dh \neq 0,$$

where dh is the Tamagawa measure on $H^V(\mathbb{A}_F)$. Assume that either one of the following two assumptions holds:

- (a) π^V is not $(G^V, \mathbf{T})$ -CAP (Definition 3.4);

(b) there is a prime v_0 of F split in E such that $\pi_{v_0}^V$ is supercuspidal.

Then weak automorphic base change of π^V (Definition 1.6), as an isobaric automorphic representation of $G'(\mathbb{A}_F) = \mathrm{GL}_n(\mathbb{A}_E) \times \mathrm{GL}_{n+1}(\mathbb{A}_E)$, exists. Moreover, if we put $\Pi := \mathrm{BC}(\pi^V) = \Pi_n \boxtimes \Pi_{n+1}$, then

- (1) the base change of $\chi_{\pi_\infty^V}$ is χ_{Π_∞} ;
- (2) in the situation (b), Π_m is cuspidal and hermitian (Definition 1.5) for $m = n, n+1$.

Proof. In the beginning of this subsection, we take $\mathfrak{V}_{(\mathbf{S})}^\circ = \{V\}$, $\chi^V = \chi_{(\pi^V)^\tau}$, and χ' to be the base change of χ^V .

Suppose that weak automorphic base change of π^V does not exist. Then χ' is not $(G', \mathbf{T})$ -CAP and $L_{\mathrm{cusp}}^2(G'(F) \backslash G'(\mathbb{A}_F)/K', 1)[\chi'] = \{0\}$. By our assumption on the non-vanishing of $\mathscr{P}$ on π^V , for every place v of F either archimedean or in $\mathbf{S}$, we can choose an element $f_v^V \in \mathcal{S}(G^V(F_v))$ of positive type (see the definition of positive type functions above [Zha14b, Prop. 2.12]) such that $J_{\pi_v^V}(f_v^V) > 0$. Moreover, we may assume that $f_{v_0}^V$ is a supercuspidal matrix coefficient in the situation of (b). For a prime v not in $\mathbf{S}$, we take $f_v^V := \mathbf{1}_{K_v^V}$. Put $f^V := \bigotimes_v f_v^V$, and shrink $K_{\mathbf{S}}^V$ if necessary so that $f^V \in \mathcal{S}(G^V(\mathbb{A}_F))_{K^V}$. By our choice of f^V , we have $J_{\pi^V}(f^V) > 0$, and $J_\pi(f^V) \geq 0$ for every $\pi \in \mathfrak{C}(G^V, 1)$. Note that the set

$$\mathfrak{C} := \{\pi \in \mathfrak{C}(G^V, 1) \mid L^2(G^V, \pi) \cap L_{\mathrm{cusp}}^2(G^V(F) \backslash G^V(\mathbb{A}_F)/K^V, 1)[\chi^V] \neq \{0\}\}$$

is finite, by a well-known result of Harish-Chandra that there are only finitely (up to isomorphism) cuspidal automorphic representations of $G^V(\mathbb{A}_F)$ with a given infinitesimal character and non-trivial K_V -invariants [HC68]. Thus, we may replace f_v^V by an element in the dense subspace $\mathcal{S}_{\mathrm{tr}}(G^V(F_v))$ for each archimedean place v non-split in E , such that

$$\sum_{\pi \in \mathfrak{C}} J_\pi(f^V) \neq 0.$$

Now we can apply Proposition 4.6(2) to f_v^V for every place v of F either archimedean or in $\mathbf{S}$, to obtain an element $f'_v \in \mathcal{S}(G'(F_v))$ as in that proposition. We may also assume that f'_{v_0} is a supercuspidal matrix coefficient in the situation (b). For a prime v not in $\mathbf{S}$, we take $f'_v := \mathbf{1}_{K'_v}$. Put $f' := \bigotimes_v f'_v$, which is an element of $\mathcal{S}(G'(\mathbb{A}_F))_{K'}$ after shrinking $K_{\mathbf{S}}'$ if necessary. Then by the relative fundamental lemma [Yun11, BP21a], we know that f' and $(f^V, 0, \dots)$ are complete smooth transfer in the sense of Definition 4.4(3).

We claim that there is a $\mathbf{T}$ -multiplier μ' of $\mathcal{S}(G'(\mathbb{A}_F))_{K'}$ satisfying $\mu'(\chi') = 1$ and such that $R(\mu' \star f')$ sends $L^2(G'(F) \backslash G'(\mathbb{A}_F)/K', 1)$ into

$$L_{\mathrm{cusp}}^2(G'(F) \backslash G'(\mathbb{A}_F)/K', 1)[\chi'].$$

In the situation (a), this follows from [Theorem 3.6](#) since χ' is not $(G', \mathbf{T})$ -CAP. In the situation (b), this follows from [Theorem 3.7](#), and the observation that $R(\mu' \star f')$ automatically annihilates the orthogonal complement of $L^2_{\text{cusp}}(G'(F) \backslash G'(\mathbb{A}_F)/K', 1)$ in $L^2(G'(F) \backslash G'(\mathbb{A}_F)/K', 1)$ since f'_{v_0} is a supercuspidal matrix coefficient. Similarly, there is a $\mathbf{T}$ -multiplier μ^V of $\mathcal{S}(G^V(\mathbb{A}_F))_{K^V}$ satisfying $\mu^V(\chi^V) = 1$, such that $R(\mu^V \star f^V)$ sends $L^2(G^V(F) \backslash G^V(\mathbb{A}_F)/K^V, 1)$ into $L^2_{\text{cusp}}(G^V(F) \backslash G^V(\mathbb{A}_F)/K^V, 1)[\chi^V]$. Moreover, by [Lemma 4.12](#), we may further assume that $\mu^V = \text{bc}^V(\mu')$. Thus, by [Proposition 4.8](#), $\mu' \star f'$ and $(\mu^V \star f^V, 0, \dots)$ are complete smooth transfer as well.

Now we run relative trace formulae from the unitary side to the general linear side. We have

$$\sum_{\pi \in \mathfrak{C}(G^V, 1)} J_{\pi}(\mu^V \star f^V) = \sum_{\pi \in \mathfrak{C}} J_{\pi}(\mu^V \star f^V) = \sum_{\pi \in \mathfrak{C}} J_{\pi}(f^V) \neq 0.$$

Since $\mu^V \star f^V$ is quasi-cuspidal, we have

$$\sum_{\delta^V \in B^V(F)} J_{\delta^V}(\mu^V \star f^V) = \sum_{\pi \in \mathfrak{C}(G^V, 1)} J_{\pi}(\mu^V \star f^V) \neq 0$$

by [Proposition 4.2](#). By [Proposition 4.7](#), we have

$$\sum_{\gamma \in B'(F)} I_{\gamma}(\mu' \star f') = \sum_{\delta^V \in B^V(F)} J_{\delta^V}(\mu^V \star f^V) \neq 0.$$

Since $\mu' \star f'$ is quasi-cuspidal, we have

$$\sum_{\tilde{\Pi} \in \mathfrak{C}(G', 1)} I_{\tilde{\Pi}}(\mu' \star f') = \sum_{\gamma \in B'(F)} I_{\gamma}(\mu' \star f') \neq 0$$

by [Proposition 4.1](#). However, this is a contradiction since the image of $R(\mu' \star f')$ is contained in $L^2_{\text{cusp}}(G'(F) \backslash G'(\mathbb{A}_F)/K', 1)[\chi']$, which is zero. Thus $\mathfrak{D}(G', 1)_{\chi'}^{\heartsuit} \neq \emptyset$; that is, weak automorphic base change of π^V exists. It remains to show the two additional claims.

For (1), it follows from the fact that $\mathfrak{D}(G', 1)_{\chi'}^{\heartsuit} \neq \emptyset$.

For (2), note that in the situation (b), we did not use the assumption that χ' is not $(G', \mathbf{T})$ -CAP in the main argument above. Thus, we should have $L^2_{\text{cusp}}(G'(F) \backslash G'(\mathbb{A}_F)/K', 1)[\chi'] \neq \{0\}$; in other words, $\Pi := \text{BC}(\pi^V)$ is cuspidal and we have $I_{\Pi}(\mu' \star f') \neq 0$. By [\(4.1\)](#),

$$\int_{Z'(\mathbb{A}_F)H'_2(F) \backslash H'_2(\mathbb{A}_F)} \phi(h_2) \eta(h_2) dh_2 \neq 0$$

for some cuspidal form ϕ in Π . Then by the main theorem of [\[Fli88\]](#), we know that $L(s, \Pi_m, \text{As}^{(-1)^{m+1}})$ has a pole at $s = 1$ for $m = n, n + 1$, which implies that Π_m is hermitian.

The proposition is proved. $\square$

The last theorem of this subsection contains results on weak automorphic base change, proved using Gan–Gross–Prasad period integrals.

THEOREM 4.14. *Let V° be a hermitian space over E with $G^\circ := \mathrm{U}(V^\circ)$, and let $\pi = \otimes_v \pi_v$ be an irreducible admissible representation of $G^\circ(\mathbb{A}_F)$.*

- (1) *If π is cuspidal automorphic satisfying that there exist infinitely many primes v of F split in E such that π_v is generic, then weak automorphic base change of π exists.*
- (2) *If π is cuspidal automorphic satisfying that there exists a prime v of F split in E such that π_v is supercuspidal, then weak automorphic base change of π exists and is cuspidal and hermitian (Definition 1.5).*
- (3) *If weak automorphic base change of π exists and is hermitian, then π is not $(G^\bullet, \mathbf{T}_0)$ -CAP (Definition 3.4) for every pure inner form $G^\bullet$ of G° .*
- (4) *If weak automorphic base change of π exists and is hermitian, then the base change of χ_{π_∞} is $\chi_{\mathrm{BC}(\pi)_\infty}$.*

Part (3) of the above theorem is consistent with the so-called CAP conjecture [Jia10, Conj. 6.1] (for unitary groups).

Proof. We prove the four statements at once via the induction on the rank of V° . The statements are all trivial when the rank of V° is 1. Assume that (1)–(4) are all known for all V° of rank at most n .

Now take a hermitian space V° of rank $n + 1$. By scaling the hermitian form, we may assume that there is an element $V = (V_n, V_{n+1}) \in \mathfrak{V}$ such that $V^\circ = V_{n+1}$. We consider (1) and (2) first. Since π is cuspidal automorphic, it gives an element $\pi_{n+1}^V \in \mathfrak{C}(G_{n+1}^V, 1)$. By the Burger–Sarnak trick [Zha14b, Prop. 2.14], we can find another element $\pi_n^V \in \mathfrak{C}(G_n^V, 1)$ such that

- $\mathscr{P}$ is non-zero on cuspidal automorphic forms in $L^2(G^V, \pi^V)$, where $\pi^V := \pi_n^V \boxtimes \pi_{n+1}^V \in \mathfrak{C}(G^V, 1)$;
- there is a prime v_0 of F split in E such that π_{n, v_0}^V is supercuspidal, and π_{n+1, v_0}^V is generic (resp. supercuspidal) in (1) (resp. (2)).

For (1), there are two cases. If π is $(G^\circ, \mathbf{T}_0)$ -CAP, then by definition there exist a proper parabolic subgroup P° of G° and a cuspidal automorphic representation σ of $M_{P^\circ}(\mathbb{A}_F)$, such that π_v is a constituent of $\mathrm{I}_{P^\circ}^{G^\circ}(\sigma_v)$ for all but finitely many primes v of F split in E . Write $M_{P^\circ} = \mathrm{U}(W^\circ) \times M'$ for some hermitian space W° of rank at most n and M' for a product of general linear groups, under which $\sigma = \sigma^\circ \boxtimes \sigma'$. Then there exist infinitely many primes v of F split in E such that σ_v° is generic. By the induction hypothesis on (1), weak automorphic base change of σ° exists, which implies that weak automorphic base change of π exists as well. If π is not $(G^\circ, \mathbf{T}_0)$ -CAP, then π_{n+1}^V is not $(G_{n+1}^V, \mathbf{T}_0)$ -CAP. By the induction hypothesis on (2) and (3), we know that π_n^V is not $(G_n^V, \mathbf{T}_0)$ -CAP. Thus, π^V is not $(G^V, \mathbf{T}_0)$ -CAP. By the situation (a) of

[Proposition 4.13](#), we know that weak automorphic base change of π^V , hence of π , exists.

For (2), by the situation (b) of [Proposition 4.13](#), we know that weak automorphic base change Π of π^V , hence of π , exists and is cuspidal and hermitian.

We prove (3) by contradiction. If π is $(G^\bullet, \mathbf{T})$ -CAP for some pure inner form $G^\bullet$ of G° , then by definition there exist a hermitian space $V^\bullet$ of rank $n + 1$ such that $G^\bullet = \mathrm{U}(V^\bullet)$, a proper parabolic subgroup $P^\bullet$ of $G^\bullet$, and a cuspidal automorphic representation σ of $M_{P^\bullet}(\mathbb{A}_F)$, such that π_v is a constituent of $\mathrm{I}_{P^\bullet}^{G^\bullet}(\sigma_v)$ for all but finitely many primes v of F split in E . Write $M_{P^\bullet} = \mathrm{U}(W^\bullet) \times M'$ for some hermitian space $W^\bullet$ of rank at most n and M' a non-trivial product of general linear groups, under which $\sigma = \sigma^\bullet \boxtimes \sigma'$. Since $\mathrm{BC}(\pi)$ is an isobaric sum of cuspidal automorphic representations with unitary central characters, $\mathrm{BC}(\pi)_w$ is generic for every place w of E , which implies that $\sigma^\bullet$ satisfies the assumption in (1). Thus, by the induction hypothesis on (1), weak automorphic base change of $\sigma^\bullet$ exists. Thus, we have $\mathrm{BC}(\pi) \simeq \mathrm{BC}(\sigma^\bullet) \boxplus \Pi \boxplus (\Pi^\vee \circ c) \boxplus \cdots$, where $c \in \mathrm{Gal}(E/F)$ is the involution, for at least one non-trivial Π . This contradicts the fact that $\mathrm{BC}(\pi)$ is an isobaric sum of mutually non-isomorphic conjugate self-dual cuspidal automorphic representations, as it is hermitian. Therefore, (3) is proved.

For (4), π is not $(G^\circ, \mathbf{T}_0)$ -CAP by (3). In particular, by the induction hypothesis, π^V is not $(G^V, \mathbf{T}_0)$ -CAP. Then (4) follows from [Proposition 4.13](#)(1).

The theorem is proved. $\square$

Remark 4.15. [Theorem 4.14](#) already follows from [\[Mok15\]](#), [\[KMSW14\]](#) (see the end of [\[KMSW14, §3.3\]](#)). However, our proof is different and does not use any knowledge from the endoscopy theory for unitary groups. Furthermore, our method can actually be used to show the local-global compatibility at all places where π_v is unramified as well, but the argument will implicitly relies on [\[Mok15\]](#), [\[KMSW14\]](#) as we will need [Lemma 4.9](#) for $\square$ containing primes inert in E (see [Remark 4.10](#)).

4.4. Gan–Gross–Prasad and Ichino–Ikeda conjectures. In this subsection, we complete the proofs of [Theorem 1.8](#) (for the Gan–Gross–Prasad conjecture), [Theorem 1.9](#), and [Theorem 1.10](#) (for the Ichino–Ikeda conjecture). We keep the setup in the previous two subsections.

We start with the following lemma as a preliminary on the descent of hermitian isobaric automorphic representations of $\mathrm{GL}_m(\mathbb{A}_E)$.

LEMMA 4.16. *Let Π be a hermitian isobaric automorphic representation of $\mathrm{GL}_m(\mathbb{A}_E)$ ([Definition 1.5](#)) for some integer $m \geq 1$. Let V be a hermitian space over E of rank m such that $\mathrm{U}(V)$ is quasi-split. Then there exists a cuspidal automorphic representation π of $\mathrm{U}(V)(\mathbb{A}_F)$ satisfying that for every prime v of*

F split in E such that Π_v is unramified, π_v is unramified, and the base change of χ_{π_v} is χ_{Π_v} . In particular, we have $\Pi \simeq \text{BC}(\pi)$ and the base change of χ_{π_∞} is χ_{Π_∞} .

Proof. The existence of π follows from the automorphic descent construction [GRS11]. The last assertion follows from Theorem 4.14(4). $\square$

Proof of Theorem 1.8. There are two directions.

(2) $\Rightarrow$ (1): We take $V = (V_n, V_{n+1}) \in \mathfrak{V}$ and $\pi^V := \pi_n \boxtimes \pi_{n+1} \in \mathfrak{C}(G^V, 1)$. Put $\Pi := \Pi_n \boxtimes \Pi_{n+1} \in \mathfrak{C}(G', 1)$. By Theorem 4.14(3), π^V is not (G^V, T_0) -CAP. Thus, by the same argument as for the situation (a) in Proposition 4.13, we obtain

$$\sum_{\tilde{\Pi} \in \mathfrak{C}(G', 1)} I_{\tilde{\Pi}}(\mu' \star f') \neq 0,$$

in which the image of $R(\mu' \star f')$ is contained in $L_{\text{cusp}}^2(G'(F) \backslash G'(\mathbb{A}_F)/K', 1)[\chi_{\Pi^\Gamma}]$. However, by the strong multiplicity one property [Ram18, Th. A], we have

$$L_{\text{cusp}}^2(G'(F) \backslash G'(\mathbb{A}_F)/K', 1)[\chi_{\Pi^\Gamma}] = L^2(G', \Pi).$$

Thus, we have $I_{\Pi}(\mu' \star f') \neq 0$. By (4.1),

$$\int_{H'_1(F) \backslash H'_1(\mathbb{A}_F)} \phi(h_1) dh_1 \neq 0$$

for some cusp form ϕ in Π . Then by [JPSS83], we have $L(\frac{1}{2}, \Pi) \neq 0$.

(1) $\Rightarrow$ (2): Again, put $\Pi := \Pi_n \boxtimes \Pi_{n+1}$. Take $\mathbf{S}$ to be the finite set of primes v of F at which either E or Π is ramified. For $m = n, n+1$, since Π_m is hermitian, $L(s, \Pi_m, \text{As}^{(-1)^{m+1}})$ has a pole at $s = 1$, which implies that Π_m is $\eta_{E/F}^{m+1}$ -distinguished by $\text{GL}_m(\mathbb{A}_F)$. In particular, Π belongs to $\mathfrak{C}(G', 1)$, and there exists a cusp form ϕ in Π that is fixed by $\prod_{v \notin \mathbf{S}} K'_v$ such that

$$\int_{Z'(\mathbb{A}_F)H'_2(F) \backslash H'_2(\mathbb{A}_F)} \phi(h_2) \eta(h_2) dh_2 \neq 0.$$

Since $L(\frac{1}{2}, \Pi) \neq 0$, by [JPSS83], there exists a cusp form ϕ' in Π that is fixed by $\prod_{v \notin \mathbf{S}} K'_v$ such that

$$\int_{H'_1(F) \backslash H'_1(\mathbb{A}_F)} \phi'(h_1) dh_1 \neq 0.$$

Together, we can find an element $f' = \bigotimes_v f'_v \in \mathcal{S}(G'(\mathbb{A}_F))$ with $f'_v \in \mathcal{S}_{\text{tr}}(G'(F_v))$ for every archimedean place v of F and $f'_v = \mathbf{1}_{K'_v}$ for every prime v of F not in $\mathbf{S}$, such that

$$I_{\Pi}(f') \neq 0$$

by (4.1). After shrinking $K'_\mathbf{S}$ if necessary, we may assume $f' \in \mathcal{S}(G'(\mathbb{A}_F))_{K'}$. Now we can apply Proposition 4.6(1) to f'_v for every place v of F that is either archimedean or in $\mathbf{S}$, to obtain elements $f_v^V \in \mathcal{S}(G^V(F_v))$ as in that

proposition. For $V \in \mathfrak{V}_{(\mathbf{S})}$ and a prime v not in $\mathbf{S}$, we put $f_v^V := \mathbf{1}_{K_v^V}$, hence obtain an element $f^V := \bigotimes_v f_v^V \in \mathcal{S}(G^V(\mathbb{A}_F))_{K^V}$ for some $K_{\mathbf{S}}^V$. For $V \notin \mathfrak{V}_{(\mathbf{S})}$, we put $f^V = 0$. By the relative fundamental lemma [Yun11, BP21a], f' and $(f^V)_{V \in \mathfrak{V}}$ are complete smooth transfer in the sense of Definition 4.4(3).

To pass to unitary groups, we consider $\mathfrak{V}_{(\mathbf{S})}^\circ = \mathfrak{V}_{(\mathbf{S})}$, $\chi' = \chi_{\Pi^\Gamma}$, and a T-character χ^V for G^V whose base change is χ' for every $V \in \mathfrak{V}_{(\mathbf{S})}$ (which is possible by Lemma 4.16), at the beginning of Section 4.3. By the strong multiplicity one property [Ram18, Th. A], we know that χ' is not G' -CAP. By Theorem 3.6, there is a T-multiplier μ' of $\mathcal{S}(G'(\mathbb{A}_F))_{K'}$ satisfying $\mu'(\chi') = 1$ and such that $R(\mu' \star f')$ sends $L^2(G'(F) \backslash G'(\mathbb{A}_F)/K', 1)$ into $L_{\text{cusp}}^2(G'(F) \backslash G'(\mathbb{A}_F)/K', 1)[\chi']$, which coincides with $L^2(G', \Pi)$. For each $V \in \mathfrak{V}_{(\mathbf{S})}$, we know that χ^V is not G^V -CAP, since otherwise π^{V^*} would be (G^V, T_0) -CAP, which contradicts Theorem 4.14(3); here $V^* \in \mathfrak{V}$ is the unique element such that G^{V^*} is quasi-split and π^{V^*} is a cuspidal automorphic representation of $G^{V^*}(\mathbb{A}_F)$ as in Lemma 4.16. Then by Theorem 3.6, for every $V \in \mathfrak{V}_{(\mathbf{S})}$, there is a T-multiplier μ^V of $\mathcal{S}(G^V(\mathbb{A}_F))_{K^V}$ satisfying $\mu^V(\chi^V) = 1$ and such that $R(\mu^V \star f^V)$ sends

$$L^2(G^V(F) \backslash G^V(\mathbb{A}_F)/K^V, 1)$$

into $L_{\text{cusp}}^2(G^V(F) \backslash G^V(\mathbb{A}_F)/K^V, 1)[\chi^V]$. Moreover, by Lemma 4.12, we may further assume that $\mu^V = \text{bc}^V(\mu')$ for every $V \in \mathfrak{V}_{(\mathbf{S})}$. To summarize, we conclude that $\mu' \star f'$ and $((\mu^V \star f^V)_{V \in \mathfrak{V}_{(\mathbf{S})}}, 0, \dots)$ are complete smooth transfer by Proposition 4.8.

Now we run relative trace formulae from the general linear side to the unitary side. We have

$$\sum_{\tilde{\Pi} \in \mathfrak{C}(G', 1)} I_{\tilde{\Pi}}(\mu' \star f') = I_{\Pi}(\mu' \star f') = I_{\Pi}(f') \neq 0.$$

Since $\mu' \star f'$ is quasi-cuspidal, we have

$$\sum_{\gamma \in B'(F)} I_{\gamma}(\mu' \star f') = \sum_{\tilde{\Pi} \in \mathfrak{C}(G', 1)} I_{\tilde{\Pi}}(\mu' \star f') \neq 0$$

by Proposition 4.1. By Proposition 4.7, we have

$$\sum_{V \in \mathfrak{V}_{(\mathbf{S})}^\circ} \sum_{\delta^V \in B^V(F)} J_{\delta^V}(\mu^V \star f^V) = \sum_{\gamma \in B'(F)} I_{\gamma}(\mu' \star f') \neq 0.$$

Thus, we can choose some $V \in \mathfrak{V}_{(\mathbf{S})}$ such that

$$\sum_{\delta^V \in B^V(F)} J_{\delta^V}(\mu^V \star f^V) \neq 0.$$

Since $\mu^V \star f^V$ is quasi-cuspidal, we have

$$\sum_{\pi \in \mathfrak{C}(G^V, 1)} J_\pi(\mu^V \star f^V) = \sum_{\delta^V \in B^V(F)} J_{\delta^V}(\mu^V \star f^V) \neq 0$$

by [Proposition 4.2](#). Therefore, by the property of $\mu^V \star f^V$, we can find some element $\pi \in \mathfrak{C}(G^V, 1)$ satisfying

$$L^2(G^V, \pi) \cap L^2_{\text{cusp}}(G^V(F) \backslash G^V(\mathbb{A}_F)/K^V, 1)[\chi^V] \neq \{0\},$$

such that

$$J_\pi(f^V) = J_\pi(\mu^V \star f^V) \neq 0.$$

In particular, the weak automorphic base change of π is isomorphic to Π , and $\mathscr{P}$ is non-vanishing on cuspidal automorphic forms in $L^2(G^V, \pi)$ by [\(4.3\)](#). Thus, (2) is achieved.

[Theorem 1.8](#) is proved. $\square$

Remark 4.17. In the proof of [Theorem 1.8](#), we actually obtain a stronger statement in the direction (1) $\Rightarrow$ (2) by further requiring in (2) that

- the base change of $\chi_{\pi_{m,\infty}}$ is $\chi_{\Pi_{m,\infty}}$ for $m = n, n+1$;
- the form $\varphi_n \otimes \varphi_{n+1}$ is fixed by a relative hyperspecial maximal subgroup at every prime v of F that is unramified in E and such that $\Pi_{n,v} \otimes \Pi_{n+1,v}$ is unramified.

Proof of [Theorem 1.9](#). Let $V = (V_n, V_{n+1}) \in \mathfrak{V}$ be the unique element such that G^V is quasi-split. Since Π_{n+1} is hermitian, by [Lemma 4.16](#), we have an element $\pi_{n+1} \in \mathfrak{C}(G^V_{n+1}, 1)$ such that $\text{BC}(\pi_{n+1}) \simeq \Pi_{n+1}$. By the Burger–Sarnak trick [[Zha14b](#), Prop. 2.14], we can find another element $\pi_n \in \mathfrak{C}(G^V_n, 1)$ that is supercuspidal at some prime of F split in E , fulfilling the situation in [Theorem 1.8\(2\)](#). By [Theorem 4.14\(2\)](#), $\text{BC}(\pi_n)$ exists and is cuspidal and hermitian, which we denote by Π_n . Moreover, we have

$$L(\tfrac{1}{2}, \Pi_n \times \Pi_{n+1}) \neq 0$$

by [Theorem 1.8](#). The theorem is proved. $\square$

Proof of [Theorem 1.10](#). We continue the proof of [Theorem 1.8](#). Using the endoscopic classification for generic packets obtained in [[Mok15](#)], [[KMSW14](#)] and the local Gan–Gross–Prasad [[BP20b](#)], we arrive at the identity

$$I_\Pi(f') = J_\pi(f^V) \neq 0$$

for some $f' = \bigotimes_v f'_v \in \mathcal{S}(G'(\mathbb{A}_F))_{K'}$ and $f^V = \bigotimes_v f_v^V \in \mathcal{S}(G^V(\mathbb{A}_F))_{K^V}$ such that f'_v and f_v^V have matching orbital integrals for every place v . The remaining argument is the same as in the proof of [[BP21b](#), Th. 5]. $\square$

Appendix A. Extending a result of Delorme to reductive groups

In this appendix, we extend a theorem of Delorme [Del86, Th. 1.7] from semisimple groups to reductive groups. Let the setup be as in Section 2.3. In particular, $G = \underline{G}(\mathbb{R})$ for a connected reductive algebraic group $\underline{G}$ over $\mathbb{R}$. Denote by $C_c^\infty(G)_{(K)}$ and $\mathcal{S}(G)_{(K)}$ the subalgebras of bi- K -finite functions in $C_c^\infty(G)$ and $\mathcal{S}(G)$, respectively. Recall from Definition 2.8 that $\mathcal{N}(\mathfrak{h}_\mathbb{C}^*)$ is the space of holomorphic functions on $\mathfrak{h}_\mathbb{C}^*$ that have rapid decay on vertical strips.

PROPOSITION A.1. *For every element $\mu \in \mathcal{N}(\mathfrak{h}_\mathbb{C}^*)$, there is a unique linear operator*

$$\mu \star : C_c^\infty(G)_{(K)} \rightarrow \mathcal{S}(G)_{(K)},$$

such that

$$\pi(\mu \star f) = \mu(\chi_\pi) \cdot \pi(f)$$

holds for every $f \in C_c^\infty(G)_{(K)}$ and every irreducible admissible representation π of G .

When G is semisimple, linear, and connected (in the analytic topology), this is exactly [Del86, Th. 1.7]. In general, we need to show that there is a bilinear map

$$(A.1) \quad \mathcal{N}(\mathfrak{h}_\mathbb{C}^*)^W \times C_c^\infty(G)_{(K)} \rightarrow \mathcal{S}(G)_{(K)}$$

sending (μ, f) to $\mu \star f$, satisfying the requirement in the proposition.

In what follows, for a real Lie group H , we denote by H^0 its neutral connected component. By the Iwasawa decomposition, the natural map $K/K^0 \rightarrow G/G^0$ is an isomorphism, which gives rise to a decomposition

$$(A.2) \quad C_c^\infty(G)_{(K)} = \bigoplus_{k \in K/K^0} C_c^\infty(kG^0)_{(K^0)}.$$

Here, $C_c^\infty(kG^0)_{(K^0)}$ denotes the space of compactly supported smooth functions on kG^0 that are bi- K^0 -finite. We first reduce the construction of (A.1) to the one for G^0 ,

$$(A.3) \quad \mathcal{N}(\mathfrak{h}_\mathbb{C}^*)^W \times C_c^\infty(G^0)_{(K^0)} \rightarrow \mathcal{S}(G^0)_{(K^0)},$$

sending (μ, f) to $\mu \star f$, satisfying $\pi(\mu \star f) = \mu(\chi_\pi) \cdot \pi(f)$ for every irreducible admissible representation π of G^0 . Indeed, once we have (A.3), we may define

$$\mu \star f := \sum_{k \in K/K^0} L(k)(\mu \star L(k^{-1})f_k)$$

for $(\mu, f) \in \mathcal{N}(\mathfrak{h}_\mathbb{C}^*)^W \times C_c^\infty(G)_{(K)}$, where $f = \sum_{k \in K/K^0} f_k$ is the decomposition of f under (A.2), and L denotes the left regular action. Since the restriction to G^0 of an irreducible admissible representation of G is a finite direct sum of irreducible admissible representations of G^0 with the same infinitesimal character,

it is easy to check that the above bilinear map satisfies the requirement in the proposition.

Now it remains to construct (A.3). Denote by $\underline{Z}$ and $\underline{G}_{\text{der}}$ the center and the derived subgroup of $\underline{G}$, respectively. Put $Z := \underline{Z}(\mathbb{R})$ and $G_{\text{der}} := \underline{G}_{\text{der}}(\mathbb{R})$. Put $K_Z^0 := K \cap Z^0$, $K_{\text{der}}^0 := K \cap G_{\text{der}}^0$, and $Z_{\text{der}}^0 := Z^0 \cap G_{\text{der}} = K_Z^0 \cap K_{\text{der}}^0$. Then Z_{der}^0 is a finite group, and the natural inclusions induce isomorphisms $(K_Z^0 \times K_{\text{der}}^0)/Z_{\text{der}}^0 \xrightarrow{\sim} K^0$ and

$$(A.4) \quad (Z^0 \times G_{\text{der}}^0)/Z_{\text{der}}^0 \xrightarrow{\sim} G^0.$$

For finite subsets $\mathfrak{T}_Z$ and $\mathfrak{T}_{\text{der}}$ of K_Z^0 -types and K_{der}^0 -types, respectively, we define $\mathfrak{T}$ to be the set of K^0 -types whose inflation to $K_Z^0 \times K_{\text{der}}^0$ is an exterior tensor product of elements in $\mathfrak{T}_Z$ and $\mathfrak{T}_{\text{der}}$. Similarly, for compact subsets Ω_Z and Ω_{der} of Z^0 and G_{der}^0 that are bi-invariant under K_Z^0 and K_{der}^0 , respectively, we let Ω be the image of $\Omega_Z \times \Omega_{\text{der}}$ under the isomorphism (A.4), which is a compact subset of G^0 bi-invariant under K^0 . Clearly, we have

$$C_c^\infty(G^0)_{(K^0)} = \bigcup_{\Omega} \bigcup_{\mathfrak{T}} C_\Omega^\infty(G^0)_{(\mathfrak{T})},$$

where $C_\Omega^\infty(G^0)_{(\mathfrak{T})}$ denotes the subspace of smooth functions supported on Ω of bi- K^0 -types in $\mathfrak{T}$. Thus, it suffices to construct (A.3) for a fixed subspace $C_\Omega^\infty(G^0)_{(\mathfrak{T})}$ of $C_c^\infty(G^0)_{(K^0)}$. The isomorphism (A.4) induces a natural isomorphism

$$(A.5) \quad (C_{\Omega_Z}^\infty(Z^0)_{(\mathfrak{T}_Z)} \widehat{\otimes} C_{\Omega_{\text{der}}}^\infty(G_{\text{der}}^0)_{(\mathfrak{T}_{\text{der}})})^{Z_{\text{der}}^0} \xrightarrow{\sim} C_\Omega^\infty(G^0)_{(\mathfrak{T})}$$

of Fréchet spaces, where $\widehat{\otimes}$ stands for the projective completed tensor product.¹²

On the other hand, let $\mathfrak{z}^*$ and $\mathfrak{h}_{\text{der}}^*$ be the real vector spaces spanned by the weight lattices of $\underline{Z}_{\mathbb{C}}$ and the abstract Cartan group of $\underline{G}_{\text{der}, \mathbb{C}}$, respectively.

LEMMA A.2. *The decomposition $\mathfrak{h}^* = \mathfrak{z}^* \oplus \mathfrak{h}_{\text{der}}^*$ induced from (A.4) induces a natural isomorphism*

$$\mathcal{N}(\mathfrak{z}_{\mathbb{C}}^*) \widehat{\otimes} \mathcal{N}(\mathfrak{h}_{\text{der}, \mathbb{C}}^*)^{\mathbb{W}} \xrightarrow{\sim} \mathcal{N}(\mathfrak{h}_{\mathbb{C}}^*)^{\mathbb{W}}$$

of Fréchet spaces. Here, $\mathcal{N}(\mathfrak{z}_{\mathbb{C}}^*)$ and $\mathcal{N}(\mathfrak{h}_{\text{der}, \mathbb{C}}^*)$ are defined similarly as for $\mathcal{N}(\mathfrak{h}_{\mathbb{C}}^*)$.

Proof. Let Z_{sp}^0 , H_{der}^0 , and H^0 be the split real analytic tori whose cotangent spaces are $\mathfrak{z}^*$, $\mathfrak{h}_{\text{der}}^*$, and $\mathfrak{h}^*$, respectively. By the classical Paley–Wiener theorem for Schwartz functions on split real analytic tori, we have canonical isomorphisms

$$\mathcal{S}(Z_{\text{sp}}^0) \simeq \mathcal{N}(\mathfrak{z}_{\mathbb{C}}^*), \quad \mathcal{S}(H_{\text{der}}^0) \simeq \mathcal{N}(\mathfrak{h}_{\text{der}, \mathbb{C}}^*), \quad \mathcal{S}(H^0) \simeq \mathcal{N}(\mathfrak{h}_{\mathbb{C}}^*)$$

¹²In general this is not correct if one does not fix the K^0 -types and the support.

induced by Fourier transforms. Since $H^0 = Z_{\text{sp}}^0 \times H_{\text{der}}^0$, the natural map $\mathcal{S}(Z_{\text{sp}}^0) \widehat{\otimes} \mathcal{S}(H_{\text{der}}^0) \rightarrow \mathcal{S}(H^0)$ is an isomorphism of Fréchet spaces by [AG10, Cor. 2.6.3]. By taking W -invariants, we obtain the isomorphism in the lemma. $\square$

Now take $\mu_1 \in \mathcal{N}(\mathfrak{z}_{\mathbb{C}}^*)$, $\mu_2 \in \mathcal{N}(\mathfrak{h}_{\text{der}, \mathbb{C}}^*)^W$, and put $\mu := \mu_1 \otimes \mu_2 \in \mathcal{N}(\mathfrak{h}_{\mathbb{C}}^*)^W$. By [Del86, Th. 1.7], there is a linear map

$$\mu_2 \star: C_{\Omega_{\text{der}}}^{\infty}(G_{\text{der}}^0)_{(\mathfrak{T}_{\text{der}})} \rightarrow \mathcal{S}(G_{\text{der}}^0)_{(\mathfrak{T}_{\text{der}})}$$

sending f to $\mu_2 \star f$, satisfying $\pi_{\text{der}}(\mu_2 \star f) = \mu_2(\chi_{\pi_{\text{der}}}) \cdot \pi_{\text{der}}(f)$ for every irreducible admissible representation π_{der} of G_{der}^0 . Similarly, by [Sak18, Th. 2.1.2], there is a linear map

$$\mu_1 \star: C_{\Omega_Z}^{\infty}(Z^0)_{(\mathfrak{T}_Z)} \rightarrow \mathcal{S}(Z^0)_{(\mathfrak{T}_Z)}$$

sending f to $\mu_1 \star f$, satisfying $\xi(\mu_1 \star f) = \mu_1(\chi_{\xi}) \cdot \xi(f)$ for every smooth character ξ of Z^0 . Moreover, by the injectivity of the (operator valued) Fourier transform and the closed graph theorem, both $\mu_1 \star$ and $\mu_2 \star$ are continuous. Therefore, $(\mu_1 \star) \otimes (\mu_2 \star)$ extends uniquely to a continuous linear map

$$\begin{aligned} \mu \star: C_{\Omega}^{\infty}(G^0)_{(\mathfrak{T})} &\simeq (C_{\Omega_Z}^{\infty}(Z^0)_{(\mathfrak{T}_Z)} \widehat{\otimes} C_{\Omega_{\text{der}}}^{\infty}(G_{\text{der}}^0)_{(\mathfrak{T}_{\text{der}})})^{Z_{\text{der}}^0} \\ &\rightarrow (\mathcal{S}(G_{\text{der}}^0)_{(\mathfrak{T}_{\text{der}})} \widehat{\otimes} \mathcal{S}(Z^0)_{(\mathfrak{T}_Z)})^{Z_{\text{der}}^0} \rightarrow \mathcal{S}(G^0)_{(\mathfrak{T})} \end{aligned}$$

by (A.5). Let π be an irreducible admissible representation of G^0 . By the decomposition (A.4), we may write the inflation of π to $Z^0 \times G_{\text{der}}^0$ as $\xi \boxtimes \pi_{\text{der}}$, where ξ is a smooth character of Z^0 and π_{der} is an irreducible admissible representation of G_{der}^0 . By construction, we have $\pi(\mu \star f) = \mu(\chi_{\pi})\pi(f)$ for every $f \in C_{\Omega_Z}^{\infty}(Z^0)_{(\mathfrak{T}_Z)} \otimes C_{\Omega_{\text{der}}}^{\infty}(G_{\text{der}}^0)_{(\mathfrak{T}_{\text{der}})}$ that is Z_{der}^0 -invariant. Since the map $f \in C_{\Omega}^{\infty}(G^0)_{(\mathfrak{T})} \mapsto \pi(f)$ is continuous when we equip the space of continuous endomorphisms of π with the topology of pointwise convergence, we deduce by density that $\pi(\mu \star f) = \mu(\chi_{\pi})\pi(f)$ holds for every $f \in C_{\Omega}^{\infty}(G^0)_{(\mathfrak{T})}$. In summary, we have constructed our desired bilinear map from $\mathcal{N}(\mathfrak{z}_{\mathbb{C}}^*) \otimes \mathcal{N}(\mathfrak{h}_{\text{der}, \mathbb{C}}^*)^W \times C_{\Omega}^{\infty}(G^0)_{(\mathfrak{T})}$ to $\mathcal{S}(G^0)_{(\mathfrak{T})}$. Now again by the injectivity of the (operator valued) Fourier transform and the closed graph theorem, we know that, for every $f \in C_{\Omega}^{\infty}(G^0)_{(\mathfrak{T})}$, the map

$$(A.6) \quad \star f: \mathcal{N}(\mathfrak{z}_{\mathbb{C}}^*) \times \mathcal{N}(\mathfrak{h}_{\text{der}, \mathbb{C}}^*)^W \rightarrow \mathcal{S}(G^0)_{(\mathfrak{T})}$$

sending (μ_1, μ_2) to $(\mu_1 \otimes \mu_2) \star f$ is a separately continuous bilinear map. Therefore, (A.6) extends uniquely to a continuous map $\star f: \mathcal{N}(\mathfrak{z}_{\mathbb{C}}^*) \widehat{\otimes} \mathcal{N}(\mathfrak{h}_{\text{der}, \mathbb{C}}^*)^W \rightarrow \mathcal{S}(G^0)_{(\mathfrak{T})}$ by [Tr 67, Th. 34.1]. By Lemma A.2, we obtain the desired bilinear map $\mathcal{N}(\mathfrak{h}_{\mathbb{C}}^*)^W \times C_{\Omega}^{\infty}(G^0)_{(\mathfrak{T})} \rightarrow \mathcal{S}(G^0)_{(\mathfrak{T})}$ sending (μ, f) to $\mu \star f$, satisfying $\pi(\mu \star f) = \mu(\chi_{\pi}) \cdot \pi(f)$ for every irreducible admissible representation π of G^0 . Taking union over all $\mathfrak{T}$ and Ω , we obtain (A.3).

Proposition A.1 is proved.

References

- [AG10] A. AIZENBUD and D. GOUREVITCH, The de-Rham theorem and Shapiro lemma for Schwartz function on Nash manifolds, *Israel J. Math.* **177** (2010), 155–188. [MR 2684417](#). [Zbl 1208.58002](#). <https://doi.org/10.1007/s11856-010-0042-9>.
- [Art81] J. ARTHUR, The trace formula in invariant form, *Ann. of Math.* (2) **114** no. 1 (1981), 1–74. [MR 0625344](#). [Zbl 0495.22006](#). <https://doi.org/10.2307/1971376>.
- [Art05] J. ARTHUR, An introduction to the trace formula, in *Harmonic Analysis, the Trace Formula, and Shimura Varieties*, *Clay Math. Proc.* **4**, Amer. Math. Soc., Providence, RI, 2005, pp. 1–263. [MR 2192011](#). [Zbl 1152.11021](#).
- [Art13] J. ARTHUR, *The Endoscopic Classification of Representations*, *Amer. Math. Soc. Colloq. Publ.* **61**, Amer. Math. Soc., Providence, RI, 2013, Orthogonal and symplectic groups. [MR 3135650](#). [Zbl 1310.22014](#). <https://doi.org/10.1090/coll/061>.
- [Bar88] W. H. BARKER, L^p harmonic analysis on $SL(2, \mathbf{R})$, *Mem. Amer. Math. Soc.* **76** no. 393 (1988), iv+110. [MR 0946617](#). [Zbl 0655.22011](#). <https://doi.org/10.1090/memo/0393>.
- [BP20a] R. BEUZART-PLESSIS, Comparison of local relative characters and the Ichino-Ikeda conjecture for unitary groups, *J. Inst. Math. Jussieu* (2020), 52 pp., published online 22 January 2020. <https://doi.org/10.1017/S1474748019000707>.
- [BP20b] R. BEUZART-PLESSIS, *A Local Trace Formula for the Gan-Gross-Prasad Conjecture for Unitary Groups: The Archimedean Case*, *Astérisque* **418**, Math. Soc. France, Paris, 2020. [MR 4146145](#). [Zbl 07269852](#). <https://doi.org/10.24033/ast>.
- [BP21a] R. BEUZART-PLESSIS, A new proof of the Jacquet-Rallis fundamental lemma, *Duke Math. J.* (2021), 10 pp., published online 16 April 2021. <https://doi.org/10.1215/00127094-2020-0090>.
- [BP21b] R. BEUZART-PLESSIS, Plancherel formula for $GL_n(F) \backslash GL_n(E)$ and applications to the Ichino-Ikeda and formal degree conjectures for unitary groups, *Invent. Math.* (2021), published online 23 February 2021. <https://doi.org/10.1007/s00222-021-01032-6>.
- [BPCZ20] R. BEUZART-PLESSIS, P.-H. CHAUDOUARD, and M. ZYDOR, The global Gan-Gross-Prasad conjecture for unitary groups: the endoscopic case, 2020. [arXiv 2007.05601](#).
- [Cas89] W. CASSELMAN, Canonical extensions of Harish-Chandra modules to representations of G , *Canad. J. Math.* **41** no. 3 (1989), 385–438. [MR 1013462](#). [Zbl 0702.22016](#). <https://doi.org/10.4153/CJM-1989-019-5>.
- [CZ21] P.-H. CHAUDOUARD and M. ZYDOR, Le transfert singulier pour la formule des traces de Jacquet-Rallis, *Compos. Math.* **157** no. 2 (2021), 303–434. [MR 4234897](#). [Zbl 07327256](#). <https://doi.org/10.1112/S0010437X20007599>.

- [Del84] P. DELORME, Multipliers for the convolution algebra of left and right K -finite compactly supported smooth functions on a semisimple Lie group, *Invent. Math.* **75** no. 1 (1984), 9–23. MR 0728136. Zbl 0536.43007. <https://doi.org/10.1007/BF01403087>.
- [Del86] P. DELORME, Formules limites et formules asymptotiques pour les multiplicités dans $L^2(G/\Gamma)$, *Duke Math. J.* **53** no. 3 (1986), 691–731. MR 0860667. Zbl 0623.22012. <https://doi.org/10.1215/S0012-7094-86-05338-X>.
- [Don82] H. DONNELLY, On the cuspidal spectrum for finite volume symmetric spaces, *J. Differential Geometry* **17** no. 2 (1982), 239–253. MR 0664496. Zbl 0494.58029. <https://doi.org/10.4310/jdg/1214436921>.
- [Fli88] Y. Z. FLICKER, Twisted tensors and Euler products, *Bull. Soc. Math. France* **116** no. 3 (1988), 295–313. MR 0984899. Zbl 0674.10026. <https://doi.org/10.24033/bsmf.2099>.
- [GGP12] W. T. GAN, B. H. GROSS, and D. PRASAD, Symplectic local root numbers, central critical L values, and restriction problems in the representation theory of classical groups, in *Sur les Conjectures de Gross et Prasad. I*, *Astérisque* **346**, Math. Soc. France, Paris, 2012, pp. 1–109. MR 3202556. Zbl 1280.22019.
- [GRS11] D. GINZBURG, S. RALLIS, and D. SOUDRY, *The descent map from automorphic representations of $GL(n)$ to classical groups*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2011. MR 2848523. Zbl 1233.11056. <https://doi.org/10.1142/9789814304993>.
- [HC51] HARISH-CHANDRA, On some applications of the universal enveloping algebra of a semisimple Lie algebra, *Trans. Amer. Math. Soc.* **70** (1951), 28–96. MR 0044515. Zbl 0042.12701. <https://doi.org/10.2307/1990524>.
- [HC66] HARISH-CHANDRA, Discrete series for semisimple Lie groups. II. Explicit determination of the characters, *Acta Math.* **116** (1966), 1–111. MR 0219666. Zbl 0199.20102. <https://doi.org/10.1007/BF02392813>.
- [HC68] HARISH-CHANDRA, *Automorphic forms on semisimple Lie groups*, *Lecture Notes in Mathematics*, No. 62, Springer-Verlag, Berlin-New York, 1968, Notes by J. G. M. Mars. MR 0232893. Zbl 0186.04702.
- [HC75] HARISH-CHANDRA, Harmonic analysis on real reductive groups. I. The theory of the constant term, *J. Functional Analysis* **19** (1975), 104–204. MR 0399356. Zbl 0315.43002. [https://doi.org/10.1016/0022-1236\(75\)90034-8](https://doi.org/10.1016/0022-1236(75)90034-8).
- [HC76] HARISH-CHANDRA, Harmonic analysis on real reductive groups. III. The Maass-Selberg relations and the Plancherel formula, *Ann. of Math.* (2) **104** no. 1 (1976), 117–201. MR 0439994. Zbl 0331.22007. <https://doi.org/10.2307/1971058>.
- [Har14] R. N. HARRIS, The refined Gross-Prasad conjecture for unitary groups, *Int. Math. Res. Not. IMRN* no. 2 (2014), 303–389. MR 3159075. Zbl 1322.11047. <https://doi.org/10.1093/imrn/rns219>.

- [III10] A. ICHINO and T. IKEDA, On the periods of automorphic forms on special orthogonal groups and the Gross-Prasad conjecture, *Geom. Funct. Anal.* **19** no. 5 (2010), 1378–1425. MR 2585578. Zbl 1216.11057. <https://doi.org/10.1007/s00039-009-0040-4>.
- [JPSS83] H. JACQUET, I. I. PIATETSKII-SHAPIRO, and J. A. SHALIKA, Rankin-Selberg convolutions, *Amer. J. Math.* **105** no. 2 (1983), 367–464. MR 0701565. Zbl 0525.22018. <https://doi.org/10.2307/2374264>.
- [JS81] H. JACQUET and J. A. SHALIKA, On Euler products and the classification of automorphic forms. II, *Amer. J. Math.* **103** no. 4 (1981), 777–815. MR 0623137. Zbl 0491.10020. <https://doi.org/10.2307/2374050>.
- [JR11] H. JACQUET and S. RALLIS, On the Gross-Prasad conjecture for unitary groups, in *On certain L-functions*, *Clay Math. Proc.* **13**, Amer. Math. Soc., Providence, RI, 2011, pp. 205–264. MR 2767518. Zbl 1222.22018.
- [Jia10] D. JIANG, On some topics in automorphic representations, in *Fourth International Congress of Chinese Mathematicians*, *AMS/IP Stud. Adv. Math.* **48**, Amer. Math. Soc., Providence, RI, 2010, pp. 177–199. MR 2744221. Zbl 1271.11058. <https://doi.org/10.2298/pim141026009j>.
- [KMSW14] T. KALETHA, A. MINGUEZ, S. W. SHIN, and P.-J. WHITE, Endoscopic classification of representations: Inner forms of unitary groups, 2014. arXiv 1409.3731.
- [Kna01] A. W. KNAPP, *Representation Theory of Semisimple Groups, Princeton Landmarks in Math.*, Princeton Univ. Press, Princeton, NJ, 2001, An overview based on examples, reprint of the 1986 original. MR 1880691. Zbl 0993.22001. <https://doi.org/10.1515/9781400883974>.
- [Kos75] B. KOSTANT, On the tensor product of a finite and an infinite dimensional representation, *J. Functional Analysis* **20** no. 4 (1975), 257–285. MR 0414796. Zbl 0355.17010. [https://doi.org/10.1016/0022-1236\(75\)90035-x](https://doi.org/10.1016/0022-1236(75)90035-x).
- [KK86] B. KOSTANT and S. KUMAR, The nil Hecke ring and cohomology of G/P for a Kac-Moody group G , *Adv. in Math.* **62** no. 3 (1986), 187–237. MR 0866159. Zbl 0641.17008. [https://doi.org/10.1016/0001-8708\(86\)90101-5](https://doi.org/10.1016/0001-8708(86)90101-5).
- [Lan76] R. P. LANGLANDS, *On the functional equations satisfied by Eisenstein series*, *Lecture Notes in Math.* **544**, Springer-Verlag, New York, 1976. MR 0579181. Zbl 0332.10018.
- [LV07] E. LINDENSTRAUSS and A. VENKATESH, Existence and Weyl’s law for spherical cusp forms, *Geom. Funct. Anal.* **17** no. 1 (2007), 220–251. MR 2306657. Zbl 1137.22011. <https://doi.org/10.1007/s00039-006-0589-0>.
- [Liu14] Y. LIU, Relative trace formulae toward Bessel and Fourier-Jacobi periods on unitary groups, *Manuscripta Math.* **145** no. 1-2 (2014), 1–69. MR 3244725. Zbl 1301.11050. <https://doi.org/10.1007/s00229-014-0666-x>.

- [LTX⁺19] Y. LIU, Y. TIAN, L. XIAO, W. ZHANG, and X. ZHU, On the Beilinson–Bloch–Kato conjecture for Rankin–Selberg motives, 2019. [arXiv 1912.11942](#).
- [MW95] C. MœGLIN and J.-L. WALDSPURGER, *Spectral Decomposition and Eisenstein Series*, *Cambridge Tracts in Math.* **113**, Cambridge Univ. Press, Cambridge, 1995, Une paraphrase de l'Écriture [A paraphrase of Scripture]. [MR 1361168](#). [Zbl 0846.11032](#). <https://doi.org/10.1017/CBO9780511470905>.
- [Mok15] C. P. MOK, Endoscopic classification of representations of quasi-split unitary groups, *Mem. Amer. Math. Soc.* **235** no. 1108 (2015), vi+248. [MR 3338302](#). [Zbl 1316.22018](#). <https://doi.org/10.1090/memo/1108>.
- [Mül89] W. MÜLLER, The trace class conjecture in the theory of automorphic forms, *Ann. of Math.* (2) **130** no. 3 (1989), 473–529. [MR 1025165](#). [Zbl 0701.11019](#). <https://doi.org/10.2307/1971453>.
- [PS79] I. I. PIATETSKI-SHAPIRO, Multiplicity one theorems, in *Automorphic Forms, Representations and L-Functions, Part 1* (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), *Proc. Sympos. Pure Math.*, XXXIII, Amer. Math. Soc., Providence, R.I., 1979, pp. 209–212. [MR 0546599](#). [Zbl 0423.22017](#).
- [PS82] I. I. PIATETSKI-SHAPIRO, Number theory related to Fermat's last theorem, in *Proceedings of the Conference held in Cambridge, Mass., September 1981* (N. KOBLITZ, ed.), *Progr. Math.* **26**, Birkhäuser, Boston, Mass., 1982, pp. 143–151. [MR 0685284](#).
- [Ram18] D. RAMAKRISHNAN, A theorem on $GL(n)$ à la Tchebotarev, 2018. [arXiv 1806.08429](#).
- [RSZ20] M. RAPOPORT, B. SMITHLING, and W. ZHANG, Arithmetic diagonal cycles on unitary Shimura varieties, *Compos. Math.* **156** no. 9 (2020), 1745–1824. [MR 4167594](#). [Zbl 1456.14031](#). <https://doi.org/10.1112/s0010437x20007289>.
- [Sak18] Y. SAKELLARIDIS, Transfer operators and Hankel transforms between relative trace formulas, I: character theory, 2018, *Adv. Math.*, to appear, 58 pp. [arXiv 1804.02383](#).
- [Trè67] F. TRÈVES, *Topological Vector Spaces, Distributions and Kernels*, *Pure Appl. Math. (Amst.)* **25**, Academic Press, Inc., New York, 1967. [MR 0225131](#). [Zbl 0171.10402](#).
- [Wal83] N. R. WALLACH, Asymptotic expansions of generalized matrix entries of representations of real reductive groups, in *Lie Group Representations, I* (College Park, Md., 1982/1983), *Lecture Notes in Math.* **1024**, Springer, Berlin, 1983, pp. 287–369. [MR 0727854](#). [Zbl 0553.22005](#). <https://doi.org/10.1007/BFb0071436>.
- [Wal92] N. R. WALLACH, *Real Reductive Groups. II*, *Pure Appl. Math.* **132**, Academic Press, Inc., Boston, MA, 1992. [MR 1170566](#). [Zbl 0785.22001](#).

- [Xue14] H. XUE, The Gan-Gross-Prasad conjecture for $U(n) \times U(n)$, *Adv. Math.* **262** (2014), 1130–1191. [MR 3228451](#). [Zbl 1301.11051](#). <https://doi.org/10.1016/j.aim.2014.06.010>.
- [Xue16] H. XUE, Fourier-Jacobi periods and the central value of Rankin-Selberg L -functions, *Israel J. Math.* **212** no. 2 (2016), 547–633. [MR 3505397](#). [Zbl 1347.11046](#). <https://doi.org/10.1007/s11856-016-1300-2>.
- [Xue19] H. XUE, On the global Gan-Gross-Prasad conjecture for unitary groups: approximating smooth transfer of Jacquet-Rallis, *J. Reine Angew. Math.* **756** (2019), 65–100. [MR 4026449](#). [Zbl 07126236](#). <https://doi.org/10.1515/crelle-2017-0016>.
- [Yun11] Z. YUN, The fundamental lemma of Jacquet and Rallis, *Duke Math. J.* **156** no. 2 (2011), 167–227, with an appendix by Julia Gordon. [MR 2769216](#). [Zbl 1211.14039](#). <https://doi.org/10.1215/00127094-2010-210>.
- [YZ17] Z. YUN and W. ZHANG, Shtukas and the Taylor expansion of L -functions, *Ann. of Math. (2)* **186** no. 3 (2017), 767–911. [MR 3702678](#). [Zbl 1385.11032](#). <https://doi.org/10.4007/annals.2017.186.3.2>.
- [Zha14a] W. ZHANG, Automorphic period and the central value of Rankin-Selberg L -function, *J. Amer. Math. Soc.* **27** no. 2 (2014), 541–612. [MR 3164988](#). [Zbl 1294.11069](#). <https://doi.org/10.1090/S0894-0347-2014-00784-0>.
- [Zha14b] W. ZHANG, Fourier transform and the global Gan-Gross-Prasad conjecture for unitary groups, *Ann. of Math. (2)* **180** no. 3 (2014), 971–1049. [MR 3245011](#). [Zbl 1322.11048](#). <https://doi.org/10.4007/annals.2014.180.3.4>.
- [Zyd20] M. ZYDOR, Les formules des traces relatives de Jacquet-Rallis grossières, *J. Reine Angew. Math.* **762** (2020), 195–259. [MR 4195660](#). [Zbl 07201152](#). <https://doi.org/10.1515/crelle-2018-0027>.

(Received: January 17, 2020)

(Revised: May 17, 2021)

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ D'AIX-MARSEILLE,
CNRS, I2M, MARSEILLE, FRANCE
E-mail: raphael.beuzart-plessis@univ-amu.fr

INSTITUTE FOR ADVANCED STUDY IN MATHEMATICS, ZHEJIANG UNIVERSITY,
HANGZHOU, CHINA
E-mail: liuyf0719@zju.edu.cn

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY,
CAMBRIDGE MA, UNITED STATES
E-mail: weizhang@mit.edu

DIVISION OF PHYSICS, MATHEMATICS AND ASTRONOMY, CALIFORNIA INSTITUTE
OF TECHNOLOGY, PASADENA CA, UNITED STATES
E-mail: xzhu@caltech.edu