

Smooth factors of projective actions of higher-rank lattices and rigidity

ALEXANDER GORODNIK
RALF SPATZIER

We study smooth factors of the standard actions of lattices in higher-rank semisimple Lie groups on flag manifolds. Under a mild condition on the existence of a single differentiable sink, we show that these factors are C^∞ –conjugate to the standard actions on flag manifolds.

37C15, 37C85

1 Introduction

Let Γ be a lattice in a connected semisimple Lie group G , and let P be a parabolic subgroup of G . We will be interested in the action of Γ on the flag manifold $F = G/P$ by left translations. The simplest example is given by the linear action of a lattice in $\mathrm{SL}_n(\mathbb{R})$ on the projective space \mathbb{P}^{n-1} . These actions have played a major role in rigidity theory. In particular, understanding their dynamics proved crucial in Margulis’s superrigidity and finiteness theorems. Margulis [18] classified all measurable factors of the Γ –action on $F = G/P$ when G has real rank at least two and the lattice Γ is irreducible. He showed that every such factor is measurably isomorphic to the Γ –action on $F' = G/Q$, where Q is a parabolic subgroup containing P . This was one of the ingredients in his proof of Margulis’s finiteness theorem, which shows that all normal subgroups of Γ are either of finite index or central. Dani [4] analysed topological factors of these actions when G has no compact or real rank-one factors. He proved that any Hausdorff factor of the action of Γ on $F = G/P$ is C^0 –conjugate to the action of Γ on $F' = G/Q$, where Q is a parabolic subgroup containing P . The aim of this paper is to establish a smooth analogue of Margulis’s and Dani’s factor theorems. Our results also complement recent work by Brown, Rodriguez Hertz and Wang [3] on low-dimensional actions of higher-rank lattices, and provide partial solutions and further evidence for their Conjecture 1.8.

The actions on flag manifolds are very different from measure-preserving actions, and one of their essential features is existence of sinks. Since analysis of dynamics in neighbourhoods of sinks will play a central role in our discussion, we give precise definitions.

Definition 1.1 Let p be a fixed point of a C^1 -map f on a manifold M .

- (i) The point p is called a *topological sink* if there exists a neighbourhood W_0 of p such that for every neighbourhood W of p and all sufficiently large n , we have $f^n(W_0) \subset W$.
- (ii) The point p is called a *differentiable sink* if all eigenvalues of $D(f)_p$ have modulus less than one, or equivalently, there exists a Riemannian metric in a neighbourhood U of p such that $\|D(f)_x\| < 1$ for all $x \in U$.

It is easy to see that a differentiable sink is always a topological sink, while the converse fails. In the case of actions on flag manifolds $F = G/P$, every \mathbb{R} -regular element $g \in G$ has a unique differentiable sink. We recall that an element $g \in G$ is called \mathbb{R} -*regular* if the number of eigenvalues of $\text{Ad}(g)$ of modulus one, counted with multiplicity, is as minimal as possible.

Let Γ be a group. Given Γ -actions on two spaces X and Y , we call a surjective map $\psi: X \rightarrow Y$ a *semiconjugacy* if ψ is Γ -equivariant. Then the main result of the paper is:

Theorem 1.2 Let G be a connected semisimple Lie group of \mathbb{R} -rank at least 2 and without compact factors. Let P be a parabolic subgroup of G with $F = G/P$ the corresponding flag manifold. Let Γ be an irreducible lattice in G . We denote by ρ_0 the standard action of Γ on F . Let ρ be a C^∞ -action of Γ on a manifold M such that for some $\gamma \in \Gamma$, the transformation $\rho(\gamma)$ has a differentiable sink in M . Suppose $\psi: F \rightarrow M$ is a C^0 -semiconjugacy between ρ_0 and ρ . Then there exist a parabolic subgroup Q containing P and a C^∞ -smooth Γ -equivariant diffeomorphism $\phi: G/Q \rightarrow M$ such that $\phi = \psi \circ \pi$, where $\pi: G/P \rightarrow G/Q$ is the canonical factor map:

$$\begin{array}{ccc} F = G/P & & \\ \psi \downarrow & \searrow \pi & \\ M & \xleftarrow[\phi]{} & G/Q \end{array}$$

It is clear that any C^0 factor of a standard action on a flag manifold has topological sinks as the original action does. However, we don't know whether the existence of a differentiable sink is automatic for C^0 factors of standard actions. There are examples of smooth lattice actions on S^1 -bundles over flag manifolds which have topological sinks that are not differentiable sinks. (see Section 7 below).

The assumption regarding existence of a differentiable sink in Theorem 1.2 is similar to the hyperbolicity assumptions that appeared in previous works on rigidity in the setting of measure-preserving actions. Our main result could be considered as an analogue of the smooth rigidity theorems for higher-rank Anosov actions; see Fisher, Kalinin and Spatzier [7], Katok and Spatzier [16] and Rodriguez Hertz and Wang [24]. There again one proves regularity of a C^0 -conjugacy under suitable uniform hyperbolicity hypotheses.

As an application of our main result, we also get local rigidity results. Given a C^∞ -action ρ_0 of a finitely generated group Γ on a compact manifold M , we call the action ρ_0 C^1 -locally rigid if any other C^∞ -action ρ of Γ on M is C^∞ -conjugate to ρ_0 provided that, for a finite set of generators S of Γ , the maps $\rho(s)$ and $\rho_0(s)$ are sufficiently close in the C^1 -topology for all $s \in S$. There is by now a long history of local rigidity results for higher-rank actions and, in particular, higher-rank lattices (see Fisher and Margulis [8], Ghys [10], Kanai [15] and Katok and Spatzier [16] amongst others and Fisher [6] for a survey). The following result was already obtained by M Kanai [15] under a more stringent closeness condition and by A Katok and R Spatzier [16].

Corollary 1.3 *Let G and P be as above. Let Γ be a uniform lattice in G . Then the Γ -action on $F = G/P$ is locally C^1 -rigid.*

We obtain this as an almost-immediate corollary of our main theorem. In fact, following Ghys [10], if Γ is cocompact, Katok and Spatzier [16] constructed a C^1 -close perturbation of the action of a Cartan subgroup A on G/Γ by left translations that corresponds to the perturbation of the Γ -action on G/P . This action is Anosov, and thus structurally stable. This in turn gives a C^0 -conjugacy between the action of Γ on the flag manifold F and its perturbation. Hence, now Corollary 1.3 follows immediately from Theorem 1.2. In a sense our argument is dual to the argument in [16].

We do not know if local rigidity holds for actions of nonuniform lattices on flag manifolds. Our argument shows that it suffices to prove structural stability for such actions.

Our results fail for cocompact lattices Γ in $\mathrm{PSL}_2(\mathbb{R})$. Indeed, the moduli space of discrete faithful cocompact representations of Γ in $\mathrm{PSL}_2(\mathbb{R})$ is nontrivial (in fact, it has positive dimension by standard results of Teichmüller theory). Let us take two representations ρ_1 and ρ_2 that are not equivalent, and consider the resulting extension of the natural actions by isometries on the hyperbolic plane \mathcal{H}^2 . The boundary circle $\partial\mathcal{H}^2$ of \mathcal{H}^2 is naturally identified with one-dimensional projective space: $\mathbb{R}P^1 \simeq \mathrm{PSL}_2(\mathbb{R})/P$, where P denotes the parabolic subgroup of $\mathrm{PSL}_2(\mathbb{R})$ consisting of upper triangular matrices. Both ρ_1 and ρ_2 define C^∞ -actions of Γ on $\partial\mathcal{H}^2$. Since $\rho_1(\Gamma)$ and $\rho_2(\Gamma)$ are quasi-isometric, their actions on the boundary are C^0 -conjugate. The conjugating homeomorphism however cannot be differentiable even at a single point, by well-known results of Ivanov [14] and Tukia [26]. Similar constructions can be made for Zariski-dense groups in higher dimension, for example, for convex cocompact groups via quasiconformal deformations in 3-dimensional hyperbolic space. Typically, one has a large moduli space of representations for Zariski-dense groups, and we expect that the above considerations generalise to give counterexamples for actions of Zariski-dense subgroups on flag manifolds. It is not clear to us whether lattices in other real rank-one groups have more rigidity. We remark that both Margulis's and Dani's theorem fail in the real rank-one case (see Margulis [18], Spatzier [25] and Zimmer [27]).

Organisation of the paper

In the next section we recall basic properties of the standard actions on flag manifolds F . Next, in Section 3 we investigate general smooth actions on manifold M which are continuously conjugate to the standard actions and establish existence of many sinks in M . In Section 4, we analyse properties of the conjugacy map $F \rightarrow M$ further and introduce projection maps to certain dynamically defined submanifolds of a sink. These maps are defined on open subsets of F and M and are intertwined by the conjugacy $F \rightarrow M$. Then in Section 5 we establish smoothness of the map $F \rightarrow M$ along a family of foliations. Finally, in Section 6 we complete the proof combining results from Sections 4 and 5. In Section 7, we give an example of a lattice action with a topological sink which is not a differentiable sink.

Acknowledgements We are grateful to David Fisher and Gopal Prasad for many pertinent discussions related to this paper. In addition we thank Tasho Kaletha and Loren Spice for discussions on Lie theory. We also thank the referee for a careful

reading of the text, and the suggestion to include irreducible lattices in products with rank-one factors in Theorem 1.2.

Gorodnik would like to thank the University of Michigan for hospitality during visits while work on this project proceeded. Spatzier thanks the University of Bristol for hospitality and support during this work. Both authors also benefited from the special semester *Dynamics on moduli spaces of geometric structures* at MSRI, and thank the Institute for its support and hospitality.

Gorodnik was supported in part by EPSRC grant EP/H000091/1, ERC grant 239606 and MSRI. Spatzier was supported in part by NSF grant DMS-1307164 and MSRI.

2 Actions on flag manifolds

We start by recalling the definition of flag manifolds and discuss basic properties of the standard actions on flag manifolds. We also discuss properties of dynamics for actions on projective spaces and existence/uniqueness of sinks. Throughout this section, G is a connected semisimple Lie group without any assumptions on its rank. Without loss of generality, we may and will assume that the centre of G is trivial. In particular, G is a real algebraic group. Indeed the centre of G will act trivially on any of the flag varieties in question.

2.1 Flag manifolds

We fix a Cartan involution θ of G . It determines the maximal compact subgroup K of G and the Cartan decomposition

$$(2-1) \quad \mathfrak{g} = \text{Lie}(K) \oplus \mathfrak{p}$$

of the Lie algebra \mathfrak{g} of G . Let $\mathfrak{a} \subset \mathfrak{p}$ be a Cartan subalgebra (ie a maximal abelian subalgebra of \mathfrak{p}). A nonzero linear form $\alpha \in \mathfrak{a}^*$ is a (restricted) root if the corresponding root space

$$\mathfrak{g}_\alpha = \{x \in \mathfrak{g} : [a, x] = \alpha(a)x \text{ for } a \in \mathfrak{a}\}$$

is nonzero. We denote by $\Phi \subset \mathfrak{a}^*$ the set of (restricted) roots. Then

$$\mathfrak{g} = \mathfrak{g}_0 \oplus \sum_{\alpha \in \Phi} \mathfrak{g}_\alpha \quad \text{and} \quad \mathfrak{g}_0 = \mathfrak{m} \oplus \mathfrak{a},$$

where $\mathfrak{m} = \text{Lie}(K) \cap \mathfrak{g}_0$. We fix a set $\Delta \subset \Phi$ of simple roots and denote by Φ^+ the corresponding subset of positive roots.

We introduce the set of standard parabolic subgroups P_I of G which are associated to subsets $I \subset \Delta$. We denote by $\Phi^I \subset \Phi$ the subset of roots that are linear combinations of elements from I . The standard parabolic subalgebra is defined by

$$\mathfrak{p}_I = \mathfrak{m}_I + \mathfrak{a} + \mathfrak{n}_I,$$

where

$$\mathfrak{m}_I = \mathfrak{m} + \mathfrak{a} + \sum_{\alpha \in \Phi^I} \mathfrak{g}_\alpha \quad \text{and} \quad \mathfrak{n}_I = \sum_{\alpha \in \Phi^+ \setminus \Phi^I} \mathfrak{g}_\alpha.$$

We also set

$$\mathfrak{n}_I^- = \sum_{\alpha \in \Phi^- \setminus \Phi^I} \mathfrak{g}_\alpha.$$

Then

$$\mathfrak{g} = \mathfrak{n}_I^- \oplus \mathfrak{p}_I.$$

The standard parabolic subgroup P_I is defined as the normaliser of \mathfrak{p}_I in \mathfrak{g} . A general parabolic subgroup is a subgroup of G which is conjugate to one of the standard parabolic subgroups P_I . The flag manifolds are the homogeneous spaces

$$F_I = G/P_I.$$

Let

$$N_I^- = \exp(\mathfrak{n}_I^-).$$

Then it follows from the Bruhat decomposition that

$$U_I = N_I^- P_I$$

is an open dense subset of F_I , and, moreover, the complement of $N_I^- P_I$ in F_I is a finite union of analytic submanifolds of lower dimensions.

Let us describe how typical elements $g \in G$ act on the flag manifolds F_I . Every $g \in G$ can be written uniquely as a commuting product

$$(2-2) \quad g = g_c g_{nc} g_u,$$

where $\text{Ad}(g_c)$ is semisimple and has all eigenvalues of modulus one, $\text{Ad}(g_{nc})$ is semisimple and has all eigenvalues real and positive, and $\text{Ad}(g_u)$ is unipotent. Note that the actions of $g \in G$ and its conjugate hgh^{-1} on F_I are smoothly conjugate. Thus we can analyse the action of g using that of hgh^{-1} on F_I . Hence, replacing g by a conjugate of g , we may assume that $g_{nc} = a \in \exp(\mathfrak{a}^+)$, where

$$\mathfrak{a}^+ = \{a \in \mathfrak{a} : \alpha(a) \geq 0 \text{ for all } \alpha \in \Delta\}$$

is the positive Weyl chamber in \mathfrak{a} . Then the action of g on the open cell $U_I \subset F_I$ can be described as follows:

$$(2-3) \quad g \cdot \exp(x) P_I = \exp\left(\sum_{\alpha \in \Phi^- \setminus \Phi^I} e^{\alpha(\log(a))} \text{Ad}(g_c g_u) x_\alpha\right) P_I$$

for $x = \sum_{\alpha \in \Phi^- \setminus \Phi^I} x_\alpha \in \mathfrak{n}_I^-$.

We recall that the element g is called \mathbb{R} –regular if the number of eigenvalues of $\text{Ad}(g)$ of modulus one, counted with multiplicity, is as minimal as possible. This condition is equivalent to a being in the interior of \mathfrak{a}^+ . We also use the following more general notion of regularity.

Definition 2.1 For $J \subset \Delta$, an element $g \in G$ is called J –regular if it is of the form (2-2) with g_{nc} being conjugated to $a \in \exp(\mathfrak{a}^+)$ such that $\alpha(\log(a)) > 0$ for all $\alpha \in J$. In particular, Δ –regular elements are precisely the \mathbb{R} –regular elements.

Now suppose that the element g in (2-3) is $(\Delta \setminus I)$ –regular. Then $\alpha(\log(a)) < 0$ for all $\alpha \in \Phi^- \setminus \Phi^I$. We observe that the map $\text{Ad}(g_c g_u)$ preserves the root spaces and has eigenvalues of absolute value one, and $\|\text{Ad}(g_c g_u)^n\|$ grows at most polynomially as $n \rightarrow \infty$. Hence, it follows that the identity coset eP_I is a differentiable sink for g , and $g^n z \rightarrow eP_I$ as $n \rightarrow \infty$ for every $z \in U_I$. Since U_I is dense, it also clear that this sink is unique. We denote by $s_g \in F_I$ the sink of the element g .

The above discussion shows that every $(\Delta \setminus I)$ –regular element has a sink in F_I . As we shall see in the next section, the converse is also true: if an element has a topological sink in F_I , then it is $(\Delta \setminus I)$ –regular.

2.2 Dynamics on projective spaces

It is convenient to study the action of G on the flag manifolds F_I by embedding F_I in a product of suitable projective spaces, endowed with the sup metric. Thus we can reduce metric and topological questions about F_I to their natural analogues in the various projective spaces. More precisely, we consider the G –equivariant embedding

$$(2-4) \quad \iota_I: F_I \rightarrow \prod_{\alpha \in \Delta \setminus I} \mathbb{P}(V_\alpha), \quad gP_I \mapsto (\sigma_\alpha(g)v_\alpha : \alpha \in \Delta \setminus I),$$

introduced in [1, Section 3], which is defined using suitable irreducible representations

$$\sigma_\alpha: G \rightarrow \text{GL}(V_\alpha), \quad \alpha \in \Delta,$$

and the highest-weight vectors $v_\alpha \in V_\alpha$. These representations have the property that the transformation $\sigma_\alpha(g)$ is proximal if and only if g is $\{\alpha\}$ -regular (see [1, Section 2.5]). We recall that a linear transformation is called *proximal* if it has a unique eigenvalue of maximal modulus, and this eigenvalue has multiplicity one.

Proposition 2.2 *Suppose that the action of an element $g \in G$ on the flag manifold F_I has a topological sink. Then g is $(\Delta \setminus I)$ -regular. In particular, every topological sink for the standard action on F_I is also a differentiable sink, and this sink is unique.*

Proof We prove the proposition by considering the action on $\iota_I(F_I)$. If the action of $(\sigma_\alpha(g))_{\alpha \in \Delta \setminus I}$ on $\iota_I(F_I)$ has a topological sink, then the actions of $\sigma_\alpha(g)$ on $\iota_\alpha(F_I)$ for $\alpha \in \Delta \setminus I$ also have topological sinks. We will show that then the $\sigma_\alpha(g)$ are proximal. In view of the above remark, this will imply the proposition.

We write the transformation $\sigma_\alpha(g)$ as a commuting product

$$\sigma_\alpha(g) = kau,$$

where k is semisimple with eigenvalues of absolute value one, a is semisimple with real positive eigenvalues, and u is unipotent. Let $s_\alpha \in \iota_\alpha(F_I)$ be a topological sink of $\sigma_\alpha(g)$. We fix a $\sigma_\alpha(K)$ -invariant metric on $\mathbb{P}(V_\alpha)$ and denote by $B_\epsilon(s_\alpha)$ the ϵ -ball centred at s_α in $\iota_\alpha(F_I)$. Then, since s_α is a sink, there exists $\epsilon_0 > 0$ such that for every $\epsilon > 0$ and all sufficiently large n , we have

$$g^n(B_{\epsilon_0}(s_\alpha)) \subset B_\epsilon(s_\alpha),$$

and hence

$$(au)^n(B_{\epsilon_0}(s_\alpha)) \subset B_\epsilon(k^{-n}s_\alpha).$$

Passing to subsequence, we may assume that $k^{-n_i}s_\alpha \rightarrow s_\alpha$. Hence, it follows that for every $\epsilon > 0$ and all sufficiently large i , we have

$$(2-5) \quad (au)^{n_i}(B_{\epsilon_0}(s_\alpha)) \subset B_{2\epsilon}(s_\alpha).$$

In this case, we say that s_α is a topological sink for the sequence of transformations $(au)^{n_i}$.

The transformation au has real positive eigenvalues. Let λ be the maximal eigenvalue of this transformation, and let $V_\alpha(\lambda)$ be the corresponding Jordan subspace. We denote by $\pi_\lambda: V_\alpha \rightarrow V_\alpha(\lambda)$ the projection map defined by the Jordan decomposition of au . We claim that $\pi_\lambda(s_\alpha) \neq 0$. Indeed, suppose that $\pi_\lambda(s_\alpha) = 0$. It follows from (2-5) that

for a sufficiently small neighbourhood \mathcal{O} of the identity in G and all $v \in \sigma_\alpha(\mathcal{O})s_\alpha$, we have

$$(au)^{n_i}v \rightarrow s_\alpha.$$

On the other hand, if $\pi_\lambda(v) \neq 0$, then, since λ is maximal, $(au)^{n_i}v$ must converge to a point in $\mathbb{P}(V_\alpha(\lambda))$, which contradicts our assumption that $\pi_\lambda(s_\alpha) = 0$. Hence, we conclude that

$$\sigma_\alpha(\mathcal{O})s_\alpha \subset \ker(\pi_\lambda).$$

Since $\iota_\alpha(F_I) = \sigma_\alpha(G)s_\alpha$ is an analytic submanifold of $\mathbb{P}(V_\alpha)$, this implies that

$$\iota_\alpha(F_I) \subset \ker(\pi_\lambda).$$

However, this contradicts irreducibility of the representation σ_α . Hence, we conclude that $\pi_\lambda(s_\alpha) \neq 0$.

The transformation au acts on $\mathbb{P}(V_\alpha(\lambda))$ as u . We can write $u = \exp(X)$ for some nilpotent $X \in \text{End}(V_\alpha)$ preserving the Jordan decomposition. Then u is contained in a unipotent one-parameter subgroup $U = \{\exp(tX)\}_{t \in \mathbb{R}}$ of $\sigma_\alpha(G)$. The projection map $\pi_\lambda: V_\alpha \rightarrow V_\alpha(\lambda)$ is equivariant with respect to the action of U . We consider the action of U on $S = \pi_\lambda(\iota_\alpha(F_I))$. Then the point $s = \pi_\lambda(s_\alpha)$ is a topological sink for the sequence u^{n_i} . It follows from Lemma 2.3 below that $S = \{s\}$. If $\dim(V_\alpha(\lambda)) > 1$, then it would follow that $\iota_\alpha(F_I)$ was contained in a proper subspace of V_α , but this contradicts irreducibility of the representation σ_α . Hence, we conclude that $\dim(V_\alpha(\lambda)) = 1$, and $\sigma_\alpha(g)$ is proximal. This implies that g is α -regular for all $\alpha \in \Delta \setminus I$, and completes the proof. \square

The following lemma was used in the proof of the previous proposition.

Lemma 2.3 *Let $U = \{u_t\}_{t \in \mathbb{R}}$ be a one-parameter unipotent group of linear transformations of a vector space V , $S \subset \mathbb{P}(V)$ a U -invariant connected analytic submanifold, and $s \in S$ a topological sink for the restriction to S of a sequence u_{n_i} satisfying $n_i \rightarrow \infty$. Then $S = \{s\}$.*

Proof Since s is a topological sink for the sequence u_{n_i} , there exists a neighbourhood W_0 of s in S such that for every neighbourhood W of s in S and all sufficiently large i ,

$$(2-6) \quad u_{n_i}(W_0) \subset W.$$

Without loss of generality, we may assume that $V = \langle S \rangle$. Moreover, since S is an analytic submanifold, it follows that $V = \langle W_0 \rangle$. We write $u_t = \exp(tX)$ for a nilpotent transformation $X \in \text{End}(V)$. We suppose that $X \neq 0$ and take $l \geq 1$ such that $X^l \neq 0$ and $X^{l+1} = 0$. Then

$$u_t = \sum_{j=0}^l \frac{t^j X^j}{j!}.$$

There exists $w \in W_0$ such that $X^l w \neq 0$. Then

$$u_t w \rightarrow [X^l w] \quad \text{in } \mathbb{P}(V) \text{ as } t \rightarrow -\infty \text{ and as } t \rightarrow +\infty.$$

In particular, it follows that

$$(2-7) \quad s = [X^l w].$$

It also follows that there exists $t_0 \in \mathbb{R}$ such that for all $t < t_0$, we have

$$u_t w \in W_0.$$

Then we deduce from (2-6) that for every neighbourhood W of s in S and all sufficiently large i ,

$$u_{n_i} \cdot u_t w = u_{n_i+t} w \in W.$$

Then, taking $t = -n_i$, we conclude that $w = s$. However, it follows from (2-7) that $Xs = 0$, so that $w \neq s$. This contradiction implies that $X = 0$ (that is, U is trivial), and since s is a sink, $S = \{s\}$. \square

3 Existence of many sinks

In this section we establish an abundance of differentiable sinks for arbitrary smooth actions which are C^0 –conjugate to the standard actions on flag manifolds. We only require existence of a single differentiable sink.

The following proposition will play a central role in the proof of our main result. It might have other applications, and we emphasise that this result is applicable to any Zariski-dense subgroup of a semisimple group without any rank assumptions.

Proposition 3.1 *Let G be a semisimple real algebraic group, $F_I = G/P_I$ a flag manifold, and $\Gamma \subset G$ a Zariski-dense subgroup of G . We denote by ρ_0 the standard action of Γ on F_I . Let ρ be a C^1 –action of Γ on a manifold M , and $\phi: F_I \rightarrow M$ is*

a C^0 –conjugacy intertwining the actions ρ_0 and ρ . Suppose that there exists $\gamma_0 \in \Gamma$ such that $\rho(\gamma_0)$ has a differentiable sink in M . Then there exists a Zariski-dense subsemigroup $S \subset \Gamma$ such that $\rho(\gamma)$ has a differentiable sink in M for every $\gamma \in S$.

We will need a quantitative version of the proximal property discussed in Section 2.2. Given a proximal linear transformation $g: V \rightarrow V$ of a vector space V , we denote by $s_g \in \mathbb{P}(V)$ the direction corresponding to the maximal eigenvalue, and by $X_g^< \subset \mathbb{P}(V)$ the set of directions corresponding to the complementary g –invariant subspace. We fix standard metrics on the projective spaces $\mathbb{P}(V)$ and set

$$b_g^\epsilon = \{x \in \mathbb{P}(V) : d(x, s_g) \leq \epsilon\} \quad \text{and} \quad B_g^\epsilon = \{x \in \mathbb{P}(V) : d(x, X_g^<) \geq \epsilon\}.$$

Definition 3.2 We call a proximal transformation $g: V \rightarrow V$ (r, ϵ) –proximal if

$$d(s_g, X_g^<) \geq r, \quad g(B_g^\epsilon) \subset b_g^\epsilon, \quad g|_{B_g^\epsilon} \text{ is } \epsilon\text{--Lipschitz.}$$

This definition is a slight variation of the notion introduced in [1, Section 2-3]. Adapting it to our setting, we say that:

Definition 3.3 For $J \subset \Delta$, an element $g \in G$ is called (J, r, ϵ) –regular if the linear transformations $\sigma_\alpha(g)$ are (r, ϵ) –proximal for all $\alpha \in J$.

We use the following lemma [1, Section 3.6]. We fix standard metrics on the projective spaces $\mathbb{P}(V_\alpha)$, which also define a metric on the flag manifolds F_I via the embedding (2-4).

Lemma 3.4 (Benoist) Let $\gamma_0 \in \Gamma$ be a $(\Delta \setminus I)$ –regular element with the sink $s_{\gamma_0} \in F_I$. Then for every sufficiently small $r, \epsilon > 0$, the set

$$\mathcal{G}(\gamma_0, r, \epsilon) = \{\delta \in \Gamma : \delta \text{ is } (\Delta \setminus I, r, \epsilon)\text{--regular and } d(s_\delta, s_{\gamma_0}) \leq \epsilon\}$$

is Zariski-dense in G .

This lemma is proved in [1] for $r = 2\epsilon$, but the same argument allows one to treat (r, ϵ) –proximal elements as well.

Proof of Proposition 3.1 Throughout the proof, to simplify notation, for $\gamma \in \Gamma$ and $x \in F_I$, we write $\rho_0(\gamma)x = \gamma x$ for the standard action Γ on F_I .

Let $m_0 \in M$ be a differentiable sink for $\rho(\gamma_0)$. We fix a Riemannian metric on M such that

$$(3-1) \quad \|D(\rho(\gamma_0))_{m_0}\| < 1.$$

Since ϕ is a homeomorphism, it follows that $\phi^{-1}(m_0)$ is a topological sink for γ_0 . Hence, it follows from Proposition 2.2 that γ_0 is $(\Delta \setminus I)$ -regular, and $\phi^{-1}(m_0) = s_{\gamma_0}$ is the differentiable sink for γ_0 .

Since γ_0 is $(\Delta \setminus I)$ -regular, there exists $r_0 = r_0(\gamma_0) > 0$ and $n_0(\lambda) > 0$ such that for every $\lambda > 0$ and every $n \geq n_0(\lambda)$, the element γ_0^n is $(\Delta \setminus I, r_0, \lambda)$ -regular. We fix r_0 as above and, moreover, assume that it is sufficiently small, so that the set $\mathcal{G}(\gamma_0, r_0, \epsilon)$ is Zariski-dense for all sufficiently small $\epsilon > 0$ (see Lemma 3.4).

Denote by $B_\epsilon(p)$ the ϵ -ball around p in F_I . Then we claim that for every $\kappa \in (0, 1)$ and $\delta \in \mathcal{G}(\gamma_0, r_0, \epsilon)$, there exists $c = c(\delta) > 0$ such that for $\epsilon \in (0, \frac{1}{3}r_0)$ as above and $n \geq n_0(\kappa, \delta)$, we have

$$(3-2) \quad \delta \gamma_0^n \delta^{-1} (B_{\epsilon/c}(\delta s_{\gamma_0})) \subset B_{\epsilon/c}(\delta s_{\gamma_0}),$$

$$(3-3) \quad \|D(\rho(\delta \gamma_0^n \delta^{-1}))_x\| \leq \kappa \quad \text{when } x \in \phi(B_{\epsilon/c}(\delta s_{\gamma_0})).$$

The quantity $n_0(\kappa, \delta)$ will be specified during the proof.

To prove (3-2), we choose $c = c(\delta) \geq 1$ so that

$$d(\delta^{-1}x, \delta^{-1}y) \leq cd(x, y) \quad \text{for all } x, y \in F_I.$$

This implies that

$$(3-4) \quad \delta^{-1} (B_{\epsilon/c}(\delta s_{\gamma_0})) \subset B_\epsilon(s_{\gamma_0}).$$

We take $\lambda = \min\{\frac{1}{c}, \frac{1}{2}r_0\}$. For every $n \geq n_0(\lambda)$, the element γ_0^n is $(\Delta \setminus I, r_0, \lambda)$ -regular. Then since $\epsilon < \frac{1}{3}r$, we have $r_0 - \epsilon \geq \lambda$, and

$$\iota_\alpha(B_\epsilon(s_{\gamma_0})) \subset B_{\sigma_\alpha(\gamma_0)}^\lambda \quad \text{for all } \alpha \in \Delta \setminus I.$$

Hence, since the transformations $\sigma_\alpha(\gamma_0^n)|_{B_{\sigma_\alpha(\gamma_0)}^\lambda}$ for $\alpha \in \Delta \setminus I$ are λ -Lipschitz, we conclude that

$$\gamma_0^n(B_\epsilon(s_{\gamma_0})) \subset B_{\lambda\epsilon}(s_{\gamma_0}) \subset B_{\epsilon/c}(s_{\gamma_0}).$$

Since $d(s_\delta, s_{\gamma_0}) \leq \epsilon$, we also have

$$B_{\epsilon/c}(s_{\gamma_0}) \subset B_{\epsilon/c+\epsilon}(s_\delta) \subset B_{2\epsilon}(s_\delta),$$

and since $\epsilon < \frac{1}{3}r_0$,

$$\iota_\alpha(B_{2\epsilon}(s_\delta)) \subset B_{\sigma_\alpha(\delta)}^\epsilon \quad \text{for all } \alpha \in \Delta \setminus I.$$

Using that the transformations $\sigma_\alpha(\delta)|_{B_{\sigma_\alpha(\delta)}^\epsilon}$ for $\alpha \in \Delta \setminus I$ are ϵ -Lipschitz, we deduce that

$$\delta(B_{\epsilon/c}(s_{\gamma_0})) \subset B_{\epsilon^2/c}(\delta s_{\gamma_0}),$$

which is contained in $B_{\epsilon/c}(\delta s_{\gamma_0})$ provided that $\epsilon \leq 1$. This completes the proof of (3-2).

Now we proceed with proving (3-3). It follows from (3-4) that

$$(3-5) \quad \rho(\delta^{-1})(\phi(B_{\epsilon/c}(\delta s_{\gamma_0}))) \subset \phi(B_\epsilon(s_{\gamma_0})).$$

We take $l_1 \geq 1$ such that $\gamma_0^{l_1}$ is $(\Delta \setminus I, r_0, \epsilon_0)$ -regular with $\epsilon_0 = \frac{1}{2} \min\{r_0, 1\}$. Then, since we have assumed that $\epsilon < \frac{1}{3}r_0$, we have $\epsilon_0 < r_0 - \epsilon$, and for all $\alpha \in \Delta \setminus I$,

$$\iota_\alpha(B_\epsilon(s_{\gamma_0})) \subset B_{\sigma_\alpha(\gamma_0)}^{\epsilon_0}.$$

This implies that

$$(3-6) \quad \gamma_0^{l_1}(B_\epsilon(s_{\gamma_0})) \subset B_{\epsilon_0}(s_{\gamma_0}).$$

Moreover, since the transformations $\sigma_\alpha(\gamma_0^{l_1})|_{B_{\sigma_\alpha(\gamma_0)}^{\epsilon_0}}$ for $\alpha \in \Delta \setminus I$ are ϵ_0 -Lipschitz,

$$(3-7) \quad \gamma_0^{l_1}(B_\theta(s_{\gamma_0})) \subset B_{\epsilon_0\theta}(s_{\gamma_0}) \quad \text{for every } \theta \in (0, \epsilon_0].$$

This implies that the sets $W_\theta = \phi(B_\theta(s_{\gamma_0}))$ give $\rho(\gamma_0^{l_1})$ -invariant neighbourhoods of m_0 for $\theta \in (0, \epsilon_0]$. We choose $\theta_0 \in (0, \epsilon_0]$ sufficiently small that

$$\eta = \sup_{x \in W_{\theta_0}} \|D(\rho(\gamma_0^{l_1}))_x\| < 1.$$

This is possible in view of (3-1). Then it follows from the chain rule that there exists $C > 0$ such that, for every $n \geq 1$,

$$(3-8) \quad \sup_{x \in W_{\theta_0}} \|D(\rho(\gamma_0^n))_x\| \leq C \eta^{\lfloor n/l_1 \rfloor}.$$

Using (3-7), we deduce that there exists $l_2 \geq 1$ such that

$$\gamma_0^{l_2}(B_{\epsilon_0}(s_{\gamma_0})) \subset B_{\theta_0}(s_{\gamma_0}).$$

Then it follows from (3-6) that

$$\gamma_0^{l_1+l_2}(B_\epsilon(s_{\gamma_0})) \subset B_{\theta_0}(s_{\gamma_0})$$

and

$$\rho(\gamma_0^{l_1+l_2})(\phi(B_\epsilon(s_{\gamma_0}))) \subset W_{\theta_0}.$$

Hence, using (3-5), we conclude that

$$\rho(\gamma_0^{l_1+l_2}\delta^{-1})(\phi(B_{\epsilon/c}(\delta s_{\gamma_0}))) \subset W_{\theta_0}.$$

Now the claim (3-3) follows the estimate (3-8) and the chain rule.

We consider the semigroup of the form

$$S = \langle \delta \gamma_0^n \delta^{-1} : \delta \in \mathcal{G}(\gamma_0, r_0, \epsilon), n \geq n_0(\kappa, \delta) \rangle.$$

It follows from property (3-2) that for every $\gamma \in S$, the transformation $\rho(\gamma)$ preserves the neighbourhood $U = \phi(B_{\epsilon/c}(\delta s_{\gamma_0}))$. Moreover, by (3-3),

$$\sup_{x \in U} \|D(\rho(\gamma))_x\| \leq \kappa.$$

When κ is sufficiently small, this implies that $\rho(\gamma)$ is a contraction with respect to the Riemannian metric on U . Hence, we conclude that the map $\rho(\gamma)$ has a fixed point in U which is a differentiable sink.

It remains to show that the semigroup S is Zariski-dense. Let \bar{S} be the Zariski closure of S . We denote by A_n the Zariski connected component of the Zariski closure of the cyclic group $\langle \gamma_0^n \rangle$. We note that since γ_0 has a sink, it must be of infinite order, so that A_n is not trivial for all n . Moreover, we may assume that the projections of γ_0 to all nontrivial simple factors of G also have infinite order. Indeed, suppose that for some nontrivial simple factor $G_i \subset G$, the G_i -component of γ_0 has finite order. Then, for some n , the transformation γ_0^n acts trivially on the submanifold $G_i s_{\gamma_0} \subset F_I$. Since s_{γ_0} is a sink for γ_0 , this implies that $G_i s_{\gamma_0} = s_{\gamma_0}$, and $G_i \subset P_I$. In this case, we can replace the group G by G/G_i . Hence, without loss of generality, the projections of γ_0 to all nontrivial simple factors of G also have infinite order.

We note that $A_m \supset A_n$ when m divides n and consider the descending sequence of Zariski-closed subgroups $B_n = A_{n!}$. For sufficiently large n , this sequence stabilises, and we denote the minimal element by B . For every $\delta \in \mathcal{G}(\gamma_0, r, \epsilon)$, we have $\delta B \delta^{-1} \subset \bar{S}$. Hence, it follows from Zariski-density of $\mathcal{G}(\gamma_0, r, \epsilon)$ that \bar{S} contains the conjugacy class B^G . Since S is a semigroup, its Zariski-closure \bar{S} is a group. We conclude that \bar{S} contains the normal subgroup generated by B . Since the projections of γ_0 to all nontrivial simple factors of G have infinite order, it follows that $\bar{S} = G$, so that S is Zariski-dense. \square

4 Projection maps

Let G be a connected semisimple Lie group, $F_I = G/P_I$ a flag manifold, and $\Gamma \subset G$ an irreducible lattice subgroup of G . We denote by ρ_0 the action of Γ on F_I . Let ρ be a smooth action of Γ on a compact manifold M . In this section we study properties of a C^0 -conjugacy map

$$\phi: F_I \rightarrow M$$

that intertwines the actions ρ_0 and ρ . We will consider an element γ with a differentiable sink in M . After a conjugation, we may and will assume that the sink for γ in F_I is the identity coset eP_I . The aim of this section is to construct a family of projections maps $\pi_0^{(\alpha)}: U_I \rightarrow U_I$ for simple roots α not in I , defined on the open cell $U_I = N_I^- P_I \subset F_I$ and the corresponding projection maps $\tilde{\pi}_0^{(\alpha)}$ for M , which are conjugated to $\pi_0^{(\alpha)}$ via ϕ . Note that we get similar projections for different $\gamma^* \in \Gamma$ with a differentiable sink. Naturally, the parabolic stabilising the sink of γ^* and the open cell will depend on γ^* .

Since the centre of G acts trivially on the flag manifold F_I , and hence on M thanks to the semiconjugacy Φ , we may assume without loss of generality that G is centre-free. It follows from the Margulis arithmeticity theorem (see [19, Chapter IX] or [28, Chapter 6]) that there exist a connected semisimple algebraic \mathbb{Q} -group \mathbf{G} and surjective homomorphism $\iota: \mathbf{G}(\mathbb{R})^\circ \rightarrow G$ such that $\ker(\iota)$ is compact and $\iota(\mathbf{G}(\mathbb{Z}) \cap \mathbf{G}(\mathbb{R})^\circ)$ is commensurable to Γ . We will make use of arithmetic properties of Γ which are easiest to state when the group G has a \mathbb{Q} -structure. Hence, now we assume that $G = \mathbf{G}(\mathbb{R})^\circ$ and Γ is a finite-index subgroup of $\mathbf{G}(\mathbb{Z}) \cap \mathbf{G}(\mathbb{R})^\circ$. This setup is a bit different from that of Theorem 1.2. Indeed, there we assume that G does not have compact factors, and explained how to interpolate between the cases with and without compact factors. Alternatively we could have allowed compact factors in Theorem 1.2, and considered only factors G/P where P contains all compact factors. Since the components of $\gamma \in \Gamma$ in any of the compact factors act trivially, this is equivalent to Theorem 1.2 as stated.

In order to have rich dynamics in a neighbourhood of a sink, we need to construct commuting elements satisfying certain independence properties. More precisely, these elements will be chosen in the centraliser $Z_\Gamma(\gamma_0)$, where γ_0 is picked from a given Zariski-dense subsemigroup S . Eventually, we apply this construction for the semigroup $S \subset \Gamma$ introduced in Proposition 3.1, but the discussion in the first part of this section applies to arbitrary Zariski-dense subsemigroups $S \subset \mathbf{G}(\mathbb{Q})$.

Our argument is based on the results established by Prasad and Rapinchuk [23]. We also refer to [22] for basic properties of regular and \mathbb{R} –regular elements. We start by introducing required notation. We denote by $Z_{\mathbf{G}}(g)^\circ$ the connected component of the centraliser of g in \mathbf{G} with respect to the Zariski topology. We recall that if $g \in G$ is a regular and \mathbb{R} –regular element, then $Z_{\mathbf{G}}(g)^\circ$ is a maximal torus in \mathbf{G} , and

$$(4-1) \quad Z_{\mathbf{G}}(g)^\circ = \mathbf{B}_g \mathbf{T}_g,$$

where \mathbf{B}_g is a torus such that $\mathbf{B}_g(\mathbb{R})$ is compact, and \mathbf{T}_g is a maximal \mathbb{R} –split torus in \mathbf{G} . We note that an \mathbb{R} –regular element is necessarily semisimple. In particular, it follows from [2, IV.11.12] that $g \in Z_{\mathbf{G}}(g)^\circ$. For a character χ of \mathbf{T}_g and $h = bt \in Z_{\mathbf{G}}(g)^\circ = \mathbf{B}\mathbf{T}_g$, we set $\chi(h) = \chi(t)$. Since $Z_{\mathbf{G}}(g)^\circ$ has finite index in $Z_{\mathbf{G}}(g)$, it follows that, for every $h \in Z_{\mathbf{G}}(g)$, $h^l \in Z_{\mathbf{G}}(g)^\circ$ for some $l \geq 1$. In that case, we set $\chi(h) = \chi(h^l)^{1/l}$. It is clear that this definition is independent of the choice of the exponent l .

We say that $g \in \mathbf{G}(\mathbb{Q})$ is *anisotropic* if the torus $Z_{\mathbf{G}}(g)^\circ$ is anisotropic over \mathbb{Q} .

The group \mathbf{G} has a decomposition as an almost direct product

$$(4-2) \quad \mathbf{G} = \mathbf{G}^{(1)} \cdots \mathbf{G}^{(r)},$$

where the $\mathbf{G}^{(i)}$ are the connected \mathbb{Q} –simple subgroups of \mathbf{G} . We say that an element $g \in \mathbf{G}$ is *without components of finite order* if with respect to this decomposition, $g = g_1 \cdots g_r$ with all g_i of infinite order. A maximal \mathbb{Q} –subtorus \mathbf{T} of \mathbf{G} is called \mathbb{Q} –*quasi-irreducible* if it does not contain any \mathbb{Q} –subtori other than almost direct products of the tori $\mathbf{T}^{(i)} = \mathbf{T} \cap \mathbf{G}^{(i)}$.

We say that commuting elements $\delta_1, \delta_2 \in \mathbf{G}(\mathbb{Q})$ are *multiplicatively independent* if the projections of δ_1 and δ_2 on every nontrivial \mathbb{Q} –simple factor of \mathbf{G} generate a subgroup isomorphic to \mathbb{Z}^2 .

Lemma 4.1 *Let S be a Zariski-dense subsemigroup of $\mathbf{G}(\mathbb{Q})$. Then there exists a regular and \mathbb{R} –regular element $\gamma_0 \in S$ which is anisotropic, without components of finite order, and such that for every commuting multiplicatively independent $\delta_1, \delta_2 \in Z_{\mathbf{G}}(\gamma_0)(\mathbb{Q})$ and every nontrivial \mathbb{R} –character χ of \mathbf{T}_{γ_0} , the real numbers $\chi(\delta_1)$ and $\chi(\delta_2)$ are multiplicatively independent.*

Proof It follows from [23, Theorem 2] that there exists a regular and \mathbb{R} –regular $\gamma_0 \in S$ such that $Z_{\mathbf{G}}(\gamma_0)^\circ$ is a \mathbb{Q} –quasi-irreducible torus in \mathbf{G} which is anisotropic over \mathbb{Q} . Let us suppose that for some nontrivial \mathbb{R} –character χ of \mathbf{T}_{γ_0} , the real numbers $\chi(\delta_1)$

and $\chi(\delta_2)$ are multiplicatively dependent, that is, there exists $(n_1, n_2) \in \mathbb{Z}^2 \setminus \{(0, 0)\}$ such that

$$\chi(\delta_1)^{n_1} \chi(\delta_2)^{n_2} = \chi(\delta_1^{n_1} \delta_2^{n_2}) = 1.$$

Replacing δ_1 and δ_2 by δ_1^l and δ_2^l for suitable $l \geq 1$, we may assume without loss of generality that $\delta_1, \delta_2 \in Z_{\mathbf{G}}(\gamma_0)^\circ$. Then the subgroup $\langle \delta_1^{n_1} \delta_2^{n_2} \rangle$ is contained in a proper subtorus of $Z_{\mathbf{G}}(\gamma_0)^\circ$. Hence, its Zariski closure gives a proper \mathbb{Q} -subtorus of $Z_{\mathbf{G}}(\gamma_0)^\circ$. Since $Z_{\mathbf{G}}(\gamma_0)^\circ$ is \mathbb{Q} -quasi-irreducible, it follows that the projection of $\delta_1^{n_1} \delta_2^{n_2}$ to one of the nontrivial \mathbb{Q} -simple factors of \mathbf{G} should have finite order. However, this contradicts the assumption that δ_1 and δ_2 are multiplicatively independent. Hence, we conclude that $\chi(\delta_1)$ and $\chi(\delta_2)$ are multiplicatively independent for all nontrivial \mathbb{R} -characters χ . \square

Given a regular and \mathbb{R} -regular $g \in G$, we denote by $\Phi_g = \Phi(\mathbf{T}_g, \mathbf{G})$ the root system arising from the action of \mathbf{T}_g on the Lie algebra of \mathbf{G} . Once an ordering on $\Phi(\mathbf{T}_g, \mathbf{G})$ is given, we define the set of positive roots Φ_g^+ , the set of negative roots Φ_g^- , and the set of simple roots $\Delta_g \subset \Phi_g^+$.

Lemma 4.2 *Let $\gamma_0 \in \Gamma$ be an element as in Lemma 4.1. We fix an ordering on Φ_{γ_0} such that*

$$(4-3) \quad \alpha(\gamma_0) < 1 \quad \text{for all } \alpha \in \Phi_{\gamma_0}^-.$$

Then, for every simple root $\alpha_0 \in \Delta_{\gamma_0}$, there exists a sequence $\delta_n \in \Gamma \cap Z_{\mathbf{G}}(\gamma_0)^\circ$ consisting of commuting \mathbb{R} -regular elements and satisfying

$$\alpha_0(\delta_n) \rightarrow 1$$

and

$$\alpha(\delta_n) \rightarrow 0 \quad \text{for all } \alpha \in \Phi_{\gamma_0}^- \text{ that are not proportional to } \alpha_0.$$

Proof Since the group $\mathbf{H} = Z_{\mathbf{G}}(\gamma_0)^\circ$ is anisotropic over \mathbb{Q} , it follows that $\mathbf{H}(\mathbb{Z})$ is a lattice in $\mathbf{H}(\mathbb{R})$. In particular, we deduce that $\Gamma \cap \mathbf{H}$ is a lattice in $\mathbf{H}(\mathbb{R})$, and it contains a subgroup $\Lambda \simeq \mathbb{Z}^r$, where $r = \dim(\mathbf{T}_{\gamma_0})$ is the \mathbb{R} -rank of \mathbf{G} . It also follows that $\gamma_0^l \in \Lambda$ for some $l \geq 1$. We have a decomposition

$$\mathbf{H} = \mathbf{H}^{(1)} \cdots \mathbf{H}^{(r)},$$

where $\mathbf{H}^{(i)} = \mathbf{H} \cap \mathbf{G}^{(i)}$ and the $\mathbf{G}^{(i)}$ are the connected \mathbb{Q} -simple normal subgroups of \mathbf{G} from (4-2). Since the $\mathbf{H}^{(i)}$ are anisotropic over \mathbb{Q} , $\mathbf{H}^{(i)}(\mathbb{Z})$ is a lattice in $\mathbf{H}^{(i)}(\mathbb{R})$ as well. It follows from our assumption on the rank of G that each of the

factors $\mathbf{G}^{(i)}$ has \mathbb{R} -rank at least two, so that

$$(4-4) \quad \mathbf{H}^{(i)} \cap \Lambda \simeq \mathbb{Z}^{r_i} \quad \text{with } r_i \geq 2.$$

We consider a collection of linear forms $L_\alpha = \log(\alpha)$ for $\alpha \in \Delta_{\gamma_0}$ on \mathbb{Z}^r that defines the negative Weyl chamber

$$C^- = \{a \in \Lambda \otimes \mathbb{R} : L_\alpha(a) < 0 \text{ for all } \alpha \in \Delta_{\gamma_0}\}$$

that contains γ_0^l . Since Λ is a lattice in $\Lambda \otimes \mathbb{R}$, there exists $\delta_0 \in \Lambda$ such that

$$(4-5) \quad \alpha(\delta_0) \leq \alpha_0(\delta_0) < 0 \quad \text{for every } \alpha \in \Delta_{\gamma_0}.$$

This, in particular, implies that δ_0 has no components of finite order. It follows from (4-4) that there exists $\delta_1 \in \Lambda$ such that δ_0 and δ_1 are multiplicatively independent. We consider the subgroup $\Lambda_0 = \langle \delta_0, \delta_1 \rangle \simeq \mathbb{Z}^2$ of Λ . We note that $C^- \cap (\Lambda_0 \otimes \mathbb{R})$ defines a nontrivial cone in $\Lambda_0 \otimes \mathbb{R} \simeq \mathbb{R}^2$. It follows from Lemma 4.1 that $\ker(L_\alpha|_{\Lambda_0}) = 0$, and forms $L_{\alpha_1}|_{\Lambda_0}$ and $L_{\alpha_2}|_{\Lambda_0}$ are proportional only when the roots α_1 and α_2 are proportional. In particular, every nontrivial element of Λ_0 is \mathbb{R} -regular. It follows from (4-5) that the line $L_{\alpha_0} = 0$ gives one of the faces of the cone $C^- \cap (\Lambda_0 \otimes \mathbb{R})$. Then there exists a sequence $\delta_n \in C^- \cap \Lambda_0$ such that $\delta_n \rightarrow \infty$ and $L_{\alpha_0}(\delta_n) \rightarrow 0$. Moreover, it is clear that $|L(\delta_n)| \rightarrow \infty$ for any linear form L on $\Lambda_0 \otimes \mathbb{R}$ which is not proportional to $L_{\alpha_0}|_{\Lambda_0 \otimes \mathbb{R}}$. In particular, $|L_\alpha(\delta_n)| \rightarrow \infty$ for any $\alpha \in \Delta_{\gamma_0} \setminus \{\alpha_0\}$. Since $\delta_n \in C^- \cap \Lambda_0$, it follows that $L_\alpha(\delta_n) \rightarrow -\infty$ for all $\alpha \in \Delta_{\gamma_0} \setminus \{\alpha_0\}$. This also implies that $L_\alpha(\delta_n) \rightarrow -\infty$ for all $\alpha \in \Phi_{\gamma_0}^-$ that are not proportional to α_0 and proves the lemma. \square

Now we apply the above results to the Zariski-dense subsemigroup S of Γ constructed in Proposition 3.1. We recall that for every element $\gamma \in S$, the map $\rho(\gamma)$ has a differentiable sink in M . We fix $\gamma_0 \in S$ as in Lemma 4.1 that determines the ordering on Φ_{γ_0} satisfying (4-3) and the corresponding set of simple roots $\Delta_{\gamma_0} \subset \Phi_{\gamma_0}$.

It will be convenient to work with the Cartan decomposition (2-1), which is compatible with $Z_{\mathbf{G}}(\gamma_0)^\circ$. We choose a Cartan involution θ such that the torus $Z_{\mathbf{G}}(\gamma_0)^\circ$ is θ -invariant. Then

$$\mathbf{B}_{\gamma_0}(\mathbb{R})^\circ \subset K \quad \text{and} \quad \text{Lie}(\mathbf{T}_{\gamma_0}(\mathbb{R})) \subset \mathfrak{p}.$$

Since \mathbf{T}_{γ_0} is a maximal \mathbb{R} -split torus, $A = \mathbf{T}_{\gamma_0}(\mathbb{R})^\circ$ gives a Cartan subgroup of G . We abuse notation and identify the root system Φ_{γ_0} of \mathbf{T}_{γ_0} with the root system of \mathfrak{a} introduced in Section 2.1.

Let $I \subset \Delta_{\gamma_0}$ and $\alpha_0 \in \Delta_{\gamma_0} \setminus I$. We recall that $U_I = N_I^- P_I$ denotes the open cell in the flag manifold $F_I = G/P_I$. We also denote by $\mathfrak{n}_I^{(\alpha_0)}$ the subalgebra of \mathfrak{n}_I^- generated by the root spaces \mathfrak{g}_α with $\alpha \in \Phi_{\gamma_0}^- \setminus \Phi_I^I$ which are proportional to α_0 , and define

$$N_I^{(\alpha_0)} = \exp(\mathfrak{n}_I^{(\alpha_0)}) \quad \text{and} \quad U_I^{(\alpha_0)} = N_I^{(\alpha_0)} P_I.$$

The $U_I^{(\alpha_0)}$ is a nontrivial submanifold of the open cell U_I . Let

$$\pi_0^{(\alpha_0)}: U_I = \exp(\mathfrak{n}_I^-) P_I \rightarrow U_I^{(\alpha_0)} = \exp(\mathfrak{n}_I^{(\alpha_0)}) P_I$$

be the natural projection map.

Lemma 4.3 *Let $\gamma_0 \in \Gamma$ be as in Lemma 4.1. Then there exists a sequence of commuting \mathbb{R} -regular elements $\delta_n^{(\alpha_0)} \in \Gamma \cap Z_G(\gamma_0)^\circ$ satisfying*

$$(4-6) \quad \alpha_0(\delta_n^{(\alpha_0)}) \rightarrow 1$$

and

$$(4-7) \quad \alpha(\delta_n^{(\alpha_0)}) \rightarrow 0 \quad \text{for all } \alpha \in \Phi_{\gamma_0}^- \text{ that are not proportional to } \alpha_0$$

such that the projection map $\pi_0^{(\alpha_0)}$ is the limit of the maps $\rho_0(\delta_n^{(\alpha_0)})$ acting on $U_I \subset F_I$.

In particular, for every $g \in N_I^{(\alpha_0)}$ and $x \in U_I$,

$$(4-8) \quad \pi_0^{(\alpha_0)}(gx) = g\pi_0^{(\alpha_0)}(x),$$

and for every $\gamma \in \Gamma \cap Z_G(\gamma_0)^\circ$ and $x \in U_I$,

$$(4-9) \quad \pi_0^{(\alpha_0)}(\rho_0(\gamma)x) = \rho_0(\gamma)\pi_0^{(\alpha_0)}(x).$$

Proof Let $\delta_n \in \Gamma \cap Z_G(\gamma_0)^\circ$ be the sequence constructed in Lemma 4.2. Since γ_0 is regular and \mathbb{R} -regular, its centraliser satisfies (4-1), so that we can write $\delta_n = k_n a_n$ with $k_n \in \mathbf{B}_{\gamma_0}(\mathbb{R})$ and $a_n \in \mathbf{T}_{\gamma_0}(\mathbb{R})$. For every $v \in \mathfrak{n}_I$,

$$\rho_0(\delta_n) \exp(v) P_I = \exp(\text{Ad}(\delta_n)v) P_I = \exp(\text{Ad}(a_n)\text{Ad}(k_n)v) P_I.$$

It follows from properties of the sequence δ_n that $\alpha_0(a_n) \rightarrow 1$ and $\alpha(a_n) \rightarrow 0$ for every $\Phi_{\gamma_0}^- \setminus \Phi_I^I$, so that for every $l \geq 1$ and $w \in \mathfrak{n}_I$,

$$(4-10) \quad \text{Ad}(a_n^l)w \rightarrow p_0^{(\alpha_0)}(w),$$

where $p_0^{(\alpha_0)}: \mathfrak{n}_I^- \rightarrow \mathfrak{n}_I^{(\alpha_0)}$ is the natural projection map. Moreover, this convergence is uniform over $l \geq 1$ and w in compact sets. We observe that the transformations

$\text{Ad}(k_n)|_{\mathfrak{n}_I}$ belong to a compact abelian group $\Omega < \text{GL}(\mathfrak{n}_I)$. Hence, passing to a subsequence, we may assume that $\text{Ad}(k_n)|_{\mathfrak{n}_I} \rightarrow \omega$ for some $\omega \in \Omega$. For every $j \geq 1$, there exists $l_j \geq 1$ such that

$$d(\omega^{l_j}, \text{id}) < j^{-1},$$

and there exists $n_j \geq 1$ such that

$$d(\text{Ad}(k_{n_j})|_{\mathfrak{n}_I}, \omega) < (l_j j)^{-1}.$$

Hence, it follows that

$$d(\text{Ad}(k_{n_j}^{l_j})|_{\mathfrak{n}_I}, \text{id}) < j^{-1}.$$

Combining this estimate with (4-10), we deduce that for $v \in \mathfrak{n}_I$,

$$\text{Ad}(\delta_{n_j}^{l_j})v = \text{Ad}(a_{n_j}^{l_j})\text{Ad}(k_{n_j}^{l_j})v \rightarrow p_0^{(\alpha_0)}(v).$$

Hence, the required sequence can be taken to be $\delta_j^{(\alpha_0)} = \delta_{n_j}^{l_j}$. This proves the first part of the lemma. The second part, (4-8), also follows because for $v \in \mathfrak{n}_I^{(\alpha_0)}$ and $x \in U_I$,

$$\rho_0(\delta_n^{(\alpha_0)}) \exp(v)x = \exp(\text{Ad}(\delta_n^{(\alpha_0)})v)\rho_0(\delta_n^{(\alpha_0)})x \rightarrow \exp(v)\pi_0^{(\alpha_0)}(x).$$

The last claim, (4-9), is immediate as well because $Z_G(\gamma_0)^\circ$ is commutative. This completes the proof. \square

Next we show that there exist dynamically defined projection maps on M which are analogues of the projection maps $\pi_0^{(\alpha_0)}$. We recall that $\rho(\gamma_0)$ has a differentiable sink $s \in M$. It is also clear that $s_0 = eP_I$ is the unique sink of the \mathbb{R} -regular element γ_0 . Since the C^0 -conjugacy $\phi: F_I \rightarrow M$ intertwines $\rho_0(\gamma_0)$ and $\rho(\gamma_0)$, it follows that $\phi(s_0) = s$.

Lemma 4.4 *There exists a neighbourhood \mathcal{O}_0 of the sink s such that $\rho(\delta_n^{(\alpha_0)})|_{\mathcal{O}_0}$ converges to a smooth map $\tilde{\pi}_0^{(\alpha_0)}: \mathcal{O}_0 \rightarrow M$ satisfying*

$$(4-11) \quad \phi \circ \pi_0^{(\alpha_0)} = \tilde{\pi}_0^{(\alpha_0)} \circ \phi$$

on $\phi^{-1}(\mathcal{O}_0)$.

To prove this lemma, we need to use the theory of polynomial normal forms for smooth diffeomorphisms, which we now recall. We refer to [11; 12] for more details. We note that we only require normal forms at a differentiable sink, rather than the more elaborate theory of contractions on vector bundle extensions developed in [11; 12]. However, we are not aware of a simpler reference for our case.

Let f be a diffeomorphism with a differentiable sink s , χ_1, \dots, χ_l the different moduli of eigenvalues of $D(f)_s$, and m_1, \dots, m_l their multiplicities. We represent the tangent space $T_s(M) \simeq \mathbb{R}^m$ as the direct sum of the spaces $\mathbb{R}^{m_1}, \dots, \mathbb{R}^{m_l}$, and let (t_1, \dots, t_l) be the corresponding coordinate representation of a vector $t \in \mathbb{R}^n$. Let

$$P: \mathbb{R}^m \rightarrow \mathbb{R}^m, \quad (t_1, \dots, t_l) \mapsto (P_1(t_1, \dots, t_l), \dots, P_l(t_1, \dots, t_l)),$$

be a polynomial map preserving the origin. We will say that the map P is of *subresonance type* if it contains only homogeneous terms in $P_i(t_1, \dots, t_l)$ with degree of homogeneity s_j in the coordinates of t_j for $i = 1, \dots, l$, for which the subresonance relations

$$\chi_i \leq \sum_{j \neq i} s_j \chi_j$$

hold. There are only finitely many subresonance relations and it is known (see [11; 12]) that polynomial maps of the subresonance type with invertible derivative at the origin generate a finite-dimensional Lie group. We will denote this group by SR_χ . The polynomial maps of subresonance type provide convenient normal forms of the diffeomorphism f and its centraliser.

Proposition 4.5 [11; 12] *There exists a coordinate chart $\omega: \mathcal{O} \rightarrow \mathcal{O}' \subset \mathbb{R}^m$ around the sink s for which $\omega \circ f \circ \omega^{-1}$ is a polynomial map of subresonance type contained in the group SR_χ . Moreover, this coordinate chart transforms into such a normal form in SR_χ any diffeomorphism which commutes with f .*

With the help of this proposition, we prove Lemma 4.4.

Proof of Lemma 4.4 By Proposition 4.5, there is a neighbourhood \mathcal{O} of $s = \phi(s_0)$ on which we have normal forms for every diffeomorphism that commutes with $\rho(\gamma_0)$ and, in particular, for the diffeomorphisms $\rho(\delta_n^{(\alpha_0)})$. Replacing \mathcal{O} by a neighbourhood $\mathcal{O}_0 \subset \mathcal{O}$, we can assume that $\rho(\gamma_0)$ and $\rho(\delta_n^{(\alpha_0)})$ map \mathcal{O}_0 into \mathcal{O} . Indeed, this follows from properties of the sequence $\delta_n^{(\alpha_0)}$ (see (4-6)–(4-7)). Then we define $\tilde{\pi}_0^{(\alpha_0)}$ on \mathcal{O}_0 as the limit of $\rho(\delta_n^{(\alpha_0)})$. It is clear that this limit exists in C^0 –topology because ϕ intertwines the map $\rho(\delta_n^{(\alpha_0)})$ with the map $\rho_0(\delta_n^{(\alpha_0)})$. Moreover, since the maps $\rho(\delta_n^{(\alpha_0)})|_{\mathcal{O}_0}$ are polynomials of bounded degree in the normal form coordinates, $\tilde{\pi}_0^{(\alpha_0)}$ is also a polynomial of the same degree, hence smooth. The equivariance property is immediate from the equivariance of ϕ with respect to the actions ρ and ρ_0 . \square

Proposition 4.6 *The local projection map $\tilde{\pi}_0^{(\alpha_0)}: \mathcal{O}_0 \rightarrow M$, defined in Lemma 4.4, extends to a smooth map $\tilde{\pi}_0^{(\alpha_0)}: \phi(U_I) \rightarrow M$ such that*

$$(4-12) \quad \phi \circ \pi_0^{(\alpha_0)} = \tilde{\pi}_0^{(\alpha_0)} \circ \phi$$

on U_I .

Proof We can extend the smooth projection map $\tilde{\pi}_0^{(\alpha_0)}$ defined on the neighbourhood of the sink \mathcal{O}_0 to the whole $\phi(U_I)$ using conjugation by $\rho(\gamma_0)$. More precisely, given any compact subset Ω of $\phi(U_I)$, there exists n such that $\rho(\gamma_0)^n(\Omega) \subset \mathcal{O}_0$. Then we set

$$\tilde{\pi}_0^{(\alpha_0)}(x) = \rho(\gamma_0)^{-n} \tilde{\pi}_0^{(\alpha_0)}(\rho(\gamma_0)^n x), \quad x \in \Omega.$$

It follows from (4-9) and (4-11) that the definition of $\tilde{\pi}_0^{(\alpha_0)}$ is consistent with $\tilde{\pi}_0^{(\alpha_0)}$ on \mathcal{O}_0 and is independent of the choice of n . This allows us to extend $\tilde{\pi}_0^{(\alpha_0)}$ to $\phi(U_I)$ so that (4-12) holds. \square

Lemma 4.7 *There exists $\gamma \in \Gamma \cap Z_G(\gamma_0)^\circ$ such that $\phi(U_I^{(\alpha_0)})$ in a neighbourhood of the sink s is equal to the strong stable manifold of $\rho(\gamma)$. In particular, $\phi(U_I^{(\alpha_0)})$ is an immersed submanifold.*

Proof As we already observed in the proof of Lemma 4.2, $\Gamma \cap Z_G(\gamma_0)^\circ$ is a lattice subgroup in $Z_G(\gamma_0)^\circ(\mathbb{R})$. Its projection to $T_{\gamma_0}(\mathbb{R})$ is also a lattice. Since the set of simple roots Δ_{γ_0} forms a basis of the dual space of $T_{\gamma_0}(\mathbb{R})^\circ$. It follows that there exists $\gamma \in \Gamma \cap Z_G(\gamma_0)^\circ$ such that $\alpha_0(\gamma) < 1$ and $\alpha(\gamma) > 1$ for all $\alpha \in \Delta_{\gamma_0} \setminus \{\alpha_0\}$. Then $U_I^{(\alpha_0)}$ is the strong stable manifold of $\rho_0(\gamma)$ at $s_0 = \phi^{-1}(s)$. Suppose y is contained in a small neighbourhood of $s = \phi(s_0)$ in M , and it has its forward $\rho(\gamma)$ -orbit in this neighbourhood. Let $y = \phi(x)$ for some $x \in U_I$. Then the forward orbit $\rho_0(\gamma)^n x$ stays in a small neighbourhood of s_0 . In particular, it follows that $x \in U_I^{(\alpha_0)}$ and $y \in \phi(U_I^{(\alpha_0)})$. Since $\rho_0(\gamma)$ is purely hyperbolic at s_0 , we see that $\rho_0(\gamma)^n y$ converges to s_0 exponentially fast. Hence, for some fixed sufficiently large n_0 , the transformation $\rho_0(\gamma^{n_0} \gamma_0^{-1})$ also contracts x to s_0 . Since ϕ is continuous, it follows that $\rho(\gamma^{n_0} \gamma_0^{-1})^n y = \rho(\gamma_0)^{-n} \rho(\gamma^{n_0})^n y$ converges to $s = \phi(s_0)$. On the other hand, s is a differentiable sink of $\rho(\gamma_0)$, so that we conclude that $\rho(\gamma^{n_0})^n y$ must converge to s exponentially fast. This proves that $\phi(U_I^{(\alpha_0)})$ is equal to the strong stable manifold of $\rho(\gamma^{n_0})$ in a neighbourhood of s . In particular, we also derive that $\phi(U_I^{(\alpha_0)})$ is a submanifold in a neighbourhood of s . We deduce that $\phi(U_I^{(\alpha_0)})$ is an immersed submanifold using the action of $\rho(\gamma_0)^{-1}$. \square

5 Smoothness along foliations

We keep the notation from the previous section and continue our investigation of the conjugacy map $\phi: F_I \rightarrow M$. In this section, we establish smoothness of ϕ restricted to open dense subsets of submanifolds $U_I^{(\alpha_0)}$ of F_I . The higher-rank assumption on G is crucial here.

Let $H_I^{(\alpha_0)}$ be a connected component of the closure of $\rho_0(Z_G(\gamma_0)^\circ \cap \Gamma)|_{U_I^{(\alpha_0)}}$. The latter can be identified with the closure of the linear group $\text{Ad}(Z_G(\gamma_0)^\circ \cap \Gamma)|_{\mathfrak{n}_I^{(\alpha_0)}}$. It follows from Lemma 4.1 that $H_I^{(\alpha_0)}$ is not trivial. Since $H_I^{(\alpha_0)}$ acts freely away from the fixed point s_0 , we get a foliation of $U_I^{(\alpha_0)} \setminus \{s_0\}$ by $H_I^{(\alpha_0)}$ -orbits. We also note the action of $H_I^{(\alpha_0)}$ commutes with the action of $\rho_0(\gamma_0)$.

Proposition 5.1 *The conjugacy map $\phi: F_I \rightarrow M$ restricted to $U_I^{(\alpha_0)}$ is C^∞ on an open dense subset.*

In the proof of this proposition, we use a technical lemma which involves equidistribution properties of flows on the homogeneous space $\Gamma \backslash G$. This requires new notation which we now introduce. Let

$$M = Z_K(\mathfrak{a})^\circ.$$

Let $\{a_t\}$ be a one-parameter subgroup of the Cartan group A such that N_I^+ is the contracting horospherical subgroup for it, namely,

$$N_I^+ = \{g \in G : a_t g a_t^{-1} \rightarrow e\}.$$

We write G as an almost direct product

$$G = G^{(1)} \cdots G^{(r)},$$

where the $G^{(i)}$ are connected normal subgroups of G such that $\Gamma \cap G^{(i)}$ are irreducible lattices in $G^{(i)}$. Then $(\Gamma \cap G^{(1)}) \cdots (\Gamma \cap G^{(r)})$ has finite index in Γ and, without loss of generality, we may assume that

$$\Gamma = (\Gamma \cap G^{(1)}) \cdots (\Gamma \cap G^{(r)}).$$

We denote by G_I the product of the $G^{(i)}$ which are contained in P_I .

Lemma 5.2 *Given arbitrary $g_0, g \in G$, $p_0 \in MA$, a neighbourhood \mathcal{O} of the identity in G , and a neighbourhood \mathcal{O}' of the identity in N_I^+ , for all sufficiently large t ,*

$$\Gamma \cap g_0 \mathcal{O} G_I (\mathcal{O}' p_0 a_t)^{-1} g^{-1} \neq \emptyset.$$

Proof We first assume for simplicity that the lattice Γ is irreducible. Then this lemma follows directly from well-known equidistribution properties of the expanding horospherical subgroups on the space $\Gamma \backslash G$ (see, for instance, [17, 2.2.1]): for arbitrary $\psi \in C_c(\Gamma \backslash G)$, $f \in C_c(N_I^+)$ and $x \in \Gamma \backslash G$,

$$(5-1) \quad \int_{N_I^+} f(p) \psi(xpa_t) dm_{N_I^+}(p) \rightarrow \left(\int_{N_I^+} f dm_{N_I^+} \right) \left(\int_{\Gamma \backslash G} \psi dm_{\Gamma \backslash G} \right)$$

as $t \rightarrow \infty$, where $m_{N_I^+}$ denotes a Haar measure on N_I^+ , and $m_{\Gamma \backslash G}$ is the Haar measure on $\Gamma \backslash G$ such that $m_{\Gamma \backslash G}(\Gamma \backslash G) = 1$. We take nonzero $f \geq 0$ with $\text{supp}(f) \subset p_0^{-1}\mathcal{O}'p_0$ (note that MA normalises N_I^+), nonzero $\psi \geq 0$ with $\text{supp}(\psi) \subset \Gamma g_0\mathcal{O}$, and $x = \Gamma g p_0$. It follows from (5-4) that

$$\int_{N_I^+} f(p) \psi(xpa_t) dm_{N_I^+}(p) > 0$$

for all sufficiently large t . This implies that there exists $p \in p_0^{-1}\mathcal{O}'p_0$ such that

$$\Gamma g p_0 p a_t \in \Gamma g_0 \mathcal{O}.$$

Hence, it follows that

$$\Gamma \cap g_0 \mathcal{O}(\mathcal{O}' p_0 a_t)^{-1} g^{-1} \neq \emptyset.$$

This proves the lemma under the assumption that Γ is irreducible.

Now we discuss the general case. We note that the proof of (5-4) is based on the mixing property on $\Gamma \backslash G$: for every $\psi_1, \psi_2 \in C_c(\Gamma \backslash G)$,

$$(5-2) \quad \int_{\Gamma \backslash G} \psi_1(z) \psi_2(zg) dm_{\Gamma \backslash G}(z) \rightarrow \left(\int_{\Gamma \backslash G} \psi_1 dm_{\Gamma \backslash G} \right) \left(\int_{\Gamma \backslash G} \psi_2 dm_{\Gamma \backslash G} \right)$$

as $g \rightarrow \infty$ in G . Although (5-2) fails in general when the lattice Γ is not irreducible, it is true provided that the projection of g to every factor $G^{(i)}$ goes to infinity. We observe that the projection of a_t to every simple factor which is not contained in G_I goes to infinity as $t \rightarrow \infty$. Hence, this flow still satisfies the mixing property on the space $(\Gamma G_I) \backslash G$, or equivalently the mixing property for G_I -invariant functions on the space $\Gamma \backslash G$. Namely, for every G_I -invariant $\psi_1, \psi_2 \in C_c(\Gamma \backslash G)$,

$$(5-3) \quad \int_{\Gamma \backslash G} \psi_1(z) \psi_2(z a_t) dm_{\Gamma \backslash G}(z) \rightarrow \left(\int_{\Gamma \backslash G} \psi_1 dm_{\Gamma \backslash G} \right) \left(\int_{\Gamma \backslash G} \psi_2 dm_{\Gamma \backslash G} \right)$$

as $t \rightarrow \infty$. This allows us to use the same argument as in [17, 2.2.1] to deduce that for a G_I -invariant $\psi \in C_c(\Gamma \backslash G)$, $f \in C_c(N_I^+)$ and $x \in \Gamma \backslash G$,

$$(5-4) \quad \int_{N_I^+} f(p) \psi(xpa_t) dm_{N_I^+}(p) \rightarrow \left(\int_{N_I^+} f dm_{N_I^+} \right) \left(\int_{\Gamma \backslash G} \psi dm_{\Gamma \backslash G} \right)$$

as $t \rightarrow \infty$. Now we can finish the proof as in the previous paragraph by taking nonzero G_I -invariant function $\psi \geq 0$ with $\text{supp}(\psi) \subset \Gamma g_0 \mathcal{O} G_I$. We deduce that for all sufficiently large t , there exists $p \in p_0^{-1} \mathcal{O}' p_0$ such that

$$\Gamma g p_0 p a_t \in \Gamma g_0 \mathcal{O} G_I.$$

This implies the lemma. \square

Proof of Proposition 5.1 First, we show that the conjugacy map $\phi: F_I \rightarrow M$ is C^∞ along the $H_I^{(\alpha_0)}$ -orbits in $U_I^{(\alpha_0)} \setminus \{s_0\}$. As before, for a sequence $\gamma_n \in Z_G(\gamma_0)^\circ \cap \Gamma$ such that $\rho_0(\gamma_n)|_{U_I^{(\alpha_0)}}$ converges to $h \in H_I^{(\alpha_0)}$, we get convergence in C^0 -topology of the maps $\rho(\gamma_n)$ on the intersection of $\phi(U_I^{(\alpha_0)} \setminus \{s_0\})$ with a small enough neighbourhood contained in a normal forms coordinate chart of the sink $s = \phi(s_0)$ of $\rho(\gamma_0)$. Since the normal forms of the $\rho(\gamma_n)$ are polynomials of fixed degree, the limiting map is also a polynomial of the same degree, and hence smooth. By [20, Section 5.1, Corollary], we get a smooth action of $H_I^{(\alpha_0)}$ on this neighbourhood which intertwines via ϕ with the action of $H_I^{(\alpha_0)}$ on $U_I^{(\alpha_0)} \setminus \{s_0\}$. Since $H_I^{(\alpha_0)}$ commutes with γ_0 , we can extend the action to a smooth action on $\phi(U_I^{(\alpha_0)} \setminus \{s_0\})$ which is again intertwined with the action of $H_I^{(\alpha_0)}$ on $U_I^{(\alpha_0)} \setminus \{s_0\}$ via ϕ . Hence, it follows that ϕ is smooth along the $H_I^{(\alpha_0)}$ -orbits in $U_I^{(\alpha_0)} \setminus \{s_0\}$.

Our next step is, starting with the orbit foliation of $H_I^{(\alpha_0)}$, to construct additional smooth foliations \mathcal{F}_i defined on open subsets of $U_I^{(\alpha_0)}$. If we show that for $x \in F_I$,

$$(5-5) \quad \phi|_{\mathcal{F}_i(x)} \text{ is smooth}$$

and

$$(5-6) \quad T_x(\mathcal{F}_1(x)) + \cdots + T_x(\mathcal{F}_r(x)) = T_x(U_I^{(\alpha_0)}),$$

then it will follow that ϕ is smooth in a neighbourhood of x in $U_I^{(\alpha_0)}$. These new foliations are constructed as

$$(5-7) \quad \mathcal{F}_\gamma(y) = \pi_0^{(\alpha_0)}(\gamma H_I^{(\alpha_0)} y), \quad y \in U_I^{(\alpha_0)} \setminus \{s_0\},$$

for suitable $\gamma \in \Gamma$. We note that since ϕ is Γ -equivariant and also equivariant with respect to $\pi_0^{(\alpha_0)}$ and $\tilde{\pi}_0^{(\alpha_0)}$ (see Proposition 4.6), it is clear that (5-5) holds. Hence, the main task is to arrange the transversality property (5-6). We construct such foliations inductively.

We take arbitrary $g_0 \in N_I^{(\alpha_0)} \setminus \{e\}$ and a neighbourhood \mathcal{O} of identity in G and consider a distribution \mathcal{E} in $T(F_I)$ defined on the neighbourhood $\pi_0^{(\alpha_0)}(g_0 \mathcal{O} P_I)$ of $g_0 P_I$ in $U_I^{(\alpha_0)}$ and contained in the tangent distribution of $U_I^{(\alpha_0)}$. We start with \mathcal{E} being the tangent distribution to the orbit foliation $H_I^{(\alpha_0)}x$ for $x \in U_I^{(\alpha_0)} \setminus \{s_0\}$. If $\dim(U_I^{(\alpha_0)}) = \dim(H_I^{(\alpha_0)})$, then the proposition follows from smoothness of ϕ along the orbit foliation, so that we assume that \mathcal{E} is not equal to the tangent distribution of $U_I^{(\alpha_0)}$. For g in the neighbourhood of g_0 in $N_I^{(\alpha_0)}$, we consider a family of subspaces

$$\mathcal{V}(g) = D(g)_{eP_I}^{-1} \mathcal{E}(gP_I)$$

of the tangent space $T_{eP_I}(F_I)$. We make the identification $T_{eP_I}(F_I) \simeq \mathfrak{g}/\mathfrak{p}_I$, so that $D(p)_{eP_I} = \text{Ad}(p)$ for $p \in P_I$.

We recall that $N_I^{(\alpha_0)} = \exp(\mathfrak{n}_I^{(\alpha_0)})$, where $\mathfrak{n}_I^{(\alpha_0)}$ is the span of root spaces for negative roots proportional to α_0 . It follows from the properties of root systems that either

$$N_I^{(\alpha_0)} = \exp(\mathfrak{g}_{-\alpha_0}) \quad \text{or} \quad N_I^{(\alpha_0)} = \exp(\mathfrak{g}_{-\alpha_0} + \mathfrak{g}_{-2\alpha_0}).$$

We treat these two cases separately.

Case 1 ($N_I^{(\alpha_0)} = \exp(\mathfrak{g}_{-\alpha_0})$) Then $N_I^{(\alpha_0)}$ is commutative, and

$$\mathcal{E}(g_0 P_I) \subseteq T_{g_0 P_I}(\exp(\mathfrak{g}_{-\alpha_0})g_0 P_I),$$

so that

$$\mathcal{V}(g_0) \subseteq D(g_0)_{eP_I}^{-1} T_{g_0 P_I}(\exp(\mathfrak{g}_{-\alpha_0})g_0 P_I) = T_{eP_I}(\exp(\mathfrak{g}_{-\alpha_0})P_I) = \mathfrak{g}_{-\alpha_0} + \mathfrak{p}_I.$$

In particular, it follows that $\text{Ad}(N_I^+)$ acts trivially on $\mathcal{V}(g_0)$, and $\text{Ad}(A)$ acts on $\mathcal{V}(g_0)$ by scalar multiples. By Lemma 5.2, given arbitrary $p_0 \in MA$, a neighbourhood $\tilde{\mathcal{O}} \subset \mathcal{O}$ of identity in G , a neighbourhood \mathcal{O}' of identity in N_I^+ and sufficiently large t , there exists $\gamma \in \Gamma$ such that

$$(5-8) \quad \gamma \in g_0 \tilde{\mathcal{O}} G_I(\mathcal{O}' p_0 a_t)^{-1} g_0^{-1}.$$

In particular, it follows that $\gamma g_0 P_I \in g_0 \mathcal{O} P_I$. We claim that there exists $\gamma \in \Gamma$ such that

$$(5-9) \quad \gamma g_0 P_I \in g_0 \mathcal{O} P_I \quad \text{and} \quad D(\pi_0^{(\alpha_0)} \circ \gamma)_{g_0 P_I} \mathcal{E}(g_0 P_I) \not\subseteq \mathcal{E}(\pi_0^{(\alpha_0)}(\gamma g_0 P_I)).$$

It follows from (5-8) that given arbitrary $p_0 \in M$, there exists a sequence $\gamma_i \in \Gamma$ such that

$$(5-10) \quad \gamma_i = g_i (u_i p_0 a_{t_i})^{-1} g_0^{-1} \quad \text{with } g_i P_I \rightarrow g_0 P_I, u_i \in N_I^+, u_i \rightarrow e, t_i \rightarrow \infty.$$

Then $\gamma_i g_0 P_I = g_i P_I \rightarrow g_0 P_I$, so that $\gamma_i g_0 P_I \in g_0 \mathcal{O} P_I$ for sufficiently large i . Suppose that for those i , we have

$$D(\pi_0^{(\alpha_0)} \circ \gamma_i)_{g_0 P_I} \mathcal{E}(g_0 P_I) \subset \mathcal{E}(\pi_0^{(\alpha_0)}(g_i P_I)).$$

This means that

$$D(\pi_0^{(\alpha_0)} \circ g_i)_{e P_I} \text{Ad}((u_i p_0 a_{t_i})^{-1}) \mathcal{V}(g_0) \subset \mathcal{E}(\pi_0^{(\alpha_0)}(g_i P_I)).$$

We have

$$\text{Ad}((u_i p_0 a_{t_i})^{-1}) \mathcal{V}(g_0) = \text{Ad}(p_0^{-1}) \mathcal{V}(g_0).$$

Hence, we conclude that for all $p_0 \in M$,

$$D(\pi_0^{(\alpha_0)} \circ g_0)_{e P_I} \text{Ad}(p_0^{-1}) \mathcal{V}(g_0) \subset \mathcal{E}(g_0 P_I).$$

Since $g_0 \in N_I^{(\alpha_0)}$, it follows from (4-8) that

$$D(g_0)_{e P_I} D(\pi_0^{(\alpha_0)})_{e P_I} \text{Ad}(p_0^{-1}) \mathcal{V}(g_0) \subset \mathcal{E}(g_0 P_I).$$

Let \mathcal{W} be the subspace generated by $\text{Ad}(p_0^{-1}) \mathcal{V}(g)$ for $p_0 \in M$. Then

$$D(g_0)_{e P_I} D(\pi_0^{(\alpha_0)})_{e P_I} \mathcal{W} \subset \mathcal{E}(g_0 P_I)$$

and

$$D(\pi_0^{(\alpha_0)})_{e P_I} \mathcal{W} \subset \mathcal{V}(g_0).$$

Since \mathcal{W} is $\text{Ad}(M)$ -invariant, it follows from Lemma 5.3 below that $\mathcal{W} = \mathfrak{g}_{-\alpha_0} + \mathfrak{p}_I$, and $\dim(\mathcal{E}) = \dim(U_I^{(\alpha_0)})$, but we have assumed that the distribution \mathcal{E} is not equal to the tangent distribution of $U_I^{(\alpha_0)}$. This contradiction shows that (5-9) holds. Therefore, we obtain a new distribution $D(\pi_0^{(\alpha_0)} \circ \gamma) \mathcal{E}$ contained in $T(U_I^{(\alpha_0)})$ and defined in a neighbourhood $\pi_0^{(\alpha_0)}(\gamma g_0 P_I) \in g_0 \mathcal{O} P_I$. Moreover, the distribution $D(\pi_0^{(\alpha_0)} \circ \gamma) \mathcal{E} + \mathcal{E}$ is strictly larger than the distribution \mathcal{E} . Now we can apply the above argument to the distribution $D(\pi_0^{(\alpha_0)} \circ \gamma) \mathcal{E} + \mathcal{E}$ defined in a neighbourhood of $\pi_0^{(\alpha_0)}(\gamma g_0 P_I)$ contained in $\pi_0^{(\alpha_0)}(g_0 \mathcal{O} P_I)$, and doing this inductively, we conclude that there exist $\gamma_1 = e, \gamma_2, \dots, \gamma_r \in \Gamma$ and $x \in \pi_0^{(\alpha_0)}(g_0 \mathcal{O} P_I)$ such that

$$(5-11) \quad D(\pi_0^{(\alpha_0)} \circ \gamma_1) \mathcal{E} + \dots + D(\pi_0^{(\alpha_0)} \circ \gamma_r) \mathcal{E} = T_x(U_I^{(\alpha_0)}).$$

Case 2(a) $(N_I^{(\alpha_0)} = \exp(\mathfrak{g}_{\alpha_0} + \mathfrak{g}_{2\alpha_0}))$ Then

$$\mathcal{E}(g P_I) \subseteq T_{g P_I}(\exp(\mathfrak{g}_{-\alpha_0} + \mathfrak{g}_{-2\alpha_0})g P_I)$$

and

$$\begin{aligned}\mathcal{V}(g) &\subseteq D(g)^{-1}_{eP_I} T_g P_I (\exp(\mathfrak{g}_{-\alpha_0} + \mathfrak{g}_{-2\alpha_0}) g P_I) = T_{eP_I} (\exp(\mathfrak{g}_{-\alpha_0} + \mathfrak{g}_{-2\alpha_0}) P_I) \\ &= \mathfrak{g}_{-\alpha_0} + \mathfrak{g}_{-2\alpha_0} + \mathfrak{p}_I.\end{aligned}$$

We also observe that when \mathcal{E} is equal to the tangent distribution of the orbit foliation of $H_I^{(\alpha_0)}$, it follows from the definition of the group $H_I^{(\alpha_0)}$ that

$$\mathcal{E}(g_0 P_I) \not\subseteq T_{g_0 P_I} (\exp(\mathfrak{g}_{-2\alpha_0}) g_0 P_I).$$

Hence,

$$\mathcal{V}(g_0) \not\subseteq D(g_0)^{-1}_{eP_I} T_{g_0 P_I} (\exp(\mathfrak{g}_{-2\alpha_0}) g_0 P_I).$$

Since $g_0 \in N_I^{(\alpha_0)} = \exp(\mathfrak{g}_{-\alpha_0} + \mathfrak{g}_{-2\alpha_0})$, we have $g_0^{-1} \exp(\mathfrak{g}_{-2\alpha_0}) = \exp(\mathfrak{g}_{-2\alpha_0}) g_0^{-1}$. Thus, we conclude that

$$(5-12) \quad \mathcal{V}(g_0) \not\subseteq T_{eP_I} (\exp(\mathfrak{g}_{-2\alpha_0}) P_I) = \mathfrak{g}_{-2\alpha_0} + \mathfrak{p}_I.$$

A similar argument also gives that

$$(5-13) \quad \mathcal{V}(g_0) \not\subseteq \mathfrak{g}_{-\alpha_0} + \mathfrak{p}_I.$$

In Case 2(a), we additionally assume that

$$(5-14) \quad \mathfrak{g}_{\alpha_0} + \mathfrak{p}_I \not\subseteq \mathcal{V}(g_0).$$

We claim that there exists $\gamma \in \Gamma$ such that

$$(5-15) \quad \gamma g_0 P_I \in g_0 \mathcal{O} P_I \quad \text{and} \quad D(\pi_0^{(\alpha_0)} \circ \gamma)_{g_0 P_I} \mathcal{E}(g_0 P_I) \not\subseteq \mathcal{E}(\pi_0^{(\alpha_0)}(\gamma g_0 P_I)).$$

As in Case 1, we show that given an arbitrary $p_0 \in MA$, there exists $\gamma_i \in \Gamma$ such that (5-10) holds. Then $\gamma_i g_0 P_I = g_i P_I \rightarrow g_0 P_I$, so that for sufficiently large i , we have $\gamma_i g_0 P_I \in g_0 \mathcal{O} P_I$. Suppose that for those i , we have

$$D(\pi_0^{(\alpha_0)} \circ \gamma_i)_{g_0 P_I} \mathcal{E}(g_0 P_I) \subset \mathcal{E}(\pi_0^{(\alpha_0)}(g_i P_I)).$$

This means that

$$D(\pi_0^{(\alpha_0)} \circ g_i)_{eP_I} \text{Ad}((u_i p_0 a_{t_i})^{-1}) \mathcal{V}(g_0) \subset \mathcal{E}(\pi_0^{(\alpha_0)}(g_i P_I)).$$

We have

$$\begin{aligned}\text{Ad}((u_i p_0 a_{t_i})^{-1}) \mathcal{V}(g_0) &= \text{Ad}(a_{t_i}^{-1}) \text{Ad}(p_0^{-1}) \text{Ad}(u_i^{-1}) \mathcal{V}(g_0) \\ &= \text{Ad}(p_0^{-1}) \text{Ad}(a_{t_i}^{-1}) \text{Ad}(u_i^{-1}) \mathcal{V}(g_0).\end{aligned}$$

We observe that for $x \in \mathfrak{n}_I^+$, $v_1 \in \mathfrak{g}_{-\alpha_0}$ and $v_2 \in \mathfrak{g}_{-2\alpha_0}$,

$$(5-16) \quad \text{Ad}(\exp(x))(v_1 + v_2 + \mathfrak{p}_I) = (v_1 + [x, v_2]) + v_2 + \mathfrak{p}_I$$

and

$$(5-17) \quad \text{Ad}(a_t^{-1})(v_1 + v_2 + \mathfrak{p}_I) = e^{-\theta_0 t} v_1 + e^{-2\theta_0 t} v_2 + \mathfrak{p}_I$$

for some $\theta_0 > 0$. It follows from (5-12) that there exists $v = v_1 + v_2 \in \mathfrak{g}_{-\alpha_0} \oplus \mathfrak{g}_{-2\alpha_0}$ with $v_1 \neq 0$ such that $v + \mathfrak{p}_I \in \mathcal{V}(g_0)$. Then, using (5-16) and (5-17), we deduce that

$$\frac{\text{Ad}(a_{t_i}^{-1})\text{Ad}(u_i^{-1})v}{\|\text{Ad}(a_{t_i}^{-1})\text{Ad}(u_i^{-1})v\|} \rightarrow \frac{v_1}{\|v_1\|} \in \mathfrak{g}_{-\alpha_0}.$$

Moreover, we observe that this limit is independent of the sequences t_i and u_i . Therefore, we conclude that every limit point of the sequence subspaces

$$\text{Ad}(a_{t_i}^{-1})\text{Ad}(u_i^{-1})\mathcal{V}(g) \rightarrow \mathcal{V}_\infty$$

contains a nontrivial subspace $\mathcal{V}_\infty \subset \mathfrak{g}_{-\alpha_0} + \mathfrak{p}_I$ independent of p_0 . We conclude that for all $p_0 \in MA$,

$$D(\pi_0^{(\alpha_0)} \circ g_0)_{eP_I} \text{Ad}(p_0^{-1})\mathcal{V}_\infty \subset \mathcal{E}(g_0 P_I).$$

Since $g_0 \in N_I^{(\alpha_0)}$, it follows from (4-8) that also

$$D(g_0)_{eP_I} D(\pi_0^{(\alpha_0)})_{eP_I} \text{Ad}(p_0^{-1})\mathcal{V}_\infty \subset \mathcal{E}(g_0 P_I).$$

Let \mathcal{W} be the subspace generated by $\text{Ad}(p_0^{-1})\mathcal{V}_\infty$, $p_0 \in MA$. Then

$$D(g_0)_{eP_I} D(\pi_0^{(\alpha_0)})_{eP_I} \mathcal{W} \subset \mathcal{E}(g_0 P_I)$$

and

$$D(\pi_0^{(\alpha_0)})_{eP_I} \mathcal{W} \subset \mathcal{V}(g_0).$$

Since $\mathcal{W} \subset \mathfrak{g}_{-\alpha_0} + \mathfrak{p}_I$ is $\text{Ad}(M)$ -invariant, it follows from Lemma 5.3 below that

$$\mathcal{W} = \mathfrak{g}_{-\alpha_0} + \mathfrak{p}_I.$$

However, this contradicts our assumption (5-14). Hence, we conclude that (5-15) holds. This gives a new distribution $D(\pi_0^{(\alpha_0)} \circ \gamma)\mathcal{E}$ contained in $T(U_I^{(\alpha_0)})$ and defined in a neighbourhood $\pi_0^{(\alpha_0)}(\gamma g_0 P_I) \in g_0 \mathcal{O} P_I$. If the distribution $D(\pi_0^{(\alpha_0)} \circ \gamma)\mathcal{E} + \mathcal{E}$ is equal to the tangent distribution of $U_I^{(\alpha_0)}$ in a neighbourhood of $\pi_0^{(\alpha_0)}(\gamma g_0 P_I)$, we obtain (5-6). Otherwise, we apply the argument of Case 2(a) to the distribution $D(\pi_0^{(\alpha_0)} \circ \gamma)\mathcal{E} + \mathcal{E}$ to construct another distribution. This is possible provided that this new distribution satisfies (5-14). Hence, it remains to consider the last case.

Case 2(b) $(N_I^{(\alpha_0)} = \exp(\mathfrak{g}_{\alpha_0} + \mathfrak{g}_{2\alpha_0}) \text{ and } \mathfrak{g}_{-\alpha_0} + \mathfrak{p}_I \subseteq \mathcal{V}(g_0))$ As in the previous cases, we show that for arbitrary $p_0 \in M$, there exists $\gamma_i \in \Gamma$ such that (5-10) holds. Let us suppose that for all sufficiently large i , we have

$$D(\pi_0^{(\alpha_0)} \circ \gamma_i)_{g_0 P_I} \mathcal{E}(g_0 P_I) \subset \mathcal{E}(\pi_0^{(\alpha_0)}(g_i P_I)).$$

Then we deduce that

$$D(\pi_0^{(\alpha_0)} \circ g_i)_{e P_I} \text{Ad}((u_i p_0 a_{t_i})^{-1}) \mathcal{V}(g_0) \subset \mathcal{E}(\pi_0^{(\alpha_0)}(g_i P_I)),$$

where

$$\text{Ad}((u_i p_0 a_{t_i})^{-1}) \mathcal{V}(g_0) = \text{Ad}(p_0^{-1}) \text{Ad}((u_i a_{t_i})^{-1}) \mathcal{V}(g_0).$$

It follows from $\mathfrak{g}_{-\alpha_0} + \mathfrak{p}_I \subseteq \mathcal{V}(g_0)$ that the space $\mathcal{V}(g_0)$ is preserved by $\text{Ad}((u_i a_{t_i})^{-1})$. Hence, we conclude that

$$D(\pi_0^{(\alpha_0)} \circ g_0)_{e P_I} \text{Ad}(p_0^{-1}) \mathcal{V}(g_0) \subset \mathcal{E}(g_0 P_I).$$

Let \mathcal{W} be the subspace generated by $\text{Ad}(p_0^{-1}) \mathcal{V}(g_0)$ for $p_0 \in M$. Then

$$D(g_0)_{e P_I} D(\pi_0^{(\alpha_0)})_{e P_I} \mathcal{W} \subset \mathcal{E}(g_0 P_I)$$

and

$$\mathcal{W} = D(\pi_0^{(\alpha_0)})_{e P_I} \mathcal{W} \subset \mathcal{V}(g_0).$$

Since \mathcal{W} is $\text{Ad}(M)$ -invariant, and $\mathfrak{g}_{-\alpha_0} + \mathfrak{p}_I \subseteq \mathcal{V}(g_0)$ holds, it follows that

$$\mathcal{W} = \mathcal{W}' + \mathfrak{g}_{-\alpha_0} + \mathfrak{p}_I,$$

where \mathcal{W}' is an $\text{Ad}(M)$ -invariant subspace of $\mathfrak{g}_{-2\alpha_0}$. Moreover, it follows from (5-13) that $\mathcal{W}' \neq 0$. Hence, by Lemma 5.3, $\mathcal{W} = \mathfrak{g}_{-2\alpha_0} + \mathfrak{g}_{-\alpha_0} + \mathfrak{p}_I$. However, we have assumed that the distribution \mathcal{E} is not equal to the tangent distribution of $U_I^{(\alpha_0)}$. This contradiction shows that (5-15) holds. Hence, we obtain a new distribution $D(\pi_0^{(\alpha_0)} \circ \gamma) \mathcal{E}$ contained in $T(U_I^{(\alpha_0)})$ and defined in a neighbourhood of $\pi_0^{(\alpha_0)}(g_0 P_I) \in g_0 \mathcal{O} P_I$ such that $D(\pi_0^{(\alpha_0)} \circ \gamma) \mathcal{E} + \mathcal{E}$ is strictly larger than the distribution \mathcal{E} . Applying this argument inductively, we conclude that there exist $\gamma_1 = e, \gamma_2, \dots, \gamma_r \in \Gamma$ and $x \in \pi_0^{(\alpha_0)}(g_0 \mathcal{O} P_I)$ such that (5-11) holds.

Now we are ready to complete the proof. The above argument shows that every open subset of F_I contains x such that for some $\gamma_1, \gamma_2, \dots, \gamma_r \in \Gamma$, the foliations $\mathcal{F}_{\gamma_1}, \dots, \mathcal{F}_{\gamma_r}$ satisfy (5-6) at x . As we already observed, ϕ is C^∞ along these foliations. This implies that ϕ is smooth on a neighbourhood of x in $U_I^{(\alpha_0)}$. This shows that ϕ , restricted to $U_I^{(\alpha_0)}$, is C^∞ on an open dense set and completes the proof of the proposition. \square

It remains to establish the following lemma, which was used in the above proof.

Lemma 5.3 *Let $\alpha_0 \in \Delta$ be a simple root. Then:*

- (i) *The representation of $\text{Ad}(M)$ on the root space $\mathfrak{g}_{-\alpha_0}$ is irreducible.*
- (ii) *The representation of $\text{Ad}(M)$ on the root space $\mathfrak{g}_{-2\alpha_0}$ is irreducible.*

Proof We may assume, without loss of generality, that \mathfrak{g} is simple. We refer to [21, Chapter 5, Section 4] for basic facts about relationship between real semisimple Lie algebras and their complexifications. In particular, we recall that either \mathfrak{g} is a complex Lie algebra considered as a real one or its complexification $\hat{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}$ is simple. We observe that in the first case, $\dim(\mathfrak{g}_{-\alpha_0}) = 2$ and $\mathfrak{g}_{-2\alpha_0} = 0$. The action of $\text{Ad}(M)$ on $\mathfrak{g}_{-\alpha_0}$ is given by the standard action of $\text{SO}(2)$. Hence, the lemma is clear in this case. Now we assume that we are in the second case.

Let \mathfrak{h}^+ be a maximal commutative subalgebra of

$$\mathfrak{m} = \text{Lie}(M) = \text{Lie}(K) \cap \mathfrak{g}_0.$$

Then $\mathfrak{h} = \mathfrak{h}^+ \oplus \mathfrak{a}$ is a maximal commutative subalgebra of \mathfrak{g} , $\hat{\mathfrak{h}} = \mathfrak{h} \otimes \mathbb{C}$ is a Cartan subalgebra in $\hat{\mathfrak{g}}$, and $\hat{\mathfrak{h}}^+ = \mathfrak{h}^+ \otimes \mathbb{C}$ is a Cartan subalgebra in $\hat{\mathfrak{m}}$. Under the natural identification $\mathfrak{a}^* \simeq \hat{\mathfrak{a}}(\mathbb{R})^*$, the root system Φ of \mathfrak{a} is identified with the root system of $\hat{\mathfrak{a}}$ in $\hat{\mathfrak{g}}$. We denote by $\hat{\Phi}$ the root system $\hat{\mathfrak{h}}$ on $\hat{\mathfrak{g}}$. Then

$$\Phi \cup \{0\} = \{\beta|_{\mathfrak{a}} : \beta \in \hat{\Phi}\}.$$

Moreover, there exists a choice of simple roots $\hat{\Delta} \subset \hat{\Phi}$ and $\Delta \subset \Phi$ such that

$$\Delta \cup \{0\} = \{\beta|_{\mathfrak{a}} : \beta \in \hat{\Delta}\}.$$

We set $\hat{\Phi}_0 = \{\beta \in \hat{\Phi} : \beta|_{\mathfrak{a}} = 0\}$ and $\hat{\Delta}_0 = \Phi \cap \hat{\Phi}_0$. Then

$$\hat{\mathfrak{m}} = \hat{\mathfrak{h}} \oplus \left(\bigoplus_{\beta \in \hat{\Phi}_0} \hat{\mathfrak{g}}_\beta \right),$$

and for $\alpha \in \Phi$,

$$(5-18) \quad \mathfrak{g}_\alpha \otimes \mathbb{C} = \bigoplus_{\substack{\beta \in \hat{\Phi} \\ \beta|_{\mathfrak{a}} = \alpha}} \hat{\mathfrak{g}}_\beta.$$

In particular, Φ_0 is the root system of the semisimple Lie algebra $\hat{\mathfrak{m}}'$ with respect to $\hat{\mathfrak{h}} \cap \hat{\mathfrak{m}}'$. Moreover, $\hat{\Delta}_0$ provides the set of simple roots for $\hat{\Phi}_0$.

Now in (5-18), let us consider the case when $\alpha = -\alpha_0$ with $\alpha_0 \in \Delta$. Let $\beta_0 \in \widehat{\Delta}$ be such that $\beta_0|_{\mathfrak{a}} = \alpha_0$. We recall (see [21, Chapter 5, Section 4.3]) that there exists an involution $\omega: \widehat{\Delta} \setminus \widehat{\Delta}_0 \rightarrow \widehat{\Delta} \setminus \widehat{\Delta}_0$ such that if $\beta|_{\mathfrak{a}} = \alpha_0$ for some $\beta \in \widehat{\Delta}$, then either $\beta = \beta_0$ or $\beta = \beta_0^\omega$, and for every $\beta \in \widehat{\Delta} \setminus \widehat{\Delta}_0$, we have a relation

$$\beta \circ \theta = -\beta^\omega - \sum_{\sigma \in \widehat{\Delta}_0} c_{\beta, \sigma} \sigma$$

for some $c_{\beta, \sigma} \geq 0$. Since $\beta|_{\mathfrak{h}^+}$ is purely imaginary and $\theta|_{\mathfrak{h}^+} = \text{id}$, it follows that

$$\overline{\beta(x)} = \beta^\omega(x) - \sum_{\sigma \in \widehat{\Delta}_0} c_{\beta, \sigma} \sigma(x), \quad x \in \mathfrak{h}^+.$$

Hence, since this equality also holds for $x \in \mathfrak{a}$, we conclude that

$$(5-19) \quad \bar{\beta} = \beta^\omega - \sum_{\sigma \in \widehat{\Delta}_0} c_{\beta, \sigma} \sigma.$$

For $\beta \in \widehat{\Phi}$, we set

$$\widehat{\Phi}^-(\beta) = \left\{ \rho \in \widehat{\Phi}^- : \rho = \beta - \sum_{\sigma \in \widehat{\Delta}_0} n_\sigma \sigma \text{ with } n_\sigma \geq 0 \right\}$$

and

$$V(\beta) = \bigoplus_{\rho \in \widehat{\Phi}^-(\beta)} \widehat{\mathfrak{g}}_\rho.$$

Then, if $\beta_0^\omega \neq \beta_0$,

$$(5-20) \quad \mathfrak{g}_{-\alpha_0} \otimes \mathbb{C} = V(-\beta_0) \oplus V(-\beta_0^\omega),$$

and if $\beta_0^\omega = \beta_0$,

$$(5-21) \quad \mathfrak{g}_{-\alpha_0} \otimes \mathbb{C} = V(-\beta_0).$$

It is clear that the action of $\widehat{\mathfrak{m}}$ preserves $V(-\beta_0)$ and $V(-\beta_0^\omega)$. We claim that these actions are irreducible. It follows from the definition of $\widehat{\Phi}^-(\beta_0)$ that $\lambda = -\beta_0|_{\mathfrak{h}^+}$ is a highest weight for the representation $\widehat{\mathfrak{m}}$ on $V(-\beta_0)$. Hence, $V(-\beta_0)$ contains an irreducible subrepresentation W of $\widehat{\mathfrak{m}}$ with the highest weight λ . Let us consider arbitrary

$$(5-22) \quad \mu = \lambda - \sum_{\sigma \in \widehat{\Delta}_0} n_\sigma \sigma \quad \text{with } n_\sigma \geq 0$$

which is dominant (that is, $\langle \mu, \sigma \rangle \geq 0$ for all $\sigma \in \widehat{\Delta}_0$). It follows from the Freudenthal multiplicity formula (see [9, Section 25.1]) that μ appears as a weight in W . Now

suppose that W' is another irreducible subrepresentation in $V(-\beta_0)$ with the highest weight μ . Then μ is dominant and of the form (5-22). This proves that if the representation \hat{m} on $V(-\beta_0)$ is not irreducible, then it has a weight μ of $\hat{\mathfrak{h}}^+$ which has multiplicity greater than one. Let $\beta \in \hat{\Phi}^-(\beta_0)$ be such that $\beta|_{\hat{\mathfrak{h}}^+} = \mu$. Since $\beta|_{\mathfrak{a}} = -\beta_0$, this implies that $\dim(\hat{\mathfrak{g}}_\beta) > 1$, but $\dim(\hat{\mathfrak{g}}_\beta) = 1$ for complex simple Lie algebras. This contradiction implies that the action of \hat{m} on $V(-\beta_0)$ is irreducible. The same argument implies that the action of \hat{m} on $V(-\beta_0^\omega)$ is also irreducible. It follows from (5-19) that the complex conjugation applied to (5-20) maps $V(-\beta_0)$ to $V(-\beta_0^\omega)$. This implies that $\mathfrak{g}_{-\alpha_0} \otimes \mathbb{C}$ contains no nontrivial \mathfrak{m} -invariant subspaces which are defined over \mathbb{R} . Hence, the action of \mathfrak{m} on $\mathfrak{g}_{-\alpha_0}$ is irreducible.

Now we consider the action of \mathfrak{m} on the root space $\mathfrak{g}_{-2\alpha_0}$. The proof obviously reduces to the case when \mathfrak{g} is simple. Looking through the classification of real simple Lie algebras \mathfrak{g} (see [21, Table 9]), we notice that the only case when the Lie algebra \mathfrak{g} has higher rank and $\dim(\mathfrak{g}_{-2\alpha_0}) > 1$ is $\mathfrak{g} = \mathfrak{sp}(p, q)$ with $p < q$. In this case, we check irreducibility directly. Let

$$S = \begin{pmatrix} 0 & \cdots & 1 \\ \vdots & & \vdots \\ 1 & \cdots & 0 \end{pmatrix} \quad \text{and} \quad J = \begin{pmatrix} 0 & 0 & S \\ 0 & I & 0 \\ S & 0 & 0 \end{pmatrix},$$

where S has dimension p and I is the identity matrix of dimension $q - p$. Then

$$\mathfrak{g} = \mathfrak{sp}(p, q) = \{X \in \mathfrak{sl}(p+q, \mathbb{H}) : X^*J + JX = 0\},$$

or more explicitly,

$$\mathfrak{g} = \left\{ \begin{pmatrix} X_{11} & X_{12} & X_{13} \\ X_{21} & X_{22} & -X_{12}^*J \\ X_{31} & -JX_{21}^* & -JX_{11}^*J \end{pmatrix} : \begin{array}{l} X_{22}^* + X_{22} = 0, \quad X_{31}^* = -JX_{31}J, \\ \text{tr}(X_{22}) = 0, \quad X_{13}^* = -JX_{13}J \end{array} \right\}.$$

Its Cartan subalgebra is

$$\mathfrak{a} = \{\text{diag}(a_1, \dots, a_p, 0, \dots, 0, -a_p, \dots, -a_1) : a_1, \dots, a_p \in \mathbb{R}\}.$$

The roots of the form $-2\alpha_0$ with simple α_0 are given by $-2a_1, \dots, -2a_p$, and the corresponding root spaces are

$$\mathfrak{g}_{-2a_i} = \{\text{diag}(0, \dots, 0, x_i, 0, \dots, 0)S : x_i \in \mathbb{H}, x_i^* = -x_i\}.$$

We observe that M contains a copy of $\text{SU}(2)^p$, and one of the $\text{SU}(2)$ factors acts on \mathfrak{g}_{-2a_i} as $x_i \mapsto gx_i g^*$ for $g \in \text{SU}(2)$. This representation is irreducible.

Finally suppose that the Lie algebra \mathfrak{g} is simple with real rank one and $\dim(\mathfrak{g}_{-2\alpha_0}) > 1$. There are only two cases to consider. In the first case, \mathfrak{g} is the Lie algebra of the isometry group of quaternionic hyperbolic space, the dimension of $\mathfrak{g}_{-2\alpha_0}$ is 3. Also, M again contains a copy of $\mathrm{SU}(2)$, and acts irreducibly on $\mathfrak{g}_{-2\alpha_0}$.

In the second case, \mathfrak{g} is the Lie algebra of the isometry group of Cayley hyperbolic space, which is a real form of F_4 . We observe that in this case $|\widehat{\Delta} \setminus \widehat{\Delta}_0| = 1$, so that $\beta_0|_{\mathfrak{a}} = \alpha_0$ for a unique $\beta_0 \in \widehat{\Delta}$. Hence, it follows that

$$\mathfrak{g}_{-2\alpha_0} \otimes \mathbb{C} = V(-2\beta_0).$$

This allows us to deduce irreducibility by the same argument as for $\mathfrak{g}_{-\alpha_0}$. \square

6 Completion of the proof

In this section we complete the proof of our main theorem (Theorem 1.2) using results established in the previous section. We note that it follows from [4] that given a C^0 –semiconjugacy $\psi: F = G/P \rightarrow M$, there exist a G –equivariant factor map $\pi: F \rightarrow F_I$, where $F_I = G/P_I$ is another flag manifold with $P_I \supset P$, and a C^0 –conjugacy $\phi: F_I \rightarrow M$ such that $\psi = \phi \circ \pi$. Hence, it remains to prove that the homeomorphism ϕ is smooth. The idea for the proof is that the conjugacy map $\phi: F_I \rightarrow M$ is smooth along open dense subsets of the submanifold $U_I^{(\alpha_0)}$ (see Proposition 5.1) and its translates by $\rho_0(\delta)$ for $\delta \in \Gamma$, and moreover we have smooth projections to these submanifolds in both F_I and M (see Proposition 4.6) that determine a point in some open set uniquely and smoothly—as strings of a *marionettes* puppet. We note that while the higher-rank assumption on G is absolutely crucial in Section 5, the marionettes argument presented here is applicable to general Zariski-dense subgroups.

Proof of Theorem 1.2 For $\alpha \in \Delta \setminus I$ and $\delta \in \Gamma$, we define

$$\pi_\delta^{(\alpha)} = \pi_0^{(\alpha)} \rho_0(\delta).$$

Each of these maps is defined on the open dense set $\rho_0(\delta)^{-1} U_I$ of F_I . By Proposition 5.1, the map ϕ , restricted to $U_I^{(\alpha)}$, is C^∞ on an open dense subset \mathcal{U}_α of $U_I^{(\alpha)}$. Since the maps $\pi_0^{(\alpha)}: U_I \rightarrow U_I^{(\alpha)}$ are open, the sets $\mathcal{V}_\alpha = (\pi_0^{(\alpha)})^{-1}(\mathcal{U}_\alpha)$ are open dense subsets of U_I and, hence, of F_I . By the Baire category theorem, the intersection of the sets $\delta^{-1}(\mathcal{V}_\alpha)$ for $\delta \in \Gamma$ and $\alpha \in \Delta \setminus I$ is nonempty. Let us pick a point x_0 that belongs to this intersection.

Let $\mathcal{K}^{(\alpha)}$ be the distribution on U_I defined by $\ker(D(\pi_0^{(\alpha)})_x)$, $x \in U_I$. We claim that

$$(6-1) \quad \bigcap_{\substack{\alpha \in \Delta \setminus I \\ \delta \in \Gamma}} D(\delta)_{x_0}^{-1} \mathcal{K}^{(\alpha)}(\delta \cdot x_0) = 0.$$

Suppose that (6-1) fails. Then there exists a nonzero vector $v \in T_{x_0}(F_I)$ such that

$$D(\delta)_{x_0} v \in \mathcal{K}^{(\alpha)}(\delta \cdot x_0) \quad \text{for all } \alpha \in \Delta \setminus I \text{ and } \delta \in \Gamma.$$

We observe that the projection maps $\pi_0^{(\alpha)}$ are algebraic with respect to the Lie coordinates on U_I , so that the distributions $\mathcal{K}^{(\alpha)}$ are also algebraic, and it follows from the Zariski-density of Γ that

$$D(g)_{x_0} v \in \mathcal{K}^{(\alpha)}(g \cdot x_0) \quad \text{for all } g \in G \text{ such that } g \cdot x_0 \in U_I.$$

Without loss of generality, we may assume that $x_0 = eP_I$. We recall from Lemma 4.3 that $\pi_0^{(\alpha)}$ is realised as a limit of the maps $\delta_n^{(\alpha)}$ on U_I . Hence, it follows that for every $g \in N_I^- P_I$,

$$(6-2) \quad D(\delta_n^{(\alpha)} g)_{eP} v \rightarrow 0.$$

We write $g = up$ with $u \in N_I^-$ and $p \in P_I$. Then $\delta_n^{(\alpha)} u (\delta_n^{(\alpha)})^{-1}$ converges to the identity in G . Hence, (6-2) is equivalent to

$$(6-3) \quad D(\delta_n^{(\alpha)} p)_{eP} v \rightarrow 0$$

for all $p \in P_I$. We make the identification $T_{eP}(F_I) \simeq \mathfrak{g}/\mathfrak{p}_I$, so that $D(p)_{eP} w = \text{Ad}(p)w$ for $p \in P_I$. Then $\langle \text{Ad}(P_I)v \rangle$ gives a nonzero $\text{Ad}(P_I)$ -invariant subspace of $\mathfrak{g}/\mathfrak{p}_I$ such that

$$\text{Ad}(\delta_n^{(\alpha)})w \rightarrow 0 \quad \text{for all } w \in \langle \text{Ad}(P_I)v \rangle.$$

This is equivalent to $\langle \text{Ad}(P_I)v \rangle$ being contained in $u_I^{(\alpha)} + \mathfrak{p}_I \subset \mathfrak{g}/\mathfrak{p}_I$, where $u_I^{(\alpha)}$ denotes the subspace of \mathfrak{n}_I^- spanned by all root spaces with roots not proportional to α . Since this property must hold for all $\alpha \in \Delta \setminus I$, we obtain a contradiction with Lemma 6.1 below. Hence, we conclude that (6-1) holds. Then there exist $(\alpha_1, \delta_1), \dots, (\alpha_l, \delta_l) \in (\Delta \setminus I) \times \Gamma$ such that

$$\bigcap_{i=1}^l D(\delta_i)_x^{-1} \mathcal{K}^{(\alpha_i)}(\delta_i \cdot x) = 0.$$

This property still holds in an open neighbourhood \mathcal{O} of x in F_I . Hence, we conclude that the map

$$\Pi = (\pi_{\delta_1}^{(\alpha_1)}, \dots, \pi_{\delta_l}^{(\alpha_l)}): \mathcal{O} \rightarrow \prod_{i=1}^l U_I^{(\alpha_i)}$$

is an immersion, and hence a local diffeomorphism onto its image. Similarly, we also get maps

$$\tilde{\pi}_{\delta_i}^{(\alpha_i)} = \tilde{\pi}_0^{(\alpha_i)} \rho(\delta_i): \phi(\mathcal{O}) \rightarrow \phi(U_I^{(\alpha_i)})$$

and define

$$\tilde{\Pi} = (\tilde{\pi}_{\delta_1}^{(\alpha_1)}, \dots, \tilde{\pi}_{\delta_l}^{(\alpha_l)}): \phi(\mathcal{O}) \rightarrow \prod_{i=1}^l \phi(U_I^{(\alpha_i)}).$$

We recall that by Lemma 4.7, the $\phi(U_I^{(\alpha_i)})$ are immersed submanifolds of $\phi(U_I)$. Let us consider the commutative diagram

$$\begin{array}{ccc} \mathcal{O} \subset F_I & \xrightarrow{\phi} & \phi(\mathcal{O}) \subset M \\ \Pi \downarrow & & \downarrow \tilde{\Pi} \\ \prod_{i=1}^l U_I^{(\alpha_i)} & \xrightarrow{\Phi} & \prod_{i=1}^l \phi(U_I^{(\alpha_i)}) \end{array}$$

where $\Phi = \prod_{i=1}^l \phi|_{U_I^{(\alpha_i)}}$. We observe that if the neighbourhood \mathcal{O} is sufficiently small,

$$\Pi(\mathcal{O}) \subset \prod_{i=1}^l \mathcal{U}_{\alpha_i}.$$

Since $\phi|_{\mathcal{U}_{\alpha_i}}$ is smooth and ϕ is a homeomorphism, it follows that $\phi|_{U_I^{(\alpha_i)}}$ is a local diffeomorphism on an open dense set. We conclude that

$$\phi^{-1} = \Pi^{-1} \Phi^{-1} \tilde{\Pi}$$

on a nonempty open subset of $\phi(\mathcal{O})$. This implies that ϕ is a local diffeomorphism on a nonempty open subset \mathcal{U} of F_I . We take an arbitrary $x \in \mathcal{U}$. Since Γ acts minimally on F_I (see [5]), there is $\gamma \in \Gamma$ such that $\rho_0(\gamma)x \in \mathcal{U}$. By equivariance, ϕ is C^∞ on $\rho_0(\gamma)^{-1}(\mathcal{U})$, which is an open neighbourhood of x . Furthermore, the rank of ϕ is maximal everywhere by minimality. Since ϕ is a homomorphism, this implies that ϕ is a diffeomorphism. \square

It remains to prove the following lemma, which was used in the above proof.

Lemma 6.1 *Let V be an $\text{Ad}(P_I)$ -invariant subspace of \mathfrak{g} which properly contains \mathfrak{p}_I . Then $V \cap \mathfrak{g}_\alpha \neq 0$ for some $\alpha \in \Delta \setminus I$. In particular, any nontrivial $\text{Ad}(P_I)$ -invariant subspace of $\mathfrak{g}/\mathfrak{p}_I$ is not contained in the intersection of $\mathfrak{u}_I^{(\alpha)} + \mathfrak{p}_I$ for $\alpha \in \Delta \setminus I$.*

Proof We first consider the case when the Lie algebra \mathfrak{g} is complex semisimple. Then the root spaces \mathfrak{g}_α for $\alpha \in \Phi$ are one-dimensional and $[\mathfrak{g}_{\alpha_1}, \mathfrak{g}_{\alpha_2}] = \mathfrak{g}_{\alpha_1 + \alpha_2}$ for all $\alpha_1, \alpha_2 \in \Phi$ such that $\alpha_1 + \alpha_2 \neq 0$ (see [13, Chapter III, Section 4]). Since V is $\text{Ad}(P_I)$ -invariant, it is also invariant under the action of the Cartan subalgebra contained in \mathfrak{p}_I , and it follows that V is equal to a sum of root spaces \mathfrak{g}_ρ for some $\rho \in \Phi$.

We claim that $\mathfrak{g}_{-\alpha_0} \subset V$ for some $\alpha_0 \in \Delta \setminus I$. To prove this claim, we take $\beta \in \Phi^+ \setminus \Phi_I$ such that $\mathfrak{g}_{-\beta} \subset V$, and β is “minimal” in the sense that $\mathfrak{g}_{-(\beta-\alpha)} \not\subset V$ for every $\alpha \in \Delta$ such that $\beta - \alpha \in \Phi^+ \setminus \Phi_I$. If $\beta \in \Delta$, let $\alpha_0 = \beta$. If not, then $\beta \notin \Delta$. Since V is $\text{Ad}(P_I)$ -invariant, it also follows that for every $\alpha \in \Delta$,

$$[\mathfrak{g}_\alpha, \mathfrak{g}_{-\beta}] = \mathfrak{g}_{-(\beta-\alpha)} \subset V.$$

Hence, according to our choice of β , either $\mathfrak{g}_{-(\beta-\alpha)} = 0$, ie $\beta - \alpha \notin \Phi$, or $\beta - \alpha \in \Phi_I$. We have

$$\beta \in \sum_{\alpha \in \Delta \setminus I} n_\alpha \alpha + \langle I \rangle$$

for some $n_\alpha \geq 0$ with $\sum_{\alpha \in \Delta \setminus I} n_\alpha > 0$. First, we consider the case $\sum_{\alpha \in \Delta \setminus I} n_\alpha \geq 2$. Then clearly, $\beta - \alpha \notin \Phi_I$ for all $\alpha \in \Delta$, and it follows that $[\mathfrak{g}_\alpha, \mathfrak{g}_{-\beta}] = 0$ for all simple roots α . This means that $\mathfrak{g}_{-\beta}$ consists of highest-weight vectors for the adjoint representation of \mathfrak{g} . In particular, it follows that $-\beta$ must be dominant, but this is impossible because $\beta \in \Phi^+$ (recall that all dominant roots are contained in Φ^+). Hence, we conclude that

$$\beta \in \alpha_0 + \langle I \rangle$$

for some $\alpha_0 \in \Delta \setminus I$. Then $\beta - \alpha \notin \Phi_I$ for all $\alpha \in \Delta \setminus \{\alpha_0\}$, and it follows as before that $[\mathfrak{g}_\alpha, \mathfrak{g}_{-\beta}] = 0$ when $\alpha \in \Delta \setminus \{\alpha_0\}$. If we also had that $[\mathfrak{g}_{\alpha_0}, \mathfrak{g}_{-\beta}] = 0$, then $\mathfrak{g}_{-\beta}$ would have consisted of the highest-weight vectors of the adjoint representation. Then we would get a contradiction as before. Therefore, since we assumed that $\beta \notin \Delta$, we conclude that

$$[\mathfrak{g}_{\alpha_0}, \mathfrak{g}_{-\beta}] = \mathfrak{g}_{-(\beta-\alpha_0)} \neq 0,$$

so that $\beta - \alpha_0$ is a root, and since $\beta \in \Phi^+$, we have $\beta - \alpha_0 \in \Phi^+$ too. Then, since V is $\text{Ad}(P_I)$ -invariant,

$$[\mathfrak{g}_{\beta-\alpha_0}, \mathfrak{g}_{-\beta}] = \mathfrak{g}_{-\alpha_0} \subset V.$$

This proves the lemma when \mathfrak{g} is a complex semisimple Lie algebra.

To treat the general case, we consider the complex semisimple Lie algebra $\hat{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}$. It contains (see [21, Chapter 5, Section 4]) a Cartan subalgebra $\hat{\mathfrak{a}}$ of the form $\hat{\mathfrak{a}} = (\mathfrak{h}^+ \oplus \mathfrak{a}) \otimes \mathbb{C}$, where \mathfrak{h}^+ is a Cartan subalgebra of \mathfrak{m} such that the root system $\hat{\Phi}$ and the system of simple roots $\hat{\Delta} \subset \hat{\Phi}$ associated to $\hat{\mathfrak{a}}$ satisfy

$$\{\alpha|_{\mathfrak{a}} : \alpha \in \hat{\Delta}\} = \Delta \cup \{0\}$$

and

$$\mathfrak{g}_\beta \otimes \mathbb{C} = \bigoplus_{\substack{\rho \in \hat{\Phi} \\ \rho|_{\mathfrak{a}} = \beta}} \hat{\mathfrak{g}}_\rho \quad \text{for all } \beta \in \Phi.$$

We set

$$\hat{I} = \{\alpha \in \hat{\Delta} : \alpha|_{\mathfrak{a}} \in I \cup \{0\}\}.$$

Then $\mathfrak{p}_I \otimes \mathbb{C} = \hat{\mathfrak{p}}_{\hat{I}}$. The previous discussion implies that $\hat{\mathfrak{g}}_{-\alpha} \subset V \otimes \mathbb{C}$ for some $\alpha \in \hat{\Delta} \setminus \hat{I}$, and hence $(\mathfrak{g}_{-\beta} \otimes \mathbb{C}) \cap (V \otimes \mathbb{C}) \neq 0$ for $\beta \in \Delta \setminus I$ such that $\beta = \alpha|_{\mathfrak{a}}$. This also implies that $\mathfrak{g}_{-\beta} \cap V \neq 0$ and proves the lemma. \square

7 On topological sinks

A key assumption for our main result (Theorem 1.2) is that the lattice action on the manifold M has at least one differentiable sink. While the existence of a topological sink is clear for continuous factors of projective actions, we note that a topological sink of a lattice action is not always a differentiable sink. In this section, we give an example of a smooth lattice action on a circle bundle over a flag manifold to illustrate this.

Let f_t be a flow on the circle $S^1 \simeq \mathbb{R}/\mathbb{Z}$ such that $0 \in \mathbb{R}/\mathbb{Z}$ is a topological sink for the maps f_t for $t > 0$, but it is not a differentiable sink. For example, we could take the flow for the vector field on \mathbb{R}/\mathbb{Z} given in local coordinates around 0 by $x^3 - x$ for $x \in \mathbb{R}$. Let G be a noncompact connected semisimple Lie group and Γ a lattice subgroup of G . We choose a Cartan subgroup A of G such that $A \cap \Gamma$ is a lattice in A . Such a subgroup exists by [22]. We fix a set Δ of simple roots and a root $\alpha \in \Delta$. This determines a minimal parabolic subgroup P containing A . The root α gives a homomorphism $\log(\alpha) : P \rightarrow \mathbb{R}$. Since $A \cap \Gamma$ is a lattice in A , there exists $\gamma_0 \in A \cap \Gamma$ such that $\alpha(\gamma_0) > 1$ for all $\alpha \in \Delta$. Then eP is a smooth sink for the action of γ on G/P . We consider the equivalence relation on $G \times S^1$ defined by

$$(g, x) \sim (gp^{-1}, f_{\log(\alpha)(p)}(x)), \quad g \in G, p \in P, x \in S^1.$$

Then

$$M = (G \times S^1)/\sim$$

is a manifold equipped with the smooth Γ -action

$$[g, x] \mapsto [\gamma g, x], \quad [g, x] \in M, \gamma \in \Gamma.$$

It is clear $[e, 0] \in M$ is fixed by γ_0 . Moreover, for $v \in \mathfrak{n}^-$ and $x \in S^1$,

$$\gamma_0 \cdot [\exp(v), x] = [\exp(\text{Ad}(\gamma_0)v), \phi_{\log(\alpha)(\gamma_0)}(x)].$$

Hence, $[e, 0]$ is a topological sink of γ_0 , but it is not a differentiable sink.

References

- [1] **Y Benoist**, *Propriétés asymptotiques des groupes linéaires*, Geom. Funct. Anal. 7 (1997) 1–47 MR
- [2] **A Borel**, *Linear algebraic groups*, 2nd edition, Graduate Texts in Mathematics 126, Springer (1991) MR
- [3] **A Brown, F Rodriguez Hertz, Z Wang**, *Invariant measures and measurable projective factors for actions of higher-rank lattices on manifolds*, preprint (2016) arXiv
- [4] **S G Dani**, *Continuous equivariant images of lattice-actions on boundaries*, Ann. of Math. 119 (1984) 111–119 MR
- [5] **S G Dani**, *Orbits of horospherical flows*, Duke Math. J. 53 (1986) 177–188 MR
- [6] **D Fisher**, *Groups acting on manifolds: around the Zimmer program*, from “Geometry, rigidity, and group actions” (B Farb, D Fisher, editors), Univ. Chicago Press (2011) 72–157 MR
- [7] **D Fisher, B Kalinin, R Spatzier**, *Global rigidity of higher rank Anosov actions on tori and nilmanifolds*, J. Amer. Math. Soc. 26 (2013) 167–198 MR
- [8] **D Fisher, G Margulis**, *Local rigidity of affine actions of higher rank groups and lattices*, Ann. of Math. 170 (2009) 67–122 MR
- [9] **W Fulton, J Harris**, *Representation theory: a first course*, Graduate Texts in Mathematics 129, Springer (1991) MR
- [10] **E Ghys**, *Rigidité différentiable des groupes fuchsiens*, Inst. Hautes Études Sci. Publ. Math. 78 (1993) 163–185 MR
- [11] **M Guysinsky**, *The theory of non-stationary normal forms*, Ergodic Theory Dynam. Systems 22 (2002) 845–862 MR
- [12] **M Guysinsky, A Katok**, *Normal forms and invariant geometric structures for dynamical systems with invariant contracting foliations*, Math. Res. Lett. 5 (1998) 149–163 MR
- [13] **S Helgason**, *Differential geometry, Lie groups, and symmetric spaces*, Graduate Studies in Mathematics 34, Amer. Math. Soc., Providence, RI (2001) MR
- [14] **N V Ivanov**, *Action of Möbius transformations on homeomorphisms: stability and rigidity*, Geom. Funct. Anal. 6 (1996) 79–119 MR

- [15] **M Kanai**, *A new approach to the rigidity of discrete group actions*, *Geom. Funct. Anal.* 6 (1996) 943–1056 MR
- [16] **A Katok, R J Spatzier**, *Differential rigidity of Anosov actions of higher rank abelian groups and algebraic lattice actions*, *Tr. Mat. Inst. Steklova* 216 (1997) 292–319 MR
Reprinted in *Proc. Steklov Inst. Math.* 216 (1997) 287–314
- [17] **D Y Kleinbock, G A Margulis**, *Bounded orbits of nonquasiunipotent flows on homogeneous spaces*, from “Sinaĭ’s Moscow Seminar on Dynamical Systems” (L A Bunimovich, B M Gurevich, Y B Pesin, editors), *Amer. Math. Soc. Transl. Ser. 2* 171, Amer. Math. Soc., Providence, RI (1996) 141–172 MR
- [18] **G A Margulis**, *Finiteness of quotient groups of discrete subgroups*, *Funktional. Anal. i Prilozhen.* 13 (1979) 28–39 MR In Russian; translated in *Funct. Anal. Appl.* 13 (1979) 178–187
- [19] **G A Margulis**, *Discrete subgroups of semisimple Lie groups*, *Ergeb. Math. Grenzgeb.* 17, Springer (1991) MR
- [20] **D Montgomery, L Zippin**, *Topological transformation groups*, Interscience, New York (1955) MR
- [21] **A L Onishchik, E B Vinberg**, *Lie groups and algebraic groups*, Springer (1990) MR
- [22] **G Prasad, M S Raghunathan**, *Cartan subgroups and lattices in semi-simple groups*, *Ann. of Math.* 96 (1972) 296–317 MR
- [23] **G Prasad, A S Rapinchuk**, *Existence of irreducible \mathbb{R} –regular elements in Zariski-dense subgroups*, *Math. Res. Lett.* 10 (2003) 21–32 MR
- [24] **F Rodriguez Hertz, Z Wang**, *Global rigidity of higher rank abelian Anosov algebraic actions*, *Invent. Math.* 198 (2014) 165–209 MR
- [25] **R J Spatzier**, *On lattices acting on boundaries of semisimple groups*, *Ergodic Theory Dynamical Systems* 1 (1981) 489–494 MR
- [26] **P Tukia**, *Differentiability and rigidity of Möbius groups*, *Invent. Math.* 82 (1985) 557–578 MR
- [27] **R J Zimmer**, *Equivariant images of projective space under the action of $\mathrm{SL}(n, \mathbb{Z})$* , *Ergodic Theory Dynamical Systems* 1 (1981) 519–522 MR
- [28] **R J Zimmer**, *Ergodic theory and semisimple groups*, Monographs in Mathematics 81, Birkhäuser, Basel (1984) MR

Department of Mathematics, University of Bristol
Bristol, United Kingdom

Department of Mathematics, University of Michigan
Ann Arbor, MI, United States

a.gorodnik@bristol.ac.uk, spatzier@umich.edu

Proposed: Benson Farb

Seconded: Anna Wienhard, Leonid Polterovich

Received: 22 September 2016

Accepted: 15 May 2017