

**Metabolomic response of early-stage wheat (*Triticum aestivum*) to surfactant-aided foliar application
of copper hydroxide and molybdenum trioxide nanoparticles**

Xiangning Huang[†] and Arturo A. Keller^{†}*

[†] Bren School of Environmental Science and Management, University of California at Santa Barbara, CA,
USA 93106

^{*} University of California Center for Environmental Implications of Nanotechnology, Santa Barbara, CA,
USA 93106

Abstract

Surfactants are commonly used in foliar applications to enhance interactions of active ingredients with plant leaves. We employed metabolomics to understand the effects of Triton™ X-100 surfactant (SA) and nanomaterials (NMs) on wheat (*Triticum aestivum*) at the molecular level. Leaves of three-week-old wheat seedlings were exposed to deionized water (DI), surfactant solution (SA), NMs-surfactant suspensions (Cu(OH)₂ NMs and MoO₃ NMs), and ionic-surfactant solutions (Cu IONs and Mo IONs). Wheat leaves and roots were evaluated via physiological, nutrient distribution, and targeted metabolomics analyses. SA had no impact on plant physiological parameters, however, 30+ dysregulated metabolites and 15+ perturbed metabolomic pathways were identified in wheat leaves and roots. Cu(OH)₂ NMs resulted in accumulation of 649.8 µg/g Cu in leaves; even with minimal Cu translocation, levels of 27 metabolites were significantly changed in roots. Due to the low dissolution of Cu(OH)₂ NMs in SA, the low concentration of Cu IONs induced minimal plant response. In contrast, given the substantial dissolution of MoO₃ NMs (35.8%), the corresponding high levels of Mo IONs resulted in significant metabolite reprogramming (30+ metabolites dysregulated). Aspartic acid, proline, chlorogenic acid, adenosine, ascorbic acid, phenylalanine, and lysine were significantly upregulated for MoO₃ NMs, yet downregulated under Mo IONs condition. Surprisingly, Cu(OH)₂ NMs stimulated wheat plant tissues more than MoO₃

NMs. The glyoxylate/dicarboxylate metabolism (in leaves) and valine/leucine/isoleucine biosynthesis (in roots) uniquely responded to $\text{Cu}(\text{OH})_2$ NMs. Findings from this study provide novel insights on the use of surfactants to enhance the foliar application of nanoagrochemicals.

Keywords: surfactants, nanomaterials, nanoagrochemicals, metabolomics, wheat

Synopsis

Surfactants enhance the delivery of NMs via foliar exposure, with minimal physiological effects. However, surfactants do elicit metabolic reprogramming. NM size, zeta potential, dissolution rate, adhesion of surfactant and NMs to plant leaves, and translocation are all important factors during NMs foliar applications. Plant molecular responses to surfactant-aided NM application should be taken into account to maximize the benefit of nanoagrochemicals and reduce negative effects.

Introduction

Nanosized materials (NMs) are increasingly being considered to improve agricultural sustainability. Nanofertilizers (e.g., Ca, Fe_2O_3 , MgO, P, CuO, MnO, MoS_2 , and ZnO)¹⁻³ and nanopesticides (e.g., Ag, single-walled carbon nanotubes (SWCNTs), CeO_2 , $\text{Cu}(\text{OH})_2$, MgO, $\text{Mg}(\text{OH})_2$, SiO_2 , and TiO_2)^{2, 4-6} have been investigated in previous studies. Some NMs (e.g., TiO_2 , SiO_2 , and SWCNTs) have also been considered to alleviate salt/drought stress or enhance plant photosynthesis.⁷⁻⁹ Due to interactions between NMs or their dissolved ions with soil constituents and root exudates,^{10, 11} they are not fully taken up by the plant from the soil, resulting in inefficient delivery of the active ingredient(s). Foliar delivery provides an efficient and scalable approach for direct interaction between NMs and plants.¹² However, the aggregation and dissolution of NMs, environmental conditions (e.g., rain and wind), as well as various leaf surface characteristics can substantially affect the efficiency of foliar applications.^{13, 14} In particular, the abundant

cuticular waxes and trichomes (hairs) on wheat (*Triticum aestivum*) leaf surfaces make them extremely difficult to wet, with little or no water adhesion.¹⁵ To improve the efficiency of foliar application, adjuvants (e.g., humectants, oils, pH buffers, and surfactants) are often employed to improve the wettability of leaf surfaces and prevent off-target drift.¹⁶ We hypothesize that the combination of nanoagrochemicals with surfactants can improve the delivery of the active ingredients via foliar applications, particularly for plants with hydrophobic or superhydrophobic leaf surface properties. The main objectives of this study were to (1) employ a surfactant to improve delivery of NMs to plants; (2) determine whether the surfactant had any effect on the perturbation of target crop metabolic pathways; (3) evaluate the delivery of nano-scale active ingredients through foliar application; and (4) determine the metabolic response of plants to surfactant-enhanced NMs foliar exposure.

Surfactants are often used to lower the surface tension between droplets and leaf surfaces, as well as to improve penetration of active ingredients through plant cuticles.¹⁷ Surfactants are usually classified as anionic, cationic, non-ionic, and amphoteric.¹⁸ Non-ionic surfactants are widely used in many agrochemical formulations.¹⁷ In the current work, Triton™ 100-X (polyethylene glycol tert-octylphenyl ether, a non-ionic surfactant) was added into NMs suspensions during the wheat foliar applications. Past work has shown that Triton™ 100-X not only enhances the wettability of wheat leaf surfaces, but also promotes nutrient uptake through foliar applications.^{15, 19, 20} Compared with two other surfactants (sodium dodecyl-sulfate and dodecyltrimethylammonium bromide), Triton™ 100-X solution ($>1 \times 10^{-5}$ mol L⁻¹) resulted in pronounced wettability and considerably decreased the contact angle ($<90^\circ$) of water drops on wheat leaves.²⁰ In another study, two commercial surfactants (3 g/L LI 700® and 1 g/L Agral®), two pure surfactants (1 g/L Genapol® X-080 and 1 g/L Triton™ 100-X), and one humectant (1 g/L glycerol) were employed to assist phosphorous (P) translocation in wheat through foliar applications.¹⁵ Except for glycerol, P uptake was greater than 70% for all surfactants; Triton™ 100-X resulted in 82.4% - 83.5% increased P uptake during early tillering and flag leaf emergence periods. Past agricultural surfactant-

related studies have mostly focused on interactions between surfactant and leaf surfaces,¹⁹⁻²¹ the development of new surfactants formulations,^{17, 22} and applications of surfactants with conventional fertilizers/pesticides.^{15, 23-25} Few studies have considered the use of surfactant during foliar applications of NMs.^{26, 27}

Many factors can affect the effectiveness of NMs foliar application processes, including NM properties (e.g., size,²⁸⁻³⁰ concentration,³¹⁻³³ zeta potential,³⁰ response duration,³¹ and coating^{29, 30}), leaf surface characteristics (e.g., wax thickness, pores, trichomes, and epidermis maturity),^{34, 35} NM uptake pathways (i.e., stomata^{32, 36, 37} vs cuticle^{29, 30}), and environmental factors.^{2,27-29} For example, Avellan et al., (2019) investigated the effects of Au NMs sizes and coatings for wheat foliar exposure without the assistance of surfactants.³⁰ Regardless of tested conditions, smaller Au NMs (3 nm) had a stronger adhesion to leaf surfaces, and 10-25% of the Au was transported to wheat roots after the rinsing process. Au NMs coated with polyvinylpyrrolidone (PVP) appeared in the mesophyll (cuticular pathway) and moved further through the plant vasculature. However, similar-sized Au NMs with a citrate coating were not present. In another short-term exposure study (24 h), wheat leaves exposed to zinc salts containing 0.05 wt% of Tween20 resulted in about six times greater Zn content in wheat leaves than the ZnO NMs-Tween20 foliar exposure. However, no significantly Zn content differences were found among ZnO NMs-Tween20 suspensions with different coatings.²⁹ The authors proposed that ionic adsorption was likely the dominant mechanism for passing through the cuticular pathway. Most of above-mentioned studies either did not include surfactants, or simply employed surfactant as the wetting agent, without analyzing cellular level plant response. To date, little is known about the metabolomics of applying surfactants on crop plants.

Metabolomics is increasingly employed to study plant molecular responses/mechanisms to different stimulus (e.g., NMs).^{4, 39-50} Given the extended use of the commercial nanopesticide Kocide 3000 (active ingredient: Cu(OH)₂ NMs), the effects of Cu(OH)₂ NMs on basil (*Ocimum basilicum*),⁴ cucumber (*Cucumis Sativus*),^{44, 46} maize (*Zea mays*),^{44, 45} and lettuce (*Lactuca Sativa*)⁴⁷ have been explored. For example, Zhao

et al., (2017) conducted a one-week $\text{Cu}(\text{OH})_2$ NMs foliar spray on corn (*Zea mays*) that systematically studied plant cellular responses.^{44,45} The general plant response was dose-dependent, with a higher dose (100 mg per plant) significantly upregulating 4-hydroxycinnamic acid (2.22 fold), myo-inositol (1.50 fold), tyrosine (1.39 fold), phenylalanine (1.24 fold), and total phenolic content (1.17 fold) in maize leaves.⁴⁵ The biological pathway analysis revealed that the most significantly disturbed pathway was inositol phosphate metabolism.⁴⁴

In the current study, the molecular response to foliar application of MoO_3 NMs was also evaluated. Molybdenum (Mo) is an essential micronutrient, often involved in reductive and oxidative reactions via specific plant enzymes.⁵¹ Mo also plays an important role in N fixation, nitrate reduction, and amino acid and protein biosynthesis processes.⁵² Excess exposure to Mo NMs and Mo^{6+} can inhibit root growth/elongation, prolong seed germination, increase nitrate reductase, and cause oxidative imbalance.⁵³⁻⁵⁷ Recently we demonstrated that corn and wheat exposed to MoO_3 NMs through a three-week root exposure (at 200 and 1000 mg /kg Mo levels) resulted in a significant response in plant roots and leaves at the molecular level.⁴⁹ Although the physiological data showed that the MoO_3 NMs had a more severe impact on corn roots and above-ground plant tissues, 53 dysregulated metabolites were found in wheat leaves compared with 21 in corn leaves. Furthermore, TCA cycle, amino acid metabolism, and pyrimidines metabolism were perturbed in wheat leaves, but not in corn. To our knowledge, there have been no studies regarding surfactant-enhanced foliar application of $\text{Cu}(\text{OH})_2/\text{MoO}_3$ NMs and the subsequent metabolomic alterations. Furthermore, due to the natural loss of NMs during the spraying process, to date, most foliar application studies have been qualitative rather than quantitative.

In this study, three-week-old wheat seedlings were exposed to deionized water (DI) only, surfactant only solution, NM-surfactant suspensions, and ionic-surfactant solutions through one-week foliar exposures. Wheat was selected since it is an important global crop, which requires crop protection from pests. Wheat responses to the surfactant (without active ingredients) were evaluated separately in order

to determine how the surfactant would affect plant response at the molecular level. At the end of the exposure duration, plants were harvested and separated into wheat roots/leaves and changes in physiological parameters (e.g., dry biomass) were recorded. Target metals (e.g., Cu and Mo), along with other macro and micronutrient levels were analyzed to track metal translocation and distribution changes. Targeted metabolites, including amino acids, antioxidants, fatty acids, nucleobase/side/tide, organic acids/phenolics, and sugar/alcohols were measured and dysregulated metabolites were identified under each exposure condition. Furthermore, a perturbed metabolic pathways analysis was also conducted and the distinct effects of the two NMs (e.g., $\text{Cu}(\text{OH})_2$ NMs and MoO_3 NMs) on wheat growth was evaluated. This work sheds insights into the effects of a surfactant on plant growth and the importance of enhancing foliar application conditions for NMs.

2. Materials and methods

2.1. Characterization and stability of $\text{Cu}(\text{OH})_2$ and MoO_3 NMs

$\text{Cu}(\text{OH})_2$ NMs (99.5% purity, US3078) and MoO_3 NMs (99.94% purity, US3330) were purchased from U.S. Research Nanomaterials Inc. (U.S.). The original powders have been fully characterized elsewhere¹¹. Briefly, the $\text{Cu}(\text{OH})_2$ NMs are nanowires (diameter: 50 nm and length: 2-5 μm) and the MoO_3 NMs are spheres (diameter: 13-80 nm) (**Figure S1**). Both ENMs have an orthorhombic crystalline structure as confirmed by X-ray diffraction (XRD) analysis, and no carbon-based coatings were detected based on X-ray photoelectron spectrometry (XPS) measurements.

TritonTM X-100 (BioXtra, T9284) was obtained from Sigma Aldrich. Based on the information provided by the manufacturer, the critical micelle concentration (CMC) is 0.2-0.9 mM (20-25 °C). A droplet dispersion test was conducted by depositing a 5 μL droplet of TritonTM X-100 solutions on three-week-old wheat leaf surfaces (**Figure S2**). The optimum application level was 200 mg/L (0.32 mM), with adequate

droplet spreading without dripping off. The applied concentration of Triton™ X-100 solution in the current work is lower than previous foliar application studies, where 500 - 2000 mg/L of Triton™ X-100 were employed.^{15, 23, 26} The influence of the surfactant on NM zeta potential, hydrodynamic diameter, and dissolution rate was measured for each NM.^{11, 58} The tests were conducted in 50 mL metal-free polypropylene tubes that contained 200 mg/L Triton™ X-100 with 100 mg/L Cu(OH)₂ or 100 mg/L MoO₃ NMs (as metal content) suspensions. After spiking the corresponding amount of NMs, the tubes were sonicated for 20 min and the hydrodynamic diameter and surface charge (zeta potential) were measured via dynamic light scattering (Zetasizer Nano ZS, Malvern). For the dissolution test, 2 ml samples were withdrawn at 0 hr and 5 hr, and centrifuged in Amicon Ultra 3 kDa MWCO tubes (Sigma-Aldrich) at 3000 rpm for 30 min.^{11, 59} Then the filtrate was acidified to 10 ml with 2% HNO₃ for inductively coupled plasma-mass spectrometry (ICP-MS) (Agilent 7900, Agilent Technologies) analysis. Three replicates were utilized for all analyses.

2.2. Wheat growth and leaf exposure assay

Wheat (*Triticum aestivum* 'Red Fife') seeds were selected for the current study. Before germination, seeds were immersed in a 1% sodium hypochlorite solution for 10 min, rinsed well with deionized water, and soaked for 24 h. Four seeds were germinated in each pot (40 g vermiculite/pot) containing water-saturated vermiculite. Five days later, seedlings were transplanted to a new pot with 2 plants per pot. Growth conditions were 150 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ light intensity (as daylight white) for 16 h daily, the temperature was set at 22°C, and relative humidity was 60%. A 10% Hoagland solution was used for watering purposes to ensure adequate nutrients for plant growth.⁴⁹ Water content in the pots was maintained between 70-90%.

Foliar exposure was initiated at the beginning of the fourth week and the exposure duration was 7 days. A preliminary test showed that without adding a surfactant, immersing wheat leaves in a 100 mg/L

MoO₃ NMs suspension resulted in little Mo accumulated on wheat leaves, with minimal translocation to roots (**Figure S3**). This finding confirmed the crucial role of surfactant during the wheat foliar application process.^{15, 19, 20} For the leaf exposure test, the second and third true leaves were exposed to metal-surfactant suspensions (two times per day) using a multichannel micropipette (2.54 ± 0.30 µl/drop, eight channels). The treatments were as follows: *DI* - deionized water only; *SA* - 200 mg/L Triton™ X-100 solution; *Cu(OH)₂ NMs* - 100 mg/L Cu(OH)₂ NMs (as Cu content) in *SA*; *Cu IONs* - 0.1 mg/L CuSO₄ · 5H₂O (as Cu content) in *SA*; *MoO₃ NMs* - 100 mg/L MoO₃ NMs (as Mo content) in *SA*; and *Mo IONs* - 35 mg/L Na₂MoO₄ · 2H₂O (as Mo content) in *SA*. Eight replicates were employed for each experimental condition. At the end of the exposure period, all plants were harvested, well rinsed,¹⁴ freeze-dried, and weighted. All freeze-dried plant tissues were stored at -80 °C for further analyses.

2.3. Plant tissue digestions and nutrient measurements

Freeze-dried plant tissues (i.e., roots and leaves) were cut into small pieces and transferred into 50 ml digestion tubes. The digestion process was conducted in an SCP Science DigiPREP hot block digestion system. First, 2 ml of plasma pure HNO₃ was added into the tube and heated at 115°C for 20 min. Then 8 ml of H₂O₂ reagent was added into the system and held at 115°C for another 60 min.^{45, 49} When the digestion process was complete, all samples were diluted to the 50 ml mark for nutrient analysis. In addition to the targeted metals (i.e., Cu and Mo), four macronutrients (Ca, K, Mg, and P) and three additional micronutrients (Fe, Mn, and Zn) were quantified through ICP-MS analysis. The dried biomass and nutrient data were analyzed by using the statistical package SPSS (Social Sciences, 22.0). More specifically, a one-way analysis of variance (ANOVA) was performed followed by a Tukey-HSD test (*p* value = 0.05).

2.4. Plant tissue extractions and targeted metabolites measurements

A total of 82 targeted metabolites (**Table S1**) were selected from previous studies, which showed good sensitivities during plant (corn, cucumber, soybean, and wheat) exposures to various NMs (Ag, Cd, Cu, and Mo).^{49, 59-62} Briefly, the freeze-dried plant samples were finely ground by using pestle and mortar with liquid nitrogen. Then around 10 mg of sample was weighed and transferred into a 2ml Eppendorf microcentrifuge tube that containing 1.2 mL of 80% methanol and 2% formic acid.^{60, 61} The tubes were first vortexed for 20 min, then sonicated for 20 min, and finally centrifuged at 2×10^4 g for another 20 min. The supernatant was analyzed through liquid chromatography - triple quadrupole mass spectrometry (LC-MS/MS) analysis (Agilent 1260) (see SI for more details).

Metabolite data was log-transformed and auto-scaled before further analyses using MetaboAnalyst 5.0. For the statistical analysis, a one-way ANOVA followed by a Fisher's least significant difference method (Fisher's LSD, p value set as 0.05) was conducted to identify the significantly altered metabolites. In addition, the PLS-DA method was applied. PLS-DA has often been used to maximize the metabolite cluster differences between treatment groups from the previous metabolomics studies.^{39, 41, 44, 59} From the PLS-DA test results, variables with importance in the projection (VIP) values greater than 1 were also considered as the featured metabolites. For the metabolomic pathway analysis, the perturbed pathways with an impact value threshold greater than 0.1 (p value threshold was 0.05) were considered.⁴¹

Results and discussion

NM characterization and dissolution

$\text{Cu}(\text{OH})_2$ and MoO_3 NMs were characterized in the 200 mg/L TritonTM X-100 suspension (SA) in terms of hydrodynamic diameter, zeta potential (ξ), and dissolution rate. Compared with their behavior in DI water, placing the NMs in SA resulted in slightly larger (but not significant) hydrodynamic diameters (**Table S2**). However, in SA ξ increased significantly for $\text{Cu}(\text{OH})_2$ NMs (4.84 mV to 15.9 mV) and MoO_3 NMs (-65.0 mV to -51.9 mV) compared to in DI water. In general, the addition of surfactant accelerated NM

dissolution (**Figure 1 (A) and (B)**). More specifically, the overall dissolved amount of $\text{Cu}(\text{OH})_2$ NMs was quite small (0.1%) in SA. In contrast, 35.8% of MoO_3 NMs was dissolved in SA with 33.5% more Mo^{6+} ions released after 5 hr. This demonstrates the importance of refreshing NM-SA suspensions after each foliar exposure to maintain a similar ionic composition. Based on the NM dissolution results, the corresponding ionic exposure doses were determined as 0.1 mg/L of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ as Cu (Cu IONS) and 35 mg/L of $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$ as Mo (Mo IONS).




Figure 1. Metal ions released from (A) 100 mg/L $\text{Cu}(\text{OH})_2$ NMs and (B) 100 mg/L MoO_3 NMs in DI water (DI) and surfactant containing suspensions (SA). PLS-DA score plots of overall metabolites in wheat (C) leaves and (D) roots after one-week foliar exposure to DI and SA solutions. Three replicates under each condition.

Plant responses to the surfactant solution

We examined the wheat responses to foliar application of 200 mg/L Triton™ X-100 solutions (SA) by quantifying the plant dry biomass, nutrient distribution, and metabolomic profile alteration. Compared with DI water foliar application, the surfactant solution did not significantly change the dry biomass, nor the majority of nutrient content (i.e., Mg, P, K, Ca, Mn, Fe, Cu, and Zn) in wheat (**Figure S4 and Table S3**). However, a significant decrease in Mo content (59.1%) was observed in wheat leaves. Mo is an essential component of nitrate reductase and N-fixing enzyme nitrogenase; decreasing Mo could potentially impact N metabolism in wheat leaves.⁶³ In addition, the metabolomic profile showed a clear separation between DI and SA treatments, which could be explained by the total variance along component 1 in wheat leaves (77.8%) and roots (71.8%) (**Figure 1 (C) and (D)**).

When wheat leaves were exposed to SA, 37 metabolites were significantly altered in wheat leaves and 34 dysregulated metabolites were discovered in wheat roots (**Table S4**). Furthermore, more than 85% of these dysregulated metabolites were significantly upregulated (with fold change >2) (**Figure 2**). The most significantly altered metabolites (with fold change >5) were certain amino acids (leucine, lysine, phenylalanine, proline, serine, and tryptophan), fatty acids (linolenic acid), nucleobase/side/tide (adenosine, guanosine, and uridine), and sugar/sugar alcohols (trehalose). Amino acids are synthesized from TCA cycle intermediates; the increased levels indicate some favoring of N metabolism over C metabolism.⁴⁴ As one of the most abundant fatty acids in plant membrane lipids, the increased concentration of linolenic acid suggests potential membrane lipid peroxidation as a defense mechanism to abiotic stress.⁶⁴ The significant changes in the abundance of adenosine, guanosine, and uridine indicate the perturbation of pyrimidine and purine metabolism. Even though the role of trehalose remains unclear, a study has shown that this carbohydrate may serve to preserve the plant's cellular integrity under stress conditions.⁶⁵ Foliar exposure to SA resulted in 17 perturbed metabolomic pathways in wheat leaves and

15 metabolomic pathways were significantly altered in roots (**Table S5**). Arginine biosynthesis and arginine/proline metabolism were only disturbed in wheat leaves, which involve the uniquely dysregulated metabolite from wheat leaves - glutamate. Glutamate plays a central role in the synthesis of γ -aminobutyric acid (GABA), arginine, and proline, and regulates ammonium assimilation.⁶⁶ The increased glutamate level may indicate a response to excess ROS.⁶⁰ On the other hand, pyruvic acid was only significantly downregulated in wheat roots (with fold change < 0.5), which contributed to the perturbation of valine, leucine and isoleucine metabolism. As a precursor for valine, leucine, and isoleucine,⁶⁷ pyruvic acid is also used as a substrate for the TCA cycle. The depletion of pyruvic acid indicates possible redistribution of N and C metabolism.




Figure 2. Significant metabolic pathway changes in wheat after one-week foliar exposure to surfactant solution. The color scale indicates the fold changes compared with DI water. The border of the box indicates whether the metabolite changes happened in wheat leaves (green) or roots (red).

Wheat leaf responses to metal-surfactant suspensions

With the assistance of the surfactant (200 mg/L TritonTM X-100), the NMs and ionic salts were able to adhere to wheat leaf surfaces more effectively, resulting in a significant increase in target metals (i.e., Cu and Mo) found on wheat leaves (**Figure S6 (A) and (B)**). Overall, there were no significant changes in wheat leaf dry biomass for all treatments (**Figure S5 (A)**). Foliar exposure to Cu(OH)₂ NMs resulted in an average of 649.8 µg/g Cu detected in wheat leaves, which is well above the Cu content found in other treatments. In addition, exposure to Cu(OH)₂ NMs also significantly increased Zn content by almost 54% in wheat leaves (**Table S6**). Previous Cu(OH)₂ NMs foliar exposure-related studies often found no significant changes in Zn content,^{4, 45, 46} however, one study did observe increased Zn in vascular tissues of lettuce (*Lactuca sativa*).⁴⁷ Zn is an essential micronutrient for plant growth and is involved in several essential processes. Plants with high Zn efficiency exhibit healthy growth status and high yield.⁶⁸ Exposure to Cu IONs had a minimal effect due to the low applied Cu²⁺ concentration. Contrary to the copper-surfactant foliar exposure results, higher Mo loading was found in wheat leaves that were exposed to Mo IONs (104.5 µg/g Mo) than those exposed to MoO₃ NMs (72.3 µg/g Mo). The properties of the NMs (i.e., surface charge, surface coating, and hydrophobicity) and plant leaf surface characteristics (i.e., trichomes and stomata) can both influence NM-leaf surface interactions and uptake.³⁴




Figure 3. Comparison of significantly changed metabolites ($|\log_2(\text{fold change})| > 1$) in wheat leaves exposed to (A) $\text{Cu}(\text{OH})_2$ NMs; (B) MoO_3 NMs; (C) Cu IONs; and (D) Mo IONs. Experimental conditions: SA - 200 mg/L TritonTM X-100 solution; $\text{Cu}(\text{OH})_2$ NMs - 100 mg/L $\text{Cu}(\text{OH})_2$ NMs (as Cu content) in SA; Cu IONs - 0.1 mg/L $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ (as Cu content) in SA; MoO_3 NMs - 100 mg/L MoO_3 NMs (as Mo content) in SA; and Mo IONs - 35 mg/L $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$ (as Mo content) in SA. The numbers in the pie chart represent the significantly changed metabolites from each metabolite category. The color of the numbers in the outer circular ring indicates fold changes of the significantly dysregulated metabolites: red (fold change > 5), orange (2 < fold change < 5), blue (0.2 < fold change < 0.5), and purple (fold change < 0.2). Three replicates under each condition.

The overall metabolite profiles in wheat leaves showed a clear separation between surfactant and metal-surfactant foliar exposures (**Figure S6 (C) and (D)**). Foliar exposures to Cu(OH)₂ NMs and Mo IONs resulted in the most significant changes of metabolite levels in wheat leaves, followed by the MoO₃ NMs; exposure to Cu IONs had the least impact (**Figure 3 (A)-(D), Table S7**). These findings correspond well with the metal accumulation results on wheat leaves (**Figure S6 (A) and (B)**), reflecting the importance of retaining metals on leaf surfaces to elicit plant's internal response. In wheat leaves, a total of 34 dysregulated metabolites were identified for the Cu(OH)₂ NMs and 16 for the Cu IONs foliar exposures (**Table S7**). When Cu(OH)₂ NMs were applied on wheat leaves, three antioxidants and fourteen amino acids were significantly changed, whereas no antioxidants or amino acids were significantly altered under the Cu IONs treatment (**Figure 3 (A) and (C)**). Thus, the wheat leaf ROS defense system and changes in N metabolism were triggered by Cu(OH)₂ NMs but not by Cu IONs. Fructose, maltose, raffinose, and sucrose from wheat leaves were significantly downregulated under both Cu(OH)₂ NMs and Cu IONs foliar applications. The decreasing levels of these soluble sugars indicate that C fixation capacity of wheat leaves was hindered in the copper foliar exposures. Zhao et al. (2017) employed a much higher foliar dosage of ionic Cu (0.15-1.5 mg CuSO₄/plant) versus 1.8-18 mg Cu(OH)₂ nanopesticide/plant on lettuce, which led to contrary findings.⁶⁹ They discovered that 30 metabolites were markedly altered in response to CuSO₄ exposure and 24 dysregulated metabolites were identified for Cu(OH)₂ nanopesticide. Ascorbic acid, which had minor changes in the current study, decreased around 10 fold at a 1.5 mg CuSO₄ dose and 10 fold at a 18 mg Cu(OH)₂ nanopesticide dose.⁶⁹


Unlike the copper materials, exposure to Mo IONs resulted in 16 more dysregulated metabolites in wheat leaves than MoO₃ NMs (**Table S7**). Even though a majority of the dysregulated metabolites were significantly downregulated (**Figure 3 (B) and (D)**), aspartic acid, proline, and chlorogenic acid were significantly upregulated (fold change>2) in wheat leaves for the MoO₃ NMs treatment. Aspartic acid is an important factor in NH₄⁺ assimilation and it can also serve as a donor for both C and N metabolism.⁷⁰ Thus,

upregulation of aspartic acid may indicate the perturbation of primary N and C metabolism. Proline is a radical scavenger and can also act as an electron sink that aids in ROS defense response.⁷¹ Dysregulation of proline has been extensively studied in plant responses to metal stressors, including metal NMs, where the proline concentration was often upregulated,^{39, 44, 45, 60, 72-74} but the level can also decrease in other cases.⁷⁵ Curcumin, as one of antioxidants, was consistently upregulated under both MoO₃ NMs and Mo IONs exposures. Dysregulation of antioxidant metabolites (chlorogenic acid and curcumin) suggests an induced oxidative stress response in wheat leaves to cope with increased Mo levels.⁷⁵

Wheat root responses to metal-surfactant suspensions

Foliar exposure of wheat to Cu(OH)₂ NMs and Cu IONs had little impact on Cu content in wheat roots (**Figure S7 (A)**), with minimal translocation. In addition, foliar exposure to all metal-surfactant suspensions did not influence root dry biomass, nor play a major role in nutrient distribution (**Figure S5 (B) and Table S6**). On the other hand, a significant amount of Mo was translocated to roots under both MoO₃ NM and Mo ION foliar treatments (**Figure S7 (B)**). The translocated fraction of MoO₃ NMs was 65.5% (137.3 µg/g Mo in roots) and 39.0% for Mo IONs (66.9 µg/g Mo in roots). The much higher MoO₃ NMs translocation than Cu(OH)₂ NMs could be explained from NM properties and translocation mechanisms. NMs with larger hydrodynamic size have shown significantly lower foliar delivery efficiency.^{26, 30} Since plant cell walls are mainly negatively charged, positively charged ions have shown a much higher accumulation in the apoplastic space than negative ions.⁷⁶ Compared with negatively charged MoO₃ NMs, the movement of positively charged metal ions (e.g., Mo ions) and NMs (e.g., Cu(OH)₂ NMs) are also likely restricted. However, the opposite result was found by Hu et al., (2020), where the positively charged NMs resulted in higher foliar transport efficiency than negative NMs in guard cells, extracellular space, and chloroplasts.²⁶ NM size and surface charge are not the only factors that drive NMs foliar uptake and translocation. Other parameters, such as NMs surface coating, leaf and NM hydrophilicity, solubility,

334 binding affinity to organic phases, and plant species are likely important, but their role remains largely
 335 unknown.³⁴



336
 337 **Figure 4.** Comparison of significantly changed metabolites ($|\log_2(\text{fold change})| > 1$) in wheat roots exposed
 338 to (A) Cu(OH)₂ NMs; (B) MoO₃ NMs; (C) Cu IONs; and (D) Mo IONs. Experimental conditions: SA - 200 mg/L
 339 TritonTM X-100 solution; Cu(OH)₂ NMs - 100 mg/L Cu(OH)₂ NMs (as Cu content) in SA; Cu IONs - 0.1 mg/L
 340 CuSO₄ · 5H₂O (as Cu content) in SA; MoO₃ NMs - 100 mg/L MoO₃ NMs (as Mo content) in SA; and Mo IONs
 341 - 35 mg/L Na₂MoO₄ · 2H₂O (as Mo content) in SA. The numbers in the pie chart represent the significantly
 342 changed metabolites from each metabolite category. The color of the numbers in the outer circular ring
 343 indicates fold changes of the significantly dysregulated metabolites: red (fold change > 5), orange (2 < fold
 344 change < 5), blue (0.2 < fold change < 0.5), and purple (fold change < 0.2). Three replicates under each
 345 condition.

The PLS-DA score plot revealed that the metabolite profiles of wheat roots exposed to metal-surfactant treatments had good separation from the surfactant-only (SA) treatment (**Figure S7 (C) and (D)**). Physiological and nutrient distribution results showed only minor differences among different treatments (**Figure S5 (B) and Table S6**). Metabolomic analysis showed that Cu(OH)₂ NMs resulted in 27 significantly changed metabolites ($|\log_2(\text{fold change})| > 1$), while there were only 3 dysregulated metabolites discovered under Cu IONs exposure (**Figure 4 (A) and (C)**). Except for curcumin, all the altered metabolites were downregulated (**Table S8**). Zhao et al., (2016) proposed that metabolites transported from leaves to roots were partially responsible for alterations of root metabolite levels.⁷⁷ In another similar study, Zhang et al. (2019) exposed spinach leaves to CeO₂ NMs and found minimal Ce content difference in roots between the control and treatment groups.⁷⁸ Nevertheless, more profound metabolomic reprogramming was discovered in spinach roots than in leaves. However, Hong et al. (2014) treated cucumber (*Cucumis sativus*) leaves with nanoceria, and found that Ce was significantly translocated to other plant sections (i.e., stems, roots, and flowers).¹⁴ This shows the complexity of NM foliar uptake, which often depends on NM properties, plant species, mode of application, and other environmental factors.

Even though foliar exposure to MoO₃ NMs resulted in 110% more Mo load in roots than exposure to Mo IONs, 12 more dysregulated metabolites were identified in wheat roots under the Mo ION treatment than to MoO₃ NMs (**Figures 4 (B) and (D)**). Compared with leaf metabolomic results, four more significantly upregulated (fold change > 2) metabolites (lysine, phenylalanine, adenosine, and guanosine) were found in wheat roots under MoO₃ NMs condition. In contrast, those upregulated metabolites were all significantly downregulated (fold change < 0.2) for Mo IONs (**Table S8**). Amino acids (lysine and phenylalanine) play a central role in the synthesis of many cellular enzymes and in-plant detoxification towards abiotic stressors (i.e., heavy metal exposures).⁷⁹ The significant changes of adenosine and

guanosine indicated the purine metabolism pathway was disturbed. In addition, xanthine dehydrogenase/oxidase is a molybdoenzyme that has been identified in legumes. These amino acids are also involved in purine catabolism and ureide biosynthesis,⁶³ which are essential functions for remobilizing nitrogen for plant growth and development. A recent study conducted by Huang et al. (2021) applied 200 mg/kg of MoO₃ NMs and the corresponding Mo ionic concentrations (70 mg/kg) to wheat roots for one week.⁴⁹ Results showed that wheat leaves accumulated 2 times more Mo under MoO₃ NM treatment than under Mo ionic exposure. Furthermore, nucleic acids and sugars in wheat leaves separated well from the control for MoO₃ NMs, but only minor differences were found when wheat roots were exposed to Mo ions. The current finding demonstrates the distinct differences in NM uptake and translocation mechanisms for the two entry pathways (i.e., roots vs foliar applications).

Comparison of wheat responses to Cu(OH)₂ NMs and MoO₃ NMs

The metabolic pathway analysis showed that both wheat leaves and roots were significantly perturbed for Cu(OH)₂ NMs foliar exposure (**Figure S8 and Table S9**). In total, 17 pathways were perturbed in wheat leaves and 17 in roots. Glyoxylate/dicarboxylate metabolism and stilbenoid/diarylheptanoid/gingerol biosynthesis were only disturbed in wheat leaves. Chlorogenic acid is a uniquely dysregulated metabolite that is involved in the later metabolic pathway. The glyoxylate and dicarboxylate pathway is related to carbohydrate metabolism, where carbohydrates are synthesized from fatty acids.⁸⁰ On the other hand, perturbation of glutathione metabolism and valine/leucine/ isoleucine biosynthesis were only found in wheat roots. These perturbed pathways are related to carbon and nitrogen metabolism and involve a uniquely altered metabolite - glutamate. Foliar exposure to Cu IONs led to 12 disturbed pathways in wheat leaves and 7 in roots (**Figure S9 and Table S10**). Compared with the Cu(OH)₂ NMs foliar exposure, fewer metabolic pathways were perturbed when wheat leaves were exposed to MoO₃ NMs (**Figure S10 and Table S9**). Citric acid (TCA cycle), methionine and serine (cysteine and methionine metabolism), raffinose (galactose metabolism), and serine (glycine, serine and threonine

metabolism) were uniquely dysregulated metabolites in wheat leaves when they were exposed to MoO₃ NMs. Tryptophan metabolism was the uniquely perturbed pathway in wheat roots after exposure to MoO₃ NMs. Tryptophan is an essential component in protein synthesis and is also a central molecule that serves as a precursor for many secondary metabolites.⁸¹ The Mo IONs treatment resulted in mostly downregulated metabolites and very similar perturbation pathways between wheat leaves and roots (**Figure S11 and Table S10**). Valine, isoleucine, and leucine were significantly downregulated under the Mo IONs treatment that led to the perturbation of valine/leucine/isoleucine biosynthesis.

In general, the Venn diagram demonstrates that exposure to Cu(OH)₂ NMs reprogramed the wheat metabolic profile more than exposure to MoO₃ NMs (**Figure 5**). This finding was quite interesting, considering only a very small amount of Cu was dissolved from Cu(OH)₂ NMs and even less was translocated to wheat roots. On the other hand, a much higher amount of Mo was dissolved from MoO₃ NMs and 65.5% of Mo was translocated to wheat roots (**Figure S7 (A) and (B)**). Compared with MoO₃ NMs exposure, there were 10 more dysregulated metabolites found in wheat leaves and 9 more discovered in roots when Cu(OH)₂ NMs applied (**Table S11**). Even though four dysregulated metabolites (i.e., ribitol, citric acid, adenine, and ascorbic acid) only responded to MoO₃ NMs, there were no uniquely perturbed pathways found when compared to Cu(OH)₂ NMs (**Table S12**). Conversely, glyoxylate/dicarboxylate metabolism from wheat leaves and valine/leucine/isoleucine biosynthesis from wheat roots uniquely responded to Cu(OH)₂ NMs.




Figure 5. Venn diagram of (A) dysregulated metabolites and (B) perturbed metabolic pathways in wheat after one-week foliar exposure to NMs suspensions. Legend: ■ wheat leaves exposed to Cu(OH)₂ NMs; ■ wheat leaves exposed to MoO₃ NMs; ■ wheat roots exposed to Cu(OH)₂ NMs; ■ wheat roots exposed to MoO₃ NMs.

Conclusions

When Triton™ X-100 surfactant (SA) was applied to wheat leaves, minimal changes were observed in physiological responses and nutrient distributions. However, metabolite profiles were significantly reprogrammed. For instance, leucine, lysine, phenylalanine, proline, serine, tryptophan, linolenic acid, adenosine, guanosine, uridine, and trehalose were significantly changed (with fold change >5) in wheat leaves and roots. In addition, 17 metabolic pathways were perturbed in wheat leaves and 15 in roots.

Adding the surfactant significantly promoted applied NMs and ionic metal deposition to wheat leaves. A significant amount of Cu (649.8 µg/g) was detected in wheat leaves under Cu(OH)₂ NMs that resulted in 34 dysregulated metabolites. Since only 0.1% of Cu(OH)₂ NMs dissolved in SA, Cu IONs had minimum effect on Cu accumulation or metabolic alterations. Even though minimum translocation was observed for Cu(OH)₂ NMs, there were 27 significantly altered metabolites discovered in wheat roots. In contrast, a large percentage of the MoO₃ NMs dissolved in SA (e.g., 35.8%), which led to 32.2 µg/g more Mo and

16 more dysregulated metabolites detected in wheat leaves under MoO₃ IONs treatment. Even MoO₃ NMs resulted in 110% more Mo loading in wheat roots than Mo IONs, 16 additional dysregulated metabolites were discovered for the Mo IONs treatment.

Due to differences in NM properties (e.g., hydrodynamic diameter, zeta potential, and dissolution extent), metabolite pathway analysis showed Cu(OH)₂ NMs reprogrammed wheat metabolites profile more than MoO₃ NMs. Even though ribitol, citric acid, adenine, and ascorbic acid only responded to MoO₃ NMs, no uniquely disturbed pathways were discovered. Conversely, Cu(OH)₂ NMs resulted in uniquely disturbed glyoxylate/dicarboxylate metabolism in wheat leaves and the perturbation valine/leucine/isoleucine metabolism in wheat roots.

Overall, this study revealed that foliar application of NM-surfactant suspensions could deposit more NMs on the leaves and significantly reprogram the metabolic profile of early stage wheat seedlings. A thorough understanding of the plant's internal status changes could benefit surfactant innovation, thus promoting more sustainable and efficient NMs foliar application in agriculture. [In future studies, other omics techniques, such as proteomics, transcriptomics, and genomics can help to illustrate plant biochemical responses to NMs exposures. Here we observed plant responses after one-week foliar exposure, as a proof-of-concept; future studies should consider plant responses with longer time durations. Furthermore, other agricultural-related surfactants and environmental conditions could be considered to better mimic field applications.](#)

Supporting Information

The Supporting Information contains detailed LC/MS approaches/parameters, along with 12 tables and 6 figures presenting the effect of surfactant-containing NMs suspensions on plant growth, nutrient distribution, dysregulated metabolites, perturbation pathways, as well as the Venn diagram data.

Conflict of interest

454 The authors declare no conflict of interest

455 **Acknowledgments**

456 This work was supported by the National Science Foundation (NSF) under cooperative agreement number
 457 NSF-1901515. The authors thank Dr. Sanghamitra Majumdar for her advice on the experimental approach,
 458 and Weiwei Li, Arthur Kao, Devin Everett, and Xiaofan Liu for their contributions to the preliminary
 459 experiment design and test. Arturo A. Keller would like to give special thanks to Agilent Technologies for
 460 their Agilent Thought Leader Award.

461 **Author Contributions**

462 Conceptualization, Xiangning Huang and Arturo A. Keller; Methodology, Xiangning Huang and Arturo A.
 463 Keller; Investigation, Xiangning Huang; Writing—original draft preparation, Xiangning Huang; Writing—
 464 review and editing, Xiangning Huang and Arturo A. Keller; Supervision, Arturo A. Keller. All authors have
 465 read and agreed to the submitted version of the manuscript.

466 **References**

- 467 1. Liu, R. and Lal, R. Potentials of engineered nanoparticles as fertilizers for increasing agronomic
 468 productions. *Sci. Total Environ.* **2015**, *514*, 131-139.
- 469 2. Zhao, L.; Lu, L.; Wang, A.; Zhang, H.; Huang, M.; Wu, H.; Xing, B.; Wang, Z.; Ji, R. Nano-Biotechnology
 470 in Agriculture: Use of Nanomaterials to Promote Plant Growth and Stress Tolerance. *J. Agr. Food*
 471 *Chem.* **2020**, *68*, (7), 1935-1947.
- 472 3. Alshaal, T. and El-Ramady, H. Foliar application: from plant nutrition to biofortification. *Envir.*
 473 *Biodivers. Soil Secur.* **2017**, *1*, (2017), 71-83.
- 474 4. Tan, W.; Gao, Q.; Deng, C.; Wang, Y.; Lee, W.-Y.; Hernandez-Viezcas, J. A.; Peralta-Videa, J. R.; Gardea-
 475 Torresdey, J. L. Foliar exposure of Cu(OH)₂ nanopesticide to basil (*Ocimum basilicum*): variety-
 476 dependent copper translocation and biochemical responses. *J. Agr. Food Chem.* **2018**, *66*, (13), 3358-
 477 3366.
- 478 5. Adisa, I. O.; Pullagurala, V. L. R.; Peralta-Videa, J. R.; Dimkpa, C. O.; Elmer, W. H.; Gardea-Torresdey, J.
 479 L.; White, J. C. Recent advances in nano-enabled fertilizers and pesticides: a critical review of
 480 mechanisms of action. *Environ. Sci. Nano.* **2019**, *6*, (7), 2002-2030.
- 481 6. Wang, X.; Liu, X.; Chen, J.; Han, H.; Yuan, Z. Evaluation and mechanism of antifungal effects of carbon
 482 nanomaterials in controlling plant fungal pathogen. *Carbon.* **2014**, *68*, 798-806.
- 483 7. Shallan, M. A.; Hassan, H. M.; Namich, A. A.; Ibrahim, A. A. Biochemical and physiological effects of
 484 TiO₂ and SiO₂ nanoparticles on cotton plant under drought stress. *Res. J. Pharm. Biol. Chem. Sci.* **2016**,
 485 *7*, (4), 1540-1551.
- 486 8. Siddiqui, M. H.; Al - Whaibi, M. H.; Faisal, M.; Al Sahli, A. A. Nano - silicon dioxide mitigates the
 487 adverse effects of salt stress on Cucurbita pepo L. *Environ. Toxicol. Chem.* **2014**, *33*, (11), 2429-2437.

9. Giraldo, J. P.; Landry, M. P.; Faltermeier, S. M.; McNicholas, T. P.; Iverson, N. M.; Boghossian, A. A.; Reuel, N. F.; Hilmer, A. J.; Sen, F.; Brew, J. A. Plant nanobionics approach to augment photosynthesis and biochemical sensing. *Nat. Mater.* **2014**, *13*, (4), 400-408.
10. Huang, Y.; Zhao, L.; Keller, A. A. Interactions, transformations, and bioavailability of nano-copper exposed to root exudates. *Environ. Sci. Technol.* **2017**, *51*, (17), 9774-9783.
11. Cervantes-Avilés, P.; Huang, X.; Keller, A. A. Dissolution and Aggregation of Metal Oxide Nanoparticles in Root Exudates and Soil Leachate: Implications for Nanoagrochemical Application. *Environ. Sci. Technol.* **2021**, *55*, (20), 13443-13451.
12. Iavicoli, I.; Leso, V.; Beezhold, D. H.; Shvedova, A. A. Nanotechnology in agriculture: Opportunities, toxicological implications, and occupational risks. *Toxicol. Appl. Pharm.* **2017**, *329*, 96-111.
13. Larue, C.; Castillo-Michel, H.; Sobanska, S.; Cécillon, L.; Bureau, S.; Barthès, V.; Ouerdane, L.; Carrière, M.; Sarret, G. Foliar exposure of the crop *Lactuca sativa* to silver nanoparticles: evidence for internalization and changes in Ag speciation. *J. Hazard. Mater.* **2014**, *264*, 98-106.
14. Hong, J.; Peralta-Videa, J. R.; Rico, C.; Sahi, S.; Viveros, M. N.; Bartonjo, J.; Zhao, L.; Gardea-Torresdey, J. L. Evidence of translocation and physiological impacts of foliar applied CeO₂ nanoparticles on cucumber (*Cucumis sativus*) plants. *Environ. Sci. Technol.* **2014**, *48*, (8), 4376-4385.
15. Peirce, C. A.; McBeath, T. M.; Priest, C.; McLaughlin, M. J. The timing of application and inclusion of a surfactant are important for absorption and translocation of foliar phosphoric acid by wheat leaves. *Front. Plant Sci.* **2019**, *10*, 1532.
16. Green, J. M. and Beestman, G. B. Recently patented and commercialized formulation and adjuvant technology. *Crop Prot.* **2007**, *26*, (3), 320-327.
17. Castro, M. J.; Ojeda, C.; Cirelli, A. F. Advances in surfactants for agrochemicals. *Environ. Chem. Lett.* **2014**, *12*, (1), 85-95.
18. Wang, P. and Keller, A. A. Partitioning of hydrophobic organic compounds within soil–water–surfactant systems. *Water Res.* **2008**, *42*, (8-9), 2093-2101.
19. Zhao, X.; Gao, Y.; Zhang, C.; Zhu, Y.; Lei, J.; Ma, Y.; Du, F. Wettability of ionic surfactants SDS and DTAB on wheat (*Triticum aestivum*) leaf surfaces. *J. Disper. Sci. Technol.* **2018**, *39*, (12), 1820-1828.
20. Zhang, C.; Zhao, X.; Lei, J.; Ma, Y.; Du, F. The wetting behavior of aqueous surfactant solutions on wheat (*Triticum aestivum*) leaf surfaces. *Soft Matter.* **2017**, *13*, (2), 503-513.
21. Peirce, C. A.; Priest, C.; McBeath, T. M.; McLaughlin, M. J. Uptake of phosphorus from surfactant solutions by wheat leaves: spreading kinetics, wetted area, and drying time. *Soft Matter.* **2016**, *12*, (1), 209-218.
22. Castro, M. J.; Ojeda, C.; Cirelli, A. F. Surfactants in agriculture. In *Green Materials for Energy, Products and Depollution*. Springer. **2013**, 287-334.
23. Januszkiewicz, K.; Mrozek-Niećko, A.; Róžański, J. Effect of surfactants and leaf surface morphology on the evaporation time and coverage area of ZnIDHA droplets. *Plant Soil.* **2019**, *434*, (1-2), 93-105.
24. Banks, M.-L. L.; Kremer, R. J.; Eivazi, F.; Motavalli, P. P.; Nelson, K. A. Effects of selected surfactants on nutrient uptake in corn (*Zea mays* L.). *J. Plant Nutr.* **2015**, *38*, (7), 1036-1049.
25. Peirce, C.; McBeath, T.; Fernández, V.; McLaughlin, M. Wheat leaf properties affecting the absorption and subsequent translocation of foliar-applied phosphoric acid fertiliser. *Plant Soil.* **2014**, *384*, (1-2), 37-51.
26. Hu, P.; An, J.; Faulkner, M. M.; Wu, H.; Li, Z.; Tian, X.; Giraldo, J. P. Nanoparticle Charge and Size Control Foliar Delivery Efficiency to Plant Cells and Organelles. *ACS Nano.* **2020**, *14*, (7), 7970-7986.
27. Zimbovskaya, M. M.; Polyakov, A. Y.; Volkov, D. S.; Kulikova, N. A.; Lebedev, V. A.; Pankratov, D. A.; Konstantinov, A. I.; Parfenova, A. M.; Zhilkibaev, O.; Perminova, I. V. Foliar Application of Humic-Stabilized Nanoferrhydrite Resulted in an Increase in the Content of Iron in Wheat Leaves. *Agron.* **2020**, *10*, (12), 1891.

28. Larue, C.; Veronesi, G.; Flank, A.-M.; Surble, S.; Herlin-Boime, N.; Carrière, M. Comparative uptake and impact of TiO₂ nanoparticles in wheat and rapeseed. *J. Toxicol. Environ. Heal. A*. **2012**, *75*, (13-15), 722-734.
29. Read, T. L.; Doolette, C. L.; Li, C.; Schjoerring, J. K.; Kopittke, P. M.; Donner, E.; Lombi, E. Optimising the foliar uptake of zinc oxide nanoparticles: Do leaf surface properties and particle coating affect absorption? *Physiol. Plantarum*. **2020**, *170*, (3), 384-397.
30. Avellan, A.; Yun, J.; Zhang, Y.; Spielman-Sun, E.; Unrine, J. M.; Thieme, J.; Li, J.; Lombi, E.; Bland, G.; Lowry, G. V. Nanoparticle size and coating chemistry control foliar uptake pathways, translocation, and leaf-to-rhizosphere transport in wheat. *ACS Nano*. **2019**, *13*, (5), 5291-5305.
31. Pérez - de - Luque, A.; Cifuentes, Z.; Beckstead, J. A.; Sillero, J. C.; Ávila, C.; Rubio, J.; Ryan, R. O. Effect of amphotericin B nanodisks on plant fungal diseases. *Pest Manag. Sci.* **2012**, *68*, (1), 67-74.
32. Rathore, I. and Tarafdar, J. Perspectives of biosynthesized magnesium nanoparticles in foliar application of wheat plant. *J. Bionanosci.* **2015**, *9*, (3), 209-214.
33. Ghafari, H. and Razmjoo, J. Effect of foliar application of nano-iron oxidase, iron chelate and iron sulphate rates on yield and quality of wheat. *Int. J. Agron. Plant Prod.* **2013**, *4*, (11), 2997-3003.
34. Avellan, A.; Yun, J.; Morais, B. P.; Clement, E. T.; Rodrigues, S. M.; Lowry, G. V. Critical Review: Role of Inorganic Nanoparticle Properties on Their Foliar Uptake and in Planta Translocation. *Environ. Sci. Technol.* **2021**, *55*, (20), 13417-13431.
35. Hong, J.; Wang, C.; Wagner, D. C.; Gardea-Torresdey, J.; He, F.; Rico, C. M. Foliar application of nanoparticles: Mechanism of absorption, transfer, and multiple impacts. *Environ. Sci. Nano*. **2021**, *8*, 1196-1210.
36. Deshpande, P.; Dapkekar, A.; Oak, M. D.; Paknikar, K. M.; Rajwade, J. M. Zinc complexed chitosan/TPP nanoparticles: A promising micronutrient nanocarrier suited for foliar application. *Carbohydr. Polym.* **2017**, *165*, 394-401.
37. Dimkpa, C. O.; Singh, U.; Adisa, I. O.; Bindraban, P. S.; Elmer, W. H.; Gardea-Torresdey, J. L.; White, J. C. Effects of manganese nanoparticle exposure on nutrient acquisition in wheat (*Triticum aestivum* L.). *Agron.* **2018**, *8*, (9), 158.
38. Su, Y.; Ashworth, V.; Kim, C.; Adeleye, A. S.; Rolshausen, P.; Roper, C.; White, J.; Jassby, D. Delivery, uptake, fate, and transport of engineered nanoparticles in plants: a critical review and data analysis. *Environ. Sci. Nano*. **2019**, *6*, (8), 2311-2331.
39. Tian, L.; Shen, J.; Sun, G.; Wang, B.; Ji, R.; Zhao, L. Foliar Application of SiO₂ Nanoparticles Alters Soil Metabolite Profiles and Microbial Community Composition in the Pakchoi (*Brassica chinensis* L.) Rhizosphere Grown in Contaminated Mine Soil. *Environ. Sci. Technol.* **2020**, *54*, (20), 13137-13146.
40. Zhao, L.; Zhang, H.; Wang, J.; Tian, L.; Li, F.; Liu, S.; Peralta-Videa, J. R.; Gardea-Torresdey, J. L.; White, J. C.; Huang, Y. C60 fullerenes enhance copper toxicity and alter the leaf metabolite and protein profile in Cucumber. *Environ. Sci. Technol.* **2019**, *53*, (4), 2171-2180.
41. Zhao, L.; Huang, Y.; Paglia, K.; Vaniya, A.; Wancewicz, B.; Keller, A. A. Metabolomics reveals the molecular mechanisms of copper induced cucumber leaf (*Cucumis sativus*) senescence. *Environ. Sci. Technol.* **2018**, *52*, (12), 7092-7100.
42. Zhang, H.; Du, W.; Peralta-Videa, J. R.; Gardea-Torresdey, J. L.; White, J. C.; Keller, A.; Guo, H.; Ji, R.; Zhao, L. Metabolomics reveals how cucumber (*Cucumis sativus*) reprograms metabolites to cope with silver ions and silver nanoparticle-induced oxidative stress. *Environ. Sci. Technol.* **2018**, *52*, (14), 8016-8026.
43. Salehi, H.; Chehregani, A.; Lucini, L.; Majd, A.; Gholami, M. Morphological, proteomic and metabolomic insight into the effect of cerium dioxide nanoparticles to *Phaseolus vulgaris* L. under soil or foliar application. *Sci. Total Environ.* **2018**, *616*, 1540-1551.

44. Zhao, L.; Huang, Y.; Keller, A. A. Comparative metabolic response between cucumber (*Cucumis sativus*) and corn (*Zea mays*) to a $\text{Cu}(\text{OH})_2$ nanopesticide. *J. Agric. Food Chem.* **2017**, *66*, (26), 6628-6636.
45. Zhao, L.; Hu, Q.; Huang, Y.; Keller, A. A. Response at genetic, metabolic, and physiological levels of maize (*Zea mays*) exposed to a $\text{Cu}(\text{OH})_2$ nanopesticide. *ACS Sustain. Chem. Eng.* **2017**, *5*, (9), 8294-8301.
46. Zhao, L.; Hu, Q.; Huang, Y.; Fulton, A. N.; Hannah-Bick, C.; Adeleye, A. S.; Keller, A. A. Activation of antioxidant and detoxification gene expression in cucumber plants exposed to a $\text{Cu}(\text{OH})_2$ nanopesticide. *Environ. Sci. Nano.* **2017**, *4*, (8), 1750-1760.
47. Zhao, L.; Huang, Y.; Hannah-Bick, C.; Fulton, A. N.; Keller, A. A. Application of metabolomics to assess the impact of $\text{Cu}(\text{OH})_2$ nanopesticide on the nutritional value of lettuce (*Lactuca sativa*): enhanced Cu intake and reduced antioxidants. *NanoImpact.* **2016**, *3*, 58-66.
48. Majumdar, S. and Keller, A. A. Omics to address the opportunities and challenges of nanotechnology in agriculture. *Crit. Rev. Environ. Sci Technol.* **2021**, *51*, (22), 2595-2636.
49. Huang, X.; Cervantes-Avilés, P.; Li, W.; Keller, A. A. Drilling into the Metabolomics to Enhance Insight on Corn and Wheat Responses to Molybdenum Trioxide Nanoparticles. *Environ. Sci. Technol.* **2021**, *55*, (20), 13452-13464.
50. Majumdar, S.; Long, R. W.; Kirkwood, J. S.; Minakova, A. S.; Keller, A. A. Unraveling Metabolic and Proteomic Features in Soybean Plants in Response to Copper Hydroxide Nanowires Compared to a Commercial Fertilizer. *Environ. Sci. Technol.* **2021**, *55*, (20), 13477-13489.
51. Hewitt, E. and Bolle-Jones, E. Molybdenum as a Plant Nutrient: II. The Effects of Molybdenum Deficiency on Some Horticultural and Agricultural Crop Plants in Sand Culture. *Int. J. Hortic. Sci.* **1952**, *27*, (4), 257-265.
52. C Gupta, U.; C Srivastava, P.; C Gupta, S. Role of micronutrients: Boron and molybdenum in crops and in human health and nutrition. *Curr. Nutr. Food Sci.* **2011**, *7*, (2), 126-136.
53. Adhikari, T.; Kundu, S.; Rao, A. S. Impact of SiO_2 and Mo nano particles on seed germination of rice (*Oryza sativa* L.). *Int. J. Agric. Food Sci. Technol.* **2013**, *4*, (8), 809-816.
54. Kanneganti, A. and Talasila, M. MoO_3 nanoparticles: synthesis, characterization and its hindering effect on germination of *Vigna Unguiculata* seeds. *J. Eng. Re.s Appl.* **2014**, *4*, (7), 116-120.
55. Mushinskiy, A. and Aminova, E. In *Effect of iron, copper and molybdenum nanoparticles on morphometric parameters of Solanum tuberosum L. plants*. IOP Pubg. **2019**, *341*, (1), 12195.
56. Abbasifar, A.; ValizadehKaji, B.; Iravani, M. A. Effect of green synthesized molybdenum nanoparticles on nitrate accumulation and nitrate reductase activity in spinach. *J. Plant Nutr.* **2020**, *43*, (1), 13-27.
57. Sharma, P. K.; Raghubanshi, A. S.; Shah, K. Examining the uptake and bioaccumulation of molybdenum nanoparticles and their effect on antioxidant activities in growing rice seedlings. *Environ. Sci. Pollut. Res.* **2021**, *28*, (11), 13439-13453.
58. Majumdar, S.; Ma, C.; Villani, M.; Zuverza-Mena, N.; Pagano, L.; Huang, Y.; Zappettini, A.; Keller, A. A.; Marmioli, N.; Dhankher, O. P. Surface coating determines the response of soybean plants to cadmium sulfide quantum dots. *NanoImpact.* **2019**, *14*, 100151.
59. Majumdar, S.; Pagano, L.; Wohlschlegel, J. A.; Villani, M.; Zappettini, A.; White, J. C.; Keller, A. A. Proteomic, gene and metabolite characterization reveal the uptake and toxicity mechanisms of cadmium sulfide quantum dots in soybean plants. *Environ. Sci. Nano.* **2019**, *6*, (10), 3010-3026.
60. Liu, W.; Majumdar, S.; Li, W.; Keller, A. A.; Slaveykova, V. I. Metabolomics for early detection of stress in freshwater alga *Poteroiochromonas malhamensis* exposed to silver nanoparticles. *Sci. Rep.* **2020**, *10*, (1), 1-13.
61. Huang, Y.; Adeleye, A. S.; Zhao, L.; Minakova, A. S.; Anumol, T.; Keller, A. A. Antioxidant response of cucumber (*Cucumis sativus*) exposed to nano copper pesticide: Quantitative determination via LC-MS/MS. *Food Chem.* **2019**, *270*, 47-52.

62. Huang, Y.; Li, W.; Minakova, A. S.; Anumol, T.; Keller, A. A. Quantitative analysis of changes in amino acids levels for cucumber (*Cucumis sativus*) exposed to nano copper. *NanoImpac.* **2018**, *12*, 9-17.
63. Kaiser, B. N.; Gridley, K. L.; Ngaire Brady, J.; Phillips, T.; Tyerman, S. D. The role of molybdenum in agricultural plant production. *Ann. Bot.* **2005**, *96*, (5), 745-754.
64. Xu, L.; Han, L.; Huang, B. Membrane fatty acid composition and saturation levels associated with leaf dehydration tolerance and post - drought rehydration in Kentucky bluegrass. *Crop Sci.* **2011**, *51*, (1), 273-281.
65. Jain, N. K. and Roy, I. Effect of trehalose on protein structure. *Protein Sci.* **2009**, *18*, (1), 24-36.
66. Forde, B. G. and Lea, P. J. Glutamate in plants: metabolism, regulation, and signalling. *J. Exp. Bot.* **2007**, *58*, (9), 2339-2358.
67. Schulze-Siebert, D.; Heineke, D.; Scharf, H.; Schultz, G. Pyruvate-derived amino acids in spinach chloroplasts: synthesis and regulation during photosynthetic carbon metabolism. *Plant Physiol.* **1984**, *76*, (2), 465-471.
68. Mahil, E. and Kumar, B. Foliar application of nanofertilizers in agricultural crops—A review. *J. Farm Sci.* **2019**, *32*, (3), 239-249.
69. Zhao, L.; Huang, Y.; Adeleye, A. S.; Keller, A. A. Metabolomics reveals Cu(OH)₂ nanopesticide-activated anti-oxidative pathways and decreased beneficial antioxidants in spinach leaves. *Environ. Sci. Technol.* **2017**, *51*, (17), 10184-10194.
70. O'Leary, B. and Plaxton, W. The central role of glutamate and aspartate in the post-translational control of respiration and nitrogen assimilation in plant cells. *Amino Acids High. Plant. CAB Int.* **2015**, 277-297.
71. Sharma, S. S. and Dietz, K.-J. The significance of amino acids and amino acid-derived molecules in plant responses and adaptation to heavy metal stress. *J. Exp. Bot.* **2006**, *57*, (4), 711-726.
72. Gong, C.; Wang, L.; Li, X.; Wang, H.; Jiang, Y.; Wang, W. Responses of seed germination and shoot metabolic profiles of maize (*Zea mays* L.) to Y₂O₃ nanoparticle stress. *RSC Adv.* **2019**, *9*, (47), 27720-27731.
73. Zhang, H.; Huang, M.; Zhang, W.; Gardea-Torresdey, J. L.; White, J. C.; Ji, R.; Zhao, L. Silver Nanoparticles Alter Soil Microbial Community Compositions and Metabolite Profiles in Unplanted and Cucumber-Planted Soils. *Environ. Sci. Technol.* **2020**, *54*, (6), 3334-3342.
74. Zhao, L.; Huang, Y.; Zhou, H.; Adeleye, A. S.; Wang, H.; Ortiz, C.; Mazer, S. J.; Keller, A. A. GC-TOF-MS based metabolomics and ICP-MS based metallomics of cucumber (*Cucumis sativus*) fruits reveal alteration of metabolites profile and biological pathway disruption induced by nano copper. *Environ. Sci. Nano.* **2016**, *3*, (5), 1114-1123.
75. Zhao, L.; Zhang, H.; White, J. C.; Chen, X.; Li, H.; Qu, X.; Ji, R. Metabolomics reveals that engineered nanomaterial exposure in soil alters both soil rhizosphere metabolite profiles and maize metabolic pathways. *Environ. Sci. Nano.* **2019**, *6*, (6), 1716-1727.
76. Marschner, H., Marschner's mineral nutrition of higher plants. Academic Press. **2011**.
77. Zhao, L.; Ortiz, C.; Adeleye, A. S.; Hu, Q.; Zhou, H.; Huang, Y.; Keller, A. A. Metabolomics to detect response of lettuce (*Lactuca sativa*) to Cu(OH)₂ nanopesticides: oxidative stress response and detoxification mechanisms. *Environ. Sci. Technol.* **2016**, *50*, (17), 9697-9707.
78. Zhang, H.; Lu, L.; Zhao, X.; Zhao, S.; Gu, X.; Du, W.; Wei, H.; Ji, R.; Zhao, L. Metabolomics reveals the "invisible" responses of spinach plants exposed to CeO₂ nanoparticles. *Environ. Sci. Technol.* **2019**, *53*, (10), 6007-6017.
79. Pratelli, R. and Pilot, G., Regulation of amino acid metabolic enzymes and transporters in plants. *J. Exp. Bot.* **2014**, *65*, (19), 5535-5556.
80. Wu, B.; Zhu, L.; Le, X. C. Metabolomics analysis of TiO₂ nanoparticles induced toxicological effects on rice (*Oryza sativa* L.). *Environ. Pollut.* **2017**, *230*, 302-310.

- 676 81. Tzin, V. and Galili, G., The biosynthetic pathways for shikimate and aromatic amino acids in
677 *Arabidopsis thaliana*. *The Arabidopsis Book*. **2010**, 8, 132.

678

679