

HITCHIN FIBRATIONS, ABELIAN SURFACES, AND THE P=W CONJECTURE

MARK ANDREA DE CATALDO, DAVESH MAULIK, AND JUNLIANG SHEN

CONTENTS

Introduction	1
1. Perverse filtrations	6
2. Moduli of one-dimensional sheaves	10
3. Splittings of perverse filtrations	24
4. Topology of Hitchin fibrations	31

INTRODUCTION

0.1. Perverse filtrations. Throughout the paper, we work over the complex numbers $\mathbb{C}$.

Let $\pi : X \rightarrow Y$ be a proper morphism with X a nonsingular algebraic variety. The perverse t -structure on the constructible derived category $D_c^b(Y)$ with rational coefficients $\mathbb{Q}$ induces a finite and increasing filtration on the rational cohomology groups $H^*(X, \mathbb{Q})$

$$(1) \quad P_0 H^*(X, \mathbb{Q}) \subset P_1 H^*(X, \mathbb{Q}) \subset \cdots \subset P_k H^*(X, \mathbb{Q}) \subset \cdots \subset H^*(X, \mathbb{Q}),$$

called the *perverse filtration* associated with π . The filtration (1) is governed by the topology of the morphism $\pi : X \rightarrow Y$. See Section 1 for a brief review of the subject.

Every cohomology class on X is indexed by the perverse filtration (1). We say that a class $\alpha \in H^d(X, \mathbb{Q})$ has *perversity* k if $\alpha = 0$, or

$$\alpha \in P_k H^d(X, \mathbb{Q}) \text{ and } \alpha \notin P_{k-1} H^d(X, \mathbb{Q}).$$

The purpose of this paper is to study perverse filtrations associated with Hitchin fibrations in view of the *P=W conjecture* by de Cataldo, Hausel, and Migliorini [11]. Our method is to use *symmetries* induced by the monodromy of moduli spaces of sheaves on abelian surfaces.

Received by the editors October 3, 2019, and, in revised form, June 4, 2021.

2020 *Mathematics Subject Classification.* Primary 14H60, 14F45, 14D20.

The first author was partially supported by NSF grants DMS 1600515 and 1901975.

The second author was partially supported by NSF FRG grant DMS-1159265.

The third author was supported by the NSF grant DMS 2134315.

©2021 American Mathematical Society

0.2. The $P=W$ conjecture. Let C be an irreducible nonsingular projective curve of genus $g \geq 2$. There are two moduli spaces which are attached to the curve C , the reductive Lie group GL_r and an integer χ with $\gcd(r, \chi) = 1$. They are the twisted versions of Simpson's Dolbeault and Betti moduli spaces [42]; see [25] for the non-abelian Hodge theory in the twisted case.

The first moduli space $\mathcal{M}_{\mathrm{Dol}}$ parametrizes stable Higgs bundles on C

$$(\mathcal{E}, \theta), \quad \theta : \mathcal{E} \rightarrow \mathcal{E} \otimes \Omega_C$$

with $\mathrm{rank}(\mathcal{E}) = r$ and $\chi(\mathcal{E}) = \chi$. The variety $\mathcal{M}_{\mathrm{Dol}}$ admits a projective morphism with connected fibers

$$(2) \quad h : \mathcal{M}_{\mathrm{Dol}} \rightarrow \Lambda = \bigoplus_{i=1}^n H^0(C, \Omega_C^{\otimes i})$$

sending $(\mathcal{E}, \theta) \in \mathcal{M}_{\mathrm{Dol}}$ to the characteristic polynomial $\mathrm{char}(\theta) \in \Lambda$. The proper morphism (2), called the *Hitchin fibration*, is Lagrangian with respect to the canonical holomorphic symplectic form on $\mathcal{M}_{\mathrm{Dol}}$ given by the hyper-Kähler metric on $\mathcal{M}_{\mathrm{Dol}}$; see [27]. The second moduli space is the (twisted) character variety $\mathcal{M}_B$. It can be described (see [25], or [40] for an alternative description) as parametrizing isomorphism classes of irreducible local systems

$$\rho : \pi_1(C \setminus \{p\}) \rightarrow \mathrm{GL}_r,$$

where ρ sends a loop around a chosen point p to $\xi_r^\chi I_r \in \mathrm{GL}_r$, with ξ_r a primitive root of unity. The character variety $\mathcal{M}_B$ is affine.

In [41], Simpson constructed a diffeomorphism between the (un-twisted) moduli spaces $\mathcal{M}_{\mathrm{Dol}}$ and $\mathcal{M}_B$, called the non-abelian Hodge theory; see [25] for the twisted case. A striking prediction, suggested by the parallel between the Relative Hard Lefschetz [5] and Curious Hard Lefschetz [24] Theorems, was formulated by de Cataldo, Hausel, and Migliorini [11]; it predicts that the perverse filtration of $\mathcal{M}_{\mathrm{Dol}}$ with respect to the Hitchin fibration (2) matches the weight filtration of the mixed Hodge structure on $\mathcal{M}_B$ under the identification

$$H^*(\mathcal{M}_{\mathrm{Dol}}, \mathbb{Q}) = H^*(\mathcal{M}_B, \mathbb{Q})$$

induced by Simpson's (twisted) non-abelian Hodge theory.

Conjecture 0.1 ([11], $P=W$). *We have*

$$P_k H^*(\mathcal{M}_{\mathrm{Dol}}, \mathbb{Q}) = W_{2k} H^*(\mathcal{M}_B, \mathbb{Q}) = W_{2k+1} H^*(\mathcal{M}_B, \mathbb{Q}), \quad k \geq 0.$$

Conjecture 0.1 establishes a surprising connection between the topology of Hitchin fibrations and the Hodge theory of character varieties. It was proven in [11] in the case of $r = 2$ for any genus $g \geq 2$. See also [12, 39, 45] for certain parabolic cases, and [23, 38] for a compact analog concerning Lagrangian fibrations on projective holomorphic symplectic manifolds.

The first main result of this paper is a proof of Conjecture 0.1 for curves of genus 2 and arbitrary rank $r \geq 1$.

Theorem 0.2. *The $P=W$ Conjecture 0.1 holds when C has genus $g = 2$.*

For an irreducible nonsingular curve C of genus $g \geq 2$, it is a general fact that $P = W$ for GL_r implies $P = W$ for PGL_r ; see (70) and (90). Hence we conclude immediately from Theorem 0.2 that the $P = W$ conjecture holds for PGL_r when the curve C has genus $g = 2$.

As explained in [11, Section 1], the curious Poincaré duality and the Curious Hard Lefschetz conjectures [24, Conjectures 4.2.4 and 4.2.7] for character varieties are consequences of the P=W Conjecture 0.1. Moreover, by [7] and [35, Section 9.3], the P=W Conjecture 0.1 implies the correspondence between Gopakumar–Vafa invariants and Pandharipande–Thomas invariants [35, Conjecture 3.13] for the local Calabi–Yau 3-fold $T^*C \times \mathbb{C}$ in the curve class $r[C]$. Hence Theorem 0.2 verifies all these conjectures for genus 2 curves.¹

0.3. Tautological classes. Assume that C has genus $g \geq 2$ and $r \geq 1$. A set of generators for the cohomology rings identified by the non-abelian Hodge diffeomorphism

$$(3) \quad H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) = H^*(\mathcal{M}_B, \mathbb{Q})$$

is described in [30] by tautological classes.

Let

$$p_C : C \times \mathcal{M}_{\text{Dol}} \rightarrow C, \quad p_{\mathcal{M}} : C \times \mathcal{M}_{\text{Dol}} \rightarrow \mathcal{M}_{\text{Dol}}$$

be the projections. We say that a triple

$$(\mathcal{U}^\alpha, \theta) = (\mathcal{U}, \theta, \alpha)$$

is a *twisted universal family* over $C \times \mathcal{M}_{\text{Dol}}$ if $(\mathcal{U}, \theta)$ is a universal family and

$$\alpha = p_C^* \alpha_1 + p_{\mathcal{M}}^* \alpha_2 \in H^2(C \times \mathcal{M}_{\text{Dol}}, \mathbb{Q}),$$

with $\alpha_1 \in H^2(C, \mathbb{Q})$ and $\alpha_2 \in H^2(\mathcal{M}_{\text{Dol}}, \mathbb{Q})$.

For a twisted universal family $(\mathcal{U}^\alpha, \theta)$, we define the *twisted Chern character* $\text{ch}^\alpha(\mathcal{U})$ as

$$\text{ch}^\alpha(\mathcal{U}) = \text{ch}(\mathcal{U}) \cup \exp(\alpha) \in H^*(C \times \mathcal{M}_{\text{Dol}}, \mathbb{Q}),$$

and we denote by

$$\text{ch}_k^\alpha(\mathcal{U}) \in H^{2k}(C \times \mathcal{M}_{\text{Dol}}, \mathbb{Q})$$

its degree k part. The class $\text{ch}^\alpha(\mathcal{U})$ is called *normalized* if

$$\text{ch}_1^\alpha(\mathcal{U})|_{p \times \mathcal{M}_{\text{Dol}}} = 0 \in H^2(\mathcal{M}_{\text{Dol}}, \mathbb{Q}), \quad \text{ch}_1^\alpha(\mathcal{U})|_{C \times q} = 0 \in H^2(C, \mathbb{Q}),$$

with $p \in \mathcal{M}_{\text{Dol}}$ and $q \in C$ points. Since two universal families differ by the pull-back of a line bundle on $\mathcal{M}_{\text{Dol}}$, a straightforward calculation shows that normalized classes on $C \times \mathcal{M}_{\text{Dol}}$ exist and are uniquely determined. We introduce the tautological classes associated with the normalized class $\text{ch}^\alpha(\mathcal{U})$ as follows.

For any $\gamma \in H^i(C, \mathbb{Q})$, let $c(\gamma, k)$ denote the *tautological class*

$$(4) \quad c(\gamma, k) := \int_\gamma \text{ch}_k^\alpha(\mathcal{U}) = p_{\mathcal{M}*}(p_C^* \gamma \cup \text{ch}_k^\alpha(\mathcal{U})) \in H^{i+2k-2}(\mathcal{M}_{\text{Dol}}, \mathbb{Q}).$$

Markman showed in [30] that the classes $c(\gamma, k)$ generate the cohomology ring (3) as a $\mathbb{Q}$ -algebra. Furthermore, Shende proved in [40] (cf. Lemma 4.6) that

$$(5) \quad c(\gamma, k) \in {}^k\text{Hdg}^{i+2k-2}(\mathcal{M}_B), \quad \forall \gamma \in H^i(C, \mathbb{Q}),$$

where

$${}^k\text{Hdg}^d(\mathcal{M}_B) = W_{2k}H^d(\mathcal{M}_B, \mathbb{Q}) \cap F^kH^d(\mathcal{M}_B, \mathbb{C}) \cap \bar{F}^kH^d(\mathcal{M}_B, \mathbb{C}),$$

with F^* and W_* the Hodge and weight filtrations, respectively. It follows that the rational cohomology $H^*(\mathcal{M}_B, \mathbb{Q})$ is split of Hodge–Tate type, and there is a

¹As we discuss later, the curious Poincaré and Lefschetz conjectures have been recently shown by Mellit [36] to hold for all genera.

canonical decomposition of the graded vector spaces (3) induced by the mixed Hodge theory of $\mathcal{M}_B$,

$$(6) \quad H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) = H^*(\mathcal{M}_B, \mathbb{Q}) = \bigoplus_{k,d} {}^k\text{Hdg}^d(\mathcal{M}_B).$$

The P=W conjecture, which can be restated as predicting that

$$(7) \quad P_k H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) = \bigoplus_{k' \leq k} {}^{k'}\text{Hdg}^*(\mathcal{M}_B),$$

is equivalent to the following statements.

Conjecture 0.3 (Equivalent version of P=W). *There exists a splitting $G_* H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q})$ of the perverse filtration associated with the Hitchin fibration (2) satisfying the following two properties.*

- (a) (Tautological classes) *Every tautological class $c(\gamma, k)$ has perversity k and in fact*

$$c(\gamma, k) \in G_k H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}), \quad \forall k \geq 0, \forall \gamma \in H^*(C, \mathbb{Q}).$$

- (b) (Multiplicativity) *The perverse decomposition is multiplicative, i.e.,*

$$\cup : G_k H^d(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) \times G_{k'} H^{d'}(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) \rightarrow G_{k+k'} H^{d+d'}(\mathcal{M}_{\text{Dol}}, \mathbb{Q}).$$

We recall that the multiplicativity of the weight filtration

$$\cup : W_k H^d(X, \mathbb{Q}) \times W_{k'} H^{d'}(X, \mathbb{Q}) \rightarrow W_{k+k'} H^{d+d'}(X, \mathbb{Q})$$

is standard from mixed Hodge theory. However, the perverse filtration associated with a proper flat morphism is not multiplicative in general; see [8, Exercise 5.6.8]. It is mysterious why the perverse filtration associated with the Hitchin fibration (2) should be multiplicative, as is predicted by the P=W conjecture. In fact, one consequence of this paper is that the full P=W conjecture can be reduced to the multiplicativity of the perverse filtration; see Theorem 0.6. This approach of analyzing the P=W conjecture via Conjecture 0.3 goes back to [11] where it is applied to prove the case of rank 2.

0.4. P=W for tautological classes. Theorem 0.2 is concerned with genus $g = 2$. In this section, we state our results for genus $g \geq 2$.

0.4.1. Even tautological classes. We consider the $\mathbb{Q}$ -subalgebra

$$R^*(\mathcal{M}_{\text{Dol}}) \subset H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) = H^*(\mathcal{M}_B, \mathbb{Q})$$

generated by all the *even* tautological classes

$$c(\gamma, k), \quad k \geq 0, \quad \gamma \in H^0(C, \mathbb{Q}) \oplus H^2(C, \mathbb{Q}).$$

Theorem 0.4 establishes the P=W conjecture in arbitrary genus and rank, but restricted to this subalgebra.

Theorem 0.4. *The P=W conjecture holds, for any genus $g \geq 2$ and rank $r \geq 1$ for $R^*(\mathcal{M}_{\text{Dol}})$, i.e.*

$$\begin{aligned} P_k H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) \cap R^*(\mathcal{M}_{\text{Dol}}) &= W_{2k} H^*(\mathcal{M}_B, \mathbb{Q}) \cap R^*(\mathcal{M}_{\text{Dol}}) \\ &= W_{2k+1} H^*(\mathcal{M}_B, \mathbb{Q}) \cap R^*(\mathcal{M}_{\text{Dol}}). \end{aligned}$$

In fact, in the proof of Theorem 0.4, we obtain a multiplicative decomposition

$$R^*(\mathcal{M}_{\text{Dol}}) = \bigoplus_{k,d} G_k R^d(\mathcal{M}_{\text{Dol}})$$

that splits the restricted perverse filtration $P_k H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) \cap R^*(\mathcal{M}_{\text{Dol}})$ and such that

$$G_k R^d(\mathcal{M}_{\text{Dol}}) = {}^k \text{Hdg}^d(\mathcal{M}_B) \cap R^d(\mathcal{M}_{\text{Dol}}).$$

We refer to Section 4.6 for more details.

0.4.2. Odd tautological classes. Theorem 0.5 concerns odd tautological classes

$$(8) \quad c(\gamma, k) \in H^{2k-1}(\mathcal{M}_{\text{Dol}}, \mathbb{Q}), \quad \forall k \geq 0, \forall \gamma \in H^1(C, \mathbb{Q}).$$

Theorem 0.5. *Any odd tautological class (8) has perversity k .*

Theorems 0.4 and 0.5 provide strong evidence for the P=W Conjecture 0.1 (and for its reformulation Conjecture 0.3) for any genus $g \geq 2$. In particular, we obtain that all tautological generators $c(\gamma, k)$ satisfy the $P = W$ match:

$$(9) \quad c(\gamma, k) \in P_k H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}), \quad \forall \gamma \in H^*(C, \mathbb{Q}).$$

0.4.3. Multiplicativity is equivalent to $P=W$. The P=W Conjecture 0.1 implies immediately that the perverse filtration $P_* H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q})$ is multiplicative, *i.e.*

$$(10) \quad \cup : P_k H^d(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) \times P_{k'} H^{d'}(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) \rightarrow P_{k+k'} H^{d+d'}(\mathcal{M}_{\text{Dol}}, \mathbb{Q}).$$

Theorem 0.6 shows that the converse is also true. It is a corollary of Theorems 0.4 and 0.5, and of the Curious Hard Lefschetz conjecture recently established by Mellit [36].

Theorem 0.6. *The $P=W$ Conjecture 0.1 is equivalent to the multiplicativity (10) of the perverse filtration.*

In fact, we deduce from (5), (9), and (10) that

$$W_{2i} H^*(\mathcal{M}_B, \mathbb{Q}) \subset P_i H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}), \quad \forall i \geq 0,$$

which further implies Conjecture 0.1 by [36, Theorem 1.5.3] and the second paragraph following [36, Corollary 1.5.4].

Remark 0.7. All the results of this paper hold for arbitrary rank $r \geq 1$. The case $r = 1$ is classical and easy: the Dolbeault moduli space is the cotangent bundle of a Jacobian of some degree of the curve and the Betti moduli space is a product of $\mathbb{G}_m^{2g}$. In this paper, starting with the assumption $\beta^2 \geq 6$ in Section 2.1, we focus on the case of rank $r \geq 2$, which corresponds to $\beta^2 \geq 8$. The case of rank one could be dealt with also by using the methods of this paper, but at the price of some easy modifications that we deemed a distraction.

0.5. Idea of the proof. The key idea in proving our results is to first study perverse filtrations for certain *compact* hyperkähler manifolds, motivated in part by earlier work of the third author and Yin [38]. That is, we consider an abelian surface A with ample curve class $\beta \in H_2(A, \mathbb{Z})$, and prove analogs of our main result for the moduli space $\mathcal{M}_{\beta, A}$ of one-dimensional sheaves with support in the class β ; see Theorem 2.1 for the precise statement. The advantage of the compact geometry is that we can use results of Markman [34] on monodromy operators for $\mathcal{M}_{\beta, A}$. These are symmetries, arising from parallel transport and Fourier-Mukai transforms, that

do not appear in the Hitchin setting and which heavily constrain tautological classes of universal families. We show that these operators relate perverse filtrations and tautological classes for different choices of β and, in particular, allow us to pass from the case of imprimitive β to primitive β , which can be studied directly using [39, 45].

In order to apply this to Hitchin fibrations, we consider the degeneration to the normal cone of an embedding of a genus g curve C into an abelian surface

$$j_C : C \hookrightarrow A,$$

and study the specialization map on cohomology of the associated moduli spaces. In general, there is a great deal of information loss in this specialization because the family is non-proper. Nevertheless, we are able to show its compatibility with tautological classes and perverse splittings. Finally we use these compatibilities to deduce our main theorems for such a curve C ; this implies our results hold for all curves in view of [13].

0.6. Outline of paper. We briefly outline the contents of this paper. In Section 1, we recall some basic facts about perverse filtrations and earlier results of [39, 45] for Hilbert schemes of points on the special abelian surface $E \times E'$. In Section 2, we pass to the setting of abelian surfaces. One key result here is the splitting Theorem 2.1. After recalling Markman's work on monodromy operators, we prove the result for primitive classes β by a direct analysis, and then we combine the primitive case with Markman's results to establish the general case. In Section 3, we formulate and prove Theorem 3.8, which is a strengthened version of the splitting of Theorem 2.1 and which is more robust for specialization arguments. Finally, in Section 4, we study specialization maps for our degeneration, and we use them to prove the main theorems.

While this paper is written in logical order, some readers may prefer to start from Section 4, and refer backwards as needed.

1. PERVERSE FILTRATIONS

1.1. Overview. We begin with the definition and some relevant properties of the perverse filtration associated with a proper surjective morphism $\pi : X \rightarrow Y$. Some references are [5, 15, 16, 43]. Throughout this section, for simplicity, we assume X and Y are nonsingular.

Following [12, 39, 45], we discuss the perverse filtration for the Hilbert scheme of n -points on an abelian surface $E \times E'$ product of two elliptic curves, induced by the natural second projection morphism

$$E \times E' \rightarrow E'.$$

This example plays a crucial role in Section 2 concerning moduli of sheaves on abelian surfaces.

1.2. Perverse sheaves. Let $D_c^b(Y)$ denote the bounded derived category of $\mathbb{Q}$ -constructible sheaves on Y , and let $\mathbb{D} : D_c^b(Y)^{\text{op}} \rightarrow D_c^b(Y)$ be the Verdier duality functor. The full subcategories

$$\begin{aligned} {}^pD_{\leq 0}^b(Y) &= \{ \mathcal{E} \in D_c^b(Y) : \dim \text{Supp}(\mathcal{H}^i(\mathcal{E})) \leq -i \}, \\ {}^pD_{\geq 0}^b(Y) &= \{ \mathcal{E} \in D_c^b(Y) : \dim \text{Supp}(\mathcal{H}^i(\mathbb{D}\mathcal{E})) \leq -i \} \end{aligned}$$

give rise to the *perverse t -structure* on $D_c^b(Y)$, whose heart is the abelian category of perverse sheaves,

$$\mathrm{Perv}(Y) \subset D_c^b(Y).$$

For $k \in \mathbb{Z}$, let ${}^p\tau_{\leq k}$ be the truncation functor associated with the perverse t -structure. Given an object $\mathcal{C} \in D_c^b(Y)$, there is a natural morphism

$$(11) \quad {}^p\tau_{\leq k}\mathcal{C} \rightarrow \mathcal{C}.$$

For the morphism $\pi : X \rightarrow Y$, we obtain from (11) the morphism

$${}^p\tau_{\leq k}R\pi_*\mathbb{Q}_X \rightarrow R\pi_*\mathbb{Q}_X,$$

which further induces a morphism of (hyper-)cohomology groups

$$(12) \quad H^{d-(\dim X - R)}\left(Y, {}^p\tau_{\leq k}(R\pi_*\mathbb{Q}_X[\dim X - R])\right) \rightarrow H^d(X, \mathbb{Q}).$$

Here

$$(13) \quad R = \dim X \times_Y X - \dim X$$

is the *defect of semismallness*. The k -th piece of the perverse filtration (1)

$$P_k H^d(X, \mathbb{Q}) \subset H^d(X, \mathbb{Q})$$

is defined to be the image of (12).²

We say that a graded vector space decomposition

$$H^*(X, \mathbb{Q}) = \bigoplus_{k,d} G_k H^d(X, \mathbb{Q})$$

splits the perverse filtration if

$$P_k H^d(X, \mathbb{Q}) = \bigoplus_{i \leq k} G_i H^d(X, \mathbb{Q}), \quad \forall k, d.$$

1.3. The decomposition theorem. Perverse filtrations can be described through the decomposition theorem [5, 15]. By applying the decomposition theorem to the morphism $\pi : X \rightarrow Y$, we obtain an isomorphism

$$(14) \quad R\pi_*\mathbb{Q}_X[\dim X - R] \simeq \bigoplus_{i=0}^{2R} \mathcal{P}_i[-i] \in D_c^b(Y)$$

with $\mathcal{P}_i \in \mathrm{Perv}(Y)$. The perverse filtration can be identified as

$$P_k H^d(X, \mathbb{Q}) = \mathrm{Im}\left\{H^{d-(\dim X - R)}\left(Y, \bigoplus_{i=0}^k \mathcal{P}_i[-i]\right) \rightarrow H^d(X, \mathbb{Q})\right\}.$$

In general, the isomorphism (14) in the decomposition theorem is not canonical. Once we fix such an isomorphism ϕ , we obtain a splitting $G_* H^*(X, \mathbb{Q})$ of the perverse filtration,

$$P_k H^d(X, \mathbb{Q}) = \bigoplus_{i \leq k} G_i H^d(X, \mathbb{Q}).$$

Here

$$G_i H^d(X, \mathbb{Q}) = \mathrm{Im}\left\{H^{d-(\dim X - R)}(Y, \mathcal{P}_i[-i]) \rightarrow H^d(X, \mathbb{Q})\right\}$$

²Here the shift $[\dim X - R]$ is to ensure that the perverse filtration is concentrated in the degrees $[0, 2R]$.

with

$$H^{d-(\dim X-R)}(Y, \mathcal{P}_i[-i]) \rightarrow H^d(X, \mathbb{Q})$$

the morphism induced by ϕ^{-1} .

1.4. Perverse filtration for projective bases. We review another description of the perverse filtration associated with

$$\pi : X \rightarrow Y,$$

when Y is projective.

We fix η to be an ample class on Y , and we consider

$$L = \pi^* \eta \in H^2(X, \mathbb{Q}).$$

The class L acts on $H^*(X, \mathbb{Q})$ as an nilpotent operator via cup product

$$L : H^*(X, \mathbb{Q}) \xrightarrow{\cup L} H^*(X, \mathbb{Q}).$$

Proposition 1.1 shows that the filtration (1) is completely described by an ample class on the base. It is stated in the special case needed in this paper.

Proposition 1.1 ([15, Proposition 5.2.4]). *Assume that*

$$\dim X = 2 \dim Y = 2R.$$

Then we have

$$P_k H^m(X, \mathbb{Q}) = \sum_{i \geq 1} (\text{Ker}(L^{R+k+i-m}) \cap \text{Im}(L^{i-1})) \cap H^m(X, \mathbb{Q}).$$

1.5. Abelian surfaces and Hilbert schemes. Let $A = E \times E'$ be an abelian surface with E and E' elliptic curves, and let $n \in \mathbb{Z}^{\geq 0}$. The natural projection

$$p : A \rightarrow E'$$

induces a morphism

$$p_n : A^{[n]} \rightarrow E'^{(n)}$$

from the Hilbert scheme of n points on A to the symmetric product of the elliptic curve E' . We briefly review the construction [39] of a *canonical* splitting

$$(15) \quad H^*(A^{[n]}, \mathbb{Q}) = \bigoplus_{i,d} \tilde{G}_i H^d(A^{[n]}, \mathbb{Q})$$

of the perverse filtration associated with p_n , which culminates with the explicit formula (18).

We start with a canonical decomposition

$$(16) \quad H^*(A, \mathbb{Q}) = \bigoplus_{i,d} \tilde{G}_i H^d(A, \mathbb{Q})$$

with $\tilde{G}_i H^d(A, \mathbb{Q})$ the Künneth factor

$$H^i(E, \mathbb{Q}) \otimes H^{d-i}(E', \mathbb{Q}) \subset H^d(A, \mathbb{Q}).$$

Since $p : A \rightarrow E'$ is a trivial fibration, (16) splits the perverse filtration associated with p . Assume that we already have decompositions of $H^*(X_1, \mathbb{Q})$ and $H^*(X_2, \mathbb{Q})$; a decomposition of $H^*(X_1 \times X_2, \mathbb{Q})$ can be constructed by using the Künneth

decomposition. In particular, we obtain the direct sum decomposition of $H^*(A^n, \mathbb{Q})$ with summands

$$\tilde{G}_k H^*(A^n, \mathbb{Q}) = \left\langle \alpha_1 \boxtimes \cdots \boxtimes \alpha_n; \alpha \in \tilde{G}_{k_i} H^*(A, \mathbb{Q}), \sum_i k_i = k \right\rangle,$$

whose $\mathfrak{S}_n$ -invariant part induces a decomposition

$$H^*(A^{(n)}, \mathbb{Q}) = \bigoplus \tilde{G}_i H^d(A^{(n)}, \mathbb{Q}).$$

In turn, by using Künneth decompositions again, this gives us canonical decompositions for

$$H^*(A^{(n_1)} \times A^{(n_2)} \times \cdots \times A^{(n_k)}, \mathbb{Q}), \quad n_i \geq 1.$$

Finally, the cohomology of the Hilbert scheme $A^{[n]}$ is related to the cohomology of symmetric products by [14, 22]. We employ exponential notation for partitions $\nu := (\nu_i)_{i=1}^l$, with $\nu_i > 0$ and $\sum_i \nu_i = n$, of a positive integer n . Namely, we let a_k be the number of times the integer k appears in the partition, so that the expression

$$1^{a_1} 2^{a_2} \cdots n^{a_n}$$

recovers the partition ν . We use $A^{(\nu)}$ to denote the variety $A^{(a_1)} \times A^{(a_2)} \times \cdots \times A^{(a_n)}$. The cohomology group $H^d(A^{[n]}, \mathbb{Q})$ admits a canonical decomposition

$$(17) \quad H^d(A^{[n]}, \mathbb{Q}) = \bigoplus_{\nu} H^{d+2l(\nu)-2n}(A^{(\nu)}, \mathbb{Q}),$$

where ν runs through all partitions of n and $l(\nu)$ is the length of ν . The desired canonical splitting (15) is then defined by setting

$$(18) \quad \tilde{G}_k H^d(A^{[n]}, \mathbb{Q}) = \bigoplus_{\nu} \tilde{G}_{k+l(\nu)-n} H^{d+2l(\nu)-2n}(A^{(\nu)}, \mathbb{Q})$$

to be the sub-vector space of $H^d(A^{[n]}, \mathbb{Q})$ under the identification (17). By [45, Proposition 4.12], this decomposition splits the perverse filtration associated with p_n .³

In view of [45, Proposition 2.1], by using the Künneth decomposition again, we obtain a canonical splitting

$$(19) \quad H^*(A \times A^{[n]}, \mathbb{Q}) = \bigoplus_{i,d} \tilde{G}_i H^d(A \times A^{[n]}, \mathbb{Q})$$

of the perverse filtration associated with

$$p \times p_n : A \times A^{[n]} \rightarrow E' \times E'^{(n)}.$$

Theorem 1.2 obtained in [39, 45] is concerned with tautological classes and multiplicativity in the context of Hilbert schemes.

Theorem 1.2. *Let $\mathcal{I}_n$ be the universal ideal sheaf on $A \times A^{[n]}$.*

(a) *(Tautological classes) We have*

$$\mathrm{ch}_k(\mathcal{I}_n) \in \tilde{G}_k H^{2k}(A \times A^{[n]}, \mathbb{Q}).$$

(b) *(Multiplicativity) The decomposition (15) is multiplicative, i.e.*

$$\cup : \tilde{G}_i H^d(A^{[n]}, \mathbb{Q}) \times \tilde{G}_{i'} H^{d'}(A^{[n]}, \mathbb{Q}) \rightarrow \tilde{G}_{i+i'} H^{d+d'}(A^{[n]}, \mathbb{Q}).$$

³We will give another interpretation of the splitting $\tilde{G}_* H^*(A^{[n]}, \mathbb{Q})$ in Corollary 3.7.

Proof. Theorem 1.2(b) was proven in [45]. Although the multiplicativity was shown only for the perverse filtration in [45], the same proof works for the decomposition (15). This was explained in [39, Proposition 1.8]. Theorem 1.2(a) follows directly from the proof of [39, Theorem 0.1] and the proof of [39, Theorem 3.3], where it is explained how to treat perverse decompositions instead of perverse filtrations. $\square$

2. MODULI OF ONE-DIMENSIONAL SHEAVES

2.1. Overview and main results. Throughout this section, we assume A is an abelian surface and fix $\chi \in \mathbb{Z}$. Let $\beta \in H_2(A, \mathbb{Z})$ be an ample curve class with $\beta^2 \geq 6$; we only consider classes β for which the vector $(0, \beta, \chi) \in H^{\text{ev}}(A, \mathbb{Z})$ is primitive. Finally, we assume that we are given a polarization H that is generic with respect to the vector $(0, \beta, \chi)$. We will typically omit χ and H from our notation.

Let $\mathcal{M}_{\beta, A}$ be the moduli space which parametrizes one-dimensional sheaves satisfying

$$(20) \quad [\text{supp}(\mathcal{F})] = \beta, \quad \chi(\mathcal{F}) = \chi,$$

that are Gieseker-stable with respect to our generic polarization. In the paper, by “support”, we mean Fitting support (see [28]), and the square bracket denotes the associated homology class. Equivalently, these are Gieseker-stable sheaves on A with Mukai vector $(0, \beta, \chi)$. By [44], the moduli space $\mathcal{M}_{\beta, A}$ is a non-empty, nonsingular projective variety of dimension $\beta^2 + 2$.

Let

$$\pi_A : A \times \mathcal{M}_{\beta, A} \rightarrow A, \quad \pi_{\mathcal{M}} : A \times \mathcal{M}_{\beta, A} \rightarrow \mathcal{M}_{\beta, A}$$

be the projections. If there exists a universal family $\mathcal{F}_{\beta}$ on $A \times \mathcal{M}_{\beta, A}$, then, in analogy to Section 0.3, we define the twisted Chern character associated with a class

$$(21) \quad \alpha = \pi_A^* \alpha_1 + \pi_{\mathcal{M}}^* \alpha_2 \in H^2(A \times \mathcal{M}_{\beta, A}, \mathbb{C})$$

by setting

$$\text{ch}^{\alpha}(\mathcal{F}_{\beta}) = \text{ch}(\mathcal{F}_{\beta}) \cup \exp(\alpha) \in H^*(A \times \mathcal{M}_{\beta, A}, \mathbb{C})$$

(the use of $\mathbb{C}$ -coefficients is for later use in Theorem 2.1), and we denote its degree $2k$ part by

$$\text{ch}_k^{\alpha}(\mathcal{F}_{\beta}) \in H^{2k}(A \times \mathcal{M}_{\beta, A}, \mathbb{C}).$$

In general, a universal family may not exist; however, as explained in [31, Section 3.1], there exists a *universal class*

$$[\mathcal{F}_{\beta}] \in K_{\text{top}}(A \times \mathcal{M}_{\beta, A}),$$

which is well-defined up to pullback of a topological line bundle on $\mathcal{M}_{\beta, A}$.⁴ As before, given a choice of α as in (21), we can take its associated twisted Chern character

$$\text{ch}^{\alpha}([\mathcal{F}_{\beta}]) \in H^*(A \times \mathcal{M}_{\beta, A}, \mathbb{C}),$$

which we still denote by $\text{ch}^{\alpha}(\mathcal{F}_{\beta})$ for notational convenience.

The moduli space $\mathcal{M}_{\beta, A}$ admits a natural Hilbert–Chow morphism

$$(22) \quad \pi_{\beta} : \mathcal{M}_{\beta, A} \rightarrow B, \quad \pi_{\beta}(\mathcal{F}) = [\text{supp}(\mathcal{F})],$$

⁴Here $K_{\text{top}}(X)$ denotes the Grothendieck ring of topological complex vector bundles on X [1].

where B is the Chow variety parametrizing effective one-cycles in the class β ; see [28, 29]. The purpose of this section is to prove Theorem 2.1, which can be viewed as an analog of Conjecture 0.3 for abelian surfaces.

Theorem 2.1. *There exists a choice of universal twisted class $\text{ch}^\alpha(\mathcal{F}_\beta)$ with α of type (21), and a splitting $\tilde{G}_*H^*(\mathcal{M}_{\beta,A}, \mathbb{C})$ of the perverse filtration associated with the Hilbert-Chow morphism π_β (22), satisfying the following properties.*

(a) *(Tautological classes) For any $\gamma \in H^*(A, \mathbb{C})$, we have*

$$\int_\gamma \text{ch}_k^\alpha(\mathcal{F}_\beta) \in \tilde{G}_{k-1}H^*(\mathcal{M}_{\beta,A}, \mathbb{C}).$$

(b) *(Multiplicativity) We have*

$$\cup : \tilde{G}_i H^d(\mathcal{M}_{\beta,A}, \mathbb{C}) \times \tilde{G}_{i'} H^{d'}(\mathcal{M}_{\beta,A}, \mathbb{C}) \rightarrow \tilde{G}_{i+i'} H^{d+d'}(\mathcal{M}_{\beta,A}, \mathbb{C}).$$

Theorem 2.1 will be deduced from Theorem 1.2 and Markman's results on the monodromy of holomorphic symplectic varieties [32, 34]. The strategy of the argument is to use Theorem 1.2 and direct computations to study the case of primitive curve classes for special abelian varieties. To pass to imprimitive curve classes (which form a distinct deformation class of Lagrangian fibrations), we require a mild extension of Markman's work to the setting of complex coefficients.

In the next three sections, we review Markman's work, and in the subsequent three sections, we prove Theorem 2.1.

2.2. Mukai lattices. Let

$$H^*(A, \mathbb{Z}) = S_A^+ \oplus S_A^-$$

be the even/odd decomposition of the cohomology $H^*(A, \mathbb{Z})$ with

$$S_A^+ = H^0(A, \mathbb{Z}) \oplus H^2(A, \mathbb{Z}) \oplus H^4(A, \mathbb{Z}), \quad S_A^- = H^1(A, \mathbb{Z}) \oplus H^3(A, \mathbb{Z}).$$

The Mukai pairing on S_A^+ is given by

$$(23) \quad \langle a, b \rangle = \int_A (a_1 \cup b_1 - a_0 \cup b_2 - a_2 \cup b_0),$$

where $a = (a_0, a_1, a_2)$ and $b = (b_0, b_1, b_2)$ are elements in S_A^+ . We have

$$(S_A^+, \langle \cdot, \cdot \rangle) \cong U^{\oplus 4}$$

with U the parabolic plane. For a coherent sheaf $\mathcal{F}$ on A , its Mukai vector

$$v(\mathcal{F}) = \text{ch}(\mathcal{F}) \cup \sqrt{\text{td}_A} \in S_A^+$$

coincides with the Chern character $\text{ch}(\mathcal{F})$.

Throughout this section, we assume $v \in S_A^+$ is a primitive vector such that the moduli space $M_{v,A}$ of Gieseker-stable sheaves $\mathcal{F}$ on A (with respect to a very general polarization H) with Mukai vector $v(\mathcal{F}) = v$ is non-empty. By [44, Theorem 0.1], the variety $M_{v,A}$ is nonsingular and projective of dimension

$$\dim(M_{v,A}) = \langle v, v \rangle + 2,$$

and it is deformation equivalent to

$$M_{(1,0,-n),A} = A^{[n]} \times \hat{A},$$

where $\hat{A}$ is the dual abelian variety of A and $2n = \langle v, v \rangle$.

A universal family on

$$A \times M_{(1,0,-n),A} = A \times A^{[n]} \times \hat{A}$$

is

$$(24) \quad \mathcal{E}_n = \pi_{12}^* \mathcal{I}_n \otimes \pi_{13}^* \mathcal{P},$$

with $\mathcal{I}_n$ the universal ideal sheaf, $\mathcal{P}$ the normalized Poincaré line bundle on $A \times \hat{A}$, and π_{ij} the corresponding projections.

2.3. Spin representations. The even cohomology S_A^+ together with the Mukai pairing is a unimodular lattice. Consider the corresponding Spin group for this lattice, denoted by $\text{Spin}(S_A^+)$. We recall certain representations of this group involved in Markman's work following [34, Section 3].

Let $H^*(A, \mathbb{Z})$ denote the total cohomology of A equipped with the pairing

$$(25) \quad (a, b)_A = \int_A \tau(a) \cup b,$$

where τ acts on $H^i(A, \mathbb{Z})$ by $(-1)^{i(i-1)/2}$. The group $\text{Spin}(S_A^+)$ admits an action by parity-preserving isometries on $H^*(A, \mathbb{Z})$

$$\text{Spin}(S_A^+) \rightarrow \text{Aut}(H^*(A, \mathbb{Z})),$$

which extends the natural representation on S_A^+ .

We consider V to be the lattice

$$V = H^1(A, \mathbb{Z}) \oplus H^1(\hat{A}, \mathbb{Z})$$

carrying a symmetric bilinear form via the canonical identification

$$H^1(\hat{A}, \mathbb{Z}) = H^1(A, \mathbb{Z})^*.$$

There exists a representation

$$\text{Spin}(S_A^+) \rightarrow \text{Aut}(V)$$

by isometries which we can extend to an action of $\text{Spin}(S_A^+)$ on $\bigwedge^k V = H^k(A \times \hat{A}, \mathbb{Z})$ for each k . This action will play a role in the proof of Theorem 2.8.

These constructions are linear-algebraic and make sense for the groups $\text{Spin}(S_A^+ \otimes \mathbb{K})$ with general coefficients

$$\mathbb{K} = \mathbb{Z}, \mathbb{Q}, \mathbb{R}, \text{ or } \mathbb{C}.$$

2.4. Monodromy representations. In this section, we recall Markman's monodromy operators [34] for abelian surfaces. They play a crucial role in the proof of Theorem 2.1. We refer to [32] for a parallel theory concerning $K3$ surfaces.

Let $H^*(A)$ be the cohomology $H^*(A, \mathbb{K})$ with coefficients $\mathbb{K} = \mathbb{Z}$ or $\mathbb{C}$. Assume that

$$g : H^*(A) \rightarrow H^*(A)$$

is a parity-preserving homomorphism induced by an element (cf. Section 2.3)

$$g \in \text{Spin}(S_A^+ \otimes \mathbb{K})$$

which fixes a primitive Mukai vector $v \in S_A^+$. Assume we are given a generic polarization H with respect to v . We explain how to construct a graded $\mathbb{K}$ -algebra automorphism

$$\gamma_{g,v} : H^*(M_{v,A}, \mathbb{K}) \rightarrow H^*(M_{v,A}, \mathbb{K}).$$

For a projective variety X , we define

$$l : \bigoplus_i H^{2i}(X, \mathbb{Q}) \rightarrow \bigoplus_i H^{2i}(X, \mathbb{Q})$$

to be the universal polynomial map which takes the exponential Chern character to its total Chern class,

$$l(r + a_1 + a_2 + \cdots) = 1 + a_1 + (a_1^2/2 - a_2) + \cdots,$$

and we define

$$(-)^\vee : \bigoplus_i H^{2i}(X, \mathbb{Z}) \rightarrow \bigoplus_i H^{2i}(X, \mathbb{Z}), \quad \omega \mapsto \omega^\vee$$

to be the dualizing automorphism which acts by $(-1)^i$ on the even cohomology $H^{2i}(X, \mathbb{Z})$.

Let $[\mathcal{E}_v]$ and $[\mathcal{E}'_v]$ be universal classes on $A \times M_{v,A}$, in the sense of Section 2.1. We denote by $x_{g,v}$ the class

$$x_{g,v} = \pi_{12}^*[(\text{id} \otimes g)\text{ch}(\mathcal{E}_v)]^\vee \cup \pi_{23}^*\text{ch}(\mathcal{E}'_v) \in H^*(M_{v,A} \times A \times M_{v,A}, \mathbb{K}),$$

where π_{ij} are the projections from the product $M_{v,A} \times A \times M_{v,A}$ to the corresponding factors. Via the formalism of correspondences, to define an endomorphism of cohomology, it suffices to give a cohomology class in the product; in our case, the morphism $\gamma_{g,v}$ is defined by the class ()

$$(26) \quad \gamma_{g,v}([\mathcal{E}_v], [\mathcal{E}'_v]) = \left([l(\pi_{13*}x_{g,v})]^{-1} \right)_{\deg 2d_M} \in H^{2d_M}(M_{v,A} \times M_{v,A}, \mathbb{K}),$$

where $d_M = \dim(M_{v,A})$, and $-\deg k$ denotes the degree k part of a cohomology class $-$.

Markman proves the following statements regarding $\gamma_{g,v}([\mathcal{E}_v], [\mathcal{E}'_v])$ when $\mathbb{K} = \mathbb{Z}$

- (a) $\gamma_{g,v} := \gamma_{g,v}([\mathcal{E}_v], [\mathcal{E}'_v])$ is independent of the choice of universal classes $[\mathcal{E}_v]$ and $[\mathcal{E}'_v]$.
- (b) $\gamma_{g,v}$ is a graded $\mathbb{K}$ -algebra automorphism.

Consider the subgroup

$$\text{Spin}(S_A^+)_v \subset \text{Spin}(S_A^+)$$

preserving v . By [34, Corollary 8.4], every $\gamma_{g,v}$ with $g \in \text{Spin}(S_A^+)_v$ is a graded ring automorphism of $H^*(M_{v,A}, \mathbb{Z})$ and the relation

$$(27) \quad \gamma_{g_1,v} \circ \gamma_{g_2,v} = \gamma_{g_1 g_2, v}$$

holds. Therefore, we have a group homomorphism (Markman's monodromy operators)

$$(28) \quad \text{mon} : \text{Spin}(S_A^+)_v \rightarrow \text{Aut}(H^*(M_{v,A}, \mathbb{Z})), \quad \text{mon}(g) = \gamma_{g,v}.$$

Theorem 2.2 is proved by Markman for $\mathbb{Z}$ coefficients in [32, 34]. The extension of (28) to $\mathbb{C}$ -coefficients is crucial in the proof of Theorem 2.1.

Theorem 2.2. *For $g \in \text{Spin}(S_A^+ \otimes \mathbb{C})_v$, the class $\gamma_{g,v} := \gamma_{g,v}([\mathcal{E}_v], [\mathcal{E}'_v])$ is independent of the choice of universal classes $[\mathcal{E}_v]$ and $[\mathcal{E}'_v]$. The morphism (28) extends to a group homomorphism*

$$\text{mon} : \text{Spin}(S_A^+ \otimes \mathbb{C})_v \rightarrow \text{Aut}(H^*(M_{v,A}, \mathbb{C})).$$

Moreover, for a fixed universal class $[\mathcal{E}_v]$ and any $g \in \text{Spin}(S_A^+ \otimes \mathbb{C})_v$, there exists $\alpha \in H^2(M_{v,A}, \mathbb{C})$ so that we have

$$(29) \quad (g \otimes \gamma_{g,v})\text{ch}(\mathcal{E}_v) = \text{ch}(\mathcal{E}_v) \cup \pi_M^* \exp(\alpha),$$

where $\pi_M : A \times M_{v,A} \rightarrow M_{v,A}$ is the second projection.

Proof. We deduce this theorem from the case of integral coefficients using the Zariski-density of integral points

$$\text{Spin}(S_A^+)_v \subset \text{Spin}(S_A^+ \otimes \mathbb{C})_v,$$

which follows from the Borel Density Theorem [6] applied to the simple group

$$\text{Spin}(S_A^+)_v \cong \text{Spin}(v^\perp), \quad v^\perp \subset S_A^+.$$

The first claim, on the independence of universal classes, expresses a Zariski-closed condition on the element g , so that it can be deduced from (a) above; see [32, Lemma 3.11].

For the remaining claims, as in the proof of [34, Corollary 8.4], we can reduce the theorem to the case $\text{rank}(v) > 0$. It follows that the class α in the equation (29) is uniquely determined by

$$\text{rank}(v) \cdot \pi_M^* \alpha = c_1([\mathcal{E}_v]) - [(g \otimes \gamma_{g,v})\text{ch}([\mathcal{E}_v])]_{\deg 2}.$$

The operator $\gamma_{g,v}$ (26) arises as a grading-preserving element in

$$\text{End}(H^*(M_{v,A}, \mathbb{C})).$$

We need to show that for any $g \in \text{Spin}(S_A^+ \otimes \mathbb{C})_v$, we have that:

- (i) $\gamma_{g,v}$ is an automorphism of graded $\mathbb{C}$ -algebras, and
- (ii) the equation (29) holds.

We first show (ii). By [34, Corollary 8.4], for every integral point $g \in \text{Spin}(S_A^+)$, the homomorphism

$$g \otimes \gamma_{g,v} : H^*(A \times M_{v,A}, \mathbb{C}) \rightarrow H^*(A \times M_{v,A}, \mathbb{C})$$

sends a universal class to a universal class in the sense of [34, Definition 3.4]. It follows that the equation (29) holds for any integral g . Since this equation expresses a Zariski-closed condition with respect to the parameter g , we obtain (29) by the density of integral points in $\text{Spin}(S_A^+ \otimes \mathbb{C})_v$.

As to (i), we deduce, again, from Zariski-density that $\gamma_{g,v}$ is a graded $\mathbb{C}$ -algebra homomorphism for any $g \in \text{Spin}(S_A^+ \otimes \mathbb{C})_v$. The endomorphism $\gamma_{g,v}$ being an automorphism is not a Zariski closed condition. However, to prove $\gamma_{g,v} \in \text{Aut}(H^*(M_{v,A}, \mathbb{C}))$, it suffices to show that

$$(30) \quad \gamma_{g,v} \circ \gamma_{g^{-1},v} = \text{id}, \quad g \in \text{Spin}(S_A^+ \otimes \mathbb{C})_v.$$

For integral g , the equation (30) follows from (27),

$$\gamma_{g,v} \circ \gamma_{g^{-1},v} = \gamma_{g \cdot g^{-1},v} = \gamma_{\text{id},v} = \text{id}.$$

Since (30) is a Zariski closed condition, we conclude that it holds for any g , and (i) follows. $\square$

2.5. Fourier–Mukai transforms. This section and Sections 2.6 to 2.7 are devoted to proving Theorem 2.1. We first treat the special case where, if we let E and E' be two elliptic curves, then we set

$$(31) \quad A = E \times E', \quad \beta = \sigma + nf,$$

where $\sigma = [\text{pt} \times E']$ and $f = [E \times \text{pt}]$ are the classes of a section and a fiber with respect to the elliptic fibration $p : A \rightarrow E'$. Note that $\beta^2 = 2n$.

By [44, Theorem 3.15], we have an isomorphism of moduli spaces

$$(32) \quad \mathcal{M}_{\beta,A} \cong M_{v_n,A} (= A^{[n]} \times \hat{A}), \quad v_n = (1, 0, -n)$$

induced by a relative Fourier–Mukai transform with respect to the elliptic fibration $p : A \rightarrow E'$. Our strategy is to compare a universal rank 1 torsion free sheaf on $A \times M_{v_n,A}$ with a universal 1-dimensional sheaf on $A \times \mathcal{M}_{\beta,A}$ under the isomorphism (32). Then we reduce the special case (31) of Theorem 2.1 to Theorem 1.2.

The proof of [44, Theorem 3.15] shows that the support of a sheaf $\mathcal{F} \in \mathcal{M}_{\beta,A}$ is the union of a section and several fibers (with multiplicities). Hence the Hilbert–Chow morphism (22) is exactly the morphism

$$(33) \quad p_n \times q : A^{[n]} \times \hat{A} \rightarrow E'^{(n)} \times E.$$

Here we use the identification between $\hat{A}$ and A for the abelian surface $A = E \times E'$, and q is the projection to the first factor. Under the identification

$$A \times A = (E \times E) \times (E' \times E'),$$

the kernel for the relative Fourier–Mukai transform associated with the elliptic fibration $p : A \rightarrow E'$ is given by

$$(34) \quad \mathcal{P}'_E = \mathcal{P}_E \boxtimes \mathcal{O}_{\Delta_{E'}}$$

with

$$\mathcal{P}_E = \mathcal{O}_{E \times E}(\Delta_E - o_E \times E - E \times o_E)$$

the normalized Poincaré line bundle. Recall the universal family $\mathcal{E}_n$ on $A \times M_{v_n,A}$ defined in (24). A universal family $\mathcal{F}_\beta$ of 1-dimensional Gieseker-stable sheaves on $A \times \mathcal{M}_{\beta,A}$ is induced by $\mathcal{E}_n$ and $\mathcal{P}'_E$ under the isomorphism (32) as follows.

For χ given in (20), we define

$$N = \mathcal{O}_A(\chi \cdot [E \times o_{E'}]) = p^* \mathcal{O}_{E'}(\chi \cdot o_{E'}) \in \text{Pic}(A).$$

By [44, Theorem 3.15] together with its proof, every closed point in $\mathcal{M}_{\beta,A}$ is given by

$$(\phi_{\mathcal{P}'_E}(I_Z \otimes L) \otimes N) [1] \in \text{Coh}(A),$$

where $\phi_{\mathcal{P}'_E}$ is the Fourier–Mukai transform with kernel (34), and $I_Z \otimes L \in M_{v_n,A} = A^{[n]} \times \hat{A}$ is a uniquely determined rank one torsion free sheaf on A . In particular, the sheaf

$$(35) \quad \mathcal{F}_\beta = R\pi_{13*}(\pi_{12}^*(\pi_1^* N \otimes \mathcal{P}'_E) \otimes^{\mathbb{L}} \pi_{23}^* \mathcal{E}_n) [1] \in \text{Coh}(A \times \mathcal{M}_{\beta,A})$$

is a universal family on $A \times \mathcal{M}_{\beta,A}$, where π_i, π_{ij} are projections from $A \times A \times M_{v_n,A}$ to the corresponding factors, and we identify $A \times M_{v_n,A}$ with $A \times \mathcal{M}_{\beta,A}$ via (32).

For notational convenience, we define the shifted normalized universal family

$$(36) \quad \mathcal{F}_\beta^n = R\pi_{13*}(\pi_{12}^* \mathcal{P}'_E \otimes^{\mathbb{L}} \pi_{23}^* \mathcal{E}_n) = \mathcal{F}_\beta \otimes \pi_A^* N^{-1}[-1]$$

on $A \times \mathcal{M}_{\beta,A}$, whose restriction over every closed point $\mathcal{F} \in \mathcal{M}_{\beta,A}$ recovers the shifted 1-dimensional coherent sheaf $(\mathcal{F} \otimes N^{-1})[-1]$ on A .

We use the universal family (35) and the class α (39) to serve as the ones in Theorem 2.1 for the special case (31). Next, we construct a decomposition of the perverse filtration

$$(37) \quad H^*(\mathcal{M}_{\beta,A}, \mathbb{Q}) = \bigoplus_{i,d} \tilde{G}_i H^d(\mathcal{M}_{\beta,A}, \mathbb{Q})$$

which serves as the splitting in Theorem 2.1.

A canonical splitting of the perverse filtration associated with

$$q : \hat{A} = A \rightarrow E$$

is given by

$$G'_k H^d(\hat{A}, \mathbb{Q}) = \bigoplus_{i+k=d} H^{d-k}(E, \mathbb{Q}) \otimes H^k(E', \mathbb{Q}).$$

We define (37) as the decomposition

$$(38) \quad \tilde{G}_k H^*(\mathcal{M}_{\beta,A}, \mathbb{Q}) = \bigoplus_{i+j=k} \tilde{G}_i H^*(A^{[n]}, \mathbb{Q}) \otimes G'_j H^*(\hat{A}, \mathbb{Q}),$$

where $\tilde{G}_* H^*(A^{[n]}, \mathbb{Q})$ was defined by (17). Since $\tilde{G}_* H^*(A^{[n]}, \mathbb{Q})$ splits the perverse filtration (18) associated with

$$p_n : A^{[n]} \rightarrow E'^{(n)},$$

the filtration (38) splits the perverse filtration associated with the Hilbert–Chow morphism (33).

Theorem 2.3 verifies Theorem 2.1 for the special case (31), where we can choose the twisting class to be

$$(39) \quad \alpha = -\pi_A^* c_1(N) \in H^2(A \times \mathcal{M}_{\beta,A}, \mathbb{Q}).$$

Theorem 2.3. *The decomposition (38) is multiplicative, and, with α as in (39), we have*

$$(40) \quad \int_{\gamma} \text{ch}_k^{\alpha}(\mathcal{F}_{\beta}) \in \tilde{G}_{k-1} H^*(\mathcal{M}_{\beta,A}, \mathbb{Q}), \quad \forall k \geq 0, \forall \gamma \in H^*(A, \mathbb{Q}).$$

Proof. The multiplicativity of (38) follows directly from Theorem 1.2(b) and the Künneth formula. We need to prove (40), which, by the very definition of α and $\mathcal{F}_{\beta}$ and by the fact that a cohomological shift only changes the signs of Chern classes, is equivalent to

$$(41) \quad \int_{\gamma} \text{ch}_k(\mathcal{F}_{\beta}^n) \in \tilde{G}_{k-1} H^*(\mathcal{M}_{\beta,A}, \mathbb{Q}), \quad \forall k \geq 0, \forall \gamma \in H^*(A, \mathbb{Q}).$$

We consider the product $A \times \mathcal{M}_{\beta,A}$ and the multiplicative splitting of its cohomology

$$(42) \quad \tilde{G}_k H^*(A \times \mathcal{M}_{\beta,A}, \mathbb{Q}) := H^*(A, \mathbb{Q}) \otimes \tilde{G}_k H^*(\mathcal{M}_{\beta,A}, \mathbb{Q}).$$

This decomposition splits the perverse filtration associated with the morphism

$$\text{id} \times \pi_{\beta} : A \times \mathcal{M}_{\beta,A} \rightarrow A \times B.$$

The statement (41) is equivalent to the following:

$$(43) \quad \text{ch}(\mathcal{F}_{\beta}^n) \in \bigoplus_{k \geq 1} \tilde{G}_{k-1} H^{2k}(A \times \mathcal{M}_{\beta,A}, \mathbb{Q}).$$

For a nonsingular projective variety X with a decomposition $G_*H^*(X, \mathbb{Q})$ on its cohomology, we say that a class $\alpha \in H^*(X, \mathbb{Q})$ is *balanced* with respect to this decomposition if

$$\alpha \in \bigoplus_k G_k H^{2k}(X, \mathbb{Q}).$$

We prove (43) by the following 3 steps.

Step 1. We consider the new decomposition

$$\begin{aligned} \overline{G}_k H^*(A \times \mathcal{M}_{\beta, A}, \mathbb{Q}) &:= \bigoplus_{i+j=k} H^i(E, \mathbb{Q}) \otimes H^*(E', \mathbb{Q}) \otimes \tilde{G}_j H^*(\mathcal{M}_{\beta, A}, \mathbb{Q}) \\ (44) \quad &= \bigoplus_{i+j=k} \tilde{G}_i H^*(A \times A^{[n]}, \mathbb{Q}) \otimes G'_j H^*(\hat{A}, \mathbb{Q}) \end{aligned}$$

of the cohomology $H^*(A \times \mathcal{M}_{\beta, A}, \mathbb{Q})$. It is a multiplicative decomposition splitting the perverse filtration associated with the morphism

$$p \times \pi_\beta : A \times \mathcal{M}_{\beta, A} \rightarrow E' \times B.$$

We first show that the class

$$\text{ch}(\mathcal{E}_n) \in H^*(A \times M_{A, v_n}, \mathbb{Q}) = H^*(A \times \mathcal{M}_{\beta, A}, \mathbb{Q})$$

is balanced with respect to the splitting (44). Here the last identification is given by (32).

Recall the expression (24) of the universal family $\mathcal{E}_n$. By Theorem 1.2(a), the class

$$\text{ch}(\mathcal{I}_n) \in H^*(A \times A^{[n]}, \mathbb{Q})$$

is balanced with respect to the splitting (19). Hence via pulling back along the projection

$$\pi_{12} : A \times \mathcal{M}_{\beta, A} = A \times A^{[n]} \times \hat{A} \rightarrow A \times A^{[n]},$$

we obtain that the class

$$\begin{aligned} \pi_{12}^* \text{ch}(\mathcal{I}_n) &= \text{ch}(\mathcal{I}_n) \boxtimes 1_{\hat{A}} \in H^*(A \times A^{[n]}, \mathbb{Q}) \otimes H^0(\hat{A}, \mathbb{Q}) \\ &\subset H^*(A \times \mathcal{M}_{\beta, A}, \mathbb{Q}) \end{aligned}$$

is also balanced with respect to (44). A direct calculation shows that the class $\pi_{13}^* c_1(\mathcal{P})$ is balanced. Hence we conclude from the multiplicativity of (44) that the class

$$\text{ch}(\mathcal{E}_n) = \pi_{12}^* \text{ch}(\mathcal{I}_n) \otimes \pi_{13}^* \text{ch}(\mathcal{P}) = \pi_{12}^* \text{ch}(\mathcal{I}_n) \otimes \pi_{13}^* \exp(c_1(\mathcal{P}))$$

is balanced.

Step 2. We further consider the multiplicative decomposition

$$(45) \quad \tilde{G}_k H^*(A \times A \times \mathcal{M}_{\beta, A}, \mathbb{Q}) := H^*(A, \mathbb{Q}) \otimes \overline{G}_k H^*(A \times \mathcal{M}_{\beta, A}, \mathbb{Q})$$

of $H^*(A \times A \times \mathcal{M}_{\beta, A}, \mathbb{Q})$, which splits the perverse filtration associated with the morphism

$$\text{id} \times p \times \pi_\beta : A \times A \times \mathcal{M}_{\beta, A} \rightarrow A \times E' \times B.$$

We show that

$$(46) \quad \text{ch}(\pi_{12}^* \mathcal{P}'_E \otimes^{\mathbb{L}} \pi_{23}^* \mathcal{E}_n) \in \bigoplus_{k \geq 1} \tilde{G}_{k-1} H^{2k}(A \times A \times \mathcal{M}_{\beta, A}, \mathbb{Q}).$$

Here π_{ij} are the projections from $A \times A \times \mathcal{M}_{\beta, A}$ to the corresponding factors; compare the notation of (36).

In fact, by Step 1, we obtain that the class

$$\pi_{23}^* \text{ch}(\mathcal{E}_n) = 1_A \boxtimes \text{ch}(\mathcal{E}_n) \in H^0(A, \mathbb{Q}) \otimes H^*(A \times \mathcal{M}_{\beta, A}, \mathbb{Q})$$

is balanced with respect to (45). The relative Poincaré sheaf (34) can be expressed as

$$(47) \quad \pi_{13}^* \text{ch}(\mathcal{P}'_E) = \pi_{13}^* \text{ch}(\mathcal{P}_E) \cup \pi_{13}^* \text{ch}(\mathcal{O}_{\Delta_{E'}}).$$

The class $\pi_{12}^* \text{ch}(\mathcal{P}_E)$ is balanced via a direct calculation, and

$$\pi_{12}^* \text{ch}(\mathcal{O}_{\Delta_{E'}}) = \pi_{12}^* c_1(\mathcal{O}_{\Delta_{E'}}) \in \tilde{G}_0 H^2(A \times A \times \mathcal{M}_{\beta, A}, \mathbb{Q}).$$

Hence by (47) and the multiplicativity of (45), we get

$$\pi_{12}^* \text{ch}(\mathcal{P}'_E) \in \bigoplus_{k \geq 1} \tilde{G}_{k-1} H^{2k}(A \times A \times \mathcal{M}_{\beta, A}).$$

This yields (46) since

$$\text{ch}(\pi_{12}^* \mathcal{P}'_E \otimes^{\mathbb{L}} \pi_{23}^* \mathcal{E}_n) = \pi_{12}^* \text{ch}(\mathcal{P}'_E) \cup \pi_{23}^* \text{ch}(\mathcal{E}_n).$$

Step 3. We finish the proof of (43).

Recall the expression (36) of $\mathcal{F}_\beta^n$. Applying the Grothendieck–Riemann–Roch formula to the projection

$$\pi_{13} : A \times A \times \mathcal{M}_{\beta, A} \rightarrow A \times \mathcal{M}_{\beta, A},$$

we obtain that

$$\text{ch}(\mathcal{F}_\beta^n) = \pi_{13*} \text{ch}(\pi_{12}^* \mathcal{P}'_E \otimes^{\mathbb{L}} \pi_{23}^* \mathcal{E}_n).$$

Equivalently, the class $\text{ch}(\mathcal{F}_\beta^n)$ corresponds to the Künneth factor of the class

$$\text{ch}(\pi_{12}^* \mathcal{P}'_E \otimes^{\mathbb{L}} \pi_{23}^* \mathcal{E}_n)$$

in the subspace

$$H^4(A, \mathbb{Q}) \otimes H^*(A \times \mathcal{M}_{\beta, A}, \mathbb{Q}) \subset H^*(A \times A \times \mathcal{M}_{\beta, A}, \mathbb{Q}).$$

Here $H^4(A, \mathbb{Q})$ is spanned by the point class in the second factor of the product $A \times A \times \mathcal{M}_{\beta, A}$. Hence (43) follows from (46). $\square$

2.6. Perverse filtrations and $H^2(\mathcal{M}_{\beta, A}, \mathbb{Q})$. In this section, we assume that A is any abelian surface and β is an ample curve class satisfying $\beta^2 = v_n^2 = 2n$. Recall from Section 2.1 that $\mathcal{M}_{\beta, A}$ is the moduli space of Gieseker-stable sheaves with respect to the primitive Mukai vector

$$(48) \quad v_\beta = (0, \beta, \chi) \in S_A^+$$

with dimension

$$\dim(\mathcal{M}_{\beta, A}) = v_\beta^2 + 2 = 2n + 2.$$

Our goal is to prove Proposition 2.5, which is a variant of Proposition 1.1 for $\mathcal{M}_{\beta, A}$. More precisely, we introduce a *canonical* class

$$(49) \quad L_\beta \in H^2(\mathcal{M}_{\beta, A}, \mathbb{Q})$$

to characterize the perverse filtration $P_* H^*(\mathcal{M}_{\beta, A}, \mathbb{Q})$ associated with the morphism

$$\pi_\beta : \mathcal{M}_{\beta, A} \rightarrow B.$$

Let $\mathcal{F}_0 \in \mathcal{M}_{\beta,A}$ be a general point on the moduli space. We consider the morphisms defined by considering determinants of coherent sheaves

$$\det : \mathcal{M}_{\beta,A} \rightarrow \hat{A}, \quad \mathcal{F} \mapsto \det(\mathcal{F}) \otimes \det(\mathcal{F}_0)^\vee$$

and

$$\hat{\det} : \mathcal{M}_{\beta,A} \rightarrow A, \quad \mathcal{F} \mapsto \det(\phi_{\mathcal{P}}(\mathcal{F})) \otimes \det(\phi_{\mathcal{P}}(\mathcal{F}))^\vee.$$

Here $\phi_{\mathcal{P}} : D^b\text{Coh}(A) \rightarrow D^b\text{Coh}(\hat{A})$ is the Fourier–Mukai transform induced by the normalized Poincaré line bundle $\mathcal{P}$ on $A \times \hat{A}$. By [44, (4.1) and Theorem 4.1], the Albanese morphism for $\mathcal{M}_{\beta,A}$ with respect to the reference point $\mathcal{F}_0$ is

$$\text{alb} = \hat{\det} \times \det : \mathcal{M}_{\beta,A} \rightarrow A \times \hat{A}.$$

The closed fiber over the origin $0 \in A \times \hat{A}$, denoted by

$$K_{\beta,A} = \text{alb}^{-1}(0) \subset \mathcal{M}_{\beta,A},$$

is a holomorphic symplectic variety of Kummer type; see [44, Theorem 0.2]. It parametrizes 1-dimensional sheaves $\mathcal{F} \in \mathcal{M}_{\beta,A}$ satisfying

$$\det(\mathcal{F}) = \det(\mathcal{F}_0) \in \hat{A}, \quad \det(\phi_{\mathcal{P}}(\mathcal{F})) = \det(\phi_{\mathcal{P}}(\mathcal{F}_0)) \in \hat{A} = A.$$

The restriction of the Hilbert–Chow morphism (22) to the subvariety $K_{\beta,A}$ is a Lagrangian fibration

$$\pi'_\beta : K_{\beta,A} \rightarrow |\mathcal{O}_A(\beta)| = \mathbb{P}^{n-1}.$$

Here $|\mathcal{O}_A(\beta)|$ denotes the linear system associated with the divisor

$$\text{supp}(\mathcal{F}_0) \subset A.$$

We consider the following commutative diagram,

$$(50) \quad \begin{array}{ccc} K_{\beta,A} \times A \times \hat{A} & \xrightarrow{h} & \mathcal{M}_{\beta,A} \\ \downarrow \pi'_\beta \times \text{pr}_A & & \downarrow \pi_\beta \\ \mathbb{P}^{n-1} \times A & \xrightarrow{h'} & B. \end{array}$$

Here the horizontal arrows are defined by

$$h(\mathcal{F}, a, L) = t_{a*}\mathcal{F} \otimes L, \quad h'(C, a) = t_{a*}C$$

with $t_a : A \xrightarrow{\cong} A$ the translation by a . Since both h and h' are finite and surjective, the map

$$h^* : H^k(\mathcal{M}_{\beta,A}, \mathbb{Q}) \rightarrow H^k(K_{\beta,A} \times A \times \hat{A}, \mathbb{Q})$$

is injective by the projection formula. Moreover, it is an isomorphism when $k = 2$ since

$$\dim H^2(\mathcal{M}_{\beta,A}, \mathbb{Q}) = \dim H^2(A^{[n]} \times \hat{A}, \mathbb{Q}) = \dim H^2(K_{\beta,A} \times A \times \hat{A}, \mathbb{Q})$$

by Göttsche's formula [21] for the Betti numbers of the Hilbert schemes and the generalized Kummer varieties.

By [44, Theorem 0.1 and (1.6)], the correspondence induced by $\text{ch}(\mathcal{F}_\beta) \in H^*(A \times \mathcal{M}_{\beta,A})$ together with the restriction map

$$H^2(\mathcal{M}_{\beta,A}, \mathbb{Q}) \rightarrow H^2(K_{\beta,A}, \mathbb{Q})$$

gives an isometry between $v_\beta^\perp \subset S_A^+ \otimes \mathbb{Q}$ and $H^2(K_{\beta,A}, \mathbb{Q})$ (endowed with the Beauville-Bogomolov-Fujiki form)

$$\theta : v_\beta^\perp \xrightarrow{\sim} H^2(K_{\beta,A}, \mathbb{Q}).$$

Therefore, the pullback h^* induces an isometry (see also [44, Theorem 4.1.(3) eq. (4.3)])

$$(51) \quad \varphi : H^2(\mathcal{M}_{\beta,A}, \mathbb{Q}) \xrightarrow{\sim} v_\beta^\perp \oplus H^2(A \times \hat{A}, \mathbb{Q}),$$

which, by [32, Section 8], is $\text{Spin}(S_A^+ \otimes \mathbb{Q})_{v_\beta}$ -equivariant.

Remark 2.4. The equivariance means that Markman's monodromy operators γ_{g,v_β} (28) acting on $H^2(\mathcal{M}_{\beta,A}, \mathbb{Q})$ are identified, via φ , with the action of g on the right-hand side of (51) described in Section 2.3. We may say that φ is compatible with g and γ_{g,v_β} .

We consider the following classes

$$(52) \quad \begin{aligned} l_\beta &= \theta((0, 0, 1) \in v_\beta^\perp) \in H^2(K_{\beta,A}, \mathbb{Q}), \\ L_\beta &= \varphi^{-1}((0, 0, 1) + (\beta \boxtimes 1_{\hat{A}})) \in H^2(\mathcal{M}_{\beta,A}, \mathbb{Q}). \end{aligned}$$

Proposition 2.5. *We have*

$$P_k H^m(\mathcal{M}_{\beta,A}, \mathbb{Q}) = \left(\sum_{i \geq 1} \text{Ker} \left(L_\beta^{(n+1)+k+i-m} \right) \cap \text{Im}(L_\beta^{i-1}) \right) \cap H^m(\mathcal{M}_{\beta,A}, \mathbb{Q}).$$

Proof. We denote by

$$\pi_K = \pi'_\beta \times \text{pr}_A : K_{\beta,A} \times A \times \hat{A} \rightarrow \mathbb{P}^{n-1} \times A$$

the left vertical map of the diagram (50).

By [33, Example 3.1], we have

$$l_\beta = \pi'_\beta{}^* \mathcal{O}_{\mathbb{P}^{n-1}}(1) \in H^2(K_{\beta,A}, \mathbb{Q}).$$

Hence

$$h^* L_\beta = l_\beta \boxtimes 1_{A \times \hat{A}} + 1_{K_{\beta,A}} \boxtimes (\beta \otimes 1_{\hat{A}}) \in H^2(K_{\beta,A} \times A \times \hat{A}, \mathbb{Q})$$

is the pullback of an ample class on the base $\mathbb{P}^{n-1} \times A$ via the morphism π_K . Since h and h' are finite, Proposition 2.5 is deduced immediately from the diagram (50) and Lemma 2.6. $\square$

Lemma 2.6. *We consider a commutative diagram*

$$\begin{array}{ccc} X & \xrightarrow{h} & X' \\ \downarrow f & & \downarrow f' \\ Y & \xrightarrow{h'} & Y' \end{array},$$

where all the varieties are nonsingular and irreducible, the horizontal morphisms are finite and surjective, and the vertical morphisms are proper. We assume

$$\dim(X) = \dim(X') = 2\dim(Y) = 2\dim(Y') = 2R$$

for some positive integer R , and that there is a class $\alpha \in H^2(X', \mathbb{Q})$ satisfying

$$h^* \alpha = f^* L$$

with $L \in H^2(Y, \mathbb{Q})$ an ample class on Y . Then the perverse filtration $P_* H^*(X', \mathbb{Q})$ associated with f' can be expressed as

$$(53) \quad P_k H^m(X', \mathbb{Q}) = \left(\sum_{i \geq 1} \text{Ker}(\alpha^{R+k+i-m}) \cap \text{Im}(\alpha^{i-1}) \right) \cap H^m(X', \mathbb{Q}).$$

Proof. The composition of the following maps is multiplication by the degree of h

$$\mathbb{Q}_{X'} \rightarrow h_* h^* \mathbb{Q}_{X'} = h_* \mathbb{Q}_X \rightarrow \mathbb{Q}_{X'},$$

where the first map is the adjunction and the last map is the trace map. This composition is nonzero since h has non-zero degree, so it yields the splitting

$$(54) \quad h_* \mathbb{Q}_X = \mathbb{Q}_{X'} \oplus \mathcal{G}, \quad \mathcal{G} \in D_c^b(X'),$$

which further induces the embedding

$$(55) \quad h^* : H^*(X', \mathbb{Q}) \hookrightarrow H^*(X, \mathbb{Q})$$

as a direct summand. After pushing forward (54) along $f' : X' \rightarrow Y'$, we obtain

$${}^p \mathcal{H}^k(\text{Rf}'_* h_* \mathbb{Q}_X) = {}^p \mathcal{H}^k(\text{Rf}'_* \mathbb{Q}_{X'}) \oplus {}^p \mathcal{H}^k(\text{Rf}'_* \mathcal{G}).$$

On the other hand, by the t -exactness of the finite morphism h' , we have

$${}^p \mathcal{H}^k(\text{Rf}'_* h_* \mathbb{Q}_X) = {}^p \mathcal{H}^k(h'_* \text{Rf}_* \mathbb{Q}_X) = h'_* {}^p \mathcal{H}^k(\text{Rf}_* \mathbb{Q}_X).$$

Hence the pullback (55) induced by (54) is filtered strict with respect to the perverse filtrations associated with f and f' , and we have

$$(56) \quad P_k H^d(X, \mathbb{Q}) \cap \text{Im}(h^*) \subset P_k H^d(X', \mathbb{Q}), \quad \forall k, d.$$

We deduce (53) from (56) and Proposition 1.1 (applied to $f : X \rightarrow Y$). $\square$

2.7. Proof of Theorem 2.1. In this section, we complete the proof of Theorem 2.1 by combining the tools developed in the previous sections.

Let A be an abelian surface and β be an ample curve class. A *perverse package* associated with the pair (A, β) is defined to be a triple

$$([\mathcal{F}_\beta], \alpha_\beta, L_\beta),$$

where $[\mathcal{F}_\beta]$ is a universal class on $A \times \mathcal{M}_{\beta, A}$, $L_\beta \in H^2(\mathcal{M}_{\beta, A}, \mathbb{C})$ is the class introduced in Section 2.6, and $\alpha \in H^2(A \times \mathcal{M}_{\beta, A}, \mathbb{C})$ is of the type (21).

An isomorphism between the perverse packages associated with (A, β) and (A', β') is the following:

- (i) An isomorphism of graded $\mathbb{C}$ -vector spaces

$$g : H^*(A, \mathbb{C}) \xrightarrow{\sim} H^*(A', \mathbb{C}).$$

- (ii) An isomorphism of graded $\mathbb{C}$ -algebras

$$\phi : H^*(\mathcal{M}_{\beta, A}, \mathbb{C}) \xrightarrow{\sim} H^*(\mathcal{M}_{\beta', A'}, \mathbb{C}).$$

- (iii) The isomorphisms g and ϕ satisfy that

$$(g \otimes \phi) \text{ch}^{\alpha_\beta}(\mathcal{F}_\beta) = \text{ch}^{\alpha_{\beta'}}(\mathcal{F}_{\beta'}),$$

$$\phi(L_\beta) = L_{\beta'}.$$

Isomorphisms of perverse packages form an equivalence relation. We call two pairs (A, β) and (A', β') *perversely isomorphic* if there is a perverse package associated with (A, β) isomorphic to a perverse package associated with (A', β') .

Proposition 2.7. *Assume that (A, β) is perversely isomorphic to (A', β') . If Theorem 2.1 holds for (A, β) , then it also holds for (A', β') .*

Proof. By Proposition 2.5, the condition

$$\phi(L_\beta) = L_{\beta'}$$

implies that the perverse filtrations on $H^*(\mathcal{M}_{\beta,A}, \mathbb{C})$ and $H^*(\mathcal{M}_{\beta',A'}, \mathbb{C})$ are identified via ϕ . Let

$$H^*(\mathcal{M}_{\beta,A}, \mathbb{C}) = \bigoplus_{k,d} \tilde{G}_k H^d(\mathcal{M}_{\beta,A}, \mathbb{C})$$

be the splitting of the perverse filtration for $\mathcal{M}_{\beta,A}$ satisfying

$$\int_\gamma \text{ch}_k^{\alpha_\beta}(\mathcal{F}_\beta) \in \tilde{G}_{k-1} H^*(\mathcal{M}_{\beta,A}, \mathbb{C}).$$

Then the decomposition

$$\tilde{G}_k H^d(\mathcal{M}_{\beta',A'}, \mathbb{C}) = \phi(\tilde{G}_k H^d(\mathcal{M}_{\beta,A}, \mathbb{C}))$$

also splits the perverse filtration for $\mathcal{M}_{\beta',A'}$, and the corresponding tautological classes lie in its correct pieces by condition (iii) in the definition of isomorphism of perverse package. $\square$

If (A, β) is perversely isomorphic to (A', β') , then condition (ii) implies that

$$(57) \quad \beta^2 = \dim(\mathcal{M}_{\beta,A}) - 2 = \dim(\mathcal{M}_{\beta',A'}) - 2 = \beta'^2.$$

Theorem 2.8, which we deduce from Theorem 2.2, establishes the converse to (57). In other words, perverse equivalence classes associated with pairs (A, β) are larger than the equivalence classes given by the deformation types of the fibrations $\pi_\beta : \mathcal{M}_{\beta,A} \rightarrow B$; see Remark 2.9.

Theorem 2.8. *Any two pairs (A, β) and (A', β') satisfying*

$$\beta^2 = \beta'^2$$

are perversely isomorphic.

Proof. As we explain at the end of the proof, it will be sufficient to find an isomorphism of graded $\mathbb{C}$ -vector spaces

$$(58) \quad g : H^*(A, \mathbb{C}) \rightarrow H^*(A', \mathbb{C})$$

satisfying the following two properties:

- (a) g preserves the intersection pairing;
- (b) g satisfies the following identities

$$(59) \quad \begin{aligned} g(0, 0, 1) &= (0, 0, 1), \\ g(1, 0, 0) &= (1, 0, 0), \\ g(0, \beta, 0) &= (0, \beta', 0), \end{aligned}$$

and a graded $\mathbb{C}$ -algebra isomorphism

$$(60) \quad \phi : H^*(\mathcal{M}_{\beta,A}, \mathbb{C}) \rightarrow H^*(\mathcal{M}_{\beta',A'}, \mathbb{C})$$

satisfying condition (iii) in the definition of isomorphism of perverse packages.

We construct (g, ϕ) in two steps.

We note that g_1, g_2 , and $g = g_2 \circ g_1$ constructed below automatically satisfy condition (a) by construction.

Firstly, since the Mukai vectors (48) v_β and $v_{\beta'}$ are primitive with the same Mukai length,

$$v_\beta^2 = v_{\beta'}^2 \in \mathbb{Z},$$

we can apply [34, Theorem 8.3] (which follows from Yoshioka [44]) to find isometries

$$g_1 : H^*(A, \mathbb{Z}) \rightarrow H^*(A', \mathbb{Z}), \quad \phi_1 : H^*(\mathcal{M}_{\beta, A}, \mathbb{Q}) \rightarrow H^*(\mathcal{M}_{\beta', A'}, \mathbb{Q})$$

such that $g_1(v_\beta) = v_{\beta'}$ and

$$(61) \quad (g_1 \otimes \phi_1) \text{ch}(\mathcal{F}_\beta) = \text{ch}(\mathcal{F}_{\beta'})$$

with $[\mathcal{F}_{\beta'}]$ a universal class on $A \times \mathcal{M}_{\beta, A}$. Moreover, the morphism ϕ_1 satisfies the evident variant of Remark 2.4, *i.e.*, the morphism

$$\varphi_{\beta'}^{-1} \phi_1 \varphi_\beta : v_\beta^\perp \oplus H^2(A \times \hat{A}, \mathbb{Q}) \rightarrow v_{\beta'}^\perp \oplus H^2(A' \times \hat{A}', \mathbb{Q})$$

is induced by the isometry $g_1 : H^*(A, \mathbb{Z}) \rightarrow H^*(A', \mathbb{Z})$; see [32, Proposition 8.5]. We may say that the identifications φ (51) for β and β' are compatible with g_1 and ϕ_1 .

If β and β' have different divisibility, then g_1 cannot satisfy the condition (b) above, so that we proceed as follows. We choose $g_2 \in \text{SO}(S_{A'}^+ \otimes \mathbb{Q})_{v_{\beta'}}$ such that the restriction

$$g_2 \circ g_1|_{S_A^+} : S_A^+ \otimes \mathbb{Q} \rightarrow S_{A'}^+ \otimes \mathbb{Q}$$

is grading-preserving and satisfies condition (b). We can lift g_2 to an element of $\text{Spin}(S_{A'}^+ \otimes \mathbb{C})_{v_{\beta'}}$, also denoted by g_2 , and take the associated isomorphisms

$$g_2 : H^*(A', \mathbb{C}) \rightarrow H^*(A', \mathbb{C}) \quad (\text{cf. Section 2.3}),$$

$$\phi_2 = \gamma_{g_2, v_{\beta'}} : H^*(\mathcal{M}_{\beta', A'}, \mathbb{C}) \rightarrow H^*(\mathcal{M}_{\beta', A'}, \mathbb{C}) \quad (\text{cf. Section 2.4}).$$

By Theorem 2.2, we have

$$(62) \quad (g_2 \otimes \phi_2) \text{ch}(\mathcal{F}_{\beta'}) = \text{ch}^{\pi^* \mathcal{M}^a}(\mathcal{F}_{\beta'}), \quad \exists a \in H^2(\mathcal{M}_{\beta', A'}, \mathbb{C}).$$

We set $g = g_2 \circ g_1$ and $\phi = \phi_2 \circ \phi_1$. Then (61) and (62) imply that

$$(g \otimes \phi) \text{ch}(\mathcal{F}_\beta) = \text{ch}^{\pi^* \mathcal{M}^a}(\mathcal{F}_{\beta'}).$$

In particular, for any class $\alpha_\beta \in H^2(A \times \mathcal{M}_{\beta, A}, \mathbb{C})$ of type (21), we have

$$(g \otimes \phi) \text{ch}^{\alpha_\beta}(\mathcal{F}_\beta) = \text{ch}^{\alpha_{\beta'}}(\mathcal{F}_{\beta'}), \quad \text{with } \alpha_{\beta'} = \pi^* \mathcal{M}^a + (g \otimes \phi) \alpha_\beta,$$

where the class $\alpha_{\beta'}$ is also of type (21). The first condition appearing in (iii) is thus met.

It remains to verify that $\phi(L_\beta) = L_{\beta'}$. In view of the compatibility of the φ 's for β and β' with g_1, ϕ_1 , and of φ for β' with g_2, ϕ_2 (cf. Remark 2.4), and in view of the construction of g_1 and g_2 , the isomorphism ϕ sends $(0, 0, 1) \in v_\beta^\perp$ to $(0, 0, 1) \in v_{\beta'}^\perp$, and sends $\beta \boxtimes 1_{\hat{A}} \in H^2(A \times \hat{A}, \mathbb{C})$ to $\beta' \boxtimes 1_{\hat{A}'} \in H^2(A' \times \hat{A}', \mathbb{C})$. In particular, we have $\phi(L_\beta) = L_{\beta'}$, and the desired condition (iii) is met in its entirety. This completes the proof of Theorem 2.8. $\square$

Theorem 2.1 follows from Theorem 2.3, Proposition 2.7, and Theorem 2.8 (with $n := \beta^2/2$, v_n and β). In Section 3.5, we prove a strengthened version of Theorem 2.1 to the effect that the decomposition $\tilde{G}_* H^*(\mathcal{M}_{A, \beta}, \mathbb{C})$ is the one induced by a variant of a construction of Deligne's.

Remark 2.9. It was shown in [13] that perverse filtrations behave well under deformations. However, for the pairs (A, β) and (A', β') such that β and β' have different divisibilities in $H^2(A, \mathbb{Z})$ and $H^2(A', \mathbb{Z})$, the fibrations π_β and $\pi_{\beta'}$ are not deformation equivalent. Therefore, in order to reduce the proof of Theorem 2.1 for any pair (A, β) to that for a special pair

$$A = E \times E', \quad \beta = \sigma + nf,$$

it is essential to consider an equivalence relation weaker than deformation equivalences of fibrations π_β . This is our motivation for introducing equivalences of perverse packages.

3. SPLITTINGS OF PERVERSE FILTRATIONS

3.1. Overview. In this section, we study splittings of the perverse filtration associated with a proper surjective morphism $\pi : X \rightarrow Y$ with X and Y nonsingular. As an application, we strengthen Theorem 2.1 by requiring that the decomposition given by said theorem

$$(63) \quad H^*(\mathcal{M}_{\beta,A}, \mathbb{C}) = \bigoplus_{i,d} \tilde{G}_i H^d(\mathcal{M}_{\beta,A}, \mathbb{C})$$

is induced by a “Lefschetz class” via the mechanisms introduced in [9, 17, 18]; see Theorem 3.8. As discussed in Remark 4.5, this is crucial in the study of the specialization morphism (82) in Section 4.

Throughout, we work with $\mathbb{Q}$ -coefficients except for Section 3.5, where we need to switch to $\mathbb{C}$ -coefficients in order to apply results in Section 2.7. However, we note that all discussions in Sections 3.2–3.4 remain valid if we replace $\mathbb{Q}$ by $\mathbb{C}$.

3.2. Lefschetz classes. We consider the perverse filtration associated with $\pi : X \rightarrow Y$,

$$(64) \quad P_0 H^*(X, \mathbb{Q}) \subset P_1 H^*(X, \mathbb{Q}) \subset \cdots \subset P_{2R} H^*(X, \mathbb{Q}) = H^*(X, \mathbb{Q}),$$

where R is the defect of semismallness of π (13). The action of a class $\eta \in H^2(X, \mathbb{Q})$ on the cohomology $H^*(X, \mathbb{Q})$ via cup product satisfies

$$(65) \quad \eta : P_k H^i(X, \mathbb{Q}) \rightarrow P_{k+2} H^{i+2}(X, \mathbb{Q}),$$

which further induces an action on

$$\mathbb{H} = \bigoplus_{*, \bullet \geq 0} \mathrm{Gr}_*^P H^\bullet(X, \mathbb{Q}) = \bigoplus_{*, \bullet \geq 0} (P_* H^\bullet(X, \mathbb{Q}) / P_{*-1} H^\bullet(X, \mathbb{Q})).$$

We say that $\eta \in H^2(X, \mathbb{Q})$ is a π -Lefschetz class if its induced action on $\mathbb{H}$ satisfies the hard Lefschetz-type condition in the sense of [9, Assumption 2.3.1], i.e., the actions

$$(66) \quad \eta^k : \mathrm{Gr}_{R-k}^P H^*(X, \mathbb{Q}) \xrightarrow{\sim} \mathrm{Gr}_{R+k}^P H^*(X, \mathbb{Q}), \quad \forall k \geq 0,$$

are isomorphisms. As a typical example, the relative Hard Lefschetz Theorem [5] with respect to $\pi : X \rightarrow Y$ implies that a relatively ample class for π is a π -Lefschetz class. Lemma 3.1 gives examples of Lefschetz classes other than relatively ample classes.

Lemma 3.1. *Let A be an abelian variety of dimension m . Any class $\eta \in H^2(A, \mathbb{Q})$ satisfying*

$$(67) \quad \eta^m \neq 0$$

is a Lefschetz class with respect to $\pi : A \rightarrow \text{pt.}$

Proof. Let $W = H^1(A, \mathbb{Q})$. We identify $H^k(A, \mathbb{Q})$ with $\wedge^k W$, and therefore $\eta \in \wedge^2 W$. The condition (67) implies that η defines a (constant) symplectic form on W^* . In particular, we can write $\eta = \sum_{i=1}^m e_i \wedge f_i$ under a basis $e_1, \dots, e_m, f_1, \dots, f_m$ of W . Hence the pair $(W \otimes \mathbb{C}, \eta)$ is the tensor product of m copies of the symplectic plane $(\mathbb{C}^2, e \wedge f)$, for which the statement is clear. $\square$

Recall the Hitchin fibration $h : \mathcal{M}_{\text{Dol}} \rightarrow \Lambda$. Proposition 3.2 is a numerical criterion for Lefschetz classes with respect to h .

Proposition 3.2. *Let F be a closed fiber of h . A class $\eta \in H^2(\mathcal{M}_{\text{Dol}}, \mathbb{Q})$ is a Lefschetz class with respect to h if and only if the following cap product with the fundamental class of the connected fiber F (counting components with multiplicities) is non-trivial:*

$$(68) \quad 0 \neq \eta^{\dim(F)} \cap [F] \in H_0(F, \mathbb{Q}) \simeq \mathbb{Q}.^5$$

Proof. We denote by $\hat{\mathcal{M}}_{\text{Dol}}$ the corresponding moduli space of stable PGL_r -Higgs bundles with

$$\hat{h} : \hat{\mathcal{M}}_{\text{Dol}} \rightarrow \hat{\Lambda}$$

the Hitchin fibration. By [25, 30], we have

$$H^2(\hat{\mathcal{M}}_{\text{Dol}}, \mathbb{Q}) = \mathbb{Q} \cdot \alpha,$$

where α is the first Chern class of an $\hat{h}$ -ample line bundle on $\hat{\mathcal{M}}_{\text{Dol}}$, and therefore is an $\hat{h}$ -Lefschetz class. By the discussion in [11, Section 2.4], we have an isomorphism

$$(69) \quad H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) = H^*(\hat{\mathcal{M}}_{\text{Dol}}, \mathbb{Q}) \otimes H^*(\text{Pic}^0(C), \mathbb{Q})$$

satisfying that

$$(70) \quad P_k H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) = \bigoplus_{i+j=k} P_i H^*(\hat{\mathcal{M}}_{\text{Dol}}, \mathbb{Q}) \otimes H^j(\text{Pic}^0(C), \mathbb{Q}).$$

Under the identification

$$H^2(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) = H^2(\hat{\mathcal{M}}_{\text{Dol}}, \mathbb{Q}) \oplus H^2(\text{Pic}^0(C), \mathbb{Q}) = \mathbb{Q} \cdot \alpha \oplus H^2(\text{Pic}^0(C)),$$

induced by (69), we can express any class $\eta \in H^2(\mathcal{M}_{\text{Dol}}, \mathbb{Q})$ as

$$(71) \quad \eta = \mu \alpha \otimes 1 + 1 \otimes \xi, \quad \mu \in \mathbb{Q}, \quad \xi \in H^2(\text{Pic}^0(C), \mathbb{Q}).$$

Lemma 3.1 combined with [9, Appendix] implies that the class η is Lefschetz if and only if

$$(72) \quad \mu \neq 0, \quad \xi^g \neq 0.$$

Therefore, it suffices to show that (72) is equivalent to the condition (68).

We first consider traceless Higgs bundles

$$\mathcal{M}_{\text{Dol}}^0 = \{(\mathcal{E}, \theta) \in \mathcal{M}_{\text{Dol}} : \text{trace}(\theta) = 0\} \subset \mathcal{M}_{\text{Dol}}.$$

⁵We note that the condition (68) does not depend on the choice of the fiber F .

Recall from [11, Proposition 2.4.3] that there is a finite morphism

$$q : \check{\mathcal{M}}_{\text{Dol}} \times \text{Pic}^0(C) \rightarrow \mathcal{M}_{\text{Dol}}^0$$

with $\check{\mathcal{M}}_{\text{Dol}}$ the corresponding Dolbeault moduli space of stable SL_r -Higgs bundles. The preimage of a closed fiber $F \subset \mathcal{M}_{\text{Dol}}^0$ of the restricted Hitchin fibration $h|_{\mathcal{M}_{\text{Dol}}^0}$ is the product

$$q^{-1}(F) = \check{F} \times \text{Pic}^0(C),$$

where $\check{F}$ is the corresponding closed fiber of the SL_n Hitchin fibration. The pullback of a class (71) along

$$\check{F} \times \text{Pic}^0(C) \hookrightarrow \check{\mathcal{M}}_{\text{Dol}} \times \text{Pic}^0(C) \xrightarrow{q} \mathcal{M}^0 \hookrightarrow \mathcal{M}_{\text{Dol}}$$

is of the type

$$\check{\eta} = \mu\check{\alpha} \otimes 1 + 1 \otimes \xi \in H^2(\check{F}, \mathbb{Q}) \oplus H^2(\text{Pic}^0(C), \mathbb{Q}) \subset H^2(\check{F} \times \text{Pic}^0(C), \mathbb{Q})$$

with $\check{\alpha}$ an ample class on $\check{F}$. Since $q : \check{F} \times \text{Pic}^0(C) \rightarrow F$ is finite and surjective, the condition (68) is equivalent to

$$0 \neq \check{\eta}^{\dim(\check{F} \times \text{Pic}^0(C))} \cap [\check{F}] = \check{\eta}^{\dim(F)} \cap [F],$$

which, in turn, is equivalent to (72). This completes the proof. $\square$

3.3. The first Deligne splitting. Given a projective morphism $\pi : X \rightarrow Y$ and a π -ample line bundle on X , Deligne [18] constructs three splittings of the direct image $R\pi_*\mathbb{Q}_X$ (resp. $R\pi_*\text{IC}_X$ if X is singular), which induce splittings of the perverse filtration on the (resp. intersection) cohomology groups $H^*(X, \mathbb{Q})$. In this paper, we need the variant [9] of this construction where one starts with a π -Lefschetz class $\eta \in H^2(X, \mathbb{Q})$, *i.e.* one that does not necessarily satisfy the relative Hard Lefschetz Theorem in the derived category $D_c^b(Y)$, but nonetheless satisfies the cohomological version (66). We need only the first of the resulting three splittings, which we name the *first Deligne splitting* [9, 17, 18].

According to [9], the first Deligne splitting of the perverse filtration (64)

$$H^*(X, \mathbb{Q}) = \bigoplus_i G_i H^*(X, \mathbb{Q})$$

associated with the π -Lefschetz class η (*i.e.* (66) holds) can be described using *only* the action of η on $H^*(X, \mathbb{Q})$. We explain this description explicitly as follows.

For $i \geq 0$, we let $\text{Gr}_i^P := P_i H^*(X, \mathbb{Q}) / P_{i-1} H^*(X, \mathbb{Q})$ denote the graded spaces, which are subquotients of cohomology, and let $G_i := G_i H^*(X, \mathbb{Q})$ denote the corresponding image of Gr_i^P via the first Deligne splitting, *i.e.* these are the summands in the last paragraph, which are subspaces of cohomology that we want to characterize.

For $0 \leq k \leq R$, let

$$\text{Gr}_k^P \supseteq Q'_k := \ker\{\eta^{R-k+1} : \text{Gr}_k^P \rightarrow \text{Gr}_{2R-k+2}^P\}$$

be the associated graded-primitive spaces (here η acts on the graded spaces of the perverse filtration on cohomology). Let $Q_k \subseteq G_k$ be the image of Q'_k via the first Deligne splitting. We have $G_k = \sum_{i \geq 0} \eta^i Q_{k-2i}$ for $0 \leq k \leq R$, and $G_{R+\kappa} = \eta^\kappa G_{R-\kappa}$ (here η acts on cohomology).

It follows that, in order to have a complete description of the first Deligne splitting that involves solely the action of η in cohomology, we only need to describe $Q_k \subseteq H^*(X, \mathbb{Q})$ in such terms.

We describe Q_k following [17, Section 2.7]. Note that its context is the one of the first Deligne splitting arising from working in the derived category, but the proof works verbatim in the present context of cohomology acted upon by a π -Lefschetz class. By (65), we have

$$\eta^j P_k H^*(X, \mathbb{Q}) \subset P_{k+2j} H^*(X, \mathbb{Q}).$$

Let $\Phi_0(\eta)$ be the composition of the morphisms

$$\Phi_0(\eta) : P_k H^*(X, \mathbb{Q}) \xrightarrow{\eta^{R-k+1}} P_{2R-k+2} H^*(X, \mathbb{Q}) \rightarrow \mathrm{Gr}_{2R-k+2}^P H^*(X, \mathbb{Q}),$$

where the first map is cup product and the second map is the projection to the graded piece. We obtain the sub-vector space

$$\mathrm{Ker}(\Phi_0(\eta)) \subset P_k H^*(X, \mathbb{Q}).$$

The morphisms $\Phi_m(\eta)$ are defined inductively for $m \geq 0$ as follows:

$$\Phi_m(\eta) : \mathrm{Ker}(\Phi_{m-1}(\eta)) \xrightarrow{\eta^{R-k+m}} P_{2R-k+2m} H^*(X, \mathbb{Q}) \rightarrow \mathrm{Gr}_{2R-k+2m}^P H^*(X, \mathbb{Q}).$$

Therefore for fixed k , we obtain a sequence of sub-vector spaces

$$\cdots \subset \mathrm{Ker}(\Phi_1(\eta)) \subset \mathrm{Ker}(\Phi_0(\eta)) \subset P_k H^*(X, \mathbb{Q}).$$

According to [17, Proposition 2.7.1], we have the desired description

$$Q_k = Q_k(\eta) = \mathrm{Ker}(\Phi_k(\eta)) \subset P_k H^*(X, \mathbb{Q}), \quad \forall 0 \leq k \leq R.$$

The description of the first Deligne splitting yields immediately the following comparison lemma, which expresses a kind of functoriality for the first Deligne splitting.

Lemma 3.3. *Let X_i ($i = 1, 2$) be nonsingular varieties with proper surjective morphisms $f_i : X_i \rightarrow Y_i$, and let $P_\star H^*(X_i, \mathbb{Q})$ be the corresponding perverse filtrations. Let*

$$\phi : H^*(X_1, \mathbb{Q}) \rightarrow H^*(X_2, \mathbb{Q})$$

be a morphism of graded $\mathbb{Q}$ -algebras satisfying

- (a) $\phi(P_k H^d(X_1, \mathbb{Q})) \subset P_k H^d(X_2, \mathbb{Q})$;
- (b) $\phi(\eta_1) = \eta_2$, with $\eta_i \in H^2(X_i, \mathbb{Q})$ an f_i -Lefschetz class for $i = 1, 2$.

Then we have

$$\phi(G_k H^*(X_1, \mathbb{Q})) \subset G_k H^*(X_2, \mathbb{Q}), \quad \forall k \geq 0,$$

with $H^(X_i, \mathbb{Q}) = \bigoplus_{k \geq 0} G_k H^*(X_i, \mathbb{Q})$ the first Deligne splitting associated with η_i , $i = 1, 2$.*

Proof. It follows from the description of the first Deligne splittings

$$H^*(X_i, \mathbb{Q}) = \bigoplus_{0 \leq k \leq r} \bigoplus_{j \geq 0} \left(\eta_i^j Q_{k-2j}(\eta_i) \bigoplus \eta_i^{r-k+j} Q_{k-2j}(\eta_i) \right) \quad i = 1, 2$$

summarized above, and the fact that the sub-vector spaces

$$\mathrm{Ker}(\Phi_k(\eta_i)) = Q_k(\eta_i) \subset P_k H^*(X_i, \mathbb{Q})$$

are preserved under the P -filtered morphism ϕ . □

Remark 3.4. Since cohomologically, the Hitchin fibration

$$h : \mathcal{M}_{\text{Dol}} \rightarrow \Lambda$$

behaves like the product of the fibrations $\hat{h} : \hat{\mathcal{M}}_{\text{Dol}} \rightarrow \hat{\Lambda}$ and $\text{Pic}^0(C) \rightarrow \text{pt}$ via the ring isomorphism

$$H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) = H^*(\hat{\mathcal{M}}_{\text{Dol}}, \mathbb{Q}) \otimes H^*(\text{Pic}^0(C), \mathbb{Q})$$

given in [11, Section 2.4], we see from the proof of Proposition 3.2 (cf. (72)) together with [9, Appendix] that the first Deligne splitting associated with any h -Lefschetz class, i.e. a class of the form $\mu\alpha \otimes 1 + 1 \otimes \xi$ with $\mu \neq 0$ and $\xi^g \neq 0$, has the form

$$(73) \quad G_k H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) = \bigoplus_{i+j=k} G_i H^*(\hat{\mathcal{M}}_{\text{Dol}}, \mathbb{Q}) \otimes H^j(\text{Pic}^0(C), \mathbb{Q}),$$

where $G_i H^*(\hat{\mathcal{M}}_{\text{Dol}}, \mathbb{Q})$ is the first Deligne splitting associated with the $\hat{h}$ -Lefschetz class

$$\alpha \in H^2(\hat{\mathcal{M}}_{\text{Dol}}, \mathbb{Q}).$$

Remark 3.5. In particular, the splitting (73) does not depend on the choice of an h -Lefschetz class.

For every genus $g \geq 2$, we expect, and actually prove in the $g = 2$ case, that (73) serves as the splitting in Conjecture 0.3.

3.4. Semismall maps and Hilbert schemes. In this section, we study the situation where our morphism $\pi : X \rightarrow Y$ can be factored as

$$X \xrightarrow{f} Z \xrightarrow{g} Y$$

with $f : X \rightarrow Z$ semismall and surjective; in particular, f is generically finite and $\dim(X) = \dim(Z)$. We further assume that there are closed irreducible subvarieties $Z_i \subset Z$ such that the decomposition theorem for f takes the form of a canonical finite direct sum decomposition

$$(74) \quad Rf_* \mathbb{Q}_X = \bigoplus_i \text{IC}_{Z_i}[-\dim(X)],$$

where each IC_{Z_i} is the (perverse) intersection cohomology complex of Z_i . Note that we have the following identity concerning intersection cohomology groups $\text{IH}^d(Z_i, \mathbb{Q}) = H^{d-\dim Z_i}(Z_i, \text{IC}_{Z_i})$. One of the Z_i is the total variety Z . The restriction of g to each Z_i yields the morphism

$$g_i : Z_i \rightarrow Y_i \subset Y.$$

We deduce from (74) a canonical decomposition of the cohomology of X

$$(75) \quad H^d(X, \mathbb{Q}) = \bigoplus_i \text{IH}^{d-c_i}(Z_i, \mathbb{Q}), \quad c_i = \dim X - \dim Z_i = \text{codim}_X Z_i.$$

Let R be the defect of semismallness of π and, for each i , let R_i be the defect of semismallness of $g_i : Z_i \rightarrow Y_i$.

Proposition 3.6. *Let $\alpha \in H^2(Z, \mathbb{Q})$ satisfy that, for every i , the restriction*

$$\alpha_i = \alpha|_{Z_i} \in H^2(Z_i, \mathbb{Q})$$

is a g_i -Lefschetz class with associated first Deligne splitting

$$\text{IH}^*(Z_i, \mathbb{Q}) = \bigoplus_k G_k \text{IH}^*(Z_i, \mathbb{Q}).$$

Then $\eta = f^*\alpha \in H^2(X, \mathbb{Q})$ is a π -Lefschetz class, whose associated first Deligne splitting is, under the identification (75), given by

$$(76) \quad G_k H^d(X, \mathbb{Q}) = \bigoplus_i G_{k-R+R_i} \mathrm{IH}^{d-c_i}(Z_i, \mathbb{Q}).$$

Proof. Recall that we have defined the perverse filtration P on $H^*(X, \mathbb{Q})$ concentrated in the interval $[0, 2R]$, and similarly, for every i , the perverse filtration P on $\mathrm{IH}^*(Z_i, \mathbb{Q})$ is concentrated in the interval $[0, 2R_i]$. It follows that, according to (74) and (75), the direct summand $\mathrm{IH}^*(Z_i, \mathbb{Q})$ contributes to $H^*(X, \mathbb{Q})$ in perversities in the interval $[R - R_i, R + R_i]$.

We apply the decomposition theorem to the composition

$$X \rightarrow Z \rightarrow Y$$

and obtain that the perverse filtration $P_* H^*(X, \mathbb{Q})$ can be expressed in terms of the perverse filtrations $P_* \mathrm{IH}^*(Z_i, \mathbb{Q})$ under (75), *i.e.*,

$$P_k H^d(X, \mathbb{Q}) = \bigoplus_i P_{k-R+R_i} \mathrm{IH}^{d-c_i}(Z_i, \mathbb{Q}).$$

By [15, Remark 4.4.3], the action of $\eta = f^*\alpha$ on the l.h.s. of (75) is the direct sum of the actions of the classes $\alpha_i = \alpha|_{Z_i}$ on the summands of the r.h.s.

Since every α_i is a g_i -Lefschetz class, we deduce that η is a π -Lefschetz class with associated first Deligne splitting decomposition (76). $\square$

Next, we show that, given a fibered abelian surface

$$p : A = E \times E' \rightarrow E'$$

as in Section 1.5, the splitting (15) of the perverse filtration on $H^*(A^{[n]}, \mathbb{Q})$ associated with the natural morphism $p_n : A^{[n]} \rightarrow E'^{(n)}$ is the first Deligne splitting induced by a p_n -Lefschetz class.

The morphism $p_n : A^{[n]} \rightarrow E'^{(n)}$ admits the natural factorization

$$(77) \quad A^{[n]} \xrightarrow{f} A^{(n)} \xrightarrow{g} E'^{(n)},$$

where the Hilbert–Chow morphism f is semismall [22]. There are canonical morphisms

$$\kappa_\nu : A^{(\nu)} \rightarrow A^{(n)}$$

together with a canonical stratification of $A^{(n)}$ indexed by the partitions ν of n ,

$$A^{(n)} = \bigcup_\nu A_\nu, \quad \overline{A_\nu} = \mathrm{Im}(\kappa_\nu) \subset A^{(n)}.$$

Note that the resulting morphism $\kappa_\nu : A^{(\nu)} \rightarrow \overline{A_\nu}$ is the normalization of the target, so that

$$(78) \quad H^*(A^{(\nu)}, \mathbb{Q}) = \mathrm{IH}^*(\overline{A_\nu}, \mathbb{Q}).$$

By [22, Theorem 3], the decomposition theorem associated with f takes the form analogous to (74),

$$Rf_* \mathbb{Q}_{A^{[n]}} = \bigoplus_\nu \mathrm{IC}_{\overline{A_\nu}}[-2n],$$

and the restriction of $g : A^{(n)} \rightarrow E'^{(n)}$ to each $\overline{A}_\nu$ is described by the commutative diagrams

$$(79) \quad \begin{array}{ccccc} A^{(\nu)} & \xrightarrow{\kappa_\nu} & \overline{A}_\nu & \hookrightarrow & A^{(n)} \\ \downarrow & & p_\nu \downarrow & & \downarrow \\ E'^{(\nu)} & \longrightarrow & \overline{E}'_\nu & \hookrightarrow & E'^{(n)}. \end{array}$$

We consider the p -ample class

$$\alpha = \text{pt} \otimes 1_{E'} \in H^2(E, \mathbb{Q}) \otimes H^0(E', \mathbb{Q}) \subset H^2(A, \mathbb{Q})$$

of $p : A \rightarrow E'$ which induces a Lefschetz class

$$(80) \quad \alpha^{(n)} \in H^2(A^{(n)}, \mathbb{Q})$$

with respect to $A^{(n)} \rightarrow E'^{(n)}$. It further induces a Lefschetz class

$$\alpha^{(\nu)} \in H^2(A^{(\nu)}, \mathbb{Q})$$

with respect to $A^{(\nu)} \rightarrow E'^{(\nu)}$ for every partition ν of n . By the diagram (79), the pullback of $\alpha^{(n)}$ to every $A^{(\nu)}$ via κ_ν coincides with $\alpha^{(\nu)}$.

By keeping in mind that $\text{codim}_{A^{(n)}} \overline{A}_\nu = 2n - 2l(\nu)$, that the defects of semismallness of the morphisms p_n and p_ν are n and $l(\nu)$, respectively, and the identity (78), we obtain Corollary 3.7 by applying Proposition 3.6 to the factorization (77).

Corollary 3.7. *The decomposition (15) is the first Deligne splitting induced by the $(p_n : A^{[n]} \rightarrow E'^{(n)})$ -Lefschetz class (f as in (77), $\alpha^{(n)}$ as in (80))*

$$\eta_{A^{[n]}} = f^* \alpha^{(n)} \in H^2(A^{[n]}, \mathbb{Q}).$$

3.5. A strengthened version of Theorem 2.1. We study the decomposition (63) constructed for Theorem 2.1. In the special case (31), we have

$$\mathcal{M}_{\beta, A} \cong A^{[n]} \times \hat{A}, \quad A = E \times E',$$

with the morphism $\pi_\beta : \mathcal{M}_{\beta, A} \rightarrow B$ given by the morphism $p_n \times q$ in (33). Corollary 3.7 implies that the decomposition (38) is the first Deligne splitting associated with the $(p_n \times q)$ -Lefschetz class

$$\eta_{A, \beta} = \eta_{A^{[n]}} \boxtimes 1_{\hat{A}} + 1_{A^{[n]}} \boxtimes (1_E \boxtimes \text{pt}) \in H^2(A^{[n]} \times \hat{A}, \mathbb{Q}) = H^2(\mathcal{M}_{\beta, A}, \mathbb{Q}).$$

By Theorem 2.8, a pair (A', β') with $\beta'^2 = 2n$ is perversely isomorphic to the special pair (A, β) given by (31). So there is a graded isomorphism

$$\phi : H^*(\mathcal{M}_{\beta, A}, \mathbb{C}) \xrightarrow{\sim} H^*(\mathcal{M}_{\beta', A'}, \mathbb{C})$$

of $\mathbb{C}$ -algebras preserving the corresponding perverse filtrations, and we obtain a decomposition (63) as

$$(81) \quad \tilde{G}_k H^d(\mathcal{M}_{\beta', A'}, \mathbb{C}) = \phi(\tilde{G}_k H^d(\mathcal{M}_{\beta, A}, \mathbb{C}));$$

see the proof of Proposition 2.7. Hence

$$\eta_{\beta', A'} = \phi(\eta_{\beta, A}) \in H^2(\mathcal{M}_{\beta', A'}, \mathbb{C})$$

is a $\pi_{\beta'}$ -Lefschetz class, and Lemma 3.3 implies that the decomposition (81) is the first Deligne splitting associated with $\eta_{\beta', A'}$.

This gives the following strengthened version of Theorem 2.1, which, we stress, is about any pair (A, β) with $v_\beta = (0, \beta, \chi)$ primitive, not just the special cases (31).

Theorem 3.8. *Theorem 2.1 holds for a splitting $\tilde{G}_*H^*(\mathcal{M}_{\beta,A}, \mathbb{C})$ given by the first Deligne splitting associated with a suitable π_β -Lefschetz class η_β , where $\pi_\beta : \mathcal{M}_{\beta,A} \rightarrow B$ is the morphism (22).*

4. TOPOLOGY OF HITCHIN FIBRATIONS

4.1. Overview. Throughout the section, we assume C is a nonsingular projective integral curve of genus g embedded into an abelian surface A . We study a kind of specialization morphism

$$(82) \quad \text{sp}^! : H^*(\mathcal{M}_{\beta,A}, \mathbb{Q}) \rightarrow H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}),$$

where $\mathcal{M}_{\text{Dol}}$ is the moduli of stable Higgs bundles of rank r and Euler characteristic χ , and

$$\beta = r[C] \in H_2(A, \mathbb{Z}).$$

Then we deduce Theorems 0.2, 0.4, and 0.5 from Theorem 2.1 via the properties of the morphism (82) which we establish hereafter.

4.2. Deformation to the normal cone. The moduli space $\mathcal{M}_{\text{Dol}}$ of stable Higgs bundles with rank r and Euler characteristic χ can be realized as the moduli space of 1-dimensional Gieseker-stable sheaves $\mathcal{F}$ on the cotangent bundle surface T^*C with (cf. see [4])

$$[\text{supp}(\mathcal{F})] = r[C] \in H_2(T^*C, \mathbb{Z}), \quad \chi(\mathcal{F}) = \chi.$$

In the following, we describe $\mathcal{M}_{\text{Dol}}$ as the “limit” of a trivial family of $\mathcal{M}_{\beta,A}$. A similar construction using $K3$ surfaces was considered in [19].

Assume $T = \mathbb{P}^1$ and $T^\circ = \mathbb{P}^1 \setminus 0 = \mathbb{C}$. Let

$$(83) \quad \mathcal{S} = \text{Bl}_{C \times 0}(A \times T) \setminus (A \times 0) \rightarrow T$$

be the total space of the deformation to the normal cone associated with the embedding $j_C : C \hookrightarrow A$. The central fiber of (83) is

$$\mathcal{S}_0 = T^*C \rightarrow 0 \in T,$$

and the restriction over T° is a trivial fibration

$$A \times T^\circ \rightarrow T^\circ \subset T.$$

We associate to β a family of homology classes

$$(84) \quad \beta_t = r[C] \in H_2(\mathcal{S}_t, \mathbb{Z}).$$

Let $\mathcal{M} \rightarrow T$ be the (coarse) relative moduli space which parametrizes, for each $t \in T$, pure one-dimensional Gieseker-stable sheaves $\mathcal{F}$ on $\mathcal{S}_t$ with $\chi(\mathcal{F}) = \chi$ and such that the support of $\mathcal{F}$ is a proper subscheme in the class β_t . Similarly, let $\mathcal{B} \rightarrow T$ denote the component of the relative Hilbert scheme which parametrizes Cartier divisors in $\mathcal{S}_t$ with proper support in the class β_t . Following [37, Section 5.3], we have a proper morphism

$$\begin{array}{ccc} \mathcal{M} & \xrightarrow{h_T} & \mathcal{B} \\ & \searrow & \swarrow \\ & T & \end{array}$$

over the base T , which, on the level of points, sends a sheaf $\mathcal{F}$ on $\mathcal{S}_t$ to its Fitting support, viewed as an element in $\mathcal{B}_t$. Clearly both $\mathcal{M}$ and $\mathcal{B}$ become trivial families when restricted over T° ,

$$\begin{array}{ccc} \mathcal{M}_{\beta,A} \times T^\circ & \xrightarrow{h_{T^\circ}} & B \times T^\circ \\ & \searrow & \swarrow \\ & T^\circ & \end{array} .$$

The fibers over $0 \in T$ recover the moduli space $\mathcal{M}_{\text{Dol}}$, the Hitchin base Λ , and the Hitchin fibration (2).

Lemma 4.1 seems standard; we include a brief proof since we are not aware of an adequate reference.

Lemma 4.1.

- (i) *Both $\mathcal{M}$ and $\mathcal{B}$ are irreducible varieties and smooth over T .*
- (ii) *There exists a universal class*

$$[\mathcal{F}_T] \in K_{\text{top}}(\mathcal{M} \times_T \mathcal{S})$$

whose restriction to each fiber gives a universal class for $\mathcal{M}_t \times \mathcal{S}_t$.

Proof. For part (i), we first argue for $\mathcal{B}$. For notational convenience, we define

$$\tilde{g} = \dim(\mathcal{B}_t) = r^2(g-1) + 1, \quad \forall t \in T.$$

We observe that the closure $\overline{\mathcal{B}^\circ}$ in $\mathcal{B}$ of

$$\mathcal{B}^\circ = B \times T^\circ$$

is irreducible of dimension $\tilde{g} + 1$, so that the intersection

$$\overline{\mathcal{B}^\circ} \cap \mathcal{B}_0 \subset \mathcal{B}_0$$

must have at least dimension $\tilde{g}$ at every closed point. This intersection is non-empty since it contains the divisor $r[C]$ which clearly deforms to the generic fiber. Since $\mathcal{B}_0$ is irreducible of dimension $\tilde{g}$, we have $\mathcal{B}_0 \subset \overline{\mathcal{B}^\circ}$. Therefore $\mathcal{B}$ is irreducible. This implies the flatness of $\mathcal{B}$ over the nonsingular curve T . Furthermore, since its fibers are all nonsingular, the flatness further implies that the morphism $\mathcal{B} \rightarrow T$ is smooth.

The same argument applies for $\mathcal{M}$. The only thing to check is that there exists a point in the central fiber $\mathcal{M}_0$ which deforms to the generic fiber. By the smoothness of $\mathcal{B}/T$, we can choose a nonsingular curve $Z_0 \subset \mathcal{S}_0$ which deforms to $Z_t \subset \mathcal{S}_t$. Any line bundle on Z_0 with Euler characteristic χ can be deformed as well and this gives the desired point in $\mathcal{M}_0$.

For part (ii), we follow the same argument from [31, Section 3.1]. We denote by $\mathfrak{M}$ the relative moduli stack of stable sheaves on $\mathcal{S}$, which is a $\mathbb{G}_m$ -gerbe over $\mathcal{M}$. It suffices to show that this gerbe is *topologically* trivializable, so that we can pull back the topological K -theory class of the universal family on $\mathfrak{M} \times_T \mathcal{S}$ to $\mathcal{M} \times_T \mathcal{S}$ via a section. To construct a trivialization, we take the nonsingular 3-fold $\mathcal{S}' = \text{Bl}_{C \times 0}(A \times T)$ which contains $\mathcal{S}$ as an open subset,

$$\mathcal{S} \subset \mathcal{S}'.$$

Let $\mathcal{M}_{\mathcal{S}'}$ be the (coarse) moduli space of stable sheaves on $\mathcal{S}'$ with Chern character given by $w = \text{ch}(i_{t*}\mathcal{F}) \in H^*(\mathcal{S}', \mathbb{Q})$, where

$$i_t : \mathcal{S}_t \xrightarrow{\sim} A \times t \subset \mathcal{S}', \quad t \neq 0$$

is the inclusion of a nonsingular fiber, and $\mathcal{F} \in \mathcal{M}$ is a coherent sheaf supported on $\mathcal{S}_t$. The moduli space $\mathcal{M}_{\mathcal{S}'}$ contains $\mathcal{M}$ as an open subvariety; therefore it is enough to show that the corresponding $\mathbb{G}_m$ -gerbe $\mathfrak{M}_{\mathcal{S}'} \rightarrow \mathcal{M}_{\mathcal{S}'}$ is topologically trivial. As in [31, Section 3.1], a trivialization can be constructed with a topological K-theory class in $\mathcal{S}'$ whose pairing with w is 1. For this, we can use the pullback to $\mathcal{S}'$ of a class on A , whose pairing with $(0, r[C], \chi)$ is 1, via the natural projection

$$\mathcal{S}' = \text{Bl}_{C \times 0}(A \times T) \rightarrow A. \quad \square$$

4.3. Specializations. Specialization morphisms with respect to perverse filtrations have been studied systematically in [10, 13]. We provide a concrete description of the specialization morphism in our setting as follows.

As before, we assume $T = \mathbb{P}^1$ and $T^\circ = \mathbb{P}^1 \setminus 0$. Let $f : W \rightarrow T$ be a smooth morphism, whose restriction over T° is a trivial product

$$f^\circ : W^\circ = W_t \times T^\circ \rightarrow T^\circ, \quad \forall t \neq 0,$$

with W_t proper. By Deligne's Global Invariant Cycle Theorem, see [16, Theorem 1.7.1] for example, the restriction

$$\text{res}_t : H^*(W, \mathbb{Q}) \rightarrow H^*(W^\circ, \mathbb{Q}) = H^*(W_t, \mathbb{Q})$$

is surjective for $t \neq 0$. Let

$$\alpha_1, \alpha_2 \in H^*(W, \mathbb{Q})$$

be two liftings of a class $\alpha \in H^*(W_t, \mathbb{Q})$ with $t \neq 0$. Since $H^*(W^\circ, \mathbb{Q}) = H^*(W_t, \mathbb{Q})$, the long exact sequences for the pairs (W, W_t) and (W, W°) are isomorphic. Since W is nonsingular, the relative cohomology for (W, W°) can be identified with the Borel-Moore homology of W_0 . It follows that

$$(85) \quad \alpha_1 - \alpha_2 = i_{0*}\gamma,$$

for some class γ in the Borel-Moore homology of W_0 . Here $i_0 : W_0 \hookrightarrow W$ is the closed embedding of the central fiber, and we identify the Borel-Moore homology and the cohomology of the nonsingular W in the equation (85). The excess intersection formula [20, Corollary 6.3], together with the triviality of the normal bundle $N_{W_0/W}$, implies that we have

$$\text{res}_0(\alpha_1) - \text{res}_0(\alpha_2) = i_0^* i_{0*} \gamma = c_1(N_{W_0/W}) \cap \gamma = 0.$$

We define the *specialization morphism*⁶

$$(86) \quad \text{sp}^! : H^*(W_t, \mathbb{Q}) \rightarrow H^*(W_0, \mathbb{Q}), \quad t \neq 0$$

as

$$(87) \quad \text{sp}^!(\alpha) = \text{res}_0(\tilde{\alpha}) \in H^*(W_0, \mathbb{Q}),$$

⁶In the literature, often, one defines a specialization morphism sp^* in the opposite direction, as it is dictated by the morphism $\sigma : i_0^* \rightarrow \psi_f$ with ψ_f nearby cycle-functor. In the present set-up, since W_0 is not proper, such an arrow sp^* does not exist; by dualizing σ , one sees that an arrow in the opposite direction always exists, but if W_0 is singular, for example, this arrow is not a familiar object.

where $\tilde{\alpha} \in H^*(W, \mathbb{Q})$ is any lifting of α . The discussion above implies that the class (87) does not depend on the lifting $\tilde{\alpha}$. Since res_0 is a homomorphism of $\mathbb{Q}$ -algebras, so is $\text{sp}^!$ (86).

Example 4.2. Let $\mathcal{S} \rightarrow T$ be the relative surface (83) associated with the embedding $j_C : C \hookrightarrow A$. We have the specialization morphism

$$(88) \quad \text{sp}^! : H^*(A, \mathbb{Q}) \rightarrow H^*(T^*C, \mathbb{Q}) = H^*(C, \mathbb{Q}).$$

By the definition of $\text{sp}^!$, the morphism (88) is given by the pullback along

$$T^*C \hookrightarrow \mathbb{P}(T^*C \oplus \mathcal{O}_C) \hookrightarrow \text{Bl}_{C \times 0}(A \times T) \rightarrow A \times T \rightarrow A.$$

In particular, we have

$$\text{sp}^!(\gamma) = j_C^* \gamma \in H^*(T^*C, \mathbb{Q}) = H^*(C, \mathbb{Q}), \quad \forall \gamma \in H^*(A, \mathbb{Q}).$$

Now we discuss the interaction between $\text{sp}^!$ and perverse filtrations. Let W/T be as above. We consider a commutative diagram

$$\begin{array}{ccc} W & \xrightarrow{h} & V \\ & \searrow & \swarrow \\ & T & \end{array}$$

where the morphism $h : W \rightarrow V$ is proper of relative dimension d . For every $t \in T$, there is a perverse filtration $P_* H^*(W_t, \mathbb{Q})$ associated with the morphism $h_t : W_t \rightarrow V_t$.

Proposition 4.3. *The specialization morphism (86) preserves⁷ the perverse filtrations, i.e.,*

$$\text{sp}^!(P_k H^*(W_t, \mathbb{Q})) \subset P_k H^*(W_0, \mathbb{Q}), \quad \forall k.$$

Proof. We show that there exist splittings of the perverse filtrations

$$H^*(W_t, \mathbb{Q}) = \bigoplus_i G_i H^*(W_t, \mathbb{Q}), \quad \forall t \in T$$

such that the splitting is constant for $t \in T^o$, and

$$\text{sp}^!(G_i H^*(W_t, \mathbb{Q})) \subset G_i H^*(W_0, \mathbb{Q}).$$

We apply the decomposition theorem [5] to the morphism $h : W \rightarrow V$, and fix an isomorphism

$$\phi : Rh_* \mathbb{Q}_W[\dim W - d] \xrightarrow{\sim} \bigoplus_{i=0}^{2d} \mathcal{P}_V^i[-i].$$

Here, the $\mathcal{P}_V^i$ are semisimple perverse sheaves on V . By the discussion in Section 1.3, the isomorphism ϕ induces a splitting of the perverse filtration on $H^*(W, \mathbb{Q})$ associated with h . By the smoothness of f and the compatibility of vanishing cycles with the derived pushforward Rh_* , we have $\varphi(Rh_* \mathbb{Q}_W) = 0$. As a result, [13, Corollary 3.1.6] shows that the restriction of $\mathcal{P}_V^i$ to a closed fiber over $t \in T$ is a shifted perverse sheaf,

$$\mathcal{P}_{V,t}^i[-1] = \text{res}_t^* \mathcal{P}_V^i[-1] \in \text{Perv}(V_t), \quad \forall t \in T.$$

⁷In fact, the proof shows more, namely that $\text{sp}^!$ is filtered strict.

Therefore, the restriction ϕ_t of the decomposition ϕ induces splittings of the perverse filtration

$$H^*(W_t, \mathbb{Q}) = \bigoplus_{i \geq 0} \phi_t(H^*(V_t, \mathcal{P}_{V,t}^i)), \quad \forall t \in T,$$

and we conclude from the definition of $\mathrm{sp}^!$ that

$$\mathrm{sp}^! (\phi_t(H^k(V_t, \mathcal{P}_{V,t}^i))) \subset \phi_0(H^k(V_0, \mathcal{P}_{V,0}^i)).$$

This completes the proof. $\square$

Now we consider the family

$$h_T : \mathcal{M} \rightarrow \mathcal{B}$$

over T constructed in Section 4.2, especially Lemma 4.1. Proposition 4.3 implies that the specialization morphism

$$\mathrm{sp}^! : H^*(\mathcal{M}_{\beta,A}, \mathbb{C}) \rightarrow H^*(\mathcal{M}_{\mathrm{Dol}}, \mathbb{C})$$

preserves the perverse filtrations. Let $\tilde{G}_* H^*(\mathcal{M}_{A,\beta}, \mathbb{Q})$ be a splitting given by Theorem 3.8. In particular, it is the first Deligne splitting induced by a $(\pi_\beta = h_{t \neq 0})$ -Lefschetz class⁸

$$\eta_{\beta,A} \in H^2(\mathcal{M}_{A,\beta}, \mathbb{C}).$$

Proposition 4.4. *The class*

$$\eta_0 = \mathrm{sp}^! (\eta_{\beta,A}) \in H^2(\mathcal{M}_{\mathrm{Dol}}, \mathbb{C})$$

is a Lefschetz class for the Hitchin fibration $h : \mathcal{M}_{\mathrm{Dol}} \rightarrow \Lambda$. Assume further that

$$H^*(\mathcal{M}_{\mathrm{Dol}}, \mathbb{C}) = \bigoplus_{k,j} \tilde{G}_k H^j(\mathcal{M}_{\mathrm{Dol}}, \mathbb{C})$$

is the first Deligne splitting associated with η_0 . Then we have (cf. Theorem 3.8)

$$(89) \quad \mathrm{sp}^! (\tilde{G}_k H^d(\mathcal{M}_{A,\beta}, \mathbb{C})) \subset \tilde{G}_k H^d(\mathcal{M}_{\mathrm{Dol}}, \mathbb{C}).^9$$

Proof. Let $F_{\beta,A} \subset \mathcal{M}_{\beta,A}$ be a closed fiber of the morphism $\pi_\beta : \mathcal{M}_{\beta,A} \rightarrow B$. We have

$$\dim(F_{\beta,A}) = \frac{1}{2} \dim(\mathcal{M}_{\beta,A}) = \tilde{g}.$$

Since the fiber class

$$[F_{\beta,A}] \in H^{2\tilde{g}}(\mathcal{M}_{\beta,A}, \mathbb{C})$$

lies in $P_0 H^{2\tilde{g}}(\mathcal{M}_{\beta,A}, \mathbb{C})$ by Proposition 1.1, and $\eta_{\beta,A}$ is a Lefschetz class, we obtain that

$$\eta_{\beta,A}^{\tilde{g}} \cap [F_{\beta,A}] = \eta_{\beta,A}^{\tilde{g}} \cup [F_{\beta,A}] \neq 0$$

by the hard Lefschetz condition. Hence after specialization, we have

$$\eta_0^{\dim(F_0)} \cap [F_0] = \eta_{\beta,A}^{\tilde{g}} \cap [F_{\beta,A}] \neq 0$$

with F_0 a closed fiber of $h : \mathcal{M}_{\mathrm{Dol}} \rightarrow \Lambda$. We conclude from Proposition 3.2 that η_0 is a Lefschetz class with respect to h . The inclusion (89) then follows from Lemma 3.3 (applied to $\phi = \mathrm{sp}^!$). $\square$

⁸Note that this class was denoted by $\eta_{\beta',A'}$ in Section 3.5, since A was special there.

⁹Here we use $\mathbb{C}$ -coefficients since the Lefschetz class $\eta_{A,\beta}$ lies in $H^2(\mathcal{M}_{\beta,A}, \mathbb{C})$; see Section 3.5.

Remark 4.5. Proposition 4.4 implies that $\mathrm{sp}^!(G_*H^*(\mathcal{M}_{\beta,A}, \mathbb{C}))$ splits the restricted perverse filtration $P_*H^*(\mathcal{M}_{\mathrm{Dol}}, \mathbb{C}) \cap \mathrm{Im}(\mathrm{sp}^!)$, *i.e.*,

$$P_kH^*(\mathcal{M}_{\mathrm{Dol}}, \mathbb{C}) \cap \mathrm{Im}(\mathrm{sp}^!) = \bigoplus_{i \leq k} \mathrm{sp}^! \left(\tilde{G}_iH^*(\mathcal{M}_{\beta,A}, \mathbb{C}) \right).$$

In general, for an arbitrary splitting $G_*H^*(\mathcal{M}_{\beta,A}, \mathbb{C})$ of the perverse filtration $P_*H^*(\mathcal{M}_{\beta,A}, \mathbb{C})$, it may not be true that

$$\mathrm{sp}^!(G_iH^*(\mathcal{M}_{\beta,A}, \mathbb{C})) \cap \mathrm{sp}^!(G_jH^*(\mathcal{M}_{\beta,A}, \mathbb{C})) = \{0\}, \quad \forall i \neq j.$$

It is thus crucial, in our approach to the results of this paper, to realize the splitting

$$H^*(\mathcal{M}_{\beta,A}, \mathbb{C}) = \bigoplus_{k,j} \tilde{G}_kH^j(\mathcal{M}_{\beta,A}, \mathbb{C})$$

as the first Deligne splitting associated with a Lefschetz class as in Theorem 3.8.

4.4. Normalized classes. Our purpose is to apply the specialization morphism $\mathrm{sp}^!$ to the tautological classes. We first discuss some properties of the normalized classes introduced in Section 0.3.

Following [24, 25, 40], the cohomology

$$(90) \quad H^*(\mathcal{M}_{\mathrm{Dol}}, \mathbb{Q}) = H^*(\mathcal{M}_B, \mathbb{Q})$$

can be understood by the associated PGL_n -character variety $\hat{\mathcal{M}}_B$. More precisely, let

$$H^*(\mathcal{M}_B, \mathbb{Q}) \cong H^*(\hat{\mathcal{M}}_B, \mathbb{Q}) \otimes H^*((\mathbb{C}^*)^{2g}, \mathbb{Q})$$

be the isomorphism established in [25, Section 1] and [24, Theorem 2.2.12]. Every class $w \in H^i(\hat{\mathcal{M}}_B, \mathbb{Q})$ can be naturally viewed as a class

$$w = w \otimes 1 \in H^i(\mathcal{M}_B, \mathbb{Q}).$$

The weights of the tautological classes associated with the universal PGL_r -bundle $\mathcal{T}$ on $C \times \mathcal{M}_B$ (induced by the universal GL_r -bundle) were calculated in [40],

$$(91) \quad \int_{\gamma} c_k(\mathcal{T}) \in {}^k\mathrm{Hdg}^{i+2k-2}(\mathcal{M}_B), \quad \forall k \geq 0, \forall \gamma \in H^i(C, \mathbb{Q}).$$

Lemma 4.6 deduces (5) from (91).

Lemma 4.6. *A twisted universal family $(\mathcal{U}^\alpha, \theta)$ on $C \times \mathcal{M}_{\mathrm{Dol}}$ satisfies*

$$(92) \quad \int_{\gamma} \mathrm{ch}_k^\alpha(\mathcal{U}) \in {}^k\mathrm{Hdg}^{i+2k-2}(\mathcal{M}_B), \quad \forall \gamma \in H^i(C, \mathbb{Q}), \forall k \geq 0,$$

if and only if $\mathrm{ch}^\alpha(\mathcal{U})$ is normalized.

Proof. We define

$${}^k\mathrm{Hdg}^d(C \times \mathcal{M}_B) = \bigoplus_{i+j=d} H^i(C, \mathbb{Q}) \otimes {}^k\mathrm{Hdg}^j(\mathcal{M}_B).$$

Then (92) is equivalent to

$$\mathrm{ch}^\alpha(\mathcal{U}) \in \bigoplus_k {}^k\mathrm{Hdg}^{2k}(C \times \mathcal{M}_B).$$

By a direct calculation using Chern roots, we have (recall that r is the rank)

$$\mathrm{ch}^\alpha(\mathcal{U}) = \mathrm{ch}(\mathcal{T}) \cup \exp\left(\frac{c_1(\mathcal{U})}{r} + \alpha\right).$$

By using the universal relations between Chern character and total Chern class, we note that (91) is equivalent to

$$\mathrm{ch}(\mathcal{T}) \in \bigoplus_k {}^k\mathrm{Hdg}^{2k}(C \times \mathcal{M}_B).$$

By [40], we have

$$H^0(\mathcal{M}_B, \mathbb{Q}) = {}^0\mathrm{Hdg}^0(\mathcal{M}_B), \quad H^2(\mathcal{M}_B, \mathbb{Q}) = {}^2\mathrm{Hdg}^2(\mathcal{M}_B),$$

which implies

$${}^1\mathrm{Hdg}^2(C \times \mathcal{M}_B) = H^1(C, \mathbb{Q}) \otimes H^1(\mathcal{M}_B, \mathbb{Q}).$$

Therefore, we obtain that (92) holds if and only if

$$\mathrm{ch}_1^\alpha(\mathcal{U}) = c_1(\mathcal{U}) + r\alpha \in H^1(C, \mathbb{Q}) \otimes H^1(\mathcal{M}_B, \mathbb{Q}).$$

This is equivalent to the condition that the class $\mathrm{ch}(\mathcal{U}^\alpha)$ is normalized. $\square$

Next, in parallel to Lemma 4.6, we give a sufficient criterion involving the perverse filtration for a class $\mathrm{ch}^\alpha(\mathcal{U})$ to be normalized.

Lemma 4.7. *Suppose we have a twisted universal family $(\mathcal{U}^\alpha, \theta)$ such that, for every $k \geq 0$ and every $\gamma \in H^{2i}(C, \mathbb{Q})$, the tautological class*

$$\int_\gamma \mathrm{ch}_k^\alpha(\mathcal{U}) \in H^{2i+2k-2}(\mathcal{M}_{\mathrm{Dol}}, \mathbb{Q})$$

has perversity k . Then $\mathrm{ch}^\alpha(\mathcal{U})$ is normalized.

Proof. We have

$$P_1 H^2(\mathcal{M}_{\mathrm{Dol}}, \mathbb{Q}) = 0, \quad P_0 H^0(\mathcal{M}_{\mathrm{Dol}}, \mathbb{Q}) = H^0(\mathcal{M}_{\mathrm{Dol}}, \mathbb{Q}).$$

The first vanishing follows from the decomposition (69), and the corresponding vanishing for $\hat{\mathcal{M}}_{\mathrm{Dol}}$ due to the fact that its second cohomology is generated by a relative ample class which must therefore have perversity 2.¹⁰ The second equality is clear in view of the isomorphism $H^0(\mathcal{M}_{\mathrm{Dol}}, \mathbb{Q}) \cong H^0(\Lambda, \mathbb{Q})$ via pull-back from the base of the Hitchin fibration (2). Since by assumption, any Künneth factor of $\mathrm{ch}_1^\alpha(\mathcal{U})$ in $H^*(\mathcal{M}_{\mathrm{Dol}}, \mathbb{Q})$ has perversity 1, we reach the desired conclusion

$$\mathrm{ch}_1^\alpha(\mathcal{U}) \in H^1(C, \mathbb{Q}) \otimes H^1(\mathcal{M}_{\mathrm{Dol}}, \mathbb{Q}). \quad \square$$

Remark 4.8. In fact, we see from the proofs of Lemmas 4.6 and 4.7 that both lemmas hold if we use $\mathbb{C}$ -coefficients for the cohomology groups and we allow α to be a $\mathbb{C}$ -class. In particular, a $\mathbb{C}$ -normalized class $\mathrm{ch}^\alpha(\mathcal{U})$ is unique and rational.

4.5. Specializations and tautological classes. Assume $j_C : C \hookrightarrow A$ is the closed embedding of the curve in an abelian surface. Let $\mathcal{S} \rightarrow T$ and $\mathcal{M} \rightarrow T$ be the family of surfaces and the relative moduli space of 1-dimensional sheaves introduced in Section 4.2. In the following, we construct a family of cohomology classes on $\mathcal{M}$ over T whose restriction to every closed fiber $\mathcal{M}_t$ is a twisted tautological class.

For our purpose, we take a relative compactification $\mathcal{S} \subset \mathcal{S}'$ over T as in the proof of Lemma 4.1(ii). Then $\mathcal{M}$ is the coarse moduli space of Gieseker-stable

¹⁰To see that a relative ample class has perversity 2, we apply [11, Theorem 1.4.8] together with the fact that an ample class does not vanish over a general fiber of the Hitchin morphism.

1-dimensional sheaves $\mathcal{F}$ on $\mathcal{S}'$ satisfying that $\text{supp}(\mathcal{F})$ is proper and contained in $\mathcal{S}_t \subset \mathcal{S}'_t$ with the numerical conditions

$$\chi(\mathcal{F}) = \chi, \quad [\text{supp}(\mathcal{F})] = \beta_t.$$

Here recall that β_t is the class (84). By the proof of Lemma 4.1(ii), there is a universal K -theory class $[\mathcal{F}_T]$ on $\mathcal{S}' \times_T \mathcal{M}$.

Let

$$\pi_{\mathcal{S}'} : \mathcal{S}' \times_T \mathcal{M} \rightarrow \mathcal{S}', \quad \pi_{\mathcal{M}} : \mathcal{S}' \times_T \mathcal{M} \rightarrow \mathcal{M}$$

be the projections. We have the Chern character

$$\text{ch}(\mathcal{F}_T) \in H^*(\mathcal{S}' \times_T \mathcal{M}, \mathbb{C})$$

defined by the universal class $[\mathcal{F}_T]$. For any class of the type

$$(93) \quad \tilde{\alpha} = \pi_{\mathcal{S}'}^* \tilde{\alpha}_1 + \pi_{\mathcal{M}}^* \tilde{\alpha}_2 \in H^2(\mathcal{S}' \times_T \mathcal{M}, \mathbb{C})$$

with $\tilde{\alpha}_1 \in H^2(\mathcal{S}', \mathbb{C})$, $\tilde{\alpha}_2 \in H^2(\mathcal{M}, \mathbb{C})$, we consider the relative twisted class

$$(94) \quad \text{ch}^{\tilde{\alpha}}(\mathcal{F}_T) = \exp(\tilde{\alpha}) \cup \text{ch}(\mathcal{F}_T),$$

whose degree $2k$ parts

$$\text{ch}_k^{\tilde{\alpha}}(\mathcal{F}_T) \in H^{2k}(\mathcal{S}' \times_T \mathcal{M}, \mathbb{C})$$

induce a relative tautological class

$$(95) \quad \int_{\tilde{\gamma}} \text{ch}_k^{\tilde{\alpha}}(\mathcal{F}_T) := \pi_{\mathcal{M}*} \left((\pi_{\mathcal{S}'}^* \tilde{\gamma} \cup \text{ch}_k^{\tilde{\alpha}}(\mathcal{F}_T)) \cap [\mathcal{S}' \times_T \mathcal{M}] \right) \in H_*^{\text{BM}}(\mathcal{M}, \mathbb{C}) = H^*(\mathcal{M}, \mathbb{C})$$

for any $\tilde{\gamma} \in H^*(\mathcal{S}', \mathbb{C})$. Here we use the Poincaré duality in the last identity, and we require the properness of $\pi_{\mathcal{M}}$ for the pushforward functor $\pi_{\mathcal{M}*} : H_*^{\text{BM}}(\mathcal{S}' \times_T \mathcal{M}, \mathbb{C}) \rightarrow H_*^{\text{BM}}(\mathcal{M}, \mathbb{C})$.

Now we check that (95) is the desired cohomology class on $\mathcal{M}$ which gives a family of twisted tautological classes over the base T .

For any closed point $t \in T$, base change [20, Proposition 1.7] implies that the restriction of (95) to a closed fiber $\mathcal{M}_t$ is a twisted universal class of the form

$$(96) \quad \int_{\gamma_t} \text{res}_t \left(\text{ch}_k^{\tilde{\alpha}}(\mathcal{F}_T) \right) \in H^*(\mathcal{M}_t, \mathbb{C}), \quad \gamma_t \in H^*(\mathcal{S}'_t, \mathbb{C}).$$

When $t \neq 0$, we have $\mathcal{S}_t = \mathcal{S}'_t$, and therefore (96) recovers a twisted universal class on $\mathcal{M}_{\beta, A}$,

$$\int_{\gamma_t} \text{ch}_k^{\alpha}(\mathcal{F}_{\beta}) \in H^*(\mathcal{M}_{\beta, A}, \mathbb{C}).$$

We now calculate the class (96) for $t = 0$. By [26, Theorem 4.6.5], there exists a coherent sheaf $\mathcal{F}_0$ supported on $T^*C \times \mathcal{M}_{\text{Dol}}$ which represents a universal class $[\mathcal{F}_0]$ on $\mathcal{S}' \times \mathcal{M}_{\text{Dol}}$. Moreover, the coherent sheaf $\mathcal{F}_0$ is supported on the universal spectral curve $\mathcal{C} \subset T^*C \times \mathcal{M}_{\text{Dol}}$ which is proper over $\mathcal{M}_{\text{Dol}}$. Hence by [3, Property 2.1], we have

$$(97) \quad \text{ch}_k(\mathcal{F}_0) \cap [\mathcal{S}'_0 \times \mathcal{M}_{\text{Dol}}] = \sum_i c_i [Z_i] \in \text{CH}_*(\mathcal{S}'_0 \times \mathcal{M}_{\text{Dol}}) \mathbb{Q},$$

where $c_i \in \mathbb{Q}$ and Z_i are closed subvarieties in $T^*C \times \mathcal{M}_{\text{Dol}}$ which are proper over $\mathcal{M}_{\text{Dol}}$ via the composition

$$(98) \quad Z_i \hookrightarrow T^*C \times \mathcal{M}_{\text{Dol}} \xrightarrow{p_{\mathcal{M}}} \mathcal{M}_{\text{Dol}}.$$

In particular, the expression (97) together with the properties for Z_i implies that the class

$$\int_{\gamma_0} \text{res}_0 \left(\text{ch}_k^{\tilde{\alpha}}(\mathcal{F}_T) \right) \in H^*(\mathcal{M}_{\text{Dol}}, \mathbb{C})$$

only depends on the restrictions of the classes γ_0 to $H^2(T^*C, \mathbb{C})$ and $\text{res}_0(\tilde{\alpha})$ to $H^2(T^*C, \mathbb{C}) \oplus H^2(\mathcal{M}_{\text{Dol}}, \mathbb{C})$.

Note that a class of the type

$$(99) \quad p_{\mathcal{M}*}(\omega \cup \text{ch}_k(\mathcal{F}_0)) := \sum_i c_i \cdot p_{\mathcal{M}*}(\omega \cap [Z_i]) \in H_*^{\text{BM}}(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) = H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q})$$

is well-defined for $\omega \in H^*(T^*C \times \mathcal{M}_{\text{Dol}}, \mathbb{C})$ due to the properness of (98). We see from the discussion above that the restriction of the class (95) to $\mathcal{M}_0 (= \mathcal{M}_{\text{Dol}})$ is given by the following:

$$(100) \quad \text{res}_0 \left(\int_{\tilde{\gamma}} \text{ch}_k^{\tilde{\alpha}}(\mathcal{F}_T) \right) = \int_{\text{sp}^!(\gamma_t)} \text{ch}_k^{\alpha_0}(\mathcal{F}_0) \in H^*(\mathcal{M}_{\text{Dol}}, \mathbb{C}).$$

Here $\text{sp}^!(\gamma_t) \in H^*(\mathcal{S}_0, \mathbb{C}) = H^*(T^*C, \mathbb{C})$, $\alpha_0 \in H^2(T^*C, \mathbb{C}) \oplus H^2(\mathcal{M}_{\text{Dol}}, \mathbb{C})$, and the class on the r.h.s. of (100) is defined by (99).

Proposition 4.9 shows that the tautological classes (4) on the Dolbeault side $\mathcal{M}_{\text{Dol}}$ are obtained by specializing certain other tautological classes on the compact geometry side $\mathcal{M}_{\beta, A}$.

Proposition 4.9. *Let*

$$\int_{\gamma} \text{ch}_k^{\alpha}(\mathcal{F}_{\beta}) \in H^*(\mathcal{M}_{\beta, A}, \mathbb{C})$$

be the classes of Theorem 2.1 with $\gamma \in H^(A, \mathbb{Q})$ a rational class on A ; then we have*

$$\text{sp}^! \left(\int_{\gamma} \text{ch}_k^{\alpha}(\mathcal{F}_{\beta}) \right) = c(k-1, j_C^* \gamma) \in H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}).$$

Proof. By the definition of $\text{sp}^!$ and (100), we have

$$\text{sp}^! \left(\text{res}_t \left(\int_{\tilde{\gamma}} \text{ch}_k^{\tilde{\alpha}}(\mathcal{F}_T) \right) \right) = \text{res}_0 \left(\int_{\tilde{\gamma}} \text{ch}_k^{\tilde{\alpha}}(\mathcal{F}_T) \right) = \int_{\text{sp}^!(\gamma_t)} \text{ch}_k^{\alpha_0}(\mathcal{F}_0)$$

for any $\tilde{\alpha}$ of the type (93).

A direct calculation by applying the Grothendieck–Riemann–Roch formula to the natural projection

$$\text{pr} : T^*C \times \mathcal{M}_{\text{Dol}} \rightarrow C \times \mathcal{M}_{\text{Dol}}$$

(see the paragraph before [34, Remark 8]) together with Example 4.2 yields

$$\int_{\text{sp}^!(\gamma)} \text{ch}_k^{\alpha_0}(\mathcal{F}_0) = \int_{j_C^* \gamma} \text{ch}_{k-1}^{\alpha_0}(\mathcal{U}).$$

Here $\mathcal{U} = \text{pr}_* \mathcal{F}_0$ is a universal family on $C \times \mathcal{M}_{\text{Dol}}$.

In conclusion, we obtain that

$$\text{sp}^! \left(\int_{\gamma} \text{ch}_k^{\alpha}(\mathcal{F}_{\beta}) \right) = \int_{j_C^* \gamma} \text{ch}_{k-1}^{\alpha_0}(\mathcal{U}).$$

Moreover, we know from Proposition 4.4 that the twisted class

$$\int_{j_C^* \gamma} \text{ch}_{k-1}^{\alpha_0}(\mathcal{U}), \quad \forall \gamma \in H^*(A, \mathbb{Q}), \quad \forall k \geq 1$$

has perversity $k - 1$. Hence Lemma 4.7 and Remark 4.8 imply, by recalling (4), that

$$\int_{j_C^* \gamma} \text{ch}_{k-1}^{\alpha_0}(\mathcal{U}) = c(k-1, j_C^* \gamma). \quad \square$$

4.6. Proofs of Theorems 0.2, 0.4, and 0.5. Since the perverse filtration with respect to the Hitchin fibration $h : \mathcal{M}_{\text{Dol}} \rightarrow \Lambda$ is locally constant when we deform the curve C [13], it suffices to prove all the three theorems for a curve which can be embedded in an abelian surface

$$j_C : C \hookrightarrow A.$$

After having done so, we can apply the specialization morphism

$$\text{sp}^! : H^*(\mathcal{M}_{\beta, A}, \mathbb{C}) \rightarrow H^*(\mathcal{M}_{\text{Dol}}, \mathbb{C})$$

introduced in Section 4.3. The following results are immediate consequences of Theorem 3.8, Proposition 4.4, and Proposition 4.9:

(i) We have

$$(101) \quad c(\gamma, k) \in \tilde{G}_k H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}), \quad \forall \gamma \in \text{Im}(j_C^*) \subset H^*(C, \mathbb{Q}).$$

(ii) The restriction of the decomposition

$$H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q}) = \bigoplus_{k,d} \tilde{G}_k H^d(\mathcal{M}_{\text{Dol}}, \mathbb{Q})$$

to the subalgebra of $H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q})$ generated by the classes (101) is multiplicative.

We first prove Theorem 0.2. The Abel-Jacobi morphism embeds a genus 2 curve C into its Jacobian

$$j_C : C \hookrightarrow \text{Jac}(C) = A.$$

Hence the restriction morphism j_C^* is surjective, and Theorem 0.2 follows from (i) and (ii) above.

The proof of Theorem 0.4 is similar. For any embedding j_C , the image of j_C^* always contains the sub-vector space

$$H^0(C, \mathbb{Q}) \oplus H^2(C, \mathbb{Q}) \subset H^*(C, \mathbb{Q}).$$

Hence the subalgebra

$$R^*(\mathcal{M}_{\text{Dol}}) \subset H^*(\mathcal{M}_{\text{Dol}}, \mathbb{Q})$$

is contained in the subalgebra generated by the classes (101), and we again conclude Theorem 0.4 by (i) and (ii).

Finally we treat the odd classes

$$c(\gamma, k) \in H^{2k-1}(\mathcal{M}_{\text{Dol}}, \mathbb{Q}), \quad \gamma \in H^1(C, \mathbb{Q})$$

and prove Theorem 0.5.

When the curve C has genus ≥ 3 , the restriction

$$(102) \quad j_C^* : H^1(A, \mathbb{Q}) \rightarrow H^1(C, \mathbb{Q})$$

is not surjective. We know from (i) that $c(\gamma, k)$ has perversity k for any γ lying in the image of (102). Since the monodromy group of the moduli space $\mathcal{M}_g$ of nonsingular genus g curves is the full symplectic group Sp_{2g} by [2], the sub-vector space

$$\text{Im}(j_C^* : H^1(A, \mathbb{Q}) \rightarrow H^1(C, \mathbb{Q})) \subset H^1(C, \mathbb{Q})$$

generates the total cohomology $H^1(C, \mathbb{Q})$ via the action of the monodromy group. We deduce Theorem 0.5 from [13].

ACKNOWLEDGMENTS

We are grateful to Tamás Hausel, Jochen Heinloth, Luca Migliorini, Vivek Shende, David Vogan, Qizheng Yin, Zhiwei Yun, and Zili Zhang for helpful discussions.

REFERENCES

- [1] M. F. Atiyah, *K-theory*, W. A. Benjamin, Inc., New York-Amsterdam, 1967. Lecture notes by D. W. Anderson. MR0224083
- [2] Norbert A'Campo, *Tresses, monodromie et le groupe symplectique* (French), Comment. Math. Helv. **54** (1979), no. 2, 318–327, DOI 10.1007/BF02566275. MR535062
- [3] Paul Baum, William Fulton, and Robert MacPherson, *Riemann-Roch for singular varieties*, Inst. Hautes Études Sci. Publ. Math. **45** (1975), 101–145. MR412190
- [4] Arnaud Beauville, M. S. Narasimhan, and S. Ramanan, *Spectral curves and the generalised theta divisor*, J. Reine Angew. Math. **398** (1989), 169–179, DOI 10.1515/crll.1989.398.169. MR998478
- [5] A. A. Beilinson, J. Bernstein, and P. Deligne, *Faisceaux pervers* (French), Analysis and topology on singular spaces, I (Luminy, 1981), Astérisque, vol. 100, Soc. Math. France, Paris, 1982, pp. 5–171. MR751966
- [6] Armand Borel, *Density properties for certain subgroups of semi-simple groups without compact components*, Ann. of Math. (2) **72** (1960), 179–188, DOI 10.2307/1970150. MR123639
- [7] W.-Y. Chuang, D.-E. Diaconescu, and G. Pan, *BPS states and the $P = W$ conjecture*, Moduli spaces, London Math. Soc. Lecture Note Ser., vol. 411, Cambridge Univ. Press, Cambridge, 2014, pp. 132–150. MR3221294
- [8] Mark Andrea de Cataldo, *Perverse sheaves and the topology of algebraic varieties*, Geometry of moduli spaces and representation theory, IAS/Park City Math. Ser., vol. 24, Amer. Math. Soc., Providence, RI, 2017, pp. 1–58. MR3752459
- [9] Mark Andrea A. de Cataldo, *Hodge-theoretic splitting mechanisms for projective maps*, J. Singul. **7** (2013), 134–156, DOI 10.5427/jsing.2013.7h. With an appendix containing a letter from P. Deligne. MR3077721
- [10] M. A. de Cataldo, *Perverse Leray filtration and specialization with applications to the Hitchin morphism*, to appear in Math. Proc. Cambridge Phil. Soc., arXiv:2102.02264, 2021.
- [11] Mark Andrea A. de Cataldo, Tamás Hausel, and Luca Migliorini, *Topology of Hitchin systems and Hodge theory of character varieties: the case A_1* , Ann. of Math. (2) **175** (2012), no. 3, 1329–1407, DOI 10.4007/annals.2012.175.3.7. MR2912707
- [12] Mark Andrea A. de Cataldo, Tamás Hausel, and Luca Migliorini, *Exchange between perverse and weight filtration for the Hilbert schemes of points of two surfaces*, J. Singul. **7** (2013), 23–38, DOI 10.5427/jsing.2013.7c. MR3033732
- [13] Mark Andrea A. de Cataldo and Daves Maulik, *The perverse filtration for the Hitchin fibration is locally constant*, Pure Appl. Math. Q. **16** (2020), no. 5, 1441–1464, DOI 10.4310/PAMQ.2020.v16.n5.a4. MR4221002
- [14] Mark Andrea A. de Cataldo and Luca Migliorini, *The Chow groups and the motive of the Hilbert scheme of points on a surface*, J. Algebra **251** (2002), no. 2, 824–848, DOI 10.1006/jabr.2001.9105. MR1919155
- [15] Mark Andrea A. de Cataldo and Luca Migliorini, *The Hodge theory of algebraic maps* (English, with English and French summaries), Ann. Sci. École Norm. Sup. (4) **38** (2005), no. 5, 693–750, DOI 10.1016/j.ansens.2005.07.001. MR2195257
- [16] Mark Andrea A. de Cataldo and Luca Migliorini, *The decomposition theorem, perverse sheaves and the topology of algebraic maps*, Bull. Amer. Math. Soc. (N.S.) **46** (2009), no. 4, 535–633, DOI 10.1090/S0273-0979-09-01260-9. MR2525735
- [17] Mark Andrea A. de Cataldo and Luca Migliorini, *Hodge-theoretic aspects of the decomposition theorem*, Algebraic geometry—Seattle 2005. Part 2, Proc. Sympos. Pure Math., vol. 80, Amer. Math. Soc., Providence, RI, 2009, pp. 489–504, DOI 10.1090/pspum/080.2/2483945. MR2483945

- [18] Pierre Deligne, *Décompositions dans la catégorie dérivée* (French), Motives (Seattle, WA, 1991), Proc. Sympos. Pure Math., vol. 55, Amer. Math. Soc., Providence, RI, 1994, pp. 115–128, DOI 10.1090/pspum/055.1/1265526. MR1265526
- [19] Ron Donagi, Lawrence Ein, and Robert Lazarsfeld, *Nilpotent cones and sheaves on K3 surfaces*, Birational algebraic geometry (Baltimore, MD, 1996), Contemp. Math., vol. 207, Amer. Math. Soc., Providence, RI, 1997, pp. 51–61, DOI 10.1090/conm/207/02719. MR1462924
- [20] William Fulton, *Intersection theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 2, Springer-Verlag, Berlin, 1984, DOI 10.1007/978-3-662-02421-8. MR732620
- [21] Lothar Göttsche, *The Betti numbers of the Hilbert scheme of points on a smooth projective surface*, Math. Ann. **286** (1990), no. 1-3, 193–207, DOI 10.1007/BF01453572. MR1032930
- [22] Lothar Göttsche and Wolfgang Soergel, *Perverse sheaves and the cohomology of Hilbert schemes of smooth algebraic surfaces*, Math. Ann. **296** (1993), no. 2, 235–245, DOI 10.1007/BF01445104. MR1219901
- [23] Andrew Harder, Zhiyuan Li, Junliang Shen, and Qizheng Yin, *$P = W$ for Lagrangian fibrations and degenerations of hyper-Kähler manifolds*, Forum Math. Sigma **9** (2021), Paper No. e50, 6, DOI 10.1017/fms.2021.31. MR4277887
- [24] Tamás Hausel and Fernando Rodriguez-Villegas, *Mixed Hodge polynomials of character varieties*, Invent. Math. **174** (2008), no. 3, 555–624, DOI 10.1007/s00222-008-0142-x. With an appendix by Nicholas M. Katz. MR2453601
- [25] Tamás Hausel and Michael Thaddeus, *Generators for the cohomology ring of the moduli space of rank 2 Higgs bundles*, Proc. London Math. Soc. (3) **88** (2004), no. 3, 632–658, DOI 10.1112/S0024611503014618. MR2044052
- [26] Daniel Huybrechts and Manfred Lehn, *The geometry of moduli spaces of sheaves*, Aspects of Mathematics, E31, Friedr. Vieweg & Sohn, Braunschweig, 1997, DOI 10.1007/978-3-663-11624-0. MR1450870
- [27] Nigel Hitchin, *Stable bundles and integrable systems*, Duke Math. J. **54** (1987), no. 1, 91–114, DOI 10.1215/S0012-7094-87-05408-1. MR885778
- [28] J. Le Potier, *Faisceaux semi-stables de dimension 1 sur le plan projectif* (French), Rev. Roumaine Math. Pures Appl. **38** (1993), no. 7-8, 635–678. MR1263210
- [29] Joseph Le Potier, *Systèmes cohérents et structures de niveau* (French, with English and French summaries), Astérisque **214** (1993), 143. MR1244404
- [30] Eyal Markman, *Generators of the cohomology ring of moduli spaces of sheaves on symplectic surfaces*, J. Reine Angew. Math. **544** (2002), 61–82, DOI 10.1515/crll.2002.028. MR1887889
- [31] Eyal Markman, *Integral generators for the cohomology ring of moduli spaces of sheaves over Poisson surfaces*, Adv. Math. **208** (2007), no. 2, 622–646, DOI 10.1016/j.aim.2006.03.006. MR2304330
- [32] Eyal Markman, *On the monodromy of moduli spaces of sheaves on K3 surfaces*, J. Algebraic Geom. **17** (2008), no. 1, 29–99, DOI 10.1090/S1056-3911-07-00457-2. MR2357680
- [33] Eyal Markman, *Lagrangian fibrations of holomorphic-symplectic varieties of $K3^{[n]}$ -type*, Algebraic and complex geometry, Springer Proc. Math. Stat., vol. 71, Springer, Cham, 2014, pp. 241–283, DOI 10.1007/978-3-319-05404-9_10. MR3278577
- [34] E. Markman, *The monodromy of generalized Kummer varieties and algebraic cycles on their intermediate Jacobians*, arXiv:1805.11574, 2018.
- [35] Daves Maulik and Yukinobu Toda, *Gopakumar-Vafa invariants via vanishing cycles*, Invent. Math. **213** (2018), no. 3, 1017–1097, DOI 10.1007/s00222-018-0800-6. MR3842061
- [36] A. Mellit, *Cell decompositions of character varieties*, arXiv:1905.10685, 2019.
- [37] David Mumford, *Geometric invariant theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete, (N.F.), Band 34, Springer-Verlag, Berlin-New York, 1965. MR0214602
- [38] J. Shen and Q. Yin, *Topology of Lagrangian fibrations and Hodge theory and Hodge theory of hyper-Kähler manifolds*, Appendix by Claire Voisin, Duke Math. J., to appear.
- [39] J. Shen and Z. Zhang, *Perverse filtrations, Hilbert schemes, and the $P=W$ conjecture for parabolic Higgs bundles*, Algebr. Geom., **8** (4), 2021, pp. 465–489.
- [40] Vivek Shende, *The weights of the tautological classes of character varieties*, Int. Math. Res. Not. IMRN **22** (2017), 6832–6840, DOI 10.1093/imrn/rnv363. MR3737322
- [41] Carlos T. Simpson, *Higgs bundles and local systems*, Inst. Hautes Études Sci. Publ. Math. **75** (1992), 5–95. MR1179076

- [42] Carlos T. Simpson, *Moduli of representations of the fundamental group of a smooth projective variety. II*, Inst. Hautes Études Sci. Publ. Math. **80** (1994), 5–79 (1995). MR1320603
- [43] Geordie Williamson, *The Hodge theory of the decomposition theorem*, Astérisque **390** (2017), Exp. No. 1115, 335–367. Séminaire Bourbaki. Vol. 2015/2016. Exposés 1104–1119. MR3666031
- [44] Kōta Yoshioka, *Moduli spaces of stable sheaves on abelian surfaces*, Math. Ann. **321** (2001), no. 4, 817–884, DOI 10.1007/s002080100255. MR1872531
- [45] Zili Zhang, *Multiplicativity of perverse filtration for Hilbert schemes of fibered surfaces*, Adv. Math. **312** (2017), 636–679, DOI 10.1016/j.aim.2017.03.028. MR3635821

DEPARTMENT OF MATHEMATICS, STONY BROOK UNIVERSITY, STONY BROOK, NY 11794, USA
Email address: `mark.decataldo@stonybrook.edu`

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, 77 MASSACHUSETTS AVE, CAMBRIDGE MA 02138, USA
Email address: `maulik@mit.edu`

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, 442 DUNHAM LAB, 10 HILLHOUSE AVE., NEW HAVEN, CT 06511, USA
Email address: `junliang.shen@yale.edu`