

Anomaly cascade in (2 + 1)-dimensional fermionic topological phases

Daniel Bulmash  and Maissam Barkeshli

Condensed Matter Theory Center and Joint Quantum Institute, Department of Physics, University of Maryland, College Park, Maryland 20472, USA



(Received 22 October 2021; revised 28 February 2022; accepted 1 March 2022; published 15 April 2022)

We develop a theory of anomalies of fermionic topological phases of matter in (2 + 1)D with a general fermionic symmetry group G_f . In general, G_f can be a nontrivial central extension of the bosonic symmetry group G_b by fermion parity $(-1)^F$. We encounter four layers of obstructions to gauging the G_f symmetry, which we dub the anomaly cascade: (i) An $\mathcal{H}^1(G_b, \mathbb{Z}_T)$ obstruction to extending the symmetry permutations on the anyons to the fermion parity gauged theory, (ii) An $\mathcal{H}^2(G_b, \ker r)$ obstruction to extending the G_b group structure of the symmetry permutations to the fermion parity gauged theory, where r is a map that restricts symmetries of the fermion parity gauged theory to the anyon theory, (iii) An $\mathcal{H}^3(G_b, \mathbb{Z}_2)$ obstruction to extending the symmetry fractionalization class to the fermion parity gauged theory, and (iv) the well-known $\mathcal{H}^4(G_b, U(1))$ obstruction to developing a consistent theory of G_b symmetry defects for the fermion parity gauged theory. We describe how the \mathcal{H}^2 obstruction can be canceled by anomaly inflow from a bulk (3 + 1)D symmetry-protected topological state (SPT) and also its relation to the Arf invariant of spin structures on a torus. If any anomaly in the above sequence is nontrivial, the subsequent ones become relative anomalies. A number of conjectures regarding symmetry actions on super-modular categories, guided by general expectations of anomalies in physics, are also presented.

DOI: [10.1103/PhysRevB.105.155126](https://doi.org/10.1103/PhysRevB.105.155126)

I. INTRODUCTION

Topological phases of matter in (2 + 1)D can exhibit the remarkable phenomenon of symmetry fractionalization: In the presence of a global symmetry group G , topologically nontrivial quasiparticles can carry fractional quantum numbers under G . Such phases are called symmetry-enriched topological phases (SETs) and are described in full generality using the mathematics of G -crossed modular tensor categories [1–4]. Remarkably, some patterns of symmetry fractionalization yield SETs that, while they are mathematically consistent in the absence of a background G gauge field, cannot be consistently coupled to a G gauge field. The physical interpretation of this inconsistency is that such SETs cannot occur in a purely (2 + 1)D system with the symmetry generated on-site [1,5–7]. In the language of quantum field theory, such SETs are said to have a 't Hooft anomaly.

Anomalies provide crucial information about a phase of matter; they are renormalization group invariants, which connect microscopic and low-energy physics, leading, for example, to generalizations of the famed Lieb-Schulz-Mattis theorem [8]. Anomalies are subject to the bulk-boundary correspondence, in the sense that the classification of anomalies in d space-time dimensions is believed to be equivalent to the classification of invertible topological phases in $(d + 1)$ space-time dimensions [9]. In particular, anomalous (2 + 1)D SETs can exist on the surface of (3 + 1)D symmetry-protected topological phases (SPTs) [5,6,10–25]. This remarkable fact begs the question of understanding anomaly inflow—can we determine physical processes on the boundary that are sensitive to the anomaly and then see how the nontrivial bulk cancels that anomaly?

For bosonic SETs, the 't Hooft anomaly is relatively well understood [1,4–7,24,26]. A bosonic (3 + 1)D G -SPT is (partially) specified by an element of the cohomology group $\mathcal{H}^4(G, U(1))$, and this element dictates the inconsistency of fusion and braiding of symmetry defects together with anyons in the (2 + 1)D boundary SET.

In contrast, the data specifying a (3 + 1)D fermionic SPT (FSPT) has much more structure [27–30], which involve the cohomology groups $\mathcal{H}^1(G_b, \mathbb{Z}_T)$, $\mathcal{H}^2(G_b, \mathbb{Z}_2)$, $\mathcal{H}^3(G_b, \mathbb{Z}_2)$, and $\mathcal{H}^4(G_b, U(1))$, where G_b is the “bosonic” symmetry group obtained by modding out fermion parity symmetry. Specifically, (3 + 1)D FSPTs form an Abelian group that corresponds to a group extension involving the groups $\mathcal{H}^1(G_b, \mathbb{Z}_T)$, $\mathcal{H}^2(G_b, \mathbb{Z}_2)/\Gamma^2$, $\mathcal{H}^3(G_b, \mathbb{Z}_2)/\Gamma^3$, and $\mathcal{H}^4(G_b, U(1))/\Gamma^4$, where Γ^i are certain subgroups of the \mathcal{H}^i .

In particular, recently [30] have shown that (3 + 1)D FSPTs can in general be characterized by a set of data $(n_1, n_2, n_3, v_4) \in Z^1(G_b, \mathbb{Z}_T) \times C^2(G_b, \mathbb{Z}_2) \times C^3(G_b, \mathbb{Z}_2) \times C^4(G_b, U(1))$, where C^k denotes k cochains, Z^k denotes k cocycles, and \mathbb{Z}_T refers to the integers with an action of G_b according to whether group elements are antiunitary, which can be thought of as involving time-reversal symmetry. The data (n_1, n_2, n_3, v_4) satisfy a complicated set of consistency equations and equivalences. These data imply that a (3 + 1)D FSPT generically determines an element in $[n_1] \in \mathcal{H}^1(G_b, \mathbb{Z}_T)$. Moreover, when we can set $n_1 = 0$, a (3 + 1)D FSPT determines an element $[n_2] \in \mathcal{H}^2(G_b, \mathbb{Z}_2)/\Gamma^2$. When we can set $n_1, n_2 = 0$, a (3 + 1)D FSPT determines an element $[n_3] \in \mathcal{H}^3(G_b, \mathbb{Z}_2)/\Gamma^3$. Finally when we can set $n_1, n_2, n_3 = 0$, a (3 + 1)D FSPT determines an element of $[v_4] \in \mathcal{H}^4(G_b, U(1))/\Gamma^4$. The subgroups Γ^i can be explicitly determined in general [30].

The main aim of this paper is to obtain a detailed and general understanding of the 't Hooft anomaly of any $(2+1)$ D fermionic SET (FSET), and in particular the appearance of the above cohomology groups. The most general description of the anomaly is that it is an obstruction to gauging the full global symmetry group G_f . Rather than gauging G_f all at once, our approach is to gauge fermion parity first, which is always possible [31]. If the $G_b = G_f/\mathbb{Z}_2^f$ symmetry is to be preserved after gauging, the data specifying symmetry fractionalization must be lifted to the parity-gauged theory. We break the process of finding a consistent lift into a sequence of physically meaningful steps, each of which may be obstructed; the obstruction to each step is given by a piece of data. The appearance of FSPT data as an inconsistency of the $(2+1)$ D boundary SET is only partially understood [32,33]. Reference [32] gives a partial understanding of the \mathcal{H}^3 anomaly of FSETs, which we will make fully general in this paper. Reference [33] gives examples, mostly with $G_b = \mathbb{Z}_2^T$, explaining how “layers” of anomalies of fermionic phases in general dimensions appear in the action of symmetries on the Hilbert space on a spatial torus, but does not extract any cohomological obstructions for the $(2+1)$ D theory. Our formalism is fully general and starts from the general algebraic data characterizing the FSET rather than the Hilbert space on a torus, although we will make direct contact with Ref. [33] in Sec. VIC. Reference [25] recently showed how in general, given any $(2+1)$ D FSET, one can identify the bulk SPT using a state sum construction. However the results of [25] do not directly explain the appearance of the above cohomology groups in terms of an inconsistency of the boundary $(2+1)$ D FSET.

A. Summary of main results

We start with a super-modular category \mathcal{C} , which is a unitary braided fusion category (UBFC) that captures the braiding and fusion properties of the anyons. A super-modular category contains a single “invisible” particle ψ , which braids trivially with all other particles and which physically corresponds to the local fermion of the system. We then consider the minimal modular extensions $\check{\mathcal{C}}_v$ of the super-modular category. A minimal modular extension is a unitary modular tensor category (UMTC) that characterizes the phase obtained by gauging fermion parity. In particular $\check{\mathcal{C}}_v$ characterizes properties of the anyons, the local fermion ψ , and the fermion parity vortices. It was recently proved that every super-modular category admits a minimal modular extension [31], and therefore, according to the “16-fold way” theorem [34], admits exactly 16 distinct minimal modular extensions labeled by $v = 0, \dots, 15$.

In the absence of symmetry, it is expected that a $(2+1)$ D fermionic topological phase can be fully specified by either a choice of (\mathcal{C}, c_-) or $(\check{\mathcal{C}}_v, c_-)$, where c_- is the chiral central charge of the theory, which, physically, determines the system's thermal Hall conductivity. \mathcal{C} determines the theory modulo a fermionic invertible phase (e.g., up to stacking with $p+ip$ superconductors), and therefore determines $c_- \bmod 1/2$. In contrast, $\check{\mathcal{C}}_v$ determines the phase modulo a bosonic invertible phase, and therefore determines $c_- \bmod 8$.

Two different minimal modular extensions $\check{\mathcal{C}}_v$ and $\check{\mathcal{C}}_{v'}$ have central charges that differ by $(v - v')/2 \bmod 8$.

Fermionic systems always have a \mathbb{Z}_2 fermion parity symmetry $(-1)^F$, which generates a symmetry group we call \mathbb{Z}_2^f . The full symmetry group G_f of a fermionic system is in general a central extension of $G_b = G_f/\mathbb{Z}_2^f$ by \mathbb{Z}_2^f , characterized by a cocycle

$$\omega_2 \in Z^2(G_b, \mathbb{Z}_2). \quad (1)$$

In this paper, we assume that we are given G_f symmetry fractionalization data on \mathcal{C} , as detailed in [35,36], and which we review in detail in Sec. IVC. Briefly, this amounts to a map

$$[\rho] : G_b \rightarrow \text{Aut}_{LR}(\mathcal{C}), \quad (2)$$

where $\text{Aut}_{LR}(\mathcal{C})$ is the group of “locality-respecting” braided autoequivalences of \mathcal{C} . $\text{Aut}_{LR}(\mathcal{C})$ is similar to the group $\text{Aut}(\mathcal{C})$ of braided autoequivalences, except it takes the locality of the fermion into account by restricting the classes of maps that are considered trivial to those that act trivially on the fermion. Here the representative maps ρ_g satisfy

$$\rho_{gh} = \kappa_{g,h} \rho_g \rho_h, \quad (3)$$

where $\kappa_{g,h}$ is a natural isomorphism [1]. Once a representative set of maps ρ_g are chosen, symmetry fractionalization is characterized by a set of phases $\eta_a(g, h) \in U(1)$ for each anyon a . These phases $\eta_a(g, h)$ are subject to a series of consistency equations, constraints, and gauge equivalences.

Given a minimal modular extension $\check{\mathcal{C}}_v$, our aim is to lift the given symmetry fractionalization data from \mathcal{C} to $\check{\mathcal{C}}_v$. That is, we wish to gauge fermion parity while preserving the G_b symmetry. We perform this procedure systematically and characterize the cascade of obstructions that appear along the way.

I. Anomaly cascade: First layer

The first layer of the anomaly cascade is an obstruction to lifting the maps $[\rho_g]$ to autoequivalences of the fermion parity-gauged theory $\check{\mathcal{C}}$.

More precisely, the first step is to define a lifted “topological symmetry” of $\check{\mathcal{C}}$, that is, a map

$$[\check{\rho}] : G_b \rightarrow \text{Aut}_{LR}(\check{\mathcal{C}}_v). \quad (4)$$

For a particular v , generically not all elements of $\text{Aut}_{LR}(\mathcal{C})$ can be lifted to elements of $\text{Aut}_{LR}(\check{\mathcal{C}}_v)$. Thus we encounter the first possible obstruction, which concerns whether there exists some v such that for every $g \in G_b$, the autoequivalence $[\rho_g] \in \text{Aut}_{LR}(\mathcal{C})$ can be lifted to an element $[\check{\rho}_g] \in \text{Aut}_{LR}(\check{\mathcal{C}}_v)$.

In the case that there does not exist a v such that one can lift every element $[\rho_g]$ to $[\check{\rho}_g] \in \text{Aut}_{LR}(\check{\mathcal{C}}_v)$, a weaker lift may be possible. In particular, it may be possible that there exists at least one pair v_1 and v_2 , such that a representative of each $[\rho_g]$ lifts to a map

$$\check{\rho}_g : \check{\mathcal{C}}_{v_1} \rightarrow \check{\mathcal{C}}_{v_2}. \quad (5)$$

By comparing with the classification of $(3+1)$ D fermion SPTs, which defines the anomaly of $(2+1)$ D fermionic topological phases, we conjecture that the existence of such a weak lift is unobstructed:

Conjecture I.1. Given a super-modular category \mathcal{C} with an action $[\rho_g] : G_b \rightarrow \text{Aut}_{LR}(\mathcal{C})$, there always exists at least one pair v_1 and v_2 such that there exists a map $\check{\rho}_g : \check{\mathcal{C}}_{v_1} \rightarrow \check{\mathcal{C}}_{v_2}$ and $\check{\rho}_g$ restricts to a representative ρ_g on \mathcal{C} .

Next, regarding lifts of the form Eq. (5), we prove the following:

Theorem I.1. Let $\rho_g : \mathcal{C} \rightarrow \mathcal{C}$ be a unitary map. Then any pair (v_1, v_2) satisfying Eq. (5) must have $v_1 = v_2$.

This follows from the fact that the number of anyons, quantum dimensions, and topological twists must be invariant under the map $\check{\rho}_g$, which fixes $v_1 = v_2$ by the Gauss sum. Furthermore, we have

Theorem I.2. Let $\rho_g : \mathcal{C} \rightarrow \mathcal{C}$ be an antiunitary map. Then the pair (v_1, v_2) satisfying Eq. (5) must necessarily satisfy $e^{2\pi i c_{v_2}/8} = e^{-2\pi i c_{v_1}/8} = e^{2\pi i (c_{v_1} + o_1(\mathbf{g})/2)/8}$, for some integer $o_1(\mathbf{g})$. If there are multiple pairs $(v_{i,1}, v_{i,2})$ satisfying Eq. (5), then $e^{2\pi i c_{v_{i,2}}/8} = e^{-2\pi i c_{v_{i,1}}/8} = e^{2\pi i (c_{v_{i,1}} + o_1(\mathbf{g})/2)/8}$, where $o_1(\mathbf{g})$ is independent of i , modulo 2.

A corollary of the above is that:

Corollary I.1. A collection of maps $[\rho_g] \in \text{Aut}_{LR}(\mathcal{C})$ for each $\mathbf{g} \in G_b$ defines an element $[o_1] \in \mathcal{H}^1(G_b, \mathbb{Z}_T)$.

We conjecture that $[o_1]$ is really the obstruction to lifting the symmetry the action, in the following sense:

Conjecture I.2. Suppose that for each $\mathbf{g} \in G_b$ we are given a lift $\check{\rho}_g^{(1)} : \check{\mathcal{C}}_{v_1} \rightarrow \check{\mathcal{C}}_{v_2}$ of $\rho_g : \mathcal{C} \rightarrow \mathcal{C}$. These maps define $[o_1] \in \mathcal{H}^1(G_b, \mathbb{Z}_T)$. Then there exists a minimal modular extension $\check{\mathcal{C}}_{v_0}$ with lifts $\check{\rho}_g^{(0)} : \check{\mathcal{C}}_{v_0} \rightarrow \check{\mathcal{C}}_{v_0}$ for every $\mathbf{g} \in G_b$ if and only if $[o_1] = 0$.

A well-known example of the above \mathcal{H}^1 anomaly occurs for the $\text{SO}(3)_3$ super-modular category, which was studied in [15] as an example of a (2 + 1)D surface theory for an odd index topological superconductor in the class DIII. (Here $\text{SO}(3)_3$ consists of the integer spin representations of the $\text{SU}(2)_6$ affine Kac-Moody algebra, which arises in $\text{SO}(3)_3$ Chern-Simons theory.) All minimal modular extensions of this theory have central charge $c = 1/4 \bmod 1/2$, and no minimal modular extension can be compatible with time-reversal symmetry. This is because a UMTC that is compatible with time-reversal symmetry must have central charge $c_- = 0 \bmod 4$.

2. Anomaly cascade: Second layer

The second layer of the anomaly cascade is an obstruction to choosing the autoequivalences of the fermion parity-gauged theory so that they compose in an appropriate way, determined by the group G_f and details of the fermion parity-gauged theory.

Suppose that the \mathcal{H}^1 anomaly vanishes. Then, according to the discussion above, for each $\mathbf{g} \in G_b$, we assume we have an invertible map

$$\check{\rho}_g : \check{\mathcal{C}}_v \rightarrow \check{\mathcal{C}}_v, \quad (6)$$

for at least some subset of the possible values of v . Unless otherwise stated, we now fix a particular choice of v and omit it from our notation.

Below we will briefly summarize the obstruction theory for the case where the extension $\check{\mathcal{C}}_v$ contains an Abelian fermion

parity vortex. The more complicated cases will be discussed in the main text.

When $\check{\mathcal{C}}_v$ contains an Abelian fermion parity vortex, $[\rho]$ is a group homomorphism, and we need to require that the lift $[\check{\rho}_g]$ is also a group homomorphism $G_b \rightarrow \text{Aut}_{LR}(\check{\mathcal{C}})$. We show that there is an obstruction $[o_2] \in \mathcal{H}^2(G_b, \ker r)$ to $[\check{\rho}_g]$ defining a group homomorphism. Here r is the restriction map,

$$r : \text{Aut}_{LR}(\check{\mathcal{C}}) \rightarrow \text{Aut}_{LR}(\check{\mathcal{C}})|_{\mathcal{C}} = \text{Aut}_{LR}(\mathcal{C}), \quad (7)$$

which has a nontrivial kernel.

We characterize $\ker r$ in the following way. First, note that a fermion parity vortex x is called v type if $x \times \psi \neq x$.

Theorem I.3. Suppose that the modular S matrix of $\check{\mathcal{C}}$, when restricted to its block involving only v -type vortices, is block diagonal with k decoupled blocks. Then if $[\rho] \in \ker r$ and x is a fermion parity vortex, $\rho(x) = x \times \psi^{q(x)}$, where $q(x) \in \{0, 1\}$ is independent of x within a block if x is a v -type vortex.

The proof of Theorem I.3 is given in Sec. VIB 3. Theorem I.3 fully characterizes all possible ways that elements of $\ker r$ can permute the vortices. In Theorem VI.1, we fully characterize $\ker r \subset \text{Aut}_{LR}(\check{\mathcal{C}})$ in cases where permutation actions of the anyons and vortices uniquely determine the Aut groups. In Sec. VIB, we conjecture that in general $\ker r = \mathbb{Z}_2$ (see Conjecture VI.1). In Sec. VIF, we provide an explicit example of a theory where $k = 2$.

In Sec. VIB, we will define a special element $[\alpha_\psi] \in \ker r$, such that $\alpha_\psi(\phi) = \phi \times \psi$ for all v -type fermion parity vortices ϕ .

In cases where $\ker r = \mathbb{Z}_2 \simeq \{[1], [\alpha_\psi]\}$, the anomaly $[o_2] \in \mathcal{H}^2(G_b, \mathbb{Z}_2)$ can be directly related to the physical origin of $\mathcal{H}^2(G_b, \mathbb{Z}_2)$ in the classification of (3 + 1)D FSPT phases. The data $n_2 \in C^2(G_b, \mathbb{Z}_2)$ that is used in specifying an FSPT can be understood as decorating each codimension-2 trijunction of G_b domain walls $\mathbf{g}, \mathbf{h}, \mathbf{gh}$ in the (3 + 1)D bulk with a (1 + 1)D Kitaev chain if $n_2(\mathbf{g}, \mathbf{h})$ is nontrivial. We show how the anomaly $[o_2]$ arising in the problem of extending the group homomorphism ρ to $\check{\rho}$ is related to the presence of Kitaev chains on domain walls in the bulk (3 + 1)D FSPT.

Recently, [33] has shown that in the case $G_b = \mathbb{Z}_2^T$, the $\mathcal{H}^2(\mathbb{Z}_2^T, \mathbb{Z}_2)$ anomaly corresponds to an anomaly in the action of T^2 on the torus Hilbert space of the fermionic topological phase. We generalize their results and show how, for $\ker r = \mathbb{Z}_2$, the nontrivial $[o_2]$ that we find implies that the action of the symmetry operators $\check{\rho}_g$ for $\mathbf{g} \in G_b$ get extended on the torus. Letting $|\Psi\rangle_s$ be a state in the Hilbert space of the topological quantum field theory on the torus with a fixed spin structure s , we find

$$\check{\rho}_g \check{\rho}_h |\Psi\rangle_s = \check{\rho}_{gh} (\omega_2(\mathbf{g}, \mathbf{h}))^F (-1)^{\tilde{o}_2(\mathbf{g}, \mathbf{h}) \text{Arf}(s)} |\Psi\rangle_s \quad (8)$$

where \tilde{o}_2 means we are interpreting $\tilde{o}_2 \in \mathbb{Z}_2 \simeq \{0, 1\}$ instead of $o_2 \in \mathbb{Z}_2 \simeq \ker r$. In fact, we show that the element $[\alpha_\psi] \in \ker r$ discussed above allows us to in general change the symmetry action $\check{\rho}_g \rightarrow \check{\rho}_g \alpha_\psi$, which has the effect of changing the symmetry action on the torus Hilbert space by

$$\check{\rho}_g |\Psi\rangle_s \rightarrow \check{\rho}_g (-1)^{\text{Arf}(s)} |\Psi\rangle_s \quad (9)$$

As an example, we show that the semion-fermion theory, $U(1)_2 \times U(1)_{-1}$, which exists at the surface of a $v = 2$

fermionic topological superconductor in Class DIII, possesses the above $\mathcal{H}^2(G_b, \mathbb{Z}_2)$ obstruction.

We note that the $[o_2]$ anomaly defined above can, a priori, depend on which particular modular extension ν we consider. To highlight this dependence we write $[o_2^{(\nu)}]$. However, we expect that the 't Hooft anomaly of the $(2+1)$ D theory depends only on \mathcal{C} and the symmetry fractionalization data and should be independent of ν . In reviewing $(3+1)$ D FSPTs in Sec. IV B, we will explicitly define a subgroup $\Gamma^2 \subset \mathcal{H}^2(G_b, \mathbb{Z}_2)$, which leads to the natural homomorphism $q_{\Gamma^2} : \mathcal{H}^2(G_b, \mathbb{Z}_2) \rightarrow \mathcal{H}^2(G_b, \mathbb{Z}_2)/\Gamma^2$. In Sec. V E we will discuss how $q([o_2^{(\nu)}])$ is expected to determine the 't Hooft anomaly of the theory and results regarding its independence of ν .

Finally, we note that there is still a remnant of the \mathcal{H}^2 anomaly if the \mathcal{H}^1 anomaly is nontrivial. Specifically, one can always define a relative \mathcal{H}^2 anomaly for two theories \mathcal{C}_1 and \mathcal{C}_2 that possess the same \mathcal{H}^1 anomaly. This can be done by considering the theory \mathcal{C}_{12} obtained by stacking \mathcal{C}_1 and $\overline{\mathcal{C}_2}$, so that the \mathcal{H}^1 anomaly of the stacked theory vanishes, and then computing the resulting \mathcal{H}^2 anomaly of \mathcal{C}_{12} . The \mathcal{H}^2 anomaly of \mathcal{C}_{12} defines the relative \mathcal{H}^2 anomaly between \mathcal{C}_1 and \mathcal{C}_2 . This agrees with the $(3+1)$ D fermion SPT classification, where the $(3+1)$ D fermion SPTs form a torsor over \mathcal{H}^2 if the \mathcal{H}^1 piece is nontrivial.

3. Anomaly cascade: Third layer

The third layer of the anomaly cascade asks whether the symmetry fractionalization data can be extended from the anyons to the fermion parity-gauged theory.

Assume that the first and second layer obstructions vanish, so that the map $[\check{\rho}_g]$ defined in Eq. (6) satisfies the appropriate group structure.

The next task is to determine if the symmetry fractionalization class for \mathcal{C} , defined by a set of U(1) phases $\eta_a : G_b \times G_b \rightarrow \text{U}(1)$ for each anyon a , can be extended to the full modular extension $\check{\mathcal{C}}$. In [32], it was shown that there is an anomaly $[o_3] \in \mathcal{H}^3(G_b, \mathbb{Z}_2)$, which quantifies the obstruction to such an extension. The analysis of [32] was restricted to the special case where $G_f = G_b \times \mathbb{Z}_2^f$, with some additional technical assumptions on the symmetry fractionalization on \mathcal{C} . Here we provide a completely general discussion, which applies to arbitrary group extensions G_f and symmetry fractionalization.

As an example, we compute the $\mathcal{H}^3(G_b, \mathbb{Z}_2)$ obstruction for the case of the doubled semion-fermion theory $[\text{U}(1)_2 \times \text{U}(1)_{-1}]^2$, which exists at the surface of a $\nu = 4$ topological superconductor in class DIII, where $\mathbf{T}^2 = (-1)^F$ so that $G_b = \mathbb{Z}_2^T$, $G_f = \mathbb{Z}_4^{T,f}$. We show that this $[o_3]$ class is nontrivial, which matches the expectation from the bulk-boundary correspondence.

As in the \mathcal{H}^2 case, a priori there may be multiple different modular extensions $\check{\mathcal{C}}_\nu$ for which we have a valid group homomorphism $[\check{\rho}_g]$, and multiple possible choices of $[\check{\rho}_g]$. Therefore in general $[o_3]$ depends on ν and $\check{\rho}$; to highlight this dependence we can write $[o_3^{(\nu, \check{\rho})}]$.

As in the case of the \mathcal{H}^2 layer, we can define the natural homomorphism $q_{\Gamma^3} : \mathcal{H}^3(G_b, \mathbb{Z}_2) \rightarrow \mathcal{H}^3(G_b, \mathbb{Z}_2)/\Gamma^3$, and we

expect that the 't Hooft anomaly is determined by $q_{\Gamma^3}([o_3^{(\nu, \check{\rho})}])$. In particular, since we expect that the 't Hooft anomaly is determined by \mathcal{C} and its symmetry fractionalization data, we expect that $q_{\Gamma^3}([o_3^{(\nu, \check{\rho})}])$ is independent of the valid choices of $\nu, \check{\rho}$. We discuss this expectation in detail in Sec. VII D.

Just like the \mathcal{H}^2 anomaly, we can always define a relative \mathcal{H}^3 anomaly between two theories \mathcal{C}_1 and \mathcal{C}_2 with identical \mathcal{H}^1 and \mathcal{H}^2 anomalies. That is, we consider \mathcal{C}_1 and \mathcal{C}_2 to have identical \mathcal{H}^1 anomaly, and vanishing relative \mathcal{H}^2 anomaly. Then, we can consider the stacked theory \mathcal{C}_{12} , which has vanishing \mathcal{H}^1 and \mathcal{H}^2 anomaly, and for which we can define an \mathcal{H}^3 anomaly. The \mathcal{H}^3 anomaly of \mathcal{C}_{12} defines the relative \mathcal{H}^3 anomaly between \mathcal{C}_1 and \mathcal{C}_2 .

4. Anomaly cascade: Fourth layer

The fourth layer of the anomaly cascade asks whether it is possible to define a theory of symmetry defects for G_b consistent with the symmetry fractionalization data on the anyons and fermion parity vortices.

If all the preceding obstructions vanish, we can define a fermion parity gauged theory described by the modular extension $\check{\mathcal{C}}$, and a notion of symmetry fractionalization on $\check{\mathcal{C}}$. Thus we can define a fully bosonic topological phase with a symmetry fractionalization class. We can then refer to the theory of G_b -crossed modular categories, for which there is an $\mathcal{H}^4(G_b, \text{U}(1))$ obstruction to gauging G_b [1, 4, 7, 26], which is now known how to explicitly compute in general [7, 24].

One may think of the first three layers of obstructions as determining the mixed anomaly between fermion parity and G_b —they are the obstructions to gauging fermion parity while preserving G_b symmetry. If the mixed anomaly vanishes, then this fourth layer characterizes the remaining pure G_b anomaly.

In general, to define the $[o_4]$ obstruction, there was a choice of ν , $\check{\rho}$, and symmetry fractionalization data $\check{\eta}$. To highlight this dependence we write $[o_4^{(\nu, \check{\rho}, \check{\eta})}]$. As in the case of the second and third layers, we have a map $q_{\Gamma^4} : \mathcal{H}^4(G_b, \mathbb{Z}_2) \rightarrow \mathcal{H}^4(G_b, \mathbb{Z}_2)/\Gamma^4$. Physically $\Gamma^4 \subset \mathcal{H}^4(G_b, \text{U}(1))$ can be thought of as the subgroup of $(3+1)$ D boson SPTs that become trivial upon introducing fermions with G_f symmetry. We expect then that the 't Hooft anomaly of the theory is $q([o_4^{(\nu, \check{\rho}, \check{\eta})}])$, and that this is independent of the choices $\nu, \check{\rho}, \check{\eta}$, as summarized in Conjecture VIII.1. We will in particular prove independence of $q_{\Gamma^4}([o_4^{(\nu, \check{\rho}, \check{\eta})}])$ under changes of $\check{\eta}$ for fixed $\nu, \check{\rho}$.

Just as in the previous cases, even if a theory possesses a nontrivial \mathcal{H}^i anomaly with $i < 4$, we can always define a relative \mathcal{H}^4 anomaly between two theories \mathcal{C}_1 and \mathcal{C}_2 that have vanishing relative \mathcal{H}^1 , \mathcal{H}^2 , and \mathcal{H}^3 anomalies.

5. Organization of paper

The rest of this paper is structured as follows. In Sec. II, we review some basic facts about super-modular and spin modular categories and the Hilbert space of a spin modular theory on a torus. In Sec. III, we give a brief review of symmetry fractionalization in bosonic systems. In Sec. IV A we summarize symmetries and symmetry fractionalization in fermionic systems and also review in detail the classification of $(3+1)$ D FSPTs due to [30]. We then consider each level

of the anomaly cascade in order. We prove Theorem I.2 about the $\mathcal{H}^1(G_b, \mathbb{Z}_T)$ obstruction in Sec. V. Section VI is devoted to a thorough discussion of the $\mathcal{H}^2(G_b, \ker r)$ obstruction and its relation to the $\mathcal{H}^2(G_b, \mathbb{Z}_2)$ part of the 't Hooft anomaly. In Sec. VII, we give a fully general discussion of the $\mathcal{H}^3(G_b, \mathbb{Z}_2)$ obstruction. In Sec. VIII we discuss the $\mathcal{H}^4(G_b, U(1))$ obstruction, in particular its dependence on the various choices made in its definition. Section IX contains additional interesting examples, and some summary and discussion appears in Sec. X.

II. SUPER-MODULAR AND SPIN MODULAR CATEGORIES AND FERMIONIC TOPOLOGICAL PHASES OF MATTER

There are two equivalent descriptions of a fermionic topological phase of matter. One description uses a super-modular tensor category, denoted \mathcal{C} , along with a chiral central charge c_- . In this description, the super-modular tensor category determines $c_- \bmod 1/2$. A super-modular tensor category [34,37,38] is a unitary braided fusion category with exactly one nontrivial invisible particle ψ , which is a fermion and satisfies \mathbb{Z}_2 fusion rules. “Invisible” means that the double braid

$$M_{a,\psi} = +1 \quad (10)$$

for all $a \in \mathcal{C}$, “fermion” means that the topological twist $\theta_\psi = -1$, and the \mathbb{Z}_2 fusion rules means $\psi \times \psi = 1$. Physically, this description tracks the topologically nontrivial quasiparticle content of the phase. The presence of ψ is used to track the presence of a fermion that is topologically trivial in the sense that the fermion can be created or annihilated by a local fermion operator. Different fermionic phases with the same quasiparticle content are distinguished by stacking with $p + ip$ superconductors, each of which changes c_- by $1/2$.

The alternate description is via a spin modular category, denoted $\check{\mathcal{C}}$, and a chiral central charge c_- , which is determined modulo 8 by $\check{\mathcal{C}}$. A spin modular category is a UMTC together with a preferred choice of fermion ψ that has \mathbb{Z}_2 fusion rules. Physically, the spin modular category describes the phase after gauging fermion parity, that is, it describes the quasiparticle content of the phase along with the fermion parity vortices. More precisely, $\check{\mathcal{C}}$ possesses a natural \mathbb{Z}_2 grading determined by double braids with ψ :

$$\check{\mathcal{C}} = \check{\mathcal{C}}_0 \oplus \check{\mathcal{C}}_1 \quad (11)$$

$$\check{\mathcal{C}}_0 \simeq \mathcal{C} \quad (12)$$

where $M_{a,\psi} = +1$ if $a \in \check{\mathcal{C}}_0$ and $M_{a,\psi} = -1$ if $a \in \check{\mathcal{C}}_1$. Fusion respects this grading. The objects in $\check{\mathcal{C}}_1$ are physically interpreted as fermion parity vortices, or equivalently symmetry defects of fermion parity symmetry.

The sector $\check{\mathcal{C}}_1$ can be further decomposed as follows:

$$\check{\mathcal{C}}_1 = \check{\mathcal{C}}_v \oplus \check{\mathcal{C}}_\sigma \quad (13)$$

where

$$\check{\mathcal{C}}_v = \{a \in \check{\mathcal{C}}_1 \mid a \times \psi \neq a\} \quad (14)$$

$$\check{\mathcal{C}}_\sigma = \{a \in \check{\mathcal{C}}_1 \mid a \times \psi = a\}. \quad (15)$$

The two descriptions are related as follows. Every super-modular category \mathcal{C} admits [31] a minimal modular extension $\check{\mathcal{C}}$, which means that $\check{\mathcal{C}}$ contains \mathcal{C} as a subcategory, the preferred fermion of $\check{\mathcal{C}}$ is the invisible fermion of \mathcal{C} , and $\check{\mathcal{C}}$ has minimal possible total quantum dimension

$$\mathcal{D}_{\check{\mathcal{C}}}^2 = 2\mathcal{D}_{\mathcal{C}}^2. \quad (16)$$

Every super-modular category has precisely 16 distinct minimal modular extensions $\check{\mathcal{C}}_v$, for $v = 0, \dots, 15$, and whose chiral central charges modulo 8 differ by $v/2$. A fermionic topological phase is described either by (\mathcal{C}, c_-) or by $(\check{\mathcal{C}}_v, c_-)$ where the chiral central charge of $\check{\mathcal{C}}_v$ is equal to $c_- \bmod 8$.

We will make use of the following fact [37] about the topological S matrix of a minimal modular extension $\check{\mathcal{C}}$:

$$S_{x,y} = \begin{cases} S_{x \times \psi, y} & y \in \check{\mathcal{C}}_0 \\ -S_{x \times \psi, y} & y \in \check{\mathcal{C}}_1 \end{cases}. \quad (17)$$

Note that the second line implies that $S_{\sigma,y} = 0$ for $\sigma \in \check{\mathcal{C}}_\sigma$, $y \in \check{\mathcal{C}}_1$.

As we will see now, the full spin modular category, not just the super-modular tensor category, is required to describe the fermionic system on nontrivial surfaces and with arbitrary spin structures.

A. Torus degeneracy and spin structure

The spin modular category determines the Hilbert space and the action of the mapping class group of the fermionic topological phase of matter on a topologically nontrivial surface.

Consider the (spatial) torus T^2 with nontrivial cycles α and β . Let us label the states of the spin modular category as $|a\rangle_\alpha$, for $a \in \check{\mathcal{C}}$ an anyon. This means that the topological charge as measured through the loop α is a . Below for ease of notation we drop the subscript α and keep it implicit in the definition of the state $|a\rangle$.

Now suppose we have a choice of spin structure (μ_α, ν_β) where $\mu = 0$ corresponds to Neveu-Schwarz (antiperiodic) boundary conditions on the loop α and $\mu = 1$ corresponds to Ramond (periodic) boundary conditions on α , with ν_β similar for the β loop.¹ We drop the α and β labels in what follows. Let $\mathcal{H}_{\mu,\nu}$ denote the Hilbert space of the fermionic topological phase on the torus with the chosen spin structure, $|\Psi\rangle_{\mu,\nu} \in \mathcal{H}_{\mu,\nu}$, and let $W_x(\gamma)$ denote the Wilson loop of a particle x around the loop γ .

The defining distinction between the different sectors (μ, ν) is in the fermion boundary conditions, which defines the eigenvalue of the fermion Wilson loop W_ψ :

$$\begin{aligned} W_\psi(\alpha)|\Psi\rangle_{\mu,\nu} &= (-1)^\mu |\Psi\rangle_{\mu,\nu} \\ W_\psi(\beta)|\Psi\rangle_{\mu,\nu} &= (-1)^\nu |\Psi\rangle_{\mu,\nu}. \end{aligned} \quad (18)$$

Note that $W_\psi(\alpha)$ has -1 eigenvalue for the sector with periodic ($\mu = 1$) boundary conditions along the α cycle. This is

¹Our convention is slightly unusual because the modular T transformation does not act linearly on the μ and ν indices. However, the formulas relevant to us like Eq. (18) are more natural.

because periodic boundary conditions occur when there is a fermion parity vortex threading the conjugate cycle, as can be derived by studying the modular matrices carefully. The -1 then arises due to the mutual statistics between ψ and v .

A basis of states for the fermionic topological phase on a torus is as follows (see e.g., [33] for a recent discussion):

$$\begin{aligned} \mathcal{H}_{0,0} : |a\rangle_{00} &= \frac{1}{\sqrt{2}}(|a\rangle + |a \times \psi\rangle) \\ \mathcal{H}_{1,0} : &\begin{cases} |v\rangle_{10} = \frac{1}{\sqrt{2}}(|v\rangle + |v \times \psi\rangle) \\ |\sigma\rangle_{10} = |\sigma\rangle \end{cases} \\ \mathcal{H}_{0,1} : |a\rangle_{01} &= \frac{1}{\sqrt{2}}(|a\rangle - |a \times \psi\rangle) \\ \mathcal{H}_{1,1} : &\begin{cases} |v\rangle_{11} = \frac{1}{\sqrt{2}}(|v\rangle - |v \times \psi\rangle) \\ |\sigma\rangle_{11} = |\sigma; \psi\rangle \end{cases}, \end{aligned} \quad (19)$$

where $a \in \check{\mathcal{C}}_0$, $v \in \check{\mathcal{C}}_v$, and $\sigma \in \check{\mathcal{C}}_\sigma$. Furthermore, $|\sigma; \psi\rangle$ denotes the state on a torus with a puncture labeled ψ .

These states on the torus can be built by gluing together states on the 3-punctured sphere, i.e., splitting spaces V_c^{ab} and their duals V_{ab}^c . Eq. (19) can be reexpressed in this language as

$$\begin{aligned} \mathcal{H}_{00} &= \left(\bigoplus_{a \in \check{\mathcal{C}}_0} V_1^{a\bar{a}} \otimes V_{a\bar{a}}^1 \right) \Big|_{\text{symm}} \\ \mathcal{H}_{01} &= \left(\bigoplus_{a \in \check{\mathcal{C}}_0} V_1^{a\bar{a}} \otimes V_{a\bar{a}}^1 \right) \Big|_{\text{anti-symm}} \\ \mathcal{H}_{1,0} &= \left(\bigoplus_{v \in \check{\mathcal{C}}_v} V_1^{v\bar{v}} \otimes V_{v\bar{v}}^1 \right) \Big|_{\text{symm}} \oplus \bigoplus_{\sigma \in \check{\mathcal{C}}_\sigma} (V_1^{\sigma\bar{\sigma}} \otimes V_{\sigma\bar{\sigma}}^1) \\ \mathcal{H}_{1,1} &= \left(\bigoplus_{v \in \check{\mathcal{C}}_v} V_1^{v\bar{v}} \otimes V_{v\bar{v}}^1 \right) \Big|_{\text{anti-symm}} \oplus \bigoplus_{\sigma \in \check{\mathcal{C}}_\sigma} (V_1^{\sigma\bar{\sigma}} \otimes V_{\sigma\bar{\sigma}}^\psi) \end{aligned} \quad (20)$$

The “(anti)-symm” notation means we restrict to the subspace that consists of (anti)-symmetric sums of states in the a and $a \times \psi$ sectors.

III. REVIEW OF SYMMETRY FRACTIONALIZATION IN BOSONIC SYSTEMS

We briefly review the formalism for symmetry fractionalization in bosonic systems.

Consider a UMTC \mathcal{B} with global symmetry group G . The basic data required to define symmetry fractionalization is the following. First, we define a group homomorphism

$$[\rho_g] : G \rightarrow \text{Aut}(\mathcal{B}) \quad (21)$$

where $\text{Aut}(\mathcal{B})$ is the group of braided autoequivalences of \mathcal{B} , modulo a set of gauge equivalences called natural isomorphisms.² A braided autoequivalence, or autoequivalence for

²We define natural isomorphisms using Eq. (22). There exists a more abstract mathematical definition, and it is unclear if this def-

short, is a map from \mathcal{B} to itself that preserves the data of the theory up to a gauge transformation. We will also use the term “autoequivalence” to refer to braided anti-autoequivalences, which, up to a gauge transformation, complex conjugate the data of the theory. A natural isomorphism Υ is an autoequivalence, which acts on fusion vertices as

$$\Upsilon(|a, b; c; \mu\rangle) = \frac{\gamma_a \gamma_b}{\gamma_c} |a, b; c; \mu\rangle \quad (22)$$

where $\gamma_a \in \text{U}(1)$. Natural isomorphisms have a redundancy

$$\gamma_a \rightarrow \gamma_a \zeta_a \quad (23)$$

where $\zeta_a \in \text{U}(1)$ obeys the fusion rules in the sense that $\zeta_a \zeta_b = \zeta_c$ whenever $N_{ab}^c > 0$. A representative ρ_g of the equivalence class $[\rho_g]$ determines a permutation of the anyons $a \rightarrow {}^g a$ and a set of unitary matrices $U_g(a, b; c)$ as follows:

$$\rho_g(|a, b; c; \mu\rangle) = \sum_v U_g({}^g a, {}^g b; {}^g c)_{\mu v} |{}^g a, {}^g b; {}^g c; v\rangle. \quad (24)$$

The statement that ρ_g is an autoequivalence means that the F and R symbols are preserved (up to complex conjugation) according to the following consistency conditions:

$$\begin{aligned} &([F_d^{abc}]_{(e,\alpha,\beta),(f,\mu,\nu)})^{\sigma(g)} \\ &= \sum_{\alpha',\beta',\mu',\nu'} U_g({}^g a, {}^g b; {}^g e)_{\alpha\alpha'} U_g({}^g e, {}^g c; {}^g d)_{\beta\beta'} \\ &\times [F_{g\bar{d}}^{{}^g a {}^g b {}^g c}]_{({}^g e,\alpha',\beta'),({}^g f,\mu',\nu')} \\ &\times (U_g({}^g b, {}^g c; {}^g f)_{\mu'\mu}^{-1} (U_g({}^g a, {}^g f; {}^g d)_{\nu'\nu}^{-1}, \quad (25) \\ &([R_c^{ab}]_{\mu\nu})^{\sigma(g)} = U_g({}^g b, {}^g a; {}^g c)_{\mu\mu'} (R_{g\bar{c}}^{{}^g a {}^g b})_{\mu'\nu'} \\ &\times [U_g({}^g a, {}^g b; {}^g c)_{\nu'\nu}^{-1}]. \end{aligned}$$

Here

$$\sigma(g) = \begin{cases} 1 & g \text{ unitary} \\ * & g \text{ antiunitary} \end{cases}. \quad (27)$$

We will also use

$$s_1(g) = \begin{cases} 0 & g \text{ unitary} \\ 1 & g \text{ antiunitary} \end{cases} \quad (28)$$

The map s_1 is a group homomorphism, i.e., $s_1 \in Z^1(G_b, \mathbb{Z}_2)$. The maps ρ_g define the natural isomorphisms

$$\kappa_{g,h} = \rho_{gh} \rho_h^{-1} \rho_g^{-1}, \quad (29)$$

which have actions on fusion vertices given by

$$\kappa_{g,h} |a, b; c; \mu\rangle = \kappa_{g,h} (a, b; c) |a, b; c; \mu\rangle. \quad (30)$$

ition is equivalent to ours. In considering equivalence classes of braided autoequivalences, we will restrict our attention to natural isomorphisms in our definition, since this appears to naturally describe SETs [1].

Here

$$\begin{aligned}\kappa_{\mathbf{g}, \mathbf{h}}(a, b; c) &= \frac{\beta_a(\mathbf{g}, \mathbf{h})\beta_b(\mathbf{g}, \mathbf{h})}{\beta_c(\mathbf{g}, \mathbf{h})} \\ &= U_{\mathbf{g}}^{-1}(a, b; c)U_{\mathbf{h}}(\bar{\mathbf{g}}a, \bar{\mathbf{g}}b; \bar{\mathbf{g}}c)^{-\sigma(\mathbf{g})}U_{\mathbf{gh}}(a, b; c)\end{aligned}\quad (31)$$

and $\beta_a(\mathbf{g}, \mathbf{h})$ are phases defining $\kappa_{\mathbf{g}, \mathbf{h}}$ as a natural isomorphism. The β_a define the phases

$$\Omega_a(\mathbf{g}, \mathbf{h}, \mathbf{k}) = \frac{\beta_{\bar{\mathbf{g}}a}^{\sigma(\mathbf{g})}(\mathbf{h}, \mathbf{k})\beta_a(\mathbf{g}, \mathbf{h}\mathbf{k})}{\beta_a(\mathbf{g}, \mathbf{h})\beta_a(\mathbf{g}\mathbf{h}, \mathbf{k})}, \quad (32)$$

which can be shown to obey the fusion rules, in the sense

$$\Omega_a\Omega_b = \Omega_c \quad \text{whenever} \quad N_{ab}^c > 0. \quad (33)$$

These phases define, for modular \mathcal{B} , an obstruction $[\mathfrak{G}] \in \mathcal{H}^3(G, \mathcal{A})$ to localizing G on the anyons, where $\mathcal{A} \subset \mathcal{B}$ is the group of Abelian anyons of the theory; see standard references, e.g., [1], for further details. If said obstruction vanishes, then one can define symmetry fractionalization on \mathcal{B} , which amounts to a choice of phases $\eta_a(\mathbf{g}, \mathbf{h}) \in \text{U}(1)$, which satisfy the following consistency conditions:

$$\eta_{\bar{\mathbf{g}}a}(\mathbf{h}, \mathbf{k})^{\sigma(\mathbf{g})}\eta_a(\mathbf{g}, \mathbf{h}\mathbf{k}) = \eta_a(\mathbf{g}, \mathbf{h})\eta_a(\mathbf{g}\mathbf{h}, \mathbf{k}), \quad (34)$$

$$\begin{aligned}\frac{\eta_c(\mathbf{k}, \mathbf{l})}{\eta_a(\mathbf{k}, \mathbf{l})\eta_b(\mathbf{k}, \mathbf{l})}U_{\mathbf{kl}}(a, b; c)_{\mu\nu} \\ = \sum_{\lambda} U_{\mathbf{l}}(\bar{\mathbf{k}}a, \bar{\mathbf{k}}b; \bar{\mathbf{k}}c)_{\mu\lambda}U_{\mathbf{k}}(a, b; c)_{\lambda\nu}.\end{aligned}\quad (35)$$

Symmetry fractionalization can equivalently be specified by a set of phases $\omega_a(\mathbf{g}, \mathbf{h})$, which obey the fusion rules and satisfy

$$\Omega_a(\mathbf{g}, \mathbf{h}, \mathbf{k}) = \frac{\omega_{\bar{\mathbf{g}}a}^{\sigma(\mathbf{g})}(\mathbf{h}, \mathbf{k})\omega_a(\mathbf{g}, \mathbf{h}\mathbf{k})}{\omega_a(\mathbf{g}, \mathbf{h})\omega_a(\mathbf{g}\mathbf{h}, \mathbf{k})} \equiv (d\omega)_a(\mathbf{g}, \mathbf{h}, \mathbf{k}). \quad (36)$$

The relationship between the η_a and ω_a descriptions is

$$\eta_a(\mathbf{g}, \mathbf{h}) = \frac{\beta_a(\mathbf{g}, \mathbf{h})}{\omega_a(\mathbf{g}, \mathbf{h})}. \quad (37)$$

In all of the above, we have taken a fixed representative ρ_g of the class $[\rho_g]$. If ρ_g is modified by a natural isomorphism given by the phases $\gamma_a(\mathbf{g})$, then we obtain gauge-equivalent data

$$U'_{\mathbf{g}}(a, b; c)_{\mu\nu} = \frac{\gamma_a(\mathbf{g})\gamma_b(\mathbf{g})}{\gamma_c(\mathbf{g})}U_{\mathbf{g}}(a, b; c)_{\mu\nu} \quad (38)$$

$$\eta'_a(\mathbf{g}, \mathbf{h}) = \frac{\gamma_a(\mathbf{g}\mathbf{h})}{[\gamma_{\bar{\mathbf{g}}a}(\mathbf{h})]^{\sigma(\mathbf{g})}\gamma_a(\mathbf{g})}\eta_a(\mathbf{g}, \mathbf{h}). \quad (39)$$

Separately, there is gauge freedom

$$\begin{aligned}\beta_a(\mathbf{g}, \mathbf{h}) &\rightarrow \beta_a(\mathbf{g}, \mathbf{h})\nu_a(\mathbf{g}, \mathbf{h}) \\ \Omega_a(\mathbf{g}, \mathbf{h}, \mathbf{k}) &\rightarrow \Omega_a(\mathbf{g}, \mathbf{h}, \mathbf{k})(d\nu)_a(\mathbf{g}, \mathbf{h}, \mathbf{k}) \\ \omega_a(\mathbf{g}, \mathbf{h}) &\rightarrow \omega_a(\mathbf{g}, \mathbf{h})\nu_a(\mathbf{g}, \mathbf{h})\end{aligned}\quad (40)$$

where $\nu_a(\mathbf{g}, \mathbf{h})$ is a phase obeying the fusion rules.

For a fixed map $[\rho_g]$, the set of symmetry fractionalization patterns form a torsor over $\mathcal{H}^2(G, \mathcal{A})$. Specifically, given consistent symmetry fractionalization data $\eta_a(\mathbf{g}, \mathbf{h})$ and an

element $\mathbf{t}(\mathbf{g}, \mathbf{h}) \in Z^2(G, \mathcal{A})$, then one obtains a new symmetry fractionalization pattern

$$\widehat{\eta}_a(\mathbf{g}, \mathbf{h}) = M_{a, \mathbf{t}(\mathbf{g}, \mathbf{h})}\eta_a(\mathbf{g}, \mathbf{h}). \quad (41)$$

One can check that up to gauge transformations, $\widehat{\eta}$ depends only on the cohomology class $[\mathbf{t}] \in \mathcal{H}^2(G, \mathcal{A})$ and that different cohomology classes produce gauge-inequivalent symmetry fractionalization patterns.

IV. SYMMETRIES IN FERMIONIC TOPOLOGICAL PHASES

In this section, we briefly review fermionic symmetries, the classification of (3 + 1)D fermionic SPTs, and the results of [35] on fermionic symmetry fractionalization in fermionic topological phases, and also define a map $\breve{\Upsilon}_\psi$ in Eq. (73), which will be useful later.³

We assume that we are describing a system whose Hilbert space decomposes into a tensor product of local Hilbert spaces, which include fermionic degrees of freedom. Further, we assume the dynamics of the system are given by a local Hamiltonian with an energy gap such that the system is in the fermionic topological phase associated to \mathcal{C} . In order for the formalism to describe anomalous fermionic SETs, we use the term “symmetry fractionalization of a fermionic topological phase” to refer only to fractionalization data on the super-modular category \mathcal{C} .

A. Fermionic symmetries

Fermionic systems always have a special symmetry, fermion parity symmetry $(-1)^F$, which generates a central \mathbb{Z}_2 subgroup \mathbb{Z}_2^f of the full symmetry group G_f . Define the “bosonic” symmetry group $G_b = G_f/\mathbb{Z}_2^f$; then the symmetry generators restricted to their action on bosonic operators form a representation of G_b . An alternate characterization of G_f is as a \mathbb{Z}_2^f central extension of G_b via the short exact sequence

$$1 \rightarrow \mathbb{Z}_2^f \rightarrow G_f \rightarrow G_b \rightarrow 1 \quad (42)$$

and a cocycle $\omega_2 \in Z^2(G_b, \mathbb{Z}_2)$.

We will reserve the notation ω_2 for a cocycle valued in $\{\pm 1\} \simeq \mathbb{Z}_2$ and use the notation $\tilde{\omega}_2$ for the $\{0, 1\}$ -valued additive parametrization of ω_2 , i.e., define

$$\omega_2(\mathbf{g}, \mathbf{h}) = (-1)^{\tilde{\omega}_2(\mathbf{g}, \mathbf{h})}. \quad (43)$$

Physically, we can write a set of operators $R_{\mathbf{g}}$ for $\mathbf{g} \in G_b$, which implement the action of G_b on the microscopic Hilbert space. Since the symmetry group on the full fermionic Hilbert space is actually G_f , these operators multiply projectively,

$$R_{\mathbf{g}}R_{\mathbf{h}} = (\omega_2(\mathbf{g}, \mathbf{h}))^F R_{\mathbf{gh}}. \quad (44)$$

The cocycle $\omega_2 \in Z^2(G_b, \mathbb{Z}_2)$ determines a decomposition $G_f = G_b \times \mathbb{Z}_2^f$ as sets, but all choices of cocycle representative in the same cohomology class $[\omega_2]$ lead to isomorphic

³The results on fermionic symmetry fractionalization reviewed here also appeared in [36], the first version of which appeared on the arXiv at the same time as both the first version of this paper and [35].

groups G_f . Physically, modifying $\omega_2 \rightarrow \omega_2 \times d\phi$ for $\phi \in C^1(G_b, \mathbb{Z}_2)$ changes

$$R_g \rightarrow (\phi(g))^F R_g, \quad (45)$$

where we are taking $\phi \in \{\pm 1\}$. Such a transformation changes the physical meaning of R_g ; for example, if $G_b = \mathbb{Z}_2^T$, the time-reversal operator T is physically distinct from $(-1)^F T$, and these operators should not be interchanged. Therefore, a microscopic realization of G_f symmetry will in general specify a cocycle representative ω_2 , not just its cohomology class $[\omega_2]$.

B. General classification of (3 + 1)D fermionic SPTs and 't Hooft anomalies in (2 + 1)D fermionic systems

The group of (3 + 1)D fermionic SPTs, which also defines 't Hooft anomalies for (2 + 1)D fermionic topological phases, has been classified in recent years through a variety of approaches. One general approach is through the cobordism classification [27,28], where (3 + 1)D fermionic SPTs are classified using bordism groups of 4-manifolds equipped with G_b gauge fields and a generalized spin structure.

An alternative general classification method was recently provided by Wang and Gu [30], based on decorating symmetry defects of varying codimension with lower dimensional fermionic invertible topological phases. Below we summarize the Wang-Gu consistency equations and equivalence relations. We note that our treatment of the equivalence relations below differs slightly from Wang-Gu, as we will explain.

The calculation of the bordism groups mentioned above can be performed by various spectral sequence methods. These methods typically give the resulting FSPT classification in terms of a group extension involving subgroups of $\mathcal{H}^1(G_b, \mathbb{Z}_T)$, $\mathcal{H}^2(G_b, \mathbb{Z}_2)$, $\mathcal{H}^3(G_b, \mathbb{Z}_2)$, and $\mathcal{H}^4(G_b, U(1))$. The Wang-Gu classification provides a partial solution to the above spectral sequence computation for general groups G_b .⁴

According to Wang and Gu, (3 + 1)D fermionic SPTs can be specified by four layers of data,

$$(n_1, n_2, n_3, v_4) \in Z^1(G_b, \mathbb{Z}_T) \times C^2(G_b, \mathbb{Z}_2) \times C^3(G_b, \mathbb{Z}_2) \times C^4(G_b, U(1)). \quad (46)$$

We will refer to the subscript of n_i and v_4 as the layer index.

This set of data is subject to consistency conditions:

$$\begin{aligned} dn_2 &= \tilde{\omega}_2 \cup n_1 + s_1 \cup n_1 \cup n_1, \\ dn_3 &= \tilde{\omega}_2 \cup n_2 + n_2 \cup n_2 + s_1 \cup (n_2 \cup_1 n_2), \\ dv_4 &= \mathcal{O}_5[n_3]. \end{aligned} \quad (47)$$

with s_1 defined in Eq. (28). The precise formula for \mathcal{O}_5 is unimportant for our purposes and can be found in [30]. Physically, the different layers of data correspond to decorating defects of various codimension with lower-dimensional invertible fermionic topological phases; for example, n_1 corresponds to decorating time-reversal domain walls with a

⁴It is a partial solution because Wang-Gu do not derive the group multiplication law for stacking FSPT phases, and thus do not fully solve the group extension problem.

$c_- = 1/2$ invertible fermionic phase (e.g., a $p_x + ip_y$ superconductor); n_2 corresponds to decorating codimension-2 junctions with Kitaev-Majorana chains; and n_3 corresponds to decorating codimension-3 junctions with fermions. The main fact about these equations that is of relevance to us in this paper is that data of the i th layer is an i -cocycle if the data of the lower layers vanish. That is, if $n_1 = 0$, then $n_2 \in Z^2(G_b, \mathbb{Z}_2)$; if $n_1, n_2 = 0$, then $n_3 \in Z^3(G_b, \mathbb{Z}_2)$, and if $n_1, n_2, n_3 = 0$, then $v_4 \in Z^4(G_b, U(1))$.

In addition to the above consistency equations, the data above is also subject to a number of equivalence relations. Each layer can change by a coboundary, and also an additional equivalence, as we explain.⁵

1. 1st layer equivalence

We can change n_1 by a coboundary:

$$(n_1, n_2, n_3, v_4) \simeq (n_1 + db_0, n'_2, n'_3, v'_4), \quad (48)$$

where $b_0 \in C^0(G_b, \mathbb{Z}_T)$. Note that when n_1 changes by a coboundary, the higher layer data may in principle also change.

If we ignore the higher layer data, then, we see that each (3 + 1)D FSPT determines an element

$$[n_1] \in \mathcal{H}^1(G_b, \mathbb{Z}_T). \quad (49)$$

2. 2nd layer equivalence and Γ^2

There are the following 2nd layer equivalences:

$$(n_1, n_2, n_3, v_4) \simeq (n_1, n_2 + \tilde{\omega}_2, n'_3, v'_4) \quad (50)$$

$$\simeq (n_1, n_2 + db_1, n''_3, v''_4) \text{ if } G_b \text{ unitary}, \quad (51)$$

where $b_1 \in C^1(G_b, \mathbb{Z}_2)$ and recalling $\tilde{\omega}_2 = (-1)^{\tilde{\omega}_2}$. Note that under changing the n_2 , the higher level data n_3 and v_4 can change to some other consistent data n'_3, v'_4 or n''_3, v''_4 , whose precise form has not been computed and is not relevant for our purposes. It is useful to denote

$$\Gamma^2 = \begin{cases} \{1, [\omega_2]\} & G_b \text{ unitary} \\ \{1\} & G_b \text{ contains antiunitary symmetries} \end{cases} \quad (52)$$

where $\Gamma^2 \subset \mathcal{H}^2(G_b, \mathbb{Z}_2)$. We also define the group homomorphism

$$q_{\Gamma^2} : \mathcal{H}^2(G_b, \mathbb{Z}_2) \rightarrow \mathcal{H}^2(G_b, \mathbb{Z}_2)/\Gamma^2, \quad (53)$$

which we will use later.

The implication of the above equivalence is that if $n_1 = 0$ and we forget about n_3, v_4 , we can define an equivalence class

$$[n_2] \in \mathcal{H}^2(G_b, \mathbb{Z}_2)/\Gamma^2. \quad (54)$$

In other words, any (3 + 1)D FSPT with $n_1 = 0$ defines a class $[n_2] \in \mathcal{H}^2(G_b, \mathbb{Z}_2)/\Gamma^2$.

⁵We note that in the equivalences summarized here for a given layer, we allow the possibility for the higher layer data to change as well. The equivalence relations of Ref. [30] imply that the higher layer data does not change under these equivalences, however Ref. [39] found that in (2 + 1)D the higher layer data does change under equivalences of a given layer. We expect a similar phenomenon to generally occur in (3 + 1)D as well.

3. 3rd layer equivalence and Γ^3

There are the following 3rd layer equivalences:

$$\begin{aligned} (n_1, n_2, n_3, v_4) &\simeq (n_1, n_2, n_3 + \chi_3, v'_4) \\ &\simeq (n_1, n_2, n_3 + db_2, v''_4), \end{aligned} \quad (55)$$

where $b_2 \in C^2(G_b, \mathbb{Z}_2)$. Here $\chi_3 \in Z^3(G_b, \mathbb{Z}_2)$ consists of 3-cocycles that satisfy

$$\chi_3 = \tilde{\omega}_2 \cup \lambda_1 + s_1 \cup \lambda_1 \cup \lambda_1 + \lfloor \lambda_0/2 \rfloor (\tilde{\omega}_2 \cup_1 \tilde{\omega}_2) \pmod{2}, \quad (56)$$

for any choice of $\lambda_1 \in Z^1(G_b, \mathbb{Z}_2)$ and $\lambda_0 \in \mathbb{Z}$.

Formally, we can define the group $\Gamma^3 \subset \mathcal{H}^3(G_b, \mathbb{Z}_2)$, where all representative 3-cocycles have the form χ_3 given in Eq. (56). It is useful to then define the group homomorphism

$$q_{\Gamma^3} : \mathcal{H}^3(G_b, \mathbb{Z}_2) \rightarrow \mathcal{H}^3(G_b, \mathbb{Z}_2)/\Gamma^3. \quad (57)$$

Similar to the case of the 2nd layer, the implication of the above equivalence is that if $n_1, n_2 = 0$ and we ignore v_4 , then we can define an equivalence class

$$[n_3] \in \mathcal{H}^3(G_b, \mathbb{Z}_2)/\Gamma^3. \quad (58)$$

In other words, any (3 + 1)D FSPT with $n_1, n_2 = 0$ defines a class $[n_3] \in \mathcal{H}^3(G_b, \mathbb{Z}_2)/\Gamma^3$.

4. 4th layer equivalence and Γ^4

There exist the following 4th layer equivalences:

$$(n_1, n_2, n_3, v_4) \simeq (n_1, n_2, n_3, v_4 \chi_4) \simeq (n_1, n_2, n_3, v_4 d \epsilon_3) \quad (59)$$

for $\epsilon_3 \in C^3(G_b, \mathbb{Z}_2)$. Here χ_4 is any 4-cocycle in a group Γ^4 with a rather involved definition that can be found in [30] but is unimportant for our present purposes. It will suffice to know that Γ^4 contains a subgroup of the form

$$\{[\lambda_2] \cup [\lambda_2] + [\omega_2] \cup [\lambda_2] \mid [\lambda_2] \in \mathcal{H}^2(G_b, \mathbb{Z}_2)\} \subset \Gamma^4. \quad (60)$$

We define the group homomorphism

$$q_{\Gamma^4} : \mathcal{H}^4(G_b, \mathrm{U}(1)) \rightarrow \mathcal{H}^4(G_b, \mathrm{U}(1))/\Gamma^4. \quad (61)$$

Physically, Γ^4 is the group of bosonic (3 + 1)D SPTs with G_b symmetry that are trivial when viewed as a fermionic SPT with G_f symmetry. Equivalently, Γ^4 is the group of “anomalous” fermionic (2 + 1)D SPTs; these have the property that the surface of a bosonic (3 + 1)D SPT characterized by a 4-cocycle in Γ^4 can have a topologically trivial gapped symmetric gapped (2 + 1)D surface, if fermions transforming under G_f symmetry are introduced to the surface. We see that any (3 + 1)D FSPT for which $n_1, n_2, n_3 = 0$ defines an element

$$[v_4] \in \mathcal{H}^4(G_b, \mathrm{U}(1))/\Gamma^4. \quad (62)$$

C. Fermionic symmetry fractionalization

The first step of defining G_f symmetry fractionalization is to assign an autoequivalence $[\rho_g]$ of \mathcal{C} to each element $g \in G_b$. Topological autoequivalences are well defined for any BFC regardless of modularity, so autoequivalences of a supermodular category \mathcal{C} are also well defined. This assignment defines the U symbols as given in Eq. (24). There is, however,

a physical constraint on the choice of autoequivalence. The symmetry operator R_g is defined on the physical, microscopic Hilbert space, and we will be seeking to localize R_g . Then if $\gamma_{i,r}$ is a basis of (Majorana) fermion operators at position r , the Hilbert space defines matrices $\tilde{U}_{ij}(\mathbf{g}, \mathbf{r})$ such that

$$R_g \gamma_{i,r} R_g^{-1} = \sum_j \tilde{U}_{ij}(\mathbf{g}, \mathbf{r}) \gamma_{j,r}, \quad (63)$$

where i and j label elements of the basis of fermionic operators.

As shown in [35], compatibility of a representative autoequivalence ρ_g of \mathcal{C} with Eq. (63) constrains

$$\rho_g(|\psi, \psi; 1\rangle) = |\psi, \psi; 1\rangle, \quad (64)$$

or equivalently, for all $\mathbf{g} \in G_b$,

$$U_{\mathbf{g}}(\psi, \psi; 1) = +1. \quad (65)$$

In the bosonic case, the topological autoequivalence may be redefined by a natural isomorphism of the form Eq. (22). Clearly we have less freedom in the fermionic case; maintaining Eq. (65) requires natural isomorphisms to have $\gamma_{\psi} \in \{\pm 1\}$. In fact, as shown in [35], modifying R_g with a natural isomorphism with $\gamma_{\psi} = -1$ amounts to redefining $R_g \rightarrow (-1)^F R_g$, which, as discussed in Sec. IV A, is a physical change to the system and not a gauge redundancy. Therefore, the redundancy in topological autoequivalences for fermionic symmetries is not given by arbitrary natural isomorphisms, but instead by “locality-respecting” natural isomorphisms, which have $\gamma_{\psi} = +1$. We denote the group of topological autoequivalences of \mathcal{C} , which obey Eq. (65), modulo locality-respecting natural isomorphisms, as $\mathrm{Aut}_{LR}(\mathcal{C})$.⁶

As such, fermionic symmetry actions are specified by a map

$$[\rho_g] : G_b \rightarrow \mathrm{Aut}_{LR}(\mathcal{C}), \quad (66)$$

which obeys

$$[\kappa_{\mathbf{g}, \mathbf{h}}][\rho_{\mathbf{g}}][\rho_{\mathbf{h}}] = [\rho_{\mathbf{gh}}] \quad (67)$$

for some (not necessarily locality-respecting) natural isomorphism $[\kappa_{\mathbf{g}, \mathbf{h}}]$. If $\mathrm{Aut}_{LR}(\mathcal{C})$ is isomorphic to $\mathrm{Aut}(\mathcal{C})$, then $[\kappa] = [\mathrm{Id}]$, where Id is the identity map, and $[\rho_g]$ is a group homomorphism.

There is some subtlety in demanding that locality-respecting natural isomorphisms have $\gamma_{\psi} = +1$ because of the redundancy Eq. (23) in natural isomorphisms. Define an Abelian group $K(\mathcal{C})$ consisting of maps ζ_a that obey the fusion rules, that is,

$$\begin{aligned} K(\mathcal{C}) = \{ &\zeta : \text{anyon labels} \rightarrow \mathrm{U}(1) \mid \zeta_a \zeta_b \\ &= \zeta_c \text{ whenever } N_{ab}^c > 0 \}. \end{aligned} \quad (68)$$

Clearly $\zeta_{\psi} \in \{\pm 1\}$, which provides a natural \mathbb{Z}_2 grading on $K(\mathcal{C})$:

$$K(\mathcal{C}) = K_+(\mathcal{C}) \oplus K_-(\mathcal{C}) \quad (69)$$

⁶It is not hard to show that every element of $\mathrm{Aut}(\mathcal{C})$ has a representative, which obeys Eq. (65). Accordingly, one can check that, depending on details, which will be discussed shortly, as a group $\mathrm{Aut}_{LR}(\mathcal{C})$ is either isomorphic to $\mathrm{Aut}(\mathcal{C})$ or double-covers $\mathrm{Aut}(\mathcal{C})$.

where $K_{\pm}(\mathcal{C})$ consists of maps with $\zeta_{\psi} = \pm 1$. It was proven in [35] that, given any minimal modular extension $\check{\mathcal{C}}$ of \mathcal{C} , every element ζ_a of $K(\mathcal{C})$ can be written

$$\zeta_a = M_{a,x} \quad (70)$$

for some $x \in \check{\mathcal{C}}$, and in particular, $x \in \mathcal{A}$ if $\zeta \in K_+(\mathcal{C})$. Hence

$$K_+(\mathcal{C}) \simeq \mathcal{A}/\{1, \psi\}, \quad (71)$$

where $\mathcal{A} \subset \mathcal{C}$ is the group of Abelian anyons of \mathcal{C} and $\{1, \psi\}$ is the \mathbb{Z}_2 subgroup generated by the transparent fermion ψ . If $K_-(\mathcal{C})$ is nonempty, then every natural isomorphism, which preserves Eq. (65) is equivalent to a locality-respecting natural isomorphism. There is a canonical natural isomorphism Υ_{ψ} of the same form as Eq. (22) with

$$\Upsilon_{\psi}(|a, b; c\rangle) = \frac{\gamma_a \gamma_b}{\gamma_c} |a, b; c\rangle$$

$$\text{with } \gamma_a = \begin{cases} -1 & a = \psi \\ 1 & \text{any other } a \in \mathcal{C} \end{cases} \quad (72)$$

We immediately see that Υ_{ψ} respects locality if and only if $K_-(\mathcal{C})$ is nonempty. If Υ_{ψ} does not respect locality, then $[\Upsilon_{\psi}] \neq [1]$ as elements of $\text{Aut}_{LR}(\mathcal{C})$.

Many of the results and casework in this paper depend on whether or not Υ_{ψ} respects locality, so it is useful to summarize some characterizations from [35] of when Υ_{ψ} respects locality. The following are equivalent:

- (1) Υ_{ψ} respects locality
- (2) $K(\mathcal{C})/K_+(\mathcal{C}) \simeq \mathbb{Z}_2$ (which means $K_-(\mathcal{C})$ is nonempty)
- (3) As groups, $\text{Aut}(\mathcal{C}) \simeq \text{Aut}_{LR}(\mathcal{C})$
- (4) There exists a minimal modular extension of \mathcal{C} that contains an Abelian fermion parity vortex
- (5) There exists a set of phases ζ_a that obey the fusion rules and have $\zeta_{\psi} = -1$.

Conversely, the following are also equivalent:

- (1) Υ_{ψ} violates locality
- (2) $K(\mathcal{C}) = K_+(\mathcal{C}) \simeq \mathcal{A}/\{1, \psi\}$
- (3) As groups, $\text{Aut}_{LR}(\mathcal{C})/\mathbb{Z}_2 \simeq \text{Aut}(\mathcal{C})$
- (4) Any set of phases ζ_a that obey the fusion rules must have $\zeta_{\psi} = +1$.

Another useful fact is that if Υ_{ψ} respects locality, then exactly half of the minimal modular extensions $\check{\mathcal{C}}$ contain only v -type vortices and half contain only σ -type vortices. Also, if elements of $\text{Aut}(\mathcal{C})$ are uniquely determined by their permutation action on the anyons (as is true in many theories of physical interest), then the same holds for $\text{Aut}_{LR}(\mathcal{C})$ if and only if Υ_{ψ} respects locality. One can also define a natural isomorphism $\check{\Upsilon}_{\psi}$ on a minimal modular extension $\check{\mathcal{C}}$ analogously:

$$\check{\Upsilon}_{\psi}(|a, b; c\rangle) = \frac{\gamma_a \gamma_b}{\gamma_c} |a, b; c\rangle$$

$$\text{with } \gamma_a = \begin{cases} -1 & a = \psi \\ 1 & \text{any other } a \in \check{\mathcal{C}} \end{cases} \quad (73)$$

Similar to the case of Υ_{ψ} , the following are equivalent: (i) the map $\check{\Upsilon}_{\psi}$ respects locality; (ii) $\check{\mathcal{C}}$ contains an Abelian fermion parity vortex; (iii) there exists a set of phases $\check{\zeta}_a$ on $\check{\mathcal{C}}$ that obey the fusion rules and have $\check{\zeta}_{\psi} = -1$. After specifying the group homomorphism, one must compute the obstruction to symmetry localization. There are two obstructions. The first is the

“bosonic” obstruction to defining any G_b symmetry fractionalization whatsoever on \mathcal{C} and is valued in $\mathcal{H}^3(G_b, K(\mathcal{C}))$; if Υ_{ψ} violates locality, then we may characterize $K(\mathcal{C}) \simeq \mathcal{A}/\{1, \psi\}$, but if Υ_{ψ} respects locality, there is nothing further to say in general. The derivation is very similar to the bosonic case; see [35] for details.

At this point one may define symmetry fractionalization data $\eta_a(\mathbf{g}, \mathbf{h})$ subject to the usual consistency conditions Eqs. (34) and (35). However, compatibility of the symmetry localization ansatz with Eq. 63 and the full G_f symmetry in Eq. (44) requires [35] the constraint

$$\eta_{\psi}(\mathbf{g}, \mathbf{h}) = \omega_2(\mathbf{g}, \mathbf{h}) \in Z^2(G_b, \mathbb{Z}_2). \quad (74)$$

Regarding autoequivalences as gauge equivalent only if they differ by *locality-respecting* natural isomorphisms means that gauge transformations preserve Eq. (74).

The existence of a symmetry fractionalization pattern obeying Eq. (74) is subject to a “fermionic” symmetry localization obstruction, which is valued in $Z^2(G_b, \mathbb{Z}_2)$ if Υ_{ψ} violates locality and is valued in $\mathcal{H}^3(G_b, K_+(\mathcal{C})) = \mathcal{H}^3(G_b, \mathcal{A}/\{1, \psi\})$ if Υ_{ψ} respects locality.

To summarize, symmetry fractionalization of a fermionic symmetry group G_f on a super-modular category \mathcal{C} is given by a homomorphism $[\rho_{\mathbf{g}}] : G_b \rightarrow \text{Aut}_{LR}(\mathcal{C})$ [which defines the U symbols $U_{\mathbf{g}}(a, b; c)$] and a choice of data $\eta_a(\mathbf{g}, \mathbf{h})$, subject to the same consistency conditions Eqs. (25), (26), (34), and (35) as in the bosonic case. This data is subject to the constraints

$$U_{\mathbf{g}}(\psi, \psi; 1) = +1 \quad (75)$$

$$\eta_{\psi}(\mathbf{g}, \mathbf{h}) = \omega_2(\mathbf{g}, \mathbf{h}), \quad (76)$$

and symmetry action gauge transformations are restricted to be locality respecting, in that they must have

$$\gamma_{\psi}(\mathbf{g}) = +1. \quad (77)$$

The existence of consistent symmetry fractionalization requires two obstructions to vanish: A bosonic symmetry localization obstruction $[\mathfrak{O}_b] \in \mathcal{H}^3(G_b, \mathcal{A}/\{1, \psi\})$ and a fermionic symmetry localization obstruction $[\mathfrak{O}_f]$, which is valued in $Z^2(G_b, \mathbb{Z}_2)$ if Υ_{ψ} violates locality and which is valued in $\mathcal{H}^3(G_b, \mathcal{A}/\{1, \psi\})$ if Υ_{ψ} respects locality.

V. $\mathcal{H}^1(G_b, \mathbb{Z}_T)$ OBSTRUCTION

We prove Theorem 1.2.

Proof. Using the Gauss sum,

$$c_{v_2} = -c_{v_1} \pmod{8}. \quad (78)$$

We have defined

$$c_{v_2} = c_{v_1} + o_1(\mathbf{g})/2 \pmod{8} \quad (79)$$

where $o_1(\mathbf{g})$ must, by the 16-fold way, be an integer. Hence

$$o_1(\mathbf{g}) = -4c_{v_1} \pmod{16}. \quad (80)$$

Hence $c_{v_1} = 0 \pmod{1/4}$. Now, suppose that (v_3, v_4) also satisfy Eq. (79) with a different integer $n'(\mathbf{g})$. Then we could run the same argument to obtain

$$o'_1(\mathbf{g}) = -4c_{v_3} \pmod{16}. \quad (81)$$

But $c_{v_3} - c_{v_1} \in \mathbb{Z}/2$ by the 16-fold way, which means

$$o'_1(\mathbf{g}) - o_1(\mathbf{g}) \in 2\mathbb{Z}. \quad (82)$$

■

It immediately follows that if $o_1(\mathbf{g}) = 1 \pmod{2}$, then it is not possible to have a lift $\rho_{\mathbf{g}} : \check{\mathcal{C}}_{v_0} \rightarrow \check{\mathcal{C}}_{v_0}$. This is a rather familiar statement because $o_1(\mathbf{g}) = 1$ implies $c_{v_i} \in \mathbb{Z} \pm 1/4$, which can never be left invariant by an antiunitary symmetry. Hence $[o_1] \in \mathcal{H}^1(G_b, \mathbb{Z}_T)$ obstructs the ability to lift $\rho_{\mathbf{g}}$ to a map that is a true symmetry of a minimal modular extension $\check{\mathcal{C}}$, rather than a map between two distinct minimal modular extensions. Conjecture 1.2 states that $[o_1]$ is the only such obstruction.

VI. $\mathcal{H}^2(G_b, \ker r)$ OBSTRUCTION

Now let us suppose that the \mathcal{H}^1 obstruction defined in the preceding section vanishes. Here we will find that there is an obstruction to lifting the maps $[\rho_{\mathbf{g}}]$ to $[\check{\rho}_{\mathbf{g}}]$ in a way that appropriately satisfies the group structure.

In what follows, we fix a choice of minimal modular extension $\check{\mathcal{C}}_v$ and suppress the v index. In general the obstruction o_2 that we will find depends on v ; we will discuss this dependence in Sec. VI E. We will assume for now that $\check{\Upsilon}_{\psi}$ [defined in Eq. (73)] respects locality.

We will discuss cases where it violates locality afterwards in Sec. VI A 1. Since $\check{\Upsilon}_{\psi}$ and therefore Υ_{ψ} both respect locality in the first part of our discussion, $[\kappa_{\mathbf{g}, \mathbf{h}}]$ is always the identity and so the map $[\rho_{\mathbf{g}}] : G_b \rightarrow \text{Aut}_{LR}(\mathcal{C})$ is a group homomorphism. Since the \mathcal{H}^1 obstruction vanishes, for each $\mathbf{g} \in G_b$, one can define a consistent lift $[\check{\rho}_{\mathbf{g}}]$ on a fixed minimal modular extension $\check{\mathcal{C}}$, that is, $r([\check{\rho}_{\mathbf{g}}]) = [\rho_{\mathbf{g}}]$, where r is the restriction map

$$r : \text{Aut}_{LR}(\check{\mathcal{C}}) \rightarrow \text{Aut}_{LR}(\check{\mathcal{C}})|_{\mathcal{C}} \subseteq \text{Aut}_{LR}(\mathcal{C}). \quad (83)$$

However, the lifted map $[\check{\rho}_{\mathbf{g}}]$ may not be a group homomorphism. Our aim is to show that there is an obstruction to finding a lift $[\check{\rho}_{\mathbf{g}}]$ that is a group homomorphism $G_b \rightarrow \text{Aut}_{LR}(\check{\mathcal{C}})$, and this obstruction is valued in $\mathcal{H}^2(G_b, \ker r)$. We then relate this obstruction to the $\mathcal{H}^2(G_b, \mathbb{Z}_2)$ part of the 't Hooft anomaly for fermionic SETs.

A. Defining the obstruction

Suppose we have a lift $[\check{\rho}_{\mathbf{g}}]$ of a general autoequivalence $[\rho_{\mathbf{g}}]$ of \mathcal{C} . In general r may have a nontrivial kernel, so $\check{\rho}_{\mathbf{g}}$ can be composed with any element of $\ker r$ to obtain another, equally valid lift. Although $\rho_{\mathbf{g}}$ and $\check{\rho}_{\mathbf{g}}$ may in general be antiunitary, elements of $\ker r$ are automatically unitary.

We will show in Sec. VI B that the permutation action of all elements of $\ker r$ on anyon labels commute with each other, although they in general need not commute with the permutation action of $[\rho_{\mathbf{g}}]$. In the case where elements of $\text{Aut}(\check{\mathcal{C}})$ are completely determined by their permutation action on the anyons, then this implies that $\ker r$ is Abelian. We will assume that $\ker r$ is Abelian in general.

Let us consider

$$o_2(\mathbf{g}, \mathbf{h}) := \check{\rho}_{\mathbf{g}\mathbf{h}} \check{\rho}_{\mathbf{h}}^{-1} \check{\rho}_{\mathbf{g}}^{-1}. \quad (84)$$

By inspection, modifying a representative lift $\check{\rho}_{\mathbf{g}}$ by a locality-respecting natural isomorphism modifies o_2 by a locality-respecting natural isomorphism, so this equation is also well defined in $\text{Aut}_{LR}(\check{\mathcal{C}})$.

We warn the reader that we will overload notation so that $o_2(\mathbf{g}, \mathbf{h})$ can refer both to a topological autoequivalence and its equivalence class in $\text{Aut}_{LR}(\check{\mathcal{C}})$ (after modding out by natural isomorphisms). We reserve $[o_2]$ for later use as a cohomology class.

Since $[\rho_{\mathbf{g}}]$ is a group homomorphism $G_b \rightarrow \text{Aut}_{LR}(\mathcal{C})$, o_2 restricts to a trivial map in $\text{Aut}_{LR}(\mathcal{C})$, i.e., $o_2(\mathbf{g}, \mathbf{h}) \in \ker r$. However, it may be a nontrivial element of $\ker r$. In general, $o_2(\mathbf{g}, \mathbf{h}) \in C^2(G_b, \ker r)$ defines a $(\ker r)$ -valued 2-cochain on G_b .

Demanding that the $\check{\rho}_{\mathbf{g}}$ be associative, we find by decomposing $\check{\rho}_{\mathbf{g}\mathbf{h}\mathbf{k}}$ in two distinct ways that

$$\check{\rho}_{\mathbf{g}\mathbf{h}\mathbf{k}} = o_2(\mathbf{g}\mathbf{h}, \mathbf{k}) \check{\rho}_{\mathbf{g}} \check{\rho}_{\mathbf{h}} \check{\rho}_{\mathbf{k}} \quad (85)$$

$$= o_2(\mathbf{g}\mathbf{h}, \mathbf{k}) o_2(\mathbf{g}, \mathbf{h}) \check{\rho}_{\mathbf{g}} \check{\rho}_{\mathbf{h}} \check{\rho}_{\mathbf{k}} \quad (86)$$

$$= o_2(\mathbf{g}, \mathbf{h}\mathbf{k}) \check{\rho}_{\mathbf{g}} \check{\rho}_{\mathbf{h}\mathbf{k}} \quad (87)$$

$$= o_2(\mathbf{g}, \mathbf{h}\mathbf{k})^g o_2(\mathbf{h}, \mathbf{k}) \check{\rho}_{\mathbf{g}} \check{\rho}_{\mathbf{h}} \check{\rho}_{\mathbf{k}}, \quad (88)$$

where we have defined

$$g o_2(\mathbf{h}, \mathbf{k}) = \check{\rho}_{\mathbf{g}} o_2(\mathbf{h}, \mathbf{k}) \check{\rho}_{\mathbf{g}}^{-1}. \quad (89)$$

For these two decompositions of $\check{\rho}_{\mathbf{g}\mathbf{h}\mathbf{k}}$ to be equal, we need $o_2(\mathbf{g}, \mathbf{h}) \in Z^2(G_b, \ker r)$.

Clearly $[\check{\rho}_{\mathbf{g}}]$ is only a group homomorphism if $o_2 = 1$. This condition is not generically satisfied, but we may obtain another lift by modifying each $[\check{\rho}_{\mathbf{g}}]$ by an element of $\ker r$. Such a modification changes $o_2(\mathbf{g}, \mathbf{h})$ by a $(\ker r)$ -valued 2-coboundary. Therefore, the lift can be modified to obtain a group homomorphism $G_b \rightarrow \text{Aut}_{LR}(\check{\mathcal{C}})$ if and only if $[o_2] \in \mathcal{H}^2(G_b, \ker r)$ is cohomologically trivial. That is, $[o_2]$ is the obstruction to lifting the permutation action of G_b on \mathcal{C} to $\check{\mathcal{C}}$.

In the case where $\ker r = \mathbb{Z}_2$, we will see that $\ker r$ commutes with all of $\text{Aut}_{LR}(\check{\mathcal{C}})$. Accordingly, $[o_2]$ defines a group extension \check{G}_b of G_b by \mathbb{Z}_2 . In fact, if we enlarge the symmetry to \check{G}_b , then there is a consistent lift to $[\check{\rho}_{\mathbf{g}}] \in \text{Aut}_{LR}(\check{\mathcal{C}})$. Let $\ker r = \{1, [\alpha_{\psi}]\}$, and with $\check{G}_b = G_b \times \mathbb{Z}_2$ as sets, define

$$\check{\rho}_{(\mathbf{g}, \mathbf{p})} = \check{\rho}_{\mathbf{g}} \alpha_{\psi}^{\mathbf{p}} \quad (90)$$

with $\mathbf{g} \in G_b$ and $\mathbf{p} \in \{0, 1\} \simeq \mathbb{Z}_2$. Then

$$[\check{\rho}_{(\mathbf{g}, \mathbf{p})}] [\check{\rho}_{(\mathbf{h}, \mathbf{q})}] = o_2(\mathbf{g}, \mathbf{h}) [\alpha_{\psi}]^{\mathbf{p}+\mathbf{q}} [\check{\rho}_{\mathbf{g}\mathbf{h}}] \quad (91)$$

$$= [\check{\rho}_{(\mathbf{g}\mathbf{h}, \mathbf{p}+\mathbf{q}+\tilde{o}_2(\mathbf{g}, \mathbf{h}))}] \quad (92)$$

$$= [\check{\rho}_{(\mathbf{g}, \mathbf{p}) \times (\mathbf{h}, \mathbf{q})}], \quad (93)$$

where $\tilde{o}_2 \in \{0, 1\} \simeq \mathbb{Z}_2$ means we are interpreting $o_2 \in \ker r \simeq \mathbb{Z}_2$ as an element of additive \mathbb{Z}_2 instead of $\ker r$. Hence these symmetry actions are a group homomorphism $[\check{\rho}_{(\mathbf{g}, \mathbf{p})}] : \check{G}_b \rightarrow \text{Aut}_{LR}(\check{\mathcal{C}})$ as claimed.

As an example, let us consider the semion-fermion theory with $G_b = \mathbb{Z}_2^T$. The appropriate modular extension of this theory is $\check{\mathcal{C}} = U(1)_2 \times U(1)_{-4}$. The simple objects of $\check{\mathcal{C}}$ can be labeled (a, b) for $a = 0, 1$ and $b = 0, 1, 2, 3$. Here, $v = (0, 1)$ is the fermion parity vortex, $\psi = (0, 2)$ is the fermion, and $s = (1, 0)$ is the semion. It is clear that $\ker r = \mathbb{Z}_2$ in this

case; its nontrivial element $[\alpha_\psi]$ takes $v \leftrightarrow v \times \psi$. Under \mathbf{T} , \mathcal{C} transforms as follows:

$$\begin{aligned} \mathbf{T}_s &= s \times \psi, \\ \mathbf{T}_\psi &= \psi. \end{aligned} \quad (94)$$

There are two possible lifts $[\check{\rho}_T]$ to the modular extension, which differ by the action of $[\alpha_\psi]$; we may take either

$$\mathbf{T}_v = s \times v \text{ or } \mathbf{T}_v = s \times \psi \times v. \quad (95)$$

For the first choice

$$\mathbf{T}(\mathbf{T}_v) = \mathbf{T}(s \times v) = (s \times \psi) \times (s \times v) = \psi \times v. \quad (96)$$

That is,

$$[\check{\rho}_T]^2 = [\alpha_\psi] = o_2(\mathbf{T}, \mathbf{T}). \quad (97)$$

One can check straightforwardly that the second choice of $[\check{\rho}_T]$ leads to the same o_2 . Therefore $[o_2] \neq +1 \in \mathcal{H}^2(\mathbb{Z}_2^T, \mathbb{Z}_2)$. Thus there is no way to have the permutations faithfully act on the modular extension as $G_b = \mathbb{Z}_2^T$. However we can have them act as $\check{G}_b = \mathbb{Z}_4^T$.

1. $\check{\Upsilon}_\psi$ violates locality

If $\check{\Upsilon}_\psi$ violates locality, the discussion above must be slightly modified.⁷

In particular, there are now two possibilities. The first possibility is that Υ_ψ does not respect locality. In this case, if the symmetry fractionalization on \mathcal{C} is unobstructed, $[\rho_g]$ generally fails to be a group homomorphism up to factors of $[\Upsilon_\psi]$ [35,36]:

$$[\Upsilon_\psi]^{\tilde{\omega}_2(\mathbf{g}, \mathbf{h})} [\rho_g] [\rho_h] = [\rho_{gh}]. \quad (98)$$

where $\tilde{\omega}_2$ is defined by Eq. (43). In this case, we must instead enforce the equation

$$[\check{\Upsilon}_\psi]^{\tilde{\omega}_2(\mathbf{g}, \mathbf{h})} [\check{\rho}_g] [\check{\rho}_h] = [\check{\rho}_{gh}] \quad (99)$$

because $r([\check{\Upsilon}_\psi]) = [\Upsilon_\psi]$ and we need Eq. (99) to reduce to Eq. (98) upon applying the restriction map r . We can thus define a modified obstruction, which must vanish if we want Eq. (99) to hold:

$$o_2(\mathbf{g}, \mathbf{h}) := \check{\rho}_{gh} \check{\rho}_h^{-1} \check{\rho}_g^{-1} \check{\Upsilon}_\psi^{\tilde{\omega}_2(\mathbf{g}, \mathbf{h})}. \quad (100)$$

The rest of the analysis showing that $[o_2] \in \mathcal{H}^2(G_b, \ker r)$ proceeds analogously to the case where $\check{\Upsilon}_\psi$ respects locality, with the additional factors of $\check{\Upsilon}_\psi$ canceling out at all stages. Proving this statement requires using the fact, discussed in Sec. VIB, that $\check{\Upsilon}_\psi$ commutes with all topological autoequivalences.

⁷The first version of this paper assumed that $[\rho_g]$ and $[\check{\rho}_g]$ is always a group homomorphism, and did not incorporate the distinction between situations where $\check{\Upsilon}_\psi$ violates locality and preserves it. This was addressed in the first version of Ref. [36], which appeared on the arXiv simultaneously with the first version of this paper. We provide an alternate treatment here.

In the case where Υ_ψ respects locality but $\check{\Upsilon}_\psi$ does not, we again need to enforce Eq. (99), but for a slightly more involved reason. We take a lift $\check{\rho}_g$ and attempt to enforce the condition

$$\check{\kappa}_{\mathbf{g}, \mathbf{h}} \check{\rho}_g \check{\rho}_h = \check{\rho}_{gh} \quad (101)$$

with $\check{\kappa}_{\mathbf{g}, \mathbf{h}}$ a (possibly locality-violating) natural isomorphism. Since $\check{\Upsilon}_\psi$ violates locality, this condition requires that the decomposition $\check{\kappa}_{\mathbf{g}, \mathbf{h}}(a, b; c)$ into anyon-dependent factors $\check{\beta}_a$ obey

$$\check{\beta}_\psi = \begin{cases} 1 & [\check{\kappa}_{\mathbf{g}, \mathbf{h}}] = [1] \\ -1 & [\check{\kappa}_{\mathbf{g}, \mathbf{h}}] = [\check{\Upsilon}_\psi] \end{cases}. \quad (102)$$

The above equation is gauge invariant. Now, in order to be compatible with G_f , any putative G_f symmetry fractionalization pattern will need to obey $\check{\eta}_\psi(\mathbf{g}, \mathbf{h}) = \omega_2(\mathbf{g}, \mathbf{h})$. If such a G_f symmetry fractionalization pattern can possibly exist, then

$$\check{\omega}_\psi(\mathbf{g}, \mathbf{h}) = \frac{\check{\beta}_\psi(\mathbf{g}, \mathbf{h})}{\check{\eta}_\psi(\mathbf{g}, \mathbf{h})} = +1 \Rightarrow \check{\beta}_\psi(\mathbf{g}, \mathbf{h}) = \omega_2(\mathbf{g}, \mathbf{h}) \quad (103)$$

[see Eq. (37)] since $\check{\Upsilon}_\psi$ violates locality and $\check{\omega}_a$ must respect the fusion rules. Combining Eqs. (101)–(103), we find that our desired lift should again obey Eq. (99). Accordingly, we should define o_2 via Eq. (100). The rest of the analysis is unchanged from the case where Υ_ψ violates locality.

Note that in Eq. (99), the brackets $[\cdot]$ correspond to taking equivalence under *locality-respecting* natural isomorphisms, so that $[\check{\rho}_g] \in \text{Aut}_{LR}(\check{\mathcal{C}})$. When we instead consider equivalence under all natural isomorphisms, $\check{\rho}$ will reduce to a group homomorphism, as expected based on the theory of bosonic SETs [1].

B. Characterizing $\ker r$

In order to understand $[o_2]$ better, we need to characterize $\ker r$. We make the following conjecture:

Conjecture VI.1. In all cases, $\ker r = \mathbb{Z}_2$.

As a consequence, $[o_2] \in \mathcal{H}^2(G_b, \mathbb{Z}_2)$. Although we cannot prove this conjecture in full generality, in the following subsections we will provide a number of concrete results that motivate the conjecture and give partial progress towards proving it. One motivation for the above conjecture is that it is $\mathcal{H}^2(G_b, \mathbb{Z}_2)$ that appears in the characterization of $(3+1)$ D FSPTs, which should classify the 't Hooft anomalies in $(2+1)$ D.

We begin by discussing a few properties of the map $[\check{\Upsilon}_\psi]$ and define an important map $[\alpha_\psi]$. We will see that $[\alpha_\psi]$ always generates a $\mathbb{Z}_2 \subseteq \ker r$.

1. $[\check{\Upsilon}_\psi]$

We defined the map $\check{\Upsilon}_\psi$ by Eq. (73). It immediately follows that $r([\check{\Upsilon}_\psi]) = [\Upsilon_\psi]$, so $[\check{\Upsilon}_\psi] \in \ker r$ if and only if Υ_ψ respects locality. If $[\check{\Upsilon}_\psi] \in \ker r$ and $\check{\Upsilon}_\psi$ violates locality, then it forms a \mathbb{Z}_2 subgroup of $\ker r$. On the other hand, if $\check{\Upsilon}_\psi$ respects locality, then it is a trivial element of $\ker r$.

Since ψ is invariant under all (fermionic) topological autoequivalences, it follows that $\check{\Upsilon}_\psi$ commutes with all topological autoequivalences.

There are three possibilities for how $\check{\Upsilon}_\psi$ and Υ_ψ behave. First, $\check{\Upsilon}_\psi$ may respect locality, in which case, Υ_ψ does as well. It was proven in [35] that all parity vortices are v type in this case (and at least one is Abelian). Second, $\check{\Upsilon}_\psi$ may violate locality but Υ_ψ may respect locality. From the list of properties in Sec. IV A, we see that this occurs when \mathcal{C} has some minimal modular extension with an Abelian fermion parity vortex, but $\check{\mathcal{C}}$ is not such a minimal modular extension. It was proven in [35] that all parity vortices in $\check{\mathcal{C}}$ are σ -type in this case. Finally, both $\check{\Upsilon}_\psi$ and Υ_ψ may violate locality.

One can check that all of the cases we considered above can actually occur. For an example where $\check{\Upsilon}_\psi$ respects locality, we can take $\mathcal{C} = \mathcal{B} \boxtimes \{1, \psi\}$ for any modular \mathcal{B} and $\check{\mathcal{C}} = \mathcal{B} \boxtimes D(\mathbb{Z}_2)$, where $D(\mathbb{Z}_2)$ is the quantum double of \mathbb{Z}_2 , also known as the toric code topological order. Here \boxtimes is the Deligne product and physically corresponds to stacking decoupled topological orders. Instead taking $\check{\mathcal{C}} = \mathcal{B} \boxtimes \text{Ising}$ gives an example where $\check{\Upsilon}_\psi$ does not respect locality but Υ_ψ does. For an example where neither $\check{\Upsilon}_\psi$ nor Υ_ψ respects locality, we can take $\mathcal{C} = \text{SO}(3)_3$; an example minimal modular extension is $\check{\mathcal{C}} = \text{SU}(2)_6$.

2. $[\alpha_\psi]$

Next, we consider a map α_ψ , with the following permutation action:

$$\begin{aligned} \alpha_\psi(a) &= a & \text{if } a \in \check{\mathcal{C}}_0 \simeq \mathcal{C} \\ \alpha_\psi(a) &= a \times \psi & \text{if } a \in \check{\mathcal{C}}_1. \end{aligned} \quad (104)$$

This permutation preserves the fusion rules, twists, and modular S matrix of the theory. One can check in a range of examples that there indeed exists a braided autoequivalence α_ψ of $\check{\mathcal{C}}$ with this permutation action. In fact, Ref. [36] (which appeared simultaneously on the arXiv with the first version of this work) gave an explicit formula for the U -symbols of exactly such a braided autoequivalence as follows:

$$\begin{aligned} U_{\alpha_\psi}(a', b'; c'; \mu, v) &= \sum_{\lambda} \left[(F_c^{a, \psi f_b, b'})^{-1} \right]_{(b, \mu), (a \times \psi f_b, \lambda)} \\ &\times R^{\psi f_b, a} \left[(F_{c'}^{\psi f_c, \psi f_b \times a, b'})^{-1} \right]_{(c, \lambda), (a', v)} \end{aligned} \quad (105)$$

where we use the shorthand $a' = \alpha_\psi(a)$ and define

$$f_x = \begin{cases} 0 & x \in \check{\mathcal{C}}_0 \\ 1 & x \in \check{\mathcal{C}}_1 \end{cases}. \quad (106)$$

Consider the case where (i) every permutation of simple objects of $\check{\mathcal{C}}$ that preserves the modular data corresponds to a unique element of $\text{Aut}(\check{\mathcal{C}})$, and similarly (ii) every permutation of simple objects in \mathcal{C} preserving the modular data corresponds to a unique element of $\text{Aut}(\mathcal{C})$. Then $[\alpha_\psi]$ generates a \mathbb{Z}_2 subgroup of $\ker r$ as long as $\check{\mathcal{C}}_v$ is nonempty. In more general situations, Ref. [36] demonstrated that as a braided autoequivalence, it is always true that $[\alpha_\psi] \in \ker r$, commutes with all of $\text{Aut}_{LR}(\check{\mathcal{C}})$, and squares to the identity.

If $\check{\mathcal{C}}_v$ is nonempty, then $[\alpha_\psi]$ is clearly nontrivial. If $\check{\mathcal{C}}_\sigma$ is nonempty, then we can calculate the gauge-invariant (in

$\text{Aut}_{LR}(\check{\mathcal{C}})$) quantity

$$U_{\alpha_\psi}(\sigma, \psi; \sigma) = (F^{\sigma 1\psi})^{-1} R^{1\sigma} (F^{\psi \sigma \psi})^{-1} \quad (107)$$

$$= (F^{\psi \sigma \psi})^{-1} = -1, \quad (108)$$

where the last equality follows from a straightforward use of the hexagon equation. Hence $[\alpha_\psi]$ is nontrivial in this case as well, and so $[\alpha_\psi]$ always generates a central \mathbb{Z}_2 subgroup of $\ker r$. Note that Eq. (105) provides a definition of a nontrivial $[\alpha_\psi]$ even when $\check{\mathcal{C}}_v$ is empty, in which case α_ψ always has trivial permutation action on the objects in $\check{\mathcal{C}}$.

For a familiar example, consider $\mathcal{C} = \{1, \psi\}$; then there is a modular extension $\check{\mathcal{C}} = \{1, \psi, e, m\}$ that is equivalent to \mathbb{Z}_2 gauge theory, where we are viewing $\check{\mathcal{C}}_v = \{e, m\}$. Then the map α_ψ permutes $e \leftrightarrow m$, implementing electric-magnetic duality.

If $\check{\mathcal{C}}_v$ is empty, then $[\alpha_\psi]$ may or may not equal $[\check{\Upsilon}_\psi]$. If Υ_ψ violates locality, then these maps cannot be equal since $[\alpha_\psi]$ is in $\ker r$ but $[\check{\Upsilon}_\psi]$ is not. In a version of Ref. [36] posted after the second version of this paper, it was proven that $[\alpha_\psi] = [\check{\Upsilon}_\psi]$ if and only if $\check{\Upsilon}_\psi$ violates locality but Υ_ψ respects locality. In other words, any time $[\check{\Upsilon}_\psi]$ is a nontrivial element in $\ker r$, it is equal to $[\alpha_\psi]$.

3. Permutation actions of elements of $\ker r$

Fully characterizing $\ker r$ is a nontrivial task in general. We can, however, determine the allowed *permutation action* of all elements of $\ker r$ by proving Theorem I.3:

Proof. We may apply the Verlinde formula to $\check{\mathcal{C}}$, which is modular:

$$N_{v, \psi}^{\check{\rho}(v)} + N_{v, 1}^{\check{\rho}(v)} = \sum_{x \in \check{\mathcal{C}}} \frac{(S_{\psi, x} + S_{1x}) S_{vx} S_{\check{\rho}(v)x}^*}{S_{1x}} \quad (109)$$

$$= \sum_{x \in \check{\mathcal{C}}_0} \frac{(S_{\psi, x} + S_{1x}) S_{vx} S_{\check{\rho}(v)x}^*}{S_{1x}} \quad (110)$$

$$+ \sum_{x \in \check{\mathcal{C}}_1} \frac{(S_{\psi, x} + S_{1x}) S_{vx} S_{\check{\rho}(v)x}^*}{S_{1x}}. \quad (110)$$

By Eq. (17), $S_{\psi, x} = \pm S_{1, x}$ with the upper sign for $x \in \check{\mathcal{C}}_0$ and the lower sign for $x \in \check{\mathcal{C}}_1$. Therefore,

$$N_{v, \psi}^{\check{\rho}(v)} + N_{v, 1}^{\check{\rho}(v)} = \sum_{x \in \check{\mathcal{C}}_0} 2 S_{vx} S_{\check{\rho}(v)x}^* + 0 \quad (111)$$

$$= \sum_{x \in \check{\mathcal{C}}_0} 2 |S_{vx}|^2 > 0, \quad (112)$$

where we have used the fact that S is invariant under $\check{\rho}$ and that $\check{\rho}(x) = x$ if $x \in \check{\mathcal{C}}_0$. Therefore either $\check{\rho}(v) = v$ or $\check{\rho}(v) = \psi \times v$, and in particular, if $v \in \check{\mathcal{C}}_\sigma$, $\check{\rho}(v) = v$. Next, suppose $v_1, v_2 \in \check{\mathcal{C}}_v$. Then

$$S_{v_1, v_2} = \check{\rho}(S_{v_1, v_2}) = S_{\check{\rho}(v_1), \check{\rho}(v_2)} = S_{v_1, v_2} (-1)^{m_1 + m_2} \quad (113)$$

where

$$m_i = \begin{cases} 0 & \check{\rho}(v_i) = v_i \\ 1 & \check{\rho}(v_i) = v_i \times \psi \end{cases}. \quad (114)$$

Hence if $S_{v_1, v_2} \neq 0$, then $m_1 = m_2$, that is, $\check{\rho}$ changes the fermion parity of both v_1 and v_2 or of neither. If $S_{v_1, v_2} = 0$ but v_1, v_2 are in the same one of the k blocks of the S_{a_v, b_v} part of the S matrix, then there exists a sequence of v -type vortices a_1, a_2, \dots, a_p such that $S_{v_1, a_1} \neq 0, S_{a_1, a_2} \neq 0, \dots, S_{a_p, v_2} \neq 0$. Applying the above argument to each pair in the sequence, we conclude that if v_1, v_2 belong to the same block of the S matrix, $\check{\rho}$ acts the same way on both vortices, i.e., it either changes the fermion parity of both or of neither.

4. $\ker r$ when permutations determine $\text{Aut}(\check{\mathcal{C}})$ and $\text{Aut}(\mathcal{C})$

In many well-studied examples, every permutation of the simple objects of a BFC \mathcal{B} uniquely determines an element of $\text{Aut}(\mathcal{B})$. We can fully characterize $\ker r$ as long as $\check{\mathcal{C}}$ and \mathcal{C} obey slightly weaker properties:

Theorem VI.1. Let k be as in Theorem I.3. Suppose that every permutation of vortices in $\check{\mathcal{C}}$ given in Theorem I.3 uniquely determines an element of $\text{Aut}(\check{\mathcal{C}})$ (but not necessarily a unique element of $\text{Aut}_{LR}(\check{\mathcal{C}})$), and further suppose that there is a unique element of $\text{Aut}(\mathcal{C})$ that does not permute anyons. Then $\ker r = \mathbb{Z}_2^{\max(k, 1)}$.

Proof. Let $[\check{\rho}] \in \ker r$. According to Theorem I.3, there are $2^{\max(k, 1)}$ possible permutation actions for $[\check{\rho}]$, given by k independent choices of whether or not $[\check{\rho}]$ changes the fermion parity of the v -type vortices in each block, and each permutation action squares to the identity. If $k = 0$, then $[\check{\rho}]$ acts as the identity permutation.

We first claim that all of these possible permutation actions commute with each other. This statement is only nontrivial for $k > 1$. To prove this claim, it suffices to show that v and $v \times \psi$ belong to the same block.

Suppose first that for some $w \neq v$, v and w belong to the same one of the k blocks of the S matrix. Then there is a sequence of v -type vortices a_1, a_2, \dots, a_p such that $S_{v, a_1} \neq 0, S_{a_1, a_2} \neq 0, \dots, S_{a_p, w} \neq 0$. But $S_{v \times \psi, a_1} = -S_{v, a_1} \neq 0$ as well. Hence $v \times \psi$ is also in the same block as w , so v and $v \times \psi$ are in the same block, namely the block containing w .

We claim that such a w must exist. Suppose by way of contradiction that no such w exists, i.e., $S_{v, w} = 0$ for all $w \in \check{\mathcal{C}}_1$, in which case v is in a block by itself. It follows that $S_{v \times \psi, w} = -S_{v, w} = 0$ for all $w \in \check{\mathcal{C}}_1$, and since $S_{v \times \psi, a} = S_{v, a}$ for all $a \in \check{\mathcal{C}}_0$, we must have $S_{v, a} = S_{v \times \psi, a}$ for all $a \in \check{\mathcal{C}}$. Hence S has two identical rows and is not invertible, which is a contradiction since $\check{\mathcal{C}}$ is modular. This proves that all of the aforementioned permutations commute.

By assumption, the permutation action of $\check{\rho}$ defines a unique element of $\text{Aut}(\check{\mathcal{C}})$, but may or may not uniquely determine an element of $\text{Aut}_{LR}(\check{\mathcal{C}})$. Whether or not it does depends on the properties of $[\check{\Upsilon}_\psi]$, which is the one possibly-nontrivial element of $\text{Aut}_{LR}(\check{\mathcal{C}})$, which does not permute anyons.

We need to consider three possible cases, depending on whether Υ_ψ and $\check{\Upsilon}_\psi$ respect locality.

Case 1: $\check{\Upsilon}_\psi$ respects locality in $\check{\mathcal{C}}$. Then Υ_ψ respects locality in \mathcal{C} , and also $\check{\mathcal{C}}$ contains an Abelian fermion parity vortex. Hence $\check{\mathcal{C}}_v$ is nonempty, $k > 0$, $\text{Aut}_{LR}(\check{\mathcal{C}}) = \text{Aut}(\check{\mathcal{C}})$, and $\text{Aut}_{LR}(\mathcal{C}) = \text{Aut}(\mathcal{C})$. Therefore, each of the 2^k anyon permutations defined above determines a unique element of

$\text{Aut}_{LR}(\check{\mathcal{C}})$ that restricts to the identity in $\text{Aut}_{LR}(\mathcal{C})$ (since the restricted permutation action is trivial). Furthermore, if $[\check{\rho}] \in \ker r$, $[\check{\rho}]^2 = [1]$ because $[\check{\rho}]^2$ does not permute anyons. Thus $\ker r = \mathbb{Z}_2^k$.

Case 2: $\check{\Upsilon}_\psi$ does not respect locality in $\check{\mathcal{C}}$, but Υ_ψ respects locality in \mathcal{C} . Then some minimal modular extension of \mathcal{C} contains an Abelian fermion parity vortex, but $\check{\mathcal{C}}$ does not; according to the list of properties in Sec. IV C, this implies that $\check{\mathcal{C}}$ has no v -type vortices, i.e., $k = 0$. According to Theorem I.3, $[\check{\rho}]$ must therefore act as the identity permutation. Exactly two elements of $\text{Aut}_{LR}(\check{\mathcal{C}})$ implement the trivial permutation, namely $[\check{\Upsilon}_\psi] \neq [1]$ in $\check{\mathcal{C}}$. In particular, $[\check{\Upsilon}_\psi]$ restricts to $[\Upsilon_\psi] = [1]$ in \mathcal{C} . Hence $[\check{\Upsilon}_\psi] \in \ker r$, and $\ker r = \mathbb{Z}_2^{\max(k, 1)} = \mathbb{Z}_2$.

Case 3: Neither $\check{\Upsilon}_\psi$ nor Υ_ψ respect locality in their respective categories. In this case, $r([\check{\Upsilon}_\psi]) = [\Upsilon_\psi] \neq [1]$, so $[\check{\Upsilon}_\psi] \notin \ker r$. But $[\alpha_\psi]$ is a nontrivial element of $\ker r$, and thus must not equal $[\check{\Upsilon}_\psi]$. Therefore, $[\alpha_\psi]$ has a nontrivial permutation action, that is, $\check{\mathcal{C}}_v$ is nonempty and $k > 0$. Now consider any braided autoequivalence $[\check{\rho}] \in \text{Aut}_{LR}(\check{\mathcal{C}})$ that implements one of the anyon permutation actions given above. $[\check{\rho}]$ must restrict to either $[1]$ or $[\Upsilon_\psi]$ on \mathcal{C} , and $[\check{\Upsilon}_\psi \check{\rho}]$ will restrict to $[\Upsilon_\psi]$ or $[1]$ respectively. Hence exactly one of $[\check{\rho}]$ and $[\check{\Upsilon}_\psi \check{\rho}]$ is in $\ker r$, that is, each of the 2^k permutation actions above defines a unique element of $\ker r$. If $[\check{\rho}] \in \ker r$, then $r([\check{\rho}]^2) = [1]$, so $[\check{\rho}]^2 = 1$ as well. Hence $\ker r = \mathbb{Z}_2^k$, concluding our proof. ■

It is true but not immediately obvious that one can have $k > 1$ decoupled blocks of the $v - v$ part of the S matrix; we consider an explicit example with $k = 2$ in Sec. VI F.

5. $\ker r$ when permutations do not uniquely determine $\text{Aut}(\check{\mathcal{C}})$ and $\text{Aut}(\mathcal{C})$

If there is not a one-to-one correspondence between elements of $\text{Aut}(\check{\mathcal{C}})$ (resp. $\text{Aut}(\mathcal{C})$) and anyon permutations, which preserve the fusion rules and modular data of $\check{\mathcal{C}}$ (resp. \mathcal{C}), then we are not generally able to give any further characterization of $\ker r$ as a group beyond the results of the previous subsections.

We do know that Theorem I.3 still applies; the anyon permutations of all elements of $\ker r$ are still restricted. However, three possibilities could further complicate the analysis of $\ker r$:

(1) There could be an anyon permutation that preserves the modular data of $\check{\mathcal{C}}$ but does not correspond to any braided autoequivalence of $\check{\mathcal{C}}$. In this case, there must be a map $\check{\rho}$ from $\check{\mathcal{C}}$ to an inequivalent theory with the same modular data as $\check{\mathcal{C}}$, that is, $\check{\mathcal{C}}$ must have a “modular isotope.” This can occur in general [40,41]. In this situation, $\ker r$ may contain fewer than 2^k distinct permutation actions.

(2) There could be a nontrivial but nonpermuting element of $\text{Aut}(\check{\mathcal{C}})$. If such an autoequivalence exists and has a representative in $\ker r$, then each allowed permutation action may determine many elements of $\ker r$, and each allowed permutation action may give rise to a subgroup of $\ker r$ that is larger

than \mathbb{Z}_2 . UMTCs can in general have nontrivial nonpermuting autoequivalences [42].

(3) There could be a nontrivial but nonpermuting element of $\text{Aut}(\mathcal{C})$. If such an autoequivalence exists, then there may be some permutation actions allowed by Theorem I.3, which, nevertheless, do not determine an element of $\ker r$ because any braided autoequivalence, which implements such a permutation action necessarily restricts to a nontrivial but nonpermuting element of $\text{Aut}_{LR}(\mathcal{C})$. In this situation, $\ker r$ may again contain fewer than 2^k distinct permutation actions.

A version of Ref. [36] posted after the second version of the present paper showed that whenever $\check{\Upsilon}_\psi$ respects locality, $\ker r = \{1, [\alpha_\psi]\} \simeq \mathbb{Z}_2$. Conjecture VI.1 remains open in the case where $\check{\Upsilon}_\psi$ violates locality.

We have not found any example where we know that a particular one of the above possibilities is relevant for characterizing $\ker r$. In Sec. VI F, we study an example involving two copies of $\text{SO}(3)_3$, for which we suspect one of the above possibilities is occurring.

C. Relation to Arf invariant on torus

In this section, we demonstrate that when $\ker r = \{1, [\alpha_\psi]\} \simeq \mathbb{Z}_2$, our obstruction o_2 leads to a modification of the action of the symmetry on the torus Hilbert space of the SET in a way which is sensitive to the Arf invariant of the spin structure on the torus. Reference [33] showed that for the special case $G_f = \mathbb{Z}_4^{\text{T},f}$, such a sensitivity appears in certain systems with a 't Hooft anomaly; where there is overlap, our results agree.

Consider the ground states of the TQFT on a spatial torus, T^2 . As described in Sec. II A, the Hilbert space breaks up into four sectors $\mathcal{H}_{\mu,\nu}$, where $\{\mu, \nu\} \in \{0, 1\}$ specify a spin structure. Here $\mu = 0$ refers to antiperiodic (Neveu-Schwarz) boundary conditions and 1 refers to periodic (Ramond) boundary conditions. These are sometimes also referred to as bounding and nonbounding spin structures, respectively.

Here $\mathcal{H}_{\mu,\nu}$ is a \mathbb{Z}_2 graded Hilbert space that includes both even and odd fermion number sectors. That is, we can think of this as the TQFT Hilbert space allowing for the possibility of a ψ puncture. We will use the basis Eq. (19) and the description Eq. (20) for the torus Hilbert space. Since the torus Hilbert space of the fermionic theory is defined via states of the minimal modular extension $\check{\mathcal{C}}$ on the 3-punctured sphere (i.e., its states are defined using simple objects in $\check{\mathcal{C}}$), only a (representative) lift $\check{\rho}_g$ has a well-defined action on the torus Hilbert space.

Our aim is to show that if there is an \mathcal{H}^2 anomaly, then the action of $\check{\rho}_g$ on the torus Hilbert space is necessarily deformed by the Arf invariant

$$\text{Arf}(\mu, \nu) = \mu\nu \quad (115)$$

of the spin structure. More precisely, given a state $|\Psi\rangle_{\mu\nu} \in \mathcal{H}_{\mu,\nu}$, the G_b group law of the lift is deformed in the sense

$$\check{\rho}_{gh}\check{\rho}_h^{-1}\check{\rho}_g^{-1}|\Psi\rangle_{\mu\nu} = (\omega_2(\mathbf{g}, \mathbf{h}))^F (-1)^{\tilde{o}_2(\mathbf{g}, \mathbf{h})\text{Arf}(\mu, \nu)}|\Psi\rangle_{\mu\nu}, \quad (116)$$

where $(-1)^F$ is the fermion parity operator. The notation $\tilde{o}_2(\mathbf{g}, \mathbf{h}) \in \mathbb{Z}_2 \simeq \{0, 1\}$ distinguishes when we are viewing o_2

as an element of \mathbb{Z}_2 as an additive group from when we view $o_2 \in \mathbb{Z}_2 \simeq \ker r$. Equation (116) extends and sharpens the results of [33], which considered only the special case $G_f = \mathbb{Z}_4^{\text{T},f}$.

From the definition,

$$\begin{aligned} & \check{\rho}_{gh}\check{\rho}_h^{-1}\check{\rho}_g^{-1}|\Psi\rangle_{\mu\nu} \\ &= \begin{cases} o_2(\mathbf{g}, \mathbf{h})|\Psi\rangle_{\mu\nu} & \check{\Upsilon}_\psi \text{ respects locality} \\ o_2(\mathbf{g}, \mathbf{h})\check{\Upsilon}_\psi^{\tilde{o}_2(\mathbf{g}, \mathbf{h})}|\Psi\rangle_{\mu\nu} & \check{\Upsilon}_\psi \text{ violates locality} \end{cases} \end{aligned} \quad (117)$$

Recall that \tilde{o}_2 is the additive parametrization of ω_2 defined by Eq. (43). One can compute directly that $\check{\Upsilon}_\psi$ acts on the torus Hilbert space as fermion parity, that is, it inserts a minus sign on states of the form $|\sigma\rangle_{11}$ and acts as the identity otherwise, so in all other sectors there is no difference whether $\check{\Upsilon}_\psi$ respects or violates locality.

Since $o_2(\mathbf{g}, \mathbf{h})$ acts trivially on fusion spaces V_{ab}^c with $a, b, c \in \check{\mathcal{C}}_0$, it is immediate that Eq. (116) holds for $\mu = 0$.

Now fix a v -type vortex v and consider $|v, \bar{v}; 1\rangle \in V_1^{v\bar{v}}$; then

$$\begin{aligned} o_2(\mathbf{g}, \mathbf{h})(|v, \bar{v}; 1\rangle) &= U_{o_2(\mathbf{g}, \mathbf{h})}(v \times \psi^{\tilde{o}_2}, \bar{v} \times \psi^{\tilde{o}_2}; 1) \\ &\quad \times |v \times \psi^{\tilde{o}_2}, \bar{v} \times \psi^{\tilde{o}_2}; 1\rangle. \end{aligned} \quad (118)$$

The U factors obtained from the action of o_2 on $V_1^{v\bar{v}}$ and its dual $V_{\bar{v}\bar{v}}^1$ are complex conjugates and thus will always cancel out, so the action of $o_2(\mathbf{g}, \mathbf{h})$ is either trivial or simply interchanges $(V_1^{v\bar{v}} \otimes V_{\bar{v}\bar{v}}^1)$ and $(V_1^{v\bar{v}, \bar{v}\bar{v}} \otimes V_{v\bar{v}, \bar{v}\bar{v}}^1)$. Said differently,

$$o_2(\mathbf{g}, \mathbf{h})|v\rangle = |v \times \psi^{\tilde{o}_2}\rangle \quad (119)$$

where $|v\rangle$ is a torus state of definite topological charge v piercing the α cycle. Therefore, using Eq. (19), we find

$$o_2(\mathbf{g}, \mathbf{h})|v\rangle_{1,0} = |v\rangle_{1,0} \quad (120)$$

$$o_2(\mathbf{g}, \mathbf{h})|v\rangle_{1,1} = (-1)^{\tilde{o}_2(\mathbf{g}, \mathbf{h})}|v\rangle_{1,1}, \quad (121)$$

which verifies Eq. (116) for the present case.

Finally, we must consider states built from a σ -type vortex, which only exist when $\check{\Upsilon}_\psi$ violates locality. The above argument immediately generalizes to the unpunctured states to show that

$$o_2(\mathbf{g}, \mathbf{h})|\sigma\rangle_{1,0} = |\sigma\rangle_{1,0}. \quad (122)$$

Running a similar argument on the states with a puncture, we find that

$$\begin{aligned} & o_2(\mathbf{g}, \mathbf{h})\check{\Upsilon}_\psi^{\tilde{o}_2(\mathbf{g}, \mathbf{h})}|\sigma; \psi\rangle \\ &= \omega_2(\mathbf{g}, \mathbf{h})U_{o_2(\mathbf{g}, \mathbf{h})}(\sigma, \bar{\sigma}; 1)U_{o_2(\mathbf{g}, \mathbf{h})}^*(\sigma, \bar{\sigma}; \psi)|\sigma; \psi\rangle \end{aligned} \quad (123)$$

Since o_2 does not permute σ , and $\sigma \times \psi = \sigma$, one can show from the consistency conditions that

$$U_{o_2(\mathbf{g}, \mathbf{h})}(\sigma, \bar{\sigma}; 1)U_{o_2(\mathbf{g}, \mathbf{h})}^*(\sigma, \bar{\sigma}; \psi) = U_{o_2(\mathbf{g}, \mathbf{h})}(\sigma, \psi; \sigma). \quad (124)$$

This last quantity was computed in Eq. (108); combining it with Eqs. (123) and (124), we obtain Eq. (116) for this last class of states.




FIG. 1. Anomaly inflow for the $\mathcal{H}^2(G_b, \mathbb{Z}_2)$ anomaly. The fermion SET lives on the surface of a $(3+1)$ D fermion SPT. A trijunction of domain walls (orange) with $o_2(\mathbf{g}, \mathbf{h}) \neq 0$ is decorated with a Kitaev chain (green) in the bulk with a Majorana fermion γ at its endpoint (grey). Braiding a fermion parity vortex v (pink) around the trijunction transforms $v \rightarrow v \times \psi$ according to the surface theory, so reannihilating the parity vortices leaves behind a fermion. The parity vortices on the surface are endpoints of a fermion parity vortex string that goes through the bulk (thick-black line). The parity vortex creation and motion is given by a membrane operator in the bulk (blue surface), which links with the Kitaev chain. The linking of the blue membrane operator with the Kitaev chain induces a \mathbb{Z}_2^f defect in the Kitaev chain, which locally changes the fermion parity, compensating for the additional fermion arising from the transformation $v \rightarrow v \times \psi$.

D. Anomaly inflow from bulk $(3+1)$ D FSPT

According to the classification of fermion SPTs reviewed in Sec. IV A, if an FSPT has $[n_1] = 0$, then there is a piece of data $[n_2] \in \mathcal{H}^2(G_b, \mathbb{Z}_2)/\Gamma^2$ in the specification of a $(3+1)$ D FSPT. Let us assume that $\ker r = \mathbb{Z}_2 = \{[1], [\alpha_\psi]\}$; then our obstruction $[o_2]$ is valued in $\mathcal{H}^2(G_b, \mathbb{Z}_2)$. Letting

$$q_{\Gamma^2} : \mathcal{H}^2(G_b, \mathbb{Z}_2) \rightarrow \mathcal{H}^2(G_b, \mathbb{Z}_2)/\Gamma^2, \quad (125)$$

we argue that the 't Hooft anomaly data $[n_2]$ of the FSET is given by

$$[n_2] = q_{\Gamma^2}([o_2]). \quad (126)$$

In support of this conjecture, we give an argument explicitly describing anomaly inflow using the decorated domain wall construction for FSPTs. This argument will work at the level of cocycles, without any quotient by coboundaries or by Γ^2 ; we discuss the Γ^2 redundancy in the 't Hooft anomaly in Sec. VIE. We will also discuss the possibility that $\ker r \neq \mathbb{Z}_2$ in Sec. VIF.

First, we note that the physical meaning of the 2-cocycle $n_2(\mathbf{g}, \mathbf{h})$ that characterizes a $(3+1)$ D bulk SPT is as follows. The $(3+1)$ D fermionic SPT state can be considered to be a superposition of all possible networks of codimension-1 domain walls. A nontrivial $n_2(\mathbf{g}, \mathbf{h})$ means that the codimension-2 junction of three codimension-1 G_b domain walls labeled \mathbf{g} , \mathbf{h} , and $\mathbf{gh} \in G_b$ are decorated with a $(1+1)$ D Kitaev chain if $n_2(\mathbf{g}, \mathbf{h})$ is nontrivial. Therefore, we can consider our system on a 3-dimensional space with boundary, and with a particular choice of domain wall junction, as shown in Fig. 1.

Second, we note that an important property of a $(1+1)$ D Kitaev chain is that it can be coupled to a \mathbb{Z}_2^f gauge field, and we can insert a \mathbb{Z}_2^f symmetry defect (flux). For the Kitaev

chain defined on a ring, the \mathbb{Z}_2^f symmetry defect changes the spin structure; that is, it changes the boundary conditions from periodic to antiperiodic (or vice versa). It is well-known that changing the spin structure changes the fermion parity of the ground state. To see this concretely, we can consider the Hamiltonian for the Kitaev chain:

$$H_K = -i \sum_j (u \gamma'_j \gamma_{j+1} \sigma_{j,j+1} + v \gamma_j \gamma'_j). \quad (127)$$

Here γ_j, γ'_j are independent Majorana fermion operators on site j (i.e., $\{\gamma_j, \gamma_k\} = \{\gamma'_j, \gamma'_k\} = 2\delta_{jk}$ and $\{\gamma_j, \gamma'_k\} = 0$). Note that this system can also be written in terms of complex fermions $c_i = \gamma_i + i\gamma'_i$. Furthermore, we have coupled the system to a \mathbb{Z}_2 gauge field $\sigma_{j,j+1} = \pm 1$ on the links $(j, j+1)$.

Setting $\sigma_{j,j+1} = 1$ for all j , the limits $u/v \ll 1$ and $u/v \gg 1$ realize topologically distinct phases. When the system originates from a model of hopping and pairing of complex fermions c_i , it is natural to identify the $u/v \ll 1$ to be the trivial phase and $u/v \gg 1$ to be the topological phase. A hallmark of the topological phase is that the the fermion parity of the ground state changes in the presence of a \mathbb{Z}_2 symmetry defect on the link $(j, j+1)$ (which corresponds to setting $\sigma_{j,j+1} = -1$). This can be seen easily in the limit $v = 0$, where the ground state is simply $i\gamma'_j \gamma_{j+1} = \sigma_{j,j+1}$. Therefore taking $\sigma_{j,j+1} \rightarrow -\sigma_{j,j+1}$ changes the local fermion parity $i\gamma'_j \gamma_{j+1}$ by one.

Next, let us consider fermion parity vortices at the $(2+1)$ D surface, which are endpoints of fermion parity vortex lines that go into the $(3+1)$ D bulk. Let us consider a process, shown in Fig. 1, where a fermion parity vortex v at the surface encircles the trijunction where the defects \mathbf{g} , \mathbf{h} , \mathbf{gh} all meet. Importantly, to come back to the original configuration, the vortex line in the bulk sweeps across a membrane that must necessarily intersect the Kitaev chain on the codimension-2 junction in the bulk. This changes the fermion parity of the Kitaev chain, which must be compensated for by a change in fermion parity on the vortex line. This can then be interpreted as a transformation $v \rightarrow v \times \psi$ in the surface theory. Therefore we see that we indeed obtain the symmetry action $\bar{\mathbf{gh}}(\mathbf{g}(\bar{\mathbf{h}}v)) \times \bar{v} = o_2(\mathbf{g}, \mathbf{h}) \times v \times \bar{v}$, where here we take $o_2(\mathbf{g}, \mathbf{h}) \in \{1, \psi\}$, which is consistent with the action $\bar{\mathbf{gh}}(\mathbf{g}(\bar{\mathbf{h}}v)) = o_2(\mathbf{g}, \mathbf{h}) \times v$. If $v \in \check{\mathcal{C}}_\sigma$, this process does not permute the vortices but instead changes the fusion channel of two σ -type vortices, which is expected because they are braiding with a Majorana zero mode.

E. Dependence of $[o_2]$ on modular extension

To define the obstruction $[o_2] \in \mathcal{H}^2(G_b, \mathbb{Z}_2)$, we have started with symmetry fractionalization defined on \mathcal{C} , picked a particular modular extension \mathcal{C}_v , and attempted to lift the symmetry action ρ_g to $\check{\rho}_g$. In principle there may be multiple different choices of v for which there exists a lift of ρ_g to $\check{\rho}_g$. As such, $[o_2]$ also has an implicit dependence on v , which we may write as $[o_2^{(v)}]$. This raises the question of how $[o_2^{(v)}]$ may change under a valid change of v , if at all.

In the case where $\ker r = \mathbb{Z}_2$, then as we discussed in the Sec. VID, we expect that $q_{\Gamma^2}([o_2]) \in \mathcal{H}^2(G_b, \mathbb{Z}_2)/\Gamma^2$ should be interpreted as the 't Hooft anomaly of the theory. In other

words, such a theory exists at the (2 + 1)D surface of a (3 + 1)D FSPT characterized by $[n_2] = q_{\Gamma^2}([o_2])$. Since the modular extension ν can be changed by layering a (2 + 1)D invertible topological phase with chiral central charge $c_- = \nu/2$, the bulk (3 + 1)D system should be independent of ν . This leads to the general expectation that $q_{\Gamma^2}([o_2^{(\nu)}])$ is independent of ν (for any valid ν).

Recently an updated version of Ref. [36], motivated by a conjecture of an earlier version of this paper,⁸ showed that for unitary symmetry groups G_b , the following result holds:

$$[o_2^{(\nu'}]) = [\alpha_\psi]^{(\nu'-\nu)[\omega_2]} [o_2^{(\nu)}], \quad (128)$$

where in the above formula we are interpreting $[\omega_2]$ as 0 or 1 depending on whether it is nontrivial in $\mathcal{H}^2(G_b, \mathbb{Z}_2)$. This proves the statement that $q_{\Gamma^2}([o_2^{(\nu)}])$ is independent of ν in the case where G_b is unitary (as long as $\check{\mathcal{C}}$ admits a lift of ρ_g for all \mathbf{g}). In what follows, we will prove that, under some assumptions⁹, $[o_2^{(\nu)}]$ is independent of modular extension in the case where G_b contains antiunitary symmetries. We will then provide additional results (already included in the first version of this paper) about how $[o_2^{(\nu)}]$ depends on modular extension when we only restrict attention to how the symmetry actions permute anyons.

Theorem VI.2. Suppose G_b contains at least one antiunitary symmetry. Assume that $\ker r = \mathbb{Z}_2$ and the nontrivial element of $\ker r$ has a nontrivial permutation action. Then the obstruction $[o_2] \in \mathcal{H}^2(G_b, \ker r)$ is identical for every ν for which there exists, for every $\mathbf{g} \in G_b$, a lift $\check{\rho}_g^{(\nu)} : \check{\mathcal{C}}_\nu \rightarrow \check{\mathcal{C}}_\nu$ of ρ_g .

Proof. Suppose that lifts of $[\rho_g]$ exist for two modular extensions $\check{\mathcal{C}}_{\nu_1}$ and $\check{\mathcal{C}}_{\nu_2}$. Let r_i be the restriction map $r_i : \text{Aut}_{LR}(\check{\mathcal{C}}_{\nu_i}) \rightarrow \text{Aut}_{LR}(\check{\mathcal{C}}_{\nu_i})|_{\mathcal{C}}$. Suppose we have a particular lift $\check{\rho}_g^{(1)}$ of a representative ρ_g to $\check{\mathcal{C}}_{\nu_1}$. Since G_b contains antiunitary symmetries, the only possible $\check{\mathcal{C}}_{\nu_2}$ are obtained from $\check{\mathcal{C}}_{\nu_1}$ by stacking the minimal modular extension $\mathcal{I}(8)$ of $\{1, \psi\}$ with central charge $c_- = 4 \pmod{8}$ and condensing the bound state of the local fermions in the two theories.

The theory $\mathcal{I}(8)$ is equivalent to the 3-fermion topological order. Stacking and condensing such a phase has the effect of changing the topological twist of the fermion parity vortices by a minus sign,

$$\theta_x \rightarrow -\theta_x \quad \text{if } x \in (\check{\mathcal{C}}_{\nu_1})_1, \quad (129)$$

while keeping the fusion rules and S matrix invariant.

Therefore, we can use the same anyon labels for $\check{\mathcal{C}}_{\nu_1}$ and $\check{\mathcal{C}}_{\nu_2}$. In defining the condensation procedure, we need to specify an action of G_b on the 3-fermion topological order $\mathcal{I}(8)$. The only autoequivalences of $\mathcal{I}(8)$ that preserve a choice of physical fermion are the trivial one and the one that permutes

⁸An earlier version of this paper conjectured that $[o_2]$ itself, and not just its image under q_{Γ^2} , would be independent of ν .

⁹The case we do not prove is when $[\alpha_\psi]$ is nonpermuting, i.e., $\check{\Upsilon}_\psi$ violates locality and Υ_ψ respects locality. This last case is proven in v4 of Ref. [36], which was posted around the same time as the revision of this paper containing the current version of Thm. VI.3.

its parity vortices. Let

$$\lambda_1(\mathbf{g}) = \begin{cases} 0 & \mathbf{g} \text{ nonpermuting on } \mathcal{I}(8) \\ 1 & \mathbf{g} \text{ permutes vortices of } \mathcal{I}(8) \end{cases}. \quad (130)$$

Since $\mathcal{I}(8)$ contains an Abelian parity vortex, λ_1 must be a group homomorphism $\lambda_1 : G_b \rightarrow \mathbb{Z}_2$. The permutation action of $\check{\rho}_g^{(2)}$ on anyon labels is therefore the same as that of $\alpha_\psi^{\lambda_1(\mathbf{g})} \check{\rho}_g^{(1)}$. Accordingly, the permutation action of $o_2^{(2)}(\mathbf{g}, \mathbf{h})$ is the same as that of $o_2^{(1)}(\mathbf{g}, \mathbf{h}) \alpha_\psi^{d\lambda_1(\mathbf{g}, \mathbf{h})}$. Since we assumed that the only nontrivial element of $\ker r$ is permuting, the permutation action of $o_2^{(2)}$ determines it as an element of $\ker r$. Hence, as cohomology classes, $[o_2^{(2)}] = [o_2^{(1)}]$. ■

In the remaining part of this section, we will provide a number of results about the dependence of $[o_2^{(\nu)}]$ on ν when we only restrict our attention to how ρ and o_2 permute the anyons. Hence we will define $P(\check{\mathcal{C}})$ to be the group of anyon permutations of $\check{\mathcal{C}}$ that preserve the modular S and T matrices of $\check{\mathcal{C}}$. Furthermore, for $\rho, o_2 \in \text{Aut}_{LR}(\check{\mathcal{C}})$, we will denote $\underline{\rho}, \underline{o_2} \in P(\check{\mathcal{C}})$ their respective permutation actions (and similarly for $P(\mathcal{C})$). Note that if we have a lift $\check{\rho} \in P(\check{\mathcal{C}})$ of $\underline{\rho} \in P(\mathcal{C})$ then $\check{\rho}$ defines a permutation $\underline{o_2}$ by Eq. (84).

Proposition VI.1. Let $\check{\rho}_g^{(1)}$ be a lift of ρ_g to $\text{Aut}_{LR}(\check{\mathcal{C}}_{\nu_1})$, which defines the obstruction $o_2^{(1)} \in \mathcal{H}^2(G_b, \ker r)$. If G_b is unitary, then there is a permutation $\check{\rho}_g^{(2)} \in P(\check{\mathcal{C}}_{\nu_2})$, which lifts ρ_g for any ν_2 . If $\delta\nu = \nu_2 - \nu_1$ is even, then $\underline{o_2^{(2)}} = \underline{o_2^{(1)}}$. If $\delta\nu$ is odd, then $\underline{o_2^{(2)}}$ is the trivial permutation.

The proof is quite technical, so we briefly sketch it here and defer the details to Appendix B. The main idea is to use the 16-fold way theorem [34], which states that a generic minimal modular extension may be derived from a given $\check{\mathcal{C}}_{\nu_1}$ by layering $\check{\mathcal{C}}_{\nu_1}$ with a minimal modular extension $\mathcal{I}(\delta\nu)$ of $\{1, \psi\}$ and condensing the bound state of the fermions in the two layers. Using results on anyon condensation [33], we can derive the modular data of a generic minimal modular extension from the given one $\check{\mathcal{C}}_{\nu_1}$. This step is straightforward when $\delta\nu$ is even because $\mathcal{I}(\delta\nu)$ is Abelian, but it is quite involved when $\delta\nu$ is odd so that $\mathcal{I}(\delta\nu)$ is non-Abelian. From the new modular data, we can explicitly construct an element of $P(\check{\mathcal{C}}_{\nu_2})$ for each $\check{\rho}_g^{(1)}$ and then calculate the permutation action of compositions of these permutations.

Proposition VI.1 has the rather surprising corollary:

Corollary VI.1. Suppose that in every minimal modular extension of \mathcal{C} , every lift of ρ_g from $P(\mathcal{C})$ to $P(\check{\mathcal{C}})$ defines a lift of the autoequivalence class $[\rho_g]$ from $\text{Aut}_{LR}(\mathcal{C})$ to $\text{Aut}_{LR}(\check{\mathcal{C}})$. Further assume that the only possible nonpermuting element of $\ker r$ is $[\check{\Upsilon}_\psi]$. Then if Υ_ψ violates locality, $[o_2^{(\nu)}] = +1$ is trivial for all ν . If Υ_ψ respects locality, then $[o_2^{(\nu)}] = +1$ is trivial for the eight minimal modular extensions with only ν -type vortices.

Proof. (Corollary VI.1) First consider the case where Υ_ψ violates locality. Then $[\check{\Upsilon}_\psi]$ is not in $\ker r$ for any minimal modular extension $\check{\mathcal{C}}_{\nu_2}$. But according to Proposition VI.1, for $\nu_2 - \nu_1$ odd, $o_2^{(2)}$ has trivial permutation action, so it must be the identity as an autoequivalence. Now repeat the argument

starting from the permutation action $\tilde{\rho}_g^{(2)}$ to see that for $v_3 - v_2$ odd, that is, $v_3 - v_1$ even, we can again construct a $\tilde{\rho}_g^{(3)}$ such that $o_2^{(3)}$ is also trivial.

If Υ_ψ respects locality, then by a property discussed in Sec. IV C, half of the modular extensions of \mathcal{C} contain only σ -type fermion parity vortices and half contain only v -type fermion parity vortices. We may assume without loss of generality that $\check{\mathcal{C}}_{v_1}$ has only σ -type fermion parity vortices; if $\check{\mathcal{C}}_{v_1}$ contains only v -type fermion parity vortices, use Proposition VI.1 to construct a symmetry action on $\check{\mathcal{C}}_{v_1+1}$ and reindex $v_1 \rightarrow v_1 - 1 \pmod{16}$. Now apply the δv odd case of Proposition VI.1. Then $o_2^{(v_1+1)}$ is a nonpermuting element of $\ker r$. Hence $o_2^{(v_1+1)}$ is the identity; the reason is that $\check{\mathcal{C}}_{v_1+1}$ has only v -type vortices, so all nontrivial elements of $\ker r$ are permuting (see Sec. V B). Using the δv even case of Proposition VI.1, we obtain a nonanomalous lift for all eight modular extensions $\check{\mathcal{C}}_v$ that have $v = v_1 + 1 \pmod{2}$. The above argument does not extend to $v = v_1 \pmod{2}$, since $\ker r$ contains a nontrivial nonpermuting element Υ_ψ .

The interplay of Eq. (128) and Corollary VI.1 is quite interesting. Consider a theory \mathcal{C} , which obeys the assumptions of Corollary VI.1. If Υ_ψ respects locality, then Eq. (128) and Corollary VI.1 combine to prove that the minimal modular extensions of \mathcal{C} with only v -type vortices have no $[o_2]$ obstruction and the minimal modular extensions with only σ -type vortices have $[o_2] = [\omega_2]$.

On the other hand, if Υ_ψ violates locality, then these two statements combine to prove that $[\omega_2] = 0$. Therefore, if Υ_ψ violates locality and the theory has unobstructed fractionalization with $[\omega_2] \neq 0$, then the assumptions of either Eq. (128) (that $\ker r = \mathbb{Z}_2$) or Corollary VI.1 must be violated. We expect in general, given Conjecture VI.1 that $\ker r = \mathbb{Z}_2$, that the latter occurs.

F. $\text{SO}(3)_3 \times \text{SO}(3)_3$: an example with $k > 1$

As discussed in Sec. V B 5, we cannot presently exclude the possibility that $\ker r$ is different from \mathbb{Z}_2 . This might happen because the number of blocks in the $v - v$ portion of the S matrix of $\check{\mathcal{C}}$ is $k > 1$, or because certain permutations of $\check{\mathcal{C}}$ that preserve the modular data do not uniquely determine an element of $\text{Aut}(\check{\mathcal{C}})$. If $\ker r \neq \mathbb{Z}_2$, it is not immediately obvious how to connect the 't Hooft anomaly $[n_2] \in \mathcal{H}^2(G_b, \mathbb{Z}_2)/\Gamma^2$ to the obstruction $[o_2] \in \mathcal{H}^2(G_b, \ker r)$. We do know, however, that $\ker r$ contains a \mathbb{Z}_2 subgroup generated by $[\alpha_\psi]$.

Our anomaly arguments in Secs. V C and V D are quite general and physical, so we expect that there should be a fully general connection between the 't Hooft anomaly $[n_2]$ and the $\mathcal{H}^2(G_b, \ker r)$ obstruction. However, our arguments only strictly hold when $\ker r = \mathbb{Z}_2$. Assuming Conjecture VI.1 neatly allows $[o_2]$ and $[n_2]$ to always be related. However, if there exists a theory $\check{\mathcal{C}}$ has $k > 1$, then there is tension; at the level of permutations, Theorem VI.1 suggests that $\ker r$ may be larger than \mathbb{Z}_2 .

To explore this issue, we now present an explicit example with $k = 2$ and which has a nontrivial 't Hooft anomaly $[n_2]$. We argue for this example that if Theorem VI.1 applies, that is, if every permutation allowed by Theorem I.3

TABLE I. UMTC data and action of time-reversal symmetry for $\check{\mathcal{C}} = \text{SU}(2)_6 \boxtimes \text{SU}(2)_6 \boxtimes \text{Ising}_{-9/2}/\{\psi\psi \sim 1\}$. The parity vortices $(1, 3, \sigma)_\pm$ are v type and interchanged by fusion with ψ ; likewise for $(3, 1, \sigma)_\pm$. The parity vortices $(1, 1, \sigma)$ and $(3, 3, \sigma)$ are σ type. The notation \pm_1 and \pm_2 indicate two independent choices of signs in the action of time reversal. All four such permutations complex conjugate the modular data.

Label	θ_a	d_a	\mathbf{t}_a
(0,0,0)	1	1	(0,0,0)
(2,0,0)	i	$1 + \sqrt{2}$	(4,0,0)
(0,2,0)	i	$1 + \sqrt{2}$	(0,4,0)
(4,0,0)	$-i$	$1 + \sqrt{2}$	(2,0,0)
(0,4,0)	$-i$	$1 + \sqrt{2}$	(0,2,0)
(2,2,0)	-1	$(1 + \sqrt{2})^2$	(2,2,0)
(2,4,0)	1	$(1 + \sqrt{2})^2$	(2,4,0)
(6,0,0)	-1	1	(6,0,0)
$(1, 3, \sigma)_+$	1	$2 + \sqrt{2}$	$(3, 1, \sigma)_{\pm_1}$
$(1, 3, \sigma)_-$	1	$2 + \sqrt{2}$	$(3, 1, \sigma)_{\mp_1}$
$(3, 1, \sigma)_+$	1	$2 + \sqrt{2}$	$(1, 3, \sigma)_{\pm_2}$
$(3, 1, \sigma)_-$	1	$2 + \sqrt{2}$	$(1, 3, \sigma)_{\mp_2}$
$(1, 1, \sigma)$	$e^{5\pi i/4}$	$2 + 2\sqrt{2}$	$(3, 3, \sigma)$
$(3, 3, \sigma)$	$e^{3\pi i/4}$	$2 + 2\sqrt{2}$	$(1, 1, \sigma)$

uniquely defines an element of $\text{Aut}(\check{\mathcal{C}})$ and $\text{Aut}(\mathcal{C})$ contains no nontrivial nonpermuting elements, then the obstruction $[o_2] \in \mathcal{H}^2(G_b, \ker r)$ is trivial. Since $[n_2]$ is nontrivial in this example, there would be no connection between $[n_2]$ and $[o_2]$. This is in tension with the physical arguments for Conjecture VI.1. We give a well-defined but difficult in practice way to test Conjecture VI.1 in this case.

Consider the theory $\mathcal{C} = \text{SO}(3)_3 \boxtimes \text{SO}(3)_3/\{\psi\psi \sim 1\}$, where the quotient means we condense the bound state of the transparent fermions from each copy of the theory. We take $G_b = \mathbb{Z}_2^T$ and $G_f = \mathbb{Z}_4^{T,f}$. It is easy to check that this theory is super-modular. For this G_f , the $(3+1)$ D FSPT classification is given by an element $\mu \in \mathbb{Z}_{16}$, and it is known that $\text{SO}(3)_3$ has a $\mu = 3$ anomaly [25]. Hence \mathcal{C} has a $\mu = 6$ anomaly, which is associated with the nontrivial element $[n_2] \in \mathcal{H}^2(\mathbb{Z}_2^T, \mathbb{Z}_2) = \mathbb{Z}_2$.

The minimal modular extension we consider is $\check{\mathcal{C}} = \text{SU}(2)_6 \boxtimes \text{SU}(2)_6 \boxtimes \mathcal{I}(-9)/\{\psi\psi \sim 1\}$, where $\mathcal{I}(-9)$ is the $c_- = -9/2$ minimal modular extension of $\{1, \psi\}$ and the quotient means we condense all pairs of Abelian fermions from the different theories. The particle content, topological twists, quantum dimensions, and possible actions of \mathbf{T} are listed in Table I, while the S matrix is given in Appendix C. The data were derived from a slight generalization of the results of [33]; the ungeneralized results are reviewed in Appendix A.

Examining the set of permutations, which preserve the modular data, one would naively conclude that $\ker r = \mathbb{Z}_2^2$ because there are $k = 2$ distinct blocks of the $\check{\mathcal{C}}_v$ part of the S matrix, namely one block formed by $(1, 3, \sigma)_\pm$ and one block formed by $(3, 1, \sigma)_\pm$. There are thus four possible lifts of the permutation action of $\rho_{\mathbf{T}}$ to $\check{\mathcal{C}}$ that preserve the modular data of $\check{\mathcal{C}}$. Two form a group homomorphism $\mathbb{Z}_2^T \rightarrow \text{Aut}_{LR}(\mathcal{C})$; their

actions are given by a choice of signs:

$$\begin{aligned} (1, 3, \sigma)_+ &\leftrightarrow (3, 1, \sigma)_\pm \\ (1, 3, \sigma)_- &\leftrightarrow (3, 1, \sigma)_\mp \\ (1, 1, \sigma) &\leftrightarrow (3, 3, \sigma), \end{aligned} \quad (131)$$

where the particle content is labeled as in Appendix C. The other two possible lifts of ρ_T form a group homomorphism $\mathbb{Z}_4^T \rightarrow \text{Aut}_{LR}(\check{\mathcal{C}})$:

$$\begin{aligned} (1, 3, \sigma)_+ &\rightarrow (3, 1, \sigma)_\pm \rightarrow (1, 3, \sigma)_- \rightarrow (3, 1, \sigma)_\mp \\ &\rightarrow (1, 3, \sigma)_+(1, 1, \sigma) \leftrightarrow (3, 3, \sigma). \end{aligned} \quad (132)$$

Within each pair, the two possible lifts differ by the permutation action of α_ψ .

Choosing a lift in Eq. (131), the permutation action of $\alpha_2(\mathbf{g}, \mathbf{h}) \in \ker r$ is trivial. One can check that Υ_ψ does not respect locality in this theory, so $\check{\Upsilon}_\psi$ is not in $\ker r$; accordingly, if we assume that permutations that preserve the modular data uniquely determine an element of $\text{Aut}(\check{\mathcal{C}})$, $[\alpha_2]$ is trivial. However, as argued above, $[n_2]$ is nontrivial. We therefore expect that permutations do not uniquely determine an element of $\text{Aut}(\check{\mathcal{C}})$ in this theory.

In the cases in Eq. (132) where the lift is not a group homomorphism $\mathbb{Z}_2^T \rightarrow \text{Aut}_{LR}(\check{\mathcal{C}})$, we see explicitly that, as permutations,

$$\check{\rho}_T^2 = \alpha_\psi. \quad (133)$$

Therefore, if only the permutations in Eq. (132) define valid autoequivalences of $\check{\mathcal{C}}$ and $\ker r = \mathbb{Z}_2$, we would indeed obtain $[\alpha_2] = [n_2]$ for $\mathcal{C} = \text{SO}(3)_3^2$ (recall that since G_b contains antiunitary symmetries, Γ^2 is trivial so q_{Γ^2} is the identity map). We speculate that this is the case; in order to check this speculation, one would need to solve for the F - and R -symbols of $\check{\mathcal{C}}$, which is a nontrivial task, and then directly attempt to solve for the U symbols for each permutation action. We expect that a solution for the U symbols exists only for the permutations in Eq. (132).

VII. $\mathcal{H}^3(G_b, \mathbb{Z}_2)$ OBSTRUCTION

Suppose that the $\mathcal{H}^2(G_b, \ker r)$ obstruction vanishes, so that we may lift a group homomorphism $[\rho_g] : G_b \rightarrow \text{Aut}_{LR}(\mathcal{C})$ to a group homomorphism $[\check{\rho}_g] : G_b \rightarrow \text{Aut}_{LR}(\check{\mathcal{C}})$. Given a symmetry fractionalization pattern on \mathcal{C} , we ask whether or not that fractionalization pattern can be lifted to $\check{\mathcal{C}}$. We will show that there is an obstruction to this process valued in $\mathcal{H}^3(G_b, \mathbb{Z}_2)$. Our discussion will very similar to that of Ref. [32], but [32] assumed $G_f = G_b \times \mathbb{Z}_2$ and made some technical assumptions that are known to fail in certain cases. We will use our general understanding of fermionic symmetry fractionalization to remove those technical assumptions.

Note also that $[\check{\rho}_g]$ defines a G_b symmetry action on a UMTG $\check{\mathcal{C}}$, so there may be an obstruction to localizing $[\check{\rho}_g]$, that is, to finding any symmetry fractionalization on $\check{\mathcal{C}}$ irrespective of whether it matches the symmetry fractionalization on \mathcal{C} . This obstruction is valued in $\mathcal{H}^3(G_b, \check{\mathcal{A}})$ and can be computed in the standard way for bosonic SETs, see [1]. We

will show that this obstruction is in fact determined by the $\mathcal{H}^3(G_b, \mathbb{Z}_2)$ obstruction.

A. $\mathcal{H}^3(G_b, \mathbb{Z}_2)$ anomaly

We start with symmetry fractionalization data, which we choose to characterize by $\omega_a(\mathbf{g}, \mathbf{h}) \in C^2(G_b, K(\mathcal{C}))$ satisfying Eq. (36). In the present language, $\Omega_a \in Z^3(G_b, K(\mathcal{C}))$. The gauge freedom ν_a appearing in Eq. (40) is an element of $K(\mathcal{C})$, so only $[\Omega_a] \in \mathcal{H}^3(G_b, K(\mathcal{C}))$ is gauge invariant.

We are also given a lift $[\check{\rho}]$ of the symmetry action ρ to $\check{\mathcal{C}}$. A representative $\check{\rho}_g$ determines $\check{\kappa}_{\mathbf{g}, \mathbf{h}}$ via

$$\check{\kappa}_{\mathbf{g}, \mathbf{h}} \check{\rho}_g \check{\rho}_{\mathbf{h}} = \check{\rho}_{\mathbf{g} \mathbf{h}}. \quad (134)$$

Restricting this equation to \mathcal{C} , we find that

$$r(\check{\kappa}_{\mathbf{g}, \mathbf{h}}) = \kappa_{\mathbf{g}, \mathbf{h}}. \quad (135)$$

Hence, given a decomposition $\check{\beta}_a(\mathbf{g}, \mathbf{h})$ of $\check{\kappa}_{\mathbf{g}, \mathbf{h}}$ as a natural isomorphism, we can simply restrict these $\check{\beta}_a$ to \mathcal{C} to obtain a valid gauge choice for the decomposition $\beta_a(\mathbf{g}, \mathbf{h})$ of $\kappa_{\mathbf{g}, \mathbf{h}}$. We will always work in this gauge where $\check{\beta}_a$ lifts β_a . In this gauge the function $\check{\Omega}_a(\mathbf{g}, \mathbf{h}, \mathbf{k}) \in Z^3(G_b, K(\check{\mathcal{C}}))$ lifts $\Omega_a(\mathbf{g}, \mathbf{h}, \mathbf{k}) \in Z^3(G_b, K(\mathcal{C}))$.

Note the logic here—we are using the a gauge-fixing (of the ν type) of $\check{\beta}_a$ to determine a gauge-fixing (again of the ν type) of β_a . In general, not every gauge choice on \mathcal{C} allows β_a and Ω_a to be lifted to $\check{\mathcal{C}}$, specifically when $\check{\Upsilon}_\psi$ violates locality but Υ_ψ respects locality.

We now ask whether or not there exists a lift $\check{\omega}_a \in C^2(G_b, K(\check{\mathcal{C}}))$ such that

$$\check{\Omega}_a = d\check{\omega}_a \quad (136)$$

for all $a \in \check{\mathcal{C}}$ and such that $r(\check{\omega}_a) = \omega_a$.

For the moment, we assume that at least one lift $\check{\omega}_a \in C^2(G_b, K(\check{\mathcal{C}}))$ of ω_a exists. We will prove that such a lift exists later; the reason depends on whether Υ_ψ and $\check{\Upsilon}_\psi$ respect or violate locality.

We next claim that there always exists exactly two lifts $\check{\omega}_a$ of ω_a [which need not, *a priori*, satisfy Eq. (136)]. There always exists an element $p_a \in K(\check{\mathcal{C}})$ defined by

$$p_a = M_{a, \psi} = \begin{cases} 1 & \text{if } a \in \check{\mathcal{C}}_0 \\ -1 & \text{if } a \in \check{\mathcal{C}}_1 \end{cases}. \quad (137)$$

Hence, given a lift $\check{\omega}_a(\mathbf{g}, \mathbf{h})$, there is always another one $\check{\omega}_a p_a^{\alpha(\mathbf{g}, \mathbf{h})}$ with $\alpha \in C^2(G_b, \mathbb{Z}_2)$ (here $\mathbb{Z}_2 = \{0, 1\}$). Also, we may always write

$$\check{\Omega}_a = M_{a, \check{\mathbf{g}}} \quad (138)$$

$$\check{\omega}_a = M_{a, \check{\mathbf{w}}} \quad (139)$$

for some $\check{\mathbf{w}}, \check{\mathbf{g}} \in \check{\mathcal{A}}$ because $\check{\mathcal{C}}$ is modular; the p_a freedom mentioned above amounts to changing $\check{\mathbf{w}} \rightarrow \check{\mathbf{w}} \times \psi$. By super-modularity of \mathcal{C} , the only element of $\check{\mathcal{C}}$ that braids trivially with all of \mathcal{C}_0 is ψ , so the two lifts defined by $\check{\mathbf{w}}$ and $\check{\mathbf{w}} \times \psi$ are the only ones that restrict to ω_a on \mathcal{C} . Hence, if a lift exists, there are exactly two such lifts. Choose one of these lifts, which we call $\check{\omega}_a$. We must ask if Eq. (136) is satisfied.

Define

$$\tilde{\Omega}_a(\mathbf{g}, \mathbf{h}, \mathbf{k}) = \check{\Omega}_a(\mathbf{g}, \mathbf{h}, \mathbf{k})(d\check{\omega}_a(\mathbf{g}, \mathbf{h}, \mathbf{k}))^{-1}. \quad (140)$$

Certainly $\tilde{\Omega}_a \in Z^3(G_b, K(\check{\mathcal{C}}))$ since both $\check{\Omega}_a$ and $d\check{\omega}_a$ are in $Z^3(G_b, K(\check{\mathcal{C}}))$. Also, Eq. (136) is satisfied for all $a \in \check{\mathcal{C}}$ if and only if $\tilde{\Omega}_a = 1 \in Z^3(G_b, K(\check{\mathcal{C}}))$. This will not generally be the case; however, by definition Eq. (136) is satisfied for all $a \in \mathcal{C}$. Hence $\tilde{\Omega}_a = 1$ for all $a \in \mathcal{C}$ and therefore

$$\tilde{\Omega}_a(\mathbf{g}, \mathbf{h}, \mathbf{k}) = M_{a, o_3(\mathbf{g}, \mathbf{h}, \mathbf{k})} \quad (141)$$

where one can check that

$$o_3(\mathbf{g}, \mathbf{h}, \mathbf{k}) = \check{\mathfrak{G}}(\mathbf{g}, \mathbf{h}, \mathbf{k}) \times \overline{d\check{\omega}(\mathbf{g}, \mathbf{h}, \mathbf{k})} \in \{1, \psi\} \cong \mathbb{Z}_2. \quad (142)$$

By straightforward computation, $d\tilde{\Omega}_a = 1$, which implies $do_3 = 1$. We have the freedom to choose a different lift $\check{\omega}_a(\mathbf{g}, \mathbf{h})p_a^{\alpha(\mathbf{g}, \mathbf{h})}$, which modifies $\tilde{\Omega}_a \rightarrow \tilde{\Omega}_a p_a^{d\alpha}$, that is, it changes o_3 by a \mathbb{Z}_2 -coboundary. Therefore, $[o_3] \in \mathcal{H}^3(G_b, \mathbb{Z}_2)$ is a well-defined cohomology class independent of the choice of lift $\check{\omega}_a$. Also, if $[o_3] = 1$, then there exists a representative $\tilde{\Omega}_a = 1$, i.e., some lift satisfies Eq. (136). We therefore see that $[o_3] \in \mathcal{H}^3(G_b, \mathbb{Z}_2)$ is the obstruction to lifting symmetry fractionalization from \mathcal{C} to all of $\check{\mathcal{C}}$.

Note that we do not need to make any explicit reference to G_f , which is encoded via the constraint $\eta_\psi = \omega_2$. The lifted symmetry fractionalization data automatically agrees with the symmetry fractionalization data on \mathcal{C} and therefore also obeys the constraint.

The above argument is essentially a reformulation of that of Ref. [32]. Both our argument and that of Ref. [32] rely on an assumption that a lift $\check{\omega}_a$ always exists. At the cochain level (i.e. in an arbitrary gauge), such a lift need not exist; for example, with $\mathcal{C} = \{1, \psi\}$ and $\check{\mathcal{C}} = \text{Ising} = \{1, \psi\}$, there is no lift of the function $\omega_a \in K(\mathcal{C})$ with $\omega_1 = 1, \omega_\psi = -1$ to $K(\check{\mathcal{C}})$; Ref. [32] observes this failure but does not go further. We now carefully prove our assumptions, specifically that there always exists some gauge in which both Ω_a and ω_a lift, proving that the above definition of $[o_3]$ is always valid.

We consider different cases depending on whether $\check{\Upsilon}_\psi$ and Υ_ψ respect locality. We will heavily rely on results from [35] summarized in Sec. IV C that relate whether or not Υ_ψ respects locality to various properties of \mathcal{C} and $\check{\mathcal{C}}$. One might ask separately if there could be an $\mathcal{H}^3(G_b, \check{\mathcal{A}})$ obstruction to defining *any* G_b symmetry fractionalization pattern on $\check{\mathcal{C}}$, whether it agrees with the symmetry fractionalization on \mathcal{C} or not. Obviously if the $\mathcal{H}^3(G_b, \mathbb{Z}_2)$ obstruction $[\tilde{\Omega}]$ vanishes, there cannot be any such obstruction. We claim more generally that $[\tilde{\Omega}]$ actually determines the $\mathcal{H}^3(G_b, \check{\mathcal{A}})$ obstruction, and in the following we explain the relationship in each case.

1. Case: Υ_ψ violates locality

If Υ_ψ violates locality, then $K(\mathcal{C}) = K_+(\mathcal{C})$, that is, every set of phases, which obey the fusion rules on \mathcal{C} are $+1$ on the fermion, and also $\check{\mathcal{A}} = \mathcal{A}$. Hence Ω_a and ω_a are in $K_+(\mathcal{C})$, so

$$\begin{aligned} \Omega_a &= M_{a, \mathfrak{G}} \\ \omega_a &= M_{a, \mathfrak{w}} \end{aligned} \quad (143)$$

for $\mathfrak{G} \in Z^3(G_b, \mathcal{A}/\{1, \psi\})$ and $\mathfrak{w} \in C^2(G_b, \mathcal{A}/\{1, \psi\})$. We can then lift Ω_a and ω_a to $\check{\mathcal{C}}$ straightforwardly by choosing one of two lifts $\check{\mathfrak{G}}$ of \mathfrak{G} to $Z^3(G_b, \mathcal{A})$ and one of two lifts $\check{\mathfrak{w}}$ of \mathfrak{w} to $C^2(G_b, \mathcal{A})$, which allows us to extend Eq. (143) to all $a \in \check{\mathcal{C}}$. The lifted symmetry action $[\check{\rho}]$ on $\check{\mathcal{C}}$, which we have already taken as a given, determines which of the two lifts $\tilde{\Omega}_a$ we must use, while there is freedom in which lift of ω_a we choose using p_a , as discussed previously.

Having proven that a lift exists, we now discuss the $\mathcal{H}^3(G_b, K(\check{\mathcal{C}})) \cong \mathcal{H}^3(G_b, \check{\mathcal{A}})$ obstruction. The short exact sequence

$$1 \rightarrow \mathbb{Z}_2 = \{1, \psi\} \rightarrow \mathcal{A} \rightarrow \mathcal{A}/\{1, \psi\} \rightarrow 1 \quad (144)$$

induces a map

$$i : \mathcal{H}^3(G_b, \mathbb{Z}_2) \rightarrow \mathcal{H}^3(G_b, \mathcal{A}) \quad (145)$$

that is part of the long exact sequence

$$\cdots \mathcal{H}^2(G_b, \mathcal{A}/\{1, \psi\}) \xrightarrow{\delta} \mathcal{H}^3(G_b, \mathbb{Z}_2) \xrightarrow{i} \mathcal{H}^3(G_b, \mathcal{A}) \xrightarrow{q} \mathcal{H}^3(G_b, \mathcal{A}/\{1, \psi\}) \rightarrow \cdots \quad (146)$$

The fact that Υ_ψ violates locality means that no minimal modular extension of \mathcal{C} contains an Abelian parity vortex; therefore, $\check{\mathcal{A}} = \mathcal{A}$. By construction, then, $i([o_3]) = [\check{\mathfrak{G}}]$, which is the actual $\mathcal{H}^3(G_b, \check{\mathcal{A}})$ obstruction to symmetry localization.

Therefore, the $\mathcal{H}^3(G_b, \mathbb{Z}_2)$ obstruction actually determines the $\mathcal{H}^3(G_b, \check{\mathcal{A}})$ obstruction of the lifted theory in the following sense. One possibility is that $[o_3] \in \ker i$, in which case symmetry localization is not obstructed on $\check{\mathcal{C}}$ even if the lift of the particular symmetry fractionalization pattern in question is obstructed. The other possibility is that $i([o_3])$ is nontrivial, in which case there is an obstruction to lifting the given symmetry fractionalization pattern on \mathcal{C} to $\check{\mathcal{C}}$ simply because there is no consistent symmetry fractionalization pattern on $\check{\mathcal{C}}$ at all.

We note that the fact that symmetry fractionalization is unobstructed on \mathcal{C} , i.e., that $[\check{\mathfrak{G}}] = q([\mathfrak{G}])$ is trivial, means that $[\check{\mathfrak{G}}] \in \ker q = \text{im } i$, which is why we could always find $[o_3] \in \mathcal{H}^3(G_b, \mathbb{Z}_2)$ with $i([o_3]) = [\check{\mathfrak{G}}]$.

2. Case: Υ_ψ and $\check{\Upsilon}_\psi$ respect locality

We show that every element of $K(\mathcal{C})$ has a lift to $K(\check{\mathcal{C}})$, which implies that Ω_a and ω_a lift to $K(\check{\mathcal{C}})$. First suppose that $\zeta_a \in K_+(\mathcal{C})$; then

$$\zeta_a = M_{a, x} \quad (147)$$

for some $x \in \mathcal{A}/\{1, \psi\}$. By simply choosing a representative of $x \in \mathcal{A}$, we obtain a lift of ζ_a by extending Eq. (147) to all $a \in \check{\mathcal{C}}$. As usual, there is another lift related by sending $x \rightarrow x \times \psi$, or equivalently by modifying the lift of ζ_a by p_a .

Suppose instead that $\zeta_a \in K_-(\mathcal{C})$. Since $\check{\Upsilon}_\psi$ respects locality, $\check{\mathcal{C}}$ contains an Abelian fermion parity vortex; call such a parity vortex v , and define $\lambda_a = M_{a, v} \in K_-(\mathcal{C})$ and $\check{\lambda}_a = M_{a, v} \in K_-(\check{\mathcal{C}})$. Then $\lambda_a^{-1} \zeta_a \in K_+(\mathcal{C})$ and, as we have already shown, has exactly two possible lifts $(\check{\lambda}\zeta)_a = M_{a, \check{x}}$, where the two possible choices of $\check{x} \in \mathcal{A}$ differ by a fermion, to $K_+(\check{\mathcal{C}})$. Hence there are exactly two (distinct) possible lifts

$\check{\lambda}_a(\check{\lambda}^{-1}\zeta)_a = M_{a,\check{x}\times v}$ of ζ_a to $K_-(\check{\mathcal{C}})$. These two lifts differ by changing $\check{x} \times v$ by a fermion, or equivalently by modifying the lift by p_a .

This proves that every element of $K(\mathcal{C})$ lifts to $K(\check{\mathcal{C}})$, as desired.

Our argument that the $\mathcal{H}^3(G_b, \mathbb{Z}_2)$ obstruction determines the $\mathcal{H}^3(G_b, \check{\mathcal{A}})$ obstruction for general symmetry fractionalization on $\check{\mathcal{C}}$ carries through from the case where Υ_ψ violates locality. The only difference is that we start from a short exact sequence

$$1 \rightarrow \mathbb{Z}_2 \rightarrow K(\check{\mathcal{C}}) = \check{\mathcal{A}} \rightarrow K(\mathcal{C}) \rightarrow 1. \quad (148)$$

Using the fact that the (bosonic) obstruction to symmetry fractionalization on \mathcal{C} is valued in $\mathcal{H}^3(G_b, K(\mathcal{C}))$, the rest of the argument carries through *mutatis mutandis*.

3. Case: Υ_ψ respects locality but $\check{\Upsilon}_\psi$ does not

This case is a bit more subtle than the others because $\check{\mathcal{C}}$ does not contain an Abelian fermion parity vortex; hence $\check{\mathcal{A}} = \mathcal{A}$, and every element of $K(\check{\mathcal{C}})$ restricts to an element of $K_+(\mathcal{C})$. As such, $K_+(\check{\mathcal{C}})/\mathbb{Z}_2 = K_+(\mathcal{C})$, where the \mathbb{Z}_2 subgroup is generated by $p_a = M_{a,\psi}$.

In general, Ω_a and ω_a take values in $K(\mathcal{C})$, not necessarily in $K_+(\mathcal{C})$, so they need not have lifts to $K(\check{\mathcal{C}})$. However, our gauge fixing on \mathcal{C} guaranteed that $\check{\Omega}_a$ lifts Ω_a . We just need to check that in this gauge, ω_a also has a lift. By Eq. (99), $[\check{\kappa}_{\mathbf{g},\mathbf{h}}] = [\check{\Upsilon}_\psi]^{\check{\omega}_2(\mathbf{g},\mathbf{h})}$. Hence

$$\check{\beta}_\psi(\mathbf{g}, \mathbf{h}) = \beta_\psi(\mathbf{g}, \mathbf{h}) = \omega_2(\mathbf{g}, \mathbf{h}) \quad (149)$$

in this gauge. Since $\eta_\psi(\mathbf{g}, \mathbf{h}) = \omega_2$ as well, we conclude using Eq. (37) that

$$\omega_\psi(\mathbf{g}, \mathbf{h}) = +1, \quad (150)$$

that is, $\omega_a \in K_+(\mathcal{C})$. Hence Eq. (143) applies with $\mathbf{w} \in C^3(G_b, \mathcal{A}/\{1, \psi\})$. As in the previous cases, then, we may lift ω_a to $K(\check{\mathcal{C}})$ by extending Eq. (143) to all of $\check{\mathcal{C}}$, with a $C^2(G_b, \mathbb{Z}_2)$ choice for $\check{\omega}_a$.

The argument that the $\mathcal{H}^3(G_b, \mathbb{Z}_2)$ obstruction determines the $\mathcal{H}^3(G_b, \check{\mathcal{A}})$ obstruction is identical to the case in which Υ_ψ violates locality; the argument in the latter case really only used the fact that $\check{\mathcal{A}} = \mathcal{A}$ for the minimal modular extension in question, which is true whenever $\check{\Upsilon}_\psi$ violates locality.

B. Dependence on symmetry fractionalization class

We can see explicitly how the $\mathcal{H}^3(G_b, \mathbb{Z}_2)$ anomaly depends on the symmetry fractionalization class on \mathcal{C} as follows. As discussed in [35], symmetry fractionalization on \mathcal{C} is an $\mathcal{H}^2(G_b, \mathcal{A}/\{1, \psi\})$ torsor, that is, changing symmetry fractionalization classes on \mathcal{C} amounts to shifting $\mathbf{w}(\mathbf{g}, \mathbf{h}) \rightarrow \mathbf{w}'(\mathbf{g}, \mathbf{h}) = \mathbf{w}(\mathbf{g}, \mathbf{h}) \times \mathbf{t}(\mathbf{g}, \mathbf{h})$ with $\mathbf{t} \in Z^2(G_b, \mathcal{A}/\{1, \psi\})$. The new symmetry fractionalization pattern depends only on the cohomology class $[\mathbf{t}]$. Given a particular lift $\check{\mathbf{w}}$ of \mathbf{w} , then, we can pick a particular lift $\check{\mathbf{t}}$ of \mathbf{t} to obtain a lift $\check{\mathbf{w}}'$ of \mathbf{w}' . A representative of the obstruction class $[\check{\mathbf{w}}'_3]$ corresponding to $\check{\mathbf{w}}'$ is thus given by

$$\check{\mathbf{w}}'_3 = \check{\mathbf{G}} \times \overline{d\check{\mathbf{w}}'} = o_3 \times \overline{d\check{\mathbf{t}}}. \quad (151)$$

Changing \mathbf{t} by a coboundary in $\mathcal{A}/\{1, \psi\}$ leaves dt invariant and thus can affect o_3 only through the choice of lift to $\check{\mathbf{t}}$. Changing the lift $\check{\mathbf{t}}$ changes o_3 by a \mathbb{Z}_2 coboundary, so $[\check{\mathbf{w}}'_3]$ is independent of the lift. Certainly $d\check{\mathbf{t}} \in B^3(G_b, \mathcal{A})$, but generically $d\check{\mathbf{t}} \notin B^3(G_b, \mathbb{Z}_2)$. As such, $[d\check{\mathbf{t}}]$ is not generally trivial in $\mathcal{H}^3(G_b, \mathbb{Z}_2)$ and thus can change the obstruction class.

C. Example: two layers of semion-fermion

We consider two layers of semion-fermion topological order with $G_f = \mathbb{Z}_4^{T,f} = \mathbb{Z}_2^T \times \mathbb{Z}_2^f$ symmetry. The particles in \mathcal{C} are generated by two Abelian semions s_1, s_2 with $\theta_{s_i} = +i$ and trivial mutual braiding and a transparent fermion ψ . Time reversal acts as $T_{s_i} = \psi s_i$.

Up to gauge transformations, one can check that, with the anyons ordered $1, s_1, s_2, s_1 s_2, \psi, \psi s_1, \psi s_2, \psi s_1 s_2$, then

$$U_{\mathbf{T}}(a, b; a \times b) = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 \\ 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 & 1 & -1 & -1 & 1 \\ 1 & -1 & -1 & 1 & 1 & -1 & -1 & 1 \\ 1 & -1 & -1 & 1 & 1 & -1 & -1 & 1 \\ 1 & 1 & -1 & -1 & 1 & 1 & -1 & -1 \\ 1 & 1 & -1 & -1 & 1 & 1 & -1 & -1 \end{pmatrix} \quad (152)$$

and $\eta_a(\mathbf{T}, \mathbf{T}) = \{1, i, i, 1, -1, -i, -i\}$ in the same order. There is also a solution with $\eta_a \rightarrow \eta_a^*$, but it behaves similarly. We can calculate κ from U by

$$\kappa_{\mathbf{T}, \mathbf{T}}(a, b; a \times b) = U_{\mathbf{T}}^*(a, b; a \times b) U_{\mathbf{T}}^*(\mathbf{T}a, \mathbf{T}b; \mathbf{T}(a \times b)) \quad (153)$$

from which we obtain $\beta_a(\mathbf{T}, \mathbf{T}) = \{1, i, i, 1, -1, -i, -i\}$ in our preferred gauge. From this we find

$$\omega_a(\mathbf{T}, \mathbf{T}) = \frac{\beta_a(\mathbf{T}, \mathbf{T})}{\eta_a(\mathbf{T}, \mathbf{T})} = +1 \text{ for all } a. \quad (154)$$

We now need to obtain \mathfrak{G} of the modular extension. The modular extension we consider is the tensor product $U(1)_2 \times U(1)_2 \times U(1)_{-2} \times U(1)_{-2}$, whose quasiparticles are generated by s_1, s_2, v , and ψv , where v is a semionic fermion-parity vortex with $\theta_v = -i$. Note that $\theta_{\psi v} = -i$ as well due to the nontrivial braiding. Coincidentally this theory is two copies of the double-semion theory, but that fact plays no role here. Note that $\check{\Upsilon}_\psi$ respects locality since there is an Abelian fermion parity vortex.

Our earlier expression for ω can now be written

$$\omega_a = M_{a,1}, \quad (155)$$

for $a \in \mathcal{C}$, up to a fermion. That is, $\mathfrak{w}(\mathbf{T}, \mathbf{T}) = 1$.

There are two options for the extension of the action of time-reversal. Note that $s_1 s_2$ is a fermion, so $v s_1 s_2$ and $v \psi s_1 s_2$ are both semions with $\theta = +i$. Either we can set $v \rightarrow v s_1 s_2$ or $v \rightarrow v \psi s_1 s_2$. We choose the latter permutation action first.

Solving the $U - F$ and $U - R$ consistency equations by computer, with the anyons ordered

$$1, s_1, s_2, s_1 s_2, v, v s_1, v s_2, v s_1 s_2, v \psi, v \psi s_1, v \psi s_2, v \psi s_1 s_2, \psi, \psi s_1, \psi s_2, \psi s_1 s_2,$$

we obtain

$$U_{\mathbf{T}}(a, b; a \times b) = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 & -1 & -1 & 1 & 1 & -1 & -1 & 1 & 1 & 1 & 1 & 1 & -1 & -1 & -1 \\ 1 & 1 & 1 & 1 & -1 & -1 & -1 & -1 & 1 & 1 & 1 & 1 & -1 & -1 & -1 & -1 & -1 & -1 \\ 1 & 1 & -1 & -1 & 1 & 1 & -1 & -1 & -1 & -1 & 1 & 1 & -1 & -1 & -1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 & 1 & 1 & -1 & -1 & -1 & -1 & -1 & 1 & 1 & -1 & -1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 & 1 & 1 & -1 & -1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & -1 & -1 & -1 & -1 & -1 & -1 & 1 & 1 & -1 & -1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 & 1 & 1 & -1 & -1 & -1 & -1 & -1 & 1 & 1 & -1 & -1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & -1 & -1 & -1 & -1 & -1 & -1 & 1 & 1 & -1 & -1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 & 1 & 1 & -1 & -1 & -1 & -1 & -1 & 1 & 1 & -1 & -1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & -1 & -1 & -1 & -1 & -1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & -1 & -1 & 1 & 1 & 1 & 1 & 1 & -1 & -1 & -1 & 1 & 1 & -1 & -1 & 1 & 1 & 1 \\ 1 & -1 & -1 & 1 & 1 & 1 & 1 & 1 & -1 & -1 & -1 & 1 & 1 & -1 & -1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 & -1 & -1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & -1 & -1 & -1 \\ 1 & 1 & 1 & -1 & -1 & -1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & -1 & -1 & -1 \end{pmatrix} \quad (156)$$

from which we obtain κ and

$$\beta_a = \{1, i, i, 1, i, -1, 1, -i, -i, 1, -1, i, 1, i, i, 1\} \quad (157)$$

up to a gauge transformation. Then

$$\Omega_a = \{1, 1, 1, 1, -1, -1, -1, -1, -1, -1, -1, 1, 1, 1, 1, 1\} = M_{a,\psi} \quad (158)$$

so $\check{\mathfrak{G}} = \psi$. Note that $\check{\mathfrak{G}}$ is not ambiguous by a fermion since it is defined for all of $\check{\mathcal{C}}$. We can now compute, whether we choose $\mathfrak{w}(\mathbf{T}, \mathbf{T}) = 1$ or ψ ,

$$\sigma_3(\mathbf{T}, \mathbf{T}, \mathbf{T}) = \check{\mathfrak{G}}(\mathbf{T}, \mathbf{T}) \overline{d\mathfrak{w}(\mathbf{T}, \mathbf{T})} = \psi \times 1 = \psi. \quad (159)$$

Hence $[\sigma_3] \in \mathcal{H}^3(\mathbb{Z}_2^T, \mathbb{Z}_2)$ is nontrivial and the symmetry fractionalization is obstructed.

One can check that choosing the other action $v \rightarrow v s_1 s_2$ under time reversal leads to the same result. There is a shortcut to see that there is an inconsistency here. We stated above that

$$\eta_{s_1 s_2} = +1 = -\theta_{s_1 s_2} \quad \text{and} \quad \eta_{\psi s_1 s_2} = -1 = -\theta_{\psi s_1 s_2}. \quad (160)$$

One can show on general grounds that [1],

$$\eta_a = \theta_a \quad \text{if } a = b \times \mathbf{T}(b) \quad (161)$$

for some b in the category in question. For $a = s_1 s_2$ or $\psi s_1 s_2$, we have $\eta_a \neq \theta_a$ for this fractionalization pattern. Equation (161) is satisfied in \mathcal{C} because, for the a in question, there is no $b \in \mathcal{C}$ for which $a = b \times \mathbf{T}(b)$. However, in the modular extension $\check{\mathcal{C}}$, such a b does exist; $s_1 s_2 = v s_1 s_2 \times \mathbf{T}(v s_1 s_2)$

if we use the permutation action $v \rightarrow v s_1 s_2$, and $\psi s_1 s_2 = v s_1 s_2 \times \mathbf{T}(v s_1 s_2)$ under the permutation action $v \rightarrow v \psi s_1 s_2$. Therefore, there is an inconsistency between the consistency of fractionalization in the modular extension and the fractionalization pattern in \mathcal{C} .

D. 't Hooft anomaly and dependence of $[\sigma_3]$ on choices

In defining $[\sigma_3]$, we have made two choices. We made a choice of modular extension $\check{\mathcal{C}}_v$ that is free of \mathcal{H}^2 obstruction and a choice of lift $\check{\rho}$. Given these choices, it then makes sense to ask whether the symmetry fractionalization data $\{\eta_a\}$ on \mathcal{C} can be lifted to $\check{\mathcal{C}}_v$, and $[\sigma_3] \in \mathcal{H}^3(G_b, \mathbb{Z}_2)$ is the obstruction to such a lift. In principle, $[\sigma_3]$ could depend on the choices v and $\check{\rho}$; to highlight this dependence, we can write $[\sigma_3^{(v, \check{\rho})}]$. In Sec. IV B, we saw that $(3+1)$ D FSPTs define an element $[n_3] \in \mathcal{H}^3(G_b, \mathbb{Z}_2)/\Gamma^3$, if the lower layer data, n_1, n_2 , vanish. Since $q_{\Gamma^3}([\sigma_3]) \in \mathcal{H}^3(G_b, \mathbb{Z}_2)/\Gamma^3$, it is therefore natural to assume that $(2+1)$ D FSETs with vanishing $[\sigma_1]$ and $[\sigma_2]$ obstructions must exist at the surface of $(3+1)$ D FSPTs characterized by $[n_3] = q_{\Gamma^3}([\sigma_3])$ (and vanishing n_1, n_2).

It is natural to expect that the super-modular category \mathcal{C} and its symmetry fractionalization data fully determine the 't Hooft anomaly, equivalently the (3 + 1)D FSPT that hosts the given theory at its surface. It would thus follow that $q_{\Gamma^3}([o_3^{(v, \check{\rho})}])$ should be independent of valid changes of v and $\check{\rho}$. Below we will examine this expectation in detail.

Given a lift $\check{\rho}$ for $\check{\mathcal{C}}_v$ and a group homomorphism $\pi \in Z^1(G_b, \mathbb{Z}_2)$ from G_b to \mathbb{Z}_2 , one can obtain a specific topological autoequivalence $\check{\rho}'_g$ on $\check{\mathcal{C}}_v$ [36]. In the case $v' - v = 0$, this amounts to modifying the lift $\check{\rho}_g$ for a fixed modular extension as follows:

$$\check{\rho}_g \rightarrow \check{\rho}'_g = \alpha_\psi^{\pi(g)} \check{\rho}_g. \quad (162)$$

Then, as we will explain, we expect the following result to hold in all cases:

$$\begin{aligned} [o_3^{(v', \check{\rho}')}] = & [o_3^{(v, \check{\rho})}] + [s_1] \cup [\pi] \cup [\pi] + [\pi] \cup [\tilde{\omega}_2] \\ & + \frac{v' - v}{2} [\tilde{\omega}_2] \cup_1 [\tilde{\omega}_2]. \end{aligned} \quad (163)$$

Note that when $v' - v$ is odd, we can only consider the change of $[o_3]$ when $[\tilde{\omega}_2]$ and $[s]$ are trivial, in which case Eq. (163) gives $[o_3^{(v', \check{\rho}')}] = [o_3^{(v, \check{\rho})}]$. This is because when $[\tilde{\omega}_2]$ is nontrivial and $v' - v$ is odd, Eq. (128) implies that either $[o_2^{(v)}]$ or $[o_2^{(v')}]$ is nontrivial, in which case $[o_3]$ would be ill defined. Also, this equation is only meaningful with $s_1 \neq 0$ when $v' - v = 0 \pmod{8}$.

Motivated by a conjecture of an earlier version of this paper,¹⁰ version 4 of Ref. [36] proved the above formula in general except for the case when Υ_ψ violates locality with $v' - v$ odd (which forces $\tilde{\omega}_2 = 0$). We reproduce¹¹ the proof for $v' - v = 0$ (with G_b allowed to be antiunitary) in Appendix D, and also discuss invariance of $[o_3]$ under various gauge transformations. The remaining case where Υ_ψ violates locality with $v' - v$ odd is technically challenging, and therefore still open, but we see no conceptual reason to expect the formula to fail.

The last unproven case notwithstanding, this result shows that $q_{\Gamma^3}([o_3^{(v, \check{\rho})}])$ is the same for all v with vanishing $[o_2^{(v)}]$ and all valid lifts $\check{\rho}$. It also shows that if $q_{\Gamma^3}([o_3^{(v, \check{\rho})}])$ is trivial in $\mathcal{H}^3(G_b, \mathbb{Z}_2)/\Gamma^3$, then there exists a choice of v and $\check{\rho}$ such that $[o_3] \in \mathcal{H}^3(G_b, \mathbb{Z}_2)$ is trivial. As such, $q_{\Gamma^3}([o_3])$ should indeed be viewed as a piece of the 't Hooft anomaly of the FSET.

VIII. $\mathcal{H}^4(G_b, \text{U}(1))$ OBSTRUCTION AND 'T HOOFT ANOMALY

Suppose that the obstructions o_1, o_2, o_3 all vanish. Then we can lift the full set of symmetry fractionalization data on \mathcal{C}

¹⁰An earlier version of this paper conjectured that $[o_3]$ itself, and not just its image under q_{Γ^3} would be independent of v .

¹¹Version 3 of Ref. [36] contained Eq. (163) restricted to the case $s_1 = 0$, $v' - v$ even, and Υ_ψ respects locality. After discussions between the present authors and the authors of Ref. [36], v4 corrected their formula for the change of $\tilde{\Omega}_a$ under change of lift when Υ_ψ violates locality and used this result to generalize Eq. (163) to all cases except when Υ_ψ violates locality with $v' - v$ odd.

to $\check{\mathcal{C}}$. That is, we now have a bosonic SET, i.e., a UMTC $\check{\mathcal{C}}$ equipped with G_b symmetry fractionalization data. One can then attempt to gauge G_b . As a first step, one must apply the standard formalism for bosonic topological phases [1] to construct a G_b -crossed extension of $\check{\mathcal{C}}$, that is, a theory that describes both the excitations in $\check{\mathcal{C}}$ and G_b symmetry defects. There is a known obstruction [1,4], which we shall call $[o_4]$, to doing so, valued in $\mathcal{H}^4(G_b, \text{U}(1))$. $[o_4]$ quantifies the failure of the consistency of fusion of the symmetry defects, that is, it quantifies an inability to define F -symbols for the defects that obey the pentagon equation. This obstruction is the last obstruction in the anomaly cascade.

The 4-cocycle o_4 depends explicitly on all of the choices required to gauge fermion parity and define symmetry fractionalization on $\check{\mathcal{C}}$. We can denote this dependence explicitly by writing $o_4^{(v, \check{\rho}, \check{\eta})}$. Therefore we have a 4th cohomology class

$$[o_4^{(v, \check{\rho}, \check{\eta})}] \in \mathcal{H}^4(G_b, \text{U}(1)). \quad (164)$$

$o_4^{(v, \check{\rho}, \check{\eta})}$ is in general a nontrivial function of its arguments.

It is straightforward to see how $[o_4]$ changes under changing the symmetry fractionalization class $\check{\eta}$. Changes of symmetry fractionalization, which are consistent with the symmetry fractionalization on \mathcal{C} are given by:

$$\check{\eta}_a(\mathbf{g}, \mathbf{h}) \rightarrow \check{\eta}_a(\mathbf{g}, \mathbf{h}) M_{a, t(\mathbf{g}, \mathbf{h})}, \quad (165)$$

with $t(\mathbf{g}, \mathbf{h}) \in \{1, \psi\} \cong \mathbb{Z}_2$. One can then use the relative anomaly formula [7] to compute

$$\begin{aligned} o_4(\mathbf{g}, \mathbf{h}, \mathbf{k}, \mathbf{l}) & \rightarrow o_4(\mathbf{g}, \mathbf{h}, \mathbf{k}, \mathbf{l}) R^{t(\mathbf{k}, \mathbf{l}), t(\mathbf{g}, \mathbf{h})} \eta_{t(\mathbf{k}, \mathbf{l})}(\mathbf{g}, \mathbf{h}) \\ & = o_4(\mathbf{g}, \mathbf{h}, \mathbf{k}, \mathbf{l}) (-1)^{t \cup t + \tilde{\omega}_2 \cup t} \end{aligned} \quad (166)$$

where $\tilde{\omega}_2$ is the additive \mathbb{Z}_2 representation of ω_2 appearing in Eq. (43). Comparing to Eq. (60), we see that o_4 changes by an element of Γ^4 .

We can also consider how $[o_4]$ depends on the choice of v and $\check{\rho}$. In general, changing v and $\check{\rho}$ may change the lower-level obstructions $[o_2]$, and $[o_3]$, in which case the change of $[o_4]$ is not well-defined. Nevertheless, we may consider changes in v and $\check{\rho}$ such that o_2 and o_3 remain trivial. It would be interesting to derive a general formula for how $[o_4]$ changes under changing v and $\check{\rho}$. It is not clear if a simple general formula exists.

While $[o_4^{(v, \check{\rho}, \check{\eta})}]$ is in general a nontrivial function of its arguments, we can consider the image of $[o_4]$ under the map q_{Γ^4} defined in Eq. (61). Below we conjecture that $q_{\Gamma^4}([o_4])$ is independent of the choices $v, \check{\rho}, \check{\eta}$.

Recall that in Sec. IV B, we explained that (3 + 1)D FSPTs, and therefore 't Hooft anomalies for (2 + 1)D FSETs, are classified by a set of data (n_1, n_2, n_3, v_4) . Furthermore, if $n_1, n_2, n_3 = 0$, then (3 + 1)D FSPTs, and therefore 't Hooft anomalies of (2 + 1)D FSETs, are characterized by an element

$$[v_4] \in \mathcal{H}^4(G_b, \text{U}(1))/\Gamma^4. \quad (167)$$

We expect that the 't Hooft anomaly of the (2 + 1)D system is entirely a property of the quasiparticles of the fermionic theory, described by the super-modular category \mathcal{C} , together with the symmetry action ρ and symmetry fractionalization data η . If this expectation is correct, then the anomaly should

be independent of all the choices involved in lifting the symmetry fractionalization data to a given modular extension. Anomaly matching therefore leads to the expectation that a $(2+1)$ D FSET with o_1, o_2, o_3 all vanishing and some $[o_4]$ can only exist at the surface of a $(3+1)$ D FSPT with n_1, n_2, n_3 vanishing and some $[v_4]$ such that

$$[v_4] = q_{\Gamma^4}([o_4]). \quad (168)$$

The above discussion then leads us to the following formal conjecture:

Conjecture VIII.1. (a) $q_{\Gamma^4}([o_4^{(v, \check{\rho}, \check{\eta})}])$ is independent of changing $v, \check{\rho}, \check{\eta}$, as long as $[o_1], [o_2], [o_3]$ all vanish.

(b) If $q_{\Gamma^4}([o_4^{(v, \check{\rho}, \check{\eta})}])$ is trivial in $\mathcal{H}^4(G_b, \text{U}(1))/\Gamma^4$, then there exists a choice of $v, \check{\rho}, \check{\eta}$ such that $[o_4^{(v, \check{\rho}, \check{\eta})}]$ is trivial in $\mathcal{H}^4(G_b, \text{U}(1))$.

IX. ADDITIONAL EXAMPLES

A. T-Pfaffian

T-Pfaffian is the surface theory for the $G_b = \text{U}(1) \rtimes \mathbb{Z}_2^{\mathbf{T}}$ topological insulator, where $G_f = [\text{U}(1) \rtimes \mathbb{Z}_4^{\mathbf{T}, f}]/\mathbb{Z}_2$. It consists of a subcategory \mathcal{C} of the Ising $\times \text{U}(1)_{-8}$ state as follows. Labeling elements of Ising $\times \text{U}(1)_{-8}$ by a_j with $a \in \{I, \psi, \sigma\}$ and $j = 0, 1, \dots, 7$, the quasiparticle content of \mathcal{C} consists of the twelve quasiparticles $\{I_{2k}, \psi_{2k}, \sigma_{2k+1}\}$ for $k = 0, 1, 2, 3$. The transparent fermion of the category is $\psi = \psi_4$ (caution with the notation; ψ alone means the physical fermion of \mathcal{C} , and ψ_k is a label in Ising $\times \text{U}(1)_{-8}$). Time reversal $[\rho_{\mathbf{T}}]$ interchanges $I_2 \leftrightarrow \psi_2$ and $I_6 \leftrightarrow \psi_6$.

It is not hard to check that Υ_{ψ} respects locality in this theory; therefore, $\text{Aut}(\mathcal{C}) = \text{Aut}_{LR}(\mathcal{C})$, and one can check that $[\rho_g]$ is determined entirely by its permutation action. In particular, writing $\mathbf{g} = (e^{i\theta_g}, \mathbf{T}^{a_g})$ for $a \in \{0, 1\}$, then $[\rho_g]$ is the identity if $a_g = 0$ and is nontrivial and equal to $[\rho_{\mathbf{T}}]$ if $a_g = 1$, independent of θ_g . The minimal modular extensions of this theory have $c_- \in \frac{1}{2}\mathbb{Z}$. The minimal modular extension $\tilde{\mathcal{C}}$ that is compatible with time reversal, i.e., with $c_- = 0$ can be written

$$\tilde{\mathcal{C}} = (\mathcal{C} \boxtimes \text{Ising}) / \{\psi \psi' \sim 1\}, \quad (169)$$

where the denominator means condensing the bound state of the physical fermion ψ in \mathcal{C} with the fermion ψ' in the additional copy of Ising. One can check that $\tilde{\mathcal{C}}$ has only σ -type fermion parity vortices and admits a permutation action of \mathbf{T} that lifts the permutation action of $\rho_{\mathbf{T}}$ as shown in Refs. [14, 16, 43]. Since \mathcal{C} has no modular isotopes [44], the aforementioned permutation must define a full autoequivalence $[\check{\rho}_{\mathbf{T}}]$ of \mathcal{C} , so the $\mathcal{H}^1(G_b, \mathbb{Z}_{\mathbf{T}})$ anomaly vanishes:

$$[o_1] = 0 \in \mathcal{H}^1(G_b, \mathbb{Z}_{\mathbf{T}}). \quad (170)$$

Since all the vortices are σ type, $\check{\Upsilon}_{\psi}$ does not respect locality and is therefore the nontrivial element of $\ker r = \mathbb{Z}_2$. This permutation action squares to the identity permutation, although this does not guarantee that the $\mathcal{H}^2(G_b, \mathbb{Z}_2)$ anomaly vanishes because $\check{\Upsilon}_{\psi}$ is nonpermuting.

Physically, we expect that the anomaly arises because the physical fermion ψ carries nontrivial $\text{U}(1)^f$ quantum numbers, but the fermion parity vortices are all σ -type, that is,

they absorb ψ . The fact that ψ carries nontrivial $\text{U}(1)^f$ quantum numbers enters through η_{ψ} , which appears at the level of the $\mathcal{H}^3(G_b, \mathbb{Z}_2)$ obstruction. We therefore conjecture the following:

Conjecture IX.1. The T-Pfaffian state has vanishing $[o_2]$, but nonvanishing $[o_3]$.

Checking this conjecture explicitly would require knowledge of the full F and R symbols of the gauged T-Pfaffian state, which we do not know in general how to compute from the decomposition Eq. (169).

We contrast the present picture with a more typical argument [45]. One can use a decorated domain wall construction wherein the bulk $(3+1)$ D topological insulator can be understood in terms of decorating \mathbf{T} domain walls with a $(2+1)$ D integer quantum Hall state with Chern number 1. This construction assumes $\text{U}(1)^f$ symmetry from the outset and requires a $c_- = 1/2 \bmod 1$ minimal modular extension on the boundary so that \mathbf{T} domain walls on the boundary carry a chiral mode arising from the integer quantum Hall state in the bulk. However, $c_- = 1/2 \bmod 1$ is manifestly incompatible with time reversal symmetry in isolation. In this context, then, the anomaly is characterized as a nontrivial element of $\mathcal{H}^1(\mathbb{Z}_2^{\mathbf{T}}, \mathbb{Z}_{\mathbf{T}})$. Another way to state this latter construction is that we gauge $\text{U}(1)^f$ symmetry first, and then attempt to lift the $G_f/\text{U}(1)^f = \mathbb{Z}_2^{\mathbf{T}}$ symmetry to the gauged theory. In this context, the T-Pfaffian anomaly appears as an $\mathcal{H}^1(G_f/\text{U}(1)^f, \mathbb{Z}_{\mathbf{T}})$ obstruction. By comparison, in our framework, we gauge fermion parity first and then attempt to lift the $G_f/\mathbb{Z}_2^f = G_b$ symmetry to the gauged theory. In the latter case, the $\mathcal{H}^1(G_b, \mathbb{Z}_{\mathbf{T}})$ anomaly vanishes, and T-Pfaffian fits into a higher level of the anomaly cascade. Both perspectives correspond to valid ways of calculating the same anomaly; they simply decompose the classification differently. Our perspective is more general, since fermionic systems always have \mathbb{Z}_2^f symmetry but need not have $\text{U}(1)^f$ symmetry.

B. $\text{Sp}(3)_3 \boxtimes \{1, \psi\}$

$\mathcal{C} = \text{Sp}(3)_3 \boxtimes \{1, \psi\}$ gives an example of a $v = 6$ phase of the \mathbb{Z}_{16} classification for $G_f = \mathbb{Z}_4^{\mathbf{T}, f}$. $\text{Sp}(3)_3$ has central charge $c = 1 \bmod 8$, so we should consider $\text{Sp}(3)_3 \times \text{U}(1)_{-1}$ to get a time-reversal invariant theory. $\text{Sp}(3)_3$ itself has 20 particles, so $\mathcal{C} = \text{Sp}(3)_3 \boxtimes \{1, \psi\}$ has 40 particles. The modular data of $\text{Sp}(3)_3$ were obtained by computer using SageMath. We tabulate the quantum dimensions and topological twists of $\text{Sp}(3)_3$ in Table II; the S matrix is large and unenlightening, so we do not write it explicitly here. The particles will be labeled from 1 to 20, where particle 1 is the identity and the others are ordered by increasing quantum dimension but otherwise arbitrarily. We can label a particle in \mathcal{C} by a pair (n, x) with n from 1 to 20 and $x \in \{1, \psi\}$. There is a unique action of time reversal that preserves the modular data up to complex conjugation, with the corresponding permutation given in Table II.

The time-reversal invariant modular extension is $\tilde{\mathcal{C}} = \text{Sp}(3)_3 \boxtimes \mathcal{I}(-2)$. We label the parity vortices of $\mathcal{I}(-2)$ by a and \bar{a} , with $a \times \psi = \bar{a}$ and $a \times \bar{a} = 1$. It is clear that $\tilde{\mathcal{C}}$ has v -type vortices, and furthermore that $\check{\Upsilon}_{\psi}$ respects locality, so $\text{Aut}(\tilde{\mathcal{C}}) = \text{Aut}_{LR}(\mathcal{C})$ and $\ker r = \{1, \alpha_{\psi}\} = \mathbb{Z}_2$ where α_{ψ}

TABLE II. Quantum dimensions, topological twists, and time-reversal actions for $\check{\mathcal{C}} = \text{Sp}(3)_3 \boxtimes \mathcal{I}(2)$. All other data can be determined from the tabulated data; e.g., $\theta_{(n,\psi)} = -\theta_{(n,1)}$, and $\mathbf{T}((n,\bar{a})) = \mathbf{T}((n,a)) \times (1,\psi)$. The action of \mathbf{T} on $\mathcal{C} = \text{Sp}(3)_3 \boxtimes \{1,\psi\}$ is uniquely determined and squares to the identity. Its lift to $\check{\mathcal{C}}$ is ambiguous by α_ψ and squares to α_ψ . Quantum dimensions can be written exactly in terms of 56th roots of unity, but we do not do so here.

n	1	2	3	4	5	6	7	8	9	10
$d_{(n,x)}$	1	1	3.49	3.49	4.49	4.49	4.49	4.49	5.60	5.60
$\theta_{(n,1)}$	1	i	i	1	$e^{-13\pi i/14}$	$e^{4\pi i/7}$	$e^{-\pi i/14}$	$e^{-4\pi i/7}$	$e^{6\pi i/7}$	$e^{-6\pi i/7}$
$\mathbf{T}((n,1))$	$(1,1)$	$(2,\psi)$	$(3,\psi)$	$(4,1)$	$(7,\psi)$	$(8,1)$	$(5,\psi)$	$(6,1)$	$(10,1)$	$(9,1)$
$\theta_{(n,a)}$	$e^{-\pi i/4}$	$e^{\pi i/4}$	$e^{\pi i/4}$	$e^{-\pi i/4}$	$e^{23\pi i/28}$	$e^{9\pi i/28}$	$e^{-9\pi i/28}$	$e^{-23\pi i/28}$	$e^{17\pi i/28}$	$e^{25\pi i/28}$
$\mathbf{T}((n,a))$	$(2,\bar{a})$	$(1,a)$	$(4,a)$	$(3,\bar{a})$	$(8,a)$	$(7,\bar{a})$	$(6,a)$	$(5,\bar{a})$	$(12,\bar{a})$	$(11,\bar{a})$
n	11	12	13	14	15	16	17	18	19	20
$d_{(n,x)}$	5.60	5.60	9.10	9.10	10.10	10.10	10.10	10.10	11.59	11.59
$\theta_{(n,1)}$	$e^{-9\pi i/14}$	$e^{-5\pi i/14}$	1	i	$e^{-2\pi i/7}$	$e^{3\pi i/14}$	$e^{2\pi i/7}$	$e^{11\pi i/14}$	$-i$	-1
$\mathbf{T}((n,1))$	$(12,\psi)$	$(11,\psi)$	$(13,1)$	$(14,\psi)$	$(17,1)$	$(18,\psi)$	$(15,1)$	$(16,\psi)$	$(19,\psi)$	$(20,1)$
$\theta_{(n,a)}$	$e^{-25\pi i/28}$	$e^{-17\pi i/28}$	$e^{-\pi i/4}$	$e^{\pi i/4}$	$e^{-15\pi i/28}$	$e^{-\pi i/28}$	$e^{\pi i/28}$	$e^{15\pi i/28}$	$e^{-3\pi i/4}$	$e^{3\pi i/4}$
$\mathbf{T}((n,a))$	$(10,a)$	$(9,\bar{a})$	$(14,\bar{a})$	$(13,a)$	$(18,\bar{a})$	$(17,a)$	$(16,\bar{a})$	$(15,a)$	$(20,a)$	$(19,\bar{a})$

fuses a fermion into each parity vortex. The modular data of this product theory can be computed straightforwardly from the modular data of the constituents. Labeling the particles in $\check{\mathcal{C}}$ as (n,x) with, again, $n = 1, 2, \dots, 20$, and this time $x \in \{1, a, \bar{a}, \psi\}$, one can check directly that there are exactly two lifts of the action of time reversal on \mathcal{C} to $\check{\mathcal{C}}$. One is tabulated in Table II, and the other is given by composing with α_ψ (which switches $a \leftrightarrow \bar{a}$). It is clear by inspection that these actions do not square to the identity; for example, under time reversal,

$$(1,a) \rightarrow (2,\bar{a}) \rightarrow (1,\bar{a}) \rightarrow (2,a) \rightarrow (1,a), \quad (171)$$

which yields a representation of \mathbb{Z}_4^T , not of \mathbb{Z}_2^T . Composition with α_ψ does not change this fact, so we conclude that, as expected, this theory has an $\mathcal{H}^2(\mathbb{Z}_2^T, \mathbb{Z}_2)$ obstruction.

X. DISCUSSION

We have systematically characterized the set of obstructions that appear in lifting symmetry fractionalization data from a super-modular category \mathcal{C} to a minimal modular extension $\check{\mathcal{C}}$. We found that this data is in good correspondence with the known classification of fermionic SPTs and provided an understanding for each obstruction:

(1) The $\mathcal{H}^1(G_b, \mathbb{Z}_T)$ piece is the obstruction to defining a lift of the autoequivalence $[\rho_g]$ to $[\check{\rho}_g] \in \text{Aut}_{LR}(\check{\mathcal{C}}_v)$ for some minimal modular extension $\check{\mathcal{C}}_v$.

(2) When the \mathcal{H}^1 obstruction vanishes and one can define lifts of the maps $[\rho_g]$ for some choice of minimal modular extension v , the $\mathcal{H}^2(G_b, \ker r)$ piece is the obstruction to choosing lifts $[\check{\rho}_g] : G_b \rightarrow \text{Aut}_{LR}(\check{\mathcal{C}}_v)$ with the appropriate group structure.

(3) When the \mathcal{H}^1 and \mathcal{H}^2 obstructions vanish and we pick a set of lifts $[\check{\rho}]$ with the appropriate group structure for a given modular extension v , the $\mathcal{H}^3(G_b, \mathbb{Z}_2)$ piece is the obstruction to lifting the symmetry fractionalization data $\{\eta_a\}$ on \mathcal{C} to $\{\check{\eta}_a\}$ on $\check{\mathcal{C}}_v$.

(4) When the \mathcal{H}^1 , \mathcal{H}^2 , \mathcal{H}^3 obstructions all vanish and we pick the lifts $\check{\rho}$ and symmetry fractionalization data $\check{\eta}$ on $\check{\mathcal{C}}_v$ for some v , the $\mathcal{H}^4(G_b, \text{U}(1))$ piece is the obstruction to lifting

$\check{\mathcal{C}}_v$, together with its symmetry fractionalization data, to a G_b -crossed modular tensor category $\check{\mathcal{C}}_{G_b}^\times$.

Our paper raises a number of open questions. First, we have defined a series of obstructions $[o_1]$, $[o_2^{(v)}]$, $[o_3^{(v,\check{\rho})}]$, and $[o_4^{(v,\check{\rho},\check{\eta})}]$. We expect, but have not shown explicitly, that the bulk $(3+1)$ D FSPT that hosts our given $(2+1)$ D theory on its surface is characterized by the image of the maps $q_{\Gamma^i} : \mathcal{H}^i \rightarrow \mathcal{H}^i / \Gamma^i$, for the finite groups Γ^i reviewed in Sec. IV B. This has led us to conjecture that the $q_{\Gamma^i}[o_i]$ are independent of the various valid choices involving v , $\check{\rho}$, and $\check{\eta}$. For the case $i = 2$, this conjecture is proven so long as $\ker r = \mathbb{Z}_2$. For the case $i = 3$, there is a loose end in proving Eq. (163) for the case where Υ_ψ violates locality with $v' - v$ is odd. For $i = 4$, we have derived how $[o_4^{(v,\check{\rho},\check{\eta})}]$ changes under changes of $\check{\eta}$, but not under valid changes of v , $\check{\rho}$.

We have also conjectured, based on the $(3+1)$ D FSPT classification, that in general $\ker r = \mathbb{Z}_2$. However we have also found an interesting example involving doubled $\text{SU}(2)_6$ where there are multiple independent permutations of fermion parity vortices that keep the modular data invariant and act trivially on the super-modular category; these point to a possibility that $\ker r$ is larger than \mathbb{Z}_2 , or that there are permutations of anyons that preserve the modular data which do not correspond to autoequivalences of the category.

While we have examples of a nontrivial \mathcal{H}^2/Γ^2 anomaly for antiunitary symmetries (namely the semion-fermion theory with $\mathbf{T}^2 = (-1)^F$), we do not have an example for unitary G_b .

We have given an anomaly inflow argument for the \mathcal{H}^2 contribution to the anomaly using the decorated domain wall construction; it would be useful to gain a similar understanding for the other contributions.

The Wang-Gu results for $(3+1)$ D FSPTs reviewed in Sec. IV B have significantly more structure than we have derived so far in our obstruction theory. There is a set of data (n_1, n_2, n_3, n_4) obeying complicated consistency equations, group multiplication laws under stacking invertible phases, and equivalences. However we have only seen how one can extract part of this data from the $(2+1)$ D fermion SET. It would be interesting to understand to what extent more

aspects of the general $(3+1)$ D FSPT characterization can be extracted. For example, it may be possible to extract, from the $(2+1)$ D data, the full group structure of the anomalies, which corresponds to a group extension of the groups \mathcal{H}^1 , \mathcal{H}^2/Γ^2 , \mathcal{H}^3/Γ^3 , and \mathcal{H}^4/Γ^4 . Alternatively, can we extract the specific consistency equations for (n_1, n_2, n_3, ν_4) derived in [30]?

Our formalism suggests a very general way of understanding mixed anomalies. We have shown that the data characterizing the anomaly, namely, a spectral sequence decomposition into the Wang-Gu data of the bordism group $\Omega_4(BG_b, \xi)$, where ξ is a kind of twisted spin structure, can be interpreted by first gauging fermion parity and then considering a sequence of obstructions to lifting the G_b symmetry to the fermion parity gauged theory. Consider instead a bosonic SET with symmetry group $G = G_1 \times G_2$. Then there is a spectral sequence decomposition of the $\mathcal{H}^4(G, U(1))$ bosonic SET anomaly via the Künneth decomposition. Our procedure might be applied to understand the mixed anomaly between G_1 and G_2 in terms of a “two-step” gauging process, where one first gauges G_1 and then determines a cascade of obstructions to lifting the G_2 symmetry to the G_1 -gauged theory. More generally, if G is given by a short exact sequence

$$1 \rightarrow G_1 \rightarrow G \rightarrow G_2 \rightarrow 1, \quad (172)$$

there is again a spectral sequence decomposition of $\mathcal{H}^4(G, U(1))$ and a similar construction may apply.

Note added. The closely related works Refs. [35,39] were developed in parallel with the present paper. Reference [35] develops a theory of fermionic symmetry fractionalization and is used as a starting point for this paper. Reference [39] develops a theory of $(2+1)$ D invertible fermionic topological phases. Reference [36] is also closely related and was developed independently except as documented elsewhere in this paper; it develops a general characterization and classification of fermionic symmetry-enriched topological phases in $(2+1)$ D, and contains many results which overlap with [35,39] and the present paper.

ACKNOWLEDGMENTS

We thank Parsa Bonderson, Meng Cheng, and Zhenghan Wang for discussions. In particular, we thank Parsa Bonderson for extensive discussions regarding multiple revisions of this paper and Ref. [36] and for sharing early versions of revised drafts of Ref. [36]. We also thank Ryohei Kobayashi and Srivatsa Tata for collaboration on related projects. This work is supported by NSF CAREER (DMR-1753240) and JQI-PFC-UMD.

APPENDIX A: S-MATRIX OF A CONDENSED THEORY

We presently reproduce, with additional detail, the derivation from [33] of the S matrix \hat{S} of the theory obtained by condensing an Abelian boson ϕ in a theory with S -matrix S . Our main addition to [33] is a careful discussion of gauge invariance. Specifically, \hat{S} should be gauge-invariant under vertex basis transformations, a fact that is in tension with the expression of \hat{S} in terms of the gauge-dependent punctured S matrix of the uncondensed theory.

Assume that $\phi^n = 1$ for some $n > 1$; we will specialize to $n = 2$ later. Then we can view the Wilson lines for ϕ as generating a \mathbb{Z}_n 1-form symmetry of the theory; condensing ϕ corresponds to gauging that symmetry. We perform this gauging to understand the Hilbert space of the condensed theory on the torus.

In the path integral picture, gauging the symmetry means that we sum over all insertions of the symmetry generator on nontrivial cycles of the 3-torus. We begin by considering insertions of a ϕ Wilson line through the time circle. Then at every spatial slice, we are summing over states with no insertion and with all possible insertions of ϕ^k , that is, we consider an expanded Hilbert space of the original theory consisting of the Hilbert space on the torus and the Hilbert spaces on the punctured torus with punctures labeled ϕ^k for all $k < n$. (Notationally, the use of k in this Appendix is completely unrelated to the k we defined in Sec. VI.)

States on the punctured Hilbert space with puncture ϕ^k are labeled $|a; \phi^k\rangle$ where $N_{a, \bar{a}}^{\phi^k} > 0$. Physically, these states correspond to ones where an a, \bar{a} pair is created from vacuum, wrapped around a given spatial cycle of the torus (which we will call cycle β for concreteness), and then fused into ϕ^k at the puncture. Under a basis transformation Γ_c^{ab} of the fusion spaces V_c^{ab} , these states transform as

$$|a; \phi^k\rangle \rightarrow \Gamma_{\phi^k}^{a\bar{a}} (\Gamma_1^{a\bar{a}})^* |a; \phi^k\rangle \quad (A1)$$

and are thus gauge-invariant (in this sense) only when $k = 0$.

Also, observe that each anyon a must have a \mathbb{Z}_n orbit of some length ℓ_a that divides n . By definition, if $|a; \phi^k\rangle$ is a nonzero state, then $N_{a, \bar{a}}^{\phi^k} = N_{a, \phi^k}^a > 0$, that is, $a \times \phi^k = a$. Hence k is a multiple of ℓ_a . Hence the total number of distinct states in the expanded Hilbert space that are associated to the anyon a is n/ℓ_a .

Next consider inserting a Wilson loop for ϕ^k around cycle α of the torus. Let $W_\alpha(\phi^k)$ be the operator that does this; then since ϕ is Abelian,

$$W_\alpha(\phi^k) |a; \phi^m\rangle = M_{a, \phi^k} |a; \phi^m\rangle \quad (A2)$$

with M the mutual statistics. Since ϕ is Abelian,

$$M_{a, \phi^k} = e^{2\pi i k q/n} \quad (A3)$$

for some integer q . Summing over all of these insertions,

$$\begin{aligned} \sum_{k=0}^{n-1} W_\alpha(\phi^k) |a; \phi^m\rangle &= \left(\sum_{k=0}^{n-1} e^{2\pi i k q/n} \right) |a; \phi^m\rangle \\ &= \delta_{q, 0} |a; \phi^m\rangle. \end{aligned} \quad (A4)$$

That is, we require a to braid trivially with ϕ to keep its corresponding states in the condensed theory.

Finally, we consider inserting a Wilson loop of ϕ^k Wilson loops around cycle β of the torus. Clearly this takes the state $|a; \phi^m\rangle \rightarrow |a \times \phi^k; \phi^m\rangle$, so the only states we should consider are

$$|[a]; \phi^m\rangle = \frac{1}{\sqrt{\ell_a}} \sum_{k=0}^{\ell_a} |a \times \phi^k; \phi^m\rangle \quad (A6)$$

where here $[a]$ labels the orbit of a under fusion with ϕ .

The Hilbert space therefore consists of states $|(a); \phi^m\rangle$ such that a braids trivially with ϕ and $m = 0, 1, \dots, n/\ell_a - 1$ (that is, there are n/ℓ_a states per anyon). In particular, there is a unique state associated to $[a]$ if a has orbit length $\ell_a = n$ but the orbit $[a]$ splits to n/ℓ_a states after condensation if $a \times \phi^{\ell_a} = a$ for some $\ell_a < n$.

Now consider the transformation of these states under modular transformations. By definition, in this basis, if \mathbf{S} and \mathbf{T} are the operators that implement S and T modular transformations,

$$\langle a; \phi^m | \mathbf{S} | b; \phi^s \rangle = \delta_{m,s} S_{ab}^{(\phi^m)} \quad (\text{A7})$$

$$\langle a; \phi^m | \mathbf{T} | b; \phi^s \rangle = \delta_{m,s} T_{ab}^{(\phi^m)} \quad (\text{A8})$$

where $S_{ab}^{(\phi^m)}$ and $T_{ab}^{(\phi^m)}$ are the punctured S – and T matrices. Note that all elements of $T^{(\phi^m)}$ and diagonal elements $S_{aa}^{(\phi^m)}$ of the punctured S matrix are gauge-invariant under vertex basis transformations, while off-diagonal elements of the punctured S matrix (for $m > 0$) are not generally gauge invariant.

It is diagrammatically straightforward to check that

$$S_{a \times \phi^j, b}^{(\phi^m)} = S_{a, b \times \phi^k}^{(\phi^m)} = S_{a, b}^{(\phi^m)} \quad (\text{A10})$$

provided $[a]$ and $[b]$ are both deconfined particles, i.e., a and b both braid trivially with ϕ , and that

$$T_{ab}^{(\phi^m)} = \theta_a \delta_{a,b}. \quad (\text{A11})$$

Hence

$$\begin{aligned} \hat{S}_{([a],m),([b],s)} &= \langle [a]; \phi^m | \mathbf{S} | [b]; \phi^s \rangle \\ &= \frac{1}{\sqrt{\ell_a \ell_b}} \sum_{a \in [a], b \in [b]} \delta_{m,s} S_{ab}^{(\phi^m)} \end{aligned} \quad (\text{A12})$$

$$= \sqrt{\ell_a \ell_b} \delta_{m,s} S_{ab}^{(\phi^m)} \quad (\text{A13})$$

$$\hat{T}_{([a],m),([b],s)} = \langle [a]; \phi^m | \mathbf{T} | [b]; \phi^s \rangle = \delta_{m,s} T_{ab}^{(\phi^m)} \quad (\text{A14})$$

are well-defined expressions independent of the representatives of $[a]$ and $[b]$ we choose and are the S and T matrices of the condensed theory in this basis for the torus Hilbert space.

We now discuss gauge freedom very carefully. In the uncondensed theory, there is a preferred basis $|a; \phi^m\rangle$ of the Hilbert space, where a is any anyon. For $m = 0$, these states are invariant under gauge transformations of the fusion spaces. However, there is still some gauge freedom, in the sense that we could send $|a; 1\rangle \rightarrow e^{i\alpha_a} |a; 1\rangle$ for some phases; this gauge

freedom modifies the S matrix. We can canonically fix this gauge freedom up to a global phase rotation (physically, a gauge choice for the vacuum state) by demanding that S_{1a} be real and positive for all a and also that S_{ab} be symmetric. Another way to say this is that we could be handed some phase-rotated states $|a; 1\rangle$ from the outset; we would observe that we have the “wrong” basis because the resulting S matrix would not have these nice properties, and this could be corrected with a (diagonal) basis transformation.

For $m > 0$, there is no such canonical basis for these states. The situation is in some sense worse, because changing the basis of the fusion spaces induces a particular change of basis on the punctured torus Hilbert space, of the form Eq. (A1), and therefore changes the punctured S matrix of the uncondensed theory. This naively seems disturbing because the S matrix of the condensed theory appears to depend on a fusion space basis in the original theory. However, from the above perspective, there is no such problem; such a change of basis of fusion spaces in the uncondensed theory is just a particular special case of changing the basis for the torus Hilbert space of the condensed theory. We simply imagine that we are handed the states $|a; \phi^n\rangle$ in some fixed but noncanonical basis, and then inspect the S matrix of the condensed theory; if it does not have the intended properties, we were handed the “wrong” basis and should perform a torus Hilbert space basis transformation to fix it.

There is one subtlety here if $1 < \ell_a < n$, which is that we must gauge-fix

$$W_\beta(\phi) |a; \phi^m\rangle = |a \times \phi; \phi^m\rangle \quad (\text{A15})$$

in the uncondensed theory. This is needed in order to ensure that the state $|(a); \phi^m\rangle$ in Eq. (A6) is indeed symmetric under insertion of ϕ Wilson loops.

Having discussed gauge freedom, we now need to check if our states in the condensed theory are in the canonical gauge. For $\ell_a = \ell_b = n$, then $\hat{S}_{[a],[b]} \propto S_{a,b}$, so \hat{S} inherits its nice properties from S ; these states are in the correct gauge. For $\ell_a < n$, however, we are certainly not in the correct gauge since in this basis

$$\hat{S}_{1,([a],k)} = \begin{cases} \sqrt{2} S_{1,a} & k = 0 \\ 0 & \text{else} \end{cases}. \quad (\text{A16})$$

On general grounds, quasiparticles must correspond to some superposition of states $|(a); \phi^m\rangle$ with fixed $[a]$, so we may restrict our attention to a fixed- $[a]$ sector.

At this point we restrict ourselves to $n = 2$. Then if $\ell_a = 1$, we have $a \times \phi = a$ and a general basis transformation is of the form

$$\begin{pmatrix} |[a]; +\rangle \\ |[a]; -\rangle \end{pmatrix} = U^{(a)} \begin{pmatrix} |[a]; 1\rangle \\ |[a]; \phi\rangle \end{pmatrix} = e^{ia_a} \begin{pmatrix} e^{-i(\beta_a + \delta_a)/2} \cos(\gamma_a/2) & -e^{-i(\beta_a - \delta_a)/2} \sin(\gamma_a/2) \\ e^{i(\beta_a - \delta_a)/2} \sin(\gamma_a/2) & e^{i(\beta_a + \delta_a)/2} \cos(\gamma_a/2) \end{pmatrix} \begin{pmatrix} |[a]; 1\rangle \\ |[a]; \phi\rangle \end{pmatrix}. \quad (\text{A17})$$

Strictly speaking $U^{(a)}$ is a diagonal block in the transformation of the entire Hilbert space; the state $|1; 1\rangle$ does not transform in this basis transformation. Accordingly,

$$(\hat{S}_{1,(a,+)}, \hat{S}_{1,(a,-)}) = (\hat{S}_{1,(a,1)}, \hat{S}_{1,(a,\phi)}) (U^{(a)})^\dagger \quad (\text{A18})$$

$$= \sqrt{2} S_{1,a} e^{-i(\alpha - \delta/2)} (e^{i\beta/2} \cos(\gamma/2), e^{-i\beta/2} \sin(\gamma/2)). \quad (\text{A19})$$




FIG. 2. Diagrammatic manipulation in the standard BFC graphical calculus (see, e.g., [44] or [1] for reviews and [40,41] for graphical representations of the punctured S matrix) leading to Eq. (A25). Blue lines are a or \bar{a} orange are b or \bar{b} , and green is ϕ . We are assuming $\phi \times \phi = 1$ and that $a \times \phi = a$, $b \times \phi = b$. “Inverse for b ” means to repeat the same process as the first three steps, but in reverse and moving the ϕ line attached to the $|b, \bar{b}; \phi\rangle$ vertex. In the step labeled “isotopy”, we use the fact that the double braid of ϕ is trivial with both a and b .

We require that

$$d_{(a,+)} + d_{(a,-)} = \frac{\hat{S}_{1,(a,+)}}{\hat{S}_{1,1}} + \frac{\hat{S}_{1,(a,-)}}{\hat{S}_{1,1}} = \frac{S_{1,a}}{S_{1,1}} = d_a \quad (\text{A20})$$

following the usual rules for preserving the quantum dimension of split particles. Using $\hat{S}_{1,1} = 2S_{1,1}$, equating the magnitudes leads to

$$\cos \beta_a \sin \gamma_a = 1. \quad (\text{A21})$$

That is, $\gamma = \pi/2$ and $\beta = 0$ or $\gamma = 3\pi/2$ and $\beta = \pi$. We now ensure that $S_{1,(a,\pm)}$ are real and positive. In the first solution, we find this requires $\alpha - \delta/2 = 0 \bmod 2\pi$, while in the second we find $\alpha - \delta/2 = 3\pi/2 \bmod 2\pi$. Substituting back, we find that these two solutions are related by switching the rows of $U^{(a)}$, so that up to a basis reordering the only solutions for $U^{(a)}$ are

$$U^{(a)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & e^{i\delta_a} \\ 1 & -e^{i\delta_a} \end{pmatrix}. \quad (\text{A22})$$

Note that δ_a is exactly the gauge freedom in $|a; \phi\rangle$.

Next, we demand that \hat{S} is symmetric. In this basis,

$$\begin{aligned} \hat{S}_{(a,\pm),(b,\pm)} &= U^{(a)} \begin{pmatrix} S_{ab} & 0 \\ 0 & S_{ab}^{(\phi)} \end{pmatrix} (U^{(b)})^\dagger \\ &= \frac{1}{2} \begin{pmatrix} S_{ab} + e^{i(\delta_a - \delta_b)} S_{ab}^{(\phi)} & S_{ab} - e^{i(\delta_a - \delta_b)} S_{ab}^{(\phi)} \\ S_{ab} - e^{i(\delta_a - \delta_b)} S_{ab}^{(\phi)} & S_{ab} + e^{i(\delta_a - \delta_b)} S_{ab}^{(\phi)} \end{pmatrix}. \end{aligned} \quad (\text{A23})$$

For $a = b$ this is clearly symmetric and independent of our remaining gauge freedom. If $a \neq b$, then symmetry of \hat{S} amounts to a choice

$$e^{2i(\delta_a - \delta_b)} = \frac{S_{ba}^{(\phi)}}{S_{ab}^{(\phi)}}. \quad (\text{A24})$$

Equivalently, we may think of this as fixing a gauge so that the *uncondensed* punctured S matrix $S_{ab}^{(\phi)}$ is symmetric in a and b . If we are already in such a gauge, then we may choose $d_a = d_b = 0$, which reduces to the results of Ref. [33].

To prove that such a gauge exists, we perform the diagrammatic manipulation in Fig. 2 to show

$$S_{ab}^{(\phi)} = \frac{d_a F_{a,\phi,1}^{a,\bar{a},a} F_{a,1,a}^{\phi,\phi,a} F_{\phi,a,1}^{\phi,a,\bar{a}}}{d_b F_{b,\phi,1}^{b,\bar{b},b} F_{b,1,b}^{\phi,\phi,b} F_{\phi,b,1}^{\phi,b,\bar{b}}} S_{ba}^{(\phi)}. \quad (\text{A25})$$

Letting

$$e^{2i\delta_a} = d_a F_{a,\phi,1}^{a,\bar{a},a} F_{a,1,a}^{\phi,\phi,a} F_{\phi,a,1}^{\phi,a,\bar{a}} \quad (\text{A26})$$

we obtain the gauge transformation, which symmetrizes the punctured S matrix.

Notice that there is some residual gauge freedom preserving the symmetry of the uncondensed punctured S matrix; we may send $e^{i\delta_a} \rightarrow -e^{i\delta_a}$. This gauge transformation simply flips the role of the rows in $U^{(a)}$ and thus corresponds to relabeling $(a, \pm) \rightarrow (a, \mp)$.

APPENDIX B: PROOF OF PROPOSITION VI.1

The proof is quite involved, so we begin with an outline of the strategy. First, for $\delta v \in \mathbb{Z}_{16}$, let $\mathcal{I}(\delta v)$ be the minimal modular extension of $\{1, \psi\}$ with chiral central charge

$c_- = \delta v/2 \bmod 8$. Then the 16-fold way tells us that, given $\check{\mathcal{C}}_{v_1}$, the minimal modular extension $\check{\mathcal{C}}_{v_2}$ is constructed as

$$\check{\mathcal{C}}_{v_2} = \check{\mathcal{C}}_{v_1} \boxtimes \mathcal{I}(\delta v)/\{\psi\psi \sim 1\} \quad (\text{B1})$$

where the quotient means that we condense the bound state of the preferred fermions in $\check{\mathcal{C}}_{v_1}$ and $\mathcal{I}(\delta v)$. The proof goes in four steps:

Step 1: Use the results of [33], which we review in expanded detail in Appendix A, to derive the anyon content and the S - and T -matrices of the condensed theory $\check{\mathcal{C}}_{v_2}$.

Step 2: Directly construct an anyon permutation $\check{\rho}_{\mathbf{g}}^{(2)}$ on $\check{\mathcal{C}}_{v_2}$ that lifts the permutation action of $\rho_{\mathbf{g}}$.

Step 3: Show that $\check{\rho}_{\mathbf{g}}^{(2)}$ preserves the S and T matrices of $\check{\mathcal{C}}_{v_2}$.

Step 4: Compute the permutation action of $o_2^{(2)}(\mathbf{g}, \mathbf{h})$.

Step 1: According to [33], the anyon content of $\check{\mathcal{C}}_{v_1+1}$ can be labeled by equivalence classes of anyons $(a, x) \in \check{\mathcal{C}}_{v_1} \boxtimes \mathcal{I}(\delta v)$ and, in some cases, a sign. In all cases, the anyon sector $(\check{\mathcal{C}}_{v_2})_0 \sim \mathcal{C}$ consists of the equivalence classes $(a_0, 1) \sim (a_0 \times \psi, \psi)$ for $a_0 \in (\check{\mathcal{C}}_{v_1})_0 = \mathcal{C}$. The behavior of the fermion parity vortex sector depends on the parity of δv .

If δv is even, then $\mathcal{I}(\delta v)$ is Abelian, and its particles can be labeled $\{1, \psi, v, v \times \psi\}$. Then the fermion parity vortices of $\check{\mathcal{C}}_{v_2}$ are as follows:

(1) $(a_v, v) \sim (a_v \times \psi, v \times \psi)$ for $a_v \in (\check{\mathcal{C}}_{v_1})_v$. These form the sector $(\check{\mathcal{C}}_{v_2})_v$.

(2) $(a_{\sigma}, v) \sim (a_{\sigma}, v \times \psi)$ for $a_{\sigma} \in (\check{\mathcal{C}}_{v_1})_{\sigma}$. These form the sector $(\check{\mathcal{C}}_{v_2})_{\sigma}$.

Notice that anyon labels in $\check{\mathcal{C}}_{v_1}$ are in one-to-one correspondence with anyon labels in $\check{\mathcal{C}}_{v_2}$.

If δv is odd, the particles of $\mathcal{I}(\delta v)$ obey Ising fusion rules. Labeling the particles by $\{1, \psi, \sigma\}$, the fermion parity vortices of $\check{\mathcal{C}}_{v_2}$ are as follows:

(1) $(a_v, \sigma) \sim (a_v \times \psi, \sigma)$ for $a_v \in (\check{\mathcal{C}}_{v_1})_v$. These form the sector $(\check{\mathcal{C}}_{v_1+1})_{\sigma}$.

(2) $(a_{\sigma}, \sigma)_{\pm}$ for $a_{\sigma} \in (\check{\mathcal{C}}_{v_1})_{\sigma}$. These form the sector $(\check{\mathcal{C}}_{v_1+1})_v$ with $(a_{\sigma}, \sigma)_+ \times \psi = (a_{\sigma}, \sigma)_-$. We say in this case that (a_{σ}, σ) splits after condensation.

The T matrix of the condensed theory is simple: the topological spin of (a, x) [including the split case $(a_{\sigma}, \sigma)_{\pm}$] is $\theta_a \theta_x$.

$$S_{(a,x)\pm, (b,y)\pm}^{\check{\mathcal{C}}_{v_2}} = e^{-2\pi i \delta v / 8} \frac{S_{a,b}^{\check{\mathcal{C}}_{v_1}, (\psi)}}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}$$

where $S_{a,b}^{\check{\mathcal{C}}_{v_1}, (\psi)}$ is the punctured S matrix of $\check{\mathcal{C}}_{v_1}$. Thanks to our gauge fixing for the punctured S matrix in the uncondensed theory, $S^{\check{\mathcal{C}}_{v_2}}$ is symmetric.

Step 2: Define the permutation action $\check{\rho}_{\mathbf{g}}^{(2)}$ as follows:

$$\begin{aligned} \check{\rho}_{\mathbf{g}}^{(2)}((a, x)) &= (\check{\rho}_{\mathbf{g}}^{(1)}(a), x) && \text{if } (a, x) \text{ does not split} \\ \check{\rho}_{\mathbf{g}}^{(2)}((a, x)_{\pm}) &= (\check{\rho}_{\mathbf{g}}^{(1)}(a), x)_{\pm} \times \psi^{z_a(\mathbf{g})} && \text{if } (a, x) \text{ splits} \end{aligned} \quad (B7)$$

where $z_a(\mathbf{g}) \in \{0, 1\} \simeq \mathbb{Z}_2$ must be defined for each $a \in \check{\mathcal{C}}_{\sigma}^{(1)}$. It is immediate that $\check{\rho}_{\mathbf{g}}^{(2)}$ lifts the permutation action of $\rho_{\mathbf{g}}$ as long as $\check{\rho}_{\mathbf{g}}^{(1)}$ does as well.

In order to define $z_a(\mathbf{g})$, first calculate directly

$$\check{\rho}_{\mathbf{g}}(S_{ab}^{\check{\mathcal{C}}_{v_1}, (\psi)}) = U_{\mathbf{g}}(\mathbf{g}a, \mathbf{g}\bar{a}; \psi) U_{\mathbf{g}}^*(\mathbf{g}a, \mathbf{g}\bar{a}; 1) U_{\mathbf{g}}^*(\mathbf{g}b, \mathbf{g}\bar{b}; \psi) U_{\mathbf{g}}(\mathbf{g}b, \mathbf{g}\bar{b}; 1) S_{\mathbf{g}a, \mathbf{g}b}^{\check{\mathcal{C}}_{v_1}, (\psi)} = S_{a,b}^{\check{\mathcal{C}}_{v_1}, (\psi)}. \quad (B8)$$

In our particular gauge, the punctured S matrix is symmetric in a and b . Hence we can write

$$\check{\rho}_{\mathbf{g}}(S_{ab}^{\check{\mathcal{C}}_{v_1}, (\psi)}) = \check{\rho}_{\mathbf{g}}(S_{ba}^{\check{\mathcal{C}}_{v_1}, (\psi)}) = U_{\mathbf{g}}(\mathbf{g}b, \mathbf{g}\bar{b}; \psi) U_{\mathbf{g}}^*(\mathbf{g}b, \mathbf{g}\bar{b}; 1) U_{\mathbf{g}}^*(\mathbf{g}a, \mathbf{g}\bar{a}; \psi) U_{\mathbf{g}}(\mathbf{g}a, \mathbf{g}\bar{a}; 1) S_{\mathbf{g}b, \mathbf{g}a}^{\check{\mathcal{C}}_{v_1}, (\psi)}. \quad (B9)$$

But $S_{\mathbf{g}b, \mathbf{g}a}^{\check{\mathcal{C}}_{v_1}, (\psi)} = S_{\mathbf{g}a, \mathbf{g}b}^{\check{\mathcal{C}}_{v_1}, (\psi)}$, so Eqs. (B8) and (B9) combine to

$$[U_{\mathbf{g}}(\mathbf{g}b, \mathbf{g}\bar{b}; \psi) U_{\mathbf{g}}^*(\mathbf{g}b, \mathbf{g}\bar{b}; 1)]^2 = [U_{\mathbf{g}}(\mathbf{g}a, \mathbf{g}\bar{a}; \psi) U_{\mathbf{g}}^*(\mathbf{g}a, \mathbf{g}\bar{a}; 1)]^2. \quad (B10)$$

Equivalently, if we fix a reference σ -type vortex r_σ , then we must have

$$z_a(\mathbf{g}) = \frac{U_{\mathbf{g}}(\mathbf{g}a, \mathbf{g}\bar{a}; 1)U_{\mathbf{g}}^*(\mathbf{g}a, \mathbf{g}\bar{a}; \psi)}{U_{\mathbf{g}}(\mathbf{g}r_\sigma, \mathbf{g}\bar{r}_\sigma; 1)U_{\mathbf{g}}^*(\mathbf{g}r_\sigma, \mathbf{g}\bar{r}_\sigma; \psi)} \in \mathbb{Z}_2 \quad (\text{B11})$$

where we are slightly abusing notation; the above defines $z_a(\mathbf{g}) \in \{1, -1\} \simeq \mathbb{Z}_2$ instead of $\{0, 1\} \simeq \mathbb{Z}_2$. We emphasize that the entire analysis above requires a be a σ -type vortex.

The above proof actually does not apply if $S_{ab}^{\check{C}_{v_1}, (\psi)} = 0$. As in Sec. VIB, we can break the S matrix of $(\check{C}_{v_2})_v$ into k blocks and apply the above argument to each block separately; we have a separate choice of r_σ for each block.

We choose r_σ to be \mathbf{g} -independent. As we will see, such a choice will lead to a permutation $o^{(2)}(\mathbf{g}, \mathbf{h})$ that is exactly the identity; instead choosing a \mathbf{g} -dependent r_σ in each block will amount to modifying the permutation action of $o^{(2)}(\mathbf{g}, \mathbf{h})$ by a $\text{ker } r$ -valued coboundary.

Step 3: It is immediately obvious that $\check{\rho}_{\mathbf{g}}^{(2)}$ preserves the T matrix provided $\check{\rho}_{\mathbf{g}}^{(1)}$ does. Also, given the fact that $\check{\rho}_{\mathbf{g}}^{(1)}$ preserves the S matrix of \check{C}_{v_1} , Eqs. (B2) and (B3) immediately imply that $\check{\rho}_{\mathbf{g}}^{(2)}$ preserves $S_{(a,x), (b,y)}^{\check{C}_{v_2}}$ when at most one of (a, x) and (b, y) split. The case where both split requires some calculation.

From Eq. (B8), we see that $\check{\rho}_{\mathbf{g}}^{(1)}(S_{a,b}^{\check{C}_{v_1}, (\psi)}) = z_a(\mathbf{g})z_b(\mathbf{g})S_{\mathbf{g}a, \mathbf{g}b}^{\check{C}_{v_1}, (\psi)} = S_{a,b}^{\check{C}_{v_1}, (\psi)}$. Hence

$$\check{\rho}_{\mathbf{g}}^{(2)}(S_{(a,x)\pm, (b,y)\pm}^{\check{C}_{v_2}}) = \frac{e^{-2\pi i \delta v/8}}{\sqrt{2}} \times z_a(\mathbf{g})z_b(\mathbf{g})S_{\mathbf{g}a \mathbf{g}b}^{\check{C}_{v_1}, (\psi)} \times \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}. \quad (\text{B12})$$

On the other hand, $\check{\rho}_{\mathbf{g}}^{(2)}((a, \sigma)_\pm) = (a, \sigma)_{\pm z_a(\mathbf{g})}$. Hence,

$$S_{\check{\rho}_{\mathbf{g}}^{(2)}((a,x)_\pm) \check{\rho}_{\mathbf{g}}^{(2)}((b,y)_\pm)}^{\check{C}_{v_2}} = S_{(\mathbf{g}a, x)_{\pm z_a(\mathbf{g})} (\mathbf{g}b, y)_{\pm z_b(\mathbf{g})}}^{\check{C}_{v_2}} = \frac{e^{-2\pi i \delta v/8}}{\sqrt{2}} S_{\mathbf{g}a \mathbf{g}b}^{\check{C}_{v_1}, (\psi)} \times z_a(\mathbf{g})z_b(\mathbf{g}) \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} = \check{\rho}_{\mathbf{g}}^{(2)}(S_{(a,x)_\pm, (b,y)_\pm}^{\check{C}_{v_2}}) \quad (\text{B13})$$

where the factor of $z_a(\mathbf{g})$ is interpreted as permuting the rows of the matrix if it is -1 and the factor of $z_b(\mathbf{g})$ should be interpreted as permuting the columns if it is -1 . Hence $\check{\rho}_{\mathbf{g}}^{(2)}$ indeed preserves the S matrix.

Step 4: By direct computation,

$$\begin{aligned} (\check{\rho}_{\mathbf{gh}}^{(2)})^{-1} \check{\rho}_{\mathbf{g}}^{(2)} \check{\rho}_{\mathbf{h}}^{(2)}((a, x)) &= (o_2^{(1)}(\mathbf{g}, \mathbf{h})(a), x) \text{ if } (a, x) \text{ does not split} \\ (\check{\rho}_{\mathbf{gh}}^{(2)})^{-1} \check{\rho}_{\mathbf{g}}^{(2)} \check{\rho}_{\mathbf{h}}^{(2)}((a, x)_\pm) &= (a, x)_\pm \times \psi^{z_a(\mathbf{g}) + z_a(\mathbf{h}) - z_a(\mathbf{gh})} \text{ if } (a, x) \text{ splits.} \end{aligned} \quad (\text{B14})$$

If δv is even, then no particles split, and $(o_2^{(1)}(\mathbf{g}, \mathbf{h})(a), x) = \psi \times (a, x)$ if and only if $o_2^{(1)}(\mathbf{g}, \mathbf{h})(a) = \psi \times a$. Hence $o_2^{(1)}$ and $o_2^{(2)}$, as permutations, are identical on the cochain level.

If δv is odd, then $(o_2^{(1)}(\mathbf{g}, \mathbf{h})(a), x) \sim (a, x)$ for all $a \in \check{C}_{v_1}$. In particular, if $o_2^{(1)}$ acts nontrivially on a , then $a \in (\check{C}_{v_1})_v$, and we saw that in such a case $(a, \sigma) \sim (a \times \psi, \sigma)$. Hence the only permutation action comes from split particles, and by inspection, modifying the reference vortex r_σ in a \mathbf{g} -dependent way changes $o^{(2)}(\mathbf{g}, \mathbf{h})$ by a $\text{ker } r$ -valued coboundary. We can directly calculate from Eq. (84) that

$$\begin{aligned} o^{(2)}(\mathbf{g}, \mathbf{h})((a_\sigma, \sigma)_\pm) &= \frac{U_{\mathbf{g}}^*(\mathbf{g}^{\mathbf{h}} a_\sigma, \mathbf{g}^{\mathbf{h}} \bar{a}_\sigma; 1)U_{\mathbf{g}}(\mathbf{g}^{\mathbf{h}} a_\sigma, \mathbf{g}^{\mathbf{h}} \bar{a}_\sigma; \psi)}{U_{\mathbf{g}}^*(\mathbf{g}^{\mathbf{h}} r_\sigma, \mathbf{g}^{\mathbf{h}} \bar{r}_\sigma; 1)U_{\mathbf{g}}(\mathbf{g}^{\mathbf{h}} r_\sigma, \mathbf{g}^{\mathbf{h}} \bar{r}_\sigma; \psi)} \times \frac{U_{\mathbf{h}}^*(\mathbf{h} a_\sigma, \mathbf{h} \bar{a}_\sigma; 1)U_{\mathbf{h}}(\mathbf{h} a_\sigma, \mathbf{h} \bar{a}_\sigma; \psi)}{U_{\mathbf{h}}^*(\mathbf{h} r_\sigma, \mathbf{h} \bar{r}_\sigma; 1)U_{\mathbf{h}}(\mathbf{h} r_\sigma, \mathbf{h} \bar{r}_\sigma; \psi)} \\ &\quad \times \frac{U_{\mathbf{gh}}(\mathbf{g}^{\mathbf{h}} a_\sigma, \mathbf{g}^{\mathbf{h}} \bar{a}_\sigma; 1)U_{\mathbf{gh}}^*(\mathbf{g}^{\mathbf{h}} a_\sigma, \mathbf{g}^{\mathbf{h}} \bar{a}_\sigma; \psi)}{U_{\mathbf{gh}}(\mathbf{g}^{\mathbf{h}} r_\sigma, \mathbf{g}^{\mathbf{h}} \bar{r}_\sigma; 1)U_{\mathbf{gh}}^*(\mathbf{g}^{\mathbf{h}} r_\sigma, \mathbf{g}^{\mathbf{h}} \bar{r}_\sigma; \psi)} \end{aligned} \quad (\text{B15})$$

$$= \frac{\kappa_{\mathbf{g}, \mathbf{h}}(\mathbf{g}^{\mathbf{h}} a_\sigma, \mathbf{g}^{\mathbf{h}} \bar{a}_\sigma; 1)\kappa_{\mathbf{g}, \mathbf{h}}^*(\mathbf{g}^{\mathbf{h}} a_\sigma, \mathbf{g}^{\mathbf{h}} \bar{a}_\sigma; \psi)}{\kappa_{\mathbf{g}, \mathbf{h}}(\mathbf{g}^{\mathbf{h}} r_\sigma, \mathbf{g}^{\mathbf{h}} \bar{r}_\sigma; 1)\kappa_{\mathbf{g}, \mathbf{h}}^*(\mathbf{g}^{\mathbf{h}} r_\sigma, \mathbf{g}^{\mathbf{h}} \bar{r}_\sigma; \psi)} \quad (\text{B16})$$

$$= +1. \quad (\text{B17})$$

where the last line comes from decomposing κ as a product of anyon-dependent factors β . Hence $\check{\rho}_{\mathbf{g}}^{(2)}$ is a group homomorphism $G_b \rightarrow P(\check{C}_{v_2})$.

APPENDIX C: DOUBLED SU(2)₆

We write down the UMTC data for $\check{C} = \text{SU}(2)_6 \times \text{SU}(2)_6 \times \text{Ising}_{-9/2}/\{\psi\psi \sim 1\}$. Here $\text{Ising}_{-9/2}$ is the minimal modular extension of $\{1, \psi\}$ with central charge $c_- = -9/2$. The quotient means that we condense pairs of preferred fermions in these three spin modular theories, i.e., we condense (6,6,0), (0,6,ψ), and (6,0,ψ), where the first two labels label particles in the two copies of $\text{SU}(2)_6$ and the third labels particles in the Ising theory.

$\check{\mathcal{C}}$ contains 14 particles. Labeling the particles of $SU(2)_6$ by integers from 0 to 6 in the usual way, and labeling elements of Ising as $\{0, \sigma, \psi\}$, the deconfined particles of the theory, their topological twists, and quantum dimensions are given in Table I.

The particle (6,0,0) is the preferred fermion of this spin modular theory. All labels are redundant under fusion with (6,6,0), (0, 6, ψ), and (6, 0, ψ) in the product theory before condensation. We have

$$\check{\mathcal{C}}_v = \{(1, 3, \sigma)_{\pm}, (3, 1, \sigma)_{\pm}\} \quad (C1)$$

$$\check{\mathcal{C}}_{\sigma} = \{(1, 1, \sigma), (3, 3, \sigma)\} \quad (C2)$$

and all other particles are in $\check{\mathcal{C}}_0$. The total quantum dimension is $\mathcal{D} = 4(2 + \sqrt{2})$.

With $d = 1 + \sqrt{2}$ and the quasiparticles ordered as in Table I, the modular data of $\check{\mathcal{C}}$ is:

$$S = \frac{1}{\mathcal{D}} \begin{pmatrix} 1 & d & d & d & d & d^2 & d^2 & 1 & \sqrt{2}d & \sqrt{2}d & \sqrt{2}d & 2d & 2d \\ d & -1 & d^2 & -1 & d^2 & -d & -d & d & \sqrt{2}d & \sqrt{2}d & -\sqrt{2}d & -\sqrt{2}d & 2d \\ d & d^2 & -1 & d^2 & -1 & -d & -d & d & -\sqrt{2}d & -\sqrt{2}d & \sqrt{2}d & \sqrt{2}d & 2d \\ d & -1 & d^2 & -1 & d^2 & -d & -d & d & -\sqrt{2}d & -\sqrt{2}d & \sqrt{2}d & \sqrt{2}d & -2d \\ d & d^2 & -1 & d^2 & -1 & -d & -d & d & \sqrt{2}d & \sqrt{2}d & -\sqrt{2}d & -\sqrt{2}d & 2d \\ d^2 & -d & -d & -d & -d & 1 & 1 & d^2 & -\sqrt{2}d & -\sqrt{2}d & -\sqrt{2}d & -\sqrt{2}d & 2d \\ d^2 & -d & -d & -d & -d & 1 & 1 & d^2 & \sqrt{2}d & \sqrt{2}d & \sqrt{2}d & \sqrt{2}d & -2d \\ 1 & d & d & d & d & d^2 & d^2 & 1 & -\sqrt{2}d & -\sqrt{2}d & -\sqrt{2}d & -\sqrt{2}d & -2d \\ \sqrt{2}d & \sqrt{2}d & -\sqrt{2}d & -\sqrt{2}d & \sqrt{2}d & -\sqrt{2}d & \sqrt{2}d & -\sqrt{2}d & 2\sqrt{2}d & -2\sqrt{2}d & 0 & 0 & 0 \\ \sqrt{2}d & \sqrt{2}d & -\sqrt{2}d & -\sqrt{2}d & \sqrt{2}d & -\sqrt{2}d & \sqrt{2}d & -\sqrt{2}d & -2\sqrt{2}d & 2\sqrt{2}d & 0 & 0 & 0 \\ \sqrt{2}d & -\sqrt{2}d & \sqrt{2}d & \sqrt{2}d & -\sqrt{2}d & -\sqrt{2}d & \sqrt{2}d & -\sqrt{2}d & 0 & 0 & 2\sqrt{2}d & -2\sqrt{2}d & 0 \\ \sqrt{2}d & -\sqrt{2}d & \sqrt{2}d & \sqrt{2}d & -\sqrt{2}d & -\sqrt{2}d & \sqrt{2}d & -\sqrt{2}d & 0 & 0 & -2\sqrt{2}d & 2\sqrt{2}d & 0 \\ 2d & 2d & 2d & -2d & -2d & 2d & -2d & -2d & 0 & 0 & 0 & 0 & 0 \\ 2d & -2d & -2d & 2d & 2d & -2d & -2d & -2d & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad (C3)$$

$$T = \text{diag}(1, i, i, -i, -i, -1, 1, -1, 1, 1, 1, 1, e^{5\pi i/4}, e^{3\pi i/4}) \quad (C4)$$

There are two permutation actions on the anyons, which lift the action of \mathbf{T} on \mathcal{C} , complex conjugate the modular data, and square to the identity. Listed as a 14×14 matrix acting on anyon labels, these permutations are

$$P_1 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad (C5)$$

$$P_2 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}. \quad (C6)$$

Two other anyon permutations lift the action on \mathcal{C} , complex conjugate the modular data, but yield a \mathbb{Z}_4 action; they are

$$P_3 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}, \quad (C7)$$

$$P_4 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}. \quad (C8)$$

APPENDIX D: CHANGE OF $[o_3]$ UNDER CHANGE IN LIFT

Fix a minimal modular extension $\check{\mathcal{C}}$, and assume $\ker r = \mathbb{Z}_2$. Given symmetry fractionalization data on \mathcal{C} , suppose we have a lift $\check{\rho}_g$ for which $o_2 = 0$. Then if $\pi : G \rightarrow \mathbb{Z}_2 \cong \{0, 1\}$ is a group homomorphism, all other valid lifts are obtained (up to locality-respecting natural isomorphism) by writing

$$\check{\rho}_g^V = \alpha_{\psi}^{\pi(g)} \circ \check{\rho}_g. \quad (D1)$$

In this Appendix, we reproduce [36] the calculation¹² for how $[o_3^{(V, \check{\rho}^V)}]$ is related to $[o_3^{(V, \check{\rho})}]$, which is a special case of Eq. (163). We will also show that o_3 is invariant under various gauge choices.

¹²We thank Parsa Bonderson for sharing additional unpublished notes.

1. Change of o_3

Recall our setup from Sec. VII; we choose a gauge where $r(\check{\rho}_g) = \rho_g$ on the nose, and we can choose a gauge where the phases β_a on \mathcal{C} are just the restriction of $\check{\beta}_a$. We are given symmetry fractionalization data $\omega_a(\mathbf{g}, \mathbf{h}) \in C^2(G_b, K(\mathcal{C}))$ on \mathcal{C} , which is guaranteed in this gauge to lift to some $\check{\omega}_a \in C^2(G_b, K(\check{\mathcal{C}}))$.

Now let us compute how this data changes for $\check{\rho}_g^V$.

First, we need the following facts about α_ψ . Let φ be an arbitrary topological (anti-)autoequivalence. Then from Eq. (105), one can compute that

$$\alpha_\psi \circ \varphi = \Upsilon_\varphi \circ \varphi \circ \alpha_\psi \quad (D2)$$

$$\alpha_\psi^2 = \xi \quad (D3)$$

where Υ_φ and ξ are natural isomorphisms defined by their actions on anyons

$$\gamma_{\Upsilon_\varphi, a_x} = U_\varphi(\psi^x, a_x, a_x \psi^x) \quad (D4)$$

$$\gamma_{\xi, a_x} = i^x. \quad (D5)$$

Here we are denoting $a_x \in \check{\mathcal{C}}_x$ for $x \in \{0, 1\}$. We also note that when treating π as an *integer*-valued function, it must obey

$$\pi(\mathbf{g}) + \pi(\mathbf{h}) = \pi(\mathbf{g}\mathbf{h}) + 2\pi(\mathbf{g})\pi(\mathbf{h}). \quad (D6)$$

Next, we need to compute how $\check{\kappa}_{\mathbf{g}, \mathbf{h}}$ changes.

$$\check{\kappa}_{\mathbf{g}, \mathbf{h}}^V = \alpha_\psi^{\pi(\mathbf{g}\mathbf{h})} \check{\rho}_{\mathbf{g}\mathbf{h}} \check{\rho}_{\mathbf{h}}^{-1} \alpha_\psi^{-\pi(\mathbf{h})} \check{\rho}_{\mathbf{g}}^{-1} \alpha_\psi^{-\pi(\mathbf{g})} \quad (D7)$$

$$= \alpha_\psi^{\pi(\mathbf{g}\mathbf{h})} \check{\rho}_{\mathbf{g}\mathbf{h}} \check{\rho}_{\mathbf{h}}^{-1} \check{\rho}_{\mathbf{g}}^{-1} \alpha_\psi^{-\pi(\mathbf{h})} \Upsilon_g^{\pi(\mathbf{h})} \alpha_\psi^{-\pi(\mathbf{g})} \quad (D8)$$

$$= \alpha_\psi^{\pi(\mathbf{g}\mathbf{h})} \check{\kappa}_{\mathbf{g}, \mathbf{h}} \alpha_\psi^{-\pi(\mathbf{h})} \alpha_\psi^{-\pi(\mathbf{g})} \Upsilon_g^{\pi(\mathbf{h})} \Upsilon_{\gamma_g^{-\pi(\mathbf{h})}}^{-\pi(\mathbf{g})} \quad (D9)$$

$$= \Upsilon_{\check{\kappa}_{\mathbf{g}, \mathbf{h}}}^{\pi(\mathbf{g}\mathbf{h})} \check{\kappa}_{\mathbf{g}, \mathbf{h}} \alpha_\psi^{\pi(\mathbf{g}\mathbf{h})} \xi^{-\pi(\mathbf{g})\pi(\mathbf{h})} \alpha_\psi^{-\pi(\mathbf{g}\mathbf{h})} \Upsilon_g^{\pi(\mathbf{h})} \Upsilon_{\gamma_g^{-\pi(\mathbf{h})}}^{-\pi(\mathbf{g})} \quad (D10)$$

$$= \Upsilon_{\check{\kappa}_{\mathbf{g}, \mathbf{h}}}^{\pi(\mathbf{g}\mathbf{h})} \xi^{-\pi(\mathbf{g})\pi(\mathbf{h})} \Upsilon_g^{\pi(\mathbf{h})} \Upsilon_{\gamma_g^{-\pi(\mathbf{h})}}^{-\pi(\mathbf{g})} \check{\kappa}_{\mathbf{g}, \mathbf{h}} \quad (D11)$$

where we have repeatedly commuted α_ψ through other maps, used the fact that

$$\alpha_\psi^{\pi(\mathbf{g})} \alpha_\psi^{\pi(\mathbf{h})} = \alpha_\psi^{\pi(\mathbf{g}\mathbf{h}) + 2\pi(\mathbf{g})\pi(\mathbf{h})}, \quad (D12)$$

and used the fact that $\alpha_\psi^2 = \xi$. Using the expressions for Υ_φ , this change in $\check{\kappa}_{\mathbf{g}, \mathbf{h}}$ can be used to calculate the change in $\check{\beta}_a$:

$$\begin{aligned} \check{\beta}'_{a_x} &= (-i)^{\pi(\mathbf{g})\pi(\mathbf{h})} \check{\kappa}_{\mathbf{g}, \mathbf{h}}(\psi^x, a_x, \psi^x a_x)^{\pi(\mathbf{g}\mathbf{h})} \\ &\quad \times \check{U}_{\mathbf{g}}(\psi^x, a_x, \psi^x a_x)^{\pi(\mathbf{h})} \\ &\quad \times \left[\frac{\check{U}_{\mathbf{g}}(\psi^x, a_x, \psi^x a_x)}{\check{U}_{\mathbf{g}}(\psi^x, \psi^x a_x, a_x)} \right]^{\pi(\mathbf{g})\pi(\mathbf{h})} \check{\beta}_{a_x}. \end{aligned} \quad (D13)$$

Note in particular that if $a \in \check{\mathcal{C}}_0$, then $\check{\beta}_a$ is completely unchanged. Furthermore, α_ψ is strictly the identity when restricted to \mathcal{C}_0 . Hence there is no gauge transformation on any of the data of \mathcal{C}_0 , and we can choose the lift

$$\check{\omega}_a^V = \check{\omega}_a. \quad (D14)$$

We can now calculate the change in o_3 directly. The calculation proceeds differently for σ -type and v -type vortices:

a. σ -type vortices

First note that $d\check{\omega}_a^V = d\check{\omega}_a$ since the permutation action of the symmetry on σ -type vortices is unaffected. All change in o_3 must therefore come from a change in $\check{\Omega}_a$.

If there are any σ -type vortices, then $\check{\Upsilon}_\psi$ must violate locality. Hence, since $o_2 = 0$, $\check{\beta}_\psi = \omega_2$. Simplifying Eq. (D13) with the fact that $a \times \psi = a$ for σ -type vortices, we can calculate directly that

$$\begin{aligned} \frac{\check{\Omega}_a^V}{\check{\Omega}_a} &= (-1)^{\pi_1 \cup \pi \cup \pi} \times \frac{\check{\beta}_\psi(\mathbf{h}, \mathbf{k})^{\sigma(\mathbf{g})\pi(\mathbf{h}\mathbf{k})} \check{\beta}_\psi(\mathbf{g}, \mathbf{h}\mathbf{k})^{\pi(\mathbf{g}\mathbf{h}\mathbf{k})}}{\check{\beta}_\psi(\mathbf{g}, \mathbf{h})^{\pi(\mathbf{g}\mathbf{h})} \check{\beta}_\psi(\mathbf{g}\mathbf{h}, \mathbf{k})^{\pi(\mathbf{g}\mathbf{h}\mathbf{k})}} \\ &\quad \times \frac{\check{U}_{\mathbf{h}}(\psi, \bar{\mathbf{g}}\mathbf{a}, \bar{\mathbf{g}}\mathbf{a})^{\sigma(\mathbf{g})\pi(\mathbf{k})} \check{U}_{\mathbf{g}}(\psi, a, a)^{\pi(\mathbf{h}\mathbf{k})}}{\check{U}_{\mathbf{g}}(\psi, a, a)^{\pi(\mathbf{h})} \check{U}_{\mathbf{g}\mathbf{h}}(\psi, a, a)^{\pi(\mathbf{k})}}. \end{aligned} \quad (D15)$$

The factors of $\check{\beta}$ come from decomposing $\check{\kappa}$. Manipulating the various factors of π , we obtain

$$\begin{aligned} \frac{\check{\Omega}_a^V}{\check{\Omega}_a} &= (-1)^{\pi_1 \cup \pi \cup \pi} \times \frac{\omega_2(\mathbf{h}, \mathbf{k})^{[\pi(\mathbf{h}\mathbf{k}) - \pi(\mathbf{g}\mathbf{h}\mathbf{k})]}}{\omega_2(\mathbf{g}, \mathbf{h})^{\pi(\mathbf{g}\mathbf{h}) - \pi(\mathbf{g}\mathbf{h}\mathbf{k})}} \\ &\quad \times \frac{\check{U}_{\mathbf{h}}(\psi, \bar{\mathbf{g}}\mathbf{a}, \bar{\mathbf{g}}\mathbf{a})^{\sigma(\mathbf{g})\pi(\mathbf{k})} \check{U}_{\mathbf{g}}(\psi, a, a)^{\pi(\mathbf{h}\mathbf{k}) - \pi(\mathbf{h})}}{\check{U}_{\mathbf{g}\mathbf{h}}(\psi, a, a)^{\pi(\mathbf{k})}} \end{aligned} \quad (D16)$$

$$= (-1)^{\pi_1 \cup \pi \cup \pi} \times \frac{\omega_2(\mathbf{h}, \mathbf{k})^{-\pi(\mathbf{g})}}{\omega_2(\mathbf{g}, \mathbf{h})^{-\pi(\mathbf{k})}} \times \check{\kappa}_{\mathbf{g}, \mathbf{h}}(\psi, a, a)^{-\pi(\mathbf{k})} \\ \times \check{U}_{\mathbf{g}}(\psi, a, a)^{-2\pi(\mathbf{h})\pi(\mathbf{k})}. \quad (D17)$$

The canonical gauge-fixing $F^{a\psi\psi} = +1$ enforces $\check{U}_{\mathbf{g}}(\psi, a, a) \in \{\pm 1\}$ for σ -type vortices, so so the above expression simplifies to

$$\frac{\check{\Omega}_a'}{\check{\Omega}_a} = (-1)^{\pi_1 \cup \pi \cup \pi + \pi \cup \omega_2}. \quad (D18)$$

b. v -type vortices

Notice first that because of the change in permutation action, if $a \in \check{\mathcal{C}}_v$, then

$$\frac{d\check{\omega}_a^V(\mathbf{g}, \mathbf{h}, \mathbf{k})}{d\check{\omega}_a(\mathbf{g}, \mathbf{h}, \mathbf{k})} = \left(\frac{\check{\omega}_{(\check{\rho}_g^V)^{-1}a}(\mathbf{h}, \mathbf{k})}{\check{\omega}_{(\check{\rho}_g)^{-1}a}(\mathbf{h}, \mathbf{k})} \right)^{\sigma(\mathbf{g})} \quad (D19)$$

$$= \left(\frac{\check{\omega}_{(\check{\rho}_g)^{-1}a \times \psi}(\mathbf{h}, \mathbf{k})}{\check{\omega}_{(\check{\rho}_g)^{-1}a}(\mathbf{h}, \mathbf{k})} \right)^{\sigma(\mathbf{g})} = \omega_\psi(\mathbf{h}, \mathbf{k})^{\pi(\mathbf{g})} \quad (D20)$$

where we have used the fact that $\check{\omega}_a$ obeys the fusion rules.

The calculation for the change in $\check{\Omega}_a$ simplifies dramatically with a convenient gauge fixing. One can check that there always exists a gauge transformation $\gamma_a(\mathbf{g})$ that fixes

$$\check{U}_{\mathbf{g}}(\psi, a, a \times \psi) = +1 \quad (D21)$$

for v -type vortices and is nontrivial only on $\check{\mathcal{C}}_1$. This gauge-fixing leads to all of the factors of \check{U} and $\check{\kappa}$ dropping out from Eq. (D13). Now we can calculate

carefully:

$$\check{\Omega}_a^V = \frac{\check{\beta}_{(\check{\rho}_g)^{-1}a}^V(\mathbf{h}, \mathbf{k})^{\sigma(g)} \check{\beta}_a^V(\mathbf{g}, \mathbf{h}\mathbf{k})}{\check{\beta}_a^V(\mathbf{g}, \mathbf{h}) \check{\beta}_a^V(\mathbf{g}\mathbf{h}, \mathbf{k})} \quad (\text{D22})$$

$$= (-1)^{s_1 \cup \pi \cup \pi} \frac{\check{\beta}_{(\check{\rho}_g)^{-1}a \times \psi^{\pi(g)}}(\mathbf{h}, \mathbf{k})^{\sigma(g)} \check{\beta}_a(\mathbf{g}, \mathbf{h}\mathbf{k})}{\check{\beta}_a(\mathbf{g}, \mathbf{h}) \check{\beta}_a(\mathbf{g}\mathbf{h}, \mathbf{k})} \quad (\text{D23})$$

$$= (-1)^{s_1 \cup \pi \cup \pi} \frac{\check{\beta}_{(\check{\rho}_g)^{-1}a \times \psi^{\pi(g)}}(\mathbf{h}, \mathbf{k})^{\sigma(g)}}{\check{\beta}_{(\check{\rho}_g)^{-1}a}(\mathbf{h}, \mathbf{k})^{\sigma(g)}} \check{\Omega}_a(\mathbf{g}, \mathbf{h}, \mathbf{k}) \quad (\text{D24})$$

Combining the above results, we see that

$$\frac{\check{\Omega}_a^V \overline{d\check{\omega}_a^V}}{\check{\Omega}_a d \check{\Omega}_a} = (-1)^{s_1 \cup \pi \cup \pi} \left(\frac{\check{\beta}_\psi(\mathbf{h}, \mathbf{k})}{\check{\omega}_\psi(\mathbf{h}, \mathbf{k})} \right)^{\pi(g)} \quad (\text{D25})$$

since $\check{\beta}_\psi = \beta_\psi$, $\check{\omega}_\psi = \omega_\psi$, and $\beta_\psi / \omega_\psi = \eta_\psi = \omega_2$. Accordingly,

$$o_3 \rightarrow o_3 + s_1 \cup \pi \cup \pi + \pi \cup \tilde{\omega}_2 \quad (\text{D26})$$

as expected.

2. Gauge invariances of $[o_3]$

We discuss the invariance of $[o_3]$ under various gauge transformations.

a. Invariance under locality-respecting natural isomorphisms on \mathcal{C}

If we modify $\rho_g \rightarrow \Upsilon_g \circ \rho_g$ with Υ_g a locality-respecting natural isomorphism, then ω_a is unchanged. Suppose that Υ_g is given by the anyon-dependent factors $\gamma_a(\mathbf{g})$ with $\gamma_\psi = +1$. Then we can lift Υ_g to a locality-respecting natural

isomorphism on $\check{\mathcal{C}}$ by defining anyon-dependent factors

$$\check{\gamma}_a(\mathbf{g}) = \begin{cases} \gamma_a(\mathbf{g}) & a \in \check{\mathcal{C}}_0 \\ +1 & a \in \check{\mathcal{C}}_1 \end{cases} \quad (\text{D27})$$

Under this transformation, $\check{\Omega}_a$ is invariant, and the condition that $\check{\beta}_a$ restricts to β_a is respected. Hence the lift $\check{\omega}_a$ is also unchanged, and o_3 is strictly invariant.

b. Invariance under locality-respecting natural isomorphisms on $\check{\mathcal{C}}$

Since we are demanding that $\check{\rho}_g$ restricts to ρ_g on the nose, such a locality-respecting natural isomorphism must be nontrivial only on $\check{\mathcal{C}}_1$ (modulo a ν -type gauge transformation, which we will deal with next). It is thus clear that the condition that $\check{\beta}_a$ restricts to β_a is respected, so none of the data on \mathcal{C} is modified. Hence the allowed lifts $\check{\omega}_a$ are unchanged. Also, $\check{\Omega}_a$ is simply invariant under locality-respecting natural isomorphisms. Hence o_3 is gauge invariant.

c. Invariance under ν -type gauge transformations on $\check{\mathcal{C}}$

Recall that ν -type gauge transformations modify

$$\check{\beta}_a(\mathbf{g}, \mathbf{h}) \rightarrow \check{\nu}_a(\mathbf{g}, \mathbf{h}) \check{\beta}_a(\mathbf{g}, \mathbf{h}) \quad (\text{D28})$$

$$\check{\omega}_a(\mathbf{g}, \mathbf{h}) \rightarrow \check{\nu}_a(\mathbf{g}, \mathbf{h}) \check{\omega}_a(\mathbf{g}, \mathbf{h}) \quad (\text{D29})$$

where $\check{\nu}_a(\mathbf{g}, \mathbf{h})$ obeys the fusion rules of $\check{\mathcal{C}}$. Under such a transformation, in order to maintain the condition that $\check{\beta}_a$ restricts to β_a and $\check{\omega}_a$ restricts to ω_a , we should also perform the restricted gauge transformation ν_a on \mathcal{C} , which is allowed since ν_a obeys the fusion rules of \mathcal{C} . We can then work directly with the gauge-transformed $\check{\Omega}_a$ and $\check{\omega}_a$. It is straightforward to check that $\check{\Omega}_a$ and $d\check{\omega}_a$ transform by the same factor under this gauge transformation, so o_3 is strictly invariant.

[1] M. Barkeshli, P. Bonderson, M. Cheng, and Z. Wang, Symmetry fractionalization, defects, and gauging of topological phases, *Phys. Rev. B* **100**, 115147 (2019).

[2] N. Tarantino, N. Lindner, and L. Fidkowski, Symmetry fractionalization and twist defects *New J. Phys.* **18**, 035006 (2016).

[3] V. Turaev, Homotopy field theory in dimension 3 and crossed group-categories, [arXiv:math/0005291](https://arxiv.org/abs/math/0005291)

[4] P. Etingof, D. Nikshych, and V. Ostrik, Fusion categories and homotopy theory, *Quantum Topol.* **1**, 209 (2010).

[5] X. Chen, F. J. Burnell, A. Vishwanath, and L. Fidkowski, Anomalous Symmetry Fractionalization and Surface Topological Order, *Phys. Rev. X* **5**, 041013 (2015).

[6] A. Kapustin and R. Thorngren, Anomalies of discrete symmetries in various dimensions and group cohomology, [arXiv:1404.3230](https://arxiv.org/abs/1404.3230).

[7] M. Barkeshli and M. Cheng, Relative anomalies in (2+1)d symmetry enriched topological states, *SciPost Phys.* **8**, 028 (2020).

[8] M. Cheng, M. Zaletel, M. Barkeshli, A. Vishwanath, and P. Bonderson, Translational Symmetry and Microscopic Constraints on Symmetry-Enriched Topological Phases: A View from the Surface, *Phys. Rev. X* **6**, 041068 (2016).

[9] D. S. Freed, Anomalies and invertible field theories, [arXiv:1404.7224](https://arxiv.org/abs/1404.7224).

[10] A. Vishwanath and T. Senthil, Physics of Three-Dimensional Bosonic Topological Insulators: Surface-Deconfined Criticality and Quantized Magnetoelectric Effect, *Phys. Rev. X* **3**, 011016 (2013).

[11] C. Wang, A. C. Potter, and T. Senthil, Gapped symmetry preserving surface state for the electron topological insulator, *Phys. Rev. B* **88**, 115137 (2013).

[12] M. A. Metlitski, C. L. Kane, and M. P. A. Fisher, Bosonic topological insulator in three dimensions and the statistical witten effect, *Phys. Rev. B* **88**, 035131 (2013).

[13] M. A. Metlitski, C. L. Kane, and M. P. A. Fisher, Symmetry-respecting topologically ordered surface phase of three-dimensional electron topological insulators, *Phys. Rev. B* **92**, 125111 (2015).

[14] X. Chen, L. Fidkowski, and A. Vishwanath, Symmetry enforced non-Abelian topological order at the surface of a topological insulator, *Phys. Rev. B* **89**, 165132 (2014).

[15] L. Fidkowski, X. Chen, and A. Vishwanath, Non-Abelian Topological Order on the Surface of a 3D Topological Superconductor from an Exactly Solved Model, *Phys. Rev. X* **3**, 041016 (2013).

[16] P. Bonderson, C. Nayak, and X.-L. Qi, A time-reversal invariant topological phase at the surface of a 3d topological insulator, *J. Stat. Mech.: Theory Exp.* (2013) P09016.

[17] C. Wang and T. Senthil, Interacting fermionic topological insulators/superconductors in three dimensions, *Phys. Rev. B* **89**, 195124 (2014).

[18] M. A. Metlitski, L. Fidkowski, X. Chen, and A. Vishwanath, Interaction effects on 3D topological superconductors: surface topological order from vortex condensation, the 16 fold way and fermionic Kramers doublets, [arXiv:1406.3032](https://arxiv.org/abs/1406.3032).

[19] G. Y. Cho, J. C. Y. Teo, and S. Ryu, Conflicting symmetries in topologically ordered surface states of three-dimensional bosonic symmetry protected topological phases, *Phys. Rev. B* **89**, 235103 (2014).

[20] N. Seiberg and E. Witten, Gapped boundary phases of topological insulators via weak coupling, [arXiv:1602.04251](https://arxiv.org/abs/1602.04251).

[21] M. Hermele and X. Chen, Flux-Fusion Anomaly Test and Bosonic Topological Crystalline Insulators, *Phys. Rev. X* **6**, 041006 (2016).

[22] Y. Qi and L. Fu, Anomalous Crystal Symmetry Fractionalization on the Surface of Topological Crystalline Insulators, *Phys. Rev. Lett.* **115**, 236801 (2015).

[23] H. Song, S.-J. Huang, L. Fu, and M. Hermele, Topological Phases Protected by Point Group Symmetry, *Phys. Rev. X* **7**, 011020 (2017).

[24] D. Bulmash and M. Barkeshli, Absolute anomalies in (2+1)d symmetry-enriched topological states and exact (3+1)d constructions, *Phys. Rev. Research* **2**, 043033 (2020).

[25] S. Tata, R. Kobayashi, D. Bulmash, and M. Barkeshli, Anomalies in (2 + 1)D fermionic topological phases and (3 + 1)D path integral state sums for fermionic SPTs, [arXiv:2104.14567](https://arxiv.org/abs/2104.14567).

[26] S. X. Cui, C. Galindo, J. Y. Plavnik, and Z. Wang, On gauging symmetry of modular categories, *Commun. Math. Phys.* **348**, 1043 (2016).

[27] A. Kapustin, R. Thorngren, A. Turzillo, and Z. Wang, Fermionic symmetry protected topological phases and cobordisms, *J. High Energy Phys.* **12** (2015) 1.

[28] A. Kapustin and R. Thorngren, Fermionic SPT phases in higher dimensions and bosonization, *J. High Energy Phys.* **10** (2017) 80.

[29] D. S. Freed and M. J. Hopkins, Reflection positivity and invertible topological phases, *Geom. Topol.* **25**, 1165 (2016).

[30] Q.-R. Wang and Z.-C. Gu, Construction and Classification of Symmetry-Protected Topological Phases in Interacting Fermion Systems, *Phys. Rev. X* **10**, 031055 (2020).

[31] T. Johnson-Freyd and D. Reutter, Minimal nondegenerate extensions, [arXiv:2105.15167](https://arxiv.org/abs/2105.15167).

[32] L. Fidkowski, A. Vishwanath, and M. A. Metlitski, Surface topological order and a new 't Hooft anomaly of interaction enabled 3+1d fermion spts, [arXiv:1804.08628](https://arxiv.org/abs/1804.08628).

[33] D. Delmastro, D. Gaiotto, and J. Gomis, Global anomalies on the Hilbert space, *J. High Energy Phys.* **11** (2021) 142.

[34] P. Bruillard, C. Galindo, T. Hagge, S.-H. Ng, J. Y. Plavnik, E. C. Rowell, and Z. Wang, Fermionic modular categories and the 16-fold way, *J. Math. Phys.* **58**, 041704 (2017).

[35] D. Bulmash and M. Barkeshli, Fermionic symmetry fractionalization in (2+1)D, *Phys. Rev. B* **105**, 125114 (2022).

[36] D. Aasen, P. Bonderson, and C. Knapp, Characterization and classification of fermionic symmetry enriched topological phases, [arXiv:2109.10911](https://arxiv.org/abs/2109.10911).

[37] P. Bruillard, C. Galindo, S.-H. Ng, J. Y. Plavnik, E. C. Rowell, and Z. Wang, Classification of super-modular categories by rank, [arXiv:1705.05293](https://arxiv.org/abs/1705.05293).

[38] P. Bonderson, E. Rowell, Z. Wang, and Q. Zhang, Congruence subgroups and super-modular categories, *Pacific J. Math.* **296**, 257 (2018).

[39] M. Barkeshli, Y.-A. Chen, P.-S. Hsin, and N. Manjunath, Classification of (2+1)d invertible fermionic topological phases with symmetry, [arXiv:2109.11039](https://arxiv.org/abs/2109.11039).

[40] X. Wen and X.-G. Wen, Distinguish modular categories and 2+1D topological orders beyond modular data: Mapping class group of higher genus manifold, [arXiv:1908.10381](https://arxiv.org/abs/1908.10381).

[41] P. Bonderson, C. Delaney, C. Galindo, E. C. Rowell, A. Tran, and Z. Wang, On invariants of modular categories beyond modular data, *J. Pure Appl. Algebra* **223**, 4065 (2019).

[42] A. Davydov, Bogomolov multiplier, double class-preserving automorphisms, and modular invariants for orbifolds, *J. Math. Phys.* **55**, 092305 (2014).

[43] M. Barkeshli, P. Bonderson, M. Cheng, C.-M. Jian, and K. Walker, Reflection and time reversal symmetry enriched topological phases of matter: Path integrals, non-orientable manifolds, and anomalies, *Commun. Math. Phys.* **374**, 1021 (2019).

[44] P. H. Bonderson, Non-Abelian anyons and interferometry, Ph.D. thesis, California Institute of Technology, 2007.

[45] We thank Meng Cheng for discussions regarding the following issue.