

Computational modeling and evolutionary implications of biochemical reactions in bacterial microcompartments

Clair A Huffine^{1,2,3,4}, Lucas C Wheeler⁵, Boswell Wing⁶ and Jeffrey C Cameron^{2,3,7}



Bacterial microcompartments (BMCs) are protein-encapsulated compartments found across at least 23 bacterial phyla. BMCs contain a variety of metabolic processes that share the commonality of toxic or volatile intermediates, oxygen-sensitive enzymes and cofactors, or increased substrate concentration for magnified reaction rates. These compartmentalized reactions have been computationally modeled to explore the encapsulated dynamics, ask evolutionary-based questions, and develop a more systematic understanding required for the engineering of novel BMCs. Many crucial aspects of these systems remain unknown or unmeasured, such as substrate permeabilities across the protein shell, feasibility of pH gradients, and transport rates of associated substrates into the cell. This review explores existing BMC models, dominated in the literature by cyanobacterial carboxysomes, and highlights potentially important areas for exploration.

Addresses

¹ BioFrontiers Institute, University of Colorado Boulder, 3415 Colorado Avenue, Boulder, CO 80309, USA

² Department of Biochemistry, University of Colorado, Boulder, CO 80309, USA

³ Renewable and Sustainable Energy Institute, University of Colorado, Boulder, CO 80309, USA

⁴ Interdisciplinary Quantitative Biology Program (IQ Biology), BioFrontiers Institute, University of Colorado, Boulder, CO 80309, USA

⁵ Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, CO 80309, USA

⁶ Department of Geological Sciences, Boulder, CO 80309, USA

⁷ National Renewable Energy Laboratory, Golden, CO 80401, USA

Corresponding author:

Cameron, Jeffrey C (Jeffrey.C.Cameron@colorado.edu)

Current Opinion in Microbiology 2022, **65**:15–23

This review comes from a themed issue on **Microbes and bioenergy**

Edited by **Tim Donohoe**

For complete overview of the section, please refer to the article collection, “[Special Section on Microbes and bioenergy](#)”

Available online 28th October 2021

<https://doi.org/10.1016/j.mib.2021.10.001>

1369-5274/© 2021 Elsevier Ltd. All rights reserved.

Introduction

Once thought of as simple ‘sacks of enzymes’, it is now clear that the internal organization of bacteria rivals that of eukaryotes. For example, photosynthetic cyanobacteria not only contain internal photosynthetic membranes responsible for producing the oxygen that has transformed Earth’s atmosphere enabling eukaryotic evolution and survival, they also contain carboxysomes. Carboxysomes are a class of protein-based organelles, termed bacterial microcompartments (BMCs), that resemble icosahedral viral capsids [1]. BMCs, defined by a common shell architecture, contain different suites of enzymes in their core and have been found across at least 23 diverse bacterial phyla [2]. They fall within two main categories, catabolic metabolosomes and anabolic carboxysomes (Figure 1). Both categories of BMCs create a chemical and physical microenvironment inside the bacterial cytoplasm. BMCs are optimized to perform chemical reactions which would otherwise be toxic to other cellular components, create volatile intermediates, or involve enzymes, coenzymes, or intermediates that are inactivated by the presence of oxygen (O₂). BMCs are of interest for their potential applications in modulating the flux and selectivity of biochemical reactions, containment of toxic intermediates from industrial reactions, antigen display for vaccine development, and creation of nanoparticles in chemotherapy targeting [3,4]. Recent reviews have delved into a wide variety of topics such as BMC positioning [5], BMC repurposing for industrial uses [6], evolutionary relationship of shell proteins across different BMCs [7], and BMC protein stoichiometry [8,9].

Despite this intense focus on synthesizing what is currently known about BMCs, many important details remain unknown about the function(s), assembly mechanisms, stability, and ultimate use in enhancing the biochemical reactions that they encapsulate. Because of their small size (40–600 nm) [14•,15] and subcellular heterogeneity (e.g. intact/functional, assembly intermediates, and degrading stages (Figure 2b)), the chemical environment inside BMCs remains challenging to experimentally measure with standard technologies. Recent work tracking the position and activity of individual BMCs for the first time through single-cell lineages addresses this heterogeneity and points towards a solution for understanding how the chemical environment of BMCs dictate activity and degradation [16••].

Figure 1





Diagram of the substrates, intermediates, and products of carboxysomes and the most studied catabolic BMCs [10–13]. The charge of each metabolite is indicated with color; positive in red, negative in blue, and neutral in grey. Nearly all the intermediates are nonpolar whereas substrates and products, which transverse the protein shell, are predominantly negatively charged or nonpolar. There is theorized inclusion of B_{12} cycling inside some catabolic BMCs as well as a common exclusion of oxygen (O_2) [11,13]. Abbreviations: BMC, bacterial microcompartments; RuBP, ribulose-1,5-bisphosphate; Pdu, 1,2-propanediol utilization; GRM, glycol radical enzyme-associated microcompartments; GRM5, fuculose-associated microcompartment; Eut, ethanamine utilization; Cut, choline-utilization; RMM, 1-amino-2-propanol utilization; 3-PGA, 3-phosphoglyceric acid.

While empirical knowledge of how BMCs function *in vivo* and *in vitro* is hard won, computational modeling can be used to develop hypotheses about the encapsulated pathways using existing knowledge of enzymatic reaction rates, metabolite concentrations, and transport phenomena. Models can be designed to focus on specific aspects of BMC function with a relatively simple framework containing the relevant components or expanded to include whole cell physiology. However, the uncertainty of these predictions increases along with the number of variables in the associated model, leading to potentially misleading conclusions [17–20]. Thus, one of the most challenging aspects of modeling is determining what should and should not be included and what assumptions are reasonable to make in simplifying the model. In this review, we synthesize existing BMC models, focusing on what assumptions were made in their construction, and emphasizing areas for further investigation. We focus on models of carboxysomes due to their long history and a recent spate of new advances. With these models we can explore the parameter space that BMCs may functionally occupy, test evolutionary pressures leading to formation, and help focus the direction of future experimental work. Ultimately, these quantitative understandings could enable the *de novo* design and engineering of novel BMCs for diverse functions [21,22].

Carboxysomes

Found in cyanobacteria [23] and chemoautotrophic bacteria [24], carboxysomes are the only known anabolic BMC and contain the enzymatic machinery to fix carbon dioxide (CO_2) from the atmosphere into organic compounds needed to generate biomass (Figure 2a). Carboxysomes are notable for their role in the cyanobacterial CO_2 -concentrating-mechanism (CCM) which allows cyanobacteria to contribute nearly 25% of fixed carbon globally [25]. The intact carboxysome shell is thought to be largely impermeable to CO_2 [26–28,29[•]] and O_2 [28,29[•]] but permeable to bicarbonate (HCO_3^-) (Figure 2). Therefore, an increased local concentration of carboxysomal CO_2 sourced from HCO_3^- can be created around ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO), the rate-limiting enzymatic step of CO_2 -fixation, alongside a simultaneous exclusion of O_2 , a competitive inhibitor. The RuBisCO oxygenation reaction produces 2-phosphoglycolate, a competitive inhibitor of triose-phosphate isomerase [30], and requires metabolism through an energetically expensive pathway called photorespiration to recover invested CO_2 and nitrogen while regenerating ribulose-1,5-bisphosphate (RuBP). While not aiding directly in sugar production, photorespiration may play an important role in preventing reactive oxygen species production, production of serine and cysteine, and assimilation of sulfur and nitrogen [31–

Figure 2



Schematic representations of the CO₂-concentrating-mechanism (CCM) and light reactions of oxygenic photosynthesis in beta-cyanobacteria. Included are the active carbon uptake complexes, NDH-I₃ and NDH-I₄, bicarbonate transporters, BCT1, BicA, and SbtA, and carboxysome enzymes, carbonic anhydrase (CA) and ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO). (a) Inorganic carbon, in the form of bicarbonate (HCO₃⁻), is actively transported via transporters (BCT1, BicA, SbtA) into the cell. Carbon dioxide (CO₂) enters the cell via passive diffusion and is then converted to HCO₃⁻ by thylakoid membrane-localized NDH-I₃ and NDH-I₄ to increase cytoplasmic HCO₃⁻ concentration. Unlike CO₂ and oxygen (O₂), HCO₃⁻ diffuses across the proteinaceous shell into the carboxysome, where it is brought into equilibrium with CO₂ by CA in a fully reversible reaction. CO₂ is then fixed by RuBisCO using ribulose-1,5-bisphosphate (RuBP) as an acceptor molecule, resulting in the production of substrates that are converted into carbohydrates through the Calvin-Benson Cycle (CB Cycle). The CB Cycle is powered with ATP and NADPH produced from the photosynthetic light reactions. RuBisCO may require inorganic phosphate (Pi) for activation. RuBisCO can also undergo a side reaction with O₂ leading to an energetically expensive carbon recovery pathway called photorespiration, but this reaction may mainly occur in broken carboxysomes (dotted shell line) due to the impermeability of O₂ to the carboxysome. (b) Three co-existing pools of RubisCO in the same cell during the carboxysome life cycle [16^{**}] can exhibit distinct metabolic activities because of encapsulation and CA activation/inactivation based on redox environment [34] that modulates substrate accessibility and concentration. Abbreviations: CA, carbonic anhydrase; RuBisCO, ribulose-1,5-bisphosphate carboxylase/oxygenase; RuBP, ribulose-1,5-bisphosphate; CB Cycle, Calvin-Benson Cycle.

33]. Recent experimental work identified three separate cellular pools of RubisCO in the same cell as a result of the dynamic assembly and degradation pathways that are hypothesized to exhibit distinct metabolic activities (Figure 2b) [16^{**}].

Carboxysome models

As the first identified and prototypical BMC, most computational modeling work has been conducted on the carboxysome with the initial models published in the 1980s [35–39]. Even the most recent models have

roots in these early versions, with more information being incorporated on the composition, structure, and activity of the carboxysome-encapsulated reactions as a result of scientific advancements over the last three decades.

Two different forms of BMCs with independent evolutionary origins are found within cyanobacterial species, alpha-carboxysomes and beta-carboxysomes. While structurally and functionally similar, including the presence of homologous shell proteins, the molecular scaffolds used in the assembly are distinct. Beta-carboxysomes form

inside-out beginning with the aggregation and polar localization of CA and RuBisCO before encapsulation via shell proteins [73]. The dynamic assembly pathway and hierarchy of protein-protein interactions during alpha-carboxysome formation *in vivo* remains to be fully elucidated. Although historically, models were based on CCMs in beta-cyanobacteria, Hopkinson *et al.* explored an alpha-carboxysome-based model founded on experimental data from *Prochlorococcus* sp. They concluded that the CCM mechanism of alpha-carboxysomes functions very similarly to the more studied beta-carboxysomes [40]. The knowledge gap in carboxysome shell permeability was overcome by fitting the model to their experimental data to estimate parameter values, but this methodology ultimately limited the applicability of this model to other organisms (Table 1). The impact of O₂ on carbon fixation was not addressed in this paper despite the relevance of O₂/CO₂ competition with RuBisCO.

The study published by Clark *et al.*, took a more inclusive modeling approach inspired by chemical engineering to systematically investigate how different beta-carboxysome counts (down to zero in a CCM-lacking mutant), size, and cellular position impacts CO₂-fixation rates [41]. They found that a consistent number of RuBisCO units within the cell was the most important parameter impacting CO₂ fixation (Table 1) [41]. An analysis using dimensionless Dahmköhler numbers was also used to identify controlling phenomena in the CCM based on transport and reaction rate considerations. Additionally, this study found that a carboxysome-lacking mutant begins to mimic a potential pre-carboxysome form of ancestral cyanobacteria, the physiology of which can be replicated with CCM knockout cell lines. Notwithstanding this insight, this paper does not explore the evolutionary implications of cells without a CCM. This model also assumes zero permeability of the carboxysome to O₂ and CO₂, although more recent evidence indicates that the transport properties of various substrates across the shell could change over the life cycle of the BMC as a result of assembly, breakage, and degradation (Figure 2b).

The most recent models have focused specifically on the pH dynamics of beta-cyanobacterial CCMs. Mangan *et al.*'s study explored how the CCM handled different species of inorganic carbon with a 'pH-aware' model and found a lower carboxysome pH relative to the cytosol was favorable for increased carbon fixation activity [42,43]. Boatman *et al.* applied this model to an experimental study of *Trichodesmium erythraeum*, a filamentous diazotrophic cyanobacteria, with notable increases in their cell and carboxysome size parameters to match *Trichodesmium* (Table 1) [44]. They determined the importance of HCO₃⁻ and pH on CO₂ fixation rate and that elevated pH impacted CO₂ fixation through processes outside of the CCM [44]. Most recently, Long *et al.* modeled the inclusion of 'proton transport' in and out of the

carboxysome via RuBP and PGA, the substrate and product of RuBisCO's carboxylase activity respectively [45^{••}]. This paper placed the carboxysome in a m³ cytosolic solution and excluded the HCO₃⁻ transport aspect of the CCM (Table 1). This study included the generation of protons from CO₂-fixation and concluded that this creates a locally lowered pH in the carboxysome which, in turn, drives CA's equilibrium reaction towards CO₂ production [45^{••}].

These most recent models used equivalent permeabilities of the carboxysome shell to HCO₃⁻ and CO₂ (Table 1). Given the potential favorability of the pores to negatively charged ions by three orders of magnitude, equal transport of these molecules seems unlikely [29^{••}]. For example, if CO₂ is allowed to pass freely across the shell, then it would be released from the cell and collapse the functionality of the CCM similar to what was seen in classic experiments demonstrating a high-CO₂-requiring mutant by expressing CA outside the carboxysome [47]. Additionally, similar questions remain on the biological feasibility of maintaining a proton gradient across the shell within the bacterial cytoplasm.

Catabolic BMCs

Of the catabolic BMCs, also termed 'metabolosomes' [74], the most well studied are 1,2-propanediol utilization (Pdu) and the ethanolamine utilization (Eut), which are found in *Salmonella* and *Escherichia coli*. Less studied metabolosomes include 1-amino-2-propanol utilization (RMM), glycyl radical enzyme-associated microcompartments (GRM), choline utilization (GRM2), and fucose and rhamnose utilization (GRM5 and PVM) BMCs [8,75]. Research remains sparse on the genes, pathways, and enzymatic reactions involved in catabolic BMCs. These pathways may also be more complex than ones in carboxysomes, being dependent on redox balancing, disproportionation, and the presence of several metals and other cofactors. One commonality in metabolosomes is the presence of oxygen sensitive metal co-factor containing enzymes, such as B₁₂ cycling enzymes and glycyl-radical enzymes, with the BMC possibly acting as an O₂ exclusion mechanism [11,13].

Catabolic BMC models

There is an overall dearth of mathematical models for the catabolic BMCs with only a pair published models [22,76] of a non-carboxysome BMC systems in recent years. Considering the special interest from industry in BMCs that could shield the cytoplasm from toxic intermediates, increase metabolic flux, or modulate reaction selectivity and product inhibition, it is valuable to explore catabolic BMC capabilities systematically [77].

A notable difference between the dynamics of the carboxysome and other BMCs, such as the Pdu BMC, is that the internal reactions of the carboxysome begins with a

Table 1

Comparison of parameters used in the CCM Models. Dashes indicate values not reported, NA indicates values not applicable to the set-up of the model, and double slashes indicate two different modeled values

Parameters	Hopkinson et al. [40]	Clark et al. [41]	Mangan and Brenner [42]	Mangan et al. [43]	Boatman et al. [44]	Long et al. [45••]
Environmental species concentrations						
HCO_3^- (μM)	0–2000	1955	14 [46]	100	100	10 – 25 000 ^a [47–49]
CO_2 (μM)	Varies	45	0.14 [46]	15	15	0.01 – 25 ^a
O_2 (μM)	–	400 [50,51,52]	260	300	300	250 – 360 ^a [53,54]
Permeability and transport rates						
Cell membrane permeability to CO_2 (cm/s)	0.65 ^a [55]	no barrier [56]	0.3 [56,57]	0.3 [56]	0.3 [56]	NA
Cell membrane permeability to HCO_3^- (cm/s)	complete barrier	complete barrier	3.00E-04 [56,57]	2.00E-07 [56]	2.00E-07 [56]	NA
Carboxysome permeability to CO_2 (cm/s)	2.37E-05 ^{a,c}	complete barrier [46]	1.00E-03	1.00E-04 [39]	3.00E-05	1.00E-04 ^a
Carboxysome permeability to HCO_3^- (cm/s)	0.71 ^a [55]	no barrier [56]	1.00E-05	1.00E-04 [39]	3.00E-05	1.00E-04 ^a
HCO_3^- active transport rate (molecules/s)	1.51E + 04 ^a	1.11E + 06 ^a [58]	1.59E + 08 ^{a,c}	1.14E + 09 ^{a,c}	1.63E + 04 ^{a,c}	NA
Compartment sizes and pH						
Carboxysome radius (cm)	8.19E-06 ^a	1.00E-05 [59]	5.00E-06	5.00E-06	1.50E-05	1E-04//5E-06 ^a
Cell/system radius (cm)	3.50E-05 ^a	1.00E-04	5.00E-05	5.00E-05	3.00E-04	62 ^a
Number of Carboxysomes	6 [62]	1–4 [50,51,52]	1	1	1	1
Carboxysome pH	7.35	8	NA	7	7	<8
Cytoplasm pH	7.35	8	NA	8	8	8
Extracellular pH	8	8.2	NA	7	8.3	8
Enzyme parameters						
RuBisCO active sites	3975 ^{a,c}	3351 [50,51,52]	2160 [63–65]	2000 [65,66]	54,000	2.52E + 07//3011 ^a [48,67]
RuBisCO k_{cat} (s ⁻¹)	10.6 [68]	11.4 ^a [50,51,52]	26 [63–65]	11.6 [68–70]	1.92	14.4 [48,67]
RuBisCO $K_{1/2}$ (μM)	263	185 [50,51,52]	270 [63–65]	340 [68–70]	145 [71]	172 [48,67]
RuBisCO specificity (S _{c/o})	–	–	–	43 [68–70]	45 [71]	40 [48,67]
CA active sites	NA	NA	80 [63–65]	100 [65]	900	NA
CA hydration k_{cat} (s ⁻¹)	1000 ^b	1.14E + 05 ^b [71,72]	8.00E + 04 [63–65]	8.00E + 04 [65]	NA	0.05 ^b
CA hydration $K_{1/2}$ (μM)	NA	NA	3.20E + 03 [63–65]	3.20E + 03 [65]	104.7 ^b	NA
CA dehydration k_{cat} (s ⁻¹)	NA	NA	4.60E + 04 [63–65]	4.60E + 04 [65]	NA	100 ^b
CA dehydration $K_{1/2}$ (μM)	NA	NA	9.30E + 03 [63–65]	9.20E + 03 [65]	NA	NA

^a Values converted to the presented unit for comparison across models.

^b Values based on establishing equilibrium between CO_2 and HCO_3^- in the carboxysome.

^c Values fitted within the model to match experimental values.

reversible reaction while the propanediol pathway starts with an irreversible reaction potentially trapping the substrate within the shell once it has been produced. Jakobson *et al.*'s study (2017) on Pdu found that selective permeability of the BMC shell can increase the overall BMC function and highlighted the important role BMCs play in metabolite flux enhancement [76]. This model was adapted by Jakobson *et al.* to be generalizable to different cellular organizational strategies and metabolic pathways in order to guide industrial design of BMCs [22]. They did not include O_2 or the B_{12} cycling pathway, which is sensitive to O_2 , in their models. Given the

diversity of shell proteins and mysteries regarding their selectivity and dynamics, it is possible that additional mechanisms may allow escape or entrance of specific metabolites, co-factors, electrons, or proteins across the shell.

Conclusions

With only a handful of BMC models proposed in the last decade [22,40–44,45••,76], there remains a great many unanswered questions and research potential within this domain.

Many critical parameters of BMCs formation, function and regulation remain unknown. These parameters include the internal pH compared to the cytosol and the composition and stoichiometry of encapsulated proteins and co-factors. In particular, the permeabilities of substrates and products across the shell into the protein core remains largely unmeasured and speculative. The pores of the major structural shell proteins studied to date with crystallographic methods display overall positive charge, which allows for negatively charged compounds to readily pass through (Figure 1) [29^{••}]. However, the molecular mechanism and level of exclusion of positively charged ions, like H⁺, or nonpolar substrates, like CO₂ and O₂, has yet to be determined experimentally [29^{••}]. There is debate on the internal pH, redox environment, and the creation and relevance of a pH gradient across the BMC shell [29^{••},45^{••},78]. Additionally, systematic errors and biases in our understanding of these mechanisms plague all BMC models, with many of the featured parameters across all these models being based on previous measurements that may need to be revisited (Table 1). Therefore, it is important to combine theory with experimental data in an iterative approach to provide robust agreement between these methods.

Several mathematical models of cyanobacterial physiology not focused on BMCs take a full-cell approach, including not only carbon fixation but also photosynthetic reactions, protein production, and gene regulation [79,80]. However, these models lack the fine details of the CCM previously described in BMC models. A robust exploration, meeting in the middle of these two approaches, could yield useful insights on how the CCM is dynamically controlled in response to changes in cellular physiological conditions. In particular, several transporters relevant to the CCM are sensitive to inorganic carbon concentrations and regulated allosterically [81[•]], potentially influencing how the CCM reacts to changing environmental conditions.

Evolutionary considerations were touched upon briefly in a few of the carboxysome modeling studies. These models could be combined with recent computational approaches in metabolic evolution [82]. This approach could allow systematic investigation on how evolutionary intermediates in CCM evolution may have responded to environmental changes during proposed times of carboxysome origination such as the Great Oxidation Event (GOE) approximately 2.3–2.5 billion years ago [83[•]]. A dramatic increase in atmospheric O₂ concentrations across the GOE relative to slowly declining CO₂ concentrations led to an ~100 million fold increase in O₂:CO₂ ratios, which could have provided a selective pressure driving the encapsulation of oxygen-sensitive enzymes, such as RuBisCO. Recent explorations of similar questions featured a limited range of oxygen concentrations, and, as a result, may be missing important dynamics of

carboxysome evolution driven by the need for oxygen exclusion [45^{••}]. Furthermore, there are undeniable structural and functional similarities between catabolic BMCs and carboxysomes, which offers an underexplored opportunity for comparative evolutionary studies. One such example is the combination of homologous shell proteins between carboxysomes and other BMCs and the frequency of highly oxygen-sensitive coenzyme B₁₂ regeneration pathways and glycyl radical enzymes being sequestered in BMCs for use in the contained reactions [11,13]. This suggests that the O₂ exclusion property of carboxysomes may be a more general property of BMCs than previously considered.

Through computational modeling of BMCs, we can relate results to experimental data, probe theoretical evolutionary pressures, and gain deeper understanding of these systems [21,77]. Given the rapidly advancing power of computation and AI in addition to new experimental techniques, we anticipate many new breakthroughs in the coming decades harnessing the capabilities of BMCs that could have major benefits for society and the environment.

Conflict of interest statement

J.C.C. has equity and is co-founder and Chief Science Advisor for Prometheus Materials Inc. All other authors declare no competing interests.

Funding

This work was supported in part by the Interdisciplinary Quantitative Biology PhD Program (IQ Biology) and the NRT Integrated Data Science Program (NSF grant number 2022138) at the BioFrontiers Institute, University of Colorado-Boulder and by the National Science Foundation under Grant No. 2054085.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Krupovic M, Koonin EV: **Cellular origin of the viral capsid-like bacterial microcompartments.** *Biol Direct* 2017, **12**:25 <http://dx.doi.org/10.1186/s13062-017-0197-y>.
2. Axen SD, Erbilgin O, Kerfeld CA: **A taxonomy of bacterial microcompartment loci constructed by a novel scoring method.** *PLOS Comput Biol* 2014, **10**:e1003898 <http://dx.doi.org/10.1371/journal.pcbi.1003898>.
3. Zhang X, Konarev PV, Petoukhov MV, Svergun DI, Xing L, Cheng RH et al.: **Multiple assembly states of lumazine synthase: a model relating catalytic function and molecular assembly.** *J Mol Biol* 2006, **362**:753-770 <http://dx.doi.org/10.1016/j.jmb.2006.07.037>.
4. Dalmau M, Lim S, Chen HC, Ruiz C, Wang S-W: **Thermostability and molecular encapsulation within an engineered caged protein scaffold.** *Biotechnol Bioeng* 2008, **101**:654-664 <http://dx.doi.org/10.1002/bit.21988>.

5. MacCready JS, Vecchiarelli AG: **Positioning the model bacterial organelle, the carboxysome.** *mBio* 2021, **12** <http://dx.doi.org/10.1128/mBio.02519-19>.
6. Kennedy NW, Mills CE, Nichols TM, Abrahamson CH, Tullman-Ercek D: **Bacterial microcompartments: tiny organelles with big potential.** *Curr Opin Microbiol* 2021, **63**:36-42 <http://dx.doi.org/10.1016/j.mib.2021.05.010>.
7. Melnicki MR, Sutter M, Kerfeld CA: **Evolutionary relationships among shell proteins of carboxysomes and metabolosomes.** *Curr Opin Microbiol* 2021, **63**:1-9 <http://dx.doi.org/10.1016/j.mib.2021.05.011>.
8. Liu L-N: **Bacterial metabolosomes: new insights into their structure and bioengineering.** *Microb Biotechnol* 2021, **14**:88-93 <http://dx.doi.org/10.1111/1751-7915.13740>.
9. Liu L-N, Yang M, Sun Y, Yang J: **Protein stoichiometry, structural plasticity and regulation of bacterial microcompartments.** *Curr Opin Microbiol* 2021, **63**:133-141 <http://dx.doi.org/10.1016/j.mib.2021.07.006>.
10. Stewart KL, Stewart AM, Bobik TA: **Prokaryotic organelles: bacterial microcompartments in *E. coli* and *Salmonella*.** *EcoSal Plus* 2020, **9** <http://dx.doi.org/10.1128/ecosalplus.ESP-0025-2019>.
11. Ferlez B, Sutter M, Kerfeld CA: **Glycyl radical enzyme-associated microcompartments: redox-replete bacterial organelles.** *mBio* 2019, **10**:e02327-18 <http://dx.doi.org/10.1128/mBio.02327-18>.
12. Mallette E, Kimber MS: **Structural and kinetic characterization of (S)-1-amino-2-propanol kinase from the aminoacetone utilization microcompartment of *Mycobacterium smegmatis*.** *J Biol Chem* 2018, **293**:19909-19918 <http://dx.doi.org/10.1074/jbc.RA118.005485>.
13. Zarzycki J, Erbilgin O, Kerfeld CA: **Bioinformatic characterization of Glycyl radical enzyme-associated bacterial microcompartments.** *Appl Environ Microbiol* 2015, **81**:8315-8329 <http://dx.doi.org/10.1128/AEM.02587-15>.
14. Kennedy NW, Hershewe JM, Nichols TM, Roth EW, Wilke CD, Mills CE et al.: **Apparent size and morphology of bacterial microcompartments varies with technique.** *PLoS One* 2020, **15**:e0226395 <http://dx.doi.org/10.1371/journal.pone.0226395>
This paper examined different preparation techniques and provides recommendations for accurate measurements of BMC size distributions, which is an important consideration when investigating composition and function.
15. Liberton M, Austin JR II, Berg RH, Pakrasi HB: **Unique thylakoid membrane architecture of a unicellular N₂-fixing cyanobacterium revealed by electron tomography.** *Plant Physiol* 2011, **155**:1656-1666 <http://dx.doi.org/10.1104/pp.110.165332>.
16. Hill NC, Tay JW, Altus S, Bortz DM, Cameron JC: **Life cycle of a cyanobacterial carboxysome.** *Sci Adv* 2020, **6**:eaba1269 <http://dx.doi.org/10.1126/sciadv.aba1269>
Utilizes single-cell tracking to observe the lifecycle of a carboxysome. This method provides a way to bypass innate BMC heterogeneity in a cell and access activity levels of a single BMC over its lifespan as well as mechanisms for signaling degradation.
17. Brown KS, Sethna JP: **Statistical mechanical approaches to models with many poorly known parameters.** *Phys Rev E* 2003, **68**:021904 <http://dx.doi.org/10.1103/PhysRevE.68.021904>.
18. Chis O-T, Banga JR, Balsa-Canto E: **Structural identifiability of systems biology models: a critical comparison of methods.** *PLoS One* 2011, **6**:e27755 <http://dx.doi.org/10.1371/journal.pone.0027755>.
19. Transtrum MK, Machta BB, Brown KS, Daniels BC, Myers CR, Sethna JP: **Perspective: sloppiness and emergent theories in physics, biology, and beyond.** *J Chem Phys* 2015, **143**:010901 <http://dx.doi.org/10.1063/1.4923066>.
20. Roosa K, Chowell G: **Assessing parameter identifiability in compartmental dynamic models using a computational approach: application to infectious disease transmission models.** *Theor Biol Med Model* 2019, **16**:1 <http://dx.doi.org/10.1186/s12976-018-0097-6>.
21. Metcalf KJ, Slininger Lee MF, Jakobson CM, Tullman-Ercek D: **An estimate is worth about a thousand experiments: using order-of-magnitude estimates to identify cellular engineering targets.** *Microb Cell Fact* 2018, **17**:135 <http://dx.doi.org/10.1186/s12934-018-0979-7>.
22. Jakobson CM, Tullman-Ercek D, Mangan NM: **Spatially organizing biochemistry: choosing a strategy to translate synthetic biology to the factory.** *Sci Rep* 2018, **8**:8196 <http://dx.doi.org/10.1038/s41598-018-26399-0>.
23. Gantt E, Conti SF: **Ultrastructure of blue-green algae.** *J Bacteriol* 1969, **97**:1486-1493 <http://dx.doi.org/10.1128/jb.97.3.1486-1493.1969>.
24. Shively JM, Ball FL, Kline BW: **Electron microscopy of the carboxysomes (Polyhedral bodies) of *Thiobacillus neapolitanus*.** *J Bacteriol* 1973, **116**:1405-1411.
25. Behrenfeld MJ, Randerson JT, McClain CR, Feldman GC, Los SO, Tucker CJ et al.: **Biospheric primary production during an ENSO transition.** *Science* 2001, **291**:2594-2597 <http://dx.doi.org/10.1126/science.1055071>.
26. Dou Z, Heinhorst S, Williams EB, Murin CD, Shively JM, Cannon GC: **CO₂ fixation kinetics of halothiobacillus neapolitanus mutant carboxysomes lacking carbonic anhydrase suggest the shell acts as a diffusional barrier for CO₂.** *J Biol Chem* 2008, **283**:10377-10384 <http://dx.doi.org/10.1074/jbc.M709285200>.
27. Cai F, Menon BB, Cannon GC, Curry KJ, Shively JM, Heinhorst S: **The pentameric vertex proteins are necessary for the icosahedral carboxysome shell to function as a CO₂ leakage barrier.** *PLoS One* 2009, **4**:e7521 <http://dx.doi.org/10.1371/journal.pone.00007521>.
28. Kinney JN, Axen SD, Kerfeld CA: **Comparative analysis of carboxysome shell proteins.** *Photosynth Res* 2011, **109**:21-32 <http://dx.doi.org/10.1007/s11120-011-9624-6>.
29. Faulkner M, Szabó I, Weetman SL, Sicard F, Huber RG, Bond PJ et al.: **Molecular simulations unravel the molecular principles that mediate selective permeability of carboxysome shell protein.** *Sci Rep* 2020, **10**:17501 <http://dx.doi.org/10.1038/s41598-020-74536-5>
This paper modeled the free energy of metabolites crossing through the CcmK2 shell protein pore. Notably, O₂ and CO₂ had the same free energy and HCO₃⁻ and 3-PGA was favored by 3 orders of magnitude. This study makes an important step towards addressing the selective permeability of BMC shell proteins.
30. González-Mondragón E, Zubillaga RA, Hernández-Arana A: **Effect of a specific inhibitor on the unfolding and refolding kinetics of dimeric triosephosphate isomerase: establishing the dimeric and similarly structured nature of the main transition states on the forward and backward reactions.** *Biophys Chem* 2007, **125**:172-178 <http://dx.doi.org/10.1016/j.bpc.2006.07.007>.
31. Shi X, Bloom A: **Photorespiration: the futile cycle?** *Plants* 2021, **10**:908 <http://dx.doi.org/10.3390/plants10050908>.
32. Bauwe H, Hagemann M, Kern R, Timm S: **Photorespiration has a dual origin and manifold links to central metabolism.** *Curr Opin Plant Biol* 2012, **15**:269-275 <http://dx.doi.org/10.1016/j.pbi.2012.01.008>.
33. Broddrick JT, Rubin BE, Welkie DG, Du N, Mih N, Diamond S et al.: **Unique attributes of cyanobacterial metabolism revealed by improved genome-scale metabolic modeling and essential gene analysis.** *Proc Natl Acad Sci U S A* 2016, **113**:E8344-53 <http://dx.doi.org/10.1073/pnas.1613446113>.
34. Peña KL, Castel SE, de Araujo C, Espie GS, Kimber MS: **Structural basis of the oxidative activation of the carboxysomal γ-carbonic anhydrase, CcmM.** *Proc Natl Acad Sci U S A* 2010, **107**:2455-2460 <http://dx.doi.org/10.1073/pnas.0910866107>.
35. Badger MR, Bassett M, Comins HN: **A model for HCO₃⁻ accumulation and photosynthesis in the cyanobacterium *Synechococcus* sp.** *Plant Physiol* 1985, **77**:465-471.
36. Reinhold L, Zivman M, Kaplan A: **Inorganic carbon fluxes and photosynthesis in cyanobacteria — a quantitative model.** In *Progress in Photosynthesis Research*. Edited by Biggins J.

Dordrecht: Springer; 1987 http://dx.doi.org/10.1007/978-94-017-0519-6_62.

37. Reinhold L, Zviman M, Kaplan A: **A quantitative model for inorganic carbon fluxes and photosynthesis in cyanobacteria.** *Plant Physiol Biochem* 1989, **27**:945-954.

38. Reinhold L, Kosloff R, Kaplan A: **A model for inorganic carbon fluxes and photosynthesis in cyanobacterial carboxysomes.** *Can J Physiol* 1991, **69**:984-988 <http://dx.doi.org/10.1139/b91-126>.

39. Fridlyand L, Kaplan A, Reinhold L: **Quantitative evaluation of the role of a putative CO₂-scavenging entity in the cyanobacterial CO₂-concentrating mechanism.** *Biosystems* 1996, **37**:229-238 [http://dx.doi.org/10.1016/0303-2647\(95\)01561-2](http://dx.doi.org/10.1016/0303-2647(95)01561-2).

40. Hopkinson BM, Young JN, Tansik AL, Binder BJ: **The minimal CO₂-concentrating mechanism of *Prochlorococcus* spp. MED4 is effective and efficient.** *Plant Physiol* 2014, **166**:2205-2217 <http://dx.doi.org/10.1104/pp.14.247049>.

41. Clark RL, Cameron JC, Root TW, Pfleger BF: **Insights into the industrial growth of cyanobacteria from a model of the carbon-concentrating mechanism.** *AIChE J* 2014, **60**:1269-1277 <http://dx.doi.org/10.1002/aic.14310>.

42. Mangan NM, Brenner MP: **Systems analysis of the CO₂ concentrating mechanism in cyanobacteria.** *eLife* 2014, **3**:e02043 <http://dx.doi.org/10.7554/eLife.02043>.

43. Mangan NM, Flamholz A, Hood RD, Milo R, Savage DF: **pH determines the energetic efficiency of the cyanobacterial CO₂ concentrating mechanism.** *Proc Natl Acad Sci U S A* 2016, **113**:E5354 <http://dx.doi.org/10.1073/pnas.1525145113>.

44. Boatman TG, Mangan NM, Lawson T, Geider RJ: **Inorganic carbon and pH dependency of photosynthetic rates in *Trichodesmium*.** *J Exp Bot* 2018, **69**:3651-3660 <http://dx.doi.org/10.1093/jxb/ery141>.

45. Long BM, Förster B, Pulsford SB, Price GD, Badger MR: **Rubisco proton production can drive the elevation of CO₂ within condensates and carboxysomes.** *Proc Natl Acad Sci U S A* 2021, **118** <http://dx.doi.org/10.1073/pnas.2014406118>
Introduces tracking protons produced by RuBisCO carboxylation and brought in to the carboxysome by RuBP³⁻ and PGA²⁻ to help drive the conversion of HCO₃⁻ to CO₂ and lower pH within the carboxysome. The authors focused on how lowered pH may have driven evolution of the carboxysome from a condensate.

46. Price GD, Badger MR, Woodger FJ, Long BM: **Advances in understanding the cyanobacterial CO₂-concentrating-mechanism (CCM): functional components, Ci transporters, diversity, genetic regulation and prospects for engineering into plants.** *J Exp Bot* 2008, **59**:1441-1461 <http://dx.doi.org/10.1093/jxb/erm112>.

47. Price GD, Badger MR: **Expression of human carbonic anhydrase in the cyanobacterium *Synechococcus* PCC7942 creates a high CO₂-requiring phenotype.** *Plant Physiol* 1989, **91**:505-513.

48. Whitehead L, Long BM, Price GD, Badger MR: **Comparing the in vivo function of α -carboxysomes and β -carboxysomes in two model cyanobacteria.** *Plant Physiol* 2014, **165**:398-411 <http://dx.doi.org/10.1104/pp.114.237941>.

49. Badger MR, Kaplan A, Berry JA: **Internal inorganic carbon pool of *Chlamydomonas reinhardtii*: evidence for a carbon dioxide-concentrating mechanism.** *Plant Physiol* 1980, **66**:407-413 <http://dx.doi.org/10.1104/pp.66.3.407>.

50. Price GD, Süttemeyer D, Klughammer B, Ludwig M, Badger MR: **The functioning of the CO₂ concentrating mechanism in several cyanobacterial strains: a review of general physiological characteristics, genes, proteins, and recent advances.** *Can J Bot* 1998, **76**:973-1002 <http://dx.doi.org/10.1139/b98-081>.

51. Long BM, Badger MR, Whitney SM, Price GD: **Analysis of carboxysomes from *Synechococcus* PCC7942 reveals multiple Rubisco complexes with carboxysomal proteins CcmM and CcaA.** *J Biol Chem* 2007, **282**:29323-29335 <http://dx.doi.org/10.1074/jbc.M703896200>.

52. Savage DF, Afonso B, Chen AH, Silver PA: **Spatially ordered dynamics of the bacterial carbon fixation machinery.** *Science* 2010, **327**:1258-1261 <http://dx.doi.org/10.1126/science.1186090>.

53. Flamholz A, Shih PM: **Cell biology of photosynthesis over geologic time.** *Curr Biol* 2020, **30**:R490-4 <http://dx.doi.org/10.1016/j.cub.2020.01.076>.

54. Badger MR, Price GD: **CO₂ concentrating mechanisms in cyanobacteria: molecular components, their diversity and evolution.** *J Exp Bot* 2003, **54**:609-622 <http://dx.doi.org/10.1093/jxb/erg076>.

55. Pasciak W, Gavis J: **Transport Limitation of Nutrient Uptake in Phytoplankton1.** 1974 <http://dx.doi.org/10.4319/lo.1974.19.6.0881>.

56. Gutknecht J, Bisson MA, Tosteson FC: **Diffusion of carbon dioxide through lipid bilayer membranes: effects of carbonic anhydrase, bicarbonate, and unstirred layers.** *J Gen Physiol* 1977, **69**:779-794 <http://dx.doi.org/10.1085/jgp.69.6.779>.

57. Missner A, Kügler P, Saparov SM, Sommer K, Mathai JC, Zeidel ML et al.: **Carbon dioxide transport through membranes.** *J Biol Chem* 2008, **283**:25340-25347 <http://dx.doi.org/10.1074/jbc.M800096200>.

58. Price GD, Woodger FJ, Badger MR, Howitt SM, Tucker L: **Identification of a SulP-type bicarbonate transporter in marine cyanobacteria.** *Proc Natl Acad Sci U S A* 2004, **101**:18228-18233 <http://dx.doi.org/10.1073/pnas.0405211101>.

59. Rae BD, Long BM, Badger MR, Price GD: **Structural determinants of the outer shell of β -carboxysomes in *Synechococcus elongatus* PCC 7942: roles for CcmK2, K3-K4, CcmO, and CcmL.** *PLoS One* 2012, **7**:e43871 <http://dx.doi.org/10.1371/journal.pone.0043871>.

60. Moronta-Barrios F, Espinosa J, Contreras A: **Negative control of cell size in the cyanobacterium *Synechococcus elongatus* PCC 7942 by the essential response regulator RpaB.** *FEBS Lett* 2013, **587**:504-509 <http://dx.doi.org/10.1016/j.febslet.2013.01.023>.

61. Xu Y, Guerra LT, Li Z, Ludwig M, Dismukes G, Bryant D: **Altered carbohydrate metabolism in glycogen synthase mutants of *Synechococcus* sp. strain PCC 7002: cell factories for soluble sugars.** *Metab Eng* 2012, **16** <http://dx.doi.org/10.1016/j.ymben.2012.12.002>.

62. Ting CS, Hsieh C, Sundararaman S, Mannella C, Marko M: **Cryo-electron tomography reveals the comparative three-dimensional architecture of *Prochlorococcus*, a globally important marine cyanobacterium.** *J Bacteriol* 2007, **189**:4485-4493 <http://dx.doi.org/10.1128/JB.01948-06>.

63. Woodger FJ, Badger MR, Price GD: **Sensing of inorganic carbon limitation in *Synechococcus* PCC7942 is correlated with the size of the internal inorganic carbon pool and involves oxygen.** *Plant Physiol* 2005, **139**:1959-1969 <http://dx.doi.org/10.1104/pp.105.069146>.

64. Süttemeyer D, Price GD, Yu J-W, Badger MR: **Characterisation of carbon dioxide and bicarbonate transport during steady-state photosynthesis in the marine cyanobacterium *Synechococcus* strain PCC7002.** *Planta* 1995, **197**:597-607 <http://dx.doi.org/10.1007/BF00191566>.

65. Heinhorst S, Williams EB, Cai F, Murin CD, Shively JM, Cannon GC: **Characterization of the carboxysomal carbonic anhydrase CsoSCA from *Halothiobacillus neapolitanus*.** *J Bacteriol* 2006, **188**:8087-8094 <http://dx.doi.org/10.1128/JB.00990-06>.

66. Iancu CV, Ding HJ, Morris DM, Dias DP, Gonzales AD, Martino A et al.: **The structure of isolated *Synechococcus* strain WH8102 carboxysomes as revealed by electron cryotomography.** *J Mol Biol* 2007, **372**:764-773 <http://dx.doi.org/10.1016/j.jmb.2007.06.059>.

67. Occhialini A, Lin MT, Andralojc PJ, Hanson MR, Parry MAJ: **Transgenic tobacco plants with improved cyanobacterial Rubisco expression but no extra assembly factors grow at near wild-type rates if provided with elevated CO₂.** *Plant J Cell Mol Biol* 2016, **85**:148-160 <http://dx.doi.org/10.1111/tpj.13098>.

68. Tcherkez GGB, Farquhar GD, Andrews TJ: **Despite slow catalysis and confused substrate specificity, all ribulose bisphosphate carboxylases may be nearly perfectly optimized.** *Proc Natl Acad Sci U S A* 2006, **103**:7246-7251 <http://dx.doi.org/10.1073/pnas.0600605103>.

69. Savir Y, Noor E, Milo R, Tlusty T: **Cross-species analysis traces adaptation of Rubisco toward optimality in a low-dimensional landscape.** *Proc Natl Acad Sci U S A* 2010, **107**:3475-3480 <http://dx.doi.org/10.1073/pnas.0911663107>.

70. Morell MK, Paul K, O'Shea NJ, Kane HJ, Andrews TJ: **Mutations of an active site threonyl residue promote beta elimination and other side reactions of the enediol intermediate of the ribulosebisphosphate carboxylase reaction.** *J Biol Chem* 1994, **269**:8091-8098.

71. Badger M, Andrews T, Whitney S, Ludwig M, Yellowlees D, Leggat W et al.: **The diversity and coevolution of Rubisco, plastids, pyrenoids, and chloroplast-based CO₂-concentrating mechanisms in algae.** *Can J Bot* 1998, **76**:1052-1071 <http://dx.doi.org/10.1139/cjb-76-6-1052>.

72. Shingles R, Moroney JV: **Measurement of carbonic anhydrase activity using a sensitive fluorometric assay.** *Anal Biochem* 1997, **252**:190-197 <http://dx.doi.org/10.1006/abio.1997.2305>.

73. Cameron JC, Wilson SC, Bernstein SL, Kerfeld CA: **Biogenesis of a bacterial organelle: the carboxysome assembly pathway.** *Cell* 2013, **155**:1131-1140 <http://dx.doi.org/10.1016/j.cell.2013.10.044>.

74. Brinsmade SR, Paldon T, Escalante-Semerena JC: **Minimal functions and physiological conditions required for growth of *Salmonella enterica* on ethanamine in the absence of the metabolosome.** *J Bacteriol* 2005, **187**:8039-8046 <http://dx.doi.org/10.1128/JB.187.23.8039-8046.2005>.

75. Stewart AM, Stewart KL, Yeates TO, Bobik TA: **Advances in the world of bacterial microcompartments.** *Trends Biochem Sci* 2021, **46**:406-416 <http://dx.doi.org/10.1016/j.tibs.2020.12.002>.

76. Jakobson CM, Tullman-Ercek D, Slininger MF, Mangan NM: **A systems-level model reveals that 1,2-Propanediol utilization microcompartments enhance pathway flux through intermediate sequestration.** *PLoS Comput Biol* 2017, **13**:e1005525 <http://dx.doi.org/10.1371/journal.pcbi.1005525>.

77. Hinzpeter F, Gerland U, Tostevin F: **Optimal compartmentalization strategies for metabolic microcompartments.** *Biophys J* 2017, **112**:767-779 <http://dx.doi.org/10.1016/j.bpj.2016.11.3194>.

78. Menon BB, Heinhorst S, Shively JM, Cannon GC: **The carboxysome shell is permeable to protons.** *J Bacteriol* 2010, **192**:5881-5886 <http://dx.doi.org/10.1128/JB.00903-10>.

79. Faizi M, Zavrel T, Loureiro C, Červený J, Steuer R: **A model of optimal protein allocation during phototrophic growth.** *Biosystems* 2018, **166**:26-36 <http://dx.doi.org/10.1016/j.biosystems.2018.02.004>.

80. Westermark S, Steuer R: **Toward multiscale models of cyanobacterial growth: a modular approach.** *Front Bioeng Biotechnol* 2016, **4** <http://dx.doi.org/10.3389/fbioe.2016.00095>.

81. Fang S, Huang X, Zhang X, Zhang M, Hao Y, Guo H et al.: **• Molecular mechanism underlying transport and allosteric inhibition of bicarbonate transporter SbtA.** *Proc Natl Acad Sci U S A* 2021, **118** <http://dx.doi.org/10.1073/pnas.2101632118>

Structural studies of the cyanobacterial inorganic carbon (Ci) transporter SbtA complexed with the allosteric regulatory protein SbtB in the presence of HCO₃⁻ and adenyl nucleotides provides mechanistic insight into the signals that regulate the CCM in cyanobacteria. This study also raises additional questions about the role of cAMP in regulating carbon flux under varying Ci availabilities.

82. Wheeler LC, Smith SD: **Computational modeling of anthocyanin pathway evolution: biases, hotspots, and trade-offs.** *Integr Comp Biol* 2019, **59**:585-598 <http://dx.doi.org/10.1093/icb/icz049>.

83. Hurley SJ, Wing BA, Jasper CE, Hill NC, Cameron JC: **Carbon isotope evidence for the global physiology of Proterozoic cyanobacteria.** *Sci Adv* 2021, **7**:eabc8998 <http://dx.doi.org/10.1126/sciadv.abc8998>

Measurements of carbon isotope fractionation in wild-type cyanobacteria and ancestral analogs lacking a CCM indicate that carboxysomes evolved in response to rising levels of molecular oxygen, rather than decreasing CO₂.