

Universality in the onset of quantum chaos in many-body systems

Tyler LeBlond,¹ Dries Sels,^{2,3} Anatoli Polkovnikov⁴, and Marcos Rigol¹

¹*Department of Physics, The Pennsylvania State University, University Park, Pennsylvania 16802, USA*

²*Center for Computational Quantum Physics, Flatiron Institute, New York, New York 10010, USA*

³*Department of Physics, New York University, New York, New York 10003, USA*

⁴*Department of Physics, Boston University, Boston, Massachusetts 02215, USA*



(Received 30 December 2020; accepted 16 November 2021; published 30 November 2021)

We show that the onset of quantum chaos at infinite temperature in two many-body one-dimensional lattice models, the perturbed spin-1/2 XXZ and Anderson models, is characterized by universal behavior. Specifically, we show that the onset of quantum chaos is marked by maxima of the typical fidelity susceptibilities that scale with the square of the inverse average level spacing, saturating their upper bound, and that the strength of the integrability- or localization-breaking perturbation at these maxima decreases with increasing system size. We also show that the spectral function below the “Thouless” energy (in the quantum-chaotic regime) diverges when approaching those maxima. Our results suggest that, in the thermodynamic limit, arbitrarily small integrability- or localization-breaking perturbations result in quantum chaos in the many-body quantum systems studied here.

DOI: [10.1103/PhysRevB.104.L201117](https://doi.org/10.1103/PhysRevB.104.L201117)

Quantum chaos and eigenstate thermalization are two intertwined fields that have been the focus of much recent attention in the context of the emergence of statistical mechanics and thermodynamics in isolated quantum systems [1–3]. Those two fields are built on foundational analytical and computational results [4–14], and they have been recently linked to typicality ideas that date back to von Neumann’s work [15–17]. When quantum-chaotic systems (which are expected to exhibit eigenstate thermalization) are taken far from equilibrium, few-body operators (observables) generically equilibrate under unitary dynamics to the predictions of traditional statistical mechanics (they “thermalize”). This has been verified in experiments with ultracold quantum gases [18–21]. The “nonthermalizing” counterparts to quantum-chaotic systems are integrable [22–26] and disorder-localized [26–29] systems, which have also been probed in experiments with ultracold quantum gases [21,30–35].

In the clean case, a deeper understanding of what happens when quantum-chaotic systems approach integrable points is still needed. In finite systems there is a crossover in which quantum chaos [36–44] and eigenstate thermalization [39,42,43,45,46] indicators worsen. In the thermodynamic limit one expects quantum chaos and eigenstate thermalization to break down only at the integrable point [36–44] but the time scale for thermalization to diverge approaching that point [1,47–52]. The latter has been seen in recent experiments [21] and can be understood in the context of Fermi’s golden rule [52,53] and of the scaling of the quantum metric tensor with system size [44]. In the disorder-localized case, localization was argued to be perturbatively stable against weak short-range interactions [54,55] and against strong interactions in one dimension (1D) [56]. Disorder-induced localization in interacting systems is known as many-body localization and has attracted much theoretical and experimental

research in the strongly interacting regime [26–29]. Recent works have argued against and in favor of the occurrence of many-body localization in that regime in the thermodynamic limit [57–62].

We explore the onset of quantum chaos at infinite temperature in perturbed integrable and noninteracting disorder-localized chains, as well as its destruction upon approaching trivial classical limits. One of our goals is to identify universal features and differences between the clean and disordered cases. We compute fidelity susceptibilities χ [44,63], which are equivalent to the diagonal components of the quantum geometric tensor [64,65] or the norm of the adiabatic gauge potential [44], and spectral functions. Fidelity susceptibilities are commonly used to detect quantum phase transitions [64–68]. We find that the departure from quantum chaos is characterized by a higher sensitivity of eigenstates to perturbations [44,61,69], which results in maxima of the typical fidelity susceptibility that scale with the square of the inverse level spacing. The shifts in the maxima’s positions with system size are consistent with, at infinite temperature in the thermodynamic limit, quantum chaos only failing to occur at the unperturbed integrable, noninteracting disorder-localized, and integrable infinite-interaction (classical) limits.

We study the (clean) extended spin-1/2 XXZ chain:

$$\hat{H}_{\text{cln}} = \sum_{i=1}^L \left[\frac{J}{2} (\hat{S}_i^+ \hat{S}_{i+1}^- + \text{H.c.}) + \Delta \hat{S}_i^z \hat{S}_{i+1}^z + \Delta' \hat{S}_i^z \hat{S}_{i+2}^z \right], \quad (1)$$

with $J = \sqrt{2}$, $\Delta = (\sqrt{5} + 1)/4$, and $\Delta' \in [10^{-4}, 10^1]$. $\hat{H}_{\text{cln}}$ is Bethe-ansatz integrable for $\Delta' = 0$, and $\hat{H}_{\text{cln}}/\Delta'$ corresponds to two disconnected Ising chains for $\Delta' = \infty$. We also study the Anderson chain with added nearest-neighbor interactions,

which we write in the spin language as

$$\hat{H}_{\text{dsr}} = \sum_{i=1}^L \left[\frac{J}{2} (\hat{S}_i^+ \hat{S}_{i+1}^- + \text{H.c.}) + h_i \hat{S}_i^z + \Delta \hat{S}_i^z \hat{S}_{i+1}^z \right], \quad (2)$$

with $J = \sqrt{2}$, $h_i \in [-h, h]$ for $h = (\sqrt{5} + 1)/4$, and $\Delta \in [10^{-3}, 10^1]$ [70]. $\hat{H}_{\text{dsr}}$ is the Anderson model for $\Delta = 0$, and $\hat{H}_{\text{dsr}}/\Delta$ is the Ising chain for $\Delta = \infty$.

To probe the eigenkets $\{|m\rangle\}$ of the models above, we compute the typical fidelity susceptibility $\chi_{\text{typ}}(O) = \exp(\overline{\ln[\chi_m(O)]})$ (in short, the susceptibility) associated with observable $\hat{O}$, where

$$\chi_m(O) = L \sum_{l \neq m} \frac{|\langle m | \hat{O} | l \rangle|^2}{(E_m - E_l)^2}. \quad (3)$$

The average $\overline{\ln[\chi_m(O)]}$ is carried out over the central 50% of eigenstates in the spectrum. We also compute the average spectral function $|f_o(\omega)|^2 = \overline{|f_m^o(\omega)|^2}$ over the same 50% of eigenstates, where

$$|f_m^o(\omega)|^2 = L \sum_{l \neq m} |\langle m | \hat{O} | l \rangle|^2 \delta(\omega - \omega_{ml}). \quad (4)$$

We replace $\delta(x) \rightarrow \mu/[2\pi(x^2 + \mu^2)]$ with $\mu = 0.9 \omega_{\min}$, where $\omega_{\min}$ is the minimum level spacing. The factor of L in Eqs. (3) and (4) accounts for the Hilbert-Schmidt norm of our translationally invariant intensive observables.

The specific observables $\hat{O}$ considered [71] are the nearest-neighbor “kinetic” $\hat{K}_{\text{nn}}$ and interaction $\hat{U}_{\text{n}}$ energies:

$$\hat{K}_{\text{n}} = \frac{1}{L} \sum_{i=1}^L (\hat{S}_i^+ \hat{S}_{i+1}^- + \text{H.c.}), \quad \hat{U}_{\text{n}} = \frac{1}{L} \sum_{i=1}^L \hat{S}_i^z \hat{S}_{i+1}^z, \quad (5)$$

and the next-nearest-neighbor kinetic energy $\hat{K}_{\text{nn}}$. As shown recently [44,72,73], in integrable systems the response of eigenstates to perturbations depends on whether the perturbations do or do not break integrability. If $\hat{U}_{\text{n}}$ ($\hat{K}_{\text{nn}}$) is added to $\hat{H}_{\text{cln}}$, integrability is preserved (destroyed), while if $\hat{K}_{\text{n}}$ ($\hat{U}_{\text{n}}$) is added to $\hat{H}_{\text{dsr}}$, localization is preserved (destroyed); keeping this in mind, in what follows we show results for $\hat{U}_{\text{n}}$ and $\hat{K}_{\text{nn}}$ ($\hat{K}_{\text{n}}$ and $\hat{U}_{\text{n}}$) when studying $\hat{H}_{\text{cln}}$ ($\hat{H}_{\text{dsr}}$).

In Fig. 1 we show χ_{typ} vs Δ' (strength of the integrability-breaking next-nearest-neighbor interaction), for $\hat{U}_{\text{n}}$ [Fig. 1(a)] and $\hat{K}_{\text{nn}}$ [Fig. 1(b)]. The susceptibilities are scaled as expected for quantum-chaotic systems, for which $\chi_{\text{typ}} \propto LD^{-1}\omega_H^{-2}$ (ω_H is the mean level spacing and D is the Hilbert space dimension [74]) because $|\langle m | \hat{O} | l \rangle|^2 \propto D^{-1}$ for $E_m - E_l \rightarrow \omega_H$ [1,73]. For all chain sizes, the scaled susceptibilities exhibit an excellent collapse for about a decade in Δ' when $\Delta' \sim 1$. The region over which the scaled susceptibilities collapse increases (both towards smaller and towards larger values of Δ') with increasing system size. This highlights a quantum-chaotic regime that is robust and is increasing its extent with increasing system size.

The quantum-chaotic regime in Fig. 1 is separated from the integrable ones at small and large Δ' by maxima in χ_{typ} [71]. As a result of the trivial nature of the $\Delta' = \infty$ model, the large- Δ' maxima are more affected by finite-size effects than the small- Δ' ones. In what follows we focus on the latter. The




FIG. 1. Typical fidelity susceptibility χ_{typ} (scaled to exhibit collapse in the quantum-chaotic regime) vs the integrability-breaking parameter Δ' for observables $\hat{U}_{\text{n}}$ (a) and $\hat{K}_{\text{nn}}$ (b) in clean periodic chains. To calculate χ_{typ} and ω_H , we average over the central 50% of the eigenstates in the even- Z_2 sector in each total quasimomentum sector considered. For $L < 24$, we report the weighted average over all $k \neq (0, \pi)$ sectors, while for $L = 24$ we report results for the $k = \pi/2$ sector. Circles on the y axis show χ_{typ} at the integrable point ($\Delta' = 0$), and diamonds show the maximal χ_{typ}^* (at $\Delta'^* = -b/2a$) obtained from polynomial fits $ax^2 + bx + c$ (black solid lines about the maxima). The dotted lines on the right of the first peaks are a guide for the eye and depict $\Delta'^{-2.55}$ behavior. Inset in (a): χ_{typ}^* vs ω_H for both observables, along with the results of power-law fits. Inset in (b): Δ'^* vs ω_H for both observables (the values of Δ'^* overlap). The dotted line depicts $\omega_H^{0.39}$ behavior.

inset in Fig. 1(a) shows that χ_{typ} at the small Δ' maxima scales as the square of the inverse average level spacing ω_H . This scaling corresponds to the maximum possible sensitivity of quantum eigenstates to a perturbation [44]. It is exponentially larger, in system size, than expected from random matrix theory. The position of the maxima, Δ'^* , appears to move towards $\Delta' = 0$ exponentially fast with increasing system size (notice the near equal shift with increasing L and the log scale in the Δ' axis). In the inset in Fig. 1(b), we plot of Δ'^* vs ω_H showing that our numerical results are consistent with $\Delta'^* \propto (\omega_H)^\alpha$, with $\alpha \sim 0.39$. We note that our results in Fig. 1 are robust; Δ'^* and the scaling of χ_{typ}^* are nearly identical for both observables [71].

The susceptibility is related to the spectral function defining the dynamical response of the system [44,65]. Indeed, it follows from Eqs. (3) and (4) that

$$\chi_m(O) = \int_{-\infty}^{\infty} \frac{|f_m^o(\omega)|^2}{\omega^2} d\omega. \quad (6)$$




FIG. 2. Spectral functions in clean periodic chains with $L = 24$ for observables $\hat{U}_n$ [(a) and (b)] and $\hat{K}_m$ [(c) and (d)] over 2 decades of the integrability-breaking parameter Δ' [see labels at the top and legends in (b) and (d)]. In (a) and (c), the top insets show $F_O = (\omega/\Delta')^2 |f_O(\omega)|^2$ vs ω/Δ' at $\Delta' = 1.58 \times 10^{-2}$, while the bottom insets show $|f_O(\omega)|^2$ vs ω/Δ' at $\Delta' = 1.58 \times 10^{-1}$, for the three largest chains studied. The insets in (b) and (d) show $|f_O^p(\omega)|^2$ vs Δ' , where $|f_O^p(\omega)|^2$ is the value of $|f_O(\omega)|^2$ at the plateaus in the main panels (and for other values of Δ' for which $|f_O(\omega)|^2$ is not shown). The dotted lines are a guide for the eye and depict $\Delta'^{-2.55}$ behavior. All computations were done as for Fig. 1.

In integrable systems, $|f_O(\omega \rightarrow 0)|^2$ vanishes for integrability-preserving perturbations [44,71–73], leading to a polynomial in L scaling of $\chi_m(O)$ [44]. Typical (integrability breaking) perturbations in contrast have $|f_O(\omega \rightarrow 0)|^2 = O(1)$ [44,71–73] resulting in an exponential-in- L , $\sim D$, scaling of the susceptibility $\chi_m(O)$ [44]. As mentioned before, in quantum-chaotic systems $\chi_m(O) \propto L/[D(\omega_H)^2] \sim D$. The faster scaling at the maxima $\chi_{\text{typ}}^* \propto 1/\omega_H^2 \sim D^2$ implies that the spectral function diverges as $|f_O(\omega_H)|^2 \sim 1/\omega_H$ around Δ^* .

Figures 2(a) and 2(c) show $|f_O(\omega)|^2$ vs ω/Δ' for different values of Δ' about Δ^* for $L = 24$. The data for both observables collapse at frequencies $\omega/\Delta' \lesssim 1$ showing that $|f_O(\omega)|^2 \sim (\Delta'/\omega)^2$ in that regime [75]. In the top insets, we plot $F_O = (\omega/\Delta')^2 |f_O(\omega)|^2$ for different chain sizes when $\Delta' < \Delta^*$. The plateaus show that the $|f_O(\omega)|^2 \sim (\Delta'/\omega)^2$ behavior is robust to changing L [71]. For $\Delta' < \Delta^*$, the susceptibilities in Figs. 2(a) and 2(c) also collapse at lower frequencies showing a nontrivial dependence of ω/Δ' [71], but this collapse gradually disappears as Δ' approaches Δ^* .

When Δ' increases beyond Δ^* and the system enters into the quantum-chaotic regime [Figs. 2(b) and 2(d)], a plateau develops in the spectral function at low frequencies [76]. The formation and growth of the plateau with increasing L , at a fixed $\Delta' \gtrsim \Delta^*$, are illustrated in the bottom insets in Figs. 2(a) and 2(c). The plateau and the $|f_O(\omega)|^2 \sim (\Delta'/\omega)^2$

behavior coexist in the regime in which $\Delta' \gtrsim \Delta^*$, which is consistent with the occurrence of thermalization with relaxation rates dictated by Fermi's golden rule [51,71]. In that regime, we find that the spectral function $|f_O(\omega)|^2$ at the plateau, $|f_O^p(\omega)|^2$, appears to diverge as $(\Delta')^{-\beta}$ with $\beta \sim 2.55$ [see insets in Figs. 2(b) and 2(d)], consistent with the divergence of χ_{typ} in Fig. 1 (see dotted lines in the main panels). Remarkably, it is possible to relate the scaling of $|f_O^p(\omega)|^2$ with Δ' with the drift of Δ^* with L : $\Delta^* \sim \omega_H^\alpha$ with $\alpha = 1/\beta \sim 0.39$ [see inset in Fig. 1(b)].

We can understand this under the following scenario: Let $|f_O(\omega)|^2 = |f_O^p(\Delta')|^2$ for $\omega < \omega_p(\Delta')$ and $|f_O(\omega)|^2 \propto (\Delta'/\omega)^\kappa$ for $\omega > \omega_p(\Delta')$, with $\omega_p(\Delta')$ playing the role of the so-called Thouless energy, and $\kappa > 1$. Then from the spectral sum rule, $\int |f_O(\omega)|^2 d\omega = O(1)$, we infer that $\omega_p(\Delta') \propto (\Delta')^\beta$, with $\beta = \kappa/(\kappa - 1)$, and that $|f_O^p(\Delta')|^2 \propto (\Delta')^{-\beta}$. The maximum of χ_{typ} then occurs when $\omega_p = \omega_H$, i.e., when the maximum of the spectral function occurs at the Heisenberg scale. This results in $\Delta^* \sim \omega_H^\alpha$ with $\alpha = 1/\beta$, and $\chi_{\text{typ}}^* \sim \omega_H^{-2}$. Currently, we do not know the origin of the values of the exponents suggested by our numerical calculations. Given our observation of $|f_O(\omega)|^2 \sim (\Delta'/\omega)^2$ behavior for Δ' below and above Δ^* , which appears to grow in extent with increasing system size [see top insets in Figs. 2(a) and 2(c)], two scenarios come to mind: (i) The exponents observed numerically are affected by finite-size effects, and for larger systems than those accessible to us, $\kappa = 2$, $\beta = 2$, and $\alpha = 1/2$; and (ii) the spectral function develops a power law with an exponent $1 < \kappa < 2$ before saturating to a constant at low frequencies so that $\beta > 2$ and $\alpha < 1/2$.

In Fig. 3, we show results for the spectral function of disordered chains in the presence of nearest-neighbor interactions. The corresponding typical fidelity susceptibilities are shown in Fig. 4. The results in Figs. 3 and 4 are similar to those in Figs. 2 and 1, respectively. The similarity is remarkable considering that the unperturbed models in both cases are strikingly different, the disordered one being a noninteracting localized model and the clean one being an interacting integrable one. The slight differences between the results in Figs. 3 and 2 include a narrower $|f_O(\omega)|^2 \sim (\Delta'/\omega)^2$ regime in Figs. 3(a) and 3(c) as compared with Figs. 2(a) and 2(c), and a narrower regime in which $|f_O^p(\omega)|^2$ is consistent with a power law scaling with Δ in Fig. 3(d). Related to the latter, in the inset in Fig. 4(b) the dynamical range for Δ^* vs ω_H is smaller than in the inset in Fig. 1(b). Consequently, and also keeping in mind that in Fig. 4 we plot typical fidelity susceptibilities while in Fig. 3 we plot raw averages of the spectral functions, we cannot establish a relationship between the scaling of $|f_O^p(\omega)|^2$ with Δ' and the drift of Δ^* with L as we did for the clean case. That said, all those differences are consistent with stronger finite-size effects, and fluctuations associated with the disorder average, in the disordered systems. For the latter, the largest chains studied have $L = 18$ vs the $L = 24$ chains considered for clean systems.

In summary, our results suggest that the onset of quantum chaos at infinite temperature in the models studied, as well as its destruction when approaching classical limits for very strong interactions, is characterized by universal behavior. We focused our analysis on the onset of quantum




FIG. 3. Spectral functions in disordered periodic chains with $L = 18$ for observables $\hat{K}_n$ [(a) and (b)] and $\hat{U}_n$ [(c) and (d)] over 2 decades of the interaction strength Δ [see labels at the top, and legends in (b) and (d)]. In (a) and (c), the top insets show $F_O = (\omega/\Delta')^2 |f_O(\omega)|^2$ vs ω/Δ' at $\Delta' = 1.58 \times 10^{-2}$, while the bottom insets show $|f_O(\omega)|^2$ vs ω/Δ' at $\Delta' = 1.58 \times 10^{-1}$, for the three largest chains studied. The insets in (b) and (d) show $|f_O^p(\omega)|^2$ vs Δ , where $|f_O^p(\omega)|^2$ is the value of $|f_O(\omega)|^2$ at the plateaus in the main panels (and for other values of Δ for which $|f_O(\omega)|^2$ is not shown). The dotted lines are a guide for the eye and depict $\Delta^{-2.4}$ behavior. To calculate $|f_O(\omega)|^2$, we average over the central 50% of the eigenstates in each chain and then over disorder realizations (200 for $L \leq 16$, 100 for $L = 17$, and 50 for $L = 18$).

chaos as finite-size effects (and fluctuations associated with disorder averages) are smaller. The main universal feature identified is the divergence of the typical fidelity susceptibilities as ω_H^{-2} when entering (exiting) the quantum-chaotic regime and the associated divergence of the spectral functions below the Thouless energy. The latter is potentially universal and diverges as $\epsilon^{-\beta}$ (ϵ being the strength of either the integrability-breaking or localization-breaking perturbation) in the quantum-chaotic regime. Also potentially universal is the shift of the position ϵ^* of the maximum of the fidelity susceptibilities as $\epsilon^* \sim \omega_H^\alpha$, as well as the relation $\alpha = 1/\beta$ between the exponents. We note that $\epsilon^* \sim \omega_H^\alpha$ supports the expectation that in clean systems in the thermodynamic limit, quantum chaos and eigenstate thermalization break down only at the integrable point [36–44], and it suggests that at infinite temperature the 1D Anderson insulator (for the parameters considered here) is unstable against adding interactions. An interesting open question is whether this relates to recent findings that many-body localization is unstable against the insertion of thermal “bubbles” if disorder is not strong enough [77,78].

Much still needs to be explored, such as what happens at finite temperatures and when one changes the parameters of the unperturbed Hamiltonians [which we selected to be $O(1)$




FIG. 4. Typical fidelity susceptibility χ_{typ} (scaled to exhibit collapse in the quantum-chaotic regime) vs the interaction strength Δ for observables $\hat{K}_n$ (a) and $\hat{U}_n$ (b) in disordered periodic chains. Circles on the y axis show χ_{typ} at the Anderson-localized point ($\Delta = 0$), and diamonds show the maximal χ_{typ} (at $\Delta^* = -b/2a$) obtained from polynomial fits $ax^2 + bx + c$ (black solid lines about the maxima). The inset in (a) shows χ_{typ} vs ω_H for both observables, along with the results of power-law fits. The error bars are the (propagated) standard deviation of the average over disorder realizations (see Ref. [71] for details) at the value of Δ (for which we carried out a calculation) that is closest to Δ^* . The inset in (b) shows Δ^* vs ω_H for both observables. The dotted line depicts $\omega_H^{0.28}$ behavior. All computations were done as for Fig. 3.

to minimize finite-size effects]. In the disordered case, two parameter regimes to be explored are the strong disorder and strong interaction regimes. The contrast between the small Δ and large Δ peaks in the fidelity susceptibilities in Fig. 4 suggests that obtaining meaningful scalings using full exact diagonalization in those regimes will be computationally very challenging. We note that the results reported in this Research Letter required about 1 000 000 CPU hours of calculations.

We acknowledge discussions with E. Altman, A. Dymarsky, S. Gopalakrishnan, D. Huse, M. Pandey, and L. Vidmar. This work was supported by the National Science Foundation under Grants No. PHY-2012145 (T.L. and M.R.), No. DMR-1813499 (A.P.), and No. DMR-2103658 (A.P.), and by the AFOSR under Grants No. FA9550-16-1-0334 (A.P.), No. FA9550-21-1-0342 (A.P.), and No. FA9550-21-1-0236 (D.S.). The computations were carried out in the Roar supercomputer in the Institute for Computational and Data Sciences (ICDS) at Penn State. The Flatiron Institute is a division of the Simons Foundation.

- [1] L. D'Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics, *Adv. Phys.* **65**, 239 (2016).
- [2] J. M. Deutsch, Eigenstate thermalization hypothesis, *Rep. Prog. Phys.* **81**, 082001 (2018).
- [3] T. Mori, T. N. Ikeda, E. Kaminishi, and M. Ueda, Thermalization and prethermalization in isolated quantum systems: a theoretical overview, *J. Phys. B: At., Mol. Opt. Phys.* **51**, 112001 (2018).
- [4] E. Wigner, Characteristic vectors of bordered matrices with infinite dimensions, *Ann. Math.* **62**, 548 (1955).
- [5] E. Wigner, Characteristic vectors of bordered matrices of infinite dimensions II, *Ann. Math.* **65**, 203 (1957).
- [6] E. Wigner, On the distribution of the roots of certain symmetric matrices, *Ann. Math.* **67**, 325 (1958).
- [7] F. J. Dyson, Statistical theory of the energy levels of complex systems. I, *J. Math. Phys. (Melville, NY)* **3**, 140 (1962).
- [8] F. J. Dyson, Statistical theory of the energy levels of complex systems. II, *J. Math. Phys. (Melville, NY)* **3**, 157 (1962).
- [9] F. J. Dyson, Statistical theory of the energy levels of complex systems. III, *J. Math. Phys. (Melville, NY)* **3**, 166 (1962).
- [10] O. Bohigas, M. J. Giannoni, and C. Schmit, Characterization of Chaotic Quantum Spectra and Universality of Level Fluctuation Laws, *Phys. Rev. Lett.* **52**, 1 (1984).
- [11] J. M. Deutsch, Quantum statistical mechanics in a closed system, *Phys. Rev. A* **43**, 2046 (1991).
- [12] M. Srednicki, Chaos and quantum thermalization, *Phys. Rev. E* **50**, 888 (1994).
- [13] M. Srednicki, The approach to thermal equilibrium in quantized chaotic systems, *J. Phys. A: Math. Gen.* **32**, 1163 (1999).
- [14] M. Rigol, V. Dunjko, and M. Olshanii, Thermalization and its mechanism for generic isolated quantum systems, *Nature (London)* **452**, 854 (2008).
- [15] J. von Neumann, Proof of the ergodic theorem and the H-theorem in quantum mechanics, *Z. Phys.* **57**, 30 (1929).
- [16] S. Goldstein, J. L. Lebowitz, R. Tumulka, and N. Zanghì, Long-time behavior of macroscopic quantum systems, *Eur. Phys. J. H* **35**, 173 (2010).
- [17] M. Rigol and M. Srednicki, Alternatives to Eigenstate Thermalization, *Phys. Rev. Lett.* **108**, 110601 (2012).
- [18] S. Trotzky, Y.-A. Chen, A. Flesch, I. P. McCulloch, U. Schollwöck, J. Eisert, and I. Bloch, Probing the relaxation towards equilibrium in an isolated strongly correlated 1D Bose gas, *Nat. Phys.* **8**, 325 (2012).
- [19] A. M. Kaufman, M. E. Tai, A. Lukin, M. Rispoli, R. Schittko, P. M. Preiss, and M. Greiner, Quantum thermalization through entanglement in an isolated many-body system, *Science* **353**, 794 (2016).
- [20] G. Clos, D. Porras, U. Warring, and T. Schaetz, Time-Resolved Observation of Thermalization in an Isolated Quantum System, *Phys. Rev. Lett.* **117**, 170401 (2016).
- [21] Y. Tang, W. Kao, K.-Y. Li, S. Seo, K. Mallayya, M. Rigol, S. Gopalakrishnan, and B. L. Lev, Thermalization near Integrability in a Dipolar Quantum Newton's Cradle, *Phys. Rev. X* **8**, 021030 (2018).
- [22] L. Vidmar and M. Rigol, Generalized Gibbs ensemble in integrable lattice models, *J. Stat. Mech.: Theory Exp.* (2016) 064007.
- [23] F. H. L. Essler and M. Fagotti, Quench dynamics and relaxation in isolated integrable quantum spin chains, *J. Stat. Mech.: Theory Exp.* (2016) 064002.
- [24] J.-S. Caux, The quench action, *J. Stat. Mech.: Theory Exp.* (2016) 064006.
- [25] P. Calabrese and J. Cardy, Quantum quenches in 1+1 dimensional conformal field theories, *J. Stat. Mech.: Theory Exp.* (2016) 064003.
- [26] R. Vasseur and J. E. Moore, Nonequilibrium quantum dynamics and transport: From integrability to many-body localization, *J. Stat. Mech.: Theory Exp.* (2016) 064010.
- [27] R. Nandkishore and D. A. Huse, Many-body localization and thermalization in quantum statistical mechanics, *Annu. Rev. Condens. Matter Phys.* **6**, 15 (2015).
- [28] E. Altman and R. Vosk, Universal dynamics and renormalization in many-body-localized systems, *Annu. Rev. Condens. Matter Phys.* **6**, 383 (2015).
- [29] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, Colloquium: Many-body localization, thermalization, and entanglement, *Rev. Mod. Phys.* **91**, 021001 (2019).
- [30] T. Kinoshita, T. Wenger, and D. S. Weiss, A quantum Newton's cradle, *Nature (London)* **440**, 900 (2006).
- [31] M. Gring, M. Kuhnert, T. Langen, T. Kitagawa, B. Rauer, M. Schreitl, I. Mazets, D. A. Smith, E. Demler, and J. Schmiedmayer, Relaxation and prethermalization in an isolated quantum system, *Science* **337**, 1318 (2012).
- [32] T. Langen, S. Erne, R. Geiger, B. Rauer, T. Schweigler, M. Kuhnert, W. Rohringer, I. E. Mazets, T. Gasenzer, and J. Schmiedmayer, Experimental observation of a generalized Gibbs ensemble, *Science* **348**, 207 (2015).
- [33] M. Schreiber, S. S. Hodgman, P. Bordia, H. P. Lüschen, M. H. Fischer, R. Vosk, E. Altman, U. Schneider, and I. Bloch, Observation of many-body localization of interacting fermions in a quasi-random optical lattice, *Science* **349**, 842 (2015).
- [34] J.-Y. Choi, S. Hild, J. Zeiher, P. Schauß, A. Rubio-Abadal, T. Yefsah, V. Khemani, D. A. Huse, I. Bloch, and C. Gross, Exploring the many-body localization transition in two dimensions, *Science* **352**, 1547 (2016).
- [35] N. Malvania, Y. Zhang, Y. Le, J. Dubail, M. Rigol, and D. S. Weiss, Generalized hydrodynamics in strongly interacting 1D Bose gases, *Science* **373**, 1129 (2021).
- [36] D. A. Rabson, B. N. Narozhny, and A. J. Millis, Crossover from Poisson to Wigner-Dyson level statistics in spin chains with integrability breaking, *Phys. Rev. B* **69**, 054403 (2004).
- [37] L. F. Santos and M. Rigol, Onset of quantum chaos in one-dimensional bosonic and fermionic systems and its relation to thermalization, *Phys. Rev. E* **81**, 036206 (2010).
- [38] M. Rigol and L. F. Santos, Quantum chaos and thermalization in gapped systems, *Phys. Rev. A* **82**, 011604(R) (2010).
- [39] L. F. Santos and M. Rigol, Localization and the effects of symmetries in the thermalization properties of one-dimensional quantum systems, *Phys. Rev. E* **82**, 031130 (2010).
- [40] R. Modak, S. Mukerjee, and S. Ramaswamy, Universal power law in crossover from integrability to quantum chaos, *Phys. Rev. B* **90**, 075152 (2014).
- [41] R. Modak and S. Mukerjee, Finite size scaling in crossover among different random matrix ensembles in microscopic lattice models, *New J. Phys.* **16**, 093016 (2014).

- [42] R. Mondaini, K. R. Fratus, M. Srednicki, and M. Rigol, Eigenstate thermalization in the two-dimensional transverse field Ising model, *Phys. Rev. E* **93**, 032104 (2016).
- [43] R. Mondaini and M. Rigol, Eigenstate thermalization in the two-dimensional transverse field Ising model. II. Off-diagonal matrix elements of observables, *Phys. Rev. E* **96**, 012157 (2017).
- [44] M. Pandey, P. W. Claeys, D. K. Campbell, A. Polkovnikov, and D. Sels, Adiabatic Eigenstate Deformations as a Sensitive Probe for Quantum Chaos, *Phys. Rev. X* **10**, 041017 (2020).
- [45] M. Rigol, Breakdown of Thermalization in Finite One-Dimensional Systems, *Phys. Rev. Lett.* **103**, 100403 (2009).
- [46] M. Rigol, Quantum quenches and thermalization in one-dimensional fermionic systems, *Phys. Rev. A* **80**, 053607 (2009).
- [47] M. Stark and M. Kollar, Kinetic description of thermalization dynamics in weakly interacting quantum systems, [arXiv:1308.1610](https://arxiv.org/abs/1308.1610).
- [48] F. H. L. Essler, S. Kehrein, S. R. Manmana, and N. J. Robinson, Quench dynamics in a model with tuneable integrability breaking, *Phys. Rev. B* **89**, 165104 (2014).
- [49] B. Bertini, F. H. L. Essler, S. Groha, and N. J. Robinson, Prethermalization and Thermalization in Models with Weak Integrability Breaking, *Phys. Rev. Lett.* **115**, 180601 (2015).
- [50] B. Bertini, F. H. L. Essler, S. Groha, and N. J. Robinson, Thermalization and light cones in a model with weak integrability breaking, *Phys. Rev. B* **94**, 245117 (2016).
- [51] K. Mallayya and M. Rigol, Quantum Quenches and Relaxation Dynamics in the Thermodynamic Limit, *Phys. Rev. Lett.* **120**, 070603 (2018).
- [52] K. Mallayya, M. Rigol, and W. De Roeck, Prethermalization and Thermalization in Isolated Quantum Systems, *Phys. Rev. X* **9**, 021027 (2019).
- [53] A. J. Friedman, S. Gopalakrishnan, and R. Vasseur, Diffusive hydrodynamics from integrability breaking, *Phys. Rev. B* **101**, 180302(R) (2020).
- [54] I. V. Gornyi, A. D. Mirlin, and D. G. Polyakov, Interacting Electrons in Disordered Wires: Anderson Localization and Low- T Transport, *Phys. Rev. Lett.* **95**, 206603 (2005).
- [55] D. Basko, I. Aleiner, and B. Altshuler, Metal-insulator transition in a weakly interacting many-electron system with localized single-particle states, *Ann. Phys. (NY)* **321**, 1126 (2006).
- [56] V. Oganessian and D. A. Huse, Localization of interacting fermions at high temperature, *Phys. Rev. B* **75**, 155111 (2007).
- [57] J. Šuntajs, J. Bonča, T. Prosen, and L. Vidmar, Quantum chaos challenges many-body localization, *Phys. Rev. E* **102**, 062144 (2020).
- [58] D. A. Abanin, J. H. Bardarson, G. D. Tomasi, S. Gopalakrishnan, V. Khemani, S. A. Parameswaran, F. Pollmann, A. C. Potter, M. Serbyn, and R. Vasseur, Distinguishing localization from chaos: Challenges in finite-size systems, *Ann. Phys. (NY)* **427**, 168415 (2021).
- [59] P. Sierant, D. Delande, and J. Zakrzewski, Thouless Time Analysis of Anderson and Many-Body Localization Transitions, *Phys. Rev. Lett.* **124**, 186601 (2020).
- [60] J. Šuntajs, J. Bonča, T. Prosen, and L. Vidmar, Ergodicity breaking transition in finite disordered spin chains, *Phys. Rev. B* **102**, 064207 (2020).
- [61] D. Sels and A. Polkovnikov, Dynamical obstruction to localization in a disordered spin chain, *Phys. Rev. E* **104**, 054105 (2021).
- [62] M. Kiefer-Emmanouilidis, R. Unanyan, M. Fleischhauer, and J. Sirker, Absence of true localization in many-body localized phases, *Phys. Rev. B* **103**, 024203 (2021).
- [63] P. Sierant, A. Maksymov, M. Kuś, and J. Zakrzewski, Fidelity susceptibility in Gaussian random ensembles, *Phys. Rev. E* **99**, 050102(R) (2019).
- [64] L. Campos Venuti and P. Zanardi, Quantum Critical Scaling of the Geometric Tensors, *Phys. Rev. Lett.* **99**, 095701 (2007).
- [65] M. Kolodrubetz, V. Gritsev, and A. Polkovnikov, Classifying and measuring geometry of a quantum ground state manifold, *Phys. Rev. B* **88**, 064304 (2013).
- [66] P. Zanardi and N. Paunković, Ground state overlap and quantum phase transitions, *Phys. Rev. E* **74**, 031123 (2006).
- [67] M. Rigol, B. S. Shastry, and S. Haas, Fidelity and superconductivity in two-dimensional $t-J$ models, *Phys. Rev. B* **80**, 094529 (2009).
- [68] M. Kolodrubetz, D. Sels, P. Mehta, and A. Polkovnikov, Geometry and non-adiabatic response in quantum and classical systems, *Phys. Rep.* **697**, 1 (2017).
- [69] T. Villazon, P. W. Claeys, M. Pandey, A. Polkovnikov, and A. Chandran, Persistent dark states in anisotropic central spin models, *Sci. Rep.* **10**, 16080 (2020).
- [70] For these parameters, the disorder width ($2h \approx 1.62$) is $\approx 57\%$ of the clean noninteracting electrons bandwidth ($2J \approx 2.83$). If one defines the localization length ξ via the single-particle density decay $n_l(x) \sim \exp(-|x - l|/\xi)$, then typical single-particle eigenstates have $\xi \approx 2.6$, while the ones with maximal ξ have $\xi_{\max} \approx 6$.
- [71] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevB.104.L201117> for additional numerical results, which include the spectral functions of the unperturbed models, and for additional parameters to the ones reported in the main text, typical fidelity susceptibilities for other observables, the ratio of consecutive energy levels, and the disorder-average standard deviations for the typical fidelity susceptibilities. We also provide a perturbation theory derivation of the $(\epsilon/\omega)^2$ behavior of the spectral function and discuss its relation to Fermi's golden rule. References [79–84] are cited therein.
- [72] M. Brenes, J. Gould, and M. Rigol, Low-frequency behavior of off-diagonal matrix elements in the integrable XXZ chain and in a locally perturbed quantum-chaotic XXZ chain, *Phys. Rev. B* **102**, 075127 (2020).
- [73] T. LeBlond and M. Rigol, Eigenstate thermalization for observables that break Hamiltonian symmetries and its counterpart in interacting integrable systems, *Phys. Rev. E* **102**, 062113 (2020).
- [74] For large systems, $\omega_H \sim \sqrt{L}D^{-1}$, because the effective width of the many-body energy spectrum is $\propto \sqrt{L}$ (D is the number of energy levels).
- [75] The difference in the extent of the $(\Delta'/\omega)^2$ regime is due to the difference in behavior of $|f_o(\omega)|^2$ for integrability-preserving ($\hat{U}_n$) vs integrability-breaking ($\hat{K}_{nn}$) operators at $\Delta' = 0$, which results in a spectral gap for the former when Δ' is very small.
- [76] For large L , a diffusive $1/\sqrt{\omega}$ regime is expected to develop at frequencies above those of the plateau [1].

- [77] W. De Roeck and F. Huveneers, Stability and instability towards delocalization in many-body localization systems, *Phys. Rev. B* **95**, 155129 (2017).
- [78] P. J. D. Crowley and A. Chandran, Avalanche induced coexisting localized and thermal regions in disordered chains, *Phys. Rev. Research* **2**, 033262 (2020).
- [79] Y. Y. Atas, E. Bogomolny, O. Giraud, and G. Roux, Distribution of the Ratio of Consecutive Level Spacings in Random Matrix Ensembles, *Phys. Rev. Lett.* **110**, 084101 (2013).
- [80] T. LeBlond, K. Mallayya, L. Vidmar, and M. Rigol, Entanglement and matrix elements of observables in interacting integrable systems, *Phys. Rev. E* **100**, 062134 (2019).
- [81] E. Khatami, G. Pupillo, M. Srednicki, and M. Rigol, Fluctuation-Dissipation Theorem in an Isolated System of Quantum Dipolar Bosons after a Quench, *Phys. Rev. Lett.* **111**, 050403 (2013).
- [82] K. Mallayya and M. Rigol, Heating Rates in Periodically Driven Strongly Interacting Quantum Many-Body Systems, *Phys. Rev. Lett.* **123**, 240603 (2019).
- [83] We assume that the microcanonical average of $\hat{O}$ is zero; otherwise it should be subtracted from O_m .
- [84] C. Schönle, D. Jansen, F. Heidrich-Meisner, and L. Vidmar, Eigenstate thermalization hypothesis through the lens of auto-correlation functions, *Phys. Rev. B* **103**, 235137 (2021).