


LAB PROTOCOL

Measuring dissolved black carbon in water via aqueous, inorganic, high-performance liquid chromatography of benzenepolycarboxylic acid (BPCA) molecular markers

Riley Barton , Sasha Wagner *

Department of Earth and Environmental Sciences, Rensselaer Polytechnic Institute, Troy, New York, United States of America

* wagnes3@rpi.edu



OPEN ACCESS

Citation: Barton R, Wagner S (2022) Measuring dissolved black carbon in water via aqueous, inorganic, high-performance liquid chromatography of benzenepolycarboxylic acid (BPCA) molecular markers. PLoS ONE 17(5): e0268059. <https://doi.org/10.1371/journal.pone.0268059>

Editor: Lee W. Cooper, University of Maryland Center for Environmental Science, UNITED STATES

Received: February 16, 2022

Accepted: April 20, 2022

Published: May 26, 2022

Copyright: © 2022 Barton, Wagner. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: This protocol does not report new results. The data shown in [Table 1](#) were previously reported by Wagner et al. (2019) in *Nature Communications* and are only shown here as exemplary DBC and BPCA concentrations for different river and oceanic water samples.

Funding: National Science Foundation grant #2017577 awarded to S.W. <https://beta.nsf.gov/funding/opportunities/chemical-oceanography>
Rensselaer Polytechnic Institute Graduate

Abstract

Dissolved black carbon (DBC) is the condensed aromatic portion of dissolved organic matter produced from the incomplete combustion of biomass and other thermogenic processes. DBC quantification facilitates the examination of the production, accumulation, cycling, transformation, and effects of biologically recalcitrant condensed aromatic carbon in aquatic environments. Due to the heterogeneous nature of DBC molecules, concentrations are difficult to measure directly. Here, the method for DBC quantification consists of oxidizing condensed aromatic carbon to benzenepolycarboxylic acids (BPCAs), which are used as proxies for the assessment of DBC in the original sample. The concentrations of oxidation products (BPCAs) are quantified using high-performance liquid chromatography. DBC concentrations are determined from the concentration of BPCAs using a previously established conversion factor. Details and full descriptions of the preparative and analytical procedures and techniques of the BPCA method are usually omitted for brevity in published method sections and method-specific papers. With this step-by-step protocol, we aim to clarify the steps of DBC analysis, especially for those adopting or conducting the BPCA method for the first time.

Introduction

The heating and partial combustion of organic matter (e.g., during wildfires and the burning of fossil fuels) fundamentally alters the molecular structure of unburned biomass [1]. Pyrogenic and other thermogenic residues are best described as a continuum of organic compounds that range from mildly altered, biolabile molecules to highly altered, biorefractory, condensed aromatic compounds that are typically formed at high charring temperatures [2, 3]. The condensed aromatic portion of partially combusted organic matter is termed “black carbon” (BC; [3]). Due to its condensed aromatic molecular structure, BC is highly resistant to biodegradation and is thus stabilized in soils for centuries to millennia [4, 5]. The ubiquity and long-term storage of BC in soils suggest that the production of BC during wildfires may serve

Research Fellowship awarded to R.B. <https://info.rpi.edu/graduate-education/funding-your-graduate-studies> The funders had and will not have a role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

to partially offset fire-derived carbon emissions [6]. BC is an important component of the global carbon cycle in terms of carbon sequestration and therefore draws research interest from a wide range of disciplines, such as biogeochemistry, climatology, oceanography, and soil science.

Research gaps that currently prevent sufficient representation of BC cycling in global models include poorly understood fluxes between major and intermediate carbon reservoirs and biogeochemical processes that control the fate of BC in the environment at local and regional scales. Some portion of the BC deposited to soils and sediments “leaks out” over time, entering rivers in the form of dissolved BC (DBC). An estimated 18–27 Tg of DBC is exported globally by rivers each year, which may reduce the presumed efficacy of long-term BC storage on land [7, 8]. The mobilization of DBC serves as a link between terrestrial and marine reservoirs and DBC concentrations are usually higher in rivers (1.5 to 140 $\mu\text{M-C}$) than in the ocean (0.08 to 2.5 $\mu\text{M-C}$; [3]). Leaching experiments indicate a high degree of variability in terms of the amount and composition of DBC leached from soils, charcoal, and pyrogenic aerosols [9–13]. Due to the heterogeneous nature of DBC compounds, which exist in an even more complex dissolved organic matter matrix, the direct quantification of these condensed aromatic molecules is analytically challenging. The benzenepolycarboxylic acid (BPCA) method is one of the most widely utilized methods for condensed aromatic DBC quantification in aquatic environments (see Wagner et al. [3, 14] for further discussion on this) and is the analytical protocol we describe here.

The BPCA method was first established for quantifying BC in soils [15] and has since been adapted for measuring DBC in water [16, 17]. While the BPCA method was originally developed for the study of pyrogenic condensed aromatic carbon, its use has since expanded to encompass non-pyrogenic condensed aromatic carbon classes such as those present in crude oil and asphaltene fractions [18, 19]. Thus, we define DBC as the condensed aromatic portion of bulk dissolved organic carbon (DOC) that passes through a filter (usually 0.1 to 0.7 μm pore size) and is detectable as BPCA molecular markers upon nitric acid oxidation (Fig 1). The oxidation of condensed aromatic DBC molecules yields a mixture of substituted BPCAs, including benzenetricarboxylic acids (B3CAs), benzenetetracarboxylic acids (B4CAs), benzenepentacarboxylic acid (B5CA), and benzenhexacarboxylic acid (B6CA) as well as nitrated BPCAs [16, 20]. It is important to note here that the BPCA method detects molecular oxidation products (not native DBC structures), which represent only a fraction of all condensed aromatic carbon present in the original sample. This fraction was originally estimated to be ~30% on a per-carbon basis for DBC; determined from the average amount of carbon retained from the oxidation of polycyclic aromatic hydrocarbon standards and the average molecular formula of marine condensed aromatic structures inferred from ultrahigh-resolution mass spectroscopy [16, 20]. Thus, we rely on conversion factors to estimate DBC concentrations from the concentrations of measured BPCAs. The conversion factors utilized by the aquatic black carbon community do vary depending on the chromatographic method employed, specific BPCAs quantified, and/or DBC standard material used [16, 20–23].

The oxidation of non-pyrogenic samples has been shown to produce the less-substituted BPCAs (B3CAs and B4CAs) potentially through carbonization reactions during oxidation of large amounts of organic carbon ($> 5\text{ mg-OC per } 2\text{ mL of nitric acid}$) [26]. Therefore, the protocol we describe here limits the amount of OC oxidized and utilizes an equation derived from the power relationship between established DBC concentrations and the sum of B5CA and B6CA only [21]. Current conversion factors to translate BPCA to DBC concentrations, including the one we employ here, are based upon the oxidation patterns of polyaromatic hydrocarbon standards [20] and ultrahigh-resolution mass spectral data [16]. Other conversion factors have been developed from the oxidation of graphene oxide [23]. However, none of these




Fig 1. Oxidation of DBC to BPCAs. Dissolved black carbon (DBC) makes up a portion of bulk dissolved organic carbon (DOC). The two examples of possible DBC molecular structures shown were proposed by Dittmar and Paeng [24] based on ultrahigh-resolution mass spectral analysis of ocean water by Dittmar and Koch [25]. When black carbon molecules are oxidized with nitric acid and heat, they produce benzenepolycarboxylic acids (BPCAs). Oxidation products from highly condensed aromatic structures would include highly substituted BPCAs; benzenehexacarboxylic acid (B6CA) and benzenepentacarboxylic acid (B5CA). Structural regions of the two exemplary DBC molecules are colored purple and green to indicate how the condensed aromatic core could oxidize to produce either B6CA or B5CA [16].

<https://doi.org/10.1371/journal.pone.0268059.g001>

condensed aromatic carbon standards or analytical approaches is capable of representing the entire suite of DBC molecules that exist in aquatic matrices. Therefore, we recognize that the relationship (the conversion factor) between the DBC compound class and its BPCA oxidation products may be altered or further refined with continued research. Therefore, we strongly encourage the publication of raw BPCA concentrations alongside calculated DBC concentrations whenever the BPCA method described here is used to produce the data. In doing so, works involving published DBC concentrations can be easily adapted if new conversion factors are established and will facilitate the synthesis and scaling of DBC data among studies. Should it be shown that non-condensed aromatic structures can yield considerable BPCA products upon oxidation, the definition of DBC and its context in global biogeochemical cycling would need to be revisited. To assist in answering these questions, we provide a detailed protocol for those looking to adopt a method for the quantification of this biogeochemically relevant, highly photolabile, yet environmentally long-lived fraction of DOC.

The protocol we use to quantify DBC in aquatic samples via BPCA analysis is summarized here. DOC is isolated from the filtered water sample via solid-phase extraction (SPE) following Dittmar et al. [27]. The adsorbed DOC is eluted from the SPE cartridges using methanol and recovered as solid-phase extracted DOC (SPE-DOC). A volume of methanol eluate equating to ~0.5 mg-DOC is placed in a glass ampule and evaporated. Concentrated nitric acid is added to the ampule containing the DOC eluate residue and is flame-sealed. The sealed ampules are oxidized in an oven for 6 hours at 160°C [28]. After oxidation, the ampules are opened and the reaction mixture is evaporated to dryness under a stream of nitrogen gas. The remaining BPCA-containing residue is re-dissolved in a phosphoric acid solution for subsequent analysis. The BPCAs are separated using an established high-performance liquid chromatography (HPLC) method and quantified using a diode array detector. This method separates B6CA and B5CA peaks, which are identified by comparing retention times to that of BPCA standards. Sample BPCA concentrations are determined by external calibration and are scaled to DBC concentrations using the Stubbins et al. [21] conversion factor.

The BPCA method described in this protocol is not new and details of method development are published elsewhere [17]. However, the many small, yet important, details and full descriptions of the preparative and analytical techniques are usually omitted for brevity in published

method sections and method-specific papers. These technical omissions make it difficult for researchers who are looking to adopt or refine the BPCA method. With this detailed laboratory protocol, we outline the step-by-step considerations and techniques omitted from previous publications. We aim to lower the barrier to entry for those who want to incorporate BPCA analysis and DBC quantification into their research but do not have a high level of access to those familiar with the method. The contribution of additional perspectives and studies examining DBC will further the understanding of the production, fate, and impact of DBC in the environment.

Materials and methods

The protocol described in this peer-reviewed article is published on protocols.io, <https://doi.org/10.17504/protocols.io.5qpvoy2b9g4o/v2>, and is included for printing as [S1 File](#) with this article.

Expected results

By following this protocol, you will obtain concentrations of B6CA and B5CA produced from the oxidation of DBC in water samples. Sample BPCA peaks are identified using absorbance spectra and/or peak retention times of BPCA standards. Chromatograms of BPCA standards should not contain extraneous peaks (besides void volume) but may have baseline variations resulting from mobile phase gradients (Fig 2A and 2B). Sample chromatograms will likely have many extraneous peaks that reflect oxidation products other than B5CA and B6CA, however, targeted BPCAs are usually baseline-separated (Fig 2C). DBC concentrations (in $\mu\text{M-C}$) are calculated from B5CA and B6CA concentrations (in nM-BPCA) using the following equation (from Stubbins et al. [21]): $[\text{DBC}] = 0.0891 \times ([\text{B6CA}] + [\text{B5CA}])^{0.9175}$




Fig 2. HPLC chromatograms of (a) mobile phase gradient as well as benzenepolycarboxylic acid separation in (b) a standard mixture and (c) a dissolved organic matter sample from a fire-affected headwater stream. As seen in each chromatogram, the initial peak is representative of void volume and the baseline variation is due to changes in mobile phase gradient. The BPCA standard peaks in **b** are indicative of B6CA, B5CA, a B4CA (1,2,4,5-B4CA), and three B3CAs (1,2,4-B3CA, 1,2,3-B3CA, 1,3,5-B3CA). B3CAs and B4CA are not quantified as part of this method but are included in chromatographic analyses to confirm complete elution of sample.

<https://doi.org/10.1371/journal.pone.0268059.g002>

Using the described protocol, we have measured and published DBC data for diverse aquatic systems, including major and minor rivers, along oceanic depth profiles, throughfall and stemflow, petroleum-contaminated groundwater, and coastal environments [13, 18, 29–32]. Although DOC analysis is not explicitly described in the current protocol, we strongly recommend that users of this method measure and share bulk DOC concentrations alongside BPCA and DBC concentrations. Ratios of DBC to DOC are often reported because they can vary between fresh and marine water, vary between types of rivers, and decrease with increased photodegradation [3, 31, 33]. Molar ratios of B6CA to B5CA are also useful as a proxy to express the degree of relative condensed aromaticity of DBC among different samples. Higher B6CA:B5CA ratios indicate DBC pools containing larger fused aromatic ring structures [34]. B6CA to B5CA ratios have been found to decrease with increased photodegradation, are usually lower in oceans compared to rivers, and have been found to be higher at high river flow than low river flow [21, 31, 33].

Here, we display a subset of DBC data originally reported by Wagner et al. [31] that was determined using this protocol. The dataset includes both oceanic and riverine samples to show examples of distinctive DBC results that may be observed for diverse aquatic systems. Considering the source and prior biogeochemical processing of a target water sample is critical for making any necessary sample volume adjustments or other accommodations during sample preparation and analysis. For example, oceanic samples have significantly lower SPE efficiencies (i.e. lower DOC recovery by SPE), lower DOC and DBC concentrations, and lower DBC:DOC and B6CA:B5CA ratios than riverine samples (Table 1). Since proportions of DBC:DOC and B6CA:B5CA are inherent for any given sample, we would conclude that higher sample volumes are needed for oceanic samples than river samples to recover sufficient amounts of DBC for robust BPCA detection and quantification. When working with new samples, where DBC:DOC and B6CA:B5CA ratios are truly unknown, we recommend oxidizing 0.5 mg SPE-DOC to gain a preliminary sense of sample composition with regards to condensed aromatic DBC. It is important to remember that, since B6CA:B5CA ratios are typically < 1 (Table 1), minimal sample volumes required for DBC analysis should be estimated from the expected yield of B6CA. The limits of detection and quantification for the BPCA method using the protocol described here are 1.5 and 5 μM -B6CA, respectively. Based upon these analytical detection limits and our prior experience [17], we recommend a lower reporting limit of 20 μM -B6CA (in vial, to be injected for HPLC separation). Oftentimes, BPCA ratios such as B6CA:B5CA are used to describe relative degree of DBC aromaticity and to trace changes in DBC sources and biogeochemistry over time [21, 34]. As B6CA is almost always present in

Table 1. A subset of DBC data originally published by Wagner et al. [31].

Site	SPE-DOC Recovery	DOC ($\mu\text{M-C}$)	DBC ($\mu\text{M-C}$)	DBC: DOC (%)	B6CA (nM)	B5CA (nM)	B6CA: B5CA
Pacific Ocean	40%	81	0.62	0.8	1.6	6.7	0.23
Atlantic Ocean	48%	63	0.78	1.2	2.1	8.6	0.24
Amazon River	78%	331	25	7.6	140	332	0.42
Congo River	64%	540	37	6.8	234	473	0.49
Northern Dvina River	64%	519	42	8.0	295	516	0.57
Kolyma River	70%	327	20	6.2	112	260	0.43
Mississippi River	61%	247	18	7.3	75	251	0.30

Water samples were collected from the Pacific and Atlantic Oceans as well as the Amazon, Congo, Northern Dvina, and Mississippi Rivers. Each sample was analyzed for dissolved organic carbon (DOC), recovery of dissolved organic carbon after solid-phase extraction (SPE-DOC), dissolved black carbon (DBC), the ratio of DBC to DOC, the concentrations of B6CA and B5CA oxidation products, and the ratio of B6CA to B5CA.

<https://doi.org/10.1371/journal.pone.0268059.t001>

lower concentrations than B5CA in aqueous samples, establishing a lower reporting limit of 20 μM -B6CA ensures the calculation of robust DBC concentrations and BPCA ratios. Furthermore, if users adhere to these suggested limits, then samples prepared following this protocol could also be used for BPCA-specific stable carbon isotopic analysis [17].

However, reaching this target B6CA concentration requires different sample volumes for biogeochemically distinct sample types, like river and open ocean water. If we assume a typical river sample has a DOC concentration of 520 μM , DBC:DOC ratio of 8%, an extraction efficiency of 65%, and a B6CA:B5CA ratio of 0.57, then a minimum of 20 mL of sample volume is required to recover sufficient carbon for DBC analysis. In contrast, if we assume a typical open ocean sample has a DOC concentration of 50 μM , DBC:DOC ratio of 1% and an extraction efficiency of 45%, and a B6CA:B5CA ratio of 0.23, then a minimum of 4.5 L of sample volume would be necessary. These estimates are conservative, to ensure robust BPCA measurement. Required sample volumes may be further reduced depending upon sample composition, extraction efficiency, injection volumes, specific detection limits of the instrument used for separation and detection, etc. Users of this method are encouraged to use the BPCA preparation and calculation spreadsheets (S2 File) to estimate sample volumes needed to conduct robust DBC analyses in the specific aquatic system under study.

Supporting information

S1 File. Protocol collection. Step-by-step protocol, also available on protocols.io. (PDF)

S2 File. Example PPL-BPCA calculations spreadsheet, also available on protocols.io. (XLSX)

Acknowledgments

We thank the two anonymous reviewers for their constructive comments.

Author Contributions

Conceptualization: Riley Barton, Sasha Wagner.

Data curation: Riley Barton.

Funding acquisition: Sasha Wagner.

Methodology: Riley Barton, Sasha Wagner.

Project administration: Sasha Wagner.

Resources: Sasha Wagner.

Supervision: Sasha Wagner.

Writing – original draft: Riley Barton.

Writing – review & editing: Riley Barton, Sasha Wagner.

References

1. Knicker H. Pyrogenic organic matter in soil: Its origin and occurrence, its chemistry and survival in soil environments. *Quat Int.* 2011; 243(2):251–63. <https://doi.org/10.1016/j.quaint.2011.02.037>
2. Masiello CA. New directions in black carbon organic geochemistry. *Mar Chem.* 2004; 92(1–4 SPEC. ISS.):201–13. <https://doi.org/10.1016/j.marchem.2004.06.043>

3. Wagner S, Jaffé R, Stubbins A. Dissolved black carbon in aquatic ecosystems. *Limnol Oceanogr Lett*. 2018; 3(3):168–85. <https://doi.org/10.1002/lol2.10076>
4. Singh BP, Cowie AL, Smernik RJ. Biochar carbon stability in a clayey soil as a function of feedstock and pyrolysis temperature. *Environ Sci Technol*. 2012; 46(21):11770–8. <https://doi.org/10.1021/es302545b> PMID: 23013285
5. Kuzyakov Y, Bogomolova I, Glaser B. Biochar stability in soil: Decomposition during eight years and transformation as assessed by compound-specific ^{14}C analysis. *Soil Biol Biochem*. 2014; 70:229–36. <https://doi.org/10.1016/j.soilbio.2013.12.021>
6. Jones MW, Santín C, van der Werf GR, Doerr SH. Global fire emissions buffered by the production of pyrogenic carbon. *Nat Geosci*. 2019; 12(9):742–7. <https://doi.org/10.1038/s41561-019-0403-x>
7. Jaffé R, Ding Y, Niggemann J, Vähätalo A V., Stubbins A, Spencer RGM, et al. Global charcoal mobilization from soils via dissolution and riverine transport to the oceans. *Science* (80-). 2013; 340(6130):345–7. <https://doi.org/10.1126/science.1231476> PMID: 23599492
8. Jones MW, Coppola AI, Santín C, Dittmar T, Jaffé R, Doerr SH, et al. Fires prime terrestrial organic carbon for riverine export to the global oceans. *Nat Commun*. 2020; 11(1):4–11. <https://doi.org/10.1038/s41467-020-16576-z>
9. Abiven S, Hengartner P, Schneider MPW, Singh N, Schmidt MWI. Pyrogenic carbon soluble fraction is larger and more aromatic in aged charcoal than in fresh charcoal. *Soil Biol Biochem*. 2011; 43(7):1615–7. <https://doi.org/10.1016/j.soilbio.2011.03.027>
10. Roebuck JA, Podgorski DC, Wagner S, Jaffé R. Photodissolution of charcoal and fire-impacted soil as a potential source of dissolved black carbon in aquatic environments. *Org Geochem*. 2017; 112:16–21. <https://doi.org/10.1016/j.orggeochem.2017.06.018>
11. Wagner S, Ding Y, Jaffé R. A new perspective on the apparent solubility of dissolved black carbon. *Front Earth Sci*. 2017; 5(September):1–16. <https://doi.org/10.3389/feart.2017.00075>
12. Wozniak AS, Goranov AI, Mitra S, Bostick KW, Zimmerman AR, Schlesinger DR, et al. Molecular heterogeneity in pyrogenic dissolved organic matter from a thermal series of oak and grass chars. *Org Geochem*. 2020; 148:104065. <https://doi.org/10.1016/j.orggeochem.2020.104065>
13. Wagner S, Harvey E, Baetge N, McNair H, Arrington E, Stubbins A. Investigating Atmospheric Inputs of Dissolved Black Carbon to the Santa Barbara Channel During the Thomas Fire (California, USA). *J Geophys Res Biogeosciences*. 2021; 126(8):1–17. <https://doi.org/10.1029/2021JG006442>
14. Wagner S, Coppola AI, Stubbins A, Dittmar T, Niggemann J, Drake TW, et al. Questions remain about the biolability of dissolved black carbon along the combustion continuum. *Nat Commun*. 2021; 12(1):10–3. <https://doi.org/10.1038/s41467-021-24477-y>
15. Glaser B, Haumaier L, Guggenberger G, Zech W. Black carbon in soils: The use of benzenecarboxylic acids as specific markers. *Org Geochem*. 1998; 29(4):811–9. [https://doi.org/10.1016/S0146-6380\(98\)00194-6](https://doi.org/10.1016/S0146-6380(98)00194-6)
16. Dittmar T. The molecular level determination of black carbon in marine dissolved organic matter. *Org Geochem*. 2008 Apr 1; 39(4):396–407. <https://doi.org/10.1016/j.orggeochem.2008.01.015>
17. Wagner S, Brandes J, Goranov AI, Drake TW, Spencer RGM, Stubbins A. Online quantification and compound-specific stable isotopic analysis of black carbon in environmental matrices via liquid chromatography-isotope ratio mass spectrometry. *Limnol Oceanogr Methods*. 2017; 15(12):995–1006. <https://doi.org/10.1002/lom3.10219>
18. Podgorski DC, Zito P, Kellerman AM, Bekins BA, Cozzarelli IM, Smith DF, et al. Hydrocarbons to carboxyl-rich alicyclic molecules: A continuum model to describe biodegradation of petroleum-derived dissolved organic matter in contaminated groundwater plumes. *J Hazard Mater*. 2021; 402:123998. <https://doi.org/10.1016/j.jhazmat.2020.123998> PMID: 33254831
19. Goranov AI, Schaller MF, Long JA, Podgorski DC, Wagner S. Characterization of Asphaltenes and Petroleum Using Benzenepolycarboxylic Acids (BPCAs) and Compound-Specific Stable Carbon Isotopes. *Energy and Fuels*. 2021; 35(22):18135–45. <https://doi.org/10.1021/acs.energyfuels.1c02374>
20. Ziolkowski LA, Chamberlin AR, Greaves J, Druffel ERM. Quantification of black carbon in marine systems using the benzene polycarboxylic acid method: A mechanistic and yield study. *Limnol Oceanogr Methods*. 2011; 9(4):140–140. <https://doi.org/10.4319/lom.2011.9.140>
21. Stubbins A, Spencer RGM, Mann PJ, Holmes RM, McClelland JW, Niggemann J, et al. Utilizing colored dissolved organic matter to derive dissolved black carbon export by arctic rivers. *Front Earth Sci*. 2015; 3(October):1–11. <https://doi.org/10.3389/feart.2015.00063>
22. Fang Y, Chen Y, Huang G, Hu L, Tian C, Xie J, et al. Particulate and Dissolved Black Carbon in Coastal China Seas: Spatiotemporal Variations, Dynamics, and Potential Implications. *Environ Sci Technol*. 2021; 55(1):788–96. <https://doi.org/10.1021/acs.est.0c06386> PMID: 33275416

23. Bostick KW, Zimmerman AR, Wozniak AS, Mitra S, Hatcher PG. Production and composition of pyrogenic dissolved organic matter from a logical series of laboratory-generated chars. *Front Earth Sci*. 2018; 6(April):1–14. <https://doi.org/10.3389/feart.2018.00043>
24. Dittmar T, Paeng J. A heat-induced molecular signature in marine dissolved organic matter. *Nat Geosci*. 2009; 2(3):175–9. <https://doi.org/10.1038/ngeo440>
25. Dittmar T, Koch BP. Thermogenic organic matter dissolved in the abyssal ocean. *Mar Chem*. 2006; 102(3–4):208–17. <https://doi.org/10.1016/j.marchem.2006.04.003>
26. Kappenberg A, Bläsing M, Lehndorff E, Amelung W. Black carbon assessment using benzene polycarboxylic acids: Limitations for organic-rich matrices. *Org Geochem*. 2016; 94:47–51. <https://doi.org/10.1016/j.orggeochem.2016.01.009>
27. Dittmar T, Koch B, Hertkorn N, Kattner G. A simple and efficient method for the solid-phase extraction of dissolved organic matter (SPE-DOM) from seawater. *Limnol Oceanogr Methods*. 2008; 6(6):230–5. <https://doi.org/10.4319/lom.2008.6.230>
28. Ding Y, Yamashita Y, Dodds WK, Jaffé R. Dissolved black carbon in grassland streams: Is there an effect of recent fire history? *Chemosphere*. 2013; 90(10):2557–62. <https://doi.org/10.1016/j.chemosphere.2012.10.098> PMID: 23219080
29. Wagner S, Cawley KM, Rosario-Ortiz FL, Jaffé R. In-stream sources and links between particulate and dissolved black carbon following a wildfire. *Biogeochemistry*. 2015; 124(1–3):145–61. <https://doi.org/10.1007/s10533-015-0088-1>
30. Drake TW, Wagner S, Stubbins A, Wabakanghanzi JN, Dinga JB, Six J, et al. Du Feu à l'Eau: Source and Flux of Dissolved Black Carbon From the Congo River. *Global Biogeochem Cycles*. 2020; 34(8):1–13. <https://doi.org/10.1029/2020GB006560>
31. Wagner S, Brandes J, Spencer RGM, Ma K, Rosengard SZ, Moura JMS, et al. Isotopic composition of oceanic dissolved black carbon reveals non-riverine source. *Nat Commun*. 2019; 10(1):1–8. <https://doi.org/10.1038/s41467-019-13111-7>
32. Wagner S, Brantley S, Stuber S, Van Stan J, Whitetree A, Stubbins A. Dissolved black carbon in throughfall and stemflow in a fire-managed longleaf pine woodland. *Biogeochemistry*. 2019; 146(2):191–207. <https://doi.org/10.1007/s10533-019-00620-2>
33. Stubbins A, Niggemann J, Dittmar T. Photo-lability of deep ocean dissolved black carbon. *Biogeosciences*. 2012; 9(5):1661–70. <https://doi.org/10.5194/bg-9-1661-2012>
34. Schneider MPW, Hilf M, Vogt UF, Schmidt MWI. The benzene polycarboxylic acid (BPCA) pattern of wood pyrolyzed between 200°C and 1000°C. *Org Geochem*. 2010; 41(10):1082–8. <https://doi.org/10.1016/j.orggeochem.2010.07.001>