

Relatively geometric actions on CAT(0) cube complexes

Eduard Einstein¹ | Daniel Groves²

¹ Department of Mathematics, University of Pittsburgh, Pittsburgh, PA, USA

² Department of Mathematics, Statistics, and Computer Science, University of Illinois at Chicago, Chicago, IL, USA

Correspondence

Daniel Groves, Department of Mathematics, Statistics, and Computer Science, University of Illinois at Chicago, 322 Science and Engineering Offices (M/C 249), 851 S. Morgan St., Chicago, IL 60607-7045, USA.

Email: groves@math.uic.edu

Funding information

NSF, Grant/Award Number: DMS-1904913

Abstract

We develop the foundations of the theory of relatively geometric actions of relatively hyperbolic groups on CAT(0) cube complexes, a notion introduced in our previous work (Einstein and Groves [Compos. Math. (2020), no. 4, 862–867]). In the relatively geometric setting, we prove full relatively quasi-convex subgroups are convex cocompact, an analog of Agol’s Theorem and a version of Haglund–Wise’s Canonical Completion and Retraction.

MSC 2020

20F67, 20F65, 57M07 (primary)

1 | INTRODUCTION

The interaction between the geometry of CAT(0) cube complexes and that of hyperbolic groups is at the center of some of the most powerful aspects of Haglund and Wise’s theory of (virtually) special cube complexes [12], which in turn was at the center of the resolution of the Virtual Haken Conjecture [1], and the Virtual Fibering Conjecture [1] in the closed case. In the case of finite-volume hyperbolic 3-manifolds, the Virtual Fibering Conjecture was resolved by Wise [22], also using virtually special cube complexes, but now using the relatively hyperbolic geometry of the fundamental group.

In search of more general results, there have been numerous papers dealing with relatively hyperbolic groups acting on CAT(0) cube complexes. See, for example, [15, 21], or [20]. These papers typically deal with proper actions, which are either cocompact, or co-sparse.

In [5], we introduced a new kind of action for a relatively hyperbolic group on a CAT(0) cube complex, a *relatively geometric action* (see Definition 1.1 below). If a relatively hyperbolic pair (G, \mathcal{P}) acts relatively geometrically on a space X , then X is quasi-isometric to the coned-off Cayley graph for (G, \mathcal{P}) , and hence in particular is δ -hyperbolic for some δ . We believe that relatively

geometric actions on CAT(0) cube complexes should have as rich a theory as hyperbolic groups acting properly and cocompactly on CAT(0) cube complexes. In this paper, we begin a systematic investigation of such actions. In a relatively geometric action, the parabolic subgroups \mathcal{P} act elliptically on the cube complex, so in contrast to either of the kinds of proper actions mentioned above, relatively geometric actions may exist even when the parabolic subgroups do not act on CAT(0) cube complexes in interesting ways. Therefore, we expect that the class of relatively hyperbolic groups acting relatively geometrically on CAT(0) cube complexes is substantially broader than the class acting geometrically, thus bringing these powerful techniques to bear in a much wider setting. A final reason that relatively geometric actions are desirable is that the techniques of Dehn filling developed by the second author and Manning in [8] are applicable to relatively geometric actions, as we used already in [5]. This is a theme we further explore in this paper.

We now state our main results. First, recall the definition of relatively geometric (see Section 2 for other definitions).

Definition 1.1. Suppose that (G, \mathcal{P}) is a group pair. A (cellular) action of G on a cell complex \tilde{X} is *relatively geometric (with respect to \mathcal{P})* if

- (1) $G \setminus \tilde{X}$ is compact;
- (2) each element of \mathcal{P} acts elliptically on \tilde{X} ; and
- (3) each stabilizer in G of a cell in \tilde{X} is either finite or else conjugate to a finite-index subgroup of an element of \mathcal{P} .

In order to state our results, we make the following standing assumption.

Assumption 1.2. Suppose that (G, \mathcal{P}) is relatively hyperbolic and that \tilde{X} is a CAT(0) cube complex which admits a relatively geometric action of G with respect to \mathcal{P} . Let $X = G \setminus \tilde{X}$.

In Sageev–Wise [21] and Haglund [11] it is proved that if a hyperbolic group G acts geometrically on a CAT(0) cube complex and H is a quasi-convex subgroup of G then there is a convex H -invariant and H -cocompact sub-complex. Thus, H also acts properly and cocompactly on a CAT(0) cube complex, and in the virtually special setting this is what allows the canonical completion and retraction construction of Haglund–Wise [12] to be applied.

Our first result, proved in Section 3, is a relatively geometric analogue of the above-mentioned results of [21] and [11]. For a relatively hyperbolic analogue in the proper and cocompact or cosparse settings, see [21].

Theorem 1.3. *Under Assumption 1.2, for any full relatively quasi-convex subgroup H of G and any compact $K \subset \tilde{X}$ there exists a convex H -invariant sub-complex \tilde{Y} of \tilde{X} so that $K \subset \tilde{Y}$ and $H \setminus \tilde{Y}$ is compact.*

In order to solve the Virtual Haken and Virtual Fibering Conjectures, Agol proved that any hyperbolic group acting geometrically on a CAT(0) cube complex is virtually special. The next result is a relatively geometric version of Agol’s Theorem, and is proved in Section 4.

Theorem 1.4. *Under Assumption 1.2, if the elements of \mathcal{P} are residually finite, then there is a finite-index subgroup G_0 of G so that $G_0 \setminus \tilde{X}$ is a special cube complex.*

Note that $G_0 \setminus \tilde{X}$ should be considered as a complex of groups, rather than merely as a space, so the underlying space being a special cube complex is not obviously as useful as in the case of Agol's Theorem. However, there are still favorable separability properties—see Theorem 1.6 and Corollary 1.7 below.

In Section 4, we consider a full relatively quasi-convex subgroup H of G and explore the behavior of the hull \tilde{Y} found in Theorem 1.3 under long Dehn fillings. In particular, a special case of what we prove is the following result, proved in Section 4. See Sections 2 and 4 for definitions of terms, and Proposition 4.1 for a description of the subgroups \mathcal{Q} .

Theorem 1.5. *Make Assumption 1.2, and suppose that elements of \mathcal{P} are residually finite. Let $H \leq G$ be a full relatively quasi-convex subgroup and let \tilde{Y} be as in Theorem 1.3. For sufficiently long $(\mathcal{Q} \cup H)$ -fillings $G \rightarrow G/K$, if $K_H = K \cap H$, then $\bar{X} = K \setminus \tilde{X}$ and $\bar{Y} = K_H \setminus \tilde{Y}$ are both CAT(0) cube complexes and the natural map $\bar{Y} \rightarrow \bar{X}$ is an embedding with convex image.*

In Section 5, we develop a relatively geometric version of the Canonical Completion and Retraction (see Theorem 5.2). An application of this construction is the following, proved in Section 5.

Theorem 1.6. *Under Assumption 1.2, if the elements of \mathcal{P} are residually finite, then for any full relatively quasi-convex subgroup H of G there is a finite-index subgroup $H_0 \leq H$ which is a retract of a finite-index subgroup of G .*

Corollary 1.7. *Under Assumption 1.2, if every $P \in \mathcal{P}$ is residually finite:*

- (1) *G is residually finite and*
- (2) *every full relatively quasiconvex subgroup of G is separable.*

The first item is an immediate consequence of Theorem 2.11 and Corollary 4.2. The second item follows from the residual finiteness of G and Theorem 1.6. The separability properties of Corollary 1.7 also follow from [9, Theorem 4.7], which works in the more general setting of ‘weakly relatively geometric’ actions.

We expect the tools in this paper to be of foundational interest for the further study of relatively geometric actions. In a future work, we plan to prove a relatively geometric version of the cubulation result of Hsu–Wise [16]. We expect such a relatively geometric analogue of the Hsu–Wise results to be the key to proving a relatively geometric version of Wise's Quasi-convex Hierarchy Theorem, one of the main results from [22]. Such a result would provide a wealth of examples of relatively geometric actions on CAT(0) cube complexes.

Convention 1.8. Conjugation of g by h , written g^h , is hgh^{-1} .

The authors thank the referee for a very careful and thoughtful report, which improved this paper.

2 | PRELIMINARIES

In this section we collect some basic definitions and results about relatively hyperbolic groups and relatively hyperbolic Dehn filling needed for this paper.

2.1 | Relatively hyperbolic groups and relatively quasi-convex subgroups

Definition 2.1 (Combinatorial horoball) [7, Definition 3.1]. Let Γ be a 1-complex. The *combinatorial horoball based on Γ* , denoted $\mathcal{H}(\Gamma)$, is the 1-complex whose vertices are $\Gamma^{(0)} \times (\{0\} \cup \mathbb{N})$, and whose edge set consists of (i) edges between $(v, 0)$ and $(w, 0)$ whenever there is an edge between v and w in Γ ; (ii) for all $k > 0$ and all $v, w \in \Gamma^{(0)}$ so that $0 < d_\Gamma(v, w) \leq 2^k$, an edge between (v, k) and (w, k) ; and (iii) edges between (v, k) and $(v, k+1)$ for all $v \in \Gamma^{(0)}$ and all $k \geq 0$.

In [7], 2 cells are added to combinatorial horoballs, but we do not need them here. According to [7, Theorem 3.8, Remark 3.9], $\mathcal{H}(\Gamma)$ is 20-hyperbolic for any connected 1-complex Γ .

Definition 2.2 (Cusped space) [7, Definition 3.12]. Let G be a group and \mathcal{P} a finite collection of subgroups of G . Further, let S be a generating set for G so that $\langle S \cap P \rangle = P$ for each $P \in \mathcal{P}$. Let $\Gamma(G, S)$ be the Cayley graph of G with respect to S . The *cusped space* for (G, \mathcal{P}) , denoted $\mathcal{C}(G, \mathcal{P}, S)$, is obtained by gluing a copy of the combinatorial horoball over the Cayley graph $\Gamma_P(S \cap P)$ of P (with respect to $S \cap P$) to each left translate of the natural copy of $\Gamma_P(S \cap P)$ in $\Gamma(G, S)$.

Definition 2.3 [7, Theorem 3.25]. Let (G, \mathcal{P}) be a group pair, with G finitely generated, \mathcal{P} finite, and each element of \mathcal{P} finitely generated. Then (G, \mathcal{P}) is *relatively hyperbolic* if for some (any) finite generating set S for G so that $\langle S \cap P \rangle = P$ for each $P \in \mathcal{P}$, the cusped space $\mathcal{C}(G, \mathcal{P}, S)$ is ν -hyperbolic, for some ν .

There are analogous definitions of relatively hyperbolic pairs (G, \mathcal{P}) when G is not finitely generated, or when \mathcal{P} is not finite (see [14, 18], for example) but we do not need them here. The hyperbolicity of the cusped space for (G, \mathcal{P}) does not depend on the choice of S , though of course the value of ν does.

Convention 2.4. Whenever (G, \mathcal{P}) is relatively hyperbolic, we assume elements of \mathcal{P} are infinite. If this is not the case, discard the finite elements. This does not affect the relative hyperbolicity, or whether or not a given action is relatively geometric.

Definition 2.5. Let (G, \mathcal{P}) be relatively hyperbolic, and let $H \leq G$. The *induced peripheral structure on H* is a collection \mathcal{D} of representatives of H -conjugacy classes of maximal infinite parabolic subgroups of H .

Suppose (G, \mathcal{P}) is relatively hyperbolic, that $H \leq G$, and let \mathcal{D} be the induced peripheral structure on H . Given $D \in \mathcal{D}$, there exists $P_D \in \mathcal{P}$ and $c_D \in G$ so that $D \leq P_D^{c_D}$. Associated to the pairs (G, \mathcal{P}) and (H, \mathcal{D}) (along with choices of generating sets) are cusped spaces X_G and X_H . As explained in [2, §3], the inclusion $\iota: H \rightarrow G$ extends to an H -equivariant Lipschitz proper map $\tilde{\iota}: X_H \rightarrow X_G$.

Definition 2.6 [10, Definition 2.9] (see also [2, Definition 3.11]). Suppose (G, \mathcal{P}) is relatively hyperbolic, that $H \leq G$ and that \mathcal{D} is the induced peripheral structure on H . The subgroup H is *relatively quasi-convex* in (G, \mathcal{P}) if $\tilde{\iota}(X_H)$ is quasi-convex in X_G .

There are many equivalent characterizations of quasi-convexity in relatively hyperbolic groups. See Hruska [14] for five other conditions (which Hruska proves are all equivalent), and see [17] and [10] for a proof that the above definition is equivalent to Hruska's.

Definition 2.7. Let (G, \mathcal{P}) be relatively hyperbolic, $H \leq G$ a subgroup and \mathcal{D} the induced peripheral structure on H . The subgroup H is *full* if for every $D \in \mathcal{D}$, D is finite-index in a maximal parabolic.

Since we assume elements of \mathcal{P} are finitely generated, and relatively quasi-convex subgroups are themselves relatively hyperbolic we have the following immediate consequence.

Lemma 2.8. *Any full relatively quasi-convex subgroup of (G, \mathcal{P}) is finitely generated.*

Definition 2.9. Let (G, \mathcal{P}) be relatively hyperbolic and C be the cusped space as in Definition 2.2.

A geodesic γ in C *penetrates a horoball B* to depth $R > 0$ if there exists a $\xi \in \gamma \cap B$ with $d(\xi, C \setminus B) \geq R$.

Proposition 2.10 [17, Proposition A.6]. *Let (G, \mathcal{P}) be relatively hyperbolic, C a cusped space for (G, \mathcal{P}) and $H \leq G$ relatively quasi-convex. There exists a constant R depending on G, C and H so that whenever a horoball B in C is R -penetrated by H we have $|\text{Stab}_G(B) \cap H| = \infty$.*

2.2 | Relatively hyperbolic Dehn filling

Suppose (G, \mathcal{P}) is a group pair and that $\mathcal{N} = \{N_P \trianglelefteq P \mid P \in \mathcal{P}\}$ is a collection of normal subgroups of the peripheral groups. The *Dehn filling* of (G, \mathcal{P}) determined by \mathcal{N} is the pair $(\overline{G}, \overline{\mathcal{P}})$, where $\overline{G} = G/K$, for K the normal closure in G of $\bigcup \mathcal{N}$, and $\overline{\mathcal{P}}$ is the image of \mathcal{P} in \overline{G} . The elements of \mathcal{N} are called *filling kernels*. We sometimes also write $\overline{G} = G(N_P \mid P \in \mathcal{P})$, when we want to make the dependence on the choice of filling kernels explicit.

The filling is *peripherally finite* if N_P is finite-index in P for all $P \in \mathcal{P}$. If $H < G$ then the filling is an *H -filling* if for every $g \in G$, $|H \cap P^g| = \infty$ implies $N_P^g \leq H$. If \mathcal{H} is a family of subgroups, an *\mathcal{H} -filling* is a filling which is an H -filling for all $H \in \mathcal{H}$.

A property P holds for all sufficiently long fillings of (G, \mathcal{P}) if there is a finite set $S \subset \bigcup \mathcal{P}$ so that P holds for any filling where $(\bigcup \mathcal{N}) \cap S = \emptyset$. There is an obvious meaning to phrases such as 'for all sufficiently long H -fillings', and so on.

The following result is the basic result of relatively hyperbolic Dehn filling, and is due to Osin [19] (see also [7]).

Theorem 2.11 [19, Theorem 1.1]. *Suppose (G, \mathcal{P}) is relatively hyperbolic, and $\mathcal{F} \subset G$ is finite. For sufficiently long Dehn fillings $G \rightarrow \overline{G} = G(N_P \mid P \in \mathcal{P})$ we have:*

- (1) *For each $P \in \mathcal{P}$, the canonical map $P/N_P \rightarrow \overline{G}$ is injective, denote the image by \overline{P} ;*
- (2) *$(\overline{G}, \{\overline{P} \mid P \in \mathcal{P}\})$ is relatively hyperbolic;*
- (3) *The map $G \rightarrow \overline{G}$ is injective on \mathcal{F} .*

In [1, 2, 10] various results have been proved controlling the image of relatively quasi-convex subgroups under Dehn filling. We follow [10].

Definition 2.12. Let (G, \mathcal{P}) be relatively hyperbolic, and $H \leq G$ be relatively quasi-convex. Let \mathcal{D} be the induced peripheral structure on H , and for each $D \in \mathcal{D}$ let $P_D \in \mathcal{P}$ and $c_D \in G$ be so $D \leq P_D^{c_D}$. Let $G \rightarrow G(N_P \mid P \in \mathcal{P})$ be a Dehn filling. For $D \in \mathcal{D}$, write $N_D = N_{P_D}^{c_D} \cap D$. The *induced filling kernels for (H, \mathcal{D})* are $\{N_D\}_{D \in \mathcal{D}}$. The *induced filling for H* is the filling $H \rightarrow H(N_D \mid D \in \mathcal{D})$.

Given (G, \mathcal{P}) , $H \leq G$, a Dehn filling $G \rightarrow G(N_P \mid P \in \mathcal{P})$, and the induced filling $H \rightarrow H(N_D \mid D \in \mathcal{D})$ as in Definition 2.12, it is clear that there is an induced map $\bar{\iota} : H(N_D \mid D \in \mathcal{D}) \rightarrow G(N_P \mid P \in \mathcal{P})$.

The following is an immediate consequence of [10, Propositions 4.5 and 4.6] and [10, Lemma 3.7] (which states that a sufficiently long H -filling is sufficiently ‘ H -wide’).

Theorem 2.13. Suppose (G, \mathcal{P}) is relatively hyperbolic and $H \leq G$ is relatively quasi-convex. For sufficiently long H -fillings $G \rightarrow \bar{G}$, the induced map $\bar{\iota} : H(N_D \mid D \in \mathcal{D}) \rightarrow G(N_P \mid P \in \mathcal{P})$ is injective, with relatively quasi-convex image.

3 | BASIC PROPERTIES OF RELATIVELY GEOMETRIC ACTIONS

Throughout this section, we make Assumption 1.2. Let $q : \tilde{X} \rightarrow G \setminus \tilde{X}$ be the quotient map.

As noted in [5], the results of Charney–Crisp [4, Theorem 5.1] immediately imply the following result.

Proposition 3.1. The space \tilde{X} is quasi-isometric to the coned-off Cayley graph of (G, \mathcal{P}) , and in particular is δ -hyperbolic for some δ .

Using one of Hruska’s characterizations of relatively quasi-convexity from [14] we obtain the following result.

Corollary 3.2. Suppose that H is a relatively quasi-convex subgroup of G , and that $x \in \tilde{X}$. The orbit $H \cdot x$ is quasi-convex in \tilde{X} .

Proof. According to [14, (QC-5)], any H -orbit in the coned-off Cayley Γ graph is quasi-convex (in fact, (QC-5) is stronger than this). Since \tilde{X} is δ -hyperbolic, quasi-isometries take quasi-convex subsets to quasi-convex subsets, so a quasi-isometry of pairs $(\Gamma, H \cdot 1) \rightarrow (\tilde{X}, H \cdot x)$ implies the result. \square

3.1 | Stabilizers of cells

Notation 3.3. Suppose G is a group and that $H_1, H_2 \leq G$. If H_1 and H_2 are commensurable in G , then we write $H_1 \sim H_2$.

Definition 3.4. Let (G, \mathcal{P}) be relatively hyperbolic and let G admit a relatively geometric action on a CAT(0) cube complex \tilde{X} . For $P \in \mathcal{P}$ define a sub-complex of \tilde{X}

$$C(P) = \bigcup_{\text{cells } \sigma} \{\sigma \mid \text{Stab}(\sigma) \sim P\}.$$

Proposition 3.5. *For any $P \in \mathcal{P}$ the set $C(P)$ is compact and convex.*

Proof. Recall that elements of \mathcal{P} are almost malnormal (see, for example, Farb [6, Example 1, p. 819]). Suppose $\sigma \subseteq C(P)$ is a cell and $g \in G \setminus P$. The stabilizer of $g \cdot \sigma$ is $\text{Stab}(\sigma)^g$, which intersects P in a finite group (since $\text{Stab}(\sigma) \sim P$), and hence $g \cdot \sigma \not\subseteq C(P)$.

It follows that the quotient map $q|_{C(P)} : C(P) \rightarrow X$ is finite-to-one, and since X is compact, $C(P)$ is compact. It is clear that if $x, y \in C(P)$ then $\text{Stab}(x) \sim P \sim \text{Stab}(y)$ and that if γ is the geodesic between x and y then $\text{Stab}(x) \cap \text{Stab}(y) \sim P$ and fixes γ pointwise. It follows that $\gamma \subseteq C(P)$, so $C(P)$ is convex, as required. \square

We now recall a definition and result from Haglund [11].

Definition 3.6 [11, Definition 2.14]. Let \tilde{Z} be a CAT(0) cube complex and A be a sub-complex of \tilde{Z} . The (combinatorial) convex hull of A , denoted $\text{Hull}(A)$, is the intersection of all convex sub-complexes containing A .

Given a CAT(0) cube complex \tilde{Z} and a hyperplane H in \tilde{Z} , denote $\tilde{Z} \setminus H$ the union of cubes of \tilde{Z} whose intersection with H is empty. This has two connected components, which are called (combinatorial) half-spaces. See [11, Definition 2.15] for more details. According to [11, Proposition 2.17], any convex sub-complex Y of \tilde{Z} is the intersection of the half-spaces containing Y . Hence, for any sub-complex A of \tilde{Z} , $\text{Hull}(A)$ is the intersection of the half-spaces containing A .

3.2 | Cocompact cores for relatively geometric groups

We continue to make Assumption 1.2. In this section, we prove Theorem 1.3. This result builds on [21]. By Proposition 3.1, \tilde{X} is δ -hyperbolic, and by Corollary 3.2, H acts ‘quasi-convexly’ on \tilde{X} , in the terminology of [21]. Thus, the existence of a convex subset of \tilde{X} preserved by H and finite distance from any given H -orbit follows immediately from [21, Proposition 3.3]. However, \tilde{X} is locally infinite, so it still takes work to prove that the action of H on this core is cocompact. Recall the statement of Theorem 1.3.

Theorem 1.3. *Under Assumption 1.2, for any full relatively quasi-convex subgroup H of G and any compact $K \subset \tilde{X}$ there exists a convex H -invariant sub-complex \tilde{Y} of \tilde{X} so that $K \subset \tilde{Y}$ and $H \setminus \tilde{Y}$ is compact.*

Proof. The subgroup H is finitely generated by Lemma 2.8. Let h_1, \dots, h_k generate H . By replacing K with the union of the cells whose interiors intersect K we may assume K is a finite sub-complex, and we may add finitely many 1-cells so K is connected. Since $\text{Hull}(K) = \text{Hull}(K^{(1)})$ we may replace K by its 1-skeleton. Enlarge $H \cdot K$ to a connected subset of $\tilde{X}^{(1)}$ by fixing $x \in K^{(0)}$ and choosing, for each $i \in \{1, \dots, k\}$, a geodesic γ_i between x and $h_i \cdot x$, and replacing $H \cdot K$ by $H \cdot K \cup \{H \cdot \gamma_i \mid 1 \leq i \leq k\}$. Denote this (connected, H -cocompact) sub-graph of $\tilde{X}^{(1)}$ by Γ_H .

It suffices to prove the combinatorial convex hull of Γ_H is H -cocompact. By Corollary 3.2, H -orbits in \tilde{X} are quasi-convex, so by [21, Proposition 3.3] there exists $D \geq 0$ so that $\text{Hull}(\Gamma_H) \subseteq \mathcal{N}_D(H \cdot K)$. For the following, let $Z := \text{Hull}(\Gamma_H)$, let $z \in Z^{(0)}$ and let $A = \text{Stab}_H(z)$. Let $\mathcal{E}_{z,H}$ denote the collection of edges e adjacent to z in $\tilde{X}^{(1)}$ for which the hyperplane W_e dual to e inter-

sects Γ_H non-trivially. Observe $\mathcal{E}_{z,H}$ is precisely the set of edges adjacent to z which lie in Z . We claim there are only finitely many A -orbits of edges in $\mathcal{E}_{z,H}$. If $\text{Stab}_G(z)$ is finite, then this is clear from the fact that $G \setminus \tilde{X}$ is compact. On the other hand, if A is infinite, then since H is full and the G -action on \tilde{X} is relatively geometric, A acts cocompactly on the edges adjacent to z , so again the claim is clear. If z has finite valence there is nothing to show.

The remaining case is that $\text{Stab}_G(z)$ is infinite, z has infinite valence, and A is finite. For simplicity, since we have fixed H and z , we simply write \mathcal{E} for $\mathcal{E}_{z,H}$.

Towards a contradiction, suppose \mathcal{E} is infinite. For $e \in \mathcal{E}$ with dual hyperplane W_e , let $x_e \in \Gamma_H \cap W_e$ and let $Q_e = \text{Stab}_G(W_e)$. By passing to an infinite subset of \mathcal{E} we may assume:

- (1) all elements of \mathcal{E} belong to the same $\text{Stab}_G(z)$ -orbit and
- (2) all x_e lie in the same H -orbit.

We may insist on the first point because $\text{Stab}_G(z)$ acts cocompactly on the set of edges adjacent to z , and the second because H acts cocompactly on Γ_H (and all of the intersection points x_e lie at the midpoint of some edge in $\tilde{X}^{(1)}$). We henceforth make the above two assumptions on \mathcal{E} (with extra assumptions to come, also ensured by passing to a further infinite subset).

We claim that for all $e \in \mathcal{E}$, $|Q_e \cap \text{Stab}_G(z)| < \infty$. Indeed, the $\text{Stab}_G(z)$ orbit of e is infinite, so $\text{Stab}_G(e)$ has infinite index in $\text{Stab}_G(z)$, and because the action is relatively geometric $\text{Stab}_G(e)$ is finite. Since $Q_e \cap \text{Stab}_G(z) \subseteq \text{Stab}_G(e)$, we have the required claim.

Fix $e_1 \in \mathcal{E}$, with dual hyperplane $W_1 = W_{e_1}$, and let $Q_1 = Q_{e_1} = \text{Stab}_G(W_1)$, and $x_1 = x_{e_1} \in W_1 \cap \Gamma_H$.

For each $e \in \mathcal{E}$, choose $p_e \in \text{Stab}_G(z)$ so $p_e \cdot e = e_1$, and note that $p_e \cdot W_e = W_1$. Since Q_1 acts cocompactly on W_1 , and we have $p_e \cdot x_e \in W_1$, there is an infinite subset $\mathcal{E}' \subset \mathcal{E}$ so that for all $f_1, f_2 \in \mathcal{E}'$ there exists $q_{f_1, f_2} \in Q_1$ so that $q_{f_1, f_2} p_{f_1} \cdot x_{f_1} = p_{f_2} \cdot x_{f_2}$, which is to say that $p_{f_2}^{-1} q_{f_1, f_2} p_{f_1} \cdot x_{f_1} = x_{f_2}$. We have already ensured that x_{f_1} and x_{f_2} lie in the same H -orbit, so let $h_{f_1, f_2} \in H$ be so that $h_{f_1, f_2} \cdot x_{f_1} = x_{f_2}$. We now see that $p_{f_2}^{-1} q_{f_1, f_2} p_{f_1} h_{f_1, f_2}^{-1} \in \text{Stab}_G(x_{f_2})$.

Now, x_{f_2} is the midpoint of an edge dual to the unique hyperplane W_{f_2} , so clearly $\text{Stab}_G(x_{f_2}) \leq Q_{f_2}$, so there exists $s_{f_1, f_2} \in Q_{f_2}$ so that

$$p_{f_2}^{-1} q_{f_1, f_2} p_{f_1} h_{f_1, f_2}^{-1} = (s_{f_1, f_2})^{-1}.$$

From this it follows immediately that

$$h_{f_1, f_2}^{-1} s_{f_1, f_2} p_{f_2}^{-1} q_{f_1, f_2} p_{f_1} = 1.$$

We now translate to the cusped space $\mathcal{C} = \mathcal{C}(G, \mathcal{P})$ for (G, \mathcal{P}) (see Definition 2.2). Recall from Definition 2.3 that the cusped space is ν -hyperbolic for some ν , and that it contains a copy of the Cayley graph of G . So elements of G are vertices in \mathcal{C} (the other vertices lie at some positive depth and lie in combinatorial horoballs stabilized by the conjugates of elements of \mathcal{P}).

Fix $f_2 \in \mathcal{E}'$, and let $p = p_{f_2}$. Let R_H, R_{Q_1} and $R_{Q_{f_2}}$ be the constants obtained by applying Proposition 2.10 to H, Q_1 and Q_{f_2} , respectively.

We now consider other edges $f \in \mathcal{E}'$ and make the following simplifying notational choices: $q_f = q_{f, f_2} \in Q_1, h_f = h_{f, f_2} \in H, s_f = s_{f, f_2} \in Q_{f_2}$. Therefore, the above equation becomes

$$h_f^{-1} s_f p^{-1} q_f p_f = 1.$$

We make some observations. First, $s_f \in Q_{f_2}$ and $h_f = h_{f,f_2}$ was chosen so that $h_f \cdot x_f = x_{f_2}$. Therefore, $h_f \cdot W_f = W_{f_2}$ and $h_f^{-1}s_f h_f \in Q_f$. As above, $Q_f \cap \text{Stab}_G(z)$ and $Q_1 \cap \text{Stab}_G(z)$ are both finite. Since $p_f^{-1} \in \text{Stab}_G(z)$, $Q_1^{p_f^{-1}} \cap \text{Stab}_G(z)$ is also finite. Finally note that $q_f \in Q_1$.

Now, consider a geodesic pentagon in C with vertices $1, h_f^{-1}, h_f^{-1}s_f, h_f^{-1}s_f p^{-1}, h_f^{-1}s_f p^{-1}q_f$, so the sides are labeled (in order) $h_f^{-1}, s_f, p^{-1}, q_f, p_f$. Let B_z be the horoball stabilized by $\text{Stab}_G(z)$. We claim there exists $R_C > 0$ not depending on f so that p_f penetrates B_z to a depth at most R_C .

Let α be any point on the edge labeled p_f . By subdividing the pentagon into 3 triangles and applying a standard hyperbolic geometry argument, α is at most 3ν from one of the other four sides of the geodesic pentagon. Therefore, to prove the claim, it suffices to show that each of the other four sides penetrate $\text{Stab}_G(z)$ to some bounded depth not depending on f . The length of the side labeled p does not depend on f . Therefore, the claim reduces to proving that the sides labeled h_f^{-1}, q_f and s_f penetrate B_z to a depth bounded independent of f .

We are in the case where $A = H \cap \text{Stab}_G(z)$ is finite, and the side of the pentagon labeled h_f^{-1} has endpoints in H . Therefore, this side penetrates B_z to a depth at most R_H .

Above, we observed $Q_1 \cap \text{Stab}_G(z)$ is finite, so a geodesic with endpoints q_f^{-1} and 1 penetrates B_z to a depth at most R_{Q_1} . Translating this geodesic by $p_f^{-1} = h_f^{-1}s_f p^{-1}q_f \in \text{Stab}_G(z)$ on the left implies that the side of the pentagon labeled q_f penetrates B_z to a depth at most R_{Q_1} .

Recall that $|Q_f \cap \text{Stab}_G(z)| < \infty$. Therefore, we have:

$$|Q_{f_2} \cap \text{Stab}_G(z)^{h_f}| = |Q_{f_2}^{h_f^{-1}} \cap \text{Stab}_G(z)| = |Q_f \cap \text{Stab}_G(z)| < \infty.$$

Thus, any geodesic with endpoints 1 and $s_f \in Q_{f_2}$ penetrates the horoball stabilized by $\text{Stab}_G(z)^{h_f}$ to a depth of at most $R_{Q_{f_2}}$. Translating by h_f^{-1} , the side of the pentagon labeled s_f penetrates $\text{Stab}_G(z)$ to a depth at most $R_{Q_{f_2}}$. Hence, we have proved the claim that the side of the pentagon labeled p_f penetrates B_z to a bounded depth not depending on the choice of f .

The bounded depth of p_f in B_z implies a bound on the distance (independent of f) between 1 and p_f in the naturally embedded copy of the Cayley graph of G in C . Since G is finitely generated, there are only finitely many possibilities for p_f . Recall $p_f \cdot W_f = W_1$, so we obtain a contradiction that implies \mathcal{E} is finite, as required.

To finish the proof of Theorem 1.3, we finally prove that the H -action on $Z = \text{Hull}(\Gamma_H)$ is cocompact.

For all $y \in Z^{(0)}$ we have $d_{\tilde{X}}(y, \Gamma_H) \leq D$. Let ν_y be a shortest path in $\tilde{X}^{(1)}$ from Γ_H to y so that $|\nu_y| \leq D$. Also, $H \setminus \Gamma_H$ is compact, so up to the H -action there are only finitely many possibilities for the starting point of ν_y . Also, $\nu_y \subseteq Z$, so by the claim each vertex of ν_y has only finitely many H -orbits of edges in Z adjacent to it. It now follows that the number of H -orbits of paths ν_y is finite, meaning that $H \setminus Z$ is compact, as required. \square

To ensure that these convex cores are compatible with the relatively hyperbolic geometry, we want to ensure they satisfy a condition similar to fullness.

Corollary 3.7. *Let (G, \mathcal{P}) be a group pair acting relatively geometrically on a CAT(0) cube complex \tilde{X} , let H be a full relatively quasi-convex subgroup. For any compact $K_0 \subseteq \tilde{X}$ there exists a convex H -invariant subset $\tilde{Y}_{H,K_0} \subset \tilde{X}$ so that (i) $K_0 \subseteq \tilde{Y}_{H,K_0}$, (ii) for each $P \in \mathcal{P}$ we have $C(P) \subset \tilde{Y}_{H,K_0}$ and (iii) $H \setminus \tilde{Y}_{H,K_0}$ is compact. Moreover, the H -action on \tilde{Y}_{H,K_0} is relatively geometric.*

Proof. There are finitely many $P \in \mathcal{P}$ and by Proposition 3.5 each $C(P)$ is compact, so we can choose $K = K_0 \cup \bigcup_{P \in \mathcal{P}} C(P)$ and apply Theorem 1.3. That the H -action is relatively geometric follows immediately from the following facts: (i) the G -action on \tilde{X} is relatively geometric, (ii) H is full relatively quasi-convex and (iii) $H \setminus \tilde{Y}_{H, K_0}$ is compact. \square

4 | DEHN FILLING AND CUBE COMPLEXES

In previous papers such as [1, 5, 8, 22] the combination of (relatively) hyperbolic groups acting on CAT(0) cube complexes and the behavior under Dehn filling has yielded very powerful tools. We continue in this theme, in the context of relatively geometric actions.

The following result is [5, Proposition 2.3], and is an immediate consequence of [8, Corollary 6.6]. It follows immediately from the definition of relatively geometric that there exists a family \mathcal{Q} as in the statement below.

Proposition 4.1 [5, Proposition 2.3]. *Suppose (G, \mathcal{P}) is relatively hyperbolic and that G admits a relatively geometric action on a CAT(0) cube complex \tilde{X} . Let \mathcal{Q} be a collection of finite-index subgroups of elements of \mathcal{P} so that any infinite cell stabilizer contains a conjugate of an element of \mathcal{Q} . For sufficiently long \mathcal{Q} -fillings*

$$G \rightarrow \overline{G} = G/K$$

of (G, \mathcal{P}) , the quotient $K \setminus \tilde{X}$ is a CAT(0) cube complex.

Corollary 4.2. *Let (G, \mathcal{P}) and \mathcal{Q} be as in Proposition 4.1. For sufficiently long \mathcal{Q} -fillings $G \rightarrow \overline{G} = G(N_P \mid P \in \mathcal{P})$, where each P/N_P is virtually special and hyperbolic, the group \overline{G} is virtually special and hyperbolic.*

In particular, for sufficiently long peripherally finite \mathcal{Q} -fillings, \overline{G} is hyperbolic and virtually special.[†]

Proof. By Theorem 2.11, for sufficiently long fillings $G \rightarrow \overline{G} = G(N_P \mid P \in \mathcal{P})$ the natural map $P/N_P \rightarrow \overline{G}$ is an embedding for each $P \in \mathcal{P}$ and the pair $(\overline{G}, \{P/N_P \mid P \in \mathcal{P}\})$ is relatively hyperbolic. Thus, if each P/N_P is hyperbolic then \overline{G} itself is a hyperbolic group. Since $(\overline{G}, \{P/N_P\})$ is relatively hyperbolic, the subgroups P/N_P of \overline{G} are quasi-convex.

By Proposition 4.1 for sufficiently long \mathcal{Q} -fillings the space $\overline{X} := K \setminus \tilde{X}$ is a CAT(0) cube complex, and $\overline{G} = G/K$ acts on \overline{X} . The quotient $\overline{G} \setminus \overline{X}$ has the same underlying space as $G \setminus \tilde{X}$, so it is compact.

The cell stabilizers for the \overline{G} -action on \overline{X} are finite-index subgroups of the parabolic subgroups, and so these cell stabilizers are also quasi-convex in \overline{G} . Therefore, if each P/N_P is hyperbolic and virtually special then \overline{G} is virtually special by [8, Theorem D]. \square

We now prove Theorem 1.4 from the introduction. Recall the statement.

[†]Recall a group is *virtually special* if it has a finite-index subgroup which admits a proper and cocompact action on a CAT(0) cube complex with quotient a special cube complex.

Theorem 1.4. *Under Assumption 1.2, if the elements of \mathcal{P} are residually finite then there is a finite-index subgroup G_0 of G so that $G_0 \setminus \tilde{X}$ is a special cube complex.*

Proof. Because the elements of \mathcal{P} are residually finite, by Proposition 4.1 and Corollary 4.2 there is a peripherally finite filling $G \rightarrow \bar{G} = G/K$ so that \bar{G} is virtually special and $\bar{X} = K \setminus \tilde{X}$ is a CAT(0) cube complex. Thus, there is a finite-index subgroup $\bar{G}_0 \leq \bar{G}$ so that $\bar{G}_0 \setminus \bar{X}$ is a special cube complex. Let G_0 be the (finite-index) pre-image of \bar{G}_0 in G , and observe that the underlying space of $G_0 \setminus \tilde{X}$ is the same as the underlying space of $\bar{G}_0 \setminus \bar{X}$. \square

4.1 | Cores map to cores under suitable fillings

Suppose (G, \mathcal{P}) acts relatively geometrically on the CAT(0) cube complex \tilde{X} with residually finite peripherals. Let Q be the set of subgroups from Proposition 4.1, so that for sufficiently long Q -fillings $G \rightarrow G/K$ the space $K \setminus \tilde{X}$ is a CAT(0) cube complex. Let H be a full relatively quasi-convex subgroup of G and let $\tilde{Y} \subset \tilde{X}$ be a convex H -invariant and H -cocompact sub-complex, the existence of which is guaranteed by Theorem 1.3.

The following summarizes a collection of known and straightforward results about the existence of certain well-controlled Dehn fillings.

Proposition 4.3. *For sufficiently long $(Q \cup \{H\})$ -fillings $G \rightarrow G/K$, the following statements hold:*

- (1) *The induced maps from each P/N_P to G/K are injective, and if \bar{P} is the image of P/N_P , then $(G/K, \{\bar{P} \mid P \in \mathcal{P}\})$ is relatively hyperbolic;*
- (2) *If $K_H \trianglelefteq H$ is the kernel of the induced filling of H , then $K_H = K \cap H$;*
- (3) *If \bar{H} denotes the image of H in G/K , and $\bar{\mathcal{D}}$ is the collection of images of elements of \mathcal{D} , then $(\bar{H}, \bar{\mathcal{D}})$ is relatively quasi-convex in $(G/K, \{\bar{P}\})$;*
- (4) *$\bar{X} = K \setminus \tilde{X}$ is a CAT(0) cube complex;*
- (5) *$\bar{Y} = K_H \setminus \tilde{Y}$ is a CAT(0) cube complex;*
- (6) *If Q_1 and Q_2 are distinct maximal parabolic subgroups, then $K \cap Q_1 \cap Q_2 = \{1\}$;*
- (7) *If S is a finite cell stabilizer, then $S \cap K = \emptyset$;*
- (8) *The induced map $\bar{Y} \rightarrow \bar{X}$ is an immersion.*

Proof. Each of the items can be shown to hold for sufficiently long $(Q \cup \{H\})$ -fillings, and then we can take a single filling to satisfy them all. Thus, we explain how to ensure each of the items individually. Item 1 follows from Theorem 2.11. Items 2 and 3 follow from Theorem 2.13. Item 4 follows from Proposition 4.1. For Item 5, note that the H -action on \tilde{Y} is relatively geometric by Corollary 3.7, so this item also follows from Proposition 4.1.

Consider Item 6. By [10, Lemma 2.6] there exists $M > 0$ so that if Q_1, Q_2 are maximal parabolics, then $Q_1 \cap Q_2$ acts freely on a subset of the Cayley graph of G of diameter at most M . Up to the action of G there are only finitely many such sub-graphs of the Cayley graph, so there are only finitely many G -conjugacy classes of intersections of maximal parabolic subgroups $Q_1 \cap Q_2$. Recall that $|Q_1 \cap Q_2| < \infty$. Therefore, ensuring Item 6 involves excluding finitely many elements from the filling kernels, so this item holds for sufficiently long fillings. For Item 7, there are finitely

many conjugacy classes of finite stabilizers by cocompactness of the action. Item 7 then follows by again excluding finitely many elements from the filling kernels.

Finally, we prove that we can ensure Item 8. Consider the set of cells $\tau \in \tilde{Y}$ so that $\text{Stab}_G(\tau)$ is infinite, but $\text{Stab}_H(\tau)$ is finite. Since $H \setminus \tilde{Y}$ is compact, there are only finitely many cells ρ_1, \dots, ρ_k in \tilde{Y} which contain τ . For each such distinct pair ρ_i, ρ_j so that $\text{Stab}_G(\rho_i)$ is finite, let $\mathcal{F}_{i,j}$ be the (possibly empty) finite set of elements $g \in \text{Stab}_G(\tau)$ so that $g \cdot \rho_i = \rho_j$. There are only finitely many H -orbits of cells in \tilde{Y} , so by Theorem 2.11 for sufficiently long fillings $G \rightarrow G/K$ we have $\mathcal{F}_{i,j} \cap K = \emptyset$ for all such i, j (and τ). Now suppose that $G \rightarrow G/K$ is such a $(\mathcal{Q} \cup \{H\})$ -filling so that also Items 2, 4 and 5 hold. Further, by taking a longer filling if necessary, we suppose that for any cell κ in X so that $\text{Stab}_G(\kappa)$ is finite we have $K \cap \text{Stab}_G(\kappa) = \emptyset$. In order to obtain a contradiction, suppose that $\bar{Y} \rightarrow \bar{X}$ is not an immersion and let $\bar{\sigma}_1$ and $\bar{\sigma}_2$ be adjacent (distinct) cells in \bar{Y} with the same image in \bar{X} . Note that $\bar{\sigma}_1 \cap \bar{\sigma}_2$ is a cell in \bar{Y} . We may lift to cells σ_1, σ_2 in \tilde{Y} with $\tau = \sigma_1 \cap \sigma_2$ a cell in \tilde{Y} . Since the images of $\bar{\sigma}_1$ and $\bar{\sigma}_2$ are equal in \bar{X} there exists $k \in K$ so that $k \cdot \sigma_1 = \sigma_2$. Since \bar{X} is a CAT(0) cube complex, we have $k \cdot \tau = \tau$, so $k \in \text{Stab}_G(\tau)$. If $\text{Stab}_G(\tau)$ is finite, then we have $K \cap \text{Stab}_G(\tau) = \emptyset$, so there is no such k . Therefore, we may assume that $\text{Stab}_G(\tau)$ is infinite. If $\text{Stab}_H(\tau)$ is also infinite, then because we have a relatively geometric action, and the filling is an H -filling, we have $K \cap \text{Stab}_G(\tau) \leq H$, which means that $\bar{\sigma}_1 = \bar{\sigma}_2$ in \bar{Y} , contrary to our choice.

We are left with the possibility that $\text{Stab}_G(\tau)$ is infinite, but $\text{Stab}_H(\tau)$ is finite. If $\text{Stab}_G(\sigma_i)$ is infinite, it is commensurable into $\text{Stab}_G(\tau)$. Then, since the filling is a \mathcal{Q} -filling, $K \cap \text{Stab}_G(\tau) \leq \text{Stab}_G(\sigma_i)$, which contradicts the equation $k \cdot \sigma_1 = \sigma_2$. Finally, suppose that $\text{Stab}_G(\sigma_1)$ and $\text{Stab}_G(\sigma_2)$ are both finite, but that $\text{Stab}_G(\tau)$ is infinite (and $\text{Stab}_H(\tau)$ is still finite). In this case, we have $\sigma_1 = \rho_i$ and $\sigma_2 = \rho_j$ for some i and j , where the ρ_i and ρ_j are as chosen above. If $\text{Stab}_G(\rho_i)$ is infinite and we clearly have $k \in \mathcal{F}_{i,j}$, contradicting the assumption about our filling that $\mathcal{F}_{i,j} \cap K = \emptyset$. This completes the proof. \square

We now prove the main result of this section, which immediately implies Theorem 1.5 from the introduction.

Theorem 4.4. *For all sufficiently long $(\mathcal{Q} \cup \{H\})$ -fillings $G \rightarrow \bar{G}$, the immersion $f : \bar{Y} \rightarrow \bar{X}$ is an embedding, and (the image of) \bar{Y} is convex in \bar{X} .*

Proof. Let \mathcal{Q} be the set of subgroups as in Proposition 4.1. Let \mathcal{D} be the induced peripheral structure for H . Let $\mathcal{Q}_H = \mathcal{Q} \cup \{H\}$.

We only consider \mathcal{Q}_H -fillings which are long enough to satisfy the conclusions of Proposition 4.3. We impose a further condition below so that if both conditions are simultaneously satisfied then \bar{Y} is convex in \bar{X} .

For a cell $\sigma \subseteq \tilde{X}$, denote the G -orbit of σ by $[\![\sigma]\!]$. Moreover, for a cell $\tau \subseteq \tilde{Y}$, let $[\![\tau]\!]_H$ denote the H -orbit of τ . Consider the set \mathcal{S} of all pairs $([\![e_1]\!]_H, [\![e_2]\!]_H)$, where e_1 and e_2 are (oriented) edges in \tilde{Y} so that there exist $e'_1 \in [\![e_1]\!]_H$ and $e'_2 \in [\![e_2]\!]_H$ so that e'_1 and e'_2 have the same initial vertex, and so that e'_1 and e'_2 bound the corner of a square in \tilde{X} . Since $H \setminus \tilde{Y}$ is compact, \mathcal{S} is finite. By rechoosing e_2 if necessary, we always assume that if $([\![e_1]\!]_H, [\![e_2]\!]_H) \in \mathcal{S}$ then the edges e_1, e_2 in \tilde{Y} share the same initial vertex.

Let \mathcal{S}' denote the set of all $\rho = ([\![e_1]\!]_H, [\![e_2]\!]_H) \in \mathcal{S}$ so that

- (1) e_1 and e_2 do not bound a square in \tilde{Y} and
- (2) $\text{Stab}_G(e_1)$ and $\text{Stab}_G(e_2)$ are finite.

Let $\rho = ([\![e_1]\!]_H, [\![e_2]\!]_H) \in S'$. Since \tilde{Y} is convex in \tilde{X} , e_1 and e_2 do not bound a square in \tilde{X} either. Furthermore, since $\text{Stab}_G(e_1)$ is finite and because the action of G on \tilde{X} is cocompact, there are only finitely many squares f_1, \dots, f_k adjacent to e_1 in \tilde{X} . For each f_i , let \hat{e}_2^i be the edge of f_i which shares the initial vertex of e_1 .

Since $\text{Stab}_G(e_2)$ is finite there are only finitely many elements $g \in G$ so that for some $1 \leq i \leq n$ we have $g \cdot \hat{e}_2^i = e_2$. Let \mathcal{F}_ρ denote the set of all such g , and let \mathcal{F} be the union of the \mathcal{F}_ρ over all $\rho \in S'$. Note that each \mathcal{F}_ρ is finite and $S' \subseteq S$ is finite, so \mathcal{F} is finite.

By Theorem 2.11 for sufficiently long fillings $G \rightarrow G/K$, $K \cap \mathcal{F} = \emptyset$. Fix now a Q_H -filling $G \rightarrow G/K$ long enough to satisfy the conclusion of Proposition 4.3, and also so that $K \cap \mathcal{F} = \emptyset$.

We claim that with such a filling, and the notation as above, the subspace \bar{Y} is convex in \bar{X} . In order to obtain a contradiction, suppose that there are edges $\bar{e}_1, \bar{e}_2 \in \bar{Y}$ so that \bar{e}_1 and \bar{e}_2 do not bound a square in \bar{Y} but they do bound a square \bar{f} in \bar{X} .

Lift \bar{e}_1 to an edge e_1 in \tilde{Y} , and \bar{e}_2 to an edge e_2 with the same initial vertex, x say, as e_1 . Let f be a lift of \bar{f} to \tilde{X} so that e_1 is an edge on the boundary of f . Since $\bar{f} \notin \bar{Y}$, we see that $f \notin \tilde{Y}$. Moreover, there is no square in \tilde{X} with e_1 and e_2 at a corner, because \tilde{Y} is convex in \tilde{X} .

Let \hat{e}_2 be the edge on the boundary of f with initial point x . Because the images of \hat{e}_2 and e_2 are both \bar{e}_2 in \bar{X} , there exists $k \in K \cap \text{Stab}_G(x)$ so $k \cdot \hat{e}_2 = e_2$.

First suppose that $\text{Stab}_G(e_1)$ is infinite. Since the G -action on \tilde{X} is relatively geometric, $\text{Stab}_G(e_1)$ is finite-index in $\text{Stab}_G(x)$, and since $G \rightarrow G/K$ is a Q_H -filling, $\text{Stab}_G(x) \cap K \leq \text{Stab}_G(e_1)$. Therefore, $k \cdot e_1 = e_1$, so $k \cdot f$ is a 2-cell which has e_1 and e_2 as a corner. This is a contradiction, so $\text{Stab}_G(e_1)$ is finite.

Now suppose that $\text{Stab}_G(e_2)$ is infinite. In this case, $K \cap \text{Stab}_G(x) \leq \text{Stab}_G(e_2)$, which contradicts the equation $k \cdot \hat{e}_2 = e_2$. Therefore, $\text{Stab}_G(e_2)$ is also finite.

Using $[\![\cdot]\!]$ to denote orbits as above, $\rho = ([\![e_1]\!]_H, [\![e_2]\!]_H) \in S'$. Then $k \in \mathcal{F}_\rho \subseteq \mathcal{F}$, contradicting $K \cap \mathcal{F} = \emptyset$. This contradiction proves the immersion $\bar{Y} \rightarrow \bar{X}$ is locally convex. Since \bar{X} is a CAT(0) cube complex, a locally convex immersion is an embedding with convex image, completing the proof. \square

5 | COMPLETION AND RETRACTION WITH COMPLEXES OF GROUPS

In this section we prove Theorem 1.6. Our approach is to prove a relatively geometric analogue of the canonical completion and retraction due to Haglund and Wise [12] (see Theorem 5.9 below). We prove this by applying a Dehn filling as in Proposition 4.3 and Theorem 4.4, applying Agol's Theorem [1, Theorem 1.1], passing to a carefully chosen finite-index subgroup, applying the Haglund–Wise construction, and then noting that the retraction of cube complexes induces a retraction of complexes of groups. For the basic theory of complexes of groups, we refer to [3, III.C].

Remark 5.1. The completion and retraction we construct in Theorem 5.9 below relies on a particular Dehn filling, and so is not as ‘canonical’ as that of Haglund–Wise.

5.1 | The canonical completion for relatively geometric complexes of groups

The following result summarizes Haglund and Wise's construction of the canonical completion and retraction for special cube complexes.

Theorem 5.2 [12, § 6], [13, § 3]. Suppose that A and B are cube complexes, that B is special, that A is compact and that $f : A \rightarrow B$ is a locally convex combinatorial map. There exists a pair of cube complexes $\mathbf{C}(A, B)$ and $\mathbf{C}_\square(A, B)$, along with

- (1) a homeomorphism $s : \mathbf{C}_\square(A, B) \rightarrow \mathbf{C}(A, B)$,
- (2) a finite (combinatorial) covering $p : \mathbf{C}(A, B) \rightarrow B$,
- (3) a (combinatorial) embedding $i : A \rightarrow \mathbf{C}_\square(A, B)$ so that soi is a (combinatorial) embedding, and $p \circ soi = f$,
- (4) a cellular retraction $r : \mathbf{C}_\square(A, B) \rightarrow A$ (so $r \circ i = \text{Id}_A$).

The following diagram commutes:

$$\begin{array}{ccc} \mathbf{C}_\square(A, B) & \xrightarrow{s} & \mathbf{C}(A, B) \\ r \swarrow \quad \nearrow i & & \downarrow p \\ A & \xrightarrow{f} & B \end{array}$$

As described in [13, Definition 3.5], $\mathbf{C}_\square(A, B)$ is obtained from $\mathbf{C}(A, B)$ by sub-dividing certain cubes (some of those outside of the image of A), and the map r maps each cube onto a face of a cube in the target by orthogonal projection.

Our goal is to set up a situation of complexes of groups so that the underlying spaces are arranged in a diagram as above. We then explain how to turn the corresponding maps into morphisms of complexes of groups, giving Theorem 1.6. We now record the set-up to our construction in the following assumption, which builds on Assumption 1.2.

Assumption 5.3. Suppose that (G, \mathcal{P}) is relatively hyperbolic, that \tilde{X} is a CAT(0) cube complex which admits a relatively geometric action of G with respect to \mathcal{P} , and let $X = G \setminus \tilde{X}$. Further, suppose all elements of \mathcal{P} are residually finite.

Let H be a full relatively quasi-convex subgroup, and let (H, D) be the peripheral structure on H induced from (G, \mathcal{P}) . If $D \in \mathcal{D}$ and $D \leq P_D^{c_D}$ for some $c_D \in G$ and $P_D \in \mathcal{P}$, let $C(P_D)$ be the sub-complex of \tilde{X} associated to P_D from Definition 3.4. Let $\tilde{Y} \subset \tilde{X}$ be a convex H -invariant and H -cocompact sub-complex as in the conclusion of Theorem 1.3 so that for each $D \in \mathcal{D}$ we have $c_D \cdot C(P_D) \subset \tilde{Y}$ (this can be ensured by Proposition 3.5 and Theorem 1.3).

Let Q be a collection of subgroups as in the hypotheses of Proposition 4.1.

Let $\pi : G \rightarrow \bar{G} = G(N_P \mid P \in \mathcal{P}) = G/K$ be a peripherally finite $(Q \cup \{H\})$ -filling which satisfies the conclusions of Proposition 4.3 and Theorem 4.4.

Let \bar{H} be the image of H in \bar{G} , and let $\bar{X} = K \setminus \tilde{X}$, $\bar{Y} = K_H \setminus \tilde{Y}$ be as in Proposition 4.3.

For the remainder of this section, we make Assumption 5.3.

Proposition 5.4. The group \bar{G} is hyperbolic and virtually special. In particular, it is residually finite and virtually torsion-free. Moreover, there is a finite-index torsion-free subgroup $\bar{G}_0 \leq \bar{G}$ so that $\bar{G}_0 \setminus \bar{X}$ is a special cube complex.

Proof. Since the filling $G \rightarrow \bar{G}$ is peripherally finite, \bar{G} is hyperbolic relative to finite groups, and hence is hyperbolic. Moreover, \bar{G} acts cocompactly on the CAT(0) cube complex \bar{X} (since $\bar{G} \setminus \bar{X}$

and $G \setminus \tilde{X}$ have the same underlying space). Because the G -action on \tilde{X} is relatively geometric, it follows that stabilizers for the \overline{G} -action on \overline{X} are finite. Thus, the hyperbolic group \overline{G} acts properly and cocompactly on the CAT(0) cube complex \overline{X} . By Agol's Theorem, \overline{G} is virtually special, and hence residually finite. It is well known that any residually finite hyperbolic group is virtually torsion-free, so there is a torsion-free finite-index subgroup $\overline{G}_0 \leq \overline{G}$ so that $\overline{G}_0 \setminus \overline{X}$ is a special cube complex, as required. \square

Let $B = \overline{G}_0 \setminus \overline{X}$. Since \overline{G}_0 is torsion-free, $\overline{G}_0 = \pi_1(B)$. Define $\overline{H}_0 = \overline{H} \cap \overline{G}_0$, and note that \overline{H}_0 is torsion-free, so acts freely on \overline{Y} . Let $A = \overline{H}_0 \setminus \overline{Y}$. The convex embedding $\overline{Y} \rightarrow \overline{X}$ from the conclusion of Theorem 4.4 descends to a locally convex immersion $f : A \rightarrow B$. Since B is special, Theorem 5.2 applies to the map $f : A \rightarrow B$, and we obtain the canonical completion $\mathbf{C}(A, B)$, and its subdivided version $\mathbf{C}_{\square}(A, B)$ as in Theorem 5.2. Let \overline{G}_1 be the finite-index subgroup of \overline{G}_0 corresponding to the finite cover $\mathbf{C}(A, B) \rightarrow B$. By the construction of the canonical completion $\overline{H}_0 \leq \overline{G}_1$.

Let $C = \mathbf{C}_{\square}(A, B)$, so $\overline{G}_1 = \pi_1(C)$ (recall $\mathbf{C}_{\square}(A, B)$ is homeomorphic to $\mathbf{C}(A, B)$). Let \tilde{C} be the CAT(0) universal cover of C . Since $A \rightarrow C$ is an inclusion as a sub-complex, \tilde{Y} is a convex sub-complex of \tilde{C} .

Let \tilde{C} be the induced sub-divided version of \tilde{X} , and let $G_1 = \pi^{-1}(\overline{G}_1)$, a finite-index subgroup of G_0 , and $H_0 = H \cap G_1$, a finite-index subgroup of H so $\pi(H_0) = \overline{H}_0$. The universal cover \tilde{C} of C is a CAT(0) cube complex which is a sub-divided version of \tilde{X} .

There is a G_1 -action on \tilde{C} and an H_0 -action on \tilde{Y} , so that the underlying space of $G_1 \setminus \tilde{C}$ is C and the underlying space of $H_0 \setminus \tilde{Y}$ is A . Observe the G_1 -action on \tilde{C} is relatively geometric, with respect to the induced peripheral structure (G_1, \mathcal{P}_1) on G_1 .

Proposition 5.5. *The following properties for the G_1 -action on \tilde{C} and the H_0 -action on \tilde{Y} hold:*

- (1) *Stabilizers in G_1 of cells in \tilde{C} are either trivial or else maximal parabolic subgroups of G_1 .*
- (2) *If $P_1 \in \mathcal{P}_1$ then the sub-complex $C(P_1)$ from Definition 3.4 has cells whose stabilizers are exactly P_1 , and $C(P_1)$ embeds in C under the quotient map.*
- (3) *If $\sigma \in \tilde{Y}$ is a cell with non-trivial H_0 -stabilizer then $\text{Stab}_{H_0}(\sigma) = \text{Stab}_{G_1}(\sigma)$.*

Proof. Stabilizers are already finite or finite-index in a maximal parabolic because the action is relatively geometric. Since \overline{G}_1 is torsion free and $\pi : G \rightarrow \overline{G}$ is injective on finite stabilizers by Proposition 4.3.(7), G_1 has no non-trivial finite cell stabilizers. The filling $\pi : G \rightarrow \overline{G}$ is a Q -filling and π is injective on the finite groups $\{P/N_P\}$, so because \overline{G}_1 is torsion free any infinite stabilizer must be maximal parabolic in G_1 . This proves (1).

Item (2) follows because all non-trivial stabilizers are maximal parabolic in G_1 and maximal parabolic subgroups are malnormal.

For Item 3, suppose $\sigma \in \tilde{Y}$ has $\text{Stab}_{H_0}(\sigma) \neq \{1\}$. By Proposition 4.3.(7) the map $H \rightarrow \overline{H}$ is injective on finite cell stabilizers. Moreover, \overline{H}_0 is torsion-free, and \overline{H}_0 is the induced filling of H_0 . Therefore, $\text{Stab}_{H_0}(\sigma)$ is infinite, and hence full parabolic. Let Q be the maximal parabolic subgroup of G_1 containing $\text{Stab}_{H_0}(\sigma)$. Then $Q = \text{Stab}_{G_1}(\sigma)$ by Item 1. Since \overline{G}_1 is torsion-free and π is an H -filling, $\text{Stab}_{H_0}(\sigma) = Q = \text{Stab}_{G_1}(\sigma)$, as required. \square

The actions of G_1 on \tilde{C} and H_0 on \tilde{Y} give rise to a pair of complexes of groups $G(\mathcal{C})$ and $H(\mathcal{A})$, with underlying scwols \mathcal{C} arising from C and \mathcal{A} arising from A , respectively. The map $i: A \rightarrow C$ gives rise to a (non-degenerate) morphism of scwols $f_i: \mathcal{A} \rightarrow \mathcal{C}$ as in [3, III.C.1.5]. A complex of groups comes with a collection of data, one part of which is ‘twisting elements’ (see [3, Definition III.C.2.1]). A complex of groups is *simple* if all the twisting elements are trivial (see [3, III.C.2.1]). To build the complexes of groups $G(\mathcal{C})$ and $H(\mathcal{A})$, follow the construction from [3, § III.C.2.9]. This construction involves some choices (of the lifts of objects, and of the elements h_a). However, we make the following observation.

Lemma 5.6. *We may make choices in the constructions of $G(\mathcal{C})$ and $H(\mathcal{A})$ so that both are simple complexes of groups.*

Proof. Suppose σ is an object in \mathcal{C} whose stabilizer is non-trivial, then we lift to (the scwol \mathcal{X} associated to) X and obtain a non-trivial cell stabilizer in G_1 . By Proposition 5.5.(2) the set of objects in \mathcal{C} whose stabilizers intersect $\text{Stab}(\sigma)$ non-trivially can be simultaneously lifted to \mathcal{X} when defining $G(\mathcal{C})$, so for these cells the twisting elements can be chosen to be trivial. For other cells, the stabilizer is trivial, so twisting elements are trivial. The proof for $H(\mathcal{A})$ is the same. \square

By [3, Corollary III.C.2.18], the H_0 -equivariant inclusion $\tilde{Y} \rightarrow \tilde{X}$ induces a morphism of complexes of groups $\phi: H(\mathcal{A}) \rightarrow G(\mathcal{C})$ over f_i . We can, and do, consider the morphism f_i to be inclusion, so that objects and arrows of \mathcal{A} are contained in \mathcal{C} . By choosing lifts of \mathcal{A} before lifts of \mathcal{C} when defining the complex of groups structures, we can ensure that if σ is an object of \mathcal{A} , and H_σ is non-trivial, then $H_\sigma = G_\sigma$, and the map $\phi_\sigma: H_\sigma \rightarrow G_\sigma$ is the identity.

Now, let $r: C \rightarrow A$ be the canonical retraction from Theorem 5.2, and let $f_r: \mathcal{C} \rightarrow \mathcal{A}$ be the associated morphism of scwols (since cells in C may be collapsed under r , the morphism f_r is probably degenerate).

We now define a morphism of complexes of groups $\theta: G(\mathcal{C}) \rightarrow H(\mathcal{A})$ over f_r . Let σ be an object of \mathcal{C} , and consider the object $f_r(\sigma) \in \mathcal{A}$.

Lemma 5.7. *Either $H_{f_r(\sigma)} = G_\sigma$ or else $H_{f_r(\sigma)} \cap G_\sigma = \{1\}$.*

Proof. The subgroup $H_{f_r(\sigma)}$ is a cell stabilizer for the action of H_0 on \tilde{Y} , so if $H_{f_r(\sigma)} \neq \{1\}$ then by Proposition 5.5.(3) $H_{f_r(\sigma)}$ is a cell stabilizer for the action of G_1 on \tilde{C} . Cell stabilizers for the action of G_1 on \tilde{C} are either trivial or maximal parabolic. Therefore, either $G_\sigma = H_{f_r(\sigma)}$ or they are distinct cell stabilizers in G_1 and have trivial intersection by Proposition 4.3.(6) and the construction of G_1 . \square

In case $H_{f_r(\sigma)} = G_\sigma$ define θ_σ to be the identity map, and in case $H_{f_r(\sigma)} \cap G_\sigma = \{1\}$ define θ_σ to be the trivial map. For each arrow $a \in \mathcal{C}$, define the element $\theta(a)$ to be the identity element of $H_{t(f_r(a))}$.

These data define the structure of a morphism of complexes of groups θ , as follows from the next result (where $\psi_{f_r(a)}$ and ψ_a are the homomorphisms in the complexes of groups $H(\mathcal{A})$ and $G(\mathcal{C})$, respectively).

Lemma 5.8. *For each arrow a of \mathcal{C} we have*

$$\psi_{f_r(a)} \theta_{i(a)} = \theta_{t(a)} \psi_a.$$

Proof. Suppose first that $G_{t(a)} \neq H_{t(f_r(a))}$. Then $G_{t(a)} \cap H_{t(f_r(a))} = \{1\}$ by Lemma 5.7. If $\theta_{i(a)}$ is non-trivial, then $G_{i(a)} = H_{i(f_r(a))}$. The local maps $\psi_a, \psi_{f_r(a)}$ are inclusions, so $G_{t(a)} \cap H_{t(f_r(a))}$ is non-trivial, a contradiction. Therefore, $\theta_{i(a)}$ must be trivial, and the lemma holds in this case.

On the other hand, suppose that $G_{t(a)} = H_{t(f_r(a))}$. Then $\theta_{t(a)}$ is the identity map. Non-trivial cell stabilizers in G_1 are maximal parabolic by Proposition 5.5.(1) and intersections of maximal parabolics are trivial by Proposition 4.3.(6). Therefore, $G_{i(a)}$ is trivial or $G_{i(a)} = G_{t(a)}$. In the first case, $\theta_{i(a)}$ is the trivial map and the lemma follows, so suppose $G_{i(a)} = G_{t(a)}$, a maximal parabolic subgroup. Note that $\theta_{i(a)}\psi_a(G_{i(a)}) = G_{i(a)}$. Also, we have (in this case) $G_{i(a)} = G_{t(a)} = H_{t(f_r(a))}$. Moreover, $H_{t(f_r(a))}$ is some maximal parabolic in H_0 , which in turn is finite-index in a maximal parabolic in H . Because we chose \tilde{Y} to contain all of the $c_D \cdot C(P_D)$ in Assumption 5.3, and because \tilde{Y} is H -invariant, we see that $C(G_{i(a)}) \subseteq \tilde{Y}$. Then $i(a)$ is a cell in the image of $H_0 \setminus \tilde{Y} \subseteq G_1 \setminus \tilde{C}$ that is fixed by f_r . Therefore, $\theta_{i(a)}$ is the identity map, and the lemma follows. \square

It follows immediately from the construction that $\theta \circ \phi = \text{Id}_{H(\mathcal{A})}$.

The induced map on π_1 from ϕ is the inclusion $\iota: H_0 \rightarrow G_1$. Moreover, if $\rho: G_1 \rightarrow H_0$ is $\rho = \pi_1(\theta)$ then the fact that $\theta \circ \phi = \text{Id}_{H(\mathcal{A})}$ implies $\rho \circ \iota = \text{Id}_{H_0}$. This proves Theorem 1.6 from the introduction.

For future use, we summarize the above construction in the following result (in the statement below, we elide the difference between a quotient space and the induced complex of groups).

Theorem 5.9. *Make Assumption 1.2, and suppose further that elements of \mathcal{P} are residually finite. Let H be a full relatively quasiconvex subgroup of G . There exist*

- a cocompact convex core $\tilde{Y} \subseteq \tilde{X}$ for H ;
- finite index subgroups $H_0 \leq H$ and $G_1 \leq G$, so $H_0 \leq G_1$; and
- a subdivision \tilde{C} of \tilde{X} with an embedding $\tilde{Y} \hookrightarrow \tilde{C}$;

together with morphisms of complexes of groups:

$$\phi: H_0 \setminus \tilde{Y} \rightarrow G_1 \setminus \tilde{C} \quad \text{and} \quad \theta: G_1 \setminus \tilde{C} \rightarrow H_0 \setminus \tilde{Y}$$

so that

- the underlying map of ϕ is an embedding of special cube complexes;
- for each cell σ of $H_0 \setminus \tilde{Y}$, either the local group H_σ is trivial or H_σ and the local group $G_{\phi(\sigma)}$ are equal and the map $\phi_\sigma: H_\sigma \rightarrow G_{\phi(\sigma)}$ is the identity map;
- $\theta \circ \phi$ is the identity morphism on $H_0 \setminus \tilde{Y}$.

ACKNOWLEDGEMENTS

The first author is supported in part by an AMS-Simons Travel Grant. The second author is supported in part by NSF grant DMS-1904913.

JOURNAL INFORMATION

The *Journal of the London Mathematical Society* is wholly owned and managed by the London Mathematical Society, a not-for-profit Charity registered with the UK Charity Commission. All surplus income from its publishing programme is used to support mathematicians and mathematics research in the form of research grants, conference grants, prizes, initiatives for early career researchers and the promotion of mathematics.

REFERENCES

1. I. Agol, *The virtual Haken conjecture*, Doc. Math. **18** (2013), 1045–1087. With an appendix by Agol, Daniel Groves, and Jason F. Manning.
2. I. Agol, D. Groves, and J. F. Manning, *Residual finiteness, QCERF and fillings of hyperbolic groups*, Geom. Topol. **13** (2009), no. 2, 1043–1073.
3. M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, volume 319 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer, Berlin, 1999.
4. R. Charney and J. Crisp, *Relative hyperbolicity and Artin groups*, Geom. Dedicata **129** (2007), 1–13.
5. E. Einstein and D. Groves, *Relative cubulations and groups with a 2-sphere boundary*, Compos. Math. **156** (2020), no. 4, 862–867.
6. B. Farb, *Relatively hyperbolic groups*, Geom. Funct. Anal. **8** (1998), no. 5, 810–840.
7. D. Groves and J. F. Manning, *Dehn filling in relatively hyperbolic groups*, Israel J. Math. **168** (2008), 317–429.
8. D. Groves and J. F. Manning, *Hyperbolic groups acting improperly*, arxiv.org:1808.02325, 2018.
9. D. Groves and J. F. Manning, *Specializing cubulated relatively hyperbolic groups*, J. Topol., to appear.
10. D. Groves and J. F. Manning, *Quasiconvexity and Dehn filling*, Amer. J. Math. **143** (2021), no. 1, 95–124.
11. F. Haglund, *Finite index subgroups of graph products*, Geom. Dedicata **135** (2008), 167–209.
12. F. Haglund and D. T. Wise, *Special cube complexes*, Geom. Funct. Anal. **17** (2008), no. 5, 1551–1620.
13. F. Haglund and D. T. Wise, *A combination theorem for special cube complexes*, Ann. of Math. (2) **176** (2012), no. 3, 1427–1482.
14. G. C. Hruska, *Relative hyperbolicity and relative quasiconvexity for countable groups*, Algebr. Geom. Topol. **10** (2010), no. 3, 1807–1856.
15. G. C. Hruska and D. T. Wise, *Finiteness properties of cubulated groups*, Compos. Math. **150** (2014), no. 3, 453–506.
16. T. Hsu and D. T. Wise, *Cubulating malnormal amalgams*, Invent. Math. **199** (2015), no. 2, 293–331.
17. J. F. Manning and E. Martínez-Pedroza, *Separation of relatively quasiconvex subgroups*, Pacific J. Math. **244** (2010), no. 2, 309–334.
18. D. V. Osin, *Relatively hyperbolic groups: intrinsic geometry, algebraic properties, and algorithmic problems*, Mem. Amer. Math. Soc. **179** (2006), no. 843, vi+100.
19. D. V. Osin, *Peripheral fillings of relatively hyperbolic groups*, Invent. Math. **167** (2007), no. 2, 295–326.
20. P. Przytycki and D. T. Wise, *Mixed 3-manifolds are virtually special*, J. Amer. Math. Soc. **31** (2018), no. 2, 319–347.
21. M. Sageev and D. T. Wise, *Cores for quasiconvex actions*, Proc. Amer. Math. Soc. **143** (2015), no. (7), 2731–2741.
22. D. T. Wise, *The structure of groups with a quasiconvex hierarchy*, Princeton University Press, Princeton, 2021.