

UNIQUENESS OF K-POLYSTABLE DEGENERATIONS OF FANO VARIETIES

HAROLD BLUM AND CHENYANG XU

ABSTRACT. We prove that K-polystable degenerations of \mathbb{Q} -Fano varieties are unique. Furthermore, we show that the moduli stack of K-stable \mathbb{Q} -Fano varieties is separated. Together with recently proven boundedness and openness statements, the latter result yields a separated Deligne-Mumford stack parametrizing all uniformly K-stable \mathbb{Q} -Fano varieties of fixed dimension and volume. The result also implies that the automorphism group of a K-stable \mathbb{Q} -Fano variety is finite.

CONTENTS

1. Introduction	1
2. Preliminaries on valuations and K-stability	8
3. Uniformly K-stable Fano	17
4. Places computing the δ -invariant	22
5. Constructing the S-equivalence	26
References	41

1. INTRODUCTION

1.1. Moduli spaces of Fano varieties. To give a general framework for intrinsically constructing moduli spaces of Fano varieties is a challenging question in algebraic geometry, especially if one wants to find a compactification. Unlike the KSBA construction in the canonically polarized case, the Minimal Model Program often provides more than one limit for a family of Fano varieties over a punctured curve. Thus, it is unclear how to find a theory that picks the right limit. In examples, people have obtained a lot of working experience on how to identify the simplest limit. On the negative side, examples such as [PP10, Section 2.2], which gives a family that isotrivially degenerates a homogeneous space to a different quasi-homogeneous space (with non-reductive automorphism group), suggest that we should not consider all smooth Fano varieties.

So when the definition of K-stability from complex geometry [Tia97] and its algebraic formulation [Don02], which were introduced to characterize when a Fano variety admits a Kähler-Einstein metric, first appeared in front of algebraic geometers, it

Date: June 5, 2022.

seemed bold to expect such a notion would be a key ingredient in constructing moduli spaces of Fano varieties. However, as the theory has developed, more and more evidence makes such an expectation believable.

We now expect that the moduli functor $\mathcal{M}_{n,V}^{\text{Kss}}$ of n -dimensional K-semistable \mathbb{Q} -Fano varieties of volume V , which sends $S \in \text{Sch}_k$ to

$$\mathcal{M}_{n,V}^{\text{Kss}}(S) = \left\{ \begin{array}{l} \text{Flat proper morphisms } X \rightarrow S, \text{ whose geometric fibers are} \\ \quad n\text{-dimensional K-semistable } \mathbb{Q}\text{-Fano varieties with} \\ \quad \text{volume } V, \text{ satisfying Kollar's condition} \end{array} \right\}$$

is represented by an Artin stack $\mathcal{M}_{n,V}^{\text{Kss}}$ of finite type and admits a projective good moduli space $\mathcal{M}_{n,V}^{\text{Kss}} \rightarrow M_{n,V}^{\text{Kps}}$ (in the sense of [Alp13]), whose closed points are in bijection with n -dimensional K-polystable \mathbb{Q} -Fano varieties of volume V . Here, Kollar's condition means that the reflexive power $\omega_{X/S}^{[m]}$ is flat over S and commutes with arbitrary base change for each $m \in \mathbb{Z}$ (see [Kol09, 24]).

While smooth Kähler-Einstein Fano manifolds with finite automorphism group are asymptotically Chow stable [Don01], examples in [OSY12, LLSW17] show that the GIT approach likely fails to treat those with infinite automorphism groups or singularities. (See [WX14] for examples where asymptotic Chow stability fails to construct compact moduli spaces in the KSBA setting.) Therefore, we need to take a more abstract approach to constructing $M_{n,V}^{\text{Kps}}$.

The construction of $M_{n,V}^{\text{Kps}}$ reduces to proving a number of concrete statements about families of \mathbb{Q} -Fano varieties. We list the main ones:

- (I) **BOUNDEDNESS:** There is a positive integer $N = N(n, V)$ such that if $X \in \mathcal{M}_{n,V}^{\text{Kss}}(k)$, then $-NK_X$ is a very ample Cartier divisor. This is settled in [Jia17] using results in [Bir16].
- (II) **ZARISKI OPENNESS:** If $X \rightarrow S$ is a family of \mathbb{Q} -Fano varieties, then the locus where the fiber is K-semistable is a Zariski open set.

Together, (I) and (II) show that $\mathcal{M}_{n,V}^{\text{Kss}}$ is an Artin stack of finite type and is a global quotient. The following statements are needed to show $\mathcal{M}_{n,V}^{\text{Kss}}$ admits a projective good moduli space.

- (III) **GOOD QUOTIENT:** The stack $\mathcal{M}_{n,V}^{\text{Kss}}$ admits a good moduli space. To prove this, it suffices to show:
 - (III.a) **REDUCTIVE AUTOMORPHISM GROUP:** If X is a K-polystable \mathbb{Q} -Fano variety X , then $\text{Aut}(X)$ is reductive.
 - (III.b) **GLUING OF LOCAL QUOTIENTS:** Near each K-polystable \mathbb{Q} -Fano variety $X \in \mathcal{M}_{n,V}^{\text{Kss}}(k)$, there exists a local atlas around $[X]$ given by an $\text{Aut}(X)$ slice. Furthermore, a point in the slice is GIT (poly/semi)stable with respect to $\text{Aut}(X)$ if and only if the corresponding \mathbb{Q} -Fano variety is K-(poly/semi)stable. To complete this step, it remains to verify that the

local GIT quotient spaces glue together to give the good quotient $M_{n,V}^{\text{Kps}}$ (e.g. the hypotheses of [AFS17, Theorem 1.2] are satisfied).

(IV) **SEPARATEDNESS:** Any two K-semistable degenerations of a family of K-semistable \mathbb{Q} -Fano varieties over a punctured curve $C^\circ = C \setminus 0$ lie in the same S -equivalence class, i.e. they degenerate to a common K-semistable \mathbb{Q} -Fano variety via special test configurations.

(V) **PROPERNESS:** Roughly speaking, any family of K-semistable Fano varieties over a punctured curve $C^\circ = C \setminus 0$ can be filled in over 0 to a family of K-semistable \mathbb{Q} -Fano varieties over C .

(VI) **PROJECTIVITY:** A sufficiently divisible multiple of the CM-line bundle yields an ample line bundle on $M_{n,V}^{\text{Kps}}$.

We note that there are subtleties related to the requirement that objects in $\mathcal{M}_{n,V}^{\text{Kss}}(S)$ satisfy Kollar's condition. Luckily, such issues are of a local nature and have all been addressed in the construction of the moduli space of KSBA stable varieties (see [Kol09, Kol19]).

Strong evidence for the above picture is that, aside from (VI) (the projectivity of $M_{n,V}^{\text{Kps}}$), the problem is completely solved in [LWX19] (see also [SSY16, Oda15]) for \mathbb{Q} -Fano varieties with a \mathbb{Q} -Gorenstein smoothing and some progress on the projectivity was made in [LWX18a]. However, these results rely heavily on the deep analytic tools established in [CDS15, Tia15]. Therefore, a completely algebraic proof is highly desirable. Such a proof would likely allow us to drop the *smoothable* assumption.

The main result in this paper gives a complete solution to (IV). In the smoothable case, this step is solved in [LWX19, SSY16] using analytic tools. The argument in this document is purely algebraic.

1.2. Separatedness result. The following statement is our main result.

Theorem 1.1. *Let $\pi : (X, \Delta) \rightarrow C$ and $\pi' : (X', \Delta') \rightarrow C$ be \mathbb{Q} -Gorenstein families of log Fano pairs over a smooth pointed curve $0 \in C$. Assume there exists an isomorphism*

$$\phi : (X, \Delta) \times_C C^\circ \rightarrow (X', \Delta') \times_C C^\circ$$

over $C^\circ = C \setminus 0$.

- (1) *K-semistable case: If (X_0, Δ_0) and (X'_0, Δ'_0) are K-semistable, then they are S -equivalent.*
- (2) *K-polystable case: If (X_0, Δ_0) and (X'_0, Δ'_0) are K-polystable, then they are isomorphic.*
- (3) *K-stable case: If (X_0, Δ_0) is K-stable and (X'_0, Δ'_0) is K-semistable, then ϕ extends to an isomorphism $(X, \Delta) \simeq (X', \Delta')$ over C .*

Remark 1.2. (1) The K-polystable case of Theorem 1.1 follows immediately from the K-semistable case and Definitions 2.5 and 2.6.

(2) By [LWX18b], the K-semistable case of Theorem 1.1 can be strengthened to say (X_0, Δ_0) and (X'_0, Δ'_0) have a common K-polystable degeneration.

(3) A special case of Theorem 1.1 was proved in [Oda12, 1.4] with the additional assumption that $\alpha(X_0, \Delta_0) > \dim(X_0)/(\dim(X_0) + 1)$ (see also [Che09, 5.7]).

Theorem 1.1 implies the following special case of Step (III.a).

Corollary 1.3. *Let (X, Δ) be log Fano pair. If (X, Δ) is K-stable, then $\text{Aut}(X, \Delta)$ is finite.*

1.3. Moduli of uniformly K-stable Fano varieties. We now specialize our study to the moduli of uniformly K-stable Fano varieties. Consider the moduli functor $\mathcal{M}_{n,V}^{\text{uKs}}$ that sends $S \in \text{Sch}_k$ to

$$\mathcal{M}_{n,V}^{\text{uKs}}(S) = \left\{ \begin{array}{l} \text{flat, proper morphisms } X \rightarrow S, \text{ whose geometric fibers} \\ \text{are } n\text{-dimensional uniformly K-stable } \mathbb{Q}\text{-Fano varieties} \\ \text{of volume } V, \text{ satisfying Kollar's condition} \end{array} \right\}.$$

Combining the following recent results:

- (I^u) **BOUNDEDNESS:** Proved in [Jia17],
- (II^u) **ZARISKI OPENNESS:** Proved in [BL18], and
- (III^u) **SEPARATEDNESS (AS A STACK):** Theorem 1.1.3,

we obtain the following corollary.

Corollary 1.4. *The functor $\mathcal{M}_{n,V}^{\text{uKs}}$ is a separated Deligne-Mumford stack of finite type, which has a coarse moduli space $M_{n,V}^{\text{uKs}}$ that is a separated algebraic space.*

One still missing property is

- (IV^u) **QUASI-PROJECTIVITY:** $M_{n,V}^{\text{uKs}}$ is quasi-projective.

Significant progress on this problem was made in [CP18].

1.4. Summary of the paper. The original definition of K-stability in [Tia97, Don02] is defined in terms of the sign of the generalized Futaki invariant on all test configurations or at least special test configurations (see [LX14]). Recently, there has been tremendous progress in reinterpreting K-stability in terms of invariants associated to valuations rather than test configurations.

More specifically, in [BHK17], the data of a test configuration was translated into the data of a filtration and it was shown that a nontrivial special test configuration yields a divisorial valuation. Then in a series of papers [Fuj16, Fuj19a, Fuj18] of K. Fujita, all divisorial valuations were studied and an invariant β was defined for each divisorial valuation. After [Li17], it became more natural to extend the setup to all valuations over the log Fano variety rather than only considering divisorial valuations (see also [LX16, BJ17]). Moreover, a characterization of K-stability notions in terms of the sign of β -invariant for divisorial valuations was proved in [Li17, Fuj19a] and lead to another characterization by the δ -invariant in [FO18, BJ17]. These interpretations of K-stability using valuations have made it easier to apply techniques from birational geometry, especially the Minimal Model Program, to the study of K-stability.

In Section 2, we will have a short discussion on the above materials. More precisely, we will provide information on the language of valuations and filtrations following [BHQ17, Fuj19b, Li17] and the invariants β and δ associated to them following [Fuj19b, FO18, BJ17]. We also discuss the normalized volume function from [Li18] and its relation with the K-stability of Fano varieties (see [Li17, LX16]).

To proceed with our discussion, let us define the above invariants. Let (X, Δ) be a log Fano pair. Given a divisor E over X (i.e. $E \subset Y$ is a prime divisor, where Y is a normal variety with a proper birational morphism $\pi : Y \rightarrow X$), the β -invariant of E is given by

$$\beta_{X, \Delta}(E) := A_{X, \Delta}(E)(-K_X - \Delta)^n - \int_0^\infty \text{vol}(\pi^*(-K_X - \Delta) - tE)dt$$

where $A_{X, \Delta}(E)$ is the log discrepancy of E . This invariant was defined in [Fuj18] and the K-(semi)stability of (X, Δ) can be phrased in terms of the positivity of $\beta_{X, \Delta}(E)$ [Fuj19a, Li17].

Next, is the δ -invariant of (X, Δ) , which, as defined in [FO18], measures log canonical thresholds of a certain classes of anti-log canonical divisors of (X, Δ) . It is shown in [BJ17] that

$$\delta(X, \Delta) = \inf_E \frac{A_{X, \Delta}(E)(-K_X - \Delta)^n}{\int_0^\infty \text{vol}(\pi^*(-K_X - \Delta) - tE)dt}. \quad (1)$$

Hence, we say that a divisor E over X computes $\delta(X, \Delta)$ if it achieves the infimum in (1). The pair (X, Δ) is uniformly K-stable (resp. K-semistable) if and only if $\delta(X, \Delta) > 1$ (resp. $\delta(X, \Delta) \geq 1$) [FO18, BJ17].

In Section 3, before attacking Theorem 1.1 in full generality, we consider the special case in which (X_0, Δ_0) is uniformly K-stable and (X'_0, Δ'_0) is K-semistable. In this case, we provide a short proof of the separatedness result by using properties of the δ -invariant to reduce the question to the well known separatedness statement for the moduli functor of klt log Calabi-Yau pairs (Proposition 3.2). This argument is more straightforward than the general case and takes a slightly different approach. We hope this perspective can be applied in other cases.

To prove Theorem 1.1 in full generality is more involved. We need to study the case when the δ -invariants of the special fibers equal one. In general, analyzing the valuation computing $\delta = 1$ is quite subtle. For instance, the following statement has been conjectured by experts.

Conjecture 1.5. *Let (X, Δ) be a log Fano pair. If $\delta(X, \Delta) \leq 1$ then $\delta(X, \Delta)$ is computed by a divisor over X and any such divisor is dreamy.*

The special case of Conjecture 1.5 when $\delta(X, \Delta) = 1$ implies that K-stability is equivalent to the apparently stronger notion of uniform K-stability. This is known for smooth Fano varieties by [BBJ15], but the proof relies on analytic tools, in particular the existence of Kähler-Einstein metrics.

In Section 4, we will prove some special cases of Conjecture 1.5 which are needed in our proof of Theorem 1.1. The first result is that if (X, Δ) is a log Fano variety with $\delta(X, \Delta) = 1$, then any divisor computing $\delta(X, \Delta)$ is necessarily dreamy and induces a special test configuration of (X, Δ) . The proof relies on the MMP techniques developed in [LWX18b], which build upon work in [Li17, LX16, LX18]. Specifically, we consider the cone over our log Fano pair and use the calculation in [Li17, LX16] which shows that $\beta_{X, \Delta}(E)$ equals the derivative of the normalized volume function on the valuation space of the cone along the path given by the interval connecting the divisorial valuation associated to the pull back of E and the canonical valuation. A careful study as in [LWX18b, Theorem 3.2] shows that E is indeed a dreamy divisor and induces a special test configuration. In Section 4.2 and 4.3, we also address the situations when the δ -invariant can be calculated by an ideal or a \mathbb{Q} -divisor. These results may be of independent interest.

Section 5 is the core of this paper and where we prove Theorem 1.1. The majority of the work in this section is to construct the S-equivalence stated in the theorem.

Step 1: We first observe that a pair of two different degenerations will induce filtrations on each other's section rings. Furthermore, the associated graded rings of the filtrations are isomorphic with a grading shift matching the calculation of β -invariant.

Let us explain the above construction in more detail. Assume we have two \mathbb{Q} -Gorenstein families of log Fano pairs $\pi : (X, \Delta) \rightarrow C$ and $\pi' : (X', \Delta') \rightarrow C$ over a smooth affine curve C and an isomorphism

$$\phi : (X, \Delta) \times_C C^\circ \rightarrow (X', \Delta') \times_C C^\circ,$$

that does not extend to an isomorphism over $C^\circ = C \setminus 0$. Fix r so that $L := -r(K_X + \Delta)$ and $L' := -r(K_{X'} + \Delta')$ are Cartier. We choose a proper birational model over X and X'

$$\begin{array}{ccc} & \widehat{X} & \\ \psi \swarrow & & \searrow \psi' \\ X & \xrightarrow{\phi} & X' \end{array}$$

and write \widetilde{X}_0 and \widetilde{X}'_0 for the birational transforms of X_0 and X'_0 on \widehat{X} . The divisor \widetilde{X}'_0 induces a filtration \mathcal{F} on the section ring of (X_0, Δ_0) defined by

$$s \in \mathcal{F}^p H^0(X_0, mL_0) \quad \text{if and only if} \quad \text{ord}_{\widetilde{X}'_0}(\tilde{s}) \geq p$$

for some (non-unique) extension $\tilde{s} \in H^0(X, mL)$. Then we define

$$\beta := (-K_{X_0} - \Delta_0)^n A_{X, \Delta+X_0}(\widetilde{X}'_0) - \lim_{m \rightarrow \infty} \int_0^\infty \frac{\dim \mathcal{F}^{mx} H^0(X_0, mL_0)}{r^{n+1} m^n / n!} dx.$$

Similarly, we can define a filtration \mathcal{F}' of the section ring of (X'_0, Δ'_0) and the value β' using the divisor \widetilde{X}_0 . The construction here can be viewed as a relative version of the one in [BHJ17, Section 5], where they consider a test configuration and a trivial family.

Next, we observe that there is an isomorphism of the associated graded rings of the filtrations

$$\bigoplus_{m \in \mathbb{N}} \bigoplus_{p \in \mathbb{Z}} \text{gr}_{\mathcal{F}}^p H^0(X_0, mL_0) \xrightarrow{\varphi} \bigoplus_{m \in \mathbb{N}} \bigoplus_{p \in \mathbb{Z}} \text{gr}_{\mathcal{F}'}^{mr(a+a')-p} H^0(X'_0, mL'_0), \quad (2)$$

where $a := A_{X, \Delta+X_0}(\tilde{X}'_0)$ and $a' := A_{X', \Delta'+X'_0}(\tilde{X}_0)$. Using this isomorphism, we deduce that $\beta + \beta' = 0$.

Now if we assume (X_0, Δ_0) and (X'_0, Δ'_0) are K-semistable, then the β -invariant of any divisor over X_0 or X'_0 is non-negative [Fuj19a, Li17]. A similar result is extended to filtrations in [BL18].¹ We can then conclude that $\beta = \beta' = 0$.

Step 2: At this point, we know X_0 and X'_0 have a common degeneration. Indeed, the Rees construction gives degenerations

$$X_0 \rightsquigarrow \mathcal{X}_0 := \text{Proj} \left(\bigoplus_{m \in \mathbb{N}} \bigoplus_{p \in \mathbb{Z}} \text{gr}_{\mathcal{F}}^p H^0(X_0, mL_0) \right)$$

and

$$X'_0 \rightsquigarrow \mathcal{X}'_0 := \text{Proj} \left(\bigoplus_{m \in \mathbb{N}} \bigoplus_{p \in \mathbb{Z}} \text{gr}_{\mathcal{F}'}^p H^0(X'_0, mL'_0) \right).$$

By (2), the degenerations \mathcal{X}_0 and \mathcal{X}'_0 are isomorphic.

An immediate concern is that the above graded rings are not necessarily finitely generated. (Note that notions of K-stability have been investigated in the setting of non-finitely generated filtrations [WN12, Szé15].) Since we aim to prove (X_0, Δ_0) and (X'_0, Δ'_0) have a common degeneration to a K-semistable log Fano pair, we must show that the filtrations \mathcal{F} and \mathcal{F}' are finitely generated and induce special test configurations with generalized Futaki invariant zero. By [LWX18b, 3.1], this will imply that the degenerations are K-semistable log Fano pairs.

To proceed, we rely on the fact that our filtrations are induced by divisors over our families. More precisely, we use that $\beta = 0$ to show that there exists an extraction $Y \rightarrow X$ of \tilde{X}'_0 and the fiber $Y_0 = V \cup W$, where V and W are the birational transforms of X_0 and X'_0 . Now, we set $E = W|_V$ and observe that E induces a filtration \mathcal{F}_E on the section ring of (X_0, Δ_0) . We then show $F := \text{Supp}(E)$ is a prime divisor and $\beta_{X_0, \Delta_0}(F) = 0$. Using Theorem 4.1, we see \mathcal{F}_E is finitely generated and the corresponding degeneration of (X_0, Δ_0) is a special test configuration with generalized Futaki invariant zero.

Next, we seek to show that the filtrations \mathcal{F} and \mathcal{F}_E are equal. This statement is equivalent to the surjectivity of certain restriction maps and is non-trivial. To achieve the result, we take the relative cone of (X, Δ) over C and run an analysis similar to the proof of Theorem 4.1. After completing this argument, we can conclude that the degenerations $(\mathcal{X}_0, \mathcal{D}_0)$ and $(\mathcal{X}'_0, \mathcal{D}'_0)$ are naturally K-semistable pairs.

¹This is also independently obtained by Chi Li and Xiaowei Wang in [LWX18].

Finally, we need to show that the isomorphism $\mathcal{X}_0 \simeq \mathcal{X}'_0$ sends the degeneration of Δ_0 to the degeneration of Δ'_0 , so that we get an isomorphism of pairs. To verify this, we choose a divisor $B \subseteq \text{Supp}(\Delta)$ and write $B' \subseteq \text{Supp}(\Delta')$ for its birational transform. Now, B_0 degenerates to a divisor on \mathcal{X}_0 that corresponds to the initial ideal $\text{in}(I_{B_0})$ in the associated graded ring. Rather than showing $\varphi(\text{in}(I_{B_0})) = \text{in}(I_{B'_0})$, we introduce auxiliary ideals I and I' such that the equality $\varphi(I) = I'$ is clear. (The ideal I is defined by restricting elements of the relative section ring that vanish to certain orders along B and \tilde{X}'_0 .) Using the relative cone construction again, we show that I and I' agree with $\text{in}(I_{B_0})$ to $\text{in}(I_{B'_0})$ at codimension one points. We can then conclude that the desired isomorphism of boundaries holds.

Acknowledgement: We are grateful to Roman Bezrukavnikov, Giulio Codogni, Tommaso de Fernex, Christopher Hacon, Mattias Jonsson, János Kollár, Chi Li, Yuchen Liu, Mircea Mustață, Yuji Odaka, Xiaowei Wang, and Jun Yu for helpful conversations and comments on previous drafts of this paper. A large part of the work on this paper was completed when CX was visiting Institut Henri Poincaré as part of the Poincaré Chair program. He thanks the institute for the wonderful environment and Claire Voisin for her hospitality. Finally, we are indebted to the anonymous referees whose comments improved this paper considerably.

HB is partially supported by NSF grant DMS-1803102. CX is partially supported by the National Science Fund for Distinguished Young Scholars (NSFC 11425101) “Algebraic Geometry”.

2. PRELIMINARIES ON VALUATIONS AND K-STABILITY

2.1. Conventions. We work over an algebraically closed field \mathbb{k} of characteristic 0. We follow the terminologies in [KM98, Kol13]. A *pair* (X, Δ) is composed of a normal variety X and an effective \mathbb{Q} -divisor Δ on X such that $K_X + \Delta$ is \mathbb{Q} -Cartier. See [KM98, 2.34] for the definitions of *klt*, *plt*, and *lc* pairs. A pair (X, Δ) is *log Fano* if X is projective, (X, Δ) is klt, and $-K_X - \Delta$ is ample. A variety X is \mathbb{Q} -*Fano* if $(X, 0)$ is log Fano.

Definition 2.1. A \mathbb{Q} -*Gorenstein family of log Fano pairs* $\pi : (X, \Delta) \rightarrow C$ over a smooth curve C is composed of a flat proper morphism $\pi : X \rightarrow C$ and an effective \mathbb{Q} -divisor Δ , not containing any fiber of π , satisfying:

- (1) π has normal, connected fibers (hence, X is normal as well)
- (2) $-K_X - \Delta$ is \mathbb{Q} -Cartier and π -ample, and
- (3) (X_t, Δ_t) is klt for all $t \in C$.

2.2. Valuations. Let X be a variety. A valuation on X will mean a valuation $v : K(X)^\times \rightarrow \mathbb{R}$ that is trivial on \mathbb{k} and has center on X . Recall, v has center on X if there exists a point $\xi \in X$ such that $v \geq 0$ on $\mathcal{O}_{X, \xi}$ and > 0 on the maximal ideal. Since X is assumed to be separated, such a point ξ is unique, and we say v has *center* $c_X(v) := \xi$. See [JM12, 3.1] for the definition of quasimonomial valuations.

Following [JM12, BdFFU15], we write Val_X for the set of valuations on X and Val_X^* for the set of non-trivial ones. To any valuation $v \in \text{Val}_X$ and $p \in \mathbb{N}$, there is an associated *valuation ideal* $\mathfrak{a}_p(v)$. For an affine open subset $U \subseteq X$, $\mathfrak{a}_p(v)(U) = \{f \in \mathcal{O}_X(U) \mid v(f) \geq p\}$ if $c_X(v) \in U$ and $\mathfrak{a}_p(v)(U) = \mathcal{O}_X(U)$ otherwise.

For an ideal $\mathfrak{a} \subseteq \mathcal{O}_X$ and $v \in \text{Val}_X$, we set

$$v(\mathfrak{a}) := \min\{v(f) \mid f \in \mathfrak{a} \cdot \mathcal{O}_{X,c_X(v)}\} \in [0, +\infty].$$

We can define $v(s)$ when \mathcal{L} is a line bundle on X and $s \in H^0(X, \mathcal{L})$. After trivializing \mathcal{L} at $c_X(v)$, we set $v(s) = v(f)$, where f is the local function corresponding to s under this trivialization (this is independent of choice of trivialization).

2.2.1. Divisors over X . Let X be a variety and $\pi : Y \rightarrow X$ be a proper birational morphism, with Y normal. A prime divisor $E \subset Y$ defines a valuation $\text{ord}_E : K(X)^\times \rightarrow \mathbb{Z}$ given by order of vanishing at E . Note that $c_X(\text{ord}_E)$ is the generic point of $\pi(E)$ and, assuming X is normal, $\mathfrak{a}_p(v) = \pi_* \mathcal{O}_Y(-pE)$.

We identify two such prime divisors on Y_1 and Y_2 as above if one is the birational transform of the other. Equivalently, they induce the same valuation of $K(X)$. A *divisor over X* is an equivalence class given by this relation.

2.2.2. Log discrepancies. Let (X, Δ) be a pair. We write

$$A_{X, \Delta} : \text{Val}_X^* \rightarrow \mathbb{R} \cup \{+\infty\}$$

for the log discrepancy function with respect to (X, Δ) as in [JM12, BdFFU15] (see [Blu18b] for the case when $\Delta \neq 0$).

When $\pi : Y \rightarrow X$ is a proper birational morphism with Y normal and $E \subset Y$ a prime divisor,

$$A_{X, \Delta}(\text{ord}_E) = 1 + \text{coeff}_E(K_Y - \pi^*(K_X + \Delta)).$$

We will often write $A_{X, \Delta}(E)$ for the above value.

The function $A_{X, \Delta}$ is homogenous of degree 1, i.e. $A_{X, \Delta}(\lambda v) = \lambda \cdot A_{X, \Delta}(v)$ for $\lambda \in \mathbb{R}_{>0}$ and $v \in \text{Val}_X$. A pair (X, Δ) is klt (resp., lc) if and only if $A_{X, \Delta}(v) > 0$ (resp., ≥ 0) for all $v \in \text{Val}_X^*$.

2.2.3. Graded sequences. A *graded sequence of ideals* $\mathfrak{a}_\bullet = (\mathfrak{a}_p)_{p \in \mathbb{Z}_{>0}}$ on a variety X is a sequence of ideals on X satisfying $\mathfrak{a}_p \cdot \mathfrak{a}_q \subseteq \mathfrak{a}_{p+q}$ for all $p, q \in \mathbb{Z}_{>0}$. By convention, $\mathfrak{a}_0 = \mathcal{O}_X$. We set $M(\mathfrak{a}_\bullet) := \{p \in \mathbb{Z}_{>0} \mid \mathfrak{a}_p \neq (0)\}$ and always assume $M(\mathfrak{a}_\bullet)$ is nonempty. If $v \in \text{Val}_X^*$, then $\mathfrak{a}_\bullet(v)$ is a graded sequence of ideals.

Let \mathfrak{a}_\bullet be a graded sequence of ideals on X and $v \in \text{Val}_X$. It follows from Fekete's Lemma that the limit

$$v(\mathfrak{a}_\bullet) := \lim_{M(\mathfrak{a}_\bullet) \ni m \rightarrow \infty} \frac{v(\mathfrak{a}_m)}{m}$$

exists and equals $\inf_{m \geq 1} \frac{v(\mathfrak{a}_m)}{m}$; see [JM12, §2.1].

Let $x \in X$ be a closed point. If \mathfrak{a}_\bullet is a graded sequence of ideals on X and each ideal \mathfrak{a}_p is \mathfrak{m}_x -primary, we set

$$\text{mult}(\mathfrak{a}_\bullet) := \lim_{p \rightarrow \infty} \frac{\dim_{\mathbb{k}}(\mathcal{O}_X/\mathfrak{a}_p)}{p^n/n!}.$$

If $v \in \text{Val}_X$ has center $\{x\}$, then $\mathfrak{a}_p(v)$ is \mathfrak{m}_x -primary for each $p > 0$. In this case, we call $\text{vol}(v) := \text{mult}(\mathfrak{a}_\bullet(v))$ the *volume* of v .

2.2.4. Log canonical thresholds. Let (X, Δ) be an lc pair. Given a nonzero ideal $\mathfrak{a} \subseteq \mathcal{O}_X$, the log canonical threshold of \mathfrak{a} is given by

$$\text{lct}(X, \Delta; \mathfrak{a}) := \sup\{c \in \mathbb{Q}_{\geq 0} \mid (X, \Delta + \mathfrak{a}^c) \text{ is lc}\}.$$

If \mathfrak{a}_\bullet a graded sequence of ideals on X , the log canonical threshold of \mathfrak{a}_\bullet is given by

$$\text{lct}(X, \Delta; \mathfrak{a}_\bullet) := \lim_{M(\mathfrak{a}_\bullet) \ni m \rightarrow \infty} m \cdot \text{lct}(X, \Delta; \mathfrak{a}_m).$$

Fekete's Lemma implies that the above limit exists and equals $\sup_m m \cdot \text{lct}(X, \Delta; \mathfrak{a}_m)$ [JM12, 2.5].

It is straightforward to show $\text{lct}(X, \Delta; \mathfrak{a}_\bullet) \leq \frac{A_{X, \Delta}(v)}{v(\mathfrak{a}_\bullet)}$, for $v \in \text{Val}_X^*$ satisfying $0 \neq A_{X, \Delta}(v) < +\infty$. Hence, if $v \in \text{Val}_X^*$ satisfies $A_{X, \Delta}(v) \neq 0$, then

$$\text{lct}(X, \Delta; \mathfrak{a}_\bullet(v)) \leq A_{X, \Delta}(v), \tag{3}$$

since $v(\mathfrak{a}_\bullet(v)) = 1$ [Blu18b, 3.4.9].

2.2.5. Extractions. Let E be a divisor over a normal variety X . We say that $\mu : X_E \rightarrow X$ is an *extraction* of E if μ is a proper birational morphism with X_E is normal, E arises as a prime divisor $E \subset X_E$, and $-E$ is μ -ample.

Note that if $\mu : X_E \rightarrow X$ is an extraction of E , then $E \supseteq \text{Exc}(\mu)$ and equality holds if $\text{codim}_X(c_X(\text{ord}_E)) \geq 2$. Indeed, Lemma 4.5 implies that if $p \in \mathbb{Z}_{>0}$ is sufficiently divisible, then μ is the blowup along $\mathfrak{a}_p(\text{ord}_E)$ and $\mathfrak{a}_p(\text{ord}_E) \cdot \mathcal{O}_Y = \mathcal{O}_Y(-pE)$.

The following technical statement gives a criterion for when an exceptional divisor may be extracted. The criterion will be used repeatedly in Section 5.

Proposition 2.2. *Let (X, Δ) be a klt pair or a plt pair such that $\lfloor \Delta \rfloor = S$ is a non-zero \mathbb{Q} -Cartier divisor. If E is a divisor over X satisfying*

$$a := A_{X, \Delta}(E) - \text{lct}(X, \Delta; \mathfrak{a}_\bullet(\text{ord}_E)) < 1,$$

then there exists an extraction $\mu : X_E \rightarrow X$ of E and $(X_E, \mu_^{-1}(\Delta) + (1 - a)E)$ is lc.*

The proposition is a consequence of [BCHM10] and properties of the log canonical threshold of a graded sequence of ideals.

Proof. See the argument in [Blu17, 1.5] for the case when (X, Δ) is klt. If (X, Δ) is plt, observe that $(X, \Delta_\varepsilon := \Delta - \varepsilon S)$ is klt for $0 < \varepsilon < 1$. If we set

$$a_\varepsilon := A_{X, \Delta_\varepsilon}(\text{ord}_E) - \text{lct}(X, \Delta_\varepsilon; \mathfrak{a}_\bullet(\text{ord}_E)),$$

then $\lim_{\varepsilon \rightarrow 0} a_\varepsilon = a$ and we may reduce to the klt case. \square

2.3. Filtrations. Let (X, Δ) be a n -dimensional log Fano pair. Fix a positive integer r such that $L := -r(K_X + \Delta)$ is Cartier and write

$$R = R(X, L) = \bigoplus_{m \in \mathbb{N}} R_m = \bigoplus_{m \in \mathbb{N}} H^0(X, \mathcal{O}_X(mL))$$

for the section ring of L . Set $M(L) := \{m \in \mathbb{N} \mid H^0(X, \mathcal{O}_X(mL)) \neq 0\}$.

Definition 2.3. A *filtration* \mathcal{F} of R we will mean the data of a family of vector subspaces $\mathcal{F}^\lambda R_m \subseteq R_m$ for $m \in \mathbb{N}$ and $\lambda \in \mathbb{R}$ satisfying

- (1) $\mathcal{F}^\lambda R_m \subseteq \mathcal{F}^{\lambda'} R_m$ when $\lambda \geq \lambda'$;
- (2) $\mathcal{F}^\lambda R_m = \bigcap_{\lambda' < \lambda} \mathcal{F}^{\lambda'} R_m$;
- (3) $\mathcal{F}^0 R_m = R_m$ and $\mathcal{F}^\lambda R_m = 0$ for $\lambda \gg 0$.
- (4) $\mathcal{F}^\lambda R_m \cdot \mathcal{F}^{\lambda'} R_{m'} \subseteq \mathcal{F}^{\lambda+\lambda'} R_{m+m'}$.

A filtration \mathcal{F} of R is called an \mathbb{N} -filtration if $\mathcal{F}^\lambda R_m = \mathcal{F}^{[\lambda]} R_m$ for all $m \in \mathbb{N}$ and $\lambda \in \mathbb{R}$. To give a \mathbb{N} -filtration \mathcal{F} , it suffices to give a family of subspaces $\mathcal{F}^p R_m \subseteq R_m$ for $m, p \in \mathbb{N}$ satisfying (1), (3), and (4).

A filtration \mathcal{F} is *linearly bounded* if there exists $C > 0$ so that $\mathcal{F}^{Cm} R_m = 0$ for all $m \in \mathbb{N}$ and *trivial* if $\mathcal{F}^\lambda R_m = 0$ for all $m \in \mathbb{N}$ and $\lambda > 0$.

2.3.1. Rees construction. Let \mathcal{F} be an \mathbb{N} -filtration of R . The *Rees algebra* of \mathcal{F} is the $\mathbb{k}[t]$ -algebra

$$\text{Rees}(\mathcal{F}) := \bigoplus_{m \in \mathbb{N}} \bigoplus_{p \in \mathbb{Z}} (\mathcal{F}^p R_m) t^{-p} \subseteq R[t, t^{-1}].$$

The *associated graded ring* of \mathcal{F} is

$$\text{gr}_{\mathcal{F}} R := \bigoplus_{m \in \mathbb{N}} \bigoplus_{p \in \mathbb{Z}} \text{gr}_{\mathcal{F}}^p R_m, \quad \text{where } \text{gr}_{\mathcal{F}}^p R_m = \frac{\mathcal{F}^p R_m}{\mathcal{F}^{p+1} R_m}.$$

Note that

$$\text{Rees}(\mathcal{F}) \otimes_{k[t]} k[t, t^{-1}] \simeq R[t, t^{-1}] \quad \text{and} \quad \frac{\text{Rees}(\mathcal{F})}{t \text{Rees}(\mathcal{F})} \simeq \text{gr}_{\mathcal{F}} R. \quad (4)$$

Hence, $\text{Rees}(\mathcal{F})$ is said to give a degeneration of R to the associated graded ring of \mathcal{F} .

An \mathbb{N} -filtration \mathcal{F} is *finitely generated* if $\text{Rees}(\mathcal{F})$ is a finitely generated $\mathbb{k}[t]$ -algebra. In this case, we set $\mathcal{X} := \text{Proj}_{\mathbb{A}^1}(\text{Rees}(\mathcal{F}))$. By (4),

$$\mathcal{X}_{\mathbb{A}^1 \setminus 0} \simeq X \times (\mathbb{A}^1 \setminus 0) \quad \text{and} \quad \mathcal{X}_0 \simeq \text{Proj}(\text{gr}_{\mathcal{F}} R).$$

We write \mathcal{D} for the \mathbb{Q} -divisor that is the closure of $\Delta \times (\mathbb{A}^1 \setminus 0)$ under the embedding of $X \times (\mathbb{A}^1 \setminus 0)$ in \mathcal{X} .

The scheme \mathcal{X} can naturally be endowed with the structure of a *test configuration* of (X, Δ) . The test configuration is called *special* if $(\mathcal{X}, \mathcal{D}) \rightarrow \mathbb{A}^1$ is a \mathbb{Q} -Gorenstein family of log Fano pairs. See [LX14, §3] and [BHJ17, §2] for information on test configurations and the generalized Futaki invariant.

With the above setup, consider a subscheme $Z \subset X$ and write $I_Z \subset R$ for the corresponding homogenous ideal. The scheme theoretic closure of $Z \times (\mathbb{A}^1 \setminus 0)$ in \mathcal{X} , denoted by \mathcal{Z} , is defined by the ideal

$$\bigoplus_{m \in \mathbb{N}} \bigoplus_{p \in \mathbb{Z}} (\mathcal{F}^p R_m \cap I_Z) t^{-p} \subseteq \text{Rees}(\mathcal{F}).$$

Indeed, the corresponding subscheme agrees with $Z \times (\mathbb{A}^1 \setminus 0)$ away from 0 and is torsion free over 0. The above description of \mathcal{Z} yields that its scheme theoretic fiber along 0 is given by the *initial ideal*

$$\text{in}(I_Z) := \bigoplus_{m \in \mathbb{N}} \bigoplus_{p \in \mathbb{Z}} \text{im}(\mathcal{F}^p R_m \cap I_Z \rightarrow \text{gr}_{\mathcal{F}}^p R_m) \subset \text{gr}_{\mathcal{F}}^p R_m.$$

2.3.2. *Volume.* Given a filtration \mathcal{F} of R , we set

$$\text{vol}(\mathcal{F}R^{(x)}) := \limsup_{m \rightarrow \infty} \frac{\dim(\mathcal{F}^{xm} R_m)}{m^n/n!}$$

for $x \in \mathbb{R}_{\geq 0}$. Assuming \mathcal{F} is linearly bounded (which implies $\text{vol}(\mathcal{F}R^{(x)}) = 0$ for $x \gg 0$), we set

$$S(\mathcal{F}) := \frac{1}{r^{n+1}(-K_X - \Delta)^n} \int_0^\infty \text{vol}(\mathcal{F}R^{(x)}) dx. \text{²}$$

By [BC11] (see also [BHK17, 5.3]),

$$S(\mathcal{F}) = \lim_{m \rightarrow \infty} \left(\frac{1}{r \dim(R_m)} \int_0^\infty \dim(\mathcal{F}^{mx} R_m) dx \right) \quad (5)$$

In particular, if \mathcal{F} is an \mathbb{N} -filtration, then $S(\mathcal{F}) = \lim_{m \rightarrow \infty} \frac{\sum_{p \geq 0} (p \dim \text{gr}_{\mathcal{F}}^p R_m)}{mr \dim R_m}$.

2.3.3. *Base ideals.* Given a filtration \mathcal{F} of R , set

$$\mathfrak{b}_{p,m} := \text{im}(\mathcal{F}^p R_m \otimes \mathcal{O}_X(-mL) \rightarrow \mathcal{O}_X).$$

for $p, m \geq 0$. We set $\mathfrak{b}_p(\mathcal{F}) := \mathfrak{b}_{p,m}$ for $m \gg 1$. The ideal $\mathfrak{b}_p(\mathcal{F})$ is well defined and $\mathfrak{b}_\bullet(\mathcal{F})$ is a graded sequence of ideals assuming \mathcal{F} is non-trivial [BJ17, 3.17-3.18].

2.3.4. *Filtrations induced by valuations.* Given $v \in \text{Val}_X^*$, we set

$$\mathcal{F}_v^\lambda R_m = \{s \in R_m \mid v(s) \geq \lambda\}$$

for each $\lambda \in \mathbb{R}$ and $m \in \mathbb{N}$. Equivalently, $\mathcal{F}_v^\lambda R_m = H^0(X, \mathcal{O}_X(mL) \otimes \mathfrak{a}_\lambda(v))$. Note that \mathcal{F}_v is a non-trivial filtration of R .

If $A_{X,\Delta}(v) < +\infty$, then \mathcal{F}_v is linearly bounded [BJ17, 3.1]. In this case, we set $S(v) := S(\mathcal{F}_v)$.

²Note that this differs from the definition of $S(\mathcal{F})$ in [BJ17, BL18] by a factor of $1/r$. Since we are interested in the polarization $-K_X - \Delta$, not L , such a convention is natural.

2.3.5. *Filtrations induced by divisors.* If E is a divisor over X , we write $\mathcal{F}_E := \mathcal{F}_{\text{ord}_E}$ and $S(E) := S(\mathcal{F}_E)$. Following [Fuj19a], we say E is *dreamy* if \mathcal{F}_E is a finitely generated filtration of R .

When E arises as a prime divisor on a proper normal model $\mu : Y \rightarrow X$,

$$\mathcal{F}_E^\lambda R_m = H^0(Y, \mathcal{O}_Y(mr\mu^*(-K_X - \Delta) - \lceil \lambda \rceil E)).$$

Therefore,

$$S(E) = \frac{1}{(-K_X - \Delta)^n} \int_0^\infty \text{vol}(\mu^*(-K_X - \Delta) - xE) dx.$$

2.4. **K-stability.** Based on the original analytic definition in [Tia97], an algebraic definition of K-(semi,poly)-stability was introduced in [Don02]. Here, we will define these notations for log Fano pairs using valuations.

2.4.1. *β -invariant.* Let (X, Δ) be an n -dimensional log Fano pair and E a divisor over X . Following [Fuj19a],

$$\beta_{X, \Delta}(E) := (-K_X - \Delta)^n (A_{X, \Delta}(E) - S(E)).$$

More generally, if $v \in \text{Val}_X$ with $A_{X, \Delta}(v) < +\infty$, we set $\beta_{X, \Delta}(v) := (-K_X - \Delta)^n (A_{X, \Delta}(v) - S(v))$.

Definition 2.4. A log Fano pair (X, Δ) is

- (1) *K-semistable* if $\beta_{X, \Delta}(E) \geq 0$ for all divisors E over X ;
- (2) *K-stable* if $\beta_{X, \Delta}(E) > 0$ for all dreamy divisors E over X ;
- (3) *uniformly K-stable* if there exists an $\varepsilon > 0$ such that

$$\beta_{X, \Delta}(E) \geq \varepsilon A_{X, \Delta}(E) (-K_X - \Delta)^n$$

for all divisors E over X .

The equivalence of the above definition with the original definitions was addressed in [Fuj19a, Fuj19b, Li17] and the arguments rely on the special degeneration theory of [LX14]. In Corollary 4.2, we will show that the wordy *dreamy* may be removed from Definition 2.4.2.

Definition 2.5. A log Fano pair (X, Δ) is *K-polystable* if it is K-semistable and any special test configuration $(\mathcal{X}, \mathcal{D}) \rightarrow \mathbb{A}^1$ of (X, Δ) with $(\mathcal{X}_0, \mathcal{D}_0)$ K-semistable satisfies $(\mathcal{X}, \mathcal{D}) \simeq (X, \Delta) \times \mathbb{A}^1$.

The equivalence of the above definition with the definition in [LX14, 6.2] relies on the following result: If (X, Δ) is a K-semistable log Fano pair and $(\mathcal{X}, \mathcal{D})$ is a special test configuration of (X, Δ) , then $\text{Fut}(\mathcal{X}, \mathcal{D}) = 0$ if and only if $(\mathcal{X}_0, \mathcal{D}_0)$ is K-semistable [LWX18b, 3.1].

Definition 2.6. Two K-semistable log Fano pairs (X, Δ) and (X', Δ') are *S-equivalent* if they degenerate to a common K-semistable log Fano pair via special test configurations.

By [LWX18b, 3.2], S-equivalent log Fano pairs degenerate to a common K-polystable pair via special test configurations. Furthermore, the K-polystable pair is uniquely determined up to isomorphism.

2.4.2. *δ -invariant.* We recall an interpretation of the above discussion using an invariant introduced in [FO18].

Let (X, Δ) be a log Fano pair. Fix a positive integer r so that $L := -r(K_X + \Delta)$ is a Cartier divisor and $H^0(X, \mathcal{O}_X(L)) \neq 0$. Given $m \in r\mathbb{N}$, we say $D \sim_{\mathbb{Q}} -K_X - \Delta$ is m -basis type if there exists a basis $\{s_1, \dots, s_{N_m}\}$ of $H^0(X, \mathcal{O}_X(-m(K_X + \Delta)))$ such that

$$D = \frac{1}{mN_m} (\{s_1 = 0\} + \dots + \{s_{N_m} = 0\}).$$

We set $\delta_m(X, \Delta) := \min\{\text{lct}(X, \Delta; D) \mid D \sim_{\mathbb{Q}} -K_X - \Delta \text{ is } m\text{-basis type}\}$. The δ -invariant (also known as the *stability threshold*) of (X, Δ) is

$$\delta(X, \Delta) = \limsup_{m \rightarrow \infty} \delta_{mr}(X, \Delta),$$

and is independent of the choice of r [BJ17, 4.5]. The invariant may also be calculated in terms of valuations or filtrations.

Theorem 2.7 ([BJ17, Theorems A,C]). *We have*

$$\delta(X, \Delta) = \inf_E \frac{A_{X, \Delta}(E)}{S(E)} = \inf_v \frac{A_{X, \Delta}(v)}{S(v)},$$

where the first infimum runs through all divisors E over X and the second through all $v \in \text{Val}_X^*$ with $A_{X, \Delta}(v) < +\infty$. Furthermore, the limit $\lim_{m \rightarrow \infty} \delta_{mr}(X, \Delta)$ exists.

Proposition 2.8 ([BL18, Proposition 4.10]). *We have*

$$\delta(X, \Delta) = \inf_{\mathcal{F}} \frac{\text{lct}(X, \Delta; \mathfrak{b}_{\bullet}(\mathcal{F}))}{S(\mathcal{F})}$$

where the infimum runs through all non-trivial linearly bounded filtrations of $R(X, L)$.

Combining Definition 2.4 and Theorem 2.7, we immediately see

Theorem 2.9 ([FO18, BJ17]). *A log Fano pair (X, Δ) is uniformly K-stable (resp., K-semistable) if and only if $\delta(X, \Delta) > 1$ (resp., ≥ 1).*

While in Section 3 we will use the definition of the δ -invariant in terms of m -basis type divisors, in Section 5 we will rely on its characterization in terms of valuations and filtrations.

2.5. Normalized volume. Now, we discuss an invariant similar to the δ -invariant, but defined in a local setting. The invariant was first introduced in [Li18] and is closely related to the K-semistability of log Fano pairs.

Let (X, Δ) be an n -dimensional klt pair and $x \in X$ a closed point. The *non-archimedean link* of X at x is defined as

$$\text{Val}_{X,x} := \{v \in \text{Val}_X \mid c_X(v) = x\} \subset \text{Val}_X.$$

Definition 2.10 ([Li18]). The *normalized volume function* $\widehat{\text{vol}}_{(X, \Delta), x} : \text{Val}_{X,x} \rightarrow (0, +\infty]$ is defined by

$$\widehat{\text{vol}}_{(X, \Delta), x}(v) = \begin{cases} A_{(X, \Delta)}(v)^n \cdot \text{vol}(v) & \text{if } A_{(X, \Delta)}(v) < +\infty; \\ +\infty & \text{if } A_{(X, \Delta)}(v) = +\infty. \end{cases}$$

The *volume of the singularity* ($x \in (X, \Delta)$) is defined as

$$\widehat{\text{vol}}(x, X, \Delta) := \inf_{v \in \text{Val}_{X,x}} \widehat{\text{vol}}_{(X, \Delta), x}(v).$$

The previous infimum is a minimum by the main result in [Blu18a].

See [LLX18] for a survey on the recent study of the normalized volume function, especially the guiding question, the Stable Degeneration Conjecture (see [Li18, 7.1] and [LLX18, 4.1]).

2.5.1. Relation to K-stability. The connection between the normalized volume function and K-semistability is via the cone construction first studied in [Li17].

Let (X, Δ) be a log Fano pair and r a positive integer so that $L := -r(K_X + \Delta)$ is a Cartier divisor. Let (Z, Γ) denote the cone over X with respect to the polarization L and $x \in Z$ denote the vertex. Specifically, $Z = \text{Spec}(R)$, where $R = R(X, L)$ and Γ is the closure of the pullback of Δ via the projection map $Z \setminus \{x\} \rightarrow X$.

There is a natural map $X_L \rightarrow Z$, where $X_L := \text{Spec}_X(\bigoplus_{p \geq 0} \mathcal{O}_X(pL))$ is the total space of the line bundle on X whose sheaf of sections is $\mathcal{O}_X(L)$. The map is an isomorphism over $Z \setminus x$ and the preimage of the vertex is the zero section $X_{\text{zs}} \subset X_L$. We call $v_0 := \text{ord}_{X_{\text{zs}}}$ the *canonical valuation* over the cone.

Theorem 2.11 ([Li17, LL19, LX16]). *The canonical valuation v_0 is a minimizer of $\widehat{\text{vol}}_{(Z, \Gamma), x}$ if and only if (X, Δ) is K-semistable.*

At first sight, using the normalized volume function to study the K-stability of log Fano pairs may seem indirect. However, this approach yields a number of new results (for example, see [LX16, LWX18b]). In this paper, the following key ingredient in the proof of Theorem 2.11 plays an important role in the proof of our main result.

Following [Li17, LX16], let E be a divisor over X that arises on a proper normal model $\mu : Y \rightarrow X$. Consider the natural birational maps

$$Y_L \rightarrow X_L \rightarrow Z,$$

where $Y_L := \text{Spec}_Y(\bigoplus_{m \geq 0} \mathcal{O}_Y(m\mu^*L))$. Let $Y_{\text{zs}} \subset Y_L$ denote the zero section and E_∞ the preimage of E under the projection $Y_L \rightarrow Y$. Setting v_t equal to the quasimonomial valuation with weights $(1, t)$ along Y_{zs} and E_∞ gives a ray of valuations

$$\{v_t \mid t \in [0, \infty)\} \subset \text{Val}_{Z,x},$$

where $v_0 = \text{ord}_{X_{\text{zs}}}$ and $v_\infty = \text{ord}_{E_\infty}$. When $k \in \mathbb{N}$, there exists a divisor E_k over Z satisfying $v_{\frac{1}{k}} = \frac{1}{k} \text{ord}_{E_k}$.

By the formula for the log discrepancy of a quasimonomial valuation [JM12, 5.1],

$$A_{Z,\Gamma}(v_t) = A_{Z,\Gamma}(\text{ord}_{X_{\text{zs}}}) + tA_{Z,\Gamma}(\text{ord}_{E_\infty}) = r^{-1} + tA_{X,\Delta}(\text{ord}_E).$$

The valuation ideals are given by, for $t > 0$,

$$\mathfrak{a}_p(v_t) = \bigoplus_{m \geq 0} \mathcal{F}_E^{(p-m)/t} R_m \subseteq R \quad \text{and} \quad \mathfrak{a}_p(v_0) = \bigoplus_{m \geq p} R_m \subseteq R.$$

To see the previous formula holds, fix a uniformizer $\omega \in \mathcal{O}_{Y,E}$ and a local section s of $\mathcal{O}_Y(\mu^*L)$ that trivializes the sheaf at the generic point of E . The choice of s induces a rational map $Y_L \dashrightarrow Y \times \mathbb{A}^1$ that is an isomorphism at the generic point of $Y_{\text{zs}} \cap E_\infty$. The birational transforms of Y_{zs} and E_∞ on $Y \times \mathbb{A}^1$ are $Y \times 0$ and $E \times \mathbb{A}^1$.

Fix $f = \sum_{m \geq 0} f_m \in R$. For m such that $f_m \neq 0$, set $g_m := f_m / \omega^{\text{ord}_E(f_m)}$, which, at the generic point of E , is a non-vanishing section of $\mathcal{O}_Y(\mu^*L)$. The image of f in $\mathcal{O}_{Y \times \mathbb{A}^1, E \times 0}$ equals $\sum_m \binom{g_m}{s^m} \omega^{\text{ord}_E(f_m)} \pi^m$, where π is the parameter for \mathbb{A}^1 . Since π and ω are local equations for $Y \times 0$ and $E \times \mathbb{A}^1$ at the generic point of $E \times 0$ and $\frac{g_m}{s^m}$ does not vanish at $E \times 0$,

$$v_t(f) := \min_m \{t \cdot \text{ord}_E(f_m) + m \mid f_m \neq 0\}$$

and the formula for $\mathfrak{a}_p(v_t)$ follows.

By the calculation in [Li17, (31, 32)] (see also the proof of [LX16, 4.5] or Lemma 2.12)

$$\frac{d}{dt} \widehat{\text{vol}}(v_t) \Big|_{t=0^+} = (n+1)\beta_{X,\Delta}(E) \tag{6}$$

This equation is a key input in our proof of Theorem 4.1. More specifically, we will follow ideas from [LWX18b] and analyze directions along which the normalized volume function has derivative equal to zero.

2.5.2. C. Li's derivative formula. In the proof of Theorem 1.1, we will need a more general version of (6). The more general formula follows from the original argument in [Li17].

Let (X, Δ) be an n -dimensional log Fano pair and $r \in \mathbb{Z}_{>0}$ so that $L := -r(K_X + \Delta)$ is Cartier. Set $R = R(X, L)$ and fix a linearly bounded filtration \mathcal{F} of R .

Associated to \mathcal{F} , we define a collection of graded sequences of ideals of R . For $t \in \mathbb{R}_{>0}$ and $j \in \mathbb{Z}_{>0}$, set

$$\mathfrak{b}_{t,j} := \bigoplus_{m \geq 0} \mathcal{F}^{(j-m)/t} R_m \subseteq R \quad \text{and} \quad \mathfrak{b}_{0,j} := \bigoplus_{m \geq j} R_m \subseteq R.$$

Note that $\mathfrak{b}_{t,\bullet}$ is a graded sequence of ideals of R for each $t \in \mathbb{R}_{\geq 0}$. Additionally, $\mathfrak{b}_{t,j}$ contains $\oplus_{m \geq j} R_m$.

Lemma 2.12. *With the above notation, fix $A > 0$ and set*

$$f(t) = (r^{-1} + At)^{n+1} \text{mult}(\mathfrak{b}_{t,\bullet})$$

for $t \in \mathbb{R}_{\geq 0}$. The following hold:

- (1) $\text{mult}(\mathfrak{b}_{t,\bullet}) = r^n(-K_X - \Delta)^n - (n+1) \int_0^\infty \text{vol}(\mathcal{F}R^{(x)}) \frac{t dx}{(1+tx)^{n+2}}$;
- (2) $\frac{df}{dt} \Big|_{t=0^+} = (n+1)(-K_X - \Delta)^n (A - S(\mathcal{F}))$.

Proof. This follows from the argument in [Li17, (18)-(25)]. For the reader's convenience, we give a brief proof. For $t \in \mathbb{R}_{>0}$, we have

$$\begin{aligned} \text{mult}(\mathfrak{b}_{t,\bullet}) &= \lim_{j \rightarrow \infty} \frac{(n+1)!}{j^{n+1}} \dim(R/\mathfrak{b}_{t,j}) \\ &= \lim_{j \rightarrow \infty} \frac{(n+1)!}{j^{n+1}} \sum_{m=0}^{\infty} \dim(R_m/\mathcal{F}^{(j-m)/t} R_m) \\ &= \lim_{j \rightarrow \infty} \frac{(n+1)!}{j^{n+1}} \sum_{m=0}^j (\dim R_m - \dim \mathcal{F}^{(j-m)/t} R_m) \\ &= \text{vol}(L) - \lim_{j \rightarrow \infty} \frac{(n+1)!}{j^{n+1}} \sum_{m=0}^j \dim \mathcal{F}^{(j-m)/t} R_m. \end{aligned}$$

Statement (1) now follows from [Li17, (25)], where $c_1 = 0$, $\alpha = \beta = \frac{1}{t}$.

For (2), compute

$$\frac{df}{dt} \Big|_{t=0^+} = (n+1)Ar^{-n} \text{mult}(\mathfrak{b}_{0,\bullet}) + r^{-n-1} \frac{d}{dt} (\text{mult}(\mathfrak{b}_{t,\bullet})) \Big|_{t=0^+}.$$

From (1), we know $\text{mult}(\mathfrak{b}_{0,\bullet}) = r^n(-K_X - \Delta)^n$ and

$$\frac{d}{dt} (\text{mult}(\mathfrak{b}_{t,\bullet})) \Big|_{t=0^+} = -(n+1) \int_0^\infty \left(\text{vol}(\mathcal{F}R^{(x)}) \left(\frac{1 - (n+1)tx}{(1+tx)^{n+3}} \right) \right) dx.$$

Since the latter simplifies to $-(n+1) \int_0^\infty \text{vol}(\mathcal{F}R^{(x)}) dx$, (2) is complete. \square

3. UNIFORMLY K-STABLE FANOS

In this section, we prove a special case of Theorem 1.1 for uniformly K-stable Fano varieties. We will then apply the result to study the moduli functor $\mathcal{M}_{n,V}^{\text{uKs}}$.

3.1. Separatedness result. The following result is a special case of Theorem 1.1 and will be reproved in Section 5. We present its proof independently, since the following argument is simpler than the proof in Section 5.

Theorem 3.1. *Let $\pi : (X, \Delta) \rightarrow C$ and $\pi' : (X', \Delta') \rightarrow C$ be \mathbb{Q} -Gorenstein families of log Fano pairs over a smooth pointed curve $0 \in C$. Assume there exists an isomorphism*

$$\phi : (X, \Delta) \times_C C^\circ \rightarrow (X', \Delta') \times_C C^\circ$$

over $C^\circ := C \setminus 0$. If (X_0, Δ_0) is uniformly K -stable and (X'_0, Δ'_0) is K -semistable, then ϕ extends to an isomorphism $(X, \Delta) \simeq (X', \Delta')$ over C .

The proof of Theorem 3.1 follows from properties of the δ -invariant and the following birational geometry fact.

Proposition 3.2. *Let $\pi : (X, \Delta) \rightarrow C$ and $\pi' : (X', \Delta') \rightarrow C$ be \mathbb{Q} -Gorenstein families of log Fano pairs over a smooth pointed curve $0 \in C$. Assume there exists an isomorphism*

$$\phi : (X, \Delta) \times_C C^\circ \rightarrow (X', \Delta') \times_C C^\circ$$

over $C^\circ := C \setminus 0$. If there exist effective horizontal³ \mathbb{Q} -divisors D and D' on X and X' satisfying

- (1) $D \sim_{\mathbb{Q}, C} -K_X - \Delta$ and $D' \sim_{\mathbb{Q}, C} -K_{X'} - \Delta'$,
- (2) D is the birational transform of D' , and
- (3) $(X_0, \Delta_0 + D_0)$ is klt and $(X'_0, \Delta'_0 + D'_0)$ is lc,

then ϕ extends to an isomorphism $(X, \Delta) \simeq (X', \Delta')$ over C .

The above proposition is well known to experts and follows from the separatedness of the moduli functor of klt log Calabi-Yau pairs (e.g. see [Oda12, Theorem 4.3], [LWX19, Theorem 5.2]). For the convenience of the reader, we prove the result.

Proof. Fix a common log resolution \widehat{X} of (X, Δ) and (X', Δ')

$$\begin{array}{ccc} & \widehat{X} & \\ \psi \swarrow & & \searrow \psi' \\ X & \xrightarrow{\phi} & X' \end{array}$$

and write \widetilde{X}_0 and \widetilde{X}'_0 for the birational transforms of X_0 and X'_0 on \widehat{X} .

First, assume $\widetilde{X}_0 = \widetilde{X}'_0$. This equality implies $\phi : X \dashrightarrow X'$ is an isomorphism in codimension one. Thus, ϕ induces an isomorphism

$$\pi_* \mathcal{O}_X(-m(K_X + \Delta)) \simeq \pi'_* \mathcal{O}_{X'}(-m(K_{X'} + \Delta'))$$

for all $m \in \mathbb{N}$. Since

$$X = \text{Proj}_C \bigoplus_{m \geq 0} \pi_* \mathcal{O}_X(-m(K_X + \Delta)) \text{ and } X' = \text{Proj}_C \bigoplus_{m \geq 0} \pi'_* \mathcal{O}_{X'}(-m(K_{X'} + \Delta')),$$

we conclude ϕ extends to an isomorphism over C .

³A \mathbb{Q} -divisor on X or X' is called *horizontal* if its support does not contain a fiber of the map to C .

We now assume $\tilde{X}_0 \neq \tilde{X}'_0$ and aim for a contradiction. Write

$$K_{\tilde{X}} + \psi_*^{-1}(\Delta + D) = \psi^*(K_X + \Delta + D) + a\tilde{X}'_0 + P$$

and

$$K_{\tilde{X}} + \psi'^{-1}(\Delta' + D') = \psi'^*(K_{X'} + \Delta' + D') + a'\tilde{X}_0 + P'$$

where the components of $\text{Supp}(P) \cup \text{Supp}(P')$ are both ψ and ψ' -exceptional. By assumption (2), $P - P'$ is supported on \tilde{X}_0 .

Inversion of adjunction and our assumption that $(X_0, \Delta_0 + D_0)$ is klt imply $(X, \Delta + D + X_0)$ is plt in a neighborhood of X_0 . Hence,

$$-1 < a(\tilde{X}'_0, X, \Delta + D + X_0) = a - \text{ord}_{\tilde{X}'_0}(X_0).$$

Since $\text{ord}_{\tilde{X}'_0}(X_0) = 1$, we see $a > 0$. The same argument, but with the assumption that $(X'_0, \Delta'_0 + D'_0)$ is lc implies $a' \geq 0$.

Observe $a\tilde{X}'_0 - a'\tilde{X}_0 + (P - P') \sim_{\mathbb{Q}, C} 0$, since $K_X + \Delta + D \sim_{\mathbb{Q}, C} 0$, $K_{X'} + \Delta' + D' \sim_{\mathbb{Q}, C} 0$, and $\psi_*^{-1}(D + \Delta) = \psi'^{-1}(D' + \Delta')$. Therefore, there exists a rational number c so that

$$a\tilde{X}'_0 - a'\tilde{X}_0 + (P - P') \sim_{\mathbb{Q}, C} c\psi^*(X_0).$$

Comparing the coefficients of \tilde{X}'_0 on the two sides implies $c > 0$, while comparing the coefficients of \tilde{X}_0 implies $c \leq 0$. This is a contradiction. \square

Lemma 3.3. *Keep the notation and setup of Theorem 3.1. If $m \in \mathbb{Z}_{>0}$ is sufficiently divisible, then there exist effective horizontal \mathbb{Q} -divisors B and B' on X and X' such that*

- (1) $B \sim_{\mathbb{Q}, C} -K_X - \Delta$ and $B' \sim_{\mathbb{Q}, C} -K_{X'} - \Delta'$,
- (2) B is the birational transform of B' , and
- (3) B_0 and B'_0 are m -basis type with respect to (X_0, Δ_0) and (X'_0, Δ'_0) .

Proof. Fix a positive integer m so that $L := -m(K_X - \Delta)$ and $L' := -m(K_{X'} - \Delta')$ are Cartier and $\pi_*\mathcal{O}_X(L)$ and $\pi'_*\mathcal{O}_{X'}(L)$ are nonzero. Since $H^i(X_t, \mathcal{O}_{X_t}(L_t))$ and $H^i(X'_t, \mathcal{O}_{X'_t}(L'_t))$ are zero for all $i > 0$ and $t \in C$ by Kawamata-Viehweg vanishing, $\pi_*\mathcal{O}_X(L)$ and $\pi'_*\mathcal{O}_{X'}(L')$ are vector bundles. Furthermore, the sheaves satisfy cohomology and base change.

Now, the birational map ϕ induces a map from local sections of $\pi_*\mathcal{O}_X(L)$ to rational sections of $\pi'_*\mathcal{O}_{X'}(L')$. After twisting by dX'_0 , where $d \gg 0$, we get a morphism

$$\pi_*\mathcal{O}_X(L) \rightarrow \pi'_*\mathcal{O}_{X'}(L' + dX'_0)$$

that is an isomorphism away from $0 \in C$. Tensoring by $\mathcal{O}_{C,0}$ gives a morphism

$$\varphi : \pi_*\mathcal{O}_X(L) \otimes_{\mathcal{O}_C} \mathcal{O}_{C,0} \rightarrow \pi'_*\mathcal{O}_{X'}(L' + dX'_0) \otimes_{\mathcal{O}_C} \mathcal{O}_{C,0},$$

of locally free $\mathcal{O}_{C,0}$ modules that is an isomorphism after tensoring with $K(C)$.

Write t for the uniformizer of $\mathcal{O}_{C,0}$. Since $\mathcal{O}_{C,0}$ is a principal ideal domain, there exist bases $\{s_1, \dots, s_N\}$ and $\{s'_1, \dots, s'_N\}$ for the above free modules so that the transformation matrix is diagonal. Hence, for each $1 \leq i \leq N$, there exist $p_i \in \mathbb{Z}_{\geq 0}$ and $a_i \in \mathcal{O}_{C,0}^\times$ so that $\varphi(s_i) = a_i t^{p_i} s'_i$.

For a sufficiently small neighborhood $0 \in U \subset C$, we may extend each s_i to a section $\tilde{s}_i \in \pi_* \mathcal{O}_X(L)(U)$ and s'_i to a section $\tilde{s}'_i \in \pi'_* \mathcal{O}_{X'}(L')(U)$. Let B and B' denote the closures of

$$\frac{1}{mN} (\{\tilde{s}_1 = 0\} + \dots + \{\tilde{s}_N = 0\}) \text{ and } \frac{1}{mN} (\{\tilde{s}'_1 = 0\} + \dots + \{\tilde{s}'_N = 0\}).$$

in X and X' . By construction, B_0 and B'_0 are both m -basis type divisors and B is the birational transform of B' . \square

Proof of Theorem 3.1. Since X_0 is uniformly K-stable and X'_0 is K-semistable,

$$\delta(X_0, \Delta_0) > 1 \quad \text{and} \quad \delta(X'_0, \Delta'_0) \geq 1.$$

Hence, we may choose $0 < \varepsilon \ll 1$ so that

$$\frac{1 - \varepsilon}{\delta(X_0, \Delta_0)} + \frac{\varepsilon}{\alpha(X_0, \Delta_0)} < 1, \tag{7}$$

where $\alpha(X_0, \Delta_0)$ is Tian's α -invariant, i.e.

$$\alpha(X_0, \Delta_0) = \inf \{ \text{lct}(X_0, \Delta_0; D) \mid 0 \leq D \sim_{\mathbb{Q}} -K_{X_0} - \Delta_0 \}.$$

Next, choose a positive integer M so that

$$\frac{1 - \varepsilon}{\delta_m(X_0, \Delta_0)} + \frac{\varepsilon}{\alpha(X_0, \Delta_0)} < 1 \text{ and } \delta_m(X'_0, \Delta'_0) > 1 - \varepsilon \tag{8}$$

for all positive integers m divisible by M . Such a choice is possible by (7), the inequality $\delta(X'_0, \Delta'_0) \geq 1$, and the fact that δ is a limit.

Now, fix a positive integer m divisible by M so that the conclusion of Lemma 3.3 holds for m and $-m(K_{X'} + \Delta')$ is relatively base point free over C . Hence, we may find \mathbb{Q} -divisors $B \sim_{\mathbb{Q}, C} -K_X - \Delta$ and $B' \sim_{\mathbb{Q}, C} -K_{X'} - \Delta'$ satisfying the conclusion of Lemma 3.3 for m . Since B_0 and B'_0 are m -basis type,

$$\text{lct}(X_0, \Delta_0; B_0) \geq \delta_m(X_0, \Delta_0) \quad \text{and} \quad \text{lct}(X'_0, \Delta'_0; B'_0) \geq \delta_m(X'_0, \Delta'_0) > 1 - \varepsilon.$$

The latter implies $(X'_0, \Delta'_0 + (1 - \varepsilon)B'_0)$ is lc.

Since $-m(K_{X'} + \Delta')$ is relatively base point free over C , after shrinking C in a neighborhood of 0 , we may apply [KM98, Lemma 5.17] to find an effective divisor $G' \in |-m(K_{X'} + \Delta')|$ in general position so that $(X'_0, \Delta'_0 + (1 - \varepsilon)B'_0 + (\varepsilon/m)G'_0)$ remains lc. Write G for the birational transform of G' on X . Note that $G \sim_{\mathbb{Q}, C} -m(K_X + \Delta)$, since the statement holds over C° . Thus, $\text{lct}(X_0, \Delta_0; (1/m)G_0) \geq \alpha(X_0, \Delta_0)$.

Now, consider the divisors

$$D := (1 - \varepsilon)B + \frac{\varepsilon}{m}G \quad \text{and} \quad D' := (1 - \varepsilon)B' + \frac{\varepsilon}{m}G'.$$

Observe that $D \sim_{\mathbb{Q},C} -K_X - \Delta$ and $D' \sim_{\mathbb{Q},C} -K_{X'} - \Delta'$. As mentioned above, $(X_0, \Delta'_0 + D'_0)$ is lc. Additionally, the pair $(X_0, \Delta_0 + D_0)$ is klt. Indeed, since

$$1/\text{lct}(D + F) \leq 1/\text{lct}(D) + 1/\text{lct}(F)$$

for any two effective \mathbb{Q} -Cartier \mathbb{Q} -divisors D and F on a klt pair, we know

$$\begin{aligned} \frac{1}{\text{lct}(X_0, \Delta_0; D_0)} &\leq \frac{1}{\text{lct}(X_0, \Delta_0; (1 - \varepsilon)B_0)} + \frac{1}{\text{lct}(X_0, \Delta_0; (\varepsilon/m)G_0)} \\ &\leq \frac{1 - \varepsilon}{\delta_m(X_0, \Delta_0)} + \frac{\varepsilon}{\alpha(X_0, \Delta_0)} \end{aligned}$$

which is < 1 by (8). Proposition 3.2 now implies ϕ extends to an isomorphism. \square

Remark 3.4. If (X_0, Δ_0) and (X'_0, Δ'_0) are only assumed to be K-semistable, then they are not necessarily isomorphic (but are S-equivalent by Theorem 1.1). Therefore, we do not expect the the above strategy to be useful in this more general case.

Recall, if (X, Δ) is a log Fano pair, then $\text{Aut}(X, \Delta)$ is the closed subgroup of $\text{Aut}(X)$ defined by

$$\text{Aut}(X, \Delta) := \{g \in \text{Aut}(X) \mid g^*\Delta = \Delta\}.$$

The following result is an immediate corollary of Theorem 3.1 and a special case of Corollary 1.3.

Corollary 3.5. *Let (X, Δ) be a log Fano pair. If (X, Δ) is uniformly K-stable, then $\text{Aut}(X, \Delta)$ is finite.*

Proof. Since $\text{Aut}(X, \Delta)$ is a linear algebraic group, it is affine. To conclude that $\text{Aut}(X, \Delta)$ is finite, it suffices to show that it is proper. To see the properness, consider a map $g : C^\circ \rightarrow \text{Aut}(X, \Delta)$, where $0 \in C$ is a smooth pointed curve and $C^\circ = C \setminus 0$. The map g induces an isomorphism

$$(X \times C, \Delta \times C) \times_C C^\circ \rightarrow (X \times C, \Delta \times C) \times_C C^\circ$$

over C° . By applying Theorem 3.1 to the above isomorphism, we see f extends to a map $\bar{g} : C \rightarrow \text{Aut}(X, \Delta)$. Hence, $\text{Aut}(X, \Delta)$ is proper, and the proof is complete. \square

In [BHJ16, Corollary E], it is shown that the polarized automorphism group of a uniformly K-stable polarized manifold (X, L) is finite. Their proof uses analytic tools.

Remark 3.6. Our proofs of Theorem 3.1 and Corollary 3.5 extend to the case of polarized klt pairs $(X, \Delta; L)$ (that is, (X, Δ) is a projective klt pair and L an ample \mathbb{Q} -Cartier divisor on X) such that $K_X + \Delta + L$ is nef and $\delta(X, \Delta; L) > 1$.

3.2. Moduli spaces.

Proof of Corollary 1.4. As previously mentioned, the result relies on [Jia17, BL18] and Theorem 3.1. Indeed, [Jia17] (see also [Che18] or [LLX18, 6.14]) states that the set of varieties $\mathcal{M}_{n,V}^{\text{uKs}}(k)$ is bounded. Hence, there exists a positive integer M so that

$-MK_X$ is a very ample Cartier divisor for all $X \in \mathcal{M}_{n,V}^{\text{uKs}}(k)$. Furthermore, the set of Hilbert functions $m \mapsto \chi(\omega_X^{[-mM]})$ with $X \in \mathcal{M}_{n,V}^{\text{uKs}}(k)$ is finite.

For such a Hilbert function h , consider the subfunctor $\mathcal{M}_h^{\text{uKs}} \subset \mathcal{M}_{n,V}^{\text{uKs}}$ parameterizing uniformly K-semistable \mathbb{Q} -Fano varieties with Hilbert function h . Note that $\mathcal{M}_{n,V}^{\text{uKs}} = \coprod_h \mathcal{M}_h^{\text{uKs}}$. Set $N := h(1) - 1$, and let $\text{Hilb}(\mathbb{P}^N)$ be the Hilbert scheme parameterizing closed subschemes of \mathbb{P}^N with Hilbert polynomial h . Write $X \rightarrow \text{Hilb}(\mathbb{P}^N)$ for the corresponding universal family.

Now, let $U \subset \text{Hilb}(\mathbb{P}^N)$ denote the open subscheme parameterizing normal, Cohen-Macaulay varieties. By [HK04, 3.11], there is a locally closed subscheme $V \subset U$ such that a map $T \rightarrow U$ factors through V if and only if there is an isomorphism $\omega_{X_T/T}^{[-M]} \simeq \mathcal{L}_T \otimes \mathcal{O}_{X_T}(1)$, where \mathcal{L}_T is the pullback of a line bundle from T . By applying [BL18] to the normalization of V , we see

$$V' := \{t \in V \mid X_{\bar{t}} \text{ is a uniformly K-stable } \mathbb{Q}\text{-Fano variety}\}$$

is open in V . Finally, we apply [Kol09, 25] or [AH11] to find a locally closed decomposition $W \rightarrow V'$ such that a morphism $T \rightarrow V'$ factors through W if and only if $X_T \rightarrow X$ satisfies Kollar's condition.

As a consequence of the above discussion, $\mathcal{M}_h^{\text{uKs}} \simeq [W/\text{PGL}(N+1)]$. Theorem 3.1 implies $\mathcal{M}_h^{\text{uKs}}$ is a separated Deligne-Mumford stack. Furthermore, we may apply [KM97] to see $\mathcal{M}_h^{\text{uKs}}$ has a coarse moduli space M_h^{uKs} , which is a separated algebraic space. \square

4. PLACES COMPUTING THE δ -INVARIANT

In this section, we will study the cases when valuations, ideals, and \mathbb{Q} -divisors compute the δ -invariant. The results proved here are related to Conjecture 1.5 and will be used in the proof of Theorem 1.1.

4.1. Divisors computing δ .

Theorem 4.1. *Let (X, Δ) be a K-semistable log Fano pair. If E is a divisor over X satisfying $\delta(X, \Delta) = \frac{A_{X, \Delta}(E)}{S(E)} = 1$, then E is dreamy and induces a non-trivial special test configuration $(\mathcal{X}, \mathcal{D})$ such that $\text{Fut}(\mathcal{X}, \mathcal{D}) = 0$. In particular, (X, Δ) is not K-stable.*

The proof follows an argument in [LWX18b, Section 3.1]. The argument will be used again in the proof of Lemma 5.10 in a relative setting.

Proof. Fix a positive integer r so that $L := -r(K_X + \Delta)$ is a Cartier divisor and set $R = R(X, L)$. Consider the cone (Z, Γ) over (X, Δ) with respect to the polarization L .

The divisor E over X induces a ray of valuations

$$\{v_t \mid t \in [0, \infty)\} \subset \text{Val}_{Z,x}$$

(see Section 2.5.1). For $k \in \mathbb{Z}_{>0}$, there is a divisor E_k over Z so that $v_{\frac{1}{k}} = \frac{1}{k} \text{ord}_{E_k}$. By (6),

$$\frac{d}{dt} \widehat{\text{vol}}(v_t) \Big|_{t=0^+} = (n+1)\beta_{X,\Delta}(E).$$

Since $A_{X,\Delta}(E) - S(E) = 0$, we know $\beta_{X,\Delta}(E) = 0$. Defining $f(t) := \widehat{\text{vol}}(v_t)$, a Taylor expansion gives

$$f(t) = f(0) + O(t^2) \quad \text{for } 0 \leq t \ll 1.$$

For a fixed positive integer k , set

$$\mathfrak{a}_{k,\bullet} := \mathfrak{a}_\bullet(\text{ord}_{E_k}) \quad \text{and} \quad c_k := \text{lct}(Z, \Gamma; \mathfrak{a}_{k,\bullet}).$$

Note that $c_k \leq A_{Z,\Gamma}(E_k)$ by (3). This implies

$$f(0) \leq c_k^{n+1} \cdot \text{mult}(\mathfrak{a}_{k,\bullet}) \leq f\left(\frac{1}{k}\right) = A_{Z,\Gamma}(E_k)^{n+1} \cdot \text{mult}(\mathfrak{a}_{k,\bullet}),$$

where the first inequality follows from [Liu18, 7] and the assumption that (X, Δ) is K-semistable. Therefore,

$$\left(\frac{1}{1+O(1/k^2)}\right)^{\frac{1}{n+1}} \leq \left(\frac{f(0)}{f(1/k)}\right)^{\frac{1}{n+1}} \leq \frac{c_k}{A_{Z,\Gamma}(E_k)} \leq 1.$$

Since $(1+O(1/k^2))^{1/(n+1)}$ is of the order $1+O(1/k^2)$, we see

$$1 - O\left(\frac{1}{k^2}\right) = \frac{c_k}{A_{Z,\Gamma}(E_k)} \leq 1.$$

Using that $A_{Z,\Gamma}(E_k) = kr^{-1} + A_{X,\Delta}(E)$,

$$\lim_{k \rightarrow \infty} (A_{Z,\Gamma}(E_k) - c_k) = \lim_{k \rightarrow \infty} \left(A_{Z,\Gamma}(E_k) \left(1 - \frac{c_k}{A_{Z,\Gamma}(E_k)}\right) \right) = 0.$$

Hence, we may fix $k \gg 0$ so that $A_{Z,\Gamma}(E_k) - c_k < 1$.

By Proposition 2.2, there exists a proper birational morphism $\mu_k: Z_k \rightarrow Z$ such that $E_k \subset Z_k$ and $-E_k$ is ample over Z . Therefore, $\bigoplus_{p \geq 0} \mu_{k*} \mathcal{O}_{Z_k}(-pE_k)$ is a finitely generated \mathcal{O}_Z -algebra. Since $\mu_{k*}(\mathcal{O}_{Z_k}(-pE_k)) = \mathfrak{a}_p(kv_{1/k})$, the latter implies

$$\bigoplus_{p \in \mathbb{N}} \bigoplus_{m \in \mathbb{N}} \mathcal{F}_E^{p-mk} R_m$$

is a finitely generated R -algebra. Therefore, $\text{Rees}(\mathcal{F}_E)$ is finitely generated as well and E is dreamy.

Let $(\mathcal{X}, \mathcal{D})$ denote the test configuration induced by \mathcal{F}_E . The test configuration is normal and non-trivial [Fuj17, 3.8] and $\text{Fut}(\mathcal{X}, \mathcal{D})$ is a multiple of $A_{X,\Delta}(E) - S(E)$ [Fuj19a, 6.12], which is zero. We conclude $(\mathcal{X}, \mathcal{D})$ is special, since otherwise there would exist a test configuration of (X, Δ) with negative Futaki invariant [LX14, 1]. \square

An immediate corollary to Theorem 4.1 is the following strengthening of [Fuj19a, 1.6] and [Li18, 3.7]. The result was expected in the arXiv version of [Li18].

Corollary 4.2. *A log Fano pair (X, Δ) is K-stable if and only if $\beta_{X, \Delta}(E) > 0$ for any divisor E over X .*

Proof. Theorem 4.1 implies the forward implication. The reverse implication was shown in [Fuj19a, 1.6] and [Li18, 3.7]. \square

4.2. Ideals computing δ . Let (X, Δ) be a log Fano pair and $\mathfrak{a} \subsetneq \mathcal{O}_X$ a nonzero ideal. Write $\pi : Y \rightarrow X$ for the normalized blowup of X along \mathfrak{a} and E for the effective Cartier divisor on Y such that $\mathfrak{a} \cdot \mathcal{O}_Y = \mathcal{O}_Y(-E)$. We set

$$S(\mathfrak{a}) := \frac{1}{\text{vol}(-K_X - \Delta)} \int_0^{+\infty} \text{vol}(\pi^*(-K_X - \Delta) - tE) dt.$$

Proposition 4.3. *If (X, Δ) is a log Fano pair and $\mathfrak{a} \subsetneq \mathcal{O}_X$ a nonzero ideal, then*

$$\frac{\text{lct}(X, \Delta; \mathfrak{a})}{S(\mathfrak{a})} \geq \delta(X, \Delta). \quad (9)$$

Furthermore, write $\pi : Y \rightarrow X$ for the normalized blowup of \mathfrak{a} and E for the Cartier divisor on Y such that $\mathfrak{a} \cdot \mathcal{O}_Y = \mathcal{O}_Y(-E)$. If (9) is an equality, then $\text{Supp}(E)$ is a prime divisor and computes $\delta(X, \Delta)$.

The above proposition is an analog of [LX16, Theorem 3.11] for the δ -invariant and is similar to [Fuj19b, Corollary 3.22].

Proof. Choose a divisor F over X computing $\text{lct}(X, \Delta; \mathfrak{a})$. By [BCHM10], there is an extraction $\rho : X_F \rightarrow X$ of F . Set $p := \text{ord}_F(\mathfrak{a})$. Hence, $A_{X, \Delta}(F)/p = \text{lct}(X, \Delta; \mathfrak{a})$ and $\mathfrak{a}^k \cdot \mathcal{O}_{X_F} \subseteq \mathcal{O}_{X_F}(-kpF)$ for all $k \in \mathbb{N}$.

By the previous inclusion, if we set $L := -K_X - \Delta$, then

$$\text{vol}(\pi^*L - tE) \leq \text{vol}(\rho^*L - tpF)$$

for all $t \in \mathbb{R}_{\geq 0}$. Hence, $S(\mathfrak{a}) \leq p^{-1}S(F)$, and we see

$$\frac{\text{lct}(\mathfrak{a})}{S(\mathfrak{a})} \geq \frac{A_{X, \Delta}(F)}{S(F)}.$$

Since $A_{X, \Delta}(F)/S(F) \geq \delta(X, \Delta)$, (9) holds.

Now assume (9) is an equality. In this case, the above argument implies F computes $\delta(X, \Delta)$. To finish the proof, it suffices to show $Y = X_F$ and $\text{Supp}(E) = F$.

Fix a positive integer k so that $-kpF$ is Cartier and choose an ideal $\mathfrak{c} \subseteq \mathcal{O}_{X_F}$ such that

$$\mathfrak{a}^k \cdot \mathcal{O}_{X_F} = \mathfrak{c} \cdot \mathcal{O}_{X_F}(-pkF).$$

Write $\tau : Z \rightarrow X_F$ for the normalized blowup of X_F along \mathfrak{c} and G for the Cartier divisor on Z such that $\mathfrak{c} \cdot \mathcal{O}_Z = \mathcal{O}_Z(-G)$. Since Z is normal and

$$\mathfrak{a}^k \cdot \mathcal{O}_Z = (\mathfrak{c} \cdot \mathcal{O}_{X_F}(-pkF)) \cdot \mathcal{O}_Z = \mathcal{O}_Z(-pk\tau^*(F) - G)$$

is locally free, $\rho \circ \tau$ factors through π :

$$\begin{array}{ccccc}
& & Z & & \\
& \swarrow \sigma & & \searrow \tau & \\
Y & & & & X_F \\
& \searrow \pi & & \swarrow \rho & \\
& & X & &
\end{array}$$

Additionally, $\sigma^*(E) = \tau^*(pF) + k^{-1}G$.

If we can show $\mathfrak{c} = \mathcal{O}_{X_F}$, the proof will be complete. Indeed, if $\mathfrak{c} = \mathcal{O}_{X_F}$, then τ is an isomorphism and $\sigma^*E = pF$. But, since $\sigma^*E = pF$ is anti-ample over X , σ must also be an isomorphism and we are done.

We claim that if $\mathfrak{c} \subsetneq \mathcal{O}_{X_F}$, then

$$\text{vol}(\pi^*L - tE) < \text{vol}(\rho^*L - tpF)$$

for $0 < t \ll 1$ and, thus, $S(\mathfrak{a}) < (1/p)S(F)$. Since, we will then have

$$\delta(X, \Delta) \leq \frac{A_{X, \Delta}(F)}{S(F)} < \frac{\text{lct}(\mathfrak{a})}{S(\mathfrak{a})} = \delta(X, \Delta),$$

a contradiction will be reached.

To prove the above claim, fix $0 < \varepsilon \ll 1/k$ so that $H := p\tau^*F + \varepsilon G$ is anti-ample over X . Note that by our choice of ε , we also have

$$\text{vol}(\pi^*L - tE) = \text{vol}(\tau^*(\rho^*L) - t\sigma^*E) \leq \text{vol}(\tau^*(\rho^*L) - tH).$$

Therefore, it suffices to show

$$\text{vol}(\tau^*(\rho^*L) - tH) < \text{vol}(\rho^*L - tpF)$$

for $0 < t \ll 1$.

Fix $0 < t \ll 1$ so that both $A_t := \rho^*L - tpF$ and $B_t := \tau^*(\rho^*L) - tH$ are both ample. Following an argument in [Fuj19b, 3.3], we note that for $0 \leq i \leq n-1$,

$$\begin{aligned}
0 &\leq \varepsilon tG \cdot (\tau^*A_t)^i \cdot B_t^{n-i-1} \\
&= (\tau^*A_t - B_t) \cdot (\tau^*A_t)^i \cdot B_t^{n-i-1},
\end{aligned}$$

since G is effective, τ^*A_t is nef, and B_t is ample. Additionally,

$$0 < (\tau^*A_t - B_t) \cdot B_t^{n-1}.$$

We now see

$$\begin{aligned}
0 &< \sum_{i=0}^{n-1} ((\tau^*A_t - B_t) \cdot (\tau^*A_t)^i \cdot B_t^{n-i-1}) \\
&= (\tau^*A_t)^n - (B_t)^n \\
&= \text{vol}(\rho^*L - tpF) - \text{vol}(\tau^*(\rho^*L) - tH),
\end{aligned}$$

and conclude $\text{vol}(\rho^*L - tpF) < \text{vol}(\tau^*(\rho^*L) - tH)$ for $0 < t \ll 1$. \square

4.3. \mathbb{Q} -divisors computing δ . Let (X, Δ) be a log Fano pair, $\mu : Y \rightarrow X$ a proper birational morphism with Y normal, and E an effective \mathbb{Q} -Cartier \mathbb{Q} -divisor on Y such that $-E$ is μ -ample. We set $\mathfrak{a}_p(E) := \mu_* \mathcal{O}_Y(-\lceil pE \rceil) \subseteq \mathcal{O}_X$ and

$$S(E) := \frac{1}{(-K_X - \Delta)^n} \int_0^\infty \text{vol}(\mu^*(-K_X - \Delta) - tE) dt.$$

Proposition 4.4. *With the above notation, we have*

$$\frac{\text{lct}(X, \Delta; \mathfrak{a}_\bullet(E))}{S(E)} \geq \delta(X, \Delta). \quad (10)$$

Furthermore, if (10) is an equality, then $\text{Supp}(E)$ is a prime divisor.

The statement is a consequence Proposition 4.3 and the following elementary lemma.

Lemma 4.5. *Let $\mu : Y \rightarrow X$ be a proper birational morphism of normal varieties and E an effective \mathbb{Q} -Cartier \mathbb{Q} -divisor on Y such that $-E$ is μ -ample. Set*

$$\mathfrak{a}_p(E) := \mu_* \mathcal{O}_Y(-\lceil pE \rceil) \subseteq \mathcal{O}_X.$$

If $p \in \mathbb{Z}_{>0}$ is sufficiently divisible, then

- (1) Y is the blowup of X along $\mathfrak{a}_p(E)$,
- (2) $\mathfrak{a}_p(E) \cdot \mathcal{O}_Y = \mathcal{O}_Y(-pE)$, and
- (3) $(\mathfrak{a}_p(E))^\ell = \mathfrak{a}_{p\ell}(E)$ for all $\ell \in \mathbb{Z}_{>0}$.

Proof. Since $-E$ is ample over X , $\bigoplus_{m \in \mathbb{N}} \mathfrak{a}_m(E)$ is a finitely generated \mathcal{O}_X -algebra and $Y \simeq \text{Proj}_X(\bigoplus_{m \in \mathbb{N}} \mathfrak{a}_m(E))$. The former statement implies that if $p \in \mathbb{Z}_{>0}$ is sufficiently divisible, then the p -th Veronese, $\bigoplus_{m \in \mathbb{N}} \mathfrak{a}_{pm}(E)$, is finitely generated in degree 1. Hence, (1) and (3) are complete. For (2), observe that the natural map $\mu^* \mu_* \mathcal{O}_X(-pE) \rightarrow \mathcal{O}_Y(-pE)$ is surjective for $p \in \mathbb{Z}_{>0}$ sufficiently divisible, since $-E$ is μ -ample. \square

Proof of Proposition 4.4. Fix $p \in \mathbb{Z}_{>0}$ satisfying (1)-(3) of Lemma 4.5 and set $\mathfrak{a} := \mathfrak{a}_p(E)$. By (1) and (2), $p \cdot S(\mathfrak{a}) = S(E)$. By (3)

$$\text{lct}(X, \Delta; \mathfrak{a}_\bullet(E)) := \lim_{m \rightarrow \infty} (mp \cdot \text{lct}(X, \Delta; \mathfrak{a}_{pm}(E))) = p \cdot \text{lct}(X, \Delta; \mathfrak{a}).$$

The result now follows immediately from Proposition 4.3. \square

5. CONSTRUCTING THE S-EQUIVALENCE

In this section, we prove Theorem 1.1. In Section 5.1 we will construct filtrations of

$$R = \bigoplus_{m \in \mathbb{N}} H^0(X_0, -mr(K_{X_0} + \Delta_0)) \quad \text{and} \quad R' = \bigoplus_{m \in \mathbb{N}} H^0(X'_0, -mr(K_{X'_0} + \Delta'_0)),$$

whose associated graded rings are isomorphic. Then in Section 5.2, we concentrate on proving that these filtrations and their associated graded rings are finitely generated.

5.1. Filtrations induced by degenerations.

Let

$$\pi : (X, \Delta) \rightarrow C \quad \text{and} \quad \pi' : (X', \Delta') \rightarrow C$$

be \mathbb{Q} -Gorenstein families of n -dimensional log Fano pairs over a smooth pointed curve $0 \in C$. Assume there exists an isomorphism

$$\phi : (X, \Delta) \times_C C^\circ \rightarrow (X', \Delta') \times_C C^\circ$$

over $C^\circ := C \setminus 0$ that does not extend to an isomorphism $(X, \Delta) \simeq (X', \Delta')$ over C . Furthermore, assume C is affine and there exists $t \in \mathcal{O}(C)$ so that $\text{div}_C(t) = 0$.

From this setup, we will construct filtrations on the section rings of the special fibers. Set $L := -r(K_X + \Delta)$ and $L' := -r(K_{X'} + \Delta')$, where r is a positive integer so that L and L' are Cartier. For each non-negative integer m , set

$$\begin{aligned} \mathcal{R}_m &:= H^0(X, \mathcal{O}_X(mL)) & \mathcal{R}'_m &:= H^0(X', \mathcal{O}_{X'}(mL')) \\ R_m &:= H^0(X_0, \mathcal{O}_{X_0}(mL_0)) & R'_m &:= H^0(X'_0, \mathcal{O}_{X'_0}(mL'_0)). \end{aligned}$$

Additionally, set

$$\mathcal{R} := \bigoplus_m \mathcal{R}_m, \quad R := \bigoplus_m R_m, \quad \mathcal{R}' := \bigoplus_m \mathcal{R}'_m, \quad \text{and} \quad R' := \bigoplus_m R'_m.$$

Observe that the natural maps

$$\mathcal{R}_m \otimes k(0) \rightarrow R_m \quad \text{and} \quad \mathcal{R}'_m \otimes k(0) \rightarrow R'_m$$

are isomorphisms. Indeed, Kawamata-Viehweg applied to the fibers of π and π' implies $R^i \pi_* \mathcal{O}_X(mL)$ and $R^i \pi'_* \mathcal{O}_{X'}(mL')$ vanish for all $i > 0$ and $m \geq 0$. Hence, $\pi_* \mathcal{O}_X(mL)$ and $\pi'_* \mathcal{O}_{X'}(mL')$ are vector bundles and their cohomology commutes with base change. Since C is affine, \mathcal{R}_m and \mathcal{R}'_m can be identified with the \mathcal{O}_C -module $\pi_* \mathcal{O}_X(mL)$ and $\pi'_* \mathcal{O}_{X'}(mL')$, and the statement follows.

Fix a common log resolution \widehat{X} of (X, Δ) and (X', Δ')

$$\begin{array}{ccc} & \widehat{X} & \\ \psi \swarrow & & \searrow \psi' \\ X & \xrightarrow{\phi} & X' \end{array}$$

and write \widetilde{X}_0 and \widetilde{X}'_0 for the birational transforms of X_0 and X'_0 on \widehat{X} . Set

$$a := A_{X, \Delta+X_0}(\widetilde{X}'_0) \quad \text{and} \quad a' := A_{X', \Delta'+X'_0}(\widetilde{X}_0). \quad (11)$$

Observe that $\widetilde{X}_0 \neq \widetilde{X}'_0$, since otherwise ϕ would extend to an isomorphism over C by the second paragraph of the proof of Proposition 3.2.

5.1.1. Definition of filtrations.

For each $p \in \mathbb{Z}$ and $m \in \mathbb{N}$, set

$$\mathcal{F}^p \mathcal{R}_m := \{s \in \mathcal{R}_m \mid \text{ord}_{\widetilde{X}'_0}(s) \geq p\}, \quad \text{and} \quad \mathcal{F}'^p \mathcal{R}'_m := \{s \in \mathcal{R}'_m \mid \text{ord}_{\widetilde{X}_0}(s) \geq p\}.$$

We define \mathbb{N} -filtrations of R and R' by setting

$$\mathcal{F}^p R_m := \text{im}(\mathcal{F}^p \mathcal{R}_m \rightarrow R_m) \quad \text{and} \quad \mathcal{F}'^p R'_m := \text{im}(\mathcal{F}'^p \mathcal{R}'_m \rightarrow R'_m),$$

where the previous maps are given by restriction of sections. It is straightforward to check that \mathcal{F} and \mathcal{F}' are filtrations of R and R' .

Observe that $R_m \simeq \mathcal{R}_m/t\mathcal{R}_m$ and $\mathcal{F}^p R_m \simeq \text{im}(\mathcal{F}^p \mathcal{R}_m \rightarrow \mathcal{R}_m/t\mathcal{R}_m) \simeq \frac{\mathcal{F}^p \mathcal{R}_m}{\mathcal{F}^p \mathcal{R}_m \cap t\mathcal{R}_m}$ and similar statement holds for \mathcal{F}' . Therefore, we have natural isomorphisms

$$\text{gr}_{\mathcal{F}}^p R_m \simeq \frac{\mathcal{F}^p \mathcal{R}_m}{(\mathcal{F}^p \mathcal{R}_m \cap t\mathcal{R}_m) + \mathcal{F}^{p+1} \mathcal{R}_m} \quad \text{gr}_{\mathcal{F}'}^p R'_m \simeq \frac{\mathcal{F}'^p \mathcal{R}'_m}{(\mathcal{F}'^p \mathcal{R}'_m \cap t\mathcal{R}'_m) + \mathcal{F}'^{p+1} \mathcal{R}'_m}. \quad (12)$$

5.1.2. *Relating the filtrations.* We aim to show that $\text{gr}_{\mathcal{F}} R$ and $\text{gr}_{\mathcal{F}'} R'$ are isomorphic up to a grading shift.

Since $\psi^*(X_0) = \psi'^*(X'_0)$ have multiplicity one along \tilde{X}_0 and \tilde{X}'_0 , we may write

$$K_{\tilde{X}} + \psi_*^{-1}(\Delta) = \psi^*(K_X + \Delta) + a\tilde{X}'_0 + P$$

and

$$K_{\tilde{X}} + \psi'^{-1}_*(\Delta') = \psi'^*(K_{X'} + \Delta') + a'\tilde{X}_0 + P',$$

where the components of $\text{Supp}(P) \cup \text{Supp}(P')$ are both ψ and ψ' -exceptional. Now,

$$\begin{aligned} \mathcal{F}^p \mathcal{R}_m &\simeq H^0\left(\tilde{X}, \mathcal{O}_{\tilde{X}}(m\psi^* L - p\tilde{X}'_0)\right) \\ &= H^0\left(\tilde{X}, \mathcal{O}_{\tilde{X}}(m\psi'^* L' + (mra - p)\tilde{X}'_0 - mra'\tilde{X}_0 + mr(P - P'))\right). \end{aligned}$$

Hence, for $s \in \mathcal{F}^p \mathcal{R}_m$, multiplying $\psi^* s$ by t^{mra-p} gives an element of

$$H^0\left(\tilde{X}, \mathcal{O}_{\tilde{X}}(m\psi'^* L' - (mr(a + a') - p)\tilde{X}_0)\right),$$

which can be identified with $\mathcal{F}'^{mr(a+a')-p} \mathcal{R}'_m$.

As described above, for each $p \in \mathbb{Z}$ and $m \in \mathbb{N}$, there is a map

$$\tilde{\varphi}_{p,m} : \mathcal{F}^p \mathcal{R}_m \longrightarrow \mathcal{F}'^{mr(a+a')-p} \mathcal{R}'_m,$$

which, when \mathcal{R}_m and \mathcal{R}'_m are viewed as submodules of $K(X)$ and $K(X')$, sends $s \mapsto t^{mra-p}(\phi^{-1})^*(s)$. Similarly, there is a map

$$\tilde{\varphi}'_{p,m} : \mathcal{F}'^p \mathcal{R}'_m \longrightarrow \mathcal{F}^{mr(a+a')-p} \mathcal{R}_m,$$

which sends $s' \mapsto t^{mra'-p}\phi^*(s')$. Observe that $\tilde{\varphi}'_{mr(a+a')-p,m} \circ \tilde{\varphi}_{p,m}$ is the identity map, since the composition corresponds to multiplication by $t^{mra' - (mr(a+a') - p)} t^{mra-p} = 1$. Hence, $\tilde{\varphi}_{p,m}$ is an isomorphism.

Lemma 5.1. *For each $p \in \mathbb{Z}$ and $m \in \mathbb{N}$,*

- (1) $\tilde{\varphi}_{p,m}(\mathcal{F}^p \mathcal{R}_m \cap t\mathcal{R}_m) = \mathcal{F}'^{mr(a+a')-p+1} \mathcal{R}'_m$;
- (2) $\tilde{\varphi}_{p,m}(\mathcal{F}^{p+1} \mathcal{R}_m) = \mathcal{F}'^{mr(a+a')-p} \mathcal{R}'_m \cap t\mathcal{R}'_m$.

Proof. To see (1), fix $s \in \mathcal{F}^p \mathcal{R}_m$. Now, $s \in t\mathcal{R}_m$ if and only if s vanishes along X_0 , which is equivalent to the condition that

$$\psi^* s \in H^0\left(\tilde{X}, \mathcal{O}_{\tilde{X}}(m\psi^* L - p\tilde{X}'_0 - \tilde{X}_0)\right).$$

Since the latter holds precisely when

$$t^{mr(a-p)}\psi^*s \in H^0\left(\widehat{X}, \mathcal{O}_{\widehat{X}}(m\psi'^*L' - (mr(a+a')-p+1)\widetilde{X}_0)\right),$$

which is identified with $\mathcal{F}'^{mr(a+a')-p+1}\mathcal{R}'_m$, (1) holds. (2) follows from a similar argument. \square

Proposition 5.2. *The collection of maps $(\tilde{\varphi}_{p,m})$ induce an isomorphism of graded rings*

$$\varphi : \bigoplus_{m \in \mathbb{N}} \bigoplus_{p \in \mathbb{Z}} \text{gr}_{\mathcal{F}}^p R_m \rightarrow \bigoplus_{m \in \mathbb{N}} \bigoplus_{p \in \mathbb{Z}} \text{gr}_{\mathcal{F}'}^{mr(a+a')-p} R'_m.$$

Hence, $\text{gr}_{\mathcal{F}}^p R_m$ and $\text{gr}_{\mathcal{F}'}^p R'_m$ vanish for $p > mr(a+a')$.

Proof. For fixed $p \in \mathbb{Z}$ and $m \in \mathbb{N}$, consider the natural maps

$$\varrho : \mathcal{F}^p \mathcal{R}_m \rightarrow \text{gr}_{\mathcal{F}}^p R_m \quad \text{and} \quad \varrho' : \mathcal{F}'^{mr(a+a')-p} \mathcal{R}'_m \rightarrow \text{gr}_{\mathcal{F}'}^{mr(a+a')-p} R'_m.$$

By (12) and Lemma 5.1, $\tilde{\varphi}_{p,m}$ sends the kernel of ϱ to the kernel ϱ' . Hence, $\tilde{\varphi}_{p,m}$ induces an isomorphism $\varphi_{p,m} : \text{gr}_{\mathcal{F}}^p R_m \rightarrow \text{gr}_{\mathcal{F}'}^{mr(a+a')-p} R'_m$.

Write $\varphi : \text{gr}_{\mathcal{F}} \mathcal{R} \rightarrow \text{gr}_{\mathcal{F}'} R'_m$ for the induced module isomorphism on the direct sums. Since

$$\tilde{\varphi}_{p,m}(\tilde{s}_1)\tilde{\varphi}_{q,\ell}(\tilde{s}_2) = \tilde{\varphi}_{p+q,m+\ell}(\tilde{s}_1 \cdot \tilde{s}_2)$$

for any $\tilde{s}_1 \in \mathcal{F}^p \mathcal{R}_m$ and $\tilde{s}_2 \in \mathcal{F}^q \mathcal{R}_\ell$, we see φ is a ring isomorphism. Since $\text{gr}_{\mathcal{F}}^p R_m$ and $\text{gr}_{\mathcal{F}'}^p R'_m$ vanish when $p < 0$, the isomorphism φ implies the vanishing when $p > mr(a+a')$. \square

5.1.3. Properties of the filtrations.

Lemma 5.3. *For each positive integer p ,*

$$\mathfrak{b}_p(\mathcal{F}) = \mathfrak{a}_p(\text{ord}_{\widetilde{X}'_0}) \cdot \mathcal{O}_{X_0} \quad \text{and} \quad \mathfrak{b}_p(\mathcal{F}') = \mathfrak{a}_p(\text{ord}_{\widetilde{X}_0}) \cdot \mathcal{O}_{X'_0}.$$

Proof. Recall that

$$\mathfrak{b}_p(\mathcal{F}) := \text{im}(\mathcal{F}^p \mathcal{R}_m \otimes \mathcal{O}_{X_0}(-mL_0) \rightarrow \mathcal{O}_{X_0})$$

for $m \gg 0$. Since $\mathcal{F}^p \mathcal{R}_m := \text{im}(\mathcal{F}^p \mathcal{R}_m \rightarrow R_m)$, we see

$$\mathfrak{b}_p(\mathcal{F}) = \text{im}(\mathcal{F}^p \mathcal{R}_m \otimes \mathcal{O}_X(-mL) \rightarrow \mathcal{O}_X) \cdot \mathcal{O}_{X_0}.$$

Therefore, proving the first equality reduces to showing

$$\mathfrak{a}_p(\text{ord}_{\widetilde{X}'_0}) = \text{im}(\mathcal{F}^p \mathcal{R}_m \otimes \mathcal{O}_X(-mL) \rightarrow \mathcal{O}_X)$$

for $m \gg 0$. Since $\mathcal{F}^p \mathcal{R}_m = H^0(X, \mathcal{O}_X(mL) \otimes \mathfrak{a}_p(\text{ord}_{\widetilde{X}'_0}))$ and L is π -ample, the latter statement holds. The argument for $\mathfrak{b}_p(\mathcal{F}')$ is the same. \square

Proposition 5.4. *The following hold:*

- (1) $a \geq \text{lct}(X, \Delta + X_0; \mathfrak{a}_\bullet(\text{ord}_{\widetilde{X}'_0})) = \text{lct}(X_0, \Delta_0; \mathfrak{b}_\bullet(\mathcal{F}))$;
- (2) $a' \geq \text{lct}(X', \Delta' + X'_0; \mathfrak{a}_\bullet(\text{ord}_{\widetilde{X}_0})) = \text{lct}(X'_0, \Delta'_0; \mathfrak{b}_\bullet(\mathcal{F}'))$.

Proof. The first pair of inequalities holds by (3). The second pair follows from Lemma 5.3 and inversion of adjunction. \square

Proposition 5.5. *The filtrations \mathcal{F} and \mathcal{F}' of R and R' are linearly bounded, non-trivial, and satisfy*

$$a + a' = S(\mathcal{F}) + S(\mathcal{F}').$$

Proof. Proposition 5.2 implies $\mathcal{F}^p R_m = 0$ and $\mathcal{F}'^p R'_m = 0$ when $m > 0$ and $p > mr(a + a')$. Therefore, \mathcal{F} and \mathcal{F}' are linearly bounded.

The base ideals $\mathfrak{b}_p(\mathcal{F})$ and $\mathfrak{b}_{p'}(\mathcal{F}')$ are non-zero for $p > 0$ by Lemma 5.3. Therefore, the filtrations cannot be trivial.

Applying Proposition 5.2, we see

$$\begin{aligned} \sum_{p \geq 0} (p \dim \text{gr}_{\mathcal{F}}^p R_m) + \sum_{p \geq 0} (p \dim \text{gr}_{\mathcal{F}'}^p R'_m) &= \sum_{p \geq 0} (p \dim \text{gr}_{\mathcal{F}}^p R_m) + \sum_{p \geq 0} (p \dim \text{gr}_{\mathcal{F}}^{mr(a+a')-p} R_m) \\ &= \sum_{p \geq 0} (mr(a + a') \dim \text{gr}_{\mathcal{F}}^p R_m) . \\ &= mr(a + a') \dim R_m. \end{aligned}$$

Combining the previous equation with (5) gives $S(\mathcal{F}) + S(\mathcal{F}') = a + a'$. \square

It is also natural to rescale the above values and set

$$\beta := (-K_{X_0} - \Delta_0)^n (a - S(\mathcal{F})) \quad \text{and} \quad \beta' := (-K_{X'_0} - \Delta'_0)^n (a' - S(\mathcal{F}')).$$

In this language, Proposition 5.5 states that $\beta + \beta' = 0$.

5.2. Proof of Theorem 1.1. The goal of this subsection is to prove Theorem 1.1. To do so, we consider the filtrations defined in Section 5.1. Under the hypothesis that (X_0, Δ_0) and (X'_0, Δ'_0) are K-semistable, we will show that the filtrations are induced by dreamy divisors.

Furthermore, we will prove that these dreamy divisors induce special test configurations $(\mathcal{X}, \mathcal{D})$ and $(\mathcal{X}', \mathcal{D}')$ of (X_0, Δ_0) and (X'_0, Δ'_0) with generalized Futaki invariant zero. Hence, the log Fano pairs cannot be K-stable. Proposition 5.2 will then be used to show that $(\mathcal{X}_0, \mathcal{D}_0) \simeq (\mathcal{X}'_0, \mathcal{D}'_0)$ and allow us to conclude that (X_0, Δ_0) and (X'_0, Δ'_0) degenerate to a common K-semistable log Fano pair.

Proof of Theorem 1.1. Assume (X_0, Δ_0) and (X'_0, Δ'_0) are both K-semistable and ϕ does not extend to an isomorphism. We must show (X_0, Δ_0) and (X'_0, Δ'_0) are S-equivalent and not K-stable. To do so, we use the filtrations \mathcal{F} and \mathcal{F}' constructed in Section 5.1.

Since (X_0, Δ_0) and (X'_0, Δ'_0) are K-semistable, Proposition 2.8 implies

$$\text{lct}(X_0, \Delta_0; \mathfrak{b}_\bullet(\mathcal{F})) \geq S(\mathcal{F}) \quad \text{and} \quad \text{lct}(X'_0, \Delta'_0; \mathfrak{b}_\bullet(\mathcal{F}')) \geq S(\mathcal{F}').$$

Combining the previous inequalities with Propositions 5.4 and 5.5, we see

$$a = \text{lct}(X, \Delta + X_0; \mathfrak{a}_\bullet(\text{ord}_{\tilde{X}'_0})) = \text{lct}(X_0, \Delta_0; \mathfrak{b}_\bullet(\mathcal{F})) = S(\mathcal{F}) \tag{13}$$

and

$$a' = \text{lct}(X', \Delta' + X'_0; \mathfrak{a}_\bullet(\text{ord}_{\tilde{X}_0})) = \text{lct}(X'_0, \Delta'_0; \mathfrak{b}_\bullet(\mathcal{F}')) = S(\mathcal{F}'). \quad (14)$$

Furthermore, $\delta(X_0, \Delta_0) = \delta(X'_0, \Delta'_0) = 1$.

By the first pair of equalities in (13) and (14), we may apply Proposition 2.2 to extract \tilde{X}'_0 over X and \tilde{X}_0 over X' . Specifically, there exist proper birational morphisms μ and μ' :

$$\begin{array}{ccc} V \cup W \subset Y & & Y' \supset V' \cup W' \\ \mu \downarrow & & \downarrow \mu' \\ X_0 \subset X & \dashrightarrow^\phi & X' \supset X'_0 \\ \pi \searrow & & \swarrow \pi' \\ & C & \end{array}$$

such that the following hold:

- (1) the fibers of Y (respectively, Y') over 0 contains two components V and W (respectively, V' and W') and they are the birational transforms of X_0 and X'_0 ;
- (2) $-W$ and $-V'$ are ample over X and X' respectively;
- (3) $(Y, V + W + \mu_*^{-1}\Delta)$ and $(Y', V' + W' + \mu'^{-1}\Delta')$ are lc.

We write

$$\mu_0 : V \rightarrow X_0 \quad \text{and} \quad \mu'_0 : W' \rightarrow X'_0$$

for the restrictions of μ and μ' to V and W' . Clearly, μ_0 and μ'_0 are proper birational morphisms.

Lemma 5.6. *The pairs $(Y, V + \mu_*^{-1}\Delta)$ and $(Y', W' + \mu'^{-1}\Delta')$ are plt. Hence, V and W' are normal.*

Proof. By inversion of adjunction, $(X, X_0 + \Delta)$ is plt. Therefore, $(Y, V + \mu_*^{-1}\Delta)$ is plt away from $\text{Exc}(\mu) = W$. Since $(Y, V + W + \mu_*^{-1}\Delta)$ is lc, $(Y, V + \mu_*^{-1}\Delta)$ cannot have lc centers in W . Therefore, $(Y, V + \mu_*^{-1}\Delta)$ is plt, and V is normal by [KM98, 5.52]. The same argument works for Y' . \square

Now, consider the restrictions of W and V' to the birational transforms of X_0 and X'_0 :

$$E := W|_V \quad \text{and} \quad E' := V'|_{W'}.$$

Since W and V' are \mathbb{Q} -Cartier, but not necessarily Cartier, E and E' may have fractional coefficients.

The \mathbb{Q} -divisors E and E' induce \mathbb{N} -filtrations on R and R' defined by

$$\mathcal{F}_E^p R_m := H^0\left(V, \mathcal{O}_V(\mu_0^*(mL_0) - \lceil pE \rceil)\right) \subseteq R_m$$

and

$$\mathcal{F}_{E'}^p R'_m := H^0\left(W', \mathcal{O}_{W'}(\mu_0'^*(mL'_0) - \lceil pE' \rceil)\right) \subseteq R'_m$$

for $p, m \geq 0$. Note that

$$\mathcal{F}^p R_m \subseteq \mathcal{F}_E^p R_m \quad \text{and} \quad \mathcal{F}'^p R'_m \subseteq \mathcal{F}_{E'}^p R'_m.$$

Therefore,

$$S(\mathcal{F}) \leq S(\mathcal{F}_E) \quad \text{and} \quad S(\mathcal{F}') \leq S(\mathcal{F}_{E'}). \quad (15)$$

Lemma 5.7. *The supports $F := \text{Supp}(E)$ and $F' := \text{Supp}(E')$ are prime divisors. Furthermore,*

- (1) F computes $\delta(X_0, \Delta_0)$ and $E = \frac{1}{d}F$ for some positive integer d ;
- (2) F' computes $\delta(X'_0, \Delta'_0)$ and $E' = \frac{1}{d'}F'$ for some positive integer d' .

Proof. Since $-W$ is ample over X , the restriction map

$$\mu_* \mathcal{O}_Y(-pW) \rightarrow \mu_{0*} \mathcal{O}_V(-pE)$$

is surjective for all positive integers p sufficiently divisible. Hence, if we set

$$\mathfrak{a}_p(E) := \mu_{0*} \mathcal{O}_V(-pE) \subseteq \mathcal{O}_{X_0},$$

then $\mathfrak{a}_p(E) = \mathfrak{a}_p(\text{ord}_W) \cdot \mathcal{O}_{X_0}$ for such p and inversion of adjunction yields

$$\text{lct}(X_0, \Delta_0; \mathfrak{a}_\bullet(E)) = \text{lct}(X, \Delta + X_0; \mathfrak{a}_\bullet(\text{ord}_W)).$$

Combining the previous equality with (13) and (15) yields

$$\text{lct}(X_0, \Delta_0; \mathfrak{a}_\bullet(E)) = S(\mathcal{F}) \leq S(\mathcal{F}_E). \quad (16)$$

Since (X_0, Δ_0) is K-semistable, Proposition 4.4 implies (16) is an equality and $F := \text{Supp}(E)$ is a prime divisor. Therefore, $S(\mathcal{F}) = S(\mathcal{F}_E)$ and $F := \text{Supp}(E)$ is a prime divisor that computes $\delta(X_0, \Delta_0)$.

To see $E = \frac{1}{d}F$ for a positive integer d , we cut by hyperplanes to reduce the statement to a surface computation. The statement then follows from the fact that $(Y, V + W + \mu_*^{-1}(\Delta))$ is lc and the classification of lc surface pairs ([Kol13, 3.32] and [Kol13, 3.35.2]). The argument for E' is identical. \square

Lemma 5.8. *For all but finitely many $x \in \mathbb{R}_{\geq 0}$,*

$$\text{vol}(\mathcal{F}_E R^{(x)}) = \text{vol}(\mathcal{F} R^{(x)}) \quad \text{and} \quad \text{vol}(\mathcal{F}_{E'} R'^{(x)}) = \text{vol}(\mathcal{F}' R'^{(x)}).$$

Proof. As shown in the proof of Lemma 5.7, $S(\mathcal{F}) = S(\mathcal{F}_E)$. Hence,

$$\frac{1}{rL^n} \int_0^\infty \text{vol}(\mathcal{F} R^{(x)}) dx = S(\mathcal{F}) = S(\mathcal{F}_E) = \frac{1}{rL^n} \int_0^\infty \text{vol}(\mathcal{F}_E R^{(x)}) dx.$$

Since $\text{vol}(\mathcal{F} R^{(x)}) \leq \text{vol}(\mathcal{F}_E R^{(x)})$ by (15) and the two functions are continuous at all but one value [BHK17, 5.3.ii], the desired equality holds. \square

Proposition 5.9. *For all $p \in \mathbb{Z}$ and $m \in \mathbb{N}$,*

$$\mathcal{F}^p R_m = \mathcal{F}_E^p R_m \quad \text{and} \quad \mathcal{F}'^p R'_m = \mathcal{F}_{E'}^p R'_m.$$

Proving this key proposition amounts to showing that the restriction map

$$H^0\left(Y, \mathcal{O}_Y(m\mu^*L - pW)\right) \rightarrow H^0\left(V, \mathcal{O}_V(m\mu_0^*L_0 - \lceil pE \rceil)\right)$$

is surjective. Since such a statement is quite subtle, we will not study this restriction map directly. Instead, we use a construction that originated in [Li17] (with a refining analysis from [LWX18b]) and work on the cone over our family of log Fano pairs.

Consider the relative cone over $(X, \Delta) \rightarrow C$ with polarization L given by

$$Z := C(X/C, L) = \text{Spec}(\mathcal{R}) \rightarrow C.$$

Write $\sigma : C \rightarrow Z$ for the section of cone points and Γ for the closure of the inverse image of Δ under the projection $Z \setminus \sigma(C) \rightarrow X$. Note that the fiber of (Z, Γ) over 0, denoted (Z_0, Γ_0) , is the cone over (X_0, Δ_0) and $Z_0 = \text{Spec}(R)$.

There is a natural proper birational morphism $Y_L \rightarrow Z$, where $Y_L := \text{Spec}_Y\left(\bigoplus_{m \geq 0} \mathcal{O}_Y(mL)\right)$, and it is the total space of the line bundle whose sheaf of sections is $\mathcal{O}_Y(mL)$. We write $Y_{\text{zs}} \subset Y_L$ for the zero section and W_∞ for the preimage of W under the projection map $Y_L \rightarrow Y$. Hence, $Y_{\text{zs}} \cap W_\infty \simeq W$.

Associated to the divisor W over X , is a ray of valuations

$$\{w_t \mid t \in [0, \infty)\} \subset \text{Val}_Z,$$

where w_t is the quasi-monomial valuation with weights $(1, t)$ along Y_{zs} and W_∞ . For each positive integer k , let W_k denote the divisor over X such that $w_{1/k} = \frac{1}{k} \text{ord}_{W_k}$.

Note that

$$\begin{aligned} A_{Z, \Gamma+Z_0}(w_t) = A_{Z, \Gamma+Z_0}(\text{ord}_{Y_{\text{zs}}}) + t A_{Z, \Gamma+Z_0}(\text{ord}_{W_\infty}) &= A_{Z, \Gamma+Z_0}(\text{ord}_{Y_{\text{zs}}}) + t A_{X, \Delta+X_0}(\text{ord}_W) \\ &= r^{-1} + ta \end{aligned}$$

and, by a local computation as in 2.5.1,

$$\mathfrak{a}_p(w_t) = \bigoplus_{m \in \mathbb{N}} \mathcal{F}^{(p-m)/t} \mathcal{R}_m \subseteq \mathcal{R}. \quad (17)$$

Therefore,

$$\mathfrak{a}_p(w_t) \cdot \mathcal{O}_{Z_0} = \bigoplus_{m \in \mathbb{N}} \mathcal{F}^{(p-m)/t} R_m \subseteq R. \quad (18)$$

We also consider a ray in the valuation space of Z_0 . Consider the natural map $V_{L_0} \rightarrow Z_0$, where $V_{L_0} = \text{Spec}_V\left(\bigoplus_{m \geq 0} \mathcal{O}_V(m\mu_0^*L_0)\right)$. We write $V_{\text{zs}} \subset V_{L_0}$ for the zero section and F_∞ for the inverse image of F under the projection $V_{L_0} \rightarrow V$. Let v_t denote the the quasi-monomial valuation with weights $(1, td)$ along V_{zs} and F_∞ . Note that

$$\mathfrak{a}_p(v_t) = \bigoplus_{m \in \mathbb{N}} \mathcal{F}_F^{(p-m)d/t} R_m = \bigoplus_{m \in \mathbb{N}} \mathcal{F}_E^{(p-m)/t} R_m \subseteq R. \quad (19)$$

For each positive integer k divisible by d , there is a divisor F_k over Z_0 such that $v_{1/k} = \frac{d}{k} \text{ord}_{F_k}$. Since F computes $\delta(X_0, \Delta_0) = 1$, the proof of Theorem 4.1 implies

that F_k may be extracted for $k \gg 0$. Let

$$\rho_k : Z_{0,F_k} \rightarrow Z_0$$

denote this extraction.

Lemma 5.10. *For $k \gg 0$, there exists an extraction $\tau_k : Z_{W_k} \rightarrow Z$ of W_k over Z such that $(Z_{W_k}, W_k + \tau_{k*}^{-1}(\Gamma + Z_0))$ is lc.*

Proof. For $t \in \mathbb{R}_{\geq 0}$, let $\mathfrak{a}_\bullet(w_t) \cdot \mathcal{O}_{Z_0}$ denote the restriction of $\mathfrak{a}_\bullet(w_t)$ to a graded sequence of ideals on Z_0 . By Lemma 2.12.1 and Lemma 5.8,

$$\text{mult}(\mathfrak{a}_\bullet(w_t) \cdot \mathcal{O}_{Z_0}) = \text{mult}(\mathfrak{a}_\bullet(v_t)) \quad (20)$$

for each $t \in \mathbb{R}_{\geq 0}$.

Set

$$f(t) := (r^{-1} + at)^{n+1} \text{mult}(\mathfrak{a}_\bullet(w_t) \cdot \mathcal{O}_{Z_0}).$$

Applying Lemma 2.12, we see

$$f(0) = (-K_{X_0} - \Delta_0)^n \quad \text{and} \quad f'(0) = (-K_{X_0} - \Delta_0)^n(a - S(\mathcal{F})) = 0.$$

Hence, a Taylor expansion gives $f(t) = f(0) + O(t^2)$ for $0 < t \ll 1$.

For each positive integer k , define

$$c_k := \text{lct}(Z, \Gamma + Z_0; \mathfrak{a}_\bullet(\text{ord}_{W_k})).$$

Note that $c_k \leq A_{Z,\Gamma+Z_0}(W_k) = kr^{-1} + a$ by (3). Additionally,

$$c_k = k \cdot \text{lct}(Z, \Gamma + Z_0; \mathfrak{a}_\bullet(w_{1/k})) = k \cdot \text{lct}(Z_0, \Gamma_0; \mathfrak{a}_\bullet(w_{1/k}) \cdot \mathcal{O}_{Z_0})$$

by inversion of adjunction and the relation $\mathfrak{a}_{\bullet k}(\text{ord}_{W_k}) = \mathfrak{a}_\bullet(w_{1/k})$. Therefore,

$$\begin{aligned} f(0) &\leq \text{lct}(Z_0, \Gamma_0; \mathfrak{a}_\bullet(w_{1/k}) \cdot \mathcal{O}_{Z_0})^{n+1} \text{mult}(\mathfrak{a}_\bullet(w_{1/k}) \cdot \mathcal{O}_{Z_0}) \\ &= \left(\frac{c_k}{k}\right)^{n+1} \text{mult}(\mathfrak{a}_\bullet(w_{1/k}) \cdot \mathcal{O}_{Z_0}) \leq f\left(\frac{1}{k}\right), \end{aligned}$$

where the first inequality follows from [Liu18, 7] and the assumption that (X_0, Δ_0) is K-semistable (see also [Li17, Theorem 3.1] and [LX16, Theorem A]).

Now, set $a_k := A_{Z,\Gamma+Z_0}(W_k) - c_k$. As in the proof of Theorem 4.1, the previous inequalities imply $\lim_{k \rightarrow \infty} a_k = 0$. Hence, if $k \gg 0$, Proposition 2.2 yields an extraction $\tau_k : Z_{W_k} \rightarrow Z$ of W_k such that the pair

$$(Z_{W_k}, \tau_{k*}^{-1}(\Gamma + Z_0) + (1 - a_k)W_k)$$

is lc. Since $\lim_{k \rightarrow \infty} (1 - a_k) = 1$, the ACC for log canonical thresholds [HMX14] implies $(Z_{W_k}, \tau_{k*}^{-1}(\Gamma + Z_0) + W_k)$ must be lc when $k \gg 0$. \square

From now on, we fix a positive integer k so that d divides k , there exist extractions

$$\rho_k : Z_{0,F_k} \rightarrow Z_0 \quad \text{and} \quad \tau_k : Z_{W_k} \rightarrow Z,$$

and $(Z_{W_k}, \tau_{k*}^{-1}(\Gamma + Z_0) + W_k)$ is lc. The argument used to prove Lemma 5.6 implies $(Z_{W_k}, \tau_{k*}^{-1}(\Gamma + Z_0))$ is plt and $\tau_{k*}^{-1}(Z_0)$ is normal.

Lemma 5.11. *We have a diagram*

$$\begin{array}{ccccccc} F_k \subset Z_{0,F_k} & \hookrightarrow & Z_{W_k} & \supset & W_k \\ \downarrow \rho_k & & \downarrow \tau_k & & \\ Z_0 & \hookrightarrow & Z & & \end{array}$$

(i.e. the birational transform of Z_0 on Z_{W_k} is the extraction of F_k). Additionally,

- (i) $W_k|_{Z_{0,F_k}} = \frac{1}{d}F_k$ and
- (ii) dW_k is Cartier at the generic point of F_k .

Proof. Since $-W_k$ and $-F_k$ are ample over Z and Z_0 , we may find a positive integer p so that

$$\mathfrak{a}_{pd}(\text{ord}_{W_k}) \subseteq \mathcal{O}_Z \quad \text{and} \quad \mathfrak{a}_p(\text{ord}_{F_k}) \subseteq \mathcal{O}_{Z_0}$$

satisfy the conclusions of Lemma 4.5. Hence, Z_{W_k} is the blowup of Z along $\mathfrak{a}_{pd}(\text{ord}_{W_k})$ and Z_{0,F_k} is the blowup of Z_0 along $\mathfrak{a}_p(\text{ord}_{F_k})$. The former statement implies $\tau_k^{-1}(Z_0)$ is the blowup of Z_0 along $\mathfrak{a}_{pd}(\text{ord}_{W_k}) \cdot \mathcal{O}_{Z_0}$.

Claim: $\text{mult}(\mathfrak{a}_{pd}(\text{ord}_{W_k}) \cdot \mathcal{O}_{Z_0}) = \text{mult}(\mathfrak{a}_p(\text{ord}_{F_k}))$

To compute these multiplicities, observe

$$\text{mult}(\mathfrak{a}_p(\text{ord}_{F_k})) = p^{n+1} \cdot \text{mult}(\mathfrak{a}_\bullet(\text{ord}_{F_k})) = (pd/k)^{n+1} \text{mult}(\mathfrak{a}_\bullet(v_{1/k})),$$

since $\mathfrak{a}_{p\ell}(\text{ord}_{F_k}) = \mathfrak{a}_p(\text{ord}_{F_k})^\ell$ for all $\ell \geq 0$ and $\frac{d}{k}\text{ord}_{F_k} = v_{1/k}$. Similar reasoning implies

$$\text{mult}(\mathfrak{a}_{pd}(\text{ord}_{W_k}) \cdot \mathcal{O}_{Z_0}) = (pd/k)^{n+1} \text{mult}(\mathfrak{a}_\bullet(w_{1/k}) \cdot \mathcal{O}_{Z_0}).$$

Equation 20 now completes the claim.

Observe $\mathfrak{a}_{pd}(\text{ord}_{W_k}) \cdot \mathcal{O}_{Z_0} \subseteq \mathfrak{a}_p(\text{ord}_{F_k})$, since

$$\mathfrak{a}_{pd}(\text{ord}_{W_k}) \cdot \mathcal{O}_{Z_0} = \bigoplus_{m \in \mathbb{N}} \mathcal{F}^{pd-mk} R_m \subseteq \bigoplus_{m \in \mathbb{N}} \mathcal{F}_E^{pd-mk} R_m = \mathfrak{a}_p(\text{ord}_{F_k}).$$

A theorem of Rees [Ree61] now implies that $\mathfrak{a}_{pd}(\text{ord}_{W_k}) \cdot \mathcal{O}_{Z_0}$ and $\mathfrak{a}_p(\text{ord}_{F_k})$ have the same integral closure.

The latter implies $\mathfrak{a}_{pd}(\text{ord}_{W_k}) \cdot \mathcal{O}_{Z_0}$ and $\mathfrak{a}_p(\text{ord}_{F_k})$ have the same normalized blowups. Since the corresponding blowups equal $\tau_k^{-1}(Z_0)$ and Z_{0,F_k} and are already normal, they must be isomorphic. The equality of the integral closures further implies $pdW_k|_{\tau_k^{-1}(Z_0)} = pF_k$, which completes (1).

To see (2), cut by $n-1$ generic hyperplanes to get a lc surface pair. The statement then follows from the fact that $(Y, V + W + \mu_*^{-1}(\Delta))$ is lc and the classification of lc surface singularities (see [Kol13, 3.32] and [Kol13, 3.35.2]). \square

Proof of Proposition 5.9. With the above results, the equality of the two filtrations is now a statement concerning valuation ideals (see Equations 18 and 19).

Let us consider the restriction sequence

$$0 \rightarrow \mathcal{O}_{Z_{W_k}}(-pdW_k - Z_{0,F_k}) \rightarrow \mathcal{O}_{Z_{W_k}}(-pdW_k) \rightarrow \mathcal{O}_{Z_{0,F_k}}(-pF_k) \rightarrow 0. \quad (21)$$

where p is a positive integer. By the proof [KM98, 5.26], the sequence is exact if pdW_k is Cartier at all codimension two points of Z_{W_k} contained in Z_{0,F_k} . Since the latter holds by Lemma 5.11, (21) is exact.

Claim: $R^1\tau_{k*}\mathcal{O}_{Z_{W_k}}(-pdW_k - Z_{0,F_k}) = 0$ for all $p > 0$.

Note that $(Z_{W_k}, \tau_{k*}^{-1}(\Gamma))$ is klt, since $(Z_{W_k}, \tau_{k*}^{-1}(\Gamma + Z_0))$ is plt. Therefore, [Kol13, 10.37] implies the desired vanishing holds as long as

$$-pdW_k - Z_{0,F_k} - (K_{Z_{W_k}} + \tau_{k*}^{-1}(\Gamma)) \quad (22)$$

is τ_k -nef. To prove the latter, observe

$$\begin{aligned} K_{Z_{W_k}} + \tau_{k*}^{-1}(\Gamma) &\sim_{\mathbb{Q}, \tau_k} K_{Z_{W_k}} + \tau_{k*}^{-1}(\Gamma) - \tau_k^*(K_Z + \Gamma + Z_0) \\ &= (A_{Z, \Gamma + Z_0}(\text{ord}_{W_k}) - 1)W_k - Z_{0,F_k}. \end{aligned}$$

Therefore, (22) is relatively \mathbb{Q} -linearly equivalent to $-(pd + A_{Z, \Gamma + Z_0}(\text{ord}_{W_k}) - 1)W_k$. Since $-W_k$ is τ_k -ample, (22) is τ_k -nef when $p > 0$ and the proof of the claim is complete.

Returning to the proof of the proposition, we apply τ_{k*} to (21) and see

$$0 \rightarrow \tau_{k*}\mathcal{O}_{Z_{W_k}}(-pdW_k - Z_{0,F_k}) \rightarrow \mathfrak{a}_{pd}(\text{ord}_{W_k}) \rightarrow \mathfrak{a}_p(\text{ord}_{F_k}) \rightarrow 0$$

is exact for all $p > 0$. The right exactness implies $\mathcal{F}^{pd-mk}R_m = \mathcal{F}_E^{pd-mk}R_m$ for all $p > 0$ and $m \geq 0$. Since k was chosen to be a multiple of d , the latter implies $\mathcal{F}^{pd}R_m = \mathcal{F}_E^{pd}R_m$ for all $p, m \geq 0$. Using the relations

$$\mathcal{F}^pR_m \subseteq \mathcal{F}_E^pR_m = \mathcal{F}_E^{[p/d]d}R_m,$$

we conclude $\mathcal{F}^pR_m = \mathcal{F}_E^pR_m$ for all $p, m \geq 0$. \square

We now return to the proof of Theorem 1.1. Recall, $F := \text{Supp}(E)$ and $F' := \text{Supp}(E')$ compute $\delta(X_0, \Delta_0)$ and $\delta(X'_0, \Delta'_0)$, which are both one. Theorem 4.1 implies F and F' are dreamy. Therefore, (X_0, Δ_0) and (X'_0, Δ'_0) are not K-stable.

It remains to show that (X_0, Δ_0) and (X'_0, Δ'_0) are S-equivalent. Consider the filtrations \mathcal{F} and \mathcal{F}' , which agree with \mathcal{F}_E and $\mathcal{F}_{E'}$ by Proposition 5.9. The filtrations \mathcal{F} and \mathcal{F}' are finitely generated (since F and F' are dreamy). Let $(\mathcal{X}, \mathcal{D})$ and $(\mathcal{X}', \mathcal{D}')$ denote the test configuration of (X_0, Δ_0) and (X'_0, Δ'_0) associated to these filtrations.

We claim that $(\mathcal{X}, \mathcal{D})$ and $(\mathcal{X}', \mathcal{D}')$ are non-trivial special test configurations and the fibers over $0 \in \mathbb{A}^1$ are K-semistable. Indeed, $(\mathcal{X}, \mathcal{D})$ is a normal non-trivial test configuration [Fuj17, 3.8] and its Futaki invariant is a multiple of $A_{X_0, \Delta_0}(F) - S(F)$ [Fuj19a, 6.12], which is zero. Therefore, $(\mathcal{X}, \mathcal{D})$ must be special, since otherwise there would exist a test configuration of (X_0, Δ_0) with negative Futaki invariant [LX14, 1]. [LWX18b, 3.1] now implies $(\mathcal{X}_0, \mathcal{D}_0)$ is K-semistable. Since the same argument may be applied to $(\mathcal{X}', \mathcal{D}')$, the claim holds.

To finish the proof of the S-equivalence, we are left to show that there is an isomorphism $(\mathcal{X}_0, \mathcal{D}_0) \simeq (\mathcal{X}'_0, \mathcal{D}'_0)$. Note that

$$\mathcal{X}_0 = \text{Proj} \left(\bigoplus_{m \in \mathbb{N}} \bigoplus_{p \in \mathbb{N}} \text{gr}_{\mathcal{F}}^p R_m \right) \quad \text{and} \quad \mathcal{X}'_0 = \text{Proj} \left(\bigoplus_{m \in \mathbb{N}} \bigoplus_{p \in \mathbb{N}} \text{gr}_{\mathcal{F}'}^p R'_m \right).$$

Therefore, the isomorphism $\varphi : \text{gr}_{\mathcal{F}} R \rightarrow \text{gr}_{\mathcal{F}'} R'$ in Proposition 5.2 induces an isomorphism $\mathcal{X}_0 \simeq \mathcal{X}'_0$. This proves Theorem 1.1 in the case when the boundaries Δ and Δ' are trivial. We claim that φ indeed induces an isomorphism of pairs $(\mathcal{X}_0, \mathcal{D}_0) \simeq (\mathcal{X}'_0, \mathcal{D}'_0)$. Proving \mathcal{D}_0 and \mathcal{D}'_0 match under the isomorphism $\mathcal{X}_0 \simeq \mathcal{X}'_0$ is quite delicate.

To proceed, fix a prime divisor $B \subset \text{Supp}(\Delta)$, and let $B' \subset \text{Supp}(\Delta')$ denote its birational transform on X' . Write $\mathcal{B} \subset \text{Supp}(\mathcal{D})$ and $\mathcal{B}' \subset \text{Supp}(\mathcal{D}')$ for the degenerations of $B_0 \subset X_0$ and $B'_0 \subset X'_0$ on \mathcal{X} and \mathcal{X}' . To complete the proof, we will show that the isomorphism $\mathcal{X}_0 \simeq \mathcal{X}'_0$ sends \mathcal{B}_0 to \mathcal{B}'_0 , where \mathcal{B}_0 and \mathcal{B}'_0 denote the divisorial parts of the scheme theoretic fibers of \mathcal{B} and \mathcal{B}' over 0.

Recall that the scheme theoretic fibers of \mathcal{B} and \mathcal{B}' over 0 are defined by the ideals

$$\text{in}(I_{B_0}) \subset \text{gr}_{\mathcal{F}} R \quad \text{and} \quad \text{in}(I_{B'_0}) \subset \text{gr}_{\mathcal{F}'} R',$$

where $I_{B_0} \subset R$ and $I_{B'_0} \subset R'$ denote the ideals defining B_0 and B'_0 . Observe that $\text{in}(I_{B_0})$ and $\text{in}(I_{B'_0})$ are homogenous with respect to the gradings by m and p . Furthermore, the graded components may be expressed as

$$\text{in}(I_{B_0})_{p,m} := \text{in}(I_{B_0}) \cap \text{gr}_{\mathcal{F}}^p R_m = \text{im} (\mathcal{F}^p R_m \cap I_{B_0} \rightarrow \text{gr}_{\mathcal{F}}^p R_m) \simeq \frac{\mathcal{F}^p R_m \cap I_{B_0}}{\mathcal{F}^{p+1} R_m \cap I_{B_0}}$$

and

$$\text{in}(I_{B'_0})_{p,m} := \text{in}(I_{B'_0}) \cap \text{gr}_{\mathcal{F}'}^p R'_m = \text{im} (\mathcal{F}'^p R'_m \cap I_{B'_0} \rightarrow \text{gr}_{\mathcal{F}'}^p R'_m) \simeq \frac{\mathcal{F}'^p R'_m \cap I_{B'_0}}{\mathcal{F}'^{p+1} R'_m \cap I_{B'_0}}.$$

Rather than showing that the isomorphism $\text{gr}_{\mathcal{F}} R \rightarrow \text{gr}_{\mathcal{F}'} R'$ sends $\text{in}(I_{B_0})$ to $\text{in}(I_{B'_0})$, we introduce auxiliary ideals defined using sections of the relative section rings that vanish along B and B' . For $p, m \geq 0$, set

$$I_{p,m} := \text{im} (\mathcal{F}^p \mathcal{R}_m \cap \mathcal{I}_B \rightarrow \text{gr}_{\mathcal{F}}^p R_m) \quad \text{and} \quad I'_{p,m} := \text{im} (\mathcal{F}'^p \mathcal{R}'_m \cap \mathcal{I}_{B'} \rightarrow \text{gr}_{\mathcal{F}'}^p R'_m),$$

where $\mathcal{I}_B \subset \mathcal{R}$ and $\mathcal{I}_{B'} \subset \mathcal{R}'$ are the ideals defining B and B' . It is straightforward to check that

$$I := \bigoplus_{m \in \mathbb{N}} \bigoplus_{p \in \mathbb{N}} I_{p,m} \subset \text{gr}_{\mathcal{F}} R \quad \text{and} \quad I' := \bigoplus_{m \in \mathbb{N}} \bigoplus_{p \in \mathbb{N}} I'_{p,m} \subset \text{gr}_{\mathcal{F}'} R'$$

are ideals and are contained in $\text{in}(I_{B_0})$ and $\text{in}(I_{B'_0})$.

The following two propositions show that the isomorphism $\mathcal{X}_0 \simeq \mathcal{X}'_0$ induced by φ sends \mathcal{B}_0 to \mathcal{B}'_0 . Indeed, Proposition 5.12 states that the isomorphism $\mathcal{X}_0 \simeq \mathcal{X}'_0$ sends $V(I)$ to $V(I')$. Since $V(I)$ and $V(I')$ agree with \mathcal{B}_0 and \mathcal{B}'_0 away from codimension two subsets by Proposition 5.13, the result follows. \square

We are left to prove the following two propositions used in the above proof.

Proposition 5.12. *The isomorphism $\varphi : \text{gr}_{\mathcal{F}} R \rightarrow \text{gr}_{\mathcal{F}'} R'$ sends I to I' .*

Proof. Observe that for $\tilde{s} \in \mathcal{F}^p \mathcal{R}_m$,

$$\tilde{s} \in \mathcal{F}^p \mathcal{R}_m \cap \mathcal{I}_B \quad \text{if and only if} \quad \tilde{\varphi}_{p,m}(\tilde{s}) \in \mathcal{F}'^{mr(a+a')-p} \mathcal{R}'_m \cap \mathcal{I}_{B'}.$$

Indeed, \tilde{s} and $\tilde{\varphi}_{p,m}(\tilde{s})$ differ by a unit away from $0 \in C$ and membership in the ideals \mathcal{I}_B and $\mathcal{I}_{B'}$ may be tested away from $0 \in C$, since B and B' are horizontal. Therefore, $\varphi(I_{p,m}) = I'_{mr(a+a')-p, m}$ and the result follows. \square

The next proposition is more difficult to prove.

Proposition 5.13. *The subschemes defined by*

- (1) $\text{in}(I_{B_0})$ and I on \mathcal{X}_0 ;
- (2) $\text{in}(I_{B'_0})$ and I' on \mathcal{X}'_0

agree away from codimension 2 subsets.

To prove the statement for (1), it suffices to show that

$$\dim \left(\bigoplus_{p \geq 0} \frac{\text{in}(I_{B_0})_{p,m}}{I_{p,m}} \right) = O(m^{n-2}). \quad (23)$$

To bound the dimension of the previous module, we return to the cone construction argument used earlier in this section.

Consider the relative cone (Z, Γ) and the extractions

$$\tau_k : Z_{W_k} \rightarrow Z \quad \text{and} \quad \rho_k : Z_{0,F_k} \rightarrow Z_0$$

used in the proof of Proposition 5.9. Let $G \subseteq \text{Supp}(\Gamma)$ denote the prime divisor defined via pulling back $B \subseteq \text{Supp}(\Delta)$. Write \tilde{G} and \tilde{G}_0 for the birational transforms of G and G_0 on Z_{W_k} and Z_{0,F_k} .

Observe that for $j \geq 0$

$$\mathfrak{a}_{jd}(\text{ord}_{W_k}) \cap \mathcal{I}_G = \bigoplus_{m \geq 0} (\mathcal{F}^{jd-mk} \mathcal{R}_m \cap \mathcal{I}_B)$$

and

$$\mathfrak{a}_j(\text{ord}_{F_k}) \cap \mathcal{I}_{G_0} = \bigoplus_{m \geq 0} (\mathcal{F}^{jd-mk} R_m \cap I_{B_0}).$$

Therefore,

$$\frac{\mathfrak{a}_j(\text{ord}_{F_k}) \cap \mathcal{I}_{G_0}}{(\mathfrak{a}_{jd}(\text{ord}_{W_k}) \cap \mathcal{I}_G) \cdot \mathcal{O}_{Z_0} + (\mathfrak{a}_{j+1}(\text{ord}_{F_k}) \cap \mathcal{I}_{G_0})} \simeq \bigoplus_{m \geq 0} \frac{\text{in}(I_{B_0})_{jd-mk,m}}{I_{jd-mk,m}}. \quad (24)$$

Lemma 5.14. *We have*

$$\frac{\mathfrak{a}_j(\text{ord}_{F_k}) \cap \mathcal{I}_{G_0}}{(\mathfrak{a}_{jd}(\text{ord}_{W_k}) \cap \mathcal{I}_G) \cdot \mathcal{O}_{Z_0} + (\mathfrak{a}_{j+1}(\text{ord}_{F_k}) \cap \mathcal{I}_{G_0})} = O(j^{n-2}).$$

A key subtlety in proving this lemma is that the divisors \tilde{G} and \tilde{G}_0 may fail to be \mathbb{Q} -Cartier. The proof we will utilize the fact that

$$(Z_{W_k}, W_k + Z_{0,F_k} + \tau_{k*}^{-1}(\Gamma))$$

is lc. The latter implies $F_k = W_k \cap Z_{0,F_k}$ is not contained in $\text{Supp}(\tau_{k*}^{-1}(\Gamma))$ by [Kol13, 2.32.2]. Hence, $F_k \not\subset \tilde{G}$.

Proof. Fix a positive integer q such that qdW_k is Cartier. For each $r \in \{0, \dots, q-1\}$, set

$$\mathcal{Q}_r := \text{coker} \left(\mathcal{O}_{Z_{W_k}}(-\tilde{G} - rdW_k) \rightarrow \mathcal{O}_{Z_{0,F_k}}(-\tilde{G}_0 - rF_k) \right),$$

where the previous map is defined via restriction.

Claim: The support of \mathcal{Q}_r is contained in the intersection of Z_{0,F_k} and the locus where \tilde{G} is not \mathbb{Q} -Cartier.

To prove the claim, it suffices to show

$$\mathcal{O}_{Z_{W_k}}(-\tilde{G} - rdW_k) \rightarrow \mathcal{O}_{Z_{0,F_k}}(-\tilde{G}_0 - rF_k) \rightarrow 0$$

is exact along

$$U := \{z \in Z_{W_k} \mid \tilde{G} \text{ is } \mathbb{Q}\text{-Cartier at } z\}.$$

The the proof of [KM98, 5.26] implies the statement holds, assuming

- (i) $(\tilde{G} + rdW_k)|_U$ and $Z_{0,F_k}|_U$ are \mathbb{Q} -Cartier and
- (ii) $(\tilde{G} + rdW_k)|_U$ is Cartier at all codimension two points of U contained in $Z_{0,F_k}|_U$.

Statement (i) is clear, since W_k and Z_{0,F_k} are \mathbb{Q} -Cartier and U is the locus where \tilde{G} is \mathbb{Q} -Cartier. For (ii), observe that $G + rdW_k$ is Cartier at the generic point of F_k , since $F_k \not\subset \tilde{G}$ and dW_k is Cartier at the generic point of F_k . Note that Z_{W_k} is regular at the remaining codimension two points contained in Z_{0,F_k} . Indeed, $Z_{W_k} \setminus W_k \simeq Z \setminus \sigma(C)$ and Z is regular along all codimension one points of Z_0 , since Z_0 is a normal Cartier divisor on Z .

We now return to the proof of the lemma. Given a positive integer j , write $j = bq + r$ where $r \in \{0, \dots, q-1\}$. Consider the exact sequence

$$\mathcal{O}_{Z_{W_k}}(-\tilde{G} - jdW_k) \rightarrow \mathcal{O}_{Z_{0,F_k}}(-\tilde{G}_0 - jF_k) \rightarrow \mathcal{Q}_r(-bqdW_k) \rightarrow 0.$$

Pushing forward the sequence by τ_{k*} , we see

$$\mathfrak{a}_{jd}(\text{ord}_{W_k}) \cap \mathcal{I}_G \rightarrow \mathfrak{a}_j(\text{ord}_{F_k}) \cap \mathcal{I}_{G_0} \rightarrow \tau_{k*} \mathcal{Q}_r(-bqdW_k) \rightarrow 0,$$

is exact for $b \gg 0$, since $-W_k$ is τ_k -ample. Hence,

$$\begin{aligned} \dim \left(\frac{\mathfrak{a}_j(\text{ord}_{F_k}) \cap \mathcal{I}_{G_0}}{(\mathfrak{a}_{jd}(\text{ord}_{W_k}) \cap \mathcal{I}_G) \cdot \mathcal{O}_{Z_0} + (\mathfrak{a}_{j+1}(\text{ord}_{F_k}) \cap \mathcal{I}_{G_0})} \right) \\ \leq \dim \left(\frac{\mathfrak{a}_j(\text{ord}_{F_k}) \cap \mathcal{I}_{G_0}}{(\mathfrak{a}_{jd}(\text{ord}_{W_k}) \cap \mathcal{I}_G) \cdot \mathcal{O}_{Z_0} + (\mathfrak{a}_{j+q}(\text{ord}_{F_k}) \cap \mathcal{I}_{G_0})} \right) \\ \leq \dim (\text{coker} (\tau_{k*} \mathcal{Q}_r(-bqdW_k) \rightarrow \tau_{k*} \mathcal{Q}_r(-(b+1)qdW_k))). \end{aligned}$$

We are now reduced to showing that the last term equals $O(b^{n-2})$.

Let D denote the effective Cartier divisor qdW_k . Consider the exact sequence

$$\mathcal{Q}_r(-(b+1)D) \rightarrow \mathcal{Q}_r(-bD) \rightarrow \mathcal{Q}_r(-bD)|_D \rightarrow 0$$

After pushing forward by τ_k , we see

$$\tau_{k*}\mathcal{Q}_r(-(b+1)D) \rightarrow \tau_{k*}\mathcal{Q}_r(-bD) \rightarrow H^0(D, \mathcal{Q}_r(-bD)|_D) \rightarrow 0.$$

is exact for $b \gg 0$. Since \mathcal{Q}_r is supported on the locus of Z_{0,F_k} where \tilde{G} is not \mathbb{Q} -Cartier, Lemma 5.15 implies $\mathcal{Q}_r|_D$ has dimension at most $\dim(Z_{0,F_k}) - 3 = n - 2$. Therefore,

$$H^0(D, \mathcal{Q}_r|_D(-bD|_D)) = O(b^{n-2})$$

and the lemma is complete. \square

The previous proof used the following property of lc pairs.

Lemma 5.15. *Let $(X, \Delta + E_1 + E_2)$ be an lc pair such that (i) (X, Δ) is klt and (ii) E_1 and E_2 are \mathbb{Q} -Cartier prime divisors. If $x \in X$ is a codimension three point and $x \in E_1 \cap E_2 \cap \text{Supp}(\Delta)$, then X is \mathbb{Q} -factorial at x .*

Proof. After taking appropriate index one covers, we can assume E_1 and E_2 are Cartier. By cutting, we can assume $\dim(X) = 3$ and x is a closed point. Since (X, Δ) is klt, E_1 is Cohen-Macaulay [KM98, 5.25].

We claim E_1 is normal at x . If not, since E_1 is S_2 , it cannot be R_1 by Serre's Theorem. Hence, E_1 is singular on a curve C passing through x . Note that $C \not\subset \text{Supp}(\Delta) \cup E_2$ by [Kol13, 2.32]. If we consider the normalization $E_1^\nu \rightarrow E_1$, we see $\text{Diff}_{E_1^\nu}(\Delta)$ has coefficient one along the divisors in the preimage of C and positive coefficient along the preimage of $\text{Supp}(\Delta)$. By [Kol13, 2.31], this implies that $(E_1^\nu, \text{Diff}_{E_1^\nu}(\Delta) + E_2|_{E_1^\nu})$ is not lc, which contradicts adjunction.

Shrinking around x , we may assume E_1 is normal. Adjunction gives $(E_1, \Delta|_{E_1} + E_2|_{E_1})$ is lc. Since $E_2|_{E_1}$ is Cartier, $(E_1, \Delta|_{E_1})$ is canonical at x . Using that $x \in \text{Supp}(\Delta|_{E_1})$, [Kol13, 2.29.2] yields that E_1 is smooth at x . Hence, X is smooth at x . \square

Proof of Proposition 5.13. To prove the statement for (1), it suffices to show

$$\sum_{p \geq 0} \dim N_{p,m} = O(m^{n-2}), \quad (25)$$

where $N_{p,m} := \text{in}(I_{B_0})_{p,m}/I_{p,m}$. The previous estimate follows from Lemma 5.14.

Indeed, by Lemma 5.14 and (24)

$$\sum_{m \geq 0} \dim N_{jd-mk,m} = O(j^{n-2}).$$

Since \mathcal{F} is linearly bounded, there exists a positive integer C so that $\text{gr}_{\mathcal{F}}^p R_m = 0$ for all $p \geq mC$. Hence, $N_{p,m} = 0$ for $p \geq mC$ and we see

$$\sum_{m=0}^M \sum_{p \geq 0} \dim N_{pd,m} \leq \sum_{j=0}^{M(C+k)/d} \sum_{m \geq 0} \dim N_{jd-mk,m} = O(M^{n-1})$$

Therefore,

$$\sum_{p \geq 0} \dim N_{pd,m} = O(m^{n-2}).$$

Observe that $\text{gr}_{\mathcal{F}}^p R_m = 0$ for all p not divisible by d by Proposition 5.9 and the fact that $E = d^{-1}F$. Therefore, the previous equation implies (25) holds. Hence, (1) holds and (2) holds by an identical argument. \square

Proof of Corollary 1.3. The proof is the same as the proof of Corollary 3.5, but with Theorem 3.1 replaced by Theorem 1.1 (3). \square

REFERENCES

- [AH11] D. Abramovich and B. Hassett, *Stable varieties with a twist*, Classification of algebraic varieties, 2011, pp. 1–38.
- [Alp13] J. Alper, *Good moduli spaces for Artin stacks*, Ann. Inst. Fourier (Grenoble) **63** (2013), no. 6, 2349–2402.
- [AFS17] J. Alper, M. Fedorchuk, and D. Smyth, *Second flip in the Hassett-Keel program: existence of good moduli spaces*, Compos. Math. **153** (2017), no. 8, 1584–1609.
- [BBJ15] R. Berman, S. Boucksom, and M. Jonsson, *A variational approach to the Yau-Tian-Donaldson conjecture*, arXiv:1509.04561 (2015).
- [Bir16] C. Birkar, *Anti-pluricanonical systems on Fano varieties*, arXiv:1603.05765, to appear in Ann. of Math. (2016).
- [BCHM10] C. Birkar, P. Cascini, C. Hacon, and J. McKernan, *Existence of minimal models for varieties of log general type*, J. Amer. Math. Soc. **23** (2010), no. 2, 405–468.
- [BC11] S. Boucksom and H. Chen, *Okounkov bodies of filtered linear series*, Compos. Math. **147** (2011), no. 4, 1205–1229.
- [BdFFU15] S. Boucksom, T. de Fernex, C. Favre, and S. Urbinati, *Valuation spaces and multiplier ideals on singular varieties*, Recent advances in algebraic geometry, 2015, pp. 29–51.
- [BHQ16] S. Boucksom, T. Hisamoto, and M. Jonsson, *Uniform K-stability, and asymptotics of energy functionals in Kähler geometry*, arXiv:1603.01026, to appear in J. Eur. Math. Soc. (2016).
- [BHQ17] ———, *Uniform K-stability, Duistermaat-Heckman measures and singularities of pairs*, Ann. Inst. Fourier (Grenoble) **67** (2017), no. 2, 743–841.
- [BJ18] S. Boucksom and M. Jonsson, *A non-Archimedean approach to K-stability*, arXiv:1805.11160 (2018).
- [BJ17] H. Blum and M. Jonsson, *Thresholds, valuations, and K-stability*, arXiv:1706.04548 (2017).
- [BL18] H. Blum and Y. Liu, *Openness of uniform K-stability in families of \mathbb{Q} -Fano varieties*, arXiv:1808.09070 (2018).
- [Blu17] H. Blum, *On divisors computing MLD's and LCT's*, arXiv:1605.09662v3 (2017).
- [Blu18a] ———, *Existence of valuations with smallest normalized volume*, Compos. Math. **154** (2018), no. 4, 820–849.

- [Blu18b] ———, *Singularities and K-stability*, PhD thesis. University of Michigan, 2018.
- [Che09] I. Cheltsov, *On singular cubic surfaces*, Asian J. Math. **13** (2009), no. 2, 191–214.
- [Che18] W. Chen, *Boundedness of varieties of Fano type with alpha-invariants and volumes bounded below*, arXiv:1810.04019 (2018).
- [CDS15] X. Chen, S. Donaldson, and S. Sun, *Kähler-Einstein metrics on Fano manifolds.*, J. Amer. Math. Soc. **28** (2015), no. 1, 183–197, 199–234, 235–278.
- [CP18] G. Codogni and Z. Patakfalvi, *Positivity of the CM line bundle for families of K-stable klt Fano*, arXiv:1806.07180 (2018).
- [Der16] R. Dervan, *Uniform stability of twisted constant scalar curvature Kähler metrics*, Int. Math. Res. Not. **15** (2016), 4728–4783.
- [Don01] S. K. Donaldson, *Scalar curvature and projective embeddings. I*, J. Differential Geom. **59** (2001), no. 3, 479–522.
- [Don02] S. Donaldson, *Scalar curvature and stability of toric varieties*, J. Differential Geom. **62** (2002), no. 2, 289–349.
- [Fuj16] K. Fujita, *On K-stability and the volume functions of \mathbb{Q} -Fano varieties*, Proc. Lond. Math. Soc. (3) **113** (2016), no. 5, 541–582.
- [Fuj17] ———, *K-stability of log Fano hyperplane arrangements*, arXiv:1709.08213 (2017).
- [Fuj18] ———, *Optimal bounds for the volumes of Kähler-Einstein Fano manifolds*, Amer. J. Math. **140** (2018), no. 2, 391–414.
- [Fuj19a] ———, *A valuative criterion for uniform K-stability of \mathbb{Q} -Fano varieties*, J. Reine Angew. Math. **751** (2019), 309–338.
- [Fuj19b] ———, *Uniform K-stability and plt blowups of log Fano pairs*, Kyoto J. Math. **59** (2019), no. 2, 399–418.
- [FO18] K. Fujita and Y. Odaka, *On the K-stability of Fano varieties and anticanonical divisors*, Tohoku Math. J. (2) **70** (2018), no. 4, 511–521.
- [HMX14] C. Hacon, J. McKernan, and C. Xu, *ACC for log canonical thresholds*, Ann. of Math. (2) **180** (2014), no. 2, 523–571.
- [HK04] B. Hassett and S. Kovács, *Reflexive pull-backs and base extension*, J. Algebraic Geom. **13** (2004), no. 2, 233–247.
- [Jia17] C. Jiang, *Boundedness of \mathbb{Q} -Fano varieties with degrees and alpha-invariants bounded from below*, arXiv:1705.02740, to appear in Ann. Sci. École Norm. Sup. (2017).
- [JM12] M. Jonsson and M. Mustață, *Valuations and asymptotic invariants for sequences of ideals*, Ann. Inst. Fourier (Grenoble) **62** (2012), no. 6, 2145–2209 (2013).
- [KM97] S. Keel and S. Mori, *Quotients by groupoids*, Ann. of Math. (2) **145** (1997), no. 1, 193–213.
- [Kol09] J. Kollar, *Hulls and Husks*, arXiv:0805.0576 (2009).
- [Kol13] ———, *Singularities of the minimal model program*, Cambridge Tracts in Math., vol. 200, Cambridge University Press, Cambridge, 2013. With the collaboration of S. Kovács.
- [Kol19] ———, *Families of varieties of general type*, Book to appear, 2019.
- [KM98] J. Kollar and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Math., vol. 134, Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti.
- [LLSW17] K. Lee, Z. Li, J. Sturm, and X. Wang, *Asymptotic Chow stability of toric Del Pezzo surfaces*, arXiv:1711.10099, to appear in Math. Res. Lett. (2017).
- [Li17] C. Li, *K-semistability is equivariant volume minimization*, Duke Math. J. **166** (2017), no. 16, 3147–3218.
- [Li18] ———, *Minimizing normalized volumes of valuations*, Math. Zeit. **289** (2018), no. 1-2, 491–513.

- [LL19] C. Li and Y. Liu, *Kähler-Einstein metrics and volume minimization*, Adv. Math. **341** (2019), 440–492.
- [LW18] C. Li and X. Wang, *Private communications* (2018).
- [LWX18a] C. Li, X. Wang, and C. Xu, *Quasi-projectivity of the moduli space of smooth Kähler-Einstein Fano manifolds*, Ann. Sci. École Norm. Sup. **51** (2018), no. 3, 739–772.
- [LWX18b] ———, *Algebraicity of the metric tangent cones and equivariant K-stability*, arXiv:1805.03393 (2018).
- [LWX19] ———, *On the proper moduli spaces of smoothable Kähler-Einstein Fano varieties*, Duke Math. J. **168** (2019), no. 8, 1387–1459.
- [LX14] C. Li and C. Xu, *Special test configuration and K-stability of Fano varieties*, Ann. of Math. (2) **180** (2014), no. 1, 197–232.
- [LX16] ———, *Stability of Valuations and Kollar Components*, arXiv:1604.05398, to appear in J. Eur. Math. Soc. (2016).
- [LX18] ———, *Stability of Valuations: Higher Rational Rank*, Peking Math. J. **1** (2018), 1–79.
- [Liu18] Y. Liu, *The volume of singular Kähler-Einstein Fano varieties*, Compos. Math. **154** (2018), no. 6, 1131–1158.
- [LLX18] C. Li, Y. Liu, and C. Xu, *A guided tour to normalized volumes*, arXiv:1806.07112 (2018).
- [Oda12] Y. Odaka, *On the moduli of Kähler-Einstein Fano manifolds*, arXiv:1211.4833 (2012).
- [Oda15] ———, *Compact moduli spaces of Kähler-Einstein Fano varieties*, Publ. Res. Inst. Math. Sci. **51** (2015), no. 3, 549–565.
- [OSY12] H. Ono, Y. Sano, and N. Yotsutani, *An example of an asymptotically Chow unstable manifold with constant scalar curvature*, Ann. Inst. Fourier (Grenoble) **62** (2012), no. 4, 1265–1287.
- [PP10] B. Pasquier and N. Perrin, *Local rigidity of quasi-regular varieties*, Math. Z. **265** (2010), no. 3, 589–600.
- [Ree61] D. Rees, *\mathfrak{a} -transforms of local rings and a theorem on multiplicities of ideals*, Proc. Cambridge Philos. Soc. **57** (1961), 8–17.
- [SSY16] C. Spotti, S. Sun, and C. Yao, *Existence and deformations of Kähler-Einstein metrics on smoothable \mathbb{Q} -Fano varieties*, Duke Math. J. **165** (2016), no. 16, 3043–3083.
- [Szé15] G. Székelyhidi, *Filtrations and test-configurations*, Math. Ann. **362** (2015), no. 1-2, 451–484. With an appendix by S. Boucksom.
- [Tia97] G. Tian, *Kähler-Einstein metrics with positive scalar curvature*, Invent. Math. **130** (1997), no. 1, 1–37.
- [Tia15] ———, *K-stability and Kähler-Einstein metrics*, Comm. Pure Appl. Math. **68** (2015), no. 7, 1085–1156.
- [Vie95] E. Viehweg, *Quasi-projective moduli for polarized manifolds*, Ergeb. Math. Grenzgeb. (3), vol. 30, Springer-Verlag, Berlin, 1995.
- [WX14] X. Wang and C. Xu, *Nonexistence of asymptotic GIT compactification*, Duke Math. J. **163** (2014), no. 12, 2217–2241.
- [WN12] D. Witt Nyström, *Test configurations and Okounkov bodies*, Compos. Math. **148** (2012), no. 6, 1736–1756.

UNIVERSITY OF UTAH, SALT LAKE CITY, UT 48112, USA
Email address: blum@math.utah.edu

BEIJING INTERNATIONAL CENTER FOR MATHEMATICAL RESEARCH, BEIJING 100871, CHINA
Email address: cyxu@math.pku.edu.cn

CURRENT ADDRESS: MIT, CAMBRIDGE, MA 02139, USA

Email address: `cyxu@math.mit.edu`