

# Wilton Ripples in Weakly Nonlinear Models of Water Waves: Existence and Computation

Benjamin Akers · David P. Nicholls

Received: date / Accepted: date

**Abstract** In this contribution we prove that small amplitude, resonant harmonic, **spatially periodic** traveling waves (Wilton **ripples**) exist in a family of weakly nonlinear PDEs which model water waves. The proof is inspired by that of Reeder & Shinbrot (1981) and complements the authors' recent, independent result proven by a perturbative technique (Akers & Nicholls, 2020). The method is based on a Banach Fixed Point Iteration and, in addition to proving that this iteration has Wilton **ripples** as a fixed point, we use it as a numerical method for simulating these solutions. The output of this numerical scheme and its performance are evaluated against a **quasi**-Newton iteration.

**Keywords** Wilton Ripples · weakly nonlinear PDEs · Whitham equation · Benjamin equation · Kawahara equation · Akers–Milewski equation.

## 1 Introduction

In this work we consider the existence of a family of resonant traveling waves, first described by Wilton [61], in a class of weakly nonlinear wave equations

---

B.A. was supported during the preparation of this manuscript by the Air Force Office of Sponsored Research and the Office of Naval Research. D.P.N. gratefully acknowledges support from the National Science Foundation through grant No. DMS-1813033.

---

Benjamin Akers  
Department of Mathematics  
Air Force Institute of Technology  
Wright Patterson Air Force Base  
Dayton, OH 45433 E-mail: Benjamin.Akers@afit.edu

David P. Nicholls  
Department of Mathematics, Statistics,  
and Computer Science  
University of Illinois at Chicago  
Chicago, IL 60607 E-mail: davidn@uic.edu

which model water waves. These waves occur in the models at certain *critical* values of the surface tension [42]. The surface tension coefficient appears as a parameter in these PDEs and, for special values of this parameter, the operator defining the linear portion of the equation has a two dimensional kernel implying that the linear problem supports *two* co-propagating resonant harmonics. Continuous branches of these near-bichromatic waves exist about special combinations of the harmonics which are called “Wilton **ripples**.” These waves have been studied for over a century now including asymptotic [61, 46, 34, 39], numerical [11, 60, 19, 14], and experimental work [45, 35, 50]. In this paper we present a rigorous treatment of the phenomena in a class of weakly nonlinear PDEs, proving existence of these resonant ripples and building a rapid and robust numerical method based directly upon the iteration we describe in our proof. We point out that, while connected to our previous work on this topic [7], our iteration does not deliver the analyticity of solution branches required of the previous numerical method of the first author [11].

Our method of proof is to apply a Banach Fixed Point Iteration (FPI) [30], however, suitable “initiation” is required to demonstrate the existence of a fixed point. In more detail, in the case of Stokes waves (which feature only a one-dimensional null space in the linear operator) the first step must be carefully described. By contrast, in the case of Wilton **ripples** the first *two* steps must be constructed in order to assure that the correct type of solution is found.

Away from special surface tension values, small amplitude traveling solutions are monochromatic so that the linear problem has a one-dimensional solution space, resulting in Stokes waves. These waves have a significant history of both asymptotic [56, 33, 23, 44] and numerical study [52, 53, 48, 3]. Their rigorous study is also well developed as Stokes waves are known to exist [43, 57], and be parametrically analytic in amplitude [21, 22, 49]. The global bifurcation problem has been described and extreme waves on branches have been characterized [13].

Wilton **ripples** have a shorter and less complete history of rigorous study. Often the resonant and non-resonant cases are studied separately due to the change in form of the wave’s asymptotics. The bifurcation structure has been described for the non-resonant case for both potential flow [10] and the hydroelastic water wave problem [58]. Reeder and Shinbrot discuss the existence and analyticity of small resonant Wilton **ripples** in potential flow [51] while the bifurcation structure of Wilton **ripples** has recently been described for the Whitham equation [29]. Although our motivation stems from water waves, here we consider Wilton **ripples** in a family of weakly nonlinear models, including the Kawaraha [40], Benjamin [15], Whitham [47] and Akers–Milewski [2] equations. In a recent paper the authors used a perturbation theoretic method to demonstrate that resonant triad ripples exist along analytic bifurcation curves in this class of model equations [7]. Although the proof in [7] gives a stronger result, we believe that the proof in this work is also of value, both for its associated numerical method and its potential to be applied to the stability

problem (perhaps to amplitudes at which the spectrum is continuously, but not analytically, connected to the flat configuration).

As a bifurcation problem, the computation of traveling waves shares similar structure to that of the computation of their spectral data. This analogy forms the inspiration for the numerical methods in [4–6]. In the setting of this analogy, resonant traveling waves correspond to eigenvalue collisions; thus, while resonant traveling waves might be considered “exotic” and “uncommon,” eigenvalue collisions are viewed as the *primary* mechanism for instability and are therefore the most interesting and important parts of the spectrum [9]. In the future, we hope to develop this proof and numerical method for traveling waves to complement traditional studies of the spectrum, both numerical [28, 27, 59] and theoretical [17, 36–38, 8]. Recently Creedon, Deconinck and Trichtchenko combined the Fourier–Floquet–Hill method from [28] with the asymptotic ideas in [8] to compute high frequency spectrum in a Boussinesq–Whitham model [25] and the Kawahara equation [26]. Applying the ideas of this paper to the spectral data would be a natural complement to the work this group is doing on high-frequency instabilities.

In this paper we consider a family of model equations whose linear phase speed allows two co-propagating resonant harmonics. In these equations we show that small amplitude Wilton **ripples** exist, and that there are branches of these waves, at least up to some maximal amplitude. The paper is organized as follows: In § 2 we state the weakly nonlinear wave equations which govern the physical phenomena of interest in this paper. We give a proof of the existence of Stokes waves in § 3, with the study of the linear problem § 3.1, a description of the iteration in § 3.2, and the result in § 3.3. Subsequent to this, we establish the existence of Wilton **ripples** in § 4, with the study of the linear and quadratic problems in § 4.1 & § 4.2, a description of the iteration in § 4.3, and the result in § 4.4. We close with numerical results in § 5 and concluding remarks in § 6.

## 2 Governing Equations

In this paper we study  $x$ -periodic solutions of weakly nonlinear dispersive wave equations of the form

$$\partial_t u + \mathcal{L} \partial_x u - \frac{1}{2} \partial_x [u^2] = 0, \quad u(x + 2\pi, t) = u(x, t), \quad (1)$$

where  $\mathcal{L}$  is a linear operator (see Table 1 for examples considered here), which model the full water wave equations [42] in the limit of long wavelength and weak nonlinearity. Among the myriad choices, we focus on those which permit co-propagating traveling waves in the linear regime. Examples are the Whitham [47], Akers–Milewski [2], Benjamin [15], and Kawahara [40] equations (see Table 1). We now turn to traveling wave solutions of (1)

$$u(x, t) = f(x - ct)$$

which satisfy

$$-c\partial_x f + \mathcal{L}\partial_x f - \frac{1}{2}\partial_x [f^2] = 0.$$

After integrating once with respect to  $x$  we find

$$(-c + \mathcal{L})f = -\frac{1}{2}f^2.$$

which we write as

$$(-c_0 + \mathcal{L})f = (c - c_0)f - \frac{1}{2}f^2,$$

where we define  $c_0$  presently. By defining

$$\mathcal{R} := -c_0 + \mathcal{L},$$

we express this as

$$\mathcal{R}f = \left( (c - c_0) - \frac{1}{2}f \right) f. \quad (2)$$

**Remark 1** *At this point we can clarify the requirements of the operators  $\mathcal{L}$  which fall into our theory. Principally we demand that they be “wave-like” in that there exists a real value  $c_0$  such that  $(-c_0 + \mathcal{L})[\cos(x)] = 0$  which delivers a sinusoidal traveling wave in the linearized equation. Furthermore, we require that the operator  $\mathcal{R}$  (equivalently  $\mathcal{L}$ ) “take” only a finite number of derivatives,  $0 \leq \tau < \infty$ , so that  $\mathcal{R} : H^{\sigma+\tau} \rightarrow H^\sigma$ , c.f., Theorems 1 and 3. Also, we point out that the nonlinearity*

$$N(u) = \frac{1}{2}\partial_x [u^2]$$

*can be generalized somewhat and still fit into our framework at the cost of a slight increase in the complexity of our later estimates.*

### 3 Stokes Waves

In order to give some notation and build intuition for the Wilton **ripple** case, we begin with the simpler case of the classical Stokes waves which result from simple bifurcation [24] in (2). For this we note that there is a speed  $c_0$  such that the operators  $\mathcal{R}$  and its adjoint  $\mathcal{R}^*$  have one-dimensional null spaces spanned by  $\phi$  and  $\psi$ , respectively, so that

$$\mathcal{R}\phi = 0, \quad \mathcal{R}^*\psi = 0, \quad \phi(x) = \psi(x) = \frac{\cos(x)}{\sqrt{\pi}},$$

and we note that

$$\langle \phi, \phi \rangle = \langle \psi, \psi \rangle = \langle \phi, \psi \rangle = 1.$$

### 3.1 Preparing the Iteration: Linear Order

We now seek a solution of (2) of the form

$$f = \varepsilon f_1 + \varepsilon^2 F, \quad c = c_0 + \varepsilon s, \quad (3)$$

where we point out that, as proven in [7],  $F$  and  $s$  depend upon  $\varepsilon$  and, furthermore,  $F = \mathcal{O}(1)$  and  $s = \mathcal{O}(\varepsilon)$ . Inserting these into (2) gives

$$\mathcal{R}[\varepsilon f_1 + \varepsilon^2 F] = \left( \varepsilon s - \frac{1}{2} \varepsilon f_1 - \frac{1}{2} \varepsilon^2 F \right) (\varepsilon f_1 + \varepsilon^2 F). \quad (4)$$

At linear order we find

$$\mathcal{R}f_1 = 0$$

which demands that  $f_1 \in \mathcal{N}(\mathcal{R})$  so that  $f_1 = \alpha \phi$ ; without loss of generality we may select  $\alpha = 1$ .

### 3.2 The Stokes Iteration

Using this choice of  $f_1$  we return to (4) and, using the properties of  $f_1$ , and dividing by  $\varepsilon^2$  we find

$$\mathcal{R}F - s f_1 = -\frac{1}{2} f_1^2 + \varepsilon \left\{ sF - f_1 F - \frac{1}{2} \varepsilon F^2 \right\}.$$

Inspired by this we conduct the following Stokes iteration

$$\mathcal{R}F_n - s_n f_1 = -\frac{1}{2} f_1^2 + \varepsilon G(F_{n-1}, s_{n-1}; \varepsilon), \quad (5)$$

where

$$G(F, s; \varepsilon) := sF - f_1 F - \frac{1}{2} \varepsilon F^2.$$

Our procedure for solving (5) at each iteration is two-step:

1. First, we solve for the speed correction  $s_n$  by projecting the right-hand-side of (5) and  $s_n f_1$  onto the range of  $\mathcal{R}$ . This is enforced by demanding that this be orthogonal to the null space of its adjoint,

$$\begin{aligned} 0 &= \left\langle s_n f_1 - \frac{1}{2} f_1^2 + \varepsilon G(F_{n-1}, s_{n-1}; \varepsilon), \psi \right\rangle \\ &= s_n \langle \phi, \psi \rangle - \frac{1}{2} \langle \phi^2, \psi \rangle + \varepsilon \langle G(F_{n-1}, s_{n-1}; \varepsilon), \psi \rangle. \end{aligned}$$

Using  $\langle \phi, \psi \rangle = 1$  and (from our choices of  $\{\phi, \psi\}$ )  $\langle \phi^2, \psi \rangle = 0$  we find that

$$s_n = -\varepsilon \langle G(F_{n-1}, s_{n-1}; \varepsilon), \psi \rangle.$$

2. With this choice of  $s_n$  we know that (5) is uniquely solvable provided that we demand orthogonality of the solution to the null space of  $\mathcal{R}$ . So, we can write

$$F_n = \mathcal{R}^{-1} \left[ s_n f_1 - \frac{1}{2} f_1^2 + \varepsilon G(F_{n-1}, s_n; \varepsilon) \right],$$

$$\langle \phi, F_n \rangle = 0.$$

This procedure can be replicated for the linearized version of this problem

$$\mathcal{R}F - s f_1 = -\frac{1}{2} f_1^2 + J,$$

with generic function  $J(x)$ , and rigorous estimates derived for the unique solution. This is summarized in the following “elliptic estimate” [7].

**Theorem 1** *For any real  $\sigma \geq 0$ , given  $J \in H^\sigma$  there exists a unique solution pair  $(F, s)$  of*

$$\mathcal{R}F - s f_1 = -\frac{1}{2} f_1^2 + J,$$

$$\langle \phi, F \rangle = 0,$$

such that  $F \in H^{\sigma+\tau}$  and

$$|s| \leq C_e \|J\|_{H^\sigma},$$

$$\|F\|_{H^{\sigma+\tau}} \leq C_e \left\| s f_1 - \frac{1}{2} f_1^2 + J \right\|_{H^\sigma}$$

for some  $C_e > 0$  and

$$\tau = \begin{cases} 1/2, & \text{Whitham (A),} \\ 1, & \text{Akers–Milewski (B),} \\ 2, & \text{Benjamin (C),} \\ 4, & \text{Kawahara (D).} \end{cases}$$

### 3.3 Banach Fixed Point Formulation

We now write this as an FPI so that we can appeal to the Banach Fixed Point Theorem [30]. For this we write (2) as

$$\mathcal{R}F - f_1 s = -\frac{1}{2} f_1^2 + \varepsilon G(F, s; \varepsilon),$$

$$\langle \phi, F \rangle = 0,$$

or

$$M \begin{pmatrix} F \\ s \end{pmatrix} = \begin{pmatrix} -(1/2) f_1^2 + \varepsilon G(F, s; \varepsilon) \\ 0 \end{pmatrix}, \quad M := \begin{pmatrix} \mathcal{R} & -f_1 \\ \langle \phi, \cdot \rangle & 0 \end{pmatrix}.$$

We now express this as  $x = T(x)$  where

$$x := \begin{pmatrix} F \\ s \end{pmatrix}, \quad T(x) := M^{-1} \begin{pmatrix} -(1/2)f_1^2 + \varepsilon G(F, s; \varepsilon) \\ 0 \end{pmatrix},$$

and seek a fixed point of the iteration

$$x_n = T(x_{n-1}).$$

From Theorem 1 we know that  $M^{-1}$  is well-defined and that the action of  $x^* = T(x)$  can be given by

$$s^* = -\varepsilon \langle G(F, s; \varepsilon), \psi \rangle, \quad (6a)$$

$$F^* = \mathcal{R}^{-1} \left[ s^* f_1 - \frac{1}{2} f_1^2 + \varepsilon G(F, s; \varepsilon) \right]. \quad (6b)$$

A fixed point of the iteration  $x_n = T(x_{n-1})$  would solve  $x = T(x)$  and represent a solution of (2). To achieve this we appeal to Banach's Fixed Point Theorem [30] which requires a Banach space,

$$X = H^{\sigma+\tau} \times \mathbf{R},$$

and a distance,

$$d(x, y)^2 = \|F_x - F_y\|_{H^{\sigma+\tau}}^2 + |s_x - s_y|^2.$$

To show that  $T$  is a contraction, we need the following classical Lemma [41, 1, 31, 30].

**Lemma 1** *Provided that  $\sigma > 1/2$ , if  $f, g \in H^\sigma$  then  $fg \in H^\sigma$  and*

$$\|fg\|_{H^\sigma} \leq K \|f\|_{H^\sigma} \|g\|_{H^\sigma},$$

for some  $K > 0$ .

With this we can establish the following crucial estimate which demonstrates that  $T$  is a contraction for  $\varepsilon$  sufficiently small.

**Lemma 2** *Provided that  $\sigma > 1/2$  and  $x, y \in B_R(0) \subset X$ , where  $B_R(0)$  is the ball of radius  $R$  centered at the origin, we have that*

$$d(T(x), T(y))^2 \leq \varepsilon^2 C(R)^2 d(x, y)^2,$$

where  $C(R)$  is a quadratic function of  $R$ .

*Proof* We begin with

$$\begin{aligned} d(T(x), T(y))^2 &= d(x^*, y^*)^2 \\ &= \|F_x^* - F_y^*\|_{H^{\sigma+\tau}}^2 + |s_x^* - s_y^*|^2. \end{aligned}$$

The first term can be estimated

$$\begin{aligned} \|F_x^* - F_y^*\|_{H^{\sigma+\tau}} &= \left\| \mathcal{R}^{-1} \left[ s_x^* f_1 - \frac{1}{2} f_1^2 + \varepsilon G(F_x, s_x; \varepsilon) \right] \right. \\ &\quad \left. - \mathcal{R}^{-1} \left[ s_y^* f_1 - \frac{1}{2} f_1^2 + \varepsilon G(F_y, s_y; \varepsilon) \right] \right\|_{H^{\sigma+\tau}} \\ &\leq \| (s_x^* - s_y^*) \phi + \varepsilon (\Delta G) \|_{H^\sigma} \\ &\leq |s_x^* - s_y^*| \|\phi\|_{H^\sigma} + \varepsilon \|\Delta G\|_{H^\sigma}, \end{aligned}$$

where we have used Theorem 1,  $f_1 = \phi$ , and defined

$$\Delta G := G(F_x, s_x; \varepsilon) - G(F_y, s_y; \varepsilon).$$

Using  $(a+b)^2 \leq 2(a^2 + b^2)$ , we now have

$$d(T(x), T(y))^2 \leq \left(1 + 2\|\phi\|_{H^\sigma}^2\right) |s_x^* - s_y^*|^2 + 2\varepsilon^2 \|\Delta G\|_{H^\sigma}^2. \quad (7)$$

Now, the difference in the speed corrections can be estimated from the solution formula (6a) and the Cauchy–Schwartz Inequality

$$\begin{aligned} |s_x^* - s_y^*| &= \varepsilon |\langle G(F_x, s_x; \varepsilon) - G(F_y, s_y; \varepsilon), \psi \rangle| \\ &\leq \varepsilon \|\Delta G\|_{H^0} \|\psi\|_{H^0} \leq \varepsilon \|\Delta G\|_{H^\sigma}, \end{aligned}$$

and we have used  $\sigma > 1/2 > 0$  and  $\|\psi\|_{H^0} = \|\psi\|_{L^2} = 1$ . With this (7) tells us that

$$d(T(x), T(y))^2 \leq \varepsilon^2 \left(3 + 2\|\phi\|_{H^\sigma}^2\right) \|\Delta G\|_{H^\sigma}^2. \quad (8)$$

Finally, we estimate the term  $\Delta G$  with the following manipulations

$$\begin{aligned} \|\Delta G\|_{H^\sigma} &\leq \left\| s_x F_x - \phi F_x - \frac{1}{2} \varepsilon F_x^2 - \left( s_y F_y - \phi F_y - \frac{1}{2} \varepsilon F_y^2 \right) \right\|_{H^\sigma} \\ &\leq |s_x - s_y| \|F_x\|_{H^\sigma} + |s_y| \|F_x - F_y\|_{H^\sigma} + \textcolor{red}{K} \|\phi\|_{H^\sigma} \|F_x - F_y\|_{H^\sigma} \\ &\quad + \frac{1}{2} \varepsilon \textcolor{red}{K} \|F_x + F_y\|_{H^\sigma} \|F_x - F_y\|_{H^\sigma}. \end{aligned}$$

Since  $x, y \in B_R(0)$  we have that

$$\max \{s_m, \|F_m\|_{H^\sigma}\} < R, \quad m \in \{x, y\},$$

and since

$$\max \{|s_x - s_y|, \|F_x - F_y\|_{H^{\sigma+\tau}}\} \leq d(x, y),$$

we discover that

$$\begin{aligned} \|\Delta G\|_{H^\sigma} &\leq |s_x - s_y| R + R \|F_x - F_y\|_{H^\sigma} + \textcolor{red}{K} \|\phi\|_{H^\sigma} \|F_x - F_y\|_{H^\sigma} \\ &\quad + \frac{1}{2} \varepsilon \textcolor{red}{K} (2R) \|F_x - F_y\|_{H^\sigma} \\ &\leq \{2R + \textcolor{red}{K} \|\phi\|_{H^\sigma} + \varepsilon \textcolor{red}{K} R\} d(x, y). \end{aligned}$$

Finally, with this, (8) now gives

$$\begin{aligned} d(T(x), T(y))^2 &\leq \varepsilon^2 \left( 3 + 2 \|\phi\|_{H^\sigma}^2 \right) \{2R + K \|\phi\|_{H^\sigma} + \varepsilon KR\}^2 d(x, y)^2 \\ &\leq \varepsilon^2 C(R)^2 d(x, y)^2, \end{aligned}$$

where

$$C(R)^2 := \left( 3 + 2 \|\phi\|_{H^\sigma}^2 \right) \{2R + K \|\phi\|_{H^\sigma} + \varepsilon KR\}^2,$$

and we are done.

Now, using Lemma 2 we can show that, for sufficiently small  $\varepsilon$ , if the sequence  $x_n = T(x_{n-1})$  starts in  $B_R(0)$  it always stays there.

**Lemma 3** *Provided that  $\sigma > 1/2$ , if  $x_0 \in B_R(0)$  and  $x_n = T(x_{n-1})$  then, for any  $R > 0$ , there exists  $\varepsilon > 0$  such that  $x_n \in B_R(0)$  for all  $n \geq 0$ .*

*Proof* We work by induction. Given any  $x_0 \in B_R(0)$  we have

$$d(x_0, 0) \leq R,$$

and we are done with the base step. Now, suppose that  $x_n \in B_R(0)$  for all  $n < N$  and consider

$$\begin{aligned} d(x_N, 0) &= d(T(x_{N-1}), 0) \\ &\leq \varepsilon C(R) d(x_{N-1}, 0) \\ &\leq \varepsilon C(R) R \\ &\leq R, \end{aligned}$$

provided that  $\varepsilon < 1/C(R)$ .

With Lemma 3 in hand we are now able to prove our main result for Stokes waves.

**Theorem 2** *Provided that  $\sigma > 1/2$ , for  $\varepsilon$  sufficiently small the mapping  $T$  is a contraction on  $B_R(0)$  implying that, for  $\varepsilon$  sufficiently small, there exists a unique Stokes wave of the form (3).*

*Proof* Consider  $0 \leq q < 1$  and suppose that  $x, y \in B_R(0)$ . From Lemma 2 we have that

$$d(T(x), T(y)) \leq \varepsilon C(R) d(x, y) \leq q d(x, y),$$

provided that  $\varepsilon < q/C(R)$ .

#### 4 Wilton Ripples

We now turn to the case of Wilton **ripples** and the situation where there is a speed  $c_0$  such that the operator  $\mathcal{R}$  and its adjoint  $\mathcal{R}^*$  have two-dimensional null spaces spanned by  $\phi_m$  and  $\psi_m$  ( $m = 1, 2$ ) respectively, so that

$$\mathcal{R}\phi_m = 0, \quad \mathcal{R}^*\psi_m = 0, \quad \phi_m(x) = \psi_m(x) = \frac{\cos(mx)}{\sqrt{\pi}}; \quad m = 1, 2,$$

where we observe that

$$\langle \phi_m, \phi_\ell \rangle = \delta_{m,\ell}, \quad \langle \psi_m, \psi_\ell \rangle = \delta_{m,\ell}, \quad \langle \phi_m, \psi_\ell \rangle = \delta_{m,\ell},$$

and  $\delta_{m,\ell}$  is the Kronecker delta function.

##### 4.1 Preparing the Iteration: Linear Order

As before, we seek a solution of (2) of the form

$$f = \varepsilon f_1 + \varepsilon^2 F, \quad c = c_0 + \varepsilon s.$$

Inserting these into (2) gives

$$\mathcal{R}[\varepsilon f_1 + \varepsilon^2 F] = \left( \varepsilon s - \frac{1}{2}\varepsilon f_1 - \frac{1}{2}\varepsilon^2 F \right) (\varepsilon f_1 + \varepsilon^2 F).$$

At linear order we find

$$\mathcal{R}f_1 = 0$$

which demands that  $f_1 \in \mathcal{N}(\mathcal{R})$  so

$$f_1 = \alpha \phi_1 + \beta \phi_2, \quad \alpha, \beta \in \mathbf{R}.$$

As before, we may, without loss of generality, assume that  $\alpha = 1$ , however,  $\beta$  is still free to be chosen. This must be done at the next order.

##### 4.2 Preparing the Iteration: Quadratic Order

In contrast to the case of Stokes waves, for Wilton **ripples** additional initiation is required to *quadratic* order rather than just *linear*. For this, we enhance the form of our solution of (2) to be of the form

$$f = \varepsilon f_1 + \varepsilon^2 f_2 + \varepsilon^3 F, \quad c = c_0 + \varepsilon c_1 + \varepsilon^2 s. \quad (9)$$

Inserting these into (2) gives

$$\begin{aligned} & \mathcal{R}[\varepsilon f_1 + \varepsilon^2 f_2 + \varepsilon^3 F] \\ &= \left( \varepsilon c_1 + \varepsilon^2 s - \frac{1}{2}\varepsilon f_1 - \frac{1}{2}\varepsilon^2 f_2 - \frac{1}{2}\varepsilon^3 F \right) (\varepsilon f_1 + \varepsilon^2 f_2 + \varepsilon^3 F). \end{aligned} \quad (10)$$

By our choice of  $f_1$  this equation is satisfied at linear order for *any* choice of  $\beta$ , and we must study the quadratic order equation

$$\mathcal{R}f_2 - c_1f_1 = -\frac{1}{2}f_1^2.$$

The most general solution of this equation can be written

$$f_2 = a\phi_1 + b\phi_2 + \Phi, \quad \langle \Phi, \phi_1 \rangle = \langle \Phi, \phi_2 \rangle = 0,$$

and we follow the lead of [7] by selecting  $a = 0$ , and finding the triple  $\{\Phi, c_1, \beta\}$  which satisfies

$$\mathcal{R}\Phi - c_1f_1 = -\frac{1}{2}f_1^2.$$

Following the procedure outlined in [7] we insist that

$$0 = \left\langle c_1f_1 - \frac{1}{2}f_1^2, \psi_m \right\rangle, \quad m = 1, 2,$$

or, expanding,

$$0 = \left\langle c_1\phi_1 + c_1\beta\phi_2 - \frac{1}{2}\phi_1^2 - \beta\phi_1\phi_2 + \frac{1}{2}\beta^2\phi_2^2, \psi_m \right\rangle, \quad m = 1, 2.$$

If we define

$$C_{jm} := \langle \phi_j, \psi_m \rangle, \quad C_{j\ell m} := \langle \phi_j \phi_\ell, \psi_m \rangle,$$

we can show (c.f., [7], Lemma 4) that (from our choices of  $\{\phi_1, \phi_2, \psi_1, \psi_2\}$ )

$$C_{11} = C_{22} = 1, \quad C_{12} = C_{21} = 0, \quad (11a)$$

$$C_{112} = C_{121} = C_{211} = 1/2, \quad (11b)$$

$$C_{111} = C_{221} = C_{212} = C_{122} = C_{222} = 0. \quad (11c)$$

We find that for  $m = 1, 2$ , respectively,

$$c_1 - (1/2)\beta = 0, \quad c_1\beta - 1/4 = 0.$$

These can be solved sequentially to give

$$\beta = \pm \frac{1}{\sqrt{2}}, \quad c_1 = \pm \frac{1}{2\sqrt{2}}, \quad (12)$$

and since we have  $(c_1f_1 - (1/2)f_1^2)$  in the range of  $\mathcal{R}$  we can solve

$$\Phi = \mathcal{R}^{-1} \left[ c_1f_1 - \frac{1}{2}f_1^2 \right].$$

Before proceeding we note that, while we lose no generality by setting  $a = 0$  we cannot yet choose  $b$ , and this is one of the unknowns of our iteration. To summarize our developments thus far, we seek solutions of (2) of the form

$$f = \varepsilon(\phi_1 + \beta\phi_2) + \varepsilon^2(b\phi_2 + \Phi) + \varepsilon^3F, \quad c = c_0 + \varepsilon c_1 + \varepsilon^2s, \quad (13)$$

where  $\{c_0, \phi_1, \phi_2, \Phi, c_1, \beta\}$  are now known, and we must determine  $\{F, s, b\}$ .

**Remark 2** *We note that the solution pair  $\{c_1, \beta\}$  depends solely upon the choice of null space (spanned by  $\{\phi_1, \phi_2\}$ ) rather than the details of the linear operator  $\mathcal{L}$ . Naturally, the higher order corrections to our solution depend upon the form of  $\mathcal{L}$  quite strongly.*

**Remark 3** *Before continuing we point out that the analysis becomes somewhat more involved if the null space is spanned by  $\{\phi_1, \phi_m\}$  for  $m > 2$ . From preliminary investigations we believe that, for instance, in the case  $m = 3$  one would need to further “prepare” the iteration at cubic order before proceeding to the Wilton iteration that we describe in the next section, § 4.3.*

**Remark 4** *Finally, we note that we are seeking one particular type of Wilton ripple consisting of the resonance between two sinusoids of commensurate frequencies,  $(1, m)$ , where  $m$  is an integer. Of course there are other possibilities including non-resonant ripples with frequencies  $(\ell, m)$  where  $\ell, m$  are integers which do not divide each other [10].*

#### 4.3 The Wilton Iteration

Using these choices we return to (10) which is now satisfied at linear and quadratic orders. Using the fact that  $f_1, b\phi_2 \in \mathcal{N}(\mathcal{R})$  and dividing by  $\varepsilon^3$  we find

$$\begin{aligned} \mathcal{R}F &= \left\{ \left( c_1 - \frac{1}{2}f_1 \right) f_2 + \left( s - \frac{1}{2}f_2 \right) f_1 \right\} \\ &\quad + \varepsilon \left\{ \left( c_1 - \frac{1}{2}f_1 \right) F + \left( s - \frac{1}{2}f_2 \right) f_2 - \frac{1}{2}f_1 F \right\} \\ &\quad + \varepsilon^2 \left\{ \left( s - \frac{1}{2}f_2 \right) F - \frac{1}{2}f_2 F \right\} + \varepsilon^3 \left\{ -\frac{1}{2}F^2 \right\} \\ &= \{(c_1 - f_1)f_2 + sf_1\} + \varepsilon G(F, s, b), \end{aligned}$$

where

$$G(F, s, b; \varepsilon) := \left\{ (c_1 - f_1)F + \left( s - \frac{1}{2}f_2 \right) f_2 \right\} + \varepsilon \{(s - f_2)F\} + \varepsilon^2 \left\{ -\frac{1}{2}F^2 \right\}.$$

Using the fact that  $f_2 = b\phi_2 + \Phi$  and simplifying we find

$$\mathcal{R}F - sf_1 - (c_1 - f_1)b\phi_2 = (c_1 - f_1)\Phi + \varepsilon G(F, s, b; \varepsilon).$$

Inspired by this we conduct the following Wilton iteration

$$\mathcal{R}F_n - s_nf_1 - (c_1 - f_1)b_n\phi_2 = (c_1 - f_1)\Phi + \varepsilon G(F_{n-1}, s_{n-1}, b_{n-1}; \varepsilon). \quad (14)$$

Our procedure for solving (14) at each iteration is two-step:

1. First, we solve for the speed corrections  $\{s_n, b_n\}$  by projecting the right-hand-side of (14) and  $s_n f_1 + (c_1 - f_1)b_n \phi_2$  onto the range of  $\mathcal{R}$ . This is enforced by demanding that this be orthogonal to the null space of its adjoint,

$$0 = \langle s_n f_1 + (c_1 - f_1)b_n \phi_2 + (c_1 - f_1)\Phi + \varepsilon G(F_{n-1}, s_{n-1}, b_{n-1}; \varepsilon), \psi_m \rangle,$$

$m = 1, 2$ . Using (11) we find that

$$\begin{pmatrix} 1 & -1/2 \\ \beta & c_1 \end{pmatrix} \begin{pmatrix} s_n \\ b_n \end{pmatrix} = - \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix},$$

where

$$Q_m := \langle (c_1 - f_1)\Phi + \varepsilon G(F_{n-1}, s_{n-1}, b_{n-1}; \varepsilon), \psi_m \rangle, \quad m = 1, 2.$$

Using the fact that  $c_1 = \beta/2$ , c.f. (12), this can be solved *explicitly* as

$$s_n = -\frac{1}{2} \left( Q_1 + \frac{Q_2}{\beta} \right) \quad (15a)$$

$$b_n = -\frac{1}{2} \left( -2Q_1 + \frac{2Q_2}{\beta} \right). \quad (15b)$$

We also know from (12) that  $\beta = \pm 1/\sqrt{2}$  so that once a choice is made, these formulas give the unique solution.

2. With this choice of  $\{s_n, b_n\}$  we know that (14) is uniquely solvable provided that we demand orthogonality to the null space of  $\mathcal{R}$ . So, we can write

$$F_n = \mathcal{R}^{-1} [s_n f_1 + (c_1 - f_1)b_n \phi_2 + (c_1 - f_1)\Phi + \varepsilon G(F_{n-1}, s_{n-1}, b_{n-1}; \varepsilon)], \quad (16a)$$

$$\langle \phi_1, F_n \rangle = 0 \quad (16b)$$

$$\langle \phi_2, F_n \rangle = 0. \quad (16c)$$

This procedure can be replicated for the linearized version of this problem

$$\mathcal{R}F - s f_1 - (c_1 - f_1)b \phi_2 = (c_1 - f_1)\Phi + J,$$

and rigorous estimates derived for the unique solution. This is summarized in the following “elliptic estimate” [7].

**Theorem 3** *For any real  $\sigma \geq 0$ , given  $J \in H^\sigma$  there exists a unique solution triple  $\{F, s, b\}$  of*

$$\mathcal{R}F - s f_1 - (c_1 - f_1)b \phi_2 = (c_1 - f_1)\Phi + J,$$

$$\langle \phi_1, F \rangle = 0,$$

$$\langle \phi_2, F \rangle = 0,$$

*such that  $F \in H^{\sigma+\tau}$  and*

$$\max \{ \|F\|_{H^{\sigma+\tau}}, |s|, |b| \} \leq C_e \| (c_1 - f_1)\Phi + J \|_{H^\sigma}$$

for some  $C_e > 0$  and

$$\tau = \begin{cases} 1/2, & \text{Whitham (A),} \\ 1, & \text{Akers–Milewski (B),} \\ 2, & \text{Benjamin (C),} \\ 4, & \text{Kawahara (D).} \end{cases}$$

#### 4.4 Banach Fixed Point Formulation

We now write this as an FPI so that we can appeal to the Banach Fixed Point Theorem [30]. To accomplish this we write (2) as

$$\begin{aligned} \mathcal{R}F - sf_1 - (c_1 - f_1)b\phi_2 &= (c_1 - f_1)\Phi + \varepsilon G(\Phi, s, b; \varepsilon), \\ \langle \phi_1, F \rangle &= 0, \\ \langle \phi_2, F \rangle &= 0, \end{aligned}$$

or

$$M \begin{pmatrix} F \\ s \\ b \end{pmatrix} = \begin{pmatrix} (c_1 - f_1)\Phi + \varepsilon G(F, s, b; \varepsilon) \\ 0 \\ 0 \end{pmatrix}, \quad M := \begin{pmatrix} \mathcal{R} & -f_1 & -(c_1 - f_1)\phi_2 \\ \langle \phi_1, \cdot \rangle & 0 & 0 \\ \langle \phi_2, \cdot \rangle & 0 & 0 \end{pmatrix}.$$

We now express this as  $x = T(x)$  where

$$x := \begin{pmatrix} F \\ s \\ b \end{pmatrix}, \quad T(x) := M^{-1} \begin{pmatrix} (c_1 - f_1)\Phi + \varepsilon G(F, s, b; \varepsilon) \\ 0 \\ 0 \end{pmatrix},$$

and seek a fixed point of the iteration

$$x_n = T(x_{n-1}).$$

From Theorem 3 we know that  $M^{-1}$  is well-defined and that the action of  $x^* = T(x)$  can be given by

$$s^* = -\frac{1}{2} \left( Q_1(F, s, b) + \frac{Q_2(F, s, b)}{\beta} \right), \quad (17a)$$

$$b^* = -\frac{1}{2} \left( -2Q_1(F, s, b) + \frac{2Q_2(F, s, b)}{\beta} \right), \quad (17b)$$

$$F^* = \mathcal{R}^{-1} [s^*f_1 + (c_1 - f_1)b^*\phi_2 + (c_1 - f_1)\Phi + \varepsilon G(F, s, b; \varepsilon)], \quad (17c)$$

where

$$Q_m(F, s, B) := \langle (c_1 - f_1)\Phi + \varepsilon G(F, s, b; \varepsilon), \psi_m \rangle, \quad m = 1, 2, \quad (17d)$$

c.f. (15) and (16). A fixed point of the iteration  $x_n = T(x_{n-1})$  would solve  $x = T(x)$  and represent a solution of (2). To achieve this we appeal to Banach's Fixed Point Theorem [30] which requires a Banach space,

$$X = H^{\sigma+\tau} \times \mathbf{R} \times \mathbf{R},$$

and a distance,

$$d(x, y)^2 = \|F_x - F_y\|_{H^{\sigma+\tau}}^2 + |s_x - s_y|^2 + |b_x - b_y|^2.$$

We are now in a position to prove the central estimate which establishes that  $T$  is a contraction for  $\varepsilon$  small enough.

**Lemma 4** *Provided that  $\sigma > 1/2$  and  $x, y \in B_R(0) \subset X$  we have that*

$$d(T(x), T(y))^2 \leq \varepsilon^2 C(R)^2 d(x, y)^2,$$

where  $C(R)$  is a quadratic function of  $R$ .

*Proof* Exactly analogous to that of Lemma 2.

Just as we did for the Stokes waves, using Lemma 4 we can show that, for sufficiently small  $\varepsilon$ , if the sequence  $x_n = T(x_{n-1})$  starts in  $B_R(0)$  it always stays there, c.f. Lemma 3.

**Lemma 5** *Provided that  $\sigma > 1/2$ , if  $x_0 \in B_R(0)$  and  $x_n = T(x_{n-1})$  then, for any  $R > 0$ , there exists  $\varepsilon > 0$  such that  $x_n \in B_R(0)$  for all  $n \geq 0$ .*

This results in our final theorem which has the same proof as that of the Stokes result, Theorem 2.

**Theorem 4** *Provided that  $\sigma > 1/2$ , for  $\varepsilon$  sufficiently small the mapping  $T$  is a contraction on  $B_R(0)$  implying that, for  $\varepsilon$  sufficiently small, there exists a unique Wilton ripple of the form (9).*

## 5 Numerical Results

Inspired by the above results, we have implemented a numerical algorithm based on the FPI (14) which, we note, converges only for amplitudes up to a *finite* value, beyond which the iteration ceases to be a contraction. Due to the flexibility of not only our formulation but also our numerical implementation, we readily computed traveling Wilton ripples for four model PDEs which appear in the literature, the Kawaraha [40], Benjamin [15], Whitham [47], and Akers–Milewski [2] equations. We refer the reader to Table 1 for a detailed prescription of each of these four PDEs, in particular the “Spectral Gap” that we define as

$$\min_{k \geq 2} |\mathcal{L}(k) - \mathcal{L}(1)|,$$

which is significant as the radii  $R$  from Theorems 2 and 4 tend to zero as this gap vanishes.

Regarding our novel numerical FPI algorithm, we implemented a Fourier collocation approach [32, 18, 16, 54, 55] where the unknown functions,  $F_n(x)$ , in (14) were approximated by

$$F_n(x) \approx F_n^{N_x}(x) = \sum_{k=-N_x/2}^{N_x/2-1} a_{n,k} e^{ikx}, \quad a_{n,k} \approx \frac{1}{2\pi} \int_0^{2\pi} F_n(x) e^{-ikx} dx,$$

| Model                    | Operator                                               | $c_0$        | Spectral Gap                                             | Asymptotic             | $\varepsilon^+$ | $\varepsilon^-$ |
|--------------------------|--------------------------------------------------------|--------------|----------------------------------------------------------|------------------------|-----------------|-----------------|
| (A) Whitham (Deep Water) | $\hat{\mathcal{L}} = \sqrt{ k ^{-1} + \frac{1}{2} k }$ | $\sqrt{3/2}$ | $\sqrt{\frac{11}{6}} - \sqrt{\frac{3}{2}} \approx 0.129$ | $\mathcal{O}(k^{1/2})$ | 0.14            | 0.22            |
| (B) Akers–Milewski       | $\hat{\mathcal{L}} =  k ^{-1} + \frac{1}{2} k $        | $3/2$        | $1/3$                                                    | $\mathcal{O}(k)$       | 0.38            | 0.61            |
| (C) Benjamin             | $\hat{\mathcal{L}} =  k  - \frac{1}{3} k ^2$           | $2/3$        | $2/3$                                                    | $\mathcal{O}(k^2)$     | 1.48            | 0.96            |
| (D) Kawahara             | $\hat{\mathcal{L}} = k^2 - \frac{1}{5}k^4$             | $4/5$        | $8$                                                      | $\mathcal{O}(k^4)$     | 3.33            | 5.10            |

**Table 1** Listing of key parameters of the **four** models simulated in this paper: The Fourier symbol of the linear operator, the phase speed at which there are Wilton **ripples** (between wavenumbers  $k = 1$  and  $k = 2$ ), the spectral gap, the high frequency asymptotics, and an estimate for the maximal  $\varepsilon$  for which each branch converged for each model.

and (14) was enforced at the equally-spaced gridpoints  $\{x_j = 2\pi j/N_x\}$  for  $0 \leq j \leq N_x - 1$ . We began by choosing  $\{c_0, \phi_1, \phi_2, \Phi, c_1, \beta\}$  as outlined in § 4.1 and § 4.2, and initiated our iteration with  $\{F_0, s_0, b_0\} = \{0, 0, 0\}$ . For  $n \geq 1$  we conducted the iteration (14) which involved two steps: Utilizing (15) to determine the next  $\{s_n, b_n\}$  iterates, and then solving (16) to find  $F_n^{N_x}$  via the Discrete Fourier Transform (DFT)—accelerated by the Fast Fourier Transform (FFT) algorithm—which diagonalizes the operator  $\mathcal{R}$ . In these formulas derivatives and Fourier multipliers were evaluated in Fourier space with the DFT, and products were implemented in physical space; we did not notice appreciable errors due to aliasing. This iteration was continued until the Cauchy error,

$$e_N := \|f_N - f_{N-1}\|_{L^2} + |s_N - s_{N-1}| + |b_N - b_{N-1}|, \quad (18)$$


reached machine precision ( $\approx 2 \times 10^{-16}$ ).

We compared the results of this novel iteration scheme with a classic **quasi**-Newton iteration of (2) given the quadratic Wilton **ripple** as a starting guess. More specifically, we once again used a Fourier collocation method [32, 18, 16, 54, 55] where the unknown function,  $F(x)$ , in (2) was approximated by

$$F(x) \approx F^{N_x}(x) = \sum_{k=-N_x/2}^{N_x/2-1} a_k e^{ikx}, \quad a_k \approx \frac{1}{2\pi} \int_0^{2\pi} F(x) e^{-ikx} dx,$$

and a numerical continuation approach [12] applied to (2) with an initial guess provided by (13) with  $\{F, s, b\}$  set to zero. We note that both approaches produced the same branches of traveling Wilton **ripples**, provided that they converged. While the **quasi**-Newton solver converged for larger solutions, it was far more expensive due to the *formation* and *factorization* of the *full* Jacobian matrix. By contrast, our FPI was much more rapid, though its utility for larger solutions was more limited.

In Figure 1 we display results of our novel iteration scheme for the numerical simulation of (2) for the case of the Akers–Milewski equation (B in Table 1). On the left we display two traveling wave profiles, the solid curve corresponding to  $\beta = 1/\sqrt{2}$ , c.f. (12), and the dashed curve comes from  $\beta = -1/\sqrt{2}$ . The speed–amplitude curve for each of these two branches is plotted on the right of this figure with both our FPI scheme and the **quasi**-Newton approach



**Fig. 1 Left:** Example profiles from the FPI, (14), with  $\varepsilon = 0.01$  (the profile on branch with  $\beta = 1/\sqrt{2}$ , (12), is the solid curve, the profile on the branch with  $\beta = -1/\sqrt{2}$  is the dashed curve). **Right:** The speed–amplitude curves for two branches of ripples in the Akers–Milewski equation are depicted. The output of the FPI (triangles) is compared to that of the **quasi**–Newton iteration (solid lines).


[20]. The FPI converges for a finite range of amplitudes which we denote as  $\varepsilon^\pm$  corresponding to  $\beta = \pm 1/\sqrt{2}$ . **More specifically, the FPI converges for  $|\varepsilon| \leq \varepsilon^+$  for the choice  $\beta = 1/\sqrt{2}$ .** By “converge” we mean that the Cauchy error in the iteration is less than  $10^{-5}$  in under 5000 iterations and a finite limit has been found. We estimated these amplitudes for the **four** PDEs listed in Table 1 and report the results there.

We close with results on the estimated errors in our computations and the time required for our new FPI versus the classical **quasi**–Newton method. For the error, as there is no exact solution, we settled on the Cauchy measure, (18). On the left panel in Figure 2 we plot this versus iteration number  $N$ , while on the right of Figure 2 we display run times for the FPI (triangles) and the **quasi**–Newton method (circles) as a function of the number of spatial points,  $N_x$ . Over this range of points the cost of the **quasi**–Newton method is dominated by the formation of the Jacobian matrix,  $\mathcal{O}(N_x^2 \log N_x)$ , while the FPI has computational complexity  $\mathcal{O}(N_x \log N_x)$  (these asymptotic curves are depicted with solid lines in the figure).

## 6 Conclusion

In this paper we have undertaken the study of weakly nonlinear dispersive wave equations which admit Wilton **ripple** solutions. Not only do we demonstrate their existence rigorously using a Banach Fixed Point Iteration, but we also utilize a numerical simulation of this iteration scheme to produce highly accurate solutions in a fraction of the cost of a classical **quasi**–Newton approach.

The Wilton **ripple** problem has very similar structure to that of the spectral stability problem near an eigenvalue collision. This similarity suggests that very similar methods to those of this paper can be used to prove their existence



**Fig. 2 Left:** The Cauchy error, (18), measured for a Wilton ripple in each of the PDEs listed in Table 1 from the FPI (14) at  $\varepsilon = 0.01$ . The markers correspond to the PDEs as follows: (A) circles, (B) x's, (C) stars, (D) squares. **Right:** The run times for our novel FPI (triangles) and the quasi-Newton method (circles) as a function of the number of spatial points. The amplitude parameter  $\varepsilon = 0.01$  was simulated for the Akers–Milewski equation. The asymptotic curves  $N_x^2 \log N_x$  and  $N_x \log N_x$  are plotted with solid lines.

and compute instabilities. The tools of this paper do not require analyticity (in contrast to our previous work [7]) and thus have the potential to compute instabilities which are merely continuous, e.g., of Benjamin–Feir type.

### Conflict of interest

The authors declare that they have no conflict of interest.

### Data availability

Data will be made available on reasonable request.

### References

1. Adams, R.A.: Sobolev spaces. Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London (1975). Pure and Applied Mathematics, Vol. 65
2. Akers, B., Milewski, P.: A model equation for wavepacket solitary waves arising from capillary-gravity flows. *Stud. Appl. Math.* **122**, 249–274 (2009)
3. Akers, B., Nicholls, D.P.: Traveling waves with gravity and surface tension. *SIAM J. Appl. Math.* **70**, 2373–2389 (2010)
4. Akers, B., Nicholls, D.P.: Spectral stability of deep two-dimensional gravity water waves : repeated eigenvalues. *SIAM J. Appl. Math.* **72**, 689–711 (2012)
5. Akers, B., Nicholls, D.P.: Spectral stability of deep two-dimensional gravity capillary water waves. *Stud. Appl. Math.* **130**, 81–107 (2013)
6. Akers, B., Nicholls, D.P.: The spectrum of finite depth water waves. *European Journal of Mechanics-B/Fluids* **46**, 181–189 (2014)
7. Akers, B., Nicholls, D.P.: Wilton ripples in weakly nonlinear dispersive models of water waves: Existence and analyticity of solution branches. *Water Waves* **3**(1), 25–47 (2021)

8. Akers, B.F.: Modulational instabilities of periodic traveling waves in deep water. *Physica D: Nonlinear Phenomena* **300**, 26–33 (2015)
9. Akers, B.F.: High-order perturbation of surfaces short course: Stability of travelling water waves. *Lectures on the Theory of Water Waves* **426**, 51 (2016)
10. Akers, B.F., Ambrose, D.M., Sulon, D.W.: Periodic travelling interfacial hydroelastic waves with or without mass ii: Multiple bifurcations and ripples. *European Journal of Applied Mathematics* **30**(4), 756–790 (2019)
11. Akers, B.F., Gao, W.: Wilton ripples in weakly nonlinear model equations. *Communications in Mathematical Sciences* **10**(3), 1015–1024 (2012)
12. Allgower, E.L., Georg, K.: Numerical continuation methods. Springer-Verlag, Berlin (1990)
13. Ambrose, D.M., Strauss, W.A., Wright, J.D.: Global bifurcation theory for periodic traveling interfacial gravity-capillary waves. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **33**(4), 1081–1101 (2016)
14. Amundsen, D., Benney, D.: Resonances in dispersive wave systems. *Studies in Applied Mathematics* **105**(3), 277–300 (2000)
15. Bona, J.L., Albert, J.P., Restrepo, J.M.: Solitary-wave solutions of the Benjamin equation. *SIAM Journal on Applied Mathematics* **59**(6), 2139–2161 (1999)
16. Boyd, J.P.: Chebyshev and Fourier spectral methods, second edn. Dover Publications Inc., Mineola, NY (2001)
17. Bronski, J.C., Hur, V.M., Johnson, M.A.: Modulational instability in equations of KdV type. In: *New approaches to nonlinear waves*, pp. 83–133. Springer (2016)
18. Canuto, C., Hussaini, M.Y., Quarteroni, A., Zang, T.A.: Spectral methods in fluid dynamics. Springer-Verlag, New York (1988)
19. Christodoulides, P., Dias, F.: Resonant capillary-gravity interfacial waves. *J. Fluid Mech.* **265**, 303–343 (1994)
20. Claassen, K.M., Johnson, M.A.: Numerical bifurcation and spectral stability of wave-trains in bidirectional Whitham models. *Studies in Applied Mathematics* **141**(2), 205–246 (2018)
21. Craig, W., Nicholls, D.P.: Traveling two and three dimensional capillary gravity water waves. *SIAM J. Math. Anal.* **32**, 323–359 (2000)
22. Craig, W., Nicholls, D.P.: Traveling gravity water waves in two and three dimensions. *Eur. J. Mech. B/Fluids* **21**, 615–641 (2002)
23. Craik, A.: George Gabriel Stokes on water wave theory. *Ann. Rev. Fluid Mech.* **37**, 23–42 (2005)
24. Crandall, M.G., Rabinowitz, P.H.: Bifurcation from simple eigenvalues. *J. Functional Analysis* **8**, 321–340 (1971)
25. Creedon, R., Deconinck, B., Trichtchenko, O.: High frequency instabilities of a Boussinesq-Whitham: A perturbative approach. preprint (2020)
26. Creedon, R., Deconinck, B., Trichtchenko, O.: High frequency instabilities of the Kawahara equation: A perturbative approach. preprint (2020)
27. Deconinck, B., Oliveras, K.: The instability of periodic surface gravity waves. *J. Fluid Mech.* **675**, 141–167 (2011)
28. Deconinck, B., Trichtchenko, O.: Stability of periodic gravity waves in the presence of surface tension. *European Journal of Mechanics-B/Fluids* **46**, 97–108 (2014)
29. Ehrnström, M., Johnson, M.A., Maehlen, O.I., Remonato, F.: On the bifurcation diagram of the capillary-gravity Whitham equation. *Water Waves* **1**, 275–313 (2019)
30. Evans, L.C.: Partial differential equations, second edn. American Mathematical Society, Providence, RI (2010)
31. Folland, G.B.: Introduction to partial differential equations. Princeton University Press, Princeton, N.J. (1976). Preliminary informal notes of university courses and seminars in mathematics, Mathematical Notes
32. Gottlieb, D., Orszag, S.A.: Numerical analysis of spectral methods: theory and applications. Society for Industrial and Applied Mathematics, Philadelphia, Pa. (1977). CBMS-NSF Regional Conference Series in Applied Mathematics, No. 26
33. Harrison, W.: The influence of viscosity and capillarity on waves of finite amplitude. *Proceedings of the London Mathematical Society* **2**(1), 107–121 (1909)
34. Haupt, S.E., Boyd, J.P.: Modeling nonlinear resonance: A modification to the Stokes perturbation expansion. *Wave Motion* **10**(1), 83–98 (1988)

35. Henderson, D.M., Hammack, J.L.: Experiments on ripple instabilities. part 1. resonant triads. *Journal of Fluid Mechanics* **184**, 15–41 (1987)
36. Hur, V.M., Johnson, M.A.: Modulational instability in the Whitham equation for water waves. *Studies in Applied Mathematics* **134**(1), 120–143 (2015)
37. Hur, V.M., Johnson, M.A.: Modulational instability in the Whitham equation with surface tension and vorticity. *Nonlinear Analysis: Theory, Methods & Applications* **129**, 104–118 (2015)
38. Hur, V.M., Pandey, A.K.: Modulational instability in a full-dispersion shallow water model. *Studies in Applied Mathematics* **142**(1), 3–47 (2019)
39. Kamesvara, R.J.: On ripples of finite amplitude. *Proc. Indian Ass. Cultiv. Sci* **6**, 175–193 (1920)
40. Kawahara, T.: Oscillatory solitary waves in dispersive media. *Journal of the physical society of Japan* **33**(1), 260–264 (1972)
41. Ladyzhenskaya, O.A., Ural'tseva, N.N.: Linear and quasilinear elliptic equations. Academic Press, New York (1968)
42. Lamb, H.: *Hydrodynamics*, sixth edn. Cambridge University Press, Cambridge (1993)
43. Levi-Civita, T.: Détermination rigoureuse des ondes permanentes d'ampleur finie. *Math. Ann.* **93**, 264–314 (1925)
44. Longuet-Higgins, M.: Some relations between Stokes' coefficients in the theory of gravity waves. *J. Inst. Math. Appl.* **22**, 261 (1978)
45. McGoldrick, L.: An experiment on second-order capillary gravity resonant wave interactions. *J. Fluid Mech.* **40**, 251–271 (1970)
46. McGoldrick, L.: On Wilton's ripples: a special case of resonant interactions. *Journal of Fluid Mechanics* **42**(1), 193–200 (1970)
47. Moldabayev, D., Kalisch, H., Dutykh, D.: The Whitham equation as a model for surface water waves. *Physica D: Nonlinear Phenomena* **309**, 99–107 (2015)
48. Nicholls, D., Reitich, F.: Stable, high-order computation of traveling water waves in three dimensions. *Eur. J. Mech. B/Fluids* **25**, 406–424 (2006)
49. Nicholls, D.P., Reitich, F.: On analyticity of traveling water waves. *Proc. Roy. Soc. Lond., A* **461**(2057), 1283–1309 (2005)
50. Perlin, M., Henderson, D., Hammack, J.: Experiments on ripple instabilities. part 2 selective amplification of resonant triads. *Journal of Fluid Mechanics* **219**, 51–80 (1990)
51. Reeder, J., Shinbrot, M.: On Wilton ripples II: Rigorous results. *Arch. Rat. Mech. Anal.* **77**, 321–347 (1981)
52. Schwartz, L., Vanden-Broeck, J.M.: Numerical solution of the exact equations for capillary-gravity waves. *J. Fluid Mech.* **95**, 119 (1979)
53. Schwartz, L.W.: Computer extension and analytic continuation of Stokes expansion for gravity waves. *Journal of Fluid Mechanics* **62**(3), 553–578 (1974)
54. Shen, J., Tang, T.: Spectral and high-order methods with applications, *Mathematics Monograph Series*, vol. 3. Science Press Beijing, Beijing (2006)
55. Shen, J., Tang, T., Wang, L.L.: Spectral methods, *Springer Series in Computational Mathematics*, vol. 41. Springer, Heidelberg (2011). Algorithms, analysis and applications
56. Stokes, G.G.: On the theory of oscillatory waves. *Trans. Camb. Phil. Soc.* **8**, 441–455 (1847)
57. Struik, D.: Détermination rigoureuse des ondes irrotationnelles périodiques dans un canal à profondeur finie. *Math. Ann.* **95**, 595–634 (1926)
58. Toland, J.F., Jones, M.: The bifurcation and secondary bifurcation of capillary-gravity waves. *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences* **399**(1817), 391–417 (1985)
59. Trichtchenko, O., Deconinck, B., Kollár, R.: Stability of periodic traveling wave solutions to the Kawahara equation. *SIAM Journal on Applied Dynamical Systems* **17**(4), 2761–2783 (2018)
60. Vanden-Broeck, J.M.: Wilton ripples generated by a moving pressure disturbance. *J. Fluid Mech.* **451**, 193–201 (2002)
61. Wilton, J.: On ripples. *Phil. Mag.* **29**, 173 (1915)