Structure formulas for wave operators under a small scaling invariant condition

Marius Beceanu¹ and Wilhelm Schlag²

Abstract. We continue our work on the structure formula for the intertwining wave operators W_{\pm} associated with $H=-\Delta+V$ in \mathbb{R}^3 , cf. [3]. We consider small potentials relative to a scaling invariant norm.

Mathematics Subject Classification (2010). 35P25, 47A40, 47A10, 81U30.

Keywords. Intertwining wave operators.

1. Introduction

In a recent paper [3] we obtained a structure formula for the intertwining wave operators for $H = -\Delta + V$ in three dimensions. We imposed the following condition on the potential V. Define B^{β} , $\beta > 0$, as L^2 functions with

$$||f||_{B^{\beta}} := ||\mathbb{1}_{[|x| \le 1]} f||_2 + \sum_{j=0}^{\infty} 2^{j\beta} ||\mathbb{1}_{[2^j \le |x| \le 2^{j+1}]} f||_2 < \infty$$
 (1.1)

Then for V real-valued, $V \in B^{\beta}(\mathbb{R}^3)$, $\beta \geq \frac{1}{2}$, the wave operators

$$W_{\pm} = \lim_{t \to \pm \infty} e^{itH} e^{-itH_0}$$

exist in the strong L^2 sense, with $H_0 = -\Delta$. These operators satisfy for continuous, bounded f on the line, $f(H)W_{\pm} = W_{\pm}f(H_0)$, and $P_c = W_{\pm}W_{\pm}^*$, where P_c is the projection onto the absolutely continuous spectral subspace of H in L^2 .

 $^{^{1}}$ M. Beceanu thanks the University of Chicago for its hospitality during the summers of 2015 and 2016.

² W. Schlag was partially supported by NSF grant DMS-1500696 during the preparation of this work.

There is no singular continuous spectrum (asymptotic completeness). Yajima [6] and [7], established the L^p boundedness of the wave operators assuming that zero energy is neither an eigenvalue nor a resonance.

In [3] we proved the following theorem. By B^{1+} we mean B^{β} for some $\beta > 1$. By \mathcal{M}_{ν} we mean the Borel measures on Euclidean space.

Theorem 1.1 ([3]). Let $V \in B^{1+}$ be real-valued and assume that $H = -\Delta + V$ admits no eigenfunction or resonance at zero energy. Then there exists $g(x, y, \omega) \in L^1_\omega \mathcal{M}_y L^\infty_x$, i.e.,

$$\int\limits_{\mathbb{S}^2}\|g(x,dy,\omega)\|_{\mathcal{M}_yL^\infty_x}\,d\omega<\infty$$

such that for any $f \in L^2$ one has the representation formula

$$(W_+f)(x) = f(x) + \int_{\mathbb{S}^2} \int_{\mathbb{R}^3} g(x, dy, \omega) f(S_{\omega}x - y) d\omega.$$

where $S_{\omega}x = x - 2(x \cdot \omega)\omega$ is a reflection. A similar result holds for W_{-} .

As an application, suppose X is any Banach space of measurable functions on \mathbb{R}^3 which is invariant under translations and reflections, and in which Schwartz functions are dense. Assume that $\|\mathbb{1}_H f\|_X \leq A\|f\|_X$ for all half spaces $H \subset \mathbb{R}^3$ and $f \in X$ with some uniform constant A. Then

$$||W_+ f||_X \le A C(V) ||f||_X \quad \text{for all } f \in X$$
 (1.2)

where C(V) is a constant depending on V alone. In particular, this recovers Yajima's L^p boundedness of the wave operators. Furthermore, [3] obtains quantitative estimates on the norm in (1.2) as well as on C(V). These bounds blow up as $\beta \to 1$ with $V \in B^{\beta}$. Nevertheless, we remark that one can obtain Theorem 1.1, albeit without quantitative control, under the condition $V \in B^1$ although the details are not worked out in [3].

The goal here is to seek a scaling invariant condition on V under which a structure formula (1.6) can be obtained. The natural scaling of the Schrödinger operator $H = -\Delta + V$ is $V \to \lambda^2 V(\lambda x)$, $\lambda > 0$ in any dimension. In the framework of the B-spaces above (1.1) the critical norm relative to this scaling is $\dot{B}^{\frac{1}{2}}(\mathbb{R}^3)$ where

$$||f||_{\dot{B}^{\beta}} := \sum_{j=-\infty}^{\infty} 2^{j\beta} ||1_{[2^{j} \le |x| \le 2^{j+1}]} f||_{2}$$
(1.3)

This norm is invariant under the aforementioned scaling provided λ is a power of 2. However, it is currently unclear whether Theorem 1.1 might hold for potentials $V \in \dot{B}^{\frac{1}{2}}$. It is possible that the threshold $\beta = 1$ could be optimal for the B^{β} spaces. It is natural to investigate the scaling invariant class for several reasons: (i) it is an optimal scenario, corresponding to the $|x|^{-2}$ decay rate which balances the Laplacian (ii) it arises in widely-studied energy critical nonlinear equations such as the u^5 wave equation in \mathbb{R}^3 :

$$u_{tt} - \Delta u - u^5 = 0$$

which admits explicit the 1-parameter stationary solutions $u(t,x) = \lambda^{\frac{1}{2}}W(\lambda x) =$: $W_{\lambda}(x), \ \lambda > 0, \ W(x) = (1+|x|^2/3)^{-\frac{1}{2}}$. In the radial class, these are the only stationary nonzero solutions of finite energy. Linearizing about W_{λ} yields the family of Schrödinger operators $H_{\lambda} = -\Delta - \lambda^2 W^4(\lambda x)$. It is therefore desirable to work with a condition on the potential that is uniform in $\lambda > 0$.

This paper presents such a norm, but currently we only consider small potentials in this norm. To formulate it, we recall some notation.

Definition 1.1. For any Schwartz function V we define $||V|| = ||L_V||_{L^1_{t,x}}$, where

$$L_V(t,\omega) = \int_0^\infty \hat{V}(-\tau\omega)e^{\frac{i}{2}t\tau} \tau d\tau$$

is as above. For any Schwartz function v in \mathbb{R}^3

$$||v||_{B} := \sup_{\Pi} \int_{-\infty}^{\infty} |||\delta_{\Pi(t)} v(x)||| dt$$
 (1.4)

where Π is a 2-dimensional plane through the origin, and $\Pi(t) = \Pi + t\vec{N}, \vec{N}$ being the unit norm to Π .

Clearly, $||v|| \le ||v||_B$. We will show below that $||v||_B < \infty$ is finite for Schwartz functions. We will do this by dominating $||v||_B$ by a stronger norm which is also scaling invariant and more explicit, cf. Lemma 3.1. This more explicit norm involves half of a derivative on 2-planes. So it is not a pure decay condition on the potential. The key analytical arguments in this paper are based on the precise norm as defined in (1.4). The main result of this paper is the following one.

Theorem 1.2. There exists $c_0 > 0$ with the following property: for any real-valued V with $\|V\|_B + \|V\|_{\dot{B}^{\frac{1}{2}}} \le c_0$, there exists $g(x, y, \omega) \in L^1_\omega \mathcal{M}_y L^\infty_x$ with

$$\int_{\mathbb{S}^2} \|g(x, dy, \omega)\|_{\mathcal{M}_y L_x^{\infty}} d\omega \lesssim c_0$$
(1.5)

such that for any $f \in L^2$ one has the representation formula

$$(W_+ f)(x) = f(x) + \int_{\mathbb{S}^2} \int_{\mathbb{R}^3} g(x, dy, \omega) f(S_\omega x - y) d\omega.$$
 (1.6)

where $S_{\omega}x = x - 2(x \cdot \omega)\omega$ is a reflection. A similar result holds for W_{-} .

The spectral properties of $H = -\Delta + V$ are irrelevant under the smallness assumption. For $L^1 \to L^\infty$ dispersive estimates of the Schrödinger evolution $e^{itH} P_c$ with a scaling-invariant condition on V in \mathbb{R}^3 without any smallness assumption, see [2]. It is not clear to the authors if there might be other scaling-invariant norms which are better suited for structure theorems for the wave operators. This question is particularly relevant with respect to large potentials and the Wiener formalism that is instrumental for Theorem 1.1.

2. The wave operators and their expansion

We now recall the formalism of the wave operator W_+ going back to Kato [5]. First, by [3, Lemma 2.2], $\dot{B}^{\frac{1}{2}} \hookrightarrow L^{\frac{3}{2},1}$ (using Lorentz space notation). If $V \in L^{\frac{3}{2},1}$, then the wave operators W_{\pm} exist, and are isometries from L^2 onto the range of P_c , the projection onto the continuous spectrum of H. Moreover, if V is small in $L^{\frac{3}{2},1}$, then H has no eigenvalues and no zero energy resonance, and the spectrum is purely absolutely continuous. In other words, W_{\pm} are unitary operators. Moreover, for any $f \in L^2(\mathbb{R}^3)$ the integral

$$W_{+}f = f + i \int_{0}^{\infty} e^{itH} V e^{-itH_{0}} f dt$$
 (2.1)

converges in the strong sense. See for example Section 4 of [3] for more details.

Expanding (2.1) iteratively by means of the Duhamel formula one has

$$W_{+}f = f + W_{1+}f + \dots + W_{n+}f + \dots,$$

$$W_{1+}f = i \int_{t>0} e^{-it\Delta}Ve^{it\Delta}f \,dt,$$

$$\vdots$$

$$W_{n+}f = i^{n} \int_{t>s_{1}>\dots>s_{n-1}>0} e^{-i(t-s_{1})\Delta}Ve^{-i(s_{1}-s_{2})\Delta}V \dots e^{-is_{n-1}\Delta}Ve^{it\Delta}f \,dt \,ds_{1}\dots ds_{n-1}$$

$$t>s_{1}>\dots>s_{n-1}>0$$
(2.2)

for $f \in L^2$. For small potentials one can actually sum this series, which will give Theorem 1.2. In addition to the operators W_{n+} , we shall work with their regularized version,

$$W_{n+}^{\varepsilon} f := i^{n} \int e^{i(t_{n} - t_{n-1})H_{0} - \varepsilon(t_{n} - t_{n-1})} V \dots e^{i(t_{2} - t_{1})H_{0} - \varepsilon(t_{2} - t_{1})} V e^{it_{1}H_{0} - \varepsilon t_{1}} V e^{-it_{n}H_{0}} f dt_{1} \dots dt_{n},$$
(2.3)

where $\varepsilon > 0$. By [3, Lemma 4.3], $W_{n+}^{\varepsilon} \to W_{n+}$ in the strong L^2 sense as $\varepsilon \to 0+$. One has the following representation formulas for each $n \ge 1$ going back to Yajima, see [3, Lemma 4.7]:

$$\langle W_{n+}^{\varepsilon} f, g \rangle = (-1)^n \int_{\mathbb{R}^9} \mathcal{F}_{x_0}^{-1} T_{n+}^{\varepsilon}(0, x, y) f(x - y) \overline{g}(x) \, dy \, dx \tag{2.4}$$

where for any $\varepsilon > 0$, $T_{1\pm}^{\varepsilon}(x_0, x_1, y)$ is defined in the sense of distributions as

$$(\mathcal{F}_{x_0}^{-1}\mathcal{F}_{x_1,y}T_{1\pm}^{\varepsilon})(\xi_0,\xi_1,\eta) := \frac{\widehat{V}(\xi_1 - \xi_0)}{|\xi_1 + \eta|^2 - |\eta|^2 \pm i\varepsilon}$$
(2.5)

and, more generally, for all $n \ge 1$ we have

$$(\mathcal{F}_{x_0}^{-1}\mathcal{F}_{x_n,y}T_{n\pm}^{\varepsilon})(\xi_0,\xi_n,\eta) := \int_{\mathbb{R}^{3(n-1)}} \frac{\prod_{\ell=1}^n \widehat{V}(\xi_{\ell} - \xi_{\ell-1}) d\xi_1 \dots d\xi_{n-1}}{\prod_{\ell=1}^n (|\xi_{\ell} + \eta|^2 - |\eta|^2 \pm i\varepsilon)}.$$
 (2.6)

Even though the first variable does not play a role in (2.4), it is essential in order to express T_{n+}^{ε} in terms of T_{1+}^{ε} by means of a convolution structure. In fact, we (formally) compose three variable kernels $T(x_0, x_1, y)$ on \mathbb{R}^9 by the rule

$$(T_1 \circledast T_2)(x_0, x_2, y) = \int_{\mathbb{R}^6} T_1(x_0, x_1, y_1) T_2(x_1, x_2, y - y_1) \, dx_1 \, dy_1. \tag{2.7}$$

Dually (on the Fourier side), ® is given by

$$(\mathcal{F}_{x_0}^{-1}\mathcal{F}_{x_2,y}(T_1 \circledast T_2))(\xi_0, \xi_2, \eta) = \int_{\mathbb{R}^3} (\mathcal{F}_{x_0}^{-1}\mathcal{F}_{x_1,y}T_1)(\xi_0, \xi_1, \eta)(\mathcal{F}_{x_1}^{-1}\mathcal{F}_{x_2,y}T_2)(\xi_1, \xi_2, \eta) d\xi_1.$$
 (2.8)

So \circledast consists of convolution in the y variable (or multiplication in the dual variable η), and composition of operators relative to the other two variables. In the dual coordinates ξ_0 , ξ_1 , and ξ_2 , composition of operators is preserved. Note the order of the variables: x_0 is the "input", x_2 the "output" variable, whereas y is the dual energy variable.

Lemma 2.1. Let V be a Schwartz potential. For all $\varepsilon > 0$ and any $n, m \ge 1$

$$T_{m+}^{\varepsilon} \circledast T_{n+}^{\varepsilon} = T_{(m+n)+}^{\varepsilon}$$

in the sense of (2.8).

Proof. By inspection

$$\mathcal{F}_{x_0}^{-1} \mathcal{F}_{x_2,y} T_{2+}^{\varepsilon}(\xi_0, \xi_2, \eta) = \int_{\mathbb{R}^3} \frac{\widehat{V}(\xi_2 - \xi_1)}{|\xi_2 + \eta|^2 - |\eta|^2 + i\varepsilon} \cdot \frac{\widehat{V}(\xi_1 - \xi_0)}{|\xi_1 + \eta|^2 - |\eta|^2 + i\varepsilon} d\xi_1
= \mathcal{F}_{x_0}^{-1} \mathcal{F}_{x_2,y} (T_{1+}^{\varepsilon} \otimes T_{1+}^{\varepsilon})(\xi_0, \xi_2, \eta)$$
(2.9)

both in the pointwise sense, as well as in the space of distributions. The general case follows by induction. \Box

In order to prove Theorem 1.2, we will show that there exists an algebra under \circledast with the norms of Definition 1.1. In [3, Section 5] it was shown that

$$W_{1+}^{\varepsilon} f(x) = \int_{\mathbb{R}^3} K_{1+}^{\varepsilon}(x, x - y) f(y) dy,$$

$$K_{1+}^{\varepsilon}(x, z) = -\lim_{R \to \infty} \int_{\mathbb{R}^6} \frac{e^{ix \cdot \xi} \hat{V}(\xi) e^{iz \cdot \eta}}{|\xi + \eta|^2 - |\eta|^2 + i\varepsilon} e^{-\frac{|\eta|^2}{2R^2}} d\xi d\eta.$$
(2.10)

Furthermore, the kernels $K_{1+}^{\varepsilon}(x,z)$ are of the form,

$$K_{1+}^{\varepsilon}(x,z) = \operatorname{const} \cdot L_{\varepsilon}(|z| - 2x \cdot \hat{z}, z)$$

where

$$L_{\varepsilon}(r,\omega) := \int_{0}^{\infty} \widehat{V}(-s\omega)e^{i\frac{rs}{2}|\omega|}e^{-\frac{\varepsilon}{2s}} s ds.$$
 (2.11)

In the previous line ω need not be a unit vector. Sending $\varepsilon \to 0$, one has

$$K_{1+}^{0}(x,z) = \operatorname{const} \cdot |z|^{-2} L(|z| - 2x \cdot \hat{z}, \hat{z})$$
 (2.12)

where for any $\omega \in \mathbb{S}^2$, and $r \in \mathbb{R}$,

$$L(r,\omega) = L_V(r,\omega) := \int_0^\infty \hat{V}(-s\omega)e^{i\frac{rs}{2}} s \, ds.$$

Here V is any Schwartz function. The following corollary from [3] shows how the structure function for W_{1+} arises easily from this formalism. It also explains how the norm $\|\cdot\|$ arises in Definition 1.1.

Corollary 2.2. Let V be a Schwartz function. Define $S_{\omega} := x - 2(\omega \cdot x)\omega$ to be the reflection about the plane ω^{\perp} . Then for all Schwartz functions f

$$(W_{1+}f)(x) = \int_{\mathbb{S}^2} \int_{\mathbb{R}^3} g_1(x, dy, \omega) f(S_{\omega}x - y) \, \sigma(d\omega). \tag{2.13}$$

For fixed $x \in \mathbb{R}^3$, $\omega \in \mathbb{S}^2$ the function $g_1(x,\cdot,\omega)$ is a measure satisfying

$$\int_{\mathbb{S}^2} \|g_1(x, dy, \omega)\|_{\mathcal{M}_y L_x^{\infty}} d\omega \le \int_{\mathbb{S}^2} \int_{\mathbb{R}} |L_V(r, \omega)| dr d\omega = ||V||$$
(2.14)

with $\|\cdot\|_{\mathfrak{M}}$ being the total variation norm for Borel measures.

Proof. From eq. (2.10) and (2.12),

$$(W_{1+}f)(x) = \int_{0}^{\infty} \int_{\mathbb{S}^2} L(r - 2\omega \cdot x, \omega) f(x - r\omega) dr d\omega$$

$$= \int_{\mathbb{S}^2} \int_{\mathbb{R}} \mathbb{1}_{[r > -2\omega \cdot x]} L(r, \omega) f(x - 2(\omega \cdot x)\omega - r\omega) dr d\omega.$$
(2.15)

Define

$$g_1(x, dy, \omega) := \mathbb{1}_{[(y+2x)\cdot\omega>0]} L(y\cdot\omega, \omega) \mathcal{H}^1_{\ell_\omega}(dy). \tag{2.16}$$

Here $\ell_{\omega} := \{r\omega \mid r \in \mathbb{R}\}$ is the line along ω , and $\mathcal{H}^1_{\ell_{\omega}}$ is the 1-dimensional Hausdorff measure on the line ℓ_{ω} . Then (2.13) holds and

$$||g_1(x, dy, \omega)||_{L_x^{\infty}} = |L(y \cdot \omega, \omega)| \mathcal{H}_{\ell_{\omega}}^1(dy)$$
(2.17)

which implies (2.14).

This result does not explain the origin of the other norm, $\|V\|_B$ in Definition 1.1. That norm is needed to bound the higher order structure functions g_n , $n \geq 2$. The remainder of this paper will be devoted to working out the details of this construction. To end this section, we recall how [3] fails to reach the scaling-invariant space $\dot{B}^{\frac{1}{2}}$ and we explain how $\|V\|_B$ is designed to circumvent the exact difficulty responsible for the loss of $\frac{1}{2}$ power in Theorem 1.1.

First, we point out the connection between $\|\cdot\|$ and $\|\cdot\|_{\dot{B}^{\frac{1}{2}}}$ as given by [3, Proposition 6.1].

Proposition 2.3. Let $L = L_V$ be as above, and V a Schwartz function. Then, with $r \in \mathbb{R}$ and $\omega \in \mathbb{S}^2$,

$$||L(r,\omega)||_{L^2_{r,\omega}} \lesssim ||V||_{L^2}$$
 (2.18)

and

$$||L(r,\omega)||_{L^{1}_{r,\omega}} \lesssim \sum_{k \in \mathbb{Z}} 2^{k/2} ||\mathbb{1}_{[2^{k},2^{k+1}]}(|r|)L(r,\omega)||_{L^{2}_{r,\omega}} \lesssim ||V||_{\dot{B}^{\frac{1}{2}}}.$$
 (2.19)

Moreover, for any $0 < \alpha < 1$,

$$\sum_{k \in \mathbb{Z}} 2^{\alpha k} \| \mathbb{1}_{[2^k, 2^{k+1}]}(|r|) L(r, \omega) \|_{L^2_{r, \omega}} \lesssim \|V\|_{\dot{B}^{\alpha}}. \tag{2.20}$$

The aforementioned loss of a $\frac{1}{2}$ power occurred in the following estimate (6.9) from [3]:

$$\|v(x)K_{1+}^{\varepsilon}(x,y)\|_{L_{v}^{1}B_{x}^{\frac{1}{2}}}\lesssim \|v\|_{B^{1}}\|V\|_{B^{\frac{1}{2}}}.$$

In view of (2.12) this is the same as (in the limit $\varepsilon \to 0$)

$$\int_{S^2} \int_{0}^{\infty} \|v(x)L_V(t-2x\cdot\omega,\omega)\|_{B^{\frac{1}{2}}} dt d\omega \lesssim \|v\|_{B^1} \|V\|_{B^{\frac{1}{2}}}.$$

We now show how to avoid this loss by means of the norm (1.4).

Lemma 2.4. For Schwartz functions v, V one has

$$\int_{S^2} \int_{0}^{\infty} |||v(x)L_V(t - 2x \cdot \omega, \omega)||| \, dt \, d\omega \lesssim ||v||_B |||V|||. \tag{2.21}$$

Proof. Writing $\omega^{\perp}(s) = \omega^{\perp} + s\omega$, we compute

$$\int_{S^{2}} \int_{0}^{\infty} \|v(x)L_{V}(t-2x\cdot\omega,\omega)\| dt d\omega$$

$$\leq \int_{S^{2}-\infty} \int_{0}^{\infty} \|\delta_{\omega^{\perp}(s)}v(x)L_{V}(t-2x\cdot\omega,\omega)\| dt ds d\omega$$

$$\leq \int_{S^{2}-\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |L_{V}(t-2s,\omega)| \|\delta_{\omega^{\perp}(s)}v(x)\| dt ds d\omega$$

$$\leq \int_{S^{2}-\infty} \int_{-\infty}^{\infty} |L_{V}(t,\omega)| dt d\omega \sup_{\omega \in \mathbb{S}^{2}-\infty} \int_{-\infty}^{\infty} \|\delta_{\omega^{\perp}(s)}v(x)\| ds$$

$$= \|V\| \|v\|_{B}$$
(2.22)

which is (2.21).

Before continuing with the main argument, the following section exhibits norms that dominate those in Definition 1.1, but which are more explicit. We also check that $\|\cdot\|_B$ is scaling invariant.

3. A closer look at the norms of Definition 1.1

The following lemma bounds the rather implicit $\|\cdot\|_B$ -norm by a more explicit Sobolev-type norm. It appears that this cannot be improved significantly.

Lemma 3.1. The norm $\|\cdot\|_B$ is scaling invariant in the sense that with $v_{\lambda}(x) := \lambda^{-2}v(x/\lambda)$, $\lambda > 0$, one has $\|v\|_B = \|v_{\lambda}\|_B$. Furthermore,

$$\|v\|_{B} \le C \sup_{\omega \in \mathbb{S}^{2}} \int_{-\infty}^{\infty} \sum_{k \in \mathbb{Z}} 2^{\frac{k}{2}} \|\psi(2^{-k}x')v(x'+s\omega)\|_{\dot{H}^{\frac{1}{2}}(\omega^{\perp})} ds =: \|v\|_{B}^{*}$$
 (3.1)

where $x' \in \omega^{\perp}$ and $\sum_{k \in \mathbb{Z}} \psi(2^{-k}x') = 1$ for $x' \in \omega^{\perp} \setminus \{0\}$ is a Littlewood–Paley partition of unity. The norm $\|v\|_B^*$ is finite on Schwartz functions, and $\|v\|_B^* = \|v_{\lambda}\|_B^*$ for $\lambda = 2^{-\ell}$, $\ell \in \mathbb{Z}$.

Proof. One has $L_V(t,\omega) = \lambda^{-1} L_{V_\lambda}(t\lambda^{-1},\omega)$ whence $||V_\lambda||| = |||V|||$. With χ a standard bump function on the line,

$$\begin{split} \|V_{\lambda}\|_{B} &= \sup_{\omega \in \mathbb{S}^{2}} \lim_{\delta \to 0} \int_{-\infty}^{\infty} \|\delta^{-1}\chi((x \cdot \omega - t)/\delta)\lambda^{-2}V(x/\lambda)\|_{L} dt \\ &= \sup_{\omega \in \mathbb{S}^{2}} \lim_{\delta \to 0} \int_{-\infty}^{\infty} \lambda^{-1} \|(\delta/\lambda)^{-1}\chi(((x/\lambda) \cdot \omega - t/\lambda)/(\delta/\lambda))\lambda^{-2}V(x/\lambda)\|_{L} dt \\ &= \sup_{\omega \in \mathbb{S}^{2}} \lim_{\delta \to 0} \int_{-\infty}^{\infty} \|\delta^{-1}\chi(((x/\lambda) \cdot \omega - t)/\delta)\lambda^{-2}V(x/\lambda)\|_{L} dt \\ &= \sup_{\omega \in \mathbb{S}^{2}} \lim_{\delta \to 0} \int_{-\infty}^{\infty} \|\delta^{-1}\chi((x \cdot \omega - t)/\delta)V(x)\|_{L} dt = \|V\|_{B} \end{split}$$

which is the scaling invariance of the *B*-norm.

To prove (3.1) we fix the plane Π to be $x_1=0$, or equivalently we set $\omega=(1,0,0)$. One has, with $v_s=\delta_{\Pi(s)}v$,

$$L_{v_s}(t,\omega) = \int_0^\infty \mathcal{F}_{x_2,x_3} v(s,-\omega_2\tau,-\omega_3\tau) e^{i\frac{\tau}{2}(t-2s\omega_1)} \tau d\tau$$

so that

$$\|v\|_{B} = \int_{-\infty}^{\infty} \left\| \int_{0}^{\infty} \mathcal{F}_{x_{2},x_{3}} v(s, -\omega_{2}\tau, -\omega_{3}\tau) e^{i\frac{\tau}{2}t} \tau \, d\tau \right\|_{L_{t,\omega}^{1}} ds$$

$$= \int_{-\infty}^{\infty} ds \int_{-\infty}^{\infty} dt \int_{0}^{\pi} \int_{0}^{2\pi} \left| \int_{0}^{\infty} \mathcal{F}_{x_{2},x_{3}} v(s, -\tau \sin \theta \sin \phi, -\tau \sin \theta \cos \phi) e^{i\frac{\tau}{2}t} \tau \, d\tau \right| \sin \theta \, d\phi d\theta$$

$$= \pi \int_{-\infty}^{\infty} ds \int_{-\infty}^{\infty} dt \int_{0}^{\pi} \left| \int_{0}^{\infty} \mathcal{F}_{x_{2},x_{3}} v(s, -\tau \sin \phi, -\tau \cos \phi) e^{i\frac{\tau}{2}t} \tau \, d\tau \right| d\phi$$

$$\leq C \sum_{k \in \mathbb{Z}_{-\infty}} \int_{0}^{\infty} 2^{\frac{k}{2}} \left(\int_{-\infty}^{\infty} \int_{0}^{\pi} \mathbb{1}_{\left[|t| \simeq 2^{k}\right]} \right. \\ \left. \left| \int_{0}^{\infty} \mathcal{F}_{x_{2}, x_{3}} v(s, -\tau \sin \phi, -\tau \cos \phi) e^{i\frac{\tau}{2}t} \tau \, d\tau \right|^{2} d\phi dt \right)^{\frac{1}{2}} ds.$$

For a Schwartz function w in \mathbb{R}^2 define the sublinear operator $A_k w$ as

$$A_k w = \left(\int_{-\infty}^{\infty} \int_{0}^{\pi} \mathbb{1}_{[|t| \simeq 2^k]} \middle| \int_{0}^{\infty} \hat{w}(\tau \sin \phi, \tau \cos \phi) e^{i\frac{\tau}{2}t} \tau d\tau \middle|^2 d\phi dt \right)^{\frac{1}{2}}.$$
 (3.2)

Then, on the one hand,

$$\sum_{k \in \mathbb{Z}} (A_k w)^2 \lesssim \int_{-\infty}^{\infty} \int_{0}^{\pi} \left| \int_{0}^{\infty} \hat{w}(\tau \sin \phi, \tau \cos \phi) e^{i\frac{\tau}{2}t} \tau d\tau \right|^2 d\phi dt$$

$$\lesssim \int_{0}^{\pi} \int_{0}^{\infty} |\hat{w}(\tau \sin \phi, \tau \cos \phi)|^2 |\tau|^2 d\tau d\phi = ||w||_{\dot{H}^{\frac{1}{2}}}^2$$
(3.3)

and, on the other hand,

$$\sum_{k \in \mathbb{Z}} 2^{2k} (A_k w)^2 \lesssim \int_{-\infty}^{\infty} \int_{0}^{\pi} \left| t \int_{0}^{\infty} \hat{w}(\tau \sin \phi, \tau \cos \phi) e^{i\frac{\tau}{2}t} \tau d\tau \right|^2 d\phi dt$$

$$\lesssim \int_{0}^{\pi} \int_{0}^{\infty} \left| \partial_{\tau}(\tau \hat{w}(\tau \sin \phi, \tau \cos \phi)) \right|^2 d\tau d\phi$$

$$\lesssim \int_{0}^{\pi} \int_{0}^{\infty} \left[\tau^{-1} |\hat{w}(\tau \sin \phi, \tau \cos \phi)|^2 + |(\partial_{r} \hat{w})(\tau \sin \phi, \tau \cos \phi)|^2 \tau \right] \tau d\tau d\phi$$

$$\lesssim \|\hat{w}\|_{\dot{H}^{\frac{1}{2}}}^2 + \||\xi|^{\frac{1}{2}} \partial_{r} \hat{w}\|_{2}^2. \tag{3.4}$$

The first term in the last line is obtained by Hardy's inequality in the ξ variable, and we bound it further by applying Hardy's inequality in the x variable:

$$\|\hat{w}\|_{\dot{H}^{\frac{1}{2}}} = \||x|^{\frac{1}{2}}w\|_{2} = \||x|^{-\frac{1}{2}}|x|w\|_{2} \lesssim \||x|w\|_{\dot{H}^{\frac{1}{2}}}.$$
 (3.5)

For second term in (3.4) we first rewrite $\partial_r \hat{w}$ as

$$\partial_{r}\hat{w}(\xi) = \frac{\xi}{|\xi|} \nabla_{\xi} \int_{\mathbb{R}^{2}} e^{-ix\cdot\xi} w(x) dx$$

$$= \int_{\mathbb{R}^{2}} e^{-ix\cdot\xi} a(x,\xi) |x| w(x) dx, \quad a(x,\xi) = \frac{x}{|x|} \cdot \frac{\xi}{|\xi|} =: \hat{x} \cdot \hat{\xi}.$$
(3.6)

Therefore,

$$\||\xi|^{\frac{1}{2}}\partial_r \hat{w}\|_2 \le \|\hat{x}|x|w\|_{\dot{H}^{\frac{1}{2}}}.$$
(3.7)

By Lemma 3.2 below one has $\|\hat{x}|x|w\|_{\dot{H}^{\frac{1}{2}}} \lesssim \||x|w\|_{\dot{H}^{\frac{1}{2}}}$. Combining this bound with (3.5) we conclude that

$$\sum_{k \in \mathbb{Z}} 2^{2k} (A_k w)^2 \lesssim \||x|w\|_{\dot{H}^{\frac{1}{2}}}^2. \tag{3.8}$$

By interpolation

$$\sum_{k \in \mathbb{Z}} 2^{\frac{k}{2}} A_k w \lesssim \|w\|_* \tag{3.9}$$

where in the notation of the real interpolation method

$$||w||_* = (||w||_{\dot{H}^{\frac{1}{2}}}, ||x|w||_{\dot{H}^{\frac{1}{2}}})_{(\frac{1}{2},1)}.$$
 (3.10)

By Lemma 3.2 the right-hand side is bounded by $\|w\|_B^*$ and (3.1) is proved. The other stated properties of $\|\cdot\|_B^*$ are immediate.

The previous proof required two technical properties which we now establish. They are special cases of more general statements, but we limit ourselves to what is needed here.

Lemma 3.2. The following two properties hold:

- For any Schwartz function f in \mathbb{R}^2 one has $\|\hat{x}f\|_{\dot{H}^{\frac{1}{2}}} \lesssim \|f\|_{\dot{H}^{\frac{1}{2}}}$ where $\hat{x} = x/|x|$.
- With $\|\cdot\|_B^*$ defined as, cf. (3.1)

$$\sum_{k \in \mathbb{Z}} 2^{\frac{k}{2}} \| \psi(2^{-k} x) w(x) \|_{\dot{H}^{\frac{1}{2}}(\mathbb{R}^2)} =: \| w \|_{B}^{*}$$

one has

$$(\|w\|_{\dot{H}^{\frac{1}{2}}}, \||x|w\|_{\dot{H}^{\frac{1}{2}}})_{(\frac{1}{2},1)} \lesssim \|w\|_{B}^{*}.$$

Proof. The first property cannot simply be obtained by interpolating between the obvious L^2 property and the corresponding \dot{H}^1 inequality. Indeed, the latter would require Hardy's inequality in \mathbb{R}^2 with an r^{-1} weight, which fails. So we proceed differently. Using polar coordinates and complex notation we expand f into a Fourier series:

$$f(re^{i\theta}) = \sum_{n \in \mathbb{Z}} a_n(r)e^{2\pi i n\theta}, \quad a_n(r) = \int_0^1 f(re^{2\pi i \theta})e^{-2\pi i n\theta} d\theta.$$

By Plancherel

$$||f||_2^2 = \operatorname{const} \cdot \sum_{n \in \mathbb{Z}} \int_0^\infty |a_n(r)|^2 r \, dr$$
 (3.11)

and

$$||f||_{\dot{H}^{1}}^{2} = ||\partial_{r} f||_{2}^{2} + ||r^{-1}\partial_{\theta} f||_{2}^{2}$$

$$= \operatorname{const} \cdot \sum_{n \in \mathbb{Z}} \int_{0}^{\infty} (|a'_{n}(r)|^{2} + \frac{n^{2}}{r^{2}} |a_{n}(r)|^{2}) r \, dr.$$
(3.12)

By interpolation,

$$||f||_{\dot{H}^{\frac{1}{2}}}^{2} \simeq \sum_{n \in \mathbb{Z}} \int_{0}^{\infty} (|(-\partial_{r}^{2})^{\frac{1}{4}} a_{n}(r)|^{2} + \frac{|n|}{r} |a_{n}(r)|^{2}) r \, dr. \tag{3.13}$$

Since $\hat{x} = e^{2\pi i \theta} =: e(\theta)$ we conclude that

$$\|e(\theta)f\|_{\dot{H}^{\frac{1}{2}}}^{2} \lesssim \sum_{n \in \mathbb{Z}} \int_{0}^{\infty} (|(-\partial_{r}^{2})^{\frac{1}{4}} a_{n}(r)|^{2} + \frac{|n+1|}{r} |a_{n}(r)|^{2}) r \, dr$$

$$\lesssim \|f\|_{\dot{H}^{\frac{1}{2}}}^{2} + \int_{0}^{\infty} \frac{|a_{0}(r)|^{2}}{r} r \, dr.$$
(3.14)

Since

$$a_0(r) = \int_0^1 f(re(\theta)) d\theta,$$

the final term in (3.14) is

$$\int_{0}^{\infty} \frac{|a_{0}(r)|^{2}}{r} r \, dr \leq \int_{0}^{\infty} \int_{0}^{1} \frac{|f(re(\theta))|^{2}}{r} r \, d\theta dr$$

$$\lesssim \|r^{-\frac{1}{2}} f\|_{2}^{2}$$

$$\lesssim \|f\|_{\dot{H}^{\frac{1}{2}}}^{2}$$
(3.15)

by Hardy, and the first claim is proved.

To prove the second claim we first dominate the weighted norm via a smooth Littlewood–Paley partition of unity, viz.

$$\||x|w\|_{\dot{H}^{\frac{1}{2}}} \lesssim \sum_{k \in \mathbb{Z}} \|\psi(2^{-k}x)|x|w\|_{\dot{H}^{\frac{1}{2}}} \lesssim \sum_{k \in \mathbb{Z}} 2^{k} \|\psi(2^{-k}x)w\|_{\dot{H}^{\frac{1}{2}}}.$$
 (3.16)

For the final inequality it suffices to verify the case k=0 by scaling. Then, by the fractional Leibnitz rule and with $\tilde{\psi}\psi=\psi$ another Littlewood–Paley function,

$$\|\psi(x)|x|w\|_{\dot{H}^{\frac{1}{2}}} \lesssim \||x|\widetilde{\psi}(x)\|_{\infty} \|\psi(x)w\|_{\dot{H}^{\frac{1}{2}}} + \||\nabla|^{\frac{1}{2}}|x|\widetilde{\psi}(x)\|_{L^{4}} \|\psi(x)w\|_{L^{4}}$$

$$\lesssim \|\psi(x)w\|_{\dot{H}^{\frac{1}{2}}},$$
(3.17)

where the final step is obtained by Sobolev embedding. Clearly,

$$\|w\|_{\dot{H}^{\frac{1}{2}}} \lesssim \sum_{k \in \mathbb{Z}} \|\psi(2^{-k}x)w\|_{\dot{H}^{\frac{1}{2}}}.$$
 (3.18)

By the real interpolation property, see [3, Section 2], [4]

$$(\|w\|_{\dot{H}^{\frac{1}{2}}}, \||x|w\|_{\dot{H}^{\frac{1}{2}}})_{(\frac{1}{2},1)} \lesssim \left(\sum_{k \in \mathbb{Z}} \|\psi(2^{-k}x)w\|_{\dot{H}^{\frac{1}{2}}}, \sum_{k \in \mathbb{Z}} 2^{k} \|\psi(2^{-k}x)w\|_{\dot{H}^{\frac{1}{2}}}\right)_{(\frac{1}{2},1)}$$

$$\simeq \sum_{k \in \mathbb{Z}} 2^{\frac{k}{2}} \|\psi(2^{-k}x)w\|_{\dot{H}^{\frac{1}{2}}} = \|w\|_{\mathcal{B}}^{*}$$

$$(3.19)$$

and we are done. \Box

4. The convolution algebra and the proof of Theorem 1.2

We now present the algebra formalism in the scaling invariant setting.

Definition 4.1. The Banach space Z of tempered distributions is defined as

$$Z := \{ T(x_0, x_1, y) \in \mathcal{S}'(\mathbb{R}^9) \mid \mathcal{F}_y T(x_0, x_1, \eta) \in L_\eta^\infty L_{x_1}^\infty L_{x_0}^\infty \}$$
 (4.1)

with norm

$$||T||_{Z} := \sup_{\eta \in \mathbb{R}^{3}} ||\mathcal{F}_{y}T(x_{0}, x_{1}, \eta)||_{L_{x_{1}}^{\infty} L_{x_{0}}^{1}}$$
(4.2)

sup being the essential supremum. We add the identity I to Z, which corresponds to the kernel $T = \delta_0(y)\delta_0(x_1 - x_0)$. The convolution \circledast on $T_1, T_2 \in Z$ is defined by

$$(T_1 \circledast T_2)(x_0, x_2, y) = \mathcal{F}_{\eta}^{-1} \left[\int_{\mathbb{R}^3} \mathcal{F}_y T_1(x_0, x_1, \eta) \mathcal{F}_y T_2(x_1, x_2, \eta) \, dx_1 \right] (y). \tag{4.3}$$

Lemma 4.1. Let Z is a Banach algebra under \circledast with identity element I. If $V \in L^{3/2,1}$ then T_{1+}^{ε} defined by (2.5) belongs to Z and $\mathfrak{F}_y T_{1+}^{\varepsilon}$ is given by

$$\mathcal{F}_{\nu} T_{1+}^{\varepsilon}(x_0, x_1, \eta) = e^{-ix_1\eta} R_0(|\eta|^2 - i\varepsilon)(x_0, x_1) V(x_0) e^{ix_0\eta}. \tag{4.4}$$

Moreover,

$$\sup_{\varepsilon>0} \|T_{1+}^{\varepsilon}\|_{Z} \lesssim \|V\|_{L^{3/2,1}} \lesssim \|V\|_{\dot{B}^{\frac{1}{2}}}.$$
 (4.5)

If, in addition, $||V||_{L^{3/2,1}}$ is sufficiently small, then T^{ε}_{+} also belongs to Z and

$$(I + T_{1+}^{\varepsilon}) \circledast (I - T_{+}^{\varepsilon}) = (I - T_{+}^{\varepsilon}) \circledast (I + T_{1+}^{\varepsilon}) = I. \tag{4.6}$$

Proof. Z is a Banach space. The expressions in (4.3) appearing in brackets satisfies

$$\sup_{\eta \in \mathbb{R}^{3}} \left\| \int_{\mathbb{R}^{3}} \mathcal{F}_{y} T_{1}(x_{0}, x_{1}, \eta) \mathcal{F}_{y} T_{2}(x_{1}, x_{2}, \eta) dx_{1} \right\|_{L_{x_{2}}^{\infty} L_{x_{0}}^{1}}$$

$$\leq \left\| \mathcal{F}_{y} T_{1} \right\|_{L_{\eta}^{\infty} L_{x_{1}}^{\infty} L_{x_{0}}^{1}} \left\| \mathcal{F}_{y} T_{2} \right\|_{L_{\eta}^{\infty} L_{x_{2}}^{\infty} L_{x_{1}}^{1}}$$

$$= \left\| T_{1} \right\|_{Z} \left\| T_{2} \right\|_{Z}$$

$$(4.7)$$

and so it is a tempered distribution in \mathbb{R}^9 . Therefore, the composition (4.3) is well-defined in Z and

$$||T_1 \circledast T_2||_Z \le ||T_1||_Z ||T_2||_Z$$

whence Z is a Banach algebra under $\|\cdot\|_Z$.

Formula (4.4) follows from (2.5) by taking Fourier transforms. By the resolvent identity

$$(I + R_0(z)V)^{-1} = I - R_V(z)V, \quad R_V(z) = (I + R_0(z)V)^{-1}R_0(z) \tag{4.8}$$

for Im z > 0. Here $R_V(z) = (H - z)^{-1}$ which exists for Im z > 0 since H is self-adjoint. For V small in $L^{\frac{3}{2},1}$ it follows that

$$\sup_{\text{Im } z > 0} \|R_0(z)Vf\| \le C\|V\|_{L^{\frac{3}{2},1}} \|f\|_{\infty} \le \frac{1}{2} \|f\|_{\infty}.$$

Hence $(I + R_0(z)V)^{-1}$ exists as a bounded operator on L^{∞} uniformly in Im z > 0, and we may also take the limit Im $z \ge 0$. In particular,

$$R_V(z): L^{\frac{3}{2},1}(\mathbb{R}^3) \longrightarrow L^{\infty}(\mathbb{R}^3)$$

From (4.8),

$$R_0(|\eta|^2 - i\varepsilon)V - R_V(|\eta|^2 - i\varepsilon)V + R_0(|\eta|^2 - i\varepsilon)VR_V(|\eta|^2 - i\varepsilon)V = 0 \quad (4.9)$$

whence, with $e^{ix\eta} f(x) =: (M_{\eta} f)(x)$,

$$M_{\eta}^{-1} R_{0}(|\eta|^{2} - i\varepsilon)(x_{0}, x_{1})V(x_{0}) M_{\eta} - M_{\eta}^{-1} R_{V}(|\eta|^{2} - i\varepsilon)(x_{0}, x_{1})V(x_{0}) M_{\eta}$$

$$+ M_{\eta}^{-1} R_{0}(|\eta|^{2} - i\varepsilon)(x_{2}, x_{1})V(x_{2}) M_{\eta} \circ M_{\eta}^{-1} R_{V}(|\eta|^{2} - i\varepsilon)(x_{0}, x_{2})V(x_{0}) M_{\eta}$$

$$= 0$$

$$(4.10)$$

where \circ signifies integration. In view of (4.4) this is tantamount to

$$0 = T_{1+}^{\varepsilon} - T_{+}^{\varepsilon} + T_{1+}^{\varepsilon} \circledast T_{+}^{\varepsilon} \tag{4.11}$$

or $(I + T_{1+}^{\varepsilon}) \otimes (I - T_{+}^{\varepsilon}) = I$. The second identity in (4.6) is valid since the resolvent identity also implies (4.9) with R_0 and R_V reversed:

$$R_0(|\eta|^2 - i\varepsilon)V - R_V(|\eta|^2 - i\varepsilon)V + R_V(|\eta|^2 - i\varepsilon)VR_0(|\eta|^2 - i\varepsilon)V = 0 \quad (4.12)$$

and so that same argument as before concludes the proof. \Box

The following spaces play a key role in the proof of Theorem 1.2. The Y-space in particular allows us to inductively bound the structure function of each W_{n+}^{ε} .

Definition 4.2. Let \mathcal{B} be the closure of the Schwartz functions in \mathbb{R}^3 under the norm $\|\cdot\|$. Fix any measurable function $v:\mathbb{R}^3\to\mathbb{R}$ which does not vanish a.e., and so that $\|v\|_B<\infty$. We introduce the following structures depending on v.

• The seminormed space

$$v^{-1}\mathcal{B} = \{ f \text{ measurable } | v(x)f(x) \in \mathcal{B} \}$$

with the seminorm $||f||_{v^{-1}\mathcal{B}} := ||vf|||$.

• Let *X* the space of two-variable kernels

$$X := \left\{ \mathfrak{X} \in \mathcal{B}(L^{\infty}, L^{\infty}) \mid (\mathfrak{X}f)(x) = \int_{\mathbb{R}^{3}} \mathfrak{X}(x, y) f(x - y) \, dy, \quad f \in L^{\infty} \right.$$

$$\left\| \mathfrak{X}(x, y) \right\|_{L_{x}^{\infty} L_{y}^{1}} < \infty, \quad \left\| \mathfrak{X}(x, y) \right\|_{L_{y}^{1} v^{-1} \mathcal{B}_{x}} < \infty \right\},$$

$$(4.13)$$

with norm (the first v factor is only for homogeneity)

$$\|\mathfrak{X}\|_{X} := \|v\|_{B} \|\mathfrak{X}\|_{L_{x}^{\infty} L_{y}^{1}} + \int_{\mathbb{R}^{3}} \|v(x)\mathfrak{X}(x, y)\|_{\mathcal{B}_{x}} dy.$$
(4.14)

• Let *Y* be the space of three-variable kernels

$$Y := \left\{ T(x_0, x_1, y) \in Z \mid \text{ for all } f \in L^{\infty} \cap v^{-1} \mathcal{B} \right.$$

$$(fT)(x, y) := \int_{\mathbb{R}^3} f(x_0) T(x_0, x_1, y) \, dx_0 \in X_{x_1, y} \right\},$$
(4.15)

with norm

$$||T||_{Y} := ||T||_{Z} + ||T||_{B(v^{-1}\mathcal{B}_{x_0}, X_{x_1, y})}$$
(4.16)

We adjoin an identity element to Y, in the form of

$$I(x_0, x_1, y) = \delta_{x_0}(x_1)\delta_0(y) = \delta_{x_1}(x_0)\delta_0(y). \tag{4.17}$$

Notice that in (4.14) we use the stronger norm $||v||_B$ rather than ||v||. The presence of $||T||_Z$ in (4.16) will require us to ensure that $||V||_B < \infty$ as well as $V \in \dot{B}^{\frac{1}{2}}$.

Lemma 4.2. Let V be a Schwartz function, and let K_{1+}^{ε} be defined in terms of V. Then uniformly in $\varepsilon > 0$,

$$||K_{1+}^{\varepsilon}(x,y)||_{L_{x}^{\infty}L_{y}^{1}} \lesssim |||V|||,$$
 (4.18)

$$||v(x)K_{1+}^{\varepsilon}(x,y)||_{L^{1}_{u}\mathcal{B}_{x}} \lesssim ||v||_{B} ||V||_{H},$$
 (4.19)

for any $v \in B^1$. With f a Schwartz function, define a kernel

$$\widetilde{K}_{1+}^{\varepsilon}(x,y) = \int_{\mathbb{R}^3} f(x_0) T_{1+}^{\varepsilon}(x_0, x, y) \, dx_0 \tag{4.20}$$

with the integral being understood as distributional duality pairing. Then uniformly in $\varepsilon > 0$,

$$\|\widetilde{K}_{1+}^{\varepsilon}(x,y)\|_{L_{y}^{\infty}L_{y}^{1}} \lesssim \|fV\|, \tag{4.21}$$

$$\|v(x)\widetilde{K}_{1+}^{\varepsilon}(x,y)\|_{L^{1}_{0}\mathcal{B}_{x}} \lesssim \|v\|_{B} \|fV\|,$$
 (4.22)

for any $v \in B^1$.

Proof. From (2.12) one has for all $\varepsilon > 0$,

$$||K_{1+}^{\varepsilon}(x,y)||_{L_{x}^{\infty}L_{y}^{1}} \leq \operatorname{const} \cdot \int_{-\infty}^{\infty} \int_{\mathbb{S}^{2}} |L(t,\omega)| \, dt \, d\omega = \operatorname{const} \cdot ||V|| \tag{4.23}$$

which is (4.18). For the second estimate (4.19) we invoke Lemma 2.4, viz.

$$\|v(x)K_{1+}^{\varepsilon}(x,y)\|_{L_{y}^{1}\mathcal{B}_{x}} = \operatorname{const} \cdot \int_{\mathbb{R}^{3}} \|v(x)|u|^{-2}L(|u| - 2\hat{u} \cdot x, \hat{u})\| du$$

$$= \operatorname{const} \cdot \int_{\mathbb{S}^{2}} \int_{0}^{\infty} \|v(x)L(t - 2\omega \cdot x, \omega)\| dt d\omega \qquad (4.24)$$

$$\lesssim \|v\|_{B} \|V\| \qquad (4.25)$$

as claimed.

Next,

$$\mathcal{F}_{x,y}\widetilde{K}_{1+}^{\varepsilon}(\xi_{1},\eta) = \int_{\mathbb{R}^{3}} \hat{f}(\xi_{0})\mathcal{F}_{x_{0}}^{-1}\mathcal{F}_{x,y}T_{1+}^{\varepsilon}(\xi_{0},\xi_{1},\eta) d\xi_{0}
= \int_{\mathbb{R}^{3}} \frac{\hat{f}(\xi_{0})\hat{V}(\xi_{1}-\xi_{0})}{|\xi_{1}+\eta|^{2}-|\eta|^{2}+i\varepsilon} d\xi_{0} = \frac{\widehat{fV}(\xi_{1})}{|\xi_{1}+\eta|^{2}-|\eta|^{2}+i\varepsilon}.$$
(4.26)

In view of (2.5), this leads to the kernel K_{1+}^{ε} associated with the potential fV. \square

Next, we define the operation of *contraction*:

Lemma 4.3. For $\mathfrak{X} \in X$, the contraction of $T \in Y$ by \mathfrak{X} is

$$(\mathfrak{X}T)(x,y) := \int_{\mathbb{R}^6} \mathfrak{X}(x_0, y_0) T(x_0, x, y - y_0) \, dx_0 \, dy_0. \tag{4.27}$$

Then $\mathfrak{X}T \in X_{x,y}$ and $\|\mathfrak{X}T\|_X \leq \|T\|_Y \|\mathfrak{X}\|_X$. We interpret the right-hand side of (4.27) relative to the Fourier variable:

$$\mathcal{F}_{\eta}^{-1} \left[\int_{\mathbb{R}^3} \mathcal{F}_{y_0} \mathfrak{X}(x_0, \eta) \mathcal{F}_{y_0} T(x_0, x, \eta) \, dx_0 \right] (y). \tag{4.28}$$

The integral is absolutely convergent and the inverse Fourier transform relative to η is a tempered distribution.

Proof. We have $\mathcal{F}_{y_0}\mathfrak{X}(x_0,\eta) \in L^{\infty}_{x_0,\eta}$ and $\mathcal{F}_{y_0}T(x_0,x,\eta) \in L^{\infty}_x L^1_{x_0}$, whence the claim about the integral in brackets. The estimate $\|\mathfrak{X}T\|_X \leq \|T\|_Y \|\mathfrak{X}\|_X$ follows from the definition of the space Y:

$$\left\| \int_{\mathbb{R}^{6}} \mathfrak{X}(x_{0}, y_{0}) T(x_{0}, x, y - y_{0}) dx_{0} dy_{0} \right\|_{X_{x,y}}$$

$$\leq \int_{\mathbb{R}^{3}} \|(\mathfrak{X}(\cdot, y_{0}) T)(x, y - y_{0})\|_{X_{x,y}} dy_{0}$$

$$= \int_{\mathbb{R}^{3}} \|(\mathfrak{X}(\cdot, y_{0}) T)(x, y)\|_{X_{x,y}} dy_{0}$$

$$\lesssim \|T\|_{Y} \int_{\mathbb{R}^{3}} \|\mathfrak{X}(\cdot, y_{0})\|_{V^{-1}\mathcal{B}_{X}} dy_{0} \leq \|T\|_{Y} \|\mathfrak{X}\|_{X}$$

and we are done.

The previous lemma allows us to prove that Y is a Banach algebra under the composition \circledast . This will allow us to prove the key property that $T_{n+}^{\varepsilon} \in Y$ starting from the case $T_{1+} \in Y$, which we now state.

Lemma 4.4. *Y defined by* (4.15) *is a Banach algebra with the operation* \otimes *defined in the ambient algebra Z.*

Proof. The fact that \circledast is associative (and non-commutative) is clear in Z, and the unit element is given by (4.17). Since $Y \subset Z$, the same is true in Y.

The definitions of X and Y imply that each contraction $\mathfrak{X}T$ (see (4.27)) is in X and $\|\mathfrak{X}T\|_X \lesssim \|\mathfrak{X}\|_X \|T\|_Y$. We have

$$\int_{\mathbb{R}^{3}} f(x_{0})T_{3}(x_{0}, x_{2}, y) dx_{0}$$

$$= \int_{\mathbb{R}^{9}} f(x_{0})T_{1}(x_{0}, x_{1}, y_{1})T_{2}(x_{1}, x_{2}, y - y_{1}) dx_{1} dy_{1} dx_{0}.$$
(4.29)

As in the case of (4.27), the *y*-integral is to be understood in the distributional Fourier sense. Integrating in x_0 , we obtain an expression of the form $\mathfrak{X}T_2$ for $\mathfrak{X} \in X$ with $\|\mathfrak{X}\|_X \lesssim \|f\|_{V^{-1}B}\|T_1\|_Y$. Then $\mathfrak{X}T_2$ belongs to X as stated above and has a norm at most $\lesssim \|f\|_{V^{-1}B}\|T_1\|_Y\|T_2\|_Y$. Thus, $T_3 = T_1 \circledast T_2 \in Y$ and

$$||T_1 \circledast T_2||_Y \le C ||T_1||_Y ||T_2||_Y$$

with some absolute constant C. Multiplying the norm by C removes this constant from the previous inequality, and so Y is an algebra under this new norm. \Box

Corollary 4.5. Let V be Schwartz and apply Definition 4.2 with v = V, the potential. Then for every $\varepsilon > 0$ we have $T_{1+}^{\varepsilon} \in Y$ and

$$\sup_{\varepsilon>0} \|T_{1+}^{\varepsilon}\|_{Y} \lesssim \|V\|_{B} + \|V\|_{\dot{B}^{\frac{1}{2}}}.$$
(4.30)

Proof. By (4.5) we have

$$\sup_{\varepsilon>0} \|T_{1+}^{\varepsilon}\|_{Z} \lesssim \|V\|_{\dot{B}^{\frac{1}{2}}}.$$

It remains to show that

$$\sup_{\varepsilon>0} \left\| \int_{\mathbb{R}^3} f(x_0) T_{1+}^{\varepsilon}(x_0, x, y) \, dx_0 \right\|_{X_{x,y}} \lesssim \|V\|_B \| f V \|. \tag{4.31}$$

In view of (4.14) this is implied by Lemma 4.2.

We are now in a position to obtain the key representation result concerning the partial wave operators W_{n+} , see (2.2). In what follows, we let B_* be the space obtained as the closure of the Schwartz functions under the norm

$$\|\cdot\|_{B_*} := \|\cdot\|_B + \|\cdot\|_{\dot{B}^{\frac{1}{2}}},$$

see (3.1). We define B_* as the space obtained as the closure of Schwartz functions under the norm $\|\cdot\|_{B_*}$.

Proposition 4.6. Let V be a Schwartz potential. Then $T_{n+}^{\varepsilon} \in Y$ for any $n \ge 1$ and $\varepsilon > 0$ and

$$\sup_{\varepsilon>0} \|T_{n+}^{\varepsilon}\|_{Y} \le C^{n} \|V\|_{B_{*}}^{n} \tag{4.32}$$

with some absolute constant C. Moreover, for all Schwartz functions f one has

$$(W_{n+}^{\varepsilon}f)(x) = \int_{\mathbb{S}^2} \int_{\mathbb{R}^3} g_n^{\varepsilon}(x, dy, \omega) f(S_{\omega}x - y) \, \sigma(d\omega) \tag{4.33}$$

where for fixed $x \in \mathbb{R}^3$, $\omega \in \mathbb{S}^2$ the expression $g_n^{\varepsilon}(x,\cdot,\omega)$ is a measure satisfying

$$\sup_{\varepsilon>0} \int_{\mathbb{S}^2} \|g_n^{\varepsilon}(x, dy, \omega)\|_{\mathcal{M}_y L_x^{\infty}} d\omega \le C^n \|V\|_{B_*}^n$$
(4.34)

where $\|\cdot\|_{\mathcal{M}}$ refers to the total variation norm of Borel measures. The same conclusion also holds if $V \in B_*$.

Proof. First, $T_{n+}^{\varepsilon} = T_{1+}^{\varepsilon} \otimes T_{(n-1)+}^{\varepsilon}$. Corollary 4.5 and the algebra property of *Y* imply (4.32) by induction. Second, we have

$$W_{n+}^{\varepsilon} = (-1)^{n} \mathbb{1}_{\mathbb{R}^{3}} T_{n+}^{\varepsilon} = (-1)^{n} \mathbb{1}_{\mathbb{R}^{3}} (T_{(n-1)+}^{\varepsilon} \circledast T_{1+}^{\varepsilon})$$

$$= -((-1)^{n-1} \mathbb{1}_{\mathbb{R}^{3}} T_{(n-1)+}^{\varepsilon}) T_{1+}^{\varepsilon} = -W_{(n-1)+}^{\varepsilon} T_{1+}^{\varepsilon}.$$
(4.35)

The notation in the second line contraction of a kernel in Y by an element of X; this follows again by induction starting from $W_{0+}^{\varepsilon} = \mathbb{1}_{\mathbb{R}^3}$ via (4.27). By the boundedness of T_{n+}^{ε} in Y it follows that the right-hand side of (4.35) is well-defined in Y. Thus, by the first equality sign in (4.35),

$$\sup_{\varepsilon>0} \|W_{n+}^{\varepsilon}\|_{X} \leq \|\mathbb{1}_{\mathbb{R}^{3}}\|_{V^{-1}\mathcal{B}} \sup_{\varepsilon>0} \|T_{n+}^{\varepsilon}\|_{Y} \leq C^{n+1} \|V\| \|V\|_{B_{*}}^{n} \\
\leq C^{n+1} \|V\|_{B_{*}}^{n+1}. \tag{4.36}$$

We denote the kernel of W_{1+}^{ε} by $\mathfrak{X}_{V}^{\varepsilon}$, where V is the potential. Thus,

$$\mathfrak{X}_V^{\varepsilon}(x,y) = -\int_{\mathbb{R}^3} T_{1+}^{\varepsilon}(x_0,x,y) \, dx_0 = -(\mathbb{1}_{\mathbb{R}^3} T_{1+}^{\varepsilon})(x,y) \in X.$$

By (4.35),

$$W_{n+}^{\varepsilon}(x,y) = -\int_{\mathbb{R}^{6}} W_{(n-1)+}^{\varepsilon}(x',y') T_{1+}^{\varepsilon}(x',x,y-y') \, dx' dy'$$

$$= \int_{\mathbb{R}^{3}} \mathfrak{X}_{f_{y'}^{\varepsilon}V}^{\varepsilon}(x,y-y') \, dy'.$$
(4.37)

Here we wrote $f_{v'}^{\varepsilon}(x') = W_{(n-1)+}^{\varepsilon}(x', y')$ and we used (4.26).

We now invoke the representation from Corollary 2.2. Specifically, by (2.13) there exists $g_{1,f_{\varepsilon}}^{\varepsilon}(x,dy,\omega)$ so that for every $\phi \in \mathcal{S}$ one has

$$(\mathfrak{X}^{\varepsilon}_{f^{\varepsilon}_{y'}V}\phi)(x) = \int\limits_{\mathbb{S}^{2}}\int\limits_{\mathbb{R}^{3}}g^{\varepsilon}_{1,f^{\varepsilon}_{y'}}(x,dy,\omega)\phi(S_{\omega}x - y)\,\sigma(d\omega)$$

where for fixed $x \in \mathbb{R}^3$, $\omega \in \mathbb{S}^2$ the expression $g_{1,f_{y'}^{\varepsilon}}^{\varepsilon}(x,\cdot,\omega)$ is a measure satisfying

$$\begin{split} \sup_{\varepsilon>0} \int\limits_{\mathbb{S}^2} \|g_{1,f_{y'}^\varepsilon}^\varepsilon(x,dy,\omega)\|_{\mathcal{M}_y L^\infty_x} \, d\omega &\leq C \, \|f_{y'}^\varepsilon V\| \\ &= C \, \|W_{(n-1)+}^\varepsilon(x',y')\|_{V^{-1}\mathcal{B}_{x'}}. \end{split}$$

Therefore, with $\phi_{y'}(\cdot) := \phi(\cdot - y')$,

$$(W_{n+}^{\varepsilon}\phi)(x) = \int_{\mathbb{R}^{3}} W_{n+}^{\varepsilon}(x,y)\phi(x-y) \, dy$$

$$= \int_{\mathbb{R}^{6}} \mathfrak{X}_{f_{y'}^{\varepsilon}V}^{\varepsilon}(x,y-y')\phi(x-y) \, dy \, dy'$$

$$= \int_{\mathbb{R}^{6}} \mathfrak{X}_{f_{y'}^{\varepsilon}V}^{\varepsilon}(x,y)\phi_{y'}(x-y) \, dy \, dy'$$

$$= \int_{\mathbb{R}^{3}} (\mathfrak{X}_{f_{y'}^{\varepsilon}V}^{\varepsilon}\phi_{y'})(x) \, dy'$$

$$= \int_{\mathbb{R}^{3}} \int_{\mathbb{S}^{2}} \int_{\mathbb{R}^{3}} g_{1,f_{y'}^{\varepsilon}}^{\varepsilon}(x,dy,\omega)\phi(S_{\omega}x-y-y') \, \sigma(d\omega) \, dy'$$

$$= \int_{\mathbb{R}^{2}} \int_{\mathbb{R}^{3}} \left[\int_{\mathbb{R}^{3}} g_{1,f_{y'}^{\varepsilon}}^{\varepsilon}(x,d(y-y'),\omega) \, dy' \right] \phi(S_{\omega}x-y) \, \sigma(d\omega).$$
(4.38)

The expressions in brackets is the structure function

$$g_n(x, dy, \omega) := \int_{\mathbb{R}^3} g_{1, f_{y'}^{\varepsilon}}^{\varepsilon}(x, d(y - y'), \omega) \, dy'. \tag{4.39}$$

In fact, it is a measure in the y-coordinate and

$$(W_{n+}^{\varepsilon}\phi)(x) = \int_{\mathbb{S}^2} \int_{\mathbb{R}^3} g_n(x, dy, \omega)\phi(S_{\omega}x - y)\,\sigma(d\omega). \tag{4.40}$$

Moreover, we have the bounds, uniformly in $\varepsilon > 0$

$$\begin{split} &\int_{\mathbb{S}^{2}} \|g_{n}^{\varepsilon}(x,dy,\omega)\|_{\mathcal{M}_{y}L_{x}^{\infty}} d\omega \\ &= \int\int_{\mathbb{S}^{2}} \|g_{1,f_{y'}^{\varepsilon}}^{\varepsilon}(x,d(y-y'),\omega)\|_{\mathcal{M}_{y}L_{x}^{\infty}} dy'd\omega \\ &= \int\int_{\mathbb{S}^{2}} \|g_{1,f_{y'}^{\varepsilon}}^{\varepsilon}(x,dy,\omega)\|_{\mathcal{M}_{y}L_{x}^{\infty}} dy'd\omega \\ &\leq C \int\limits_{\mathbb{R}^{3}} \|W_{(n-1)+}^{\varepsilon}(x',y')\|_{V^{-1}\mathcal{B}_{x'}} dy' \\ &= C \|W_{(n-1)+}^{\varepsilon}(x',y')\|_{L_{y'}^{1}V^{-1}\mathcal{B}_{x'}} \\ &\leq C \|W_{(n-1)+}^{\varepsilon}\|_{X} \\ &\leq C^{n} \|V\|_{B_{*}}^{n} \end{split}$$

by (4.36). This concludes the argument under the assumption that $f_{y'}^{\varepsilon}(x')$ is a Schwartz function. To remove this assumption, we can make

$$\|W_{(n-1)+}^{\varepsilon}(x',y')-\tilde{f}_{y'}^{\varepsilon}(x')\|_X$$

arbitrarily small with a Schwartz function $\tilde{f}^{\varepsilon}_{y'}(x')$ in \mathbb{R}^6 . Then the previous calculation shows that

$$\int_{\mathbb{S}^2} \|g_n^{\varepsilon}(x, dy, \omega) - \tilde{g}_n^{\varepsilon}(x, dy, \omega)\|_{\mathcal{M}_y L_x^{\infty}} d\omega$$

can be made as small as we wish where $\tilde{g}_n^{\varepsilon}(x, dy, \omega)$ is the function generated by $\tilde{f}_{v'}^{\varepsilon}(x')$. Passing to the limit concludes the proof.

To remove the assumption that V be a Schwartz function, we approximate $V \in B_*$ by Schwartz functions in the norm $\|\cdot\|_{B_*}$. We achieve convergence of of the functions g_n by means of (4.34) and of the kernels W_{n+}^{ε} themselves by means of (4.36). To be specific, denoting by $\widetilde{W}_{n+}^{\varepsilon}$ and \widetilde{g}_n the quantities corresponding to the potential \widetilde{V} , taking differences yields

$$\|\widetilde{W}_{n+}^{\varepsilon} - W_{n+}^{\varepsilon}\|_{X} + \int_{\mathbb{S}^{2}} \|g_{n}^{\varepsilon}(x, dy, \omega) - \tilde{g}_{n}^{\varepsilon}(x, dy, \omega)\|_{\mathcal{M}_{y}L_{x}^{\infty}} d\omega$$

$$\leq C^{n} \|V - \tilde{V}\|_{B_{*}} (\|V\|_{B_{*}}^{n-1} + \|\tilde{V}\|_{B_{*}}^{n-1})$$

uniformly in $\varepsilon > 0$.

To prove Theorem 1.2 we now simply sum the series $\sum_{n=1}^{\infty} g_n$ which can be done in view of the previous proposition, provided c_0 is sufficiently small.

References

- [1] M. Beceanu, New estimates for a time-dependent Schrödinger equation. *Duke Math. J.* **159** (2011), no. 3, 417–477. MR 2831875 Zbl 1229.35224
- [2] M. Beceanu and M. Goldberg, Schrödinger dispersive estimates for a scaling-critical class of potentials. *Comm. Math. Phys.* 314 (2012), no. 2, 471–481. MR 2958960 Zbl 1250.35047
- [3] M. Beceanu and W. Schlag, Structure formulas for wave operators. Preprint, 2016. arXiv:1612.07304 [math.AP]
- [4] J. Bergh and J. Löfström, *Interpolation spaces. An introduction*. Grundlehren der Mathematischen Wissenschaften, 223. Springer-Verlag, Berlin etc., 1976. MR 0482275 Zbl 0344.46071
- [5] T. Kato, Wave operators and similarity for some non-selfadjoint operators. *Math. Ann.* **162** (1966) 258–279. MR 0190801 Zbl 0139.31203
- [6] K. Yajima, The $W^{k,p}$ -continuity of wave operators for Schrödinger operators. *Proc. Japan Acad. Ser. A Math. Sci.* **69** (1993), no. 4, 94–98. MR 1222831 Zbl 0810.35100
- [7] Y, Yajima, The $W^{k,p}$ -continuity of wave operators for Schrödinger operators. textitJ. Math. Soc. Japan 47 (1995), no. 3, 551–581. MR 1331331 Zbl 0837.35039

Received May 4, 2017

Marius Beceanu, Department of Mathematics, University at Albany, State University of New York, 1400 Washington Avenue, Albany, NY 12222, USA e-mail: mbeceanu@albany.edu

Wilhelm Schlag, Department of Mathematics, Yale University, 10 Hillhouse Avenue, New Haven, CT 06511, USA

e-mail: wilhelm.schlag@yale.edu